## Lagrangian formalism and Noether-type theorems for second-order delay ODEs

Vladimir Dorodnitsyn<sup>*a*</sup>, Roman Kozlov<sup>*b*</sup>, Sergey Meleshko<sup>*c*</sup>

#### 10.02.2023

<sup>a</sup> Keldysh Institute of Applied Mathematics, Russian Academy of Science, Miusskaya Pl. 4, Moscow, 125047, Russia;
e-mail: Dorodnitsyn@Keldysh.ru, dorod2007@gmail.com
<sup>b</sup> Department of Business and Management Science, Norwegian School of Economics, Helleveien 30, 5045, Bergen, Norway;
e-mail: Roman.Kozlov@nhh.no
<sup>c</sup> School of Mathematics, Institute of Science, Suranaree University of Technology, 30000, Thailand;

e-mail: sergey@math.sut.ac.th

#### Abstract

The Lagrangian formalism for variational problem for second-order delay ordinary differential equations (DODEs) is developed. The Noether-type operator identities and theorems for DODEs of second order are presented. Algebraic construction of integrals for DODEs based on symmetries are demonstrated by examples.

### 1 Introduction

Lie group analysis is efficient tool for studying ordinary and partial differential equations since its introduction in the classical work of Sophus Lie in [1, 2]. A symmetry transformation of a differential equation maps a solution into a solution. This property is used to obtain new solutions from known ones, and to classify equations into equivalence classes. It can also be used to obtain exact analytic solutions that are invariant under some subgroup of the symmetry group ('group invariant solutions'). Applications of Lie groups to differential equations is the topic of many books and articles [3–8]. Since the fundamental work of E.Noether [9] the symmetry group becomes a starting point to obtain first integrals and conservation laws for differential equations, which possess Lagrangian or Hamiltonian formulation. Recently, the connections between a symmetry group and conservation laws for differential equations that do not have a variational formulation (and hence they have neither a Lagrangian nor a Hamiltonian) were developed in [5, 10].

More recently applications of Lie groups of transformations have been extended to finitedifference, difference, discrete equations and integral equations [11–29]. This article is part of a study that aims to extend group analysis application to another type of equations: delay differential equations. In the recent article [30] there was presented the Lie group classification of first-order delay ordinary differential equations. Linear first-order delay ordinary differential equations were considered in article [31]. Lie group classification of delay second-order ordinary differential equations was studied in [32]. Exact solutions for delay PDEs were constructed in [33–35].

The purpose of this article is to construct Lagrangian formalism and the Noether-type theorems for second-order delay ordinary differential equations.

As shown below, the variational problem leads to an ODE with a delay of second-order (in the sense of the order of a differential equation) with two delays. Thus, we have to set an initial-value problem for delay ODEs with *two* delays.

We consider the formulation of the initial-value problem for a second-order ordinary delay differential equation with two constant delays  $\tau$ 

$$\ddot{u} = F(\ddot{u}^{--}, \ddot{u}^{-}, \dot{u}^{-}, \dot{u}, \dot{u}^{--}, u^{-}, u, u^{--}, t, t^{-}, t^{--}), \qquad x \in I,$$
(1.1)

where  $I \subset \mathbb{R}$  is some finite or semifinite interval,  $t^- = t - \tau$ ,  $t^{--} = t - 2\tau$ . Functions  $u^- = u(t - \tau), u^{--} = u(t - 2\tau)$ , while function F satisfies the condition

$$\left(\frac{\partial F}{\partial u^{--}}\right)^2 + \left(\frac{\partial F}{\partial \dot{u}^{--}}\right)^2 + \left(\frac{\partial F}{\partial \ddot{u}^{--}}\right)^2 \neq 0.$$
(1. 2)

The variable t varies continuously over the entire interval, where Eq. (1, 1) is defined.

In order to solve DODEs on some interval I one must add initial conditions to the DODE (1. 1). Contrary to the case of ordinary differential equations, the initial condition must be given by a smooth function  $\varphi(t)$  and it's derivative  $\dot{\varphi}(t)$  on an initial interval  $I_0 \subset \mathbb{R}$ , e.g.

$$u(t) = \varphi(t), \qquad \dot{u}(t) = \dot{\varphi}(t), \qquad t \in [t_{-2}, t_0], \quad t_0 - t_{-2} = 2\tau, \quad t_0 - t_{-1} = \tau.$$
(1.3)

For a constant  $\tau$  this leads to the *method of steps* [36] for solving the DODE either analytically or numerically. Hence, for  $t_0 \leq t \leq t_1 = t_0 + \tau$  we replace

$$u^{--} = u(t^{--}) = \varphi(t - 2\tau), \quad \dot{u}^{--} = u(t - 2\tau) = \dot{\varphi}(t - 2\tau), \quad \ddot{u}^{--} = \ddot{u}(t - 2\tau) = \ddot{\varphi}(t - 2\tau),$$
$$u^{-} = u(t^{-}) = \varphi(t - \tau), \quad \dot{u}^{-} = u(t - \tau) = \dot{\varphi}(t - \tau), \quad \ddot{u}^{-} = \ddot{u}(t - \tau) = \ddot{\varphi}(t - \tau),$$

and this reduces DODE(1, 1) to the ODE

$$\ddot{u} = F(\ddot{\varphi}(t-2\tau), \dot{\varphi}(t-2\tau), \varphi(t-2\tau), \ddot{\varphi}(t-\tau), \dot{\varphi}(t-\tau), \varphi(t-\tau), u, \dot{u}, t, t^{-}, t^{--}), \quad (1.4)$$

which is solved with the initial conditions

$$u(t_0) = \varphi(t_0), \quad \dot{u}(t_0) = \dot{\varphi}(t_0).$$

On the second step we consider the same procedure: we solve the DODE (1. 1) where  $u(u - \tau), u(u - 2\tau)$ , it's derivatives and the initial conditions are known from the first step. Thus, one continues at each step solving the ODE with data from the previous step. This procedure provides a solution that is in general continuous at the points  $t_n = t_0 + n\tau$ .

The method of steps for solving the initial value problem introduces a natural sequence of intervals

$$[t_n, t_{n+1}], \qquad n = -1, 0, 1, 2, \dots \tag{1.5}$$

We should indicate that peculiarity of the initial value problem for DODEs with *two* delays is that the initial conditions should be given on a double length interval  $[t_{-2}, t_0]$ .

This paper is devoted to analytic technique of integration of a second-order DODE, based on its symmetry and Lagrangian formalism. The Noether-type theorems for invariant delay ordinary differential equations of a second-order are presented. The integration technique based on a symmetry of an DODE is demonstrated by examples.

The paper is organized as follows. In Section 2 we describe how to apply symmetry generators to delay ODEs. The analog of the first Noether's theorem of the invariance condition for delay functional is proved in Section 3. In Section 4 we show that the stationary value of delay functional along a given group orbit at an arbitrary point t, u is achieved on so called local extremal equation, special case of which is an equation named the Elsgolts equation. The invariance of the Elsgolts equation considered in Section 5. In section 6 we introduce the first integrals for DODE, which are sufficiently different from those for the ODE. The main operator identities, which allow us to formulate a Noether-type theorem for the conservation law and symmetries are given in section 7. In section 8 we demonstrate application of the developed methodology by examples. Short observation of the obtained results are considered in the Conclusion.

### 2 Point symmetries of delay ODEs

We consider a second-order delay ODE (1. 1) and its point symmetries defined by a generator in the standard form:

$$X = \xi(t, u)\frac{\partial}{\partial t} + \eta(t, u)\frac{\partial}{\partial u} + ..., \qquad (2. 1)$$

which we prolong for  $\dot{u}$ ,  $\ddot{u}$  using standard formulas, and for the shifted values  $(t^-, u^-, \dot{u}^-), (t^{--}, u^{--}, \dot{u}^{--})$ :

$$\begin{split} \tilde{X} &= \xi(t, u) \frac{\partial}{\partial t} + \eta(t, u) \frac{\partial}{\partial u} + \xi^-(t^-, u^-) \frac{\partial}{\partial t^-} + \eta^-(t^-, u^-) \frac{\partial}{\partial u^-} \\ &+ (D(\eta) - \dot{u}D(\xi)) \frac{\partial}{\partial \dot{u}} + (D^-(\eta^-) - \dot{u}^- D^-(\xi^-)) \frac{\partial}{\partial \dot{u}^-} + \dots \\ &+ (\xi(t, u) - \xi^-(t^-, u^-)) \frac{\partial}{\partial \tau} + \dots, \end{split}$$
(2. 2)

where  $\tau = t - t^{-} = t^{-} - t^{--} = const$ , and

$$D = \frac{\partial}{\partial t} + \dot{u}\frac{\partial}{\partial u} + \dots, \qquad D^- = \frac{\partial}{\partial t^-} + \dot{u}^-\frac{\partial}{\partial u^-} + \dots$$

are the operators of the total derivative at points t and  $t^-$ , respectively.

We also use the operators

$$D^{+} = \frac{\partial}{\partial t^{+}} + \dot{u}^{+} \frac{\partial}{\partial u^{+}} + \ddot{u}^{+} \frac{\partial}{\partial \dot{u}^{+}} + \dots,$$

and the following sum of operators

$$\bar{D} = D^- + D + D^+.$$

We split the operator of the total derivative into three different operators for preserving their point character. In this case they can be multiplied (from the left) by any functions  $\xi(t, u)$ ,

 $\xi^{-}(t^{-}, u^{-})$  and  $\xi^{+}(t^{+}, u^{+})$ , respectively, without loss the nature of its tangent conditions (for belonging to a higher order group or Lie-Bäcklund group, see [37]).

We also use a pair of shift operators to the right and left hand-sides, respectively:

$$S_{+}(f(t,u)) = f(t+\tau, u^{+}) = f^{+}, \qquad S^{-}(f(t,u)) = f(t-\tau, u^{-}) = f^{-}.$$
 (2. 3)

The shift operators can be expressed using the operators of the total derivative D:

$$S_{+} = \sum_{s \ge 0} \frac{\tau^{s}}{s!} D^{s}, \quad S^{-} = \sum_{s \ge 0} \frac{(-\tau)^{s}}{s!} (D^{-})^{s}.$$

Here and below, it is assumed that the delay parameter  $\tau = const$ . To keep regular intervals invariant with respect to generator (2. 1) we should consider those of them, which satisfy the condition (see [17]):

$$(S_{+} - 1)(S_{-} - 1)[\xi(t, u)] = 0, \qquad (2. 4)$$

or

$$\xi^+ - 2\xi + \xi^- = 0. \tag{2.5}$$

Thus, the criterion of the invariance of equation (1. 1) reads:

$$\tilde{X}(\ddot{u} - F(\ddot{u}^{-}, \dot{u}^{-}, \dot{u}, u, u^{-}, t, ...))|_{(1.1)} = 0,$$
(2. 6)

provided by condition (2.5).

## 3 The invariance of delay Lagrangians

Let a one-parameter group of point transformations be:

$$t_* = f(t, u, a), \quad u_* = g(t, u, a), \quad t_*^- = f^-(t^-, u^-, a), \quad u_*^- = g^-(t^-, u^-, a), \tag{3. 1}$$

which are prolonged for derivatives with the help of standard formulas of Lie group analysis [3–5]. We need also a group action on the differential:

$$dt_* = Df(t, u, a)da. \tag{3. 2}$$

Considering delay functional

$$L = \int_{a}^{b} \mathcal{L}(t, u, u^{-}, \dot{u}, \dot{u}^{-}) dt, \qquad (3. 3)$$

we chose any inner interval  $a + \tau < t_1 < t_2 < b$  for the analysis:

$$L = \int_{t_1}^{t_2} \mathcal{L}(t, u, u^-, \dot{u}, \dot{u}^-) dt.$$
(3. 4)

The functional is referred as invariant, if

$$L = \int_{t_1}^{t_2} \mathcal{L}(t, u, u^-, \dot{u}, \dot{u}^-) dt = \int_{t_1^*}^{t_2^*} \mathcal{L}(t_*, u_*, u_*^-, \dot{u}_*, \dot{u}_*^-) dt^*.$$
(3. 5)

The transformed variables and the interval of integration are represented by means of the initial values and the group parameter

$$\int_{t_1}^{t_2} \mathcal{L}(t, u, u^-, \dot{u}, \dot{u}^-) dt = \int_{t_1}^{t_2} \mathcal{L}(f, g, g^-, \dot{g}, \dot{g}^-, ) J(\frac{f}{t}) dt, \qquad (3. 6)$$

where  $J(\frac{f}{t})$  is the Jacobian of the transformation (3. 1).

As the interval is arbitrary one can omit the integration:

$$\mathcal{L}(t, u, u^{-}, \dot{u}, \dot{u}^{-})dt = \mathcal{L}(f, g, g^{-}, \dot{g}, \dot{g}^{-}, )J(\frac{f}{t})dt.$$
(3. 7)

Hence, the variational integral is invariant if and only if the elementary action is invariant with respect to (3, 7).

Differentiating (3. 7) with respect to a group parameter a, and setting a = 0, we obtain a criterion for a delay functional to be invariant.

**Theorem 3.1.** (First Noether's theorem for delay ODEs)

Functional (3. 3) is invariant if and only if

$$\xi \frac{\partial \mathcal{L}}{\partial t} + \xi^{-} \frac{\partial \mathcal{L}}{\partial t^{-}} + \eta \frac{\partial \mathcal{L}}{\partial u} + \eta^{-} \frac{\partial \mathcal{L}}{\partial u^{-}} +$$

$$+ (D(\eta) - \dot{u}D(\xi)) \frac{\partial \mathcal{L}}{\partial \dot{u}} + (D^{-}(\eta^{-}) - \dot{u}^{-}D^{-}(\xi^{-})) \frac{\partial \mathcal{L}}{\partial \dot{u}^{-}} + \mathcal{L}D(\xi) = 0.$$
(3.8)

We should note that, in contrast to the classical case of ODEs, there are *two* group orbits passing through the elementary action  $\mathcal{L}(t, u, u^-, \dot{u}, \dot{u}^-)dt$ . When  $\tau \to 0$  the criterion (3. 8) becomes the classical first Noether's theorem.

*Remark* 1. Paper [38] (see also [39]) contains wrong formula for the invariance of a delay Lagrangian. Therefore, the subsequent formulas and formulations are also incorrect.

# 4 The Elsgolts extremal equation and local extremal equation

Consider the delay functional (3, 3)

$$L = \int_{a}^{b} \mathcal{L}(t, t^{-}, u, u^{-}, \dot{u}, \dot{u}^{-}) dt$$
(4. 9)

and apply the variational operator along a fixed curve (for considerations of directional variation see [40]).

The variation of all variables is applied *only* along the group orbit at the internal point (t, u), a < t < b:

 $\delta t = \xi \delta a, \quad \delta u = \eta \delta a, \quad \delta \dot{u} = (D(\eta) - \dot{u} D(\xi)) \delta a, \quad \delta(dt) = D(\xi) dt \delta a,$ 

where  $\delta a$  is the variation of the group parameter a.

Variation of the functional only along a group orbit yields:

$$\delta L = \int_{a}^{b} \left[ \left( \frac{\partial \mathcal{L}}{\partial t} \delta t + \frac{\partial \mathcal{L}^{+}}{\partial t} \delta t + \frac{\partial \mathcal{L}}{\partial u} \delta u + \frac{\partial \mathcal{L}^{+}}{\partial u} \delta u + \frac{\partial \mathcal{L}}{\partial \dot{u}} \delta \dot{u} + \frac{\partial \mathcal{L}^{+}}{\partial \dot{u}} \delta \dot{u} \right) dt + \mathcal{L}\delta(dt) \right] = \\ = \int_{a}^{b} \left( \xi \left( \frac{\partial \mathcal{L}}{\partial t} + \frac{\partial \mathcal{L}^{+}}{\partial t} \right) + \eta \left( \frac{\partial \mathcal{L}}{\partial u} + \frac{\partial \mathcal{L}^{+}}{\partial u} \right) + (D(\eta) - \dot{u}D(\xi)) \left( \frac{\partial \mathcal{L}}{\partial \dot{u}} + \frac{\partial \mathcal{L}^{+}}{\partial \dot{u}} \right) + \mathcal{L}D(\xi) \right) dt \delta a.$$

$$(4. 10)$$

Integration by parts gives (here boundary values are not varied):

$$\delta L = \int_{a}^{b} \xi \left( \frac{\partial \mathcal{L}}{\partial t} + \frac{\partial \mathcal{L}^{+}}{\partial t} + D \left( \dot{u} \frac{\partial \mathcal{L}}{\partial \dot{u}} \right) + D \left( \dot{u} \frac{\partial \mathcal{L}^{+}}{\partial \dot{u}} \right) - D(\mathcal{L}) \right) + \\ + \eta \left( \frac{\partial \mathcal{L}}{\partial u} + \frac{\partial \mathcal{L}^{+}}{\partial u} - D \left( \frac{\partial \mathcal{L}}{\partial \dot{u}} \right) - D \left( \frac{\partial \mathcal{L}^{+}}{\partial \dot{u}} \right) \right) dt \delta a.$$
(4. 11)

The integral is equal to zero for arbitrary interval of integration if the integrand is zero. Hence, we arrive at *the local extremal equation*:

$$\xi \left( \frac{\partial \mathcal{L}}{\partial t} + \frac{\partial \mathcal{L}^+}{\partial t} + D\left( \dot{u} \frac{\partial \mathcal{L}}{\partial \dot{u}} \right) + D\left( \dot{u} \frac{\partial \mathcal{L}^+}{\partial \dot{u}} \right) - D(\mathcal{L}) \right) + \eta \left( \frac{\partial \mathcal{L}}{\partial u} + \frac{\partial \mathcal{L}^+}{\partial u} - D\left( \frac{\partial \mathcal{L}}{\partial \dot{u}} \right) - D\left( \frac{\partial \mathcal{L}^+}{\partial \dot{u}} \right) \right) = 0,$$

$$(4. 12)$$

which *explicitly depends* on  $\xi$ ,  $\eta$ , i.e. on a given group.

Notice, that a local extremal equation contains  $\mathcal{L}$  and  $\mathcal{L}^+$ , and therefore, the equation contains *two delays*. The operator D is applied at the point (t, u), and the directional variation only occurs along the group orbit passing through the point (t, u).

"Vertical variation", i.e. variation by u when  $\xi = 0$ , yields the following extremal delay equation

$$\frac{\partial \mathcal{L}}{\partial u} - D(\frac{\partial \mathcal{L}}{\partial \dot{u}}) + \frac{\partial \mathcal{L}^+}{\partial u} - D(\frac{\partial \mathcal{L}^+}{\partial \dot{u}}) = \frac{\delta}{\delta u}(\mathcal{L} + \mathcal{L}^+) = 0, \qquad (4. 13)$$

This equation is known since Elsgolts [36] (see also [45], [46]). We will call it *the Elsgolts equation*.

We call the Elsgolts variational derivative the following operator

$$\frac{\delta}{\delta u_{(E)}} = \frac{\partial}{\partial u} - D\frac{\partial}{\partial \dot{u}} + S_+ \left(\frac{\partial}{\partial u^-} - D\frac{\partial}{\partial \dot{u}^-}\right), \qquad (4. 14)$$

which one can apply to a delay Lagrangian  $\mathcal{L}$ . Notice, that the shift operator  $S_+$  acts on all arguments.

The Elsgolts equation operates with  $\mathcal{L}$  and  $\mathcal{L}^+$ , while the criterion of invariance of a Lagrangian only involves  $\mathcal{L}$ . We should stress that in contrast to a case of ODEs and PDEs, the Elsgolts equation is generally not an equation, on a solution of which the Lagrangian achieves its extremal value.

We also separate out "horizontal" variation

$$\frac{\delta}{\delta t}(\mathcal{L} + \mathcal{L}^{+}) = \frac{\partial \mathcal{L}}{\partial t} + \frac{\partial \mathcal{L}^{+}}{\partial t} + D\left(\dot{u}\frac{\partial \mathcal{L}}{\partial \dot{u}}\right) + D\left(\dot{u}\frac{\partial \mathcal{L}^{+}}{\partial \dot{u}}\right) - D(\mathcal{L}) = 0, \quad (4. 15)$$

which yields a variational equation for the case  $\eta=0$ .

The horizontal variational operator is:

$$\frac{\delta}{\delta t} = \frac{\partial}{\partial t} + D\left(\dot{u}\frac{\partial}{\partial \dot{u}}\right) + S_{+}\left(\frac{\partial}{\partial t^{-}} + D\left(\dot{u}\frac{\partial}{\partial \dot{u}^{-}}\right)\right) - D.$$
(4. 16)

Now one can rewrite the local extremal equation as

$$\xi \frac{\delta L}{\delta t} + \eta \frac{\delta L}{\delta u_{(E)}} = 0. \tag{4. 17}$$

The local extremal equation gives a necessary condition for a Lagrangian to achieve extremal value on the direction of a given group. The invariance of a Lagrangian is not needed (for directional variation see [40]). Connection of the invariance of a Lagrangian with local extremal equation is the subject of the next section.

### 5 Invariance of the Elsgolts equation

Consider the generator

$$X = \xi(t, u)\partial_t + \eta(t, u)\partial_u$$

preserving constant delay.

**Lemma 5.1.** Let the coefficient  $\xi(t)$  of the generator

$$X = \xi(t)\partial_t + \eta(t, u)\partial_u$$

satisfy condition (2. 5) and  $\dot{\xi}_t = \dot{\xi}_{t+1}^+$ . The following identity holds:

$$\frac{\delta}{\delta u_{(E)}} \left( X \mathcal{L} + \mathcal{L} D(\xi) \right) = X \left( \frac{\delta \mathcal{L}}{\delta u_{(E)}} \right) + \left( \eta_u + D(\xi) - \dot{u} \xi_u \right) \frac{\delta \mathcal{L}}{\delta u_{(E)}}.$$
(5. 18)

The proof can be obtained by direct calculations.

In more details relation (5. 18) can be rewritten as

$$\left(\frac{\partial}{\partial u} - \bar{D}\frac{\partial}{\partial \dot{u}}\right) \left(X\mathcal{L} + \mathcal{L}\bar{D}(\xi) + X\mathcal{L}^{+} + \mathcal{L}^{+}\bar{D}(\xi^{+})\right)$$

$$= X\left(\left(\frac{\partial}{\partial u} - \bar{D}\frac{\partial}{\partial \dot{u}}\right) \left(\mathcal{L} + \mathcal{L}^{+}\right)\right) + \left(\left(\frac{\partial}{\partial u} - \bar{D}\frac{\partial}{\partial \dot{u}}\right) \left(\mathcal{L} + \mathcal{L}^{+}\right)\right) \left(\eta_{u} + \bar{D}(\xi) + \dot{u}\xi_{u}\right).$$

$$(5. 19)$$

From identity (5. 18) one can establish the relation between the invariance of a Lagrangian and the invariance of the corresponding Elsgolts equation.

Considering Eq.(5. 18) on a solution of the Elsgolts equation one has

$$\frac{\delta}{\delta u_{(E)}} \left( X\mathcal{L} + \mathcal{L}D(\xi) \right) \big|_{\frac{\delta \mathcal{L}}{\delta u_{(E)}} = 0} = X \left( \frac{\delta \mathcal{L}}{\delta u_{(E)}} \right) \big|_{\frac{\delta \mathcal{L}}{\delta u_{(E)}} = 0} = 0.$$

**Theorem 5.2.** Let the generator X satisfy property (2. 5) and  $\dot{\xi}_t = \dot{\xi}_{t+t+}$ . Then the Elsgolts equation is invariant if and only if the following condition holds:

$$\frac{\delta}{\delta u_{(E)}} \left( X \mathcal{L} + \mathcal{L} D(\xi) \right) \big|_{\frac{\delta \mathcal{L}}{\delta u_{(E)}} = 0} = 0$$
(5. 20)

for any solution of the Elsgolts equation.

Note that Theorem 5.2 well matches with the analogous theorem for ODE [41].

**Example 5.1.** Consider the Lie algebra  $L_{4.13}$  [43] of the generators

$$X_1 = \frac{\partial}{\partial u}, \quad X_2 = \frac{\partial}{\partial t}, \quad X_3 = t \frac{\partial}{\partial t}, \quad X_4 = u \frac{\partial}{\partial u},$$
 (5. 21)

which preserves the uniformity of delay parameter  $\tau$ .

The complete set of invariants of this Lie algebra consists of the invariants

$$I_1 = \frac{u - u^-}{\dot{u}(t - t^-)}, \quad I_2 = \frac{\dot{u}^-}{\dot{u}}, \quad I_3 = (t - t^-)\frac{\ddot{u}}{\dot{u}}.$$

As all generators (5. 21) satisfy property (5. 20), the Elsgolts equation with the Lagrangian  $\mathcal{L} = \mathcal{L}(I_1, I_2)$  admits the generators (5. 21). Meanwhile the Lagrangian  $\mathcal{L} = \mathcal{L}(I_1, I_2)$  does not admit the scaling generator  $X_3 = t \frac{\partial}{\partial t}$ .

## 6 First integrals for DODEs

The standard form of a solution (integral) of an ODE

$$I(t,u) = A, (6. 22)$$

can not satisfy any initial data which are given on an initial interval for DODEs. Hence, we have to introduce another form of an integral.

DODEs (1. 1) can contain the dependent variable and its derivatives at three points  $t^+$ , t and  $t^-$ . It is possible to consider two types of conserved quantities: differential first integrals and difference first integrals.

**Definition 6.1.** Quantity

$$I(t^+, t, t^-, u^+, u, u^-, \dot{u}^+, \dot{u}, \dot{u}^-) = A_0$$
(6. 23)

is called a *differential first integral* of DODE (1, 1) if it holds constant on the solutions of the DODE.

Consideration of differential first integrals needs differentiation and requires to chose constant delays. We have to define differentiation of functions, which depend on the dependent variables for several values of the independent variable. Differentiating I at some point, for example, at the point t, we get

$$\bar{D}(I) = (I_{t^+} + I_{u^+} \dot{u}^+ + I_{\dot{u}^+} \ddot{u}^+) \frac{dt^+}{dt} + (I_t + I_u \dot{u} + I_{\dot{u}} \ddot{u}) + (I_{t^-} + I_{u^-} \dot{u}^- + I_{\dot{u}^-} \ddot{u}^-) \frac{dt^-}{dt},$$

where

$$\bar{D} = \frac{\partial}{\partial t} + \dot{u}\frac{\partial}{\partial u} + \ddot{u}\frac{\partial}{\partial \dot{u}} + \dots + \frac{\partial}{\partial t^{-}} + \dot{u}^{-}\frac{\partial}{\partial u^{-}} + \ddot{u}^{-}\frac{\partial}{\partial \dot{u}^{-}} + \dots + \frac{\partial}{\partial t^{+}} + \dot{u}^{+}\frac{\partial}{\partial u^{+}} + \ddot{u}^{+}\frac{\partial}{\partial \dot{u}^{+}} + \ddot{u}^{+}\frac{\partial}{\partial \dot{u}^{+}} + \dots \quad (6. 24)$$

Generally, the delay can depend on the independent variable t as well as on the solution u(t) of the DODE. Then, the values  $t^+$  and  $t^-$  have the same dependence. It makes sense to restrict the considered delays by the conditions

$$\frac{dt^+}{dt} = 1, \qquad \frac{dt^-}{dt} = 1$$
 (6. 25)

for all considered t. In this case the differentiation does not depend on a chosen point, and we obtain that the differential first integrals should satisfy the equation

$$\bar{D}(I) = I_{t^+} + I_t + I_{t^-} + I_{u^+} \dot{u}^+ + I_u \dot{u} + I_{u^-} \dot{u}^- + I_{\dot{u}^+} \ddot{u}^+ + I_{\dot{u}} \ddot{u} + I_{\dot{u}^-} \ddot{u}^- = 0, \qquad (6.\ 26)$$

which should hold on the solutions of the considered DODE.

Conditions (6. 25) imply that the delays should be solution independent. Moreover, integrating these conditions, we obtain the delay equations that justifies the choice of the constant delays for Eq. (1. 1). If we tried to consider nonconstant delays, we would have difficulties with defining differentiation and differential first integrals.

#### **Definition 6.2.** A function

$$J(t, t^{-}, u, u^{-}, \dot{u}, \dot{u}^{-}, \ddot{u}, \ddot{u}^{-})$$
(6. 27)

is called a difference first integral of DODE (1, 1) if it satisfies the equation

$$(S_+ - 1)J = 0 (6. 28)$$

on solutions of the DODE.

We illustrate the definitions of the first integrals on a simple example.

#### Example 6.1. DODE

has differential first integral

 $J = \ddot{u} - \ddot{u}^{-}$ .

Differential first integrals are constant on the solutions of DODEs. In this case, the difference first integral should not be constant on DODE solutions, they can be periodic functions with period  $\tau$ , where  $\tau$  is the delay parameter.

### 7 The Noether-type identity for delay Lagrangians

In this Section we introduce the main operator identity, which relates together the invariance of a Lagrangian and first integral of local extremal equation.

We consider a Lagrangian function

$$\mathcal{L} = \mathcal{L}(t, t^{-}, u, u^{-}, \dot{u}, \dot{u}^{-}), \qquad (7. 29)$$

which depends on  $t, t^-$ , and we associate its invariance with first integral.

```
\ddot{u}^+ = \ddot{u}^-
```

 $I = \dot{u}^+ - \dot{u}^-.$ 

**Lemma 7.1.** The following identity holds  $(\tau = const)$ :

$$\begin{split} \xi \frac{\partial \mathcal{L}}{\partial t} + \xi^{-} \frac{\partial \mathcal{L}}{\partial t^{-}} + \eta \frac{\partial \mathcal{L}}{\partial u} + \eta^{-} \frac{\partial \mathcal{L}}{\partial u^{-}} + (D(\eta) - \dot{u}D(\xi)) \frac{\partial \mathcal{L}}{\partial \dot{u}} + \\ + (D^{-}(\eta^{-}) - \dot{u}^{-}D^{-}(\xi^{-})) \frac{\partial \mathcal{L}}{\partial \dot{u}^{-}} + \mathcal{L}D(\xi) \equiv \\ \equiv \eta \left( \frac{\partial \mathcal{L}}{\partial u} + \frac{\partial \mathcal{L}^{+}}{\partial u} - D\left(\frac{\partial \mathcal{L}}{\partial \dot{u}}\right) - D\left(\frac{\partial \mathcal{L}^{+}}{\partial \dot{u}}\right) \right) + \\ + \xi \left( \frac{\partial \mathcal{L}}{\partial t} + \frac{\partial \mathcal{L}^{+}}{\partial t} + D\left(\dot{u}\frac{\partial \mathcal{L}}{\partial \dot{u}}\right) + D\left(\dot{u}\frac{\partial \mathcal{L}^{+}}{\partial \dot{u}}\right) - D(\mathcal{L}) \right) + \\ + \bar{D} \left[ \xi \mathcal{L} + (\eta - \dot{u}\xi) \left( \frac{\partial \mathcal{L}}{\partial \dot{u}} + \frac{\partial \mathcal{L}^{+}}{\partial \dot{u}} \right) \right] + \\ + (1 - S_{+}) \left( \xi^{-} \frac{\partial \mathcal{L}}{\partial t^{-}} + \eta^{-} \frac{\partial \mathcal{L}}{\partial u^{-}} + (D^{-}(\eta^{-}) - \dot{u}^{-}D^{-}(\xi^{-})) \frac{\partial \mathcal{L}}{\partial \dot{u}^{-}} \right) = 0. \end{split}$$

$$(7. 30)$$

Identity (7. 30) allows one to specify different DODEs on the right-hand side and formulate various appropriate theorems (a similar idea was implemented for ODEs in [42]).

#### Theorem 7.2. (Delay version of Noether's theorem for a local extremal equation)

Let a functional  $\mathcal{L}$  be invariant with respect to a one-parameter group of transformations. Then the invariance of the Lagrangian gives the integral

$$\bar{D}\left[\xi\mathcal{L} + (\eta - \dot{u}\xi)\left(\frac{\partial\mathcal{L}}{\partial\dot{u}} + \frac{\partial\mathcal{L}^+}{\partial\dot{u}}\right)\right] = 0$$
(7. 31)

for each solution of a local extremal equation (4. 17), provided by the difference relation

$$\xi^{-}\frac{\partial \mathcal{L}}{\partial t^{-}} + \eta^{-}\frac{\partial \mathcal{L}}{\partial u^{-}} + (D^{-}(\eta^{-}) - \dot{u}^{-}D^{-}(\xi^{-}))\frac{\partial \mathcal{L}}{\partial \dot{u}^{-}} =$$

$$= \xi\frac{\partial \mathcal{L}^{+}}{\partial t} + \eta\frac{\partial \mathcal{L}^{+}}{\partial u} + (D(\eta) - \dot{u}D(\xi))\frac{\partial \mathcal{L}^{+}}{\partial \dot{u}}.$$
(7. 32)

The difference integral (or periodic condition) relates corresponding values at two points and is not a constant for each point of the considered interval.

*Remark* 2. It should be noted that the local extremal equation (4. 17) depends on second-order derivatives, and usually contains *two* delays.

#### Consequences of Lemma 7.1 and Theorem 7.2

The 7.2 theorem has an obvious generalization for the following cases:

1. If an action of a symmetry operator on an elementary Lagrangian action  $\mathcal{L}dt$  gives  $\overline{D}(A(t, t^-, u, u^-, \dot{u}, \dot{u}^-))$ , then one can add the corresponding function (-A) into the expression of local extremal equation. This is a generalization similar to that for ODE [44].

2. If an action of a symmetry operator on an elementary Lagrangian action  $\mathcal{L}dt$  gives  $(S_+ - 1)(B(t, t^-, u, u^-, \dot{u}, \dot{u}^-))$ , then one can add the corresponding function (-B) into the expression of difference integral (periodic condition).

Both cases can exist in one action of a symmetry operator on an elementary Lagrangian action  $\mathcal{L}dt$ .

3. It may happens that expression (7. 32) is equivalent to some divergent term  $\overline{D}(C(t, t^-, u, u^-, \dot{u}, \dot{u}^-))$ :

$$\xi^{-} \frac{\partial \mathcal{L}}{\partial t^{-}} + \eta^{-} \frac{\partial \mathcal{L}}{\partial u^{-}} + (D^{-}(\eta^{-}) - \dot{u}^{-} D^{-}(\xi^{-})) \frac{\partial \mathcal{L}}{\partial \dot{u}^{-}} -$$
(7. 33)  
$$-\xi \frac{\partial \mathcal{L}^{+}}{\partial t} - \eta \frac{\partial \mathcal{L}^{+}}{\partial u} - (D(\eta) - \dot{u} D(\xi)) \frac{\partial \mathcal{L}^{+}}{\partial \dot{u}} = \bar{D}(C),$$

then we arrive to the case 1 (see examples of the linear oscillator below).

Below we consider two special cases of Noether's theorem for two special cases of admitted group, which are in the examples.

#### Corollary. (First special version of Noether's theorem for delay ODEs)

Let a functional  $\mathcal{L}$  be invariant with respect to a one-parameter group of transformations with  $\xi = 0$ . Then the invariance of the Lagrangian gives the integral

$$\bar{D}\left[\eta\left(\frac{\partial\mathcal{L}}{\partial\dot{u}} + \frac{\partial\mathcal{L}^+}{\partial\dot{u}}\right)\right] = 0 \tag{7.34}$$

for each solution of the Elsgolts equation

$$\frac{\partial \mathcal{L}}{\partial u} + \frac{\partial \mathcal{L}^+}{\partial u} - \bar{D} \left( \frac{\partial \mathcal{L}}{\partial \dot{u}} \right) - \bar{D} \left( \frac{\partial \mathcal{L}^+}{\partial \dot{u}} \right) = 0, \qquad (7. 35)$$

under the following periodic condition (difference integral)

$$\eta^{-} \frac{\partial \mathcal{L}}{\partial u^{-}} + (D^{-}(\eta^{-}) - \dot{u}^{-} D^{-}(\xi^{-})) \frac{\partial \mathcal{L}}{\partial \dot{u}^{-}} =$$

$$= \eta \frac{\partial \mathcal{L}^{+}}{\partial u} + (D(\eta) - \dot{u} D(\xi)) \frac{\partial \mathcal{L}^{+}}{\partial \dot{u}}.$$
(7. 36)

#### Corollary. (Second special version of Noether's theorem for delay ODEs)

Let a functional  $\mathcal{L}$  be invariant with respect to a one-parameter group of transformations with  $\eta = 0$ . Then the invariance of Lagrangian yields the integral

$$\bar{D}\left[\xi\mathcal{L} + \dot{u}\xi\left(\frac{\partial\mathcal{L}}{\partial\dot{u}} + \frac{\partial\mathcal{L}^+}{\partial\dot{u}}\right)\right] = 0$$
(7. 37)

for each solution of the following equation

$$\frac{\partial \mathcal{L}}{\partial t} + \frac{\partial \mathcal{L}^+}{\partial t} + \bar{D}\left(\dot{u}\frac{\partial \mathcal{L}}{\partial \dot{u}}\right) + \bar{D}\left(\dot{u}\frac{\partial \mathcal{L}^+}{\partial \dot{u}}\right) - \bar{D}(\mathcal{L}) = 0, \qquad (7. 38)$$

under the following periodic condition (difference integral)

$$\xi^{-}\frac{\partial \mathcal{L}}{\partial t^{-}} - \dot{u}^{-}D^{-}(\xi^{-})\frac{\partial \mathcal{L}}{\partial \dot{u}^{-}} = \xi\frac{\partial \mathcal{L}^{+}}{\partial t} - \dot{u}D(\xi)\frac{\partial \mathcal{L}^{+}}{\partial \dot{u}}.$$
(7. 39)

*Remark* 3. It should be noted that the Theorems formulated above yield restriction (periodic conditions) on the set of solutions of the corresponding equations (7.35) and (7.38).

Remark 4. It may happens that expression (7. 36) or (7. 39) is equivalent to some divergent term  $\overline{D}(C)$ , then we add the term  $-\overline{D}(C)$  to integrals (7. 34), (7. 37). In these cases we have no periodic restrictions for the set of solutions.

## 8 Examples of applications

#### 1. Linear oscillator 1

Consider the following delay Lagrangian

$$\mathcal{L} = uu^- - \dot{u}\dot{u}^-, \tag{8. 40}$$

which is divergently invariant with respect to the generator  $X_1 = \cos t \frac{\partial}{\partial u}$ :

$$X_1(\mathcal{L}) = u^- \cos t + u \cos t^- + \dot{u}^- \sin t + \dot{u} \sin t^- = \bar{D}[u^- \sin t + u \sin t^-].$$

Notice that this symmetry allows one to consider the constant  $\tau$  for the whole interval. According to Theorem 7.2, we get the following difference integral

$$u\cos t^{-} + \dot{u}\sin t^{-} = u^{+}\cos t + \dot{u}^{+}\sin t.$$
(8. 41)

Taking into account divergence term in the action of  $X_1$  on the Lagrangian, we get the following differential first integral

$$\bar{D}(\cos t(\dot{u}^+ + \dot{u}^-) + u\sin t^- + u^-\sin t) = 0, \qquad (8.42)$$

for solutions of the Elsgolts equation

$$\frac{\delta \mathcal{L}}{\delta u_{(E)}} = \ddot{u}^+ + \ddot{u}^- + u^+ + u^- = 0.$$
(8. 43)

The first integral (8. 42) is equivalent to the Elsgolts equation (8. 43) by virtue of difference relation (8. 41). It is easy to check that eq. (8. 43) is invariant with respect to  $X_1$ .

The delay Lagrangian (8. 40) is divergently invariant with respect to the generator  $X_2 = \sin t \frac{\partial}{\partial u}$ :

$$X_2(\mathcal{L}) = u^{-} \sin t + u \sin t^{-} - \dot{u}^{-} \cos t - \dot{u} \cos t^{-} = -\bar{D}[u^{-} \cos t + u \cos t^{-}].$$

According to Theorem 7.2, we derive the difference integral

$$u\sin t^{-} - \dot{u}\cos t^{-} = u^{+}\sin t - \dot{u}^{+}\cos t.$$
(8. 44)

Taking into account the divergent term in the action of  $X_2$  on the Lagrangian elementary action, we get the following differential first integral

$$\bar{D}\left(-\sin t(\dot{u}^{+}+\dot{u}^{-})+u\cos t^{-}+u^{-}\cos t\right)=0,$$
(8. 45)

for a solution of the same Elsgolts eq. (8. 43). Notice that eq. (8. 43) is invariant with respect to  $X_2$ .

The first integral (8. 66) is equivalent to Elsgolts eq. (8. 43) by virtue of difference relation (8. 44).

Thus, for Elsgolts eq. (8. 43) we derived two differential delay integrals

$$\cos t(\dot{u}^{+} + \dot{u}^{-}) + u\sin t^{-} + u^{-}\sin t = A, \qquad (8.46)$$

$$-\sin t(\dot{u}^{+} + \dot{u}^{-}) + u\cos t^{-} + u^{-}\cos t = B,$$

from which one can obtain the integral of the form (6. 23)

$$u\cos\tau + u^- = A\sin t + B\cos t. \tag{8.47}$$

Notice that we have used both differential and difference integrals.

Remark 5. Consider an oscillating solution of the Elsgolts eq. (8.54) of the form

$$u = \alpha \sin t + \beta \cos t, \quad \alpha, \beta = const, \tag{8.48}$$

that means that we also take

$$u^{-} = \alpha \sin(t - \tau) + \beta \cos(t - \tau) \tag{8.49}$$

as the initial data together with (8. 48). Hence, u and  $u^-$  are given as initial conditions on the intervals  $[-2\tau, -\tau]$  and  $[-\tau, 0]$  respectively, while  $u^+$  is a solution to be found.

One can show that in this case the solution

$$u^{+} = \alpha \sin(t+\tau) + \beta \cos(t+\tau)$$
(8. 50)

solves the Elsgolts eq. (8. 43). Substituting the solutions (8. 48), (8. 49) into integral (8. 47), one obtains the relations for the constants:

$$2\alpha\cos\tau + \beta\sin\tau = A,\tag{8.51}$$

 $2\beta\cos\tau + \alpha\sin\tau = B.$ 

The Elsgolts eq. (8, 43) also possesses infinitely many solutions besides special oscillating solution (8, 48).

#### 2. An alternative approach to linear oscillator

Consider the same Lagrangian  $\mathcal{L} = uu^- - \dot{u}\dot{u}^-$ , which is divergently invariant with respect to the generator  $X_1 = \cos t \frac{\partial}{\partial u} + \cos t^- \frac{\partial}{\partial u^-} - \sin t \frac{\partial}{\partial \dot{u}} - \sin t^- \frac{\partial}{\partial u^-}$ .

The difference expression

$$u\cos t^{-} + \dot{u}\sin t^{-} - u^{+}\cos t - \dot{u}^{+}\sin t = \bar{D}(u\sin t^{-} - u^{+}\sin t)$$
(8. 52)

is equivalent to exact divergence of some expression for the underlined case. We add this divergent term into the differential first integral

$$-\bar{D}(\cos t(\dot{u}^{+} + \dot{u}^{-}) + \sin t(u^{-} + u^{+})) =$$

$$-\cos t(\ddot{u}^{+} + \ddot{u}^{-} + u^{+} + u^{-}) = 0$$
(8. 53)

for the Elsgolts equation

$$\frac{\delta \mathcal{L}}{\delta u_{(E)}} = \ddot{u}^+ + \ddot{u}^- + u^+ + u^- = 0.$$
(8. 54)

The delay Lagrangian (8. 40) is also divergently invariant with respect to the generator  $X_2 = \sin t \frac{\partial}{\partial u}$ :

$$X_2(\mathcal{L}) = u^{-} \sin t + u \sin t^{-} - \dot{u}^{-} \cos t - \dot{u} \cos t^{-} = -\bar{D}[u^{-} \cos t + u \cos t^{-}].$$

The difference expression is

$$u\sin t^{-} - \dot{u}\cos t^{-} - u^{+}\sin t + \dot{u}^{+}\cos t = \bar{D}(u^{+}\cos t - u\cos t^{-})$$
(8. 55)

Taking into account the latter divergent term, we get the following differential first integral:

$$\bar{D}\left(-\sin t(\dot{u}^{+}+\dot{u}^{-})+\cos t(u^{-}+u^{+})\right) = -\sin t(\ddot{u}^{+}+\ddot{u}^{-}+u^{+}+u^{-}) = 0, \qquad (8.56)$$

which is equivalent to Elsgolts eq. (8.54).

Thus, for Elsgolts eq. (8.54) we derived two differential integrals

$$\cos t(\dot{u}^{+} + \dot{u}^{-}) + \sin t(u^{-} + u^{+}) = A,$$
$$-\sin t(\dot{u}^{+} + \dot{u}^{-}) + \cos t(u^{-} + u^{+}) = B,$$

from which one can obtain the integral of the form (6. 23)

$$u^{+} + u^{-} = A\sin t + B\cos t. \tag{8.57}$$

Notice that we have no restrictions for integral (8. 57).

#### 3. Nonlinear oscillator 1

Lagrangian (8. 40) is also invariant with respect to  $X_3 = \frac{\partial}{\partial t}$  and for the local extremal equation

$$\ddot{u}\dot{u}^{+} + \ddot{u}^{+}\dot{u} + \dot{u}u^{-} + \dot{u}^{-}u = 0 \tag{8.58}$$

we get zero difference integral and the following differential one

$$\bar{D}[\dot{u}\dot{u}^+ + uu^-] = 0.$$

Thus, the symmetry  $X_3$  yields another *nonlinear* local extremal equation and corresponding differential integral without periodic restriction for the set of solutions of Eq. (8. 58).

#### 4. Nonlinear oscillator 2

Consider the delay Lagrangian

$$\mathcal{L} = uu^- - \dot{u}\dot{u}^-,$$

which is divergently invariant with respect to the sum of the generators  $X_1 + X_3 = \frac{\partial}{\partial t} + \cos t \frac{\partial}{\partial u}$ :

$$(X_1 + X_3)\mathcal{L} = u^- \cos t + u \cos t^- + \dot{u}^- \sin t + \dot{u} \sin t^- = \bar{D}[u^- \sin t + u \sin t^-].$$
(8. 59)

In this case, the local extremal equation takes the form of the *first-order* DODE:

$$\sin t(\dot{u}^{+} + \dot{u}^{-}) + \cos t(u^{-} + u^{+}) = 0.$$
(8. 60)

According to Theorem 7.2, we get the following difference integral

$$u\cos t^{-} + \dot{u}\sin t^{-} + u^{+}\cos t - \dot{u}^{+}\sin t = 0.$$
(8. 61)

Taking into account divergent term in action of  $X_1 + X_3$  on the Lagrangian, we can rewrite the local extremal equation in the form

$$D(u^{+}\sin t - u\sin t^{-}) = 0, \qquad (8.62)$$

which is equivalent to the difference integral.

Thus, a solution can be found from the relation

$$u^{+}\sin t - u\sin t^{-} = A_0 \tag{8.63}$$

and initial data.

#### 5. Linear oscillator 2

Consider the following delay Lagrangian

$$\mathcal{L} = \frac{(\dot{u} + \dot{u}^{-})^2}{2} - \frac{(u + u^{-})^2}{2}, \qquad (8. 64)$$

which is divergently invariant with respect to the generator  $X_1 = \cos t \frac{\partial}{\partial u}$ :

$$X_1(\mathcal{L}) = -(u+u^-)(\cos t + \cos t^-) - (\dot{u}^- + \dot{u})(\sin t + \sin t^-) = -\bar{D}[(u+u^-)(\sin t + \sin t^-)].$$

According to Theorem 7.2, we obtain the following difference integral

$$(u+u^{-})\cos t^{-} + (\dot{u}+\dot{u}^{-})\sin t^{-} = (u+u^{+})\cos t + (\dot{u}+\dot{u}^{+})\sin t.$$
(8. 65)

Taking into account the divergence term in the action of  $X_1$  on the Lagrangian, we get the following differential first integral

$$\bar{D}[\cos t(\dot{u}^+ + 2\dot{u} + \dot{u}^-) + (u + u^-)(\sin t^- + \sin t)] = 0, \qquad (8.66)$$

for a solution of the Elsgolts equation

$$\frac{\delta \mathcal{L}}{\delta u_{(E)}} = \ddot{u}^+ + 2\ddot{u} + \ddot{u}^- + u^+ + 2u + u^- = 0.$$
(8. 67)

The first integral (8. 66) is equivalent to the Elsgolts eq. (8. 67) by virtue of difference relation (8. 65). One can check that Eq. (8. 67) is invariant with respect to  $X_1$ .

The same delay Lagrangian is divergently invariant with respect to the generator  $X_2 = \sin t \frac{\partial}{\partial u}$ :

$$X_2(\mathcal{L}) = (\dot{u}^- + \dot{u})(\cos t + \cos t^-) - (u + u^-)(\sin t + \sin t^-) = \bar{D}[u + u^-)(\cos t + \cos t^-)].$$

Theorem 7.2 yields the following difference integral

$$-\sin t^{-}(u^{-}+u) + \cos t^{-}(\dot{u}+\dot{u}^{-}) = -\sin t(u^{+}+u) + \cos t(\dot{u}+\dot{u}^{+}).$$
(8. 68)

Taking into account the divergent term in the action of  $X_2$  on the Lagrangian elementary action, we get the following differential first integral

$$\bar{D}[\sin t(\dot{u}^+ + 2\dot{u} + \dot{u}^-) - (u + u^-)(\cos t^- + \cos t)] = 0, \qquad (8.69)$$

for a solution of the same Elsgolts Eq. (8. 67). Eq. (8. 67) is also invariant with respect to  $X_2$ .

The first integral (8.69) is equivalent to the Elsgolts Eq. (8.67) by virtue of the difference relation (8.68).

Thus, for the Elsgolts Eq. (8. 67) we have got two differential delay integrals

$$\cos t(\dot{u}^{+} + 2\dot{u} + \dot{u}^{-}) + (u + u^{-})(\sin t^{-} + \sin t) = A, \qquad (8.70)$$

$$\sin t(\dot{u}^{+} + 2\dot{u} + \dot{u}^{-}) - (u + u^{-})(\cos t^{-} + \cos t) = B, \qquad (8.71)$$

from which one can obtain the integral of the form

$$(u+u^{-})(1+\cos\tau) = A\sin t - B\cos t.$$
(8. 72)

Again, we have used both differential and difference integrals.

*Remark* 6. There exists a special oscillating solution of the Elsgolts eq. (8. 67)

$$u = \alpha \sin t + \beta \cos t, \quad \alpha, \beta = const, \tag{8.73}$$

with the corresponding initial values and the relations for the constants:

$$(\alpha + \alpha \cos \tau + \beta \sin \tau)(1 + \cos \tau) = A,$$

$$(\beta + \beta \cos \tau - \alpha \sin \tau)(1 + \cos \tau) = -B.$$
(8. 74)

#### 6. Nonlinear DODEs

Consider the following delay Lagrangian

$$\mathcal{L} = \frac{(u - u^{-})^2}{\dot{u}\dot{u}^{-}},\tag{8.75}$$

which is invariant with respect to the following generators:

$$X_1 = \frac{\partial}{\partial u}; \quad X_2 = u \frac{\partial}{\partial u}; \quad X_3 = u^2 \frac{\partial}{\partial u}; \quad X_3 = \frac{\partial}{\partial t}.$$
 (8. 76)

1. Using the generator  $X_1$ , and according to Theorem 7.2, we have the following difference relation:

$$\frac{u-u^{-}}{\dot{u}\dot{u}^{-}} = \frac{u^{+}-u}{\dot{u}\dot{u}^{+}},\tag{8.77}$$

and the following differential first integral

$$\bar{D}\left(\frac{(u-u^{-})^{2}}{\dot{u}^{2}\dot{u}^{-}} + \frac{(u^{+}-u)^{2}}{\dot{u}^{2}\dot{u}^{+}}\right) = 0, \qquad (8.78)$$

for a solution of the Elsgolts equation

$$\frac{\delta \mathcal{L}}{\delta u_{(E)}} = \frac{2(u-u^{-})}{\dot{u}\dot{u}^{-}} - \frac{2(u^{+}-u)}{\dot{u}\dot{u}^{+}} + \bar{D}\left(\frac{(u-u^{-})^{2}}{\dot{u}^{2}\dot{u}^{-}} + \frac{(u^{+}-u)^{2}}{\dot{u}^{2}\dot{u}^{+}}\right) = 0.$$
(8. 79)

The first integral (8.78) is equivalent to Elsgolts eq. (8.79) by virtue of difference relation (8.77), which provides restrictions for a set of solutions of Elsgolts eq. (8.79).

One can check that eq. (8. 79) is invariant with respect to  $X_1$ .

2. The delay Lagrangian (8. 75) admits the generator  $X_2 = u \frac{\partial}{\partial u}$ . According to Theorem 7.2, we derive the difference relation:

$$u^{-}\frac{2(u-u^{-})}{\dot{u}^{2}\dot{u}^{-}} + \frac{(u-u^{-})^{2}}{\dot{u}\dot{u}^{-}} = u\frac{2(u^{+}-u)}{\dot{u}^{2}\dot{u}^{+}} + \frac{(u^{+}-u)^{2}}{\dot{u}\dot{u}^{+}},$$
(8. 80)

and the following differential first integral

$$\bar{D}\left(\frac{(u-u^{-})^{2}}{\dot{u}^{2}\dot{u}^{-}} + \frac{(u^{+}-u)^{2}}{\dot{u}^{2}\dot{u}^{+}}\right) = 0,$$
(8. 81)

for solutions of the Elsgolts equation (8. 79).

By virtue of difference relation (8. 80), the first integral (8. 81) is equivalent to the Elsgolts eq. (8. 79).

One can check that eq. (8. 79) is invariant with respect to  $X_2$ .

3. The delay Lagrangian (8. 75) also admits the generator  $X_3 = u^2 \frac{\partial}{\partial u}$ , which gives the difference relation:

$$(u^{-})^{2}\frac{(u-u^{-})}{\dot{u}^{-}} + u^{-}\frac{(u-u^{-})}{\dot{u}^{-}} = u^{2}\frac{(u^{+}-u)}{\dot{u}} + u\frac{(u^{+}-u)}{\dot{u}}, \qquad (8.82)$$

and the following differential first integral

$$\bar{D}\left(u^2\left(\frac{(u-u^-)^2}{\dot{u}^2\dot{u}^-} + \frac{(u^+-u)^2}{\dot{u}^2\dot{u}^+}\right)\right) = 0,$$
(8. 83)

for a solution of the Elsgolts equation (8. 79) multiplied by  $u^2$ . The first integral (8. 83) is equivalent to Elsgolts eq. (8. 79) by virtue of difference relation (8. 82), which gives constrains for a set of solutions of the Elsgolts equation.

One can check that eq. (8. 79) is invariant with respect to  $X_3$ .

4. The Lagrangian (8. 75) is also invariant with respect to  $X_4 = \frac{\partial}{\partial t}$ . In this case we have another local extremal equation

$$\bar{D}\left(\frac{2(u-u^{-})^{2}}{\dot{u}\dot{u}^{-}} + \frac{(u^{+}-u)}{\dot{u}\dot{u}^{+}}\right) = 0, \qquad (8.84)$$

which also admits  $X_4$ . In this case difference relation vanishes. The following differential integral holds

$$\frac{2(u-u^{-})^{2}}{\dot{u}\dot{u}^{-}} + \frac{(u^{+}-u)}{\dot{u}\dot{u}^{+}} = const.$$
(8. 85)

Thus, the symmetry  $X_4$  yields another nonlinear local extremal equation and corresponding differential integral, which is valid for all solutions of eq. (8. 84),

### 9 Conclusion

Lagrangian formalism for variational problem of second-order delay differential equations (DO-DEs) was considered in the present paper. The invariance condition for delay functional was established and the delay analogue of first Noether's theorem was proved. It was shown that the variation of a functional along the only group orbit gives some equation (named a local extremal equation), which explicitly depends on a group. For 'vertical variation' a local extremal equation degenerates to the Elsgolts equation named after mathematician, who was the first to derive it. For 'horizontal' variation one gets some other DODEs. The necessary and sufficient condition for the Elsgolts equation to be invariant is established. The local extremal equation and the Elsgolts equation have in general *two* delay parameters and it is far from common consideration of DODEs in the literature. The appropriate initial value problem for a DODE with two delays was formulated. The generalization of the term a 'first integral' for DODEs has been given.

The basic Noether-type operator identity and delay analogues of the Noether theorem for invariant of second-order delay ordinary differential equations are presented. The integration technique of DODEs based on symmetry is demonstrated by examples.

### Dedication

The article is dedicated to the memory of Pavel Winternitz in recognition of his diverse contributions to applications of Lie group methods. The authors were happy to collaborate with him for many years.

## References

- S. Lie. Klassifikation und Integration von gewöhnlichen Differentialgleichungen zwischen x, y, die eine Gruppe von Transformationen gestatten. I, II. Math. Ann., 32:213–281, 1888. Gesammelte Abhandlungen, vol. 5, B.G. Teubner, Leipzig, 1924, pp. 240–310.
- [2] S. Lie. Gruppenregister. Gesammelte Abhandlungen, 5:767–773, 1924.
- [3] L. V. Ovsiannikov. Group Analysis of Differential Equations. Nauka, Moscow, 1978. English translation, Ames, W.F., Ed., published by Academic Press, New York, 1982.
- [4] P. J. Olver. Applications of Lie Groups to Differential Equations. Springer-Verlag, New York, 1986.
- [5] N. H. Ibragimov. Transformation Groups Applied to Mathematical Physics. Nauka, Moscow, 1983. English translation, Reidel, D., Ed., Dordrecht, 1985.
- [6] S. C. Anco and G. Bluman. Direct construction of conservation laws from field equations. *Phys. Rev. Lett.*, 78(3):2869–2873, 1997.
- [7] N. H. Ibragimov, editor. CRC Handbook of Lie Group Analysis of Differential Equations, volume 1, 2, 3. CRC Press, Boca Raton, 1994, 1995, 1996.
- [8] G. Gaeta. Nonlinear Symmetries and Nonlinear Equations. Kluwer, Dordrecht, 1994.
- [9] E. Noether. Invariante variations problem. Nachr. d. Königlichen Gesellschaft der Wissenschaften zu Göttingen, Nachrichten, Mathematisch-Physikalische Klasse Heft 2, pages 235–257, 1918. English translation : Transport Theory and Statist. Phys., 1(3), 1971, 183-207, (arXiv:physics/0503066 [physics.hist-ph]).
- [10] G. W. Bluman and S. C. Anco. Symmetry and Integration Methods for Differential Equations. Springer, New York, 2002.
- [11] V. A. Dorodnitsyn. Transformation groups in net spaces. Journal of Soviet Mathematics, 55(1):1490–1517, 1991.
- [12] V. A. Dorodnitsyn. Finite-difference models entirely inheriting symmetry of original differential equations. In *Modern Group Analysis: Advanced Analytical and Computational Methods in Mathematical Physics*, volume 191. Kluwer Academic Publishers, Boston, 1993.
- [13] V. Dorodnitsyn, R. Kozlov, and P. Winternitz. Continuous symmetries of lagrangians and exact solutions of discrete equations. J. Math. Phys., 45(1):336–359, 2004.
- [14] Decio Levi and Pavel Winternitz. Continuous symmetries of difference equations. Journal of Physics A: Mathematical and General, 39(2):R1–R63, dec 2005.

- [15] V. Dorodnitsyn, R. Kozlov, and P. Winternitz. Lie group classification of second-order ordinary difference equations. J. Math. Phys., 41(1):480–504, 2000.
- [16] G. R. W. Quispel, H. W. Capel, and R. Sahadevan. Continuous symmetries of differentialdifference equations. *Phys. Lett.*, 170A:379–383, 1992.
- [17] V. A. Dorodnitsyn. Applications of Lie Groups to Difference Equations. CRC Press, Boca Raton, 2011.
- [18] P. E. Hydon. *Difference Equations by Differential Equation Methods*. Cambridge University Press, Cambridge, 2014.
- [19] V.A.Dorodnitsyn, E.I.Kaptsov, R.V.Kozlov, and P.Winternitz. The adjoint equation method for constructing first integrals of difference equations. Journal of Physics A: Mathematical and Theoretical, 48(5):055202, 01 2015.
- [20] V. A. Dorodnitsyn and E. I. Kaptsov. Shallow water equations in Lagrangian coordinates: Symmetries, conservation laws and its preservation in difference models. Communications in Nonlinear Science and Numerical Simulation, 89:105343, 2020.
- [21] V. A. Dorodnitsyn and E. I. Kaptsov. Discrete shallow water equations preserving symmetries and conservation laws. Journal of Mathematical Physics, 2021, J. Math. Phys.
- [22] V. A. Dorodnitsyn, E. I. Kaptsov, and S. V. Meleshko. Symmetries, conservation laws, invariant solutions and difference schemes of the one-dimensional Green-Naghdi equations. Journal of Nonlinear Mathematical Physics, Vol. 28(1); (2021), pp. 90-107.
- [23] V. A. Dorodnitsyn, R. V. Kozlov, and P. Winternitz. Continuous symmetries of Lagrangians and exact solutions of discrete equations. Journal of Mathematical Physics, 45(1):336-359, 2004.
- [24] A.F.Cheviakov, V.A.Dorodnitsyn, E.I.Kaptsov. Invariant conservation law-preserving discretizations of linear and nonlinear wave equations. Journal of Mathematical Physics, 61(8):081504, 2020.
- [25] Dorodnitsyn V.A. and Kaptsov E.I., "Invariant Finite-Difference Schemes for Plane One-Dimensional MHD Flows That Preserve Conservation Laws" Mathematics, MDPI, 2022.
- [26] Dorodnitsyn V., Kaptsov E., Meleshko S., Conservative Invariant Finite-Difference Schemes for the Modified Shallow Water Equations in Lagrangian Coordinates, Studies in Applied Mathematics, 2022.
- [27] Dorodnitsyn V. A., Kaptsov E. I., Invariant finite-difference schemes for plane onedimensional MHD flows that preserve conservation laws. Mathematics, 2022.
- [28] S. V. Meleshko. Methods for Constructing Exact Solutions of Partial Differential Equations. Mathematical and Analytical Techniques with Applications to Engineering. Springer, New York, 2005.
- [29] Yu. N. Grigoriev, N. H. Ibragimov, V. F. Kovalev, and S. V. Meleshko. Symmetries of integro-differential equations and their applications in mechanics and plasma physics. Lecture Notes in Physics, Vol. 806. Springer, Berlin / Heidelberg, 2010.

- [30] V. A. Dorodnitsyn, R. Kozlov, S. V. Meleshko, and P. Winternitz. Lie group classification of first-order delay ordinary differential equations. *Journal of Physics A: Mathematical* and Theoretical, 51(205202), 2018.
- [31] V. A. Dorodnitsyn, R. Kozlov, S. V. Meleshko, and P. Winternitz. Linear or linearizable first-order delay ordinary differential equations and their lie point symmetries. *Journal of Physics A: Mathematical and Theoretical*, 51(205203), 2018.
- [32] V. A. Dorodnitsyn, R. Kozlov, S. V. Meleshko, and P. Winternitz. Second-order delay ordinary differential equations, their symmetries and application to a traffic problem. *Journal* of Physics A: Mathematical and Theoretical. IOP Publishing Ltd Volume 54, Number 10, 2021.
- [33] Polyanin A.D., Zhurov A.I., Exact separable solutions of delay reaction-diffusion equations and other nonlinear partial functional-differential equations, Commun. Nonlinear Sci. Numer. Simul., 2014, Vol. 19, 409-416.
- [34] Polyanin A.D., Zhurov A.I., Nonlinear delay reaction-diffusion equations with varying transfer coefficients: Exact methods and new solutions, Appl. Math. Lett., 2014, Vol. 37, 43-48.
- [35] Polyanin A.D., Sorokin V.G., Reductions and exact solutions of nonlinear wave-type PDEs with proportional and more complex delays, Mathematics, 2023, Vol. 11, 516.
- [36] L. E. Elsgolts. Qualative Methods in Mathematical Analysis. GITTL, Moscow, 1955. Translation: American Mathematical Society, Providence ,1964.
- [37] V. A. Dorodnitsyn. The finite-difference analogy of Noether's theorem. Doklady RAN, 328(6):678–682, 1993. Translation in English in Phys. Dokl. 38 (2) 66–68.
- [38] S. F. Frederico and F. M. Torres. Noether's symmetry theorem for variational and optimal control problems with time delay. *Numerical Algebra, Control and Optimization (NACO)*, 2(3):619–630, 2012.
- [39] Agnieszka B. Malinowska and Tatiana Odzijewicz (2016): Second Noether's theorem with time delay, Applicable Analysis, DOI: 10.1080/00036811.2016.1192136
- [40] B. L. Rozhdestvenskii and N. N. Yanenko. Systems of quasilinear equations and their applications to gas dynamics, 2nd ed. Nauka, Moscow, 1978. English translation published by Amer. Math. Soc., Providence, RI, 1983.
- [41] V. A. Dorodnitsyn and R. Kozlov. Lagrangian and Hamiltonian formalism for discrete equations: Symmetries and first integrals. In Symmetries and Integrability of Difference Equations, pages 7–49. Cambridge University Press, Cambridge, 2011. London Mathematical Society Lecture Notes.
- [42] V. Dorodnitsyn and N.Ibragimov. An extension of the Noether theorem: accompanying equations possessing conservation laws. Communications in Nonlinear Science and Numerical Simulation, 2014, v.19, N2, 328-336.
- [43] A. Gonzalez-Lopez, N. Kamran, and P. J. Olver. Lie algebras of vector fields in the real plane. Proc. London Math. Soc., 64:339–368, 1992.

- [44] E. Bessel-Hagen. Über die Erhaltungssatze der Elektrodynamik. Math. Ann., 84:258–276, 1921.
- [45] L. E. Elsgolts. Variational problems with retarded arguments. Vestnik Moskovskogo Universiteta. Seriya 1. Matematika. Mekhanika, 10:57–62, 1952. (in Russian).
- [46] L. E. Elsgolts. Variational problems with retarded arguments. Uspekhi Matematicheskikh Nauk, 12(1(73)):257–258, 1957. (in Russian).