On Structural and Spectral Properties of Distance Magic Graphs

Himadri Mukherjee¹, Ravindra Pawar¹, and Tarkeshwar Singh¹

¹Department of Mathematics, BITS Pilani K K Birla Goa Campus, Goa, India.

Abstract

A graph G = (V, E) is said to be distance magic if there is a bijection f from a vertex set of G to the first |V(G)| natural numbers such that for each vertex v, its weight given by $\sum_{u \in N(v)} f(u)$ is constant, where N(v) is an open neighborhood of a vertex v. In this paper, we introduce the concept of p-distance magic labeling and establish the necessary and sufficient condition for a graph to be distance magic. Additionally, we introduce necessary and sufficient conditions for a connected regular graph to exhibit distance magic properties in terms of the eigenvalues of its adjacency and Laplacian matrices. Furthermore, we study the spectra of distance magic graphs, focusing on singular distance magic graphs. Also, we show that the number of distance magic labelings of a graph is, at most, the size of its automorphism group.

Keywords: Distance magic labeling, Laplacian Matrix, Moore-Penrose Inverse, *p*-distance magic labeling.

AMS Subject Classification 2021: 05C78.

1 Introduction

Throughout this article, we assume that G = (V, E) denotes a simple, undirected, finite graph on *n* vertices. Let $N(u) = \{v \in V : uv \in E\}$ be the neighborhood of a vertex $u \in V(G)$. We denote the degree of a vertex *v* by deg(v) and is given by deg(v) = |N(v)|. Let K_n denote a complete graph on *n* vertices.

A bijective function $f: V(G) \to \{1, 2, ..., n\}$ is called the *vertex labeling* of a graph G. If the sum $\sum_{u \in N(v)} f(u)$ is constant k, for all $v \in V$, then the vertex labeling f is called a *distance* magic labeling, k is called a magic constant. Any graph G that admits the distance magic labeling is called a *distance* magic graph. It is known that the magic constant of a graph

himadrim@goa.bits-pilani.ac.in

p20200020@goa.bits-pilani.ac.in

tksingh@goa.bits-pilani.ac.in

G, if it exists, is unique, that is, independent of distance magic labeling [2, 12]. Since the general characterization of a graph as distance magic is not known, researchers have studied the existence of distance magic labeling for many families of graphs [5]. Although there are numerous necessary conditions for a graph to possess a distance magic labeling [2, 12, 15], the elusive sufficient condition for this phenomenon remains undiscovered. In this paper, we give several necessary and sufficient conditions for a graph to be a distance magic.

A multiset is a set in which repetition of elements is allowed. We formally define the multiset of our interest.

Definition 1.1. Given an integer $p \ge 1$, by $\{1, 2, ..., n\}_p$, we mean a *multiset* obtained by reducing all numbers 1, 2, ..., n to modulo p and replacing 0s if any by p.

For example, with p = 4 and n = 9, we have the multiset $\{1, 2, 3, 4, 1, 2, 3, 4, 1\}_4$ obtained by reducing the elements 1, 2, 3, 4, 5, 6, 7, 8, 9 to modulo 4 and replacing 0s by 4. Note that for $p > n, \{1, 2, ..., n\}_p = \{1, 2, ..., n\}$.

We introduce the concept of p-distance magic labeling as a generalization of distance magic labeling of graphs.

Definition 1.2. Let G be a graph on n number of vertices and given an integer $p \ge 1$. Consider a multiset $T = \{1, 2, ..., n\}_p$. We call graph G a p-distance magic if there is a bijective map f from the vertex set V to a multiset T called a p-distance magic labeling such that the weight of each vertex $x \in V$, denoted by w(x), is equal to the same element $\mu(\text{mod } p)$, where $w(x) = \sum_{y \in N(x)} f(y)$.

For example, consider a graph G in Figure 1 on 11 vertices. With p = 2, we have a multiset $\{1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2,$




Figure 1: A 2-distance magic labeling of a graph

The *p*-distance magic labeling also constitute an infinite class of necessary conditions that become sufficient when true for all p. In the context of a graph with n vertices, we simplify the labeling process by reducing the labels $1, 2, \ldots, n$ modulo p, creating a multiset where

the repetition of elements is allowed. We then calculate the weights modulo p in the usual manner. This approach reduces the number of unique labels used, allowing the calculations to be streamlined. It is important to emphasize that p-distance magic labeling differs from group distance magic labeling or Γ -distance magic labeling [4], particularly when $\Gamma = \mathbb{Z}_n$. In the former, the requirement for the number of distinct labels to equal the number of vertices is relaxed.

It is well known that there cannot be a subgraph avoidance criterion for the graphs that admit magic labeling (since any connected graph can be embedded in a distance magic graph [7]). As a result, the problem of magic labeling does not allow for the construction of magic labeling from the labeling on a subgraph. One can say that the problem of distance magic labeling is a difficult problem with little room for "construction" in the underlying graph because it does not allow "cutting and pasting" type techniques. However, *p*-distance magic labeling allows such cutting and pasting techniques, as is evident from applying the Chinese remainder theorem (see Theorem 2.5). The simplification through the existence of a *p*-magic labeling and the patching up of two different such labeling through the Chinese remainder theorem gives us some constructive approach. Such possibilities give us hope that to tackle the problem of distance magic labeling, this "construction" idea through *p*-distance magic labeling will prove to be very important. With sufficiently larger values of *p*, a *p*-distance magic labeling becomes \mathbb{Z}_n -distance magic labeling, and with p = n, *p*-distance magic labeling becomes \mathbb{Z}_n -distance magic labeling.

We explore certain linear combinations of matrices associated with distance magic graphs to establish a condition that is necessary and sufficient for a graph to be considered distance magic. Specifically, we prove that if a graph G possesses a distance magic property with a magic constant k, then k is an eigenvalue of a particular linear transformation associated with G. Furthermore, a connected r-regular graph is distance magic if and only if r^2 is an eigen value of $L^2 + A^2$, where A is an adjacency matrix and L is the Laplacian matrix of G. Lastly, we obtain one necessary condition by proving that if a graph is distance magic, then AA^{\dagger} is doubly stochastic, where A is the adjacency matrix of G and A^{\dagger} is a Moore-Penrose inverse of A.

We have established bounds on the number of distance magic labelings of a distance magic graph based on an automorphism group of the graph. Moreover, we present several examples of distance magic graph where the number of distance magic labelings matches the cardinality of the automorphism group.

As mentioned earlier, forbidden subgraph characterisation is not possible for distance magic graphs. However, we have the existence of some subgraphs as a necessary condition for a graph to be distance magic. It can be proved easily using the following theorem.

Theorem 1.1. [7, 15] A graph G is not distance magic if there are vertices x, y in G such that $|N(x) \triangle N(y)| = 1$ or 2.

Theorem 1.2. Every distance magic graph contains P_3 or C_4 as a subgraph or P_3 as an induced subgraph.

Proof. Let G be a distance magic graph. Then by Theorem 1.1, for any two vertices x, y in G we must have $|N(x) \triangle N(y)| = 0$ or $|N(x) \triangle N(y)| \ge 3$. That is, for any two vertices

in G, their neighborhoods are identical or symmetric difference of their neighborhoods is at least 3. Consider the following two cases.

Case i. Suppose there exists two vertices x, y such that N(x) = N(y). If deg(x) = 1 (and hence deg(y) = 1) then there is an unique vertex z adjacent to both x and y. Therefore, there is a path P_3 induced by vertices x, y, and z.

Case ii. Suppose that $N(x) \neq N(y)$ for any two vertices x, y in G. Then, for every pair of vertices x, y in G, we have $|N(x) \triangle N(y)| \ge 3$. That is $|N(x) \cup N(y)| - |N(x) \cap N(y)| \ge 3$. Note that $|N(x) \cap N(y)| = 0$ for all $x, y \in V(G)$ is not possible. Otherwise, G is a disjoint union of K_2 components, which is not distance magic. Therefore, there exist two vertices x, y in G such that $|N(x) \cap N(y)| \ge 1$. Let $|N(x) \cap N(y)| = 1$. If $N(x) \cap N(y) = \{z\}$ then G contains P_3 as a subgraph. Note that P_3 need not be an induced subgraph on the vertices x, y, and z. If $|N(x) \cap N(y)| \ge 2$ then G contains a C_4 (and hence P_3) as a subgraph.

For a set $D \,\subset \mathbb{N}$, let $ND(v) = \{u \in V(G) : d(u, v) \in D\}$, where d(u, v) denotes a distance between vertices u and v. Let W be a multiset of real numbers. A graph G is said to be (D, W)-vertex magic if there exists a bijection $g : V(G) \to W$ such that for all $v \in V(G)$, $\sum_{u \in N(v)} g(u)$ is a constant [12]. In [14], Slater proved that the problem of determining whether a graph is (D, W)-vertex magic is NP-complete. Also, it is known that this problem is NP-complete in special classes of graphs like 3-regular graphs where labeling set is a multiset with entries from the set of positive integers [6]. This shows that the problem of distance magic labeling is NP-complete when labeling set is a multiset with entries from the set of positive integers.

Organisation of paper:

In Section 2, we present the results on p-distance magic labeling and the necessary and sufficient conditions for a graph to be considered distance magic in terms of p-distance magic labeling. In section 3, we show that there are infinitely many distance magic graphs having 0 as an eigenvalue. Also, we derive the conditions for a graph to be considered distance magic based on eigenvalues of its adjacency and Laplacian matrices. In Section 4, we analyse the action of the automorphism group of a graph on the set of its distance magic labelings. We estimate the lower bounds on the number of distance magic labelings for the graph.

For undefined terminology and notation related to graph theory, we refer to [16], and for the survey on distance magic labeling, we refer to [1, 5].

2 *p*-distance magic graphs

For a distance magic graph, G of order n with distance magic labeling f and the magic constant k, the sum of the weights of all vertices is nk (see [11,15]). That is $\sum_{x \in V(G)} w(x) = \sum_{x \in V(G)} f(x) deg(x) = nk$. The maximum possible degree in G is n-1. Therefore, we obtain $nk \leq (n-1) \sum_{x \in V(G)} f(x) = \frac{(n-1)n(n+1)}{2}$. This implies, $k \leq \frac{n^2-1}{2}$.

Theorem 2.1. A graph G is distance magic if and only if it is p-distance magic for infinitely many values of p.

Proof. Suppose that G is a distance magic graph of order n with magic constant k and distance magic labeling f. Let S be an infinite subset of those natural numbers p for which G is p-distance magic. Let $p \in S$ be arbitrary. Define $f_p : V \to \{1, 2, \ldots, n\}_p$ by $f_p(v) = f(v) \pmod{p}$. Therefore, the weight $w_{f_p}(x)$ of each vertex x in V is equal to the same number $k_p \equiv k \pmod{p}$. This proves the necessary part. Conversely, suppose that G is p-distance magic for all $p \in S$. The set S being infinite guarantees the existence of q in S such that $q > \frac{(n^2-1)}{2}$ and G induce a q-distance magic labeling. Then w(x) = k' for some $k' \in \mathbb{Z}_q$ and $k' \leq \frac{(n^2-1)}{2} < q$. Therefore, q-distance magic graph.

Corollary 2.1.1. A graph G on n vertices is distance magic if and only if it is p-distance magic for all $p \ge 1$.

Corollary 2.1.2. A graph G on n vertices is distance magic if and only if it is p-distance magic for all prime numbers p.

It follows from the Theorem 2.1 that the *p*-distance magic labeling coincides with distance magic labeling whenever p is sufficiently large. This shows that *p*-distance magic labeling is a generalization of distance magic labeling. Also, some structural characterisations are possible with p = 2. Since p = 2 is the least possible, we focus on 2-distance magic labelings.

Let G be a graph whose vertices are labeled with the labeling f using the numbers in multiset $T = \{1, 2, ..., n\}_p$ for some $p \ge 1$. Let $V_i = \{v \in V : f(v) \equiv i \pmod{p}\}$, for all i = 1, 2, ..., p. By G_i , we mean a subgraph of G induced on V_i .

Recall that a graph G is *Eulerian* if and only if it has at most one non-trivial component and its vertices all have even degrees, and a *matching* in a graph is an independent subset of a set of edges.

Theorem 2.2. Let G be a graph on n vertices that admits 2-distance magic labeling. If

- 1. a magic constant is 0, then the subgraph G_1 is a union of Eulerian components.
- 2. a magic constant is 1 then the subgraph G_1 contains a matching.

Proof. Suppose that a graph G is 2-distance magic. We prove the theorem in the following two cases.

Case *i*. Let magic constant be 0. If G itself is an Eulerian, there is nothing to prove. Suppose G is not Eulerian. For a magic constant to be 0, each vertex in G_1 must have an even number of neighbors in V_1 and any number of neighbors in V_2 . Therefore, each vertex in G_1 is of even degree. Therefore, G_1 is a union of Eulerian components.

Case ii. Let magic constant be 1. Let $v \in V_1$. For w(v) = 1, v must be adjacent to the odd number of vertices in V_1 . Now, we construct a matching, say M. Let $v_1 \in V_1$ such that $vv_1 \in E(G_1)$. We add this edge vv_1 to the matching M. We repeat the process, which must end after a finite number of steps since G_1 is a finite graph. Therefore, we obtain a maximal matching M. This proves the theorem.

Figure 2 shows the 2-distance magic labeling of two graphs. A graph in Figure 2(a) has magic constant 0, and the subgraph G_1 induced by V_1 is depicted in dotted edges and hollow




Figure 2: 2-magic graphs with an Eulerian subgraph or a matching.

vertices is an Eulerian subgraph. A graph in Figure 2(b) has a magic constant 1, and G_1 contains a matching depicted in dotted edges. Note that in Figure 2(b), matching in G_1 is not unique, but it is a perfect matching. However, it is not the case that we always obtain a perfect matching. Also, in Figure 2(b), none of the vertices in G_1 are of even degree. Therefore, obtaining an Eulerian circuit is not possible.

Now we introduce the concept *modulo regularity* of graphs and use to construct different *p*-distance magic labelings from the given *p*-distance magic labeling of a graph.

Definition 2.1. We call graph G an r-modulo p regular if $deg(u) \equiv r \pmod{p}$ and we write G is $r \pmod{p}$ -regular.

Theorem 2.3. If G is $r \pmod{p}$ -regular, p-distance magic graph on n vertices with magic constant k and p-distance magic labeling f, then for each positive integer i, the map f' = f + i is also a p-distance magic labeling with magic constant $k' = (k + ir) \pmod{n}$.

Proof. Let G be $r \pmod{p}$ -regular, p-distance magic graph with magic constant k and labeling f and let $i \in \mathbb{Z}^+$. Define p-distance magic labeling g by g(u) = i + f(u). Then, for $u \in V$,

$$w(u) = \sum_{uv \in E} g(v)$$

= $\sum_{uv \in E} (i + f(v))$
= $i \ deg(u) + \sum_{uv \in E} f(v)$
= $(ir + k) \pmod{p}$.

This completes the proof.

Theorem 2.3 shows that a *p*-magic constant need not be unique. Nevertheless, in some cases, one can obtain the uniqueness of the same as described in the following theorem. (The ring \mathbb{Z}_n is considered with the usual addition and multiplication modulo n.)

Theorem 2.4. Let G be a graph on n vertices. If a graph G is a p-distance magic with magic constant k and if $\frac{n(n+1)}{2}$ is a unit in the ring \mathbb{Z}_p , then k is unique.

Proof. Let G be a graph on n vertices $\{x_1, x_2, \ldots, x_n\}$ having two p-distance magic labelings f and g with respective magic constants k and l. Let 1 be the column vector with all n entries 1. Let A be an adjacency matrix of G and put $X = (f(x_1), \ldots, f(x_n))^{\mathsf{T}}$ and $Y = (g(x_1), \ldots, g(x_n))^{\mathsf{T}}$. Since f and g are distance magic labelings with magic constants k and l, it follows that AX = k1 and AY = l1. Since, $X^{\mathsf{T}}AY$ is 1×1 matrix, we have $X^{\mathsf{T}}AY = (X^{\mathsf{T}}AY)^{\mathsf{T}} = Y^{\mathsf{T}}AX$. This gives

$$lX^{\mathsf{T}}\mathbb{1} = kY^{\mathsf{T}}\mathbb{1}$$
$$\implies l(1+2+\dots+n) = k(1+2+\dots+n).$$

Since $(1 + 2 + \dots + n)$ is a unit in \mathbb{Z}_p , by the cancellation laws we have l = k.

Corollary 2.4.1. If G is a 2-distance magic graph of order $n \ge 3$, where $n \equiv 1$ or $2 \pmod{4}$ with magic constant k, then k is unique.

Proof. Let G be 2-distance magic graph of order $n \ge 3$. Suppose $n \equiv 1 \pmod{4}$, that is, n = 4j + 1, for some integer j. Then $\frac{n(n+1)}{2} = (4j + 1)(2j + 1)$ which is unit in \mathbb{Z}_2 . Similarly, when n = 4j + 2, for some integer j, then $\frac{n(n+1)}{2} = (2j + 1)(4j + 3)$ which again an unit in \mathbb{Z}_2 . This completes the proof.

Theorem 2.5. If G is a p-distance magic as well as q-distance magic graph on n vertices for some relatively prime integers p and q such that $pq \le n$, then G is pq-distance magic.

Proof. Let G be a graph with vertex set $\{x_1, x_2, \ldots, x_n\}$ that is both p-distance magic and q-distance magic for some relatively prime integers p and q such that $pq \leq n$. Let f_p, f_q be a p-distance and q-distance magic labelings, and k_p, k_q be corresponding magic constants. For each $1 \leq i \leq n$, let $f_p(x_i) \equiv a_i \pmod{p}$ and $f_q(x_i) \equiv b_i \pmod{q}$. Since p and q are co-prime, by the Chinese remainder theorem, the system of congruences

$$x \equiv a_i \pmod{p}$$
$$x \equiv b_i \pmod{q}$$

has unique solution say $y_i \pmod{pq}$, for each $i(1 \le i \le n)$. Now we define new labeling f_{pq} by $f_{pq}(x_i) = y_i$, for each $i(1 \le i \le n)$. The uniqueness of the labels y_i s guarantees that f_{pq} is a bijective map. Now we calculate the weight of a vertex $x_i \in V$ under f_{pq} .

$$w(x_i) = \sum_{x_j \in N(x_i)} f_{pq}(x_j)$$
$$\equiv \begin{cases} k_p \pmod{p} \\ k_q \pmod{q}. \end{cases}$$

For each $1 \leq i \leq n$, again we solve the system of congruences

$$w(x_i) \equiv k_p \pmod{p}$$

$$w(x_i) \equiv k_q \pmod{q}$$
(1)

by the Chinese remainder theorem, and we conclude that the system of congruences (1) has a unique solution, denoted by $k_{pq} \pmod{pq}$. This proves that f_{pq} is the required labeling and G is a pq-distance magic graph.

Given a graph G on n vertices which is both p-distance magic and q-distance magic for some relatively prime integers p and q such that $pq \leq n$ then using the Theorem 2.5, we can find pq-distance magic labeling for G. When pq > n, we need not always obtain a pq-distance magic graph. For example, consider a cycle $C_4 = v_1, v_2, v_3, v_4, v_1$. Define 2-magic labeling of C_4 by $f_2(v_1) = 1$, $f_2(v_2) = 2$, $f_2(v_3) = 2$, $f_2(v_4) = 1$ and a 3-magic labeling by $f_3(v_1) = 2$, $f_3(v_2) = 1$, $f_3(v_3) = 3$, $f_3(v_4) = 1$ as shown in Figure 3. In Figure 3, the graph G_1 shows the 2-distance magic labeling of C_4 with magic constant 1, and the graph G_2 shows a 3-distance magic labeling with magic constant 2 of the same graph C_4 . For each $i(1 \leq i \leq 4)$, we solve the system

$$x \equiv f_2(x_i) \pmod{2}$$

$$x \equiv f_3(x_i) \pmod{3}$$
(2)

using the Chinese remainder theorem. For each $i(1 \le i \le 4)$, the unique solution of the system (2) is 2, 4, 6, 1 respectively. We label the vertices of C_4 using these solutions: $f_6(v_1) = 2$, $f_6(v_2) = 4$, $f_6(v_3) = 6$, $f_6(v_4) = 1$ as shown in the graph G_3 of Figure 3. Observe that f_6 is not a map from $V(C_4)$ to $\{1, 2, 3, 4\}_6$. Thus, in this case, we cannot obtain the 6-distance magic graph using the given 2-distance magic and 3-distance magic graph. This does not contradict the Theorem 2.5 because $(p = 2) \times (q = 3) = 6 > 4 = n$. Therefore, condition pq < n cannot be ignored in the statement of Theorem 2.5. However, it may happen that in some cases, each label $y \pmod{pq}$ obtained using the Chinese remainder theorem as described in the above theorem satisfies $1 \le y \le n$. We call such labeling a *consistent* labeling. In such cases, the labeling is indeed a pq-distance magic labeling. We state the following proposition




Figure 3: *p*-distance magic labelings of C_4

without proof, as its proof is quite similar to Theorem 2.5.

Proposition 2.1. If G is a p-distance magic as well as a q-distance magic graph on n vertices for some relatively prime integers p and q such that pq > n and if the new labeling obtained as described in the proof of Theorem 2.5 is consistent then the graph G is pq-distance magic.

3 Spectra of the linear operators related to distance magic graphs

For a given graph G, let $\mu_1, \mu_2, \ldots, \mu_t$ be distinct eigenvalues of its adjacency matrix A. We call μ an eigenvalue of G whenever it is an eigenvalue of its adjacency matrix A. A graph is considered *singular* if 0 is the eigenvalue of G and it is not singular otherwise. We always take eigenvalues in non-increasing order $\mu_1 \ge \mu_2 \ge \cdots \ge \mu_t$. The largest eigenvalue that is, μ_1 , is called *index* of G. The eigenvalue is called *simple* if its algebraic multiplicity is 1. As the adjacency matrix A of a graph G is real symmetric, it is diagonalisable. Therefore, its eigenvectors form an orthonormal basis of \mathbb{R}^n . Let $\mathcal{E}(\mu_i)$ be the eigenspace corresponding to the eigenvalue μ_i . For $a \in \mathbb{R}^n$, let $P_i(a) = \sum_b \langle a, b \rangle b$, where b is an orthonormal eigenvector corresponding to the eigenvalue μ_i and \langle , \rangle is a standard inner product on \mathbb{R}^n . Let $\mathbb{1}$ be a column vector whose all entries are 1. The *main angles* of G, denoted by β_i are given by $\beta_i = \frac{\|P_i\mathbb{1}\|}{\sqrt{n}}$. Let J_n be an $n \times n$ matrix whose all entries are 1.

Let G be a distance magic graph on n vertices v_1, v_2, \ldots, v_n with distance magic labeling f and the magic constant k. Let $x = (f(x_1), f(x_2), \ldots, f(x_n))^{\mathsf{T}}$. Define a transformation $T : \mathbb{R}^n \to \mathbb{R}^n$ by $T(e_i) = e_i + e_{i+1} + \cdots + e_n$, where e_1, e_2, \ldots, e_n is a standard basis of \mathbb{R}^n . Then, for some suitable permutation matrix P, we have $x = PT\mathbb{1}$. We assume these notations throughout this section.

The following theorem gives the necessary and sufficient condition for a graph to be distance magic in terms of the eigenvalues of the linear transformation APT.

Theorem 3.1. A graph G is distance magic if and only if 1 is an eigenvector of APT corresponding to the eigenvalue k for some suitable permutation matrix P.

Proof. Let G be the distance magic graph. Then, we have $Ax = k\mathbb{1}$. Choose P such that $PT\mathbb{1} = x$. This implies $APT\mathbb{1} = Ax = k\mathbb{1}$ as required. Conversely, suppose that APT has an eigenvalue k corresponding to eigenvector $\mathbb{1}$. Then $APT\mathbb{1} = k\mathbb{1}$. Taking $x = PT\mathbb{1}$, we get $Ax = k\mathbb{1}$. This proves that G is a distance magic graph.

It is known that odd regular graphs can not be distance magic [11]. In 2-regular graphs, only disjoint unions of C_4 is distance magic [11]. Distance magic labeling of C_4 is shown in Figure 5. Miklavič et al. [9, 10] gave a characterisation of 4-regular and 6-regular graphs. They proved the following general result based on a spectrum of a graph, giving necessary and sufficient conditions for the even regular graph to be distance magic.

Lemma 3.2. [9] Let G be an even regular graph of order n. Then G is distance magic if and only if 0 is an eigenvalue of G and there exists an eigenvector w for the eigenvalue 0 with the property that a certain permutation of its entries results in the arithmetic sequence

$$\frac{1-n}{2}, \frac{3-n}{2}, \frac{5-n}{2}, \dots, \frac{n-1}{2}.$$

In particular, if G is distance magic, then 0 is an eigenvalue for G, and there exists a corresponding eigenvector, all of whose entries are pairwise distinct.

Let G be a graph with vertex set $\{v_1, v_2, \ldots, v_n\}$. Let $D = diag(deg(v_1), deg(v_2), \ldots, deg(v_n))$ be a degree diagonal matrix of G. Laplacian matrix of a graph G is given by L = D - A. Now we prove another simple necessary and sufficient condition for connected even regular graphs to be distance magic in terms of the eigenvalues of the matrix $L^2 + A^2$.

Theorem 3.3. Let G be a connected r-regular graph, where r is even. A graph G is distance magic graph with distance magic labeling f if and only if $(L^2 + A^2)x = r^2x$, where $x = (f(x_1), \ldots, f(x_n))^{\intercal}$.

Proof. Let G be connected r-regular distance magic graph with distance magic labeling f and magic constant k. Then $Ax = k\mathbb{1}$. We know L = D - A. Hence,

$$L^{2} + A^{2} = D^{2} - 2rA + 2A^{2} = r^{2}I - 2rA + 2A^{2}.$$
(3)

Then

$$(L^{2} + A^{2})x = (r^{2}I - 2rA + 2A^{2})x$$

= $r^{2}x - 2rk\mathbb{1} + 2rk\mathbb{1}$
= $r^{2}x$.

Conversely suppose that $(L^2 + A^2)x = r^2x$. Then, after expanding the left-hand side as in Equation (3) and canceling out the terms, we obtain $A^2x = rAx$, that is, (A-rI)Ax = 0. Now, by the regularity of G, the sum of entries of each row in A - rI is 0, and by connectedness of G, r-1 is an eigenvalue of A - rI of multiplicity 1. Therefore, $\{1\}$ forms a basis for $\mathcal{E}(r-1)$. Note that each entry in Ax is an integer. Therefore, we must have Ax = k1 for some positive integer k. Therefore, G is a distance magic graph.

Now we prove that there are infinitely many singular distance magic graphs. Recall that the *cone cover* of a graph G, denoted by $K_1 \bigtriangledown G$, is a graph obtained by adding a vertex to G and joining it to all other vertices of G. Kamatchi [8] characterised the distance magic graphs G with $\Delta(G) = n - 1$.

Theorem 3.4. [8] Let G be any graph of order n with $\Delta = n - 1$. Then G is a distance magic graph if and only if n is odd and G is isomorphic to the cone cover $K_1 \bigtriangledown (K_{n-1} - M)$ of $K_{n-1} - M$, where M is a perfect matching in K_{n-1} .

Theorem 3.5. For any positive integer $n \ge 3$, there is a singular distance magic graph of order n.

Proof. Let G be a graph of order n.

Case i: Let $n \ge 4$ be even and let $V(G) = \{1, 2, ..., n\}$. We may assume that the label of the vertex *i* is *i*. For each $1 \le i \le n$, let $N(i) = V(G) - \{i, n+1-i\}$. Hence $w(i) = \frac{n(n+1)}{2} - (n+1)$ for all i = 1, 2, ..., n. Therefore, we conclude that *G* is a distance magic graph. Also, by construction, it is clear that N(1) = N(n). Therefore, the rows corresponding to 1 and *n* in

the adjacency matrix of G are identical. This proves that G must be singular.

Case ii: Let $n \ge 3$ be odd. For n = 3, we know that P_3 is distance magic (see Figure 5) and it is a singular graph. Suppose n > 3. Then there is a distance magic graph $G = K_{n-1} - M + K_1$, where M is a perfect matching in K_{n-1} . If $uv \in M$ then $N_{K_{n-1}}(u) = N_{K_{n-1}}(v)$. Therefore, $N_G(u) = N_G(v)$. Proving that G is singular.

A graph is said to *integral* if all of its eigenvalues are integers and it is not integral otherwise. We shortly prove that the cone cover of $K_{n-1} - M$ for odd n is not integral.

We will calculate the characteristic polynomial of such graphs. There is a relation between the characteristic polynomial of G and its cone cover in terms of the eigenvalues and the main angles of G:

Theorem 3.6. [3] The cone cover of G has characteristic polynomial $P_{K_1 \nabla G}(y) = P_G(y) \left(y - \sum_{i=1}^k \frac{n\beta_i^2}{y - \mu_i}\right)$, where μ_i and β_j are eigen values and main angles of G respectively.

Therefore, to calculate the characteristic polynomial of $K_1 \bigtriangledown (K_{n-1} - M)$, we need the information about the spectra and main angles of $K_{n-1} - M$.

Proposition 3.1. Let $n \ge 4$ be even, and let K_n be the complete graph on the *n* vertices 1, 2, ..., n. Let *M* be a perfect matching in K_n such that if $e \in M$ then $e = (i, i + \frac{n}{2})$. Then

spec
$$(K_n - M) = \begin{pmatrix} n-2 & 0 & -2 \\ 1 & \frac{n}{2} & \frac{n}{2} - 1 \end{pmatrix}.$$

Proof. Deletion of a maximum matching in a complete graph K_n $(n \ge 4)$ does not disconnect the graph. Therefore $G = K_n - M$ is a connected (n-2)-regular graph. Therefore, (n-2)is a simple eigenvalue of G. Also, in G, $N(i) = N(i + \frac{n}{2})$. Therefore, the corresponding rows in the adjacency matrix are identical, and hence, the eigenvalue 0 has multiplicity $\frac{n}{2}$. All $\frac{n}{2}$ eigenvectors in the basis of $\mathcal{E}(0)$ are given by v_j , where for each $1 \le j \le \frac{n}{2}$, ith coordinate of the vector v_j is given by

$$(v_j)_i = \begin{cases} -1 & \text{if } i = j \\ 1 & \text{if } i = j + \frac{n}{2} \\ 0 & \text{otherwise.} \end{cases}$$

For each $2 \leq j \leq \left(\frac{n}{2}\right)$, consider a vector w_j whose *i*th coordinate is given by

$$(w_j)_i = \begin{cases} -1 & \text{if } i = 1, \frac{n}{2} + 1\\ 1 & \text{if } i = j, \frac{n}{2} + j\\ 0 & \text{otherwise.} \end{cases}$$

It is easy to verify that these $\frac{n}{2}-1$ vectors w_j are linearly independent and are the eigenvectors corresponding to eigenvalue -2. Therefore, the multiplicity of -2 is at least $\frac{n}{2}-1$. We already found $\frac{n}{2}+1$ other eigenvalues of G. Therefore algebraic multiplicity of the eigenvalue -2 must be $\frac{n}{2}-1$. This completes the proof.

From the spectra of the graph $K_n - M$ as discussed previously, we can write its characteristic polynomial, $P_{K_n-M}(y) = y^{\frac{n}{2}}(y+2)^{\frac{n}{2}-1}(y-n+2)$. Now we calculate the main angles β_i of $K_n - M$.

Let $y = (y_1, y_2, \ldots, y_n)^{\mathsf{T}} \in \mathbb{R}^n$. Recall that to calculate $P_i(y)$ we take orthonormal vectors from $\mathcal{E}(\mu_i)$. The only eigenvector corresponding to eigenvalue (n-2) is 1. Therefore,

$$P_1(y) = \left(y, \frac{1}{\sqrt{n}}\right) \frac{1}{\sqrt{n}}$$

= $\frac{1}{n} (y_1 + y_2 + \dots + y_n, y_1 + y_2 + \dots + y_n, \dots, y_1 + y_2 + \dots + y_n).$

Therefore, the matrix representation of P_1 with respect to the standard basis of \mathbb{R}^n is the matrix J_n . Therefore, $||P_1\mathbb{1}|| = 1$ and hence $\beta_1 = \frac{1}{\sqrt{n}}$. Now for any $v \in \mathcal{E}(0)$, $||v|| = \sqrt{2}$.

$$P_{2}(y) = \frac{1}{2} \sum_{v \in \mathcal{E}(0)} \langle y, v \rangle v$$

= $\frac{1}{2} \sum_{j=1}^{\frac{n}{2}} (-y_{1} + y_{j+\frac{n}{2}}) v$
= $\frac{1}{2} (y_{1} - y_{1+\frac{n}{2}}, y_{2} - y_{2+\frac{n}{2}}, \dots, y_{\frac{n}{2}} - y_{\frac{n}{2}+\frac{n}{2}}, -y_{1} + y_{1+\frac{n}{2}}, -y_{2} + y_{2+\frac{n}{2}}, \dots, -y_{\frac{n}{2}} + y_{\frac{n}{2}+\frac{n}{2}}).$

Observe that the sum of elements in each row of the matrix representation of P_2 with respect to the standard basis of \mathbb{R}^n is 0. Therefore $||P_2\mathbb{1}|| = 0$ and hence $\beta_2 = 0$. For any $v \in \mathcal{E}(-2)$, $||v|| = \frac{1}{2}$.

$$P_{3}(y) = \frac{1}{4} \sum_{w \in \mathcal{E}(-2)} \langle y, w \rangle w$$

= $\frac{1}{4} \sum_{j=2}^{\frac{n}{2}} (-y_{1} + y_{j} - y_{1+\frac{n}{2}} + y_{j+\frac{n}{2}}) w_{j} = w,$

where *i*th coordinate of w is given by

$$w = \begin{cases} \left(\frac{n}{2} - 1\right)y_1 - y_2 - \dots - y_{\frac{n}{2}} + \left(\frac{n}{2} - 1\right)y_{\frac{n}{2}+1} - y_{\frac{n}{2}+2} - \dots - y_n & \text{if } i = 1, \frac{n}{2} + 1\\ -y_1 + y_i - y_{\frac{n}{2}+1} + y_{\frac{n}{2}+i} & \text{otherwise} \end{cases}$$

Again, observe that the sum of elements in each row of matrix representation of P_3 with respect to the standard basis of \mathbb{R}^n is 0. Therefore $||P_3\mathbb{1}|| = 0$ and hence $\beta_3 = 0$.

If n = 3, then $K_1 \bigtriangledown (K_{n-1} - M) \cong P_3$ and

spec
$$(P_3) = \begin{pmatrix} \sqrt{2} & 0 & -\sqrt{2} \\ 1 & 1 & 1 \end{pmatrix}$$
.

Therefore, P_3 is not integral. Let $n \ge 5$ be an odd integer. By Theorem 3.6, we have,

$$P_{K_1 \bigtriangledown (K_n - M)}(y) = P_{K_n - M}(y) \left(y - \sum_{i=1}^k \frac{n\beta_i^2}{y - \mu_i} \right).$$
(4)

By Proposition 3.1, distinct eigenvalues of $K_m - M$, where m = n - 1 are m - 2, 0 and -2. Also the main angles of $K_m - M$ are, $\beta_1 = \frac{1}{\sqrt{m}}, \beta_2 = \beta_3 = 0$ and $P_{K_m - M}(y) = y^{\frac{m}{2}}(y+2)^{\frac{m}{2}-1}(y-m+2)$. Therefore, the Equation (4) becomes

$$P_{K_{1} \bigtriangledown (K_{m}-M)}(y) = y^{\frac{m}{2}}(y+2)^{\frac{m}{2}-1}(y-m+2)\left(y-\frac{m\left(\frac{1}{\sqrt{m}}\right)^{2}}{y-(m-2)}\right)$$
$$= y^{\frac{m}{2}}(y+2)^{\frac{m}{2}-1}(y^{2}+(2-m)y-1).$$

Proposition 3.2. If G is a distance magic graph of order n with $\Delta(G) = n - 1$, then G is not integral.

Proof. We know the if G is distance magic graph of order n with $\Delta(G) = n - 1$ then n is odd and G is isomorphic to a cone cover of $K_m - M$, where m = n - 1 and its characteristic polynomial is $P_{K_1 \bigtriangledown (K_m - M)}(y) = y^{\frac{m}{2}}(y+2)^{\frac{m}{2}-1}(y^2 + (2-m)y-1)$. Consider the factor $y^{\frac{m}{2}}(y+2)^{\frac{m}{2}-1}$. If n = 3, we know that P_3 is not integral as $\sqrt{2}$ is its eigenvalue. Let us assume that $n \ge 5$. With $c = m - 2 \ge 2$ even, we rewrite the polynomial $y^2 - (m-2)y - 1$ as $y^2 - cy - 1$. The roots of this polynomial are: $y = \frac{c \pm \sqrt{c^2+1}}{2}$. Again, since c is even, we put c = 2d. Then the roots become $y = d \pm \sqrt{4d^2 + 1}$. But $4d^2 + 1$ is not a perfect square. If $4d^2 + 1 = a^2$ for some integer a, then we get (2d-a)(2d+a) = 1, which is not possible as a and d both are integers. Therefore, the cone cover of G has a non-integer eigenvalue as required. This completes the proof.

Up to the isomorphism, there are only 23 distance magic graphs of order up to 10 [13,17]. It can be manually checked that in any distance magic graph G of order up to 10, there exist vertices u, v such that N(u) = N(v). Therefore, the rows in the adjacency matrix of G corresponding to the vertices u and v are identical. From this, we conclude that such a graph G must be singular. Therefore, every distance magic graph of an order up to 10 is singular. The nonsingular distance magic graph of the smallest order (on the 11 vertices) is shown in Figure 4. Also, as discussed earlier, there is at least one singular distance magic graph of order n for each $n \ge 3$. These observations shifts our attention to the generalised inverse of the adjacency matrix of distance magic graphs.




Figure 4: Non-singular distance magic graph of the smallest order

A matrix $B^{\dagger} \in \mathbb{R}^{n \times m}$ is said to be the *Moore-Penrose* (or the generalised) inverse of a matrix $B \in \mathbb{R}^{m \times n}$ if $BB^{\dagger}B = B$, $B^{\dagger}BB^{\dagger} = B^{\dagger}$, $(BB^{\dagger})^* = BB^{\dagger}$ and $(B^{\dagger}B)^* = B^{\dagger}B$. We give another necessary condition for a graph to be distance magic in terms of the generalised inverse of its adjacency matrix. Recall that a *doubly stochastic matrix* is a matrix whose each row and each column sum to one.

Theorem 3.7. If G is distance magic, then AA^{\dagger} is doubly stochastic, where A^{\dagger} is the generalised inverse of an adjacency matrix A of G.

Proof. Let G be the distance magic graph with magic constant k. Then $Ax = k\mathbb{1}$.

$$\implies AA^{\dagger}Ax = kAA^{\dagger}\mathbb{1}$$
$$\implies Ax = kAA^{\dagger}\mathbb{1}$$
$$\implies k\mathbb{1} = kAA^{\dagger}\mathbb{1}$$
$$\implies \mathbb{1} = AA^{\dagger}\mathbb{1}.$$

This shows that each row in AA^{\dagger} sum to 1. Since AA^{\dagger} is symmetric, column sums are also 1. This proves the theorem.

The converse of the above theorem is not true. For example, take $G = K_{1,3}$ and let A be its adjacency matrix. Then

$A^{\dagger}=$	$\left(\frac{1}{3}\right)$	0	0	0)	(1	0	0	0)
	$\frac{1}{3}$	0	0	0		$\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{3}$
	$\frac{1}{2}$	0	0	0	$AA^{\dagger} = \begin{pmatrix} 1\\0\\0\\0 \end{pmatrix}$	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{2}$
	(ő	$\frac{1}{3}$	$\frac{1}{3}$	$\left(\frac{1}{3}\right)$		$\frac{1}{3}$	$\frac{1}{3}$	$\left(\frac{1}{3}\right)$

and 1 is an eigenvector of AA^{\dagger} corresponding to eigenvalue 1, but G is not distance magic.

4 Action of the automorphism group of the graph on its labeling set

Let G be a graph on n vertices, and let Aut(G) be the group of automorphisms of G. Let $\mathcal{M}(G) = \{f : f \text{ is distance magic labeling of } G\}$ be set of distance magic labelings of G.

Theorem 4.1. A distance magic graph G has at least |Aut(G)| distance magic labelings.

Proof. Let G be a graph with $V(G) = \{1, 2, ..., n\}$ and \mathcal{M} be a set of all distance magic labelings of a graph G. Define action of Aut(G) on \mathcal{M} by $(\sigma \cdot f)(i) = f\sigma^{-1}(i)$, for all $f \in \mathcal{M}$ and for all $i \in V$. First, we will show that $\sigma \cdot f \in \mathcal{M}$ for all f and all σ . For $v \in V$, let $N(v) = \{u_1, u_2, ..., u_t\}$. Since $f \in \mathcal{M}$, $w_f(v) = k$ and since $\sigma \in Aut(G)$, $u_i \in N(v)$ implies $\sigma^{-1}(u_i) \in N(\sigma^{-1}(v)).$

$$w_{\sigma \cdot f}(v) = \sum_{u \in N(v)} \sigma \cdot f(u)$$
$$= \sum_{u \in N(v)} f \sigma^{-1}(u)$$
$$= \sum_{x \in N(\sigma^{-1}(v))} f(x)$$
$$= k.$$

Now it is easy to check that Aut(G) acts on \mathcal{M} under the above action. For $f \in \mathcal{M}$, let \mathcal{S}_f denotes the stabiliser of f and let \mathcal{O}_f denotes the orbit of f. Let $\sigma \cdot f = f$ for some $\sigma \in Aut(G)$. Then $f\sigma^{-1} = f$ i.e. $f\sigma^{-1}(x) = f(x)$ for all $x \in V$. Since both f and σ are bijective functions, we conclude that $\sigma \equiv 1$, the identity automorphism. This proves that $|\mathcal{S}_f| = 1$. Therefore by orbit stabiliser theorem, $|Aut(G)| = |\mathcal{O}_f|$. As orbits of elements of \mathcal{M} partitions \mathcal{M} , each of which has same size |Aut(G)|, |Aut(G)| divides $|\mathcal{M}|$. This completes the proof.




Figure 5: Distance magic labeling of graphs P_3 and C_4

For a given distance magic graph G, we know that $|Aut(G)| \leq |\mathcal{M}|$. For simplicity, for a given graph G, we will denote its vertex labeling as $(f(v_1), f(v_2), \ldots, f(v_n))$ or simply $(f(v_1)f(v_2)\ldots f(v_n))$ when there is no confusion. Now, we give some examples showing that the lower bound obtained on the size of distance magic labelings \mathcal{M} is tight for several graphs. The equality is observed for the distance magic graphs P_3 and C_4 . The distance magic labelings of P_3 and C_4 with magic constants 3 and 5, respectively, are shown in Figure 5.

Recall that $|Aut(P_3)| = 2$. Let $P_3 : v_1, v_2, v_3$ be v_1, v_3 -path. It is easy to verify that $\mathcal{M}(P_3) = \{(132), (231)\}$ is the complete set of distance magic labelings of P_3 . Hence, $|Aut(P_3)| = |\mathcal{M}(P_3)|$. For a cycle $C_4 : v_1, v_2, v_3, v_4, v_1; \mathcal{M}(C_4) =$ $\{(1243), (3124), (4312), (2431), (2134), (1342), (3421), (4213)\}$ and $|Aut(C_4)| = 8$. Therefore, $|Aut(C_4)| = |\mathcal{M}(C_4)|$.

For the distance magic graph $P_3 \cup C_4$, the inequality is strict. We know that $|Aut(P_3 \cup C_4)| = 16$. From each of the distance magic labeling schemes M_1 , M_2 and M_3 , shown in Figure 6, 16 different distance magic labelings of $P_3 \cup C_4$ can be obtained using automorphisms of $P_3 \cup C_4$. Therefore, $P_3 \cup C_4$ has at least 48 different distance magic labelings.




Figure 6: Various distance magic labeling schemes of $P_3 \cup C_4$

5 Conclusion and further directions in research

The graph-theoretic properties and the spectrum of the class 'distance magic graphs are unexplored. In this paper, we have studied several graph-theoretic properties of this class. We raise the following problems:

Problem 1. Given a graph G, which is not distance magic, characterise all the primes $p \leq \frac{n^2-1}{2}$ such that G is not p-distance magic graph.

Problem 2. Characterise the distance magic graphs in terms of the spectra of the matrices related to the graphs.

Problem 3. Determine the class of singular as well as integral distance magic graphs.

Problem 4. Characterisation of distance magic graphs G for which $|Aut(G)| = |\mathcal{M}(G)|$.

Similar bounds as given in Theorem 4.1 can be obtained for other variants of graph labelings: for example, say *antimagic labeling* (a graph G is said to be antimagic if all of its vertex weights under the edge labeling $f : E(G) \to \{1, 2, ..., |E(G)|\}$ are distinct. For more details see [5]).

Problem 5. Characterise the distance magic graphs of order n with maximum degree (n-2).

Problem 6. Given a graph G and a positive integer p, study the computational complexity of the decision problem: whether G is p-distance magic?

References

- S. Arumugam, D. Froncek, and N. Kamatchi. Distance magic graphs—A survey. J. Indones. Math. Soc., (Special edition):11–26, 2011.
- [2] S. Arumugam, N. Kamatchi, and G. R. Vijayakumar. On the uniqueness of D-vertex magic constant. Discuss. Math. Graph Theory, 34(2):279–286, 2014.
- [3] D. Cvetković, P. Rowlinson, and S. Simić. An Introduction to the Theory of Graph Spectra. Cambridge University Press, 2010.
- [4] D. Froncek. Group distance magic labeling of Cartesian product of cycles. Australas. J. Combin., 55:167–174, 2013.
- [5] J. A. Gallian. A dynamic survey of graph labeling (25th edition). *Electron. J. Combin.*, 5:Dynamic Survey 6, 43, 2022.

- [6] S. Gupta and M. Zehavi. Multivariate analysis of scheduling fair competitions. arXiv:2102.03857, 2021.
- [7] M. Jinnah. On Σ-labelled graphs. In B.D. Acharya and S.M. Hedge, editors, *Techni-cal Proceedings of Group Discussion on Graph Labeling Problems*, pages 71–77. NITK Surathkal, 1999.
- [8] N. Kamatchi. Distance Magic and Distance Antimagic Labelings of Graphs. Phd thesis, Kalasalingam Academy of Research and Education, Krishnankoil, Srivilliputhur, Tamil Nadu, June 2012.
- [9] Š. Miklavič and P. Šparl. Classification of tetravalent distance magic circulant graphs. Discrete Math., 344(11):Paper No. 112557, 8, 2021.
- [10] S. Miklavič and P. Sparl. On distance magic circulants of valency 6. Discrete Appl. Math., 329:35–48, 2023.
- [11] M. Miller, C. Rodger, and R. Simanjuntak. Distance magic labelings of graphs. Australas. J. Combin., 28:305–315, 2003.
- [12] A. O'Neal and P. J. Slater. Uniqueness of vertex magic constants. SIAM J. Discrete Math., 27(2):708–716, 2013.
- [13] R. Pawar, T. Singh, H. Mukherjee, and J. Bagga. A complete characterization of all magic constants arising from distance magic graphs. arXiv:2311.10330, 2023.
- [14] P. Slater. It is all labeling. In G. Ralucca, L. Craig, and H. Stephen, editors, *Graph The-ory: Favourite conjectures and open problems-1*, pages 231–252. Springer International Publishing, 2016.
- [15] V. Vilfred. Σ-Labelled Graphs and Circulant Graphs. Phd thesis, University of Kerala, Trivandrum, Kerala, India, 1994.
- [16] D. B. West. Introduction to Graph Theory. Prentice Hall, 2nd edition, September 2000.
- [17] F. Yasin and R. Simanjuntak. A heuristic for distance magic labeling. Procedia Computer Science, 74:100–104, 2015. "The 2nd International Conference of Graph Theory and Information Security".