Geometric Barriers for Stable and Online Algorithms for Discrepancy Minimization

David Gamarnik* Eren C. Kızıldağ † Will Perkins ‡ Changji Xu § February 14, 2023

Abstract

For many computational problems involving randomness, intricate geometric features of the solution space have been used to rigorously rule out powerful classes of algorithms. This is often accomplished through the lens of the multi Overlap Gap Property $(m\text{-}\mathrm{OGP})$, a rigorous barrier against algorithms exhibiting input stability. In this paper, we focus on the algorithmic tractability of two models: (i) discrepancy minimization, and (ii) the symmetric binary perceptron (SBP), a random constraint satisfaction problem as well as a toy model of a single-layer neural network.

Our first focus is on the limits of online algorithms. By establishing and leveraging a novel geometrical barrier, we obtain sharp hardness guarantees against online algorithms for both the SBP and discrepancy minimization. Our results match the best known algorithmic guarantees, up to constant factors. Our second focus is on efficiently finding a constant discrepancy solution, given a random matrix $\mathcal{M} \in \mathbb{R}^{M \times n}$. In a smooth setting, where the entries of \mathcal{M} are i.i.d. standard normal, we establish the presence of m-OGP for $n = \Theta(M \log M)$. Consequently, we rule out the class of stable algorithms at this value. These results give the first rigorous evidence towards a conjecture of Altschuler and Niles-Weed [ANW21, Conjecture 1].

Our methods use the intricate geometry of the solution space to prove tight hardness results for online algorithms. The barrier we establish is a novel variant of the m-OGP. Furthermore, it regards m-tuples of solutions with respect to correlated instances, with growing values of m, $m = \omega(1)$. Importantly, our results rule out online algorithms succeeding even with an exponentially small probability.

^{*}Sloan School of Management, Massachusetts Institute of Technology; e-mail: gamarnik@mit.edu.

Department of Statistics, Columbia University; e-mail: eck2170@columbia.edu.

^{\$\}frac{1}{2}\$School of Computer Science, Georgia Institute of Technology; e-mail: math@willperkins.org.

[§]Center of Mathematical Sciences and Applications, Harvard University; e-mail: cxu@cmsa.fas.harvard.edu.

Contents

1	\mathbf{Intr}	$\operatorname{roduction}$	3
	1.1	Discrepancy Minimization	3
	1.2	Symmetric Binary Perceptron (SBP)	
	1.3	Connections between Discrepancy Theory and the SBP	5
	1.4	Main Results	
	1.5	Background and Prior Work	7
	1.6	Open Problems	
2	Tig	ht Hardness Guarantees for Online Algorithms	10
3	Algorithmic Barriers in Finding Constant Discrepancy Solutions		12
	3.1	Technical Preliminaries	12
	3.2	Ensemble m -OGP in Discrepancy Minimization	13
	3.3	<i>m</i> -OGP Implies Failure of Stable Algorithms	13
4	Proofs		14
	4.1	Proof of Theorem 2.3	15
	4.2	Proof of Theorem 2.4	18
	4.3	Proof Sketch for Theorem 2.5	
	4.4	Proof of Theorem 3.2	

1 Introduction

In this paper, we study the discrepancy minimization problem and the perceptron model. Combinatorial discrepancy theory [Spe85, Mat99] is a central topic at the intersection of combinatorics, probability, and algorithms. Given a matrix $\mathcal{M} \in \mathbb{R}^{M \times n}$, the central task in discrepancy theory is computing or bounding the quantity

$$\mathcal{D}(\mathcal{M}) \triangleq \min_{\boldsymbol{\sigma} \in \Sigma_n} \left\| \mathcal{M} \boldsymbol{\sigma} \right\|_{\infty},$$

known as the discrepancy of \mathcal{M} .

The perceptron is a toy one-layer neural network model storing random patterns as well as a very natural high-dimensional probabilistic model, see [JH60, Win61, Wen62, Cov65] for early works on it. Given random patterns $X_i \in \mathbb{R}^n$, $1 \leq i \leq M$, storage is achieved if one finds a $\sigma \in \mathbb{R}^n$ 'consistent' with all X_i : $\langle \sigma, X_i \rangle \geq 0$ for $1 \leq i \leq M$. The vector σ is interpreted as synaptic weights; it can either lie on the sphere in \mathbb{R}^n , $\|\boldsymbol{\sigma}\|_2 = \sqrt{n}$, or have binary entries, $\sigma \in \Sigma_n = \{-1,1\}^n$. The former is dubbed as the spherical perceptron, see [Gar88, ST03, Sto13, Tal11, AS20 for relevant work. In this paper, we only focus on the latter, dubbed as the binary perceptron. A fundamental object studied in the perceptron literature is the storage capacity: the maximum number of (random) patterns that can be stored with a suitable σ , see [Gar87, Gar88, GD88]. Krauth and Mézard [KM89] gave a detailed though non-rigorous characterization of the storage capacity. More recently, perceptron models with an activation function $U: \mathbb{R} \to \{0,1\}$ are considered, where a pattern X_i is stored with respect to (w.r.t.) U if $U(\langle \boldsymbol{\sigma}, X_i \rangle) = 1$. Of particular interest to us is the activation $U(x) = \mathbf{1}_{|x| < \kappa \sqrt{n}}$ which defines the symmetric binary perceptron (SBP) model proposed by Aubin, Perkins, and Zdeborová [APZ19] (see also [BNSX21, NS23] for results on more general perceptron models). As we see below, the SBP is closely related to discrepancy minimization.

1.1 Discrepancy Minimization

The discrepancy literature pertains to both worst-case and average-case \mathcal{M} . In the worst-case, minimal structure is assumed on \mathcal{M} , whereas in the average-case, the entries of \mathcal{M} are random, e.g. i.i.d. Bernoulli, Rademacher, or standard normal. Moreover, both existential as well as algorithmic results are sought in discrepancy theory.

Concerning the worst-case analysis, a landmark result in the area is due to Spencer [Spe85]: $\mathcal{D}(\mathcal{M}) \leq 6\sqrt{n}$ if $\mathcal{M} \in \mathbb{R}^{n \times n}$ with $|M_{ij}| \leq 1$ for $1 \leq i, j \leq n$ ('six standard deviations suffice'). The significance of this result is the improvement over the discrepancy guaranteed by the basic probabilistic method: the discrepancy incurred by a random signing is of order $\Theta(\sqrt{n \log n})$ which is substantially larger than $O(\sqrt{n})$. It is worth noting that Spencer's result is worst-case and non-constructive, but recent works [Ban10, LM15, LRR17, Rot17] has given efficient algorithms to find such low discrepancy solutions.

In this paper we focus on average-case discrepancy. Suppose $\mathcal{M} \in \mathbb{R}^{M \times n}$ has i.i.d. $\mathcal{N}(0,1)$ entries and M = o(n). In this case, a line of work initiated in [KKLO86] (for M = 1) and subsequently continued in [Cos09, TMR20] (for $M \geq 2$) established that $\mathcal{D}(\mathcal{M}) = \Theta(\sqrt{n}2^{-n/M})$ w.h.p. Algorithmic results in this regime are found in [KK82, Yak96, TMR20]. In the special case of M = 1, [GK21a] gave rigorous evidence that finding a $\boldsymbol{\sigma}$ with $\|\mathcal{M}\boldsymbol{\sigma}\|_{\infty} = 2^{-\omega(\sqrt{n\log n})}$

may be algorithmically intractable. On the other hand when $M = \Theta(n)$, then it turns out $\mathcal{D}(\mathcal{M}) = \Theta(\sqrt{n})$ and this case is closely related to the SBP, see Section 1.3 for more details.

Next suppose the entries of \mathcal{M} are i.i.d. binary, e.g. Rademacher or Bernoulli(p). In this case, while still $\mathcal{D}(\mathcal{M}) = \Theta(\sqrt{n})$ w.h.p. when $M = \Theta(n)$, but it turns out that constant discrepancy, in fact $\mathcal{D}(\mathcal{M}) = 1$, is possible when n is much larger than M. The sharpest possible result to this end is due to Altschuler and Niles-Weed [ANW21] who completely resolved the question of exactly when $\mathcal{D}(\mathcal{M}) \leq 1$: $\mathcal{D}(\mathcal{M}) \leq 1$ if \mathcal{M} consists of Bernoulli(p) entries with arbitrary p and p and p arbitrary constant greater than p and p are covers in particular the sparse regime, p = o(1), and is the sharpest possible as p and p is needed for p and p to be p and p are constant discrepancy solutions exist. Earlier results towards this direction are found in [HR19, FS20, Pot18]. Equipped with this existential guarantee from [ANW21], a natural algorithmic question is whether one can find such a constant discrepancy solution in polynomial time. This task is conjecturally hard, see [ANW21, Conjecture 1]. The algorithmic tractability of this problem is a main focuses of the present paper.

1.2 Symmetric Binary Perceptron (SBP)

Fix $\kappa > 0$, $\alpha > 0$, and set $M = \lfloor n\alpha \rfloor \in \mathbb{N}$. Let $X_i \sim \mathcal{N}(0, I_n)$, $1 \leq i \leq M$, be i.i.d. random vectors, where $\mathcal{N}(0, I_n)$ is the centered multivariate normal distribution in \mathbb{R}^n with identity covariance. Consider the (random) set

$$S_{\alpha}(\kappa) = \left\{ \boldsymbol{\sigma} \in \Sigma_n : |\langle \boldsymbol{\sigma}, X_i \rangle| \le \kappa \sqrt{n}, 1 \le i \le M \right\} = \left\{ \boldsymbol{\sigma} \in \Sigma_n : \left\| \mathcal{M} \boldsymbol{\sigma} \right\|_{\infty} \le \kappa \sqrt{n} \right\},$$
 (1)

where $\mathcal{M} \in \mathbb{R}^{M \times n}$ with rows X_1, \ldots, X_M . The word symmetric refers to the fact $\sigma \in S_{\alpha}(\kappa)$ iff $-\sigma \in S_{\alpha}(\kappa)$. The SBP was put forth by Aubin, Perkins, and Zdeborová [APZ19] as a symmetric counterpart to the asymmetric binary perceptron (ABP), where the constraints are instead of form $\langle \sigma, X \rangle \geq \kappa \sqrt{n}$, $1 \leq i \leq M$. The ABP turns out very challenging mathematically, see [KM89, KR98, Tal99, Xu19, DS19, PX21, ALS21a, GKPX22, Kız22] for relevant work and more details. The SBP, on the other hand, retains pertinent structural properties conjectured for the ABP [BDVLZ20], while being more amenable to rigorous analysis thanks to the symmetry.

The SBP undergoes a *sharp phase transition*, conjectured in [APZ19] and subsequently proven independently by Perkins and Xu [PX21] and Abbe, Li, and Sly [ALS21b]. Let

$$\alpha_c(\kappa) \triangleq -\frac{1}{\log_2 \mathbb{P}[|Z| \le \kappa]}, \text{ where } Z \sim \mathcal{N}(0, 1).$$
 (2)

Then

$$\lim_{n \to \infty} \mathbb{P} \big[S_{\alpha}(\kappa) \neq \varnothing \big] = \begin{cases} 0, & \text{if } \alpha > \alpha_c(\kappa) \\ 1, & \text{if } \alpha < \alpha_c(\kappa) \end{cases} . \tag{3}$$

The part $\alpha > \alpha_c(\kappa)$ is due to [APZ19] and established via an application of the first moment method: $\mathbb{E}[|S_{\alpha}(\kappa)|] = o(1)$ if $\alpha > \alpha_c(\kappa)$, so $S_{\alpha}(\kappa) = \emptyset$ w.h.p. by Markov's inequality. The same paper also studies the case $\alpha < \alpha_c(\kappa)$ and establishes, through the second moment method, that $\liminf_{n\to\infty} \mathbb{P}[S_{\alpha}(\kappa) \neq \emptyset] > 0$. Boosting this to a high probability guarantee requires more powerful tools, see [PX21] for a delicate martingale argument and [ALS21b] for an argument

based on a fully connected analog of the small subgraph conditioning method. Furthermore, very recently, the critical window around $\alpha_c(\kappa)$ was shown to be of constant width, see [Alt22, SS23]. These facts highlight that the first moment 'prediction' for the precise location of the phase transition is correct, and the transition is very sharp.

Given that $S_{\alpha}(\kappa)$ is w.h.p. non-empty if $\alpha < \alpha_c(\kappa)$, a natural follow-up question is algorithmic: can a solution $\sigma \in S_{\alpha}(\kappa)$ be found efficiently? And does the existence of efficient algorithms depend on α ? Efficient algorithms at small densities α were given in [KR98] and [ALS21a] for the ABP and SBP respectively while on the negative side, [GKPX22] studied the limits of efficient algorithms (see details below). The works [BBBZ07, BIL+15, BDVLZ20] put forth possible explanations for the success of efficient algorithms: while almost all solutions are totally frozen (conjectured in [MMZ05, HK14]), efficient algorithms access rare solutions lying in large clusters. Recent works including [PX21, ALS21b, ALS21a] have studied these structural predictions.

1.3 Connections between Discrepancy Theory and the SBP

In order to explicate the connection between discrepancy minimization and the SBP, we focus on the proportional regime, i.e. $M = \Theta(n)$. The discrepancy viewpoint is to take an $\mathcal{M} \in \mathbb{R}^{M \times n}$ with a fixed aspect ratio $\alpha = M/n$, and to seek a $\boldsymbol{\sigma}$ such that $\|\mathcal{M}\boldsymbol{\sigma}\|_{\infty}$ is as small as possible. The perceptron viewpoint, on the other hand, is the inverse: fix a $\kappa > 0$ first and seek the largest α for which a solution $\boldsymbol{\sigma}$ with $\|\mathcal{M}\boldsymbol{\sigma}\|_{\infty} \leq \kappa \sqrt{n}$ exists. Furthermore, the asymptotic value of the average-case discrepancy in the proportional regime immediately follows from the sharp threshold result for the SBP (3): $\mathcal{D}(\mathcal{M}) = (1 + o(1))f(\alpha)\sqrt{n}$ w.h.p., where $f(\alpha)$ is the 'inverse' of $\alpha_c(\kappa)$ (2).

Algorithmic Connections The connection between the SBP and discrepancy theory further extends to algorithmic domain: the best known efficient algorithm for the SBP comes from the discrepancy literature. Suppose $\mathcal{M} \in \mathbb{R}^{M \times n}$ has i.i.d. Rademacher entries. Bansal and Spencer [BS20] devised an efficient online algorithm that finds a $\sigma_{ALG} \in \Sigma_n$ such that $\|\mathcal{M}\sigma_{ALG}\|_{\infty} = O(\sqrt{M})$ w.h.p. if $n \geq M = \omega(1)$. Informally, an algorithm is online if the t^{th} coordinate of the output σ_{ALG} depends only on first t columns of \mathcal{M} , see Definition 2.1 for a formal definition. As an immediate corollary, this yields an efficient algorithm for the SBP that finds a solution $\sigma \in S_{\alpha}(\kappa)$ w.h.p. if $\alpha = O(\kappa^2)$, see [GKPX22, Corollary 3.6]. In fact, this is the best known algorithmic guarantee both for the SBP and for discrepancy in the random proportional regime, see [GKPX22, Section 3.3].

In light of these existential and algorithmic results, it appears that the SBP may exhibit a striking statistical-to-computational gap (SCG): the density below which solutions exist w.h.p., i.e. $\alpha_c(\kappa)$, is substantially larger than those below which polynomial-time search algorithms work. Further, this SCG is most profound when $\kappa \to 0$. While the Bansal-Spencer algorithm works only when $\alpha = O(\kappa^2)$, solutions do exist w.h.p. below $\alpha_c(\kappa)$ which, per (2), is asymptotically $\frac{1}{\log_2(1/\kappa)}$. Origins of this SCG were investigated in [GKPX22], where it was shown that the SBP exhibits an intricate geometrical property called the multi Overlap Gap Property (m-OGP) when $\alpha = \Omega(\kappa^2 \log_2 \frac{1}{\kappa})$ and consequently stable algorithms fail to find a satisfying solution for $\alpha = \Omega(\kappa^2 \log_2 \frac{1}{\kappa})$. It is worth noting, though, that stable algorithms need not include online algorithms, which achieve the computational threshold for the SBP. What the limits of online algorithms are is an open question we undertake in this paper.

In addition to the SBP, the discrepancy minimization problem — in particular the algorithmic problem of efficiently finding a constant discrepancy solution when such solutions exist w.h.p. — also exhibits a similar SCG. To recall, when $\mathcal{M} \in \mathbb{R}^{M \times n}$, then constant discrepancy solutions exist w.h.p. as soon as $n = \Omega(M \log M)$. On the other hand, the best known polynomial-time algorithm succeeds at a dramatically smaller value $M = o(\log n)$ [Ban22], highlighting another striking SCG. This is our second focus in the present paper.

1.4 Main Results

Suppose \mathcal{M} per (1) consists of i.i.d. $\mathcal{N}(0,1)$ entries. Our first main result establishes that online algorithms fail to find a satisfying solution for the SBP at densities $\alpha = \Omega(\kappa^2)$.

Theorem 1.1 (Informal, see Theorem 2.3). For densities $\alpha = \Omega(\kappa^2)$, online algorithms fail to find a solution for the SBP w.p. greater than $e^{-\Theta(n)}$.

Our next result extends Theorem 1.1 to the discrepancy minimization problem when $\mathcal{M} \in \mathbb{R}^{M \times n}$ consists of i.i.d. Rademacher or i.i.d. Bernoulli(p) entries.

Theorem 1.2 (Informal, see Theorems 2.4-2.5). There exists c > 0 such that online algorithms fail to return a solution of discrepancy at most $c\sqrt{M}$ w.p. greater than $e^{-\Theta(M)}$.

If the entries of \mathcal{M} are Rademacher, taking c=1/24 suffices. For Bernoulli case, the implied constant depends on p: it suffices to take $c \triangleq c_p = \sqrt{p-p^2}/24$. Taken together, Theorems 1.1 and 1.2 collectively yield that among the class of online algorithms, Bansal-Spencer algorithm [BS20] is optimal up to constants for both models. Our proof is based on a novel version of m-OGP: we show the non-existence of tuples of solutions agreeing on first $1-\Delta$ fraction of coordinates for a suitable $\Delta \in (0,1)$, for a collection of m correlated instances, see below for details. This barrier is more restricted than m-OGP, which asserts the non-existence of tuples of solutions at a prescribed distance. Additionally, for Theorem 1.2, one has to consider m-tuples with growing values of m, $m = \omega(1)$; this idea is originally due to Gamarnik and Kızıldağ [GK21a] for lowering the m-OGP threshold.

To the best of our knowledge, Theorems 1.1-1.2 are the first (up-to-constants) tight hardness guarantees via geometrical barriers against classes beyond stable algorithms, see Section 1.5 for details. Furthermore, unlike prior work [GJW20, Wei20, HS21, GK21a, GKPX22], the algorithms ruled out need not succeed w.h.p. or even with a constant probability: an exponentially small success probability suffices. This is made possible by using a clever application of Jensen's inequality, originally due to Gamarnik and Sudan [GS17b].

Proof Sketch We sketch the proof of Theorem 1.1, which is based on a new version of m-OGP coupled with a contradiction argument. Suppose that such an online algorithm \mathcal{A} with a success probability of p_s exists. Let $\mathcal{M}_1 \in \mathbb{R}^{M \times n}$ with i.i.d. $\mathcal{N}(0,1)$ entries. Fix an $m \in \mathbb{N}$ and a $\Delta \in (0,1)$, generate random matrices $\mathcal{M}_i \in \mathbb{R}^{M \times n}$, $2 \leq i \leq m$, by independently resampling the last Δn columns of \mathcal{M}_1 . Running \mathcal{A} on each \mathcal{M}_i , we obtain solutions $\sigma_i \triangleq \mathcal{A}(\mathcal{M}_i) \in \Sigma_n$, $1 \leq i \leq m$. An application of Jensen's inequality then reveals $\|\mathcal{M}_i \sigma_i\|_{\infty} \leq \kappa \sqrt{n}$, $1 \leq i \leq m$, w.p. at least p_s^m . Furthermore, since \mathcal{A} is online, it is the case that any σ_i and σ_j necessarily have identical first $n - \Delta n$ coordinates. Namely, if such an \mathcal{A} exists, then w.p. at least p_s^m , there exists an m-tuple $(\sigma_1, \ldots, \sigma_m)$ of satisfying solutions that agree on first $n - \Delta n$ coordinates.

We then establish, using the *first moment method*, that for suitably chosen m, Δ ; the probability that such an m-tuple exists is in fact strictly less than p_s^m . This is a contradiction. The proof of Theorem 1.2 is similar, though it requires additional technical steps. In particular, one needs an anti-concentration argument for signed sums of binary variables via Berry-Esseen Theorem.

Our next focus is on the algorithmic problem of efficiently finding a constant discrepancy solution, given a random $\mathcal{M} \in \mathbb{R}^{M \times n}$. To recall, such solutions exist w.h.p. as soon as $n = \Omega(M \log M)$ [ANW21], while the best known polynomial-time algorithm works only when $M = o(\log n)$ [Ban22]. Further, it was conjectured in [ANW21] that this task is algorithmically hard. Towards this conjecture, we focus on a smooth setting where the entries of \mathcal{M} are i.i.d. $\mathcal{N}(0,1)$. Our next main result shows the presence of m-OGP with m = O(1) when $n = \Theta(M \log M)$, giving a rigorous evidence of hardness at the 'boundary' $n = \Theta(M \log M)$.

Theorem 1.3 (Informal, see Theorem 3.2). For $n = \Theta(M \log M)$, the set of constant discrepancy solutions exhibits m-OGP (with constant m) for suitably chosen parameters.

The regime $\log n \ll M \ll n/\log n$ as well as extensions beyond Gaussian disorder—in particular to the Bernoulli or Rademacher case—are among the open problems we discuss in Section 1.6.

Our final main result leverages the m-OGP to show that stable algorithms fail to find a constant discrepancy solution when $n = \Theta(M \log M)$. Informally, an algorithm is stable if a small perturbation of its inputs induces only a small change in its output σ , see Definition 3.3 for a formal statement. The class of stable algorithm has been shown to capture powerful classes of algorithms including low-degree polynomials [GJW20, BH21], Approximate Message Passing (AMP) [GJ21], and Boolean circuits of low-depth [GJW21].

Theorem 1.4 (Informal, see Theorem 3.4). For $n = \Theta(M \log M)$, stable algorithms fail to find a constant discrepancy solution w.p. greater than a certain constant.

The proof of Theorem 1.4 is based on a Ramsey-theoretic argument developed in [GK21a] and also used in [GKPX22] coupled with the m-OGP result, Theorem 3.2; it rules out stable algorithms succeeding with a constant probability.

1.5 Background and Prior Work

Statistical-to-Computational Gaps (SCGs) Both the SBP and discrepancy minimization exhibit an SCG: known efficient algorithms perform strictly worse than the existential guarantee. Such gaps are a universal feature of many average-case algorithmic problems arising from random combinatorial structures and high-dimensional statistical inference. A partial list of problems with an SCG include random CSPs [MMZ05, ART06, ACO08, GS17b, BH21], optimization over random graphs [GS14, COE15, Wei20], spin glasses [GJ21, HS21], planted clique [DM15, BHK+19], and tensor decomposition [Wei22], see also the surveys by Gamarnik [Gam21] and Gamarnik, Moore, and Zdeborová [GMZ22]. Unfortunately, standard computational complexity theory is often useless due to the average-case nature of such problems¹. Nevertheless, a very promising line of research proposed various frameworks that provide rigorous evidence of hardness. These frameworks include average-case reductions—often from the planted clique [BR13, BBH18, BB19]—as well as unconditional lower bounds against restricted classes of algorithms,

¹Modulo a few exceptions, see e.g. [Ajt96, BABB21, GK21b].

including the statistical query algorithms [DKS17, FGR⁺17, FPV18], low-degree polynomials [Hop18, KWB22, Wei22], sum-of-squares hierarchy [HSS15, HKP⁺17, RSS18, BHK⁺19], AMP [ZK16, BPW18], and Monte Carlo Markov Chain (MCMC) methods [Jer92, DFJ02]. Yet another such approach is based on the intricate geometry of the solution space through the Overlap Gap Property.

Intricate Geometry and the Overlap Gap Property (OGP) Prior work [MMZ05, ART06, ACO08] discovered a very intriguing connection between intricate geometry and algorithmic hardness in the context of random CSPs: the onset of algorithmic hardness roughly coincides with the emergence of an intricate geometry in the solution space. The OGP framework leverages insights from statistical physics to rigorously link the intricate geometry to formal hardness. In the context of random optimization, the OGP informally states that (w.h.p. over the randomness) any two near-optima are either 'close' or 'far': there exists $0 < \nu_1 < \nu_2 < 1$ such that $n^{-1} \langle \boldsymbol{\sigma}, \boldsymbol{\sigma}' \rangle \in [0, \nu_1] \cup [\nu_2, 1]$ for any pair of near-optima $\boldsymbol{\sigma}, \boldsymbol{\sigma}' \in \Sigma_n$. Namely, the region of normalized overlaps is topologically disconnected; no pairs of near-optima at intermediate distances can be found. The OGP is a rigorous barrier for large classes of algorithms exhibiting input stability—see below. See [Gam21] for a survey on OGP.

Algorithmic Implications of OGP The first work establishing and leveraging OGP to rule out algorithms is due to Gamarnik and Sudan [GS14, GS17a]. Their focus is on the problem finding a large independent set in sparse random graphs on n vertices with average degree d, which exhibits an SCG: the largest such independent set is of size $2\frac{\log d}{d}n$ [FL92, BGT10], whereas the best known efficient algorithm finds an independent set of size $\frac{\log d}{d}n$. They establish that any pair of independent sets of size larger than $(1+1/\sqrt{2})\frac{\log d}{d}n$ exhibit the OGP. By leveraging the OGP, they then show that local algorithms fail to find an independent set of size larger than $(1+1/\sqrt{2})\frac{\log d}{d}n$. Subsequent research established and leveraged the OGP to rule out other classes of algorithms (e.g., AMP [GJ21], low-degree polynomials [GJW20, Wei20, BH21], Langevin dynamics [GJW20, HS21], low-depth circuits [GJW21]) for various other models (e.g., random graphs [RV17, GJW20, Wei20], spin glass models [CGPR19, GJ21, HS21], random CSPs [GS17b, BH21, GKPX22]). A very important feature found across the algorithms ruled out by the OGP and other versions of intricate geometry is input stability, similar to Definition 3.3 (apart from the failure of MCMC in planted models, see e.g. [Jer92, AWZ20, GJS21]). Our work marks the first instance of intricate geometry yielding tight algorithmic hardness against classes beyond stable algorithms.

Multi OGP (m-OGP) The prior work [GS14, GS17a] discussed above establish the failure of local algorithms at value $(1+1/\sqrt{2})\frac{\log d}{d}n$. By considering a certain overlap pattern involving many large independent sets, Rahman and Virág [RV17] subsequently removed the additional $1/\sqrt{2}$ term; they showed that the onset of OGP precisely coincides with the algorithmic $\frac{\log d}{d}n$ value. That is, one can potentially lower the onset of the OGP and rule out algorithms for a broader range of parameters through more intricate overlap patterns. In a similar vein, Gamarnik and Sudan [GS17b] studied the Not-All-Equal k-SAT problem and showed the presence of the OGP for the m-tuples of nearly equidistant satisfying assignments. Consequently, they obtained nearly tight hardness guarantees against sequential local algorithms. A similar m-OGP was also employed in [GK21a, GKPX22], and is also our focus here.

Recently, m-OGP for more intricate patterns were proposed. These forbidden patterns regard m-tuples of solutions where for any $2 \le i \le m$, the $i^{\rm th}$ solution has 'intermediate' overlap with the first i-1 solutions. By doing so, tight hardness guarantees against low-degree polynomials were obtained for finding independent sets in sparse random graphs by Wein [Wei20] and for the random k-SAT by Bresler and Huang [BH21]. Similarly, Huang and Sellke [HS21] construct a very intricate forbidden structure consisting of an ultrametric tree of solutions dubbed as the Branching OGP. By leveraging the branching OGP, they obtain tight hardness guarantees against Lipschitz algorithms for the p-spin model. Moreover, these papers establish the Ensemble m-OGP which regards m-tuples that are near-optimal w.r.t. correlated instances. The Ensemble OGP emerged in [CGPR19]; it has been instrumental in ruling out stable algorithms since. The investigation of m-tuples of solutions w.r.t. correlated instances is at the core of our paper. Furthermore, we inspect m-tuples with super-constant m, $m = \omega(1)$, to rule out online algorithms in Theorems 2.4-2.5. This idea originated in [GK21a] for further lowering the m-OGP threshold.

1.6 Open Problems

Geometrical Barriers for other Classes of Algorithms Prior work on OGP showed that intricate geometry is a signature of algorithmic hardness, and gave lower bounds against stable algorithms. Theorems 2.3, 2.4 and 2.5 extend this beyond stable algorithms; they leverage intricate geometry to rule out online algorithms. It would be very interesting to rule out other classes of algorithms via similar geometrical barriers.

Discrepancy Minimization beyond Gaussian Disorder Theorem 3.2 shows that, for $\mathcal{M} \in \mathbb{R}^{M \times n}$ with i.i.d. $\mathcal{N}(0,1)$ entries and $n = \Theta(M \log M)$, the set of constant discrepancy solutions exhibits the m-OGP. A very interesting question is whether m-OGP still holds when the entries of \mathcal{M} are binary. Prior work established OGP both for models with discrete disorder (e.g., random k-SAT [GS17b, BH21], random graphs [GS14, GS17a, GJW20, Wei20]) as well as for models with continuous disorder (e.g., spin glass models [GJ21, HS21], number partitioning [GK21a], the SBP [GKPX22]). These results suggest that OGP exhibits universality: the distributional details of the disorder are immaterial². In light of these, we make the following conjecture:

Conjecture 1.5. For $\mathcal{M} \in \mathbb{R}^{M \times n}$ with i.i.d. Rademacher or Bernoulli(p) entries and $n = \Theta(M \log M)$, set of constant discrepancy solutions exhibits m-OGP with suitable parameters.

Resolving Conjecture 1.5 may require understanding a probability term of the form $\mathbb{P}[M\mathbf{v} = \mathbf{x}]$ for a random \mathbf{v} with i.i.d. binary entries and a deterministic $M \in \{-1, 1\}^{m \times n}$ whose rows $\sigma_1, \ldots, \sigma_m$ satisfy $n^{-1} \langle \sigma_i, \sigma_j \rangle = \beta$ for some fixed β and every $1 \leq i < j \leq m$. One direction is to employ local limit arguments; we leave this as an open problem for future investigation.

Discrepancy Minimization beyond $n = \Theta(M \log M)$ Recall that constant discrepancy solutions exist as soon as $n = \Theta(M \log M)$, i.e. when $M = O(n/\log n)$, while the best known polynomial-time algorithm works only when $M = o(\log n)$. In light of these, Theorem 3.2 provides rigorous evidence of hardness, yet only at the 'boundary'. The regime $\log n \ll M \ll n/\log n$ is an interesting direction left for future work. A potential avenue would be to consider

²In fact, such a universality result has already been established for the SBP, see [GKPX22, Theorem 5.2].

a more intricate overlap pattern, such as those in [Wei20, HS21] or the branching OGP [HS21]. To this end, we discover an intriguing phase transition (proof omitted):

Theorem 1.6. Let $\mathcal{M} \in \mathbb{R}^{M \times n}$ with i.i.d. $\mathcal{N}(0,1)$ entries. Fix a K > 0 and let $S(m, \delta, K)$ be the set of $(\boldsymbol{\sigma}_1, \ldots, \boldsymbol{\sigma}_m)$ such that $d_H(\boldsymbol{\sigma}_i, \boldsymbol{\sigma}_j) = \delta$ and $\max_{i \leq m} \|\mathcal{M}\boldsymbol{\sigma}_i\|_{\infty} \leq K$.

(a) If
$$M = \omega(\log n)$$
 then $\mathbb{E}[|S(n \log^{-O(1)} n, \log^{O(1)} n, K)|] = o(1)$.

(b) If
$$M = o(\log n)$$
, then $\mathbb{E}[|S(n \log^{-O(1)} n, \log^{O(1)} n, K)|] = \omega(1)$.

Namely, the value $M = \log n$ is the threshold at which the (expected) number of m-tuples of constant discrepancy solutions at distance $\log^{O(1)} n$ with $m = \widetilde{\Theta}(n)$ undergoes a phase transition. Whether this phase transition at value $\log n$ is coincidental or a signature of algorithmic hardness is an open problem left for future work.

Paper Organization and Notation The rest of the paper is organized as follows. In Section 2, we formalize the class of online algorithms and state our hardness results. Section 3 is devoted to the algorithmic problem of finding constant discrepancy solutions. See Section 3.1 for preliminaries and the definition of set of m-tuples regarding OGP, Section 3.2 for the main m-OGP result and Section 3.3 for the hardness result against stable algorithms. Finally, see Section 4 for complete proofs. Our notation is fairly standard, see the beginning of Section 4 for details.

2 Tight Hardness Guarantees for Online Algorithms

In this section, we explore the limits of *online algorithms* in the context of our two-models: the SBP and average-case discrepancy. We begin by formalizing the class of online algorithms in the context of these two models.

Definition 2.1.

- (SBP) Fix a $\kappa > 0$ and an $\alpha < \alpha_c(\kappa)$. Let $\mathcal{M} \in \mathbb{R}^{\alpha n \times n}$ with i.i.d. $\mathcal{N}(0,1)$ entries.
- (Discrepancy) Fix $n \geq M$, let $\mathcal{M} \in \mathbb{R}^{M \times n}$ has i.i.d. Rademacher or Bernoulli(p) entries.

Fix a $p_s > 0$ and a K > 0. An algorithm \mathcal{A} is (p_s, α) -online for the SBP or (p_s, K) online for discrepancy if it satisfies the following. Let $\mathcal{A}(\mathcal{M}) \triangleq \boldsymbol{\sigma}_{ALG} = (\boldsymbol{\sigma}_{ALG}(i) : 1 \leq i \leq n) \in \Sigma_n$.

• (Success) We have

$$\mathbb{P}[\|\mathcal{M}\boldsymbol{\sigma}_{ALG}\|_{\infty} \leq \kappa \sqrt{n}] \geq p_s \quad (\textit{for the SBP})$$

$$\mathbb{P}[\|\mathcal{M}\boldsymbol{\sigma}_{ALG}\|_{\infty} \leq K] \geq p_s \quad (\textit{for discrepancy}).$$

• (Online) Let C_1, \ldots, C_n be the columns of \mathcal{M} . There exists deterministic functions f_1, \ldots, f_n such that for $1 \leq t \leq n$, $\sigma_{ALG}(t) = f_t(C_1, \ldots, C_t) \in \{-1, 1\}$.

The parameter p_s is the success guarantee of the algorithm, where the probability is taken w.r.t. the randomness in \mathcal{M} . The online nature of the algorithm admits the following interpretation. Columns \mathcal{C}_i arrive at a time. At the end of round t-1, the signs $\boldsymbol{\sigma}(i) \in \{-1,1\}$, $1 \leq i \leq t-1$ are assigned, and a new column \mathcal{C}_t arrives. The sign $\boldsymbol{\sigma}(t)$ then depends only on the previous decisions $\boldsymbol{\sigma}(i)$, $1 \leq i \leq t-1$ and \mathcal{C}_t . That is, $\boldsymbol{\sigma}(t)$ depends only on \mathcal{C}_i , $1 \leq i \leq t$. This abstraction captures, in particular, the Bansal-Spencer algorithm:

Theorem 2.2. [BS20, Theorem 3.4] Let $n \geq M$ and $\mathcal{M} \in \{-1,1\}^{M \times n}$ has i.i.d. Rademacher entries. Then, there exists absolute constants C > 0 and $\gamma < 1$, and an online algorithm \mathcal{A} admitting \mathcal{M} as its input and returning a $\boldsymbol{\sigma} \triangleq \mathcal{A}(\mathcal{M})$ such that

$$\mathbb{P}[\|\mathcal{M}\boldsymbol{\sigma}\|_{\infty} \le C\sqrt{M}] \le 1 - e^{-\Theta(M^{\gamma})}.$$

Theorem 2.2 immediately yields an efficient algorithm for the SBP when $\alpha \leq \kappa^2/C^2$, see [GKPX22, Corollary 4.6]. As mentioned in the introduction, the Bansal-Spencer algorithm is the best known polynomial-time algorithm both for the SBP and for the discrepancy minimization in random proportional regime. In the sense of Definition 2.1, it is a $(1 - e^{-\Theta(n^{\gamma})}, \kappa^2/C^2)$ -online algorithm for the SBP and a $(1 - e^{-\Theta(M^{\gamma})}, C\sqrt{M})$ -online algorithm for the discrepancy.

Online Algorithms for the SBP Our first main result focuses on the SBP in the regime $\kappa \to 0$ and establishes the following hardness for the class of online algorithms.

Theorem 2.3. Fix any small enough $\kappa > 0$ and any $\alpha \ge 4\kappa^2$. Then there exists an $n_0 \in \mathbb{N}$ and an absolute constant c > 0 such that the following holds. For any $n \ge n_0$, there exists no (e^{-cn}, α) -online algorithm for the SBP in the sense of Definition 2.1.

We prove Theorem 2.3 in Section 4.1. Several remarks are in order.

Theorem 2.3 establishes that in the regime $\kappa \to 0$, online algorithms fail to find a satisfying solution for the SBP for densities $\alpha = \Omega(\kappa^2)$. This substantially improves upon an earlier result in [GKPX22, Theorem 7.4], which showed the failure of online algorithms only when α is sufficiently close to the satisfiability threshold $\alpha_c(\kappa)$. Further, in light of Theorem 2.2, Theorem 2.3 is the sharpest possible: Bansal-Spencer algorithm is optimal (up to constants) among online algorithms; no online algorithm, in the sense of Definition 2.1, can improve upon it.

The algorithms Theorem 2.3 rules out need not succeed w.h.p. or even with a constant probability: an exponentially small success guarantee suffices. This is a particular strength of Theorem 2.3; we are unaware of any similar hardness guarantees for algorithms that succeed w.p. o(1). This is based on a clever application of Jensen's inequality that is originally due to Gamarnik and Sudan [GS17b].

Online Algorithms for the Discrepancy Minimization Our second main result extends Theorem 2.3 to discrepancy minimization for the case when the entries of \mathcal{M} are binary.

Theorem 2.4. Let c < 1/2 be arbitrary, $n \ge M = \omega(1)$, and $\mathcal{M} \in \{-1, 1\}^{M \times n}$ with i.i.d. Rademacher entries. Then there exists an $M_0 \in \mathbb{N}$ such that the following holds. For every $M \ge M_0$, there exists no $(e^{-cM}, \sqrt{M}/24)$ -online algorithm for discrepancy in the sense of Definition 2.1.

Furthermore, Theorem 2.4 remains valid even when the entries of \mathcal{M} are Bernoulli(p).

Theorem 2.5. Let c < 1/2 be arbitrary, $n \ge M = \omega(1)$, and $\mathcal{M} \in \{0,1\}^{M \times n}$ with i.i.d. Bernoulli(p) entries. Then there exists an $M_0 \in \mathbb{N}$ such that the following holds. For every $(p-p^2)M \ge M_0$, there exists no $(e^{-cM}, \sqrt{M(p-p^2)}/24)$ -online algorithm for discrepancy in the sense of Definition 2.1.

We prove Theorem 2.4 in Section 4.2 and give the extension to Theorem 2.5 in Section 4.3. Theorems 2.4-2.5 collectively establish that up to constant factors the Bansal-Spencer algorithm is optimal within the class of online algorithms for the discrepancy minimization problem. Once again, the algorithms ruled out can succeed even with an exponentially small probability.

At a technical level, Theorems 2.4-2.5 are established by showing the non-existence of certain m-tuples of solutions described earlier with growing values of m, $m = \omega_M(1)$. The idea of considering the 'landscape' of m-tuples with $m = \omega(1)$ was introduced in the context of random number partitioning problem [GK21a]. By doing so, the authors subsequently lowered the m-OGP threshold and ruled out stable algorithms for a broader range of parameters than what one can get for constant m. Ours is the first work leveraging such a barrier with growing values of m beyond stable algorithms; it further illustrates the potential gain of considering superconstant tuples for random computational problems. Another key ingredient of our proof is an anti-concentration inequality for signed sum of Bernoulli/Rademacher variables, via the Berry-Esseen Theorem.

3 Algorithmic Barriers in Finding Constant Discrepancy Solutions

In this section, we focus on the algorithmic problem of finding a constant discrepancy solution. More concretely, given a random $\mathcal{M} \in \mathbb{R}^{M \times n}$ we ask the following question: for what values of M and n, can a solution $\sigma \in \Sigma_n$ of constant discrepancy, $\|\mathcal{M}\sigma\|_{\infty} = O(1)$, be found efficiently?

To begin with, a simple first-moment calculation shows that $n = \Omega(M \log M)$ is necessary for such solutions to exist. This condition turns out to be sufficient, as well; [ANW21] showed that if \mathcal{M} has i.i.d. Bernoulli(p) entries then $\mathcal{D}(\mathcal{M}) \leq 1$ w.h.p. if $n \geq CM \log M$ where C is any arbitrary constant greater than $(2 \log 2)^{-1}$. On the other hand, the best known polynomial-time algorithm finding such a solution works only when $M = o(\log n)$ [Ban22]. This highlights a striking statistical-to-computational gap (SCG).

In this section, we study the nature of this SCG in a smooth setting where the entries of \mathcal{M} are i.i.d. $\mathcal{N}(0,1)$, near the existential boundary $n = \Theta(M \log M)$. We first focus on the 'landscape' of the set of constant discrepancy solutions, and show the presence of Ensemble m-OGP, an intricate geometrical property. We then leverage m-OGP to rule out the class of stable algorithms.

3.1 Technical Preliminaries

We formalize the set of tuples of constant discrepancy solutions under investigation.

Definition 3.1. Fix a K > 0, an $m \in \mathbb{N}$, $0 < \eta < \beta < 1$, and $\mathcal{I} \subset [0, \pi/2]$. Let $\mathcal{M}_i \in \mathbb{R}^{M \times n}$, $0 \le i \le m$, be i.i.d. random matrices, each having i.i.d. $\mathcal{N}(0,1)$ entries. Denote by $\mathcal{S}(K, m, \beta, \eta, \mathcal{I})$ the set of all m-tuples $\sigma_i \in \Sigma_n$, $1 \le i \le m$, satisfying the following:

• (Pairwise Overlap Condition) For any $1 \le i < j \le m, \ \beta - \eta \le n^{-1} \ \langle \sigma_i, \sigma_j \rangle \le \beta$.

• (Constant Discrepancy Condition) There exists $\tau_1, \ldots, \tau_m \in \mathcal{I}$ such that $\max_{1 \leq i \leq m} \|\mathcal{M}_i(\tau_i)\boldsymbol{\sigma}_i\|_{\infty} \leq K$, where $\mathcal{M}_i(\tau_i) = \cos(\tau_i)\mathcal{M}_0 + \sin(\tau_i)\mathcal{M}_i \in \mathbb{R}^{M \times n}$.

Definition 3.1 concerns tuples of solutions of discrepancy at most K. The parameter m is the size of tuples under consideration, and β and η collectively control the (forbidden) region of overlaps. Finally, the set \mathcal{I} is employed for generating correlated instances; this is necessary for establishing the Ensemble m-OGP to rule out stable algorithms, see below for details.

3.2 Ensemble *m*-OGP in Discrepancy Minimization

Our next main result shows that the set of constant discrepancy solutions exhibits the m-OGP.

Theorem 3.2. Fix arbitrary constants $C_1 > c_2 > 0$ and a K > 0, suppose that $C_1 M \log_2 M \ge n \ge c_2 M \log_2 M$. Then, there exists an $m \in \mathbb{N}$, a c > 0 and $0 < \eta < \beta < 1$ such that the following holds. Fix any $\mathcal{I} \subset [0, \pi/2]$ with $|\mathcal{I}| \le 2^{cn}$. Then, $\mathbb{P}[\mathcal{S}(K, m, \beta, \eta, \mathcal{I}) \ne \varnothing] \le 2^{-\Theta(n)}$.

We prove Theorem 3.2 in Section 4.4. Several remarks are in order. Theorem 3.2 shows that for any K > 0 and throughout the entire regime $n = \Theta(M \log M)$, the set of solutions with discrepancy at most K exhibits the m-OGP, for suitable m, β and η . In light of prior work discussed earlier, this gives some rigorous evidence for algorithmic hardness at the boundary $n = \Theta(M \log M)$, and constitutes a first step towards [ANW21, Conjecture 1].

Our proof is based on the first moment method: we show that the expected number of such m-tuples is exponentially small for suitably chosen m, β, η and apply Markov's inequality. Further, our proof reveals that $\beta \gg \eta$: Theorem 3.2 rules out m-tuples of constant discrepancy solutions that are nearly equidistant. Moreover, the solutions need not be of constant discrepancy w.r.t. the same instance: $\mathcal{M}_i(\tau_i)$ appearing in Definition 3.1 are potentially correlated. This is known as the Ensemble m-OGP and instrumental in ruling out stable algorithms in Theorem 3.4.

3.3 m-OGP Implies Failure of Stable Algorithms

In this section, we show that the Ensemble m-OGP established in Theorem 3.2 implies the failure of $stable\ algorithms$ in finding a constant discrepancy solution. We begin by elaborating on the algorithmic setting and formalizing the class of stable algorithms we investigate.

Algorithmic Setting An algorithm \mathcal{A} is a mapping between $\mathbb{R}^{M \times n}$ and Σ_n , where randomization is allowed: we assume there exists a probability space $(\Omega, \mathbb{P}_{\omega})$ such that $\mathcal{A} : \mathbb{R}^{M \times n} \times \Omega \to \Sigma_n$. For any $\omega \in \Omega$ and $\mathcal{M} \in \mathbb{R}^{M \times n}$, we want $\|\mathcal{M}\boldsymbol{\sigma}_{ALG}\|_{\infty} = O(1)$, where $\boldsymbol{\sigma}_{ALG} = \mathcal{A}(\mathcal{M}, \omega) \in \Sigma_n$. The class of stable algorithms is formalized as follows.

Definition 3.3. Fix a K > 0. An algorithm $\mathcal{A} : \mathbb{R}^{M \times n} \times \Omega \to \Sigma_n$ is called $(K, \rho, p_f, p_{\text{st}}, f, L)$ -stable (for discrepancy minimization) if it satisfies the following for all sufficiently large M.

• (Success) For $\mathcal{M} \in \mathbb{R}^{M \times n}$ with i.i.d. $\mathcal{N}(0,1)$ entries,

$$\mathbb{P}_{(\mathcal{M},\omega)}[\|\mathcal{M}\mathcal{A}(\mathcal{M},\omega)\|_{\infty} \leq K] \geq 1 - p_f.$$

• (Stability) Let $\mathcal{M}, \overline{\mathcal{M}} \in \mathbb{R}^{M \times n}$ be random matrices, each with i.i.d. $\mathcal{N}(0,1)$ entries, such that $\mathbb{E}[\mathcal{M}_{ij}\overline{\mathcal{M}}_{ij}] = \rho$ for $1 \leq i \leq M$ and $1 \leq j \leq n$. Then,

$$\mathbb{P}_{(\mathcal{M},\overline{M},\omega)}\left[d_H\left(\mathcal{A}(\mathcal{M},\omega),\mathcal{A}(\overline{\mathcal{M}},\omega)\right) \leq f + L\|\mathcal{M} - \overline{\mathcal{M}}\|_F\right] \geq 1 - p_{\mathrm{st}}.$$

Definition 3.3 is the same as [GKPX22, Definition 3.1]. W.p. at least $1-p_f$, \mathcal{A} finds a solution of discrepancy below K. \mathcal{A} can tolerate an input correlation value of ρ ; and the parameters f and L quantify the sensitivity of the output of \mathcal{A} to changes in its input. The stability guarantee is probabilistic—w.r.t. both $\mathcal{M}, \overline{\mathcal{M}}$ and to the randomness ω of \mathcal{A} —holding w.p. at least $1-p_{\rm st}$. Finally, the term f makes our negative result only stronger: \mathcal{A} is allowed to make f flips even when \mathcal{M} and $\overline{\mathcal{M}}$ are 'too close'. Our final main result is as follows.

Theorem 3.4. Fix a K > 0, $C_1 > c_2 > 0$ and $a \mathcal{L} > 0$. Suppose $C_1 M \log_2 M \ge n \ge c_2 M \log_2 M$. Let $m \in \mathbb{N}$ and $0 < \eta < \beta < 1$ be the m-OGP parameters prescribed by Theorem 3.2. Set

$$C = \frac{\eta^2}{1600}, \quad Q = \frac{4800\mathcal{L}\pi}{\eta^2}, \quad and \quad T = \exp_2\left(2^{4mQ\log_2 Q}\right).$$
 (4)

Then, there exists an $n_0 \in \mathbb{N}$ such that the following holds. For every $n \geq n_0$, there exists no randomized algorithm $\mathcal{A} : \mathbb{R}^{M \times n} \times \Omega \to \Sigma_n$ which, in the sense of Definition 3.3, is

$$\left(K,\cos\left(\frac{\pi}{2Q}\right),\frac{1}{9(Q+1)T},\frac{1}{9Q(T+1)},Cn,\mathcal{L}\sqrt{\frac{n}{M}}\right)-stable.$$

The proof of Theorem 3.4 is almost identical to that of [GKPX22, Theorem 3.2], and omitted for brevity. Several remarks are in order.

Firstly, there is no restriction on the running time of \mathcal{A} : Theorem 3.4 rules out any \mathcal{A} that is stable with suitable parameters in the sense of Definition 3.3. Secondly, observe that $m, \beta, \eta, \mathcal{L}$ are all O(1) (as $n \to \infty$). Hence, C, Q, T per (4) are all O(1), as well. This is an important feature of our result: Theorem 3.4 rules out algorithms with a constant success/stability guarantee. Lastly, since C = O(1), \mathcal{A} is still allowed to make $\Theta(n)$ bit flips even when \mathcal{M} and $\overline{\mathcal{M}}$ are nearly identical.

4 Proofs

Additional Notation We commence this section with an additional list of notation. For any set A, |A| denotes its cardinality. Given any event E, denote its indicator by $\mathbb{1}\{E\}$. For any $v = (v(i): 1 \le i \le n) \in \mathbb{R}^n$ and p > 0, $||v||_p = \left(\sum_{1 \le i \le n} |v(i)|^p\right)^{1/p}$ and $||v||_{\infty} = \max_{1 \le i \le n} |v(i)|$. For $v, v' \in \mathbb{R}^n$, $\langle v, v' \rangle \triangleq \sum_{1 \le i \le n} v(i)v'(i)$. For any $\boldsymbol{\sigma}, \boldsymbol{\sigma}' \in \Sigma_n \triangleq \{-1, 1\}^n$, $d_H(\boldsymbol{\sigma}, \boldsymbol{\sigma}')$ denotes their Hamming distance: $d_H(\boldsymbol{\sigma}, \boldsymbol{\sigma}') \triangleq \sum_{1 \le i \le n} \mathbb{1}\{\boldsymbol{\sigma}(i) \ne \boldsymbol{\sigma}'(i)\}$. For any r > 0, $\log_r(\cdot)$ and $\exp_r(\cdot)$ denote, respectively, the logarithm and exponential functions base r; when r = e, we omit the subscript. For $p \in [0, 1]$, $h_b(p) \triangleq -p \log_2 p - (1-p) \log_2 (1-p)$. Denote by I_k the $k \times k$ identity matrix, and by $\boldsymbol{1}$ the vector of all ones whose dimension will be clear from the context. Given $\boldsymbol{\mu} \in \mathbb{R}^k$ and $\boldsymbol{\Sigma} \in \mathbb{R}^{k \times k}$, denote by $\boldsymbol{\mathcal{N}}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ the multivariate normal distribution in \mathbb{R}^k with mean $\boldsymbol{\mu}$ and covariance $\boldsymbol{\Sigma}$. Given a matrix $\boldsymbol{\mathcal{M}}$, $||\boldsymbol{\mathcal{M}}||_F$, $||\boldsymbol{\mathcal{M}}||_2$, and $|\boldsymbol{\mathcal{M}}|$ denote, respectively, the Frobenius norm, the spectral norm, and the determinant of $\boldsymbol{\mathcal{M}}$.

We employ standard Bachmann-Landau asymptotic notation throughout, e.g. $\Theta(\cdot)$, $O(\cdot)$, $o(\cdot)$, $O(\cdot)$, where the underlying asymptotics will often be clear from the context. In certain cases where a confusion is possible, we reflect the underlying asymptotics as a subscript, e.g. $\Theta_{\kappa}(\cdot)$. All floor/ceiling operators are omitted for the sake of simplicity.

4.1 Proof of Theorem 2.3

Let $\kappa, \alpha > 0$, $M = n\alpha$, $m \in \mathbb{N}$, and $\Delta \in (0, \frac{1}{2})$. Suppose $\mathcal{M}_1 \in \mathbb{R}^{M \times n}$ has i.i.d. $\mathcal{N}(0, 1)$ entries and let $\mathcal{M}_2, \ldots, \mathcal{M}_m \in \mathbb{R}^{M \times n}$ be random matrices obtained from \mathcal{M}_1 by independently resampling the last Δn columns of \mathcal{M}_1 . Denote by $\Xi(m, \Delta)$ the set of all m-tuples satisfying the following:

- $\max_{1 \leq i \leq m} \| \mathcal{M}_i \boldsymbol{\sigma}_i \|_{\infty} \leq \kappa \sqrt{n}$.
- For $1 \le i < j \le n$ and $1 \le k \le n \Delta n$, $\sigma_i(k) = \sigma_j(k)$.

We establish the following proposition.

Proposition 4.1. Fix any $\kappa > 0$ small enough and let $\alpha \geq 4\kappa^2$. Then, there exists an $m \in \mathbb{N}$ and $a \Delta \in (0, 1/2)$ such that

$$\mathbb{P}\big[\Xi(m,\Delta) = \varnothing\big] \ge 1 - e^{-\Theta(n)}.$$

We first assume Proposition 4.1 and show how to deduce Theorem 2.3. Fix a c > 0 and suppose, for the sake of contradiction, that an (e^{-cn}, α) -online \mathcal{A} exists. For $\mathcal{M}_1, \ldots, \mathcal{M}_m \in \mathbb{R}^{M \times n}$ described above, set

$$\sigma_i \triangleq \mathcal{A}(\mathcal{M}_i) \in \Sigma_n, \quad 1 \le i \le m.$$
 (5)

Note that for any $1 \leq i < j \leq m$, the first $n - \Delta n$ columns of \mathcal{M}_i and \mathcal{M}_j are identical. Consequently,

$$\sigma_i(k) = \sigma_j(k)$$
 for $1 \le i < j \le m$ and $1 \le k \le n - \Delta n$.

Next, we establish the following probability guarantee.

Lemma 4.2.

$$\mathbb{P}\left[\max_{1\leq i\leq m} \left\|\mathcal{M}_i \boldsymbol{\sigma}_i\right\|_{\infty} \leq \kappa \sqrt{n}\right] \geq p_s^m.$$

Proof of Lemma 4.2. Our argument is based on a clever application of Jensen's inequality, due to [GS17b, Lemma 5.3]. Denote by ζ the first $(1-\Delta)n$ columns of \mathcal{M}_1 . That is, ζ is the 'common randomness' shared by $\mathcal{M}_1, \ldots, \mathcal{M}_m$. Set

$$I_i = \mathbb{1}\{\|\mathcal{M}_i\boldsymbol{\sigma}_i\|_{\infty} \leq \kappa\sqrt{n}\}.$$

Then,

$$\mathbb{P}\left[\max_{1\leq i\leq m} \left\| \mathcal{M}_i \sigma_i \right\|_{\infty} \leq \kappa \sqrt{n} \right] = \mathbb{E}\left[I_1 \cdots I_m\right].$$

We then complete the proof by noticing

$$\mathbb{E}\big[I_1\cdots I_m\big] = \mathbb{E}_{\zeta}\Big[\mathbb{E}\big[I_1\cdots I_m|\zeta\big]\Big] = \mathbb{E}_{\zeta}\left[\mathbb{E}[I_1|\zeta]^m\right] \ge \big(\mathbb{E}_{\zeta}\big[\mathbb{E}[I_1|\zeta]\big]\big)^m = \mathbb{E}[I_1]^m = p_s^m,$$

where we used fact that I_1, \ldots, I_m are independent conditional on ζ and Jensen's inequality. \square

Note that clearly $(\boldsymbol{\sigma}_1, \dots, \boldsymbol{\sigma}_m) \in \Xi(m, \Delta)$, in particular $\Xi(m, \Delta)$ is non-empty w.p. at least e^{-cmn} . Using Proposition 4.1, we obtain

$$e^{-\Theta(n)} \ge \mathbb{P}[\Xi(m, \Delta) \ne \varnothing] \ge e^{-cmn}.$$

If m is constant and c > 0 is sufficiently small, this is a contradiction for all large enough n. Therefore, it suffices to establish Proposition 4.1.

Proof of Proposition 4.1. Our proof is based on the first moment method. Let

$$S = \{(\boldsymbol{\sigma}_1, \dots, \boldsymbol{\sigma}_m) : \boldsymbol{\sigma}_i(k) = \boldsymbol{\sigma}_j(k), 1 \le i < j \le m, 1 \le k \le n - \Delta n\}.$$

Observe that

$$\left|\Xi(m,\Delta)\right| = \sum_{(\boldsymbol{\sigma}_1,\dots,\boldsymbol{\sigma}_m)\in\mathcal{S}} \mathbb{1}\left\{\max_{1\leq i\leq m} \|\mathcal{M}_i\boldsymbol{\sigma}_i\|_{\infty} \leq \kappa\sqrt{n}\right\}.$$
 (6)

In what follows, we show that for a suitable $m \in \mathbb{N}$ and $\Delta \in (0, 1/2)$,

$$\mathbb{E}\big[|\Xi(m,\Delta)|\big] \le e^{-\Theta(n)}.$$

Counting Estimate We bound |S|. There are 2^n choices for $\sigma_1 \in \Sigma_n$. Having chosen a σ_1 , there are $2^{\Delta n}$ choices for any σ_i , $1 \le i \le m$. So,

$$|\mathcal{S}| \le 2^n \left(2^{\Delta n}\right)^{m-1} \le \exp_2\left(n + nm\Delta\right) \tag{7}$$

Probability Estimate Fix any $(\sigma_1, \ldots, \sigma_m) \in \mathcal{S}$. Denote by $R_1, \ldots, R_m \in \mathbb{R}^n$ the first rows of $\mathcal{M}_1, \ldots, \mathcal{M}_m$, respectively; and set

$$Z_i = \frac{1}{\sqrt{n}} \langle R_i, \boldsymbol{\sigma}_i \rangle \stackrel{d}{=} \mathcal{N}(0, 1), \quad 1 \leq i \leq m.$$

Observe that if $k \neq k'$ or $n - \Delta n + 1 \leq k = k' \leq n$, $\mathbb{E}[R_i(k)R_j(k')] = 0$. Using this fact, we immediately conclude that $\mathbb{E}[Z_iZ_j] = 1 - \Delta$. In particular, $(Z_1, \ldots, Z_m) \in \mathbb{R}^m$ is a centered multivariate normal random vector with covariance Σ , where

$$\Sigma = \Delta I_m + (1 - \Delta) \mathbf{1} \mathbf{1}^T \in \mathbb{R}^{m \times m}$$

where $\mathbf{1} \in \mathbb{R}^m$ is the vector of all ones. In particular, the spectrum of Σ consists of the eigenvalue $\Delta + (1 - \Delta)m$ with multiplicity one and the eigenvalue Δ with multiplicity m - 1. We then obtain

$$\mathbb{P}\left[\max_{1\leq i\leq m}\|\mathcal{M}_{i}\boldsymbol{\sigma}_{i}\|_{\infty}\leq \kappa\sqrt{n}\right] \leq \mathbb{P}\left[\max_{1\leq i\leq m}\left|\langle R_{i},\boldsymbol{\sigma}_{i}\rangle\right| \leq \kappa\sqrt{n}\right]^{\alpha n} \left((2\pi)^{-\frac{m}{2}}|\Sigma|^{-\frac{1}{2}}\int_{\boldsymbol{z}\in[-\kappa,\kappa]^{m}}\exp\left(-\frac{\boldsymbol{z}^{T}\Sigma^{-1}\boldsymbol{z}}{2}\right)\right)^{\alpha n} \leq \left((2\pi)^{-\frac{m}{2}}\left(\Delta+(1-\Delta)m\right)^{-\frac{1}{2}}\Delta^{-\frac{m-1}{2}}(2\kappa)^{m}\right)^{\alpha n}.$$
(8)

Estimating $\mathbb{E}[|\Xi(\Delta, m)|]$ We now combine (7) and (8) to arrive at

$$\mathbb{E}[|\Xi(\Delta, m)|] \le \exp_2(n\Psi(\Delta, m, \alpha)), \tag{9}$$

where

$$\Psi(\Delta, m, \alpha) = 1 + m\Delta - \frac{\alpha m}{2}\log_2(2\pi) + \alpha m\log_2(2\kappa) - \frac{\alpha(m-1)}{2}\log_2\Delta - \frac{\alpha}{2}\log_2(\Delta + (1-\Delta)m).$$

Using the fact $\log_2 \frac{1}{\Delta} > 0$ if $\Delta < \frac{1}{2}$, we further arrive at the bound

$$\Psi(\Delta, m, \alpha) \le m \left(\frac{1}{m} - \frac{\alpha}{2m} \log_2(\Delta + (1 - \Delta)m) + \Upsilon(\Delta, \alpha) \right), \tag{10}$$

for

$$\Upsilon(\Delta,\alpha) = \Delta - \frac{\alpha}{2}\log_2(2\pi) + \alpha\log_2(2\kappa) - \frac{\alpha}{2}\log_2\Delta.$$

Analyzing $\Upsilon(\Delta, \alpha)$ We set $\Delta = (2\kappa)^2$, so that

$$\alpha \log_2(2\kappa) - \frac{\alpha}{2} \log_2 \Delta = 0.$$

Next, fix any $\alpha \geq 4\kappa^2$. Then,

$$\Upsilon(\Delta, \alpha) = \Delta - \frac{\alpha}{2} \log_2(2\pi) \le 4\kappa^2 - 2\kappa^2 \log_2(2\pi) = -\kappa^2 \left(2\log_2(2\pi) - 4\right) = -\Theta_{\kappa}(\kappa^2). \tag{11}$$

Combining everything For fixed small $\kappa > 0$, $\alpha \ge 4\kappa^2$, and $\Delta = (2\kappa)^2$; we have $\Upsilon(\Delta, \alpha) = -\Theta_{\kappa}(\kappa^2) < 0$. Furthermore,

$$\frac{1}{m} - \frac{\alpha}{2m} \log_2(\Delta + (1 - \Delta)m) = o_m(1)$$

as $m \to \infty$. Note that $\Upsilon(\Delta, \alpha)$ depends only on α, κ . So, for $m \in \mathbb{N}$ sufficiently large, (11) yields

$$\frac{1}{m} - \frac{\alpha}{2m} \log_2(\Delta + (1 - \Delta)m) + \Upsilon(\Delta, \alpha) < 0.$$

Hence, combining (9) and (10), we get

$$\mathbb{E}\big[\big|\Xi(m,\Delta)\big|\big] \le e^{-\Theta(n)}.$$

From here, we conclude by Markov's inequality as

$$\mathbb{P}\big[\big|\Xi(\Delta,m)\big|\geq 1\big]\leq \mathbb{E}\big[\big|\Xi(\Delta,m)\big|\big]=\exp\big(-\Theta(n)\big).$$

4.2 Proof of Theorem 2.4

The proof of Theorem 2.4 is similar to that of Theorem 2.3. We first establish the following proposition.

Proposition 4.3. Let $n \geq M = \omega(1)$, $\mathcal{M}_1 \in \{-1,1\}^{M \times n}$ with i.i.d. Rademacher entries and $m = \lceil \frac{2n}{M} \rceil$. Generate $\mathcal{M}_2, \ldots, \mathcal{M}_m \in \{-1,1\}^{M \times n}$ by independently resampling the last M columns of \mathcal{M}_1 . Denote by $\Xi_d(m,M)$ the set of all m-tuples $\sigma_1, \ldots, \sigma_m \in \Sigma_n$ satisfying the following:

- $\max_{1 \le i \le m} \|\mathcal{M}_i \boldsymbol{\sigma}_i\| \le C_u \sqrt{M}$, where $C_u \triangleq \frac{1}{24}$.
- For $1 \le i < j \le m$ and $1 \le k \le n M$, $\sigma_i(k) = \sigma_j(k)$.

Then,

$$\mathbb{P}\big[\Xi_d(m,M) = \varnothing\big] \ge 1 - e^{-n}.$$

Before proving Proposition 4.3, we highlight that if $n = \omega(M)$ then $m = \omega_M(1)$ and the fraction $\Delta = M/n$ of the resampled columns is vanishing. We first show how Proposition 4.3 yields Theorem 2.4. Suppose, for the sake of contradiction, that an $\mathcal{A}: \{-1,1\}^{M\times n} \to \Sigma_n$ which is $(e^{-cM}, \sqrt{M}/24)$ -optimal (with c < 1/2 arbitrary) exists. For \mathcal{M}_i , $1 \le i \le m$, as in the proposition, define

$$\sigma_i \triangleq \mathcal{A}(\mathcal{M}_i) \in \Sigma_n, \quad 1 \leq i \leq n$$

and observe that

$$\sigma_i(k) = \sigma_j(k)$$
, for all $1 \le i < j \le m$ and $1 \le k \le n - M$

as \mathcal{A} is online. We then establish

Lemma 4.4.

$$\mathbb{P}\left[\max_{1\leq i\leq m}\|\mathcal{M}_i\boldsymbol{\sigma}_i\|\leq \frac{\sqrt{M}}{24}\right]\geq \left(e^{-cM}\right)^m\geq e^{-2cn}.$$

Proof of Lemma 4.4 is identical to Lemma 4.2. So, under the assumption that such an \mathcal{A} exists, we obtain $\Xi_d(m,\Delta) \neq \emptyset$ w.p. at least e^{-2cn} . Finally using Proposition 4.3,

$$e^{-n} \ge \mathbb{P}[\Xi_d(m, M) \ne \varnothing] \ge e^{-2cn}$$

which is a contradiction since c < 1/2. Hence, it suffices to establish Proposition 4.3.

Proof of Proposition 4.3. The proof of Proposition 4.3 is similar to Proposition 4.1; it is based in particular on the first moment method. Let

$$\bar{S} = \{(\sigma_1, \dots, \sigma_m) : \sigma_i(k) = \sigma_j(k), 1 \le i < j \le m, 1 \le k \le n - M\}.$$

Note that

$$\left|\Xi_d(m,M)\right| = \sum_{(\sigma_1,\dots,\sigma_m)\in\bar{\mathcal{S}}} \mathbb{1}\left\{\max_{1\leq i\leq m} \|\mathcal{M}_i\boldsymbol{\sigma}_i\|_{\infty} \leq C_u\sqrt{M}\right\}, \quad \text{where} \quad C_u = \frac{1}{24}.$$
 (12)

Counting term We bound $|\bar{S}|$. There are 2^n choices for σ_1 and having fixed it, there are 2^M choices for any σ_i , $2 \le i \le m$. So,

$$|\bar{S}| \le 2^n (2^M)^{m-1} \le \exp_2(n + mM).$$
 (13)

Probability term. Fix an arbitrary $(\sigma_1, \ldots, \sigma_m) \in \bar{S}$. Let $R_i \in \{\pm 1\}^n$, $1 \leq i \leq m$, denote respectively the first rows of \mathcal{M}_i , $1 \leq i \leq m$. For each fixed i, the rows of \mathcal{M}_i are independent. So,

$$\mathbb{P}\left[\max_{1\leq i\leq m}\|\mathcal{M}_{i}\boldsymbol{\sigma}_{i}\|_{\infty}\leq C_{u}\sqrt{M}\right]=\mathbb{P}\left[\max_{1\leq i\leq m}|\langle R_{i},\boldsymbol{\sigma}_{i}\rangle|\leq C_{u}\sqrt{M}\right]^{M}.$$
(14)

Next, let

$$R_i = (R_{ik} : 1 \le k \le n), \quad 1 \le i \le m.$$

Fix any $1 \le i < j \le m$. Observe that the random vectors

$$(R_{ik}: 1 \le k \le n-M)$$
 and $(R_{jk}: 1 \le k \le n-M)$

are identical. For this reason, we drop the first index and use $(R_k : 1 \le k \le n - M)$ instead. Next, fix any $\mathbf{v} = (v_1, \dots, v_{n-M}) \in \{-1, 1\}^{n-M}$ and define

$$\Delta_i(\boldsymbol{v}) \triangleq \sum_{1 \le k \le n-M} v_k \boldsymbol{\sigma}_i(k) \quad \text{and} \quad \Sigma_i \triangleq \sum_{n-M+1 \le k \le n} R_{ik} \boldsymbol{\sigma}_i(k).$$
 (15)

Our goal is to control the right hand side in (14). To that end, our strategy is to condition on R_1, \ldots, R_{n-M} and apply Berry-Esseen inequality for Σ_i . We establish the following auxiliary result.

Lemma 4.5. Let Z_1, \ldots, Z_M be i.i.d. Rademacher random variables, $\epsilon_i \in \{-1, 1\}$, $1 \le i \le M$, be deterministic signs, and $I \subset \mathbb{R}$ be an interval of length $|I| = \omega_M(1)$. Then

$$\mathbb{P}\big[Z_1\epsilon_1 + \dots + Z_M\epsilon_M \in I\big] \le \frac{3|I|}{\sqrt{M}}$$

for every large enough M.

Proof of Lemma 4.5. Let $\frac{1}{\sqrt{M}}I$ denotes the set $\{c/\sqrt{M}:c\in I\}$. By the Central Limit Theorem,

$$\frac{1}{\sqrt{M}} \sum_{1 \le i \le M} Z_i \epsilon_i \Rightarrow \mathcal{N}(0, 1)$$

in distribution, where the speed of convergence is controlled by the Berry-Esseen inequality:

$$\left| \mathbb{P} \left[\sum_{1 \le i \le M} Z_i \epsilon_i \in I \right] - \mathbb{P} \left[\mathcal{N}(0, 1) \in \frac{1}{\sqrt{M}} I \right] \right| \le \frac{C_{\text{be}}}{\sqrt{M}}. \tag{16}$$

Here, $C_{be} > 0$ is an absolute constant. Furthermore, we have

$$\mathbb{P}\left[\mathcal{N}(0,1) \in \frac{1}{\sqrt{M}}I\right] = \frac{1}{\sqrt{2\pi}} \int_{u \in \frac{1}{\sqrt{M}}I} \exp(-u^2/2) \ du \le \frac{|I|}{\sqrt{2\pi M}}.\tag{17}$$

Combining (16) and (17) via triangle inequality, we obtain that for all large enough M,

$$\mathbb{P}\left[\sum_{1 \le i \le M} Z_i \epsilon_i \in I\right] \le \frac{1}{\sqrt{M}} \left(\mathcal{C}_{be} + \frac{|I|}{\sqrt{2\pi}}\right) \le \frac{3|I|}{\sqrt{M}},\tag{18}$$

where we recalled $|I| = \omega_M(1)$ and $C_{be} = O_M(1)$. This establishes Lemma 4.5.

Next, fix any $\boldsymbol{v} \in \{-1,1\}^{n-M}$ and set

$$I_i(\boldsymbol{v}) = \left[-C_u \sqrt{M} - \Delta_i(\boldsymbol{v}), C_u \sqrt{M} - \Delta_i(\boldsymbol{v}) \right],$$

where we recall $\Delta_i(\boldsymbol{v})$ from (15). In particular,

$$|I_i(\boldsymbol{v})| = 2C_u\sqrt{M}$$
, for all $1 \le i \le m$ and $v \in \{-1, 1\}^{n-M}$. (19)

Next fix a $1 \le i \le m$ and recall Σ_i per (15). Applying Lemma 4.5, we conclude that

$$\max_{1 \le i \le m} \max_{\boldsymbol{v} \in \{-1,1\}^{n-M}} \mathbb{P}\Big[\Sigma_i \in I_i(\boldsymbol{v})\Big] \le 6C_u. \tag{20}$$

We are ready to bound the probability term (14) by conditioning on R_1, \ldots, R_{n-M} .

$$\mathbb{P}\left[\max_{1\leq i\leq m} |\langle R_i, \boldsymbol{\sigma}_i \rangle| \leq C_u \sqrt{M}\right] \\
= \sum_{\boldsymbol{v}\in\{-1,1\}^{n-M}} \mathbb{P}\left[\Sigma_i \in I_i(\boldsymbol{v}), 1\leq i\leq m \middle| (R_1, \dots, R_{n-M}) = \boldsymbol{v}\right] \underbrace{\mathbb{P}\left[(R_1, \dots, R_{n-M}) = \boldsymbol{v}\right]}_{=2^{-(n-M)}} \tag{21}$$

$$= 2^{-(n-M)} \sum_{\mathbf{v} \in \{-1,1\}^{n-M}} \mathbb{P} \Big[\Sigma_i \in I_i(\mathbf{v}), 1 \le i \le m \Big]$$

$$= 2^{-(n-M)} \sum_{\mathbf{v} \in \{-1,1\}^{n-M}} \prod_{1 \le i \le m} \mathbb{P} \big[\Sigma_i \in I_i(\mathbf{v}) \big]$$
(23)

$$=2^{-(n-M)}\sum_{v\in\{-1,1\}^{n-M}}\prod_{1\leq i\leq m}\mathbb{P}\big[\Sigma_i\in I_i(\boldsymbol{v})\big]$$
(23)

$$\leq (6C_u)^m.

(24)$$

We now justify the lines above. Equation (21) follows by conditioning on the 'common randomness' R_1, \ldots, R_{n-M} and recalling that they are uniform over $\{-1, 1\}^{n-M}$. Equation (22) uses the fact for any fixed $1 \leq i \leq m$, Σ_i is independent of R_1, \ldots, R_{n-M} , and (23) uses the fact $\Sigma_1, \ldots, \Sigma_m$ is also a collection of independent random variables. Finally, (24) uses (20).

Combining (14) with (24), we thus conclude

$$\max_{(\boldsymbol{\sigma}_1,\dots,\boldsymbol{\sigma}_m)\in\bar{\mathcal{S}}} \mathbb{P}\left[\max_{1\leq i\leq m} \|\mathcal{M}_i\boldsymbol{\sigma}_i\|_{\infty} \leq C_u\sqrt{M}\right] \leq (6C_u)^{mM}.$$
 (25)

Bounding $\mathbb{E}[|\Xi_d(m,M)|]$ We are ready to estimate $\mathbb{E}[|\Xi_d(m,M)|]$. Using (12), (13) and (25),

$$\mathbb{E}[|\Xi_d(m, M)|] \le \exp_2\left(n + mM - mM\log_2\frac{1}{6C_u}\right).$$

Inserting the values $C_u = 1/24$ and $m \ge 2n/M$, we obtain

$$n + mM - mM \log_2 \frac{1}{6C_u} \le -n,$$

so that $\mathbb{E}[|\Xi_d(m,M)|] \leq e^{-n}$. Finally, we conclude by applying Markov's inequality:

$$\mathbb{P}\big[\Xi_d(m,M) \neq \varnothing\big] = \mathbb{P}\big[\big|\Xi_d(m,M)\big| \ge 1\big] \le \mathbb{E}\big[\big|\Xi_d(m,M)\big|\big] \le e^{-n}.$$

4.3 Proof Sketch for Theorem 2.5

The proof of Theorem 2.5 is quite similar to Theorem 2.4; we only highlight the necessary changes. Let \mathcal{M}_1 consists of i.i.d. Bernoulli(p) entries. Suppose that there exists an $\mathcal{A}: \mathbb{R}^{M \times n} \to \Sigma_n$ that is $(e^{-cM}, C'_u \sqrt{M})$ -online in the sense of Definition 2.1, where c < 1/2 is arbitrary and

$$C_u' = \frac{\sqrt{p - p^2}}{24}.$$

We set m = 2n/M and show how to adapt Proposition 4.3 to this case. Once this is done, the rest follows verbatim from Theorem 2.4. First, all instances of C_u in the proof of Theorem 2.4 are replaced with $C'_u = C_u \sqrt{p - p^2}$. Next, the counting estimate per (13) remains intact. Lemma 4.5, on the other hand, is replaced with the following.

Lemma 4.6. Let Z_1, \ldots, Z_M be i.i.d. Bernoulli(p) random variables, $\epsilon_i \in \{-1, 1\}$, $1 \le i \le M$, be deterministic signs, and $I \subset \mathbb{R}$ be an interval of length $|I| = \omega_{(p-p^2)M}(1)$. Then

$$\mathbb{P}\big[Z_1\epsilon_1 + \dots + Z_M\epsilon_M \in I\big] \le \frac{3|I|}{\sqrt{M(p-p^2)}},$$

for every large enough M.

Proof of Lemma 4.6. Observe that $\mathbb{E}[Z_i \epsilon_i] = p \epsilon_i$ and

$$Var(Z_i\epsilon_i) = \mathbb{E}[Z_i^2\epsilon_i^2] - p^2\epsilon_i^2 = p - p^2,$$

as $\epsilon_i \in \{-1, 1\}$. Thus by the CLT,

$$\frac{1}{\sqrt{(p-p^2)M}} \left(\sum_{1 \le i \le M} Z_i \epsilon_i - p \left\langle \mathbf{1}, \boldsymbol{\epsilon} \right\rangle \right) \Rightarrow \mathcal{N}(0, 1)$$

in distribution, where $\mathbf{1} \in \mathbb{R}^M$ is the vector of all ones and $\boldsymbol{\epsilon} = (\epsilon_i : 1 \leq i \leq M) \in \{-1, 1\}^M$. Further, by the Berry-Esseen inequality, we have that

$$\left| \mathbb{P} \left[\sum_{1 \le i \le M} Z_i \epsilon_i \in I \right] - \mathbb{P} \left[\mathcal{N}(0, 1) \in \frac{I - p \langle \mathbf{1}, \boldsymbol{\epsilon} \rangle}{\sqrt{(p - p^2)M}} \right] \right| \le \frac{C'_{\text{be}}}{\sqrt{(p - p^2)M}}$$

for some absolute constant $C'_{be} > 0$. Here,

$$\frac{I - p \langle \mathbf{1}, \boldsymbol{\epsilon} \rangle}{\sqrt{(p - p^2)M}} = \left\{ \frac{c - p \langle \mathbf{1}, \boldsymbol{\epsilon} \rangle}{\sqrt{(p - p^2)M}} : c \in I \right\},\,$$

so that

$$\left| \frac{I - p \langle \mathbf{1}, \boldsymbol{\epsilon} \rangle}{\sqrt{(p - p^2)M}} \right| = \frac{|I|}{\sqrt{(p - p^2)M}},$$

using the translation invariance of Lebesgue measure. From here, proceeding in the exact same way as in the proof of Lemma 4.5, we establish Lemma 4.6.

Equipped with Lemma 4.6 and using the exact same notation, (20) modifies to (26) where

$$\max_{1 \le i \le m} \max_{\mathbf{v} \in \{-1,1\}^{n-M}} \mathbb{P}\Big[\Sigma_i \in I_i(\mathbf{v})\Big] \le \frac{6C_u'}{\sqrt{p-p^2}} = \frac{1}{4}.$$
 (26)

We now proceed analogously to lines (21)-(24). Note that for any arbitrary $\mathbf{v} \in \{0,1\}^{n-M}$,

$$\mathbb{P}\Big[\Sigma_i \in I_i(\boldsymbol{v}), 1 \le i \le m \Big| (R_1, \dots, R_{n-M}) = \boldsymbol{v} \Big] \le 2^{-2m}, \tag{27}$$

using (26). Hence,

$$\mathbb{P}\left[\max_{1\leq i\leq m} |\langle R_i, \boldsymbol{\sigma}_i \rangle| \leq C'_{\boldsymbol{u}} \sqrt{M}\right] \\
\leq \sum_{\boldsymbol{v}\in\{0,1\}^{n-M}} \mathbb{P}\left[\Sigma_i \in I_i(\boldsymbol{v}), 1\leq i\leq m \middle| (R_1, \dots, R_{n-M}) = \boldsymbol{v}\right] \mathbb{P}\left[(R_1, \dots, R_{n-M}) = \boldsymbol{v}\right] \\
\leq 2^{-2m} \sum_{\boldsymbol{v}\in\{0,1\}^{n-M}} \mathbb{P}\left[(R_1, \dots, R_{n-M}) = \boldsymbol{v}\right] \\
= 2^{-2m},$$

where we used (27) in the penultimate line. This is precisely the same bound as (24), so the rest of the proof remains intact. This completes the proof of Theorem 2.5.

4.4 Proof of Theorem 3.2

Fix a K > 0, $C_1 > c_2 > 0$, and suppose

$$C_1 M \log_2 M \ge n \ge c_2 M \log_2 M. \tag{28}$$

We establish our result via the first-moment method. Notice that by Markov's inequality,

$$\mathbb{P}[\mathcal{S}(K, m, \beta, \eta, \mathcal{I}) \neq \varnothing] = \mathbb{P}[|\mathcal{S}(K, m, \beta, \eta, \mathcal{I})| \geq 1] \leq \mathbb{E}[|\mathcal{S}(K, m, \beta, \eta, \mathcal{I})|].$$

So, it suffices to prove that

$$\mathbb{E}[|\mathcal{S}(K, m, \beta, \eta, \mathcal{I})|] \le 2^{-\Theta(n)}.$$

We now estimate $\mathbb{E}[|\mathcal{S}(K, m, \beta, \eta, \mathcal{I})|]$.

Counting term Fix $m \in \mathbb{N}$, $0 < \eta < \beta < 1$ and denote by $M(m, \beta, \eta)$ the number of m-tuples $(\sigma_i \in \Sigma_n : 1 \le i \le m)$ such that $\beta - \eta \le n^{-1} \langle \sigma_i, \sigma_j \rangle \le \beta$ for $1 \le i < j \le m$. We establish

Lemma 4.7. For m = O(1) as $n \to \infty$,

$$M(m,\beta,\eta) \le \exp_2\left(n + n(m-1)h_b\left(\frac{1-\beta+\eta}{2}\right) + O(\log_2 n)\right).$$

Lemma 4.7 is verbatim from [GKPX22, Lemma 6.7], we include the proof for completeness.

Proof of Lemma 4.7. Observe that $\langle \boldsymbol{\sigma}, \boldsymbol{\sigma}' \rangle = n - 2d_H(\boldsymbol{\sigma}, \boldsymbol{\sigma}')$ for any $\boldsymbol{\sigma}, \boldsymbol{\sigma}' \in \Sigma_n$. There are 2^n choices for $\boldsymbol{\sigma}_1$. Having fixed a $\boldsymbol{\sigma}_1$, there are

$$\sum_{\substack{\rho: \frac{1-\beta}{2} \le \rho \le \frac{1-\beta+\eta}{2} \\ n \in \mathbb{N}}} \binom{n}{n \rho} \le \binom{n}{n \frac{1-\beta+\eta}{2}} n^{O(1)},$$

choices for any σ_i , $2 \le i \le m$, under the constraint $\beta - \eta \le n^{-1} \langle \sigma_1, \sigma_i \rangle \le \beta$. Next, for any $\rho \in (0,1)$, $\binom{n}{n\rho} = \exp_2(nh(\rho) + O(\log_2 n))$ by Stirling's approximation. Combining these and recalling $m = O_n(1)$, we obtain Lemma 4.7.

Probability estimate Fix any $(\sigma_1, \ldots, \sigma_m)$ with

$$\frac{1}{n} \langle \boldsymbol{\sigma}_i, \boldsymbol{\sigma}_j \rangle = \beta - \eta_{ij}, \quad 1 \le i < j \le m.$$

Clearly $0 \le \eta_{ij} \le \eta$. Further, let $\boldsymbol{\eta} = (\eta_{ij} : 1 \le i < j \le m) \in \mathbb{R}^{m(m-1)/2}$. Then $\|\boldsymbol{\eta}\|_{\infty} \le \eta$. Our eventual choice of parameters β, η and m will ensure

$$\eta = \frac{1 - \beta}{2m}.\tag{29}$$

We now control the probability term.

Lemma 4.8. Let $\Sigma(\eta) \in \mathbb{R}^{m \times m}$ with unit diagonal entries such that for $1 \leq i < j \leq m$,

$$(\Sigma(\boldsymbol{\eta}))_{ij} = (\Sigma(\boldsymbol{\eta}))_{ji} = \beta - \eta_{ij}.$$

Then, the following holds.

- (a) $\Sigma(\eta)$ is positive definite (PD) if η satisfies (29).
- (b) Suppose that η satisfies (29). Then,

$$\mathbb{P}\left[\exists \tau_1, \dots, \tau_m \in \mathcal{I} : \max_{1 \le i \le m} \left\| \mathcal{M}_i(\tau_i) \boldsymbol{\sigma}_i \right\|_{\infty} \le K \right] \le |\mathcal{I}|^m (2\pi)^{-\frac{mM}{2}} \left(\frac{1-\beta}{2}\right)^{-\frac{Mm}{2}} \left(\frac{2K}{\sqrt{n}}\right)^{Mm}.$$

Proof of Lemma 4.8.

Part (a) Let $E \in \mathbb{R}^{m \times m}$ such that $E_{ii} = 0$ and $E_{ij} = E_{ji} = -\eta_{ij}$ for $1 \le i < j \le m$. Then,

$$\Sigma(\boldsymbol{\eta}) = (1 - \beta)I + \beta \mathbf{1} \mathbf{1}^T + E.$$

Note that the smallest eigenvalue of $(1 - \beta)I + \beta \mathbf{1}\mathbf{1}^T$ is $1 - \beta$ and $||E||_2 \le ||E||_F < \eta m$. So, $\Sigma(\boldsymbol{\eta})$ is invertible if $\eta < (1 - \beta)/m$. Recalling the fact it is a covariance matrix, so in particular positive semidefinite, we establish part (a).

Part (b) As a first step, we take a union bound over \mathcal{I} to obtain

$$\mathbb{P}\left[\exists \tau_1, \dots, \tau_m \in \mathcal{I} : \max_{1 \le i \le m} \left\| \mathcal{M}_i(\tau_i) \boldsymbol{\sigma}_i \right\|_{\infty} \le K \right] \le |\mathcal{I}|^m \max_{\tau_i \in \mathcal{I}, 1 \le i \le m} \mathbb{P}\left[\max_{1 \le i \le m} \left\| \mathcal{M}_i(\tau_i) \boldsymbol{\sigma}_i \right\|_{\infty} \le K \right]. \tag{30}$$

Next, denote by $R_i \sim \mathcal{N}(0, I_n)$ the first row of $\mathcal{M}_i(\tau_i) \in \mathbb{R}^{M \times n}$, $1 \leq i \leq m$. Observe that using the fact each $\mathcal{M}_i(\tau_i)$ has independent rows,

$$\mathbb{P}\left[\max_{1\leq i\leq m}\left\|\mathcal{M}_{i}(\tau_{i})\boldsymbol{\sigma}_{i}\right\|_{\infty}\leq K\right]\leq \mathbb{P}\left[\max_{1\leq i\leq m}n^{-\frac{1}{2}}\left|\langle R_{i},\boldsymbol{\sigma}_{i}\rangle\right|\leq \frac{K}{\sqrt{n}}\right]^{M}.$$
(31)

Next, we consider the multivariate normal random vector $(n^{-1/2} \langle R_i, \boldsymbol{\sigma}_i \rangle : 1 \leq i \leq m)$ consisting of standard normal coordinates. Let $\overline{\Sigma}$ denotes its covariance matrix, which depends on the choice of τ_1, \ldots, τ_m . Observe that for $1 \leq i < j \leq m$,

$$\overline{\Sigma}_{ij} = \frac{1}{n} \mathbb{E} \left[\langle R_i, \boldsymbol{\sigma}_i \rangle \langle R_j, \boldsymbol{\sigma}_j \rangle \right] = \frac{1}{n} (\boldsymbol{\sigma}_i)^T \underbrace{\mathbb{E} \left[R_i R_j^T \right]}_{=\cos(\tau_i)\cos(\tau_j)I_m} \boldsymbol{\sigma}_j = \cos(\tau_i)\cos(\tau_j)(\beta - \eta_{ij}).$$

We now remove the dependence on τ_i by relying on a Gaussian comparison inequality, due to Sidák [Sid68, Corollary 1]. The version below is reproduced from [GKPX22, Theorem 6.5].

Theorem 4.9. Let $(X_1, \ldots, X_k) \in \mathbb{R}^k$ be a centered multivariate normal random vector. Suppose that its covariance matrix $\Sigma \in \mathbb{R}^{k \times k}$ has unit diagonal entries has the following form: there exists $0 \le \lambda_i \le 1$, $1 \le i \le k$, such that for any $1 \le i \ne j \le k$, $\Sigma_{ij} = \lambda_i \lambda_j \rho_{ij}$ where $(\rho_{ij} : 1 \le i \ne j \le k)$ is a fixed arbitrary covariance matrix. Fix values $c_1, \ldots, c_k > 0$, and set

$$P(\lambda_1, \ldots, \lambda_k) = \mathbb{P}[|X_1| < c_1, |X_2| < c_2, \ldots, |X_k| < c_k].$$

Then, $P(\lambda_1, \ldots, \lambda_k)$ is a non-decreasing function of each λ_i , $i = 1, 2, \ldots, k$, $0 \le \lambda_i \le 1$. That is,

$$P(\lambda_1, \lambda_2, \dots, \lambda_k) \leq P(1, 1, \dots, 1).$$

We now let (Z_1, \ldots, Z_m) to be a centered multivariate normal random vector with covariance $\Sigma(\boldsymbol{\eta})$. Observe that

$$\max_{\tau_{1},\dots,\tau_{m}\in\mathcal{I}}\mathbb{P}\left[\max_{1\leq i\leq m}n^{-\frac{1}{2}}\left|\langle R_{i},\boldsymbol{\sigma}_{i}\rangle\right|\leq\frac{K}{\sqrt{n}}\right]\leq\mathbb{P}\left[\max_{1\leq i\leq m}\left|Z_{i}\right|\leq\frac{K}{\sqrt{n}}\right] \qquad (32)$$

$$=(2\pi)^{-\frac{m}{2}}|\Sigma(\boldsymbol{\eta})|^{-\frac{1}{2}}\int_{\boldsymbol{z}\in\left[-\frac{K}{\sqrt{n}},\frac{K}{\sqrt{n}}\right]^{m}}\exp\left(-\frac{\boldsymbol{z}^{T}\Sigma(\boldsymbol{\eta})^{-1}\boldsymbol{z}}{2}\right)\,d\boldsymbol{z}$$

$$\leq(2\pi)^{-\frac{m}{2}}|\Sigma(\boldsymbol{\eta})|^{-\frac{1}{2}}\left(\frac{2K}{\sqrt{n}}\right)^{m},$$
(33)

where (32) follows from Theorem 4.9 and (33) follows from the trivial fact $\exp\left(-\frac{z^T\Sigma(\eta)^{-1}z}{2}\right) \leq 1$. We lastly bound $|\Sigma(\eta)|$. For this, we rely on the following tool from matrix analysis.

Theorem 4.10 (Hoffman-Wielandt Inequality). Let $A \in \mathbb{R}^{m \times m}$ and $A + E \in \mathbb{R}^{m \times m}$ be two symmetric matrices with eigenvalues

$$\lambda_1(A) \ge \cdots \ge \lambda_m(A)$$
 and $\lambda_1(A+E) \ge \cdots \ge \lambda_m(A+E)$.

Then,

$$\sum_{1 \le i \le m} (\lambda_i(A+E) - \lambda_i(A))^2 \le ||E||_F.$$

See [HJ12, Corollary 6.3.8] for a reference, and [HW53] for the original paper. We apply Theorem 4.10 to $\Sigma(\eta)$. Let $A = (1-\beta)I + \beta \mathbf{1}\mathbf{1}^T$ with eigenvalues $\lambda_1 = 1 - \beta + \beta m > \lambda_2 = \cdots = \lambda_m = 1 - \beta$ and E be as above. Suppose that the eigenvalues of A + E are $\mu_1 \geq \cdots \geq \mu_m$. Fix any $2 \leq i \leq m$. Theorem 4.10 yields

$$|\mu_i - (1 - \beta)| \le ||E||_F \le \eta m = \frac{1 - \beta}{2},$$

yielding

$$\mu_i \ge \frac{1-\beta}{2}, \quad 2 \le i \le m.$$

Furthermore, this bounds extends to μ_1 , too, as

$$\mu_1 \ge 1 - \beta + \beta m - \frac{1 - \beta}{2} > \frac{1 - \beta}{2}.$$

Since $\eta \in \mathbb{R}^{m(m-1)/2}$ is arbitrary with $\|\eta\|_{\infty} \leq \eta \leq \frac{1-\beta}{2m}$, we obtain

$$\inf_{\substack{\boldsymbol{\eta} \in \mathbb{R}^{m(m-1)/2} \\ \|\boldsymbol{\eta}\|_{\infty} \leq \frac{1-\beta}{2}}} \left| \Sigma(\boldsymbol{\eta}) \right| = \prod_{1 \leq i \leq m} \mu_i \geq \left(\frac{1-\beta}{2}\right)^m. \tag{34}$$

Finally, combining (30), (31), (33), and (34) we establish the proof of part (b).

Estimating the expectation Let $\mathcal{F}(m, \beta, \eta)$ be the set of all *m*-tuples $(\boldsymbol{\sigma}_1, \dots, \boldsymbol{\sigma}_m)$ such that $\beta - \eta \leq n^{-1} \langle \boldsymbol{\sigma}_i, \boldsymbol{\sigma}_j \rangle \leq \beta, \ 1 \leq i < j \leq m$. Then

$$\left| \mathcal{S}(K, m, \beta, \eta, \mathcal{I}) \right| = \sum_{(\boldsymbol{\sigma}_1, \dots, \boldsymbol{\sigma}_m) \in \mathcal{F}(m, \beta, \eta)} \mathbb{1} \left\{ \exists \tau_1, \dots, \tau_m \in \mathcal{I} : \max_{1 \le i \le m} \left\| \mathcal{M}_i(\tau_i) \boldsymbol{\sigma}_i \right\|_{\infty} \le K \right\}.$$

Using linearity of expectation, Lemma 4.7, Lemma 4.8, and the fact $\log_2 |\mathcal{I}| \leq cn$, we obtain

$$\mathbb{E}[\left|\mathcal{S}(K, m, \beta, \eta, \mathcal{I})\right|] \le \exp_2\left(\Psi(m, \beta, \eta, c) + O(\log_2 n)\right),\tag{35}$$

where

$$\Psi(m, \beta, \eta, c) = n + mnh_b \left(\frac{1 - \beta + \eta}{2}\right) + cmn + \frac{mM}{2}\log_2\frac{4K^2}{\pi(1 - \beta)} - \frac{Mm}{2}\log_2 n.$$
 (36)

We set η and c as

$$\eta = \frac{1 - \beta}{2m} \quad \text{and} \quad c = \frac{1}{m},\tag{37}$$

where we recalled η from (29); parameters β and m are to be tuned soon. We now recall the scaling on n from (28). In particular,

$$\log_2 n \ge \log_2 c_2 + \log_2 M + \log_2 \log_2 M \ge \log_2 c_2 + \log_2 M$$
.

With this, we arrive at

$$\Psi\left(m, \beta, \frac{1-\beta}{2m}, \frac{1}{m}\right) \leq 2C_1 M \log_2 M + mC_1 M \log_2 M \cdot h_b \left(\frac{1-\beta}{2} + \frac{1-\beta}{4m}\right) + \frac{mM}{2} \log_2 \frac{4K^2}{\pi(1-\beta)c_2} - \frac{Mm}{2} \log_2 M.$$
(38)

Note that if $\beta \in (1/2, 1)$ and $m \in \mathbb{N}$, we clearly have

$$h_b\left(\frac{1-\beta}{2} + \frac{1-\beta}{4m}\right) \le h_b(1-\beta).$$

We choose $\beta^* > 1/2$ such that

$$h_b(1-\beta^*) = \min\left\{\frac{1}{4C_1}, \frac{1}{2}\right\}.$$

So,

$$mC_1 M \log_2 M \cdot h_b \left(\frac{1 - \beta^*}{2} + \frac{1 - \beta^*}{4m} \right) \le \frac{Mm}{4} \log_2 M.$$
 (39)

Combining (38) and (39), we further upper bound

$$\Psi\left(m, \beta^*, \frac{1-\beta^*}{2m}, \frac{1}{m}\right) \le 2C_1 M \log_2 M - \frac{Mm}{4} \log_2 M + \Theta(mM). \tag{40}$$

Finally, taking $m = m^* = \max\{2, 16C_1\}$, we get

$$\Psi\left(m^*, \beta^*, \frac{1 - \beta^*}{2m^*}, \frac{1}{m^*}\right) = -\Theta(M \log_2 M). \tag{41}$$

Combining (35) with the fact $O(\log_2 n) = O(\log_2 M) = o(M \log_2 M)$ as $M = \omega(1)$, we conclude that

$$\mathbb{E}\big[\big|\mathcal{S}(K, m^*, \beta^*, \eta^*, \mathcal{I})\big|\big] \leq \exp_2\left(\Psi\left(m^*, \beta^*, \frac{1-\beta^*}{2m^*}, \frac{1}{m^*}\right) + o(M\log_2 M)\right) = 2^{-\Theta(M\log M)} = 2^{-\Theta(n)}.$$

This completes the proof of Theorem 3.2.

Acknowledgments

The first author is supported in part by NSF grant DMS-2015517. The second author is supported by a Columbia University, Distinguished Postdoctoral Fellowship in Statistics. The third author is supported in part by NSF grant DMS-1847451.

References

- [ACO08] Dimitris Achlioptas and Amin Coja-Oghlan, Algorithmic barriers from phase transitions, 2008 49th Annual IEEE Symposium on Foundations of Computer Science, IEEE, 2008, pp. 793–802.
- [Ajt96] Miklós Ajtai, Generating hard instances of lattice problems, Proceedings of the twenty-eighth annual ACM symposium on Theory of computing, 1996, pp. 99–108.
- [ALS21a] Emmanuel Abbe, Shuangping Li, and Allan Sly, Binary perceptron: efficient algorithms can find solutions in a rare well-connected cluster, arXiv preprint arXiv:2111.03084 (2021).
- [ALS21b] _____, Proof of the contiguity conjecture and lognormal limit for the symmetric perceptron, arXiv preprint arXiv:2102.13069 (2021).
- [Alt22] Dylan J Altschuler, Fluctuations of the symmetric perceptron, arXiv preprint arXiv:2205.02319 (2022).
- [ANW21] Dylan J Altschuler and Jonathan Niles-Weed, *The discrepancy of random rectangular matrices*, Random Structures & Algorithms (2021).
- [APZ19] Benjamin Aubin, Will Perkins, and Lenka Zdeborová, Storage capacity in symmetric binary perceptrons, Journal of Physics A: Mathematical and Theoretical **52** (2019), no. 29, 294003.
- [ART06] Dimitris Achlioptas and Federico Ricci-Tersenghi, On the solution-space geometry of random constraint satisfaction problems, Proceedings of the thirty-eighth annual ACM symposium on Theory of computing, 2006, pp. 130–139.
- [AS20] Ahmed El Alaoui and Mark Sellke, Algorithmic pure states for the negative spherical perceptron, arXiv preprint arXiv:2010.15811 (2020).
- [AWZ20] Gérard Ben Arous, Alexander S Wein, and Ilias Zadik, Free energy wells and overlap gap property in sparse pca, Conference on Learning Theory, PMLR, 2020, pp. 479–482.
- [BABB21] Enric Boix-Adserà, Matthew Brennan, and Guy Bresler, *The average-case complex-ity of counting cliques in Erdös-Rényi hypergraphs*, SIAM Journal on Computing (2021), no. 0, FOCS19–39.
- [Ban10] Nikhil Bansal, Constructive algorithms for discrepancy minimization, 2010 IEEE 51st Annual Symposium on Foundations of Computer Science, IEEE, 2010, pp. 3–10.
- [Ban22] _____, personal communication, 2022.
- [BB19] Matthew Brennan and Guy Bresler, Optimal average-case reductions to sparse pca: From weak assumptions to strong hardness, arXiv preprint arXiv:1902.07380 (2019).

- [BBBZ07] Carlo Baldassi, Alfredo Braunstein, Nicolas Brunel, and Riccardo Zecchina, Efficient supervised learning in networks with binary synapses, Proceedings of the National Academy of Sciences 104 (2007), no. 26, 11079–11084.
- [BBH18] Matthew Brennan, Guy Bresler, and Wasim Huleihel, Reducibility and computational lower bounds for problems with planted sparse structure, arXiv preprint arXiv:1806.07508 (2018).
- [BDVLZ20] Carlo Baldassi, Riccardo Della Vecchia, Carlo Lucibello, and Riccardo Zecchina, Clustering of solutions in the symmetric binary perceptron, Journal of Statistical Mechanics: Theory and Experiment 2020 (2020), no. 7, 073303.
- [BGT10] Mohsen Bayati, David Gamarnik, and Prasad Tetali, Combinatorial approach to the interpolation method and scaling limits in sparse random graphs, Proceedings of the forty-second ACM symposium on Theory of computing, 2010, pp. 105–114.
- [BH21] Guy Bresler and Brice Huang, The algorithmic phase transition of random k-sat for low degree polynomials, arXiv preprint arXiv:2106.02129 (2021).
- [BHK⁺19] Boaz Barak, Samuel Hopkins, Jonathan Kelner, Pravesh K Kothari, Ankur Moitra, and Aaron Potechin, *A nearly tight sum-of-squares lower bound for the planted clique problem*, SIAM Journal on Computing **48** (2019), no. 2, 687–735.
- [BIL⁺15] Carlo Baldassi, Alessandro Ingrosso, Carlo Lucibello, Luca Saglietti, and Riccardo Zecchina, Subdominant dense clusters allow for simple learning and high computational performance in neural networks with discrete synapses, Physical review letters 115 (2015), no. 12, 128101.
- [BNSX21] Erwin Bolthausen, Shuta Nakajima, Nike Sun, and Changji Xu, Gardner formula for Ising perceptron models at small densities, arXiv preprint arXiv:2111.02855 (2021).
- [BPW18] Afonso S Bandeira, Amelia Perry, and Alexander S Wein, *Notes on computational-to-statistical gaps: predictions using statistical physics*, Portugaliae Mathematica **75** (2018), no. 2, 159–186.
- [BR13] Quentin Berthet and Philippe Rigollet, Computational lower bounds for sparse PCA, arXiv preprint arXiv:1304.0828 (2013).
- [BS20] Nikhil Bansal and Joel H. Spencer, On-line balancing of random inputs, Random Structures and Algorithms 57 (2020), no. 4, 879–891 (English (US)).
- [CGPR19] Wei-Kuo Chen, David Gamarnik, Dmitry Panchenko, and Mustazee Rahman, Suboptimality of local algorithms for a class of max-cut problems, The Annals of Probability 47 (2019), no. 3, 1587–1618.
- [COE15] Amin Coja-Oghlan and Charilaos Efthymiou, On independent sets in random graphs, Random Structures & Algorithms 47 (2015), no. 3, 436–486.
- [Cos09] Kevin P Costello, *Balancing gaussian vectors*, Israel Journal of Mathematics **172** (2009), no. 1, 145–156.

- [Cov65] Thomas M Cover, Geometrical and statistical properties of systems of linear inequalities with applications in pattern recognition, IEEE transactions on electronic computers (1965), no. 3, 326–334.
- [DFJ02] Martin Dyer, Alan Frieze, and Mark Jerrum, On counting independent sets in sparse graphs, SIAM Journal on Computing 31 (2002), no. 5, 1527–1541.
- [DKS17] Ilias Diakonikolas, Daniel M Kane, and Alistair Stewart, Statistical query lower bounds for robust estimation of high-dimensional Gaussians and Gaussian mixtures, 2017 IEEE 58th Annual Symposium on Foundations of Computer Science (FOCS), IEEE, 2017, pp. 73–84.
- [DM15] Yash Deshpande and Andrea Montanari, Improved sum-of-squares lower bounds for hidden clique and hidden submatrix problems, Conference on Learning Theory, 2015, pp. 523–562.
- [DS19] Jian Ding and Nike Sun, Capacity lower bound for the Ising perceptron, Proceedings of the 51st Annual ACM SIGACT Symposium on Theory of Computing, 2019, pp. 816–827.
- [FGR⁺17] Vitaly Feldman, Elena Grigorescu, Lev Reyzin, Santosh S Vempala, and Ying Xiao, Statistical algorithms and a lower bound for detecting planted cliques, Journal of the ACM (JACM) **64** (2017), no. 2, 1–37.
- [FL92] Alan M Frieze and T Luczak, On the independence and chromatic numbers of random regular graphs, Journal of Combinatorial Theory, Series B **54** (1992), no. 1, 123–132.
- [FPV18] Vitaly Feldman, Will Perkins, and Santosh Vempala, On the complexity of random satisfiability problems with planted solutions, SIAM Journal on Computing 47 (2018), no. 4, 1294–1338.
- [FS20] Cole Franks and Michael Saks, On the discrepancy of random matrices with many columns, Random Structures & Algorithms 57 (2020), no. 1, 64–96.
- [Gam21] David Gamarnik, The overlap gap property: A topological barrier to optimizing over random structures, Proceedings of the National Academy of Sciences 118 (2021), no. 41.
- [Gar87] Elizabeth Gardner, Maximum storage capacity in neural networks, EPL (Europhysics Letters) 4 (1987), no. 4, 481.
- [Gar88] _____, The space of interactions in neural network models, Journal of physics A: Mathematical and general **21** (1988), no. 1, 257.
- [GD88] Elizabeth Gardner and Bernard Derrida, Optimal storage properties of neural network models, Journal of Physics A: Mathematical and general 21 (1988), no. 1, 271.

- [GJ21] David Gamarnik and Aukosh Jagannath, The overlap gap property and approximate message passing algorithms for p-spin models, The Annals of Probability 49 (2021), no. 1, 180–205.
- [GJS21] David Gamarnik, Aukosh Jagannath, and Subhabrata Sen, *The overlap gap property* in principal submatrix recovery, Probability Theory and Related Fields **181** (2021), 757–814.
- [GJW20] David Gamarnik, Aukosh Jagannath, and Alexander S Wein, Low-degree hardness of random optimization problems, 2020 IEEE 61st Annual Symposium on Foundations of Computer Science (FOCS), IEEE, 2020, pp. 131–140.
- [GJW21] _____, Circuit lower bounds for the p-spin optimization problem, arXiv preprint arXiv:2109.01342 (2021).
- [GK21a] David Gamarnik and Eren C Kızıldağ, Algorithmic obstructions in the random number partitioning problem, arXiv preprint arXiv:2103.01369 (2021).
- [GK21b] David Gamarnik and Eren C. Kızıldağ, Computing the partition function of the Sherrington-Kirkpatrick model is hard on average, The Annals of Applied Probability **31** (2021), no. 3, 1474 1504.
- [GKPX22] David Gamarnik, Eren C Kızıldağ, Will Perkins, and Changji Xu, Algorithms and barriers in the symmetric binary perceptron model, arXiv preprint arXiv:2203.15667 (2022).
- [GMZ22] David Gamarnik, Cristopher Moore, and Lenka Zdeborová, *Disordered systems insights on computational hardness*, Journal of Statistical Mechanics: Theory and Experiment **2022** (2022), no. 11, 114015.
- [GS14] David Gamarnik and Madhu Sudan, Limits of local algorithms over sparse random graphs, Proceedings of the 5th conference on Innovations in theoretical computer science, 2014, pp. 369–376.
- [GS17a] _____, Limits of local algorithms over sparse random graphs, Ann. Probab. 45 (2017), no. 4, 2353–2376.
- [GS17b] _____, Performance of sequential local algorithms for the random NAE-K-SAT problem, SIAM Journal on Computing 46 (2017), no. 2, 590–619.
- [HJ12] Roger A Horn and Charles R Johnson, *Matrix analysis*, Cambridge University Press, 2012.
- [HK14] Haiping Huang and Yoshiyuki Kabashima, Origin of the computational hardness for learning with binary synapses, Physical Review E **90** (2014), no. 5, 052813.
- [HKP⁺17] Samuel B Hopkins, Pravesh K Kothari, Aaron Potechin, Prasad Raghavendra, Tselil Schramm, and David Steurer, *The power of sum-of-squares for detecting hidden structures*, 2017 IEEE 58th Annual Symposium on Foundations of Computer Science (FOCS), IEEE, 2017, pp. 720–731.

- [Hop18] Samuel Brink Klevit Hopkins, Statistical inference and the sum of squares method.
- [HR19] Rebecca Hoberg and Thomas Rothvoss, A fourier-analytic approach for the discrepancy of random set systems, Proceedings of the Thirtieth Annual ACM-SIAM Symposium on Discrete Algorithms, SIAM, 2019, pp. 2547–2556.
- [HS21] Brice Huang and Mark Sellke, Tight lipschitz hardness for optimizing mean field spin glasses, arXiv preprint arXiv:2110.07847 (2021).
- [HSS15] Samuel B Hopkins, Jonathan Shi, and David Steurer, *Tensor principal component analysis via sum-of-square proofs*, Conference on Learning Theory, 2015, pp. 956–1006.
- [HW53] AJ Hoffman and HW Wielandt, The variation of the spectrum of a normal matrix, Duke Mathematical Journal **20** (1953), no. 1, 37–39.
- [Jer92] Mark Jerrum, Large cliques elude the metropolis process, Random Structures & Algorithms 3 (1992), no. 4, 347–359.
- [JH60] Roger David Joseph and Louise Hay, The number of orthants in n-space intersected by an s-dimensional subspace, Tech. report, CORNELL AERONAUTICAL LAB INC BUFFALO NY, 1960.
- [Kız22] Eren C Kızıldağ, Algorithms and algorithmic barriers in high-dimensional statistics and random combinatorial structures, Ph.D. thesis, Massachusetts Institute of Technology, 2022.
- [KK82] Narendra Karmarkar and Richard M Karp, *The differencing method of set partition*ing, Computer Science Division (EECS), University of California Berkeley, 1982.
- [KKLO86] Narendra Karmarkar, Richard M Karp, George S Lueker, and Andrew M Odlyzko, Probabilistic analysis of optimum partitioning, Journal of Applied probability 23 (1986), no. 3, 626–645.
- [KM89] Werner Krauth and Marc Mézard, Storage capacity of memory networks with binary couplings, Journal de Physique **50** (1989), no. 20, 3057–3066.
- [KR98] Jeong Han Kim and James R Roche, Covering cubes by random half cubes, with applications to binary neural networks, Journal of Computer and System Sciences 56 (1998), no. 2, 223–252.
- [KWB22] Dmitriy Kunisky, Alexander S Wein, and Afonso S Bandeira, Notes on computational hardness of hypothesis testing: Predictions using the low-degree likelihood ratio, Mathematical Analysis, its Applications and Computation: ISAAC 2019, Aveiro, Portugal, July 29–August 2, Springer, 2022, pp. 1–50.
- [LM15] Shachar Lovett and Raghu Meka, Constructive discrepancy minimization by walking on the edges, SIAM Journal on Computing 44 (2015), no. 5, 1573–1582.

- [LRR17] Avi Levy, Harishchandra Ramadas, and Thomas Rothvoss, Deterministic discrepancy minimization via the multiplicative weight update method, International Conference on Integer Programming and Combinatorial Optimization, Springer, 2017, pp. 380–391.
- [Mat99] Jiri Matousek, Geometric discrepancy: An illustrated guide, vol. 18, Springer Science & Business Media, 1999.
- [MMZ05] Marc Mézard, Thierry Mora, and Riccardo Zecchina, Clustering of solutions in the random satisfiability problem, Physical Review Letters **94** (2005), no. 19, 197205.
- [NS23] Shuta Nakajima and Nike Sun, Sharp threshold sequence and universality for ising perceptron models, Proceedings of the 2023 Annual ACM-SIAM Symposium on Discrete Algorithms (SODA), SIAM, 2023, pp. 638–674.
- [Pot18] Aditya Potukuchi, *Discrepancy in random hypergraph models*, arXiv preprint arXiv:1811.01491 (2018).
- [PX21] Will Perkins and Changji Xu, Frozen 1-RSB structure of the symmetric Ising perceptron, Proceedings of the 53rd Annual ACM SIGACT Symposium on Theory of Computing, 2021, pp. 1579–1588.
- [Rot17] Thomas Rothvoss, Constructive discrepancy minimization for convex sets, SIAM Journal on Computing 46 (2017), no. 1, 224–234.
- [RSS18] Prasad Raghavendra, Tselil Schramm, and David Steurer, *High-dimensional esti*mation via sum-of-squares proofs, arXiv preprint arXiv:1807.11419 **6** (2018).
- [RV17] Mustazee Rahman and Balint Virag, Local algorithms for independent sets are half-optimal, The Annals of Probability 45 (2017), no. 3, 1543–1577.
- [Sid68] Zbynek Sidák, On multivariate normal probabilities of rectangles: their dependence on correlations, The Annals of Mathematical Statistics **39** (1968), no. 5, 1425–1434.
- [Spe85] Joel Spencer, Six standard deviations suffice, Transactions of the American mathematical society **289** (1985), no. 2, 679–706.
- [SS23] Ashwin Sah and Mehtaab Sawhney, Distribution of the threshold for the symmetric perceptron, arXiv preprint arXiv:2301.10701 (2023).
- [ST03] Mariya Shcherbina and Brunello Tirozzi, Rigorous solution of the Gardner problem, Communications in mathematical physics **234** (2003), no. 3, 383–422.
- [Sto13] Mihailo Stojnic, Another look at the Gardner problem, arXiv preprint arXiv:1306.3979 (2013).
- [Tal99] Michel Talagrand, *Intersecting random half cubes*, Random Structures & Algorithms **15** (1999), no. 3-4, 436–449.
- [Tal11] _____, Mean field models for spin glasses: Advanced replica-symmetry and low temperature, Springer, 2011.

- [TMR20] Paxton Turner, Raghu Meka, and Philippe Rigollet, Balancing Gaussian vectors in high dimension, Conference on Learning Theory, PMLR, 2020, pp. 3455–3486.
- [Wei20] Alexander S Wein, Optimal low-degree hardness of maximum independent set, arXiv preprint arXiv:2010.06563 (2020).
- [Wei22] _____, Average-case complexity of tensor decomposition for low-degree polynomials, arXiv preprint arXiv:2211.05274 (2022).
- [Wen62] James G Wendel, A problem in geometric probability, Mathematica Scandinavica 11 (1962), no. 1, 109–111.
- [Win61] Robert O Winder, Single stage threshold logic, 2nd Annual Symposium on Switching Circuit Theory and Logical Design (SWCT 1961), IEEE, 1961, pp. 321–332.
- [Xu19] Changji Xu, Sharp threshold for the Ising perceptron model, arXiv preprint arXiv:1905.05978 (2019).
- [Yak96] Benjamin Yakir, The differencing algorithm ldm for partitioning: a proof of a conjecture of karmarkar and karp, Mathematics of Operations Research 21 (1996), no. 1, 85–99.
- [ZK16] Lenka Zdeborová and Florent Krzakala, Statistical physics of inference: Thresholds and algorithms, Advances in Physics 65 (2016), no. 5, 453–552.