Generalized Turán Problem for Complete Hypergraphs

Levente Bodnár

February 20, 2023

Abstract

Write $K_n^{(k)}$ for the complete k-graph on n vertices. For $2 \le k \le g < r$ integers, let $\pi\left(n, K_g^{(k)}, K_r^{(k)}\right)$ be the maximum density of $K_g^{(k)}$ in n vertex $K_r^{(k)}$ -free k-graphs. The main contribution of this paper is the upper bound: $\pi\left(n, K_g^{(k)}, K_r^{(k)}\right) \leq \left(1+O\left(n^{-1}\right)\right) \prod_{m=k}^g$ $\left(1-\frac{\binom{m-1}{k-1}}{\binom{r-1}{k}}\right)$ $\binom{r-1}{k-1}$ $\big)$. The graph case $(k = 2)$ is the first known generalized Turán question, investigated by Erdős. The $k = g$ case is the hypergraph Turán problem where the best known general upper bound is by de Caen. The result proved here matches both bounds asymptotically, while any triple k, q, r with $2 < k < q < r$ provides a new upper bound. The proof uses techniques from the theory of flag algebras to derive linear relations between different densities. These relations can be combined with linear algebraic methods. Additionally a simple flag algebraic certificate will be given for $\lim_{n\to\infty} \pi\left(n, K_4^{(3)}, K_5^{(3)}\right) = 3/8.$

1 Introduction

1.1 Hypergraph Turán Problems

Given a k-graph H, write $\pi(n, H)$ for the maximum density of k-uniform edges among H-free hypergraphs with size n, and let $\pi(H) = \lim_{n \to \infty} \pi(n, H)$. It is known that the limit always exists. Let $K_n^{(k)}$ be the complete k-graph with n vertices. A landmark result by Turán determined the values $\pi\left(n, K_r^{(2)}\right)$ exactly, with the unique graphs attaining the maximum.

Theorem 1 ([\[Tur41\]](#page-19-0)). $\pi\left(n, K_r^{(2)}\right)$ is uniquely attained at the balanced complete $(r-1)$ -partite graph on n vertices.

Following this, Erdős and Stone found more generally the value $\pi(H)$ for all graph H .

Theorem 2 ([\[ES46\]](#page-18-0)). Suppose H is a graph with chromatic number $\chi(H)$, then

$$
\pi(H) = 1 - \frac{1}{\chi(H) - 1}.
$$

The corresponding question, when $k > 2$, is still open and seems to be much more difficult. There are sporadic results for various k-graphs, but no $\pi\left(K_r^{(k)}\right)$ value is known. The best general upper bound comes from de Caen.

Theorem 3 ([\[dC83\]](#page-18-1)).

$$
\pi\left(n, K_r^{(k)}\right) \le 1 - \left(1 + \frac{r - k}{n - r + 1}\right) \frac{1}{\binom{r - 1}{k - 1}}.
$$

For an extensive survey, focusing on the $\pi\left(n, K_r^{(k)}\right)$ problem, with various lower and upper bounds, see [\[Sid95\]](#page-19-1). More recent coverage of the question with different *k*-graphs can be found in [\[Kee11\]](#page-18-2).

1.2 Generalized Turán Problems

As a possible generalization of the Turán question, one can ask the maximum density of a given k-graph F , instead of the k-edges. For F , H given k-graphs, write $\pi(n, F, H)$ for the maximum density of F among n sized H-free k-graphs and use $\pi(F, H) = \lim_{n \to \infty} \pi(n, F, H)$. For complete graphs, this was initially investigated by Erdős [\[Erd62,](#page-18-3) [Erd84\]](#page-18-4).

Theorem 4 ([\[Erd62\]](#page-18-3)). For $2 \leq g < r$ integers $\pi\left(n, K_g^{(2)}, K_r^{(2)}\right)$ is uniquely attained at the balanced complete $(r-1)$ -partite graph on n vertices.

Note that this gives asymptotically that $\pi\left(K_g^{(2)}, K_r^{(2)}\right) = \prod_{m=2}^g \left(1 - \frac{m-1}{r-1}\right)$. The generalized Turán problem for graphs was systematically investigated by Alon and Shikhelman, obtaining a result similar to Erdős-Stone.

Theorem 5 ([\[AS16\]](#page-18-5)). For any graph H, with chromatic number $\chi(H)$, the following holds

$$
\pi\left(K_g^{(2)}, H\right) = \prod_{m=2}^g \left(1 - \frac{m-1}{\chi(H)-1}\right).
$$

In addition, $[AS16]$ investigates degenerate generalized Turán questions – the rate of convergence of $\pi(n, F, H)$ when $\pi(F, H) = 0$. [\[XZG21\]](#page-19-2) finds various bounds for several degenerate generalized hypergraph Turán problems.

The generalized Turán problem for complete k -graphs corresponds with the separation of different layers of the boolean hypercube using a k-CNF. This idea appears for example in [\[Sid87\]](#page-19-3) and will be further explored in a different paper. [\[FGT22\]](#page-18-6) gives new insights into the set of satisfying assignments of CNFs using a variant of the VC dimension. Bounds on this variant of the VC dimension turn out to be equivalent to a generalized Turán-type conjecture.

1.3 Flag Algebras

The theory of flag algebras [\[Raz07\]](#page-19-4) provides a systematic approach to studying extremal combinatorial problems and the tools available for solving them. It gives a common ground for combinatorial ideas, by expressing them as linear operators, acting between flag algebras. Linearity means the different techniques can be easily combined with linear programming/linear algebra.

A large part of the theory can be automated with state-of-the-art optimization algorithms, providing spectacular improvements in density bounds. There has been significant progress in the famous tetrahedron problem [\[dC83,](#page-18-1) [CL99\]](#page-18-7) with the previous best bound being $\pi\left(K_4^{(3)}\right) \leq \frac{3+\sqrt{7}}{12} < 0.59360$ while flag algebraic calculations improved it to the following bound:

Theorem 6 ([\[Raz10\]](#page-19-5) verified in [\[FRV11,](#page-18-8) [Bab11\]](#page-18-9)). $\pi \left(K_4^{(3)} \right) \leq 0.56167$.

Note that the best known lower bound is $5/9 \leq \pi \left(K_4^{(3)} \right)$. For excluded $K_5^{(3)}$ the calculations give the following:

Theorem 7 ([\[Bab11\]](#page-18-9)). $\pi\left(K_{5}^{(3)}\right) \leq 0.76954$

with best known lower bound $3/4 \leq \pi \left(K_5^{(3)} \right)$.

For a list of results provided by flag algebraic calculations, see [\[Raz10,](#page-19-5) [FRV11\]](#page-18-8). The power of flag algebra has been illustrated in a wide range of other combinatorial questions [\[LP21,](#page-18-10) [SS18\]](#page-19-6). Unfortunately, the computer-generated proofs lack insight and scale-ability compared to classical, hand-crafted arguments. They often only work in a small enough parameter range (for example, bounding $\pi(H)$ for H with at most 7 vertices). A survey by Razborov [\[Raz13\]](#page-19-7) calls such applications plain.

One of the goals of this paper is to show that the powerful plain flag algebra method, in this case, can be performed by hand, resulting in a general and scale-able theorem. The main ideas and proof steps, therefore, correspond with a plain application of flag algebra and were heavily inspired by it. During the proofs, relevant parts of the flag algebra theory will be highlighted. While the asymptotic result can be fully proved with flag algebraic manipulations, the bound with finite n is only attainable with a more precise bounding of the errors. The theory is not explained here, for a quick introduction see [\[SFS16\]](#page-19-8) or the original text [\[Raz07\]](#page-19-4).

1.4 Overview of the Result

In this paper, the generalized Turán problem for complete hypergraphs will be investigated, with the following contribution:

Theorem 8. For integers $1 < k \leq g < r$ and any $n > (r-1)\left(1 + \left(\frac{(r-k)}{k-1}\right)^{r-1}\right)$ $k-1$ $\Big)^2 \Big),$

$$
\pi\left(n, K_g^{(k)}, K_r^{(k)}\right) \le
$$

$$
\leq \left(1 + \frac{(r-1)(r-k)^2}{(k-1)^2n - (r-1)\left(2k^2 - 2k(r+1) + r^2 + 1\right)}\right) \prod_{m=k}^g \left(1 - \frac{\binom{m-1}{k-1}}{\binom{r-1}{k-1}}\right).
$$

Note this means asymptotically that

Corollary 9. For integers $1 < k \leq g < r$

$$
\pi\left(K_g^{(k)}, K_r^{(k)}\right) \le \prod_{m=k}^g \left(1 - \frac{\binom{m-1}{k-1}}{\binom{r-1}{k-1}}\right).
$$

The asymptotic bound is known to be tight when $k = 2$, with the matching, balanced $(r - 1)$ -partite construction. Additionally, it agrees with the best-known general hypergraph Turán bound by de Caen [\[dC83\]](#page-18-1) which is conjectured to not be tight.

[\[Sid95\]](#page-19-1) describes various lower bound constructions for the $2 < k = g$ case. A simple construction (when $k-1$ divides $r-1$) splits the vertex set into $\frac{r-1}{k-1}$ equal groups and includes each k set that is not fully contained in a group. Using $l = \frac{r-1}{k-1}$ and an inclusion-exclusion calculation, this gives the asymptotic bound

$$
\sum_{s=0}^{\lfloor g/k\rfloor}(-1)^s\binom{l}{s}\sum_{\substack{k\leq i_1,\ldots,i_s\\i_1+\cdots+i_s\leq g}}\binom{g}{i_1,\ldots,i_s,g-i_1-\cdots-i_s}l^{-i_1-\cdots-i_s}\leq \pi\left(K_g^{(k)},K_r^{(k)}\right).
$$

While it is known that this construction is not optimal when $k = g$, in the $k \ll g$ regime, where $K_g^{(k)}$ appears more if the edges are "grouped", it provides a stronger bound. Section [6](#page-15-0) shows that this is asymptotically the best construction for $\pi\left(K_4^{(3)}, K_5^{(3)}\right)$. In general $g = r - 1$ gives

$$
\sqrt{2\pi r} e^{\frac{\log(2\pi k)r}{2k}} \approx
$$

\n
$$
\approx \left(\frac{r-1}{k-1, k-1, \dots, k-1}\right) l^{-(r-1)} \le
$$

\n
$$
\leq \pi \left(K_{r-1}^{(k)}, K_r^{(k)}\right) \le
$$

\n
$$
\leq \prod_{m=k}^{r-1} \left(1 - \frac{\binom{m-1}{k-1}}{\binom{r-1}{k-1}}\right) \le
$$

\n
$$
\leq e^{(k-r)/k}.
$$

Given a hypergraph G , write $d(H, G)$ for the induced density of H in G. The main tool used in the proof of theorem [8](#page-2-0) is:

Lemma 10. For all $0 \leq x$ and integers $k \leq m < n$, if G is an n vertex k-graph then

$$
\geq \begin{pmatrix} -\frac{1-\frac{k-1}{m}}{x} \\ 2-\frac{k-1}{mx}-\frac{1}{(n-m)x} \end{pmatrix} \begin{pmatrix} d\left(K_{m+1}^{(k)},G\right) \\ d\left(K_{m}^{(k)},G\right) \\ d\left(K_{m-1}^{(k)},G\right) \end{pmatrix} + d\left(K_{m-1}^{(k)},G\right).
$$

Note that the densities $d\left(K_m^{(k)}, G\right)$ only appear linearly in the expression. For different k, g, r parameters, a convex combination of the expressions ap-pearing in lemma [10,](#page-4-0) with suitable x, m values substituted in yields theorem [8.](#page-2-0) As a comparison, [\[dC83\]](#page-18-1) utilizes similar ideas, but with a more complicated (non-linear) expression,

$$
f_{m+1} \ge \frac{m^2 f_m}{(m-k+1)(n-m)} \left(\frac{f_m(n-m+1)}{f_{m-1}m} - \frac{(k-1)(n-m)+m}{m^2} \right)
$$

where $f_m = d\left(K_m^{(k)}, G\right)$ for short.

 $\overline{0}$

1.5 Outline of the Paper

Section [2](#page-4-1) summarizes the important notations and conventions throughout the paper. The proof of theorem [8](#page-2-0) is included in section [3](#page-5-0) using two important components: lemma [10,](#page-4-0) which is proved in section [4;](#page-7-0) and a technical calculation (lemma [12\)](#page-6-0), that is included in section [5.](#page-11-0) The short section [6](#page-15-0) includes a certificate for $\pi\left(K_4^{(3)}, K_5^{(3)}\right) = 3/8$. The paper finishes with a few concluding remarks in section [7,](#page-17-0) the limitations of this approach and possible directions.

2 Notation and Conventions

2.1 Basic Notation

For a set V, the collection of subsets with size k is denoted by $\binom{V}{k}$. The hypergraphs are identified with their edge sets; $G \subseteq {V(G) \choose k}$ is a k-graph with $V(G)$ vertex set. $K_n^{(k)}$ is the complete k-graph with n vertices. For $S \subseteq V(G)$ the induced sub-hypergraph is $G \upharpoonright_S = G \cap {S \choose k}$. Hypergraph isomorphism is represented by $G \simeq H$. $\mathcal{H}_n^{(k)}$ is the collection of non-isomorphic k-graphs having n vertices.

Bold symbols indicate random variables. The uniform distribution from a set V is represented by Unif (V) . The density of H in G is defined to be $d(H, G) = \mathbb{P}[G \mid s \simeq H]$ where $S \sim$ Unif $\binom{V(G)}{|H|}$. Notice that this is the induced density, corresponding more with the flag algebraic approach, rather than the classical sub-hypergraph inclusion (referenced in the introduction). Write $d_s(H, G) = \mathbb{P}[G \mid s \simeq F, H \subseteq F]$ for the classical inclusion with the same $\mathbf{S} \sim \text{Unif } \binom{V(G)}{|H|}$. When H is a complete hypergraph, the two notions are equivalent. The generalized Turán problem is to determine the value

$$
\pi(n, F, H) = \max \left\{ d_s(F, G) : G \in \mathcal{H}_n, d_s(H, G) = 0 \right\}.
$$

The asymptotic problem asks $\pi(F, G) = \lim_{n \to \infty} \pi(n, F, G)$ (it is known that the limit always exists).

The quantity

$$
x_{m,r}^{(k)} = 1 - \frac{\binom{m-1}{k-1}}{\binom{r-1}{k-1}}
$$

will be important, these are the terms appearing in the product.

2.2 Flag Notation

The hypergraphs represent the corresponding flags with empty type. $T_n^{(k)}$ is the complete type with n vertices and all k -uniform edges. Type is indicated as a superscript. In particular $K_n^{(k),T_m^{(k)}}$ is the unique complete flag on n vertices with a type having m vertices. For a type T, T also represents the flag with type T and no extra vertices/edges. The averaging operator, transforming a T-typed flag F^T into a flag with empty type is $\llbracket F^T \rrbracket_T$.

2.3 Conventions

In most of the proofs, the symbol k is fixed, and the appearing statements concern k -graphs. For this reason, k superscripts from the notations are often dropped. Additionally, g, r, n symbols are reserved. They are integer parameters of the main question; determining the value of $\pi(n, K_q, K_r)$.

3 Proof of Main Theorem

In this short section theorem [8](#page-2-0) will be proved with the use of lemma [10](#page-4-0) and lemma [12,](#page-6-0) a technical result included in section [5.](#page-11-0) Let's recall the main theorem, with the introduced $x_{m,r}^{(k)}$ notation.

Theorem [8.](#page-2-0) Given integers $1 < k \leq g < r$ and $n > (r-1)\left(1 + \left(\frac{(r-k)}{k-1}\right)\right)$ $k-1$ $\Big)^2 \bigg),$

then

$$
\pi\left(n, K_g^{(k)}, K_r^{(k)}\right)
$$

$$
\leq \left(1 + \frac{(r-1)(r-k)^2}{(k-1)^2n - (r-1)\left(2k^2 - 2k(r+1) + r^2 + 1\right)}\right) \prod_{m=k}^g x_{m,r}^{(k)}.
$$

In the following, the superscript k is dropped from the notations for easier readability. The following claim gives bounds on the $x_{m,r}$ values. It follows easily after expanding the definition of $x_{m,r}$, therefore the proof is not included.

Claim 11. When $k - 1 \le m \le r$,

$$
0 \le x_{m,r}^{(k)} \le 1,
$$

with equality at $m = r$ and $m = k - 1$ respectively. The smallest nonzero value is $x_{r-1,r} = \frac{k-1}{r-k}$

Proof of theorem [8.](#page-2-0) Choose any $G \in \mathcal{H}_n$ (irrespective of the value of $d(K_r, G)$), with

$$
n > (r-1)\left(1 + \left(\frac{(r-k)}{k-1}\right)^2\right),
$$

and for short write $f_m = d(K_m, G)$. In the range $m \in \{k, k+1, ..., r-1\}, x_{m,r}$ is positive, therefore lemma [10,](#page-4-0) with $x = x_{m,r}$ holds.

$$
0 \ge -\frac{1 - \frac{k-1}{m}}{x_{m,r}} f_{m+1} + \left(2 - \frac{k-1}{mx_{m,r}} - \frac{1}{(n-m)x_{m,r}}\right) f_m - x_{m,r} f_{m-1}
$$

The value $(n - m)x_{m,r}$ is minimal at $m = r - 1$. Use E_m for the above expression but $\frac{1}{(n-m)x_{m,r}}$ replaced with $\frac{r-k}{(n-r+1)(k-1)}$.

$$
E_m = -\frac{1 - \frac{k-1}{m}}{x_{m,r}} f_{m+1} + \left(2 - \frac{k-1}{mx_{m,r}} - \frac{r-k}{(n-r+1)(k-1)}\right) f_m - x_{m,r} f_{m-1}
$$

The replacement decreases the value, giving that each E_m is still nonpositive. This gives that any $\delta_k, \delta_{k+1}, ..., \delta_{r-1}$ sequence with all $0 \leq \delta_m$ results in $0 \ge \sum_{m=k}^{r-1} \delta_m E_m$. For a lower bound it is enough to find coefficients $0 \le \delta_m$ satisfying

$$
0 \geq \sum_{m=k}^{r-1} \delta_m E_m = -\delta_k x_{k,r} f_{k-1} + f_g - \delta_{r-1} \frac{1 - \frac{k-1}{r-1}}{x_{r-1,r}} f_r.
$$
 (1)

With the assumption that $f_r = d(K_r, G) = 0$ and the simple observation that $f_{k-1} = d(K_{k-1}, G) = 1$, one can deduce from eq. [\(1\)](#page-6-1) that $\pi(n, K_g, K_r) \leq$ $\delta_k x_{k,r}$. Notice that finding δ_m corresponds with solving eq. [\(1\)](#page-6-1), a system of linear equations. The technical section [5](#page-11-0) includes a way to approximate this system of linear equations. The following lemma summarizes the result, concluding the proof of theorem [8.](#page-2-0)

Lemma 12. If $n > (r - 1) \left(1 + \left(\frac{(r-k)}{k-1} \right) \right)$ $k-1$ $\binom{2}{1}$ then the solution to the linear equations eq. [\(1\)](#page-6-1) satisfies that $\delta_m \geq 0$ and that

$$
\delta_k = \left(1 + \frac{(r-1)(r-k)^2}{(k-1)^2n - (r-1)\left(2k^2 - 2k(r+1) + r^2 + 1\right)}\right) \prod_{m=k+1}^g x_{m,r}.
$$

4 Linear Density Relations

This section covers the main combinatorial calculations involved in the proof of lemma [10.](#page-4-0) The main purpose of lemma [10](#page-4-0) is to act as a building block. Not only the validity is easier to verify, but it also involves the densities $d(K_m, G)$ linearly, therefore it can be combined easily; as illustrated in theorem [8.](#page-2-0) The small claims in this section follow closely core results from the flag algebra theory. The connection will be highlighted in remark [16.](#page-9-0) The connection between the plain flag algebra application and this proof is discussed in remark [17.](#page-10-0)

In the upcoming proofs, the k superscript will not be included. S is any subset of $V(G)$, while S_m is an m element subset of $V(G)$. Write $q(S)$ for the indicator function that is 1 when S induces a complete hypergraph in G and 0 otherwise. $l(S)$ is the number of $v \in V(G) \backslash S$ where $S+v$ is complete in G. The corresponding probability is $r(S_m) = \frac{l(S_m)}{n-m}$. Similarly $rr(S)$ is the probability that two different vertex extensions are both complete.

$$
rr(S_m) = \frac{\binom{l(S_m)}{2}}{\binom{n-m}{2}}
$$

When $H \in \mathcal{H}_n$, then $s(H)$ is used for the size of the intersection of nonedges in H. In particular $s(K_n) = n$ and $s(K_n^-) = k$ where K_n^- represents the hypergraph on n vertices that has exactly one edge missing.

In the proof of lemma [10,](#page-4-0) the fact, that $q(S)(r(S) - x)^2$ is always positive, will be exploited. Understanding the terms in the square is done by the following short claims. First, $r(S)^2$ and $rr(S)$ are related.

Claim 13.

$$
r(S_{m-1})^2 \le rr(S_{m-1}) + r(S_{m-1})\frac{1}{n-m}
$$

This simply follows from

$$
r(S_{m-1})^2 - rr(S_{m-1}) = r(S_{m-1}) \left(\frac{l(S_{m-1})}{n-m+1} - \frac{l(S_{m-1})-1}{n-m} \right) \le r(S_{m-1}) \frac{1}{n-m}.
$$

Second, linear equality between densities is shown.

Claim 14. Suppose $m \leq l \leq n$ with $F \in \mathcal{H}_m$ and $G \in \mathcal{H}_n$ then

$$
d(F, G) = \sum_{H \in \mathcal{H}_l} d(F, H) d(H, G),
$$

in particular

$$
d(K_m, G) = \sum_{H \in \mathcal{H}_{m+1}} \frac{s(H)}{m+1} d(H, G).
$$

Proof. Note that a uniform m sized subset of $V(G)$ can be sampled by first choosing $\mathbf{S}_l \sim$ Uniform $\binom{V(G)}{l}$ and then $\mathbf{S}_m \sim$ Uniform $\binom{\mathbf{S}_l}{m}$. The claim follows from the law of total probability. The events $\{G \mid_{\mathbb{S}_l} \simeq H : H \in \mathcal{H}_l\}$ partition the probability space, therefore

$$
d(F, G) = \mathbb{P}\Big[G\upharpoonright_{\mathbf{S}_m} \simeq F\Big]
$$

= $\sum_{H \in \mathcal{H}_l} \mathbb{P}\Big[G\upharpoonright_{\mathbf{S}_m} \simeq F \mid G\upharpoonright_{\mathbf{S}_l} \simeq H\Big]\mathbb{P}\Big[G\upharpoonright_{\mathbf{S}_l} \simeq H\Big]$
= $\sum_{H \in \mathcal{H}_l} d(F, H) d(H, G).$

The special case follows from $d(K_m, H) = \frac{s(H)}{m+1}$ when $H \in \mathcal{H}_{m+1}$.

 \Box

In the proof of lemma [10,](#page-4-0) S_{m-1} is chosen uniformly from the possible $m-1$ sized sets. The final claim connects the expected values arising in the terms of $q(S)(p(S_{m-1}) - x)^2$ with densities of various k-graphs in G.

Claim 15. Suppose \mathbf{S}_{m-1} is chosen uniformly randomly from the set $\binom{V(G)}{m-1}$ then

1.

$$
\mathbb{E}\left[q(\mathbf{S}_{m-1})r(\mathbf{S}_{m-1})\right] = d\left(K_m, G\right),
$$

2.

$$
\mathbb{E}\left[q(\mathbf{S}_{m-1})rr(\mathbf{S}_{m-1})\right] = \sum_{H \in \mathcal{H}_{m+1}} \frac{\binom{s(H)}{2}}{\binom{m+1}{2}} d(H, G).
$$

Proof. Similar to the proof of claim [14,](#page-7-1) first choosing $\mathbf{S}_l \sim$ Uniform $\binom{V(G)}{l}$ and then $\mathbf{S}_{m-1} \sim$ Uniform $\begin{pmatrix} \mathbf{S}_l \\ m-1 \end{pmatrix}$ results in uniformly distributed \mathbf{S}_{m-1} . Note that $r(\mathbf{S}_{m-1})$ and $rr(\mathbf{S}_{m-1})$ corresponds with choosing 1 and 2 additional vertices accordingly, and then checking a condition on the extended set.

In particular, for \mathbf{S}_m , \mathbf{S}_{m-1} pair, let $R(\mathbf{S}_m, \mathbf{S}_{m-1})$ be the event that $G\upharpoonright_{\mathbf{S}_m}\simeq$ K_m (and therefore G $\succeq K_{m-1}$). The claim follows from the law of total expectation, by conditioning on the shape of $G \, | \mathbf{s}_m$

$$
\mathbb{E}\left[q(\mathbf{S}_{m-1})r(\mathbf{S}_{m-1})\right]
$$

=
$$
\sum_{H \in \mathcal{H}_m} \mathbb{P}\left[R(\mathbf{S}_m, \mathbf{S}_{m-1}) \mid G\!\upharpoonright_{\mathbf{S}_m} = H\right] \mathbb{P}\left[G\!\upharpoonright_{\mathbf{S}_m} = H\right]
$$

= $d(K_m, G)$.

Since only $H = K_m$ contains a suitable $m - 1$ sized subset that satisfies $R_{\mathbf{S}_m,\mathbf{S}_{m-1}}.$

For the second part, write $RR(\mathbf{S}_{m+1}, \mathbf{S}_{m-1})$ for the event that there are two copies of K_m inside G $\upharpoonright_{\mathbf{S}_{m+1}}$ intersecting exactly at \mathbf{S}_{m-1} (again this implies G \mid $\mathbf{S}_{m-1} = K_{m-1}$. The calculation in this case gives

$$
\mathbb{E}\left[q(\mathbf{S}_{m-1})rr(\mathbf{S}_{m-1})\right]
$$
\n
$$
= \sum_{H \in \mathcal{H}_{m+1}} \mathbb{P}\left[RR(\mathbf{S}_{m+1}, \mathbf{S}_{m-1}) \mid G\!\upharpoonright_{\mathbf{S}_{m+1}} = H\right] \mathbb{P}\left[G\!\upharpoonright_{\mathbf{S}_{m+1}} = H\right]
$$
\n
$$
= \sum_{H \in \mathcal{H}_{m+1}} \frac{\binom{s(H)}{2}}{\binom{m+1}{2}} d\left(H, G\right)
$$

since in a given $G \, | \mathbf{s}_{m+1} \simeq H$, a randomly chosen \mathbf{S}_{m-1} satisfies $RR(\mathbf{S}_{m+1}, \mathbf{S}_{m-1})$ with probability $\frac{\binom{s(H)}{2}}{\binom{m+1}{2}}$ $\frac{\binom{2}{2}}{\binom{m+1}{2}}$.

Remark 16. The above claims all correspond to parts of the general flag algebra theory [\[Raz07\]](#page-19-4).

1. claim [14](#page-7-1) corresponds to the chain rule (Lemma 2.2). In the language of flags it gives

$$
K_m = \sum_{H \in \mathcal{H}_{m+1}} \frac{s(H)}{m+1} H.
$$

2. claim [13](#page-7-2) corresponds to products (Lemma 2.3). It is more or less equivalent with

$$
p\left(K_m^{T_{m-1}};G^{T_{m-1}}\right)^2 - p\left(K_m^{T_{m-1}},K_m^{T_{m-1}};G^{T_{m-1}}\right) = O(|G|^{-1}).
$$

3. claim [15](#page-8-0) corresponds to averaging (Theorem 2.5). It is a restatement of

$$
\left[\!\left[K^{T_{m-1}}_m\right]\!\right]_{T_{m-1}}=K_m
$$

and

$$
\[K_m^{T_{m-1}}\Big)^2\]_{T_{m-1}} = \sum_{H \in \mathcal{H}_{m+1}} \frac{\binom{s(H)}{2}}{\binom{m+1}{2}} H.
$$

With these claims, the main lemma follows easily.

Lemma [10.](#page-4-0) For all $x > 0$ and integers $k \leq m < n$, if $G \in \mathcal{H}_n$ then the following holds

$$
0 \geq \left(-\frac{1-\frac{k-1}{m}}{x}\right) d(K_{m+1}, G) +
$$

$$
\left(2 - \frac{k-1}{mx} - \frac{1}{(n-m)x}\right) d(K_m, G) +
$$

$$
(-x) d(K_{m-1}, G).
$$

Proof. When $x \in \mathbb{R}$ the quantity $q(S)(r(S)-x)^2$ is always non-negative. Therefore choosing $\mathbf{S}_{m-1} \sim \text{Uniform} \binom{V(G)}{m-1}$ the following is true

$$
0 \leq \mathbb{E}\left[q(\mathbf{S}_{m-1})(r(\mathbf{S}_{m-1})-x)^2\right].
$$

Expanding the terms and applying claim [13](#page-7-2) gives

$$
0 \leq \mathbb{E}\left[q(\mathbf{S}_{m-1})rr(\mathbf{S}_{m-1})+\left(\frac{1}{n-m}-2x\right)q(\mathbf{S}_{m-1})r(\mathbf{S}_{m-1})+x^2q(\mathbf{S}_{m-1})\right].
$$

The substitution from claim [15](#page-8-0) yields the following expression, without expected values:

$$
0 \leq \sum_{H \in \mathcal{H}_{m+1}} \frac{\binom{s(H)}{2}}{\binom{m+1}{2}} d(H, G) + \left(\frac{1}{n-m} - 2x \right) d(K_m, G) + x^2 d(K_m, G).
$$

Notice that $s(H)$ is maximal on K_{m+1} , otherwise it is at most k. This observation gives

$$
0 \leq \frac{k-1}{m} \sum_{\substack{H \in \mathcal{H}_{m+1} \\ H \neq K_{m+1}}} \frac{s(H)}{(m+1)} d(H, G)
$$

+ $d(K_{m+1}, G)$
+ $\left(\frac{1}{n-m} - 2x\right) d(K_m, G)$
+ $x^2 d(K_{m-1}, G)$.

Finally expanding $\frac{k-1}{m}$ d(K_m , G) using claim [14](#page-7-1) results in

$$
0 \leq \left(1 - \frac{k-1}{m}\right) \qquad d(K_{m+1}, G) + \left(\frac{k-1}{m} + \frac{1}{(n-m)} - 2x\right) \frac{d(K_m, G)}{d(K_{m-1}, G)} + \frac{d(K_{m-1}, G)}{d(K_{m-1}, G)}.
$$

Since $x \geq 0$, note that lemma [10](#page-4-0) is a $-1/x$ multiple of the above, and the proof is complete.

 \Box

Remark 17. This lemma can be easily stated as

$$
0 \le \left(1 - \frac{k-1}{m}\right)K_{m+1} + \left(\frac{k-1}{m} - 2x\right)K_m + x^2K_{m-1}
$$

in the language of flags. The proof uses the expansion of the simple square

$$
0 \le \left[\left(K_m^{T_{m-1}} - x T_{m-1} \right)^2 \right]_{T_{m-1}}
$$

In a plain application of flag algebra, the computer finds a conic combination of squares, similar to the above expression. Lemma [10](#page-4-0) provides squares in a form that is easy to handle later (they only involve a small number of $d(K_m, G)$) values). The following section shows that the target expression

$$
K_g \le \prod_{m=k}^g x_{m,r}^{(k)} + cK_r
$$

for some c constant, lies in the conic combination of the squares.

5 The Associated Tridiagonal Matrix

Definition 18. Given $k < r$ integers, the problem has an associated tridiagonal matrix $D_r^{(k)}$ with entries $d_{l,m}$ indexed by the range $k \leq l, m < r$

$$
d_{l,m} = \begin{cases} -x_{m,r}^{(k)} & \text{if } l = m-1\\ 2 - \frac{k-1}{mx_{m,r}^{(k)}} & \text{if } l = m\\ -\frac{1 - \frac{k-1}{m}}{x_{m,r}^{(k)}} & \text{if } l = m+1\\ 0 & \text{otherwise} \end{cases}
$$

Write $\epsilon = \frac{r-k}{(n-r+1)(k-1)}$ and notice that $D_r^{(k)} - \epsilon I$ has column values equal to the coefficients in E_m . With the new notation, recall lemma [12,](#page-6-0) which was used to finish the proof of theorem [8.](#page-2-0)

Lemma [12.](#page-6-0) Let $\Delta_r^{(k)}(\epsilon) = \left(D_r^{(k)} - \epsilon I\right)^{-1}$ be the inverse of the associated tridiagonal matrix, with entries $\delta_{m,g}(\epsilon)$. If $0 \leq \epsilon < \frac{k-1}{(r-1)(r-k)}$, then the values $\delta_{m,q}(\epsilon)$ are all positive, and

$$
\delta_{k,g}(\epsilon) \le \frac{1}{1 - \epsilon \frac{(r-1)(r-k)}{k-1}} \prod_{m=k}^{g} x_{m+1,r}
$$

Notice that $\epsilon = 0$ corresponds with the inverse of $D_r^{(k)}$, and the asymptotic question when $n \to \infty$. This section is devoted to the proof of lemma [12,](#page-6-0) but it is illuminating and helpful for the proof to first calculate the inverse of $D_r^{(k)}$.

5.1 The inverse of $D_r^{(k)}$

Lemma 19. Let $\Delta_r^{(k)} = \left(D_r^{(k)}\right)^{-1}$ be the inverse of the associated tridiagonal matrix with entries $\delta_{m,g}$. Then $\delta_{m,g}$ are all positive and

$$
\delta_{k,g} = \prod_{m=k+1}^{g} x_{m,r}^{(k)}.
$$

In the following proofs, the k superscripts are omitted to increase readability. The complete inverse can be calculated following the method described in [\[Usm94\]](#page-19-9). The value θ_m represents the determinant of the rows and columns indexed by the set $\{k, k+1, ..., m\}$, while ϕ_m is the determinant for the rows and columns indexed by $\{m, m+1, ..., r-1\}.$

From the cofactor calculation of determinants and inverses, the following claim is easy to verify.

Claim 20.

1. For all $k \leq m < r$ the induction

$$
\theta_m = d_{m,m}\theta_{m-1} - d_{m-1,m}d_{m,m-1}\theta_{m-2}
$$

holds with initial values $\theta_{k-1} = 1$ and $\theta_{k-2} = 0$

2. For all $k \leq m < r$ the reverse induction

$$
\phi_m = d_{m,m}\phi_{m+1} - d_{m+1,m}d_{m,m+1}\phi_{m+2}
$$

holds with initial values $\phi_r = 1$ and $\phi_{r+1} = 0$

3. For all $k \leq m < r$ the determinant can be calculated

$$
Det (Dr) = \thetar-1 = \phik = \thetam \phim+1 - dm,m+1 dm+1,m \thetam-1 \phim+2
$$

4. The entries of the inverse matrix are

$$
\delta_{m,g} = \frac{(-1)^{m+g+r-k}}{\text{Det}(D_r)} \begin{cases} \theta_{m-1}\phi_{g+1} \prod_{i=m}^{g-1} d_{i,i+1} & \text{if } m \le g \\ \theta_{g-1}\phi_{m+1} \prod_{i=g}^{m-1} d_{i+1,i} & \text{otherwise} \end{cases}
$$

In the corresponding $k \leq m, g < r$ range.

First the ϕ_m values will be calculated using the recursive expression above. Claim 21. $\phi_m = 1$ in the range $k \leq m \leq r$

Proof. By reverse induction. The claim holds for $m = r$ and notice

$$
\phi_{r-1} = d_{r-1,r-1} = 2 - \frac{\frac{k-1}{r-1}}{1 - \frac{\binom{r-2}{k-1}}{\binom{r-1}{k-1}}} = 1
$$

Using point 2 from claim [20](#page-12-0)

$$
\phi_m = d_{m,m} - d_{m+1,m}d_{m,m+1}
$$

=2 - $\frac{k-1}{mx_{m,r}} - \left(1 - \frac{k-1}{m}\right) \frac{x_{m+1,r}}{x_{m,r}}$
=2 - $\frac{k-1}{m(1-u)} - \left(1 - \frac{k-1}{m}\right) \frac{1 - \frac{mu}{m-k+1}}{1-u}$
=1

Where $u = \frac{\binom{m-1}{k-1}}{\binom{r-1}{k}}$ $\frac{\binom{m-1}{k-1}}{\binom{r-1}{k-1}} = 1 - x_{m,r}$ and $u \frac{m}{m-k+1} = \frac{\binom{m}{k-1}}{\binom{r-1}{k-1}}$ $\frac{\binom{k-1}{k-1}}{\binom{r-1}{k-1}} = 1 - x_{m+1,r}$ substitution \Box was used to simplify the calculation.

This gives that Det $(D_r) = \theta_{r-1} = \phi_k = 1$.

Claim 22. In the $k - 1 \le m < r$ range, $Sign(\theta_m) = 1$

Proof. By induction, note that the claim holds for $\theta_{k-1} = 1$. Then using point 3 from claim [20](#page-12-0) and the value for the determinant,

$$
\theta_m = 1 + d_{m,m+1}d_{m+1,m}\theta_{m-1}
$$

Using claim [11,](#page-6-2) note that the values $d_{m,m+1} = -x_{m+1,r}$ and $d_{m+1,m} = -\frac{1-\frac{k-1}{m}}{x_m}$ are both negative. Therefore their product; and by induction, θ_{m-1} , are positive.

Proof of lemma [19.](#page-12-1) It claims two things,

1. The entries $\delta_{m,g}$ are all positive: claim [20](#page-12-0) point 4 gives that

$$
Sign(\delta_{m,g}) = (-1)^{m+g+r-k} \begin{cases} \prod_{i=m}^{g-1} \text{Sign}(d_{i,i+1}) & \text{if } m \le g\\ \prod_{i=g}^{m-1} \text{Sign}(d_{i+1,i}) & \text{otherwise.} \end{cases}
$$

using $\text{Sign}(\theta_m) = 1$ from claim [22](#page-13-0) and that $\phi_m = 1$ from claim [21.](#page-12-2) Since all $d_{i,i+1}, d_{i+1,i}$ are negative, the inverse of D_r has only positive entries.

2. $\delta_{k,g} = \prod_{m=k}^{g} x_{m+1,r}$: Again substituting the values $d_{i,i+1} = -x_{i+1,r}$ and $Det(D_r) = \phi_{g+1} = \theta_{k-1} = 1$ into claim [20](#page-12-0) point 4 gives the stated value for $\delta_{k,q}$.

Interestingly, the values $\delta_{k,g}$ are easy enough to calculate exactly. In contrast, $\delta_{m,q}$ requires the value of some θ_m which is difficult to find in general with the recursive expression. The sign of θ_m is easy to find, exactly what is needed for the proof.

 \Box

5.2 The inverse of $D_r^{(k)} - \epsilon I$

Consider the same calculation but with $D_r - \epsilon I$. The value $\phi_m(\epsilon)$ is the determinant for the rows and columns indexed by $\{m, m+1, ..., r-1\}$ of $D_r - \epsilon I$. The next claim shows that $\phi_m(\epsilon)$ is increasing in m when $\epsilon > 0$. A notation for the increments will be useful, write $\zeta_m(\epsilon) = \phi_{m+1}(\epsilon) - \phi_m(\epsilon)$.

Claim 23. When $k \leq m < r$ and $0 \leq \epsilon \leq \frac{k-1}{(r-1)(r-m)}$,

$$
0 \le \zeta_m(\epsilon) \le \epsilon \frac{r-1}{k-1} \left(1 - \left(1 - \frac{k-1}{r-1} \right)^{r-m} \right)
$$

and correspondingly

$$
1 \ge \phi_m(\epsilon) \ge 1 - \epsilon \frac{(r-1)(r-m)}{k-1}
$$

Proof. Use point 2 from claim [20.](#page-12-0) The initial value is $\zeta_r(\epsilon) = 0$ and by reverse induction take

$$
\phi_m(\epsilon) = (d_{m,m} - \epsilon) \phi_{m+1}(\epsilon) - d_{m+1,m}d_{m+1,m}\phi_{m+2}(\epsilon)
$$

= $\left(2 - \frac{k-1}{mx_{m,r}} - \epsilon\right) \phi_{m+1}(\epsilon) - \left(1 - \frac{k-1}{mx_{m,r}}\right) (\phi_{m+1}(\epsilon) + \zeta_{m+1}(\epsilon))$
= $\phi_{m+1}(\epsilon) - \zeta_{m+1}(\epsilon) \left(1 - \frac{k-1}{mx_{m,r}}\right) - y\phi_{m+1}(\epsilon).$

Therefore

$$
\zeta_m(\epsilon) = \zeta_{m+1}(\epsilon) \left(1 - \frac{k-1}{m x_{m,r}} \right) + \epsilon \phi_{m+1}(\epsilon). \tag{2}
$$

Note that

$$
0 \le d_{m,m+1}d_{m+1,m} = \left(1 - \frac{k-1}{mx_{m,r}}\right) \le \left(1 - \frac{k-1}{r-1}\right)
$$

in the $k \leq m < r$ range. As $\epsilon \leq \frac{k-1}{(r-1)(r-m)} < \frac{k-1}{(r-1)(r-m-1)}$ by reverse induction it holds that $0 \leq \phi_{m+1}(\epsilon)$ and $0 \leq \zeta_{m+1}(\epsilon)$ giving the required lower bound $0 \le \zeta_m(\epsilon)$. This implies the upper bound $\phi_m(\epsilon) \le 1$.

For the $\zeta_m(\epsilon)$ upper bound, in eq. [\(2\)](#page-14-0) bound each term: $mx_{m,r} \le (r-1)$ and $\phi_{m+1}(\epsilon) \leq 1$. This gives the intermediate result

$$
\zeta_m(\epsilon) \leq \zeta_{m+1}(\epsilon) \left(1 - \frac{k-1}{r-1}\right) + \epsilon.
$$

which, by iterated application and $\zeta_r(\epsilon) = 0$ initial value, implies

$$
\zeta_m(\epsilon) \leq \epsilon \frac{r-1}{k-1} \left(1 - \left(1 - \frac{k-1}{r-1} \right)^{r-m} \right).
$$

A summation formula for the upper and lower $\zeta_m(\epsilon)$ bounds combined with the initial $\phi_r(\epsilon) = 1$ value gives

$$
1 \ge \phi_m(\epsilon) \ge 1 - \epsilon \frac{(r-1)^2}{(k-1)^2} \left(\frac{(k-1)(r-m)}{r-1} + \left(1 - \frac{k-1}{r-1} \right)^{r-m} - 1 \right).
$$

This provides a tighter bound but for simplicity use

$$
1 - \epsilon \frac{(r-1)^2}{(k-1)^2} \left(\frac{(k-1)(r-m)}{r-1} + \left(1 - \frac{k-1}{r-1} \right)^{r-m} - 1 \right) \ge 1 - \epsilon \frac{(r-1)(r-m)}{k-1}.
$$

This gives a simple linear bound for the determinant. When $0 < \epsilon <$ $\frac{k-1}{(r-1)(r-k)},$

$$
1 - \epsilon \frac{(r-1)(r-k)}{k-1} \le \phi_k(\epsilon) = \text{Det}(D_r - \epsilon I) \le 1.
$$

If $0 \leq \epsilon < \frac{k-1}{(r-1)(r-k)}$ then the determinant is strictly positive, bounding the smallest eigenvalue of D_r .

Proof of lemma [12.](#page-6-0) Again it claims two things.

1. The entries $\delta_{m,q}(\epsilon)$ are all positive: By assumption, ϵ is smaller than the smallest eigenvalue of D_r . Therefore the expansion

$$
(D_r - \epsilon I)^{-1} = D_r^{-1} + \epsilon D_r^{-2} + \epsilon^2 D_r^{-3} ...
$$

holds. Lemma [19](#page-12-1) shows that the entries in D_r^{-1} (and in D_r^{-i} for $1 > i$ correspondingly) are all positive, giving the required positivity of $\delta_{m,g}(\epsilon)$.

2.

$$
\delta_{k,g}(\epsilon) \leq \frac{1}{1 - \epsilon \frac{(r-1)(r-k)}{k-1}} \prod_{m=k}^{g} x_{m+1,r}.
$$

This follows from substituting the bounds $\phi_m(\epsilon) \leq 1$ and $1 - \epsilon \frac{(r-1)(r-k)}{k-1} \leq$ Det($D_r - \epsilon I$) into claim [20](#page-12-0) point 4.

 \Box

$$
6 \quad \pi\left(K_4^{(3)}, K_5^{(3)}\right) = 3/8
$$

The combination of more sophisticated, but still simple squares can provide tight bounds for $\pi\left(K_4^{(3)}, K_5^{(3)}\right) = 3/8$. For an easier description of the flags, consider the complement question. If E_n is the 3-graph with n vertices and no edges then

$$
\pi\left(K_4^{(3)}, K_5^{(3)}\right) = \lim_{n \to \infty} \max\left\{d(E_4, G) : G \in \mathcal{H}_n^{(3)}, \ d(E_5) = 0\right\}.
$$

Use P_n for the corresponding type with n vertices and no edges. Note $E_n^{P_m}$ is unique for any $m < n$ pair. Define the further flags:

- 1. $L_a^{P_2}$ is the flag with vertex set $\{0, 1, 2, 3\}$, edge set $\{(0, 2, 3)\}$ and type formed from the vertices 0, 1. Similarly, write $L_b^{P_2}$ for the flag with the same vertex set and type but $\{(1, 2, 3)\}$ edge set.
- 2. Use $M_a^{P_3}$ for the flag with vertices $\{0, 1, 2, 3\}$, edges $\{(1, 2, 3)\}$ and type from 0, 1, 2. Symmetrically, with the same vertex and type set use $M_b^{P_3}$ for the edge set $\{(0,1,3)\}\.$
- 3. N^{Q_4} is the flag with vertices $\{0, 1, 2, 3, 4\}$, edges $\{(0, 1, 2)\}$ and type formed by 0, 1, 2, 3. Note that Q_4 does not agree with any of the T_n or P_n types.
- 4. $O_a^{T_4}$ has vertex set $\{0, 1, 2, 3, 4\}$, edge set $\{(0, 1, 4)\}$ and type formed from $0, 1, 2, 3$. Additionally write $O_b^{T_4}$ for the flag with the same vertex and type set but $\{(2,3,4)\}\)$ edges.

Then the following inequality holds on E_5 -free hypergraphs:

$$
0 \leq \frac{2}{3} \left[\left(E_3^{P_1} - \frac{3}{4} P_1 \right)^2 \right]_{P_1} + \frac{1}{6} \left[\left(L_a^{P_2} - L_b^{P_2} \right)^2 \right]_{P_2} +
$$

$$
\frac{13}{12} \left[\left(M_a^{P_3} + M_b^{P_3} + M_c^{P_3} - \frac{1}{2} P_3 \right)^2 \right]_{P_3} + \frac{11}{12} \left[\left(E_4^{P_3} - \frac{1}{2} P_3 \right)^2 \right]_{P_3} + (3)
$$

$$
2 \left[\left(N^{Q_4} - \frac{1}{2} Q_4 \right)^2 \right]_{Q_4} + \frac{1}{2} \left[\left(O_a^{T_4} - O_b^{T_4} \right)^2 \right]_{P_4} \leq \frac{3}{8} - E_4.
$$

So far the only verification of eq. [\(3\)](#page-16-0) requires a tedious (computer assisted) checking of all the 2102 hypergraphs in $\mathcal{H}_6^{(3)}$ without E_5 . This can be found in the supplement. The corresponding lower bound is attained at $G_n = K_{\lfloor n \rfloor}^{(3)}$ $\frac{1}{\lfloor n/2 \rfloor} \bigsqcup K_{\lceil n/3 \rceil}^{(3)}$ $\frac{1}{\lceil n/2 \rceil}$. Note that $d(E_5, G_n) = 0$ while $\lim_{n \to \infty} d(E_4, G_n) = \frac{3}{8}$.

7 Concluding Remarks

This paper investigated a natural extension of the generalized Turán problem to hypergraphs. The result matches the best-known general bounds for k-graphs but fails to provide tight bounds when $k > 3$.

The main combinatorial insight comes from the simple inequality

$$
0 \le \left[\left(K_m^{T_{m-1}} - x T_{m-1} \right)^2 \right]_{T_{m-1}}, \tag{4}
$$

combined with a close approximation of $\left\| \left(K_{m}^{T_{m-1}} \right)^{2} \right\|$ T_{m-1} . As shown in the paper, the convex combination of these squares includes difficult results for the generalized hypergraph Turán problem.

The long list of questions improved by the plain flag algebraic method indicates that finding more sophisticated squares can greatly improve the available density bounds. It would be interesting to identify other families of simple linear density relations (like the one described in lemma [10\)](#page-4-0) whose conic combination includes new bounds for extremal hypergraph problems, even better if the bounds are tight. The provided certificate for $\pi\left(K_4^{(3)}, K_5^{(3)}\right) = 3/8$ can perhaps be generalized to larger cases. It is interesting that for the smallest k, g, r tuple, which is not already known ($k = 2$) and is not a classical hypergraph Turán problem $(k = q)$, the exact solution follows from flag algebraic calculations. It also highlights the limitations of computer assisted searches: calculations for problems with higher parameters are infeasible.

7.1 Finding Squares

There is an easy to describe reason why eq. [\(4\)](#page-17-1) fails to provide tight bounds for k-graphs where $k > 2$ but is asymptotically exact when $k = 2$. The extremal configuration for $\pi\left(n, K_g^{(2)}, K_r^{(2)}\right)$ is a unique balanced $(r-1)$ -partite graph, call it $G_r(n)$, and say $G_r^{T_{m-1}}(n)$ is the same structure with a complete $(m-1)$ -tuple marked as a type. Any choice of T_{m-1} results in the same $\lim_{n\to\infty} d\left(K_m^{T_{m-1}}, G_r^{T_{m-1}}(n)\right)$ value, which is $x_{m,r}^{(2)} = 1 - \frac{m-1}{r-1}$. In contrast, the conjectured optimal constructions when $k > 2$ give different values for different T_{m-1} choices, therefore no $x \in \mathbb{R}$ exists with

$$
\[K_m^{T_{m-1}} - xT_{m-1}\big)^2\]_{T_{m-1}} = 0
$$

on the conjectured optimal constructions. This slackness gives the difference between the conjectured optimal constructions and the proved bounds here. The values $x_{m,r}^{(k)}$ are chosen optimally, any asymptotically significant improvement must utilize a different combinatorial insight.

References

- [AS16] Noga Alon and Clara Shikhelman. Many T copies in H-free graphs. Journal of Combinatorial Theory, Series B, 121:146–172, 2016. Fifty years of The Journal of Combinatorial Theory.
- [Bab11] Rahil Baber. Some Results in Extremal Combinatorics. PhD thesis, University College London, 2011.
- [CL99] Fan Chung and Linyuan Lu. An upper bound for the turán number T3(n, 4). J. Comb. Theory Ser. A, 87(2):381–389, aug 1999.
- [dC83] Dominique de Caen. Extension of a theorem of Moon and Moser on complete subgraphs, 1983.
- [Erd62] Paul Erdős. On the number of complete subgraphs contained in certain graphs. Magyar Tud. Akad. Mat. Kut. Int. Kozl, 7:459–474, 1962.
- [Erd84] Paul Erdős. On some problems in graph theory, combinatorial analysis and combinatorial number theory. Graph theory and combinatorics (Cambridge, 1983), pages 1–17, 1984.
- [ES46] Paul Erdős and Arthur H. Stone. On the structure of linear graphs. Bulletin of the American Mathematical Society, $52(12):1087 - 1091$, 1946.
- [FGT22] Peter Frankl, Svyatoslav Gryaznov, and Navid Talebanfard. A Variant of the VC-Dimension with Applications to Depth-3 Circuits. In ITCS 2022, volume 215 of LIPIcs, pages 72:1–72:19, Dagstuhl, Germany, 2022. Schloss Dagstuhl.
- [FRV11] Victor Falgas-Ravry and Emil R. Vaughan. On applications of Razborov's flag algebra calculus to extremal 3-graph theory. $arXiv$, 2011.
- [Kee11] Peter Keevash. Hypergraph Turán problems, page 83-140. London Mathematical Society Lecture Note Series. Cambridge University Press, 2011.
- [LP21] Bernard Lidick´y and Florian Pfender. Semidefinite programming and Ramsey numbers. SIAM Journal on Discrete Mathematics, 35(4):2328–2344, 2021.
- [Raz07] Alexander A. Razborov. Flag Algebras. The Journal of Symbolic Logic, 72(4):1239–1282, 2007.
- [Raz10] Alexander A. Razborov. On 3-Hypergraphs with Forbidden 4-Vertex Configurations. SIAM J. Discret. Math., 24(3):946–963, August 2010.
- [Raz13] Alexander A. Razborov. Flag Algebras: An Interim Report. In The Mathematics of Paul Erdős II, 2013.
- [SFS16] Marcel K. de Carli Silva, Fernando Mário de Oliveira Filho, and Cristiane Maria Sato. Flag Algebras: A First Glance, 2016.
- [Sid87] A. F. Sidorenko. Precise values of turan numbers. Mathematical Notes of the Academy of Sciences of the USSR, 42(5):913–918, November 1987.
- [Sid95] Alexander Sidorenko. What we know and what we do not know about Turán numbers. Graphs and Combinatorics, 11:179-199, 06 1995.
- [SS18] Jakub Sliacan and Walter Stromquist. Improving bounds on packing densities of 4-point permutations. Discrete Mathematics & Theoretical Computer Science, Vol. 19 no. 2, Permutation Patterns 2016, February 2018.
- [Tur41] Pál Turán. Eine extremalaufgabe aus der graphentheorie. Mat. és Fiz. Lapok, 48:436–452, 1941.
- [Usm94] Riaz A. Usmani. Inversion of a tridiagonal jacobi matrix. Linear Algebra and its Applications, 212-213:413–414, 1994.
- [XZG21] Zixiang Xu, Tao Zhang, and Gennian Ge. Some extremal results on hypergraph Turán problems. Science China Mathematics, 65:1765– 1774, 2021.