

Mixed Semi-Supervised Generalized-Linear-Regression with applications to Deep-Learning and Interpolators

Oren Yuval*

Department of Statistics, University of Tel-Aviv

and

Saharon Rosset

Department of Statistics, University of Tel-Aviv

May 29, 2024

Abstract

We present a methodology for using unlabeled data to design semi supervised learning (SSL) methods that improve the prediction performance of supervised learning for regression tasks. The main idea is to design different mechanisms for integrating the unlabeled data, and include in each of them a *mixing* parameter α , controlling the weight given to the unlabeled data. Focusing on Generalized Linear Models (GLM) and linear interpolators classes of models, we analyze the characteristics of different mixing mechanisms, and prove that in all cases, it is invariably beneficial to integrate the unlabeled data with some nonzero mixing ratio $\alpha > 0$, in terms of predictive performance. Moreover, we provide a rigorous framework to estimate the best mixing ratio α^* where mixed SSL delivers the best predictive performance, while using the labeled and unlabeled data on hand. The effectiveness of our methodology in delivering substantial improvement compared to the standard supervised models, in a variety of settings, is demonstrated empirically through extensive simulation, in a manner that supports the theoretical analysis. We also demonstrate the applicability of our methodology (with some intuitive modifications) to improve more complex models, such as deep neural networks, in real-world regression tasks.

Keywords: Predictive modeling; Semi-supervised regression; Generalized linear model, Deep learning

*The authors gratefully acknowledge *the Israeli Science Foundation grant 2180/20* and support from *the Israel Council for Higher Education Data-Science Centers*

1 Introduction

In numerous applications in the domain of machine learning, acquiring labeled data to train a supervised model is often more challenging than obtaining unlabeled data, which is generally linked to unsupervised learning. Given a scenario where there is an abundance of unlabeled data but a limited amount of labeled data, there is a compelling reason to leverage the unlabeled data to enhance the predictive accuracy of a supervised learning model through the implementation of a suitable semi-supervised learning (SSL) strategy. Over the years, some SSL methods and their effectiveness in improving predictive accuracy have been described by Zhou and Belkin [2014], by Singh et al. [2009] and Zhu [2005], primarily for classification tasks.

More recent works have shown promising results in improving deep neural network (DNN) by applying SSL methods, especially in the field of image classification [Sohn et al., 2020, Sun and Ge, 2020, Han et al., 2020]. These methods often rely on the clustering assumptions (i.e., low data density at decision boundaries, as discussed in Chapelle and Zien [2005]), which are generally not applicable to regression problems. Some works in the literature [Wasserman and Lafferty, 2007, Brouard et al., 2011, Niyogi, 2013, Xu et al., 2022] have explored the efficacy of utilizing unlabeled data to improve metric-based regression models, such as kernel methods. These approaches take advantage of semi-supervised smoothness and manifold assumptions (as discussed in Wasserman and Lafferty [2007]), which are less relevant to the parametric models that are the main focus of this paper.

Compared to classification and kernel methods for regression, limited attention has been paid to semi-supervised parametric regression models. In this context, one line of work [Tarpey et al., 2014, Chakraborty and Cai, 2018, Zhang et al., 2019, Azriel et al., 2022] has studied the usefulness of unlabeled data to improve ordinary least squares linear regression (OLS). In keeping with statistical tradition, these works are primarily focused on inference regarding the linear coefficients, rather than prediction performance. Another related line of work [Javanmard and Montanari, 2014, 2018, Bellec and Zhang, 2022, Bellec et al., 2018], discusses the usefulness of unlabeled data in Lasso-regularized sparse linear regression. Taking a broad look at both of these classes of related works, their results suggest that the use of unlabeled data improves the linear model’s robustness to miss-

specification and noisy data, but predictive performance improvement is not guaranteed in general.

Following the recent success in semi-supervised deep learning (SSDL) methods for classification, several recent works propose methods for applying SSDL for regression tasks [Jean et al., 2018, Rezagholiradeh and Haidar, 2018, Olmschenk et al., 2019, He et al., 2022]. A major aspect of these methods is adding a term to the loss function, that is a function of the unlabeled data, naturally named "unsupervised loss". In particular, Jean et al. [2018] suggested using the predictive variance over the unlabeled data as a regularization term that should be added to the objective function, during the training phase, multiplied by a properly selected regularization parameter. On the one hand, the intuition behind such a methodology is well explained and justified as an instance of *regularized Bayesian inference* (formalized earlier by Zhu et al. [2014]). On the other hand, theoretical analysis regarding the performance of such models is hard to obtain and there is no clear statement about the superiority of the semi-supervised model over the supervised equivalent. Instead, the authors demonstrate the empirical benefits of their suggested SSDL methods by systematic experiments on various examples of real-world data sets.

The methodology and theory presented in this paper improve and expand our previous work (Yuval and Rosset [2022]), which proposed a general methodology for semi-supervised empirical risk minimization, by adding a semi-supervised component to standard loss functions, and analyzed its effectiveness in the common class of generalized linear models (GLM). A major contribution of that work was a characterization of the relative performance of the supervised and semi-supervised estimators, and a consequent methodology for choosing between them, according to appropriate data-based estimate of the noise and signal of the model. That work provided preliminary evidence that SSL can be effective in the OLS model as well as in a one-layer neural network (NN) with ELU activation function, even when the linear model is well specified and the data is not very noisy.

In this paper, we propose and study a more sophisticated SSL methodology that examines the value of mixing between the semi-supervised and supervised estimators, rather than selecting between them. Our theoretical analysis of such mixed estimators provides a clear statement regarding the superiority of these estimators over the supervised ones and over the selection approach of Yuval and Rosset [2022]. Moreover, we provide a frame-

work for selecting the proper mixing ratio for the GLM and linear interpolator classes of models and demonstrate it empirically through extensive simulations. We also demonstrate the applicability of our methodology (with some intuitive modifications) to improve state-of-the-art deep-supervised models in a real-world regression task of facial landmark localization.

A major contribution of this work is in unifying the treatment of SSL in classical parametric models, where theoretical analysis is possible, and in modern complex models such as DNN, where mostly heuristic arguments and empirical analysis are possible. Specifically, our theoretical insights about the effectiveness of mixed estimators in OLS and GLM provide a motivation for existing techniques for SSDL in regression tasks, and lead to suggested enhancement and modifications which prove effective.

1.1 Notations, assumptions, and main idea

In the supervised framework, a training sample is given by $T = (X, Y)$, where $X \in \mathbb{R}^{n \times p}$ with rows x_1, \dots, x_n , and $Y = (y_1, \dots, y_n)^T \in \mathbb{R}^n$ such that $(x_1, y_1), \dots, (x_n, y_n)$ are i.i.d. according to some joint distribution P_{xy} . In the semi-supervised setting, the training sample is given by $T = (X, Y, Z)$, where the additional term $Z \in \mathbb{R}^{m \times p}$ represents the set of unlabeled data with z_1, \dots, z_m i.i.d. observations drawn from the marginal distribution P_x . In both cases, we are interested in defining a statistical learning process that fits a predictor $\hat{f}_T : \mathbb{R}^p \rightarrow \mathbb{R}$ that maps from a covariate vector $x \in \mathbb{R}^p$ to a predicted response $\hat{y} \in \mathbb{R}$, based on a training data set T . Throughout this paper, we evaluate the prediction errors of models trained using our proposed methods in the semi-supervised setting against those trained in the supervised framework. In the first part of this paper, we focus on the cases where $p < n$, and later in Section 5, we analyze interpolating models where $p > n$. For simplicity, we assume that the distribution P_x is centered around zero, that is, $\mathbb{E}[x] = \mathbf{0}_p$, and also that the distribution $P_{y|x}$ satisfies a constant conditional variance, meaning that: $y|x = f(x) + \epsilon = \mathbb{E}[y|x] + \epsilon$, where $\text{Var}(\epsilon) = \text{Var}(y|x) = \sigma^2$, and σ^2 is referred as the "noise" of the underlying model of the data.

Consider an independent sample (x_0, y_0) from the distribution P_{xy} , and a linear predictor β_T that is learned by the training set T , such that $\hat{f}_T(x_0) = x_0^T \beta_T$. Within the framework

of GLM, the goal is to reduce the out-of-sample *Canonical* prediction error, expressed as:

$$R(\beta_T) = \mathbb{E}_{T, x_0, y_0} [G(x_0^T \beta_T) - x_0^T \beta_T y_0],$$

where $G' = g$, with g being a known, monotonically increasing *link function*. This predictive accuracy criterion usually assumes that $\mathbb{E}[y|x] = g(x^T \beta)$ for some $\beta \in \mathbb{R}^p$. Assuming exchangeability between derivative and expectation, a premise accepted in this paper, it is straightforward to demonstrate that $R(\beta_T)$ is minimized at the unique point $\beta_T = \beta$.

The supervised Empirical Risk Minimization (ERM) approach in this context, such as in linear and logistic regression, suggests to minimize the target loss function over the training data, by solving the following optimization problem:

$$\hat{\beta} = \underset{\beta \in \mathbb{R}^p}{\operatorname{argmin}} \left\{ \hat{L}(\beta; X, Y) \right\} = \underset{\beta \in \mathbb{R}^p}{\operatorname{argmin}} \left\{ \frac{1}{n} \sum_{i=1}^n G(x_i^T \beta) - x_i^T \beta y_i \right\}. \quad (1)$$

As discussed before, the motivation for SSL arises when the amount of unlabelled data is much larger than the amount of the labeled data, meaning that $m \gg n, p$. In this case, it is natural to assume that we have the ability to accurately estimate the expectation $\mathbb{E}[\varphi(x)]$ for any given function φ , averaging it over the set Z as follows: $\mathbb{E}_x[\varphi(x)] \approx \frac{1}{m} \sum_{i=1}^m \varphi(z_i)$, and the approximation is arbitrarily good. In order to simplify the analysis in this paper and focus on key results, we adopt the *total information* scenario that assumes a diverging amount of unlabeled data, $m \rightarrow \infty$. Thus, we study methods for effectively using the exact knowledge of $\mathbb{E}_x[\varphi(x)]$ for some well-defined functions φ , to improve the predictive performance of a statistical learning process.

While the assumption of total information appears crucial, we emphasize its importance mainly for clarity and succinctness. Appendix B presents a brief analysis of a case without this assumption, demonstrating that the core arguments of this paper remain valid provided that the amount of unlabeled data exceeds that of the labeled data. This extension is accessible to curious readers and serves as a foundation for future research, which can potentially extend to various scenarios. Moreover, in empirical simulations, we estimate expected values $\mathbb{E}[\varphi(x)]$ by averaging φ on a large but finite set of unlabeled data, showing that breaking the assumption of total information does not significantly affect practical results. We note that in the total information scenario, the assumption that $\mathbb{E}[x] = \mathbf{0}_p$ is not limiting, since we can center the covariates as needed, by knowing $\mathbb{E}[x]$ precisely.

Taking advantage of the total information scenario, two semi-supervised estimators for β were introduced in Yuval and Rosset [2022]. These estimators integrate the knowledge of P_x into the optimization problem, resulting in a semi-supervised ERM strategy, which is detailed below:

$$\tilde{\beta} = \underset{\beta \in \mathbb{R}^p}{\operatorname{argmin}} \left\{ \tilde{L}(\beta; X, Y) \right\} = \underset{\beta \in \mathbb{R}^p}{\operatorname{argmin}} \left\{ \mathbb{E}_x [G(x^T \beta)] - \frac{1}{n} \sum_{i=1}^n x_i^T \beta y_i \right\}, \quad (2)$$

$$\begin{aligned} \check{\beta} &= \underset{\beta \in \mathbb{R}^p}{\operatorname{argmin}} \left\{ \check{L}(\beta; X, Y) \right\} \\ &= \underset{\beta \in \mathbb{R}^p}{\operatorname{argmin}} \left\{ \mathbb{E}_x [G(x^T \beta)] - \mathbb{E}_x [x^T \beta] \bar{Y} - \widehat{\operatorname{Cov}}(X\beta, Y) \right\}, \end{aligned} \quad (3)$$

where,

$$\widehat{\operatorname{Cov}}(X\beta, Y) = \sum_{i=1}^n (x_i^T \beta - \bar{X}\beta) (y_i - \bar{Y}) / n; \quad \bar{Y} = \frac{1}{n} \sum_{i=1}^n y_i; \quad \bar{X}\beta = \frac{1}{n} \sum_{i=1}^n x_i^T \beta.$$

The superiority of $\check{\beta}$ over $\tilde{\beta}$ was proven for the classical OLS model with Gaussian covariates, and also demonstrated in more general scenarios. This phenomenon is also observed in all the setups in this paper, and therefore we focus on $\check{\beta}$. As a result of the assumption that $\mathbb{E}[x] = \mathbf{0}_p$, we can omit the middle term of the objective function $\check{L}(\beta; X, Y)$.

From the analysis of the estimators above, Yuval and Rosset [2022] concluded that $\tilde{\beta}$ and $\check{\beta}$ outperform $\hat{\beta}$ only when the noise σ^2 is large enough. However, prediction performance can also be improved in a low noise scenario by choosing between $\hat{\beta}$ and $\check{\beta}$, adaptively, based on some data-driven estimates. In this work, we take this idea one step further and suggest using the knowledge of P_x in a more sophisticated manner, resulting in a better prediction performance. The main idea is to "mix" between $\hat{\beta}$ and $\check{\beta}$, rather than choosing between them, as we describe next.

The first suggested estimator is the linear-mixed estimator which is essentially a linear mixing between the two estimators:

$$\dot{\beta}_\alpha = (1 - \alpha)\hat{\beta} + \alpha\check{\beta}, \quad (4)$$

for some mixing ratio $\alpha \in [0, 1]$. The second learning process suggested is to mix the two

loss functions as follows:

$$\begin{aligned} \ddot{\beta}_\alpha &= \underset{\beta \in \mathbb{R}^p}{\operatorname{argmin}} \left\{ L_\alpha^M(\beta, X, Y) \right\} = \underset{\beta \in \mathbb{R}^p}{\operatorname{argmin}} \left\{ (1 - \alpha) \hat{L}(\beta, X, Y) + \alpha \check{L}(\beta, X, Y) \right\}, \\ &= \underset{\beta \in \mathbb{R}^p}{\operatorname{argmin}} \left\{ \frac{1}{n} \sum_{i=1}^n (1 - \alpha) G(x_i^T \beta) - x_i^T \beta y_i + \alpha (\mathbb{E}_x[G(x^T \beta)] + \overline{X\beta} \cdot \overline{Y}) \right\} \end{aligned} \quad (5)$$

In both learning procedures, the mixing ratio α represents the magnitude of integration of the unlabeled data in the learning process. By the definition of $\ddot{\beta}_\alpha$, the mixed loss function, L_α^M , is equivalent to $\hat{L} + \frac{\alpha}{1-\alpha} \check{L}$, for $\alpha < 1$, thus it can be viewed as a regularized ERM with the regularization term determined by the unlabeled data. This idea is related to some of the methods that were previously suggested in the domain of SSDL, in particular, to the idea of penalizing the predictive variance. However, the suggested regularization term \check{L} also takes into account the sample covariance between $X\beta$ and Y , and is more general in the sense that it can be applied to different link functions g . In fact, the predictive variance penalty is a special case of the first penalty term in L_α^M , where g is the identity, and therefore: $\mathbb{E}_x[G(x^T \beta)] = 0.5 \operatorname{Var}_x(f(x))$. Throughout this work, we analyze the prediction error of the suggested estimators, $\ddot{\beta}_\alpha$ and $\dot{\beta}_\alpha$, concluding that, in general, it is invariably beneficial to integrate the unlabeled data into the learning process (i.e., choose $\alpha > 0$), with a magnitude that increases in relation to the noise of the data, σ^2 . We note here that taking $\alpha > 0$, does not mean discarding any label y_i , since the whole term $\frac{1}{n} \sum_{i=1}^n x_i^T \beta y_i$ is taken into account in the objective function of the semi-supervised estimators.

1.2 Outline of the paper

In Section 2, we focus on the special case of the OLS model where the link function is the identity, and establish an explicit formula for α^* , the mixing ratio where $R(\dot{\beta}_\alpha)$ achieves its minimum. We also analyze the prediction error of $\ddot{\beta}_\alpha$ and prove that it has characteristics similar to $\dot{\beta}_\alpha$, with an advantage in a noisy environment. In Section 6.1, empirical simulations show that a data-driven estimate of α^* , or a grid search for the best mixing ratio of $\ddot{\beta}_\alpha$, can be used to achieve a significant improvement over the standard supervised model in the linear case. In Section 3, we analyze the semi-supervised GLM-ERM procedure for a general link function. We use a quadratic approximation to extend the

results from the OLS case, and establish an explicit formula for $\hat{\alpha}$, the mixing ratio where the approximation of $R(\hat{\beta}_\alpha)$ achieves its minimum. In Section 6.2, empirical simulations show that the quadratic approximation is accurate in terms of determining the best mixing ratio, and that the mixed-SSL is very effective in improving predictions in this general domain.

In Section 4 we demonstrate the applicability of the idea of mixed loss function to improve deep NN models for regression tasks, on a large-scale data set of images. We do so by viewing the last layer of the NN as a linear model that can be improved by applying the same methodology outlined in Section 2. We then optimize the entire neural network using the mixed loss function, which has been shown to be effective in practice. In Section 5, we broaden our approach to interpolating models by introducing the *mixed semi-supervised linear interpolator*. We demonstrate, both theoretically and empirically, that it is better than the standard minimum-norm interpolator.

2 Mixed Semi-supervised OLS

In the OLS model, the supervised and semi-supervised predictors (equations (1) and (2) respectively) can be written explicitly as follows:

$$\hat{\beta} = (X^T X)^{-1} X^T Y ; \check{\beta} = H^{-1} n \widehat{\text{Cov}}(X, Y) = H^{-1} (X^T Y - n \bar{X} \cdot \bar{Y}), \quad (6)$$

where $H = \mathbb{E}[X^T X]$, $[\bar{X}]_j = \frac{1}{n} \sum_{i=1}^n x_{ij}$ and $[\widehat{\text{Cov}}(X, Y)]_j = \widehat{\text{Cov}}([X]_j, Y)$, for $j = 1 \dots p$. From the explicit expressions above, we conclude that the mixing ratio α of the estimator $\hat{\beta}_\alpha$ is mainly related to the mixing between the known matrix H^{-1} (from the unlabeled data) and the observed matrix $(X^T X)^{-1}$ from the labeled data.

In order to compare between the estimator $\hat{\beta}$ and the semi-supervised estimators in terms of mean prediction error, we must take the following distributional assumption:

Assumption 1. *The distribution P_x satisfies that $(X^T X)^{-1}$ exists with probability 1, and also that $\mathbb{E}_X [(X^T X)^{-1}]$ exists.*

If Assumption 1 is violated, then $R(\hat{\beta}) = \infty$ or is not defined, making the semi-supervised alternative more attractive. In order to analyze the prediction error of $\check{\beta}$ and

the mixed estimators $\hat{\beta}_\alpha$ and $\check{\beta}_\alpha$, we use the following observation:

$$n\bar{X}\bar{X}^T = \frac{1}{n}X^T X + C_X; [C_X]_{jk} = \frac{1}{n} \sum_{i \neq l} x_{ij} x_{lk}, \quad (7)$$

$$\mathbb{E}[C_X] = \mathbb{E}[X^T X C_X] = \mathbb{E}[(X^T X)^{-1} C_X] = \mathbf{0}_{p \times p}.$$

As a result, the term C_X cancels out in many of the expectations that follow, unless it is squared.

Under the assumption of true linear model (meaning that $\mathbb{E}[y|x] = x^T \beta$ for some $\beta \in \mathbb{R}^p$), the prediction error of these predictors (up to the irreducible term $0.5\beta^T H \beta/n$), can be summarized into the following bias-variance decomposition:

$$\begin{aligned} R(\hat{\beta}) &= \frac{\sigma^2}{n} \text{tr} \left(\mathbb{E}_X [(X^T X)^{-1}] H \right) := \sigma^2 v_l, \\ R(\check{\beta}) &= \underbrace{\frac{1}{n} \text{tr} \left(H^{-1} \text{Var}_X \left(\widehat{\text{Cov}}(X, X\beta) \right) \right)}_{B(\check{\beta})} + \underbrace{\left(1 - \frac{1}{n} \right) \frac{\sigma^2 p}{n}}_{V(\check{\beta})} := B(\check{\beta}) + \sigma^2 v_u. \end{aligned}$$

The term $B(\check{\beta})$ is the bias component of $R(\check{\beta})$, and $V(\check{\beta}) = \sigma^2 v_u$ is its variance component. In the same manner, the term $\sigma^2 v_l$ is the variance component of $R(\hat{\beta})$, since its bias component is zero. As discussed in Yuval and Rosset [2022], v_l is always larger than v_u , and therefore we have a bias-variance trade-off between the two estimators. The *adaptive estimator* in Yuval and Rosset [2022], denoted by β^D , is equal to $\check{\beta}$ if $\sigma^2 > B(\check{\beta})/(v_l - v_u)$ according to some data-based estimates of the relevant quantities, and equal to $\hat{\beta}$ otherwise.

Using the same notations, the prediction error of the linear-mixed estimator $\dot{\beta}_\alpha$, as a function of α , can be summarized as follows:

$$\begin{aligned} R(\dot{\beta}_\alpha) &= \underbrace{\mathbb{E}_{X, x_0} \left(\mathbb{E} \left[x_0^T \dot{\beta}_\alpha | X, x_0 \right] - x_0^T \beta \right)^2}_{B(\dot{\beta}_\alpha)} + \underbrace{\mathbb{E}_{X, x_0} \left[\text{Var} \left(x_0^T \dot{\beta}_\alpha | X, x_0 \right) \right]}_{V(\dot{\beta}_\alpha)} \\ &= \alpha^2 B(\check{\beta}) + \alpha^2 \sigma^2 v_u + (1 - \alpha)^2 \sigma^2 v_l \\ &\quad + 2\alpha(1 - \alpha) \mathbb{E}_{X, x_0} \left[\text{Cov} \left(x_0^T \hat{\beta}, x_0^T \check{\beta} | X, x_0 \right) \right]. \end{aligned}$$

In the above-described equation, $B(\dot{\beta}_\alpha)$ represents the bias term of $R(\dot{\beta}_\alpha)$, and $V(\dot{\beta}_\alpha)$ represents the variance term. Analysis of the above formula leads us to the following theorem regarding the characteristics of $\dot{\beta}_\alpha$:

Theorem 2.1. *For any distribution P_X satisfying Assumption 1, and fixed vector β , the unique global minimizer of $R(\dot{\beta}_\alpha)$, denoted by α^* , is in the open interval $(0, 1)$. Moreover, α^* admits the following explicit formula:*

$$\alpha^* = \frac{\sigma^2(v_l - v_u)}{B(\check{\beta}) + \sigma^2(v_l - v_u)},$$

and the minimum value of $R(\dot{\beta}_\alpha)$ denoted by $R(\dot{\beta}_{\alpha^})$, admits the following explicit formula:*

$$R(\dot{\beta}_{\alpha^*}) = \sigma^2 v_l - \frac{\sigma^4(v_l - v_u)^2}{B(\check{\beta}) + \sigma^2(v_l - v_u)} < R(\hat{\beta}).$$

The proof of Theorem 2.1 is provided in Appendix A.1. From Theorem 2.1, we conclude that $\dot{\beta}_{\alpha^*}$ is a better estimator than both $\hat{\beta}$ and $\check{\beta}$, under the assumption of a true linear model, and any noise σ^2 and real signal β . This theoretical result differs significantly from the main conclusion in previous related works [Yuval and Rosset, 2022, Azriel et al., 2022, Chakraborty and Cai, 2018] that discussed the usefulness of unlabeled data in the OLS model and stated that, in theory, the unlabeled data is useful only when the model is biased, nonlinear, or extremely noisy. On the other hand, the fact that α^* (and also $R(\hat{\beta}) - R(\dot{\beta}_{\alpha^*})$) increases with σ^2 , giving more weight to the unlabeled data (through the knowledge of H), coincides with previous observations on the usefulness of unlabeled data.

In practice, it is necessary to estimate α^* from the data in hand to decide on the mixing ratio. We propose to estimate σ^2 by using the standard unbiased estimator $\hat{\sigma}^2 = RSS(\hat{\beta})/(n-p)$, where $RSS(\hat{\beta}) = \|X\hat{\beta} - Y\|_2^2$ is the residual sum of squares of the estimator $\hat{\beta}$, and to assess $B(\check{\beta})$ using the estimated vector $\check{\beta}$. The term v_l can be estimated precisely from the unlabeled data. The linear mixed estimator with the estimated mixing ratio $\hat{\alpha}$ is denoted by $\dot{\beta}_{\hat{\alpha}}$. In Section 2.1 (Proposition 2.3), we investigate the performance of the adaptive mixed estimator $\dot{\beta}_{\hat{\alpha}}$ in a more specific context and find that it exhibits very similar performance to the oracle estimator $\dot{\beta}_{\alpha^*}$.

We now turn our attention to the estimator $\ddot{\beta}_\alpha$, and analyze the first and second derivatives of $L_\alpha^M(\beta, X, Y)$ with respect to β . It is straightforward to show that L_α^M is convex in β , and that in the OLS case, the optimization problem in (5) has the following explicit solution:

$$\begin{aligned} \ddot{\beta}_\alpha &= (\alpha H + (1 - \alpha)X^T X)^{-1} (X^T Y - \alpha n \bar{X} \cdot \bar{Y}) \\ &:= S_\alpha(X^T Y - \alpha n \bar{X} \cdot \bar{Y}), \end{aligned}$$

where $S_\alpha = (\alpha H + (1 - \alpha)X^T X)^{-1}$. Moreover, the integrated risk of $\ddot{\beta}_\alpha$ can be summarized in the following bias-variance decomposition:

$$\begin{aligned} R(\ddot{\beta}_\alpha) &= \mathbb{E}_{X, x_0} \left(\mathbb{E} \left[x_0^T \ddot{\beta}_\alpha | X, x_0 \right] - x_0^T \beta \right)^2 + \mathbb{E}_{X, x_0} \left[\text{Var} \left(x_0^T \ddot{\beta}_\alpha | X, x_0 \right) \right] \\ &= \underbrace{\frac{1}{n} \text{tr} \left(\mathbb{E}_X \left[\beta \beta^T \Delta_\alpha^T H \Delta_\alpha \right] \right)}_{B(\ddot{\beta}_\alpha)} + \underbrace{\frac{\sigma^2 \xi_\alpha}{n} \text{tr} \left(H \mathbb{E}_X \left[S_\alpha X^T X S_\alpha \right] \right)}_{V(\ddot{\beta}_\alpha)}, \end{aligned} \quad (8)$$

where,

$$\Delta_\alpha = S_\alpha (X^T X - \alpha n \bar{X} \bar{X}^T) - I ; \quad \xi_\alpha = 1 - (2\alpha - \alpha^2)/n \quad (9)$$

In the displayed equation (8), $B(\ddot{\beta}_\alpha)$ represents the bias term of $R(\ddot{\beta}_\alpha)$, and $V(\ddot{\beta}_\alpha)$ represents the variance term. Unlike the estimator $\dot{\beta}_\alpha$, finding the exact value of α that minimizes the prediction error of $\ddot{\beta}_\alpha$ might not be possible even in the linear model. However, the following theorem shows that, similarly to the case of $\dot{\beta}_\alpha$, it is necessarily beneficial to mix the loss functions \hat{L} and \check{L} , and consequently take advantage of the knowledge of H obtained from the unlabeled data.

Theorem 2.2. *For any distribution P_X satisfying Assumption 1, and fixed vector β , the integrated risk of $\ddot{\beta}_\alpha$ achieves its (not necessarily unique) minimum in the open interval $\alpha \in (0, 1)$.*

The proof of Theorem 2.2 is provided in Appendix A.2, and from it we can deduce that the behavior of $R(\ddot{\beta}_\alpha)$ is analogous to that of $R(\dot{\beta}_\alpha)$ with respect to the mixing ratio α . However, since we are not able to explicitly find the best mixing ratio for $\ddot{\beta}_\alpha$, we suggest two alternatives for choosing the mixing ratio in practice:

1. Choosing the same estimated value $\hat{\alpha}$ as that used for $\dot{\beta}_\alpha$, as a heuristic measure for the "right" mixing ratio in the optimization problem. The resulting estimator is denoted by $\ddot{\beta}_{\hat{\alpha}}$.
2. Performing a grid search on the $(0, 1)$ interval for α that minimizes the estimate of $R(\ddot{\beta}_\alpha)$, while using the estimators $\hat{\sigma}^2$ and $\check{\beta}$ combined with the unlabeled data set at any point on the grid, according to formulas (8) and (9). The resulting estimator is denoted by $\ddot{\beta}_{\hat{\alpha}}$.

While we are not able to directly compare the minimum value of $R(\ddot{\beta}_\alpha)$ with $R(\dot{\beta}_{\alpha^*})$ or $R(\hat{\beta})$, the next theorem focuses on variance components $V(\ddot{\beta}_\alpha)$ and $V(\dot{\beta}_\alpha)$, and shows that $\ddot{\beta}_\alpha$ has a better robustness to noisy data compared to $\dot{\beta}_\alpha$, for any given mixing ratio α .

Theorem 2.3. *For any distribution P_X satisfying Assumption 1, fixed vector β , and given mixing ratio α , the following inequality holds:*

$$V(\ddot{\beta}_\alpha) \leq V(\dot{\beta}_\alpha).$$

The proof Theorem 2.3 is provided in Appendix A.3. From Theorem 2.3 we conclude that $V(\ddot{\beta}_{\alpha^*}) \leq V(\dot{\beta}_{\alpha^*})$, and when the variance term is dominant over the bias term, we expect $\ddot{\beta}_{\alpha^*}$ to perform better than $\dot{\beta}_{\alpha^*}$. In same manner, we expect that the adaptive estimators $\ddot{\beta}_{\hat{\alpha}}$ and $\ddot{\beta}_{\check{\alpha}}$ will have better robustness to noisy data compared to $\dot{\beta}_{\hat{\alpha}}$. We also emphasize that the variance term $V(\ddot{\beta}_\alpha)$ is strictly convex in α , since its second derivative is positive, unlike the bias term $B(\ddot{\beta}_\alpha)$, which is more difficult to analyze. The analysis of $B(\ddot{\beta}_\alpha)$ is done by a third-order polynomial approximation, under the following distributional assumption, which is closely aligned with that proposed in El Karoui [2009].

Assumption 2. *The distribution P_x satisfies the following generating mechanism: $x = \Sigma^{1/2}\tilde{x}$, where $\tilde{x} \in \mathbb{R}^p$ is a random vector with i.i.d. entries that have zero mean, unit variance, and a finite fourth moment q_4 , a finite sixth moment q_6 , and Σ is a positive definite covariance matrix.*

Under Assumption 2, we can show that up to a third-order polynomial approximation, the bias term of $\ddot{\beta}_\alpha$ is smaller than the of $\dot{\beta}_\alpha$, as stated in the following theorem.

Theorem 2.4. *Consider any distribution P_x satisfying Assumption 2, fixed vector β , and let $\ddot{b}(\alpha)$ be a function of α of the form $\ddot{b}(\alpha) = b_0 + b_1\alpha + b_2\alpha^2 + b_3\alpha^3$, that satisfies the following constraints:*

$$\begin{aligned} \ddot{b}(0) &= B(\ddot{\beta}_{\alpha=0}) ; \quad \ddot{b}'(0) = \frac{\partial}{\partial \alpha} B(\ddot{\beta}_\alpha)|_{\alpha=0}, \\ \ddot{b}(1) &= B(\ddot{\beta}_{\alpha=1}) ; \quad \ddot{b}'(1) = \frac{\partial}{\partial \alpha} B(\ddot{\beta}_\alpha)|_{\alpha=1}. \end{aligned}$$

Then, for any given mixing ratio α , the following inequality holds:

$$\ddot{b}(\alpha) \leq B(\dot{\beta}_\alpha).$$

The proof Theorem 2.4 is provided in Appendix A.4. From Theorem 2.4 we conclude that the third-order polynomial approximation of $B(\ddot{\beta}_\alpha)$ is uniformly smaller than $B(\dot{\beta}_\alpha)$ for $0 < \alpha < 1$, in a wide class of distributions for the covariates. If α^* is small, the higher order terms of $B(\ddot{\beta}_\alpha)$ are negligible and therefore $B(\ddot{\beta}_{\alpha^*}) \lesssim B(\dot{\beta}_{\alpha^*})$. On the other hand, a large value of α^* yields that the variance term is dominant, and together with Theorem 2.3, we can expect that the estimator $\ddot{\beta}_{\alpha^*}$ outperforms the estimators $\dot{\beta}_{\alpha^*}$, $\hat{\beta}$ and $\check{\beta}$. This can be summarized by the following chain of inequalities:

$$\min_{\alpha} \left\{ R(\ddot{\beta}_\alpha) \right\} \leq R(\ddot{\beta}_{\alpha^*}) \lesssim R(\dot{\beta}_{\alpha^*}) \leq R(\hat{\beta}).$$

In practice, if the estimation of α^* and the grid search are reliable, we can expect the following to hold:

$$R(\ddot{\beta}_{\hat{\alpha}}) \lesssim R(\ddot{\beta}_{\hat{\alpha}}) \lesssim R(\dot{\beta}_{\hat{\alpha}}) \lesssim R(\hat{\beta}).$$

In Section 6.1, we demonstrate empirically that the above chain of inequalities holds in simulations of the OLS model, with considerably small differences between the two terms on the left side.

We also analyze the suggested mixed semi-supervised estimators in a different scenario, where the vector β is also random. Let us assume that with every training sample $T = (X, Y)$, β is drawn independently from a prior distribution P_β such that $\mathbb{E}[\beta] = \mathbf{0}$ and $\mathbb{E}[\beta\beta^T] = \tau^2 I_p$. In this case, the mean out-of-sample prediction error is averaged over all possible β 's, and can be written (up to the irreducible term $0.5\tau^2\text{tr}(H)/n$) as follows:

$$\begin{aligned} R(\check{\beta}) &= \mathbb{E}_\beta \left[R(\check{\beta}) \right] = \frac{\tau^2}{n} \text{tr} \left(\mathbb{E}_X [H\Delta_1\Delta_1^T] \right) + \frac{(n-1)\sigma^2 p}{n^2} \\ &:= \tau^2 b_u + \sigma^2 v_u := B(\check{\beta}) + V(\check{\beta}), \end{aligned}$$

where $\Delta_1 = \Delta_{\alpha=1} = H^{-1}(X^T X - n\bar{X}\bar{X}^T) - I$, as defined in equation (9). We note that in this case, we use the term b_u that is associated with the bias term $B(\check{\beta})$, and can be estimated precisely with the unlabeled data.

While the setting of random- β is widely used and realistic in many practical instances, it simplifies the problem in two aspects. First, the estimation of the bias term reduces to the estimation of τ^2 by the labeled data, and b_u by the unlabeled data. Second, the mathematical analysis simplifies, and stronger theoretical results can be obtained, as we show in Section 2.1. Moreover, since all the results so far relate to the mean prediction error

and were obtained for any given vector β , they hold in expectation over β . In particular, we can use the identity $B(\check{\beta}) = \tau^2 b_u$, and adjust the main result regarding α^* and $R(\dot{\beta}_{\alpha^*})$ to this scenario, as follows:

$$\alpha^* = \frac{\sigma^2(v_l - v_u)}{\tau^2 b_u + \sigma^2(v_l - v_u)} ; \quad R(\dot{\beta}_{\alpha^*}) = \sigma^2 v_l - \frac{\sigma^4(v_l - v_u)^2}{\tau^2 b_u + \sigma^2(v_l - v_u)}. \quad (10)$$

Thus, in this case, our problem reduces to estimating the noise σ^2 and the signal τ^2 from the labeled data in order to estimate α^* . Moreover, we can write the bias term of $\ddot{\beta}_\alpha$ in this scenario as follows:

$$B(\ddot{\beta}_\alpha) = \frac{\tau^2}{n} \text{tr} \left(\mathbb{E}_X [H \Delta_\alpha \Delta_\alpha^T] \right).$$

We note that for any given α , the term $\text{tr} \left(\mathbb{E}_X [H \Delta_\alpha \Delta_\alpha^T] \right)$ depends only on P_X and can be estimated precisely with the unlabeled data. Thus, we can use the above expression to estimate $R(\ddot{\beta}_\alpha)$ at any given α based on estimates of σ^2 and τ^2 , and choose the best estimated mixing ratio for the estimator $\ddot{\beta}_{\hat{\alpha}}$. For the task of estimating the noise and the signal, we suggest the following estimators:

$$\hat{\tau}^2 = \max \left\{ \left(\frac{\sum_{i=1}^n y_i^2}{n} - \hat{\sigma}^2 \right) / \text{tr} \left(\mathbb{E} [xx^T] \right), 0 \right\}, \quad \text{where } \hat{\sigma}^2 = \frac{RSS(\hat{\beta})}{n - p}.$$

With the estimators above, we can fit the mixed semi-supervised estimators $\dot{\beta}_{\hat{\alpha}}$, $\ddot{\beta}_{\hat{\alpha}}$, and $\ddot{\beta}_{\hat{\alpha}}$, suitable for this scenario. In Section 6.1, Figures 4 and 5 present a summary of the numerical experiment on the prediction error of all the estimators under discussion. These results align with the theoretical results outlined in this section. In the next subsection, we analyze the usefulness of $\dot{\beta}_{\alpha^*}$ and $\dot{\beta}_{\hat{\alpha}}$ in this scenario, in an asymptotic setting of n and p .

2.1 Asymptotic usefulness of the mixed SSL

The primary theoretical finding concerning the usefulness of unlabeled data in enhancing predictions suggests that $R(\dot{\beta}_{\alpha^*}) < R(\hat{\beta})$ holds for any sample size n . Our aim is to investigate how the sample size impacts the difference between these two quantities and to confirm that the advantage gained from utilizing unlabeled data does not diminish as n diverges. By defining η as the relative prediction error between $\dot{\beta}_{\alpha^*}$ and $\hat{\beta}$ for a finite sample, we can confirm that the SSL approach remains beneficial in the asymptotic setting by showing that η stays strictly below 1 as n diverges.

In the random- β scenario, the expression for $R(\hat{\beta}_\alpha)$ from Equation (10) can be employed to derive the following representation for η :

$$\eta = \frac{R(\beta^{M^*})}{R(\hat{\beta})} = 1 - \frac{\sigma^2(v_l - v_u)^2}{\tau^2 v_l b_u + \sigma^2 v_l (v_l - v_u)}. \quad (11)$$

For a specific pair of (σ^2, τ^2) values, the formula above indicates that the advantage gained from using the mixed semi-supervised estimator is significant as long as the disparity $v_l - v_u$ is substantial relative to the values of b_u and v_l . To thoroughly examine the asymptotic properties of η , we investigate an asymptotic scenario where $p/n \rightarrow \gamma \in (0, 1)$ as $n, p \rightarrow \infty$, and the covariates' distribution adheres to the following assumption.

Assumption 3. *The distribution P_x satisfies $\text{Var}(x) = \Sigma$, where Σ is (a sequence of) invertible $p \times p$ matrix, and also $\text{tr}(\Sigma) \rightarrow c^2$ for some constant $c \in \mathbb{R}$, as $p \rightarrow \infty$.*

The following proposition describes the asymptotic behavior of η and α^* under the conditions of Assumption 3, in the case where the covariates follow the multidimensional normal distribution. We denote by $x \sim MN(\mathbf{0}_p, \Sigma)$, a covariate p -dimensional random vector from multivariate normal distribution with mean $\mathbf{0}_p$ and covariance matrix Σ .

Proposition 2.1. *Consider the asymptotic setting $p/n \rightarrow \gamma \in (0, 1)$ as $n, p \rightarrow \infty$, and assume that $x \sim MN(\mathbf{0}_p, \Sigma)$, where Σ follows the conditions of Assumption 3, then as $n, p \rightarrow \infty$:*

$$\begin{aligned} \eta &\rightarrow \eta_\infty = 1 - \frac{\gamma^4 \sigma^2}{(1 - \gamma)\gamma^2 \tau^2 c^2 + \gamma^3 \sigma^2}, \\ \alpha^* &\rightarrow \alpha_\infty^* = \frac{\gamma^2 \sigma^2}{(1 - \gamma)\gamma \tau^2 c^2 + \gamma^2 \sigma^2}. \end{aligned}$$

The proof of Proposition 2.1 can be found in Appendix A.5, which indicates that the limit of η is less than 1 for any positive γ . Additionally, insights can be gained regarding the impact of dimensionality on the relative advantage of utilizing unlabeled data. As γ approaches 0, η tends towards 1 and α towards 0, indicating no advantage in leveraging unlabeled data. Conversely, as γ approaches 1, η tends towards 0 and α towards 1, suggesting that unlabeled data is highly beneficial.

The key insight from Proposition 2.1 about the usefulness of the unlabeled data in an asymptotic setting, when Gaussian covariates are considered, can be generalized to

encompass a broader range of covariate distributions. The following proposition describes the asymptotic behavior of η and α^* under the generating mechanism of Assumption 2.

Proposition 2.2. *Consider the asymptotic setting as described above, and assume that Assumptions 1, 2 and 3 hold, then for any $\epsilon > 0$, there exists an $n(\epsilon)$ such that:*

$$\eta \leq \eta_\infty + \epsilon \text{ and } \alpha^* \geq \alpha_\infty^* - \epsilon, \quad \forall n > n(\epsilon).$$

The proof of Proposition 2.2 can be found in Appendix A.6, and from it we conclude that, under a wide class of distributions, the limiting value of η lies in the interval $(0, \eta_\infty]$, which means that $\dot{\beta}_{\alpha^*}$ significantly outperforms $\hat{\beta}$. Consider the scenario where τ^2 is known, but we need to estimate σ^2 in order to evaluate the best mixing ratio α^* . Denoting by $\dot{\beta}_{\hat{\alpha}(\tau)}$ the estimator that uses known τ^2 and the estimator $\hat{\sigma}^2$, in estimating the best mixing ratio, the following proposition indicates that we should expect a comparable prediction error between $\dot{\beta}_{\hat{\alpha}(\tau)}$ and $\dot{\beta}_{\alpha^*}$.

Proposition 2.3. *Consider the asymptotic setting described above and assume that the distribution P_x satisfies that $b_u \rightarrow \gamma c^2$, and $v_l \rightarrow \gamma/(1 - \gamma)$ as $n \rightarrow \infty$, then the random difference between the predictions made using $\dot{\beta}_{\hat{\alpha}(\tau)}$ and $\dot{\beta}_{\alpha^*}$ satisfies:*

$$x_0^T \dot{\beta}_{\alpha^*} - x_0^T \dot{\beta}_{\hat{\alpha}(\tau)} \xrightarrow{P} 0, \text{ as } n \rightarrow \infty.$$

The proof of Proposition 2.3 can be found in Appendix A.7. Proposition 2.3 offers an understanding of the similarity between the data-driven estimator $\dot{\beta}_{\hat{\alpha}(\tau)}$ and the oracle estimator $\dot{\beta}_{\alpha^*}$ when τ^2 is known. On the other hand, if τ^2 is unknown, the estimator $\hat{\alpha}$ is not consistent, since the variance of $\hat{\tau}^2$ is $2\tau^4 \text{tr}(\Sigma^2)/(\text{tr}(\Sigma))^2$, which does not vanish with n in general (depending on the structure of Σ). In this case, the estimator $\dot{\beta}_{\hat{\alpha}}$ can diverge from $\dot{\beta}_{\alpha^*}$, resulting in a different prediction error. However, in practice, estimating τ^2 appears to be better than knowing it, as will be demonstrated in the next subsection. We attribute this to the fact that the estimation of τ^2 takes into account the specific realization of β , which leads to a better mixing ratio per training sample. Figure 6 presents a summary of the numerical findings on the asymptotic trends of the prediction error for the mixed semi-supervised estimators compared to the supervised estimator across various scenarios. These results align with the theoretical results outlined in propositions 2.1 and 2.3.

3 Mixed Semi-supervised GLM

In this section we analyze the mixed semi-supervised GLM-ERM for general monotone increasing link function g . We do it under the assumption of a well-specified model, meaning that $\mathbb{E}[y|x] = g(x^T\beta)$, for some real vector $\beta \in \mathbb{R}^p$. For this case, we already defined the estimators $\hat{\beta}$, $\check{\beta}$, $\dot{\beta}_\alpha$ and $\ddot{\beta}_\alpha$ in Equations (1), (3), (4) and (5), respectively, through optimization problems. However, as described in Yuval and Rosset [2022], by using the quadratic approximation of the convex loss functions $\hat{L}(\beta; X, Y)$ and $\check{L}(\beta; X, Y)$ around the real vector β , the solutions for $\hat{\beta}$ and $\check{\beta}$ can be written in the following approximated explicit formulas:

$$\hat{\beta} \approx \beta - (X^T DX)^{-1} X^T (\mu - Y) := \beta - \hat{a}, \quad (12)$$

$$\check{\beta} \approx \beta - H^{-1} \left(\mathbb{E}_X[X^T \mu] - n \widehat{\text{Cov}}(X, Y) \right) := \beta - \check{a}, \quad (13)$$

where $H = \mathbb{E}_X[X^T DX]$, D is a $n \times n$ diagonal matrix with the terms $D_{ii} = g'(x_i \beta)$, and $\mu \in \mathbb{R}^n$, with $\mu_i = g(x_i^T \beta)$. The approximate equivalences above describe the linear difference between every estimator and the true vector β , assuming that the loss functions were quadratic in β . Furthermore, the approximate prediction error of these estimators (up to multiplication by a constant and addition of an irreducible term), can be summarized as follows:

$$\begin{aligned} R(\hat{\beta}) &\approx \mathbb{E}_T [\hat{a}^T H \hat{a}] \\ &= \sigma^2 \text{tr} \left(\mathbb{E}_X [(X^T DX)^{-1} X^T X (X^T DX)^{-1}] H \right) := \sigma^2 v_l, \\ R(\check{\beta}) &\approx \mathbb{E}_T [\check{a}^T H \check{a}] = \text{tr} \left(H^{-1} \text{Var}_X(n \widehat{\text{Cov}}(X, \mu)) \right) \\ &+ \frac{n-1}{n} \sigma^2 \text{tr} (H^{-1} \mathbb{E}_X[X^T X]) := B(\check{\beta}) + \sigma^2 v_u. \end{aligned} \quad (14)$$

From the above, we can see that as long as $\infty > v_l > v_u$, and the quadratic approximation error is negligible in the context of comparing between the $R(\hat{\beta})$ and $R(\check{\beta})$, there is a bias-variance trade-off between the supervised and the semi-supervised estimators. The condition $v_l > v_u$ is also a key condition for the effectiveness of the mixed estimators, as we will show next. Although we are unable to prove that $v_l > v_u$ in general, we can do it under the following distributional assumption.

Assumption 4. For any fixed vector β , the distribution P_x and the link function g satisfy the following: $(X^TDX)^{-1}$ exists with probability 1, $v_l < \infty$, and $\mathbb{E}[g'(x^T\beta)|xx^T] = \delta$, for some constant $\delta > 0$, which is independent of x .

Assumption 4 states that the knowledge of xx^T has no effect on the expected value of $g'(x^T\beta)$. While it might not seem a mild assumption, we can show that a sufficient condition is that P_x is symmetric around 0, and that $g'(z) = \delta + \varphi(z)$ for some odd function φ , bounded between $-\delta$ and δ . It follows from the fact that the conditional distribution of $\varphi(x^T\beta)$ is symmetric around 0 (and therefore $\mathbb{E}[\varphi(x^T\beta)|xx^T] = 0$):

$$\begin{aligned} P_{\varphi(x^T\beta)=d|xx^T} &= \frac{P_x(\varphi(x^T\beta) = d, xx^T)}{P_x(xx^T)} \\ &= \frac{P_x(\varphi(x^T\beta) = -d, xx^T)}{P_x(xx^T)} = P_{\varphi(x^T\beta)=-d|xx^T}. \end{aligned}$$

We also note that the widely used Relu and Leaky-ReLU functions comply with the above condition on the link function, as well as the identity function.

Proposition 3.1. Under Assumption 4, For any fixed vector β , the inequality $v_l > v_u$ holds.

The proof of Proposition 3.1 can be found in Appendix A.8. For the sake of our theoretical analysis, we will assume that Assumption 4 holds throughout this Section, and consequently $v_l > v_u$, as shown by Proposition 3.1. However, we point to the fact that empirical simulations suggest that this inequality holds in many scenarios where Assumption 4 is violated, supporting the conjecture that it holds in general. Moreover, in practice, the quantities v_l and v_u need to be estimated from the data on hand, to obtain an estimate of the best mixing ratio, which will be 0 unless the estimated value of $v_l - v_u$ is positive. Therefore, while the theoretical results are restricted to the conditions of Assumption 4, the suggested methodology is not. Finally, in Section 7, we show that under a different model for conditional variance $\text{Var}[y|x]$, the terms v_l and v_u get slightly different forms and the inequality $v_l > v_u$ is guaranteed to always hold.

Given formulas (12) and (13) above, the linear-mixed estimator $\dot{\beta}_\alpha$ can be written in the following approximated explicit formula:

$$\dot{\beta}_\alpha \approx \beta - [(1 - \alpha)\hat{a} + \alpha\check{a}] := \beta - \dot{a}_\alpha,$$

and further more, the prediction error of $\dot{\beta}_\alpha$, can be approximated by the function $\dot{r}(\alpha)$ as follows:

$$\begin{aligned} R(\dot{\beta}_\alpha) &\approx \mathbb{E}_T [\dot{a}_\alpha^T H \dot{a}_\alpha] := \dot{r}(\alpha) \\ &= \underbrace{\alpha^2 B(\check{\beta})}_{\dot{B}(\alpha)} + \underbrace{\sigma^2 (\alpha^2 v_u + (1 - \alpha)^2 v_l + 2\alpha(1 - \alpha)\mathbb{E}_T [\hat{a}^T H \check{a}])}_{\dot{V}(\alpha)}. \end{aligned}$$

In the equation presented above, $\dot{B}(\alpha)$ and $\dot{V}(\alpha)$ correspond to the bias and variance components of $\dot{r}(\alpha)$, respectively. Assuming that the quadratic approximation error is negligible in the context of determining the best mixing ratio, α^* , we suggest analyzing the characteristics of the approximate prediction error, $\dot{r}(\alpha)$, in order to gain insights into $R(\dot{\beta}_\alpha)$ in the GLM case, and determine the proper mixing ratio. The following proposition describes the properties of $\text{argmin}\{\dot{r}(\alpha)\}$, which is denoted by $\hat{\alpha}$.

Proposition 3.2. *Under Assumption 4, for any fixed vector β , the unique global minimizer of $\dot{r}(\alpha)$, $\hat{\alpha}$, is in the open interval $(0, \infty)$. Moreover, $\hat{\alpha}$ admits the following explicit formula:*

$$\hat{\alpha} = \frac{\sigma^2(v_l - v_s)}{B(\check{\beta}) + \sigma^2(v_l + v_u - 2v_s)},$$

where,

$$v_s = \frac{n-1}{n} \text{tr}(\mathbb{E}_X [X^T X (X^T D X)^{-1}]),$$

and the following inequality holds: $v_l + v_u - 2v_s > 0$. Moreover, the minimum value $\dot{r}(\hat{\alpha})$, admits the following explicit formula:

$$\dot{r}(\hat{\alpha}) = \sigma^2 v_l - \frac{\sigma^4 (v_l - v_u)^2}{B(\check{\beta}) + \sigma^2 (v_l + v_u - 2v_s)}.$$

The proof of Proposition 3.2 can be found in Appendix A.9. From Proposition 3.2 we conclude that, based on the quadratic approximation, it is always beneficial to slightly move from the supervised estimator towards the semi-supervised one. However, if $v_s > v_u$ (which is true under Assumption 4) and σ^2 is extremely large, $\hat{\alpha}$ can be greater than 1. This characteristic of $\dot{r}(\alpha)$ might be an undesirable "side effect" of the quadratic approximation used in all the derivations. However, the mixing ratio is limited by definition to the $[0, 1]$ interval, as is our estimator of $\hat{\alpha}$, which is practically set to be the minimum between 1

and the estimated value of the above formula. In order to estimate $\hat{\alpha}$ from the data, we use the following approximately unbiased estimator of σ^2 :

$$\hat{\sigma}^2 = RSS(\hat{\beta}) / (n - 2p + \text{tr}(\mathbb{E}_X [X^T X H_X^{-1} X^T D^2 X H_X^{-1}])) ,$$

where $D_{ii} = g'(x_i^T \check{\beta})$. It should be noted that, unlike the OLS model, the v -terms are also evaluated at the estimated vector $\check{\beta}$, utilizing the unlabeled data.

In order to analyze the predictive performance of $\check{\beta}_\alpha$, we adopt the concept of quadratic approximation of the loss function to equation (5), and obtain the following approximated explicit formula:

$$\begin{aligned} \check{\beta}_\alpha &\approx \beta - S_\alpha (\alpha(\mathbb{E}_X[X^T \mu] + n\bar{X} \cdot \bar{Y}) + (1 - \alpha)X^T \mu - X^T Y) \\ &:= \beta - \check{a}_\alpha, \end{aligned}$$

where $S_\alpha = (\alpha H + (1 - \alpha)X^T D X)^{-1}$. Furthermore, the quadratic approximation of the prediction error, $R(\check{\beta}_\alpha)$ is proportional to the quantity $\ddot{r}(\alpha) = \mathbb{E}_T [\check{a}_\alpha^T H \check{a}_\alpha]$, up to an irreducible term. The quantity $\ddot{r}(\alpha)$ can be decomposed into the following variance and bias terms:

$$\ddot{r}(\alpha) = \underbrace{\alpha^2 \text{tr}(\mathbb{E}_X [S_\alpha H S_\alpha \zeta \zeta^T])}_{\ddot{B}(\alpha)} + \underbrace{\xi_\alpha \sigma^2 \text{tr}(H \mathbb{E}_X [S_\alpha X^T X S_\alpha])}_{\ddot{V}(\alpha)},$$

where $\zeta = \mathbb{E}_X[X^T \mu] - n\widehat{\text{Cov}}(X, \mu) \in \mathbb{R}^p$.

The following proposition uses the above quantities to describe the characteristics of $\ddot{r}(\alpha)$, which in turn tells us about the characteristics of $R(\check{\beta}_\alpha)$.

Proposition 3.3. *Under Assumption 4, for any vector β , the quantity $\ddot{r}(\alpha)$ achieves its minimum in the open interval $(0, 1)$. Moreover, at any given mixing ratio α , the following inequality holds: $\ddot{V}(\alpha) \leq \dot{V}(\alpha)$.*

The proof of Proposition 3.3 can be found in Appendix A.10. From Proposition 3.3 we learn that $\ddot{r}(\alpha)$ has a behavior similar to $\dot{r}(\alpha)$, but with a smaller variance term, making $\check{\beta}(\alpha)$ better than $\dot{\beta}(\alpha)$ in high-variance scenarios. Taking a wider perspective on the theoretical analysis of this section, our main observation, that mixing between the estimators improves prediction, is generalized from the linear model (Theorem 2.1 and Theorem 2.2)

to the GLM class of models. As long as the quadratic approximation error is negligible, we can adopt the estimation methodology of $\dot{\alpha}$ as the desired mixing ratio for $\dot{\beta}_\alpha$ and $\ddot{\beta}_\alpha$, and use it to fit the estimators $\dot{\beta}_{\dot{\alpha}}$ and $\ddot{\beta}_{\dot{\alpha}}$ respectively. We can also perform a grid search to find an estimate of $\ddot{\alpha}$, which is the point on the $(0, 1)$ interval where $\ddot{r}(\alpha)$ is minimal, and use that to fit $\dot{\beta}_{\ddot{\alpha}}$. In Section 6.2, we demonstrate by simulation that $\dot{\alpha}$ and $\ddot{\alpha}$ are indeed very good approximations of the best mixing ratios that minimize $R(\dot{\beta}_\alpha)$ and $R(\ddot{\beta}_\alpha)$ respectively. Moreover, we find it very practical to estimate $\dot{\alpha}$ and $\ddot{\alpha}$ from the data on hand using the suggested formulas, resulting in substantial and significant improvement in the predictions.

4 Application to large-scale Deep Models

In many state-of-the-art deep models, including models from the domain of image processing, the very last layer is linear with relatively low dimension and uses squared loss. We suggest to improve predictive performance of these kind of regression deep models by applying the mixed-loss-function methodology, that was proven to be effective in linear models. The main idea is to view the last layer of the model as an isolated linear model with random features, similar to the Conjugate Kernel method of Hu and Huang [2021]. According to the results of the previous sections, this linear model can be improved by taking advantage of the unlabeled data, and a suitable mixing ratio can be found. The suggested method is described in the following algorithm.

1. Train the network over the labeled data to achieve a supervised model \hat{f} .
2. Forward-feed the model \hat{f} with the labeled and unlabeled data, **up to the last layer**, resulting in a new semi-supervised data set $(X^{\hat{f}}, Y, Z^{\hat{f}})$.
3. Fit a supervised linear model, \hat{w} , and estimate the noise and signal as follows:

$$\hat{\tau}^2 = \hat{w}^T \Sigma \hat{w} / \text{tr}(\Sigma) ; \hat{\sigma}^2 = \widehat{Var}(Y) - \hat{\tau}^2 \text{tr}(\Sigma),$$

where Σ is empirical covariance matrix of $Z^{\hat{f}}$.

4. Use the unlabeled data $Z^{\hat{f}}$ to evaluate the unsupervised statistics, and calculate an estimate $\hat{\alpha}$ of α^* , the desired mixing ratio, according to equation (10).

5. Train the model $\check{f}_{\hat{\alpha}}$ on the original semi-supervised data set $T = (X, Y, Z)$, starting from the model \hat{f} and training a few more epochs with the mixed loss function:

$$L^M = (1 - \hat{\alpha})\hat{L} + \hat{\alpha}\check{L},$$

where \hat{L} is the regular squared loss over the labeled data, and \check{L} is equivalent to the semi-supervised loss in (3), and can be written as follows:

$$\check{L} = \frac{1}{m} \sum_{i=1}^m (f(z_i) - \bar{Y})^2 - 2\widehat{\text{Cov}}(f(X), Y).$$

The described methodology has the advantage that the unlabeled data are used only in the last few epochs, thus saving computational resources during training. Moreover, it can be implemented in a stochastic manner by using mixed batches of labeled and unlabeled data. We note that in Step (5) the whole net’s parameters are being updated during the training with the semi-supervised loss.

4.1 Implementation on CelebA DataSet

We demonstrate the above algorithm in the task of landmark localization in CelebA facial images data set (Liu et al. [2018]), containing $2 \cdot 10^5$ images. We train the same Convolution NN (CNN) as in Simchoni and Rosset [2021], on a small fraction of the data designated as a labeled training set with additional 10 semi-supervised epochs where the rest of the samples are used as unlabeled set. The authors in Simchoni and Rosset [2021] reported an MSE of **1.68** in the Nose-X prediction task with a training set of size $1.6 \cdot 10^5$ labeled images. For comparison, in our experiment, with only $2 \cdot 10^4$ labeled data, we obtain an MSE of **2.46** in a supervised learning and an MSE as low as **1.71** in the semi-supervised learning process. In order to achieve valid inference, we split the whole data into ten disjoint test folds while the unlabeled set is fixed and contains the whole data-set. For each test fold, the labeled training set is taken to be the following (or the first) n images, where n is the size of the training set. In figure 1, we compare between the supervised model and the semi-supervised, for different values of n . The majority of the hyper-parameters are fixed for any value of n , but the size of the batches and number of supervised epochs vary with the training size n to ensure the stability of the supervised learning. We present two learning curves of mean

MSE, one for each learning method, and also one curve for the mean difference in MSE between the two methods, along with its 95% confidence interval. In the results, we can see that our suggested SSL method delivers substantial and significant improvement over the standard supervised learning, and that the magnitude of the difference increases as n decreases.




Figure 1: Mean prediction errors of supervised and semi-supervised CNN on CelebA dataset, for various values of labeled training set size n , out of $2 \cdot 10^5$ total images (see text for discussion).

5 Semi-supervised Interpolators

In this section we study a similar methodology for SSL in the over-parameterized regime, focusing on *interpolators* — estimators that achieve zero training error, including deep neural networks (DNN). This type of model has attracted growing attention in recent years in the field of machine learning. This is mainly because state-of-the-art DNN appear to be models of this type, as well as the fact that recently, a double-descent curve of the out-of-sample error as a function of the number of parameters was observed [Belkin et al., 2019, Geiger et al., 2019, Hastie et al., 2022]. In the last paper, the authors analyze the minimum l_2 norm interpolation in high-dimensional least squares regression, and show that double descent can occur in this model under a variety of settings.

The basic principle in this type of models is to use a large number of parameters $p \gg n$, and to interpolate the training data, which means that $\hat{f}(X) = Y$, while using an appropriate criterion to choose the "best" interpolator, typically based on norm minimization. Consider a learning model with parameters $w \in \mathbb{R}^p$ and output function f_w , the minimal-norm-solution \hat{w} can be written as:

$$\hat{w} = \underset{w \in \mathbb{R}^p}{\operatorname{argmin}} \{ \|w\|, \text{ s.t. } f_w(X) = Y \},$$

for some norm $\|\cdot\|$. In practice, deep learning models are fitted by first-order methods such as gradient descent, which, when initialized appropriately, generally converges to an interpolator \hat{w} with a nearly minimal distance from the initial point (see Oymak and Soltanolkotabi [2019]). Moreover, as shown by Vardi and Shamir [2021], while the implicit regularization of a ReLU neuron is not expressible by w , it is approximately the l_2 norm. These properties ensure that models trained using first-order methods can achieve reliable learning in modern over-parameterized regimes. For our purposes, these characteristics suggest that a semi-supervised method that improves the minimal l_2 norm interpolator could also be advantageous in deep learning models.

The main question we would like to address, is how to use unlabeled data to improve the performance of interpolation models. We address this question by developing a methodology for the standard linear interpolator model, supported by theoretical results and simulation. In the sequel, we also demonstrate how this methodology can be implemented in more complex interpolation models to improve performance. As an analogy to the variance penalty and mixing approach from the previous sections, the main idea we suggest here, is to use the unlabeled data to fit an interpolator that minimizes (or at least reduces) the variance of predictions, and mix it with the minimum-norm interpolator in order to achieve an interpolating model with better predictive performance.

5.1 The Linear Interpolator model

In the special case of fitting a linear predictor, $\hat{f}(x) = \hat{w}^T x$, the minimum-norm interpolator takes the following explicit form:

$$\hat{w} = X^T (X X^T)^{-1} Y.$$

Assuming also that $\text{Var}(x) = \Sigma$ is known from the unlabeled data, the semi-supervised estimator suggested in this case is the *minimum-variance interpolator*, which can be written as follows:

$$\begin{aligned}\tilde{w} &= \underset{w \in \mathbb{R}^p}{\text{argmin}} \{w^T \Sigma w, \text{ s.t. } Xw = Y\} \\ &= \Sigma^{-1} X^T (X \Sigma^{-1} X^T)^{-1} Y.\end{aligned}$$

To compare between the estimators \hat{w} and \tilde{w} in terms of mean prediction error, we require the following distributional assumption:

Assumption 5. *The distribution P_x satisfies that $(XX^T)^{-1}$ and $(X\Sigma X^T)^{-1}$ exist with probability 1, and also that their expectations over X exist.*

We also assume the scenario of a true linear model $f(x) = w^T x$, scaled covariates $\mathbb{E}_{x_0}[x_0] = \mathbf{0}$, and random coefficients vector w , such that $\mathbb{E}[w] = \mathbf{0}$ and $\mathbb{E}[ww^T] = \tau^2 I_p$. Analyzing the prediction error of \hat{w} and \tilde{w} in this scenario, we are able to write explicit expressions for $R(\hat{w})$ and $R(\tilde{w})$ as follows:

$$\begin{aligned}R(\hat{w}) &= \tau^2 \text{tr} (\Sigma - \Sigma \mathbb{E}_X [X^T (XX^T)^{-1} X]) \\ &\quad + \sigma^2 \text{tr} (\Sigma \mathbb{E}_X [X^T (XX^T)^{-2} X]) := \tau^2 b_l + \sigma^2 v_l, \\ R(\tilde{w}) &= \tau^2 \text{tr} (\Sigma - \mathbb{E}_X [X^T (X\Sigma^{-1} X^T)^{-1} X]) \\ &\quad + \sigma^2 \text{tr} (\mathbb{E}_X [(X\Sigma^{-1} X^T)^{-1}]) := \tau^2 b_u + \sigma^2 v_u.\end{aligned}$$

The term b_l (v_l) is associated with the bias (variance) component of $R(\hat{w})$. In the same manner, the terms b_u and v_u are associated with $R(\tilde{w})$. The following proposition describes the relations between these quantities:

Proposition 5.1. *For any distribution P_X satisfying Assumption 5, the following hold: $b_l \leq b_u$ **and** $v_l \geq v_u$.*

The proof of Proposition 5.1 is provided in A.11. We note that the equality holds in the edge case where $\hat{w} = \tilde{w}$ for any given realization of (w, X, Y) , which occurs when $\Sigma = \lambda I_p$, for some $\lambda > 0$. In general, Proposition 5.1 exposes a bias-variance trade-off between the two estimators, concluding that \tilde{w} outperforms \hat{w} if and only if the signal-to-noise ratio, τ^2/σ^2 (or SNR), is smaller than $(v_l - v_u)/(b_u - b_l)$. However, as shown in previous sections,

mixing between the two estimators might be effective in improving predictive performance under any SNR conditions. By analogy to $\hat{\beta}_\alpha$, let us denote the linear-mixed estimator in this case by $\dot{w}_\alpha = (1 - \alpha)\hat{w} + \alpha\tilde{w}$, for some mixing ratio $\alpha \in [0, 1]$. Due to the convexity of the interpolation set, it is clear that \dot{w}_α is also an interpolator. Analyzing the prediction error of \dot{w}_α , we find that $R(\dot{w}_\alpha)$ as a function of α , can be written as follows:

$$R(\dot{w}_\alpha) = R(\hat{w}) + \alpha^2 [\tau^2(b_u - b_l) + \sigma^2(v_l - v_u)] - 2\alpha\sigma^2(v_l - v_u).$$

Analysis of the above formula leads us to the following theorem regarding the characteristics of \dot{w}_α :

Theorem 5.1. *For any distribution P_X satisfying Assumption 5, and given parameters (σ^2, τ^2) , the unique global minimum of $R(\dot{w}_\alpha)$, denoted by α^* , is in the open interval $(0, 1)$. Moreover, α^* admits the following explicit formula:*

$$\alpha^* = \frac{\sigma^2(v_l - v_u)}{\tau^2(b_u - b_l) + \sigma^2(v_l - v_u)},$$

and the minimum value of $R(\dot{w}_\alpha)$ denoted by $R(\dot{w}_{\alpha^*})$, admits the following explicit formula:

$$R(\dot{w}_{\alpha^*}) = \tau^2 b_l + \sigma^2 v_l - \frac{\sigma^4(v_l - v_u)^2}{\tau^2(b_u - b_l) + \sigma^2(v_l - v_u)} < R(\hat{w}).$$

The proof of Theorem 5.1 is provided in A.12. From Theorem 5.1, we conclude that \dot{w}_α at α^* , denoted by \dot{w}_{α^*} is a better estimator than both \hat{w} and \tilde{w} , and that α^* increases with the noise, giving more weight to the unlabeled data (with the knowledge of Σ). However, it is necessary to estimate α^* from the data on hand. Noting that the quantities b_l, b_u, v_l, v_u can be estimated precisely with the unlabeled data, it is only left to estimate σ^2 and τ^2 from the labeled data to define an estimator of α^* . We suggest a methodology for estimating σ^2 and τ^2 , and in the sequel we show that it is practical to use this methodology in order to define the empirically-tuned mixed interpolator, $\dot{w}_{\hat{\alpha}}$, which is uniformly better than the standard minimum-norm interpolator. First, we preset an unbiased estimator of the noise σ^2 , based on the characteristics of the minimum-norm interpolator model, for the case where τ^2 is known.

Proposition 5.2. *The following estimator is an unbiased estimator of σ^2 :*

$$\hat{\sigma}^2 = \frac{Y^T (XX^T)^{-2} Y - \tau^2 \text{tr}((XX^T)^{-1})}{\text{tr}((XX^T)^{-2})}. \quad (15)$$

The proof of Theorem 5.2 is provided in A.13. We suggest using this unbiased estimator in order to define the empirically-tuned mixed interpolator $\hat{w}_{\hat{\alpha}(\tau)}$ in the known τ scenario. In the general scenario where τ^2 is unknown, it must be estimated together with σ^2 . In this case, we suggest to perform the following iterative calculation:

$$\begin{aligned}\hat{\tau}_{(0)}^2 &= \hat{w}^T \Sigma \hat{w} / \text{tr}(\Sigma), \\ \hat{\sigma}_{(t+1)}^2 &= \max \left\{ \frac{Y^T (X X^T)^{-2} Y - \hat{\tau}_{(t)}^2 \text{tr}((X X^T)^{-1})}{\text{tr}((X X^T)^{-2})}, 0 \right\}, \\ \hat{\tau}_{(t+1)}^2 &= \max \left\{ \left(\frac{\sum y_i^2}{n} - \hat{\sigma}_{(t+1)}^2 \right) / \text{tr}(\Sigma), 0 \right\}.\end{aligned}$$

We note that typically, it only requires a few iterations in order to converge to stable estimates $(\hat{\sigma}^2, \hat{\tau}^2)$, which can be used to calculate the estimator $\hat{\alpha}$ of α^* . As far as we know, this methodology is novel and can be implemented in other cases where the estimation of the noise and signal of an over-parameterized underlying model is required.

5.2 Asymptotic usefulness of the mixed Interpolator

Similarly to the OLS case, the relative prediction error between the oracle-tuned estimator, \hat{w}_{α^*} , and the standard supervised estimator, \hat{w} , can be written as follows:

$$\eta = \frac{R(\hat{w}_{\alpha^*})}{R(\hat{w})} = 1 - \frac{\sigma^4 (v_l - v_u)^2}{(\tau^2 (b_u - b_l) + \sigma^2 (v_l - v_u)) (\tau^2 b_l + \sigma^2 v_l)}.$$

The above expression reveals that the benefit of using the semi-supervised estimator is non-negligible as long as the differences $v_l - v_u$ and $b_u - b_l$ are not negligible compared to the quantities b_u , b_l and v_l . Detailed analysis of the characteristics of the quantities (v_l, v_u, b_u, b_l) as well as the estimator $\hat{\alpha}$, under different settings, especially regarding P_x , is beyond the scope of this paper. For the sake of brevity, and to highlight the advantage of the semi-supervised interpolator over the supervised one, we consider the following distributional assumption.

Assumption 6. *The distribution P_x satisfies $x \sim MN(\mathbf{0}, \Sigma)$, where Σ is (a sequence of) $p \times p$ diagonal matrix with entrees $\Sigma_{jj} = c_1^2(n)$ for $1 \leq j \leq \tilde{p}$ and $\Sigma_{jj} = c_2^2(n)$ for $\tilde{p} < j \leq p$.*

Moreover, the asymptotic setup satisfies the following:

$$p/n \rightarrow \gamma \in (1, \infty) ; \tilde{p}/n \rightarrow \tilde{\gamma} \in (1, \gamma)$$

$$\tilde{p}c_1^2(n) = \sum_{j=1}^{\tilde{p}} \Sigma_{jj} \rightarrow c^2 ; (p - \tilde{p})c_2^2(n) = \sum_{j=\tilde{p}}^p \Sigma_{jj} \rightarrow 0,$$

as $n, p, \tilde{p} \rightarrow \infty$, for some constant $c \in \mathbb{R}$.

The setting of Assumption 6 highlights the benefit of using the available unlabeled data, which leads to the inference of the unbalanced structure of Σ and the subsequent adjustment of the linear interpolator. Moreover, in this setting, we can find explicit expressions for the limiting values of v_l , v_u , b_l and b_u , and also for α^* and η . The following theorem summarizes the asymptotic characteristics of the mix semi-supervised interpolator, and shows that η is strictly smaller than 1 also when the sample size grows to infinity.

Theorem 5.2. *Under the conditions of Assumption 6, as $n \rightarrow \infty$, the following limiting values exist:*

$$v_l \rightarrow \frac{1}{\tilde{\gamma} - 1} ; b_l \rightarrow c^2 \left(1 - \frac{1}{\tilde{\gamma}}\right) ; v_u \rightarrow \frac{1}{\gamma - 1} ; b_u \rightarrow c^2 \left(1 - \frac{1}{\gamma}\right),$$

$$\eta \rightarrow \eta_\infty = 1 - \frac{(\gamma - \tilde{\gamma})\sigma^4}{(\gamma - 1)^2(\tilde{\gamma} - 1)^2 \left[\frac{\tau^2 c^2}{\gamma \tilde{\gamma}} + \frac{\sigma^2}{(\gamma - 1)(\tilde{\gamma} - 1)} \right] \left[\frac{\tau^2 c^2(\tilde{\gamma} - 1)}{\tilde{\gamma}} + \frac{\sigma^2}{\tilde{\gamma} - 1} \right]},$$

$$\alpha^* \rightarrow \alpha_\infty^* = \frac{\sigma^2}{(\gamma - 1)(\tilde{\gamma} - 1) \left[\frac{\tau^2 c^2}{\gamma \tilde{\gamma}} + \frac{\sigma^2}{(\gamma - 1)(\tilde{\gamma} - 1)} \right]}.$$

The proof of Theorem 5.2 is provided in Section A.14, and from it we conclude that the prediction error of \hat{w}_{α^*} is strictly lower than the one of \hat{w} as long as $\tilde{\gamma} < \gamma$. Moreover, we can see that η increases with $\tilde{\gamma}$ and equal to 1 when $\tilde{\gamma} = \gamma$. On the other hand, η decreases with γ and approaches 0 when $\tilde{\gamma}$ approaches 1. While we are not able to exactly identify the characteristics of the estimates $(\hat{\sigma}^2, \hat{\tau}^2)$, the typical behavior we observed in our simulations is that if τ^2 is known, the variance of $\hat{\sigma}^2$ is of order $1/n$. Moreover, if σ^2 is known, the variance of $\hat{\tau}^2$ vanishes as $n \rightarrow \infty$, since $\text{tr}(\Sigma^2)$ approaches zero. Therefore, we expect the empirically tuned estimators, $\hat{w}_{\hat{\alpha}}$ and $\hat{w}_{\hat{\alpha}(\tau)}$, to perform similarly to \hat{w}_{α^*} . In Section 6.3 we demonstrate that this is indeed the typical behavior.

5.3 Application to Random Fourier Features on real data

We demonstrate the applicability of the above methodology, on a common model in the domain of over-parametrized machine learning, called Random Fourier Features (RFF, Rahimi and Recht [2007]). The RFF model can be viewed as a two-layer neural network, with the first layer being fixed (generated once pre-training), with non-linear activation. The second (output) layer is essentially a linear interpolator. In particular, we adopt the following formulation for the output function of the RFF model:

$$f_{\hat{w}}(x) = \hat{w}^T \varphi(Cx) ; C \in \mathbb{R}^{H \times p}, H > n ; C_{ij} \sim N(0, 1),$$

where φ is a non-linear function acting component-wise. Thus, we can fit a supervised or semi-supervised linear interpolator over the semi-supervised training set:

$$T = (\varphi(XC^T), Y, \varphi(ZC^T)).$$

We empirically study the predictive performance of the suggested mixed semi-supervised linear interpolators as well as the OLS estimators from Section 2, on a sub-dataset of the Netflix Prize data (Bennett et al. [2007]), available at Kaggle. [2005]¹. From the full dataset, we extracted the $N = 12,931$ users who rated the movie "Miss Congeniality". The outcome y is the rating of that movie by a specific user, and the covariates vector x is the user's ratings of $p = 40$ other movies. In order to achieve valid inference, we randomly split the data into 20 disjoint folds, and each of them is split into train and test sets. This is done for different sizes of training sets n , while the unlabeled data set $Z \in \mathbb{R}^{N \times p}$ is fixed and contains the rating data of all N users, without the outcome column. For each of the training sets, we fit the OLS estimators $(\hat{\beta}, \check{\beta}, \check{\beta}_{\check{\alpha}}, \check{\beta}_{\check{\alpha}})$. We also generate an RFF model with $H = 5n$ nodes in the hidden layer, and an activation function $\varphi : \mathbb{R} \rightarrow \mathbb{R}^3$ as follows: $\varphi(c) = (\tanh(c), \text{Sigmoid}(c), \text{ELU}(c))^T$, resulting in a second layer of size $15n$. We also apply standard scaling among $\varphi(ZC^T)$, and then fit the interpolators $(\hat{w}, \tilde{w}, \dot{w}_{\check{\alpha}})$ in the same manner as described in Section 6.3. The mean squared error over the test set of the relevant fold is evaluated for every fitted estimator and averaged over the 20 folds. The outcome is seven curves describing the prediction error, changing with n for each of the estimators. We also evaluate the pairwise variances of the differences between any two

¹<https://www.kaggle.com/datasets/netflix-inc/netflix-prize-data>

estimators, for any value of n , to find the minimum difference required for significance at any desired level.

In the results (Figure 2), we can see that the OLS estimators behave with the same characteristics as in the simulated data of Section 6.1, with $\hat{\beta}_{\hat{\alpha}}$ outperforming the others, supporting the analysis of Section 2. Regarding the RFF model, we can see that compared to the OLS model, the RFF model is less sensitive to the size of the training set. More importantly, the mixed-RFF interpolator $\hat{w}_{\hat{\alpha}}$ is very effective in reducing prediction error, and outperforms all other models. Any difference greater than 0.04 is significant at level 0.05.




Figure 2: Mean prediction errors of supervised and semi-supervised OLS and RFF on Netflix data, for various values of n (see text for discussion).

5.4 Application to Fully-Connected Interpolating Neural Networks

We present a methodology for adapting the idea of mixed interpolators to fully connected deep neural networks. The main idea is to mix the predictions between two neural networks (models) as follows:

1. Supervised-trained interpolating model, denoted by \hat{f} , which is related to the minimum norm interpolator.
2. Semi-supervised-trained model, \tilde{f} , that aims to reduce the following variance term:

$\mathbb{E}_{x_0} \left(\tilde{f}(x_0) - \bar{Y} \right)^2$, while approximately maintains the interpolation constraint over the labeled data.

The supervised model \hat{f} can be trained by applying the standard gradient descent method with squared loss over the labeled data. The semi-supervised model \tilde{f} , can be trained by solving the following unconstrained optimization problem:

$$\begin{aligned} \max_{\lambda \in \mathbb{R}^n} \min_w \{J(\lambda, w)\} = \\ \max_{\lambda \in \mathbb{R}^n} \min_w \left\{ \frac{1}{m} \sum_{i=1}^m \left(\tilde{f}(z_i) - \bar{Y} \right)^2 - \lambda^T \left(Y - \tilde{f}(X) \right) \right\}, \end{aligned} \quad (16)$$

where w represents the model's parameters. Training \tilde{f} can be done by a "simple" gradient descent ascent (GDA) method, minimizing J w.r.t w , and maximizing w.r.t λ . However, the following issues need to be addressed in order to implement such an algorithm:

- Starting point. We use the supervised interpolating model \hat{f} as a starting point for training the model \tilde{f} . This is an analogy to moving from the minimum-norm solution towards the minimum-variance solution. More important, this is much more efficient than starting from a random point and converging to such an interpolating solution.
- Stopping criterion. The suggested GDA method might "over-fit" the unlabeled data Z , when applied with a high-capacity model, resulting in poor predictions. Therefore, we define a minimal value of $\frac{1}{m} \sum_{i=1}^m \left(\tilde{f}(z_i) - \bar{Y} \right)^2$, where the GDA algorithm should be stopped upon reaching this value.

The suggested methodology for the *Mixed-Semi-Supervised Deep Interpolating model* is described in the following algorithm.

1. Train the desired neural network over the labeled data to achieve near-interpolating model \hat{f} .
2. Forward-feed the model \hat{f} with the labeled and unlabeled data, **up to last layer**, and extract a new semi-supervised data set $\mathcal{T} = (X^{\hat{f}}, Y, Z^{\hat{f}})$ of high dimension $H > n$, with scaled covariates.

- Based on \mathcal{T} , fit the supervised and semi-supervised linear interpolating models (\hat{w} and \tilde{w}), and calculate an estimate of α^* (denoted by $\hat{\alpha}$), according to formula (25), combined with the following estimates of the noise and the signal:

$$\hat{\tau}^2 = \hat{w}^T \Sigma \hat{w} / \text{tr}(\Sigma) \quad ; \quad \hat{\sigma}^2 = \widehat{\text{Var}}(Y) - \hat{\tau}^2 \text{tr}(\Sigma),$$

where Σ is empirical covariance matrix of $Z^{\hat{f}}$.

- Train the semi-supervised model, \tilde{f} , by applying the GDA method with the objective J (Equation 16), starting from \hat{f} , and converging to an interpolation model with a variance term as low as $\tilde{w}^T \Sigma \tilde{w} + \bar{Y}^2$.
- The mixed-semi-supervised deep model, denoted by $\dot{f}_{\hat{\alpha}}$, is defined follows:

$$\dot{f}_{\hat{\alpha}}(x) = \hat{\alpha} \tilde{f}(x) + (1 - \hat{\alpha}) \hat{f}(x).$$

We empirically study the predictive performance of the suggested Mixed Semi Supervised Deep Interpolating model on the same Netflix data as in Section 5.3, for a smaller range of training sizes n . For any training sample, we train the model $\dot{f}_{\hat{\alpha}}$ as described above. We use a fully connected neural network with two hidden layers, containing $H_1 = 5n$ and $H_2 = 15n$ nodes, respectively, and Relu activation. The mean square error over the unseen labeled in each fold is evaluated for both \hat{f} and $\dot{f}_{\hat{\alpha}}$, and averaged over the 20 folds.

In Figure 3, we present the results and compare them with the performance curves of the mixed OLS and RFF models. We can see that the deep model is better than the other two models in both the supervised and semi-supervised settings. More importantly, the mixed-semi-supervised deep model, $\dot{f}_{\hat{\alpha}}$, improves over the supervised model \hat{f} , and delivers the best performance among all models. Any difference greater than 0.05 is significant at level 0.05.

6 Results of synthetic data simulations

We conduct an empirical study regarding the predictive performance of the various estimators presented throughout the paper. The notation of all estimators and their corresponding definitions are summarized in Table 1.

Notation	Definition
Overview	
$\hat{\beta}$	The standard supervised estimator.
$\check{\beta}$	The semi-supervised estimator.
β^D	The adaptive estimator that chooses between $\hat{\beta}$ and $\check{\beta}$.
$\dot{\beta}_\alpha$	The linear mixed estimator $\alpha\check{\beta} + (1 - \alpha)\hat{\beta}$ with mixing ratio α .
$\ddot{\beta}_\alpha$	The estimator the minimizes the mixed loss function $\alpha\check{L} + (1 - \alpha)\hat{L}$.
OLS	
$\dot{\beta}_{\alpha^*}$	The linear mixed estimator $\dot{\beta}_\alpha$ at the oracle best mixing ratio $\alpha^* = \text{agrmin}\{R(\dot{\beta}_\alpha)\}$.
$\ddot{\beta}_{\alpha^{**}}$	The mixed estimator $\ddot{\beta}_\alpha$ at the oracle best mixing ratio $\alpha^{**} = \text{agrmin}\{R(\ddot{\beta}_\alpha)\}$.
$\dot{\beta}_{\hat{\alpha}}$	The linear mixed estimator $\dot{\beta}_\alpha$ at the estimated best mixing ratio $\hat{\alpha}$.
$\dot{\beta}_{\hat{\alpha}(\tau)}$	The linear mixed estimator $\dot{\beta}_\alpha$ at the estimated best mixing ratio, considering known τ .
$\ddot{\beta}_{\hat{\alpha}}$	The mixed estimator $\ddot{\beta}_\alpha$ at the estimated best mixing ratio for $\dot{\beta}_\alpha, \alpha^*$.
$\ddot{\beta}_{\tilde{\alpha}}$	The mixed estimator $\ddot{\beta}_\alpha$ at $\tilde{\alpha}$ the best mixing ratio obtained by the grid search.
GLM	
$\dot{\beta}_{\hat{\alpha}}$	The estimator $\dot{\beta}_\alpha$ at the oracle approximate best mixing ratio $\hat{\alpha} = \text{agrmin}\{\dot{r}(\alpha)\}$
$\ddot{\beta}_{\tilde{\alpha}}$	The estimator $\ddot{\beta}_\alpha$ at the oracle approximate best mixing ratio $\tilde{\alpha} = \text{agrmin}\{\ddot{r}(\alpha)\}$.
$\dot{\beta}_{\hat{\alpha}}$	The estimator $\dot{\beta}_\alpha$ at the estimated value of $\hat{\alpha}$.
$\ddot{\beta}_{\hat{\alpha}}$	The estimator $\ddot{\beta}_\alpha$ at the estimated value of $\hat{\alpha}$.
$\dot{\beta}_{\tilde{\alpha}}$	The estimator $\dot{\beta}_\alpha$ at the estimated value of $\tilde{\alpha}$ obtained by the grid search.
$p > n$	
\hat{w}	The standard supervised minimum l_2 interpolator.
\tilde{w}	The semi-supervised minimum-variance interpolator.
\dot{w}_{α^*}	The linear mixed interpolator $\dot{w}_\alpha = \alpha\tilde{w} + (1 - \alpha)\hat{w}$ at the oracle best mixing ratio α^*
$\dot{w}_{\hat{\alpha}}$	The linear mixed estimator at the estimated best mixing ratio $\hat{\alpha}$.
$\dot{w}_{\tilde{\alpha}}$	The linear mixed estimator at the estimated best mixing ratio, considering known τ .

Table 1: Summary of estimator notations




Figure 3: Mean prediction errors of supervised and semi-supervised OLS, RFF, and fully connected NN on Netflix data, for various values of n (see text for discussion).

6.1 Simulations for OLS model

We empirically study the predictive performance of the OLS model and all its SSL variants under discussion, in two different generating models, by two-step simulations as follows:

- Unsupervised estimation.** Evaluating relevant statistics that depend only on the distribution of the covariates, such as v_l , v_u , H , etc., and consequently evaluating the oracle values of the best mixing ratios for $\hat{\beta}_\alpha$ (denoted by α^*) and $\check{\beta}_\alpha$ (denoted by α^{**}), under any combination of noise and signal in our study, as if they were known. This is done based on a large fixed data-set \mathcal{Z} of $m = 5 \cdot 10^4$ unlabeled observations $x \sim MN(\mathbf{0}_p, \Sigma)$, where $p = 25$, and Σ is block-diagonal, containing five blocks such that all variables in each block have pairwise correlation $\rho = 0.9$.
- Supervised simulation.** Generating $K = 5,000$ random training sets of $n = 50$ labeled observations (X, Y) where X is randomly sampled from \mathcal{Z} , and $Y = E[Y|X] + \epsilon_n$, where $\epsilon_n \sim MN(\mathbf{0}_n, \sigma^2 I_n)$, for various values of σ^2 . For each of the training sets, combined with the statistics from the unsupervised estimation, we fit the supervised and semi-supervised OLS estimators $(\hat{\beta}, \check{\beta}, \beta^D)$. as well as the oracle estimators, $\hat{\beta}_{\alpha^*}$ and $\check{\beta}_{\alpha^{**}}$. We then estimate from the data the best mixing ratios for $\hat{\beta}_\alpha$ and $\check{\beta}_\alpha$, and fit the mixed semi-supervised estimators $(\dot{\beta}_{\hat{\alpha}}, \ddot{\beta}_{\hat{\alpha}}, \ddot{\beta}_{\check{\alpha}})$. The mean prediction error over the data-set \mathcal{Z} is evaluated for every fitted estimator and averaged over the K training samples. The outcome is eight curves that describe the prediction error, R ,

changing with σ^2 for each of the estimators. We also evaluate the pairwise variances of the differences between any two estimators, for any value of σ^2 , in order to provide a valid statement on the significance of the results.

The two generating models for the mean function $f(x) = \mathbb{E}[y|x]$ are as follows:

1. *Constant- β* . $f(x) = \beta \sum_{j=1}^p x_j$, $\beta = 1.5$.
2. *Random- β* . $f(x) = x^T \beta$, where $\beta \sim MN(\mathbf{0}, I_p)$ is generated with each training set (X, Y) .

In the results (Figure 4 and Figure 5), we can see that the mixed semi-supervised learning is very effective in reducing the mean prediction error. In particular, the estimator $\ddot{\beta}_{\hat{\alpha}}$ outperforms all other estimators, except for the oracle estimator $\ddot{\beta}_{\alpha^{**}}$, with a benefit that increases with the noise. We can see that the estimator $\ddot{\beta}_{\hat{\alpha}}$ is also very effective despite the fact that it uses the mixing ratio aimed at $\dot{\beta}_{\hat{\alpha}}$. In the constant- β scenario (Figure 4), we can see that the oracle estimators are only slightly better than $\dot{\beta}_{\hat{\alpha}}$ and $\ddot{\beta}_{\hat{\alpha}}$ respectively, meaning that there is not much benefit in knowing the best mixing ratio, rather than estimating it. Moreover, in the random- β scenario (Figure 5), the data-based estimator $\dot{\beta}_{\hat{\alpha}}$ is slightly better than the oracle estimator $\dot{\beta}_{\alpha^*}$, which means that it is actually beneficial to estimate the noise and signal from the training data, rather than knowing them. In both figures, all variances of pairwise mean differences are tiny, hence any difference larger than the marker's size is significant at level 0.05 in a paired t -test.

In Figure 6 we evaluate the prediction error of $\dot{\beta}_{\hat{\alpha}}$, $\dot{\beta}_{\alpha^*}$ and $\dot{\beta}_{\hat{\alpha}(\tau)}$ for various values of n and $p = 0.5n$, while $\text{tr}(\Sigma) = \sigma^2 = 25$, and $\tau^2 = 1$. The outcome is three curves that describe the relative prediction error, $R(\tilde{\beta})/R(\hat{\beta})$ for $\tilde{\beta} \in \{\dot{\beta}_{\hat{\alpha}}, \dot{\beta}_{\alpha^*}, \dot{\beta}_{\hat{\alpha}(\tau)}\}$, changing the size of the training sample. For reference, we mark the theoretical bound of η in this setting, η_{∞} , by a dotted horizontal black line, and the estimated value of η according to the unsupervised estimate with $n = 500$ by a dashed horizontal gray line. In the results, we can see that the relative prediction error of both $\dot{\beta}_{\alpha^*}$ and $\dot{\beta}_{\hat{\alpha}(\tau)}$ converges to the expected limiting value of $1 - 0.5^2$, which supports the theoretical analysis. Interestingly, the prediction error of the most realistic estimator, $\dot{\beta}_{\hat{\alpha}}$, converges to a significantly lower value, supporting the conjecture that the non-vanishing variance of the estimator $\hat{\tau}^2$, leads to a better estimation

of the mixing ratio, rather than knowing the real value of τ . Table 1 provides a summary of the definitions for each estimator.




Figure 4: Mean prediction errors of true linear model with constant- β mechanism, for various values of σ^2 (see text for discussion).




Figure 5: Mean prediction errors of true linear model with random- β mechanism, for various values of σ^2 (see text for discussion).

6.2 Simulations for GLM-ERM

We perform similar experiments as in Section 6.1, but with the ELU function Clevert et al. [2015] for the link function g , as follows:

$$\mathbb{E}[y|x] = \min \left\{ e^{x^T \beta} - 1, \max(0, x^T \beta) \right\} \quad ; \quad \beta = 2 \cdot \mathbf{1}_p,$$




Figure 6: Relative prediction errors of the random- β mechanism for various values of n , with $\gamma = 0.5$, and $\text{tr}(\Sigma) = \sigma^2 = 25$ (see text for discussion).

and we set $p = 10$. We note that this model does not comply with Assumption 4, but the key condition $v_l > v_u$ holds. In this scenario, we perform unsupervised estimation using a large fixed data-set \mathcal{Z} only in order to evaluate the oracle values for $\dot{\alpha}$ and $\ddot{\alpha}$. The supervised simulation works as follows:

We generate $K = 4000$ random training sets of $n = 50$ labeled observations (X, Y) and $m = 5000$ unlabeled observations (Z) , for various values of σ^2 . For each of the training sets, we fit the estimators $\hat{\beta}$ and $\check{\beta}$. Then, by sub-sampling Z , at the fitted vector $\check{\beta}$, we evaluate $\hat{\sigma}^2$ and all other unsupervised statistics in order to obtain estimates of $\dot{\alpha}$ and $\ddot{\alpha}$, denoted by $\hat{\alpha}$ and $\tilde{\alpha}$ respectively, and use them to fit the mixed estimators $(\dot{\beta}_{\hat{\alpha}}, \ddot{\beta}_{\hat{\alpha}}, \dot{\beta}_{\tilde{\alpha}})$. The method for fitting $\dot{\beta}_{\hat{\alpha}}$ and $\dot{\beta}_{\tilde{\alpha}}$ is summarized by the following Newton-Raphson update step, where $\alpha \in \{\hat{\alpha}, \tilde{\alpha}\}$ respectively:

$$\beta_{LM}^{(t+1)} = \beta_{LM}^{(t)} - (S_{\alpha}^{(t)})^{-1} (\alpha \zeta_m^{(t)} + (1 - \alpha) \zeta_n^{(t)}),$$

where,

$$\begin{aligned} S_{\alpha}^{(t)} &= \alpha H_m^{(t)} + (1 - \alpha) H_n^{(t)}, \\ H_m^{(t)} &= \frac{1}{m} Z^T D_m^{(t)} Z ; [D_m^{(t)}]_{ii} = g'(z_i^T \beta_{LM}^{(t)}), \\ H_n^{(t)} &= \frac{1}{m} X^T D_n^{(t)} X ; [D_n^{(t)}]_{ii} = g'(x_i^T \beta_{LM}^{(t)}), \\ \zeta_m^{(t)} &= \frac{1}{m} Z^T g(Z \beta_{LM}^{(t)}) - \bar{Z} \cdot \bar{Y} - \widehat{\text{Cov}}(X, Y) ; \zeta_n^{(t)} = \frac{1}{n} X^T (g(X \beta_{LM}^{(t)}) - Y). \end{aligned}$$




Figure 7: Mean prediction errors of ELU model, for various values of σ^2 (see text for discussion).

We also fit the oracle estimators $\dot{\beta}_{\hat{\alpha}}$ and $\ddot{\beta}_{\check{\alpha}}$, based on the oracle mixing ratios for $\dot{r}(\alpha)$ and $\ddot{r}(\alpha)$ respectively, from the unsupervised simulation. The mean prediction error over the data-set \mathcal{Z} is evaluated for every fitted estimator and averaged over the K training samples. The outcome is eight curves that describe the prediction error R , changing with σ^2 for each of the estimators.

In the results (Figure 7), we can see that mixed semi-supervised learning is effective in reducing the prediction error, and the estimator $\dot{\beta}_{\hat{\alpha}}$ has the best performance (together with the oracle estimator $\ddot{\beta}_{\check{\alpha}}$). Moreover, we can see that the oracle model $\dot{\beta}_{\hat{\alpha}}$ performs a little worse than $\dot{\beta}_{\hat{\alpha}}$. We can also see that $\ddot{\beta}_{\check{\alpha}}$ performs very much like $\dot{\beta}_{\hat{\alpha}}$ despite the fact that it uses the estimate of $\hat{\alpha}$ as the mixing ratio. Any difference greater than the marker size is significant at level 0.05 in a paired t -test.

In Figure 8 we evaluate the mean prediction error of $\dot{\beta}_{\alpha}$ and $\ddot{\beta}_{\alpha}$ for various values of α in the interval $[0, 1]$, specifically for $\sigma^2 = 25$. The outcome is two curves that describe the prediction error, $R(\dot{\beta}_{\alpha})$ and $R(\ddot{\beta}_{\alpha})$, changing with α . For reference, we mark the oracle evaluations of $\hat{\alpha}$ and $\check{\alpha}$ in vertical lines, and the prediction error of the other mixed estimators in horizontal lines. In the results, we can see that $R(\ddot{\beta}_{\alpha}) < R(\dot{\beta}_{\alpha})$ and also that $R(\dot{\beta}_{\hat{\alpha}}) < R(\dot{\beta}_{\alpha})$ for any value of α . More importantly, we can see that both $\hat{\alpha}$ and $\check{\alpha}$ (especially) are very good approximations for determining the best mixing ratios of $\dot{\beta}_{\alpha}$ and $\ddot{\beta}_{\alpha}$ respectively, despite the quadratic approximation and the fact that, by definition, they are the minimizers of the r -quantities instead of the R -quantities. The differences

between $R(\dot{\beta}_{\hat{\alpha}})$ and $R(\dot{\beta}_{\alpha})$ are significant for any value of α , differences between $R(\ddot{\beta}_{\hat{\alpha}})$ and $R(\dot{\beta}_{\alpha})$ are significant for $\alpha \in [0.05, 0.95]$, and the differences between $R(\dot{\beta}_{\hat{\alpha}})$ and $R(\ddot{\beta}_{\alpha})$ are significant for $\alpha \notin (0.35, 0.5)$. In the next Section, we demonstrate how this successful methodology can be implemented to improve deep neural networks.




Figure 8: Mean prediction errors of ELU model, for various values of α , with $\sigma^2 = 25$ (see text for discussion).

6.3 Simulation for linear interpolators

We perform a similar experiment to the one in Section 6.1, with random coefficients, $p = 100$, $n = 50$, and Σ is a diagonal matrix with components $\Sigma_{ii} = 1$ for $i \leq 0.8p$, and $\Sigma_{ii} = 1/n$ for $i > 0.8p$. For each of the training sets, combined with the unsupervised statistics, we fit the interpolators (\hat{w}, \tilde{w}) . We then estimate the noise and the signal according to the described methodology, and fit w^M . We also fit a semi-oracle estimator, $w^{M(\tau)}$, which uses τ^2 as if it were known and only estimates the noise with formula (15), and one oracle estimator, w^{M^*} , using the oracle mixing ratios as if both τ^2 and σ^2 were known. We note that all the models under comparison are interpolating models.

In the results (Figure 9), we can see that w^M outperforms all non-mixed interpolators, suggesting that our mixing methodology is indeed effective in the over-parameterized regime as well. We can also see that there is no substantial benefit in knowing the real noise and signal rather than estimating them. The differences are significant whenever the markers do not overlap.

In Figure 10 we evaluate the relative prediction error of w^M , w^{M^*} and $w^{M(\tau)}$ for various values of n and $p = 2n$, while maintaining the described mechanism for Σ with scaling, such that $\text{tr}(\Sigma) = \sigma^2 = 25$. For reference, we mark the theoretical boundary of η in this setting by a dotted horizontal black line, and the estimated value of η according to the unsupervised estimation with $n = 300$ by a dashed horizontal gray line. In the results, we can see that the relative prediction errors of all three estimators converge to the expected limiting value, supporting the theoretical analysis.




Figure 9: Mean prediction errors of supervised and semi-supervised linear interpolating model, for various values of SNR (see text for discussion).




Figure 10: Relative prediction errors of linear interpolating model Linear interpolating model for various values of n , with $\gamma = 2$ (see text for discussion).

7 Discussion

This work provides new insight about the usefulness of unlabeled data in improving prediction models, with a focus on the class of GLM. We show theoretically that the unlabeled data can always be integrated in the learning process in a manner that ensures improvement in prediction. Furthermore, the optimal portion of integration or mixing (denoted by α) can be well estimated in practice by leveraging the unlabeled data on hand, leading to substantial improvement. This kind of observation is very different from the conveyed message in many previous works about the effectiveness of SSL. Due to space limits, we study and analyze the methodology of mixed semi-supervised GLM-and-interpolators under a specific set of assumptions and conditions that we found most important and interesting. However, the suggested methodology can also be applied under a variety of scenarios such as non-constant conditional variance, different predictive quality criteria for model evaluation, and different mixing mechanisms. Taking a step back, the general idea behind the studied methodology can be described as follows:

1. Define the mixed-predictor f_α^M that combines the unlabeled data during the learning process, with a portion that determined by α .
2. Analyze the quantity of interest $R(f_\alpha^M)$ according to the assumptions on the true model, and derive an approximate or heuristic formula of it, denoted by $r(\alpha)$.
3. Identify and calculate an estimation $\hat{\alpha}$ of the optimal portion α^* , where $r(\alpha)$ achieves its minimum, using estimates of the relevant statistics from the available labeled and unlabeled data.
4. Fit the model $f_{\hat{\alpha}}^M$.

As another illustration of the described approach, and a possible extension to Section 3, let us consider the common assumption in GLM theory that $\text{Var}(y|x) = g'(x^T\beta)\sigma^2$, where σ^2 is the dispersion parameter (naturally equal to 1). Moreover, let us assume that the predictive quality criterion is the out-of-sample squared error, $\mathbb{E}_{T,x_0,y_0} \left[(f_\alpha^M(x_0) - y_0)^2 \right]$, and not the Canonical loss used in the learning process of $\hat{\beta}$ and $\check{\beta}$, for the purpose of convexity. Finally, let us be bold and choose to use the following mixed predictor:

$$f_\alpha^M(x) = (1 - \alpha)g(x^T\hat{\beta}) + \alpha g(x^T\check{\beta}).$$

From an analysis of the quadratic approximation of $R(f_\alpha^M)$ in this setting, we can find that:

$$\alpha^* = \frac{\sigma^2(v_l - v_u)}{B(\check{\beta}) + \sigma^2(v_l - v_u)},$$

where $B(\check{\beta})$ is the same quantity as in (14), and we can write the other statistics as follows:

$$\begin{aligned} v_u &= \frac{n-1}{n} \text{tr}(H_2 H^{-1}) ; \quad v_l = \text{tr}(H_2 \mathbb{E}_X [(X^T D X)^{-1}]), \\ H &= \mathbb{E}_X [X^T D X] ; \quad H_2 = \mathbb{E}_X [X^T D^2 X]. \end{aligned}$$

The result by Groves and Rothenberg [1969] implies that the matrix $\mathbb{E} [(X^T D X)^{-1}] - H^{-1}$ is positive semi-definite. Thus we have: $v_l > v_u$, and we conclude that $\alpha^* \in (0, 1)$ in this setting, and mixed-SSL is effective in improving predictions. This example demonstrates the flexibility of the suggested methodology to fit different scenarios, while following Steps 1 to 4 as mentioned above.

Based on the theoretical results in this paper, combined with the illustrated flexibility above, we believe that the general approach of mixed-SSL can be relevant for improving many real-life predictive models. The transition from GLM to more complex models, however, is not a minor matter, especially in the context of obtaining rigorous theoretical insights. While this is not the main focus of this work, we find it important to demonstrate the applicability of our method to improve modern deep models in Sections 4,5. This was done by looking at the last layer of the model as a "remote" linear model with some best mixing ratio that can be estimated with the formulas in this paper. In our experiments, this estimation also turned out to be a useful heuristic measure for the desired mixing ratio of the whole deep model. We note that this heuristic approach was inspired by the fact that in Section 2 the estimator $\beta_{L\bar{M}}$ was very effective, although it used the best estimated mixing ratio of β^M , as a heuristic measure. Nevertheless, more detailed theoretical analysis of the characteristics of mixed-SSDL is an important topic for future studies. In particular, it is interesting to study different adjustments of unlabeled data integration mechanisms to different classes of models, and identify how the performance changes with the portion of integration. We believe that both the theoretical and empirical results in this paper shed some light on this domain of research, and motivate follow-up work.

A Proofs

A.1 Proof of Theorem 2.1

Using the explicit expressions of $\hat{\beta}$ and $\check{\beta}$ in (6), we can write the integrated risk $R(\dot{\beta}_\alpha)$ as a function of α , as follows:

$$\begin{aligned}
R(\dot{\beta}_\alpha) &= \alpha^2 B_u + \alpha^2 \sigma^2 v_u + (1 - \alpha)^2 \sigma^2 v_l \\
&\quad + 2\alpha(1 - \alpha) \mathbb{E}_{X, x_0} \left[\text{Cov} \left(x_0^T \hat{\beta}, x_0^T \check{\beta} \mid X, x_0 \right) \right] \\
&= \alpha^2 B(\check{\beta}) + \alpha^2 \sigma^2 v_u + (1 - \alpha)^2 \sigma^2 v_l \\
&\quad + \frac{2}{n} \alpha(1 - \alpha) \text{tr} \left(\mathbb{E}_X \left[(X^T X)^{-1} X^T \text{Cov} (Y, X^T Y - n\bar{X} \cdot \bar{Y} \mid X, x_0) \right] \right) \\
&= \alpha^2 B(\check{\beta}) + \alpha^2 \sigma^2 v_u + (1 - \alpha)^2 \sigma^2 v_l + 2\alpha(1 - \alpha) \sigma^2 v_u \\
&= \alpha^2 \left(B(\check{\beta}) + \sigma^2 (v_l - v_u) \right) - 2\alpha \sigma^2 (v_l - v_u) + \sigma^2 v_l. \tag{17}
\end{aligned}$$

Calculating the first and second derivatives with respect to α , we get the following.

$$\begin{aligned}
\frac{\partial}{\partial \alpha} R(\dot{\beta}_\alpha) \Big|_{\alpha=0} &= -2\sigma^2 (v_l - v_u) < 0 ; \quad \frac{\partial}{\partial \alpha} R(\dot{\beta}_\alpha) \Big|_{\alpha=1} = 2B(\check{\beta}) > 0, \\
\frac{\partial}{\partial \alpha^2} R(\dot{\beta}_\alpha) &= 2 \left[B(\check{\beta}) + \sigma^2 (v_l - v_u) \right] > 0.
\end{aligned}$$

From the above, we conclude that it is always beneficial to slightly move from each of the estimators towards the other. Moreover, the first derivative equals to 0 at α^* as follows:

$$\alpha^* = \frac{\sigma^2 (v_l - v_u)}{B(\check{\beta}) + \sigma^2 (v_l - v_u)}.$$

By plugging in the explicit expression of α^* into equation 17, we the following explicit expression of $R(\dot{\beta}_{\alpha^*})$:

$$\begin{aligned}
R(\dot{\beta}_{\alpha^*}) &= \frac{\sigma^4 (v_l - v_u)^2}{B(\check{\beta}) + \sigma^2 (v_l - v_u)} - 2 \frac{\sigma^4 (v_l - v_u)^2}{B(\check{\beta}) + \sigma^2 (v_l - v_u)} + \sigma^2 v_l \\
&= \sigma^2 v_l - \frac{\sigma^4 (v_l - v_u)^2}{B(\check{\beta}) + \sigma^2 (v_l - v_u)} = R(\hat{\beta}) - \frac{\sigma^4 (v_l - v_u)^2}{B(\check{\beta}) + \sigma^2 (v_l - v_u)} \quad \square
\end{aligned}$$

A.2 Proof of Theorem 2.2

The integrated risk of $\ddot{\beta}_\alpha$ can be summarized into the following bias-variance decomposition:

$$\begin{aligned}
R(\ddot{\beta}_\alpha) &= \mathbb{E}_{X, x_0} \left(\mathbb{E} \left[x_0^T \ddot{\beta}_\alpha \mid X, x_0 \right] - x_0^T \beta \right)^2 \\
&\quad + \mathbb{E}_{X, x_0} \left[\text{Var} \left(x_0^T \ddot{\beta}_\alpha \mid X, x_0 \right) \right] = B(\ddot{\beta}_\alpha) + V(\ddot{\beta}_\alpha),
\end{aligned}$$

and according to equation 8, the bias term can be written as follow:

$$B(\ddot{\beta}_\alpha) = \frac{1}{n} \text{tr} \left(\mathbb{E}_X [\beta \beta^T \Delta_\alpha^T H \Delta_\alpha] \right),$$

where,

$$\Delta_\alpha = S_\alpha (X^T X - \alpha n \bar{X} \bar{X}^T) - I ; S_\alpha = (\alpha H + (1 - \alpha) X^T X)^{-1}.$$

We can see that when $\alpha = 0$, we get $\Delta_\alpha = \mathbf{0}_{p \times p}$, and therefore:

$$B(\ddot{\beta}_\alpha)|_{\alpha=0} = \frac{\partial}{\partial \alpha} B(\ddot{\beta}_\alpha)|_{\alpha=0} = 0.$$

In order to analyze the first derivative of the bias term $B(\ddot{\beta}_\alpha)$ at $\alpha = 1$, we use the fact that $\frac{\partial}{\partial \alpha} S_\alpha|_{\alpha=1} = H^{-1} (X^T X - H) H^{-1}$, and adopt the notation introduced in 7. This allows us to express:

$$\begin{aligned} H \cdot \left(\frac{\partial}{\partial \alpha} \Delta_\alpha|_{\alpha=1} \right) &= H H^{-1} (X^T X - H) H^{-1} \left(\frac{n-1}{n} X^T X - C_X \right) \\ &\quad - H H^{-1} \left(\frac{1}{n} X^T X + C_X \right) \\ &= \left(\frac{n-1}{n} X^T X H^{-1} - I \right) X^T X + X^T X H^{-1} C_X \\ &= X^T X \left(\frac{n-1}{n} H^{-1} X^T X - I \right) + X^T X H^{-1} C_X \\ &= X^T X \Delta_{1,1} + X^T X H^{-1} C_X, \end{aligned}$$

where $\Delta_{1,1} = \left(\frac{n-1}{n} H^{-1} X^T X - I \right)$. Furthermore, it is straightforward that:

$$\Delta_{\alpha=1} = \frac{n-1}{n} H^{-1} X^T X - I + H^{-1} C_X = \Delta_{1,1} + H^{-1} C_X.$$

Combining all of the above with the fact that C_X is not correlated with $X^T X$ and $(X^T X)^2$, we get the following result.

$$\begin{aligned} \frac{\partial B(\ddot{\beta}_\alpha)}{\partial \alpha}|_{\alpha=1} &= \frac{2}{n} \mathbb{E}_X [\text{tr} (\beta \beta^T \Delta_{1,1}^T X^T X \Delta_{1,1})] \\ &\quad + \frac{2}{n} \mathbb{E}_X [\text{tr} (\beta \beta^T C_X H^{-1} X^T X H^{-1} C_X)] \\ &= \frac{2}{n} \beta^T \mathbb{E}_X [\Delta_{1,1}^T X^T X \Delta_{1,1} + C_X H^{-1} X^T X H^{-1} C_X] \beta. \end{aligned} \quad (18)$$

In the expression above, both terms in the expectation are positive semi-definite for every realization of X due to the fact that each of them is squared form. Therefore, the

expectation is positive definite, and we conclude that $\frac{\partial}{\partial \alpha} B(\ddot{\beta}_\alpha)|_{\alpha=1} \geq 0$, and the only instance in which the equality holds is when β is equal to the zero vector.

Examining the variance element $V(\ddot{\beta}_\alpha)$, we can obtain the following equations:

$$\begin{aligned} V(\ddot{\beta}_\alpha) &= \frac{\sigma^2 \xi_\alpha}{n} \text{tr} \left(H \mathbb{E}_X [S_\alpha X^T X S_\alpha] \right), \\ \frac{\partial}{\partial \alpha} V(\ddot{\beta}_\alpha) &= \frac{2\sigma^2 \xi_\alpha}{n} \text{tr} \left(H \mathbb{E}_X [S_\alpha X^T X S_\alpha (X^T X - H) S_\alpha] \right) \\ &\quad - \frac{\sigma^2}{n} \cdot \frac{2 - 2\alpha}{n} \text{tr} \left(H \mathbb{E}_X [S_\alpha X^T X S_\alpha] \right), \end{aligned} \quad (19)$$

where $\xi_\alpha = 1 - (2\alpha - \alpha^2)/n$. By taking the derivative at the extreme points, we obtain the following.

$$\begin{aligned} \frac{\partial}{\partial \alpha} V(\ddot{\beta}_\alpha)|_{\alpha=0} &= \frac{2\sigma^2}{n} \text{tr} \left(\frac{n-1}{n} H \mathbb{E}_X [(X^T X)^{-1}] - \mathbb{E}_X \left[(H(X^T X)^{-1})^2 \right] \right) \\ &< \frac{2\sigma^2}{n} \text{tr} \left(H \mathbb{E}_X [(X^T X)^{-1}] - \mathbb{E}_X \left[(H^{1/2} (X^T X)^{-1} H^{1/2})^2 \right] \right) \\ &< \frac{2\sigma^2}{n} \text{tr} \left(H \mathbb{E}_X [(X^T X)^{-1}] - (\mathbb{E}_X [H^{1/2} (X^T X)^{-1} H^{1/2}])^2 \right) \\ &= -\frac{2\sigma^2}{n} \text{tr} \left(H \mathbb{E}_X [(X^T X)^{-1}] H (\mathbb{E}_X [(X^T X)^{-1}] - H^{-1}) \right) \leq 0 \end{aligned}$$

$$\begin{aligned} \frac{\partial}{\partial \alpha} V(\ddot{\beta}_\alpha)|_{\alpha=1} &= \frac{2\sigma^2(n-1)}{n^2} \text{tr} \left(\mathbb{E}_X [(X^T X H^{-1})^2] - I_p \right) \\ &= \frac{2\sigma^2(n-1)}{n^2} \text{tr} \left(\mathbb{E}_X [(H^{-1/2} X^T X H^{-1/2})^2] - I_p \right) \\ &> \frac{2\sigma^2(n-1)}{n^2} \text{tr} \left((\mathbb{E}_X [X^T X H^{-1}])^2 - I_p \right) = 0. \end{aligned}$$

In the above derivations, we use the general results of Rahman and Ahsanullah [1973] which implies that for a given symmetric and positive definite random matrix A and any integer k , the matrix $\mathbb{E}[A^k] - (\mathbb{E}[A])^k$ is positive semi-definite. In particular, we use the fact that $\text{tr}(\mathbb{E}[A^2]) > \text{tr}((\mathbb{E}[A])^2)$ as long as A is not deterministic. From the above, we conclude that:

$$\frac{\partial}{\partial \alpha} R(\ddot{\beta}_\alpha)|_{\alpha=0} < 0 ; \quad \frac{\partial}{\partial \alpha} R(\ddot{\beta}_\alpha)|_{\alpha=1} > 0. \quad \square$$

A.3 Proof of Theorem 2.3

Focusing on the variance term of the linear-mixed estimator, $\dot{\beta}_\alpha$, we can write it as follows:

$$\begin{aligned} V(\dot{\beta}_\alpha) &= \frac{\sigma^2}{n} \text{tr} \left(H \mathbb{E}_X \left[\tilde{S}_\alpha X^T X \tilde{S}_\alpha \right] \right) - \frac{\sigma^2 p}{n} \cdot \frac{2\alpha - \alpha^2}{n} \\ &:= \frac{\sigma^2}{n} \left(\dot{V}_1(\alpha) - \dot{V}_2(\alpha) \right), \end{aligned} \quad (20)$$

where $\tilde{S}_\alpha = \alpha H^{-1} + (1 - \alpha)(X^T X)^{-1}$. By analyzing the first term \dot{V}_1 , we can show that its minimum is achieved at $\alpha = 1$ and the minimum value is p . In detail:

$$\begin{aligned} \frac{\partial}{\partial \alpha} \dot{V}_1(\alpha) &= 2 \text{tr} \left(H \mathbb{E}_X \left[\tilde{S}_\alpha X^T X (H^{-1} - (X^T X)^{-1}) \right] \right) \\ &= 2 \text{tr} \left(\mathbb{E}_X \left[(\alpha X^T X + (1 - \alpha)H) (H^{-1} - (X^T X)^{-1}) \right] \right) \\ &= 2 \text{tr} \left(\alpha (\mathbb{E}_X [X^T X] H^{-1} - I) + (1 - \alpha) (I_p - H \mathbb{E}_X [(X^T X)^{-1}]) \right) \\ &= -2(1 - \alpha) \text{tr} \left(H \mathbb{E}_X [(X^T X)^{-1} - H^{-1}] \right), \\ \frac{\partial^2}{\partial \alpha^2} \dot{V}_1(\alpha) &= 2 \text{tr} \left(H \mathbb{E}_X \left[(H^{-1} - (X^T X)^{-1}) X^T X (H^{-1} - (X^T X)^{-1}) \right] \right) > 0. \end{aligned}$$

From the above we conclude that both $\dot{V}_1(\alpha)$ and $V(\dot{\beta}_\alpha)$ are convex in α , and their global minimum is at $\alpha = 1$, where the first derivative is equal to zero. Moreover, the minimum value of $\dot{V}_1(\alpha)$ is:

$$\dot{V}_1(\alpha = 1) = \text{tr} \left(H \mathbb{E}_X [H^{-1} X^T X H^{-1}] \right) = \text{tr} (I_p) = p.$$

By putting this observation back in equation (20), we conclude that:

$$V(\dot{\beta}_\alpha) \geq \frac{\sigma^2 \xi_\alpha}{n} \text{tr} \left(H \mathbb{E}_X \left[\tilde{S}_\alpha X^T X \tilde{S}_\alpha \right] \right). \quad (21)$$

We conclude that the term in

In order to compare between $V(\dot{\beta}_\alpha)$ and $V(\ddot{\beta}_\alpha)$, we use the fact that $H \tilde{S}_\alpha X^T X$ is a symmetric matrix, and the following identity holds:

$$\text{tr} \left(H \tilde{S}_\alpha X^T X S_\alpha \right) = \text{tr} \left(X^T X \tilde{S}_\alpha H S_\alpha \right) = \text{tr} \left(H S_\alpha X^T X \tilde{S}_\alpha \right).$$

Using the above, combined with equations (19) and (21), we can write the difference between $V(\dot{\beta}_\alpha)$ and $V(\ddot{\beta}_\alpha)$ as follows.

$$\begin{aligned} V(\dot{\beta}_\alpha) - V(\ddot{\beta}_\alpha) &\geq \frac{\sigma^2 \xi_\alpha}{n} \text{tr} \left(H \mathbb{E}_X \left[\left(\tilde{S}_\alpha - S_\alpha \right) X^T X \left(\tilde{S}_\alpha + S_\alpha \right) \right] \right) \\ &= \frac{\sigma^2 \xi_\alpha}{n} \mathbb{E}_X \left[\text{tr} \left(\left(\tilde{S}_\alpha - S_\alpha \right) X^T X \left(\tilde{S}_\alpha + S_\alpha \right) H \right) \right]. \end{aligned}$$

In the expression above, we note that $\tilde{S}_\alpha - S_\alpha$ is a positive definite matrix, due to the convexity of the inverse operator, combined with the fact that both H and $X^T X$ are positive definite matrices, as stated by Nordström [2011]. Moreover, the matrix $X^T X \left(\tilde{S}_\alpha + S_\alpha \right) H$ is also positive definite since it can be written as a sum of positive definite matrices as follows:

$$X^T X \left(\tilde{S}_\alpha + S_\alpha \right) H = \alpha X^T X + (1 - \alpha)H + \left(\alpha(X^T X)^{-1} + (1 - \alpha)H^{-1} \right)^{-1}.$$

We conclude that the term inside the trace is a product of two positive definite matrices, and therefore the trace is non-negative for every realization of X , and the expectation is also non-negative. We conclude that for any given mixing ratio α , the following inequality holds:

$$V(\ddot{\beta}_\alpha) \leq V(\dot{\beta}_\alpha). \quad \square$$

A.4 Proof of Theorem 2.4

We write the function $\ddot{b}(\alpha)$ as follows:

$$B(\ddot{\beta}_\alpha) \approx \ddot{b}(\alpha) = b_0 + b_1\alpha + b_2\alpha^2 + b_3\alpha^3.$$

Regardless of the distribution P_x , $B(\ddot{\beta}_\alpha)$ satisfies the following:

$$\begin{aligned} B(\ddot{\beta}_\alpha)|_{\alpha=0} &= \frac{\partial}{\partial \alpha} B(\ddot{\beta}_\alpha)|_{\alpha=0} = 0, \\ B(\ddot{\beta}_\alpha)|_{\alpha=1} &= B(\dot{\beta}_\alpha)|_{\alpha=1} = B(\check{\beta}). \end{aligned}$$

By restricting $\ddot{b}(\alpha)$ to satisfy the same conditions as above, we get $b_0 = b_1 = 0$, and $b_2 + b_3 = B(\check{\beta})$. We also know that $B(\dot{\beta}_\alpha) = \alpha^2 B(\check{\beta})$, and therefore:

$$\frac{\partial}{\partial \alpha} B(\dot{\beta}_\alpha)|_{\alpha=1} = 2B(\check{\beta}).$$

We can also write the explicit expressions for the first derivatives of $B(\dot{\beta}_\alpha)$ and $B(\ddot{\beta}_\alpha)$ in $\alpha = 1$, for a general distribution P_X , up to a factor of $((n-1)/n)^2$, as follows.

$$\begin{aligned} \frac{\partial B(\dot{\beta}_\alpha)}{\partial \alpha} \Big|_{\alpha=1} &= \frac{2}{n} \text{tr} \left(\beta \beta^T \mathbb{E}_X \left[X^T X H^{-1} X^T X - H + C_X H^{-1} C_X \right] \right) \\ \frac{\partial B(\ddot{\beta}_\alpha)}{\partial \alpha} \Big|_{\alpha=1} &= \frac{2}{n} \text{tr} \left(\beta \beta^T \mathbb{E}_X \left[(X^T X H^{-1})^2 X^T X - 2X^T X H^{-1} X^T X + H \right] \right) \\ &\quad + \frac{2}{n} \text{tr} \left(\beta \beta^T \mathbb{E}_X \left[C_X H^{-1} C_X \right] \right) \end{aligned}$$

In order to analyze the difference between the values of the derivatives at $\alpha = 1$, we consider Assumption 2, and denote by $\tilde{X} \in \mathbb{R}^{n \times p}$ the random matrix with rows $\{\tilde{x}_i\}_{i=1}^n$, such that $X = \tilde{X}\Sigma^{1/2}$, and the following identities hold:

$$\begin{aligned} X^T X &= \Sigma^{1/2} \tilde{X}^T \tilde{X} \Sigma^{1/2} ; \quad \mathbb{E} \left[\tilde{X}^T \tilde{X} \right] = n I_p ; \quad H = n \Sigma, \\ \mathbb{E} \left[(\tilde{X}^T \tilde{X})^2 \right] &= n(n + p + q_4 - 2) I_p, \\ \mathbb{E} \left[(\tilde{X}^T \tilde{X})^3 \right] &= I_p n \left(q_6 + 3(p - 1)q_4 + p(p - 3) \right. \\ &\quad \left. + 3(n - 1)(q_4 + p - 1) + n(n - 3) + 4 \right). \end{aligned}$$

Using the above properties, we can show that the following holds:

$$\begin{aligned} \frac{\partial B(\ddot{\beta}_\alpha)}{\partial \alpha} - \frac{\partial B(\dot{\beta}_\alpha)}{\partial \alpha} \Big|_{\alpha=1} &= 2\beta^T \left(\Sigma \mathbb{E}_X \left[\frac{1}{n^3} (\tilde{X}^T \tilde{X})^3 - \frac{3}{n^2} (\tilde{X}^T \tilde{X})^2 \right] + 2\Sigma \right) \beta \\ &= 2\beta^T \Sigma \beta \left[\frac{3}{n} + \frac{p(p - 6) + 3q_4(p - 2) + 7 + q_6}{n^2} \right] > 0. \end{aligned}$$

From the above, we conclude that the first derivative of $B(\ddot{\beta}_\alpha)$ at $\alpha = 1$ is greater than the one of $B(\dot{\beta}_\alpha)$, under a wide class of distributions. We plug this constraint into $\ddot{b}(\alpha)$ and we get that necessarily $b_2 < B(\check{\beta})$. Therefore, we can write:

$$\begin{aligned} \ddot{b}(\alpha) &= b_2 \alpha^2 + \left(B(\check{\beta}) - b_2 \right) \alpha^3 = b_2(\alpha^2 - \alpha^3) + B(\check{\beta})\alpha^3 \\ &\leq B(\check{\beta})(\alpha^2 - \alpha^3 + \alpha^3) = \alpha^2 B(\check{\beta}) = B(\dot{\beta}_\alpha) \quad \square \end{aligned}$$

A.5 Proof of Proposition 2.1

By the definitions of η and α^* , their limiting behavior can be expressed by the limiting behavior of v_l , v_u and b_u . Regardless of the distribution of the covariates, the quantity v_u satisfies $v_u \rightarrow \gamma$ as $n \rightarrow \infty$. We use the fact that $X^T X$ follows the Wishart distribution and satisfies the following properties:

$$\mathbb{E} \left[(X^T X)^2 \right] = n(n + 1)\Sigma^2 + n\Sigma \text{tr}(\Sigma) ; \quad \mathbb{E} \left[(X^T X)^{-1} \right] = \frac{1}{n - p - 1} \Sigma^{-1}.$$

using the above properties, we can obtain the following limiting expressions of v_l and b_u :

$$b_u = \frac{\frac{n-1}{n} \left(p + \frac{n-1}{n} \right)}{n} \text{tr}(\Sigma) \rightarrow \gamma c^2 ; \quad v_l = \frac{p}{n - p - 1} \rightarrow \frac{\gamma}{1 - \gamma}.$$

The limiting expressions η_∞ and α_∞^* are determined by placing the above limiting values of v_l , v_u and b_u in Equations (11) and (10) respectively. \square

A.6 Proof of Proposition 2.2

Let us denote by $\tilde{X} \in \mathbb{R}^{n \times p}$ the random matrix with rows $\{\tilde{x}_i\}_{i=1}^n$, and by $C_{\tilde{X}}$ the matrix of cross terms as in (7), with components \tilde{x} . Using the fact that $X = \tilde{X}\Sigma^{1/2}$, we derive the explicit expression of b_u as follows:

$$\begin{aligned} b_u &= \frac{(n-1)^2}{n^4} \text{tr} \left(\left[\mathbb{E}(\tilde{X}^T \tilde{X})^2 - \left(\mathbb{E}[\tilde{X}^T \tilde{X}] \right)^2 \right] \Sigma \right) \\ &\quad + \frac{1}{n^2} \text{tr} \left(\mathbb{E}(C_{\tilde{X}})^2 \Sigma \right) \\ &= \left[\frac{(n-1)^2(p+q-2) + (n-1)p}{n^3} \right] \text{tr}(\Sigma). \end{aligned}$$

From the above expression we conclude that $b_u \rightarrow \gamma c^2$, as $n \rightarrow \infty$, under this generating mechanism, as in the Gaussian case. Moreover, it is clear that $v_u \rightarrow \gamma$ due to the fact that the quantity v_u is indifferent to the distribution P_X .

Regarding the quantity v_l , we denote the conditional variance component by $v_l(X)$, such that $v_l = \mathbb{E}_X[v_l(X)]$, and write as follows:

$$v_l(X) = \text{tr} \left(\Sigma (X^T X)^{-1} \right) = \text{tr} \left(\Sigma^{\frac{1}{2}} \left(\tilde{X}^T \tilde{X} \right)^{-1} \Sigma^{-\frac{1}{2}} \right) = \frac{1}{n} \text{tr} \left(\left(\frac{1}{n} \tilde{X}^T \tilde{X} \right)^{-1} \right).$$

We now note that according to the Marchenko-Pastur theorem [Marchenko and Pastur, 1967, Silverstein, 1995], under the asymptotic setting, the spectral measure of $\frac{1}{n} \tilde{X}^T \tilde{X}$ converges weakly, almost surely, to the Marchenko-Pastur law, depending only on γ , and bounded away from 0. Therefore, we can write that as $n, p \rightarrow \infty$, almost surely:

$$v_l(X) = \frac{1}{n} \text{tr} \left(\left(\frac{1}{n} \tilde{X}^T \tilde{X} \right)^{-1} \right) \rightarrow \gamma s(\gamma),$$

where $s(\gamma)$ can be obtained by the Marchenko-Pastur law. In the proof of Hastie et al. [2022], Proposition 2, it was shown that $s(\gamma) = 1/(1-\gamma)$, concluding that $v_l(X)$, converges almost surely in X to $\gamma/(1-\gamma)$, which is also the expected value of $v_l(X)$ in the Gaussian case. The almost sure convergence combined with the fact that $v_l(X) \geq 0$ for all possible realizations of X , yield that for every $\tilde{\epsilon} > 0$, there exists an $n(\tilde{\epsilon})$ such that:

$$v_l = \mathbb{E}_X[v_l(X)] \geq \frac{\gamma}{1-\gamma} - \tilde{\epsilon}, \quad \forall n > n(\tilde{\epsilon}).$$

Thus, as long as Assumption 1 holds, v_l either converges to a fixed value that is greater than or equal to the one in the Gaussian case, or diverges to large values. Noting that η

decreases as v_l increases (while b_u and v_u are fixed), we conclude that η either converges to a fixed value that is smaller than or equal to η_∞ , or fluctuates between values in the interval $(0, \eta_\infty]$ as $n \rightarrow \infty$. In the same way, α^* increases as v_l increases, and therefore lies in the interval $[\alpha_\infty^*, 1)$ as $n \rightarrow \infty$. \square

A.7 Proof of Proposition 2.3

For any given training size n , the estimator $\hat{\sigma}^2$ is an unbiased estimator of σ^2 , and its total variance over $T = (X, Y)$ is $\text{Var}(\hat{\sigma}^2) = 2\sigma^4/(n(1 - \gamma))$. Therefore, $\hat{\sigma}^2$ converges in probability to σ^2 , and also $\hat{\alpha}$ converges in probability to α^* . By the definitions of $\beta^{M(\tau)}$ and β^{M^*} , the difference between the predictions can be written as follows.

$$x_0^T \beta^{M^*} - x_0^T \beta^{M(\tau)} = (\hat{\alpha} - \alpha^*) x_0^T (\hat{\beta} - \check{\beta}).$$

Under the distributional assumptions taken over P_x , the total variance of $x_0^T (\hat{\beta} - \check{\beta})$ converges to the following fixed value as $n \rightarrow \infty$:

$$\text{Var} \left(x_0^T (\hat{\beta} - \check{\beta}) \right) \rightarrow \frac{\sigma^2 \gamma^2}{1 - \gamma} + \gamma c^2 \tau^2.$$

From the fact that $\hat{\alpha} - \alpha^*$ converges in probability to 0, and the variance of $x_0^T (\hat{\beta} - \check{\beta})$ is asymptotically bounded, it is elementary that the product between the two terms also converges in probability to 0. \square

A.8 Proof of Proposition 3.1

Denote by $\Psi \in \mathbb{R}^{n \times p \times p}$, $\Psi_{ijk} = x_{ij} x_{ij}$, the tensor that collects the data $x_i x_i^T$, $i \in \{1 \dots n\}$, for a specific realization of the covariates matrix X . Under Assumption 4, we have that $\mathbb{E}[X^T D X | \Psi] = \sum_i \mathbb{E}[g'(x_i^T \beta) x_i x_i^T | \Psi_i = x_i x_i^T] = \delta X^T X$, and therefore the following chain of equalities is straight forward:

$$\begin{aligned} \mathbb{E}[X^T X] H^{-1} &= \mathbb{E}[X^T X] (\delta \mathbb{E}[X^T X])^{-1} = I_p \delta^{-1} \\ &= \mathbb{E} \left[X^T X (\mathbb{E}[X^T D X | \Psi])^{-1} \right]. \end{aligned} \tag{22}$$

Due to the $(n - 1)/n$ factor in equation (14), the difference $v_l - v_u$ is strictly larger than the following expression:

$$\begin{aligned} v_l - v_u &> \text{tr} \left(H\mathbb{E} \left[H_X^{-1} X^T X H_X^{-1} - H^{-1} X^T X H^{-1} \right] \right) \\ &= \text{tr} \left(H\mathbb{E} \left[(H_X^{-1} - H^{-1}) X^T X (H_X^{-1} - H^{-1}) \right] \right) \\ &+ \text{tr} \left(H\mathbb{E} \left[H^{-1} X^T X H_X^{-1} + H_X^{-1} X^T X H^{-1} - 2X^T X H^{-2} \right] \right), \end{aligned}$$

where $H_X = X^T D X$. The first term in the expression above is a trace of product of two positive definite matrices and therefore positive. Using equation (22), the second term can be written as follows:

$$2\mathbb{E} \left[\text{tr} \left(X^T X \left(\mathbb{E}[(X^T D X)^{-1} | \Psi] - (\mathbb{E}[X^T D X | \Psi])^{-1} \right) \right) \right].$$

The result by Groves and Rothenberg [1969] implies that the matrix $\mathbb{E}[(X^T D X)^{-1} | \Psi] - (\mathbb{E}[X^T D X | \Psi])^{-1}$ is positive semi-definite for any realization of Ψ , and therefore the trace is always non-negative. We conclude that the above expectation is non-negative and therefore $v_l - v_u > 0$. \square

A.9 Proof of Proposition 3.2

By the definition of $\dot{r}(\alpha)$, we can write:

$$\begin{aligned} \dot{r}(\alpha) &= \mathbb{E}_T [\dot{a}_\alpha^T H \dot{a}_\alpha] = \alpha^2 (B(\check{\beta}) + \sigma^2 v_u) + (1 - \alpha)^2 \sigma^2 v_l \\ &+ 2\alpha(1 - \alpha) \mathbb{E}_T [\hat{a}^T H \check{a}] \\ &= \alpha^2 (B(\check{\beta}) + \sigma^2 v_u) + (1 - \alpha)^2 \sigma^2 v_l \\ &+ 2\alpha(1 - \alpha) \frac{n-1}{n} \sigma^2 \text{tr} (\mathbb{E}_X [X^T X (X^T D X)^{-1}]) \\ &:= \alpha^2 \left(B(\check{\beta}) + \sigma^2 v_u \right) + (1 - \alpha)^2 \sigma^2 v_l + 2\alpha(1 - \alpha) \sigma^2 v_s \\ &= \alpha^2 \left(B(\check{\beta}) + \sigma^2 (v_l + v_u - 2v_s) \right) - 2\alpha \sigma^2 (v_l - v_s) + \sigma^2 v_l, \end{aligned}$$

first and second derivatives of $\dot{r}(\alpha)$ with respect to α , we get that:

$$\begin{aligned} \frac{\partial}{\partial \alpha} \dot{r}(\alpha)|_{\alpha=0} &= -2\sigma^2 (v_l - v_s) ; \quad \frac{\partial}{\partial \alpha} \dot{r}(\alpha)|_{\alpha=1} = 2 \left(B(\check{\beta}) + \sigma^2 (v_u - v_s) \right), \\ \frac{\partial}{\partial \alpha^2} \dot{r}(\alpha) &= 2 \left(B(\check{\beta}) + \sigma^2 (v_l + v_u - 2v_s) \right). \end{aligned}$$

Looking at the term $v_l + v_u - 2v_s$, we can see that it can be written as follows:

$$\frac{n-1}{n} \text{tr} \left(H \mathbb{E}_X \left[(H_X^{-1} - H^{-1})^T X^T X (H_X^{-1} - H^{-1}) \right] \right) + \frac{v_l}{n},$$

which is an expectation of quadratic form, plus a positive term. Therefore, $v_l + v_u - 2v_s > 0$, which implies that the second derivative of $\dot{r}(\alpha)$ is strictly positive, and the unique global minimum is:

$$\dot{\alpha} = \frac{\sigma^2(v_l - v_s)}{B(\check{\beta}) + \sigma^2(v_l + v_u - 2v_s)}.$$

Using the fact that $v_l > v_u$, we conclude that $v_l > v_s$ and therefore $\frac{\partial}{\partial \alpha} \dot{r}(\alpha)|_{\alpha=0} < 0$ and necessarily: $\dot{\alpha} > 0$. \square

A.10 Proof of Proposition 3.3

Let us first analyze the variance term associated with $\dot{r}(\alpha)$ as follows:

$$\begin{aligned} \dot{V}(\alpha) &= \sigma^2 \text{tr} \left(H \mathbb{E}_X \left[\tilde{S}_\alpha X^T X \tilde{S}_\alpha \right] \right) \\ &\quad - \frac{\sigma^2}{n-1} (\alpha^2 v_u + 2\alpha(1-\alpha)v_s) \end{aligned} \tag{23}$$

$$= \sigma^2 [(1-\alpha)^2 v_l + \alpha^2 v_u + 2\alpha(1-\alpha)v_s], \tag{24}$$

where $\tilde{S}_\alpha = \alpha H^{-1} + (1-\alpha)(X^T D X)^{-1}$. Using the properties that $v_l + v_u - 2v_s > 0$ and $v_l > v_u$ we can show that the following inequality holds:

$$\frac{\alpha^2 v_u + 2\alpha(1-\alpha)v_s}{n-1} \leq (1-\xi_\alpha) \text{tr} \left(H \mathbb{E}_X \left[\tilde{S}_\alpha X^T X \tilde{S}_\alpha \right] \right),$$

and together with (23), we conclude that:

$$\dot{V}(\alpha) \geq \xi_\alpha \sigma^2 \text{tr} \left(H \mathbb{E}_X \left[\tilde{S}_\alpha X^T X \tilde{S}_\alpha \right] \right).$$

We now show the following identity for any α and realization of X :

$$\begin{aligned} \tilde{S}_\alpha X^T X S_\alpha &= \left(\alpha^2 H (X^T X)^{-1} H + (1-\alpha)^2 H_X (X^T X)^{-1} H_X \right. \\ &\quad \left. + \alpha(1-\alpha) H X H_X + \alpha(1-\alpha) H_X X H \right)^{-1} = S_\alpha X^T X \tilde{S}_\alpha. \end{aligned}$$

Using the above, we can write:

$$\dot{V}(\alpha) - \ddot{V}(\alpha) \geq \xi_\alpha \sigma^2 \text{tr} \left(H \mathbb{E}_X \left[\left(\tilde{S}_\alpha - S_\alpha \right) X^T X \left(\tilde{S}_\alpha + S_\alpha \right) \right] \right).$$

In the expression above, we note that $\tilde{S}_\alpha - S_\alpha$ is a positive definite matrix, due to the convexity of the inverse operator, combined with the fact that both H and $X^T X$ are positive definite matrices, as stated by Nordström [2011]. Moreover, the matrix

$$X^T X \left(\tilde{S}_\alpha - S_\alpha \right) X^T X \left(\tilde{S}_\alpha + S_\alpha \right)$$

is semi positive definite since it is a symmetric product of positive definite matrices. We conclude that the term inside the trace is a product of two semi positive definite matrices, and therefore the trace is non negative for every realization of X , and the expectation is also non negative. We conclude that for any given mixing ratio α , the following inequality holds:

$$\dot{V}(\alpha) \geq \ddot{V}(\alpha).$$

On the other hand, from equation (24) we find that:

$$\frac{\partial}{\partial \alpha} \dot{V}(\alpha)|_{\alpha=0} = 2\sigma^2(v_s - v_l) < 0.$$

Since $\dot{V}(\alpha = 0) = \ddot{V}(\alpha = 0)$, together with the fact that $\ddot{V}(\alpha) \leq \dot{V}(\alpha)$, we conclude that necessarily $\frac{\partial}{\partial \alpha} \ddot{V}(\alpha)|_{\alpha=0} < 0$. By analyzing the bias term of $\ddot{r}(\alpha)$, we can immediately find that $\frac{\partial}{\partial \alpha} \ddot{B}(\alpha)|_{\alpha=0} = 0$. Thus, we have $\frac{\partial}{\partial \alpha} \ddot{r}(\alpha)|_{\alpha=0} < 0$.

Now, let us look at the following expression of the first derivative at $\alpha = 1$:

$$\begin{aligned} \frac{\partial}{\partial \alpha} \ddot{r}(\alpha)|_{\alpha=1} &= \frac{n-1}{n} 2\sigma^2 \text{tr} \left(H^{-1} \mathbb{E}_X [H_X H^{-1} X^T X - X^T X] \right) \\ &\quad + 2\mathbb{E}_X \left[\text{tr} \left(H^{-1} H_X H^{-1} \zeta \zeta^T \right) \right]. \end{aligned}$$

Under Assumption 4, the first term (variance) above is positive, since the expression in the trace can be written as:

$$\delta \left(\mathbb{E} \left[\left(H^{-1/2} X^T X H^{-1/2} \right)^2 \right] - \left(\mathbb{E} \left[H^{-1/2} X^T X H^{-1/2} \right] \right)^2 \right).$$

The second term (bias) is necessarily positive, since we have a product of two positive definite matrices inside the trace. We conclude that the first derivative of $\ddot{r}(\alpha)$ is positive at $\alpha = 1$, and $\text{argmin}\{\ddot{r}(\alpha)\}$ is in the open interval $(0, 1)$. \square

A.11 Proof of Proposition 5.1

Let us denote by $\underline{\epsilon} \in \mathbb{R}^n$, the vector of random noise, such that $y_i = x_i^T w + \underline{\epsilon}_i$ for $i = 1 \cdots n$. By definition, for any given realization of $(w, X, Y = Xw + \underline{\epsilon})$:

$$\tilde{w}^T \Sigma \tilde{w} \leq \hat{w}^T \Sigma \hat{w}.$$

Therefore, for any setting of (P_X, σ^2, τ^2) , the integrated expectations hold that:

$$\mathbb{E} [\tilde{w}^T \Sigma \tilde{w}] \leq \mathbb{E} [\hat{w}^T \Sigma \hat{w}].$$

The derivation of the above expectations shows that:

$$\begin{aligned} & \tau^2 \text{tr} (\mathbb{E}_X [X^T (X \Sigma^{-1} X^T)^{-1} X]) + \sigma^2 v_u \\ & \leq \tau^2 \text{tr} (\Sigma \mathbb{E}_X [X^T (X X^T)^{-1} X]) + \sigma^2 v_l, \end{aligned}$$

which holds if and only if:

$$\sigma^2 v_u - \tau^2 b_u \leq \sigma^2 v_l - \tau^2 b_l.$$

Since the above inequality holds for any setting of (σ^2, τ^2) , it necessarily holds that $b_l \leq b_u$ and $v_l \geq v_u$. \square

A.12 Proof of Theorem 5.1

Calculating first and second derivatives of $R(\dot{w}_\alpha)$ with respect to α , we get:

$$\begin{aligned} \frac{\partial}{\partial \alpha} R(\dot{w}_\alpha) &= 2\alpha [\tau^2(b_u - b_l) + \sigma^2(v_l - v_u)] - 2\sigma^2(v_l - v_u), \\ \frac{\partial}{\partial \alpha} R(\dot{w}_\alpha)|_{\alpha=0} &= -2\sigma^2(v_l - v_u) < 0, \quad ; \quad \frac{\partial}{\partial \alpha} R(\dot{w}_\alpha)|_{\alpha=1} = 2\tau^2(b_u - b_l) > 0, \\ \frac{\partial}{\partial \alpha^2} R(\dot{w}_\alpha) &= 2 [\tau^2(b_u - b_l) + \sigma^2(v_l - v_u)] > 0. \end{aligned}$$

From the above, we conclude that for **any** SNR, it is beneficial to slightly move from each one of the estimators toward the other. Moreover, the best mixed interpolator is achieved where the first derivative equals to 0 as follows:

$$\alpha^* = \frac{\sigma^2(v_l - v_u)}{\tau^2(b_u - b_l) + \sigma^2(v_l - v_u)}. \quad \square \quad (25)$$

A.13 Proof of Proposition 5.2

By taking iterative expectation over X , w , and Y , we get:

$$\mathbb{E}[\hat{\sigma}^2] = \mathbb{E}_X \left[\frac{\mathbb{E}_w [\text{tr}((XX^T)^{-2} \mathbb{E}_Y [YY^T | X, w])] - \tau^2 \text{tr}((XX^T)^{-1})}{\text{tr}((XX^T)^{-2})} \right],$$

and since the underlying model satisfies $\mathbb{E}_Y [YY^T | X, w] = I_n \sigma^2 + Xww^T X^T$, we can write:

$$\mathbb{E}[\hat{\sigma}^2] = \mathbb{E}_X \left[\frac{\text{tr}((XX^T)^{-2} \sigma^2 + X^T (XX^T)^{-2} X \mathbb{E}_w [ww^T]) - \tau^2 \text{tr}((XX^T)^{-1})}{\text{tr}((XX^T)^{-2})} \right].$$

Now, since $\mathbb{E}_w [ww^T] = I_p \tau^2$, the middle and right terms in the numerator cancel, and we are left with:

$$\mathbb{E}[\hat{\sigma}^2] = \mathbb{E}_X \left[\frac{\text{tr}((XX^T)^{-2}) \sigma^2}{\text{tr}((XX^T)^{-2})} \right] = \sigma^2. \quad \square$$

A.14 Proof of Theorem 5.2

For a given covariance matrix Σ , let us denote by U , the random matrix $U = X\Sigma^{-0.5}$. The properties of the Gaussian distribution imply that the rows of the matrix $U = X\Sigma^{-0.5}$ have a scaled Gaussian distribution with i.i.d. components. Using the connection between U and X , we can write the terms v_u and b_u as follows:

$$v_u = \text{tr}(\mathbb{E}[(UU^T)^{-1}]) \quad ; \quad b_u = \text{tr}(\Sigma) - \text{tr}(\mathbb{E}[\Sigma^{1/2} U^T (UU^T)^{-1} U \Sigma^{1/2}]).$$

Taking advantage of the intrinsic properties of the matrix U , including its rotational invariance, and the attributes of the generalized inverse of the Wishart matrix (as outlined in Cook and Forzani [2011]), we can deduce exact expressions for v_u and b_u , as well as their asymptotic values, as described below.

$$v_u = \frac{1}{p-n-1} \text{tr}(I_n) = \frac{n}{p-n-1} \rightarrow \frac{1}{\gamma-1},$$

$$b_u = \text{tr}(\Sigma) - \frac{\text{tr}(\Sigma)}{p} \text{tr}(\mathbb{E}[U^T (UU^T)^{-1} U]) = \text{tr}(\Sigma) \left(1 - \frac{n}{p}\right) \rightarrow c^2 \left(1 - \frac{1}{\gamma}\right).$$

For the manner of analyzing the terms v_l and b_l , we adopt the conclusion that arised by Hastie et al. [2022] and also by Bartlett et al. [2020], that the estimator \hat{w} is affected by the vanishing eigenvalues of Σ in a manner that only the \tilde{p} non-vanishing components of Σ are taken into account. Therefore, we can analyze the prediction error of \hat{w} as is done

with \tilde{p} -dimensional i.i.d. Gaussian covariates, by taking into account $\tilde{\gamma}$ in place of γ . More precisely, we can assume that $x \sim MN(\mathbf{0}_{\tilde{p}}, c_1^2(n)I_{\tilde{p}})$ and the rows of the matrix $U = \frac{1}{c_1(n)}X$ have a scaled Gaussian distribution with i.i.d. components. Therefore we can write the terms v_l and b_l as follows:

$$v_l = \text{tr} \left(\mathbb{E} [U^T (UU^T)^{-2} U] \right) = \frac{1}{\tilde{p} - n - 1} \text{tr} (I_n) \rightarrow \frac{1}{\tilde{\gamma} - 1},$$

$$b_l = \text{tr} \left(c_1^2(n)I_{\tilde{p}} - c_1^2(n)I_{\tilde{p}} \mathbb{E}_X [X^T (X X^T)^{-1} X] \right) = c_1^2(n)(\tilde{p} - n) \rightarrow c^2 \left(1 - \frac{1}{\tilde{\gamma}} \right).$$

For any given combination of $(\sigma^2, \tau^2, c^2, \gamma, \tilde{\gamma})$, we can plug in the above limiting values into the formulas of η and α^* and get the following limiting expressions:

$$\eta \rightarrow \eta_\infty = 1 - \frac{(\gamma - \tilde{\gamma})\sigma^4}{(\gamma - 1)^2(\tilde{\gamma} - 1)^2 \left[\frac{\tau^2 c^2}{\gamma \tilde{\gamma}} + \frac{\sigma^2}{(\gamma - 1)(\tilde{\gamma} - 1)} \right] \left[\frac{\tau^2 c^2 (\tilde{\gamma} - 1)}{\tilde{\gamma}} + \frac{\sigma^2}{\tilde{\gamma} - 1} \right]},$$

$$\alpha^* \rightarrow \alpha_\infty^* = \frac{\sigma^2}{(\gamma - 1)(\tilde{\gamma} - 1) \left[\frac{\tau^2 c^2}{\gamma \tilde{\gamma}} + \frac{\sigma^2}{(\gamma - 1)(\tilde{\gamma} - 1)} \right]}.$$

□

B Challenging the total information assumption

In this Section we analyze the case where the amount of the unlabeled data, m is comparable to the amount of the labeled data, n , and the dimension of the data, p . In this case, the variability in the estimation of parameters like $\mathbb{E}[xx^T]$ is not negligible and affects the prediction error of the semi-supervised estimators. In subsequent analysis, we show that, on the one hand, as anticipated, the prediction error surpasses that observed in the total information scenario. On the other hand, we show that the central argument of Theorem 2.1 holds as long as the quantity of unlabeled data exceeds that of the labeled data. Additionally, for Gaussian covariates specifically, we can precisely express both the optimal mixing ratio α^* , and the ratio η that compares the prediction error of the optimally adjusted semi-supervised estimator to that of the supervised estimator.

Denoting the set of unlabeled data by $Z \in \mathbb{R}^{m \times p}$, the linear mixed semi-supervised estimator under discussion can be written as follows:

$$\hat{\beta}_\alpha = (1 - \alpha)\hat{\beta} + \alpha\tilde{\beta} ; \quad \tilde{\beta} = \frac{m}{n} (Z^T Z)^{-1} X^T Y. \quad (26)$$

We note that in the estimator $\tilde{\beta}$ above, we estimate the matrix $H = \mathbb{E}[X^T X]$ by $\frac{n}{m} Z^T Z$, which is now a random matrix and not fixed as in the total information case. For the sake of brevity, we focus on the properties of the estimator $\dot{\beta}_\alpha$ under the random- β scenario and assume that the rows of Z are independent of the rows of X . In this setting, we can write the prediction error of $\tilde{\beta}$, as follows:

$$\begin{aligned} R(\tilde{\beta}) &= \frac{\tau^2}{n} \text{tr} \left(H \mathbb{E} \left[\left(I - \frac{m}{n} (Z^T Z)^{-1} X^T X \right) \left(I - \frac{m}{n} (Z^T Z)^{-1} X^T X \right)^T \right] \right) \\ &\quad + \frac{\sigma^2 m^2}{n^3} \text{tr} \left(H \mathbb{E} \left[(Z^T Z)^{-1} X^T X (Z^T Z)^{-1} \right] \right) \\ &= \tau^2 b_{\tilde{u}} + \sigma^2 v_{\tilde{u}} := B(\tilde{\beta}) + V(\tilde{\beta}). \end{aligned}$$

Utilizing the above expression of $R(\tilde{\beta})$, the prediction error of the mixed estimator $\dot{\beta}_\alpha$, as a function of α , can be written as follows:

$$\begin{aligned} R(\dot{\beta}_\alpha) &= \alpha^2 \tau^2 b_{\tilde{u}} + \alpha^2 \sigma^2 v_{\tilde{u}} + (1 - \alpha)^2 \sigma^2 v_l + 2\alpha(1 - \alpha) \mathbb{E}_{\mathcal{X}} \left[\text{Cov} \left(x_0^T \hat{\beta}, x_0^T \tilde{\beta} \mid \mathcal{X} \right) \right] \\ &:= \alpha^2 \tau^2 b_{\tilde{u}} + \alpha^2 \sigma^2 v_{\tilde{u}} + (1 - \alpha)^2 \sigma^2 v_l + 2\alpha(1 - \alpha) v_{\tilde{s}}, \end{aligned} \quad (27)$$

where: $\mathcal{X} = (X, Z, x_0)$ and $v_{\tilde{s}} = \frac{m}{n^2} \text{tr} \left(\mathbb{E} \left[(Z^T Z)^{-1} \right] H \right)$.

When we compute the first and second derivatives of $R(\dot{\beta}_\alpha)$ with respect to α , it results in:

$$\begin{aligned} \frac{\partial}{\partial \alpha} R(\dot{\beta}_\alpha) \Big|_{\alpha=0} &= -2\sigma^2 (v_l - v_{\tilde{s}}), \\ \frac{\partial}{\partial \alpha} R(\dot{\beta}_\alpha) \Big|_{\alpha=1} &= 2 \left[\tau^2 b_{\tilde{u}} + \sigma^2 (v_{\tilde{u}} - v_{\tilde{s}}) \right], \\ \frac{\partial^2}{\partial \alpha^2} R(\dot{\beta}_\alpha) &= 2 \left[\tau^2 b_{\tilde{u}} + \sigma^2 (v_l + v_{\tilde{u}} - 2v_{\tilde{s}}) \right]. \end{aligned}$$

We note that the term $v_l + v_{\tilde{u}} - 2v_{\tilde{s}}$ is non-negative, since it can be written as an expectation of a trace of squared form as follows:

$$v_l + v_{\tilde{u}} - 2v_{\tilde{s}} = \mathbb{E} \left[\text{tr} \left(H^{1/2} Q Q^T H^{1/2} \right) \right],$$

where:

$$Q = \frac{1}{n^{0.5}} (X^T X)^{-1/2} - \frac{m}{n^{1.5}} (Z^T Z)^{-1} (X^T X)^{1/2}.$$

Based on the preceding analysis, it is determined that the second derivative of $R(\dot{\beta}_\alpha)$ is positive, which confirms the convexity of $R(\dot{\beta}_\alpha)$ with respect to α .

We now point to the fact that if the key condition $v_{\bar{s}} < v_l$ holds, it is always beneficial to slightly move from supervised estimator, $\hat{\beta}$, towards $\tilde{\beta}$. If the condition $v_{\bar{s}} < v_{\bar{u}}$ holds as well, the unique global minimizer of $R(\hat{\beta}_\alpha)$, is necessarily in the open interval $(0, 1)$. In general, α^* admits the following explicit formula:

$$\alpha^* = \frac{\sigma^2(v_l - v_{\bar{s}})}{\tau^2 b_{\bar{u}} + \sigma^2(v_l + v_{\bar{u}} - 2v_{\bar{s}})}. \quad (28)$$

The key condition $v_{\bar{s}} < v_l$ holds if and only if $\mathbb{E} \left[\left(\frac{1}{n} X^T X \right)^{-1} \right] - \mathbb{E} \left[\left(\frac{1}{m} Z^T Z \right)^{-1} \right]$ is a positive definite matrix, which seems to be a mild requirement in any case of $m > n$. Indeed, we are able to show that this condition holds without any distributional assumption other than Assumption 1, when $m = Kn$, for $K = 2, 3, \dots$, as follows:

$$\begin{aligned} \mathbb{E} \left[\left(\frac{1}{n} X^T X \right)^{-1} \right] &= \frac{1}{K} \sum_{k=1}^K \mathbb{E} \left[\left(\frac{1}{n} X^T X \right)^{-1} \right] = \mathbb{E} \left[\sum_{k=1}^K \frac{1}{Kn} \left(\sum_{i=s(k)}^{kn} z_i z_i^T \right)^{-1} \right] \\ &\geq \mathbb{E} \left[\left(\sum_{k=1}^K \frac{1}{Kn} \sum_{i=s(k)}^{kn} z_i z_i^T \right)^{-1} \right] = \mathbb{E} \left[\left(\frac{1}{m} Z^T Z \right)^{-1} \right]. \end{aligned}$$

In the above derivation, $s(k) = 1 + (k - 1)n$, and the symbol \geq denotes the relationship between positive definite (P.D.) matrices. This is valid because of the fact that the expected value of the sum of inverse matrices is (in the sense of P.D.) larger than the expected value of the inverse of the sum of matrices. Thus, we conclude that the mixed estimator is beneficial in general, without the total information assumption.

Let us also assume that the distribution P_x satisfies the conditions of Assumption 2, and denote by $\tilde{Z} \in \mathbb{R}^{m \times p}$ the random matrix with scaled i.i.d. entrees $\{z_{ij}\}$, such that $Z = \tilde{Z}\Sigma^{1/2}$. In this case, we can show that the condition $v_{\bar{s}} < v_{\bar{u}}$ also holds, as follows:

$$\begin{aligned} v_{\bar{u}} &= \frac{m^2}{n^3} \text{tr} \left(H \mathbb{E} \left[(Z^T Z)^{-1} X^T X (Z^T Z)^{-1} \right] \right) = \frac{m^2}{n} \text{tr} \left(\mathbb{E} \left[(\tilde{Z}^T \tilde{Z})^{-2} \right] \right) \\ &\geq \frac{m^2}{n} \text{tr} \left(\left(\mathbb{E} \left[(\tilde{Z}^T \tilde{Z})^{-1} \right] \right)^2 \right) \geq \frac{m^2}{n} \text{tr} \left(\mathbb{E} \left[(\tilde{Z}^T \tilde{Z})^{-1} \right] \left(\mathbb{E} \left[(\tilde{Z}^T \tilde{Z}) \right] \right)^{-1} \right) \\ &= \frac{m}{n} \text{tr} \left(\mathbb{E} \left[(\tilde{Z}^T \tilde{Z})^{-1} \right] \right) = \frac{m}{n} \text{tr} \left(\mathbb{E} \left[(Z^T Z)^{-1} \right] \Sigma \right) = v_{\bar{s}}. \end{aligned}$$

From the above, we conclude that Theorem 2.1 is valid for a wide class of distributions, without the requirement of the total information assumption.

In the more specific case of Gaussian covariates, we can explicitly write all relevant terms: $b_{\bar{u}}$, $v_{\bar{u}}$, $v_{\bar{s}}$, and v_l . Therefore, we can get more insights on how the mixed semi-

supervised estimator is effected by the size of the unlabeled data. In particular, we can compare between v_l and v_s as follows:

$$v_l - v_s = p \left(\frac{1}{n - p - 1} - \frac{1}{n - p \frac{m}{n} - 1} \right),$$

which is positive as long as $m > n$, revealing that the SSL is beneficial when the amount of unlabeled data exceeds that of the labeled data. In order to obtain a succinct view on the properties of $\dot{\beta}_\alpha$, we consider the asymptotic setting in which $p/n \rightarrow \gamma \in (0, 1)$ and $p/m \rightarrow \tilde{\gamma}$ as $n, m, p \rightarrow \infty$, and assumption 3 is satisfied. In this context, we are able to show that:

$$b_{\tilde{u}} \rightarrow c^2 \frac{1 + (1 - \tilde{\gamma})^3 - 2(1 - \tilde{\gamma})^2 + \gamma}{(1 - \tilde{\gamma})^3},$$

$$v_{\tilde{u}} \rightarrow \frac{\gamma}{(1 - \tilde{\gamma})^3} ; v_s \rightarrow \frac{\gamma}{1 - \tilde{\gamma}}.$$

From the above, it is clear that $R(\tilde{\beta})$ progressively increases with $\tilde{\gamma}$ and matches $R(\check{\beta})$ when $\tilde{\gamma} = 0$. Similarly, $R(\dot{\beta}_{\alpha^*})$ aligns with the total information scenario as $\tilde{\gamma}$ approaches 0. Moreover, these limiting values are useful for defining the limiting values of α^* in a specific configuration of $(\gamma, \tilde{\gamma})$, as outlined in Equation (28) and η , integrating all relevant elements in Equation (27).

References

- D. Azriel, L. D. Brown, M. Sklar, R. Berk, A. Buja, and L. Zhao. Semi-supervised linear regression. *Journal of the American Statistical Association*, 117(540):2238–2251, 2022.
- P. L. Bartlett, P. M. Long, G. Lugosi, and A. Tsigler. Benign overfitting in linear regression. *Proceedings of the National Academy of Sciences*, 117(48):30063–30070, 2020.
- M. Belkin, D. Hsu, S. Ma, and S. Mandal. Reconciling modern machine-learning practice and the classical bias–variance trade-off. *Proceedings of the National Academy of Sciences*, 116(32):15849–15854, 2019.
- P. C. Bellec and C. H. Zhang. De-biasing the lasso with degrees-of-freedom adjustment. *Bernoulli*, 28(2):713–743, 2022.

- P. C. Bellec, A. S. Dalalyan, E. Grappin, and Q. Paris. On the prediction loss of the lasso in the partially labeled setting. *Electronic Journal of Statistics*, 12(2):3443–3472, 2018.
- J. Bennett, S. Lanning, et al. The netflix prize. In *Proceedings of KDD cup and workshop*, volume 2007, page 35. New York, 2007.
- C. Brouard, F. d’Alché Buc, and M. Szafranski. Semi-supervised penalized output kernel regression for link prediction. In *28th International Conference on Machine Learning (ICML 2011)*, pages 593–600, 2011.
- A. Chakraborty and T. Cai. Efficient and adaptive linear regression in semi-supervised settings. *The Annals of Statistics*, 46(4):1541–1572, 2018.
- O. Chapelle and A. Zien. Semi-supervised classification by low density separation. In *International workshop on artificial intelligence and statistics*, pages 57–64. PMLR, 2005.
- D. A. Clevert, T. Unterthiner, and S. Hochreiter. Fast and accurate deep network learning by exponential linear units (elus). *arXiv preprint arXiv:1511.07289*, 2015.
- R. D. Cook and L. Forzani. On the mean and variance of the generalized inverse of a singular Wishart matrix. *Electronic Journal of Statistics*, 5(none):146 – 158, 2011. doi: 10.1214/11-EJS602. URL <https://doi.org/10.1214/11-EJS602>.
- N. El Karoui. Concentration of measure and spectra of random matrices: Applications to correlation matrices, elliptical distributions and beyond. *The Annals of Applied Probability*, 19(6):2362–2405, 2009. doi: 10.1214/08-AAP548.
- M. Geiger, S. Spigler, S. d’Ascoli, L. Sagun, M. Baity-Jesi, G. Biroli, and M. Wyart. Jamming transition as a paradigm to understand the loss landscape of deep neural networks. *Physical Review E*, 100(1):012115, 2019.
- T. Groves and T. Rothenberg. A note on the expected value of an inverse matrix. *Biometrika*, 56(3):690–691, 1969.
- T. Han, J. Gao, Y. Yuan, and Q. Wang. Unsupervised semantic aggregation and deformable template matching for semi-supervised learning. *Advances in Neural Information Processing Systems*, 33:9972–9982, 2020.

- T. Hastie, A. Montanari, S. Rosset, and R. J. Tibshirani. Surprises in high-dimensional ridgeless least squares interpolation. *The Annals of Statistics*, 50(2):949–986, 2022.
- R. He, Z. Tian, and M. J. Zuo. A semi-supervised gan method for rul prediction using failure and suspension histories. *Mechanical Systems and Signal Processing*, 168:108657, 2022.
- Z. Hu and H. Huang. On the random conjugate kernel and neural tangent kernel. In *International Conference on Machine Learning*, pages 4359–4368. PMLR, 2021.
- A. Javanmard and A. Montanari. Confidence intervals and hypothesis testing for high-dimensional regression. *The Journal of Machine Learning Research*, 15(1):2869–2909, 2014.
- A. Javanmard and A. Montanari. Debiasing the lasso: Optimal sample size for gaussian designs. *Annals of Statistics*, 46(6A):2593–2622, 2018.
- N. Jean, S. M. Xie, and S. Ermon. Semi-supervised deep kernel learning: Regression with unlabeled data by minimizing predictive variance. *Advances in Neural Information Processing Systems*, 31, 2018.
- Kaggle. Dataset from netflix’s competition to improve their recommendation algorithm. In <https://www.kaggle.com/datasets/netflix-inc/netflix-prize-data>, 2005.
- Z. Liu, P. Luo, X. Wang, and X. Tang. Large-scale celebfaces attributes (celeba) dataset. *Retrieved August*, 15(2018):11, 2018.
- V. A. Marchenko and L. A. Pastur. Distribution of eigenvalues for some sets of random matrices. *Matematicheskii Sbornik*, 114(4):507–536, 1967.
- P. Niyogi. Manifold regularization and semi-supervised learning: Some theoretical analyses. *Journal of Machine Learning Research*, 14(5), 2013.
- K. Nordström. Convexity of the inverse and moore–penrose inverse. *Linear algebra and its applications*, 434(6):1489–1512, 2011.

- G. Olmschenk, Z. Zhu, and H. Tang. Generalizing semi-supervised generative adversarial networks to regression using feature contrasting. *Computer Vision and Image Understanding*, 186:1–12, 2019.
- S. Oymak and M. Soltanolkotabi. Overparameterized nonlinear learning: Gradient descent takes the shortest path? In *International Conference on Machine Learning*, pages 4951–4960, 2019.
- A. Rahimi and B. Recht. Random features for large-scale kernel machines. *Advances in neural information processing systems*, 20, 2007.
- M. Rahman and M. Ahsanullah. A note on the expected values of powers of a matrix. *The Canadian Journal of Statistics/La Revue Canadienne de Statistique*, pages 123–125, 1973.
- M. Rezagholiradeh and M. A. Haidar. Reg-gan: Semi-supervised learning based on generative adversarial networks for regression. In *2018 IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP)*, pages 2806–2810. IEEE, 2018.
- J. W. Silverstein. Strong convergence of the empirical distribution of eigenvalues of large dimensional random matrices. *Journal of Multivariate Analysis*, 55(2):331–339, 1995.
- G. Simchoni and S. Rosset. Using random effects to account for high-cardinality categorical features and repeated measures in deep neural networks. *Advances in Neural Information Processing Systems*, 34, 2021.
- A. Singh, R. Nowak, and J. Zhu. Unlabeled data: Now it helps, now it doesn’t. In *Advances in neural information processing systems*, pages 1513–1520, 2009.
- K. Sohn, D. Berthelot, N. Carlini, Z. Zhang, H. Zhang, C. A. Raffel, E. D. Cubuk, A. Kurakin, and C.-L. Li. Fixmatch: Simplifying semi-supervised learning with consistency and confidence. *Advances in neural information processing systems*, 33:596–608, 2020.
- Q. Sun and Z. Ge. Deep learning for industrial KPI prediction: When ensemble learning meets semi-supervised data. *IEEE Transactions on Industrial Informatics*, 2020.

- T. Tarpey, R. T. Ogden, E. Petkova, and R. Christensen. A paradoxical result in estimating regression coefficients. *The American Statistician*, 68(4):271–276, 2014.
- G. Vardi and O. Shamir. Implicit regularization in relu networks with the square loss. In *Conference on Learning Theory*, pages 4224–4258. PMLR, 2021.
- L. Wasserman and J. Lafferty. Statistical analysis of semi-supervised regression. *Advances in Neural Information Processing Systems*, 20, 2007.
- L. Xu, C. Hu, and K. Mei. Semi-supervised regression with manifold: A bayesian deep kernel learning approach. *Neurocomputing*, 497:76–85, 2022.
- O. Yuval and S. Rosset. Semi-supervised empirical risk minimization: Using unlabeled data to improve prediction. *Electronic Journal of Statistics*, 16(1):1434–1460, 2022.
- A. Zhang, L. D. Brown, and T. T. Cai. Semi-supervised inference: General theory and estimation of means. *The Annals of Statistics*, 47(5):2538–2566, 2019.
- X. Zhou and M. Belkin. Semi-supervised learning. In *Academic Press Library in Signal Processing*, volume 1, pages 1239–1269. Elsevier, 2014.
- J. Zhu, N. Chen, and E. P. Xing. Bayesian inference with posterior regularization and applications to infinite latent svms. *The Journal of Machine Learning Research*, 15(1): 1799–1847, 2014.
- X. J. Zhu. Semi-supervised learning literature survey. Technical report, University of Wisconsin-Madison Department of Computer Sciences, 2005.