On the uniqueness of co-circular four body central configurations

Manuele Santoprete*

February 21, 2023

Abstract

We study central configurations lying on a common circle in the Newtonian four-body problem. Using a topological argument we prove that there is at most one co-circular central configuration for each cyclic ordering of the masses on the circle.

Contents

1	Introduction	1
2	Central Configurations of the Newtonian <i>n</i> -body problem	4
3	Central Configurations in terms of distances	5
4	Co-circular Central Configurations	7
5	Uniqueness of Co-circular configurations	11

1 Introduction

The Newtonian n-body problem is the study of the dynamics of n point particles with positive masses, moving according to Newton's laws of motion.

^{*}Department of Mathematics, Wilfrid Laurier University E-mail: msantopr@wlu.ca

A central configuration (c.c.) of the *n*-body problem is a configuration of *n* bodies where the acceleration vector of each body is a common scalar multiple of its position vector with respect to the center of mass. The study of central configurations in the Newtonian *n*-body problem has a long history dating back to Euler and Lagrange, and has become an active sub-field of celestial mechanics. While the relative equilibria of the three-body problem have long been known, a complete classification is not known for n > 3. Even the finiteness of central configurations is a hard problem and it was only established in the four-body problem by Hampton and Moeckel [18] and in the five-body problem (except for masses in a codimension two subvariety) by Albouy and Kaloshin [6]. A related problem is the study of central configurations for point vortices. A classification was obtained in the four-vortex problem in the case some of the vorticities are equal [19, 26].

In this paper we focus on a subset of the four-body convex central configurations. A configuration is convex if no body lies inside or on the convex hull formed by the other three bodies. MacMillan and Bartky [21] proved that for any four positive masses and any assigned order, there is at least one convex planar central configuration of the 4-body problem with that order. See also Xia [31] and Moeckel [22] for simpler proofs. Yoccoz [33] conjectured that there is only one such configuration.

Conjecture 1 (Simó-Yoccoz). There is a unique convex planar central configuration of the 4-body problem for each ordering of the masses in the boundary of its convex hull.

This conjecture is also implicit in Simó's paper [28] and, according to Alain Albouy, it is likely the result of several conversations between Simó and Yoccoz, and thus it is reasonable to call it Simó-Yoccoz conjecture. The conjecture was first published, as far as we know, by Albouy and Fu [4], see also (see also [5, 25]), and was included in the well known list of open problems on the classical *n*-body problems compiled by Albouy, Cabral and Santos [3]. Results related to this conjecture were obtained by either putting restrictions on the geometry or restrictions on the masses. In 1932, MacMillan and Bartky [21] already proved uniqueness in the particular case of isosceles trapezoid configurations with two pairs of equal masses located at adjacent vertices of a trapezoid. Similar results were also obtained by Xie [32]. The conjecture is also known to be true if all the masses are equal [1, 2], if two pairs of masses are equal [25, 5, 13], and for the case of three small masses [9]. Some of these results also hold for homogeneous power-law potentials [5, 14].

Our goal in this work is to prove prove the conjecture for the four-body *co-circular central configurations* (c.c.c's), namely those four-body c.c's which lie on a common circle. Specifically we prove the following theorem

Theorem 1. There is at most one co-circular central configuration of four bodies for each cyclic ordering of the masses.

It is known, however, that for most values of the masses there are no co-circular configuration [10]. Although the co-circular four body problem may seem somewhat far-fetched, it is hoped that this work will prove useful in understanding the conjecture for general four-body convex configurations. Furthermore, the co-circular problem has already attracted some attention [10, 16, 17, 20, 11, 1, 7], in part because of the following conjecture, proposed by Alain Chenciner in 2001 [20, 3].

Conjecture 2. Is the regular n-gon with equal masses the unique central configuration such that all the bodies lie on a circle, and the center of mass coincides with the center of the circle?

Note that requiring the bodies to lie on a common circle effectively restricts each body to a one dimensional manifold, allowing for a more straightforward investigation. This situation is reminiscent of the collinear n-body problem, where the bodies lie on a straight line.

The first result concerning uniqueness of central configurations was obtained by Moulton for the collinear problem [23]. Moulton proved that there is a unique collinear central configuration for each ordering of the masses on the line. A topological proof of this fact that uses Morse theory was then provided by Smale [29]. The approach to uniqueness we take in this paper is inspired by the topological approach of Smale.

We will show that, in the four body co-circular problem, the critical points of the potential restricted to a certain subset (that will be described in a subsequent section) are also local minima of the potential. This result is general and not restricted to the case where the center of mass coincides with the center of the circle. The key idea here is to use Ptolemy's theorem to characterize co-circular configuration as done by Cors and Roberts [10]. Once we know that the critical points are local minima, Morse theory can be used to prove Theorem 1. The paper is organized as follows. In Section 2 we introduce the n-body problem and define central configurations. In Section 3 we write central configurations in terms of mutual distances between the bodies. In Section 4 we define co-circular configurations and find their equations following the approach of Cors and Roberts [10]. In particular we view such configurations as critical points of the potential restricted to a certain space that we call \mathcal{M}^+ . In Section 5 we prove Theorem 1 using Morse theory. This is done in four steps. In Proposition 3 we show that all the critical points are nondegenerate local minima. In Lemma 5 we show that the space \mathcal{M}^+ is contractible and we obtain its Euler characteristic. In Lemma 6 we use Morse theory and the Euler characteristic of \mathcal{M}^+ to prove that the potential restricted to \mathcal{M}^+ has a unique critical point. We then use this last result to prove Theorem 1.

2 Central Configurations of the Newtonian *n*-body problem

Let $P_1, P_2, P_3, \ldots, P_n$ be *n* points in \mathbb{R}^d with position vectors $\mathbf{q}_1, \mathbf{q}_2, \ldots, \mathbf{q}_n$. Let $r_{ij} = \|\mathbf{q}_i - \mathbf{q}_j\|$, be the Euclidean distance between the point P_i and P_j , and let $\mathbf{r} = (r_{12}, \ldots, r_{n-1n})$ be the vector of mutual distances. The Newtonian *n*-body problem concerns the motion of *n* particles with positive masses $m_i > 0$ and positions $\mathbf{q}_i \in \mathbb{R}^d$, where $i = 1, \ldots, n$. The center of mass of the system is fixed at the origin of the coordinate systems, that is, we have $\mathbf{q}_{CM} = \frac{1}{M} \sum_{i=1}^n m_i \mathbf{q}_i = 0$, where $M = m_1 + \ldots + m_n$ is the total mass. The motion is governed by Newton's law of motion

$$m_i \ddot{\mathbf{q}}_i = \sum_{i \neq j} \frac{m_i m_j (\mathbf{q}_j - \mathbf{q}_i)}{r_{ij}^3} = \frac{\partial U}{\partial \mathbf{q}_i}, \quad 1 \le i \le n$$
(1)

where $\tilde{U}(\mathbf{q})$ is the Newtonian potential

$$\tilde{U}(\mathbf{q}) = \sum_{i < j} \frac{m_i m_j}{\|\mathbf{q}_i - \mathbf{q}_j\|}, \quad 1 \le i \le n.$$
(2)

Let us denote the Newtonian potential by $U(\mathbf{r})$ when viewed as a function of \mathbf{r} . A central configuration (c.c.) of the *n*-body problem is a configuration $\mathbf{q} \in \mathbb{R}^{nd}$ which satisfies the algebraic equations

$$\lambda m_i \mathbf{q}_i = \sum_{i \neq j} \frac{m_i m_j (\mathbf{q}_j - \mathbf{q}_i)}{r_{ij}^3}, \quad 1 \le i \le n.$$
(3)

The central configuration equation (3) is invariant under rotations, reflections and dilations. It is standard to say that two configurations \mathbf{q} and \mathbf{q}' are equivalent if there is a non-zero constant $k \in \mathbb{R}$ and an orthogonal matrix R such that $\mathbf{q}'_i = kR\mathbf{q}_i$, $i = 1, \ldots n$. This defines an equivalence relation \sim' , and thus one can speak of equivalence classes of central configurations. By convention, when counting central configurations it is standard to count the number of equivalence classes with respect to the equivalence relation \sim' . This convention is also used in the statement of Conjecture 1 and of Theorem 1.

Let $I(\mathbf{q})$ denote the moment of inertia, that is,

$$\tilde{I}(\mathbf{q}) = \frac{1}{2} \sum_{i=1}^{n} m_i \|\mathbf{q}_i\|^2$$

and let $I(\mathbf{r}) = \frac{1}{2M} \sum_{1 \le i < j \le n}^{n} m_i m_j r_{ij}^2$ denote the moment of inertia when viewed as a function of \mathbf{r} .

Using the moment of inertia we can write equation (3) as

$$\nabla_{\mathbf{q}}\tilde{U}(\mathbf{q}) = \lambda \,\nabla_{\mathbf{q}}\tilde{I}(\mathbf{q}),\tag{4}$$

where $\nabla_{\mathbf{q}} = \left[\frac{\partial}{\partial \mathbf{q}_1}, \dots, \frac{\partial}{\partial \mathbf{q}_n}\right]^T$. Viewing λ as a Lagrange multiplier, we have that \mathbf{q} is a central configuration if and only if it is a critical point (with respect to \mathbf{q}) of $\tilde{U}(\mathbf{q})$ subject to the constraint $\tilde{I} = \tilde{I}_0$.

For any configuration \mathbf{q} the vectors \mathbf{q}_j span a subspace $\mathcal{C}(\mathbf{q})$ of \mathbb{R}^d called the *centered position space* [22]. It is natural to define the dimension of a configuration to be dim(\mathbf{q}) = dim($\mathcal{C}(\mathbf{q})$). We say that \mathbf{q} is a *Dziobek configurations* if dim(\mathbf{q}) = n - 2 [22]. In the four-body problem \mathbf{q} is a *Dziobek central configuration* if it is a central configuration with dim(\mathbf{q}) = 2. The set of four-body Dziobek configurations coincides with the set of planar, non-collinear, central configurations.

3 Central Configurations in terms of distances

For four bodies it is convenient to recast the equations defining Dziobek central configuration, so that the variables are the distances between the particles rather than their coordinates. Since the mutual distances determine the configuration up to rotation and reflection symmetry, this choice not only reduces the number of variables but also removes the rotational and reflectional degeneracy. The dilational degeneracy can then be eliminated by fixing the size of the configuration with the restriction I = 1.

Let $\mathbf{r} = (r_{12}, r_{13}, r_{14}, r_{23}, r_{24}, r_{34}) \in (\mathbb{R}^+)^6$ be a vector of non-negative mutual distances, and let the Cayley–Menger determinant of four points P_1, \ldots, P_4 be

$$H(\mathbf{r}) = 288V^2 = \begin{vmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & r_{12}^2 & r_{13}^2 & r_{14}^2 \\ 1 & r_{12}^2 & 0 & r_{23}^2 & r_{24}^2 \\ 1 & r_{13}^2 & r_{23}^2 & 0 & r_{34}^2 \\ 1 & r_{14}^2 & r_{24}^2 & r_{34}^2 & 0 \end{vmatrix}$$

where V is the volume of the configuration. Not all vectors \mathbf{r} realize actual configurations of four bodies in \mathbb{R}^3 (see [10] for some examples). A necessary and sufficient condition for a given vector \mathbf{r} to correspond to an actual configuration of four bodies is that $H(\mathbf{r}) \geq 0$ and all strict triangle inequalities be satisfied. As a consequence we consider the sets

$$\mathcal{G} = \{ \mathbf{r} \in (\mathbb{R}^+)^6 | H(\mathbf{r}) \ge 0 \text{ and } r_{ij} + r_{jk} > r_{ik} \text{ for all } (i, j, k) \text{ where } i \neq j \neq k \}.$$

and

$$\mathcal{N} = \{ \mathbf{r} \in \mathcal{G} | I(\mathbf{r}) - 1 = 0, \quad H(\mathbf{r}) = 0 \}$$

We say that a vector of mutual distances \mathbf{r} is geometrically realizable if $\mathbf{r} \in \mathcal{G}$ and that \mathbf{r} is a normalized Dziobek configuration if $\mathbf{r} \in \mathcal{N}$.

Thus we have the following characterization of planar four body central configurations given by Dziobek:

Proposition 1. Let \mathbf{q} be a Dziobek configuration, let $\mathbf{r} \in \mathcal{N}$ be its corresponding normalized Dziobek configuration, and let $U|_{\mathcal{N}} : \mathcal{N} \to \mathbb{R}$ be the restriction of the Newtonian potential U to \mathcal{N} . Then, \mathbf{q} is a Dziobek central configuration if and only if \mathbf{r} is a critical point of $U|_{\mathcal{N}}$ with respect to \mathbf{r} .

Since equations (3) are invariant under rotations, dilations and reflections in the plane, we can consider two relative equilibria as equivalent if they are related by these symmetry operations. This defines an equivalence relation \sim , different from the more standard one introduced in section 2. Let X be the set of equivalence classes with respect to \sim , then the set of equivalence classes X is in a one-to-one correspondence with the set $c(U|_{\mathcal{N}})$ of critical points of the function $U(\mathbf{r})|_{\mathcal{N}}$.

4 Co-circular Central Configurations

In this section we study co-circular central configurations. For a planar configuration we say that the bodies are *ordered counterclockwise* (*clockwise*) if they are numbered consecutively while traversing the boundary of the quadrilateral in a countercklockwise (clockwise) direction. Since we use mutual distances as coordinates, we cannot distinguish between bodies ordered counterclockwise and bodies ordered clockwise. Hence, we introduce the following terminology: we say that the bodies are *ordered sequentially* if they are numbered consecutively while traversing the boundary of the quadrilateral in any direction.

Without loss of generality, we may assume that any cyclic quadrilateral is ordered sequentially so that r_{13} and r_{24} are the lengths of the diagonals. This is justified because we can always relabel the bodies so that they are ordered sequentially. Denote

$$P(\mathbf{r}) = r_{12}r_{34} + r_{14}r_{23} - r_{13}r_{24}.$$

Ptolemy's theorem states that if a quadrilateral is sequentially ordered and cyclic then P = 0. More in general, Ptolemy's inequality says that $P \ge 0$ for any convex quadrilateral ordered sequentially and for any tetrahedron [8]. Equality holds if and only if the four bodies are co-circular. Let \mathcal{P} be the set of geometrically realizable \mathbf{r} satisfying $P(\mathbf{r}) = 0$, that is

$$\mathcal{P} = \{ \mathbf{r} \in \mathcal{G} | P(\mathbf{r}) = 0 \},\$$

let \mathcal{M} be the set of **r** which satisfy I = 1 and P = 0

$$\mathcal{M} = \{ \mathbf{r} \in \mathbb{R}^6 | I(\mathbf{r}) - 1 = 0, \quad P(\mathbf{r}) = 0 \},$$

and let \mathcal{M}^+ be the set obtained from \mathcal{M} by reintroducing the restrictions on the mutual distances

$$\mathcal{M}^+ = \{ \mathbf{r} \in (\mathbb{R}^+)^6 | I(\mathbf{r}) - 1 = 0, \quad P(\mathbf{r}) = 0 \}.$$

It is easy to see that \mathcal{M} is a smooth four-dimensional manifold, since the gradients of I and P are independent, and that \mathcal{M}^+ is a manifold with boundary. In a later section we will show that $\mathcal{M} \approx S^2 \times S^2$ and that \mathcal{M}^+ is homeomorphic to a closed ball. We denote the boundary of \mathcal{M}^+ by $\partial \mathcal{M}^+$ and observe that $U|_{\partial \mathcal{M}^+} = \infty$.

One last set that will play an important role in this paper is \mathcal{D} , which is defined as follows:

$$\mathcal{D} = \{ \mathbf{r} \in \mathcal{G} | I(\mathbf{r}) - 1 = 0, P(\mathbf{r}) = 0, H(\mathbf{r}) = 0 \}$$
$$= \{ \mathbf{r} \in \mathcal{M}^+ \cap \mathcal{G} | H(\mathbf{r}) = 0 \}.$$

There is an interesting relationship between the conditions $P(\mathbf{r}) = 0$ and $H(\mathbf{r}) = 0$. The following Lemma sheds some light on this relationship, further insight is given in Lemma 2.

Lemma 1. If $\mathbf{r} \in \mathcal{P}$, then H(r) = 0. In other words on the set of geometrically realizable vectors for which P = 0 the configuration of four bodies is coplanar. Moreover, we have that $\mathcal{D} = \mathcal{M}^+ \cap \mathcal{G}$.

Proof. Pech showed [24] that the Cayley-Menger determinant can be written as follows

$$\frac{1}{2}H(\mathbf{r}) = P(\mathbf{r}) \cdot Q(\mathbf{r}) - K^2(\mathbf{r})$$

where

$$Q(\mathbf{r}) = r_{12}r_{34}(-r_{12}^2 - r_{34}^2 + r_{23}^2 + r_{14}^2 + r_{13}^2 + r_{24}^2) + r_{14}r_{23}(r_{12}^2r_{34}^2 - r_{23}^2 - r_{14}^2 + r_{13}^2 + r_{24}^2) - r_{13}r_{24}(r_{12}^2 + r_{34}^2 + r_{23}^2 + r_{14}^2 - r_{13}^2 - r_{24}^2)$$

and

$$K(\mathbf{r}) = r_{12}r_{13}r_{23} - r_{12}r_{14}r_{24} + r_{13}r_{14}r_{34} - r_{23}r_{24}r_{34}$$

If P = 0, then

$$\frac{1}{2}H(\mathbf{r}) = -(K(\mathbf{r}))^2 \le 0.$$

Since $\mathbf{r} \in \mathcal{G}$ implies $H(\mathbf{r}) \ge 0$ it follows that $K(\mathbf{r}) = H(\mathbf{r}) = 0$.

A similar relationship exists between $H(\mathbf{r}) = 0$ and the condition required of four points to form a trapezoid. This relationship was exploited in [27] to obtain equations for trapezoidal central configurations.

Since co-circular central configurations are Dziobek configuration we can give the following definition

Definition 1. The configuration vector \mathbf{q} is a sequentially ordered cyclic four-body central configuration if and only if its corresponding distance vector \mathbf{r} belongs to \mathcal{D} and it is a critical point of $U|_{\mathcal{N}}$ with respect to \mathbf{r} .

In terms of Lagrange multipliers this means that $\mathbf{r} \in \mathcal{D}$ is a sequentially ordered cyclic four body central configuration if and only if it is a critical point of the function

$$U(\mathbf{r}) + \lambda M(I(\mathbf{r}) - 1) + \eta H(\mathbf{r}),$$

satisfying I - 1 = 0, P = 0 and H = 0, where λ , and η are Lagrange multipliers. The downside of this approach is that H and its derivatives are fairly complicated. Using the following lemma however, it is possible to find simpler equations for the co-circular configurations. This lemma was proven in [10], here we provide a different proof.

Lemma 2. For any $\mathbf{r} \in \mathcal{P}$

$$\nabla_{\mathbf{r}} H(\mathbf{r}) = 2Q(\mathbf{r})\nabla_{\mathbf{r}} P(\mathbf{r})$$

where $\nabla_{\mathbf{r}} = \left[\frac{\partial}{\partial r_{12}}, \ldots, \frac{\partial}{\partial r_{34}}\right]^T$. In other words on the set of geometrically realizable vectors for which P vanish, the gradients of H and P are parallel.

Proof. Since $\frac{1}{2}H(\mathbf{r}) = P(\mathbf{r}) \cdot Q(\mathbf{r}) - K^2(\mathbf{r})$ we have that

$$\frac{1}{2}\nabla_{\mathbf{r}}H(\mathbf{r}) = Q(\mathbf{r})\nabla_{\mathbf{r}}P(\mathbf{r}) + P(\mathbf{r})\nabla_{\mathbf{r}}Q(\mathbf{r}) - 2K(\mathbf{r})\nabla_{\mathbf{r}}K(\mathbf{r}).$$

Since $\mathbf{r} \in \mathcal{P}$, then H = P = 0. It follows that K = 0 as well. Hence, $\frac{1}{2} \nabla_{\mathbf{r}} H(\mathbf{r}) = Q(\mathbf{r}) \nabla_{\mathbf{r}} P(\mathbf{r}).$

Comparing this result with the corresponding lemma in [10] we find that if $\mathbf{r} \in \mathcal{P}$ then

$$2Q(\mathbf{r}) = \left(\frac{4}{r_c^2} \prod_{i < j} r_{ij}\right) \neq 0,$$

where r_c is the circumradius of the cyclic quadrilateral.

We now have the following characterization of co-circular configurations due to Cors and Roberts ([10]):

Proposition 2. Let $\mathbf{r} \in \mathcal{D}$, then \mathbf{r} is a critical point of $U|_{\mathcal{N}}$, the restriction of U to \mathcal{N} , if and only if \mathbf{r} is a critical point of the function $U|_{\mathcal{M}^+} : \mathcal{M}^+ \to \mathbb{R}$. Therefore the vector \mathbf{q} is a sequentially ordered cyclic four-body c.c. if and

only if the corresponding distance vector $\mathbf{r} \in \mathcal{D}$ is a critical point of the Lagrangian function

$$L(\mathbf{r}; \lambda, \sigma) = U(\mathbf{r}) + \lambda M \left(I(\mathbf{r}) - 1 \right) + \sigma P(\mathbf{r})$$

satisfying I - 1 = 0, P = 0 and H = 0, where λ and σ are Lagrange multipliers.

Proof. Recall that $\nabla_{\mathbf{r}} U|_{\mathcal{M}^+}$ is the orthogonal projection of $\nabla_{\mathbf{r}} U(\mathbf{r})$ onto the tangent space $T_{\mathbf{r}}(\mathcal{M}^+)$, and similarly $\nabla_{\mathbf{r}} U|_{\mathcal{N}}$ is the orthogonal projection of $\nabla_{\mathbf{r}} U(\mathbf{r})$ onto the tangent space $T_{\mathbf{r}}(\mathcal{N})$, with

$$T_{\mathbf{r}}\mathcal{N} = \{ \mathbf{v} \in \mathbb{R}^6 \, | \, \nabla_{\mathbf{r}}(I(\mathbf{r}) - 1) \cdot \mathbf{v} = 0, \nabla_{\mathbf{r}}H(\mathbf{r}) \cdot \mathbf{v} = 0 \}$$

and

$$T_{\mathbf{r}}\mathcal{M}^{+} = \{ \mathbf{v} \in \mathbb{R}^{6} \, | \, \nabla_{\mathbf{r}}(I(\mathbf{r}) - 1) \cdot \mathbf{v} = 0, \nabla_{\mathbf{r}}P(\mathbf{r}) \cdot \mathbf{v} = 0 \}.$$

Since $\mathbf{r} \in \mathcal{D}$, by Lemma 2, $\nabla_{\mathbf{r}} H(\mathbf{r}) = 2Q(\mathbf{r})\nabla_{\mathbf{r}} P(\mathbf{r})$. It follows that, if $\mathbf{r} \in \mathcal{D}$, then $T_{\mathbf{r}}\mathcal{M}^+ = T_{\mathbf{r}}\mathcal{N}$, and hence $\nabla_{\mathbf{r}} U|_{\mathcal{N}} = \nabla_{\mathbf{r}} U|_{\mathcal{M}^+}$ for any $\mathbf{r} \in \mathcal{D}$. Then $\nabla_{\mathbf{r}} U|_{\mathcal{M}^+} = 0$ if and only if $\nabla_{\mathbf{r}} U|_{\mathcal{N}} = 0$, that is, \mathbf{r} is a critical point of $U|_{\mathcal{N}}$ if and only if \mathbf{r} is a critical point of the function $U|_{\mathcal{M}^+}$.

Proposition 2 says that if $\mathbf{r} \in \mathcal{D}$, we can find the critical points of $U|_{\mathcal{N}}$ by finding the critical points of $U|_{\mathcal{M}^+}$. The equations of the critical points of $U|_{\mathcal{M}^+} : \mathcal{M}^+ \to \mathbb{R}$, are given by $\nabla_{\mathbf{r}} L(\mathbf{r}; \lambda, \sigma) = \nabla_{\mathbf{r}} U + \lambda M \nabla_{\mathbf{r}} I + \sigma \nabla_{\mathbf{r}} P$, the gradient of the Lagrangian L. Explicitly, we have

$$m_1 m_2 (r_{12}^{-3} - \lambda) = \sigma \, \frac{r_{34}}{r_{12}} \qquad \qquad m_3 m_4 (r_{34}^{-3} - \lambda) = \sigma \, \frac{r_{12}}{r_{34}} \tag{5}$$

$$m_1 m_3 (r_{13}^{-3} - \lambda) = -\sigma \frac{r_{24}}{r_{13}} \qquad m_2 m_4 (r_{24}^{-3} - \lambda) = -\sigma \frac{r_{13}}{r_{24}} \qquad (6)$$

$$m_1 m_4 (r_{14}^{-3} - \lambda) = \sigma \frac{r_{23}}{r_{14}} \qquad m_2 m_3 (r_{23}^{-3} - \lambda) = \sigma \frac{r_{14}}{r_{23}}.$$
 (7)

It is important to observe that these equations hold for $\mathbf{r} \in \mathcal{M}^+$, and not just for $\mathbf{r} \in \mathcal{D}$, however, if $\mathbf{r} \notin \mathcal{D}$ then these equations do not give central configurations. Since $\mathbf{r} \in \mathcal{M}^+$, the constraints I - 1 = 0 and P = 0 must be satisfied, but H = 0 is not required. When $\mathbf{r} \in \mathcal{D} \subset \mathcal{M}^+$, however, H = 0and the solutions of these equations give co-circular central configurations. The equations have been grouped in pairs so that when they are multiplied together the product of the right-hand sides is σ^2 . This yields the well-known relation of Dziobek [12]

$$(r_{12}^{-3} - \lambda)(r_{34}^{-3} - \lambda) = (r_{13}^{-3} - \lambda)(r_{24}^{-3} - \lambda) = (r_{14}^{-3} - \lambda)(r_{23}^{-3} - \lambda), \quad (8)$$

which is required of any 4-body Dziobek central configuration. From equations (5),(6) and (7) we obtain three equations for σ^2 :

$$\sigma^2 = m_1 m_2 m_3 m_4 (r_{12}^{-3} - \lambda) (r_{34}^{-3} - \lambda)$$
(9)

$$\sigma^2 = m_1 m_2 m_3 m_4 (r_{14}^{-3} - \lambda) (r_{23}^{-3} - \lambda) \tag{10}$$

$$\sigma^2 = m_1 m_2 m_3 m_4 (r_{13}^{-3} - \lambda) (r_{24}^{-3} - \lambda).$$
(11)

5 Uniqueness of Co-circular configurations

In this section we want to prove Theorem 1. We break down the proof in four steps, which we summarize here.

- 1. We show that if $\mathbf{r} \in \mathcal{M}^+$ is a critical point of $U|_{\mathcal{M}^+}$, then it is necessarily a nondegenerate local minimum. This is proved in Proposition 3. Lemma 3 is a technical lemma required to prove Proposition 3.
- 2. We study the topology of \mathcal{M} and \mathcal{M}^+ . In Lemma 4 we show that $\mathcal{M} \approx S^2 \times S^2$. In Lemma 5 we show that \mathcal{M}^+ is contractible and the Euler characteristic $\chi(\mathcal{M}^+)$ of \mathcal{M}^+ is 1.
- 3. We use Morse theory to prove that the function $U|_{\mathcal{M}^+}$ has a unique critical point on \mathcal{M}^+ . This is done in Lemma 6.
- 4. We prove the theorem.

We start with the following technical lemma which is needed in the proof of Proposition 3.

Lemma 3. If $\mathbf{r}^* \in \mathcal{M}^+$ is a critical point of $U|_{\mathcal{M}^+}$ then $\lambda > 0$.

Proof. Suppose, for the sake of contradiction, that $\lambda \leq 0$. By the first of the two equation (5) we find that

$$\sigma \frac{r_{34}}{r_{12}} = m_1 m_2 (r_{12}^{-3} - \lambda) > 0$$

and hence $\sigma > 0$, since $r_{12}, r_{34} > 0$ in \mathcal{M}^+ . By the first of the two equation (6) we find that

$$-\sigma \frac{r_{24}}{r_{13}} = m_1 m_3 (r_{13}^{-3} - \lambda) > 0$$

and hence $\sigma < 0$, which contradicts the fact that $\lambda \leq 0$. Hence, $\lambda > 0$.

Note that the second derivative $D^2 L(\mathbf{r}; \lambda, \sigma)$ of $L(\cdot; \lambda, \sigma)$ with respect to the variable \mathbf{r} is the matrix

$$D^{2}L(\mathbf{r};\lambda,\sigma) = D^{2}U(\mathbf{r}) + \lambda M D^{2}I(\mathbf{r}) + \sigma D^{2}P(\mathbf{r}).$$

If **r** is a critical point of $U|_{\mathcal{M}^+}$ this second derivative, with appropriate choices of λ and σ , is the second derivative of $U|_{\mathcal{M}^+}$, the restriction of U to \mathcal{M}^+ . We can now prove the following proposition

Proposition 3. If $\mathbf{r}^* \in \mathcal{M}^+$ is a critical point of $U|_{\mathcal{M}^+}$ then \mathbf{r}^* is a nondegenerate minimum point for $U|_{\mathcal{M}^+}$.

Proof. The second derivative of L is the matrix

$$D^{2}L(\mathbf{r};\lambda,\sigma) = \operatorname{diag}(f_{12}(\mathbf{r}), f_{13}(\mathbf{r}), f_{14}(\mathbf{r}), f_{23}(\mathbf{r}), f_{24}(\mathbf{r}), f_{34}(\mathbf{r}))$$

+ adiag(\sigma, -\sigma, \sigma, -\sigma, \sigma)

where $f_{ij}(\mathbf{r}) = m_i m_j (2r_{ij}^{-3} + \lambda)$. Here, diag $(f_{12}, f_{13}, f_{14}, f_{23}, f_{24}, f_{34})$ denotes the 6 × 6 diagonal matrix whose diagonal entries are $f_{12}, f_{13}, f_{14}, f_{23}, f_{24}, f_{34}$. Similarly, adiag $(\sigma, -\sigma, \sigma, \sigma, -\sigma, \sigma)$ denotes the 6 × 6 anti-diagonal matrix whose anti-diagonal entries, starting from the upper right corner, are $\sigma, -\sigma, \sigma, \sigma, -\sigma, \sigma, \sigma, -\sigma, \sigma$.

Let $\Delta_k(\mathbf{r})$ be the principal minor of order k of $D^2L(\mathbf{r}; \lambda, \sigma)$. We first prove that if \mathbf{r}^* satisfies equations (5-7), then $\Delta_k(\mathbf{r}^*) > 0$ for k = 1, ...6. Since $\lambda > 0$ by Lemma 3 it is easy to see that the first three principal minors are always positive

$$\begin{split} \Delta_1(\mathbf{r}) &= \frac{(\lambda r_{12}^3 + 2)m_1m_2}{r_{12}^3} > 0\\ \Delta_2(\mathbf{r}) &= \frac{(\lambda r_{12}^3 + 2)(\lambda r_{13}^3 + 2)m_1^2m_2m_3}{r_{12}^3r_{13}^3} > 0\\ \Delta_3(\mathbf{r}) &= \frac{(\lambda r_{12}^3 + 2)(\lambda r_{13}^3 + 2)(\lambda r_{14}^3 + 2)m_1^3m_2m_3m_4}{r_{12}^3r_{13}^3r_{14}^3} > 0. \end{split}$$

Let

$$A_{0}(\mathbf{r}) = m_{1}m_{2}m_{3}m_{4}(\lambda^{2}r_{12}^{3}r_{34}^{3} + 2\lambda r_{12}^{3} + 2\lambda r_{34}^{3} + 4) - r_{12}^{3}r_{34}^{3}\sigma^{2}$$

$$A_{1}(\mathbf{r}) = m_{1}m_{2}m_{3}m_{4}(\lambda^{2}r_{14}^{3}r_{23}^{3} + 2\lambda r_{14}^{3} + 2\lambda r_{23}^{3} + 4) - r_{14}^{3}r_{23}^{3}\sigma^{2}$$

$$A_{2}(\mathbf{r}) = m_{1}m_{2}m_{3}m_{4}(\lambda^{2}r_{13}^{3}r_{24}^{3} + 2\lambda r_{13}^{3} + 2\lambda r_{24}^{3} + 4) - r_{13}^{3}r_{24}^{3}\sigma^{2}$$

then the remaining principal minors are:

$$\Delta_4(\mathbf{r}) = \frac{m_1^2 m_2 m_3}{r_{13}^3 r_{14}^3 r_{23}^3} (2r_{12}^{-3} + \lambda) (\lambda r_{13}^3 + 2) A_1$$

$$\Delta_5(\mathbf{r}) = \frac{m_1 m_2}{r_{13}^3 r_{14}^3 r_{23}^3 r_{24}^3} (2r_{12}^{-3} + \lambda) A_1 A_2$$

$$\Delta_6(\mathbf{r}) = \frac{1}{r_{12}^3 r_{13}^3 r_{14}^3 r_{23}^3 r_{24}^3 r_{34}^3} A_0 A_1 A_2.$$

Since $\lambda > 0$, eliminating σ^2 from A_0 using (9), from A_1 using (10), and from A_2 using (11), yields

$$A_0(\mathbf{r}^*) = 3m_1m_2m_3m_4(\lambda r_{12}^3 + \lambda r_{34}^3 + 1) > 0$$

$$A_1(\mathbf{r}^*) = 3m_1m_2m_3m_4(\lambda r_{14}^3 + \lambda r_{23}^3 + 1) > 0$$

$$A_2(\mathbf{r}^*) = 3m_1m_2m_3m_4(\lambda r_{13}^3 + \lambda r_{24}^3 + 1) > 0.$$

Consequently, we have that $\Delta_k(\mathbf{r}^*) > 0$ for k = 4, 5, 6. Since all the principal minors are positive it follows that $D^2L(\mathbf{r}^*, \lambda, \sigma)$ is positive definite and \mathbf{r}^* is a nondegenerate local minimum of $U|_{\mathcal{M}^+}$.

Remark. By Proposition 2 we see that the gradient of $U|_{\mathcal{M}^+}$ and the gradient of $U|_{\mathcal{N}}$ coincide for any $\mathbf{r} \in \mathcal{D}$. However, the second derivative of $U|_{\mathcal{M}^+}$ is in general different from the second derivative of $U|_{\mathcal{N}}$, because the fact that P and H are tangent at the critical points does not ensure that the quadratic approximation at those points is the same. Furthermore, the property concerning the gradients of $U|_{\mathcal{M}^+}$ and $U|_{\mathcal{N}}$ given in Proposition 2 holds on \mathcal{D} and not on the larger set \mathcal{M}^+ . Consequently, when looking at the second derivatives it is important to be careful to consider carefully the various restrictions of U.

We now turn to study the topology of \mathcal{M} .

Lemma 4. $\mathcal{M} \approx \operatorname{Gr}_+(2,4) \approx S^2 \times S^2$.

Proof. Consider the following change of coordinates:

$$r_{ij} = \left(\frac{2M}{(m_i m_j)}\right)^{1/2} p_{ij}, \quad i, j \in \{1, 2, 3, 4 \text{ with } i < j\}$$

In these coordinates the equations $I(\mathbf{r}) = 1$ and $P(\mathbf{r}) = 0$ take the form

$$p_{12}^2 + p_{13}^2 + p_{14}^2 + p_{23}^2 + p_{24}^2 + p_{34}^2 = 1$$
(12)

$$\left(\frac{2M}{\sqrt{m_1 m_2 m_3 m_4}}\right) \left(p_{12} p_{34} - p_{13} p_{24} + p_{14} p_{23}\right) = 0.$$
(13)

These are the equations of the oriented Grassmanian

$$Gr_+(2,4) = SO(4)/(SO(2) \times SO(2)),$$

(see [30] for details). Equations (12-13) are equivalent to the system

$$(p_{12} + p_{34})^2 + (p_{13} - p_{24})^2 + (p_{14} + p_{23})^2 = 1$$
(14)

$$(p_{12} - p_{34})^2 + (p_{13} + p_{24})^2 + (p_{14} - p_{23})^2 = 1,$$
(15)

which shows that \mathcal{M} is diffeomorphic to $S^2 \times S^2$.

We can now better understand the topology of \mathcal{M}^+ . Discussions with Shengda Hu were very helpful with this next lemma.

Lemma 5. \mathcal{M}^+ is contractible and its Euler characteristic $\chi(\mathcal{M}^+)$ is 1.

Proof. After the change of variable

$$\begin{aligned} v_1 &= p_{12} + p_{34} & v_2 &= p_{13} - p_{24} & v_3 &= p_{14} + p_{23} \\ w_1 &= p_{12} - p_{34} & w_2 &= p_{13} + p_{24} & w_3 &= p_{14} - p_{23}. \end{aligned}$$

equations (14) and (15) can be rewritten in the form

$$S_1 = \{ v = (v_1, v_2, v_3) \in \mathbb{R}^3 : v_1^2 + v_2^2 + v_3^2 = 1 \}, S_2 = \{ w = (w_1, w_2, w_3) \in \mathbb{R}^3 : w_1^2 + w_2^2 + w_3^2 = 1 \}.$$

Clearly the set \mathcal{M}^+ is homeomorphic to E, the subset of $S_1 \times S_2$ defined by the following inequalities

$$p_{12} = \frac{v_1 + w_1}{2} \ge 0 \qquad p_{13} = \frac{v_2 + w_2}{2} \ge 0 \qquad p_{14} = \frac{v_3 + w_3}{2} \ge 0$$
$$p_{34} = \frac{v_1 - w_1}{2} \ge 0 \qquad p_{24} = \frac{w_2 - v_2}{2} \ge 0 \qquad p_{23} = \frac{v_3 - w_3}{2} \ge 0.$$

These inequalities can be expressed in a more compact form as

$$v_1 \ge |w_1|, \quad v_3 \ge |w_3|, \quad w_2 \ge |v_2|,$$

and on $S_1 \times S_2$ they reduce to

$$v_1 \ge |w_1|, \quad v_3 \ge |w_3|, \quad w_2 \ge 0.$$
 (16)

This can be explained as follows. The first two inequalities in (16) imply that $|v_1|^2 + |v_3|^2 \ge |w_1|^2 + |w_3|^2$, which gives

$$|w_2| = \sqrt{1 - |w_1|^2 - |w_3|^2} \ge \sqrt{1 - |v_1|^2 - |v_3|^2} = |v_2|.$$

Hence $w_2 > |v_2|$ since $w_2 \ge 0$.

The last inequality in (16), namely $w_2 \ge 0$, selects the closed upper hemisphere H_2^+ of S_2 . The hemisphere H_2^+ is homeomorphic to a closed disk and any point on it can be represented with coordinates (w_1, w_3) in $\{(w_1, w_3) \in \mathbb{R}^2 | w_1^2 + w_3^2 \le 1\}$.

Corresponding to each point of coordinates (w_1, w_3) , there is a region F of the sphere S_1 determined by the inequalities

$$v_1 \ge |w_1|, \quad v_3 \ge |w_3|.$$

If $(w_1, w_3) = (\pm 1, 0)$ then $(v_1, v_2, v_3) = (1, 0, 0)$. If $(w_1, w_3) = (0, \pm 1)$ then $(w_1, w_2, w_3) = (0, 0, 1)$. Hence, in these cases the region F reduces to a point. For any other value of (w_1, w_3) the region F is homeomorphic to a closed 2-disk. The restriction of the projection $\tilde{p} : (v_1, v_2, v_3, w_1, w_2, w_3) \rightarrow (w_1, w_2, w_3)$, induces a fibration $p : E \rightarrow H_2^+$ with base space H_2^+ and fibers given by F. Thus, the projection p is a fibration with contractible fibers. Since H_2^+ is also contractible, we see that the space E is contractible. In particular, $\chi(\mathcal{M}^+) = \chi(E) = 1$.

Remark. The previous Lemma, and a much more general thorem, seems to follow from a recent result by Galashin, Karp, and Lam [15]. Let Gr(k, n) denote the Grassmannian of k-planes in \mathbb{R}^n , its totally nonnegative part $Gr_{\geq}(k, n)$ is defined to be the set of $x \in Gr(k, n)$ whose Plücker coordinates are all nonnegative. It has been shown that $Gr_{\geq}(k, n)$ is homeomorphic to a k(n-k) dimensional closed ball [15]. Using this result it should be possible to show that the oriented Grassmanian $Gr_+(n, k)$ with all the Plücker coordinates nonnegative, which we may call totally nonnegative oriented Grassmanian, is a k(n-k) dimensional closed ball. This would generalize the previous lemma to any oriented Grassmanian $Gr_+(n, k)$.

Since we have determined the topology of \mathcal{M}^+ we can now use Morse theory to prove the following Lemma

Lemma 6. The function $U|_{\mathcal{M}^+}$ has a unique critical point on \mathcal{M}^+ .

Proof. By Proposition 3 any critical point $\mathbf{r} \in \mathcal{M}^+$ is a nondegenerate local minimum of the function $U|_{\mathcal{M}^+}$, and hence $U|_{\mathcal{M}^+}$ is a Morse function that approaches $+\infty$ as \mathbf{r} approaches $\partial \mathcal{M}^+$, the boundary of \mathcal{M}^+ . Therefore, the function $U|_{\mathcal{M}^+}$ admits a global minimum value in the interior of \mathcal{M}^+ . Suppose there are several global minimum points where the function obtains its least possible value. By Proposition 3 any of such point must be a nondegenerate local minimum point. By Lemma 5, the Euler characteristic of \mathcal{M}^+ is $\chi(\mathcal{M}^+) = 1$. By Morse theory we have

$$1 = \chi(\mathcal{M}^+) = \sum (-1)^{\gamma} C^{\gamma} \tag{17}$$

where the sum is over the critical points, γ is the Morse index of the critical points and C^{γ} is the number of critical points of index γ . We know that there is at least one local minimum, and that all the critical points of $U|_{\mathcal{M}^+}$ are local minimum points and hence have index 0. However, this function cannot have more than one minimum point since otherwise, equation (17) would imply the existence of at least one non-minimum critical point, contradicting Proposition 3.

We are now in a position to prove Theorem 1, our main result

Proof of Theorem 1. Recall that, by Proposition 3, co-circular central configurations correspond to distance vectors $\mathbf{r} \in \mathcal{D}$ that are critical points of the function $U|_{\mathcal{M}^+}$. Lemma 6 shows that $U|_{\mathcal{M}^+}$ has a unique critical point on \mathcal{M}^+ . Since $\mathcal{D} \subset \mathcal{M}^+$, there is at most one critical point on \mathcal{D} . Recall that if \mathbf{q} and \mathbf{q}' can be transformed one into the other with a reflection than they are mapped to the same distance vector \mathbf{r} . Hence, we have shown that there is a most one equivalence class (with respect to the equivalence relation \sim') of co-circular central configurations for each ordering of the masses, and the theorem follows.

Acknowledgments

I would like to thank Alain Albouy, Shengda Hu, Steven Karp, Santiago López de Medrano, and Alessandro Portaluri for interesting discussions on this work. This work was supported by an NSERC discovery grant.

References

- Alain Albouy. Symétrie des configurations centrales de quatre corps. Comptes rendus de l'Académie des sciences. Série 1, Mathématique, 320(2):217-220, 1995.
- [2] Alain Albouy. The symmetric central configurations of four equal masses. *Contemporary Mathematics*, 198:131–136, 1996.
- [3] Alain Albouy, Hildeberto E Cabral, and Alan A Santos. Some problems on the classical n-body problem. *Celestial Mechanics and Dynamical Astronomy*, 113(4):369–375, 2012.
- [4] Alain Albouy and Yanning Fu. Euler configurations and quasipolynomial systems. *Regular and Chaotic Dynamics*, 12(1):39–55, 2007.
- [5] Alain Albouy, Yanning Fu, and Shanzhong Sun. Symmetry of planar four-body convex central configurations. In *Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, volume 464, pages 1355–1365. The Royal Society, 2008.
- [6] Alain Albouy and Vadim Kaloshin. Finiteness of central configurations of five bodies in the plane. *Annals of Mathematics*, pages 535–588, 2012.
- [7] Martha Alvarez-Ramírez, Alan Almeida Santos, and Claudio Vidal. On co-circular central configurations in the four and five body-problems for homogeneous force law. *Journal of Dynamics and Differential Equations*, 25(2):269–290, 2013.
- [8] Tom M Apostol. Ptolemy's inequality and the chordal metric. *Mathe-matics Magazine*, 40(5):233-235, 1967.
- [9] Montserrat Corbera, Josep Cors, Jaume Llibre, and Richard Moeckel. Bifurcation of relative equilibria of the (1+ 3)-body problem. SIAM Journal on Mathematical Analysis, 47(2):1377–1404, 2015.
- [10] Josep M Cors and Gareth E Roberts. Four-body co-circular central configurations. *Nonlinearity*, 25(2):343, 2012.
- [11] Yiyang Deng, Bingyu Li, and Shiqing Zhang. Some notes on four-body co-circular central configurations. *Journal of Mathematical Analysis and Applications*, 453(1):398–409, 2017.

- [12] Otto Dziobek. Über einen merkwürdigen fall des vielkörperproblems. Astron. Nach, 152:33–46, 1900.
- [13] Antonio Carlos Fernandes, Jaume Llibre, and Luis Fernando Mello. Convex central configurations of the 4-body problem with two pairs of equal adjacent masses. Archive for Rational Mechanics and Analysis, 226(1):303–320, 2017.
- [14] Antonio Carlos Fernandes, Luis Fernando Mello, and Claudio Vidal. On the uniqueness of the isosceles trapezoidal central configuration in the 4-body problem for power-law potentials. *Nonlinearity*, 33(1):388–407, dec 2019.
- [15] Pavel Galashin, Steven N Karp, and Thomas Lam. The totally nonnegative grassmannian is a ball. arXiv preprint arXiv:1707.02010, 2017.
- [16] Marshall Hampton. Co-circular central configurations in the four-body problem. In EQUADIFF 2003, pages 993–998. World Scientific, 2005.
- [17] Marshall Hampton. Splendid isolation: local uniqueness of the centered co-circular relative equilibria in the n-body problem. *Celestial Mechanics* and Dynamical Astronomy, 124(2):145–153, 2016.
- [18] Marshall Hampton and Richard Moeckel. Finiteness of relative equilibria of the four-body problem. *Inventiones Mathematicae*, 163(2):289–312, 2006.
- [19] Marshall Hampton, Gareth E Roberts, and Manuele Santoprete. Relative equilibria in the four-vortex problem with two pairs of equal vorticities. *Journal of Nonlinear Science*, 24(1):39–92, 2014.
- [20] Jaume Llibre and Claudia Valls. The co-circular central configurations of the 5-body problem. Journal of Dynamics and Differential Equations, 27(1):55–67, 2015.
- [21] WD MacMillan and Walter Bartky. Permanent configurations in the problem of four bodies. Transactions of the American Mathematical Society, 34(4):838–875, 1932.
- [22] Richard Moeckel. Central Configurations, pages 105–167. Springer Basel, Basel, 2015.

- [23] Forest Ray Moulton. The straight line solutions of the problem of n bodies. The Annals of Mathematics, 12(1):1–17, 1910.
- [24] Pavel Pech. On equivalence of conditions for a quadrilateral to be cyclic. In Beniamino Murgante, Osvaldo Gervasi, Andrés Iglesias, David Taniar, and Bernady O. Apduhan, editors, *Computational Science and Its Applications - ICCSA 2011*, pages 399–411, Berlin, Heidelberg, 2011. Springer Berlin Heidelberg.
- [25] Ernesto Perez-Chavela and Manuele Santoprete. Convex four-body central configurations with some equal masses. Archive for Rational Mechanics and Analysis, 185(3):481–494, 2007.
- [26] Ernesto Perez-Chavela, Manuele Santoprete, and Claudia Tamayo. Symmetric relative equilibria in the four-vortex problem with three equal vorticities. Dynamics of Continuous, Discrete and Impulsive Systems Series A: Mathematical Analysis, pages 189–209, 2015.
- [27] Manuele Santoprete. Four-body central configurations with one pair of opposite sides parallel. Journal of Mathematical Analysis and Applications, 464(1):421–434, 2018.
- [28] Carles Simo. Relative equilibrium solutions in the four body problem. Celestial Mechanics, 18(2):165–184, 1978.
- [29] Steven Smale. Topology and mechanics. II. Inventiones Mathematicae, 11(1):45-64, 1970.
- [30] OY Viro, DB Fuchs, SP Novikov, and VA Rokhlin. Topology ii: Homotopy and homology. classical manifolds. encyclopaedia of mathematical sciences 24, 2004.
- [31] Zhihong Xia. Convex central configurations for the n-body problem. Journal of Differential Equations, 200(2):185–190, 2004.
- [32] Zhifu Xie. Isosceles trapezoid central configurations of the Newtonian four-body problem. Proceedings of the Royal Society of Edinburgh Section A: Mathematics, 142(3):665–672, 2012.
- [33] Jean-Christophe Yoccoz. Description conjecturale des configurations centrales dans le probléme planaire des 4 corps. unpublished, 1986.