

THE LEFSCHETZ PROPERTY FOR AN ALGEBRA DEFINED BY MATCHINGS

YASUHIDE NUMATA

ABSTRACT. In this article, we consider the weighted generating function of matchings consisting of k edges in the complete graph. We define an Artinian Gorenstein algebra as the quotient ring of a polynomial ring by the annihilator of the generating function. We show the strong Lefschetz property of the algebra.

1. INTRODUCTION

The Lefschetz property of Artinian algebras is an algebraic abstraction of the property of the cohomology rings of compact Kähler manifolds known as the Hard Lefschetz Theorem, see also [1, 5]. For a graded algebra $A = \bigoplus_{d=0}^s A^{(d)}$ with $A^{(d)} \neq 0$, we say that A has the *strong Lefschetz property* with a Lefschetz element $l \in A^{(1)}$ if the multiplication map

$$\begin{aligned} \times l^{s-2d}: A^{(d)} &\rightarrow A^{(s-d)} \\ f &\mapsto fl^{s-2d} \end{aligned}$$

is bijective for each d with $2d \leq s$. For the strong Lefschetz property of Artinian Gorenstein algebras, a criterion using higher Hessian matrices is known [8, 4], see also Lemma 2.1. The strong Lefschetz property is studied from the viewpoint of theory of not only algebra but also combinatorics. Recently, some authors study Artinian Gorenstein algebras defined by combinatorial polynomials and calculate the determinants of higher Hessian matrices of the polynomials to show the strong Lefschetz property of the algebras. For example, the basis generating functions for matroids are studied in [3, 11, 10, 7, 6], and the weighted generating functions of facets of Platonic solids are studied in [9]. In this article, we consider the generating function of matchings, i.e., sets of edges which does not share vertices, consisting of k edges in the complete graphs. We show the strong Lefschetz property of the Artinian Gorenstein algebras defined by the weighted generating function.

2020 *Mathematics Subject Classification.* 05C70, 05A15, 13F70.

Key words and phrases. Artinian Gorenstein algebras; Poincaré duality algebras; weighted generating function.

The author was partially supported by JSPS KAKENHI Grant Number JP18K03206.

This article organized as follows: In Section 2, we recall definitions and know facts of Lefschetz property of Artinian Gorenstein algebras. In Section 3, we define the generating function of matchings, and calculate higher Hessian matrices to show the strong Lefschetz property of the Artinian Gorenstein algebra defined by the generating function of matchings.

2. LEFSCHETZ PROPERTIES AND HIGHER HESSIANS

Let \mathbb{K} be a field with $\text{char}(\mathbb{K}) = 0$. Consider the polynomial ring $P = \mathbb{K}[x_i | i \in U]$ in variables indexed by a finite set U with $\deg(x_i) = 1$. For a homogeneous polynomial $\Phi \in P$, we define the annihilator $\text{Ann}(\Phi)$ of Φ by

$$\text{Ann}(\Phi) = \{ f \in P \mid f(\partial)\Phi = 0 \},$$

where $f(\partial)$ is a partial derivative operator obtained from f by substituting $\partial_i = \frac{\partial}{\partial x_i}$ to x_i for each $i \in U$. Since the annihilator $\text{Ann}(\Phi)$ is a homogeneous ideal, the quotient ring $A = P/\text{Ann}(\Phi)$ is a graded ring. Let $A^{(d)}$ be the homogeneous component of degree d . Assume that $\deg(\Phi) = s$. Then we have $A^{(s)} \simeq A^{(0)} \simeq \mathbb{K}$ and $A^{(d)} = 0$ for $d > s$. We call the quotient ring $A = P/\text{Ann}(\Phi)$ for some homogeneous polynomial an *Artinian Gorenstein algebra* with standard grading. It is known that the bilinear map, called the *Poincaré duality*, defined by

$$\begin{aligned} A^{(d)} \times A^{(s-d)} &\rightarrow \mathbb{K} \\ (f, g) &\mapsto f(\partial)g(\partial)\Phi \end{aligned}$$

is non-degenerate. Sometimes, the algebra A is also called a *Poincaré duality algebra*. For an Artinian Gorenstein algebra with standard grading, the following criterion for the bijectivity of the multiplication map $\times l^{s-2d}: A^{(d)} \rightarrow A^{(s-d)}$ is known.

Lemma 2.1 ([8, 4]). *Let $\Phi \in P$ be a homogeneous polynomial of degree s . Let $A^{(d)}$ be the set of homogeneous elements in the quotient ring $A = P/\text{Ann}(\Phi)$ of degree d . Assume that the image of $B \subset P$ is a basis for the \mathbb{K} -vector space $A^{(d)}$. Let H be the matrix $(F(\partial)F'(\partial)\Phi)_{F, F' \in B}$. For $l = \sum_{j \in U} a_j x_j \in A^{(1)}$, the multiplication map $\times l^{s-2d}: A^{(d)} \rightarrow A^{(s-d)}$ is bijective if and only if $\det(H)|_{\forall i, x_i = a_i} \neq 0$.*

We call the matrix H a d -th *Hessian matrix* with respect to the basis B . By Lemma 2.1, if the determinant of a d -th Hessian matrix is a nonzero polynomial, then the set of elements $l \in A^{(1)}$ which induces the bijective multiplication map $\times l^{s-2d}: A^{(d)} \rightarrow A^{(s-d)}$ is an open dense subset in $A^{(1)}$. Hence A has the strong Lefschetz property if and only if the d -th Hessian matrix are nonzero polynomials for all d with $2d \leq s$.

In [3], the Lefschetz property of Artinian Gorenstein algebras defined by the basis generating functions of matroids are studied. In the extended abstract [2] of [3], the elementary symmetric polynomial, which

is the basis generating function of a uniform matroid, is considered as an example.

Example 2.2. For a set U , we define $\binom{U}{k}$ to be the set of subsets of U consisting of k elements. We define the k -th *elementary symmetric polynomial* $e_k(x_i|i \in U)$ by

$$e_k(x_i|i \in U) = \sum_{S \in \binom{U}{k}} x^S,$$

where $x^S = \prod_{i \in S} x_i$. By [3, Theorem 4.3], $A = P / \text{Ann}(e_k(x_i|i \in U)) = \bigoplus_{i=0}^k A^{(i)}$ has the strong Lefschetz property. Moreover, if $2d \leq k$, then the set of images of all monomials of degree d is a basis for the \mathbb{K} -vector space $A^{(d)}$. Hence the Hilbert series $h = (h_0, \dots, h_k)$ is

$$\left(\binom{k}{0}, \binom{k}{1}, \dots, \binom{k}{k} \right)$$

where h_d is the dimension of the \mathbb{K} -vector space $A^{(d)}$. The following matrix is a d -th Hessian matrix

$$\left(\partial^S \partial^{S'} e_k(x_i|i \in U) \right)_{S, S' \in \binom{U}{d}},$$

where $\partial^S = \prod_{i \in S} \partial_i$. Since the algebra A has the strong Lefschetz property, the determinant is a nonzero polynomial. If $k = 2d$, then we have

$$\partial^S \partial^{S'} e_{2d}(x_i|i \in U) = \begin{cases} 1 & (S \cap S' = \emptyset) \\ 0 & (S \cap S' \neq \emptyset) \end{cases}$$

for $S, S' \in \binom{U}{d}$. In this case, the determinant of the d -th Hessian matrix, which is also a representation matrix of the Poincaré duality $A^{(d)} \times A^{(d)} \rightarrow \mathbb{K}$, is a nonzero constant.

3. WEIGHTED GENERATING FUNCTIONS OF MATCHINGS

In this article, we consider only simple nondirected graphs. Hence an edge of a graph with vertex set U is regard as an element in $\binom{U}{2}$. Moreover we assume that the vertex set U is a nonempty finite subset of the set \mathbb{Z} of integers. We define the *complete graph* K_U to be the graph with the vertex set U and the edge set $\binom{U}{2}$. For a graph, a set of edges is called a *matching* in the graph if any distinct edges in the set does not share an vertex. In other words, a subset M of the edge set of a graph is a matching if $e, e' \in M$ and $e \neq e'$ implies $e \cap e' = \emptyset$. We define $\mathcal{M}(U)$ to be the set of matchings in K_U . We also define the set $\mathcal{M}(U, k)$ to be the set $\{ M \in \mathcal{M}(U) \mid |M| = k \}$ consisting of matchings with k edges. For a finite set U with $|U| = u$, we have $|\mathcal{M}(U, k)| = \binom{u}{2k} (2k-1)!!$. For a set $V \subset U$ with $V = \{ v_1 < \dots < v_{2d} \}$, we define $M(V)$ to be the matching $\{ \{ v_{2i-1}, v_{2i} \} \mid i = 1, \dots, d \} \in \mathcal{M}(V, d) \subset \mathcal{M}(U, d)$.

We define P_U to be the polynomial ring $\mathbb{K}[x_e | e \in \binom{U}{2}]$ in variables corresponding to edges in the complete graph K_U . We define the weighted generating function $\Phi_{U,k}$ of $\mathcal{M}(U, k)$ by

$$\Phi_{U,k} = \sum_{M \in \mathcal{M}(U,k)} x^M \in P_U.$$

By definition $\Phi_{U,k}$ is a square-free symmetric homogeneous polynomial of degree k . We define $A_{\Phi_{U,k}}$ to be the quotient ring $P_U / \text{Ann}(\Phi_{U,k})$. Our main result is the following:

Theorem 3.1 (Main result). *For a finite set U , the Artinian Gorenstein algebra $A_{\Phi_{U,k}}$ has the strong Lefschetz property with a Lefschetz element $l = \sum_{e \in \binom{U}{2}} x_e$.*

To show the main theorem, we show some lemma:

Lemma 3.2. *Let V be a subset of U with $|V| = 2d$. For a matching $M \in \mathcal{M}(V, d)$, $\partial^M \Phi_{U,k} = \Phi_{U \setminus V, k-d}$.*

Proof. Since the map

$$\begin{aligned} \mathcal{M}(U \setminus V, k-d) &\rightarrow \left\{ \tilde{M} \in \mathcal{M}(U, k) \mid M \subset \tilde{M} \right\} \\ M' &\mapsto M' \cup M \end{aligned}$$

is bijective, we obtain the equation by direct calculation. \square

Lemma 3.3. *Let*

$$\begin{aligned} G_0 &= \left\{ x_e^2 \mid e \in \binom{U}{2} \right\}, \\ G_1 &= \left\{ x^S \mid S \subset \binom{U}{2} \text{ is not a matching in } K_U \right\}, \\ G_2 &= \left\{ x^M - x^{M'} \mid M, M' \in \mathcal{M}(V, d) \text{ for some } V \in \binom{U}{2d} \right\}. \end{aligned}$$

The annihilator $\text{Ann}(\Phi_{U,k})$ contains $G = G_0 \cup G_1 \cup G_2$.

Proof. Since $\Phi_{U,k}$ is a square-free polynomial, it is obvious that the annihilator $\text{Ann}(\Phi_{U,k})$ contains G_0 . Since a subset of matching is also matching, if S is not a matching, then there does not exist a matching containing S . Hence the annihilator $\text{Ann}(\Phi_{U,k})$ contains G_1 . For $M, M' \in \mathcal{M}(V, d)$, it follows from Lemma 3.2 that

$$(\partial^M - \partial^{M'}) \Phi_{U,k} = \Phi_{U \setminus V, k-d} - \Phi_{U \setminus V, k-d} = 0$$

for $V \in \binom{U}{2d}$ and $M, M' \in \mathcal{M}(V, d)$. Hence the annihilator $\text{Ann}(\Phi_{U,k})$ contains G_2 . \square

By Lemma 3.3, if $2d \leq k$, then the image of $\{x^{M(V)} \mid V \in \binom{U}{2d}\}$ generates the \mathbb{K} -vector space $A_{V,k}^{(d)}$. To show that the set is linearly independent, we calculate the determinant of the following matrix:

$$H_{U,k,d} = \left(\partial^{M(V)} \partial^{M(V')} \Phi_{U,k} \right)_{V, V' \in \binom{U}{2d}}.$$

Lemma 3.4. *Let $V, V' \in \binom{U}{2d}$ with $2d \leq k$ and $|U| = u$. For $M \in \mathcal{M}(V, d)$ and $M' \in \mathcal{M}(V', d)$,*

$$\partial^M \partial^{M'} \Phi_{U,k} = \begin{cases} \Phi_{U \setminus (V \cup V'), k-2d} & (V \cap V' = \emptyset) \\ 0 & (V \cap V' \neq \emptyset). \end{cases}$$

Hence

$$\begin{aligned} \partial^{M'} \partial^M \Phi_{U,k} |_{\forall e, x_e=1} &= \begin{cases} \binom{u-2d}{2k-4d} (2k-4d-1)!! & (V \cap V' = \emptyset) \\ 0 & (V \cap V' \neq \emptyset) \end{cases} \\ &= \binom{u-2d}{2k-4d} (2k-4d-1)!! \partial^{V'} \partial^V e_{2d}(x_i | i \in U). \end{aligned}$$

Proof. By Lemma 3.2, we have

$$\partial^M \Phi_{U,k} = \Phi_{U \setminus V, k-d}.$$

If $V \cap V' \neq \emptyset$, then $U \setminus V$ does not contain V' . Hence any matching $\tilde{M} \in \mathcal{M}(U \setminus V)$ does not contain $M' \in \mathcal{M}(V', 2k)$. In this case, $\partial^M \partial^{M'} \Phi_{U,k} = 0$. If $V \cap V' = \emptyset$, then $\partial^{M'} \Phi_{U \setminus V, k-d} = \Phi_{U \setminus (V \cup V'), k-2d}$ by Lemma 3.2. Since $|\mathcal{M}(S, k)| = \binom{|S|}{2k} (2k-1)!!$ for a finite set S , we have $\partial^{M'} \partial^M \Phi_{U,k} |_{\forall e, x_e=1} = \binom{u-2d}{2k-4d} (2k-4d-1)!!$ in this case. Since

$$\partial^S \partial^{S'} e_{2d}(x_i | i \in U) = \begin{cases} 1 & (S \cap S' = \emptyset) \\ 0 & (S \cap S' \neq \emptyset) \end{cases}$$

for $S, S' \in \binom{U}{d}$, we have

$$\partial^{M'} \partial^M \Phi_{U,k} |_{\forall e, x_e=1} = \binom{u-2d}{2k-4d} (2k-4d-1)!! \partial^{V'} \partial^V e_{2d}(x_i | i \in U)$$

for $M \in \mathcal{M}(V, d)$ and $M' \in \mathcal{M}(V', d)$. \square

By Lemma 3.4, we have

$$\det(H_{U,k,d}) |_{\forall e, x_e=1} = c \det \left(\partial^{V'} \partial^V e_{2d}(x_i | i \in U) \right)_{V, V' \in \binom{U}{2d}},$$

where $c = \left(\binom{u-2d}{2k-4d} (2k-4d-1)!! \right) \binom{u}{2d}$ and $u = |U|$. Since the determinant $\det \left(\partial^{V'} \partial^V e_{2d}(x_i | i \in U) \right)_{V, V' \in \binom{U}{2d}}$ is not zero as in Example 2.2, the determinant $\det(H_{U,d,k}) |_{\forall e, x_e=1}$ is a nonzero constant. Since $\det(H_{U,d,k})$ is a nonzero polynomial, the image of $\{x^{M(V)} \mid V \in \binom{U}{2d}\}$ is linearly independent. Hence we have the following corollary.

Corollary 3.5. *The Hilbert series of $A_{U,k}$ with $2k \leq |U|$ is*

$$\left(\binom{2k}{0}, \binom{2k}{2}, \binom{2k}{4}, \dots, \binom{2k}{2k} \right)$$

Since the image of the set $\{x^{M(V)} \mid V \in \binom{U}{2d}\}$ is a basis for the \mathbb{K} -vector space $A_{V,k}^{(d)}$, the matrix $H_{U,d,k}$ is a d -th Hessian matrix for $A_{V,k}^{(d)}$ for d with $2d \leq k$. By Lemma 2.1, we have the strong Lefschetz property for $A_{V,k}$, which is our main result Theorem 3.1.

REFERENCES

- [1] Tadahito Harima, Toshiaki Maeno, Hideaki Morita, Yasuhide Numata, Akihito Wachi, and Junzo Watanabe, *The Lefschetz properties*, Lecture Notes in Mathematics, vol. 2080, Springer, Heidelberg, 2013, URL <https://doi.org/10.1007/978-3-642-38206-2>. MR 3112920
- [2] Toshiaki Maeno and Yasuhide Numata, *Sperner property, matroids and finite-dimensional Gorenstein algebras*, Tropical geometry and integrable systems, Contemp. Math., vol. 580, Amer. Math. Soc., Providence, RI, 2012, pp. 73–84, URL <https://doi.org/10.1090/conm/580/11496>. MR 2985388
- [3] ———, *Sperner property and finite-dimensional Gorenstein algebras associated to matroids*, J. Commut. Algebra **8** (2016), no. 4, 549–570, URL <https://doi.org/10.1216/JCA-2016-8-4-549>. MR 3566530
- [4] Toshiaki Maeno and Junzo Watanabe, *Lefschetz elements of Artinian Gorenstein algebras and Hessians of homogeneous polynomials*, Illinois J. Math. **53** (2009), no. 2, 591–603, URL <http://projecteuclid.org/euclid.ijm/1266934795>. MR 2594646
- [5] Juan Migliore and Uwe Nagel, *Survey article: a tour of the weak and strong Lefschetz properties*, J. Commut. Algebra **5** (2013), no. 3, 329–358, URL <https://doi.org/10.1216/JCA-2013-5-3-329>. MR 3161738
- [6] Satoshi Murai, Takahiro Nagaoka, and Akiko Yazawa, *Strictness of the log-concavity of generating polynomials of matroids*, J. Combin. Theory Ser. A **181** (2021), Paper No. 105351, 22, URL <https://doi.org/10.1016/j.jcta.2020.105351>. MR 4223331
- [7] Takahiro Nagaoka and Akiko Yazawa, *Strict log-concavity of the Kirchhoff polynomial and its applications to the strong Lefschetz property*, J. Algebra **577** (2021), 175–202, URL <https://doi.org/10.1016/j.jalgebra.2021.01.037>. MR 4234203
- [8] Junzo Watanabe, *A remark on the Hessian of homogeneous polynomials*, The curves seminar at Queen’s, vol. XIII, Queen’s Papers in Pure and Appl. Math., vol. 119, Queen’s Univ., Kingston, ON, 2000, pp. 171–178.
- [9] Akiko Yazawa, *On artinian gorenstein algebras associated to the face posets of regular polyhedra*, 2020, Eprint [arXiv:2011.13732](https://arxiv.org/abs/2011.13732).
- [10] Akiko Yazawa, *The eigenvalues of Hessian matrices of the complete and complete bipartite graphs*, J. Algebraic Combin. **54** (2021), no. 4, 1137–1157, URL <https://doi.org/10.1007/s10801-021-01041-x>. MR 4348920
- [11] ———, *The eigenvalues of the Hessian matrices of the generating functions for trees with k components*, Linear Algebra Appl. **631** (2021), 48–66, URL <https://doi.org/10.1016/j.laa.2021.08.017>. MR 4308087

(Y. Numata) DEPARTMENT OF MATHEMATICS, HOKKAIDO UNIVERSITY, SAPPORO, JAPAN.

Email address: nu@math.sci.hokudai.ac.jp