

Translation of “Simplizialzerlegungen von Beschränkter  
Flachheit” by Hans Freudenthal, *Annals of Mathematics*,  
Second Series, Volume 43, Number 3, July 1942, Pages  
580-583.\*

Translated by:  
Mathijs Wintraecken, Inria Sophia-Antipolis, Université Côte d’Azur  
and Institute of Science and Technology Austria †

We answer a question by Brouwer about the construction of an infinite series of subdivisions of a polytope, such that the next element in the sequence is a subdivision of the previous one and such that the subsimplices that arise do not become arbitrarily flat, that is the quotient

$$c^r/v$$

( $c$  = diameter = the longest edge of the simplex,  $v$  = volume of the simplex) is uniformly bounded for all  $r$  dimensional simplices. Such tillings are useful in analysis and the interface of analysis and topology.

The construction, that we give here, is analogous to our subdivision by simplices of the Cartesian product of two simplices.<sup>1</sup>

## 1

Simplices will be denoted by a fixed order of vertices, parallelepiped will be given by a marked vertex and an ordered set of edges that emanate from this vertex.

The simplex  $T$  has vertices

$$e_0, \dots, e_r.$$

We can also describe  $T$  by the vectors

$$x_0 = e_0, \quad x_i = e_i - e_{i-1} \quad (i = 1, \dots, r).$$

---

\*Freudenthal added the following as a footnote to the title: This note is essentially the same as a note that was submitted to *Fundamenta Mathematicae* in March 1939. However this note never appeared in print. The result of was also used in a different paper “Die Triangulation der differenzierbaren Mannigfaltigkeiten” *Nederlandse Akademie voor Wetenschappen, Proceedings*, 42 (1939), 880–901, <https://dwc.knaw.nl/DL/publications/PU00014650.pdf>. [More extensive reference inserted by translator]

†The translator has been supported by the European Research Council (ERC) under the European Union’s Seventh Framework Programme (FP/2007-2013) / ERC Grant Agreement No. 339025 GUDHI (Algorithmic Foundations of Geometry Understanding in Higher Dimensions), the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 754411, and the Austrian science fund (FWF) under grant agreement M-3073.

<sup>1</sup>Freudenthal, “Eine Simplizialzerlegung des Cartesischen Produktes zweier Simplexe” *Fundamenta Mathematicae*, 29 (1937), 138-144, <https://bibliotekanauki.pl/articles/1383987>.

Let  $\pi$  be a permutation  $p_1, \dots, p_r$  of the numbers  $1 \dots r$ . The points<sup>2</sup>

$$e_i = x_0 + \sum_{\nu=1}^i x_{p_\nu} \quad (i = 1, \dots, r)$$

(seen as vertices) yield a simplex  $T^\pi$ . If  $\pi_0$  the identity permutation is of  $1, \dots, r$  then  $T^{\pi_0} = T$ . The  $r!$  simplices  $T^\pi$  are said to be the conjugates of each other.

$T^\pi$  is the set<sup>3</sup>

$$T^\pi = \left\{ \sum_{\nu=1}^r \lambda_\nu e_{p_\nu} \mid \lambda_\nu \geq 0, \sum_{\nu=1}^r \lambda_\nu = 1 \right\},$$

or what boils down to the same thing,<sup>4</sup>

$$T^\pi = \left\{ x_0 + \sum_{\nu=1}^r \alpha_\nu x_{p_\nu} \mid 1 \geq \alpha_1 \geq \dots \geq \alpha_r \geq 0 \right\}.$$

It follows that: The  $T^\pi$  are the images of a subdivision of the parallelepiped  $P$ , given by,

$$P = \left\{ x_0 + \sum_{\nu=1}^r \alpha_\nu x_\nu \mid 1 \geq \alpha_\nu \geq 0 \right\}.$$

Conversely the simplex  $T^\pi$  is unambiguously determined by the parallelepiped  $P$ , that is given by the vertex  $x_0$  and the edge vectors  $x_1, \dots, x_r$  and by the permutation  $\pi$ .

## 2

Cutting all the edges of the parallelepiped in half all, yields [note translator:  $2^r$ ] parallelepipeds

$$P_\sigma = \left\{ x_{\sigma,0} + \frac{1}{2} \sum_{\nu=1}^r \alpha_\nu x_\nu \mid 1 \geq \alpha_\nu \geq 0 \right\}$$

where<sup>5</sup>  $\sigma = (s_1, \dots, s_r) \in \{0, 1\}^r$ , and

$$x_{\sigma,0} = x_0 + \frac{1}{2} \sum_{\nu=1}^r s_\nu x_\nu$$

In the same way that  $P$  is subdivided by  $T^\pi$ , is  $P_\sigma$  subdivided by  $(T_\sigma)^\pi$ . Note that  $T^\pi$  and  $(T_\sigma)^\pi$  are homothetic [with the ratio]  $(1 : 2)$ .

Let  $\pi$  again be a permutation  $p_1 \dots p_r$  of  $1, \dots, r$  and let  $\pi^* = \pi^\sigma$  be the permutation  $p_1^* \dots p_r^*$  that arises if one first (using the order given by  $\pi$ ) takes all  $p_\nu$  with  $s_{p_\nu} = 1$  and then all with  $s_{p_\nu} = 0$ , so

- if  $s_a > s_b$  then  $a$  comes before  $b$  in  $\pi^*$
- if  $s_a = s_b$  and  $a$  is before  $b$  in  $\pi$  then  $a$  also comes before  $b$  in  $\pi^*$ .

<sup>2</sup>Note translator: There was a typo in the sum.

<sup>3</sup>Note translator: Here I updated the notation quite significantly.

<sup>4</sup>Note translator: This is not entirely straightforward, the argument (or a slight generalization thereof) is spelled out in [BKW21].

<sup>5</sup>Note translator: Here I updated the notation quite significantly.

We claim that  $(T_\sigma)^\pi$  lies completely in  $T^{\pi^\sigma}$ . Indeed we have,

$$\begin{aligned} (T_\sigma)^\pi &= \left\{ x_0 + \frac{1}{2} \sum_{\nu=1}^r s_\nu x_\nu + \frac{1}{2} \sum_{\nu=1}^r \alpha_\nu x_{p_\nu} \mid 1 \geq \alpha_1 \geq \dots \geq \alpha_r \geq 0 \right\} \\ &= \left\{ x_0 + \sum_{\nu \in \{\nu \mid s_{p_\nu}=1\}} \frac{1}{2} (\alpha_\nu + 1) x_{p_\nu} + \frac{1}{2} \sum_{\nu \in \{\nu \mid s_{p_\nu}=0\}} \alpha_\nu x_\nu \mid 1 \geq \alpha_1 \geq \dots \geq \alpha_r \geq 0 \right\} \\ &= \left\{ x_0 + \sum_{\nu=1}^r \beta_\nu x_{p_\nu} \mid 1 \geq \beta_1 \geq \dots \geq \beta_u \geq \frac{1}{2} \geq \beta_{u+1} \geq \dots \geq \beta_r \geq 0 \right\}, \end{aligned}$$

where  $u$  is  $\#\{\nu \mid s_{p_\nu} = 1\}$ . Our claim now follows. Moreover, we have that the  $(T_\sigma)^\pi$  such that  $\pi^\sigma = \pi^*$  form a simplicial subdivision  $Z(T)$  of  $T$ .

**3**<sup>6</sup>

Let  $e_{ij}^{\pi^*}$  be the midpoint of the edge  $e_i^{\pi^*} e_j^{\pi^*}$ , where we define  $e_{ii}^{\pi^*} = e_i^{\pi^*}$ . With this definition, and assuming without loss of generality that  $i \leq j$ , we have<sup>7</sup>

$$\begin{aligned} e_{ij}^{\pi^*} &= \frac{1}{2} \left( x_0 + \sum_{\nu=1}^i x_{p_\nu} + x_0 + \sum_{\nu=1}^j x_{p_\nu} \right) \\ &= x_0 + \sum_{\nu=1}^i x_{p_\nu} + \frac{1}{2} \sum_{\nu=i+1}^j x_{p_\nu} \end{aligned}$$

We define  $k'$  to be the number of  $\nu$  with  $s_{p_\nu} = 1$ , and  $k''$  to be the number of  $\nu$  with  $s_{p_\nu} = 0$ .

We say that the  $k$ th vertex of  $T_\sigma^\pi$  is given by setting  $\alpha_1 = \dots = \alpha_k = 1$  and  $\alpha_{k+1} = \dots = \alpha_r = 0$ . The  $k$ th vertex of  $T_\sigma^\pi$  can be written in terms of the midpoints we defined above as<sup>8</sup>

$$x_0 + \sum_{\nu=1}^{k'} x_{p_\nu} + \sum_{\nu=k'+1}^{k'+u} x_{p_\nu} = e_{ij}^{\pi^*},$$

where  $i = k'$  and  $j = k'' + u$ .

From this it follows that for every subsimplex  $T_\sigma^\pi$  of  $(T^\pi)^\sigma$ :

- The 0th vertex is  $e_{0u}^{\pi^\sigma}$
- the vertex  $e_{ij}^{\pi^\sigma}$  is followed by either  $e_{i+1,j}^{\pi^\sigma}$  or  $e_{i,j+1}^{\pi^\sigma}$ .
- the first index is bounded by  $u$  and the second by  $r$ .

Conversely are the subsimplices  $(T_\sigma)^\pi$  of  $T^{\pi^*}$  characterized by these properties.

**4**

We now concentrate on  $T = T_0^\pi$  and drop the index  $\pi_0$  in the following. We can now also describe the subdivision  $Z(T)$  of  $T$  in the following way:

<sup>6</sup>Note translator: I personally find the subdivision easier to see, by first observing that it is a hyperplane arrangement and then do a rescaling, compare to [BKW21], also see the references mentioned there.

<sup>7</sup>Note translator: Here too there was a typo, I doubt that Freudenthal ever saw the proofs of this paper, because of the Nazi occupation of the Netherlands and he being Jewish.

<sup>8</sup>[Literally:] this therefore gives

- The vertices of  $Z(T)$  are the  $e_{ij}$ ;
- The  $e_{ij}$  and  $e_{i'j'}$  form a one dimensional simplex if and only if the pairs  $ij$  and  $i'j'$  do not separate each other;
- a set of  $e_{ij}e_{ij}$  form a simplex of  $Z(T)$  is and only if the elements form pairwise simplices of  $Z(T)$ ;
- the vertices in the simplices of  $Z(T)$  are ordered in ascending order of  $i + j$ .

We also note that: The simplices of  $Z(T)$  are conjugated with  $T$ .

Let now a finite polytope  $R$  be given. We impose an ordering on the vertices and form the division  $Z(R)$ , where we use the process  $Z$  on every simplex of  $R$ . The edges of  $Z(R)$  are ordered lexicographically. If we repeat the process as many times as we want we preserve a order of subdivision. All simplices that occur are similar to conjugated simplices of  $R$  and because similar simplices have the same quality

$$c^r/v$$

we see that the flatness of the simplices remains bounded.

**Amsterdam**

**Acknowledgements** The translator thanks the editorial board of the Annals of Mathematics for their permission to make a translation of this paper public. The Annals of Mathematics has the copyright to the original German text.

## References

- [BKW21] Jean-Daniel Boissonnat, Siargey Kachanovich, and Mathijs Wintraecken. Tracing Isomanifolds in  $\mathbb{R}^d$  in Time Polynomial in  $d$  Using Coxeter-Freudenthal-Kuhn Triangulations. In Kevin Buchin and Éric Colin de Verdière, editors, *37th International Symposium on Computational Geometry (SoCG 2021)*, volume 189 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 17:1–17:16, Dagstuhl, Germany, 2021. Schloss Dagstuhl – Leibniz-Zentrum für Informatik. Journal version to be published in the SIAM Journal on Computing.