Smallest and Largest Block Palindrome Factorizations

Daniel Gabric* and Jeffrey Shallit[†]

Abstract

A palindrome is a word that reads the same forwards and backwards. A block palindrome factorization (or BP-factorization) is a factorization of a word into blocks that becomes palindrome if each identical block is replaced by a distinct symbol. We call the number of blocks in a BP-factorization the width of the BP-factorization. The largest BP-factorization of a word w is the BP-factorization of w with the maximum width. We study words with certain BP-factorizations. First, we give a recurrence for the number of length-n words with largest BP-factorization of width t. Second, we show that the expected width of the largest BP-factorization of a word tends to a constant. Third, we give some results on another extremal variation of BP-factorization, the smallest BP-factorization. A border of a word w is a non-empty word that is both a proper prefix and suffix of w. Finally, we conclude by showing a connection between words with a unique border and words whose smallest and largest BP-factorizations coincide.

1 Introduction

Let Σ_k denote the alphabet $\{0, 1, \dots, k-1\}$. The length of a word w is denoted by |w|. A border of a word w is a non-empty word that is both a proper prefix and suffix of w. A word is said to be bordered if it has a border. Otherwise, the word is said to be unbordered. For example, the French word entente is bordered, and has two borders, namely entered entered.

It is well-known [1] that the number u_n of length-n unbordered words over Σ_k satisfies

$$u_{n} = \begin{cases} 1, & \text{if } n = 0; \\ ku_{n-1} - u_{n/2}, & \text{if } n > 0 \text{ is even;} \\ ku_{n-1}, & \text{if } n \text{ is odd.} \end{cases}$$
 (1)

A palindrome is a word that reads the same forwards as it does backwards. More formally, letting $w^R = w_n w_{n-1} \cdots w_1$ where $w = w_1 w_2 \cdots w_n$ and all w_i are symbols, a palindrome is

^{*}Department of Math/Stats, University of Winnipeg, Winnipeg, MB R3B 2E9, Canada; d.gabric@uwinnipeg.ca.

[†]School of Computer Science, University of Waterloo, Waterloo, ON N2L 3G1, Canada shallit@uwaterloo.ca.

a word w such that $w = w^R$. The definition of a palindrome is quite restrictive. The second half of a palindrome is fully determined by the first half. Thus, compared to all length-n words, the number of length-n palindromes is vanishingly small. But many words exhibit palindrome-like structure. For example, take the English word marjoram. It is clearly not a palindrome, but it comes close. Replacing the block jo with a single letter turns the word into a palindrome. In this paper, we consider a generalization of palindromes that incorporates this kind of palindromic structure.

In the 2015 British Olympiad [2], the concept of a block palindrome factorization was first introduced. Let w be a non-empty word. A block palindrome factorization (or BP-factorization) of w is a factorization $w = w_m \cdots w_1 w_0 w_1 \cdots w_m$ of a word such that w_0 is a possibly empty word, and every other factor w_i is non-empty for all i with $1 \le i \le m$. We say that a BP-factorization $w_m \cdots w_1 w_0 w_1 \cdots w_m$ is of width t where t = 2m + 1 if w_0 is non-empty and t = 2m otherwise. In other words, the width of a BP-factorization is the number of non-empty blocks in the factorization. The largest BP-factorization¹ [3] of a word w is a BP-factorization $w = w_m \cdots w_1 w_0 w_1 \cdots w_m$ where m is maximized (i.e., where the width of the BP-factorization is maximized). See [4, 5] for more on the topic of BP-factorizations and block reversals. Kolpakov and Kucherov [6] studied a special case of BP-factorizations, the gapped palindrome. If w_0 is non-empty and $|w_i| = 1$ for all i with $1 \le i \le m$, then w is said to be a gapped palindrome. Régnier [7] studied something similar to BP-factorizations, but in her paper she was concerned with borders of borders. See [8, 9] for results on factoring words into palindromes.

Example 1. We use the centre dot \cdot to denote the separation between blocks in the BP-factorization of a word.

Consider the word abracadabra. It has the following BP-factorizations:

```
abracadabra,
abra·cad·abra,
a·bracadabr·a,
a·br·acada·br·a,
a·br·a·cad·a·br·a.
```

The last BP-factorization is of width 7 and has the longest width; thus it is the largest BP-factorization of abracadabra.

Let w be a length-n word. Suppose $w_m \cdots w_1 w_0 w_1 \cdots w_m$ is the largest BP-factorization of w. Goto et al. [3] showed that w_i is the shortest border of $w_i \cdots w_1 w_0 w_1 \cdots w_i$ where $i \geq 1$. This means that we can compute the largest BP-factorization of w by greedily "peeling off" the shortest borders of central factors until you hit an unbordered word or the empty word.

The rest of the paper is structured as follows. In Section 2 we give a recurrence for the number of length-n words with largest BP-factorization of width t. In Section 3 we

¹Largest BP-factorizations also appear in https://www.reddit.com/r/math/comments/ga2iyo/i_just_defined_the_palindromity_function_on/.

show that the expected width of the largest BP-factorization of a length-n word tends to a constant. In Section 4 we consider *smallest BP-factorizations* in the sense that one "peels off" the longest non-overlapping border. We say a border u of a word w is *non-overlapping* if $|u| \leq |w|/2$; otherwise u is *overlapping*. Finally, in Section 5 we present some results on words with a unique border and show that they are connected to words whose smallest and largest BP-factorizations are the same.

2 Counting largest BP-factorizations

In this section, we prove a recurrence for the number $LBP_k(n,t)$ of length-n words over Σ_k with largest BP-factorization of width t. See Table 1 for sample values of $LBP_2(n,t)$ for small n, t. For the following theorem, recall the definition of u_n from Equation 1.

Theorem 2. Let $n, t \ge 0$, and $k \ge 2$ be integers. Then

$$\mathrm{LBP}_k(n,t) = \begin{cases} \sum_{i=1}^{(n-t)/2+1} u_i \, \mathrm{LBP}_k(n-2i,t-2), & \text{if n, t even;} \\ \sum_{i=1}^{(n-t+1)/2} u_{2i} \, \mathrm{LBP}_k(n-2i,t-1), & \text{if n even, t odd;} \\ 0, & \text{if n odd, t even;} \\ \sum_{i=1}^{(n-t)/2+1} u_{2i-1} \, \mathrm{LBP}_k(n-2i+1,t-1), & \text{if n, t odd.} \end{cases}$$

where

$$LBP_k(0,0) = 1,$$

$$LBP_k(2n,2) = u_n,$$

$$LBP_k(n,1) = u_n.$$

Proof. Let w be a length-n word whose largest BP-factorization $w_m \cdots w_1 w_0 w_1 \cdots w_m$ is of width t. Clearly LBP_k(0,0) = 1. We know that each block in a largest BP-factorization is unbordered, since each block is a shortest border of some central factor. This immediately implies LBP_k $(n,1) = u_n$ and LBP_k $(2n,2) = u_n$.

Now we take care of the other cases.

• Suppose n, t are even. Then by removing both instances of w_1 from w, we get $w' = w_m \cdots w_2 w_2 \cdots w_m$, which is a length- $(n-2|w_1|)$ word whose largest BP-factorization is of width t-2. This mapping is clearly reversible, since all blocks in a largest BP-factorization are unbordered, including w_1 . Thus summing over all possible w_1 and all length- $(n-2|w_1|)$ words with largest BP-factorization of width t-2 we have

LBP_k(n,t) =
$$\sum_{i=1}^{(n-t)/2+1} u_i LBP_k(n-2i, t-2).$$

• Suppose n is even and t is odd. Then by removing w_0 from w, we get $w' = w_m \cdots w_1 w_1 \cdots w_m$, which is a length- $(n - |w_0|)$ word whose largest BP-factorization is of width t - 1. This mapping is reversible for the same reason as in the previous case. The word w' is of even length since $|w'| = 2|w_1 \cdots w_m|$. Since n is even and |w'| is even, we must have that $|w_0|$ is even as well. Thus summing over all possible w_0 and all length- $(n - |w_0|)$ words with largest BP-factorization of width t - 1, we have

$$LBP_{k}(n,t) = \sum_{i=1}^{(n-t+1)/2} u_{2i} LBP_{k}(n-2i,t-1).$$

- Suppose n is odd and t is even. Then the length of w is $2|w_1 \cdots w_m|$, which is even, a contradiction. Thus $LBP_k(n,t) = 0$.
- Suppose n, t are odd. Then by removing w_0 from w, we get $w' = w_m \cdots w_1 w_1 \cdots w_m$, which is a length- $(n |w_0|)$ word whose largest BP-factorization is of width t 1. This mapping is reversible for the same reasons as in the previous cases. Since n is odd and |w'| is even (proved in the previous case), we must have that $|w_0|$ is odd. Thus summing over all possible w_0 and all length- $(n |w_0|)$ words with largest BP-factorization of width t 1, we have

$$\sum_{i=1}^{(n-t)/2+1} u_{2i-1} \operatorname{LBP}_k(n-2i+1,t-1).$$

n	1	2	3	4	5	6	7	8	9	10
10	284	12	224	40	168	72	96	64	32	32
11	568	0	472	0	416	0	336	0	192	0
12	1116	20	856	88	656	176	448	224	224	160
13	2232	0	1752	0	1488	0	1248	0	896	0
14	4424	40	3328	176	2544	432	1856	640	1152	640
15	8848	0	6736	0	5440	0	4576	0	3584	0
16	17622	74	13100	372	9896	984	7408	1744	5088	2080
17	35244	0	26348	0	20536	0	16784	0	13664	0
18	70340	148	51936	760	38824	2248	29152	4416	21088	6240
19	140680	0	104168	0	79168	0	62800	0	51008	0
20	281076	284	206744	1592	153344	4992	114688	10912	84704	17312

Table 1: Some values of LBP₂(n,t) for n,t where $10 \le n \le 20$ and $1 \le t \le 10$.

3 Expected width of largest BP-factorization

In this section, we show that the expected width $E_{n,k}$ of the largest BP-factorization of a length-n word over Σ_k is bounded by a constant. From the definition of expected value, it follows that

$$E_{n,k} = \frac{1}{k^n} \sum_{i=1}^n i \cdot LBP_k(n,i).$$

Table 2 shows the behaviour of $\lim_{n\to\infty} E_{n,k}$ as k increases.

Lemma 3. Let $k \geq 2$ and $n \geq t \geq 1$ be integers. Then

$$\frac{\mathrm{LBP}_k(n,t)}{k^n} \le \frac{1}{k^{t/2-1}}.$$

Proof. Let w be a length-n word whose largest BP-factorization $w_m \cdots w_1 w_0 w_1 \cdots w_m$ is of width t. Since w_i is non-empty for every $1 \le i \le m$, we have that $LBP_k(n,t) \le k^{n-m} \le k^{n-t/2+1}$. So

$$\frac{\text{LBP}_k(n,t)}{k^n} \le \frac{1}{k^{t/2-1}}$$

for all $n \ge t \ge 1$.

Theorem 4. The limit $E_k = \lim_{n \to \infty} E_{n,k}$ exists for all $k \ge 2$.

Proof. Follows from the definition of $E_{n,k}$, Lemma 3, and the direct comparison test for convergence.

Interpreting E_k as a power series in k^{-1} , we empirically observe that E_k is approximately equal to

$$1 + \frac{2}{k} + \frac{4}{k^2} + \frac{6}{k^3} + \frac{10}{k^4} + \frac{16}{k^5} + \frac{24}{k^6} + \frac{38}{k^7} + \frac{58}{k^8} + \frac{88}{k^9} + \cdots$$

We conjecture the following about E_k .

Conjecture 5. Let $k \geq 2$. Then

$$E_k = 1 + \sum_{i=1}^{\infty} a_i k^{-i}$$

where the sequence $(a_n/2)_{n\geq 1}$ is $\underline{A274199}$ in the On-Line Encyclopedia of Integer Sequences (OEIS) [10].

k	$\approx E_k$
2	6.4686
3	2.5908
4	1.9080
5	1.6314
6	1.4827
7	1.3902
8	1.3272
9	1.2817
10	1.2472
:	:
100	1.0204

Table 2: Asymptotic expected width of a word's largest BP-factorization.

Cording et al. [11] proved that the expected length of the longest unbordered factor in a word is $\Theta(n)$. Taking this into account, it is not surprising that the expected length of the largest BP-factorization of a word tends to a constant.

4 Smallest BP-factorization

A word w, seen as a block, clearly satisfies the definition of a BP-factorization. Thus, taken literally, the smallest BP-factorization for all words is of width 1. But this is not very interesting, so we consider a different definition instead. A border u of a word w is non-overlapping if $|u| \leq |w|/2$; otherwise u is overlapping. We say that the smallest BP-factorization of a word w is a BP-factorization $w = w_m \cdots w_1 w_0 w_1 \cdots w_m$ where each w_i is the longest non-overlapping border of $w_i \cdots w_1 w_0 w_1 \cdots w_i$, except w_0 , which is either empty or unbordered. For example, going back to Example 1, the smallest BP-factorization of abracadabra is $abra \cdot cad \cdot abra$ and the smallest BP-factorization of reappear is $r \cdot ea \cdot p \cdot p \cdot ea \cdot r$.

A natural question to ask is: what is the maximum possible width $f_k(n)$ of the smallest BP-factorization of a length-n word? Through empirical observation, we arrive at the following conjectures:

- We have $f_2(8n+i) = 6n+i$ for i with $0 \le i \le 5$ and $f_2(8n+6) = f_2(8n+7) = 6n+5$.
- We have $f_k(n) = n$ for $k \ge 3$.

To calculate $f_k(n)$, two things are needed: an upper bound on $f_k(n)$, and words that witness the upper bound.

Theorem 6. Let $l \ge 0$ be an integer. Then $f_2(8l+i) = 6l+i$ for i with $0 \le i \le 5$ and $f_2(8l+6) = f_2(8l+7) = 6l+5$.

Proof. Let $n \ge 0$ be an integer. We start by proving lower bounds on $f_2(n)$. Suppose n = 8l for some $l \ge 0$. Then the width of the smallest BP-factorization of

$$(0101)^l(1001)^l$$

is 6l, so $f_2(8l) \ge 6l$. To see this, notice that the smallest BP-factorization of 01011001 is $01 \cdot 0 \cdot 1 \cdot 1 \cdot 0 \cdot 01$, and therefore is of width 6. Suppose n = 8l + i for some i with $1 \le i \le 7$. Then one can take $(0101)^l(1001)^l$ and insert either 0, 00, 010, 0110, 01010, 010110, or 0110110 to the middle of the word to get the desired length.

Now we prove upper bounds on $f_2(n)$. Let $t \leq n$ be a positive integer. Let w be a length-n word whose largest BP-factorization $w_m \cdots w_1 w_0 w_1 \cdots w_m$ is of width t. One can readily verify that $f_2(0) = 0$, $f_2(1) = 1$, $f_2(2) = 2$, $f_2(3) = 3$, $f_2(4) = 4$, and $f_2(5) = f_2(6) = f_2(7) = 5$ through exhaustive search of all binary words of length < 8. Suppose $m \geq 4$, so $n \geq t \geq 8$. Then we can write $w = w_m w_{m-1} w_{m-2} \cdots w_{m-2} w_{m-1} w_m$ where $|w_{m-2}|, |w_{m-1}|, |w_m| > 0$. It is easy to show that $|w_{m-2}w_{m-1}w_m| \geq 4$ by checking that all binary words of length < 8 do not admit a smallest BP-factorization of width 6. In the worst case, we can peel off prefixes and suffixes of length 4 while accounting for the 6 blocks they add to the BP-factorization until we hit the middle core of length < 8. Thus, we have $f_2(8l+i) \leq 6l+j$ where j is the width of the smallest BP-factorization of the middle core, which is of length i. We have already computed $f_2(i)$ for $0 \leq i \leq 7$, so the upper bounds follow.

Theorem 7. Let $n \geq 0$ and $k \geq 3$ be integers. Then $f_k(n) = n$.

Proof. Clearly $f_k(n) \leq n$. We prove $f_k(n) \geq n$. If n is divisible by 6, then consider the word $(012)^{n/6}(210)^{n/6}$. If n is not divisible by 6, then take $(012)^{\lfloor n/6\rfloor}(210)^{\lfloor n/6\rfloor}$ and insert either 0, 00, 010, 0110, or 01010 in the middle of the word. When calculating the smallest BP-factorization of the resulting words, it is easy to see that at each step we are removing a border of length 1. Thus, their largest BP-factorization is of width n.

5 Equal smallest and largest BP-factorizations

Recall back to Example 1, that abracadabra has distinct smallest and largest BP-factorizations, namely abra·cad·abra and a·br·a·cad·a·br·a. However, the word alfalfa has the same smallest and largest BP-factorizations, namely a·lf·a·lf·a. Under what conditions are the smallest and largest BP-factorizations of a word the same? Looking at unique borders seems like a good place to start, since the shortest border and longest non-overlapping border coincide when a word has a unique border. However, the converse is not true—just consider the previous example alfalfa. The shortest border and longest non-overlapping border are both a, but a is not a unique border of alfalfa.

In Theorem 8 we characterize all words whose smallest and largest BP-factorization coincide.

Theorem 8. Let $m, m' \geq 1$ be integers. Let w be a word with smallest BP-factorization $w'_{m'} \cdots w'_1 w'_0 w'_1 \cdots w'_{m'}$ and largest BP-factorization $w_m \cdots w_1 w_0 w_1 \cdots w_m$. Then m = m'

and $w_i = w_i'$ for all $i, 0 \le i \le m$ if and only if for all $i \ne 2, 0 < i \le m$, we have that w_i is the unique border of $w_i \cdots w_1 w_0 w_1 \cdots w_i$ and for i = 2 we have that either

- 1. w_2 is the unique border of $w_2w_1w_0w_1w_2$, or
- 2. $w_2w_1w_0w_1w_2 = w_0w_1w_0w_1w_0$ where w_0 is the unique border of $w_0w_1w_0$.

Proof.

 \implies : Let i be an integer such that $0 < i \le m$. Let $u_i = w_i \cdots w_1 w_0 w_1 \cdots w_i$. Since w_i is both the shortest border and longest non-overlapping border of u_i (i.e., $w_i = w_i'$), we have that u_i has exactly one border of length $\leq |u_i|/2$. Thus, either w_i is the unique border of u_i , or u_i has a border of length $> |u_i|/2$. If w_i is the unique border of u_i , then we are done. So suppose that u_i has a border of length $> |u_i|/2$. Let v_i be the shortest such border. We have that w_i is both a prefix and suffix of v_i . In fact, w_i must be the unique border of v_i . Otherwise we contradict the minimality of v_i , or the assumption that w_i is both the shortest border and longest non-overlapping border of u_i . Since w_i is unbordered, it cannot overlap itself in v_i and w_i . So we can write $v_i = w_i y w_i$ for some word y where $u_i = w_i y w_i y w_i$, or $u_i = w_i x w_i x' w_i x'' w_i$ such that $y = x w_i x' = x' w_i x''$. If $u_i = w_i x w_i x' w_i x'' w_i$, then we see that $w_i x'$ is a suffix of y and $x' w_i$ is a prefix of y, implying that $w_i x' w_i$ is a new smaller border of u_i . This either contradicts the assumption that v_i is the shortest border of length $> |u_i|/2$, or the assumption that u_i has exactly one border of length $\leq |u_i|/2$. Thus, we have that $u_i = w_i y w_i y w_i$. The shortest border and longest non-overlapping border of $y w_i y$ must be y, by assumption. Additionally, w_i is unbordered, so u_i is of width 5 and i=2. This implies that $w_i = w_2 = w_0 \text{ and } y = w_1.$

 \Leftarrow : Let i be an integer such that $0 < i \le m$. We omit the case when i = 0, since proving $w_i = w_i'$ for all other i is sufficient. Since w_i is the unique border of $u_i = w_i \cdots w_1 w_0 w_1 \cdots w_i$, we have that the shortest border and longest non-overlapping border of u_i is w_i . In other words, we have that $w_i = w_i'$. Suppose i = 2 and $u_2 = w_2 w_1 w_0 w_1 w_2 = w_0 w_1 w_0 w_1 w_0$ where w_0 is the unique border of $w_0 w_1 w_0$. Since w_0 is the unique border of $w_0 w_1 w_0$, it is also the shortest border of u_2 . Additionally, the next longest border of u_2 is $w_0 w_1 w_0$, which is overlapping. So w_0 is also the longest non-overlapping border of u_2 . Thus $w_2 = w_2'$.

Just based on this characterization, finding a recurrence for the number of words with a coinciding smallest and largest BP-factorization seems hard. So we turn to a different, related problem: counting the number of words with a unique border.

5.1 Unique borders

Harju and Nowotka [12] counted the number $B_k(n)$ of length-n words over Σ_k with a unique border, and the number $B_k(n,t)$ of length-n words over Σ_k with a length-t unique border. However, through personal communication with the authors, a small error in one of the proofs leading up to their formula for $B_k(n,t)$ was discovered. Thus, the formula for $B_k(n,t)$ as stated in their paper is incorrect. In this section, we present the correct recurrence for the

number of length-n words with a length-t unique border. We also show that the probability a length-n word has a unique border tends to a constant. See <u>A334600</u> in the OEIS [10] for the sequence $(B_2(n))_{n>0}$.

Suppose w is a word with a unique border u. Then u must be unbordered, and |u| must not exceed half the length of w. If either of these were not true, then w would have more than one border. By combining these ideas, we get Theorem 9 and Theorem 10.

Theorem 9. Let $n > t \ge 1$ be integers. Then the number of length-n words with a unique length-t border satisfies the recurrence

$$B_k(n,t) = \begin{cases} 0, & \text{if } n < 2t; \\ u_t k^{n-2t} - \sum_{i=2t}^{\lfloor n/2 \rfloor} B_k(i,t) k^{n-2i}, & \text{if } n \ge 2t \text{ and } n+t \text{ odd;} \\ u_t k^{n-2t} - B_k((n+t)/2,t) - \sum_{i=2t}^{\lfloor n/2 \rfloor} B_k(i,t) k^{n-2i}, & \text{if } n \ge 2t \text{ and } n+t \text{ even.} \end{cases}$$

Proof. Let w be a length-n word with a unique length-t border u. Since u is the unique border of w, it is unbordered. Thus, we can write w = uvu for some (possibly empty) word v. For n < 2t, we have that $B_k(n,t) = 0$ since u is unbordered and thus cannot overlap itself in w.

Suppose $n \geq 2t$. Let $\overline{B_k}(n,t)$ denote the number of length-n words that have a length-n unbordered border and have a border of length n. Clearly n0 n1 n2 n3 suppose n4 has another border n4 of length n5. Furthermore, suppose that there is no other border n5 with n6 n7 n8 have n7 and n8 have n8 have n9 have n9 have n9. But we could possibly have n9 have n9. The only possible way for n9 for n9 have n

$$\overline{B_k}(n,t) = \sum_{i=2t}^{\lfloor n/2 \rfloor} B_k(i,t) k^{n-2i}.$$

When n+t is even, we compute $\overline{B_k}(n,t)$ in the same fashion, except we also include the case where |u'| = (n+t)/2. This gives us the recurrence,

$$\overline{B_k}(n,t) = B_k((n+t)/2,t) + \sum_{i=2t}^{\lfloor n/2 \rfloor} B_k(i,t)k^{n-2i}.$$

Theorem 10. Let $n \geq 2$ be an integer. Then the number of length-n words with a unique border is

$$B_k(n) = \sum_{t=1}^{\lfloor n/2 \rfloor} B_k(n,t).$$

5.2 Limiting values

We show that the probability that a random word of length n has a unique border tends to a constant. Table 3 shows the behaviour of this probability as k increases.

Let $P_{n,k}$ be the probability that a random word of length n has a unique border. Then

$$P_{n,k} = \frac{B_k(n)}{k^n} = \frac{1}{k^n} \sum_{i=1}^{\lfloor n/2 \rfloor} B_k(n,i).$$

Lemma 11. Let $k \geq 2$ and $n \geq 2t \geq 2$ be integers. Then

$$\frac{B_k(n,t)}{k^n} \le \frac{1}{k^t}.$$

Proof. Let w be a length-n word. Suppose w has a unique border of length t. Since $t \le n/2$, we can write w = uvu for some words u and v where |u| = t. But this means that $B_k(n, t) \le k^{n-t}$, and the lemma follows.

Theorem 12. Let $k \geq 2$ be an integer. Then the limit $P_k = \lim_{n \to \infty} P_{n,k}$ exists.

Proof. Follows from the definition of $P_{n,k}$, Lemma 11, and the direct comparison test for convergence.

k	$\approx P_k$
2	0.5155
3	0.3910
4	0.2922
5	0.2302
6	0.1890
7	0.1599
8	0.1384
9	0.1219
10	0.1089
	:
100	0.0101

Table 3: Probability that a word has a unique border.

References

[1] P. T. Nielsen. A note on bifix-free sequences. *IEEE Trans. Inform. Theory*, IT-19:704–706, 1973.

- [2] The 2015 British Informatics Olympiad (Round 1 Question 1). https://olympiad.org.uk/2015/index.html.
- [3] K. Goto, I. Tomohiro, H. Bannai, and S. Inenaga. Block palindromes: A new generalization of palindromes. In T. Gagie, A. Moffat, G. Navarro, and E. Cuadros-Vargas, editors, *String Processing and Information Retrieval*, volume 11147 of *Lecture Notes in Computer Science*, pages 183–190, Cham, 2018. Springer International Publishing.
- [4] K. Mahalingam, A. Maity, P. Pandoh, and R. Raghavan. Block reversal on finite words. *Theoret. Comput. Sci.*, 894:135–151, 2021.
- [5] K. Mahalingam, A. Maity, and P. Pandoh. Rich words in the block reversal of a word, 2023. arXiv:2302.02109.
- [6] R. Kolpakov and G. Kucherov. Searching for gapped palindromes. *Theoret. Comput. Sci.*, 410(51):5365–5373, 2009.
- [7] M. Régnier. Enumeration of bordered words, le langage de la vache-qui-rit. RAIRO-Theor. Inf. Appl., 26(4):303–317, 1992.
- [8] A. E. Frid, S. Puzynina, and L. Q. Zamboni. On palindromic factorization of words. *Adv. in Appl. Math.*, 50(5):737–748, 2013.
- [9] O. Ravsky. On the palindromic decomposition of binary words. *J. Autom. Lang. Comb.*, 8(1):75–83, 2003.
- [10] N. J. A. Sloane et al. OEIS Foundation Inc. (2022), The On-Line Encyclopedia of Integer Sequences, https://oeis.org.
- [11] P. H. Cording, T. Gagie, M. B. T. Knudsen, and T. Kociumaka. Maximal unbordered factors of random strings. *Theoret. Comput. Sci.*, 852:78–83, 2021.
- [12] T. Harju and D. Nowotka. Counting bordered and primitive words with a fixed weight. *Theoret. Comput. Sci.*, 340(2):273–279, 2005.