

LOG-CONCAVITY OF INFINITE PRODUCT AND INFINITE SUM GENERATING FUNCTIONS

BERNHARD HEIM AND MARKUS NEUHAUSER

ABSTRACT. We expand on the remark by Andrews on the importance of infinite sums and products in combinatorics. Let $\{g_d(n)\}_{d \geq 0, n \geq 1}$ be the double sequences $\sigma_d(n) = \sum_{\ell|n} \ell^d$ or $\psi_d(n) = n^d$. We associate double sequences $\{p^{g_d}(n)\}$ and $\{q^{g_d}(n)\}$, defined as the coefficients of

$$(0.1) \quad \sum_{n=0}^{\infty} p^{g_d}(n) t^n := \prod_{n=1}^{\infty} (1 - t^n)^{-\frac{\sum_{\ell|n} \mu(\ell) g_d(n/\ell)}{n}},$$

$$(0.2) \quad \sum_{n=0}^{\infty} q^{g_d}(n) t^n := \frac{1}{1 - \sum_{n=1}^{\infty} g_d(n) t^n}.$$

These coefficients are related to the number of partitions $p(n) = p^{\sigma_1}(n)$, plane partitions $\text{pp}(n) = p^{\sigma_2}(n)$ of n , and Fibonacci numbers $F_{2n} = q^{\psi_1}(n)$. Let $n \geq 3$ and let $n \equiv 0 \pmod{3}$. Then the coefficients are log-concave at n for almost all d in the exponential (0.1) and geometric (0.2) cases. The coefficients are not log-concave for almost all d in both cases, if $n \equiv 2 \pmod{3}$. Let $n \equiv 1 \pmod{3}$. Then the log-concave property flips for almost all d .

2020 *Mathematics Subject Classification*. Primary 05A17, 11P82; Secondary 05A20.
Key words and phrases. Generating functions, Log-concavity, Partition numbers.

1. INTRODUCTION

In this paper, we study log-concave properties of families of sequences related to infinite product and infinite sum generating functions [Br89, St89, HN22a].

Log-concavity is an important property. For polynomials with positive coefficients, real-rootedness entails log-concavity of all internal coefficients, which implies unimodality. Recent breakthrough works by Huh and his collaborators, using methods in algebraic geometry, have proven the Mason and Heron–Rota–Welsh conjecture on the log-concavity of the chromatic polynomials of graphs, and finally the characteristic polynomials of matroids [AHK18, BHMPW22, Hu12]. We refer to the survey by Kalai [Ka22] on the work by Huh. Note that Zhang [Zh22] proved that the coefficients of the Nekrasov–Okounkov polynomials are almost all unimodal, building on the work by Odlyzko and Richmond [OR85] and Hong and Zhang [HZ21].

We offer an approach for sequences associated with generating functions, where in general, not all coefficients are log-concave. For example, it is well-known that the partition numbers $p(n)$ are log-concave for $n \geq 26$. We encounter $p(n)$ and the number of plane partition numbers $pp(n)$ of n [An98, Kr16] and Fibonacci numbers F_n . A sequence $\{a_n\}_n$ is called log-concave at n_0 , if

$$a_{n_0}^2 - a_{n_0-1} a_{n_0+1} \geq 0.$$

Let $\{g_d(n)\}_{d \geq 0, n \geq 1}$ be a double sequence of positive integers. We examine the coefficients of the associated generating functions of exponential (1.1) and geometric type (1.2):

$$(1.1) \quad \sum_{n=0}^{\infty} p^{g_d}(n) t^n := \exp\left(\sum_{n=1}^{\infty} g_d(n) \frac{t^n}{n}\right) = \prod_{n=1}^{\infty} (1 - t^n)^{-\frac{\alpha_d(n)}{n}},$$

$$(1.2) \quad \sum_{n=0}^{\infty} q^{g_d}(n) t^n := \frac{1}{1 - \sum_{n=1}^{\infty} g_d(n) t^n}.$$

Here $\alpha_d(n) = \sum_{\ell|n} \mu(\ell) g_d(n/\ell)$, where μ is the Möbius function.

The approach offered in this paper, is incited by Andrews' remark ([An98], chapter 6, page 99) in the context of Meinardus' theorem: "Unfortunately not much is known about problems when a series rather than a product is involved". We call n an exception related to a sequence $\{a_n\}_n$, if

$$a_n^2 - a_{n-1} a_{n+1} < 0.$$

The set of all exceptions is denoted by E^a .

To this point only the exponential cases have been studied in the literature. Let $g_d = \sigma_d$. For fixed $d = 1$, we have the number of partitions $p(n) = p^{\sigma_1}(n)$.

Nicolas [Ni78] proved in 1978, that the partition function $p(n)$ is log-concave, if and only if n is not an element of the finite set

$$E^{p^{\sigma_1}} = \{2k + 1 \mid 0 \leq k \leq 12\}.$$

This was proved again by DeSalvo and Pak [DP15]. Both proofs utilize the Rademacher formula for $p(n)$. In [HNT22], we have proven that the plane partition function $pp(n) = p^{\sigma_2}(n)$ is log-concave for almost all n . Finally, based on numerical experiments, we conjectured that

$$E^{p^{\sigma_2}} = \{2k + 1 \mid 0 \leq k \leq 5\}.$$

Recently, the conjecture was proven by Ono, Pujahari, and Rolin [OPR22].

In this paper, we study the similarities between log-concavity properties of the coefficients obtained by the generating function of exponential (1.1) and geometric type (1.2).

1.1. Landscape of Exceptions in the Exponential Cases. We consider log-concavity for $\{p^{g_d}(n)\}$. We recall the results obtained in [HN22a] and [HN22b]. Note, the information on $d = 0$ is new. Numerical investigations indicate that

$$E^{p^{\sigma_0}} = \{2k + 1 \mid 0 \leq k \leq 371\} \setminus \{717, 723, 729, 735, 741\}$$

tested up to $n = 2500$. Further, for $0 \leq d \leq 5$ the cardinality of $E^{p^{\sigma_d}}$ seems to be decreasing: $367 > 13 > 6 > 4 > 2 \geq 2$. But $|E^{p^{\sigma_6}}| \geq 3$. We refer to Table 1. The case $g_d = \psi_d$, if we see Table 2, reveals the similar pattern.

$n \setminus d$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
1	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
2																		
3	•	•	•	•														
4							•	•	•	•	•	•	•	•	•	•	•	•
5	•	•	•	•	•	•	•	•	•	•								
6																		
7	•	•	•	•								•	•	•	•	•	•	•
8										•	•	•	•	•	•	•	•	•
9	•	•	•															
10																	•	•
11	•	•	•										•	•	•	•	•	•
12																		
13	•	•																
14																•	•	•

TABLE 1. Exceptions for $p^{\sigma_d}(n)$ for $0 \leq d \leq 17$ and $1 \leq n \leq 14$.

$n \backslash d$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
1		•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
2																		
3																		
4					•	•	•	•	•	•	•	•	•	•	•	•	•	•
5																		
6																		
7												•	•	•	•	•	•	•
8									•	•	•	•	•	•	•	•	•	•
9																		
10																	•	•
11													•	•	•	•	•	•
12																		
13																		
14																•	•	•

TABLE 2. Exceptions for $p^{\psi_d}(n)$ for $0 \leq d \leq 17$ and $1 \leq n \leq 14$.

Now, fixing n and studying log-concavity, reveals a new phenomenon. Let $n \geq 3$. Let $g_d(n) = \sigma_d(n)$ or $\psi_d(n)$. Then the set of all exceptions for all d is finite, if and only if $n \equiv 0 \pmod{3}$. More generally [HN22b], let $\{g_d(n)\}_{d \geq 1, n \geq 1}$ be positive real numbers satisfying $g_d(1) = 1$ and

$$0 \leq g_d(n) - n^d \leq (g_1(n) - n) (n - 1)^{d-1}.$$

Let $n \geq 3$. Then for almost all d , $p^{g_d}(n)$ is log-concave at n , if and only if n is divisible by 3. Moreover, explicit bounds are given. It would be interesting to examine the results of this paper in the context of generalized Laguerre-Pólya functions and Jensen polynomials [Wa22].

1.2. Landscape of Exceptions in the Geometric Cases. At first glance, the *geometric* case, see Table 3 and Table 4, seems not to reveal much structure. Nevertheless, we recall that $q^{\psi_1}(n) = F_{2n}$ can be identified with the sequence of the $2n$ th Fibonacci numbers, which is log-concave for $n > 1$. This follows from the fact that $q^{\psi_1}(n) = U_{n-1}(\frac{3}{2})$, where $U_n(x)$ is the n th Chebyshev polynomial of the second kind. Thus, we have some kind of analogue to Nicolas' result. Thus far, for $d = 2$ and $\psi_2(n)$, we expect infinitely many exceptions. Nevertheless, by fixing $n \geq 3$ we obtain the following new result.

We have the *geometric* cases for $g_d(n) = \sigma_d(n)$ in Table 3 and $g_d(n) = n^d$ in Table 4.

$n \setminus d$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
1	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
2																		
3	•	•	•	•	•													
4						•	•	•	•	•	•	•	•	•	•	•	•	•
5	•	•	•	•	•	•												
6																		
7	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
8																		
9	•	•	•	•	•													
10						•	•	•	•	•	•	•	•	•	•	•	•	•
11	•	•	•	•	•	•	•											
12																		
13	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
14						•	•											

TABLE 3. Exceptions for $q^{\sigma_d}(n)$ for $0 \leq d \leq 17$ and $1 \leq n \leq 14$.

$n \setminus d$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
1	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
2																		
3																		
4			•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
5			•	•														
6			•															
7					•	•	•	•	•	•	•	•	•	•	•	•	•	•
8				•	•	•												
9				•														
10			•			•	•	•	•	•	•	•	•	•	•	•	•	•
11			•		•	•	•											
12			•	•														
13				•		•	•	•	•	•	•	•	•	•	•	•	•	•
14					•	•	•											

TABLE 4. Exceptions for $q^{\psi_d}(n)$ for $0 \leq d \leq 17$ and $1 \leq n \leq 14$.

1.3. Main Results. In this paper, we prove the following:

Theorem 1.1. *Let $\{g_d(n)\}_{d \geq 0, n \geq 1}$ be a double sequence of positive real numbers with $g_d(1) = 1$ for all d and*

$$(1.3) \quad 0 \leq g_d(n) - n^d \leq (g_0(n) - 1) (n - 1)^d.$$

Suppose there is an $0 < r \leq 1$, such that $q^{g_0}(n) \leq r^{-n}$. Let $n \geq 3$ and $D^g(n)$ be defined by

$$(1.4) \quad D^g(n) := -2 \log_{9/8}(r) n,$$

$$(1.5) \quad D^g(n) := \log_{9/8}(3) - 2n \log_{9/8}(r) - \log_{9/8}(n + 2)$$

for $n \equiv 0 \pmod{3}$ in (1.4) and $n \equiv 1 \pmod{3}$ in (1.5). Further, let $n \equiv 2 \pmod{3}$ and $n \neq 5$. Then $D^g(n)$ is defined as

$$\log_{9/8}(2) - (n + 1) \log_{9/8}(r).$$

Moreover, $D^g(5) := \log_{9/8}(2q^{g_0}(4)q^{g_0}(6))$.

Let $d > D^g(n)$. Then

$$(1.6) \quad \frac{(q^{g_d}(n))^2}{q^{g_d}(n-1) q^{g_d}(n+1)} < 1, \text{ if and only if } n \equiv 1 \pmod{3}.$$

The double sequences $\{g_d(n)\}$ given by $\psi_d(n) = n^d$ and $\sigma_d(n) = \sum_{\ell|n} \ell^d$ satisfy (1.3). In the case $\psi_0(n)$, we have $\frac{t}{1-t} = \sum_{n=1}^{\infty} t^n$ and $\frac{1-t}{1-t^2} = 1 + \sum_{n=1}^{\infty} 2^{n-1} t^n$. Therefore, $q^{\psi_0}(n) \leq 2^n$ for $n \geq 1$. We can apply Theorem 1.1 with $r = \frac{1}{2}$.

Let

$$D^\psi(n) := \begin{cases} 2n \log_{9/8}(2), & n \equiv 0 \pmod{3}, \\ 2n \log_{9/8}(2) + \log_{9/8}(3) - \log_{9/8}(n + 2), & n \equiv 1 \pmod{3}, \\ (n + 2) \log_{9/8}(2), & n \equiv 2 \pmod{3}, n \neq 5, \\ \log_{9/8}(512), & n = 5. \end{cases}$$

Corollary 1.2. *Let $n \geq 3$. Let $d > D^\psi(n)$. Then*

$$(1.7) \quad \frac{(q^{\psi_d}(n))^2}{q^{\psi_d}(n-1) q^{\psi_d}(n+1)} < 1, \text{ if and only if } n \equiv 1 \pmod{3}.$$

For $g_0 = \sigma_0$, we obtain $\frac{1}{1 - \sum_{n=1}^{\infty} \sigma_0(n) t^n} = 1 + t + 3t^2 + 7t^3 + 18t^4 + 43t^5 + 108t^6 + \dots$. Obviously, $\sigma_0(n) \leq n$. Then $\sum_{n=1}^{\infty} n t^n = \frac{t}{(1-t)^2}$ and the radius of convergence of the series expansion of $\frac{1-t}{1 - \frac{t}{(1-t)^2}}$ is $(3 - \sqrt{5})/2 > 1/3$. Analyzing the coefficients shows that we can choose any $0 < r < (3 - \sqrt{5})/2$. For simplicity, we take $r = \frac{1}{3}$ and

obtain $q^{\sigma_0}(n) \leq 3^n$. We define $D^\sigma(n)$ for $n \geq 3$ and $n \neq 5$ by

$$\begin{cases} 2 \log_{9/8}(3) n, & n \equiv 0 \pmod{3}, \\ (2n+1) \log_{9/8}(3) - \log_{9/8}(n+2), & n \equiv 1 \pmod{3}, \\ \log_{9/8}(2) + (n+1) \log_{9/8}(3), & n \equiv 2 \pmod{3}. \end{cases}$$

Further, $D^\sigma(5) := \log_{9/8}(3888)$.

Corollary 1.3. *Let $n \geq 3$. Let $d > D^\sigma(n)$. Then*

$$(1.8) \quad \frac{(q^{\sigma_d}(n))^2}{q^{\sigma_d}(n-1) q^{\sigma_d}(n+1)} < 1, \text{ if and only if } n \equiv 1 \pmod{3}.$$

2. PROOF OF THEOREM 1.1

Let g_d be fixed satisfying (1.3). To simplify notation, we put $q_d(n) = q^{g_d}(n)$. We have

$$\frac{1}{1 - \sum_{n=1}^{\infty} g_d(n) t^n} = 1 + \sum_{n=1}^{\infty} \sum_{k \leq n} \sum_{\substack{m_1, \dots, m_k \geq 1 \\ m_1 + \dots + m_k = n}} g_d(m_1) \cdots g_d(m_k) t^n = \sum_{n=0}^{\infty} q_d(n) t^n.$$

Therefore,

$$q_d(n) = \sum_{k \leq n} \sum_{\substack{m_1, \dots, m_k \geq 1 \\ m_1 + \dots + m_k = n}} g_d(m_1) \cdots g_d(m_k)$$

for $n \geq 1$.

2.1. Two Lemmata. It is known [HN22a] that:

Lemma 2.1. *Let $n \geq 2$. Then*

$$\max_{\substack{m_1, \dots, m_k \geq 1 \\ m_1 + \dots + m_k = n}} m_1 \cdots m_k = \begin{cases} 3^{n/3}, & n \equiv 0 \pmod{3}, \\ 4 \cdot 3^{(n-4)/3}, & n \equiv 1 \pmod{3}, \\ 2 \cdot 3^{(n-2)/3}, & n \equiv 2 \pmod{3}. \end{cases}$$

For $n \geq 6$, $n \not\equiv 2 \pmod{3}$, the second largest products are

$$\begin{aligned} 8 \cdot 3^{(n-6)/3}, n &\equiv 0 \pmod{3}, \\ 10 \cdot 3^{(n-7)/3}, n &\equiv 1 \pmod{3} \end{aligned}$$

and 3 for $n = 4$.

Further, we provide an extension of a result from [HN22a, HN22b].

Lemma 2.2. For $n \geq 2$

$$\begin{aligned} 3^{dn/3} &\leq q_d(n) \leq 3^{dn/3} q_0(n), n \equiv 0 \pmod{3}, \\ (4 \cdot 3^{(n-4)/3})^d \frac{(n-1)(n+8)}{18} &\leq q_d(n) \leq (4 \cdot 3^{(n-4)/3})^d q_0(n), n \equiv 1 \pmod{3}, \\ (2 \cdot 3^{(n-2)/3})^d \frac{n+1}{3} &\leq q_d(n) \leq (2 \cdot 3^{(n-2)/3})^d q_0(n), n \equiv 2 \pmod{3}. \end{aligned}$$

Additionally, for $n \geq 6$, $n \not\equiv 2 \pmod{3}$, we have

$$\begin{aligned} q_d(n) &\leq 3^{nd/3} + (8 \cdot 3^{n/3-2})^d q_0(n), n \equiv 0 \pmod{3}, \\ q_d(n) &\leq (4 \cdot 3^{(n-4)/3})^d \frac{(n-1)(n+8)}{18} + (10 \cdot 3^{(n-7)/3})^d q_0(n), n \equiv 1 \pmod{3}, \end{aligned}$$

and $q_d(4) \leq 2 \cdot 4^d + 3^d q_0(4)$.

Proof. Since $m_1 \cdots m_k \leq \max_{\substack{m_1, \dots, m_k \geq 1 \\ m_1 + \dots + m_k = n}} m_1 \cdots m_k$, the upper bounds should be obvious as $g_d(n) \leq n^d + (g_0(n) - 1)(n - 1)^d \leq g_0(n) n^d$. For the lower bounds, we obtain

$$\sum_{k \leq n} \sum_{\substack{m_1, \dots, m_k \geq 1 \\ m_1 + \dots + m_k = n}} (m_1 \cdots m_k)^d \geq S(n) \max_{k \leq n} \max_{\substack{m_1, \dots, m_k \geq 1 \\ m_1 + \dots + m_k = n}} (m_1 \cdots m_k)^d$$

where $S(n)$ is the number of $m_1 + \dots + m_k = n$, which yield the maximal product. Therefore,

$$S(n) = \begin{cases} 1, & n \equiv 0 \pmod{3}, \\ \frac{n-1}{3} + \binom{(n+2)/3}{2} = \frac{(n-1)(n+8)}{18}, & n \equiv 1 \pmod{3}, \\ \frac{n+1}{3}, & n \equiv 2 \pmod{3}. \end{cases}$$

For the refined upper bounds, we consider $g_d(n) \leq n^d + (g_0(n) - 1)(n - 1)^d$. Then $g_d(m_1) \cdots g_d(m_k) \leq g_0(m_1) \cdots g_0(m_k) (m_1 \cdots m_k)^d$ and for the maximal values

$$\begin{aligned} &g_d(m_1) \cdots g_d(m_k) \\ &\leq \left(m_1^d + (g_0(m_1) - 1)(m_1 - 1)^d \right) \cdots \left(m_k^d + (g_0(m_k) - 1)(m_k - 1)^d \right). \end{aligned}$$

Therefore,

$$q_d(n) \leq \left(\max_{k \leq n} \max_{\substack{m_1, \dots, m_k \geq 1 \\ m_1 + \dots + m_k = n}} m_1 \cdots m_k \right)^d S(n) + (Z(n))^d q_0(n)$$

where $Z(n)$ is the second largest product $m_1 \cdots m_k$ of all $m_1 + \dots + m_k = n$. \square

2.2. Proof of Theorem 1.1. We consider the cases $n \equiv 0, 1, 2 \pmod{3}$ and $n = 5$ separately.

2.2.1. Let $n \equiv 0 \pmod{3}$. Then

$$\begin{aligned} \frac{(q_d(n))^2}{q_d(n-1)q_d(n+1)} &\geq \frac{3^{2dn/3}}{(2 \cdot 3^{(n-3)/3} \cdot 4 \cdot 3^{(n-3)/3})^d q_0(n-1)q_0(n+1)} \\ &\geq \left(\frac{9}{8}\right)^d \frac{1}{r^{-2n}} \geq 1 \end{aligned}$$

for $d \geq -2 \log_{9/8}(r)n$.

2.2.2. Let $n \equiv 1 \pmod{3}$. Then

$$\begin{aligned} \frac{(q_d(n))^2}{q_d(n-1)q_d(n+1)} &\leq \frac{\left((4 \cdot 3^{(n-4)/3})^d q_0(n)\right)^2}{3^{(n-1)d/3} \cdot (2 \cdot 3^{(n-1)/3})^d \frac{n+2}{3}} \\ &\leq \left(\frac{8}{9}\right)^d \frac{3r^{-2n}}{n+2} < 1 \end{aligned}$$

for $d > \log_{9/8}(3) - 2n \log_{9/8}(r) - \log_{9/8}(n+2)$.

2.2.3. Let $n \equiv 2 \pmod{3}$ and $n \neq 5$. Then

$$(2.1) \quad \frac{(q_d(n))^2}{q_d(n-1)q_d(n+1)} \geq \frac{\left((2 \cdot 3^{(n-2)/3})^d \frac{n+1}{3}\right)^2}{A(d,n) B(d,n)},$$

where

$$\begin{aligned} A(d,n) &= (4 \cdot 3^{(n-5)/3})^d \frac{(n-2)(n+7)}{18} + (10 \cdot 3^{(n-8)/3})^d q_0(n-1), \\ B(d,n) &= 3^{(n+1)d/3} + (8 \cdot 3^{(n-5)/3})^d q_0(n+1). \end{aligned}$$

Then as a lower bound for the expression on the right hand side of (2.1) we obtain the following:

$$\begin{aligned} &\frac{\left(\frac{n+1}{3}\right)^2}{\left(\frac{(n-2)(n+7)}{18} + \left(\frac{5}{6}\right)^d r^{1-n}\right) \left(1 + \left(\frac{8}{9}\right)^d r^{-n-1}\right)} \\ &\geq \frac{\left(\frac{n+1}{3}\right)^2}{\left(\frac{(n-2)(n+7)}{18} + \frac{4}{9}\right) \frac{3}{2}} \geq \frac{4(n+1)^2}{3(n+6)(n-1)} = 1 + \frac{\left(n - \frac{7}{2}\right)^2 + \frac{39}{4}}{3(n+6)(n-1)} \geq 1 \end{aligned}$$

for

$$\begin{aligned} d &\geq \max \left\{ \log_{6/5}(9/4) - (n-1) \log_{6/5}(r), \log_{9/8}(2) - (n+1) \log_{9/8}(r) \right\} \\ &= \log_{9/8}(2) - (n+1) \log_{9/8}(r) \end{aligned}$$

for $0 < r \leq 1$ as $0 \leq -\log_{6/5}(r) \leq -\log_{9/8}(r)$ and $\log_{6/5}(9/4) < \log_{9/8}(2)$.

2.2.4. Let $n = 5$. We have

$$\begin{aligned} \frac{(q_d(5))^2}{q_d(4)q_d(6)} &\geq \frac{4 \cdot 36^d}{(2 \cdot 4^d + 3^d q_0(4))(9^d + 8^d q_0(6))} \\ &\geq \frac{4 \cdot 36^d}{2 \cdot 36^d + 4 \cdot 32^d q_0(4)q_0(6)} \geq 1 \end{aligned}$$

for $d \geq \log_{9/8}(2q_0(4)q_0(6))$.

3. FINAL REMARKS

Let us examine $\{q^{g_d}(n)\}$. There are no exceptions for $g_d = \sigma_d$ or ψ_d for $n = 2$, since

$$(q^{g_d}(2))^2 - q^{g_d}(1)q^{g_d}(3) = (g_d(2))^2 - g_d(3).$$

Challenge 1. We consider the exponential case for $d = 0$ and σ_0 . We expect $E^{p^{\sigma_0}}$ to be finite. Moreover, numerical experiments (tested up to $n = 2500$) suggest that

$$E^{p^{\sigma_0}} = \{2k + 1 \mid 0 \leq k \leq 371\} \setminus \{717, 723, 729, 735, 741\}.$$

Challenge 2. We consider the geometric case. We have $E^{q^{\psi_0}} = E^{q^{\psi_1}} = \{1\}$, since $q^{\psi_0}(n) = 2^{n-1}$ and $q^{\psi_1}(n) = F_{2n}$. For $E^{q^{\psi_2}}$, we expect infinitely many exceptions and non-exceptions.

Challenge 3 (Geometric case). Let σ_d for $0 \leq d \leq 4$ be given. Then all the odd numbers n up to 10^4 are exceptions. Note that for $d = 5$ some even numbers also appear as exceptions. For example,

$$(q^{\sigma_5}(10))^2 - q^{\sigma_5}(9)q^{\sigma_5}(11) < 0.$$

Nevertheless, it seems that the set of exceptions for each σ_d is infinite.

REFERENCES

- [AHK18] K. Adiprasito, J. Huh, and E. Katz: *Hodge theory for combinatorial geometries*. *Annals of Mathematics* **188** (2018), 381–452.
- [An98] G. E. Andrews: *The theory of partitions*. Cambridge Univ. Press, Cambridge (1998).
- [BHMPW22] T. Braden, J. Huh, J. P. Matherne, N. Proudfoot, B. Wang: *Singular Hodge theory for combinatorial geometries*. arXiv:2010.06088v3.
- [Br89] F. Brenti: *Unimodal, log-concave and Pólya frequency sequences in combinatorics*. *Mem. Am. Math. Soc.* **413** (1989).
- [DP15] S. DeSalvo, I. Pak: *Log-concavity of the partition function*. *Ramanujan J.* **38** (2015), 61–73.
- [HN22a] B. Heim, M. Neuhauser: *Log-concavity of infinite product generating functions*. *Res. Number Theory* **8** No. 3 (2022), Paper No. 53, 14 pp.
- [HN22b] B. Heim, M. Neuhauser: *Turán inequalities for infinite product generating functions*. arXiv:2207.09409v1 [math.CO] 19 Jul 2022.

- [HNT22] B. Heim, M. Neuhauser, R. Tröger: *Inequalities for plane partitions*. Annals of Combinatorics (2022), 22 pp. doi:10.1007/s00026-022-00604-4.
- [HZ21] L. Hong, S. Zhang: *Towards Heim and Neuhauser’s unimodality conjecture on the Nekrasov–Okounkov polynomials*. Res. Number Theory **7** No. 1 (2021), Paper No. 17, 11 pp.
- [Hu12] J. Huh: *Milnor numbers of projective hypersurfaces and the chromatic polynomial of graphs*. Journal of the American Mathematical Society **25** (2012), 907–927.
- [Ka22] G. Kalai: *The work of June Huh*. Proc. Int. Cong. Math. 2022. Vol. 1. Preliminary version.
- [Kr16] C. Krattenthaler: *Plane partitions in the work of Richard Stanley and his school*. In: P. Hersh, T. Lam, P. Pylyavskyy, V. Reiner (eds.) The mathematical legacy of Richard Stanley. Amer. Math. Soc., R. I. (2016), 246–277.
- [Ni78] J.-L. Nicolas: *Sur les entiers N pour lesquels il y a beaucoup des groupes abéliens d’ordre N* . Ann. Inst. Fourier **28** No. 4 (1978), 1–16.
- [OR85] A. M. Odlyzko, L. B. Richmond: *On the unimodality of high convolutions of discrete distributions*. Ann. Probab. **13** No. 1 (1985), 299–306.
- [OPR22] K. Ono, S. Pujahari, L. Rolin: *Turán inequalities for the plane partition function*. Advances in Mathematics **409** Part B (2022). doi:10.1016/j.aim.2022.108692.
- [St89] R. Stanley: *Log-concave and unimodal sequences in algebra, combinatorics, and geometry*. In: M. F. Capobianco, M. G. Guan, D. F. Hsu, F. Tian (eds.) Graph theory and its applications: East and West. Proceedings of the first China–USA international conference, held in Jinan, China, June 9–20, 1986. Ann. New York Acad. Sci. **576**, New York Acad. Sci., New York (1989), 500–535.
- [St99] R. Stanley: *Enumerative combinatorics*. Vol. 2. With a foreword by Gian-Carlo Rota and appendix 1 by Sergey Fomin, Cambridge Studies in Advanced Mathematics **62**, Cambridge University Press, Cambridge (1999).
- [Wa22] I. Wagner: *On a new class of Laguerre–Pólya type functions with applications in number theory*. Pacific Journal of Mathematics Vol. **320** No. 1 (2022), 177–192.
- [Zh22] S. Zhang: *Log-concavity in powers of infinite series close to $(1 - z)^{-1}$* . Res. Number Theory **8** No. 1 (2022), Paper No. 66, 17 pp.

FACULTY OF MATHEMATICAL AND NATURAL SCIENCES, MATHEMATICAL INSTITUTE, UNIVERSITY OF COLOGNE, WEYERTAL 86–90, 50931 COLOGNE, GERMANY

Email address: bheim@math.uni-koeln.de

LEHRSTUHL A FÜR MATHEMATIK, RWTH AACHEN UNIVERSITY, 52056 AACHEN, GERMANY

Email address: bernhard.heim@rwth-aachen.de

KUTAISI INTERNATIONAL UNIVERSITY, 5/7, YOUTH AVENUE, KUTAISI, 4600 GEORGIA

LEHRSTUHL A FÜR MATHEMATIK, RWTH AACHEN UNIVERSITY, 52056 AACHEN, GERMANY

Email address: markus.neuhauser@kiu.edu.ge