

THE CHANGE-MAKING PROBLEM FOR SIX COIN VALUES AND BEYOND

CORNELIA A. VAN COTT AND QIYU ZHANG

ABSTRACT. The change-making problem asks: given a positive integer v and a collection \mathcal{C} of integer coin values $c_1 = 1 < c_2 < c_3 < \dots < c_n$, what is the minimum number of coins needed to represent v with coin values from \mathcal{C} ? For some coin systems \mathcal{C} , the greedy algorithm finds a representation with a minimum number of coins for all v . We call such coin systems *orderly*. However, there are coin systems where the greedy algorithm fails to always produce a minimal representation. Over the past fifty years, progress has been made on the change-making problem, including finding a characterization of all orderly coin systems with 3, 4, and 5 coin values. We characterize orderly coin systems with 6 coin values, and we make generalizations to orderly coin systems with n coin values.

1. INTRODUCTION

As customers, we trust store cashiers to give us our change using the minimum possible numbers of coins. Most of the time, cashiers get this done in a heartbeat. The secret to their success is that they use the so-called *greedy algorithm* to make change. Given a positive integer v , this algorithm works as follows. Find the largest coin value less than or equal to v . Take as many coins of that value as possible without the sum of the coin values exceeding v . Then move to the next largest coin denomination and repeat the process until the entire collection's value sums up to v . For both the United States' coin system $(1, 5, 10, 25)$ and the EU coin system $(1, 2, 5, 10, 20, 50, 100, 200)$, the greedy algorithm always produces the minimum number of coins.

Since the greedy algorithm serves us so well with these familiar coin systems, it often comes as a surprise that the greedy algorithm fails to be optimal in other settings. Suppose, for example, that the United States replaced the dime and quarter with 15 and 20 cent coins, making the coin system $(1, 5, 15, 20)$. If we needed 30 cents change, the greedy algorithm will produce a solution with three coins: $20 + 5 + 5$. But we can get the job done with just two coins: $15 + 15$. What algorithm should we use in general to get a representation with a minimum number of coins? This is the so-called *change-making problem*.

Stated formally, the change-making problem is as follows. Given a positive integer v and a coin system $\mathcal{C} = (c_1, c_2, c_3, \dots, c_n)$ with $c_1 = 1 < c_2 < c_3 < \dots < c_n$, what is the minimum number of coins needed to represent v with coins from \mathcal{C} ? Denote this minimum number via $\text{opt}_{\mathcal{C}}(v)$. We assume that there are an unlimited number of each type of coin available. The fact that the smallest coin value is always $c_1 = 1$ ensures that each positive integer v has at least one representation.

A collection of coins with total value v is called an *optimal solution* if the collection has a minimum number of coins. An optimal solution need not be unique.

For instance, suppose the coin system is $\mathcal{C} = (1, 3, 4)$. Then there are two optimal solutions for $v = 9$ as follows: $3 + 3 + 3$ and $4 + 4 + 1$.

Let $grd_{\mathcal{C}}(v)$ denote the number of coins used when creating v using the greedy algorithm with the coin system \mathcal{C} . For some coin systems, the change-making problem is solved via the greedy algorithm. We give a term to these coin systems.

Definition 1.1. *A coin system $\mathcal{C} = (1, c_2, c_3, \dots, c_n)$ is **orderly**¹ if for all $v > 0$, $grd_{\mathcal{C}}(v) = opt_{\mathcal{C}}(v)$.*

The United States' coin system and the EU coin system are both orderly (we will justify this later), while the coin system $(1, 5, 15, 20)$ is not orderly since the greedy algorithm failed to be optimal for $v = 30$. For a non-orderly coin system \mathcal{C} , any value v such that $grd_{\mathcal{C}}(v) \neq opt_{\mathcal{C}}(v)$ is called a **counterexample**.

The change-making problem is a special case of the knapsack problem. In this more general setting, suppose you have n types of objects available to pack into a knapsack, and the monetary value of each object is denoted by c_1, c_2, \dots, c_n , respectively. These objects have weights denoted by w_1, w_2, \dots, w_n , respectively. The knapsack problem asks how we can pack the knapsack in such a way that the monetary value of the content equals some value v and the total weight of the packed knapsack is *minimized*. The change-making problem, then, is the special case in which the items have weight $w_i = 1$ for all i . For a discussion of the knapsack problem, see survey articles [2, 3] and books [12, 15].

In Section 2, we review the progress made on the change-making problem over the last several decades and the primary tools used to study coin systems in general. Notably, characterizations of orderly coin systems with 3, 4, and 5 coin values have been established (Theorems 2.8, 2.9, and 2.10). One goal of our work here is to give a characterization of orderly coin systems with 6 coin values.

A useful way to study coin systems is through considering their prefix coin systems. Given any orderly coin system $\mathcal{C} = (1, c_2, c_3, \dots, c_n)$, a **prefix coin system** of \mathcal{C} is defined as $\mathcal{C}' = (1, c_2, c_3, \dots, c_i)$ where $1 \leq i \leq n$. The system \mathcal{C}' for $1 \leq i < n$ may or may not be orderly. We can associate to any n -value coin system a string of n symbols where the i th symbol is $+$ if the length i prefix coin system is orderly and is $-$ otherwise. So for instance, the coin system $(1, 2, 5, 6, 10)$ has pattern $(+ + + - +)$, because all the prefix coin systems are orderly except for the fourth $(1, 2, 5, 6)$ which has a counterexample of 10. Every coin system with 4 or more coin values has a pattern that begins with $(+ + +)$ (Theorem 2.5). Therefore, orderly coin systems with 6 coin values have one of four possible $+/-$ patterns. Our result is as follows.

Theorem 1.2. *A coin system $\mathcal{C} = (1, c_2, c_3, c_4, c_5, c_6)$ is orderly if and only if \mathcal{C} is one of the following:*

- (1) *Coin systems with pattern $(+ + + + - +)$.*
 - (a) $(1, 2, 3, a, a + 1, 2a)$, where $a \geq 5$
 - (b) $(1, a, 2a, b, b + a, 2b)$,
where $a \geq 2$, $b \geq 3a - 1$, $b \neq 3a$, and $grd_{\mathcal{C}}(2ma) \leq m$ where $m = \lceil \frac{b}{2a} \rceil$
 - (c) $(1, a, 2a - 1, b, b + a - 1, 2b - 1)$,
where $a \geq 2$, $b \geq 3a - 1$, $grd_{\mathcal{C}}(m'(2a - 1)) \leq m'$ where $m' = \lceil \frac{b}{2a - 1} \rceil$

¹Other terms used in place of *orderly* in the literature are canonical, greedy, and standard.

- (2) *Coin systems with pattern $(+++--)$.*
 (a) $(1, a, 2a - 1, m(2a - 1) - (a - 1), m(2a - 1), (2m - 1)(2a - 1))$,
 where $1 < m < a$.
 (b) $(1, a, 2a, m(2a - 1) - (a - 1), m(2a - 1) + 1, (2m - 1)(2a - 1) + 1)$,
 where $1 < m \leq a$.
- (3) *Coin systems with pattern $(++++)$ or $(+++--)$.*
 $(1, c_2, c_3, c_4, c_5)$ is orderly and $\text{grd}_{\mathcal{C}}(mc_5) \leq m$ where $m = \lceil \frac{c_6}{c_5} \rceil$.

While orderly coin systems of size 3, 4, and 5 provide a foundation for understanding the structure of orderly coin systems, they yield limited intuition about larger coin systems, because they are so constrained. Thus the result here for coin systems with 6 coin values is critical in understanding the general structure of orderly coin systems.

In Section 4, we generalize to coin systems with n values with pattern $(+++ - \dots -)$. All coin systems we observed with this pattern have a particular structure; we call them **fixed gap coin systems** (Definition 4.1). We give three infinite families of fixed gap coin systems \mathcal{D} , \mathcal{E} , and \mathcal{F} (see Theorems 4.3, 4.5, and 4.7). Coin systems in \mathcal{D} have $3r + 2$ coin types where $r \geq 1$, while \mathcal{E} and \mathcal{F} both have $3r$ coin types where $r \geq 2$. Every coin system with pattern $(+++ - \dots -)$ that we observed belonged to one of these three infinite families, so we make the following conjecture.

Conjecture 1.3. *A coin system has pattern $(+++ - \dots -)$ only if it is a member of one of the families \mathcal{D} , \mathcal{E} , or \mathcal{F} (defined in Theorems 4.3, 4.5, and 4.7). In particular, there are no coin systems with pattern $(+++ - \dots -)$ that have $3r + 1$ coin values where $r \geq 2$.*

In the final stages of our work, we were made aware of another paper recently posted that also characterizes orderly coin systems with 6 coin values (see Proposition 1 in [17]). The two characterizations (Theorem 1.2 here in this paper and Proposition 1 of [17]) have different perspectives and use different tools. Thus we believe that our result here stands side-by-side with the work of [17], each shedding a complementary light on the change-making problem.

2. BACKGROUND

A priori, checking whether an arbitrary coin system \mathcal{C} is orderly is an infinite task, requiring us to check that $\text{grd}_{\mathcal{C}}(v) = \text{opt}_{\mathcal{C}}(v)$ for every $v > 0$. The following result narrows down the location of the smallest counterexample of a coin system, if it exists.

Theorem 2.1. [13] *If the coin system $\mathcal{C} = (1, c_2, c_3, \dots, c_n)$ is not orderly, then the smallest counterexample v lies in the range: $c_3 < v < c_{n-1} + c_n$.*

The following result improves on the previous one and enables us to determine whether a coin system is orderly in $O(n^3)$ time. We first set up some notation and terminology.

Say we have a coin system $\mathcal{C} = (1, c_2, c_3, \dots, c_n)$, and suppose we found a way to represent the value v via coins as follows: $v = x_1 + x_2c_2 + \dots + x_nc_n$ where x_i are each positive integers. We express this representation of v by coins in \mathcal{C} via a vector of the x_i 's as follows: $(x_1, x_2, \dots, x_n)_{\mathcal{C}}$. Such a vector representing v need not be unique, and we can order all such representations as follows.

Definition 2.2. Let v be a positive integer and suppose that $\mathbf{x} = (x_1, x_2, \dots, x_n)_C$ and $\mathbf{x}' = (x'_1, x'_2, \dots, x'_n)_C$ are two representations of the value v with respect to the coin system $\mathcal{C} = (1, c_2, c_3, \dots, c_n)$. The representation \mathbf{x} is **lexicographically smaller** than \mathbf{x}' if $x_i = x'_i$ for $1 \leq i \leq k < n$ for some k and $x_{k+1} < x'_{k+1}$.

Theorem 2.3. [18] Let $\mathcal{C} = (1, c_2, c_3, \dots, c_n)$ be a non-orderly coin system. Let w be the minimum counterexample for \mathcal{C} . Suppose that the lexicographically smallest optimal solution for w in \mathcal{C} is

$$(0, 0, \dots, 0, x_i, x_{i+1}, \dots, x_j, 0, 0, \dots, 0)_C$$

where $1 \leq i \leq j < n$ and $x_i, x_j > 0$. Then the greedy representation for $c_{j+1} - 1$ in \mathcal{C} is

$$(y_1, y_2, \dots, y_{i-1}, x_i - 1, x_{i+1}, \dots, x_j, 0, 0, \dots, 0)_C$$

where each y_i is a nonnegative integer.

Now we discuss properties and results related to prefix coin systems. We have a term for orderly coin systems where all prefix coin systems are orderly.

Definition 2.4. A coin system $\mathcal{C} = (1, c_2, c_3, \dots, c_n)$ is **totally orderly** if each coin system $(1, c_2, c_3, \dots, c_i)$ is orderly for all $i = 1, 2, \dots, n$.

Sometimes knowing information about a prefix coin system yields information about the coin system as a whole.

Theorem 2.5. [1] If $\mathcal{C} = (1, c_2, c_3, \dots, c_n)$ is orderly, then the coin system $(1, c_2, c_k)$ is orderly for all $3 \leq k \leq n$. In particular, $(1, c_2, c_3)$ is orderly.

For example, consider the coin system $(1, 3, 4)$. The value 6 is a counterexample for this coin system (the greedy solution is $4 + 1 + 1$ while an optimal solution is $3 + 3$). Thus the pattern for this coin system is $(+ + -)$. One might think that adding new coin values larger than 4 to the coin system could “fix” it and make it orderly. But Theorem 2.5 tells us that this is impossible. There is no collection of higher coin values which one could add to the coin system $(1, 3, 4)$ which would cause the resulting coin system to be orderly. A more involved example involving 7-coins is as follows.

Theorem 2.6. [1] There are no coin systems with the pattern $(+ + + - + - +)$.

The following theorem (sometimes called the *One Point Theorem*) gives a necessary and sufficient condition to determine whether a coin system $(1, c_2, \dots, c_{n-1}, c_n)$ is orderly, if one already knows that the prefix coin system $(1, c_2, \dots, c_{n-1})$ is orderly.

Theorem 2.7 (One Point Theorem). [1, 10, 14] Suppose that $\mathcal{C}' = (1, c_2, \dots, c_{n-1})$ is orderly and $c_{n-1} < c_n$. Let $m = \lfloor \frac{c_n}{c_{n-1}} \rfloor$. The following are equivalent:

- (1) The coin system $\mathcal{C} = (1, c_2, \dots, c_{n-1}, c_n)$ is orderly.
- (2) $\text{grd}_{\mathcal{C}}(mc_{n-1}) = \text{opt}_{\mathcal{C}}(mc_{n-1})$
- (3) $\text{grd}_{\mathcal{C}}(mc_{n-1}) \leq m$

The known results about orderly coin systems with only a few coin values are as follows. Any coin system with two coin values $\mathcal{C} = (1, c)$ where $c > 1$ is orderly. The orderly 3-value coin systems are then characterized by the One Point Theorem (Theorem 2.7). An equivalent characterization is as follows.

Theorem 2.8. [1, 13] *The coin system $\mathcal{C} = (1, c_2, c_3)$ is orderly if and only if $c_3 - c_2 \in \mathcal{A}$, where \mathcal{A} is defined as follows*

$$\mathcal{A} = \{c_2 - 1, c_2\} \cup \{2c_2 - 2, 2c_2 - 1, 2c_2\} \cup \cdots \cup \{mc_2 - m, \dots, mc_2\} \cup \cdots$$

Theorem 2.9. [1, 8] *The coin system $\mathcal{C} = (1, c_2, c_3, c_4)$ is orderly if and only if it is totally orderly.*

Theorem 2.10. [1, 8] *The coin system $\mathcal{C} = (1, c_2, c_3, c_4, c_5)$ is orderly if and only if one of the following holds:*

- (1) $(1, c_2, c_3, c_4, c_5) = (1, 2, a, a+1, 2a)$ for some $a \geq 4$, in which case $(1, c_2, c_3, c_4)$ is not orderly,
- (2) \mathcal{C} is totally orderly.

Notice that Theorems 2.7 and 2.8 imply that the United States' coin system $(1, 5, 10, 25)$ is totally orderly. For the EU coin system $(1, 2, 5, 10, 20, 50, 100, 200)$, Theorem 2.8 together with repeated use of Theorem 2.7 show that it is also totally orderly.

Next, we state constraints on the differences between consecutive coin values for orderly coin systems. The first is simple.

Theorem 2.11. [1] *If $\mathcal{C} = (1, c_2, c_3, \dots, c_n)$ is orderly, then for all $i = 2, 3, \dots, n$,*

$$c_i - c_{i-1} \geq c_2 - 1.$$

The next result is more involved. Roughly speaking, if two consecutive differences between adjacent coin values are big, then *all* subsequent differences between adjacent coins are also big.

Theorem 2.12. [1] *Suppose $\mathcal{C} = (1, c_2, c_3, \dots, c_n)$ is orderly. If $m \geq 3$ and*

$$c_{m-1} > 2c_{m-2} \quad \text{and} \quad c_m > 2c_{m-1},$$

then for every $t \geq m$, we have $c_{t+1} - c_t \geq c_m - c_{m-1}$.

Now we give properties of non-orderly coin systems. Consider again the coin system $A = (1, 3, 4)$. The smallest counterexample for this coin system was $w = 6$. The greedy solution is $6 = 4 + 1 + 1$, while an optimal solution is $6 = 3 + 3$. Observe that the set of coin denominations used in creating the greedy solution $\{1, 4\}$ was disjoint from the coin denominations used to create the optimal solution $\{3\}$. This is true in general, as the result below states.

Proposition 2.13. [6] *Suppose a coin system $\mathcal{C} = (1, c_2, c_3, \dots, c_n)$ is not orderly. Let w be the smallest counterexample, and suppose that the greedy solution for w is given by $w = x_1 \cdot 1 + x_2 \cdot c_2 + x_3 \cdot c_3 + \cdots + x_n \cdot c_n$. Moreover, suppose the following is any optimal solution for w : $w = y_1 \cdot 1 + y_2 \cdot c_2 + y_3 \cdot c_3 + \cdots + y_n \cdot c_n$. Then $x_i y_i = 0$ for all i .*

The final result considers the situation where a non-orderly coin system has another prefix coin system that is also non-orderly. This result requires us to define a new term.

Definition 2.14. *A coin system $\mathcal{C} = (1, c_2, c_3, \dots, c_n)$ is **tight** if it has no counterexample smaller than c_n .*

Theorem 2.15. [4] *Let $\mathcal{C}_1 = (1, c_2, c_3)$, $\mathcal{C}_2 = (1, c_2, c_3, \dots, c_m)$ and $\mathcal{C}_3 = (1, c_2, c_3, \dots, c_m, c_{m+1})$ be three tight coin systems such that \mathcal{C}_1 is orderly but \mathcal{C}_2 is not. If \mathcal{C}_3 is not orderly, then there is a counterexample $x = c_i + c_j > c_{m+1}$ of \mathcal{C}_3 with $1 < c_i \leq c_j \leq c_m$.*

There are several perspectives and extensions of the change-making problem which we leave untouched, since they do not pertain to our goals. Interested readers might like to seek them out [5, 7, 9, 19]. We note that [11] contains apparently elegant necessary and sufficient conditions for a coin system to be orderly, but the result was later shown false in [16].

3. ORDERLY COIN SYSTEMS WITH SIX COIN VALUES

We are ready to prove Theorem 1.2, which gives a characterization of orderly coin systems with 6 coin values. In light of Theorem 2.5, each such coin system has one of the following four patterns:

- | | |
|-------------------|-------------------|
| (1) (+ + + + - +) | (3) (+ + + + + +) |
| (2) (+ + + - - +) | (4) (+ + + - + +) |

Part (1) of the theorem considers coin systems with pattern (+ + + + - +) and is proved in Section 3.1. Part (2) of the theorem considers the pattern (+ + + - - +) and is proved in Section 3.2. Part (3), which considers the remaining two patterns, directly follows from the One Point Theorem (Theorem 2.7).

3.1. Coin systems with pattern: (+ + + + - +).

In this section, we prove part (1) of Theorem 1.2. We restate the result here for clarity.

Theorem 3.1. *A coin system has pattern (+ + + + - +) if and only if it is one of the following:*

- a. $(1, 2, 3, a, a + 1, 2a)$, where $a \geq 5$.
- b. $(1, a, 2a, b, b + a, 2b)$,
where $a \geq 2$, $b \geq 3a - 1$ and $b \neq 3a$ and $\text{gcd}(2ma) \leq m$ where $m = \lceil \frac{b}{2a} \rceil$.
- c. $(1, a, 2a - 1, b, b + a - 1, 2b - 1)$,
where $a \geq 2$, $b \geq 3a - 1$ and $\text{gcd}(m'(2a - 1)) \leq m'$ where $m' = \lceil \frac{b}{2a-1} \rceil$.

Proof. Suppose a coin system has the form (+ + + + - +). Since $(1, c_2, c_3, c_4)$ is orderly and $(1, c_2, c_3, c_4, c_5)$ is not orderly, we know by the One Point Theorem that mc_4 is a counterexample for $(1, c_2, c_3, c_4, c_5)$ where $m = \lceil \frac{c_5}{c_4} \rceil$. Moreover, it must be the case that $c_6 \leq mc_4$, for otherwise \mathcal{C} would not be orderly.

Now observe that

$$c_6 \leq mc_4 = \lceil \frac{c_5}{c_4} \rceil c_4 < (\frac{c_5}{c_4} + 1)c_4 = c_5 + c_4 < c_5 + c_5.$$

Therefore

$$\begin{aligned} c_6 &< c_5 + c_5, \\ c_6 &< c_5 + c_4. \end{aligned}$$

Now since $(1, c_2, c_3, c_4, c_5, c_6)$ is orderly, there must exist coins c_i and c_j where $1 \leq i < j \leq 4$ such that

$$(1) \quad c_6 + c_j = c_5 + c_5,$$

$$(2) \quad c_6 + c_i = c_5 + c_4.$$

Subtracting these two equations, we have

$$c_5 - c_4 = c_j - c_i < c_j \leq c_4$$

Therefore $c_5 \leq 2c_4$. We conclude that $1 < \frac{c_5}{c_4} \leq 2$, which implies that $m = \lceil \frac{c_5}{c_4} \rceil = 2$.

We now have three main cases to consider for the possible values of j (and within each case, we will consider the values $i < j$): Case (1) $j = 4$, Case (2) $j = 3$, and Case (3) $j = 2$.

Case 1: $j = 4$. Plugging in and combining Equations 1 and 2 above, we have $c_5 + c_i = 2c_4$. This shows that $2c_4$ is not a counterexample of $(1, c_2, c_3, c_4, c_5)$, which is a contradiction. Therefore, $j = 4$ is impossible.

Case 2a: $j = 3$ and $i = 1$. Equations 1 and 2 above become:

$$c_6 + c_3 = c_5 + c_5,$$

$$c_6 + 1 = c_5 + c_4.$$

Observe that $c_5 + c_4 = c_6 + 1 > 2c_4 \geq c_6$ which implies that $2c_4 = c_6$. Setting $c_4 = a$, it follows that $c_6 = 2a$, $c_5 = a + 1$, and $c_3 = 2$. However, since $1 < c_2 < c_3$, we are left without any possible value for c_2 , which is a contradiction. Therefore it is impossible to have $j = 3$ and $i = 1$.

Case 2b: $j = 3$ and $i = 2$. Equations 1 and 2 become:

$$c_6 + c_3 = c_5 + c_5,$$

$$c_6 + c_2 = c_5 + c_4.$$

Subtracting these two equations. we have

$$c_5 - c_4 = c_3 - c_2 = d$$

where we define d is the gap between the coins. Simplifying the equations, we conclude that

$$c_3 = c_2 + d$$

$$c_5 = c_4 + d$$

$$c_6 = 2c_4 - c_2 + d$$

So then, the coin system is as follows,

$$(1, c_2, c_3, c_4, c_5, c_6) = (1, c_2, c_2 + d, c_4, c_4 + d, 2c_4 - c_2 + d)$$

By Theorem 2.11, we know $d \geq c_2 - 1$. On the other hand, because $c_6 \leq 2c_4$, it follows that $d \leq c_2$. Therefore, there are only two possibilities: $d = c_2$ or $d = c_2 - 1$. So the coin system has one of two possible forms:

$$\mathcal{C} = (1, c_2, 2c_2, c_4, c_4 + c_2, 2c_4) \quad \text{or} \quad \mathcal{C}' = (1, c_2, 2c_2 - 1, c_4, c_4 + c_2 - 1, 2c_4 - 1).$$

There are further constraints on c_4 . First, Theorem 2.11 tells us that the skip between the third and fourth coins is at least $c_2 - 1$. Therefore the coin system \mathcal{C} satisfies $c_4 \geq 3c_2 - 1$, and the coin system \mathcal{C}' satisfies $c_4 \geq 3c_2 - 2$.

Because $2c_4$ is a counterexample for the 5-value coin system, the difference $2c_4 - c_5$ is not a coin value in the coin system. Therefore for \mathcal{C} we know $c_4 \neq 3c_2$, and \mathcal{C}' satisfies $c_4 \neq 3c_2 - 2$.

Finally, since the 4-value coin system is orderly, we know by the One Point Theorem that the coin system \mathcal{C} satisfies $grd_{\mathcal{C}}(2mc_2) \leq m$ where $m = \lceil \frac{c_4}{2c_2} \rceil$, and the coin system \mathcal{C}' satisfies $grd_{\mathcal{C}'}(m'(2c_2 - 1)) \leq m'$ where $m' = \lceil \frac{c_4}{2c_2 - 1} \rceil$.

Setting $c_2 = a$ and $c_4 = b$, the coin systems \mathcal{C} and \mathcal{C}' coincide with parts (b) and (c) in the statement of the Theorem, respectively.

Case 3: $j = 2$. In this case, we must have $i = 1$. Equations 1 and 2 become

$$\begin{aligned} c_6 + c_2 &= c_5 + c_5, \\ c_6 + 1 &= c_5 + c_4. \end{aligned}$$

Now observe that $c_5 + c_4 = c_6 + 1 > 2c_4 \geq c_6$. This forces $2c_4 = c_6$. Setting $c_4 = a$, and simplifying all these equations, we conclude that

$$c_2 = 2 \qquad c_5 = a + 1 \qquad c_6 = 2a$$

So then, gathering everything together, we conclude that

$$(1, c_2, c_3, c_4, c_5, c_6) = (1, 2, b, a, a + 1, 2a)$$

for some integers a and b with $b < a$.

We now use Theorem 2.12 to show that the values of b and a cannot both be large. Suppose that $b > 4$ and $a > 2b$. Then Theorem 2.12 implies that

$$(a + 1) - a \geq a - b$$

which simplifies to $a \leq b + 1$, but this contradicts $a > 2b$. Therefore we cannot have both $b > 4$ and $a > 2b$, so it must be the case that $b \leq 4$ or $a \leq 2b$.

Suppose that $b > 4$ and $b < a \leq 2b$. We can rule out this possibility by observing that the resulting coin system $(1, 2, b, a, a + 1, 2a)$ will have $c = a + b$ as a counterexample, showing the coin system is not orderly.

Finally, suppose that $b \leq 4$, which implies that $b = 3$ or $b = 4$. We can immediately rule out the possibility that $b = 4$, since in that case our coin system is $(1, 2, 4, a, a + 1, 2a)$ and the value $c = a + 4$ is a counterexample, making our coin system non-orderly. If $b = 3$, we obtain the coin system $(1, 2, 3, a, a + 1, 2a)$. If $a = 4$ the system is totally orderly, so we must have $a \geq 5$.

Conversely, suppose that we have a coin system that is in one of the three forms stated in the theorem. We check that the coin systems have the pattern $(++++-+)$.

We first show the coin system $(1, 2, 3, a, a + 1, 2a)$ where $a \geq 5$ has pattern $(++++-+)$. The coin system $(1, 2, 3)$ is orderly by Theorem 2.8. The fact that $(1, 2, 3, a)$ is orderly for $a \geq 5$ follows from Theorem 2.1 and the fact that $(1, 2, 3)$ is orderly. The 5-value prefix coin system $(1, 2, 3, a, a + 1)$ for $a \geq 5$ is not orderly because $2a$ is a counterexample (this is where we need the assumption that $a \geq 5$).

All that remains is to show that the 6-value coin system is orderly. Theorem 2.3 enables us to find the minimum counterexample (if it exists) as follows. First, set $\mathcal{C} = (1, c_2, c_3, c_4, c_5, c_6) = (1, 2, 3, a, a + 1, 2a)$. We find the greedy representation of $c_k - 1$ for $1 < k \leq 6$ and express it as a vector with respect to \mathcal{C} . By Theorem 2.3, the minimum counterexample (if it exists) is found as follows. For each vector corresponding to $c_k - 1$ and for each p where $1 \leq p < k - 1$, set the first p entries of the vector to 0, and add 1 to the $p + 1^{\text{th}}$ entry. The resulting values are the candidates for minimal counterexample.

Doing this process to the vector associated to $c_2 - 1$ is not possible we would have to have $1 \leq p < 1$ (indeed, this is always the case). Doing this process for $c_3 - 1 = (0, 1, 0, 0, 0, 0)_C$ with $p = 1$ yields the vector $(0, 2, 0, 0, 0, 0)_C$, which corresponds to the value 4. We know 4 is not a counterexample, since the greedy solution for 4 uses only 2 coins.

Next we consider $c_4 - 1 = a - 1$. The greedy solution for this value can be expressed in vector form as $(x_1, x_2, x_3, 0, 0, 0)_C$ where $x_1 + 2x_2 + 3x_3 = a - 1$ and $x_1, x_2 \in \{0, 1\}$ and $x_1x_2 = 0$. With $p = 1$, we have $(0, x_2 + 1, x_3, 0, 0, 0)_C$, which corresponds to the value:

$$2(x_2 + 1) + 3x_3 = \begin{cases} a & \text{if } x_1 = 1 \\ a + 1 & \text{if } x_1 = 0 \end{cases}$$

For $p = 2$, we have $(0, 0, x_3 + 1, 0, 0, 0)_C$, which corresponds to the value:

$$3(x_3 + 1) = \begin{cases} a & \text{if } x_1 = 0, x_2 = 1 \\ a + 1 & \text{if } x_1 = 1, x_2 = 0 \\ a + 2 & \text{if } x_1 = 0, x_2 = 0 \end{cases}$$

None of these values a , $a + 1$, or $a + 2$ are counterexamples, because each has a greedy solution with 1 or 2 coins.

Next up, the value $c_5 - 1 = a$ has greedy representation $(0, 0, 0, 1, 0, 0)_C$ and yields three candidates for counterexample (corresponding to $p = 1, 2, 3$):

$$\begin{aligned} (0, 1, 0, 1, 0, 0)_C &= a + 2 \\ (0, 0, 1, 1, 0, 0)_C &= a + 3 \\ (0, 0, 0, 2, 0, 0)_C &= 2a \end{aligned}$$

Again, the values $a + 2$, $a + 3$, and $2a$ are not counterexamples, since each has a greedy solution with 1 or 2 coins.

Finally, we consider the vector associated to $c_6 - 1 = 2a - 1$. The greedy solution has vector representation $(y_1, y_2, y_3, 0, 1, 0)_C$ where $y_1 + 2y_2 + 3y_3 = a - 2$ and $y_1, y_2 \in \{0, 1\}$ and $y_1y_2 = 0$. With $p = 1$, we have the vector $(0, y_2 + 1, y_3, 0, 1, 0)_C$, which corresponds to the value:

$$2(y_2 + 1) + 3y_3 + a + 1 = \begin{cases} 2a & \text{if } y_1 = 1 \\ 2a + 1 & \text{if } y_1 = 0 \end{cases}$$

For $p = 2$, we have the vector $(0, 0, y_3 + 1, 0, 1, 0)_C$, which corresponds to the value:

$$3(y_3 + 1) = \begin{cases} 2a & \text{if } y_1 = 0, y_2 = 1 \\ 2a + 1 & \text{if } y_1 = 1, y_2 = 0 \\ 2a + 2 & \text{if } y_1 = 0, y_2 = 0 \end{cases}$$

For $p = 3$ and $p = 4$, we have:

$$\begin{aligned} (0, 0, 0, 1, 1, 0)_C &= 2a + 1 \\ (0, 0, 0, 0, 2, 0)_C &= 2a + 2 \end{aligned}$$

Altogether from these four cases, we generate three possible candidates for counterexample: $2a$, $2a + 1$, and $2a + 2$. None of these are counterexamples, since each has a greedy solution with 1 or 2 coins.

Because the process found no minimum counterexample, it must be the case that $(1, 2, 3, a, a + 1, 2a)$ is orderly for $a \geq 5$, as claimed.

Now let \mathcal{B} and \mathcal{B}' denote coin systems in the form of (b) and (c) in the statement of the theorem, respectively. Theorems 2.7 and 2.8 imply that the 4-value prefix coin systems of \mathcal{B} and \mathcal{B}' are both totally orderly. Furthermore, the 5-value prefix coin systems for \mathcal{B} and for \mathcal{B}' have counterexample $2b$, so they are not orderly. Hence, all that remains is to show that the 6-value coin systems \mathcal{B} and \mathcal{B}' are orderly.

Using Theorem 2.3, it follows that $2b$ is the minimum counterexample of the 5-value prefix coin systems of both \mathcal{B} and \mathcal{B}' . Thus the 5-value prefix coin systems are both tight. Now suppose that \mathcal{B} or \mathcal{B}' is not orderly. Then by Theorem 2.15, it follows that the coin system has a counterexample of the form $c_i + c_j$ for some coin values c_i and c_j in \mathcal{B} or in \mathcal{B}' , where $1 \leq i \leq j \leq 5$. However, a straightforward check of these values reveals that $c_i + c_j$ is never a counterexample. Hence it must be the case that the coin systems \mathcal{B} and \mathcal{B}' are orderly, as claimed. \square

3.2. Coin systems with pattern: $(+++--)$.

In this section, we prove part (2) of Theorem 1.2. We first prove a lemma.

Lemma 3.2. *If $\mathcal{C} = (1, c_2, c_3, c_4, c_5, c_6)$ has pattern $(+++--)$, then the coin system has one of the following forms:*

- (1) $(1, a, 2a - 1, b, a + b - 1, 2b - 1)$ where $a \geq 3$ and $b > 2a - 1$.
- (2) $(1, a, 2a, b, a + b, 2b)$ where $a \geq 2$ and $b > 2a$.

Proof. Because the 6-value coin system is orderly, both the 4-coin and 5-value coin systems are tight. By Theorem 2.15, there must exist a counterexample for the 5-value coin system $(1, c_2, c_3, c_4, c_5)$ of the form $c_s + c_t$ for some $1 \leq s, t \leq 4$. And because the 6-value coin system is orderly, it follows that $c_6 \leq c_s + c_t \leq c_4 + c_4$.

Thus there exists $0 \leq i < j < k \leq 4$ such that

$$\begin{aligned} c_4 + c_4 &= c_6 + c_i \\ c_4 + c_5 &= c_6 + c_j \\ c_5 + c_5 &= c_6 + c_k \end{aligned}$$

(We set $c_0 = 0$.) Subtracting adjacent equations, we find

$$c_5 - c_4 = c_k - c_j = c_j - c_i.$$

The possible values for j are: 1, 2, and 3. We will rule out the possibility that $j = 1$ or $j = 3$ first, and then we will study the case $j = 2$.

Case 1: $j = 1$. If $j = 1$, it follows that $i = 0$ and hence $c_k = 2c_1 - c_0 = 2$. Thus it must be the case that $k = 2$. Substituting the values $(i, j, k) = (0, 1, 2)$ into the above system of equations, we find the following relationships: $c_5 = c_4 + 1$ and $c_6 = 2c_4$. So the coin system is $(1, 2, c_3, c_4, c_4 + 1, 2c_4)$. Since the coin system is orderly, the value $c_3 + c_4$ should have a greedy solution with just 2 coins. Hence $(c_3 + c_4) - (c_4 + 1)$ must be a value in the coin system. This forces either $c_3 = 2$ or $c_3 = 3$. The former option leaves no possible value for c_2 . For the latter option, the first three coin values are $(1, 2, 3)$. Thus, no matter the value for c_4 , the 4-value coin system $(1, 2, 3, c_4)$ is orderly, which is a contradiction. Hence $j = 1$ is not possible.

Case 2: $j = 3$. In this case, we must have $k = 4$. Considering the possibilities $i = 0, 1, 2$, we know $c_4 - c_3$ equals one of the elements in the set $\{c_3, c_3 - 1, c_3 - c_2\}$.

This forces the value of c_4 to be one of the following: $2c_3$ or $2c_3 - 1$ or $2c_3 - c_2$. However, using the One Point Theorem, if the 3-value coin system $(1, c_2, c_3)$ is orderly, the resulting coin system $(1, c_2, c_3, c_4)$ is orderly as well, if c_4 is any of the options above. Therefore it is impossible to have $j = 3$.

Case 3: $j = 2$. In this case, the possibilities for i and k and the subsequent equation $c_2 - c_i = c_k - c_2$ are:

- (1) $i = 0, k = 3$, and $c_2 = c_3 - c_2$ (3) $i = 0, k = 4$, and $c_2 = c_4 - c_2$
(2) $i = 1, k = 3$, and $c_2 - 1 = c_3 - c_2$ (4) $i = 1, k = 4$, and $c_2 - 1 = c_4 - c_2$

Option (1) above implies that $c_3 = 2c_2$. Plugging the values for i, j, k into the system of equations, we also have $c_5 = c_4 + c_2$ and $c_6 = 2c_4$. Thus the coin system is in the form of the second possibility in the statement of the lemma.

Option (2) above implies that $c_3 = 2c_2 - 1$. Moreover, observe that $c_3 = 2c_2 - 1$ is possible only if $c_2 \geq 3$. For, if $c_2 = 2$, then there is no value of c_4 for which the 4-value coin system is not orderly. Plugging the values for i, j, k into the system of equations, we also have $c_5 = c_4 + c_2 - 1$ and $c_6 = 2c_4 - 1$. Thus the coin system is in the form of the first possibility in the statement of the lemma.

Options (3) and (4) cannot occur. For, (3) and (4) imply that $c_4 = 2c_2$ or $c_4 = 2c_2 - 1$, respectively. On the other hand, we know from Theorem 2.11 that $c_4 \geq 3c_2 - 2$. Applying this inequality, we conclude that $c_2 \leq 2$ and $c_2 \leq 1$, respectively. The former option implies $c_2 = 2$ and $c_4 = 4$, which forces $c_3 = 3$, and the 4-value coin system is necessarily orderly, which is a contradiction. The latter option ($c_2 \leq 1$) is impossible, since $c_2 > 1$. Thus neither (3) nor (4) above are possible. This proves the lemma. \square

Now we are ready to prove part (2) of Theorem 1.2. We restate the result here for clarity.

Theorem 3.3. *A coin system has pattern $(+++--)$ if and only if it is one of the following:*

- (1) $\mathcal{C} = (1, a, 2a - 1, m(2a - 1) - a + 1, m(2a - 1), (2m - 1)(2a - 1))$
where $1 < m < a$.
(2) $\mathcal{C}' = (1, a, 2a, m(2a - 1) - a + 1, m(2a - 1) + 1, (2m - 1)(2a - 1) + 1)$
where $1 < m \leq a$.

Proof. First we prove that any coin system with the pattern $(+++--)$ is in the form of \mathcal{C} or \mathcal{C}' as stated in the theorem. From the previous lemma, we know that the coin system can be in two different forms. First consider the form: $\mathcal{B} = (1, a, 2a - 1, b, a + b - 1, 2b - 1)$ where $a \geq 3$ and $b > 2a - 1$. From the One Point Theorem, we know that the coin system $(1, a, 2a - 1, b)$ has a counterexample of the form $m(2a - 1)$ where $m = \lceil \frac{b}{2a-1} \rceil$.

We claim that this number $m(2a - 1)$ is not a counterexample of the 5-value prefix coin system of \mathcal{B} . Suppose for contradiction that $m(2a - 1)$ is such a counterexample. Because the 6-value coin system is orderly, we know that

$$c_6 = 2b - 1 \leq m(2a - 1) = \lceil \frac{b}{2a-1} \rceil (2a - 1) \leq (\frac{b}{2a-1} + 1)(2a - 1) = b + 2a - 1.$$

This implies that $b \leq 2a$. Since we already know $b > 2a - 1$, we conclude $b = 2a$. By Theorem 2.11, we know that the differences between adjacent coin values is at least $a - 1$, and because the third and fourth coin values differ by 1, we have $a = 2$,

and the coin system is $(1, 2, 3, 4, 5, 7)$. However, this coin system is totally orderly, which is a contradiction. Hence $m(2a - 1)$ is not a counterexample for the 5-value coin system of \mathcal{B} , as claimed.

Because $m(2a - 1)$ is a counterexample for the 4-value prefix coin system, if we subtract the value of the 4th coin (which has value b) from $m(2a - 1)$, the remaining value must use more than $m - 1$ coins in its greedy solution. Whereas, because $m(2a - 1)$ is *not* a counterexample for the 5-value prefix coin system, if we subtract the value of the 5th coin (which has value $a + b - 1$) from $m(2a - 1)$, the remaining value uses at most $m - 1$ coins in its greedy solution.

Letting $z = m(2a - 1) - (a + b - 1)$ and $y = m(2a - 1) - b$, we can express the above statements as an inequality: $grd_{\mathcal{B}}(z) \leq m - 1 < grd_{\mathcal{B}}(y)$. Moreover, observe that the value of z is small:

$$z = m(2a - 1) - (a + b - 1) \leq \left(\frac{b}{2a-1} + 1\right)(2a - 1) - (a + b - 1) = a$$

Also, notice that $y = z + a - 1$.

Collecting these statements together, we are looking for a value z with two relationships: (1) $z \in [0, a]$ and (2) $grd_{\mathcal{B}}(z) \leq m - 1 < grd_{\mathcal{B}}(z + a - 1)$. Let's take a look at the function $grd_{\mathcal{B}}(x)$ at values from $x = 0$ up to $x = 2a - 1$.

$$grd_{\mathcal{B}}(x) = \begin{cases} x & \text{if } 0 \leq x \leq a - 1 \\ x - (a - 1) & \text{if } a \leq x \leq 2a - 2 \\ 1 & \text{if } x = 2a - 1 \end{cases}$$

Observe that for all $1 \leq x \leq a$, the function satisfies $grd_{\mathcal{B}}(x) = grd_{\mathcal{B}}(x + a - 1)$. So z cannot be any integer from 1 to a , because the inequality (2) given above is not satisfied. As $z \in [0, a]$, the only possible value for z is $z = 0$. Replacing $z = 0$ in the original expressions for y and z , we find the values of c_4, c_5 , and c_6 .

$$\begin{aligned} c_4 &= b = m(2a - 1) - a + 1 \\ c_5 &= a + b - 1 = m(2a - 1) \\ c_6 &= 2b - 1 = (2m - 1)(2a - 1) \end{aligned}$$

Keep in mind that by the definition of m , we know $m > 1$, and because $m - 1 < grd_{\mathcal{B}}(z + a - 1) = grd_{\mathcal{B}}(a - 1) = a - 1$, we also have that $m < a$. This proves that the coin system \mathcal{B} has the form of \mathcal{C} in the statement of the theorem.

Now consider the second coin system form in Lemma 3.2:

$$\mathcal{B}' = (1, a, 2a, b, a + b, 2b)$$

where $a \geq 2$ and $b > 2a$. We will show that \mathcal{B}' has the form of \mathcal{C}' given in the statement of the theorem. Initially, the argument for \mathcal{B}' mirrors what we just did for \mathcal{B} , but then it diverges in subtle ways, so we start from the beginning.

From the One Point Theorem, we know that the coin system $(1, a, 2a, b)$ has a counterexample of the form $2am$ where $m = \lceil \frac{b}{2a} \rceil$. Suppose that this value $2am$ is also a counterexample of the 5-value prefix coin system of \mathcal{B}' . Since the 6-value coin system \mathcal{B}' is orderly, we know that

$$c_6 = 2b \leq 2am = (2a) \lceil \frac{b}{2a} \rceil \leq (2a) \left(\frac{b}{2a} + 1\right) = b + 2a.$$

This implies that $b \leq 2a$, a contradiction. We conclude that $2am$ is not a counterexample for the 5-value prefix coin system of \mathcal{B}' .

Therefore, starting with the value $2am$, if we subtract the value of the 4th coin, the remaining value must require more than $m - 1$ coins using the greedy algorithm. Whereas, starting with the value $2am$, if we subtract the value of the 5th coin, the remaining value must require at most $m - 1$ coins using the greedy algorithm. Letting $z = 2am - (a + b)$ and $y = 2am - b$, we can express the above thought as an inequality: $grd_{\mathcal{B}'}(z) \leq m - 1 < grd_{\mathcal{B}'}(y)$. Also observe that $z \in [0, a]$ because:

$$0 \leq z = 2am - (a + b) \leq \left(\frac{b}{2a} + 1\right)(2a) - (a + b) = a.$$

Moreover, observe that $y = z + a$. Collecting these statements together, we are looking an integer z with two relationships: (1) $z \in [0, a]$ and (2) $grd_{\mathcal{B}'}(z) \leq m - 1 < grd_{\mathcal{B}'}(z + a)$. Let us take a look at the function $grd_{\mathcal{B}'}(x)$ at values from $x = 0$ to $x = 2a$, which the same as the previous case except at the last two values ($x = 2a - 1$ and $x = 2a$).

$$grd_{\mathcal{B}'}(x) = \begin{cases} x & \text{if } 0 \leq x \leq a - 1 \\ x - (a - 1) & \text{if } a \leq x \leq 2a - 1 \\ 1 & \text{if } x = 2a \end{cases}$$

Observe that $grd_{\mathcal{B}'}(x) = grd_{\mathcal{B}'}(x + a) - 1$ for all $x \in [0, a - 1]$. This implies that if z is any integer in the interval $[0, a - 1]$, the inequality (2) above will be satisfied. In addition, if z is any value on this interval, the inequality from (2) above simplifies to $z \leq m - 1 < z + 1$. Thus $z = m - 1$.

Replacing $z = m - 1$ in the original expressions for y and z , we find the expression for c_4 , c_5 , and c_6 as follows.

$$\begin{aligned} c_4 &= b = 2am - (z + a) = m(2a - 1) - a + 1 \\ c_5 &= a + b = m(2a - 1) + 1 \\ c_6 &= 2b = 2(m(2a - 1) - a + 1) = (2m - 1)(2a - 1) + 1 \end{aligned}$$

Notice that by the construction of m , we know that $m > 1$, and since $m - 1 = grd_{\mathcal{B}'}(z) \leq a - 1$, it follows that $m \leq a$. So then \mathcal{B}' has the same form as \mathcal{C}' , as desired.

Conversely, we must prove that the coin systems \mathcal{C} and \mathcal{C}' in the theorem's statement have the pattern $(+++- -)$. We prove a stronger result, in fact. In Theorem 4.5, we prove that an infinite family of coin systems denoted \mathcal{E} (for which \mathcal{C} is a special case) has the pattern $(+++ - \dots - +)$. In Theorem 4.7, we prove that an infinite family of coin system \mathcal{F} (for which \mathcal{C}' is a special case) has the pattern $(+++ - \dots - +)$. We thus delay this argument to the next section. \square

4. ORDERLY COIN SYSTEMS WITH n COIN VALUES

We now study orderly coin systems with n coin values. The only $+/-$ coin system pattern which cannot be tackled via first characterizing an orderly coin system for a value of i with $3 < i < n$ is the pattern where every subsystem of size i with $3 < i < n$ fails to be orderly. Such a coin system has the pattern $(+++ - \dots - +)$. We focus on these coin systems.

In [1], the authors found a single coin system with pattern $(+++ - \dots - +)$ and with n types of coins for each $n = 3r$ where $r \geq 2$ and for each $n = 3r + 2$ where $r \geq 1$. Interestingly, the authors wrote that they suspected that there are no coin systems with $3r + 1$ coin values with this pattern where $r \geq 2$, but they were unable to prove their conjecture.

We searched extensively for coin systems with pattern $(+++ \cdots - +)$ using a computer program. All observed examples had a common structure. We call these coin systems **fixed gap coin systems**, which we define below. Furthermore, the examples we found divided into three distinct infinite families \mathcal{D} , \mathcal{E} , and \mathcal{F} . The coin systems in family \mathcal{D} had $3r+2$ types of coin where $r \geq 1$, while the coin systems in \mathcal{E} and \mathcal{F} had $3r$ types of coins where $r \geq 2$. In line with the conjecture [1], we found no coin systems with $3r+1$ coins where $r \geq 2$. We prove that these three infinite families of coin systems \mathcal{D} , \mathcal{E} , and \mathcal{F} have the pattern $(+++ \cdots - +)$. We conjecture that they are the only such coin systems (Conjecture 1.3). The conjecture holds true for $n = 5$ and $n = 6$.

A fixed-gap coin system with n types of coins starts off with two initial coin values $(c_1, c_2) = (1, x)$. Then all the remaining coin values are generated in two stages. First choose two positive integers Δ_1 and Δ_2 where $\Delta_1 \neq \Delta_2$.

- **Stage 1:** Alternately add Δ_1 and Δ_2 to the previous coin value to produce the next coin value. We continue this until we have ℓ coin values for some $3 < \ell < n$.
- **Stage 2:** For coin values $c_{\ell+1}$ through c_n , add $\Delta_1 + \Delta_2$ to the previous coin value to produce the next coin value.

We make this process precise as follows.

Definition 4.1. *Let n, ℓ, x, Δ_1 , and Δ_2 be positive integers with $3 < \ell < n, x \geq 2$, and $\Delta_1 \neq \Delta_2$. The coin system $\mathcal{C} = (1, c_2, c_3, \dots, c_n)$ is a **fixed gap coin system** if c_i are defined as follows.*

$$(3) \quad c_i = \begin{cases} 1 & \text{if } i = 1 \\ x & \text{if } i = 2 \\ c_{i-1} + \Delta_1 & \text{if } i \text{ is odd and } 3 \leq i \leq \ell \\ c_{i-1} + \Delta_2 & \text{if } i \text{ is even and } 3 < i \leq \ell \\ c_{i-1} + \Delta_1 + \Delta_2 & \text{if } \ell < i \leq n \end{cases}$$

For most fixed-gap coin system \mathcal{C} as above, the coin system has several consecutive prefix coin systems that fail to be orderly. We prove this in general in the following lemma, before focusing on the three specific families of fixed gap coin systems \mathcal{D} , \mathcal{E} , and \mathcal{F} .

Lemma 4.2. *Let \mathcal{C} be a fixed-gap coin system. If ℓ is odd, then each prefix coin system of \mathcal{C} of length k with $5 \leq k \leq \frac{1}{2}(3\ell - 5)$ is not orderly. If ℓ is even and $x + \Delta_1 > \Delta_2 + 1$, then each prefix coin system of \mathcal{C} of length k with $5 \leq k \leq \frac{1}{2}(3\ell - 4)$ is not orderly.*

Proof. Throughout this proof, we use the same strategy multiple times to show that a given coin system $(1, x, c_3, \dots, c_k)$ is not orderly. The outline is as follows. We prove:

- (1) $2c_{k-1} - c_k > 0$, and
- (2) $2c_{k-1} - c_k \neq c_i$ for any $i < k$.

This implies that $2c_{k-1}$ is a counterexample because on one hand the greedy solution has more than 2 coins, while the optimal solution has 2 coins.

To begin, we prove that the prefix coin system $(1, x, c_3, \dots, c_k)$ is not orderly for all k such that $5 \leq k \leq \ell$.

Suppose that k is even. We show $2c_{k-1} - c_k$ is positive:

$$2c_{k-1} - c_k = 2c_{k-1} - (c_{k-1} + \Delta_2) = c_{k-1} - \Delta_2 = (c_{k-3} + \Delta_1 + \Delta_2) - \Delta_2 = c_{k-3} + \Delta_1 > 0$$

Now we show that $2c_{k-1} - c_k$ is not a coin value c_i for any i . Observe that if $\Delta_1 < \Delta_2$, then the value $2c_{k-1} - c_k = c_{k-3} + \Delta_1$ falls between consecutive coin values:

$$(4) \quad c_{k-3} < c_{k-3} + \Delta_1 < \underbrace{c_{k-3} + \Delta_2}_{c_{k-2}}$$

And if $\Delta_2 < \Delta_1$, we similarly have:

$$(5) \quad \underbrace{c_{k-3} + \Delta_2}_{c_{k-2}} < c_{k-3} + \Delta_1 < \underbrace{c_{2j-3} + \Delta_1 + \Delta_2}_{c_{k-1}}$$

So no matter the relative sizes of Δ_1 and Δ_2 , the value $2c_{k-1} - c_k$ falls between consecutive coin values, and therefore the value is not itself a coin value. We conclude that $2c_{k-1}$ is a counterexample for the prefix coin system $(1, c_2, c_3, \dots, c_k)$ where k is even and $5 < k \leq \ell$.

Now suppose that k is odd and $5 \leq k \leq \ell$. Similar to before, we show that $2c_{k-1}$ is a counterexample for the prefix coin system $(1, c_2, c_3, \dots, c_k)$. First, we simplify the expression $2c_{k-1} - c_k$ and conclude that it must be positive:

$$2c_{k-1} - c_k = 2c_{k-1} - (c_{k-1} + \Delta_1) = c_{k-1} - \Delta_1 = (c_{k-3} + \Delta_1 + \Delta_2) - \Delta_1 = c_{k-3} + \Delta_2 > 0$$

Similar to the previous case, observe that if $\Delta_1 < \Delta_2$, then the value $2c_{k-1} - c_k = c_{k-3} + \Delta_2$ falls between consecutive coin values:

$$(6) \quad \underbrace{c_{k-3} + \Delta_1}_{c_{k-2}} < c_{k-3} + \Delta_2 < \underbrace{c_{k-3} + \Delta_1 + \Delta_2}_{c_{k-1}}$$

And if $\Delta_2 < \Delta_1$, we have:

$$(7) \quad c_{k-3} < c_{k-3} + \Delta_2 < \underbrace{c_{k-3} + \Delta_1}_{c_{k-2}}$$

So in either case, the value $2c_{k-1} - c_k = c_{k-3} + \Delta_2$ falls between consecutive coin values, and therefore the value is not itself a coin value. We conclude that $2c_{k-1}$ is a counterexample for the prefix coin system $(1, c_2, c_3, \dots, c_k)$ where k is odd and $5 \leq k \leq \ell$.

It remains to show that all prefix subcurrency systems $(1, c_2, c_3, \dots, c_k)$ where $\ell < k \leq t$ also fail to be orderly, where $t = \frac{1}{2}(3\ell - 5)$ if ℓ is odd and $t = \frac{1}{2}(3\ell - 4)$ if ℓ is even.

Suppose $\ell = 2r + 1$ for some $r \geq 2$. We show that $(1, c_2, c_3, \dots, c_k)$ where $\ell < k \leq 3r - 1$ fail to be orderly by showing $2c_{2r}$ is a counterexample.

$$\begin{aligned} 2c_{2r} - c_k &= 2c_{2r} - (c_{2r+1} + (k - 2r - 1)(\Delta_1 + \Delta_2)) \\ &= 2c_{2r} - (c_{2r} + \Delta_1 + (k - 2r - 1)(\Delta_1 + \Delta_2)) \\ &= c_{2r} - \Delta_1 - (k - 2r - 1)(\Delta_1 + \Delta_2) \\ &= c_{2r-2(k-2r)} + (k - 2r)(\Delta_1 + \Delta_2) - \Delta_1 - (k - 2r - 1)(\Delta_1 + \Delta_2) \\ &= c_{6r-2k} + \Delta_2 \end{aligned}$$

We previously proved that a value of form $c_{2i} + \Delta_2$ where $1 \leq i \leq r - 1$ falls between adjacent coin values (Inequalities 6 and 7). Therefore, $2c_{2r}$ is a counterexample for the prefix coin system $(1, c_2, c_3, \dots, c_k)$ where $\ell < k \leq 3r - 1$ and ℓ is odd.

Finally, suppose ℓ is even. That is, $\ell = 2r + 2$ for some $r \geq 1$. And suppose $x + \Delta_1 > \Delta_2 + 1$. We show that $(1, c_2, c_3, \dots, c_k)$ where $\ell < k \leq 3r + 1$ fails to be orderly by showing $2c_{2r+1}$ is a counterexample. First we do the special case that $k = 3r + 1$.

$$\begin{aligned}
2c_{2r+1} - c_{3r+1} &= 2c_{2r+1} - (c_{2r+2} + (r-1)(\Delta_1 + \Delta_2)) \\
&= 2c_{2r+1} - (c_{2r+1} + \Delta_2 + (r-1)(\Delta_1 + \Delta_2)) \\
&= c_{2r+1} - \Delta_2 - (r-1)(\Delta_1 + \Delta_2) \\
&= c_{2r+1-2(r-1)} + (r-1)(\Delta_1 + \Delta_2) - \Delta_2 - (r-1)(\Delta_1 + \Delta_2) \\
&= c_3 - \Delta_2 \\
&= x + \Delta_1 - \Delta_2
\end{aligned}$$

By our assumption, $x + \Delta_1 > \Delta_2 + 1$. Thus $1 < x + \Delta_1 - \Delta_2 < c_3$ and moreover $x + \Delta_1 - \Delta_2$ cannot equal $c_2 = x$. Therefore, this value $x + \Delta_1 - \Delta_2$ is not a coin value and $2c_{2r+1}$ is a counterexample for the prefix coin system $(1, x, c_3, \dots, c_{3r+1})$.

Now suppose $\ell < k < 3r + 1$. We prove that this prefix coin system is not orderly by showing $2c_{2r+1}$ is a counterexample.

$$\begin{aligned}
2c_{2r+1} - c_k &= 2c_{2r+1} - (c_{2r+2} + (k-2r-2)(\Delta_1 + \Delta_2)) \\
&= 2c_{2r+1} - (c_{2r+1} + \Delta_2 + (k-2r-2)(\Delta_1 + \Delta_2)) \\
&= c_{2r+1} - \Delta_2 - (k-2r-2)(\Delta_1 + \Delta_2) \\
&= c_{2r+1-2(k-2r-2)} + (k-2r-2)(\Delta_1 + \Delta_2) - \Delta_2 - (k-2r-2)(\Delta_1 + \Delta_2) \\
&= c_{6r-2k+5} - \Delta_2 \\
&= c_{6r-2k+3} + \Delta_1
\end{aligned}$$

We previously proved that a value of the form $c_{2i+1} + \Delta_1$ where $1 \leq i < r - 1$ falls between adjacent coin values (Inequalities 4 and 5.). Therefore, $2c_{2r+1}$ is a counterexample for the coin system $(1, c_2, c_3, \dots, c_k)$ where $\ell < k < 3r + 2$. \square

Now we define the three infinite families of fixed-gap coin systems (denoted by \mathcal{D} , \mathcal{E} , and \mathcal{F} , respectively) and prove that they have pattern $(+++ - \dots - +)$.

Theorem 4.3. *Let $r \geq 1$ and $a \geq 2$. Define $\mathcal{D} = (1, d_2, \dots, d_{3r+2})$ as follows:*

$$d_k = \begin{cases} 1 & \text{if } k = 1 \\ 2 & \text{if } k = 2 \\ d_{k-1} + a & \text{if } k \text{ is odd and } 3 \leq k < 2r + 2 \\ d_{k-1} + 1 & \text{if } k \text{ is even and } 3 < k \leq 2r + 2 \\ d_{k-1} + a + 1 & \text{if } 2r + 2 < k \leq 3r + 2 \end{cases}$$

The coin system \mathcal{D} has pattern $(+++ - \dots - +)$.

Example 4.4. *For $r = 3$ and $a = 3$, the coin system is*

$$\mathcal{D} = (1, 2, 5, 6, 9, 10, 13, 14, 18, 22, 26).$$

Proof. The 3-coin prefix coin system $(d_1, d_2, d_3) = (1, 2, a + 2)$ is totally orderly for all values of $a \geq 2$ by Theorem 2.8 where $d_3 - d_2 = a$. The 4-coin prefix coin system $(1, 2, a + 2, a + 3)$ is not orderly because $2a + 4$ is a counterexample. From Lemma 4.2, we know all prefix coin systems of length i where $5 \leq i \leq 3r + 1$ are not orderly.

It remains is to show the full coin system \mathcal{D} is orderly. Suppose that the coin system is not orderly. We use Theorem 2.3 to find the smallest counterexample. First, we find the greedy solution for $d_k - 1$ for all $k \geq 2$ and express the value as a vector with respect to the coin system \mathcal{D} . In the vectors below, an asterisk (*) denotes that the value is in the $k - 1$ position. For k even and $2 \leq k \leq 2r + 2$,

$$d_k - 1 = d_{k-1} = (0, 0, \dots, 1^*, \dots, 0)_{\mathcal{D}}.$$

For k odd and $2 < k < 2r + 2$,

$$d_k - 1 = d_{k-1} + a - 1 = (x_1, x_2, 0, \dots, 1^*, \dots, 0)_{\mathcal{D}}$$

where x_1 and x_2 are integers such that $a - 1 = 2x_2 + x_1$, $x_2 \geq 0$ and $x_1 \in \{0, 1\}$. For $2r + 1 < k \leq 3r + 2$,

$$d_k - 1 = d_{k-1} + a = (y_1, y_2, 0, \dots, 1^*, 0, \dots, 0)_{\mathcal{D}}$$

where y_1 and y_2 are integers such that $a = 2y_2 + y_1$, $y_2 \geq 1$, and $y_1 \in \{0, 1\}$.

Now for each $1 \leq p < k - 1$, we set the first p entries of the vectors above to 0 and add 1 to the $p + 1$ entry. The set of the resulting vectors gives all the candidates for the smallest counterexample. When $p = 1$, the resulting vector in each case represents one of the following values: d_i , $d_i + 1$, or $d_i + 2$. None of these values are counterexamples, since their greedy solutions have either 1 or 2 coins. When $p \geq 2$, the resulting vector in each case represents a value of the form $d_i + d_j$ where $2 \leq i, j < 3r + 2$. We compute the greedy solution for $d_i + d_j$ for all i, j .

First we consider the sum $d_i + d_j$ where both of the coin values are belong to Stage 1 of the construction. That is, $3 \leq i, j \leq 2r + 2$.

$$d_i + d_j = \begin{cases} d_2 + d_{i+j-2} & \text{if } i \text{ and } j \text{ are even and } i + j \leq 2r + 4 \\ d_2 + d_{(i+j+2r)/2} & \text{if } i \text{ and } j \text{ are even and } i + j > 2r + 4 \\ d_{i+j} & \text{if } i \text{ and } j \text{ are odd and } i + j \leq 2r + 2 \\ d_{r+1+(i+j)/2} & \text{if } i \text{ and } j \text{ are odd and } i + j > 2r + 2 \\ d_1 + d_{i+j-1} & \text{if } i \text{ is odd and } j \text{ is even and } i + j \leq 2r + 2 \\ d_1 + d_{(i+j+2r+1)/2} & \text{if } i \text{ is odd and } j \text{ is even and } i + j > 2r + 2 \end{cases}$$

Next we compute the greedy solution for $d_i + d_j$ where at least one coin value (say, d_j) belongs to Stage 2 of the construction. That is, $2r + 2 < j < 3r + 2$. Notice that we need not consider the case $j = 3r + 2$ because in that case, the greedy solution of $d_i + d_{3r+2}$ is simply itself.

$$d_i + d_j = \begin{cases} d_{3r+2} + d_{i+2j-6r-4} & \text{if } 2 < i < 2r + 2 \text{ is odd and } \frac{1}{2}(i-1) + j \geq 3r + 2 \\ d_1 + d_{j+(i-1)/2} & \text{if } 2 < i < 2r + 2 \text{ is odd and } \frac{1}{2}(i-1) + j < 3r + 2 \\ d_{3r+2} + d_{i+2j-8r-8} & \text{if } 2 \leq i \leq 2r + 2 \text{ is even and } \frac{1}{2}(i-2) + j \geq 3r + 2 \\ d_2 + d_{j+(i-2)/2} & \text{if } 2 \leq i \leq 2r + 2 \text{ is even and } \frac{1}{2}(i-2) + j < 3r + 2 \\ d_{3r+2} + d_{2(i+j-4r-3)} & \text{if } 2r + 2 < i < 3r + 2 \text{ and } i + j \leq 5r + 4 \\ d_{3r+2} + d_{i+j-3r-2} & \text{if } 2r + 2 < i < 3r + 2 \text{ and } i + j > 5r + 4 \end{cases}$$

None of these values $d_i + d_j$ are counterexamples because each greedy solution has only 1 or 2 coins. Therefore the coin system \mathcal{D} does not have a minimum counterexample, and the coin system is orderly. \square

Theorem 4.5. *Let $r \geq 2$ and $1 < m < a$. Define $\mathcal{E} = (1, e_2, \dots, e_{3r})$ as follows:*

$$e_k = \begin{cases} 1 & \text{if } k = 1 \\ a & \text{if } k = 2 \\ e_{k-1} + a - 1 & \text{if } k \text{ is odd and } 3 \leq k \leq 2r + 1 \\ e_{k-1} + (m-1)(2a-1) - (a-1) & \text{if } k \text{ is even and } 3 < k < 2r + 1 \\ e_{k-1} + (m-1)(2a-1) & \text{if } 2r + 1 < k \leq 3r \end{cases}$$

The coin system \mathcal{E} has pattern $(+++ \dots - +)$.

Example 4.6. *For $r = 4$, $m = 3$, and $a = 4$, the coin system is*

$$\mathcal{E} = (1, 4, 7, 18, 21, 32, 35, 46, 49, 63, 77, 91).$$

Proof. The 3-value prefix coin system $(e_1, e_2, e_3) = (1, a, 2a-1)$ is totally orderly by Theorem 2.8 where $e_3 - e_2 = a - 1$. The 4-coin prefix coin system $(1, a, 2a-1, m(2a-1) - (a-1))$ where $1 < m < a$ is not orderly because $m(2a-1)$ is a counterexample. From Lemma 4.2, we know all prefix coin systems of length i where $5 \leq i \leq 3r-1$ are not orderly.

Now we show that the full coin system \mathcal{E} is orderly using Theorem 2.3. First, we find the greedy solution for $e_k - 1$ for all $k \geq 2$ and express the value as a vector with respect to the coin system \mathcal{E} . In the vectors below, the asterisk (*) denotes that the value is in the $k-1^{\text{st}}$ entry.

$$(8) \quad e_k - 1 = \begin{cases} (a-1, 0, \dots, 0)_{\mathcal{E}} & \text{if } k = 2 \\ (a-2, 0, \dots, 1^*, \dots, 0)_{\mathcal{E}} & \text{if } k \text{ is odd, } 3 \leq k \leq 2r + 1 \\ (a-1, 0, m-1, 0, \dots, 0)_{\mathcal{E}} & \text{if } k = 4 \\ (a-1, 0, m-2, 0, \dots, 1^*, \dots, 0)_{\mathcal{E}} & \text{if } k \text{ is even, } 6 \leq k < 2r + 1 \\ (a-2, 1, m-2, 0, \dots, 1^*, \dots, 0)_{\mathcal{E}} & \text{if } 2r + 1 < k \leq 3r \end{cases}$$

Now for each $1 \leq p < k-1$, we set the first p entries of the vectors above to 0 and add 1 to the $p+1$ entry. In the first vector (when $k=2$), there is nothing to check since in this case $1 \leq p$ but also $p < k-1 = 1$. In most cases, the resulting vector corresponds to the value $e_i + e_j$ where $2 \leq i, j < 3r$ (specifically, this is the case for the second vector and all values of p , and for the final two vectors in the case that $3 \leq p < k-1$). So we must check whether a value of this form is a counterexample.

Accordingly, we find the greedy solution for $e_i + e_j$ for all $2 \leq i, j < 3r$. First, suppose $2 \leq i, j \leq 2r+1$ and $m > 2$.

$$e_i + e_j = \begin{cases} e_{i+j-1} + e_1 & \text{if } i \text{ and } j \text{ even and } i + j \leq 2r + 2 \\ e_{(i+j)/2+r} + e_1 & \text{if } i \text{ and } j \text{ even and } i + j > 2r + 2 \\ e_{i+j-3} + e_3 & \text{if } i \text{ and } j \text{ odd and } i + j \leq 2r + 4 \\ e_{(i+j)/2+r-1} + e_3 & \text{if } i \text{ and } j \text{ odd and } i + j > 2r + 4 \\ e_{i+j-2} + e_2 & \text{if } i \text{ is odd and } j \text{ is even and } i + j \leq 2r + 4 \\ e_{r+(i+j-1)/2} + e_2 & \text{if } i \text{ is odd and } j \text{ is even and } i + j > 2r + 4 \end{cases}$$

Next suppose $2r + 1 < j < 3r$ and $m > 2$. Then

$$e_i + e_j = \begin{cases} e_{j+(i-3)/2} + e_3 & \text{if } 2 < i \leq 2r + 1 \text{ is odd and } i + 2j \leq 6r + 3 \\ e_{3r} + e_{i+2j-6r} & \text{if } 2 < i \leq 2r + 1 \text{ is odd and } i + 2j > 6r + 3 \\ e_{j-1+i/2} + e_2 & \text{if } 2 \leq i \leq 2r + 1 \text{ is even and } i + 2j \leq 6r + 2 \\ e_{3r} + e_{i+2j-6r} & \text{if } 2 \leq i \leq 2r + 1 \text{ is even and } i + 2j > 6r + 2 \\ e_{3r} + e_{2i+2j-8r-1} & \text{if } 2r + 1 < i < 3r, \text{ and } i + j \leq 5r + 1 \\ e_{3r} + e_{i+j-3r} & \text{if } 2r + 1 < i < 3r, \text{ and } i + j > 5r + 1 \end{cases}$$

In the special case $m = 2$, the above equations require three adjustments. If $m = 2$, then $e_3 + e_k = e_{k+2}$ for $2 \leq k < 2r$ and $e_3 + e_k = e_{k+1}$ for $2r + 1 \leq k < 3r + 1$. Thus if $m = 2$, anytime we see $e_k + e_3$ above where $k \geq 2$ and $k \neq 2r$, we eliminate the sum and replace it with a single coin value as the greedy solution. Observe that in all cases, the greedy solution for $e_i + e_j$ contains either 1 or 2 coins. Therefore $e_i + e_j$ is not a counterexample.

Now, we consider one by one the last three vectors from Equation 8 where $p = 1$ and $p = 2$. Begin with $e_4 - 1$. For $p = 1$ and $p = 2$, respectively, we have the following two vectors, associated candidates for counterexample, and greedy solutions.

Vector representation	Candidate for counterexample	Greedy solution
$(0, 1, m - 1, 0, \dots, 0)_{\mathcal{E}}$	$2am - m - a + 1$	e_4
$(0, 0, m, 0, \dots, 0)_{\mathcal{E}}$	$m(2a - 1)$	e_5

The greedy solutions above both use 1 coin, so neither of these values are counterexamples.

Now consider $e_k - 1$ where k is even, $6 \leq k < 2r + 1$ (see Equation 8 for the vector representation). For $p = 1$ and $p = 2$, respectively, we have the following results. (Recall that an asterisk (*) denotes that the value is in the $k - 1^{\text{st}}$ entry.)

Vector representation	Candidate for counterexample	Greedy solution
$(0, 1, m - 2, 0, \dots, 1^*, \dots, 0)_{\mathcal{E}}$	$e_{k-1} + 2am - 3a - m + 2$	e_k
$(0, 0, m - 1, 0, \dots, 1^*, \dots, 0)_{\mathcal{E}}$	$e_{k-1} + 2am - 2a - m + 1$	e_{k+1}

Again, the greedy solutions above both use 1 coin, so neither of these values are counterexamples.

Finally, consider $e_k - 1$ where $2r + 1 < k \leq 3r$ (see Equation 8). As before, for $p = 1$ and $p = 2$, respectively, we give the following two vectors, candidates for counterexample, and associated greedy solutions:

Vector representation	Candidate for counterexample	Greedy solution
$(0, 2, m - 2, 0, \dots, 1^*, \dots, 0)_{\mathcal{E}}$	$e_{k-1} + 2a(m - 1) - (m - 2)$	$e_k + e_1$
$(0, 0, m - 1, 0, \dots, 1^*, \dots, 0)_{\mathcal{E}}$	$e_{k-1} + (2a - 1)(m - 1)$	e_k

Once more, the greedy solutions above use either 1 or 2 coins, so neither of these values are counterexamples. Thus the coin system \mathcal{E} is orderly. \square

Theorem 4.7. *Let $r \geq 2$ and $1 < m \leq a$. Define $\mathcal{F} = (1, f_2, \dots, f_{3r})$ as follows:*

$$f_k = \begin{cases} 1 & \text{if } k = 1 \\ a & \text{if } k = 2 \\ f_{k-1} + a & \text{if } k \text{ is odd and } 3 \leq k \leq 2r + 1 \\ f_{k-1} + (m-1)(2a-1) - a & \text{if } k \text{ is even and } 3 < k < 2r + 1 \\ f_{k-1} + (m-1)(2a-1) & \text{if } 2r + 1 < k \leq 3r \end{cases}$$

The coin system \mathcal{F} has pattern $(+++ \dots - +)$.

Example 4.8. *For $r = 4$, $m = 3$, and $a = 4$, the coin system is*

$$\mathcal{F} = (1, 4, 8, 18, 22, 32, 36, 46, 50, 64, 78, 92).$$

Proof. The 3-coin prefix coin system $(f_1, f_2, f_3) = (1, a, 2a)$ is totally orderly by Theorem 2.8 where $f_3 - f_2 = a$. The 4-coin prefix coin system $(1, a, 2a, (m-1)(2a-1) + a)$ where $1 < m \leq a$ is not orderly because $2am$ is a counterexample. From Lemma 4.2, we know all prefix coin systems of length i where $5 \leq i \leq 3r-1$ are not orderly.

Finally, we show that the full coin system \mathcal{F} is orderly. Suppose that the coin system is not orderly. We use Theorem 2.3 to find the smallest counterexample. We find the greedy solution for $f_k - 1$ for all $k \geq 2$ and express it as a vector with respect to the coin system \mathcal{F} . In the vectors below, the asterisk (*) denotes that the value is in the $k-1^{\text{st}}$ entry.

$$(9) \quad f_k - 1 = \begin{cases} (a-1, 0, \dots, 0)_{\mathcal{F}} & \text{if } k = 2 \\ (a-1, 0, 0, \dots, 1^*, \dots, 0)_{\mathcal{F}} & \text{if } k \text{ is odd, } 3 \leq k \leq 2r + 1 \\ (a-m, 0, m-1, 0, \dots, 0)_{\mathcal{F}} & \text{if } k = 4 \\ (a-m, 0, m-2, 0, \dots, 1^*, \dots, 0)_{\mathcal{F}} & \text{if } k \text{ is even, } 6 \leq k < 2r + 1 \\ (a-m, 1, m-2, 0, \dots, 1^*, \dots, 0)_{\mathcal{F}} & \text{if } 2r + 1 < k \leq 3r \end{cases}$$

As before, for each $1 \leq p \leq k-1$, we set the first p entries of the above vectors to 0 and add 1 to the $p+1$ entry. Similar to the previous theorem, the case $k=2$ yields nothing. For the second vector, the resulting vector has only two 1's for all such p , which corresponds to the value $f_i + f_j$. The same is true for the last three vectors when $p \geq 3$. To settle these cases, we find the greedy solution for $f_i + f_j$ for all $2 \leq i, j < 3r$.

Suppose $2 \leq i, j \leq 2r+1$ and $m > 2$. The greedy solution for $f_i + f_j$ is:

$$(10) \quad f_i + f_j = \begin{cases} f_{i+j-1} & \text{if } i \text{ and } j \text{ even and } i+j \leq 2r+2 \\ f_{r+(i+j)/2} & \text{if } i \text{ and } j \text{ even and } i+j > 2r+2 \\ f_{i+j-3} + f_3 & \text{if } i \text{ and } j \text{ odd and } i+j \leq 2r+4 \\ f_{r-1+(i+j)/2} + f_3 & \text{if } i \text{ and } j \text{ odd and } i+j > 2r+4 \\ f_{i+j-2} + f_2 & \text{if } i \text{ is even and } j \text{ is odd and } i+j \leq 2r+3 \\ f_{r+(i+j-1)/2} + f_2 & \text{if } i \text{ is odd and } j \text{ is even and } i+j > 2r+3 \end{cases}$$

Suppose $2r + 1 < j < 3r$ and $m > 2$. Then we have:

$$(11) \quad f_i + f_j = \begin{cases} f_{j+(i-3)/2} + f_3 & \text{if } 2 \leq i \leq 2r + 1 \text{ is odd and } i + 2j \leq 6r + 3 \\ f_{3r} + f_{i+2j-6r} & \text{if } 2 \leq i \leq 2r + 1 \text{ is odd and } i + 2j > 6r + 3 \\ f_{j-1+i/2} + f_2 & \text{if } 2 \leq i \leq 2r + 1 \text{ is even and } i + 2j \leq 6r + 2 \\ f_{3r} + f_{i+2j-6r} & \text{if } 2 \leq i \leq 2r + 1 \text{ is even and } i + 2j > 6r + 2 \\ f_{3r} + f_{2i+2j-8r-1} & \text{if } 2r + 1 < i < 3r \text{ and } i + j \leq 5r + 1 \\ f_{3r} + f_{i+j-3r} & \text{if } 2r + 1 < i < 3r \text{ and } i + j > 5r + 1 \end{cases}$$

From this, we see that if $m > 2$, then $f_i + f_j$ is not a counterexample no matter the values of $3 \leq i, j < 3r$, because the greedy solutions given above always has 1 or 2 coins.

In the special case $m = 2$, the above equations require three adjustments. The greedy solution of $f_k + f_3$ is $f_{k+2} + f_1$ if $3 \leq k < 2r + 1$. And if $2r + 1 \leq k < 3r$, the greedy solution is $f_{k+1} + f_1$. So the expressions (a) $f_{i+j-3} + f_3$, (b) $f_{(i+j)/2+r-1} + f_3$ and (c) $f_{j+(i-3)/2} + f_3$ above need appropriate adjustments. Everything else remains as is, and again we conclude that $f_i + f_j$ is not a counterexample.

Now we consider the remaining special cases. Namely, we consider the last three vectors for $f_k - 1$ listed in Equation 9 in the case that $p = 1$ and $p = 2$. First we simply work out what candidates for minimum counterexample are generated from them. In fact, almost all the candidates will a greedy solution in the format of $f_k + (m - 1)f_1$ for some k .

We begin with $f_4 - 1$. (See Equation 9 for the vector representation of $f_4 - 1$.) The vectors associated to $p = 1$ and $p = 2$ and the resulting candidates for counterexamples are below.

Vector representation	Candidate for counterexample	Greedy solution
$(0, 1, m - 1, 0, \dots, 0)_{\mathcal{F}}$	$2am - a$	$f_4 + (m - 1)f_1$
$(0, 0, m, 0, \dots, 0)_{\mathcal{F}}$	$2am$	f_6 , if $m = a = 2$ else $f_5 + (m - 1)f_1$

Next, consider $f_k - 1$ where k is even and $6 \leq k < 2r + 1$. (Again, see Equation 9 for the vector representation of $f_k - 1$.) The vectors associated to $p = 1$ and $p = 2$ and the resulting candidates for counterexamples are below. An asterisk (*) denotes that the value is in the $k - 1^{st}$ entry.

Vector representation	Candidate for counterexample	Greedy solution
$(0, 1, m - 2, 0, \dots, 1^*, \dots, 0)_{\mathcal{F}}$	$f_{k-1} + 2am - 3a$	$f_k + (m - 1)f_1$
$(0, 0, m - 1, 0, \dots, 1^*, \dots, 0)_{\mathcal{F}}$	$f_{k-1} + 2am - 2a$	f_{k+2} , if $m = a = 2$, else $f_{k+1} + (m - 1)f_1$

Now consider $f_k - 1$ where $2r + 1 < k < 3r$. The vectors associated to $p = 1$ and $p = 2$ and the resulting candidates for counterexamples are below.

Vector representation	Candidate for counterexample	Greedy solution
$(0, 2, m - 2, 0, \dots, 1^*, \dots, 0)_{\mathcal{F}}$	$f_{k-1} + 2am - 2a$	$f_k + (m - 1)f_1$
$(0, 0, m - 1, 0, \dots, 1^*, \dots, 0)_{\mathcal{F}}$	$f_{k-1} + 2am - 2a$	$f_k + (m - 1)f_1$

We now show that none of the values given above as candidates for the minimum counterexample are, in fact, counterexamples. If $m = 2$, then the greedy solutions in each of the three cases are optimal, since each solution has only 1 or 2 coins. For $m > 2$, a total of m coins are in the greedy solution, and we must work carefully to show that the solution is optimal. Observe that all the greedy solutions above are of the form $f_k + (m - 1)f_1$ for some value $4 \leq k < 3r$. So it remains to show that an optimal solution for $f_k + (m - 1)f_1$ has m coins.

Suppose that there exists a k such that an optimal solution $f_k + (m - 1)f_1$ has fewer than m coins. Take the smallest such value, which we denote by $v = f_k + (m - 1)f_1$. By Theorem 2.13, an optimal solution for v does not use f_1 or f_k .

We know that an optimal solution for v uses at most one coin f_j where $2r + 1 < j < 3r$ because the sum of any two such coins is larger than $f_{3r} + m - 1$ and $v \leq f_{3r} + m - 1$.

An optimal solution for v has at most one even-indexed coin f_j where $2 \leq j < 2r + 1$, because if there were two such coins f_i and f_j , then they could be replaced with a single coin (either f_{i+j-1} or $f_{r+(i+j)/2}$, see Equation 10), which contradicts the optimality of the solution.

If an optimal solution for v contains an even indexed coin f_j where $3 < j < 2r + 1$, then by combining that coin with any other coin in the solution, we can trade the coin f_j and the other coin f_i for f_2 and a larger indexed coin (see Equations 10 and 11). Therefore, without loss of generality, the optimal solution for v either has no even-indexed coins f_j where $2 \leq j < 2r + 1$ or else contains only f_2 .

Next, if an optimal solution contains two coins f_i and f_j where i and j are odd and $3 < i, j \leq 2r + 1$, then we can trade them in for f_3 and another coin f_r (again, see Equations 10 and 11). Therefore without loss of generality, the optimal solution has at most one coin f_i with odd index $3 < i \leq 2r + 1$.

If an optimal solution for v has both a coin of value f_i where i is odd $3 \leq i \leq 2r + 1$ and a coin of value f_j where $2r + 1 < j \leq 3r$, we can exchange the pair of coins for a different pair of coins f_q and f_3 where $2r + 1 < q \leq 3r$ (Equations 11). Therefore without loss of generality, the optimal solution for v has at most one coin with index $3 < j \leq 3r$.

Putting all of these observations together, we conclude that any optimal solution for v can be exchanged for another optimal solution fitting one of the following four scenarios:

- (1) $v = xf_3$, where $0 < x \leq m - 1$
- (2) $v = f_2 + yf_3$, where $0 < y \leq m - 2$
- (3) $v = f_2 + xf_3 + f_j$, where j is odd, $3 \leq j \leq 2r + 1$, and $0 < x \leq m - 3$.
- (4) $v = yf_3 + f_j$, where j is odd, $3 \leq j \leq 2r + 1$, and $0 < y \leq m - 2$.
- (5) $v = f_2 + xf_3 + f_j$, where $2r + 1 < j \leq 3r$, and $0 < x \leq m - 3$.
- (6) $v = yf_3 + f_j$, where $2r + 1 < j \leq 3r$, and $0 < y \leq m - 2$.

Now for (1)–(2), we compute:

$$\begin{aligned}
f_3 < v &\leq f_3(m - 1) \\
&\leq f_j + (2a - 1)(m - 1) + a + m - 1 - a \\
&= f_4 + m - 1 - a \\
&< f_4.
\end{aligned}$$

And for (3)–(6), we compute:

$$\begin{aligned} f_j &< v \leq f_j + 2a(m - 2) \\ &\leq f_j + (2a - 1)(m - 1) - 2a + m - 1 \\ &< f_{j+1}. \end{aligned}$$

In all cases, v falls between consecutive coin values, we conclude that f_3 (respectively, f_j) is a coin value in the greedy solution for v . This is a contradiction because on one hand v was the minimum counterexample (and as such, the optimal solutions for v use different coin values from the greedy solution), but then on the other hand, we showed that a coin in an optimal solution for v is also in the greedy solution. Hence there is not a minimal counterexample. Therefore the coin system \mathcal{F} is orderly. \square

REFERENCES

- [1] Anna Adamaszek and Michal Adamaszek. Combinatorics of the change-making problem. *European J. Combin.*, 31(1):47–63, 2010.
- [2] Valentina Cacchiani, Manuel Iori, Alberto Locatelli, and Silvano Martello. Knapsack problems – an overview of recent advances. Part I: Single knapsack problems. *Comput. Oper. Res.*, 143:Paper No. 105692, 13, 2022.
- [3] Valentina Cacchiani, Manuel Iori, Alberto Locatelli, and Silvano Martello. Knapsack problems – an overview of recent advances. Part II: Multiple, multidimensional, and quadratic knapsack problems. *Comput. Oper. Res.*, 143:105693, 2022.
- [4] Xuan Cai. Canonical coin systems for change-making problems. In *2009 Ninth International Conference on Hybrid Intelligent Systems*, volume 1, pages 499–504, 2009.
- [5] Timothy M. Chan and Qizheng He. More on change-making and related problems. *Journal of Computer and System Sciences*, 124:159–169, 2022.
- [6] Lena Chang and James F. Korsh. Canonical coin changing and greedy solutions. *J. ACM*, 23(3):418–422, jul 1976.
- [7] S. K. Chang and A. Gill. Algorithmic solution of the change-making problem. *J. ACM*, 17(1):113–122, jan 1970.
- [8] L. J. Cowen, Robert Cowen, and Arthur Steinberg. Totally greedy coin sets and greedy obstructions. *Electron. J. Combin.*, 15(1):Research Paper 90, 13 pages, 2008.
- [9] Steffen Goebbels, Frank Gurski, Jochen Rethmann, and Eda Yilmaz. Change-making problems revisited: a parameterized point of view. *J. Comb. Optim.*, 34(4):1218–1236, 2017.
- [10] T. C. Hu and M. L. Lenard. Optimality of a heuristic solution for a class of knapsack problems. *Operations Res.*, 24(1):193–196, 1976.
- [11] John Dewey Jones. Orderly currencies. *Amer. Math. Monthly*, 101(1):36–38, 1994.
- [12] Hans Kellerer, Ulrich Pferschy, and David Pisinger. *Knapsack problems*. Springer-Verlag, Berlin, 2004.
- [13] Dexter Kozen and Shmuel Zaks. Optimal bounds for the change-making problem. *Theoret. Comput. Sci.*, 123(2):377–388, 1994.
- [14] M. J. Magazine, G. L. Nemhauser, and L. E. Trotter, Jr. When the greedy solution solves a class of knapsack problems. *Operations Res.*, 23(2):207–217, 1975.
- [15] Silvano Martello and Paolo Toth. *Knapsack problems: algorithms and computer implementations*. John Wiley & Sons, Inc., 1990.
- [16] Stephen B. Maurer. Comment: “Orderly currencies” [Amer. Math. Monthly **101** (1994), no. 1, 36–38] by J. D. Jones. *Amer. Math. Monthly*, 101(5):419, 1994.
- [17] Ryuhei Miyashiro and Yuma Suzuki. Characterization of canonical systems with six types of coins for the change-making problem. URL: <https://arxiv.org/abs/2111.12392>, 2021.
- [18] David Pearson. A polynomial-time algorithm for the change-making problem. *Oper. Res. Lett.*, 33(3):231–234, 2005.
- [19] B. N. Tien and T. C. Hu. Error bounds and the applicability of the greedy solution to the coin-changing problem. *Operations Res.*, 25(3):404–418, 1977.