

# Fast convergence to non-isolated minima: four equivalent conditions for $C^2$ functions

Quentin Rebjock and Nicolas Boumal\*

March 2, 2023

## Abstract

Optimization algorithms can see their local convergence rates deteriorate when the Hessian at the optimum is singular. These singularities are inescapable when the optima are non-isolated. Yet, under the right circumstances, several algorithms preserve their favorable rates even when optima form a continuum (e.g., due to over-parameterization). This has been explained under various structural assumptions, including the Polyak–Łojasiewicz inequality, Quadratic Growth and the Error Bound. We show that, for cost functions which are twice continuously differentiable ( $C^2$ ), those three (local) properties are equivalent. Moreover, we show they are equivalent to the Morse–Bott property, that is, local minima form differentiable submanifolds, and the Hessian of the cost function is positive definite along its normal directions. We leverage this insight to improve local convergence guarantees for safe-guarded Newton-type methods under any (hence all) of the above assumptions. First, for adaptive cubic regularization, we secure quadratic convergence even with inexact subproblem solvers. Second, for trust-region methods, we argue convergence can fail with an exact subproblem solver, then proceed to show linear convergence with an inexact one (Cauchy steps).

## 1 Introduction

We consider local convergence of algorithms for unconstrained optimization problems of the form

$$\min_{x \in \mathcal{M}} f(x),$$

where  $\mathcal{M}$  is a Riemannian manifold<sup>1</sup> and  $f: \mathcal{M} \rightarrow \mathbb{R}$  is at least  $C^1$  (continuously differentiable).

When  $f$  is  $C^2$  (twice continuously differentiable), the most classical local convergence results ensure favorable rates for standard algorithms *provided* they converge to a non-singular local minimum  $\bar{x}$ , that is, one such that the Hessian  $\nabla^2 f(\bar{x})$  is positive definite. And indeed, those rates can degrade if the Hessian is merely positive semidefinite. For example, with  $f(x) = x^4$ , gradient descent (with an appropriate step-size) converges only sublinearly to the minimum, and Newton’s method converges only linearly.

This is problematic if the minimizers of  $f$  are not isolated, because in that case the Hessian cannot be positive definite there. This situation arises commonly in applications for structural reasons such as over-parameterization, redundant parameterizations and symmetry—see Section 1.2.

Notwithstanding, algorithms often exhibit good local behavior near non-isolated minimizers. As early as the 1960s, this has prompted investigations into properties that such cost functions may satisfy and which lead to fast local rates despite singular Hessians. We study four such properties.

---

\*Ecole Polytechnique Fédérale de Lausanne (EPFL), Institute of Mathematics. This work was supported by the Swiss State Secretariat for Education, Research and Innovation (SERI) under contract number MB22.00027.

<sup>1</sup>The contributions are relevant for  $\mathcal{M} = \mathbb{R}^n$  too. We treat the more general manifold case as it involves only mild overhead in notation, summarized in Table 2.

In all that follows, we are concerned with the behavior of algorithms in the vicinity of its local minima. Since we do not assume that they are isolated, rather than selecting one local minimum  $\bar{x}$ , we select all local minima of the same value. Formally, given a local minimum  $\bar{x}$ , let

$$\mathcal{S} = \{x \in \mathcal{M} : x \text{ is a local minimum of } f \text{ and } f(x) = f_{\mathcal{S}}\} \quad (1)$$

denote the set of all local minima with a given value  $f_{\mathcal{S}} = f(\bar{x})$ .

For  $f$  of class  $C^2$ , it is particularly favorable if  $\mathcal{S}$  is a differentiable submanifold of  $\mathcal{M}$  around  $\bar{x}$ . In that case the set  $\mathcal{S}$  has a *tangent space*  $T_{\bar{x}}\mathcal{S}$  at  $\bar{x}$ . It is easy to see that each vector  $v \in T_{\bar{x}}\mathcal{S}$  must be in the kernel of the Hessian  $\nabla^2 f$  at  $\bar{x}$  because the gradient  $\nabla f$  is constant (zero) on  $\mathcal{S}$ . Thus,  $T_{\bar{x}}\mathcal{S} \subseteq \ker \nabla^2 f(\bar{x})$ . Since  $\bar{x}$  is a local minimum, we also know that  $\nabla^2 f(\bar{x})$  is positive semidefinite. Then, in the spirit of asking the Hessian to be “as positive definite as possible”, the best we can hope for is that the kernel of  $\nabla^2 f(\bar{x})$  is exactly  $T_{\bar{x}}\mathcal{S}$ , in which case the restriction of  $\nabla^2 f(\bar{x})$  to the *normal space*  $N_{\bar{x}}\mathcal{S}$ , that is, the orthogonal complement of  $T_{\bar{x}}\mathcal{S}$  in  $T_{\bar{x}}\mathcal{M}$ , is positive definite.

We call this the *Morse–Bott* property (MB), and we write  $\mu$ -MB to indicate that the positive eigenvalues are at least  $\mu > 0$ . The definition requires  $f$  to be twice differentiable.

**Definition 1.1.** *Let  $\bar{x}$  be a local minimum of  $f$  with associated set  $\mathcal{S}$  (1). We say  $f$  satisfies the Morse–Bott property at  $\bar{x}$  if*

$$\mathcal{S} \text{ is a } C^1 \text{ submanifold around } \bar{x} \quad \text{and} \quad \ker \nabla^2 f(\bar{x}) = T_{\bar{x}}\mathcal{S}. \quad (\text{MB})$$

*If also  $\langle v, \nabla^2 f(\bar{x})[v] \rangle \geq \mu \|v\|^2$  for some  $\mu > 0$  and all  $v \in N_{\bar{x}}\mathcal{S}$  then we say  $f$  satisfies  $\mu$ -MB at  $\bar{x}$ .*

At first, a reasonable objection to the above is that one may not want to assume that  $\mathcal{S}$  is a submanifold. Perhaps for that reason, it is far more common to encounter other assumptions in the optimization literature. We focus on three: *Polyak–Lojasiewicz* (PL), *error bound* (EB) and *quadratic growth* (QG). The first goes back to the 1960s [Polyak, 1963]. The latter two go back at least to the 1990s [Luo and Tseng, 1993, Bonnans and Ioffe, 1995].

Below, the first two definitions (as stated) require  $f$  to be differentiable. The distance to a set is defined as usual:  $\text{dist}(x, \mathcal{S}) = \inf_{y \in \mathcal{S}} \text{dist}(x, y)$  where  $\text{dist}(x, y)$  is the Riemannian distance on  $\mathcal{M}$ .

**Definition 1.2.** *Let  $\bar{x}$  be a local minimum of  $f$  with associated set  $\mathcal{S}$  (1). We say  $f$  satisfies*

- *the Polyak–Lojasiewicz inequality with constant  $\mu > 0$  ( $\mu$ -PL) around  $\bar{x}$  if*

$$f(x) - f_{\mathcal{S}} \leq \frac{1}{2\mu} \|\nabla f(x)\|^2; \quad (\text{PL})$$

- *the error bound with constant  $\mu > 0$  ( $\mu$ -EB) around  $\bar{x}$  if*

$$\mu \text{dist}(x, \mathcal{S}) \leq \|\nabla f(x)\|; \quad (\text{EB})$$

- *quadratic growth with constant  $\mu > 0$  ( $\mu$ -QG) around  $\bar{x}$  if*

$$f(x) - f_{\mathcal{S}} \geq \frac{\mu}{2} \text{dist}(x, \mathcal{S})^2; \quad (\text{QG})$$

*all understood to hold for all  $x$  in some neighborhood of  $\bar{x}$ .*

Note that all the definitions are *local* around a *point*  $\bar{x}$ . Two observations are immediate: (i) QG implies that  $\bar{x}$  is a strict minimum relatively to  $\mathcal{S}$  (1), meaning that  $f(x) > f_{\mathcal{S}}$  for all  $x \notin \mathcal{S}$  close enough to  $\bar{x}$ , and (ii) both EB and PL imply that critical points and  $\mathcal{S}$  coincide around  $\bar{x}$ . Thus, both EB and PL rule out existence of saddle points near  $\bar{x}$ . By extension, we say that  $f$  satisfies any of these four properties around a *set* of local minima if it holds around each point of that set.

	Statement	$f$ is $C^p$	Constants	Global?	Comments
PL $\Rightarrow$ QG	Prop. 2.2	$p \geq 1$	$\mu' = \mu$	Yes	<a href="#">Bolte et al. [2010]</a>
MB $\Rightarrow$ QG	Prop. 2.3	$p \geq 2$	$\mu' < \mu$	n/a	Taylor expansion
EB $\Rightarrow$ PL	Prop. 2.4	$p \geq 2$	$\mu' < \mu$	No	<a href="#">Karimi et al. [2016]</a>
	Rmrk 2.5	$p \geq 1$	$\mu' = \frac{\mu^2}{L}$	Yes	
QG $\Rightarrow$ EB	Prop. 2.8	$p \geq 2$	$\mu' < \mu$	No	Fails for $p = 1$ , Rmrk 2.9
PL $\Rightarrow$ EB	Rmrk 2.10	$p \geq 1$	$\mu' = \mu$	Yes	<a href="#">Karimi et al. [2016]</a>
PL $\Rightarrow$ MB	Cor. 2.17	$p \geq 2$	$\mu' = \mu$	n/a	Fails for $p = 1$ , Rmrk 2.19

Table 1: Summary of implications. For a row  $A \Rightarrow B$ , we mean that  $A$  with constant  $\mu$  implies  $B$  (in a possibly different neighborhood) with constant  $\mu'$ . The condition  $\mu' < \mu$  means  $\mu'$  can be taken arbitrarily close to  $\mu$  by shrinking the neighborhood. For  $EB \Rightarrow PL$  with  $p = 1$ , we let  $L$  denote the Lipschitz constant of  $\nabla f$ . If  $B$  holds globally when  $A$  holds globally, we write “Yes” in the column “Global?” Implications we found in the literature are marked with a reference.

## 1.1 Contributions

A number of relationships between PL, EB and QG are well known already for  $f$  of class  $C^1$ : see Table 1 and Section 1.3. Our first main contribution in this paper is to show that:

*If  $f$  is of class  $C^2$ , then PL, EB, QG and MB are essentially equivalent.*

Here, “essentially” means that the constant  $\mu$  may degrade (arbitrarily little) and the neighborhoods where properties hold may shrink. Notably, we show that if  $f$  is  $C^p$  with  $p \geq 2$ , then PL, EB and QG all imply that  $\mathcal{S}$  is locally smooth (at least  $C^{p-1}$ ). We also give counter-examples when  $f$  is only  $C^1$ . Explicitly, in Section 2 we summarize known results for  $f$  of class  $C^1$  and we contribute the following:

- Theorem 2.16 shows PL at  $\bar{x}$  implies  $\mathcal{S}$  is a  $C^{p-1}$  submanifold around  $\bar{x}$  if  $f$  is  $C^p$  with  $p \geq 2$ . Remark 2.19 provides counter-examples if  $f$  is only  $C^1$ . If  $f$  is analytic, so is  $\mathcal{S}$ , as already shown by [Feehan \[2020\]](#).
- Corollary 2.17 deduces that  $\mu$ -PL implies  $\mu$ -MB if  $f$  is  $C^2$ .
- Proposition 2.8 shows  $\mu$ -QG implies  $\mu'$ -EB with  $\mu' < \mu$  arbitrarily close if  $f$  is  $C^2$ . Remark 2.9 provides a counter-example if  $f$  is only  $C^1$ .
- Proposition 2.4 shows  $\mu$ -EB implies  $\mu'$ -PL with  $\mu' < \mu$  arbitrarily close if  $f$  is  $C^2$ . If  $f$  is only  $C^1$  with  $L$ -Lipschitz continuous  $\nabla f$ , [Karimi et al. \[2016\]](#) showed the same with  $\mu' = \mu^2/L$ .

We then proceed to study local convergence rates of optimization algorithms when minimizers are not isolated but the  $C^2$  cost function  $f$  satisfies any (hence all) of the above conditions. We aim for super-linear convergence, since linear convergence is well understood. To this end, in Section 4, we first illustrate how Newton’s method can fail even if initialized arbitrarily close to the manifold of solutions. Accordingly, we turn to the usual safe-guarded versions of Newton’s method:

- Cubic regularization enjoys super-linear convergence under PL, as shown by [Nesterov and Polyak \[2006\]](#). [Yue et al. \[2019\]](#) further showed quadratic convergence under EB. (This illustrates how the observation that PL and EB are equivalent can help algorithm analysis.) Both references assume an exact subproblem solver. We contribute by showing quadratic convergence still holds with inexact subproblem solvers (Theorem 4.1).
- For the trust-region method with exact subproblem solver, we were surprised to find that even basic capture-type convergence properties can fail in the presence of non-isolated local minima (Section 4.2). Notwithstanding, common implementations of the trust-region method use a truncated conjugate gradients (tCG) subproblem solver, and those do, empirically, exhibit super-linear convergence under the favorable conditions discussed above. We discuss this

further in Remark 4.14, and we show a partial result in Theorem 4.9, namely, that using the Cauchy step (i.e., the first iterate of tCG) yields linear convergence.

As is classical, to prove the latter results, we rely on capture theorems and Lyapunov stability. Those hold under assumptions of vanishing step-sizes and bounded path length. In Section 3, we state those building blocks succinctly, adapted to accommodate non-isolated local minima.

## 1.2 Non-isolated minima in applications

We now illustrate how optimization problems with continuous sets of minima occur in applications. In all three scenarios below, we can cast the cost function  $f: \mathcal{M} \rightarrow \mathbb{R}$  as a composition of some other




Figure 1: Optimization through the map  $\varphi$ .

function  $g: \mathcal{N} \rightarrow \mathbb{R}$  through a map  $\varphi: \mathcal{M} \rightarrow \mathcal{N}$ , where  $\mathcal{N}$  is a smooth manifold (see Figure 1 and [Levin et al., 2022b]). If  $g$  and  $\varphi$  are twice differentiable, then the Morse–Bott property (MB) for  $f = g \circ \varphi$  can come about as follows. Consider a local minimum  $\bar{y}$  for  $g$ . The set  $\mathfrak{X} = \varphi^{-1}(\bar{y})$  consists of local minima for  $f$ . Pick a point  $\bar{x} \in \mathfrak{X}$ . Assume  $x \mapsto \text{rank } D\varphi(x)$  is constant in a neighborhood of  $\bar{x}$ . Then, the set  $\mathfrak{X}$  is an embedded submanifold of  $\mathcal{M}$  around  $\bar{x}$  with tangent space  $\ker D\varphi(\bar{x})$ . Moreover, the Hessians of  $f$  and  $g$  at  $\bar{x}$  are related by

$$\nabla^2 f(\bar{x}) = D\varphi(\bar{x})^* \circ \nabla^2 g(\varphi(\bar{x})) \circ D\varphi(\bar{x}). \quad (2)$$

Therefore, if  $\nabla^2 g(\varphi(\bar{x}))$  is positive definite, then  $\ker \nabla^2 f(\bar{x}) = T_{\bar{x}}\mathfrak{X}$  and  $\nabla^2 f(\bar{x})$  is positive definite along the orthogonal complement. In other words:  $f$  satisfies the Morse–Bott property (MB) at  $\bar{x}$ . We present below a few concrete examples of optimization problems where this can happen.

**Over-parameterization and non-linear regression.** Consider minimizing  $f(x) = \frac{1}{2}\|F(x) - b\|^2$  with  $F: \mathbb{R}^m \rightarrow \mathbb{R}^n$  a  $C^2$  function. We cast this as above with  $g(y) = \frac{1}{2}\|y - b\|^2$  and  $\varphi = F$ . Suppose  $\mathfrak{X} = \varphi^{-1}(b) = \{x : F(x) = b\}$  is non-empty (interpolation regime), which is typical in deep learning. This is the set of global minimizers of  $f$ . If  $\text{rank } DF(x)$  is equal to a constant  $r$  in a neighborhood of  $\mathfrak{X}$ , then  $\mathfrak{X}$  is a smooth submanifold of  $\mathbb{R}^m$  of dimension  $m - r$ . If additionally the problem is over-parameterized, that is,  $m > n \geq r$ , then  $\mathfrak{X}$  has positive dimension (see Figure 2 for an illustration). The discussion above immediately implies that  $f$  satisfies MB on  $\mathfrak{X}$ . See also Nesterov and Polyak [2006, §4.2] who argue that PL holds in this setting.

**Redundant parameterizations and submersions.** Say we want to minimize  $g: \mathcal{N} \rightarrow \mathbb{R}$  constrained to  $\tilde{\mathcal{N}} \subseteq \mathcal{N}$ . If  $\tilde{\mathcal{N}}$  is complicated, and if we have access to a parameterization  $\varphi$  for that set (so that  $\varphi(\mathcal{M}) = \tilde{\mathcal{N}}$ ), it may be advantageous to minimize  $f = g \circ \varphi$  instead. If the parameterization is redundant, this can cause  $f$  to have non-isolated minima, even if the minima of  $g$  are isolated.

As an example, consider minimizing  $g: \mathbb{R}^{m \times n} \rightarrow \mathbb{R}$  over the bounded-rank matrices  $\tilde{\mathcal{N}} = \{Y \in \mathbb{R}^{m \times n} : \text{rank } Y \leq r\}$ . A popular approach consists in lifting the search space to  $\mathcal{M} = \mathbb{R}^{m \times r} \times \mathbb{R}^{n \times r}$  and minimizing  $f = g \circ \varphi$ , where  $\varphi: \mathcal{M} \rightarrow \mathcal{N}$  is defined as  $\varphi(L, R) = LR^\top$ . The parameterization is redundant because  $\varphi(LJ^{-1}, RJ^\top) = \varphi(L, R)$  for all invertible  $J$ . In particular, given a local minimum  $Y \in \tilde{\mathcal{N}}$  of  $g$ , the fiber  $\varphi^{-1}(Y)$  is unbounded, which hinders convergence analyses (see [Levin et al., 2022a]). However, if  $Y$  is of maximal rank  $r$  then  $D\varphi$  has constant rank in a neighborhood of  $\varphi^{-1}(Y)$ . From the discussion above, it follows that  $f$  satisfies MB on  $\varphi^{-1}(Y)$  if the (Riemannian) Hessian of  $g$  (restricted to the manifold of rank- $r$  matrices) is positive definite.




Figure 2: (Left) Submanifold of minima where MB holds. (Middle) A  $C^1$  function that satisfies QG but not PL. (Right) A  $C^1$  function that satisfies PL and whose solution set is a cross.

Similarly, Burer and Monteiro [2003, 2005] introduced a popular approach to minimize a function  $g$  over the set of positive semidefinite matrices of bounded rank through the map  $\varphi: Y \mapsto YY^\top$ . The resulting function  $f = g \circ \varphi$  can have non-isolated minima. However, we can ensure that MB holds on the maximal rank fiber with the same arguments as above, for example if  $g$  is strongly convex [Zhang, 2022]. This further extends to tensors [Li and Li, 2022].

**Symmetries and quotients.** Some optimization problems have intrinsic symmetries. For example, in estimation problems, if the measurements are invariant under particular transformations of the signal, then the signal can only be retrieved up to those transformations. The likelihood function then has symmetries, and possibly a continuous set of optima as a result. Sometimes, factoring these symmetries out (that is, passing to the quotient) yields a quotient manifold, and we can investigate optimization on that manifold [Absil et al., 2008]. In the notation of our general framework above,  $\varphi$  is then the quotient map. In particular,  $\varphi$  is a submersion, so that if  $\bar{y} \in \mathcal{N}$  is a non-singular minimum of  $g$  then  $f$  satisfies MB on  $\varphi^{-1}(\bar{y})$  (which is a submanifold of dimension  $\dim \mathcal{M} - \dim \mathcal{N}$ ). See also [Boumal, 2023, §9.9] for the case where  $\mathcal{N}$  is a Riemannian quotient of  $\mathcal{M}$ .

### 1.3 Related work

**Historical note.** Discussions about convergence to singular minima appear in the literature at least as early as [Polyak, 1987, §6.1]. Luo and Tseng [1993] introduced the EB inequality explicitly to study gradient methods around singular minima. The QG property is arguably as old as optimization, though the earliest work we could locate is by Bonnans and Ioffe [1995]. They employed QG to understand complicated landscapes with non-isolated minima. Lojasiewicz [1963, 1982] introduced his inequalities and used them subsequently to analyze gradient flow trajectories. Specifically, he proved that for analytic functions the trajectories either converge to a point or diverge. Concurrently, Polyak [1963] introduced what became known as the Polyak–Lojasiewicz (PL) variant (also “gradient dominance”) to study both gradient flows and discrete gradient methods. Later, Kurdyka [1998] developed generalizations now known as Kurdyka–Lojasiewicz (KL) inequalities. They are satisfied by most functions encountered in practice, as discussed in [Attouch et al., 2010, §4]. In contrast, the Morse–Bott property has received little attention in the optimization literature. Early work by Shapiro [1988] analyzes perturbations of optimization problems assuming a property similar to MB. There is also a mention of gradient flow under MB in [Helmke and Moore, 1996, Prop. 12.3].

**Relationships between properties.** Several articles have explored the interplay between MB, PL, EB and QG in the last decades. The implication  $PL \Rightarrow QG$  has a complex history. It can be

obtained as a corollary from [Ioffe, 2000, Basic lemma] (based on Ekeland’s variational principle; see also [Drusvyatskiy et al., 2015, Lem. 2.5]). It also follows from Łojasiewicz-type arguments that consist in bounding the length of gradient flow trajectories. For example, Bolte et al. [2010] study growth under KL inequalities, with PL  $\Rightarrow$  QG as a special case. Assuming that the function is only lower-semicontinuous, Drusvyatskiy et al. [2013, Cor. 3.2] show that  $\mu$ -EB  $\Rightarrow$   $\mu'$ -QG with arbitrary  $\mu' < \mu$ . They also show  $\mu$ -QG  $\Rightarrow$   $\mu'$ -EB but without control of  $\mu'$ . The equivalence between EB and QG was also explored in [Drusvyatskiy and Lewis, 2018] for composite non-smooth *convex* functions. More recently, Karimi et al. [2016] established implications between several properties encountered in the optimization literature. In particular, they show that PL and EB are equivalent, and that they both imply QG. Bolte et al. [2017] proved a similar result for convex and potentially *non-smooth* functions (see also [Zhang, 2017]). Implications between PL and EB are also reported in [Drusvyatskiy et al., 2021, Thm. 3.7, Prop. 3.8] for non-smooth functions and under general conditions. In the context of functional analysis, Feehan [2020] proved that MB and PL are equivalent for *analytic* functions defined on Banach spaces. The work of [Wojtowysch, 2021a, Ex. 2.9] also mentions that MB implies PL for  $C^2$  functions. A more general implication is given by Arbel and Mairal [2022, Prop. 1] for parameterized optimization problems. Previously, Bonnans and Ioffe [1995] had exhibited sufficient conditions (similar to MB) for QG to hold. As a side note, Marteau-Ferey et al. [2022] proved that MB is a sufficient condition to ensure that a function is locally decomposable as a sum of squares of smooth functions.

**Convergence guarantees.** The error bound approach of Luo and Tseng [1993] has proven to be fruitful as multiple analyses based on this condition followed. Notably, Tseng [2000] proved local super-linear convergence rates for some Newton-type methods applied to systems of non-linear equations. They relied specifically on EB and did not assume isolated solutions. Later, Yamashita and Fukushima [2001] employed EB to establish capture theorems and quadratic convergence rates for the Levenberg–Marquardt method, and Fan and Yuan [2005] generalized their results. More recently, Bellavia and Morini [2015] found that two adaptive regularized methods converge quadratically for non-linear least-squares problems (assuming that EB holds).

An early work of Anitescu [2000] combines the QG property with other conditions to ensure isolated minima in constrained optimization, then deducing convergence results. Later, QG has found applications mainly in the context of convex optimization: see [Liu and Wright, 2015] for coordinate descent, [Necoara et al., 2019] for various gradient methods, and [Drusvyatskiy and Lewis, 2018] for the proximal gradient method. It is also worth mentioning that the definition of QG does not require differentiability of the function. For this reason, QG is valuable to study algorithms in non-smooth optimization too [Davis and Jiang, 2022, Lewis and Tian, 2022].

The literature about convergence results based on Łojasiewicz inequalities is vast and we touch here on some particularly relevant references. Absil et al. [2005] discretized the arguments from [Łojasiewicz, 1982] and obtained capture results for a broad class of optimization algorithms. Later, such arguments have been used in many contexts to prove algorithmic convergence guarantees, among which [Attouch et al., 2010, Bolte et al., 2014] are particularly influential works. Moreover, Attouch et al. [2013] proposed a general abstract framework based on KL to derive capture results and convergence rates, and Frankel et al. [2015] extended their statements. See also [Necoara and Lupu, 2020] for a framework that encompasses higher-order methods. Li and Pong [2018] studied the preservation of Łojasiewicz inequalities under function transformations (such as sums and compositions). See also [Bassily et al., 2018, Terjék and González-Sánchez, 2022] for the preservation of PL through function compositions. Interestingly, the PL inequality is known to be a necessary condition for gradient descent to converge linearly [Abbaszadehpeivasti et al., 2022, Thm. 5]. Recently, Yue et al. [2022] proved that acceleration is impossible to minimize globally PL functions: gradient descent is optimal absent further structure. Assuming PL, Polyak et al. [2022] formulated stopping criteria for gradient methods when the gradient is corrupted with noise. Using KL inequalities, Noll and Rondepierre [2013] and Khanh et al. [2022] analyzed the convergence of line-search gradient descent and trust-region methods. Łojasiewicz inequalities have also proved relevant for the study of second-order algorithms and super-linear convergence rates. A prominent example is the regularized Newton algorithm that converges super-linearly when PL holds, as shown in [Nesterov

	$\text{Exp}_x(s)$	$\text{Log}_x(y)$	$\Gamma_x^y, \Gamma_v$	$\text{dist}(x, y)$	$\text{inj}(x)$
$\mathcal{M} = \mathbb{R}^n$	$x + s$	$y - x$	identity	$\ x - y\ $	$+\infty$

Table 2: Simplifications in the case where  $\mathcal{M}$  is a Euclidean space. Here,  $\Gamma_x^y$  denotes parallel transport along the minimizing geodesic between  $x$  and  $y$  (assuming they are close enough).

and Polyak, 2006]. More recently, Zhou et al. [2018], Yue et al. [2019] provided finer analyses of this algorithm, respectively assuming Lojasiewicz inequalities and EB. Qian and Pan [2022] extended the abstract framework of Attouch et al. [2013] to establish super-linear convergence rates.

Stochastic algorithms have also been extensively studied through Lojasiewicz inequalities and we briefly mention a few references here. Dereich and Kassing [2019, 2021] analyzed stochastic gradient descent (SGD) in the presence of non-isolated minima, using Lojasiewicz inequalities among other things. Local analyses of SGD using PL inequalities are given by Li et al. [2021] and Wojtowytsch [2021a]. Ko and Li [2022] studied the local stability and convergence of SGD in the presence of a compact set of minima with a condition that is weaker than PL. As for second-order algorithms, Masiha et al. [2022] proved that a stochastic version of regularized Newton has fast convergence under PL.

Lojasiewicz inequalities are also particularly suited to analyze the convergence of flows. Notably, Lojasiewicz [1982] bounded the path length of gradient flow trajectories and Polyak [1963] derived linear convergence of flows assuming PL. Related results for flows but under MB are claimed in [Helmke and Moore, 1996, Prop. 12.3]. More recently, Apidopoulos et al. [2021] considered the Heavy-Ball differential equation and deduced convergence guarantees from PL. Wojtowytsch [2021b] studied a continuous model for SGD and the impact of the noise on the trajectory.

Finally, we found only few convergence results based on MB in the optimization literature. Fehrman et al. [2020] derive capture theorems and asymptotic sub-linear convergence rates for gradient descent assuming that MB holds on the solution set. They also provide probabilistic bounds for stochastic variants. In order to solve systems of non-linear equations, Zeng [2021] proposes a Newton-type method that is robust to non-isolated solutions. This leads to local quadratic convergence assuming that the solution set satisfies an MB-type property. The algorithm requires the knowledge of the dimension of the solution set.

**Applications.** Non-isolated minima arise in all sorts of optimization problems. It is common for non-convex inverse problems to have continuous symmetries, hence non-isolated minima (see [Zhang et al., 2020]). In the context of deep learning, Cooper [2018] proved that the set of global minima of a sufficiently over-parameterized neural network is a smooth manifold. In the last decade, there has been a renewed interest in Lojasiewicz inequalities because they are compatible with these complicated non-convex landscapes. In particular, a whole line of research exploits them to understand deep learning problems specifically. As an example, Oymak and Soltanolkotabi [2019] employed PL to analyze the path taken by (stochastic) gradient descent in the vicinity of minimizers. Several other works suggested that non-convex machine learning loss landscapes can be understood in over-parameterized regimes through the lens of Lojasiewicz inequalities [Bassily et al., 2018, Belkin, 2021, Liu et al., 2022, Terjék and González-Sánchez, 2022]. Specifically, they argue that PL holds on a significant part of the search space and analyze (stochastic) gradient methods. Chatterjee [2022] also establishes local convergence results for a large class of neural networks with PL inequalities.

## 1.4 Notation and geometric preliminaries

This section anchors notation and some basic geometric facts. In the important case where  $\mathcal{M} = \mathbb{R}^n$ , several objects reduce as summarized in Table 2.

We let  $\langle \cdot, \cdot \rangle$  denote the inner product on  $T_x\mathcal{M}$ —it may depend on  $x \in \mathcal{M}$ , but the base point is always clear from context. The associated norm is  $\|v\| = \sqrt{\langle v, v \rangle}$ . The map  $\text{dist}: \mathcal{M} \times \mathcal{M} \rightarrow \mathbb{R}_+$  is the Riemannian distance on  $\mathcal{M}$ . We let  $B(x, \delta)$  denote the open ball of radius  $\delta$  around  $x \in \mathcal{M}$ . The tangent bundle is  $T\mathcal{M} = \{(x, v) : x \in \mathcal{M} \text{ and } v \in T_x\mathcal{M}\}$ .

Moving away from  $x \in \mathcal{M}$  along the geodesic with (sufficiently small) initial velocity  $v \in T_x\mathcal{M}$  for unit time produces the point  $\text{Exp}_x(v) \in \mathcal{M}$  (Riemannian exponential). The injectivity radius

at  $x$  is  $\text{inj}(x) > 0$ . It is defined such that, given  $y \in \text{B}(x, \text{inj}(x))$ , there exists a unique smallest vector  $v \in \text{T}_x \mathcal{M}$  for which  $\text{Exp}_x(v) = y$ . We denote this  $v$  by  $\text{Log}_x(y)$  (Riemannian logarithm). Additionally, given  $x \in \mathcal{M}$  and  $y \in \text{B}(x, \text{inj}(x))$ , we let  $\Gamma_x^y: \text{T}_x \mathcal{M} \rightarrow \text{T}_y \mathcal{M}$  denote parallel transport along the unique minimizing geodesic between  $x$  and  $y$ . If  $v = \text{Log}_x(y)$ , we also let  $\Gamma_v = \Gamma_x^y$ .

Let  $\mathfrak{X}$  be a subset of  $\mathcal{M}$ . We need the notions of tangent and normal cones to  $\mathfrak{X}$ , defined below.

**Definition 1.3.** *The tangent cone to a set  $\mathfrak{X}$  at  $x \in \mathfrak{X}$  is the closed set*

$$\text{T}_x \mathfrak{X} = \left\{ \lim_{k \rightarrow +\infty} \frac{1}{t_k} \text{Log}_x(x_k) \mid x_k \in \mathfrak{X}, t_k > 0 \text{ for all } k, x_k \rightarrow x, t_k \rightarrow 0 \right\}.$$

We also let  $\text{N}_x \mathfrak{X} = \{w \in \text{T}_x \mathcal{M} : \langle w, v \rangle \leq 0 \text{ for all } v \in \text{T}_x \mathfrak{X}\}$  denote the normal cone to  $\mathfrak{X}$  at  $x$ .

When  $\mathfrak{X}$  is a submanifold of  $\mathcal{M}$  around  $x$ , the cones  $\text{T}_x \mathfrak{X}$  and  $\text{N}_x \mathfrak{X}$  reduce to the tangent and normal spaces of  $\mathfrak{X}$  at  $x$ . (By ‘‘submanifold’’, we always mean *embedded* submanifold.)

Given  $x \in \mathcal{M}$ , we let  $\text{dist}(x, \mathfrak{X}) = \inf_{y \in \mathfrak{X}} \text{dist}(x, y)$  denote the distance of  $x$  to  $\mathfrak{X}$ . We further let  $\text{proj}_{\mathfrak{X}}(x)$  denote the set of solutions of the optimization problem  $\min_{y \in \mathfrak{X}} \text{dist}(x, y)$ . If this set is non-empty (which is the case in particular if  $\mathfrak{X}$  is closed), then we have:

$$\forall \bar{x} \in \mathfrak{X}, y \in \text{proj}_{\mathfrak{X}}(x), \quad \text{dist}(y, \bar{x}) \leq 2 \text{dist}(x, \bar{x}). \quad (3)$$

Indeed, the triangle inequality yields  $\text{dist}(y, \bar{x}) \leq \text{dist}(x, y) + \text{dist}(x, \bar{x})$ , and  $\text{dist}(x, y) = \text{dist}(x, \mathfrak{X}) \leq \text{dist}(x, \bar{x})$ . Moreover, if  $y \in \text{proj}_{\mathfrak{X}}(x)$  with  $\text{dist}(x, y) < \text{inj}(y)$  then  $\text{Log}_y(x) \in \text{N}_x \mathfrak{X}$ .

The set of local minima  $\mathcal{S}$  defined in (1) may not be closed: consider for example the function  $f(x) = \text{sgn}(x) \exp(-\frac{1}{x^2})(1 + \sin(\frac{1}{x^2}))$  with  $f_{\mathcal{S}} = 0$ . It follows that the projection onto  $\mathcal{S}$  may be empty. Notwithstanding, the following holds:

**Lemma 1.4.** *Around each  $\bar{x} \in \mathcal{S}$  there exists a neighborhood in which  $\text{proj}_{\mathcal{S}}$  is non-empty.*

*Proof.* Let  $\mathcal{U}$  be an open neighborhood of  $\bar{x}$  such that  $f(x) \geq f_{\mathcal{S}}$  for all  $x \in \mathcal{U}$ . Let  $\mathcal{V}_1, \mathcal{V}_2 \subset \mathcal{U}$  be two closed balls around  $\bar{x}$  of radii  $\delta > 0$  and  $\frac{1}{4}\delta$  respectively. Then  $\mathcal{S} \cap \mathcal{V}_1 = f^{-1}(f_{\mathcal{S}}) \cap \mathcal{V}_1$ , showing that  $\mathcal{S} \cap \mathcal{V}_1$  is closed and the projection onto this set is non-empty. Let  $x \in \mathcal{V}_2$  and  $y \in \text{proj}_{\mathcal{S} \cap \mathcal{V}_1}(x)$ . Then  $\text{dist}(x, y) \leq \frac{1}{4}\delta$ . Moreover, for all  $y' \in \mathcal{S} \setminus \mathcal{V}_1$  we have  $\text{dist}(x, y') \geq \frac{3}{4}\delta$ . It follows that  $\text{proj}_{\mathcal{S}}(x) = \text{proj}_{\mathcal{S} \cap \mathcal{V}_1}(x)$ , and this is non-empty.  $\square$

From these considerations we deduce that the projection onto  $\mathcal{S}$  is always locally well behaved.

**Lemma 1.5.** *Let  $\bar{x} \in \mathcal{S}$ . There exists a neighborhood  $\mathcal{U}$  of  $\bar{x}$  such that for all  $x \in \mathcal{U}$  the set  $\text{proj}_{\mathcal{S}}(x)$  is non-empty, and for all  $y \in \text{proj}_{\mathcal{S}}(x)$  we have  $\text{dist}(x, y) < \text{inj}(y)$ . In particular,  $v = \text{Log}_y(x)$  is well defined and  $v \in \text{N}_y \mathcal{S}$ .*

*Proof.* Let  $\mathcal{U}$  be a neighborhood of  $\bar{x}$  where  $\text{proj}_{\mathcal{S}}$  is non-empty (given by Lemma 1.4). Given  $\bar{\delta} < \text{inj}(\bar{x})$ , the ball  $\bar{\text{B}}(\bar{x}, \delta)$  is compact for all  $\delta < \bar{\delta}$ . Define  $h(\delta) = \inf_{x \in \bar{\text{B}}(\bar{x}, 2\delta)} \text{inj}(x)$  on the interval  $[0, \frac{\bar{\delta}}{2}]$ . The function  $h$  is continuous with  $h(0) = \text{inj}(\bar{x}) > 0$  so we can pick  $\delta > 0$  such that  $\delta \leq h(\delta)$ . Let  $x \in \text{B}(\bar{x}, \delta) \cap \mathcal{U}$  and  $y \in \text{proj}_{\mathcal{S}}(x)$ . By definition of the projection we have  $\text{dist}(x, y) \leq \text{dist}(x, \bar{x}) < \delta$ . Moreover, inequality (3) yields  $\text{dist}(y, \bar{x}) \leq 2 \text{dist}(x, \bar{x}) \leq 2\delta$  so  $h(\delta) \leq \text{inj}(y)$ . It follows that  $\text{dist}(x, y) < \delta \leq h(\delta) \leq \text{inj}(y)$  and  $v = \text{Log}_y(x)$  is well defined. The fact that  $v$  is in the normal cone follows from optimality conditions of projections.  $\square$

Given a self-adjoint linear map  $H$ , we let  $\lambda_i(H)$  denote the  $i$ th largest eigenvalue of  $H$ , and  $\lambda_{\min}(H)$  and  $\lambda_{\max}(H)$  denote the minimum and maximum eigenvalues respectively.

## 2 Four equivalent properties

In this section, we establish that MB, PL, EB and QG (see Definitions 1.1 and 1.2) are equivalent around a local minimum  $\bar{x}$  when  $f$  is  $\text{C}^2$ . Specifically, we show the implication graph in Figure 3.

It is well known that PL implies QG around minima: see references in Section 1.3. Perhaps the most popular argument relies on the bounded length of gradient flow trajectories under the more general Lojasiewicz inequality [Lojasiewicz, 1963, 1982, Absil et al., 2005, Bolte et al., 2010].






Figure 3: Implication graph when  $f$  is  $C^2$ .

**Definition 2.1.** Let  $\bar{x}$  be a local minimum of  $f$  with associated set  $\mathcal{S}$  (1). We say  $f$  satisfies the Lojasiewicz inequality with constants  $\theta \in [0, 1)$  and  $\mu > 0$  around  $\bar{x}$  if

$$|f(x) - f_{\mathcal{S}}|^{2\theta} \leq \frac{1}{2\mu} \|\nabla f(x)\|^2 \quad (\text{Loja})$$

for all  $x$  in some neighborhood of  $\bar{x}$ .

Notice that if  $f$  is Lojasiewicz with exponent  $\theta$  then it is Lojasiewicz with exponent  $\theta'$  for all  $\theta \leq \theta' < 1$  (though possibly in a different neighborhood). The case  $\theta = \frac{1}{2}$  is exactly the (PL) property.

**Proposition 2.2** (PL  $\Rightarrow$  QG). Suppose that  $f$  satisfies (Loja) around  $\bar{x} \in \mathcal{S}$ . Then  $f$  satisfies

$$f(x) - f_{\mathcal{S}} \geq ((1 - \theta)\sqrt{2\mu})^{\frac{1}{1-\theta}} \text{dist}(x, \mathcal{S})^{\frac{1}{1-\theta}}$$

for all  $x$  sufficiently close to  $\bar{x}$ . In particular, if  $\theta = \frac{1}{2}$ , this shows  $\mu$ -(PL)  $\Rightarrow$   $\mu$ -(QG).

We include a classical proof in Appendix A for completeness, with care regarding neighborhoods.

## 2.1 Two straightforward implications

In this section we show that MB  $\Rightarrow$  QG and EB  $\Rightarrow$  PL. These implications are known and direct. We give succinct proofs for completeness. The first one follows immediately from a Taylor expansion.

**Proposition 2.3** (MB  $\Rightarrow$  QG). Suppose that  $f$  is  $C^2$  and satisfies  $\mu$ -(MB) at  $\bar{x} \in \mathcal{S}$ . Then  $f$  satisfies  $\mu'$ -(QG) around  $\bar{x}$  for all  $\mu' < \mu$ .

*Proof.* Let  $\mathcal{U}$  be a neighborhood of  $\bar{x}$  as in Lemma 1.5. Let  $d$  be the codimension of  $\mathcal{S}$  (around  $\bar{x}$ ). Given  $\mu' < \mu$ , pick  $\varepsilon \in (0, \mu - \mu')$  and shrink  $\mathcal{U}$  so that for all  $x \in \mathcal{U}$  and  $y \in \text{proj}_{\mathcal{S}}(x)$  we have  $\lambda_d(\nabla^2 f(y)) \geq \mu' + \varepsilon$ . Given  $x \in \mathcal{U}$  and  $y \in \text{proj}_{\mathcal{S}}(x)$ , a Taylor expansion around  $y$  gives

$$f(x) - f_{\mathcal{S}} = \frac{1}{2} \langle v, \nabla^2 f(y)[v] \rangle + o(\|v\|^2) \geq \frac{\mu' + \varepsilon}{2} \text{dist}(x, \mathcal{S})^2 + o(\text{dist}(x, \mathcal{S})^2),$$

where  $v = \text{Log}_y(x)$  is normal to  $\mathcal{S}$ . We get the desired inequality for all  $x$  sufficiently close to  $\bar{x}$ .  $\square$

**Proposition 2.4** (EB  $\Rightarrow$  PL). Suppose that  $f$  is  $C^2$  and satisfies  $\mu$ -(EB) around  $\bar{x} \in \mathcal{S}$ . Then  $f$  satisfies  $\mu'$ -(PL) around  $\bar{x}$  for all  $\mu' < \mu$ .

*Proof.* Let  $\mathcal{U}$  be the intersection of two neighborhoods of  $\bar{x}$ : one where  $\mu$ -(EB) holds, and the other provided by Lemma 1.5. Given  $x \in \mathcal{U}$  and  $y \in \text{proj}_{\mathcal{S}}(x)$ , a Taylor expansion around  $y$  yields

$$f(x) - f_{\mathcal{S}} = \frac{1}{2} \langle v, \nabla^2 f(y)[v] \rangle + o(\|v\|^2) \quad \text{and} \quad \nabla f(x) = \Gamma_v \nabla^2 f(y)[v] + o(\|v\|),$$

where  $v = \text{Log}_y(x)$ . Using the Cauchy-Schwarz inequality and the triangle inequality, it follows that

$$f(x) - f_{\mathcal{S}} \leq \frac{1}{2} \|v\| \|\nabla^2 f(y)[v]\| + o(\|v\|^2) \leq \frac{1}{2} \|v\| \|\nabla f(x)\| + o(\|v\|^2).$$

Finally, EB gives that  $\|v\| \leq \frac{1}{\mu} \|\nabla f(x)\|$  so  $f(x) - f_{\mathcal{S}} \leq \frac{1}{2\mu} \|\nabla f(x)\|^2 + o(\|\nabla f(x)\|^2)$ . We get the desired inequality for all  $x$  sufficiently close to  $\bar{x}$ .  $\square$

**Remark 2.5.** Suppose that  $f$  is only  $C^1$  and  $\nabla f$  is locally  $L$ -Lipschitz continuous around  $\bar{x}$ . If  $\mu$ -EB holds around  $\bar{x}$  then  $f$  satisfies PL with constant  $\frac{\mu^2}{L}$  around  $\bar{x}$ , as shown by [Karimi et al. \[2016\]](#) (notice that this constant is not as sharp as the one in [Proposition 2.4](#)). To see this, let  $\mathcal{V}$  be a geodesic ball centered on  $\bar{x}$  and where  $\nabla f$  is  $L$ -Lipschitz continuous. Let  $\mathcal{U} \subseteq \mathcal{V}$  be a neighborhood of  $\bar{x}$  as in [Lemma 1.5](#) and shrink it so that for all  $x \in \mathcal{U}$  we have  $\text{proj}_{\mathcal{S}}(x) \subseteq \mathcal{V}$  (this is possible using the bound [\(3\)](#)). Given  $x \in \mathcal{U}$  and  $y \in \text{proj}_{\mathcal{S}}(x)$ , the local Lipschitz continuity of  $\nabla f$  implies that  $f(\text{Exp}_y(v)) - f_{\mathcal{S}} \leq \frac{L}{2}\|v\|^2$ , where  $v = \text{Log}_y(x)$  [[Boumal, 2023, Prop. 10.53](#)]. Written differently, we find that  $f(x) - f_{\mathcal{S}} \leq \frac{L}{2} \text{dist}(x, \mathcal{S})^2$  for all  $x \in \mathcal{U}$ . Combining with [\(EB\)](#) yields  $f(x) - f_{\mathcal{S}} \leq \frac{L}{2\mu^2} \|\nabla f(x)\|^2$ .

In the context of non-smooth optimization, [Li and Pong \[2018\]](#) also showed that EB implies PL under an additional level set separation assumption, though with no control on the PL constant.

## 2.2 Quadratic growth implies error bound

In this section, we show that QG implies EB for  $C^2$  functions. Other works proving this implication either assume that  $f$  is convex (see [[Karimi et al., 2016](#)] and [[Drusvyatskiy and Lewis, 2018, Cor. 3.6](#)]) or do not provide control on the constants (see [[Drusvyatskiy et al., 2013, Cor. 3.2](#)]). For this, we first characterize a distance growth rate when we move from  $\mathcal{S}$  in a normal direction (see [Definition 1.3](#)).

**Lemma 2.6.** Let  $\bar{x} \in \mathcal{S}$  and  $v \in N_{\bar{x}}\mathcal{S}$  unitary. Then  $\text{dist}(\text{Exp}_{\bar{x}}(tv), \mathcal{S}) = t + o(t)$  as  $t \rightarrow 0$ ,  $t \geq 0$ .

*Proof.* Let  $\mathcal{U}$  be a neighborhood of  $\bar{x}$  as in [Lemma 1.5](#). Shrink  $\mathcal{U}$  so that for all  $x \in \mathcal{U}$  and  $y \in \text{proj}_{\mathcal{S}}(x)$  we have  $\text{dist}(y, \bar{x}) < \text{inj}(\bar{x})$ . Given a small parameter  $t > 0$ , define  $x(t) = \text{Exp}_{\bar{x}}(tv)$  and let  $y(t) \in \text{proj}_{\mathcal{S}}(x(t))$ . From [\(3\)](#) we have  $\text{dist}(y(t), \bar{x}) \leq 2t$  and it follows that  $y(t) \rightarrow \bar{x}$  as  $t \rightarrow 0$ . Define  $u(t) = \text{Log}_{y(t)}(x(t))$  and  $w(t) = \text{Log}_{\bar{x}}(y(t))$ . Then

$$\text{dist}(x(t), \mathcal{S})^2 = \|u(t)\|^2 = \|tv - w(t) + \Gamma_{y(t)}^{\bar{x}} u(t) - tv + w(t)\|^2 = \|tv - w(t)\|^2 + o(t^2)$$

as  $t \rightarrow 0$  because  $\|\Gamma_{y(t)}^{\bar{x}} u(t) - tv + w(t)\| = o(t)$  as  $t \rightarrow 0$  [[Sun et al., 2019, Eq. \(6\)](#)]. In particular, for all  $t$  sufficiently small we have

$$\text{dist}(x(t), \mathcal{S})^2 = t^2 - 2t\langle w(t), v \rangle + \|w(t)\|^2 + o(t^2) \geq t^2 - 2t\langle w(t), v \rangle + o(t^2).$$

Let  $I \subseteq \mathbb{R}_{>0}$  be the times where  $y(t) \neq \bar{x}$ . If  $\inf I > 0$  then the final claim holds because  $y(t) = \bar{x}$  for small enough  $t$ . Suppose now that  $\inf I = 0$ . Define  $r(t) = \frac{w(t)}{\|w(t)\|}$  on  $I$ , and let  $\mathcal{A}$  be the set of accumulation points of  $r(t)$  as  $t \rightarrow 0$ ,  $t \in I$ . Then  $\mathcal{A}$  is included in the unit sphere and  $\mathcal{A} \subseteq T_{\bar{x}}\mathcal{S}$  by definition. Given  $a \in \mathcal{A}$  and  $t \in I$ , we can use  $\langle a, v \rangle \leq 0$  to find that

$$\text{dist}(x(t), \mathcal{S})^2 \geq t^2 - 2t\|w(t)\|\langle r(t) - a + a, v \rangle + o(t^2) \geq t^2 - 4t^2\|r(t) - a\| + o(t^2).$$

It follows that  $\text{dist}(x(t), \mathcal{S})^2 \geq t^2 - 4t^2 \text{dist}(r(t), \mathcal{A}) + o(t^2) = t^2 + o(t^2)$  because  $\text{dist}(r(t), \mathcal{A}) \rightarrow 0$  as  $t \rightarrow 0$ .  $\square$

Using this rate, we now find that QG implies the following bounds for  $\nabla^2 f$  in the normal cones of  $\mathcal{S}$ . Note that the following proposition does not yet show QG  $\Rightarrow$  MB, because to establish MB we still need to argue that  $\mathcal{S}$  is a smooth set.

**Proposition 2.7.** Suppose  $f$  is  $C^2$  and satisfies [\(QG\)](#) with constant  $\mu$  around  $\bar{x} \in \mathcal{S}$ . Then for all  $y \in \mathcal{S}$  sufficiently close to  $\bar{x}$  and all  $v \in N_y\mathcal{S}$  we have

$$\langle v, \nabla^2 f(y)[v] \rangle \geq \mu\|v\|^2 \quad \text{and} \quad \|\nabla^2 f(y)[v]\| \geq \mu\|v\|.$$

*Proof.* Let  $\mathcal{U}$  be an open neighborhood of  $\bar{x}$  where  $\mu$ -QG holds. Let  $y \in \mathcal{S} \cap \mathcal{U}$  and  $v \in N_y\mathcal{S}$ . For all small enough  $t > 0$ , we obtain from a Taylor expansion that

$$f(\text{Exp}_y(tv)) - f_{\mathcal{S}} = \frac{t^2}{2} \langle v, \nabla^2 f(y)[v] \rangle + o(t^2) \geq \frac{\mu}{2} \text{dist}(\text{Exp}_y(tv), \mathcal{S})^2 = \frac{\mu}{2} t^2 \|v\|^2 + o(t^2),$$

where the inequality comes from QG and the following equality comes from [Lemma 2.6](#). Take  $t \rightarrow 0$  to get the first inequality. The other inequality follows by Cauchy–Schwarz.  $\square$

We deduce from this that, under QG, the gradient norm is locally bounded from below and from above by the distance to  $\mathcal{S}$ , up to some constant factors. This notably secures EB.

**Proposition 2.8** (QG  $\Rightarrow$  EB). *Suppose  $f$  is  $C^2$  and satisfies  $\mu$ -(QG) around  $\bar{x} \in \mathcal{S}$ . For all  $\mu^b < \mu$  and  $\lambda^\sharp > \lambda_{\max}(\nabla^2 f(\bar{x}))$  there exists a neighborhood  $\mathcal{U}$  of  $\bar{x}$  such that for all  $x \in \mathcal{U}$  we have*

$$\mu^b \operatorname{dist}(x, \mathcal{S}) \leq \|\nabla f(x)\| \leq \lambda^\sharp \operatorname{dist}(x, \mathcal{S}).$$

*Proof.* Let  $\mathcal{U}$  be a neighborhood of  $\bar{x}$  as in Lemma 1.5. Shrink  $\mathcal{U}$  so that for all  $x \in \mathcal{U}$  and  $y \in \operatorname{proj}_{\mathcal{S}}(x)$  the inequalities of Proposition 2.7 hold. Now let  $x \in \mathcal{U}$  and  $y \in \operatorname{proj}_{\mathcal{S}}(x)$ . Define  $v = \operatorname{Log}_y(x)$  and  $\gamma(t) = \operatorname{Exp}_y(tv)$ , so that  $y = \gamma(0)$  and  $x = \gamma(1)$ . Then a Taylor expansion of  $\nabla f$  around  $y$  yields

$$\Gamma_v^{-1} \nabla f(x) = \nabla^2 f(y)[v] + r(x) \quad \text{where} \quad r(x) = \int_0^1 (\Gamma_{\tau v}^{-1} \circ \nabla^2 f(\gamma(\tau)) \circ \Gamma_{\tau v} - \nabla^2 f(y))[v] d\tau.$$

The Hessian is continuous so  $r(x) = o(\|v\|)$  as  $x \rightarrow \bar{x}$ . Moreover, Proposition 2.7 provides that  $\|\nabla^2 f(y)[v]\| \geq \mu\|v\|$  so using the triangle inequality and the reverse triangle inequality we get

$$\|v\|(\mu - o(1)) \leq \|\nabla f(x)\| \leq \|v\|(\lambda_{\max}(\nabla^2 f(y)) + o(1))$$

as  $x \rightarrow \bar{x}$ . We get the result if we choose  $x$  close enough to  $\bar{x}$ .  $\square$

The first inequality in the previous proposition is exactly EB. In Proposition 2.4 we showed that EB implies PL when  $f$  is  $C^2$ . Thus, it also holds that QG implies PL.

**Remark 2.9.** *When  $f$  is only  $C^1$  it is not true that QG implies EB or PL. To see this, consider the function  $f(x) = 2x^2 + x^2 \sin(1/\sqrt{|x|})$  (see Figure 2). It is  $C^1$  and satisfies QG around the minimum  $\bar{x} = 0$  because  $f(x) \geq x^2$ . However, there are other local minima arbitrarily close to zero. Those are critical points with function value strictly larger than  $f(\bar{x})$ , which disqualifies EB and PL around  $\bar{x} = 0$ .*

**Remark 2.10.** *Combining Propositions 2.2 and 2.8 ( $\mu$ -PL  $\Rightarrow$   $\mu$ -QG and  $\mu$ -QG  $\Rightarrow$   $\mu'$ -EB), we find that  $\mu$ -PL implies  $\mu'$ -EB for  $C^2$  functions. In fact, Karimi et al. [2016] show that  $\mu$ -PL implies  $\mu$ -EB for  $C^1$  functions, globally so if PL holds globally. Indeed, for  $x$  sufficiently close to  $\bar{x}$ , we have*

$$\frac{\mu}{2} \operatorname{dist}(x, \mathcal{S})^2 \leq f(x) - f_{\mathcal{S}} \leq \frac{1}{2\mu} \|\nabla f(x)\|^2,$$

where the second inequality is PL, and the first inequality is QG (implied by PL).

Combining all previous implications, we obtain that MB implies PL.

**Corollary 2.11** (MB  $\Rightarrow$  PL). *Suppose that  $f$  is  $C^2$  and satisfies  $\mu$ -(MB) at  $\bar{x} \in \mathcal{S}$ . Let  $0 < \mu' < \mu$ . Then  $f$  satisfies  $\mu'$ -(PL) around  $\bar{x}$ .*

As a consequence, if  $f$  is  $\mu$ -MB on  $\mathcal{S}$  then for all  $\mu' < \mu$  there exists a neighborhood of  $\mathcal{S}$  where  $f$  is  $\mu'$ -PL (and this does not require  $\mathcal{S}$  to be compact). When  $f$  is  $C^3$  the size of the neighborhood where PL holds can be controlled (to some extent) with the third derivative. A version of this corollary appears in [Wojtowytsch, 2021a, Ex. 2.9] with a different trade-off between control of the neighborhood and the constant  $\mu'$ . Feehan [2020, Thm. 6] shows a similar result in Banach spaces assuming that the function is  $C^3$ .

### 2.3 PL implies a smooth solution set and MB

The MB property is particularly strong because it presupposes a smooth solution set and it implies PL, EB, and QG. It raises a natural question: are these properties also enforcing some structure to the solution set? In this section we show that the answer is yes for  $C^2$  functions: if PL (or EB or QG) holds around  $\bar{x} \in \mathcal{S}$  then  $\mathcal{S}$  must be a submanifold around  $\bar{x}$ . To get a sense of why this is true, suppose that  $\mathcal{M} = \mathbb{R}^2$  and that around  $\bar{x}$  the solution set  $\mathcal{S}$  is the union of two orthogonal lines (a

cross) that intersect at  $\bar{x}$ . Then it must be that  $\nabla^2 f(\bar{x}) = 0$  because the gradient is zero along both lines. However, if we assume PL then the spectrum of  $\nabla^2 f(x)$  must contain at least one positive eigenvalue bigger than  $\mu$  for all points  $x \in \mathcal{S} \setminus \{\bar{x}\}$  close to  $\bar{x}$ , owing to the quadratic growth property. We obtain a contradiction because the eigenvalues of  $\nabla^2 f$  are continuous. To make this intuition precise, we first show that the PL inequality induces a lower-bound on the positive eigenvalues of  $\nabla^2 f$ .

**Proposition 2.12.** *Suppose  $f$  is  $C^2$  and  $\mu$ -**(PL)** around  $\bar{x} \in \mathcal{S}$ . If  $\lambda$  is a non-zero eigenvalue of  $\nabla^2 f(\bar{x})$  then  $\lambda \geq \mu$ .*

*Proof.* Let  $\lambda > 0$  be an eigenvalue of  $\nabla^2 f(\bar{x})$  with associated unit eigenvector  $v$ . Then

$$f(\text{Exp}_{\bar{x}}(tv)) - f_{\mathcal{S}} = \frac{\lambda}{2}t^2 + o(t^2) \quad \text{and} \quad \nabla f(\text{Exp}_{\bar{x}}(tv)) = \lambda t \Gamma_{tv} v + o(t).$$

The PL inequality implies  $\frac{\lambda}{2}t^2 + o(t^2) \leq \frac{1}{2\mu}\lambda^2 t^2 + o(t^2)$ , which gives the result as  $t \rightarrow 0$ .  $\square$

The latter argument is inconclusive when  $\lambda = 0$ . Still, we do get control of the Hessian's rank.

**Corollary 2.13.** *Suppose  $f$  is  $C^2$  and  $\mu$ -**(PL)** around  $\bar{x} \in \mathcal{S}$ . Then  $\text{rank}(\nabla^2 f(x)) = \text{rank}(\nabla^2 f(\bar{x}))$  for all  $x \in \mathcal{S}$  close enough to  $\bar{x}$ .*

*Proof.* Since  $f$  is  $C^2$  the eigenvalues of  $\nabla^2 f$  are continuous and the map  $x \mapsto \text{rank}(\nabla^2 f(x))$  is lower semi-continuous, that is, if  $x \in \mathcal{M}$  is close enough to  $\bar{x}$  then  $\text{rank} \nabla^2 f(x) \geq d$ , where  $d = \text{rank} \nabla^2 f(\bar{x})$ . Furthermore, if  $y \in \mathcal{S}$  is sufficiently close to  $\bar{x}$  then  $\lambda_{d+1}(\nabla^2 f(y)) < \mu$  by continuity of eigenvalues, and Proposition 2.12 then implies  $\lambda_{d+1}(\nabla^2 f(y)) = 0$ .  $\square$

This allows us to show that  $\nabla f$  aligns locally in a particular way with the eigenspaces of  $\nabla^2 f$ .

**Lemma 2.14.** *Suppose  $f$  is  $C^2$  and satisfies **(PL)** around  $\bar{x} \in \mathcal{S}$ . Let  $d = \text{rank}(\nabla^2 f(\bar{x}))$ . Then the orthogonal projector  $P(x)$  onto the top  $d$  eigenspace of  $\nabla^2 f(x)$  is well defined when  $x$  is sufficiently close to  $\bar{x}$ , and (with  $I$  denoting identity)*

$$\|(I - P(x))\nabla f(x)\| = o(\text{dist}(x, \mathcal{S})) = o(\|\nabla f(x)\|)$$

as  $x \rightarrow \bar{x}$ . Additionally, if  $\nabla^2 f$  is locally Lipschitz continuous around  $\bar{x}$  then  $\|(I - P(x))\nabla f(x)\| = O(\text{dist}(x, \mathcal{S})^2) = O(\|\nabla f(x)\|^2)$  as  $x \rightarrow \bar{x}$ .

*Proof.* Given a point  $x \in \mathcal{M}$ , let  $P(x): \mathbb{T}_x \mathcal{M} \rightarrow \mathbb{T}_x \mathcal{M}$  denote the orthogonal projector onto the top  $d$  eigenspace of  $\nabla^2 f(x)$ . This is well defined provided  $\lambda_d(\nabla^2 f(x)) > \lambda_{d+1}(\nabla^2 f(x))$ . Let  $\mathcal{U}$  be a neighborhood of  $\bar{x}$  as in Lemma 1.5. By continuity of the eigenvalues of  $\nabla^2 f$ , we can shrink  $\mathcal{U}$  so that for all  $x \in \mathcal{U}$  and  $y \in \text{proj}_{\mathcal{S}}(x)$  the projectors  $P(x)$  and  $P(y)$  are well defined. Given  $x \in \mathcal{U}$ , we let  $y \in \text{proj}_{\mathcal{S}}(x)$  and  $v = \text{Log}_y(x)$ . Now define  $\gamma(t) = \text{Exp}_y(tv)$ . A Taylor expansion of the gradient around  $y$  gives that

$$\nabla f(x) = \Gamma_v(\nabla^2 f(y)[v] + r(x)) \quad \text{where} \quad r(x) = \int_0^1 (\Gamma_{\tau v}^{-1} \circ \nabla^2 f(\gamma(\tau)) \circ \Gamma_{\tau v} - \nabla^2 f(y))[v] d\tau.$$

By Corollary 2.13, the rank of  $\nabla^2 f$  is locally constant on  $\mathcal{S}$  (equal to  $d$ ) so  $\nabla^2 f(y) = P(y)\nabla^2 f(y)$  whenever  $x$  is sufficiently close to  $\bar{x}$  (using the bound  $\text{dist}(y, \bar{x}) \leq 2\text{dist}(x, \bar{x})$  from (3)). It follows that

$$(I - P(x))\nabla f(x) = (I - P(x))\Gamma_v(P(y)\nabla^2 f(y)[v] + r(x)).$$

Notice that  $\Gamma_v \rightarrow I$  as  $x \rightarrow \bar{x}$  and  $P(y) \rightarrow P(\bar{x})$  as  $x \rightarrow \bar{x}$  so  $(I - P(x))\Gamma_v P(y) \rightarrow 0$  as  $x \rightarrow \bar{x}$ . It follows that  $(I - P(x))\nabla f(x) = o(\|v\|)$  as  $x \rightarrow \bar{x}$ . The claim follows by noting that  $\|v\| = \text{dist}(x, \mathcal{S})$  and the fact that  $\text{dist}(x, \mathcal{S})$  is commensurate  $\|\nabla f(x)\|$  (as shown in Proposition 2.8).

Suppose now that  $\nabla^2 f$  is locally Lipschitz continuous around  $\bar{x}$ . Then  $P$  is also locally Lipschitz continuous around  $\bar{x}$  [Vannieuwenhoven, 2022, Cor. 1] and  $(I - P(x))\Gamma_v P(y) = (I - P(x))\Gamma_v(P(y) - \Gamma_v^{-1}P(x)) = O(\|v\|)$ . Moreover, we have  $r(x) = O(\|v\|^2)$  so it follows that  $(I - P(x))\nabla f(x) = O(\|v\|^2)$  as  $x \rightarrow \bar{x}$ . We conclude again with Proposition 2.8.  $\square$

The lemma below exhibits a submanifold  $\mathcal{Z}$  that contains  $\bar{x}$ . It need *not* coincide with the solution set  $\mathcal{S}$ . However, they *do* coincide if we assume that PL holds. This is the main argument to show that  $\mathcal{S}$  is locally a submanifold whenever  $f$  is PL and sufficiently regular.

**Lemma 2.15.** *Suppose  $f$  is  $C^p$  with  $p \geq 2$ . Let  $\bar{x} \in \mathcal{S}$  and define  $\mathcal{U} = \{x \in \mathcal{M} : \text{dist}(x, \bar{x}) < \text{inj}(\bar{x})\}$ . Let  $P(\bar{x})$  denote the orthogonal projector onto the image of  $\nabla^2 f(\bar{x})$ . Then the set*

$$\mathcal{Z} = \{x \in \mathcal{U} : P(\bar{x})\Gamma_{\bar{x}}^{\bar{x}}\nabla f(x) = 0\}$$

*is a  $C^{p-1}$  embedded submanifold locally around  $\bar{x}$ . If  $f$  is analytic then  $\mathcal{Z}$  is also analytic.*

*Proof.* We build a local defining function for  $\mathcal{Z}$ . Let  $d$  be the rank of  $\nabla^2 f(\bar{x})$  and  $u_1, \dots, u_d$  be a set of orthonormal eigenvectors of  $\nabla^2 f(\bar{x})$  with associated eigenvalues  $\lambda_1, \dots, \lambda_d > 0$ . We define  $h: \mathcal{U} \rightarrow \mathbb{R}^d$  as  $h_i(x) = \langle u_i, \Gamma_{\bar{x}}^{\bar{x}}\nabla f(x) \rangle$ . Clearly,  $h(x) = 0$  if and only if  $x \in \mathcal{Z}$ . The function  $h$  is  $C^{p-1}$  if  $f$  is  $C^p$ , and analytic if  $f$  is analytic. For all  $\dot{x} \in T_{\bar{x}}\mathcal{M}$  we have

$$Dh_i(\bar{x})[\dot{x}] = \langle u_i, \nabla^2 f(\bar{x})[\dot{x}] \rangle = \langle \nabla^2 f(\bar{x})[u_i], \dot{x} \rangle = \lambda_i \langle u_i, \dot{x} \rangle.$$

It follows that  $Dh(\bar{x})$  has full rank. Thus,  $\mathcal{Z}$  is a submanifold around  $\bar{x}$  with the stated regularity.  $\square$

A result similar to Lemma 2.15 is presented in [Chill, 2006, Lem. 1] for Banach spaces. We are now ready for one of our main theorems, regarding the regularity of the set of local minimizers  $\mathcal{S}$  (1).

**Theorem 2.16.** *Suppose  $f$  is  $C^p$  with  $p \geq 2$  and satisfies (PL) around  $\bar{x} \in \mathcal{S}$ . Then  $\mathcal{S}$  is a  $C^{p-1}$  submanifold of  $\mathcal{M}$  locally around  $\bar{x}$ . If  $f$  is analytic then  $\mathcal{S}$  is also analytic.*

*Proof.* We let  $d$  denote the rank of  $\nabla^2 f(\bar{x})$ . By Corollary 2.13,  $\text{rank } \nabla^2 f(y) = d$  for all  $y \in \mathcal{S}$  sufficiently close to  $\bar{x}$ . We let  $\mathcal{U} \subseteq B(\bar{x}, \text{inj}(\bar{x}))$  be a neighborhood of  $\bar{x}$  such that for all  $x \in \mathcal{U}$  the orthogonal projector  $P(x): T_x\mathcal{M} \rightarrow T_x\mathcal{M}$  onto the top  $d$  eigenspace of  $\nabla^2 f(x)$  is well defined (it exists because eigenvalues of  $\nabla^2 f$  are continuous). Lemma 2.15 ensures that

$$\mathcal{Z} = \{x \in \mathcal{U} : P(\bar{x})\Gamma_{\bar{x}}^{\bar{x}}\nabla f(x) = 0\}$$

is a submanifold around  $\bar{x}$ . Clearly,  $\mathcal{S} \cap \mathcal{U} \subseteq \mathcal{Z}$  holds. We now show the other inclusion to obtain that  $\mathcal{S}$  and  $\mathcal{Z}$  coincide around  $\bar{x}$ . From Lemma 2.14 we have  $\|\nabla f(x)\| \leq \|P(x)\nabla f(x)\| + o(\|\nabla f(x)\|)$  as  $x \rightarrow \bar{x}$ . Moreover, the triangle inequality gives  $\|P(x)\nabla f(x)\| \leq \|(P(x) - \Gamma_{\bar{x}}^{\bar{x}}P(\bar{x})\Gamma_{\bar{x}}^{\bar{x}})\nabla f(x)\| + \|P(\bar{x})\Gamma_{\bar{x}}^{\bar{x}}\nabla f(x)\|$ . By continuity of  $P$  we have  $P(x) - \Gamma_{\bar{x}}^{\bar{x}}P(\bar{x})\Gamma_{\bar{x}}^{\bar{x}} = o(1)$  as  $x \rightarrow \bar{x}$  so it follows that

$$\|\nabla f(x)\| \leq \|P(\bar{x})\Gamma_{\bar{x}}^{\bar{x}}\nabla f(x)\| + o(\|\nabla f(x)\|)$$

as  $x \rightarrow \bar{x}$ . We conclude that  $P(\bar{x})\Gamma_{\bar{x}}^{\bar{x}}\nabla f(x) = 0$  implies  $\nabla f(x) = 0$  for all  $x$  sufficiently close to  $\bar{x}$ . This confirms that  $\mathcal{Z} \subseteq \mathcal{S} \cap \mathcal{U}$  around  $\bar{x}$  because all critical points near  $\bar{x}$  are in  $\mathcal{S}$  by PL.  $\square$

The codimension of  $\mathcal{S}$  is equal to the rank of  $\nabla^2 f$  on  $\mathcal{S}$ , as expected. A similar result holds for Banach spaces when the function is assumed analytic [Feehan, 2020, Thm. 1]. Around  $\bar{x}$ , the set of all minima of  $f$  and  $\mathcal{S}$  coincide when PL holds. Hence Theorem 2.16 implies that the set of minima of  $f$  is a submanifold around  $\bar{x}$ . Using the quadratic growth property we now deduce that PL implies MB.

**Corollary 2.17.** *Suppose  $f$  is  $C^2$  and  $\mu$ -(PL) around  $\bar{x} \in \mathcal{S}$ . Then  $f$  satisfies  $\mu$ -(MB) at  $\bar{x}$ .*

*Proof.* We apply Theorem 2.16 to get that  $\mathcal{S}$  is locally a  $C^1$  submanifold around  $\bar{x}$ . Then we apply Proposition 2.2 to get the (QG) property around  $\bar{x}$ . We finally apply Proposition 2.7 to normal eigenvectors of  $\nabla^2 f(\bar{x})$ . This yields that the normal eigenvalues are at least  $\mu$ .  $\square$

Note that this result also holds if we assume EB or QG instead of PL since these conditions are equivalent when  $f$  is  $C^2$ .

**Remark 2.18** (Connections to the distance function). *Given a closed set  $\mathfrak{X} \subseteq \mathcal{M}$ , the function  $f(x) = \frac{1}{2} \text{dist}_{\mathfrak{X}}(x)^2$  clearly satisfies the quadratic growth property. If  $f$  is  $C^p$  with  $p \geq 2$  in a neighborhood of  $\mathfrak{X}$  then Theorem 2.16 applies, telling us that  $\mathfrak{X}$  is a  $C^{p-1}$  submanifold. This question is of independent interest, see for example [Bellettini et al., 2007] for a proof when assuming  $p \geq 3$ .*

**Remark 2.19** (Structure when  $f$  is only  $C^1$ ). *To apply Theorem 2.16 we need  $f$  to be at least  $C^2$ . And indeed, if  $f$  is only  $C^1$  the solution set may not be a submanifold. As an example, the function  $f(x, y) = \frac{x^2 y^2}{x^2 + y^2}$  (see Figure 2) is  $C^1$  and locally PL around the origin, yet its set of minima is a cross.<sup>2</sup> More generally, let  $\mathfrak{X} \subseteq \mathcal{M}$  be a closed set and suppose that the distance function  $\text{dist}_{\mathfrak{X}}$  is  $C^1$  around  $\mathfrak{X}$  (such a set is called proximally smooth [Clarke et al., 1995]). We define  $f(x) = \frac{1}{2} \text{dist}_{\mathfrak{X}}(x)^2$  and find that  $\nabla f(x) = x - \text{proj}_{\mathfrak{X}}(x)$ , meaning that  $f$  is PL around  $\mathfrak{X}$  with constant  $\mu = 1$ . This holds in particular for all closed convex sets, yet many such sets fail to be  $C^0$  submanifolds (e.g., consider a closed square in  $\mathcal{M} = \mathbb{R}^2$ ). This provides further examples of  $C^1$  functions satisfying PL yet for which the solution set is not  $C^0$ .*

**Remark 2.20** (Restricted secant inequality). *It is possible to show equivalences with even more properties. For example, as in [Zhang and Yin, 2013], we say  $f$  satisfies the restricted secant inequality (RSI) with constant  $\mu$  around  $\bar{x} \in \mathcal{S}$  if  $\langle \nabla f(x), v \rangle \geq \mu \text{dist}(x, \mathcal{S})^2$  for all  $x$  in a neighborhood of  $\bar{x}$ , where  $v = \text{Log}_y(x)$  and  $y \in \text{proj}_{\mathcal{S}}(x)$ . From simple Taylor expansion arguments, we find that  $\mu$ -MB implies  $\mu'$ -RSI for all  $\mu' < \mu$ . By Cauchy–Schwarz, we also find that  $\mu$ -RSI implies  $\mu$ -EB for  $C^1$  functions (see [Karimi et al., 2016]). It follows that for  $C^2$  functions RSI is also equivalent to the four properties that we consider.*

**Remark 2.21** (Other Lojasiewicz exponents). *The PL inequality is exactly (Loja) with exponent  $\theta = \frac{1}{2}$ . We comment here about other values of  $\theta$ . First, suppose  $f$  is  $C^1$  and  $\nabla f$  locally Lipschitz continuous. If  $f$  is non-constant around  $\bar{x} \in \mathcal{S}$  then it cannot satisfy (Loja) with an exponent  $\theta < \frac{1}{2}$  around  $\bar{x}$ . This is because these assumptions are incompatible with the growth property from Proposition 2.2. See also [Abbaszadehpeivasti et al., 2022, Thm. 4] for an algorithmic perspective on this. Now suppose  $f$  is  $C^2$  and satisfies (Loja) with exponent  $\theta$  around  $\bar{x} \in \mathcal{S}$ . If  $\nabla^2 f$  is Lipschitz continuous around  $\bar{x}$  and  $\theta \in [\frac{1}{2}, \frac{2}{3})$  then PL holds around  $\bar{x}$ . Furthermore, if  $f$  is  $C^3$  and  $\theta \in [\frac{1}{2}, \frac{2}{3}]$  then PL holds around  $\bar{x}$ . We include a proof of these observations in Appendix B.*

### 3 Stability of minima and linear rates

In this section, we consider two types of algorithmic questions: the stability of minima and local convergence rates. We review some classical arguments, taking this opportunity to generalize and adapt some of them to accommodate non-isolated minima. This will serve us well to analyze algorithms in Section 4.

#### 3.1 Capture result for non-isolated minima

Typically, global convergence analyses of optimization algorithms merely guarantee that iterates accumulate only at critical points. The set of accumulation points may be empty (when the iterates diverge). Worse, it may even be infinite when minima are not isolated. See [Absil et al., 2005, §3.2.1] and [Bolte and Pauwels, 2021, §5.3] for examples of pathological functions for which reasonable algorithms (such as gradient descent) produce iterates with continuous sets of accumulation points. The latter issue cannot occur when minima are isolated.

What kind of stability results still hold when minima are not isolated? Consider an algorithm generating iterates as  $x_{k+1} = F_k(x_k, \dots, x_0)$ , where  $F_k$  is a *descent mapping*: it satisfies  $f(F_k(x_k, \dots, x_0)) \leq f(x_k)$  for all  $x_k, \dots, x_0 \in \mathcal{M}$ . Many deterministic algorithms fall in this category, including gradient descent and trust-region methods under suitable hypotheses. The standard capture theorem asserts that if the iterates generated by such descent mappings get sufficiently close

<sup>2</sup>We thank Christopher Criscitiello who found this function.

to an *isolated* local minimum then the sequence eventually converges to it (under a few weak assumptions) [Bertsekas, 1995, Prop. 1.2.5], [Absil et al., 2008, Thm. 4.4.2]. The result can be easily extended to a compact set of non-isolated local minima that satisfies several properties that we define now.

**Definition 3.1.** We say  $\mathfrak{X} \subseteq \mathcal{M}$  is isolated from critical points if there exists a neighborhood  $\mathcal{U}$  of  $\mathfrak{X}$  such that  $x \in \mathcal{U}$  and  $\nabla f(x) = 0$  imply  $x \in \mathfrak{X}$ .

Note that the points in  $\mathfrak{X}$  do *not* need to be isolated: the set  $\mathfrak{X}$  may be a continuum of critical points. It is clear that if  $\mathfrak{X} \subseteq \mathcal{S}$  (1) is isolated from critical points then there exists a neighborhood  $\mathcal{U}$  of  $\mathfrak{X}$  such that  $f(x) > f_{\mathcal{S}}$  for all  $x \in \mathcal{U} \setminus \mathfrak{X}$ , where  $f_{\mathcal{S}}$  is the value of  $f$  on  $\mathfrak{X}$ . The capture result below, based on [Absil et al., 2008], states that if the set of minima  $\mathfrak{X} \subseteq \mathcal{S}$  is both compact and isolated from critical points then it traps the iterates generated by all reasonable descent algorithms. A key hypothesis is that the steps have to be small around local minima: that is typically the case.

**Definition 3.2.** An algorithm which generates sequences on  $\mathcal{M}$  has the vanishing steps property on a set  $\mathfrak{X}$  if there exists a neighborhood  $\mathcal{U}$  of  $\mathfrak{X}$  and a continuous function  $\eta: \mathcal{M} \rightarrow \mathbb{R}_+$  with  $\eta(\mathfrak{X}) = 0$  such that, if  $x_k$  is an iterate in  $\mathcal{U}$ , then the next iterate  $x_{k+1}$  satisfies

$$\text{dist}(x_k, x_{k+1}) \leq \eta(x_k). \quad (\text{VS})$$

We say that the algorithm has the (VS) property at a point  $\bar{x} \in \mathcal{S}$  if it holds on the set  $\{\bar{x}\}$ .

**Proposition 3.3** (Capture of iterates). Let  $\mathfrak{X}$  be a compact subset of  $\mathcal{S}$  isolated from critical points. Consider an algorithm that produces iterates as  $x_{k+1} = F_k(x_k, \dots, x_0)$ , where  $F_k$  is a descent mapping. Assume that it satisfies the (VS) property on  $\mathfrak{X}$ . Also suppose that the sequences generated by this algorithm accumulate only at critical points of  $f$ . There exists a neighborhood  $\mathcal{U}$  of  $\mathfrak{X}$  such that if a sequence enters  $\mathcal{U}$  then all subsequent iterates are in  $\mathcal{U}$  and  $\text{dist}(x_k, \mathfrak{X}) \rightarrow 0$ .

*Proof.* There exists a compact neighborhood  $\mathcal{V}$  of  $\mathfrak{X}$  such that  $\mathcal{V} \setminus \mathfrak{X}$  does not contain any critical point and  $f(x) > f_{\mathcal{S}}$  for all  $x \in \mathcal{V} \setminus \mathfrak{X}$ . The (VS) property implies that there exists an open neighborhood  $\mathcal{W}$  of  $\mathfrak{X}$  included in  $\mathcal{V}$  such that for all  $k \in \mathbb{N}$ ,  $x_k \in \mathcal{W}$  and  $x_{k-1}, \dots, x_0 \in \mathcal{M}$  we have  $F_k(x_k, \dots, x_0) \in \mathcal{V}$ . The set  $\mathcal{V} \setminus \mathcal{W}$  is compact and we let  $\alpha > f_{\mathcal{S}}$  denote the minimum of  $f$  on this set. We define  $\mathcal{U} = \{x \in \mathcal{V} : f(x) < \alpha\}$ , which is included in  $\mathcal{W}$  by minimality of  $\alpha$ . Now let  $K \in \mathbb{N}$ ,  $x_K \in \mathcal{U}$  and  $x_{K-1}, \dots, x_0 \in \mathcal{M}$ . Then we have  $F_K(x_K, \dots, x_0) \in \mathcal{V}$  by definition of  $\mathcal{W}$ . Moreover,  $F_K$  is a descent mapping so  $f(F_K(x_K, \dots, x_0)) < \alpha$  and it implies that  $F_K(x_K, \dots, x_0) \in \mathcal{U}$ . It follows that  $x_{K+1}$  is in  $\mathcal{U}$  and all subsequent iterates are also in  $\mathcal{U}$ . Now we show that  $\{x_k\}$  converges to  $\mathfrak{X}$ . The sequence  $\{x_k\}$  eventually stays in a compact set (because  $x_k \in \mathcal{U} \subseteq \mathcal{V}$  for all  $k \geq K$ ) so it has a non-empty and compact set of accumulation points that we denote by  $\mathcal{A}$ . Then  $\mathcal{A} \subseteq \mathfrak{X}$  because  $\mathcal{A} \subseteq \mathcal{V}$  and the only critical points in  $\mathcal{V}$  are in  $\mathfrak{X}$ . The set of accumulation points of  $\{\text{dist}(x_k, \mathcal{A})\}$  is  $\{0\}$  (because  $\{x_k\}$  is bounded). So we deduce that  $\lim_{k \rightarrow +\infty} \text{dist}(x_k, \mathfrak{X}) \leq \lim_{k \rightarrow +\infty} \text{dist}(x_k, \mathcal{A}) = 0$ .  $\square$

This statement does *not* guarantee that the iterates converge to a specific point, but merely that  $\text{dist}(x_k, \mathfrak{X}) \rightarrow 0$ . Notice that we do not require any particular structure for  $\mathfrak{X}$  nor any form of function growth. The assumptions on the mappings  $F_k$  are also mild. However, we need  $\mathfrak{X}$  to be compact and this cannot be relaxed. Indeed, consider the function  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  defined by  $f(x, y) = \exp(xy)^2$ . Then  $\mathfrak{X} = \{(x, y) \in \mathbb{R}^2 : y = 0\}$  is a submanifold of global minima and it contains all the critical points of  $f$ . Consider the update rule  $(x_{k+1}, y_{k+1}) = (x_k - y_k^2, y_k)$ . It satisfies the descent condition because  $f(x_{k+1}, y_{k+1}) = \exp(-y_k^2)f(x_k, y_k)$  for all  $k$ . The distance assumption (VS) also holds because  $\text{dist}((x_{k+1}, y_{k+1}), (x_k, y_k)) = \text{dist}((x_k - y_k^2, y_k), (x_k, y_k)) = y_k^2$ . However, the sequence  $\{\text{dist}((x_k, y_k), \mathfrak{X})\}$  is constant (and not converging to zero) even if we initialize the algorithm arbitrarily close to  $\mathfrak{X}$  (but not exactly in  $\mathfrak{X}$ ).

The vanishing steps property is a reasonable assumption. When  $\mathcal{M} = \mathbb{R}^n$ , many optimization algorithms have an update rule of the form  $x_{k+1} = x_k + s_k$  and the vector  $s_k$  is small if  $x_k$  is close to a local minimum (e.g.,  $s_k = -\alpha_k \nabla f(x_k)$  with bounded  $\alpha_k$ ). For a general manifold  $\mathcal{M}$ , algorithms produce iterates as  $x_{k+1} = R_{x_k}(s_k)$ , where  $R: \text{T}\mathcal{M} \rightarrow \mathcal{M}: (x, s) \mapsto R_x(s)$  is a retraction [Boumal,

2023, Def. 3.47]. Specifically, for all  $(x, s) \in \text{T}\mathcal{M}$  the curve  $c(t) = \text{R}_x(ts)$  satisfies  $c(0) = x$  and  $c'(0) = s$ . To help ensure vanishing steps, we assume there is a constant  $c_r > 0$  such that

$$\text{dist}(\text{R}_x(s), x) \leq c_r \|s\| \tag{4}$$

for all  $(x, s) \in \text{T}\mathcal{M}$ . This clearly holds with  $c_r = 1$  when  $\text{R} = \text{Exp}$ .

### 3.2 Lyapunov stability and convergence to a single point

Pathological behavior such as a continuous set of accumulation points can be ruled out assuming Lojasiewicz inequalities. In this case, local minima are stable for a variety of algorithms, even without compactness hypothesis (in contrast to Proposition 3.3). This in turn ensures that the iterates converge to a single point of the solution set. In this section, we review arguments from [Polyak, 1963, Thm. 4], [Absil et al., 2005], [Attouch et al., 2013, Lem. 2.6], and [Bolte et al., 2017, Thm. 14]. The stability results in [Absil et al., 2005] and [Attouch et al., 2013] assume a sufficient decrease condition. Here, we abstract out the main arguments of their proofs and show stability results without assuming this decrease condition.

A central property to obtain convergence to a single point is a bound on the path length of the iterates. We make this precise in the following definition.

**Definition 3.4.** *An algorithm which generates sequences on  $\mathcal{M}$  has the bounded path length property on a set  $\mathfrak{X} \in \mathcal{M}$  if the following is true. There exist a neighborhood  $\mathcal{U}$  of  $\mathfrak{X}$  and a continuous function  $\gamma: \mathcal{M} \rightarrow \mathbb{R}_+$  with  $\gamma(\mathfrak{X}) = 0$  such that, if  $x_L, \dots, x_K \in \mathcal{M}$  are consecutive points generated by the algorithm and which are all in  $\mathcal{U}$ , then*

$$\sum_{k=L}^{K-1} \text{dist}(x_k, x_{k+1}) \leq \gamma(x_L). \tag{BPL}$$

We say that the algorithm has the (BPL) property at a point  $\bar{x} \in \mathcal{S}$  if it does so on the set  $\{\bar{x}\}$ .

Here we think of the algorithm as an optimization method with some fixed hyper-parameters. The definition is given for a generic set  $\mathfrak{X}$  but the bounded path length property is usually only satisfied around local minima. Combined with the vanishing steps property, bounded path length ensures stability of local minima.

**Proposition 3.5** (Lyapunov stability). *Suppose that an algorithm satisfies the (VS) and (BPL) properties at  $\bar{x} \in \mathcal{M}$ . Given a neighborhood  $\mathcal{U}$  of  $\bar{x}$ , there exists a neighborhood  $\mathcal{V}$  of  $\bar{x}$  such that if a sequence generated by this algorithm enters  $\mathcal{V}$  then all subsequent iterates stay in  $\mathcal{U}$ .*

*Proof.* The set  $\mathcal{U}$  contains an open ball of radius  $\delta_u > 0$  around  $\bar{x}$  in which the (BPL) and (VS) properties are satisfied with some functions  $\gamma$  and  $\eta$ . By continuity of  $\eta$  there exists an open ball  $\mathcal{W}$  centered on  $\bar{x}$  of radius  $\delta_w$  that satisfies  $\delta_w + \eta(x) < \delta_u$  for all  $x \in \mathcal{W}$ . Likewise, by continuity of  $\gamma$ , there exists an open ball  $\mathcal{V} \subset \mathcal{W}$  of radius  $\delta_v > 0$  around  $\bar{x}$  such that for all  $x \in \mathcal{V}$  we have  $\delta_v + \gamma(x) < \delta_w$ . Suppose that  $x_L \in \mathcal{V}$ . For contradiction, let  $K \geq L$  be the first index such that  $x_{K+1} \notin \text{B}(\bar{x}, \delta_u)$ . We deduce from the triangle inequality and the (BPL) property that

$$\text{dist}(x_K, \bar{x}) \leq \text{dist}(x_L, \bar{x}) + \sum_{k=L}^{K-1} \text{dist}(x_k, x_{k+1}) \leq \delta_v + \gamma(x_L) < \delta_w.$$

It follows that  $x_K \in \mathcal{W}$ . Using again the triangle inequality we find  $\text{dist}(x_{K+1}, \bar{x}) \leq \text{dist}(x_K, \bar{x}) + \text{dist}(x_K, x_{K+1}) \leq \delta_w + \eta(x_K) < \delta_u$ . This implies that  $x_{K+1}$  is in  $\text{B}(\bar{x}, \delta_u)$ : a contradiction.  $\square$

In particular, this excludes that the iterates diverge. We can also guarantee that accumulation points are actually limit points.

**Corollary 3.6.** *Suppose that an algorithm satisfies the (VS) and (BPL) properties at  $\bar{x} \in \mathcal{M}$ . If it generates a sequence that accumulates at  $\bar{x}$  then the sequence converges to  $\bar{x}$ .*



*Proof.* Let  $\mathcal{U}$  be a neighborhood of  $\bar{x}$ . From Proposition 3.5 there is a neighborhood  $\mathcal{V}$  such that if an iterate is in  $\mathcal{V}$  then all subsequent iterates are in  $\mathcal{U}$ . Since  $\bar{x}$  is an accumulation point we know that such an iterate exists. Repeat with a sequence of smaller and smaller neighborhoods of  $\bar{x}$ .  $\square$

Many optimization algorithms generate sequences that accumulate only at critical points. In that scenario, we can deduce that the sequence converges to a point, provided that it gets close enough to a set where (VS) and (BPL) hold.

**Corollary 3.7.** *Let  $\mathfrak{X}$  be a set of critical points and  $\bar{x} \in \mathfrak{X}$ . Let  $\mathcal{U}$  be a neighborhood of  $\bar{x}$  such that if  $x \in \mathcal{U}$  is a critical point then  $x \in \mathfrak{X}$ . Consider an algorithm that satisfies the (VS) and (BPL) properties on  $\mathfrak{X}$ . Also suppose that the sequences generated by this algorithm accumulate only at critical points of  $f$ . There exists a neighborhood  $\mathcal{V}$  of  $\bar{x}$  such that if a sequence enters  $\mathcal{V}$  then it converges to some  $x_\infty \in \mathcal{U} \cap \mathfrak{X}$ .*

*Proof.* The set  $\mathcal{U}$  contains a compact neighborhood  $\mathcal{B}$  of  $\bar{x}$  such that (i) if  $x \in \mathcal{B}$  is a critical point then  $x \in \mathfrak{X}$  and (ii) the (VS) and (BPL) properties hold on  $\mathcal{B} \cap \mathfrak{X}$ . From Proposition 3.5 there exists a neighborhood  $\mathcal{V}$  of  $\bar{x}$  such that if  $\{x_k\}$  enters  $\mathcal{V}$  then it stays in  $\mathcal{B}$ . The set  $\mathcal{B}$  is compact so  $\{x_k\}$  has at least one accumulation point  $x_\infty \in \mathcal{B}$ . Since accumulation points of  $\{x_k\}$  are critical we have  $x_\infty \in \mathfrak{X}$ . We conclude with Corollary 3.6.  $\square$

The conclusions of Corollary 3.7 are similar to the ones in [Absil et al., 2005, Prop 3.3] and [Attouch et al., 2013, Thm. 2.10].<sup>3</sup>

We describe below the argument that Absil et al. [2005] used to show that many gradient descent algorithms satisfy (BPL) around points where  $f$  is Lojasiewicz. We say that the sequence  $\{x_k\}$  satisfies the *strong decrease property* around  $\bar{x} \in \mathcal{M}$  if there exists  $\sigma > 0$  such that

$$f(x_k) - f(x_{k+1}) \geq \sigma \|\nabla f(x_k)\| \text{dist}(x_k, x_{k+1}) \quad \text{and} \quad x_k \in \mathcal{S} \Rightarrow x_{k+1} = x_k \quad (5)$$

whenever  $x_k$  is sufficiently close to  $\bar{x}$ , as introduced by Absil et al. [2005].

**Lemma 3.8.** *Suppose that  $f$  satisfies (Loja) around  $\bar{x} \in \mathcal{S}$  with constants  $\theta$  and  $\mu$ . If an algorithm generates sequences  $\{x_k\}$  that satisfy (5) around  $\bar{x}$  then it satisfies the (BPL) property at  $\bar{x}$  with*

$$\gamma(x) = \frac{1}{\sigma(1-\theta)\sqrt{2\mu}} |f(x) - f_{\mathcal{S}}|^{1-\theta}.$$

We include a proof of this statement in Appendix C for completeness. In fact, the algorithm would still satisfy the (BPL) property under the more general Kurdyka–Lojasiewicz assumption (see [Absil et al., 2005, §3.2.3]). In practice, many first-order algorithms (including gradient descent with constant step-sizes or with line-search) generate sequences with the strong decrease condition (5), as shown in [Absil et al., 2005, §4].

### 3.3 Asymptotic convergence rate

To conclude this section, we briefly review classical linear convergence results for gradient methods under the PL assumption, as needed for Section 4. Proofs are in Appendix D for completeness. It is well known that gradient descent with appropriate step-sizes converges linearly to a minimum when  $f$  satisfies PL globally and has a Lipschitz continuous gradient [Polyak, 1963]. The same arguments lead to an asymptotic linear convergence rate when PL holds only locally. We say that the sequence  $\{x_k\}$  satisfies the *sufficient decrease property* with constant  $\omega > 0$  if

$$f(x_k) - f(x_{k+1}) \geq \omega \|\nabla f(x_k)\|^2. \quad (6)$$

for all large enough  $k$ . The classical result below follows from that inequality [Polyak, 1963].

<sup>3</sup>Notice that Absil et al. [2005] do *not* assume vanishing steps, but in that case the property can fail when  $\bar{x}$  is not a global minimum. The statement of Corollary 3.7 accounts for this technical point.

**Proposition 3.9.** *Let  $\{x_k\}$  be a sequence of iterates satisfying (6) and converging to some  $\bar{x} \in \mathcal{S}$ . Suppose  $f$  satisfies (PL) around  $\bar{x}$  with constant  $\mu > 0$ . Then the sequence  $\{f(x_k)\}$  converges linearly to  $f_{\mathcal{S}}$  with rate  $1 - 2\omega\mu$ . Moreover,  $\{\|\nabla f(x_k)\|\}$  and  $\{\text{dist}(x_k, \mathcal{S})\}$  converge linearly to zero with rate  $\sqrt{1 - 2\omega\mu} \leq 1 - \omega\mu$ .*

In the case where  $\mathcal{M}$  is a Euclidean space and  $R_x(s) = x + s$ , it is well known that the sufficient decrease condition (6) holds for many first-order algorithms when  $\nabla f$  is Lipschitz continuous. This is also true for a general manifold  $\mathcal{M}$  and retraction  $R$  as we briefly describe now. We say that  $f$  and the retraction  $R$  locally satisfy a *Lipschitz-type* property around  $\bar{x} \in \mathcal{S}$  if there exists  $L > 0$  such that

$$f(R_x(s)) \leq f(x) + \langle \nabla f(x), s \rangle + \frac{L}{2} \|s\|^2 \quad (7)$$

for all  $x$  close enough to  $\bar{x}$  and  $s$  small enough. Note that if  $f \circ R$  is  $C^2$  then the inequality (7) is always (locally) satisfied. It is a classical result that (7) implies sufficient decrease for gradient descent with constant step-sizes. This yields the following statement.

**Proposition 3.10.** *Suppose  $f$  satisfies  $\mu$ -(PL) around  $\bar{x} \in \mathcal{S}$ . Also assume that (7) holds around  $\bar{x}$ . Let  $\{x_k\}$  be a sequence of iterates generated by gradient descent with constant step-size  $\gamma \in (0, \frac{2}{L})$ , that is,  $x_{k+1} = R_{x_k}(-\gamma \nabla f(x_k))$ . Given a neighborhood  $\mathcal{U}$  of  $\bar{x}$ , there exists a neighborhood  $\mathcal{V}$  of  $\bar{x}$  such that if an iterate enters  $\mathcal{V}$  then the sequence converges linearly to some  $x_{\infty} \in \mathcal{U} \cap \mathcal{S}$  with rate  $\sqrt{1 - 2\mu(\gamma - \frac{L}{2}\gamma^2)}$ .*

## 4 Aiming for super-linear convergence

Under fairly general assumptions, the PL inequality (which is compatible with non-isolated minima) ensures stability of minima and linear convergence for first-order methods, as recalled in the previous section. We now assume  $f$  is  $C^2$  and investigate super-linear convergence to non-isolated minima.

A natural starting point is Newton's method which, in spite of terrible global behavior [Jarre and Toint, 2016], enjoys quadratic convergence to a *non-singular* minimum, provided the method is initialized sufficiently close. Unfortunately, this does not extend to non-isolated minima.

We exhibit here an example showing that the MB property is in general not sufficient to ensure such a strong convergence behavior. The update rule is  $x_{k+1} = x_k - \nabla^2 f(x_k)^{-1} [\nabla f(x_k)]$  (we may use the pseudo-inverse instead). Consider the cost function  $f(x, y) = \frac{1}{2}(x^2 + 1)y^2$ , whose set of minima is the line  $\mathcal{S} = \{(x, y) \in \mathbb{R}^2 : y = 0\}$ . The gradient and Hessian of  $f$  are given by

$$\nabla f(x, y) = \begin{bmatrix} xy^2 \\ (x^2 + 1)y \end{bmatrix} \quad \text{and} \quad \nabla^2 f(x, y) = \begin{bmatrix} y^2 & 2xy \\ 2xy & x^2 + 1 \end{bmatrix}.$$

One can check that  $f$  satisfies (MB). To see how Newton's method behaves on  $f$ , notice that

$$\nabla^2 f(x, y)^{\dagger} \nabla f(x, y) = \frac{1}{3x^2 - 1} \begin{bmatrix} x^3 + x \\ (x^2 - 1)y \end{bmatrix}$$

whenever  $3x^2 \neq 1$ . Let  $x(t) = \sqrt{\frac{1-t}{3}}$  and  $y(t) = \sqrt{t}$ . We can choose  $t \in (0, 1)$  as small as desired to make the point  $(x(t), y(t))$  arbitrarily close to  $\mathcal{S}$ . Yet computing the Newton step at  $(x(t), y(t))$  results in a new point at a distance  $\frac{2}{3} \frac{1-t}{\sqrt{t}}$  from the optimal set  $\mathcal{S}$ : that is arbitrarily far away. The failure of Newton's method stems from a misalignment between the gradient and some eigenspaces of the Hessian.

The usual fix for Newton's method is to regularize it. This yields two classes of algorithms in particular: regularized Newton with cubics and trust-region methods. We will show that cubic regularization enjoys satisfying local convergence properties, even in the presence of non-isolated minima. In contrast, the picture is less clear for trust-region methods.

Throughout, given a point  $\bar{x} \in \mathcal{S}$ , we manipulate two local assumptions as stated below. For the retraction  $R = \text{Exp}$  they are equivalent (with the same constant). This is true in particular when  $\mathcal{M}$  is a Euclidean space and  $R_x(s) = x + s$ .

**A1.** The Hessian  $\nabla^2 f$  is locally  $L_H$ -Lipschitz continuous around  $\bar{x}$  for some  $L_H \geq 0$ .

**A2.** There exists a constant  $L'_H \geq 0$  such that the Lipschitz-type inequality

$$f(\mathbb{R}_x(s)) - f(x) - \langle s, \nabla f(x) \rangle - \frac{1}{2} \langle s, \nabla^2 f(x)[s] \rangle \leq \frac{L'_H}{6} \|s\|^3 \quad (8)$$

holds for all  $x$  close enough to  $\bar{x}$  and all  $s$  small enough.

These assumptions typically hold locally. We further assume that the retraction satisfies the bound (4).

## 4.1 Adaptive regularized Newton

The regularized Newton method using cubics was introduced by Griewank [1981] and later revisited by Nesterov and Polyak [2006]. An adaptive version of this algorithm was proposed by Cartis et al. [2011a,b], with versions on manifolds proposed by Qi [2011], Zhang and Zhang [2018] and Agarwal et al. [2021]. The adaptive variants update the penalty weight automatically: they are called ARC. We consider those variants, and more specifically an algorithm that generates sequences  $\{(x_k, s_k, \varsigma_k)\}$ , where  $\varsigma_k$  is the cubic penalty weight, and  $s_k$  is the step at iteration  $k$ , such that  $x_{k+1} = \mathbb{R}_{x_k}(s_k)$ . At each iteration  $k$ , we define a linear operator  $H_k: \mathbb{T}_{x_k} \mathcal{M} \rightarrow \mathbb{T}_{x_k} \mathcal{M}$  and the step  $s_k$  is chosen to approximately minimize the regularized second-order model

$$m_k(s) = f(x_k) + \langle s, \nabla f(x_k) \rangle + \frac{1}{2} \langle s, H_k[s] \rangle + \frac{\varsigma_k}{3} \|s\|^3 \quad (9)$$

in a way that we make precise below. We require the map  $H_k$  to be close to  $\nabla^2 f(x_k)$ : given a point  $\bar{x} \in \mathcal{S}$ , we assume that there is a constant  $\beta_H \geq 0$  such that

$$\|H_k - \nabla^2 f(x_k)\| \leq \beta_H \|\nabla f(x_k)\| \quad (10)$$

whenever the iterate  $x_k$  is close enough to  $\bar{x}$ .

In the literature, there are a number of super-linear (but non-quadratic) convergence results for such algorithms. Assuming a PL inequality, Nesterov and Polyak [2006] showed that regularized Newton generates sequences that converge super-linearly, with exponent  $4/3$ . Later, assuming a Lojasiewicz inequality with exponent  $\theta \in [0, 1)$ , Zhou et al. [2018] characterized the convergence speed of regularized Newton depending on  $\theta$ . In particular they also show that the PL inequality implies super-linear convergence.<sup>4</sup> More recently, Qian and Pan [2022] developed an abstract framework that encompasses these super-linear convergence results, and Cartis et al. [2022, §5.3] reviewed super-linear convergence rates of ARC under Lojasiewicz inequalities.

There is also a quadratic convergence result: Yue et al. [2019] employed a local error bound assumption to show quadratic convergence for the regularized Newton method. As discussed in Section 2, this assumption is equivalent to local PL, making their result an improvement over the super-linear rates from the aforementioned references. This underlines one of the benefits of recognizing the equivalence of the four conditions MB, PL, EB and QG, as some may more readily lead to a sharp analysis than others.

Note that the results in [Yue et al., 2019] assume that the subproblem is solved exactly, meaning that  $s_k \in \arg \min_s m_k(s)$ . Several authors proposed weaker conditions on  $s_k$  (only requiring an approximate solution to the subproblem) to ensure convergence guarantees: this is important because we cannot find an exact solution in practice. Agarwal et al. [2021] for example, following Birgin et al. [2017], establish global convergence guarantees assuming only that  $m_k(s_k) \leq m_k(0)$  and  $\|\nabla m_k(s_k)\| \leq \kappa \|s_k\|^2$  for some  $\kappa \geq 0$ . For their results to hold, they require  $H_k = \nabla^2 \hat{f}_k(0)$ , where  $\hat{f}_k = f \circ \mathbb{R}_x$  is the pullback of  $f$  at  $x$ . This choice of  $H_k$  is compatible with (10) for retractions with bounded initial acceleration (which is typical).

<sup>4</sup>In fact, Zhou et al. [2018] show a super-linear convergence rate only assuming that  $\theta \in (0, \frac{2}{3})$  and that  $\nabla^2 f$  is Lipschitz continuous. However, these assumptions imply PL as noted in Remark 2.21.

We revisit the results of Yue et al. [2019] to obtain an asymptotic quadratic convergence rate for ARC under the PL assumption, even with approximate solutions to the subproblem. Specifically, throughout this section we suppose that the steps  $s_k$  satisfy

$$m_k(s_k) \leq m_k(0) \quad \text{and} \quad \|\nabla m_k(s_k)\| \leq \kappa \|s_k\| \|\nabla f(x_k)\| \quad (11)$$

for some  $\kappa \geq 0$ . At each iteration  $k$ , we define the ratio

$$\varrho_k = \frac{f(x_k) - f(\mathbf{R}_{x_k}(s_k))}{m_k(0) - m_k(s_k) + \frac{\varsigma_k}{3} \|s_k\|^3} \quad (12)$$

(as do Birgin et al. [2017]), which measures the adequacy of the local model. Iteration  $k$  is said to be *successful* when  $\varrho_k$  is larger than some fixed parameter  $\varrho_c \in (0, 1)$ . In this case, we set  $x_{k+1} = \mathbf{R}_{x_k}(s_k)$  and decrease the penalty weight so that  $\varsigma_{k+1} \leq \varsigma_k$ . The update mechanism ensures that  $\varsigma_k \geq \varsigma_{\min}$  for all  $k$ , where  $\varsigma_{\min} > 0$  is a fixed parameter. Conversely, the step is *unsuccessful* when  $\varrho_k < \varrho_c$ : we set  $x_{k+1} = x_k$  and increase the penalty so that  $\varsigma_{k+1} > \varsigma_k$ . The explicit updates for  $\varsigma_{k+1}$  are stated in [Agarwal et al., 2021]. We prove the following result for this algorithm.

**Theorem 4.1.** *Suppose A1, A2 and (PL) hold around  $\bar{x} \in \mathcal{S}$ . We run ARC with inexact subproblem solver satisfying (10) and (11). Given any neighborhood  $\mathcal{U}$  of  $\bar{x}$ , there exists a neighborhood  $\mathcal{V}$  of  $\bar{x}$  such that if an iterate enters  $\mathcal{V}$  then the sequence converges quadratically to some  $x_\infty \in \mathcal{U} \cap \mathcal{S}$ .*

We first adapt an argument from [Agarwal et al., 2021, Lem. 6] to show that ARC satisfies the vanishing steps property (VS) defined in Section 3.1.

**Lemma 4.2.** *The step-sizes have norms bounded as  $\|s_k\| \leq \tilde{\eta}(x_k, \varsigma_k) \leq \tilde{\eta}(x_k, \varsigma_{\min})$ , where*

$$\tilde{\eta}(x, \varsigma) = \sqrt{\frac{3\|\nabla f(x)\|}{\varsigma}} + \frac{3}{2\varsigma} \Lambda(x) \quad \text{and} \quad \Lambda(x) = \max\left(0, \beta_H \|\nabla f(x)\| - \lambda_{\min}(\nabla^2 f(x))\right).$$

In particular, ARC has the (VS) property around second-order critical points with  $\eta(x) = c_r \tilde{\eta}(x, \varsigma_{\min})$ , where  $c_r$  controls possible retraction distortion as in (4).

*Proof.* The model decrease in (11) and the model accuracy (10) ensure

$$\begin{aligned} \varsigma_k \|s_k\|^3 &\leq -3 \left\langle s_k, \nabla f(x_k) + \frac{1}{2} \nabla^2 f(x_k)[s_k] + \frac{1}{2} (H_k - \nabla^2 f(x_k))[s_k] \right\rangle \\ &\leq 3 \|s_k\| \left( \|\nabla f(x_k)\| + \frac{1}{2} \Lambda(x_k) \|s_k\| \right). \end{aligned}$$

Divide by  $\|s_k\|$  and solve the quadratic inequality for  $\|s_k\|$  to get the result, recalling  $\varsigma_k \geq \varsigma_{\min}$ .  $\square$

The function  $\eta$  is indeed continuous with value zero on  $\mathcal{S}$ , as required in (VS). To obtain the vanishing steps property we only relied on the decrease requirement  $m_k(s_k) \leq m_k(0)$ . Assuming a PL inequality and a locally Lipschitz continuous Hessian, we now derive sharper bounds for the steps. We rely on the fact that that gradient of the model  $m_k$  (9)

$$\nabla m_k(s_k) = \nabla f(x_k) + H_k[s_k] + \varsigma_k \|s_k\| s_k \quad (13)$$

is small. We will exploit the particular alignment of  $\nabla f$  given in Lemma 2.14. In the following statements, given a point  $\bar{x} \in \mathcal{S}$  and  $d = \text{rank}(\nabla^2 f(\bar{x}))$ , we let  $P(x): \mathbb{T}_x \mathcal{M} \rightarrow \mathbb{T}_x \mathcal{M}$  denote the orthogonal projector onto the top  $d$  eigenspace of  $\nabla^2 f(x)$ . This is always well defined in a neighborhood of  $\bar{x}$  by continuity of eigenvalues. Additionally, we let  $Q(x) = I - P(x)$  be the projector onto the orthogonal complement.

**Lemma 4.3.** *Suppose that  $f$  is  $\mu$ -(PL) and satisfies A1 around  $\bar{x} \in \mathcal{S}$ . Given  $\varepsilon > 0$ ,  $\mu^b < \mu$  and  $\lambda^\sharp > \lambda_{\max}(\nabla^2 f(\bar{x}))$ , there exists a neighborhood  $\mathcal{U}$  of  $\bar{x}$  and a constant  $L_q \geq 0$  such that if  $x_k$  is an iterate in  $\mathcal{U}$  and  $s_k$  is a step satisfying (11) then*

$$(1 - \varepsilon) \frac{\|\nabla f(x_k)\|}{\lambda^\sharp + \varsigma_k \|s_k\|} \leq \|P(x_k) s_k\| \leq (1 + \varepsilon) \frac{\|\nabla f(x_k)\|}{\mu^b + \varsigma_k \|s_k\|} \quad \text{and} \quad (14)$$

$$\|Q(x_k) s_k\| \leq \frac{1}{\varsigma_k} \left( (\kappa + \beta_H) \|\nabla f(x_k)\| + (L_H + L_q \sqrt{\varsigma_k}) \text{dist}(x_k, \mathcal{S}) \right). \quad (15)$$

*Proof.* Let  $\mathcal{U}$  be a neighborhood of  $\bar{x}$  where the orthogonal projector  $P(x)$  is well defined for all  $x \in \mathcal{U}$ . Let  $x_k \in \mathcal{U}$  and  $s_k$  be a step that satisfies (11). We first bound the term  $P(x_k)s_k$ . Multiply (13) by  $P(x_k)$  and use commutativity of  $P(x_k)$  and  $\nabla^2 f(x_k)$  to get

$$(\nabla^2 f(x_k) + \varsigma_k \|s_k\| I) P(x_k) s_k = P(x_k) \left( -\nabla f(x_k) + \nabla m_k(s_k) - (H_k - \nabla^2 f(x_k))[s_k] \right).$$

If we apply (10) and (11), we find that  $\|\nabla m_k(s_k) - (H_k - \nabla^2 f(x_k))[s_k]\| \leq (\kappa + \beta_H) \|\nabla f(x_k)\| \|s_k\|$  when  $x_k$  is close enough to  $\bar{x}$  (shrink  $\mathcal{U}$  as needed). Consequently, the previous equality yields

$$\frac{\|P(x_k) \nabla f(x_k)\| - (\kappa + \beta_H) \|\nabla f(x_k)\| \|s_k\|}{\lambda_1(\nabla^2 f(x_k)) + \varsigma_k \|s_k\|} \leq \|P(x_k) s_k\| \leq \frac{\|P(x_k) \nabla f(x_k)\| + (\kappa + \beta_H) \|\nabla f(x_k)\| \|s_k\|}{\lambda_d(\nabla^2 f(x_k)) + \varsigma_k \|s_k\|}.$$

Lemma 2.14 gives that  $\|P(x) \nabla f(x)\| = \|\nabla f(x)\| + o(\|\nabla f(x)\|)$  as  $x \rightarrow \bar{x}$ . Moreover, the steps  $s_k$  vanish (as shown in Lemma 4.2), so we obtain the bound (14) when  $x_k$  is sufficiently close to  $\bar{x}$ . We now let  $Q(x) = I - P(x)$  and consider the term  $Q(x_k)s_k$ . Multiply (13) by  $Q(x_k)$  to obtain

$$Q(x_k) \nabla m_k(s_k) = Q(x_k) \nabla f(x_k) + \nabla^2 f(x_k) Q(x_k) s_k + Q(x_k) (H_k - \nabla^2 f(x_k))[s_k] + \varsigma_k \|s_k\| Q(x_k) s_k.$$

Taking the inner product of this expression with  $Q(x_k)s_k$ , applying the Cauchy–Schwarz inequality, dividing by  $\|s_k\|$ , and using (10) yields

$$\varsigma_k \|Q(x_k) s_k\|^2 \leq \|Q(x_k) \nabla f(x_k)\| + \left( \kappa \|\nabla f(x_k)\| + \Lambda(x_k) \right) \|Q(x_k) s_k\|,$$

where  $\Lambda$  is as in Lemma 4.2. Solving the quadratic inequality gives

$$\|Q(x_k) s_k\| \leq \frac{1}{\varsigma_k} \left( \sqrt{\varsigma_k \|Q(x_k) \nabla f(x_k)\|} + \kappa \|\nabla f(x_k)\| + \Lambda(x_k) \right).$$

Local Lipschitz continuity of  $\nabla^2 f$  provides  $\Lambda(x_k) \leq \beta_H \|\nabla f(x_k)\| + L_H \text{dist}(x_k, \mathcal{S})$  when  $x_k$  is close to  $\bar{x}$ . Via Lemma 2.14, it also provides a constant  $L_q \geq 0$  such that  $\sqrt{\|Q(x_k) \nabla f(x_k)\|} \leq L_q \text{dist}(x_k, \mathcal{S})$  when  $x_k$  is sufficiently close to  $\bar{x}$ . This is enough to secure (15).  $\square$

Using these bounds, we now show that  $\|P(x_k)s_k\|$  cannot be too small compared to  $\|s_k\|$  when  $x_k$  is close to a minimum where PL holds.

**Lemma 4.4.** *Suppose that  $f$  is  $\mu$ -PL and satisfies A1 around  $\bar{x} \in \mathcal{S}$ . Given  $\varepsilon > 0$  and  $\lambda^\# > \lambda_{\max}(\nabla^2 f(\bar{x}))$ , there exists a neighborhood  $\mathcal{U}$  of  $\bar{x}$  and a constant  $L_q \geq 0$  such that if  $x_k$  is an iterate in  $\mathcal{U}$  and  $s_k$  is a step satisfying (11) then  $\|P(x_k)s_k\| \geq r \|s_k\|$  where  $r > 0$  is a constant given by*

$$r = \sqrt{1 - \frac{1}{1 + \tilde{r}^2}} \quad \text{with} \quad \tilde{r} = \frac{(1 - \varepsilon) \varsigma_{\min}}{(\lambda^\# + \varepsilon(1 + \sqrt{\varsigma_{\min}})) \left( \kappa + \beta_H + \frac{L_H + L_q \sqrt{\varsigma_{\min}}}{\mu} \right)}.$$

*Proof.* Assume  $x_k$  is sufficiently close to  $\bar{x}$  for the projectors  $P(x_k)$  and  $Q(x_k)$  to be well defined. Define  $\nu_p = \|P(x_k)s_k\|$ ,  $\nu_q = \|Q(x_k)s_k\|$  and  $\xi = \frac{\nu_p}{\nu_q}$  (consider  $\nu_q \neq 0$  as otherwise the claim is clear). We compute that  $\nu_p^2 = (1 - \frac{1}{1 + \xi^2}) \|s_k\|^2$  and find a lower-bound for  $\xi$ . Remark 2.10 gives that  $\text{dist}(x_k, \mathcal{S}) \leq \frac{1}{\mu} \|\nabla f(x_k)\|$  when  $x_k$  is sufficiently close to  $\bar{x}$ . Together with the bound on  $\nu_q$  in Lemma 4.3, this gives

$$\nu_q \leq \frac{\|\nabla f(x_k)\|}{\varsigma_k} \left( \kappa + \beta_H + \frac{L_H + L_q \sqrt{\varsigma_k}}{\mu} \right)$$

Combining this with the lower-bound on  $\nu_p$  in Lemma 4.3, we find

$$\xi \geq \frac{(1 - \varepsilon) \varsigma_k}{(\lambda^\# + \varsigma_k \|s_k\|) \left( \kappa + \beta_H + \frac{L_H + L_q \sqrt{\varsigma_k}}{\mu} \right)}.$$

With Lemma 4.2 we can upper-bound  $\varsigma_k \|s_k\| \leq \sqrt{3\varsigma_k \|\nabla f(x_k)\|} + \frac{3}{2} \Lambda(x_k)$ , and consequently,  $\varsigma_k \|s_k\| \leq \varepsilon(1 + \sqrt{\varsigma_k})$  whenever  $x_k$  is sufficiently close to  $\bar{x}$ . We finally notice that the resulting lower-bound on  $\xi$  is an increasing function of  $\varsigma_k$ . Therefore, we get the desired inequality by using  $\varsigma_k \geq \varsigma_{\min}$ .  $\square$

From this we deduce a lower-bound on the quadratic term of the model.

**Lemma 4.5.** *Suppose that  $f$  is  $\mu$ -(PL) and satisfies A1 around  $\bar{x} \in \mathcal{S}$ . Given  $\mu^b < \mu$ , there exists a constant  $r > 0$  (provided by Lemma 4.4) and a neighborhood  $\mathcal{U}$  of  $\bar{x}$  such that if  $x_k \in \mathcal{U}$  then the step satisfies*

$$\langle s_k, \nabla^2 f(x_k)[s_k] \rangle \geq \mu^b r^2 \|s_k\|^2.$$

*Proof.* Let  $r > 0$  be a constant and  $\mathcal{U}$  a neighborhood of  $\bar{x}$  as in Lemma 4.4. Shrink  $\mathcal{U}$  for the projectors  $P$  and  $Q$  to be well defined in  $\mathcal{U}$ . If  $x_k$  is in  $\mathcal{U}$  we compute

$$\begin{aligned} \langle s_k, \nabla^2 f(x_k)[s_k] \rangle &= \langle P(x_k)s_k, \nabla^2 f(x_k)[P(x_k)s_k] \rangle + \langle Q(x_k)s_k, \nabla^2 f(x_k)[Q(x_k)s_k] \rangle \\ &\geq \lambda_d (\nabla^2 f(x_k)) r^2 \|s_k\|^2 + \lambda_{\min}(\nabla^2 f(x_k)) \|Q(x_k)s_k\|^2, \end{aligned}$$

where we used  $\|P(x_k)s_k\| \geq r\|s_k\|$ . We obtain the result by noticing that the second term is lower-bounded by  $\max(0, -\lambda_{\min}(\nabla^2 f(x_k)))\|s_k\|^2$ .  $\square$

We now show that the ratio  $\varrho_k$  is large when  $x_k$  is close to a local minimum where PL holds. The upshot is that iterations near  $\mathcal{S}$  are successful.

**Lemma 4.6.** *Suppose that  $f$  is  $\mu$ -(PL) and satisfies A1, A2 around  $\bar{x} \in \mathcal{S}$ . For all  $\varepsilon > 0$  there exists a neighborhood  $\mathcal{U}$  of  $\bar{x}$  such that if  $x_k \in \mathcal{U}$  then  $\varrho_k \geq 1 - \varepsilon$ .*

*Proof.* Let  $\tau_k(s) = f(x_k) + \langle s, \nabla f(x_k) \rangle + \frac{1}{2} \langle s, H_k[s] \rangle$  be the second-order component of the model  $m_k$ . Then,

$$1 - \varrho_k = \frac{f(\mathbf{R}_{x_k}(s_k)) - \tau_k(s_k)}{\tau_k(0) - \tau_k(s_k)} \leq \frac{1}{6} \frac{L'_H \|s_k\|^3 + 3\beta_H \|s_k\|^2 \|\nabla f(x_k)\|}{\tau_k(0) - \tau_k(s_k)},$$

where the bound on the numerator comes from A2 and (10). The denominator is given by

$$\begin{aligned} \tau_k(0) - \tau_k(s_k) &= -\langle s_k, \nabla f(x_k) \rangle - \frac{1}{2} \langle s_k, H_k[s_k] \rangle \\ &= -\langle s_k, \nabla m_k(s_k) \rangle + \frac{1}{2} \langle s_k, (H_k - \nabla^2 f(x_k))[s_k] \rangle + \frac{1}{2} \langle s_k, \nabla^2 f(x_k)[s_k] \rangle + \varsigma_k \|s_k\|^3, \end{aligned}$$

where we used identity (13) for the second equality. With the Cauchy–Schwarz inequality, (10) and (11), we bound the two first terms as

$$|\langle s_k, \nabla m_k(s_k) \rangle| \leq \kappa \|\nabla f(x_k)\| \|s_k\|^2 \quad \text{and} \quad |\langle s_k, (H_k - \nabla^2 f(x_k))[s_k] \rangle| \leq \beta_H \|\nabla f(x_k)\| \|s_k\|^2.$$

If we combine these bounds with Lemma 4.5, we find that given  $\mu^b < \mu$ , there exists  $r > 0$  and a neighborhood  $\mathcal{U}$  of the local minimum  $\bar{x}$  such that

$$x_k \in \mathcal{U} \Rightarrow \tau_k(0) - \tau_k(s_k) \geq \mu^b r^2 \|s_k\|^2,$$

and therefore  $1 - \varrho_k \leq \frac{L'_H \|s_k\| + 3\beta_H \|\nabla f(x_k)\|}{6\mu^b r^2}$ . Owing to Lemma 4.2 the steps  $s_k$  vanish around second-order critical points so we can guarantee  $1 - \varrho_k$  becomes as small as desired.  $\square$

We can now show that the iterates produced by ARC satisfy the strong decrease property (5) around minima where PL holds.

**Proposition 4.7.** *Suppose  $f$  is  $\mu$ -(PL) and satisfies A1, A2 around  $\bar{x} \in \mathcal{S}$ . Given  $\mu^b < \mu$  and  $\lambda^\sharp > \lambda_{\max}(\nabla^2 f(\bar{x}))$ , there exists a neighborhood of  $\bar{x}$  where ARC satisfies the strong decrease property (5) with constant  $\sigma = \frac{r\mu^b}{2c_r\lambda^\sharp}$ , where  $c_r$  is defined in (4) and  $r > 0$  is provided by Lemma 4.4.*

*Proof.* From Lemma 4.6, there exists a neighborhood  $\mathcal{U}$  of  $\bar{x}$  in which all the steps are successful. Given an iterate  $x_k$  in  $\mathcal{U}$ , success implies  $x_{k+1} = \mathbf{R}_{x_k}(s_k)$  and therefore, by definition of  $\varrho_k$  (12),

$$f(x_k) - f(x_{k+1}) = \varrho_k \left( m_k(0) - m_k(s_k) + \frac{\varsigma_k}{3} \|s_k\|^3 \right) = -\varrho_k \left( \langle s_k, \nabla f(x_k) \rangle + \frac{1}{2} \langle s_k, H_k[s_k] \rangle \right).$$

Also, taking the inner product of (13) with  $s_k$  and using (11) yields

$$\langle s_k, H_k[s_k] \rangle \leq -\langle s_k, \nabla f(x_k) \rangle + \kappa \|\nabla f(x_k)\| \|s_k\|^2.$$

Multiply the latter by  $-\frac{\varrho_k}{2}$  and plug into the former to deduce

$$f(x_k) - f(x_{k+1}) \geq -\frac{\varrho_k}{2} \left( \langle s_k, \nabla f(x_k) \rangle + \kappa \|\nabla f(x_k)\| \|s_k\|^2 \right). \quad (16)$$

We now bound the inner product  $\langle s_k, \nabla f(x_k) \rangle$ . Let  $d = \text{rank } \nabla^2 f(\bar{x})$  and restrict the neighborhood  $\mathcal{U}$  if need be to ensure  $\lambda_d(\nabla^2 f(x_k)) > 0$  and  $\lambda_d(\nabla^2 f(x_k)) > \lambda_{d+1}(\nabla^2 f(x_k))$ . In particular, the orthogonal projector  $P(x_k)$  onto the top  $d$  eigenspace of  $\nabla^2 f(x_k)$  is well defined. Let  $Q(x_k) = I - P(x_k)$ . Decompose  $s_k = P(x_k)s_k + Q(x_k)s_k$  and apply the Cauchy–Schwarz inequality to obtain

$$\langle s_k, \nabla f(x_k) \rangle \leq \langle P(x_k)s_k, \nabla f(x_k) \rangle + \|Q(x_k)\nabla f(x_k)\| \|s_k\|. \quad (17)$$

The second term is small owing to Lemma 2.14. Let us focus on the first term. To this end, multiply (13) by  $P(x_k)$  to verify the following (recall that  $\nabla^2 f(x_k)$  and  $P(x_k)$  commute):

$$P(x_k)\nabla f(x_k) = -\nabla^2 f(x_k)P(x_k)s_k + P(x_k)\nabla m_k(s_k) - P(x_k)(H_k - \nabla^2 f(x_k))s_k - \varsigma_k \|s_k\| P(x_k)s_k.$$

On the one hand, we can use it with (10) and (11) to lower-bound the norm of  $P(x_k)s_k$ , through:

$$\|P(x_k)\nabla f(x_k)\| \leq \left( \lambda_1(\nabla^2 f(x_k)) + \varsigma_k \|s_k\| \right) \|P(x_k)s_k\| + (\kappa + \beta_H) \|\nabla f(x_k)\| \|s_k\|.$$

On the other hand, we can use it to upper-bound  $\langle s_k, P(x_k)\nabla f(x_k) \rangle$ , also with (10) and (11) and using the fact that  $P(x_k)s_k$  lives in the top- $d$  eigenspace of  $\nabla^2 f(x_k)$ , like so:

$$\langle s_k, P(x_k)\nabla f(x_k) \rangle \leq -\left( \lambda_d(\nabla^2 f(x_k)) + \varsigma_k \|s_k\| \right) \|P(x_k)s_k\|^2 + (\kappa + \beta_H) \|\nabla f(x_k)\| \|s_k\| \|P(x_k)s_k\|.$$

Combine the two inequalities above as follows: use the former to upper-bound one of the first factors  $\|P(x_k)s_k\|$  in the latter. Also using  $\frac{\lambda_d(\nabla^2 f(x_k))}{\lambda_1(\nabla^2 f(x_k))} \leq \frac{\lambda_d(\nabla^2 f(x_k)) + \varsigma_k \|s_k\|}{\lambda_1(\nabla^2 f(x_k)) + \varsigma_k \|s_k\|} \leq 1$ , this yields:

$$\begin{aligned} \langle s_k, P(x_k)\nabla f(x_k) \rangle &\leq -\frac{\lambda_d(\nabla^2 f(x_k))}{\lambda_1(\nabla^2 f(x_k))} \|P(x_k)\nabla f(x_k)\| \|P(x_k)s_k\| \\ &\quad + 2(\kappa + \beta_H) \|\nabla f(x_k)\| \|s_k\| \|P(x_k)s_k\|. \end{aligned} \quad (18)$$

We now plug this back into (17). Using Lemma 2.14 for its second term and also Lemma 4.2 which asserts  $\|s_k\|$  is arbitrarily small for  $x_k$  near  $\bar{x}$ , we find that for all  $\varepsilon > 0$  we can restrict the neighborhood  $\mathcal{U}$  in order to secure

$$\langle s_k, \nabla f(x_k) \rangle \leq -\frac{\lambda_d(\nabla^2 f(x_k))}{\lambda_1(\nabla^2 f(x_k))} \|P(x_k)\nabla f(x_k)\| \|P(x_k)s_k\| + \varepsilon \|\nabla f(x_k)\| \|s_k\|. \quad (19)$$

Lemma 4.4 provides a (possibly smaller) neighborhood and a positive  $r$  such that  $\|P(x_k)s_k\| \geq r\|s_k\|$ . Also, for any  $\delta > 0$ , Lemma 2.14 ensures  $\|P(x_k)\nabla f(x_k)\| \geq (1 - \delta)\|\nabla f(x_k)\|$  upon appropriate neighborhood restriction. Thus,

$$\langle s_k, \nabla f(x_k) \rangle \leq -\left( \frac{\lambda_d(\nabla^2 f(x_k))}{\lambda_1(\nabla^2 f(x_k))} (1 - \delta)r - \varepsilon \right) \|\nabla f(x_k)\| \|s_k\|. \quad (20)$$

Now plugging back into (16) and possibly restricting the neighborhood again,

$$f(x_k) - f(x_{k+1}) \geq \frac{\varrho_k}{2} \left( \frac{\lambda_d(\nabla^2 f(x_k))}{\lambda_1(\nabla^2 f(x_k))} (1 - \delta)r - \varepsilon - \kappa \|s_k\| \right) \|\nabla f(x_k)\| \|s_k\|. \quad (21)$$

The result now follows since we can arrange to make  $\varepsilon, \delta$  and  $\|s_k\|$  arbitrarily small; to make  $\lambda_d(\nabla^2 f(x_k))$  and  $\lambda_1(\nabla^2 f(x_k))$  arbitrarily close to  $\mu$  and  $\lambda_{\max}(\nabla^2 f(\bar{x}))$  (respectively); and to make  $\varrho_k$  larger than a number arbitrarily close to 1 (Lemma 4.6). The final step is to account for potential distortion due to a retraction (4): this adds the factor  $c_r$ .  $\square$

If we combine this result with Lemma 3.8 we obtain that ARC satisfies the bounded path length property (BPL). In Lemma 4.2 we found that it also satisfies the vanishing steps property (VS). As a result, Corollary 3.7 applies to ARC: if an iterate gets close enough to a point where PL holds then the sequence has a limit. We conclude this section with the quadratic convergence rate of ARC.

**Proposition 4.8.** *Suppose that  $\{x_k\}$  converges to some  $\bar{x} \in \mathcal{S}$  and that  $f$  is (PL) around  $\bar{x}$ . Also assume that A1 and A2 hold around  $\bar{x}$ . Then  $\text{dist}(x_k, \mathcal{S})$  converges quadratically to zero.*

*Proof.* From Lemma 4.6 all the steps are eventually successful. In particular, the penalty weights eventually stop increasing: there exists  $\varsigma_{\max} > 0$  such that  $\varsigma_k \leq \varsigma_{\max}$  for all  $k$ . Let  $\mu^\flat < \mu$  (where  $\mu$  is the PL constant) and  $\lambda^\sharp > \lambda_{\max}(\nabla^2 f(\bar{x}))$ . We first apply the Pythagorean theorem with the upper-bounds from Lemma 4.3. Together with the upper-bound in Proposition 2.8, it implies for all large enough  $k$  that

$$\|s_k\|^2 \leq c_1^2 \text{dist}(x_k, \mathcal{S})^2 \quad \text{where} \quad c_1^2 = \left( \frac{\lambda^\sharp}{\mu^\flat} \right)^2 + \frac{1}{\varsigma_{\min}^2} \left( (\kappa + \beta_H) \lambda^\sharp + L_H + L_q \sqrt{\varsigma_{\min}} \right)^2. \quad (22)$$

We now let  $v_k = s_k - \text{Log}_{x_k}(x_{k+1})$ , which is always well defined for large enough  $k$ . Using EB (given by Remark 2.10), we obtain that for all large enough  $k$  we have

$$\begin{aligned} \text{dist}(x_{k+1}, \mathcal{S}) &\leq \frac{1}{\mu} \|\nabla f(x_{k+1})\| \\ &= \frac{1}{\mu} \left\| \Gamma_{x_{k+1}}^{x_k} \nabla f(x_{k+1}) - \nabla f(x_k) - \nabla^2 f(x_k) [\text{Log}_{x_k}(x_{k+1})] \right. \\ &\quad \left. - \nabla^2 f(x_k) [v_k] - (H_k - \nabla^2 f(x_k)) [s_k] - \varsigma_k \|s_k\| s_k + \nabla m_k(s_k) \right\|, \end{aligned}$$

where we used identity (13) for  $\nabla m_k$ . Now the triangle inequality, (10) and (11) give

$$\text{dist}(x_{k+1}, \mathcal{S}) \leq \frac{1}{\mu} \left( \frac{L_H}{2} \text{dist}(x_k, x_{k+1})^2 + \lambda^\sharp \|v_k\| + \varsigma_k \|s_k\|^2 + (\kappa + \beta_H) \|\nabla f(x_k)\| \|s_k\| \right). \quad (23)$$

Notice that  $\text{dist}(x_k, x_{k+1}) \leq c_r \|s_k\|$  using (4). We now bound the quantity  $\|v_k\|$ . For all  $x \in \mathcal{M}$ , since  $\text{DR}_x(0) = I$ , the inverse function theorem implies that  $\text{R}_x$  is locally invertible and  $\text{DR}_x^{-1}(x) = I$ . It follows that there exists a neighborhood  $\mathcal{U} \subseteq \text{B}(\bar{x}, \text{inj}(\bar{x}))$  of  $\bar{x}$  such that for all  $x, y \in \mathcal{U}$  the quantity  $\text{R}_x^{-1}(y)$  is well defined and satisfies

$$\text{R}_x^{-1}(y) = \text{R}_x^{-1}(x) + \text{DR}_x^{-1}(x) [\text{Log}_x(y)] + O(\text{dist}(x, y)^2) = \text{Log}_x(y) + O(\text{dist}(x, y)^2).$$

In particular, using the identity  $s_k = \text{R}_{x_k}^{-1}(x_{k+1})$ , we find that there exists a constant  $c_2$  such that  $\|v_k\| \leq c_2 \text{dist}(x_k, x_{k+1})^2$  holds for large enough  $k$ . Combining this with (23), we obtain

$$\text{dist}(x_{k+1}, \mathcal{S}) \leq \frac{1}{\mu} \left( \left( \frac{L_H}{2} c_r + \lambda^\sharp c_2 c_r + \varsigma_k \right) \|s_k\|^2 + (\kappa + \beta_H) \|\nabla f(x_k)\| \|s_k\| \right).$$

Finally, using (22) and the upper-bound from Proposition 2.8, we conclude that

$$\text{dist}(x_{k+1}, \mathcal{S}) \leq c_q \text{dist}(x_k, \mathcal{S})^2 \quad \text{where} \quad c_q = \frac{c_1^2}{\mu} \left( \frac{L_H}{2} c_r + \lambda^\sharp c_2 c_r + \varsigma_{\max} \right) + \frac{\lambda^\sharp}{\mu} (\kappa + \beta_H) c_1. \quad \square$$

The quadratic convergence rates of the sequences  $\{f(x_k)\}$  and  $\{\|\nabla f(x_k)\|\}$  follow immediately by QG and EB. Theorem 4.1 is a direct consequence of Corollary 3.7 and Proposition 4.8.



## 4.2 Trust-region algorithms

In this section we analyze Riemannian trust-region algorithms (RTR), which embed Newton iterations in safeguards to ensure global convergence guarantees [Absil et al., 2007]. They produce sequences  $\{(x_k, \Delta_k, s_k)\}$  where  $x_k$  is the current iterate and  $\Delta_k$  is known as the trust-region radius. At iteration  $k$ , we define the trust-region model as

$$m_k(s) = f(x_k) + \langle s, \nabla f(x_k) \rangle + \frac{1}{2} \langle s, \nabla^2 f(x_k)[s] \rangle. \quad (24)$$

The step  $s_k$  is chosen by (usually approximately) solving the trust-region subproblem

$$\min_{s_k \in T_{x_k} \mathcal{M}} m_k(s_k) \quad \text{subject to} \quad \|s_k\| \leq \Delta_k. \quad (\text{TRS})$$

The point  $x_k$  and radius  $\Delta_k$  are then updated depending on how good the model is, as measured by the ratio

$$\rho_k = \frac{f(x_k) - f(\mathbb{R}_{x_k}(s_k))}{m_k(0) - m_k(s_k)}.$$

(If the denominator is zero, we let  $\rho_k = 1$ .) Specifically, given parameters  $\rho' \in (0, \frac{1}{4})$  and  $\bar{\Delta} > 0$ , the update rules for the state are

$$x_{k+1} = \begin{cases} \mathbb{R}_{x_k}(s_k) & \text{if } \rho_k > \rho', \\ x_k & \text{otherwise,} \end{cases} \quad \Delta_{k+1} = \begin{cases} \frac{1}{4} \Delta_k & \text{if } \rho_k < \frac{1}{4}, \\ \min(2\Delta_k, \bar{\Delta}) & \text{if } \rho_k > \frac{3}{4} \text{ and } \|s_k\| = \Delta_k, \\ \Delta_k & \text{otherwise.} \end{cases}$$

**Shortcomings of trust-region with exact subproblem solver.** There exist algorithms to efficiently solve the subproblem exactly (up to some accuracy). Can we provide strong guarantees in the presence of non-isolated minima using an exact subproblem solver? We recall [Nocedal and Wright, 2006, Thm. 4.1] that a vector  $s \in T_{x_k} \mathcal{M}$  is a global solution of the subproblem (TRS) if and only if  $\|s\| \leq \Delta_k$  and there exists a scalar  $\lambda \geq 0$  such that

$$(\nabla^2 f(x_k) + \lambda I)s = -\nabla f(x_k), \quad \lambda(\Delta_k - \|s\|) = 0 \quad \text{and} \quad \nabla^2 f(x_k) + \lambda I \succeq 0. \quad (25)$$

As mentioned in [Wojtowytsch, 2021a,b, Liu et al., 2022], if  $f$  is convex in a neighborhood of  $\bar{x}$  then  $\mathcal{S}$  is locally convex, hence affine. Assuming that  $\mathcal{S}$  is not flat around  $\bar{x}$ , it follows that  $\nabla^2 f$  must have a negative eigenvalue in any neighborhood of  $\bar{x}$ . Consequently, we can arrange for an iterate  $x_k$  to be arbitrarily close to  $\bar{x}$  and for  $\nabla^2 f(x_k)$  to have a negative eigenvalue. From (25), this implies  $\lambda > 0$  and hence  $\|s\| = \Delta_k$ , meaning that  $s$  is at the border of the trust region. Consequently, we cannot hope to show a capture result in the same vein as Corollary 3.7 for this algorithm. Indeed, given an initial radius  $\Delta_0 > 0$ , we can design a cost function with a non-isolated minimum  $\bar{x}$  such that, even if  $x_0$  is arbitrarily close to  $\bar{x}$ , the next iterate is far away.

As an example, consider  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  defined by  $f(x, y) = (x^2 + y^2 - 1)^2$ . The set of solutions is the unit circle and  $f$  satisfies MB. Given  $t > 0$ , define  $x(t) = 0$  and  $y(t) = 1 - t$ . For small enough  $t$ , there are two solutions to (TRS) at  $[x(t) \ y(t)]^\top$  with parameter  $\Delta$ :

$$s = \begin{bmatrix} \pm \sqrt{\Delta^2 - \frac{t^2(t-2)^2}{4(t-1)^2}} \\ \frac{t(t-2)}{2(t-1)} \end{bmatrix}.$$

To check optimality, we can use criterion (25) with  $\lambda = 4t(2-t)$ . We find that  $s \rightarrow [\pm\Delta \ 0]^\top$  as  $t \rightarrow 0$ : the tentative step is far even when  $t$  is small. We could arrange for that step to be accepted by adjusting the function value at the tentative iterate. That rules out even basic capture-type theorems. This type of behavior does not happen with an isolated minimum and positive definite Hessian.

**Trust-region with Cauchy steps.** As just discussed, RTR with an exact subproblem solver can fail in the face of non-isolated minima. However, practical implementations of RTR typically solve the subproblem only approximately. We set out to investigate the robustness of such mechanisms to non-isolated minima.

Our investigation is prompted by the empirical observation that RTR with a popular approximate subproblem solver known as truncated conjugate gradients (tCG, see [Conn et al., 2000, Absil et al., 2007]) seems to enjoy super-linear convergence under PL, even with non-isolated minima. We are actively investigating this, but it appears to require significant additional machinery.

As a stepping stone, we show the following theorem, regarding RTR with a crude subproblem solver that computes *Cauchy steps* (see (28) below). It is relevant in particular because tCG generates a sequence of increasingly good tentative steps, the first of which is the Cauchy step.

**Theorem 4.9.** *Suppose  $f$  satisfies  $\mu$ -(PL) and A2 around  $\bar{x} \in \mathcal{S}$ , and let  $\mathcal{U}$  be a neighborhood of  $\bar{x}$ . There exists a neighborhood  $\mathcal{V}$  of  $\bar{x}$  such that if a sequence of iterates generated by RTR with Cauchy steps enters  $\mathcal{V}$  then the sequence converges linearly to some  $x_\infty \in \mathcal{U} \cap \mathcal{S}$  with rate  $\sqrt{1 - \frac{\mu}{\lambda_{\max}}}$ , where  $\lambda_{\max} = \lambda_{\max}(\nabla^2 f(x_\infty))$ .*

Note that a local convergence analysis of RTR with Cauchy steps is given in [Muoi and Yen, 2017] for *non-singular* local minima. Here, we prove that the favorable convergence properties also hold if we only assume PL.

To prove Theorem 4.9, we first establish a number of intermediate results only assuming the subproblem solver satisfies the properties (26) and (27) defined below. Then we conclude using more specific properties which we could only secure for Cauchy steps at this time. First, given a local minimum  $\bar{x} \in \mathcal{S}$ , we assume that the step  $s_k$  satisfies the *sufficient decrease* condition

$$m_k(0) - m_k(s_k) \geq c_p \|\nabla f(x_k)\| \min\left(\Delta_k, \frac{\|\nabla f(x_k)\|^3}{|\langle \nabla f(x_k), \nabla^2 f(x_k) [\nabla f(x_k)] \rangle|}\right) \quad (26)$$

whenever the iterate  $x_k$  is sufficiently close to  $\bar{x}$ . (If the denominator is zero, consider the rightmost expression to be infinite.) This condition holds for many practical subproblem solvers and ensures global convergence guarantees in particular (see [Absil et al., 2008, §7.4] and [Boumal, 2023, §6.4]). Second, given a local minimum  $\bar{x} \in \mathcal{S}$ , we assume that there exists a constant  $c_s \geq 0$  such that

$$\|s_k\| \leq c_s \|\nabla f(x_k)\| \quad (27)$$

when  $x_k$  is sufficiently close to  $\bar{x}$ .

We find that the ratios  $\{\rho_k\}$  are large around minima where these two conditions hold. This is because they imply that the trust-region model is an accurate approximation of the local behavior of  $f$ . It follows that the steps  $\{s_k\}$  decrease  $f$  nearly as much as predicted by the model.

**Proposition 4.10.** *Suppose that A2 holds around  $\bar{x} \in \mathcal{S}$ . Also assume that the steps  $s_k$  satisfy (26) and (27) around  $\bar{x}$ . For all  $\varepsilon > 0$  there exists a neighborhood  $\mathcal{U}$  of  $\bar{x}$  such that if an iterate  $x_k$  is in  $\mathcal{U}$  then  $\rho_k \geq 1 - \varepsilon$ .*

*Proof.* We follow and adapt some arguments from [Absil et al., 2008, Thm. 7.4.11], which is stated there assuming  $\nabla^2 f(\bar{x}) \succ 0$ . We can dismiss the case where  $\nabla f(x_k) = 0$  because it implies  $\rho_k = 1$ . Using the definitions of  $m_k$  and  $\rho_k$  we have

$$1 - \rho_k = \frac{f(\mathbf{R}_{x_k}(s_k)) - m_k(s_k)}{m_k(0) - m_k(s_k)}.$$

We bound the numerator as  $f(\mathbf{R}_{x_k}(s_k)) - m_k(s_k) \leq \frac{L'_H}{6} \|s_k\|^3 \leq \frac{L'_H c_s^2}{6} \|s_k\| \|\nabla f(x_k)\|^2$  using A2 and inequality (27). Combining this with the sufficient decrease (26) gives

$$1 - \rho_k \leq \frac{L'_H c_s^2 \|s_k\| \|\nabla f(x_k)\|}{6 c_p \min\left(\Delta_k, \frac{\|\nabla f(x_k)\|}{\|\nabla^2 f(x_k)\|}\right)}.$$

If  $\Delta_k$  is active in the denominator then we obtain  $1 - \rho_k \leq \frac{L'_H c_s^2 \|\nabla f(x_k)\|}{6c_p}$  because  $\|s_k\| \leq \Delta_k$ . Otherwise, we obtain  $1 - \rho_k \leq \frac{L'_H c_s^3 \|\nabla f(x_k)\| \|\nabla^2 f(x_k)\|}{6c_p}$ . In both cases this yields the result.  $\square$

This result notably implies that the trust-region radius does not decrease in the vicinity of the minimum  $\bar{x}$ . It means that the trust region eventually becomes inactive when the iterates converge to  $\bar{x}$ . We now employ the particular alignment between the gradient and the top eigenspace of the Hessian induced by PL (see Lemma 2.14) to derive bounds on the inner products  $\langle \nabla f(x), \nabla^2 f(x) [\nabla f(x)] \rangle$ .

**Proposition 4.11.** *Suppose that  $f$  is  $\mu$ -(PL) around  $\bar{x} \in \mathcal{S}$ . Let  $\mu^\flat < \mu$  and  $\lambda^\sharp > \lambda_{\max}(\nabla^2 f(\bar{x}))$ . Then there exists a neighborhood  $\mathcal{U}$  of  $\bar{x}$  such that for all  $x \in \mathcal{U}$  we have*

$$\mu^\flat \|\nabla f(x)\|^2 \leq \langle \nabla f(x), \nabla^2 f(x) [\nabla f(x)] \rangle \leq \lambda^\sharp \|\nabla f(x)\|^2.$$

*Proof.* By continuity of eigenvalues, there exists a neighborhood  $\mathcal{U}$  of  $\bar{x}$  such that for all  $x \in \mathcal{U}$  we have  $\lambda_{\max}(\nabla^2 f(x)) \leq \lambda^\sharp$ . The upper-bound  $\langle \nabla f(x), \nabla^2 f(x) [\nabla f(x)] \rangle \leq \lambda^\sharp \|\nabla f(x)\|^2$  follows immediately. We now prove the lower-bound. Let  $d$  be the rank of  $\nabla^2 f(\bar{x})$ . Given  $x$  sufficiently close to  $\bar{x}$ , we let  $P(x): T_x \mathcal{M} \rightarrow T_x \mathcal{M}$  denote the orthogonal projector onto the top  $d$  eigenspace of  $\nabla^2 f(x)$ . From Lemma 2.14 we have  $\|(I - P(x))\nabla f(x)\|^2 = o(\|\nabla f(x)\|^2)$  as  $x \rightarrow \bar{x}$ . If we write  $Q(x) = I - P(x)$ , we obtain

$$\begin{aligned} \langle \nabla f(x), \nabla^2 f(x) [\nabla f(x)] \rangle &= \langle P(x)\nabla f(x), \nabla^2 f(x)P(x)\nabla f(x) \rangle + \langle Q(x)\nabla f(x), \nabla^2 f(x)Q(x)\nabla f(x) \rangle \\ &\geq \lambda_d(\nabla^2 f(x))\|\nabla f(x)\|^2 + o(\|\nabla f(x)\|^2) \end{aligned}$$

as  $x \rightarrow \bar{x}$ .  $\square$

Combining these results guarantees a linear rate of convergence.

**Proposition 4.12.** *Suppose  $f$  satisfies  $\mu$ -(PL) and A2 around  $\bar{x} \in \mathcal{S}$ . Let  $\{x_k\}$  be a sequence of iterates produced by RTR converging to  $\bar{x}$ . Assume that the steps  $s_k$  satisfy (26) and (27) around  $\bar{x}$ . Then the iterates converge at least linearly with rate  $\sqrt{1 - \frac{2c_p\mu}{\lambda_{\max}}}$ , where  $\lambda_{\max} = \lambda_{\max}(\nabla^2 f(\bar{x}))$ .*

*Proof.* We can assume that  $\nabla f(x_k)$  is non-zero for all  $k$  (otherwise the sequence converges in a finite number of steps). We show that the sequence satisfies the sufficient decrease property (6). Given  $\mu^\flat < \mu$  and  $\lambda^\sharp > \lambda_{\max}(\nabla^2 f(\bar{x}))$ , Proposition 4.11 ensures that

$$\frac{1}{\lambda^\sharp} \leq \frac{\|\nabla f(x_k)\|^2}{\langle \nabla f(x_k), \nabla^2 f(x_k) [\nabla f(x_k)] \rangle} \leq \frac{1}{\mu^\flat}$$

for all large enough  $k$ . We let  $0 < \varepsilon < \frac{3}{4}$  and Proposition 4.10 implies that  $\rho_k \geq 1 - \varepsilon$  for all large enough  $k$ . In particular, the radii  $\{\Delta_k\}$  are bounded away from zero (because the update mechanism does not decrease the radius when  $\rho_k \geq \frac{1}{4}$ ). Combining the definition of  $\rho_k$  and the sufficient decrease (26) gives

$$f(x_k) - f(x_{k+1}) = \rho_k (m_k(0) - m_k(s_k)) \geq \frac{(1 - \varepsilon)c_p}{\lambda^\sharp} \|\nabla f(x_k)\|^2$$

for all large enough  $k$ . We can now conclude with Proposition 3.9.  $\square$

We are now in a position to prove Theorem 4.9. The Cauchy step at iteration  $x_k$  is defined as the minimum of (TRS) with the additional constraint that  $s_k \in \text{span}(\nabla f(x_k))$ . We can find an explicit expression for it: when  $\nabla f(x_k) \neq 0$ , the Cauchy step is  $s_k^c = -t_k^c \nabla f(x_k)$ , where

$$t_k^c = \begin{cases} \min\left(\frac{\|\nabla f(x_k)\|^2}{\langle \nabla f(x_k), \nabla^2 f(x_k) [\nabla f(x_k)] \rangle}, \frac{\Delta_k}{\|\nabla f(x_k)\|}\right) & \text{if } \langle \nabla f(x_k), \nabla^2 f(x_k) [\nabla f(x_k)] \rangle > 0, \\ \frac{\Delta_k}{\|\nabla f(x_k)\|} & \text{otherwise.} \end{cases} \quad (28)$$

Cauchy steps notably satisfy the sufficient decrease property (26) globally with  $c_p = \frac{1}{2}$  (see [Conn et al., 2000, Thm. 6.3.1]). We now prove that they also satisfy (27) around minima where PL holds.

**Proposition 4.13.** *Suppose that  $f$  satisfies  $\mu$ -(PL) around  $\bar{x} \in \mathcal{S}$ . Given  $0 < \mu^b < \mu$ , there exists a neighborhood  $\mathcal{U}$  of  $\bar{x}$  such that if an iterate  $x_k$  is in  $\mathcal{U}$  then the Cauchy step satisfies  $\|s_k^c\| \leq \frac{1}{\mu^b} \|\nabla f(x_k)\|$ .*

*Proof.* Given  $\mu^b < \mu$ , Proposition 4.11 yields that  $\langle \nabla f(x_k), \nabla^2 f(x_k)[\nabla f(x_k)] \rangle \geq \mu^b \|\nabla f(x_k)\|^2$  if  $x_k$  is sufficiently close to  $\bar{x}$ . It implies that the step-sizes are bounded as  $t_k^c \leq \frac{1}{\mu^b}$ , which gives the result.  $\square$

In particular, this proposition shows that the RTR algorithm with Cauchy steps satisfies the (VS) property at  $\bar{x}$  with  $\eta(x) = \frac{c_r}{\mu^b} \|\nabla f(x)\|$ , where  $c_r$  is as in (4). Furthermore, Cauchy steps satisfy the model decrease

$$m_k(0) - m_k(s_k^c) \geq \frac{1}{2} \|\nabla f(x_k)\| \|s_k^c\|,$$

as shown in [Absil et al., 2005, Lem. 4.3]. It implies that RTR with Cauchy steps generates sequences that satisfy the strong decrease property (5) with  $\sigma = \frac{\rho'}{2c_r}$ , where  $c_r$  is as in (4), and  $\rho'$  is defined in the algorithm description in Section 4.2. See [Absil et al., 2005, Thm. 4.4] for details on this. As a direct consequence, Lemma 3.8 applies: the RTR algorithm with Cauchy steps satisfies the bounded path length property (BPL) at points where a Lojasiewicz inequality holds. We can finally combine the statements from Corollary 3.7 and Proposition 4.12 to obtain Theorem 4.9.

**Remark 4.14.** *The model decrease (26) is not a sufficient condition for the strong decrease property (5) to hold. As a result, it is not straightforward to determine whether the bounded path length property (BPL) holds for a given subproblem solver: see [Absil et al., 2005, §4.2] for a discussion of this.*

## Acknowledgments

We thank Dmitriy Drusvyatskiy and Christopher Criscitiello for insightful discussions and pointers to literature.

## References

- Hadi Abbaszadehpousti, Etienne de Klerk, and Moslem Zamani. Conditions for linear convergence of the gradient method for non-convex optimization. *arXiv preprint arXiv:2204.00647*, 2022.
- P-A Absil, Christopher G Baker, and Kyle A Gallivan. Trust-region methods on Riemannian manifolds. *Foundations of Computational Mathematics*, 7(3):303–330, 2007.
- P-A Absil, Robert Mahony, and Rodolphe Sepulchre. *Optimization algorithms on matrix manifolds*. Princeton University Press, 2008.
- Pierre-Antoine Absil, Robert Mahony, and Benjamin Andrews. Convergence of the iterates of descent methods for analytic cost functions. *SIAM Journal on Optimization*, 16(2):531–547, 2005.
- Naman Agarwal, Nicolas Boumal, Brian Bullins, and Coralia Cartis. Adaptive regularization with cubics on manifolds. *Mathematical Programming*, 188(1):85–134, 2021.
- Mihai Anitescu. Degenerate nonlinear programming with a quadratic growth condition. *SIAM Journal on Optimization*, 10(4):1116–1135, 2000.
- Vassilis Apidopoulos, Nicolò Ginatta, and Silvia Villa. Convergence rates for the Heavy-Ball continuous dynamics for non-convex optimization, under Polyak–Lojasiewicz conditioning. *Journal of Global Optimization*, 84:563 – 589, 2021.

- Michael Arbel and Julien Mairal. Non-convex bilevel games with critical point selection maps. *Advances in Neural Information Processing Systems*, 35, 2022.
- Hédy Attouch, Jérôme Bolte, Patrick Redont, and Antoine Soubeyran. Proximal alternating minimization and projection methods for nonconvex problems: An approach based on the Kurdyka–Lojasiewicz inequality. *Mathematics of operations research*, 35(2):438–457, 2010.
- Hedy Attouch, Jérôme Bolte, and Benar Fux Svaiter. Convergence of descent methods for semi-algebraic and tame problems: proximal algorithms, forward–backward splitting, and regularized Gauss–Seidel methods. *Mathematical Programming*, 137(1):91–129, 2013.
- Raef Bassily, Mikhail Belkin, and Siyuan Ma. On exponential convergence of SGD in non-convex over-parametrized learning. *arXiv preprint arXiv:1811.02564*, 2018.
- Mikhail Belkin. Fit without fear: remarkable mathematical phenomena of deep learning through the prism of interpolation. *Acta Numerica*, 30:203–248, 2021.
- S Bellavia and B Morini. Strong local convergence properties of adaptive regularized methods for nonlinear least squares. *IMA Journal of Numerical Analysis*, 35(2):947–968, 2015.
- Giovanni Bellettini, M Masala, and Matteo Novaga. A conjecture of de giorgi on the square distance function. *Journal of Convex Analysis*, 14(2):353, 2007.
- D.P. Bertsekas. *Nonlinear Programming*. Athena Scientific, 1995.
- Ernesto G Birgin, JL Gardenghi, José Mario Martínez, Sandra Augusta Santos, and Ph L Toint. Worst-case evaluation complexity for unconstrained nonlinear optimization using high-order regularized models. *Mathematical Programming*, 163(1):359–368, 2017.
- Jérôme Bolte and Edouard Pauwels. Curiosities and counterexamples in smooth convex optimization. *Mathematical Programming*, pages 1–51, 2021.
- Jérôme Bolte, Aris Daniilidis, Olivier Ley, and Laurent Mazet. Characterizations of Lojasiewicz inequalities: subgradient flows, talweg, convexity. *Transactions of the American Mathematical Society*, 362(6):3319–3363, 2010.
- Jérôme Bolte, Shoham Sabach, and Marc Teboulle. Proximal alternating linearized minimization for nonconvex and nonsmooth problems. *Mathematical Programming*, 146(1):459–494, 2014.
- Jérôme Bolte, Trong Phong Nguyen, J. Peypouquet, and Bruce W. Suter. From error bounds to the complexity of first-order descent methods for convex functions. *Mathematical Programming*, 165:471–507, 2017.
- Joseph Frédéric Bonnans and Alexander Ioffe. Second-order sufficiency and quadratic growth for nonisolated minima. *Mathematics of Operations Research*, 20(4):801–817, 1995.
- Nicolas Boumal. *An introduction to optimization on smooth manifolds*. Cambridge University Press, 2023.
- Samuel Burer and Renato DC Monteiro. A nonlinear programming algorithm for solving semidefinite programs via low-rank factorization. *Mathematical Programming*, 95(2):329–357, 2003.
- Samuel Burer and Renato DC Monteiro. Local minima and convergence in low-rank semidefinite programming. *Mathematical programming*, 103(3):427–444, 2005.
- Coralia Cartis, Nicholas IM Gould, and Philippe L Toint. Adaptive cubic regularisation methods for unconstrained optimization. Part I: motivation, convergence and numerical results. *Mathematical Programming*, 127(2):245–295, 2011a.

- Coralia Cartis, Nicholas IM Gould, and Philippe L Toint. Adaptive cubic regularisation methods for unconstrained optimization. Part II: worst-case function-and derivative-evaluation complexity. *Mathematical programming*, 130(2):295–319, 2011b.
- Coralia Cartis, Nicholas IM Gould, and Philippe L Toint. Evaluation complexity of algorithms for nonconvex optimization: Theory, computation and perspectives, 2022.
- Sourav Chatterjee. Convergence of gradient descent for deep neural networks. *arXiv preprint arXiv:2203.16462*, 2022.
- Ralph Chill. The Lojasiewicz–Simon gradient inequality in Hilbert spaces. In *Proceedings of the 5th European-Maghrebian Workshop on Semigroup Theory, Evolution Equations, and Applications (MA Jendoubi, ed.)*, pages 25–36. Citeseer, 2006.
- Francis H Clarke, RJ Stern, and PR Wolenski. Proximal smoothness and the lower-C2 property. *J. Convex Anal.*, 2(1-2):117–144, 1995.
- Andrew R Conn, Nicholas IM Gould, and Philippe L Toint. *Trust region methods*. SIAM, 2000.
- Yaim Cooper. The loss landscape of overparameterized neural networks. *arXiv preprint arXiv:1804.10200*, 2018.
- Damek Davis and Liwei Jiang. A nearly linearly convergent first-order method for nonsmooth functions with quadratic growth. *arXiv preprint arXiv:2205.00064*, 2022.
- Steffen Dereich and Sebastian Kassing. Central limit theorems for stochastic gradient descent with averaging for stable manifolds. *arXiv preprint arXiv:1912.09187*, 2019.
- Steffen Dereich and Sebastian Kassing. Convergence of stochastic gradient descent schemes for Lojasiewicz landscapes. *arXiv preprint arXiv:2102.09385*, 2021.
- Dmitriy Drusvyatskiy and Adrian S Lewis. Error bounds, quadratic growth, and linear convergence of proximal methods. *Mathematics of Operations Research*, 43(3):919–948, 2018.
- Dmitriy Drusvyatskiy, Boris S Mordukhovich, and Tran TA Nghia. Second-order growth, tilt stability, and metric regularity of the subdifferential. *arXiv preprint arXiv:1304.7385*, 2013.
- Dmitriy Drusvyatskiy, Alexander D Ioffe, and Adrian S Lewis. Curves of descent. *SIAM Journal on Control and Optimization*, 53(1):114–138, 2015.
- Dmitriy Drusvyatskiy, Alexander D Ioffe, and Adrian S Lewis. Nonsmooth optimization using Taylor-like models: error bounds, convergence, and termination criteria. *Mathematical Programming*, 185(1):357–383, 2021.
- Jin-yan Fan and Ya-xiang Yuan. On the quadratic convergence of the levenberg-marquardt method without nonsingularity assumption. *Computing*, 74(1):23–39, 2005.
- Paul Feehan. On the Morse–Bott property of analytic functions on Banach spaces with Lojasiewicz exponent one half. *Calculus of Variations and Partial Differential Equations*, 59(2):1–50, 2020.
- Benjamin Fehrman, Benjamin Gess, and Arnulf Jentzen. Convergence rates for the stochastic gradient descent method for non-convex objective functions. *Journal of Machine Learning Research*, 21:136, 2020.
- Pierre Frankel, Guillaume Garrigos, and Juan Peypouquet. Splitting methods with variable metric for Kurdyka–Lojasiewicz functions and general convergence rates. *Journal of Optimization Theory and Applications*, 165(3):874–900, 2015.
- Andreas Griewank. The modification of Newton’s method for unconstrained optimization by bounding cubic terms. Technical report, Technical report NA/12, 1981.

- U. Helmke and J.B. Moore. *Optimization and Dynamical Systems*. Springer, 2nd edition, 1996.
- Aleksandr Davidovich Ioffe. Metric regularity and subdifferential calculus. *Russian Mathematical Surveys*, 55(3):501, 2000.
- Florian Jarre and Philippe L Toint. Simple examples for the failure of Newton’s method with line search for strictly convex minimization. *Mathematical Programming*, 158(1):23–34, 2016.
- Hamed Karimi, Julie Nutini, and Mark Schmidt. Linear convergence of gradient and proximal-gradient methods under the Polyak–Lojasiewicz condition. In *Joint European Conference on Machine Learning and Knowledge Discovery in Databases*, pages 795–811. Springer, 2016.
- Pham Duy Khanh, Boris S Mordukhovich, and Dat Ba Tran. Inexact reduced gradient methods in smooth nonconvex optimization. *arXiv preprint arXiv:2204.01806*, 2022.
- Taehee Ko and Xiantao Li. A local convergence theory for the stochastic gradient descent method in non-convex optimization with non-isolated local minima. *arXiv preprint arXiv:2203.10973*, 2022.
- Krzysztof Kurdyka. On gradients of functions definable in o-minimal structures. In *Annales de l’institut Fourier*, volume 48, pages 769–783, 1998.
- John M Lee. *Introduction to Riemannian manifolds*. Springer, 2018.
- E. Levin, J. Kileel, and N. Boumal. Finding stationary points on bounded-rank matrices: A geometric hurdle and a smooth remedy. *Mathematical Programming*, 2022a. doi: 10.1007/s10107-022-01851-2.
- E. Levin, J. Kileel, and N. Boumal. The effect of smooth parametrizations on nonconvex optimization landscapes. *arXiv 2207.03512*, 2022b.
- Adrian Lewis and Tonghua Tian. Identifiability, the KL property in metric spaces, and subgradient curves, 2022.
- Guoyin Li and Ting Kei Pong. Calculus of the exponent of Kurdyka–Lojasiewicz inequality and its applications to linear convergence of first-order methods. *Foundations of computational mathematics*, 18(5):1199–1232, 2018.
- Shuang Li and Qiuwei Li. Local and global convergence of general burer-monteiro tensor optimizations. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 36, 2022.
- Zhiyuan Li, Tianhao Wang, and Sanjeev Arora. What happens after SGD reaches zero loss?—a mathematical framework. *arXiv preprint arXiv:2110.06914*, 2021.
- Chaoyue Liu, Libin Zhu, and Mikhail Belkin. Loss landscapes and optimization in over-parameterized non-linear systems and neural networks. *Applied and Computational Harmonic Analysis*, 2022.
- Ji Liu and Stephen J Wright. Asynchronous stochastic coordinate descent: Parallelism and convergence properties. *SIAM Journal on Optimization*, 25(1):351–376, 2015.
- Stanislaw Lojasiewicz. Une propriété topologique des sous-ensembles analytiques réels. *Les équations aux dérivées partielles*, 117:87–89, 1963.
- Stanislaw Lojasiewicz. Sur les trajectoires du gradient d’une fonction analytique. *Seminari di geometria*, 1983:115–117, 1982.
- Zhi-Quan Luo and Paul Tseng. Error bounds and convergence analysis of feasible descent methods: a general approach. *Annals of Operations Research*, 46(1):157–178, 1993.
- Ulysse Marteau-Ferey, Francis Bach, and Alessandro Rudi. Second order conditions to decompose smooth functions as sums of squares. *arXiv preprint arXiv:2202.13729*, 2022.

- Saeed Masiha, Saber Salehkaleybar, Niao He, Negar Kiyavash, and Patrick Thiran. Stochastic second-order methods provably beat SGD for gradient-dominated functions. *arXiv preprint arXiv:2205.12856*, 2022.
- Bui Ngoc Muoi and Nguyen Dong Yen. Local stability and local convergence of the basic trust-region method. *Journal of Optimization Theory and Applications*, 172(2):578–593, 2017.
- Ion Necoara and Daniela Lupu. General higher-order majorization-minimization algorithms for (non) convex optimization. *arXiv preprint arXiv:2010.13893*, 2020.
- Ion Necoara, Yu Nesterov, and Francois Glineur. Linear convergence of first order methods for non-strongly convex optimization. *Mathematical Programming*, 175:69–107, 2019.
- Yurii Nesterov and Boris T Polyak. Cubic regularization of Newton method and its global performance. *Mathematical Programming*, 108(1):177–205, 2006.
- Jorge Nocedal and Stephen J Wright. *Numerical Optimization*. Springer New York, 2006.
- Dominikus Noll and Aude Rondepierre. Convergence of linesearch and trust-region methods using the Kurdyka–Lojasiewicz inequality. In *Computational and analytical mathematics*, pages 593–611. Springer, 2013.
- Samet Oymak and Mahdi Soltanolkotabi. Overparameterized nonlinear learning: Gradient descent takes the shortest path? In *International Conference on Machine Learning*, pages 4951–4960. PMLR, 2019.
- Boris T Polyak. Gradient methods for the minimisation of functionals. *USSR Computational Mathematics and Mathematical Physics*, 3(4):864–878, 1963.
- Boris T Polyak. Introduction to optimization. *Optimization Software*, 1987.
- Boris T Polyak, Iliia A Kuruzov, and Fedor S Stonyakin. Stopping rules for gradient methods for non-convex problems with additive noise in gradient. *arXiv preprint arXiv:2205.07544*, 2022.
- Chunhong Qi. Numerical optimization methods on Riemannian manifolds. 2011.
- Yitian Qian and Shaohua Pan. A superlinear convergence iterative framework for Kurdyka–Lojasiewicz optimization and application. *arXiv preprint arXiv:2210.12449*, 2022.
- Alexander Shapiro. Perturbation theory of nonlinear programs when the set of optimal solutions is not a singleton. *Applied Mathematics and Optimization*, 18(1):215–229, 1988.
- Yue Sun, Nicolas Flammarion, and Maryam Fazel. Escaping from saddle points on Riemannian manifolds. *Advances in Neural Information Processing Systems*, 32, 2019.
- Dávid Terjék and Diego González-Sánchez. A framework for overparameterized learning. *arXiv preprint arXiv:2205.13507*, 2022.
- Paul Tseng. Error bounds and superlinear convergence analysis of some newton-type methods in optimization. In *Nonlinear Optimization and Related Topics*, pages 445–462. Springer, 2000.
- Nick Vannieuwenhoven. The condition number of singular subspaces. *ArXiv*, abs/2208.08368, 2022.
- Stephan Wojtowytsch. Stochastic gradient descent with noise of machine learning type. Part I: Discrete time analysis. *arXiv preprint arXiv:2105.01650*, 2021a.
- Stephan Wojtowytsch. Stochastic gradient descent with noise of machine learning type. Part II: Continuous time analysis. *arXiv preprint arXiv:2106.02588*, 2021b.
- Nobuo Yamashita and Masao Fukushima. On the rate of convergence of the levenberg-marquardt method. In *Topics in numerical analysis*, pages 239–249. Springer, 2001.



- Man-Chung Yue, Zirui Zhou, and Anthony Man-Cho So. On the quadratic convergence of the cubic regularization method under a local error bound condition. *SIAM Journal on Optimization*, 29(1):904–932, 2019.
- Pengyun Yue, Cong Fang, and Zhouchen Lin. On the lower bound of minimizing Polyak–Lojasiewicz functions. *arXiv preprint arXiv:2212.13551*, 2022.
- Zhonggang Zeng. A newton’s iteration converges quadratically to nonisolated solutions too. *ArXiv*, abs/2101.09180, 2021.
- Hui Zhang. The restricted strong convexity revisited: analysis of equivalence to error bound and quadratic growth. *Optimization Letters*, 11:817–833, 2017.
- Hui Zhang and Wotao Yin. Gradient methods for convex minimization: better rates under weaker conditions. *arXiv preprint arXiv:1303.4645*, 2013.
- Junyu Zhang and Shuzhong Zhang. A cubic regularized Newton’s method over Riemannian manifolds. *arXiv preprint arXiv:1805.05565*, 2018.
- Richard Y Zhang. Improved global guarantees for the nonconvex Burer–Monteiro factorization via rank overparameterization. *arXiv preprint arXiv:2207.01789*, 2022.
- Yuqian Zhang, Qing Qu, and John Wright. From symmetry to geometry: Tractable nonconvex problems. *ArXiv*, abs/2007.06753, 2020.
- Yi Zhou, Zhe Wang, and Yingbin Liang. Convergence of cubic regularization for nonconvex optimization under KL property. *Advances in Neural Information Processing Systems*, 31, 2018.

## A Łojasiewicz and function growth

We review here the classical arguments at the basis of Proposition 2.2. Given an initial point  $x_0 \in \mathcal{M}$ , we let  $x: I \rightarrow \mathcal{M}$  denote a solution of the negative gradient flow

$$x'(t) = -\nabla f(x(t)) \quad \text{with} \quad x(0) = x_0 \quad (\text{GF})$$

on the maximum interval  $I$ . The following is classical [Łojasiewicz, 1982], restated succinctly to highlight neighborhood assumptions.

**Lemma A.1.** *Suppose  $f$  satisfies (Loja) with constants  $\theta \in [0, 1)$  and  $\mu > 0$  in a neighborhood  $\mathcal{U}$  of  $\bar{x} \in \mathcal{S}$ . Also suppose that  $f(x) > f_{\mathcal{S}}$  for all  $x \in \mathcal{U} \setminus \mathcal{S}$ . Let  $x$  be a solution to (GF) for some  $x_0 \in \mathcal{U} \setminus \mathcal{S}$ . Suppose that for all  $t \in (0, T)$  we have  $x(t) \in \mathcal{U}$  and  $\nabla f(x(t)) \neq 0$ . Then the path length is bounded as*

$$\int_0^T \|x'(t)\| dt \leq \frac{1}{(1-\theta)\sqrt{2\mu}} |f(x_0) - f_{\mathcal{S}}|^{1-\theta}.$$

*Proof.* Define  $h(t) = \frac{1}{(1-\theta)\sqrt{2\mu}} |f(x(t)) - f_{\mathcal{S}}|^{1-\theta}$ . For all  $t \in [0, T)$ , the Łojasiewicz inequality provides:

$$\|x'(t)\| = \|\nabla f(x(t))\| = \frac{\|\nabla f(x(t))\|^2}{\|\nabla f(x(t))\|} \leq \frac{\|\nabla f(x(t))\|^2}{\sqrt{2\mu}(f(x(t)) - f_{\mathcal{S}})^\theta} = -h'(t).$$

It follows that  $h(0) \geq h(0) - h(T) = \int_0^T -h'(t) dt \geq \int_0^T \|x'(t)\| dt$ . □

As shown below, this bound implies that the trajectories are trapped and have a limit point if  $x_0$  is close enough to  $\mathcal{S}$ .

**Proposition A.2** (Lyapunov stability). *Suppose that  $f$  satisfies (Loja) around  $\bar{x} \in \mathcal{S}$  and let  $\mathcal{U}$  be a neighborhood of  $\bar{x}$ . There exists a neighborhood  $\mathcal{V}$  of  $\bar{x}$  such that if  $x_0 \in \mathcal{V}$  then the solution  $x$  to (GF) is defined on  $[0, +\infty)$  and  $x(t) \in \mathcal{U}$  for all  $t \geq 0$ .*

*Proof.* The set  $\mathcal{U}$  contains a ball  $\mathcal{B}$  centered on  $\bar{x}$  of radius  $\delta_1$  such that (i)  $f$  satisfies (Loja) with constants  $\theta$  and  $\mu$  in  $\mathcal{B}$ , and (ii)  $f(y) > f_{\mathcal{S}}$  for all  $y \in \mathcal{B} \setminus \mathcal{S}$ . By continuity of  $f$  there exists an open ball  $\mathcal{V} \subseteq \mathcal{B}$  of radius  $\delta_2$  around  $\bar{x}$  such that for all  $y \in \mathcal{V}$  we have

$$\frac{1}{(1-\theta)\sqrt{2\mu}}|f(y) - f_{\mathcal{S}}|^{1-\theta} + \delta_2 < \delta_1.$$

Given  $x_0 \in \mathcal{V}$ , let  $x: I \rightarrow \mathcal{M}$  be the maximal solution to (GF). Suppose that  $\{t \in I : t \geq 0 \text{ and } x(t) \notin \mathcal{B}\}$  is non-empty and let  $T$  be the infimum of this set. Then for all  $0 \leq t < T$  we have  $x(t) \in \mathcal{B}$ . Suppose first that there exists  $t \in [0, T)$  such that  $\nabla f(x(t)) = 0$ . Then for all  $t' > t$  we have  $x(t') = x(t)$  which is impossible. So for all  $t \in [0, T)$  we have  $\nabla f(x(t)) \neq 0$ . It follows that the assumptions of Lemma A.1 are satisfied and the path length is bounded as

$$\int_0^T \|x'(t)\| dt \leq \frac{1}{(1-\theta)\sqrt{2\mu}}|f(x_0) - f_{\mathcal{S}}|^{1-\theta}.$$

This implies that

$$\text{dist}(x(T), \bar{x}) \leq \int_0^T \|x'(t)\| dt + \text{dist}(x_0, \bar{x}) < \delta_1,$$

and hence that  $x(T) \in \mathcal{B}$ . This is a contradiction and it follows that for all  $t \in I$  we have  $x(t) \in \mathcal{B}$ . Therefore, the total path length of  $x$  is bounded, and the escape lemma [Lee, 2018, Lem. A.43] implies that  $x$  is defined for all  $t \geq 0$ .  $\square$

**Corollary A.3.** *Suppose that  $f$  satisfies (Loja) around  $\bar{x} \in \mathcal{S}$ . There exists a neighborhood  $\mathcal{U}$  of  $\bar{x}$  such that for all  $x_0 \in \mathcal{U}$  the solution to (GF) is defined on  $[0, +\infty)$  and has a limit  $\lim_{t \rightarrow +\infty} x(t) \in \mathcal{S}$ .*

*Proof.* Let  $\mathcal{U}$  be a compact neighborhood of  $\bar{x}$  in which  $f$  satisfies (Loja) and  $f(x) > f_{\mathcal{S}}$  for all  $x \in \mathcal{U} \setminus \mathcal{S}$ . Let  $\mathcal{V}$  be a neighborhood of  $\bar{x}$  as in Proposition A.2 and  $x: [0, +\infty) \rightarrow \mathcal{M}$  be a solution to (GF) with  $x_0 \in \mathcal{V}$ . Then for all  $t \geq 0$  we have  $x(t) \in \mathcal{U}$ . The set  $\mathcal{U}$  is compact so  $x$  has an accumulation point  $x_{\infty} \in \mathcal{U}$ . This point must satisfy  $\nabla f(x_{\infty}) = 0$  so  $x_{\infty} \in \mathcal{S}$ .

Let  $\mathcal{W}$  be a neighborhood of  $x_{\infty}$  and  $\mathcal{V}'$  be a neighborhood as in Proposition A.2. Since  $x_{\infty}$  is an accumulation point of  $x$  there exists a time  $t$  such that  $x(t) \in \mathcal{V}'$ . So for all  $t' \geq t$  we have  $x(t') \in \mathcal{W}$ . This holds for all  $\mathcal{W}$ , and hence  $\lim x(t) = x_{\infty}$ .  $\square$

From this we can deduce that the Łojasiewicz inequality implies the local growth of  $f$  announced in Proposition 2.2.

*Proof of Proposition 2.2.* Let  $\mathcal{U}$  be a neighborhood of  $\bar{x}$  where (Loja) holds with constants  $\mu$  and  $\theta$ . By Corollary A.3 we can find a neighborhood  $\mathcal{V}$  of  $\bar{x}$  such that for all  $x_0 \in \mathcal{V}$  the solution to (GF) is defined on  $[0, \infty)$ , with  $x(t) \rightarrow x_{\infty} \in \mathcal{S}$ . Then, Lemma A.1 provides the first inequality in

$$\frac{1}{(1-\theta)\sqrt{2\mu}}|f(x_0) - f_{\mathcal{S}}|^{1-\theta} \geq \int_0^{\infty} \|x'(t)\| dt \geq \text{dist}(x_0, x_{\infty}) \geq \text{dist}(x_0, \mathcal{S}),$$

which concludes the proof.  $\square$

## B Other Łojasiewicz exponents

We prove the statements of Remark 2.21. From Proposition 2.2, there exists a neighborhood  $\mathcal{U}$  of  $\bar{x}$  and  $c > 0$  such that  $f$  satisfies the growth

$$f(x) - f_{\mathcal{S}} \geq c \text{dist}(x, \mathcal{S})^{\frac{1}{1-\theta}} \tag{29}$$

for all  $x \in \mathcal{U}$ . Assume first that  $f$  is  $C^1$  and  $\nabla f$   $L$ -Lipschitz continuous around  $\bar{x}$ . By [Boumal, 2023, Cor. 10.54], we find that  $f(x) - f_S \leq \frac{L}{2} \text{dist}(x, \mathcal{S})^2$  for all  $x$  sufficiently close to  $\bar{x}$ . This inequality is incompatible with (29) if  $\theta < \frac{1}{2}$  if  $f$  is non-constant around  $\bar{x}$ .

Now assume that  $f$  is  $C^2$  and that  $\nabla^2 f$  is  $L$ -Lipschitz continuous around  $\bar{x} \in \mathcal{S}$ . Let  $h: (y, v) \mapsto \langle v, \nabla^2 f(y)[v] \rangle$ . Lipschitz continuity of  $\nabla^2 f$  gives that [Boumal, 2023, Cor. 10.56]

$$f(\text{Exp}_y(tv)) - f_S - \frac{t^2}{2} h(y, v) \leq \frac{L}{6} t^3$$

for all  $y \in \mathcal{S}$  close enough to  $\bar{x}$ , all unitary  $v \in N_y \mathcal{S}$  and  $t > 0$  small enough. Using Lemma 2.6, we see that  $h(y, v)$  must be positive for this inequality to be compatible with the function growth (29) when  $\theta \in [\frac{1}{2}, \frac{2}{3}]$ . We conclude that there is a compact neighborhood  $\mathcal{V}$  of  $\bar{x}$  such that  $h(y, v) > 0$  for all  $y \in \mathcal{V} \cap \mathcal{S}$  and  $v \in N_y \mathcal{S}$  unitary. The function  $h$  is continuous and the set  $\mathcal{D} = \{(y, v) : y \in \mathcal{V} \cap \mathcal{S}, v \in N_y \mathcal{S} \text{ unitary}\}$  is compact so  $\mu = \inf_{(y,v) \in \mathcal{D}} h(y, v)$  is positive. Now let  $\mathcal{W}$  be a neighborhood of  $\bar{x}$  as in Lemma 1.5 such that for all  $x \in \mathcal{W}$  the projection  $\text{proj}_{\mathcal{S}}(x)$  is included in  $\mathcal{V} \cap \mathcal{S}$ . Then for all  $x \in \mathcal{W}$  we have

$$f(x) - f_S = h(y, v) + o(\|v\|^2) \geq \mu \text{dist}(x, \mathcal{S})^2 + o(\text{dist}(x, \mathcal{S})^2),$$

where  $y \in \text{proj}_{\mathcal{S}}(x)$  and  $v = \text{Log}_y(x)$ . We conclude that QG holds around  $\bar{x}$ , which implies PL, as shown in Section 2.2. Now assume that  $f$  is  $C^3$ . The argument is similar in this case. Let  $y \in \mathcal{S}$  and  $v \in N_y \mathcal{S}$  unitary and assume that  $h(y, v) = 0$ . Then a Taylor expansion gives

$$f(\text{Exp}_x(tv)) - f_S = o(t^3)$$

because the third-order term vanishes. This is incompatible with the function growth (29) when  $y$  is close to  $\bar{x}$  and  $\theta \in [\frac{1}{2}, \frac{2}{3}]$ . So  $h(y, v)$  must be positive and we conclude with the same arguments as above.

## C Lojasiewicz and bounded path length

*Proof of Lemma 3.8.* Let  $\mathcal{U}$  be an open neighborhood of  $\bar{x}$  such that (i)  $f$  satisfies (Loja) with constants  $\theta \in [0, 1)$  and  $\mu > 0$  in  $\mathcal{U}$ , (ii)  $f(x) \geq f_S$  for all  $x \in \mathcal{U}$  and (iii) condition (5) holds in  $\mathcal{U}$ . For all  $x \in \mathcal{U}$  we have

$$0 \leq |f(x) - f_S|^{2\theta} \leq \frac{1}{2\mu} \|\nabla f(x)\|^2 \quad \text{so} \quad \|\nabla f(x)\| \geq \sqrt{2\mu} |f(x) - f_S|^\theta.$$

Let  $K \geq 0$  and suppose that for all  $0 \leq k \leq K$  we have  $x_k \in \mathcal{U}$ . Given  $k \in \mathbb{N}$ , either  $\nabla f(x_k) = 0$  and  $\text{dist}(x_k, x_{k+1}) = 0$ , or  $\nabla f(x_k) \neq 0$  and  $f(x_k) > f_S$ . In this case combining the inequality  $\|\nabla f(x)\| \geq \sqrt{2\mu} |f(x) - f_S|^\theta$  above with the strong decrease condition in (5) we find that

$$\text{dist}(x_k, x_{k+1}) \leq \frac{1}{\sigma \sqrt{2\mu}} \frac{f(x_k) - f(x_{k+1})}{|f(x_k) - f_S|^\theta} \leq \frac{1}{\sigma(1-\theta)\sqrt{2\mu}} ((f(x_k) - f_S)^{1-\theta} - (f(x_{k+1}) - f_S)^{1-\theta}),$$

where we used

$$\begin{aligned} \frac{f(x_k) - f(x_{k+1})}{|f(x_k) - f_S|^\theta} &= \int_{f(x_{k+1})}^{f(x_k)} \frac{1}{|f(x) - f_S|^\theta} dt \\ &\leq \int_{f(x_{k+1})}^{f(x_k)} \frac{1}{|t - f_S|^\theta} dt \\ &= \frac{1}{1-\theta} ((f(x_k) - f_S)^{1-\theta} - (f(x_{k+1}) - f_S)^{1-\theta}). \end{aligned}$$

We get the result as we sum this inequality over  $0, \dots, K-1$ .  $\square$

## D Łojasiewicz and convergence rate

*Proof of Proposition 3.9.* Let  $\mathcal{U}$  be a neighborhood where PL is satisfied: for all large enough  $k$  we have

$$f(x_{k+1}) - f(x_k) \leq -\omega \|\nabla f(x_k)\|^2 \leq -2\mu\omega(f(x_k) - f_{\mathcal{S}})$$

where the first inequality is the sufficient decrease 6 and the second one is local PL. By adding  $f(x_k) - f_{\mathcal{S}}$  we get

$$f(x_{k+1}) - f_{\mathcal{S}} \leq (1 - 2\omega\mu)(f(x_k) - f_{\mathcal{S}}).$$

We now prove the rate for  $\{\|\nabla f(x_k)\|\}$  and  $\{\text{dist}(x_k, \mathcal{S})\}$ . For all large enough  $k$  we have

$$\omega \|\nabla f(x_k)\|^2 \leq f(x_k) - f(x_{k+1}) \leq f(x_k) - f_{\mathcal{S}} \quad \text{so} \quad \|\nabla f(x_k)\| \leq \sqrt{\frac{1}{\omega}(f(x_k) - f_{\mathcal{S}})}.$$

From Proposition 3.9 we get that  $\{\|\nabla f(x_k)\|\}$  converges linearly to zero with rate  $\sqrt{1 - 2\omega\mu}$ . We now prove the convergence rate for  $\{\text{dist}(x_k, \mathcal{S})\}$  using the local quadratic growth. From Proposition 2.2 there exists a neighborhood  $\mathcal{U}$  of  $\bar{x}$  such that for all  $x \in \mathcal{U}$  we have

$$f(x) - f_{\mathcal{S}} \geq \frac{\mu}{2} \text{dist}(x, \mathcal{S})^2 \quad \text{so} \quad \text{dist}(x_k, \mathcal{S}) \leq \sqrt{\frac{2}{\mu}(f(x_k) - f_{\mathcal{S}})}.$$

We conclude that  $\text{dist}(x_k, \mathcal{S})$  converges linearly to zero with rate  $\sqrt{1 - 2\omega\mu}$ .  $\square$

*Proof of Proposition 3.10.* When  $x_k$  is sufficiently close to  $\bar{x}$  the bound (7) implies

$$f(x_k) - f(x_{k+1}) = f(x_k) - f(\mathbb{R}_x(s_k)) \geq -\left(\langle \nabla f(x_k), s_k \rangle + \frac{L}{2} \|s_k\|^2\right). \quad (30)$$

By plugging  $s_k = -\gamma \nabla f(x_k)$  we obtain  $f(x_k) - f(x_{k+1}) = (1 - \frac{L}{2}\gamma) \|\nabla f(x_k)\| \|s_k\|$ . The constraint  $\gamma \in (0, \frac{2}{L})$  implies that  $1 - \frac{L}{2}\gamma > 0$  so this algorithm satisfies the strong decrease property (5) around  $\bar{x}$  with constant  $\sigma = \frac{1}{c_r}(1 - \frac{L}{2}\gamma)$ . Lemma 3.8 implies that it also satisfies the (BPL). By Corollary 3.7 we find that there exists a neighborhood  $\mathcal{V}$  of  $\bar{x}$  such that if an iterate enters  $\mathcal{V}$  then the sequence converges to some  $x_{\infty} \in \mathcal{U} \cap \mathcal{S}$ .

Now assume that this is the case and that  $\mu$ -(PL) and (6) hold around  $x_{\infty}$  (we can always ensure this by shrinking  $\mathcal{V}$ ). Equation (30) gives  $f(x_k) - f(x_{k+1}) \geq (\gamma - \frac{L}{2}\gamma^2) \|\nabla f(x_k)\|^2$ . The number  $\gamma - \frac{L}{2}\gamma^2$  is positive because  $\gamma \in (0, \frac{2}{L})$ . It follows that the sequence satisfies the sufficient decrease (6) with  $\omega = \gamma - \frac{L}{2}\gamma^2$ . We conclude with Proposition 3.9.  $\square$

Using performance estimation, it is possible to derive sharper convergence rates for gradient descent with constant step-sizes under the PL assumption [Abbaszadehpeivasti et al., 2022, Thm. 3]. The Lipschitz-type property (7) also implies the sufficient decrease (6) for gradient descent with backtracking line-search: see [Boumal, 2023, Lem. 4.12] for the Riemannian case.

**Proposition D.1.** *Suppose that  $f$  and  $\mathbb{R}$  satisfy (7) around  $\bar{x} \in \mathcal{S}$ . Let  $\{x_k\}$  be a sequence of iterates generated by gradient descent with Armijo backtracking line-search converging to  $\bar{x}$ . Then  $\{x_k\}$  satisfies (6) with*

$$\omega = \sigma \min\left(\bar{\alpha}, \frac{2\beta(1 - \sigma)}{L}\right).$$

*Proof.* See [Boumal, 2023, Lem. 4.12].  $\square$

See also [Khanh et al., 2022] for more exhaustive results on line-search.