

ASYMPTOTIC TOTAL ERGODICITY FOR ACTIONS OF $\mathbb{F}[t]$ AND POLYNOMIAL CONFIGURATIONS OVER FINITE FIELDS AND RINGS

ETHAN ACKELSBURG AND VITALY BERGELSON

ABSTRACT. We obtain new combinatorial results about polynomial configurations over finite fields and rings by utilizing the phenomenon of *asymptotic total ergodicity* (previously studied for actions of \mathbb{Z} on modular rings $\mathbb{Z}/N\mathbb{Z}$ in [BB23]) in the context of actions of the polynomial ring $\mathbb{F}[t]$ over a finite field \mathbb{F} . We prove that a sequence of quotient rings $\mathbb{F}[t]/Q_n(t)\mathbb{F}[t]$, $Q_n(t) \in \mathbb{F}[t]$, is asymptotically totally ergodic if and only if the minimal degree of the irreducible factors of Q_n diverges to infinity as $n \rightarrow \infty$. We then show that asymptotic total ergodicity of a sequence of quotient rings $\mathbb{F}[t]/Q_n(t)\mathbb{F}[t]$ leads to asymptotic equidistribution of polynomial sequences in subgroups of $(\mathbb{F}[t]/Q_n(t)\mathbb{F}[t], +)$. This has several combinatorial consequences:

- (1) We prove a power saving bound for the Furstenberg–Sárközy theorem over finite fields of fixed characteristic: given an *intersective* polynomial $P(x) \in (\mathbb{F}_p[t])[x]$, there exists $\gamma = \gamma(P) > 0$ such that if $A \subseteq \mathbb{F}_q$, $q = p^n$, does not contain distinct $a, b \in A$ with $b - a = P(y)$ for some $y \in \mathbb{F}_q$, then $A \ll_P q^{1-\gamma}$. This complements recent work of Li and Sauermaun [LS22], where they obtain power saving bounds under the assumption $P(0) = 0$.
- (2) Under a natural equidistribution condition on $P(x) \in (\mathbb{F}[t])[x]$, we prove the following enhancement of the Furstenberg–Sárközy theorem in the presence of asymptotic total ergodicity. Suppose the sequence of quotient rings $(\mathbb{F}[t]/Q_n(t)\mathbb{F}[t])_{n \in \mathbb{N}}$ is asymptotically totally ergodic. Then for any $\delta > 0$ and any two sets $A, B \subseteq \mathbb{F}[t]/Q_n(t)\mathbb{F}[t]$ with $|A||B| \geq \delta |\mathbb{F}[t]/Q_n(t)\mathbb{F}[t]|^2$ and n sufficiently large, there exist $x, y \in \mathbb{F}[t]/Q_n(t)\mathbb{F}[t]$ with $x \in A$ and $x + P(y) \in B$. Moreover, the values $x, y \in \mathbb{F}[t]/Q_n(t)\mathbb{F}[t]$ for which $x \in A$ and $x + P(y) \in B$ obey the “correct” statistical behavior as $n \rightarrow \infty$:

$$\frac{|\{(x, y) \in (\mathbb{F}[t]/Q_n(t)\mathbb{F}[t])^2 : x \in A, x + P(y) \in B\}|}{|\mathbb{F}[t]/Q_n(t)\mathbb{F}[t]|^2} = \frac{|A||B|}{|\mathbb{F}[t]/Q_n(t)\mathbb{F}[t]|^2} + o_{n \rightarrow \infty}(1).$$

We also show that, in the absence of asymptotic total ergodicity and an equidistribution condition on the polynomial P , one cannot hope for such a refinement of the Furstenberg–Sárközy theorem.

- (3) We establish partition regularity of families of polynomial equations over finite fields. For example, we are able to prove: if $P(x) \in \mathbb{Z}[x]$ with $P(0) = 0$, then for any $r \in \mathbb{N}$, there exists $N = N(P, r)$ and $c = c(P, r) > 0$ such that if $q > N$ and $\mathbb{F}_q = \bigcup_{i=1}^r C_i$, then there are at least cq^2 monochromatic solutions to the equation $P(x) + P(y) = P(z)$.

Interactions between finitary mathematics (combinatorics in finite fields and rings) and infinitary mathematics (ergodic theory, equidistribution, and Loeb measure spaces) play a central role throughout the paper.

CONTENTS

1. Introduction	2
2. Asymptotic total ergodicity	11
3. Asymptotic projection theorem	12
4. Power saving bounds for the Furstenberg–Sárközy theorem in characteristic p	17
5. Proof of equivalences	18

Date: September 12, 2023.

2020 Mathematics Subject Classification. 11B30 (11T06, 37A25, 05D10).

Key words and phrases. Finite fields, Furstenberg–Sárközy theorem, total ergodicity, equidistribution, partition regularity, Loeb measure.

6. Partition regularity of polynomial equations over finite fields	22
Acknowledgements	30
References	30

1. INTRODUCTION

Our starting point is the following classical result:

Theorem 1.1 (Furstenberg–Sárközy theorem [F77, S78]). *Let $P(x) \in \mathbb{Z}[x]$ be a nonzero polynomial with $P(0) = 0$. For any $\delta \in (0, 1)$, there exists $N = N(P, \delta) \in \mathbb{N}$ such that any subset $A \subseteq \{1, \dots, N\}$ of size $|A| \geq \delta N$ contains distinct elements $a, b \in A$ with $b - a = P(n)$ for some $n \in \mathbb{Z}$.*

The assumption that $P(0) = 0$ can be weakened as follows. Call a polynomial P *intersective* if P has a root mod q for every $q \in \mathbb{N}$. It follows from the work of Kamae and Mendès France [KM78, Example 3] that the conclusion of Theorem 1.1 holds if and only if P is intersective.

A version of the Furstenberg–Sárközy theorem holds also in a finite characteristic setting. The following result is a consequence of [BL16, Theorem 9.2] together with (a corrected version of) the remark¹ following Theorem 9.5 in [BL16]:

Theorem 1.2. *Let \mathbb{F} be a finite field with q elements. The following are equivalent for a polynomial $P(x) \in (\mathbb{F}[t])[x]$:*

- (i) P is intersective: for any $Q(t) \in \mathbb{F}[t] \setminus \{0\}$, there exists $n \in \mathbb{F}[t]$ such that $P(n) \equiv 0 \pmod{Q}$;
- (ii) for any $\delta \in (0, 1)$, there exists $N = N(P, \delta) \in \mathbb{N}$ such that any subset $A \subseteq \mathbb{F}[t]$ of elements of degree $< N$ with $|A| \geq \delta q^N$ contains distinct $a, b \in A$ with $b - a = P(n)$ for some $n \in \mathbb{F}[t]$.

A consequence of Theorem 1.2 is a Furstenberg–Sárközy type theorem over large finite fields:

Corollary 1.3. *Let p be a prime number, and let $P(x) \in (\mathbb{F}_p[t])[x]$ be an intersective polynomial. Then for any $\delta \in (0, 1)$, there exists $N = N(P, \delta) \in \mathbb{N}$ such that if $k \geq N$ and $A \subseteq \mathbb{F}_{p^k}$ with $|A| \geq \delta p^k$, then A contains distinct $a, b \in A$ with $b - a = P(n)$ for some $n \in \mathbb{F}_{p^k}$.*

Recent work of Li and Sauermann [LS22], building on earlier work of Green [G17] using the Croot–Lev–Pach [CLP17] polynomial method, establishes quantitative improvements of Theorem 1.2 and Corollary 1.3 under the assumption $P(0) = 0$.

Theorem 1.4 ([LS22], Theorem 1.4 and Corollary 1.5). *For $q, d \in \mathbb{N}$, let*

$$t_{q,d} = \inf_{0 < x < 1} \frac{1 + x + \dots + x^{q-1}}{x^{\frac{1}{2}(q-1)(1-1/(dd'))}},$$

where

$$d' = \min\{d, (q-1)(1 + \log_q d)\}.$$

- (i) Fix q and a polynomial $P(x) \in \mathbb{F}_q[x]$ of degree d with $P(0) = 0$. If $A \subseteq \mathbb{F}_q[t]$ is a set of polynomials of degree less than N and A does not contain distinct $a, b \in A$ with $b - a = P(x)$

¹In the remark following Theorem 9.5 in [BL16], intersective polynomials are defined as polynomials $P(x) \in (\mathbb{F}[t])[x]$ such that for any finite index subgroup $\Lambda \leq (\mathbb{F}[t], +)$, there exists $m \in \mathbb{F}[t]$ such that $P(mn) \in \Lambda$ for every $n \in \Lambda$. The definition of intersective given in item (i) of Theorem 1.2 is different and deals with a wider class of polynomials but is the correct notion in order to get the desired “if and only if” conclusion. An example of an intersective polynomial that does not fit the condition in [BL16] is $P(x) = x + t$. There is no multiple $m \in \mathbb{F}[t]$ for which $P(mn)$ always belongs to the subgroup $t^2\mathbb{F}[t]$ of index $|\mathbb{F}|^2$. However, $P(-t) = 0$, so P is intersective (according to our definition).

for some $x \in \mathbb{F}_q[t]$, then²

$$|A| \ll_{q,d} t_{q,d}^N.$$

(ii) Fix a prime p and a polynomial $P(x) \in \mathbb{F}_p[x]$ of degree d with $P(0) = 0$. If $A \subseteq \mathbb{F}_{p^k}$ does not contain distinct $a, b \in A$ with $b - a = P(x)$ for some $x \in \mathbb{F}_{p^k}$, then

$$|A| \ll_{p,d} t_{p,d}^k.$$

In this paper, we produce power saving bounds for the Furstenberg–Sárközy theorem over finite fields by a different method. The bounds we obtain are different from those in Theorem 1.4. In some cases, our bounds are stronger, while in other cases, ours are weaker; see Remark 1.13 below. Our approach draws inspiration from infinitary sources. These are: equidistributional results for polynomial sequences defined over $\mathbb{F}[t]$ and the phenomenon of *asymptotic total ergodicity* for actions of $\mathbb{F}[t]$. As a consequence of our approach, our results apply in a more general setting than finite fields of characteristic p , including quotient rings $\mathbb{F}[t]/Q(t)\mathbb{F}[t]$ under some conditions on $Q(t) \in \mathbb{F}[t]$. The phenomenon of asymptotic total ergodicity for \mathbb{Z} -actions was previously explored in [BB23], where similar Furstenberg–Sárközy-type results are proved in the setting of modular rings $\mathbb{Z}/N\mathbb{Z}$ when all prime factors of N are sufficiently large. The results of this paper are natural analogues of the results in [BB23]. Where appropriate, we note the correspondences between our setting (dealing with $\mathbb{F}[t]$ -actions and quotient rings $\mathbb{F}[t]/Q(t)\mathbb{F}[t]$) and the more familiar setting of \mathbb{Z} -actions and modular rings $\mathbb{Z}/N\mathbb{Z}$. However, some caution is needed, as our finite characteristic setting introduces new complications. Namely, the distributional behavior of a polynomial whose degree d exceeds the characteristic p is more sophisticated than the behavior of polynomials over \mathbb{Z} (see, e.g., Theorem 1.9 below), and this creates additional difficulties in our analysis.

Before stating our results, we fix some notation. Let \mathbb{F} be a finite field of characteristic p . We denote the set of monic polynomials over \mathbb{F} by $\mathbb{F}[t]^+$. Every element $Q(t) \in \mathbb{F}[t]^+$ has a unique factorization (up to reordering) into monic irreducibles $Q(t) = Q_1(t)^{s_1} \dots Q_r(t)^{s_r}$. We denote the quotient ring $\mathbb{F}[t]/Q(t)\mathbb{F}[t]$ by $\mathbb{F}[t]_Q$, and we have an isomorphism

$$(1.1) \quad \mathbb{F}[t]_Q \cong \mathbb{F}[t]_{Q_1^{s_1}} \times \dots \times \mathbb{F}[t]_{Q_r^{s_r}}$$

by the Chinese remainder theorem. Note that when Q is irreducible, $\mathbb{F}[t]_Q$ is a finite field of characteristic p . Moreover, any finite field of characteristic p can be obtained as $\mathbb{F}_p[t]_Q$ for some irreducible element $Q(t) \in \mathbb{F}_p[t]^+$. The decomposition of the ring $\mathbb{F}[t]_Q$ for general $Q(t) \in \mathbb{F}[t]^+$ given in (1.1) is parallel to the situation with modular rings, where the Chinese remainder theorem gives an isomorphism

$$\mathbb{Z}/N\mathbb{Z} \cong \mathbb{Z}/p_1^{s_1}\mathbb{Z} \times \dots \times \mathbb{Z}/p_r^{s_r}\mathbb{Z}$$

for $N = p_1^{s_1} \dots p_r^{s_r}$.

We define an absolute value on $\mathbb{F}[t]$ by $|c_d t^d + \dots + c_1 t + c_0| = |\mathbb{F}|^d$ if $c_d \neq 0$. Equivalently, for any $Q(t) \in \mathbb{F}[t]$, $|Q|$ is the cardinality of the quotient ring $\mathbb{F}[t]_Q$. For $Q = Q_1^{s_1} \dots Q_r^{s_r} \in \mathbb{F}[t]^+$, we set $\text{lpf}(Q) = \min_{1 \leq i \leq r} |Q_i|$ to be the size of the least prime factor of Q .

For a finite set S and a function $f : S \rightarrow \mathbb{C}$, we write

$$\mathbb{E}_{x \in S} f(x) = \frac{1}{|S|} \sum_{x \in S} f(x).$$

For $r \geq 1$, we define the L^r -norm on S by

$$\|f\|_{L^r(S)} := \left(\mathbb{E}_{x \in S} |f(x)|^r \right)^{1/r}.$$

²The notation $a(N) \ll b(N)$ means that there is a constant $C > 0$ such that $|a(N)| \leq C|b(N)|$ for all $N \in \mathbb{N}$. Subscripts on \ll denote on which parameters the constant C depends.

Recall that a measure-preserving \mathbb{Z} -system³ (X, \mathcal{X}, μ, T) is *totally ergodic* if for every $m \in \mathbb{Z} \setminus \{0\}$, T^m is ergodic. By analogy, we say that a measure-preserving $\mathbb{F}[t]$ -system $(X, \mathcal{X}, \mu, (T_n)_{n \in \mathbb{F}[t]})$ is *totally ergodic* if for every $m \in \mathbb{F}[t] \setminus \{0\}$, the action $(T_{mn})_{n \in \mathbb{F}[t]}$ is ergodic. Our first result provides a finitization of the phenomenon of total ergodicity. A similar result for \mathbb{Z} -actions and the quotient rings $\mathbb{Z}/N\mathbb{Z}$ appears in [BB23].

Theorem 1.5. *Let $(Q_n)_{n \in \mathbb{N}}$ be a sequence in $\mathbb{F}[t]^+$. The following are equivalent:*

(i) *The sequence of quotient rings $\mathbb{F}[t]_{Q_n}$ is asymptotically totally ergodic: for any $m \in \mathbb{F}[t] \setminus \{0\}$,*

$$\sup_{f_n: \mathbb{F}[t]_{Q_n} \rightarrow \mathbb{D}} \left\| \mathbb{E}_{y \in \mathbb{F}[t]_{Q_n}} f_n(x + my) - \mathbb{E}_{z \in \mathbb{F}[t]_{Q_n}} f_n(z) \right\|_{L^2(\mathbb{F}[t]_{Q_n})} \xrightarrow{n \rightarrow \infty} 0.$$

(ii) $\text{lpf}(Q_n) \rightarrow \infty$.

We prove Theorem 1.5 in Section 2.

Using the spectral theorem for unitary actions of $\mathbb{F}[t]$ and equidistributional results for polynomial sequences over $\mathbb{F}[t]$, one may show the following:

Theorem 1.6. *Let $P(y) \in (\mathbb{F}[t])[y]$ be a polynomial with zero constant term. Then for any totally ergodic system $(X, \mathcal{X}, \mu, (T_n)_{n \in \mathbb{F}[t]})$, any Følner sequence⁴ $(\Phi_N)_{N \in \mathbb{N}}$, and any $f \in L^2(\mu)$,*

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \in \Phi_N} T_{P(n)} f = \pi_P(f),$$

where π_P is the orthogonal projection onto the space

$$\{g \in L^2(\mu) : T_{P(n)} g = g \text{ for all } n \in \mathbb{F}[t]\}.$$

Remark 1.7. (1) A similar result with $\mathbb{F}[t]$ replaced by a countably infinite field was obtained by Larick in [L98, Theorem 1.1.1].

(2) For \mathbb{Z} -actions, the corresponding version of Theorem 1.6 is simpler. Namely, for any polynomial $P(y) \in \mathbb{Z}[y]$, any totally ergodic \mathbb{Z} -system (X, \mathcal{X}, μ, T) , any Følner sequence $(\Phi_N)_{N \in \mathbb{N}}$ in \mathbb{Z} , and any $f \in L^2(\mu)$,

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \in \Phi_N} T^{P(n)} f = \int_X f d\mu.$$

The presence of the projection π_P in Theorem 1.6 rather than $\int_X f d\mu$ is a reflection of the more intricate distributional behavior of polynomials over $\mathbb{F}[t]$. The situation where the projection $\pi_P f$ is equal to $\int_X f d\mu$ can be characterized by an equidistributional assumption on the polynomial $P(y) \in (\mathbb{F}[t])[y]$; see Proposition 1.14 below.

Our goal is to prove an asymptotic version of Theorem 1.6 and to deduce from it new combinatorial results over finite fields (and rings of the form $\mathbb{F}[t]_Q$). Before formulating our result, we sketch a proof of Theorem 1.6, which will serve as a model for our finitary results.

By the spectral theorem for actions of $\mathbb{F}[t]$ by unitary operators on a Hilbert space, we may work with the Hilbert space $\mathcal{H} = L^2(\widehat{\mathbb{F}[t]}, \sigma)$, where σ is a positive Borel measure on the dual group

³Given an abelian group Γ , a *measure-preserving Γ -system* is a quadruple $(X, \mathcal{X}, \mu, (T_g)_{g \in \Gamma})$, where (X, \mathcal{X}, μ) is a probability space, and $(T_g)_{g \in \Gamma}$ is an action of Γ on (X, \mathcal{X}, μ) by measure-preserving transformations.

⁴A *Følner sequence* in $\mathbb{F}_p[t]$ is a sequence $(\Phi_N)_{N \in \mathbb{N}}$ of finite subsets of $\mathbb{F}_p[t]$ such that, for any $n \in \mathbb{F}_p[t]$,

$$\lim_{N \rightarrow \infty} \frac{|(\Phi_N + n) \Delta \Phi_N|}{|\Phi_N|} = 0.$$

Examples include $\Phi_N = \{c_{N-1}t^{N-1} + \cdots + c_1t + c_0 : c_i \in \mathbb{F}_p\}$ (the set of all polynomials over \mathbb{F}_p of degree $< N$) and $\Phi'_N = \{t^N + c_{N-1}t^{N-1} + \cdots + c_1t + c_0 : c_i \in \mathbb{F}_p\} = \Phi_N + t^N$ (the set of all monic polynomials of degree N).

$\widehat{\mathbb{F}[t]}$, and the unitary action $(T_n)_{n \in \mathbb{F}[t]}$ is represented by the multiplication operators $(U_n h)(\chi) = \chi(n)h(\chi)$ for $h \in \mathcal{H}$ and $\chi \in \widehat{\mathbb{F}[t]}$.

Rather than working with $\widehat{\mathbb{F}[t]}$ as the abstract dual group of $\mathbb{F}[t]$, it will be convenient to work with the dual group in a more concrete form. Let $\mathbb{F}(t)$ be the field of rational functions $\mathbb{F}(t) = \{ \frac{m}{n} : m, n \in \mathbb{F}[t], n \neq 0 \}$. Extending the absolute value we defined on $\mathbb{F}[t]$ to $\mathbb{F}(t)$ by $|\frac{m}{n}| = \frac{|m|}{|n|}$, the completion of $\mathbb{F}(t)$ is the field $\mathbb{F}((t^{-1})) = \left\{ \sum_{j=-\infty}^N c_j t^j : N \in \mathbb{Z}, c_j \in \mathbb{F} \right\}$. We think of $\mathbb{F}(t)$ and $\mathbb{F}((t^{-1}))$ as natural analogues of the rational numbers \mathbb{Q} and the real numbers \mathbb{R} , respectively.

In the characteristic zero setting, the dual group of the integers is isomorphic to the torus $\mathbb{T} = \mathbb{R}/\mathbb{Z}$. A similar result is true in our setting: the dual group $\widehat{\mathbb{F}[t]}$ is isomorphic to $\mathbb{F}((t^{-1}))/\mathbb{F}[t]$. In particular, every character $\chi : \mathbb{F}[t] \rightarrow \mathbb{C}$ takes the form

$$\chi(n) = e(nx)$$

for some $x \in \mathbb{F}((t^{-1}))/\mathbb{F}[t]$, where

$$e \left(\sum_{j=-\infty}^N c_j t^j \right) = \chi_0(c_{-1})$$

and χ_0 is a fixed nontrivial character on \mathbb{F} .

A word of caution: with the objects discussed above, we have an isomorphism $\mathbb{F}((t^{-1})) \cong \mathbb{F}[t] \oplus (\mathbb{F}((t^{-1}))/\mathbb{F}[t])$. The corresponding statement in the more familiar characteristic zero setting is not true: $\mathbb{R} \not\cong \mathbb{Z} \oplus \mathbb{T}$.

To prove Theorem 1.6, it then suffices to show: for σ -a.e. $x \in \mathbb{F}((t^{-1}))/\mathbb{F}[t]$,

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \in \Phi_N} e(P(n)x) = \begin{cases} 1, & \text{if } e(P(n)x) = 1 \text{ for every } n \in \mathbb{F}[t]; \\ 0, & \text{otherwise.} \end{cases}$$

For \mathbb{Z} -action, total ergodicity is equivalent to the absence of rational spectrum. Similarly, the assumption that $(T_n)_{n \in \mathbb{N}}$ is a totally ergodic $\mathbb{F}[t]$ -action means that

$$\sigma((\mathbb{F}(t)/\mathbb{F}[t]) \setminus \{0\}) = 0$$

Therefore, Theorem 1.6 reduces to studying equidistribution of the sequences $(P(n)\alpha)_{n \in \mathbb{F}[t]}$ for irrational $\alpha \in \mathbb{F}((t^{-1})) \setminus \mathbb{F}(t)$.

A general Weyl-type equidistribution theorem for polynomials over $\mathbb{F}[t]$ was established in [BL16], and we can use the result to finish the proof of Theorem 1.6. First, we need some definitions for polynomials in finite characteristic:

Definition 1.8.

- (1) A polynomial $P(y) \in \mathbb{F}((t^{-1}))[y]$ is called *separable* if $P(y) = a_0 + \sum_{i=1}^k a_i y^{r_i}$ and $p \nmid r_i$ for $i \in \{1, \dots, k\}$.
- (2) A polynomial $\eta(y) \in \mathbb{F}((t^{-1}))[y]$ is *additive* if $\eta(x+y) = \eta(x) + \eta(y)$ for any $x, y \in \mathbb{F}((t^{-1}))$.
- (3) For $u \in \mathbb{F}((t^{-1}))$ and $f : \mathbb{F}((t^{-1})) \rightarrow \mathbb{F}((t^{-1}))$, define the differencing operator $\partial_u f(x) = f(x+u) - f(x)$. Then define $\partial_{u_1, \dots, u_k}$ recursively by $\partial_{u_1, \dots, u_k} = \partial_{u_k} \partial_{u_1, \dots, u_{k-1}}$. The *derivational degree* (abbreviated d-deg) of a polynomial $P(y) \in \mathbb{F}((t^{-1}))[y]$ is the minimum $d \geq 0$ such that $\partial_{u_1, \dots, u_{d+1}} P(y) = 0$ for any $u_1, \dots, u_{d+1}, y \in \mathbb{F}((t^{-1}))$.

Note that $\text{d-deg } y^p = 1$, since $(u+v)^p = u^p + v^p$ in characteristic p . More generally, for $r = \sum_{i=0}^n a_i p^i$ with $a_i \in \{0, \dots, p-1\}$, we have $\text{d-deg } y^r = \sum_{i=0}^n a_i$. Any polynomial can be written in the form $P(y) = a_0 + \sum_{i=1}^n \eta_i(y^{r_i})$, where $a_0 \in \mathbb{F}((t^{-1}))$, $\eta_1, \dots, \eta_n \in \mathbb{F}((t^{-1}))[y]$ are additive polynomials, and y^{r_i} are distinct separable monomials.

We say that $a : \mathbb{F}_p[t] \rightarrow \mathbb{F}_p((t^{-1}))$ is *well-distributed mod* $\mathbb{F}_p[t]$ if

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \in \Phi_N} f(a(n)) = \int_{\mathbb{F}_p((t^{-1}))/\mathbb{F}_p[t]} f \, dm$$

for every continuous function $f : \mathbb{F}_p((t^{-1}))/\mathbb{F}_p[t] \rightarrow \mathbb{C}$ and every Følner sequence $(\Phi_N)_{N \in \mathbb{N}}$ in $\mathbb{F}_p[t]$. A more refined notion of equidistribution is as follows. A function $a : \mathbb{F}_p[t] \rightarrow \mathbb{F}_p((t^{-1}))$ is *well-distributed mod* $\mathbb{F}_p[t]$ in a subgroup $H \subseteq \mathbb{F}_p((t^{-1}))/\mathbb{F}_p[t]$ if

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \in \Phi_N} f(a(n)) = \int_H f \, dm_H$$

for every continuous function $f : \mathbb{F}_p((t^{-1}))/\mathbb{F}_p[t] \rightarrow \mathbb{C}$ and every Følner sequence $(\Phi_N)_{N \in \mathbb{N}}$ in $\mathbb{F}_p[t]$. For a subgroup H and a finite set $F \subseteq \mathbb{F}_p((t^{-1}))/\mathbb{F}_p[t]$, we say that $a : \mathbb{F}_p[t] \rightarrow \mathbb{F}_p((t^{-1}))$ is *well-distributed in the components of* $H + F$ if there exists $m \in \mathbb{F}_p[t] \setminus \{0\}$ such that, for every $k \in \mathbb{F}_p[t]$, the sequence $(a(mn + k))_{n \in \mathbb{F}_p[t]}$ is well-distributed in $H + x$ for some $x \in F$.

Theorem 1.9 ([BL16], Theorem 0.3). *An additive polynomial $\eta(y) \in (\mathbb{F}((t^{-1})))[y]$ is well distributed in the subgroup⁵ $\overline{\eta(\mathbb{F}[t])} = \mathcal{F}(\eta) + \eta(K)$, where $K \subseteq \mathbb{F}((t^{-1}))/\mathbb{F}[t]$ is a finite subgroup. For any polynomial $P(y) = \alpha_0 + \sum_{i=1}^n \eta_i(y^{r^i})$, the orbit closure $\mathcal{O}(P) = \overline{P(\mathbb{F}[t])}$ is of the form $\mathcal{F}(P) + P(K)$, where $\mathcal{F}(P) = \sum_{i=1}^n \mathcal{F}(\eta_i)$ and K is a finite subset of $\mathbb{F}[t]$, and $P(y)$ is well-distributed in the components $\mathcal{F}(P) + P(k)$, $k \in K$.*

For an additive polynomial $\eta(y) \in \mathbb{F}[t]$ and irrational $\alpha \in \mathbb{F}((t^{-1})) \setminus \mathbb{F}(t)$, the orbit closure $\{\eta(y)\alpha : y \in \mathbb{F}[t]\}$ is equal to the subtorus $\mathcal{F}(\eta\alpha)$ rather than a union of finitely many shifts of $\mathcal{F}(\eta\alpha)$. It follows that for $P(y) \in \mathbb{F}[t]$ with $P(0) = 0$ and $\alpha \in \mathbb{F}((t^{-1})) \setminus \mathbb{F}(t)$, the sequence $(P(n)\alpha)_{n \in \mathbb{F}[t]}$ is well-distributed in the subtorus $\mathcal{F}(P)$; see [BL16, Theorem 8.1] for more details. Thus, for any $\alpha \in \mathbb{F}((t^{-1})) \setminus \mathbb{F}(t)$,

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \in \Phi_N} e(P(n)\alpha) = \begin{cases} 1, & \text{if } e(P(n)\alpha) = 1 \text{ for every } n \in \mathbb{F}[t]; \\ 0, & \text{otherwise.} \end{cases}$$

This completes the proof of Theorem 1.6.

We can now state our main result, which is an asymptotic version of Theorem 1.6 with quantitative bounds:

Theorem 1.10. *Let $P(y) \in (\mathbb{F}[t])[y]$ be a nonconstant polynomial of degree d and derivational degree k . Write $P(y) = a_0 + \sum_{i=1}^n \eta_i(y^{r^i})$. Let $H_i = \eta_i(\mathbb{F}[t])$ and $H = \sum_{i=1}^n H_i$. Then for any $Q(t) \in \mathbb{F}[t]^+$ and any $f : \mathbb{F}[t]_Q \rightarrow \mathbb{C}$,*

$$\left\| \mathbb{E}_{y \in \mathbb{F}[t]_Q} f(x + P(y)) - \mathbb{E}_{z \in H_Q} f(x + a_0 + z) \right\|_{L^2(\mathbb{F}[t]_Q)} \leq \left(p^{2 \lfloor \log_p d \rfloor} \frac{k-1}{\text{lpf}(Q)} \right)^{1/2^{k-1}} \|f\|_{L^2(\mathbb{F}[t]_Q)},$$

where $H_Q = \{z \pmod{Q} : z \in H\}$.

Remark 1.11. (1) In the case $a_0 = P(0) = 0$, we get that the average $\mathbb{E}_{y \in \mathbb{F}[t]_Q} f(x + P(y))$ is approximated in $L^2(\mathbb{F}[t]_Q)$ by the function

$$\tilde{f}(x) = \mathbb{E}_{z \in H_Q} f(x + z),$$

which is the projection of f onto the space of H_Q -invariant functions, so Theorem 1.10 can indeed be seen as a finitary version of Theorem 1.6.

⁵In [BL16], the subgroup $\mathcal{F}(\eta)$ is called a Φ -subtorus of level $\leq \log_p d$

(2) The phenomenon encompassed by Theorem 1.10 is simpler in the low degree situation ($d < p$). In the context of finite fields ($\mathbb{F}[t]_Q$ with Q irreducible in our notation), a closely related result was previously established in [BBI21, Lemma 3]. In particular, it is shown that for any q , any subsets $A, B \subseteq \mathbb{F}_q$, and any polynomial $P(y) \in \mathbb{F}_q[y]$ with degree $d < p$,

$$\left| \mathbb{E}_{x,y \in \mathbb{F}_q} \mathbb{1}_A(x) \mathbb{1}_B(x + P(y)) - \frac{|A||B|}{q^2} \right| \leq \left(\frac{q^{d-1} - (q-1)^{d-1}}{q^{d-1}} \right)^{1/2^{d-1}} \ll q^{-1/2^{d-1}};$$

see the statement of Lemma 3 in [BBI21] and the formula for $\mathcal{E}(q, d)$ at the top of page 713.

As a consequence of Theorem 1.10, we obtain the following power savings for the Furstenberg–Sárközy theorem:

Corollary 1.12. *Let $P(y) \in (\mathbb{F}[t])[y]$ be an intersective polynomial of degree d and derivational degree k . Let $Q(t) \in \mathbb{F}[t]^+$. If $A \subseteq \mathbb{F}[t]_Q$ does not contain distinct $a, b \in A$ with $b - a = P(y)$ for some $y \in \mathbb{F}[t]_Q$, then*

$$|A| \ll_P |Q| \cdot \text{lpf}(Q)^{-1/2^{k-1}}.$$

In particular, if Q is irreducible (so that $\mathbb{F}[t]_Q$ is a field with $|Q|$ elements), then

$$|A| \ll_P |Q|^{1-1/2^{k-1}}.$$

Remark 1.13. (1) If we restrict to $\text{lpf}(Q)$ being sufficiently large (so that $P(y)$ does not reduce to the zero polynomial mod Q), then the implicit constant in the conclusion of Corollary 1.12 depends only on the degree d and the derivational degree k .

(2) The bound given in Theorem 1.4 is difficult to compute in general and to compare with the bound in Corollary 1.12. We can, however, highlight some general features of the different bounds. The power savings obtained in Corollary 1.12 depends only on the derivational degree k of the polynomial P and applies to all intersective polynomials with coefficients in $\mathbb{F}[t]$. By contrast, the Li–Sauer mann bound depends on the degree d of the polynomial P and on the characteristic p of the field \mathbb{F}_q and applies only to polynomials with zero constant term and with coefficients in \mathbb{F}_p .

The disadvantage of our bound is that the power saving $\frac{1}{2^{k-1}}$ decays exponentially with the derivational degree. How the quantity $t_{p,d}$ appearing in Theorem 1.4 depends on p and d is not immediately clear from the definition. However, a related bound due to Green [G17, Theorem 1.2] (which Li and Sauer mann optimize) gives power savings of

$$c'(p, d) = \frac{1}{2d^2(p-1)^2(1 + \log_p d)^2 \log p}$$

for a degree d polynomial $P(y) \in \mathbb{F}_p[y]$ with $P(0) = 0$. That is, for any $q = p^k$, the largest subset of \mathbb{F}_q with no nontrivial pattern $\{x, x + P(y)\}$ has cardinality

$$|A| \ll_{p,d} q^{1-c'(p,d)}.$$

Therefore, for fixed p , there is a regime of sufficiently high degree polynomials for which the Li–Sauer mann bound beats ours. On the other hand, the quantity $c'(p, d)$ decays as $p \rightarrow \infty$, so the bound in Corollary 1.12 will outperform this bound in sufficiently high characteristic (for fixed degree d).

Thus, neither of the power saving bounds is universally better than the other, and both methods have their advantages and disadvantages. The main goal of our work is not to produce the best possible power saving bounds but to provide a heuristic backing for why any power saving bound should hold at all and to place the Furstenberg–Sárközy theorem over finite fields within the appropriate general framework.

Combining Theorems 1.9 and 1.10, we may deduce several finitary combinatorial statements from an infinitary statement about equidistribution. Say that a polynomial $P(y) \in (\mathbb{F}[t])[y]$ is *good for irrational equidistribution* if $(P(n)\alpha)_{n \in \mathbb{F}[t]}$ is well distributed for every irrational $\alpha \in \mathbb{F}((t^{-1})) \setminus \mathbb{F}(t)$.

Proposition 1.14. *A polynomial $P(y)$ is good for irrational equidistribution if and only if for any totally ergodic system $(X, \mathcal{X}, \mu, (T_n)_{n \in \mathbb{F}[t]})$, any Følner sequence $(\Phi_N)_{N \in \mathbb{N}}$ in $\mathbb{F}[t]$, and any $f \in L^2(\mu)$,*

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \in \Phi_N} T_{P(n)} f = \int_X f d\mu.$$

Proposition 1.14 can be proved along the lines of Theorem 1.6 outlined earlier using the spectral theorem for unitary actions of $\mathbb{F}[t]$. For the “only if” direction, upon replacing T_n by the multiplication operators $(U_n h)(x) = e(nx)h(x)$ on $L^2(\mathbb{F}((t^{-1}))/\mathbb{F}[t], \sigma)$, we use the fact that $P(y)$ is good for irrational equidistribution to conclude

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \in \Phi_N} U_{P(n)} h(x) = \lim_{N \rightarrow \infty} \mathbb{E}_{n \in \Phi_N} e(P(n)x)h(x) = \begin{cases} h(0), & \text{if } x = 0; \\ 0, & \text{otherwise} \end{cases}$$

in $L^2(\sigma)$. This corresponds to the desired convergence result

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \in \Phi_N} T_{P(n)} f = \int_X f d\mu.$$

For the “if” direction: suppose $P(y)$ is not good for irrational equidistribution, and let $\alpha \in \mathbb{F}((t^{-1})) \setminus \mathbb{F}(t)$ such that $\lim_{N \rightarrow \infty} \mathbb{E}_{n \in \Phi_N} e(P(n)\alpha) \neq 0$. We then take as our totally ergodic system $X = \mathbb{F}((t^{-1}))/\mathbb{F}[t]$, $\mathcal{X} = \text{Borel}(X)$, $\mu = m_X$, and $T_n x = x + n\alpha$. For the function $f(x) = e(x)$, we have $\int_X f(x) dx = 0$, since $x \mapsto e(x)$ is a nontrivial character on X , while

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \in \Phi_N} T_{P(n)} f = \lim_{N \rightarrow \infty} \mathbb{E}_{n \in \Phi_N} e(P(n)\alpha) f \neq 0.$$

Remark 1.15. The naive analogue of Proposition 5.1 for \mathbb{Z} -actions is true. That is, a polynomial $P(y) \in \mathbb{Z}[y]$ is good for irrational equidistribution (meaning that $(P(n)\alpha)$ is well-distributed mod 1 for every irrational $\alpha \in \mathbb{R} \setminus \mathbb{Q}$) if and only if for any totally ergodic system (X, \mathcal{X}, μ, T) , any Følner sequence $(\Phi_N)_{N \in \mathbb{N}}$ in \mathbb{Z} , and any $f \in L^2(\mu)$,

$$\lim_{N \rightarrow \infty} \mathbb{E}_{n \in \Phi_N} T^{P(n)} f = \int_X f d\mu.$$

However, this result is far less meaningful in the setting of \mathbb{Z} -actions, since every nonconstant integer polynomial is good for irrational equidistribution by Weyl’s equidistribution theorem. This is far from the case in the setting of $\mathbb{F}[t]$; see the examples below.

Example 1.16. (1) Every nonconstant separable polynomial is good for irrational equidistribution; see [BL16, Corollary 0.5].

(2) If η_1, \dots, η_n are additive polynomials such that $\sum_{i=1}^n c_i \eta_i(y) = ay$ for some $c_1, \dots, c_n \in \mathbb{Z}$ and $a \in \mathbb{F}[t]$, then for any distinct $r_1, \dots, r_n \in \mathbb{N}$ not divisible by p , $P(y) = \sum_{i=1}^n \eta_i(y^{r_i})$ is good for irrational equidistribution. This follows from Theorem 1.9, since the condition $\sum_{i=1}^n c_i \eta_i(y) = ay$ ensures that the orbit closure $\sum_{i=1}^n \mathcal{F}(\alpha \eta_i)$ contains the orbit $\{ay\alpha : y \in \mathbb{F}[t]\}$, which is dense mod $\mathbb{F}[t]$ for irrational $\alpha \in \mathbb{F}((t^{-1})) \setminus \mathbb{F}(t)$.

(3) The polynomial $P(y) = y^{p^2} + y^{2p} - y$ is good for irrational equidistribution. This follows from Theorem 1.18 below. Indeed, upon writing $P(y) = \eta_1(y) + \eta_2(y^2)$ with $\eta_1(y) = y^{p^2} - y$ and $\eta_2(y) = y^p$, we see that $P(y)$ satisfies condition (iii) of Theorem 1.18 for $\zeta_1(y) = -y$ and $\zeta_2(y) = y^p$.

(4) The polynomial $P(y) = y^p$ is not good for irrational equidistribution: for any α of the form $\alpha = \beta^p$, the orbit closure $\overline{\{P(y)\alpha : y \in \mathbb{F}[t]\}}$ is contained in the infinite index subgroup $\{x^p : x \in \mathbb{F}((t^{-1}))/\mathbb{F}[t]\} \subseteq \mathbb{F}((t^{-1}))/\mathbb{F}[t]$.

(5) The polynomial $P(y) = y^{2p} - y^2$ is not good for irrational equidistribution. Write $P(y) = \eta(y^2)$ with $\eta(y) = y^p - y$. Then clearly $P(\mathbb{F}[t]) \subseteq \eta(\mathbb{F}[t])$. For any $\alpha \in \mathbb{F}((t^{-1}))/\mathbb{F}[t]$ of the form

$$\alpha = \sum_{j=0}^{\infty} \alpha_j t^{-(j+1)}$$

satisfying $\alpha_{pj} = \alpha_j$ for $j \geq 0$, one can check by direct calculation that $e(\eta(y)\alpha) = 1$ for every $y \in \mathbb{F}[t]$. Hence, for any such α , $(P(n)\alpha)_{n \in \mathbb{F}[t]}$ is not well distributed mod $\mathbb{F}[t]$. Moreover, the set of all such α is uncountable so contains irrational elements.

(6) An additive polynomial $P(y) = \sum_{i=0}^k a_i y^{p^i}$ is good for irrational equidistribution if and only if $P(y) = a_0 y$; see Proposition 5.4.

The following theorem summarizes the main achievements in this paper. In particular, it emphasizes the role of equidistribution properties in obtaining finitary combinatorial results over the rings $\mathbb{F}[t]_Q$. Note that item (v) below strengthens the conclusion of Corollary 1.12 under the assumption that the polynomial $P(y)$ is good for irrational equidistribution.

Theorem 1.17. *Let $P(y) \in (\mathbb{F}[t])[y]$ be a nonconstant polynomial. The following are equivalent:*

(i) *for any $Q(t) \in \mathbb{F}[t]^+$,*

$$\sup_{\|f\|_{L^2(\mathbb{F}[t]_Q)}=1} \left\| \mathbb{E}_{y \in \mathbb{F}[t]_Q} f(x + P(y)) - \mathbb{E}_{z \in \mathbb{F}[t]_Q} f(z) \right\|_{L^2(\mathbb{F}[t]_Q)} = o_{\text{lpf}(Q) \rightarrow \infty}(1);$$

(ii) *there exist $C_1, C_2, \gamma > 0$ such that for any $Q(t) \in \mathbb{F}[t]^+$ with $\text{lpf}(Q) \geq C_1$, one has*

$$\sup_{\|f\|_{L^2(\mathbb{F}[t]_Q)}=1} \left\| \mathbb{E}_{y \in \mathbb{F}[t]_Q} f(x + P(y)) - \mathbb{E}_{z \in \mathbb{F}[t]_Q} f(z) \right\|_{L^2(\mathbb{F}[t]_Q)} \leq C_2 \cdot \text{lpf}(Q)^{-\gamma};$$

(iii) *there exists $C > 0$ such that if $Q(t) \in \mathbb{F}[t]^+$ and $\text{lpf}(Q) \geq C$, then $H + Q\mathbb{F}[t] = \mathbb{F}[t]$, where $H \leq (\mathbb{F}[t], +)$ is the group generated by $\{P(y) - P(0) : y \in \mathbb{F}[t]\}$.*

(iv) *for any $\delta > 0$, there exists $N > 0$ such that if $Q(t) \in \mathbb{F}[t]^+$ has $\text{lpf}(Q) \geq N$ and $A, B \subseteq \mathbb{F}[t]_Q$ are subsets with $|A||B| \geq \delta|Q|^2$, then there exist $x, y \in \mathbb{F}[t]_Q$ such that $x \in A$ and $x + P(y) \in B$;*

(v) *there exist $C_1, C_2, \gamma > 0$ such that for any $Q(t) \in \mathbb{F}[t]^+$ with $\text{lpf}(Q) \geq C_1$, one has*

$$\left| |\{(x, y) \in \mathbb{F}[t]_Q^2 : x \in A, x + P(y) \in B\}| - |A||B| \right| \leq C_2 |A|^{1/2} |B|^{1/2} |Q| \cdot \text{lpf}(Q)^{-\gamma}.$$

Moreover, if $P(y)$ is good for irrational equidistribution, then each of the properties (i)-(v) holds.

By restricting the coefficients of the polynomial P , we can prove a stronger version of Theorem 1.17:

Theorem 1.18. *Let $P(y) \in \mathbb{F}_p[y]$. Let $\eta_1, \dots, \eta_n \in \mathbb{F}_p[y]$ be additive polynomials and $r_1, \dots, r_n \in \mathbb{N}$ distinct positive integers not divisible by p so that $P(y) = \sum_{i=1}^n \eta_i(y^{r_i})$. The following are equivalent:*

(i) *$P(y)$ is good for irrational equidistribution;*

(ii) *for any totally ergodic system $(X, \mathcal{X}, \mu, (T_y)_{y \in \mathbb{F}[t]})$, any Følner sequence $(\Phi_N)_{N \in \mathbb{N}}$ in $\mathbb{F}[t]$, and any $f \in L^2(\mu)$,*

$$\lim_{N \rightarrow \infty} \mathbb{E}_{y \in \Phi_N} T_{P(y)} f = \int_X f d\mu;$$

(iii) *there exist additive polynomials $\zeta_1, \dots, \zeta_n \in \mathbb{F}_p[y]$ and $a \in \mathbb{F}_p^\times$ such that*

$$\sum_{i=1}^n (\eta_i \circ \zeta_i)(y) = ay;$$

(iv) for any $Q(t) \in \mathbb{F}[t]^+$,

$$\sup_{\|f\|_{L^2(\mathbb{F}[t]_Q)}=1} \left\| \mathbb{E}_{y \in \mathbb{F}[t]_Q} f(x + P(y)) - \mathbb{E}_{z \in \mathbb{F}[t]_Q} f(z) \right\|_{L^2(\mathbb{F}[t]_Q)} = o_{\text{lpf}(Q) \rightarrow \infty}(1);$$

(v) there exist $C_1, C_2, \gamma > 0$ such that for any $Q(t) \in \mathbb{F}[t]^+$ with $\text{lpf}(Q) \geq C_1$, one has

$$\sup_{\|f\|_{L^2(\mathbb{F}[t]_Q)}=1} \left\| \mathbb{E}_{y \in \mathbb{F}[t]_Q} f(x + P(y)) - \mathbb{E}_{z \in \mathbb{F}[t]_Q} f(z) \right\|_{L^2(\mathbb{F}[t]_Q)} \leq C_2 \cdot \text{lpf}(Q)^{-\gamma};$$

(vi) there exists $C > 0$ such that if $Q(t) \in \mathbb{F}[t]^+$ and $\text{lpf}(Q) \geq C$, then $H_Q = \mathbb{F}[t]$, where $H_Q = \sum_{i=1}^n H_{i,Q}$, $H_{i,Q} = \eta_i(\mathbb{F}[t]_Q)$.

(vii) for any $\delta > 0$, there exists $N > 0$ such that if $Q(t) \in \mathbb{F}[t]^+$ has $\text{lpf}(Q) \geq N$ and $A, B \subseteq \mathbb{F}[t]_Q$ are subsets with $|A||B| \geq \delta|Q|^2$, then there exist $x, y \in \mathbb{F}[t]_Q$ such that $x \in A$ and $x + P(y) \in B$;

(viii) there exist $C_1, C_2, \gamma > 0$ such that for any $Q(t) \in \mathbb{F}[t]^+$ with $\text{lpf}(Q) \geq C_1$, one has

$$\left| \left| \{(x, y) \in \mathbb{F}[t]_Q^2 : x \in A, x + P(y) \in B\} \right| - |A||B| \right| \leq C_2 |A|^{1/2} |B|^{1/2} |Q| \cdot \text{lpf}(Q)^{-\gamma}.$$

Combining Furstenberg–Sárközy-type results with the technology of Loeb measures on ultra-products, we are able to establish partition regularity of families of polynomial equations over finite fields, such as the following:

Theorem 1.19. *Let $P(x) \in \mathbb{F}_p[x]$ be a nonconstant polynomial, and let $Q(x) \in \mathbb{F}_p[x]$ be good for irrational equidistribution. For any $r \in \mathbb{N}$, there exists $K = K(P, Q, r) \in \mathbb{N}$ and $c = c(P, Q, r) > 0$ such that for any $k \geq K$ and any r -coloring $\mathbb{F}_{p^k} = \bigcup_{i=1}^r C_i$, there are at least $c(p^k)^2$ monochromatic solutions to the equation $P(x) - P(y) = Q(z)$. That is,*

$$\left| \left\{ (x, y, z) \in \mathbb{F}_{p^k}^3 : P(x) - P(y) = Q(z) \text{ and } \{x, y, z\} \subseteq C_i \text{ for some } i \in \{1, \dots, r\} \right\} \right| \geq c(p^k)^2.$$

We are in fact able to prove partition regularity under a weaker (but more technically cumbersome) condition on Q ; see Corollary 6.7. Under this weaker assumption, one application of note is a polynomial Schur theorem over finite fields:

Corollary 1.20. *Let $P(x) \in \mathbb{Z}[x]$ with $P(0) = 0$. Then for any $r \in \mathbb{N}$, there exists $N = N(P, r) \in \mathbb{N}$ and $c = c(P, r) > 0$ such that if $q > N$ and $\mathbb{F}_q = \bigcup_{i=1}^r C_i$, then there are at least cq^2 monochromatic solutions to the equation $P(x) + P(y) = P(z)$. In particular, if the coefficients of P are not all divisible by the characteristic of \mathbb{F}_q , then there are $\gg_{P,r} q^2$ monochromatic solutions with $P(x), P(y), P(z) \neq 0$.*

Remark 1.21. In the case $P(x) = x^d$, Corollary 1.20 corresponds to the Fermat equation $x^d + y^d = z^d$. Schur proved the existence of solutions to the Fermat equation in all prime fields of sufficiently high characteristic using his eponymous partition regularity theorem in [S16]. (This was in fact the original motivation for Schur’s theorem.) The much stronger property of partition regularity of the Fermat equation was established previously in the context of prime fields in [CGS12, Theorem 4] and generalized to a family of related polynomial equations in [L18]. We complete the picture here by extending the partition regularity property to arbitrary finite fields of sufficiently large order (with no assumption on the characteristic).

Related density results for Pythagorean pairs and triples in finite fields were obtained in [DLMS23, Section 6], where they also show that a density version (“density regularity”) of Corollary 1.20 fails already for the Pythagorean equation $x^2 + y^2 = z^2$.

The paper is organized as follows. In Section 2, we prove Theorem 1.5, showing that $\text{lpf}(Q)$ appropriately captures the phenomenon of asymptotic total ergodicity. Section 3 is dedicated to

the proof of Theorem 1.10. The final three sections concern applications of Theorem 1.10: Section 4 deals with power saving bounds in the Furstenberg–Sárközy theorem for intersective polynomials in the presence of asymptotic total ergodicity (Corollary 1.12); Section 5 with further enhancements of the Furstenberg–Sárközy theorem for polynomials with good equidistributional behavior (Theorems 1.17 and 1.18); and Section 6 with partition regularity of polynomial equations over finite fields. The relevant tools are introduced in the corresponding sections as needed.

2. ASYMPTOTIC TOTAL ERGODICITY

In this section, we prove that the quantity $\text{lpf}(Q)$ captures the phenomenon of asymptotic total ergodicity. Recall Theorem 1.5:

Theorem 1.5. *Let $(Q_n)_{n \in \mathbb{N}}$ be a sequence in $\mathbb{F}[t]^+$. The following are equivalent:*

(i) *The sequence of quotient rings $\mathbb{F}[t]_{Q_n}$ is asymptotically totally ergodic: for any $m \in \mathbb{F}[t] \setminus \{0\}$,*

$$\sup_{f_n: \mathbb{F}[t]_{Q_n} \rightarrow \mathbb{D}} \left\| \mathbb{E}_{y \in \mathbb{F}[t]_{Q_n}} f_n(x + my) - \mathbb{E}_{z \in \mathbb{F}[t]_{Q_n}} f_n(z) \right\|_{L^2(\mathbb{F}[t]_{Q_n})} \xrightarrow{n \rightarrow \infty} 0.$$

(ii) $\text{lpf}(Q_n) \rightarrow \infty$.

Proof. (ii) \implies (i). Fix $m \in \mathbb{F}[t] \setminus \{0\}$. If $Q \in \mathbb{F}[t]^+$ has $\text{lpf}(Q) > |m|$, then m is an element of the multiplicative group $\mathbb{F}[t]_Q^\times$. Hence, for any $x \in \mathbb{F}[t]_Q$ and any $f: \mathbb{F}[t]_Q \rightarrow \mathbb{C}$,

$$\mathbb{E}_{y \in \mathbb{F}[t]_Q} f(x + my) = \mathbb{E}_{z \in \mathbb{F}[t]_Q} f(z).$$

(i) \implies (ii). Let $Q \in \mathbb{F}[t]^+$, and let m be an irreducible factor of Q . Enumerate $\mathbb{F}[t]_m = \{x_0, \dots, x_{|m|-1}\}$, and let $f_0: \mathbb{F}[t]_m \rightarrow \mathbb{D}$ be the function $f_0(x_k) = e^{2\pi i k / |m|}$. Define $f: \mathbb{F}[t]_Q \rightarrow \mathbb{D}$ by $f(x) = f_0(x \bmod m)$. Then

$$\mathbb{E}_{y \in \mathbb{F}[t]_Q} f(x + my) = f(x)$$

for every $x \in \mathbb{F}[t]_Q$. On the other hand,

$$\mathbb{E}_{z \in \mathbb{F}[t]_Q} f(z) = \frac{|m|-1}{|m|} \mathbb{E}_{k=0}^{|m|-1} \mathbb{E}_{z \equiv x_k} f(z) = \frac{|m|-1}{|m|} \mathbb{E}_{k=0}^{|m|-1} f_0(x_k) = 0.$$

Therefore,

$$\left\| \mathbb{E}_{y \in \mathbb{F}[t]_Q} f(x + my) - \mathbb{E}_{z \in \mathbb{F}[t]_Q} f(z) \right\|_{L^2(\mathbb{F}[t]_Q)}^2 = \mathbb{E}_{x \in \mathbb{F}[t]_Q} |f(x)|^2 = 1.$$

Now suppose $\text{lpf}(Q_n) \not\rightarrow \infty$. Taking a subsequence if necessary, we may assume $\text{lpf}(Q_n)$ is bounded. By the pigeonhole principle, we may then take a further subsequence and assume that there is a common irreducible factor m of every Q_n , $n \in \mathbb{N}$. The above calculation shows that we may find $f_n: \mathbb{F}[t]_{Q_n} \rightarrow \mathbb{D}$ with

$$\left\| \mathbb{E}_{y \in \mathbb{F}[t]_{Q_n}} f_n(x + my) - \mathbb{E}_{z \in \mathbb{F}[t]_{Q_n}} f_n(z) \right\|_{L^2(\mathbb{F}[t]_{Q_n})} = 1$$

for each $n \in \mathbb{N}$, contradicting (i). \square

3. ASYMPTOTIC PROJECTION THEOREM

We now turn to proving Theorem 1.10 with Fourier analysis. Characters on $\mathbb{F}[t]_Q$ take the form $\chi(x) = e(sx/Q)$ for some $s \in \mathbb{F}[t]_Q$. For a function $f : \mathbb{F}[t]_Q \rightarrow \mathbb{C}$, we therefore define its Fourier transform $\widehat{f} : \mathbb{F}[t]_Q \rightarrow \mathbb{C}$ by

$$\widehat{f}(s) = \mathbb{E}_{x \in \mathbb{F}[t]_Q} f(x)e(-sx/Q).$$

We state some basic properties of the Fourier transform

Proposition 3.1. *For any $f : \mathbb{F} \rightarrow \mathbb{C}$, one has*

- *Fourier inversion:*

$$f(x) = \sum_{s \in \mathbb{F}[t]_Q} \widehat{f}(s)e(sx/Q).$$

- *Parseval's identity:*

$$\mathbb{E}_{x \in \mathbb{F}[t]_Q} |f(x)|^2 = \sum_{s \in \mathbb{F}[t]_Q} |\widehat{f}(s)|^2.$$

Define $F(x) := \mathbb{E}_{y \in \mathbb{F}[t]_Q} f(x + P(y)) - \mathbb{E}_{z \in H_Q} f(x + a_0 + z)$. Then

$$\widehat{F}(s) = \left(\mathbb{E}_{y \in \mathbb{F}[t]_Q} e(sP(y)/Q) - e(sa_0/Q) \mathbb{1}_{H_Q^\perp}(s) \right) \widehat{f}(s),$$

where $H_Q^\perp = \{s \in \mathbb{F}[t]_Q : e(sz) = 1 \text{ for all } z \in H_Q\}$ is the annihilator of the subgroup H_Q . Hence, by Parseval's identity

$$\begin{aligned} \mathbb{E}_{x \in \mathbb{F}[t]_Q} |F(x)|^2 &= \sum_{s \in \mathbb{F}[t]_Q} \left| \mathbb{E}_{y \in \mathbb{F}[t]_Q} e(sP(y)/Q) - e(sa_0/Q) \mathbb{1}_{H_Q^\perp}(s) \right|^2 |\widehat{f}(s)|^2 \\ &\leq \left(\sup_{s \in \mathbb{F}[t]_Q} \left| \mathbb{E}_{y \in \mathbb{F}[t]_Q} e(sP(y)/Q) - e(sa_0/Q) \mathbb{1}_{H_Q^\perp}(s) \right| \right)^2 \mathbb{E}_{x \in \mathbb{F}[t]_Q} |f(x)|^2. \end{aligned}$$

All that remains to show is the inequality:

Proposition 3.2. *Let $P(y) \in (\mathbb{F}[t])[y]$ be a nonconstant polynomial of degree d and derivational degree k . Write $P(y) = a_0 + \sum_{i=1}^n \eta_i(y^{r_i})$. Then for any $Q(t) \in \mathbb{F}[t]^+$ and any $s \in \mathbb{F}[t]_Q$,*

$$\left| \mathbb{E}_{y \in \mathbb{F}[t]_Q} e(sP(y)/Q) - e(sa_0/Q) \mathbb{1}_{H_Q^\perp}(s) \right|^{2^{k-1}} \leq p^{2 \lfloor \log_p d \rfloor} \frac{k-1}{\text{lpf}(Q)}.$$

A key ingredient in Proposition 3.2 is the following van der Corput-type inequality. We do not use any ring structure for this result, so we state and prove it in the setting of an arbitrary finite abelian group G . For a function $f : G \rightarrow \mathbb{C}$, define a multiplicative differencing operator by $\Delta_v f(u) = f(u+v)\overline{f(u)}$, and let $\Delta_{v_1, \dots, v_k} f = \Delta_{v_k} (\Delta_{v_1, \dots, v_{k-1}} f)$ for $k \in \mathbb{N}$ and $v_1, \dots, v_k \in G$.

Lemma 3.3. *Let G be a finite abelian group, and let $H \leq G$ be a subgroup. For any function $f : G \rightarrow \mathbb{C}$ and any $k \in \mathbb{N}$,*

$$\left| \mathbb{E}_{x \in G} f(x) \right|^{2^k} \leq \mathbb{E}_{v_1, \dots, v_k \in H} \mathbb{E}_{u \in G} \Delta_{v_1, \dots, v_k} f(u)$$

Remark 3.4. It is worth commenting on two special cases of Lemma 3.3. When $H = G$, the quantity on the right hand side is equal to $\|f\|_{U^k(G)}^{2^k}$, so the conclusion of Lemma 3.3 reduces to the inequality $\|f\|_{U^1(G)} \leq \|f\|_{U^k(G)}$, which is a special case of monotonicity for the Gowers (semi)norms.

On the other hand, when $H = \{0\}$, the right hand side is equal to $\mathbb{E}_{u \in G} |f(u)|^{2^k}$, so the conclusion of Lemma 3.3 follows by Jensen's inequality. The general case can be seen as interpolating between these two extremes.

Proof. Suppose $k = 1$. Note that

$$\mathbb{E}_{x \in G} f(x) = \mathbb{E}_{x \in G} \mathbb{E}_{h \in H} f(x+h).$$

Therefore, by Jensen's inequality,

$$\left| \mathbb{E}_{x \in G} f(x) \right|^2 = \left| \mathbb{E}_{x \in G} \mathbb{E}_{h \in H} f(x+h) \right|^2 \leq \mathbb{E}_{x \in G} \left| \mathbb{E}_{h \in H} f(x+h) \right|^2 = \mathbb{E}_{x \in G} \mathbb{E}_{h_1, h_2 \in H} f(x+h_1) \overline{f(x+h_2)}.$$

Interchanging the order of averaging and making the substitutions $v = h_1 - h_2$, $u = x - h_2$, we obtain the desired inequality

$$\left| \mathbb{E}_{x \in G} f(x) \right|^2 \leq \mathbb{E}_{v \in H} \mathbb{E}_{u \in G} f(u+v) \overline{f(u)}.$$

Suppose the inequality holds for $k - 1$. Then

$$\left| \mathbb{E}_{x \in G} f(x) \right|^{2^k} = \left(\left| \mathbb{E}_{x \in G} f(x) \right|^{2^{k-1}} \right)^2 \leq \left| \mathbb{E}_{v_1, \dots, v_{k-1} \in H} \mathbb{E}_{u \in G} \Delta_{v_1, \dots, v_{k-1}} f(u) \right|^2,$$

which is in turn bounded above by

$$\mathbb{E}_{v_1, \dots, v_{k-1} \in H} \left| \mathbb{E}_{u \in G} \Delta_{v_1, \dots, v_{k-1}} f(u) \right|^2.$$

For fixed v_1, \dots, v_{k-1} , applying the $k = 1$ case with the function $\Delta_{v_1, \dots, v_{k-1}} f$ gives

$$\left| \mathbb{E}_{u \in G} \Delta_{v_1, \dots, v_{k-1}} f(u) \right|^2 \leq \mathbb{E}_{v_k \in H} \mathbb{E}_{u \in G} \Delta_{v_k} \Delta_{v_1, \dots, v_{k-1}} f(x) = \mathbb{E}_{v_k \in H} \mathbb{E}_{u \in G} \Delta_{v_1, \dots, v_{k-1}, v_k} f(x).$$

Putting everything together,

$$\left| \mathbb{E}_{x \in G} f(x) \right|^{2^k} \leq \mathbb{E}_{v_1, \dots, v_k \in H} \mathbb{E}_{u \in G} \Delta_{v_1, \dots, v_k} f(u).$$

□

Proof of Proposition 3.2. We first make a reduction to separable polynomials. If $s \in H_Q^\perp$, then $e(sP(y)/Q) = e(sa_0/Q)e(s(P(y) - a_0)/Q) = e(sa_0/Q)$ for every $y \in \mathbb{F}[t]_Q$, since $P(y) - a_0 \in H_Q$. Suppose now that $s \notin H_Q^\perp$. We want to show

$$\left| \mathbb{E}_{y \in \mathbb{F}[t]_Q} e(sP(y)/Q) \right|^{2^{k-1}} \leq p^{2 \lfloor \log_p d \rfloor} \frac{k-1}{\text{lpf}(Q)}.$$

Noting that $H_Q^\perp = \bigcap_{i=1}^n H_{i,Q}^\perp$, we have

$$I = \left\{ 1 \leq i \leq n : s \notin H_{i,Q}^\perp \right\} \neq \emptyset.$$

Moreover, for any $y \in \mathbb{F}[t]_Q$,

$$e(sP(y)/Q) = e(sa_0/Q) e \left(s \sum_{i \in I} \eta_i(y^{r_i})/Q \right).$$

Now, for each $i \in I$, the function $\chi_i(x) = e(s\eta_i(x)/Q)$ is a nontrivial character on $\mathbb{F}[t]_Q$, so there exists $s_i \neq 0$ such that $\chi_i(x) = e(s_i x/Q)$. It therefore suffices to prove the following: for any nonconstant separable polynomial $P(y) \in (\mathbb{F}[t]_Q)[y]$,

$$\left| \mathbb{E}_{y \in \mathbb{F}[t]_Q} e(P(y)/Q) \right|^{2^{k-1}} \leq p^{2 \lfloor \log_p d \rfloor} \frac{k-1}{\text{lpf}(Q)}.$$

Suppose $k = 1$. Then $P(y) = s_1 y + s_0 \in (\mathbb{F}[t]_Q)[y]$ with $s_1 \neq 0$. Therefore,

$$\mathbb{E}_{y \in \mathbb{F}[t]_Q} e(P(y)/Q) = e(s_0/Q) \mathbb{E}_{y \in \mathbb{F}[t]_Q} e(s_1 y/Q) = 0.$$

Now suppose $k \geq 2$. Let $P(y) = \sum_{i=1}^n s_i y^{r_i} + P'(y)$, where $\text{d-deg } y^{r_i} = k$ and $\text{d-deg } P'(y) \leq k-1$. By Lemma 3.3,

$$\left| \mathbb{E}_{y \in \mathbb{F}[t]_Q} e(P(y)/Q) \right|^{2^{k-1}} \leq \mathbb{E}_{v_1, \dots, v_{k-1} \in K} \mathbb{E}_{u \in \mathbb{F}[t]_Q} e(\partial_{v_1, \dots, v_{k-1}} P(u)/Q)$$

for any subgroup $K \leq (\mathbb{F}[t]_Q, +)$. (We will take a convenient choice for K later.) We now wish to obtain an expression for $\partial_{v_1, \dots, v_k} P(u)$ that will allow us to bound the average

$$\mathbb{E}_u e(s \partial_{v_1, \dots, v_{k-1}} P(u)/Q).$$

For $v_1, \dots, v_{k-1} \in \mathbb{F}[t]_Q$, one has that $\partial_{v_1, \dots, v_{k-1}} P'(u)$ is constant (as a function of u), since $\text{d-deg } P'(y) \leq k-1$, so we can pull the constant $e(s \partial_{v_1, \dots, v_{k-1}} P'(u))$ outside of the average.

Let $m = \lfloor \log_p d \rfloor$ so that $p^m \leq d$ and $p^{m+1} > d$. For each $i = 1, \dots, n$, we may write $r_i = \sum_{j=0}^m c_{i,j} p^j$ with $c_{i,j} \in \{0, \dots, p-1\}$ and $\sum_{j=0}^m c_{i,j} = k$. Since P is separable by assumption, we have $c_{i,0} \neq 0$ for $i \in \{1, \dots, n\}$. Then

$$\partial_{v_1, \dots, v_{k-1}}(y^{r_i}) = b_i \sum_{j=1}^m S_{i,j}(v_1, \dots, v_{k-1}) u^{p^j} + R_i(v_1, \dots, v_{k-1}),$$

where $b_i = \prod_{j=1}^m c_{i,j}!$, $S_{i,j}(v_1, \dots, v_{k-1})$ is the sum of all monomials of the form

$$\prod_{l=1}^{k-1} v_l^{p^{j_l}}$$

with

$$|\{1 \leq l \leq k-1 : j_l = j'\}| = \begin{cases} c_{i,j'}, & \text{if } j' \neq j \\ c_{i,j} - 1, & \text{if } j' = j \end{cases}$$

and R_i is a symmetric polynomial in $k-1$ variables. (If $c_{i,j} = 0$, then $S_{i,j} = 0$.) We can therefore write

$$\begin{aligned} \partial_{v_1, \dots, v_{k-1}} P(y) &= \sum_{i=1}^n s_i b_i \sum_{j=1}^m S_{i,j}(v_1, \dots, v_{k-1}) u^{p^j} + R(v_1, \dots, v_{k-1}) \\ &= \sum_{j=0}^m \left(\sum_{i=1}^n s_i b_i S_{i,j}(v_1, \dots, v_{k-1}) \right) u^{p^j} + R(v_1, \dots, v_{k-1}), \end{aligned}$$

where $R = \sum_{i=1}^n R_i$.

Let

$$\eta_{v_1, \dots, v_{k-1}}(u) = \sum_{j=0}^m \left(\sum_{i=1}^n s_i b_i S_{i,j}(v_1, \dots, v_{k-1}) \right) u^{p^j}.$$

Note that $\eta_{v_1, \dots, v_{k-1}}$ is a group homomorphism $(\mathbb{F}[t]_Q, +) \rightarrow (\mathbb{F}[t]_Q, +)$. It follows that

$$\left| \mathbb{E}_u e(\partial_{v_1, \dots, v_{k-1}} P(u)/Q) \right| = \left| \mathbb{E}_u e(\eta_{v_1, \dots, v_{k-1}}(u)/Q) \right| = 0$$

whenever $e(\eta_{v_1, \dots, v_{k-1}}(\cdot)/Q)$ is a nonzero function. Noting that $e(\eta_{v_1, \dots, v_{k-1}}(\cdot)/Q)$ is a character on $\mathbb{F}[t]_Q$, it may be written in the form $e(\varphi(v_1, \dots, v_{k-1})u/Q)$ for some $\varphi(v_1, \dots, v_{k-1}) \in \mathbb{F}[t]_Q$. We have thus obtained the bound

$$\left| \mathbb{E}_{y \in \mathbb{F}[t]_Q} e(P(y)/Q) \right|^{2^{k-1}} \leq \frac{|\{(v_1, \dots, v_{k-1}) \in K^{k-1} : \varphi(v_1, \dots, v_{k-1}) = 0\}|}{|K|^{k-1}}.$$

The remainder of the proof consists of two main steps. First, we show that, for a convenient choice of K , the function φ becomes (after a change of coordinates) a polynomial in $k-1$ variables. Next, we establish a bound on the number of roots of multivariable polynomials mod Q .

Recall $m = \lfloor \log_p d \rfloor$. Let $K = \{x^{p^m} : x \in \mathbb{F}[t]_Q\}$. This is a subgroup, since the function $x \mapsto x^{p^m}$ is a homomorphism. For $0 \leq j \leq m$, let $C_j = \sum_{i=1}^n s_i b_i S_{i,j}$ so that

$$\eta_{v_1, \dots, v_{k-1}}(u) = \sum_{j=0}^m C_j(v_1, \dots, v_{k-1}) u^{p^j},$$

and each of the polynomials $C_j(v_1, \dots, v_{k-1})$ is an additive polynomial of degree at most p^m in each coordinate. In particular, $v_i \mid C_j(v_1, \dots, v_{k-1})$ for each $i \in \{1, \dots, k-1\}$. Making the substitution $v_i = w_i^{p^m}$, we therefore have

$$\eta_{w_1^{p^m}, \dots, w_{k-1}^{p^m}}(u) = \sum_{j=0}^m C_j(w_1^{p^m}, \dots, w_{k-1}^{p^m}) u^{p^j} = \sum_{j=0}^m \tilde{C}_j(w_1, \dots, w_{k-1}) w_1^{p^j} \dots w_{k-1}^{p^j} u^{p^j}$$

for some \tilde{C}_j .

For each $j \geq 0$, the function $\chi(u) = e(u^{p^j}/Q)$ is a character on $\mathbb{F}[t]_Q$, so there exists $z_j \in \mathbb{F}[t]_Q$ such that $\chi(u) = e(z_j u/Q)$. Hence, defining $\psi(w_1, \dots, w_{k-1}) = \varphi(w_1^{p^m}, \dots, w_{k-1}^{p^m})$, we have

$$\psi(w_1, \dots, w_{k-1}) = \sum_{j=0}^m z_j \tilde{C}_j(w_1, \dots, w_{k-1}) w_1 \dots w_{k-1}.$$

That is, ψ is a polynomial of degree at most p^{2m} in each coordinate.

We claim that ψ is not the zero polynomial. By definition, $z_0 = 1$. We also have

$$C_0(v_1, \dots, v_{k-1}) = \sum_{i=1}^n s_i b_i S_{i,0}(v_1, \dots, v_{k-1}).$$

The coefficients b_i are integers coprime to p , so $b_i \in \mathbb{F}[t]_Q^\times$. Hence, $s_i b_i \neq 0$. Now, $S_{i,0}$ is a sum of terms of the form

$$\prod_{l=1}^{k-1} v_l^{p^i}$$

with the property $\sum_{l=1}^{k-1} p^{jl} = r_i - p$. Therefore, the monomials appearing in $S_{i,0}$ are distinct from the monomials appearing in $S_{i',0}$ for $i \neq i'$. It follows that C_0 is not the zero polynomial. Thus,

$$z_0 \tilde{C}_0(w_1, \dots, w_{k-1}) w_1 \dots w_{k-1} = C_0(w_1^{p^m}, \dots, w_{k-1}^{p^m})$$

is not the zero polynomial, and each monomial appearing has degree divisible by p^m . Finally, for $j \neq 0$, we have

$$\tilde{C}_j(w_1, \dots, w_{k-1}) w_1 \dots w_{k-1} = \frac{C_j(w_1^{p^m}, \dots, w_{k-1}^{p^m})}{\prod_{l=1}^{k-1} w_l^{p^{j-1}}},$$

which consists of monomials in which each variable has degree congruent to 1 mod p . This proves that ψ is not the zero polynomial.

The final step is to show that ψ has only a small number of zeros.

Lemma 3.5. *Let $l \in \mathbb{N}$, and let $T(y_1, \dots, y_l) \in (\mathbb{F}[t]_Q)[y_1, \dots, y_l]$ be a nonzero polynomial of degree d_i in the variable y_i for $i = 1, \dots, d$. Then*

$$\left| \left\{ (y_1, \dots, y_l) \in \mathbb{F}[t]_Q^l : T(y_1, \dots, y_l) = 0 \right\} \right| \leq \left(\sum_{i=1}^l d_i \right) \frac{|Q|^l}{\text{lpf}(Q)}.$$

Proof of Lemma. Let us first consider the case $l = 1$. Write $T(y) = \alpha_d y^d + \dots + \alpha_1 y + \alpha_0$. We view $\alpha_0, \dots, \alpha_d$ as elements of $\mathbb{F}[t]$ with $|\alpha_i| < |Q|$. Let $\alpha = \gcd(\alpha_0, \dots, \alpha_d, Q)$, $\tilde{\alpha}_i = \frac{\alpha_i}{\alpha}$, and $\tilde{T}(y) = \tilde{\alpha}_d y^d + \dots + \tilde{\alpha}_1 y + \tilde{\alpha}_0$. Fix $Q' \in \mathbb{F}[t]^+$ irreducible such that $Q' \mid \frac{Q}{\alpha}$. For some $i \in \{0, \dots, d\}$, we have $Q' \nmid \tilde{\alpha}_i$. Since $\mathbb{F}[t]_{Q'}$ is a field, and \tilde{T} reduces to a nonzero polynomial of degree $\leq d \bmod Q'$, we have

$$\left| \{y' \in \mathbb{F}[t]_{Q'} : \tilde{T}(y') \equiv 0 \pmod{Q'}\} \right| \leq d.$$

Now suppose $T(y) = 0$. Then $\alpha \tilde{T}(y) = 0$. That is, $Q \mid \alpha \tilde{T}(y)$, so $\frac{Q}{\alpha} \mid \tilde{T}(y)$. Hence, $Q' \mid \tilde{T}(y)$. Equivalently, $\tilde{T}(y) \equiv 0 \pmod{Q'}$. Therefore,

$$|\{y \in \mathbb{F}[t]_Q : T(y) = 0\}| \leq d \frac{|Q|}{|Q'|} \leq d \frac{|Q|}{\text{lpf}(Q)}.$$

Suppose $l \geq 2$. If $\text{lpf}(Q) \leq \sum_{i=1}^l d_i$, then there is nothing to prove, so assume $\text{lpf}(Q) > \sum_{i=1}^l d_i$. Fix $y \in \mathbb{F}[t]_Q$, and let $T_y(y_1, \dots, y_{l-1}) = T(y_1, \dots, y_{l-1}, y)$. If $T_y(y_1, \dots, y_{l-1}) \in (\mathbb{F}[t]_Q)[y_1, \dots, y_{l-1}]$ is not the zero polynomial, then by the induction hypothesis,

$$\left| \left\{ (y_1, \dots, y_{l-1}) \in \mathbb{F}[t]_Q^{l-1} : T_y(y_1, \dots, y_{l-1}) = 0 \right\} \right| \leq \left(\sum_{i=1}^{l-1} d_i \right) \frac{|Q|^{l-1}}{\text{lpf}(Q)}.$$

Hence,

$$\begin{aligned} & \left| \left\{ (y_1, \dots, y_l) \in \mathbb{F}[t]_Q^l : T(y_1, \dots, y_l) = 0 \right\} \right| \\ &= |Q|^{l-1} |\{y \in \mathbb{F}[t]_Q : T_y = 0\}| + |\{(y_1, \dots, y_{l-1}, y) : T_y \neq 0, T_y(y_1, \dots, y_{l-1}) = 0\}| \\ &\leq |Q|^{l-1} |\{y \in \mathbb{F}[t]_Q : T_y = 0\}| + \left(\sum_{i=1}^{l-1} d_i \right) \frac{|Q|^l}{\text{lpf}(Q)}. \end{aligned}$$

It therefore suffices to prove

$$|\{y \in \mathbb{F}[t]_Q : T_y = 0\}| \leq d_l \frac{|Q|}{\text{lpf}(Q)}.$$

Fix $y_1, \dots, y_{l-1} \in \mathbb{F}[t]_Q$, and let $T^{y_1, \dots, y_{l-1}}(y) = T_y(y_1, \dots, y_{l-1}) = T(y_1, \dots, y_{l-1}, y)$. If $T_y = 0$, then $T^{y_1, \dots, y_{l-1}}(y) = 0$. By Lemma 3.5, it follows that

$$|\{y \in \mathbb{F}[t]_Q : T_y = 0\}| \leq |\{y \in \mathbb{F}[t]_Q : T^{y_1, \dots, y_{l-1}}(y) = 0\}| \leq d_l \frac{|Q|}{\text{lpf}(Q)},$$

unless $T^{y_1, \dots, y_{l-1}}$ is the zero polynomial. So, it remains to find y_1, \dots, y_{l-1} such that $T^{y_1, \dots, y_{l-1}}$ is a nonzero polynomial. Note that the coefficients of $T^{y_1, \dots, y_{l-1}}$ are polynomial expressions in y_1, \dots, y_{l-1} of degree at most d_i in the variable y_i . Since T is not the zero polynomial, there is at least one coefficient that is a nonzero polynomial $C(y_1, \dots, y_{l-1}) \in (\mathbb{F}[t]_Q)[y_1, \dots, y_{l-1}]$. By the induction hypothesis,

$$\left| \left\{ (y_1, \dots, y_{l-1}) \in \mathbb{F}[t]_Q^{l-1} : C(y_1, \dots, y_{l-1}) = 0 \right\} \right| \leq \left(\sum_{i=1}^{l-1} d_i \right) \frac{|Q|^{l-1}}{\text{lpf}(Q)}.$$

Since $\text{lpf}(Q) > \sum_{i=1}^l d_i \geq \sum_{i=1}^{l-1} d_i$ by assumption, it follows that $C(y_1, \dots, y_{l-1}) \neq 0$ for some $(y_1, \dots, y_{l-1}) \in \mathbb{F}[t]_Q^{l-1}$. For this choice of y_1, \dots, y_{l-1} , the polynomial $T^{y_1, \dots, y_{l-1}}(y) \in (\mathbb{F}[t]_Q)[y]$ is not the zero polynomial, so we are done. \square

Applying Lemma 3.5 to ψ , we get the bound

$$\left| \left\{ (w_1, \dots, w_{k-1}) \in \mathbb{F}[t]_Q^{k-1} : \psi(w_1, \dots, w_{k-1}) = 0 \right\} \right| \leq p^{2m}(k-1) \frac{|Q|^{k-1}}{\text{lpf}(Q)}.$$

Thus,

$$\begin{aligned} \left| \mathbb{E}_{y \in \mathbb{F}[t]_Q} e(P(y)/Q) \right|^{2^{k-1}} &\leq \frac{|\{(v_1, \dots, v_{k-1}) \in K^{k-1} : \varphi(v_1, \dots, v_{k-1}) = 0\}|}{|K|^{k-1}} \\ &= \frac{|\{(w_1, \dots, w_{k-1}) \in \mathbb{F}[t]_Q^{k-1} : \psi(w_1, \dots, w_{k-1}) = 0\}|}{|Q|^{k-1}} \\ &\leq \frac{p^{2m}(k-1)}{\text{lpf}(Q)}. \end{aligned}$$

\square

4. POWER SAVING BOUNDS FOR THE FURSTENBERG–SÁRKÖZY THEOREM IN CHARACTERISTIC p

We now prove Corollary 1.12, restated below:

Corollary 1.12. *Let $P(y) \in (\mathbb{F}[t])[y]$ be an intersective polynomial of degree d and derivational degree k . Let $Q(t) \in \mathbb{F}[t]^+$. If $A \subseteq \mathbb{F}[t]_Q$ does not contain distinct $a, b \in A$ with $b - a = P(y)$ for some $y \in \mathbb{F}[t]_Q$, then*

$$|A| \ll_P |Q| \cdot \text{lpf}(Q)^{-1/2^{k-1}}.$$

In particular, if Q is irreducible (so that $\mathbb{F}[t]_Q$ is a field with $|Q|$ elements), then

$$|A| \ll_P |Q|^{1-1/2^{k-1}}.$$

Proof of Corollary 1.12. Since P is intersective, there exists $y \in \mathbb{F}[t]_Q$ with $P(y) = 0$. Hence, $-a_0 = P(y) - a_0 \in H_Q$. Therefore, applying Theorem 1.10 with $f = \mathbb{1}_A$, we have

$$\left| \mathbb{E}_{x,y \in \mathbb{F}[t]_Q} \mathbb{1}_A(x) \mathbb{1}_A(x + P(y)) - \mathbb{E}_{x \in \mathbb{F}[t]_Q, z \in H_Q} \mathbb{1}_A(x) \mathbb{1}_A(x + z) \right| \leq \frac{|A|}{|Q|} \left(p^{2 \lfloor \log_p d \rfloor} \frac{k-1}{\text{lpf}(Q)} \right)^{1/2^{k-1}}$$

by the Cauchy–Schwarz inequality.

On the one hand, if A contains no nontrivial patterns $\{x, x + P(y)\}$, then by Lemma 3.5,

$$\mathbb{E}_{x,y \in \mathbb{F}[t]_Q} \mathbb{1}_A(x) \mathbb{1}_A(x + P(y)) = \frac{|A|}{|Q|} \frac{|\{y \in \mathbb{F}[t]_Q : P(y) = 0\}|}{|Q|} \leq \frac{|A|}{|Q|} \frac{d}{\text{lpf}(Q)},$$

as long as $\text{lpf}(Q)$ is large enough so that P is not the zero polynomial mod Q . On the other hand,

$$\mathbb{E}_{x \in \mathbb{F}[t]_Q, z \in H_Q} \mathbb{1}_A(x) \mathbb{1}_A(x + z) \geq \left(\frac{|A|}{|Q|} \right)^2$$

by Lemma 3.3. Therefore,

$$\left(\frac{|A|}{|Q|} \right)^2 - \frac{|A|}{|Q|} \frac{d}{\text{lpf}(Q)} \leq C \frac{|A|}{|Q|} \text{lpf}(Q)^{-1/2^{k-1}},$$

where $C = \left(p^{2 \lfloor \log_p d \rfloor} (k-1) \right)^{1/2^{k-1}}$. Multiplying both sides by $\frac{|Q|^2}{|A|}$, we get the desired bound

$$|A| \leq C|Q| \cdot \text{lpf}(Q)^{-1/2^{k-1}} + d \frac{|Q|}{\text{lpf}(Q)} \ll |Q| \cdot \text{lpf}(Q)^{-1/2^{k-1}}.$$

□

5. PROOF OF EQUIVALENCES

The goal of this section is to prove Theorem 1.17, restated here for convenience:

Theorem 1.17. *Let $P(y) \in (\mathbb{F}[t])[y]$ be a nonconstant polynomial. The following are equivalent:*

(i) *for any $Q(t) \in \mathbb{F}[t]^+$,*

$$\sup_{\|f\|_{L^2(\mathbb{F}[t]_Q)}=1} \left\| \mathbb{E}_{y \in \mathbb{F}[t]_Q} f(x + P(y)) - \mathbb{E}_{z \in \mathbb{F}[t]_Q} f(z) \right\|_{L^2(\mathbb{F}[t]_Q)} = o_{\text{lpf}(Q) \rightarrow \infty}(1);$$

(ii) *there exist $C_1, C_2, \gamma > 0$ such that for any $Q(t) \in \mathbb{F}[t]^+$ with $\text{lpf}(Q) \geq C_1$, one has*

$$\sup_{\|f\|_{L^2(\mathbb{F}[t]_Q)}=1} \left\| \mathbb{E}_{y \in \mathbb{F}[t]_Q} f(x + P(y)) - \mathbb{E}_{z \in \mathbb{F}[t]_Q} f(z) \right\|_{L^2(\mathbb{F}[t]_Q)} \leq C_2 \cdot \text{lpf}(Q)^{-\gamma};$$

(iii) *there exists $C > 0$ such that if $Q(t) \in \mathbb{F}[t]^+$ and $\text{lpf}(Q) \geq C$, then $H + Q\mathbb{F}[t] = \mathbb{F}[t]$, where $H \leq (\mathbb{F}[t], +)$ is the group generated by $\{P(y) - P(0) : y \in \mathbb{F}[t]\}$.*

(iv) *for any $\delta > 0$, there exists $N > 0$ such that if $Q(t) \in \mathbb{F}[t]^+$ has $\text{lpf}(Q) \geq N$ and $A, B \subseteq \mathbb{F}[t]_Q$ are subsets with $|A||B| \geq \delta|Q|^2$, then there exist $x, y \in \mathbb{F}[t]_Q$ such that $x \in A$ and $x + P(y) \in B$;*

(v) *there exist $C_1, C_2, \gamma > 0$ such that for any $Q(t) \in \mathbb{F}[t]^+$ with $\text{lpf}(Q) \geq C_1$, one has*

$$\left| |\{(x, y) \in \mathbb{F}[t]_Q^2 : x \in A, x + P(y) \in B\}| - |A||B| \right| \leq C_2 |A|^{1/2} |B|^{1/2} |Q| \cdot \text{lpf}(Q)^{-\gamma}.$$

Moreover, if $P(y)$ is good for irrational equidistribution, then each of the properties (i)–(v) holds.

First we prove that irrational equidistribution implies condition (iii).


Proposition 5.1. *Suppose $P(y)$ is good for irrational equidistribution, and let $H \leq (\mathbb{F}[t], +)$ be the group generated by $\{P(y) - P(0) : y \in \mathbb{F}[t]\}$. Then there exists $C > 0$ such that if $Q(t) \in \mathbb{F}[t]^+$ satisfies $\text{lpf}(Q) \geq C$, then $H + Q\mathbb{F}[t] = \mathbb{F}[t]$. That is, (iii) holds.*

Proof. We prove the contrapositive. Suppose (iii) fails. Then there is a sequence $(Q_n)_{n \in \mathbb{N}}$ in $\mathbb{F}[t]^+$ such that $\text{lpf}(Q_n) \rightarrow \infty$ and $H + Q_n \mathbb{F}[t] \neq \mathbb{F}[t]$ for $n \in \mathbb{N}$. Equivalently,

$$H^\perp \cap (Q_n \mathbb{F}[t])^\perp = (H + Q_n \mathbb{F}[t])^\perp \neq \{0\}.$$

Since $(Q_n \mathbb{F}[t])^\perp \cong \widehat{\mathbb{F}[t]}_{Q_n}$, it follows that $\frac{s_n}{Q_n} \in H^\perp$ for some $s_n \not\equiv 0 \pmod{Q_n}$. If $\frac{s_n}{Q_n} = \frac{s'_n}{Q'_n}$ in reduced terms (i.e., s'_n and Q'_n are coprime), then $\text{lpf}(Q'_n) \geq \text{lpf}(Q_n)$, so we may assume without loss of generality that s_n and Q_n are coprime. The sequence $\left(\frac{s_n}{Q_n}\right)_{n \in \mathbb{N}}$ then consists of distinct elements, so H^\perp is infinite. Every infinite compact group is uncountable, and there are only countable many rational points, so H^\perp must contain an irrational element. That is, for some irrational α , $e(P(y)\alpha) = e(P(0)\alpha)$ for every $y \in \mathbb{F}[t]$. Hence, $(P(n)\alpha)_{n \in \mathbb{N}}$ is not well distributed, so $P(y)$ is not good for irrational equidistribution. \square

We will now prove the equivalences in Theorem 1.17 by showing the implications illustrated in the following diagram:



Condition (ii) is a quantitative refinement of condition (i), so we immediately have the implication (ii) \implies (i). By Theorem 1.10, we have the additional implications (i) \implies (iii) \implies (ii).

Condition (v) follows from (ii) by a straightforward application of the Cauchy–Schwarz inequality.

Proposition 5.2. $(v) \implies (iv)$.

Proof. Let $\delta > 0$. Let C_1 , C_2 , and γ be as in (v). Let $Q(t) \in \mathbb{F}[t]^+$ with $\text{lpf}(Q) \geq C_1$. Let $A, B \subseteq \mathbb{F}[t]_Q$ with $|A||B| \geq \delta|Q|^2$. Then by (v),

$$\begin{aligned}
 |\{(x, y) \in \mathbb{F}[t]_Q^2 : x \in A, x + P(y) \in B\}| &\geq |A||B| - C_2|A|^{1/2}|B|^{1/2}|Q| \cdot \text{lpf}(Q)^{-\gamma} \\
 &= |A|^{1/2}|B|^{1/2} \left(|A|^{1/2}|B|^{1/2} - C_2|Q| \cdot \text{lpf}(Q)^{-\gamma} \right) \\
 &\geq \delta^{1/2} \left(\delta^{1/2} - C_2 \cdot \text{lpf}(Q)^{-\gamma} \right) |Q|^2.
 \end{aligned}$$

Thus, if

$$\text{lpf}(Q) \geq N(\delta) = \max \left\{ C_1, \left(\frac{C_2}{\delta^{1/2}} \right)^{1/\gamma} \right\},$$

then we can find $x, y \in \mathbb{F}[t]_Q$ with $x \in A$ and $x + P(y) \in B$. \square

We now prove the final implication to complete the proof of Theorem 1.17:

Proposition 5.3. $(iv) \implies (iii)$.

Proof. We prove the contrapositive. Suppose (iii) fails. Then there is a sequence $(Q_n)_{n \in \mathbb{N}}$ in $\mathbb{F}[t]^+$ with $\text{lpf}(Q_n) \rightarrow \infty$ such that $\{P(y) - P(0) : y \in \mathbb{F}[t]_{Q_n}\} \subseteq H_{Q_n} \not\subseteq \mathbb{F}[t]_{Q_n}$ for every $n \in \mathbb{N}$. The subgroup H_{Q_n} has index p^{k_n} for some $k_n \in \mathbb{N}$. Let A_n be a union of $\lfloor \frac{p^{k_n}}{2} \rfloor$ cosets of H_{Q_n} , and let $B_n = \mathbb{F}[t]_{Q_n} \setminus (A_n + P(0))$. For any $x \in A_n$ and $y \in \mathbb{F}[t]_{Q_n}$, we have

$$x + P(y) = x + P(0) + (P(y) - P(0)) \in A_n + P(0) + H_{Q_n} = A_n + P(0).$$

That is, if $x \in A_n$ and $y \in \mathbb{F}[t]_{Q_n}$, then $x + P(y) \notin B_n$. Moreover,

$$\frac{|A_n|}{|Q_n|} = \frac{\lfloor \frac{p^{k_n}}{2} \rfloor}{p^{k_n}} \geq \frac{1}{3} \quad \text{and} \quad \frac{|B_n|}{|Q_n|} \geq \frac{1}{2}.$$

Therefore, property (iv) fails for $\delta = \frac{1}{6}$. \square

Now we proceed to prove the remaining equivalences in Theorem 1.18. As a first step, we have the following characterization of irrational equidistribution for additive polynomials:

Proposition 5.4. *Let $\eta(y) \in (\mathbb{F}[t])[y]$ be an additive polynomial. The following are equivalent:*

- (i) $\eta(y)$ is good for irrational equidistribution;
- (ii) there exists $C > 0$ such that if $Q(t) \in \mathbb{F}[t]^+$ and $\text{lpf}(Q) \geq C$, then $\eta(\mathbb{F}[t]_Q) = \mathbb{F}[t]_Q$;
- (iii) $\eta(y) = ay$ for some $a \in \mathbb{F}[t] \setminus \{0\}$.

Proof. (i) \implies (ii). See Proposition 5.1.

(ii) \implies (iii). We prove the contrapositive. Suppose $\eta(y) = \sum_{i=0}^k a_i y^{p^i}$ with $a_k \neq 0$, $k \geq 1$. We consider two cases separately.

Case 1: $a_0 = 0$.

In this case, we may write $\eta(y) = \eta'(y)^p$, where $\eta'(y) = \sum_{i=1}^k a_i y^{p^{i-1}}$. Hence, for any $Q(t) \in \mathbb{F}[t]^+$, $\eta(\mathbb{F}[t]_Q) \subseteq \{y^p : y \in \mathbb{F}[t]_Q\}$. If $Q = Q_0^2$, then $Q_0 \not\equiv 0 \pmod{Q}$, while $Q_0^p \equiv 0 \pmod{Q}$. Therefore, the homomorphism $y \mapsto y^p$ has a nontrivial kernel mod Q , so $\{y^p : y \in \mathbb{F}[t]_Q\}$ is a proper subgroup of $\mathbb{F}[t]_Q$. Taking Q_0 to be an arbitrarily large irreducible element of $\mathbb{F}[t]^+$, this shows that (ii) does not hold.

Case 2: $a_0 \neq 0$.

Write $\eta(y) = yg(y)$ with $g(y) = a_0 + \sum_{i=1}^k a_i y^{p^i-1}$. Since $g(y)$ is a nonconstant polynomial, the set

$$R = \{Q(t) \in \mathbb{F}[t]^+ : Q \text{ is irreducible and } g \text{ has a root mod } Q\}$$

is infinite. Indeed, for any finite collection of irreducibles $Q_1, \dots, Q_r \in \mathbb{F}[t]^+$, consider

$$(5.1) \quad g(a_0 Q_1 \dots Q_r y) = a_0 \left(1 + Q_1 \dots Q_r y \sum_{i=1}^k a_i (a_0 Q_1 \dots Q_r y)^{p^i-2} \right).$$

Since g is a nonzero polynomial, there exists $y_0 \in \mathbb{F}[t]$ such that $g(a_0 Q_1 \dots Q_r y_0) \neq 0$. From the expression on the right hand side of (5.1), we have $a_0 \mid g(a_0 Q_1 \dots Q_r y_0)$, and $\frac{g(a_0 Q_1 \dots Q_r y_0)}{a_0} \equiv 1 \pmod{Q_i}$ for each $i \in \{1, \dots, r\}$. Therefore, there is some irreducible $Q \in \mathbb{F}[t]^+ \setminus \{Q_1, \dots, Q_r\}$ such that $Q \mid g(a_0 Q_1 \dots Q_r y_0)$. Hence, R is infinite as claimed.

Suppose $Q \in R$ and $Q \nmid a_0$. Let $y \in \mathbb{F}[t]_Q$ such that $g(y) = 0$. Then $\eta(y) = yg(y) = 0$, but $y \neq 0$, since $y \mid a_0$. Hence, η has a nontrivial kernel mod Q for infinitely many irreducibles Q , which contradicts condition (ii).

(iii) \implies (i). See [BL16, Theorem 0.1]. \square

The following lemma is the key tool to reduce equidistributional properties of polynomials to the additive case with which we have just dealt.

Lemma 5.5. *Let $\eta_1, \eta_2 \in \mathbb{F}_p[y]$ be additive polynomials, and let $H_i = \eta_i(\mathbb{F}[t])$ for $i = 1, 2$. There exists an additive polynomial $\eta \in \mathbb{F}_p[y]$ such that $\eta(\mathbb{F}[t]) = H_1 + H_2$. Moreover, $\eta = \eta_1 \circ \zeta_1 + \eta_2 \circ \zeta_2$, where $\zeta_1, \zeta_2 \in \mathbb{F}_p[y]$ are additive polynomials.*

Proof. If $\eta_i = 0$ for some i , then take $\eta = \eta_j$ with $j \neq i$.

Suppose now that η_1 and η_2 are both nonzero. Let $H = H_1 + H_2$. Write $\eta_1(y) = \sum_{i=0}^k a_i y^{p^i}$ and $\eta_2(y) = \sum_{j=0}^l b_j y^{p^j}$. Without loss of generality, $k \geq l$. Define

$$(5.2) \quad \eta'_1(y) = b_l \eta_1(y) - a_k \eta_2(y^{p^{k-l}}) = \eta_1(b_l y) + \eta_2(-a_k y^{p^{k-l}}),$$

and let $H'_1 = \eta'_1(\mathbb{F}[t])$. Then $\deg \eta'_1 < \deg \eta_1$.

Claim: $H'_1 + H_2 = H_1 + H_2$.

For any $y \in \mathbb{F}[t]$, (5.2) expressed $\eta'_1(y)$ as a sum of an element of H_1 and an element of H_2 . Hence, $H'_1 \subseteq H_1 + H_2$. Rearranging (5.2), we have

$$\eta_1(y) = b_l^{-1} \eta'_1(y) + b_l^{-1} a_k \eta_2(y^{p^{k-l}}).$$

Thus, $H_1 \subseteq H'_1 + H_2$. This proves the claim.

We have shown that, given any nonzero additive polynomials $\eta_1, \eta_2 \in \mathbb{F}_p[y]$, we may find $\eta'_1, \eta'_2 \in \mathbb{F}_p[y]$ with $\eta'_1(\mathbb{F}[t]) + \eta'_2(\mathbb{F}[t]) = \eta_1(\mathbb{F}[t]) + \eta_2(\mathbb{F}[t])$ such that $\deg \eta'_1 + \deg \eta'_2 < \deg \eta_1 + \deg \eta_2$, and η'_1 and η'_2 are of the appropriate form. Repeating this process finitely many times, we eventually reduce to the situation that one of the additively polynomials is zero. We then take η to be the remaining nonzero polynomial. \square

The argument in the proof of Lemma 5.5 provides an algorithm for obtaining η that bears a strong resemblance with the Euclidean algorithm. We work through a few simple examples to see more concretely how the algorithm works.

Example 5.6. (1) $\eta_1(y) = y^{p^2} - y$, $\eta_2(y) = y^{p^3} + y^p$. The polynomial η_2 has larger degree, so we shift the exponents of η_1 to match the degree of η_2 and subtract:

$$\eta'_2(y) = \eta_2(y) - \eta_1(y^p) = 2y^p.$$

If $p = 2$, then $\eta'_2(y) = 0$, so we stop, and the resulting polynomial η is simply η_1 . (Note that when $p = 2$, η_1 may be rewritten as $\eta_1(y) = y^{p^2} + y$, and then it is clear that $\eta_2(y) = \eta_1(y^p)$, so the range of η_2 is manifestly a subset of the range of η_1 .) Suppose $p > 2$. Then $\deg \eta_1 > \deg \eta'_2$, so we shift the exponents of η'_2 and subtract:

$$\eta'_1(y) = 2\eta_1(y) - \eta'_2(y^p) = -2y.$$

Since $p > 2$, the element $-2 \in \mathbb{F}_p$ is invertible, so the image of η'_1 is all of $\mathbb{F}[t]$, and we are done: $\eta(y) = \eta'_1(y) = -2y$. (One can check that applying one more step of the algorithm would result in $\eta''_2 = 0$, indicating that the process has terminated.)

(2) $\eta_1(y) = y^{p^3} + y^{p^2} + y^p$, $\eta_2(y) = y^{p^2}$. First, shifting η_2 and subtracting, we have

$$\eta'_1(y) = \eta_1(y) - \eta_2(y^p) = y^{p^2} + y^p.$$

Next, subtracting η_2 without any shifting gives

$$\eta''_1(y) = \eta'_1(y) - \eta_2(y) = y^p.$$

Shifting η''_1 and subtracting from η_2 produces $\eta'_2 = 0$, so we are done and $\eta(y) = \eta''_1(y) = y^p$.

The following proposition completes the proof of Theorem 1.18:

Proposition 5.7. *Let $P(y) \in \mathbb{F}_p[y]$ and write $P(y) = a_0 + \sum_{i=1}^n \eta_i(y^{r_i})$ with $\eta_1, \dots, \eta_n \in \mathbb{F}_p[y]$ additive polynomials and $r_1, \dots, r_n \in \mathbb{N}$ distinct positive integers not divisible by p . The following are equivalent:*

- (i) $P(y)$ is good for irrational equidistribution;

- (ii) there exists $C > 0$ such that if $Q(t) \in \mathbb{F}[t]^+$ and $\text{lpf}(Q) \geq C$, then $H + Q\mathbb{F}[t] = \mathbb{F}[t]$, where $H \leq (\mathbb{F}[t], +)$ is the group generated by $\{P(y) - P(0) : y \in \mathbb{F}[t]\}$;
- (iii) there exist additive polynomials $\zeta_1, \dots, \zeta_n \in \mathbb{F}_p[y]$ and $a \in \mathbb{F}_p^\times$ such that

$$\sum_{i=1}^n (\eta_i \circ \zeta_i)(y) = ay.$$

Proof. (i) \implies (ii). See Theorem 1.17.

(ii) \implies (iii). For $Q(t) \in \mathbb{F}[t]^+$ and $i \in \{1, \dots, n\}$, let $H_{i,Q} = \eta_i(\mathbb{F}[t]_Q)$. By (ii), we have that $H_Q = \mathbb{F}[t]_Q$ whenever $\text{lpf}(Q) \geq C$. By Lemma 5.5, there is an additive polynomial of the form

$$\eta = \sum_{i=1}^n \eta_i \circ \zeta_i$$

such that $H_Q = \eta(\mathbb{F}[t]_Q)$ for every $Q(t) \in \mathbb{F}[t]^+$. In particular, $\eta(\mathbb{F}[t]_Q) = \mathbb{F}[t]_Q$ for all Q with $\text{lpf}(Q) \geq C$. Hence, by Proposition 5.4, $\eta(y) = ay$. That is, (iii) holds.

(iii) \implies (i). Let $\alpha \in \mathbb{F}((t^{-1})) \setminus \mathbb{F}(t)$. Let

$$\mathcal{F}_i = \overline{\eta_i(\mathbb{F}[t])\alpha}$$

By Theorem 1.9, $(P(y)\alpha)_{y \in \mathbb{F}[t]}$ is well distributed if and only if $\mathcal{F} = \sum_{i=1}^n \mathcal{F}_i$ is the full ‘torus’ $\mathbb{F}((t^{-1}))/\mathbb{F}[t]$. By (iii),

$$\mathcal{F} \ni \sum_{i=1}^n \eta_i(\zeta_i(y))\alpha = ay\alpha$$

for every $y \in \mathbb{F}[t]$. But $(ay\alpha)_{y \in \mathbb{F}[t]}$ is well-distributed (in particular, it is dense) mod $\mathbb{F}[t]$ (see [BL16, Theorem 0.1]), so $\mathcal{F} = \mathbb{F}((t^{-1}))/\mathbb{F}[t]$. Thus, $P(y)$ is good for irrational equidistribution. \square

6. PARTITION REGULARITY OF POLYNOMIAL EQUATIONS OVER FINITE FIELDS

In this section, we deduce additional applications of Theorem 1.10 to partition regularity of polynomial equations over finite fields. As a first step, we observe the following criterion for the existence of solutions to polynomial equations of the form $P_1(x) + P_2(y) + P_3(z) = c$ over finite fields:

Proposition 6.1. *Let $P_1, P_2, P_3 \in (\mathbb{F}[t])[x]$ be nonconstant polynomials with $P_i(0) = 0$. Let $d_i = \deg P_i$ and $k_i = d - \deg P_i$. Let $H_{q,i} = \langle P_i(x) : x \in \mathbb{F}_q \rangle \leq (\mathbb{F}_q, +)$, and let $H_q = H_{q,1} + H_{q,2} + H_{q,3}$. Then for any q and any $c \in H_q$,*

$$\left| \{(x, y, z) \in \mathbb{F}_q^3 : P_1(x) + P_2(y) + P_3(z) = c\} \right| = \frac{q^3}{|H_q|} + O_{q \rightarrow \infty; d_1, d_2, d_3, k_1, k_2, k_3} (q^{2-\gamma}),$$

where $\gamma = \min \left\{ \frac{1}{2^{k_i-1}} : i \in \{1, 2, 3\} \right\} > 0$. In particular, if q is sufficiently large, then

$$P_1(\mathbb{F}_q) + P_2(\mathbb{F}_q) + P_3(\mathbb{F}_q) = H_q.$$

Remark 6.2. The main term, $\frac{q^3}{|H_q|}$, expresses an asymptotic equidistribution property. There are q^3 choices of $(x, y, z) \in \mathbb{F}_q^3$ and $|H_q|$ possible values $c \in H_q$ that are free from obvious algebraic obstructions to solvability of $P_1(x) + P_2(y) + P_3(z) = c$. Proposition 6.1 states that, asymptotically, the polynomial $P_1(x) + P_2(y) + P_3(z)$ takes on each such value of c with roughly equal frequency.

Proof. For any prime power q , any $c \in \mathbb{F}_q$, and any functions $f_1, \dots, f_r : \mathbb{F}_q \rightarrow \mathbb{F}_q$, let $N(q, c; f_1, \dots, f_r)$ denote the number of solutions $(x_1, \dots, x_r) \in \mathbb{F}_q^r$ to the equation $\sum_{i=1}^r f_i(x_i) = c$. Our goal is to show

$$N(q, c; P_1, P_2, P_3) = \frac{q^3}{|H_q|} + O_{q \rightarrow \infty}(q^{2-\gamma})$$

for $c \in H_q$.

For each $i \in \{1, 2, 3\}$, let η_i be an additive polynomial with $\eta_i(\mathbb{F}_q) = H_{q,i}$.

Claim: $N(q, c; P_1, P_2, P_3) = N(q, c; P_1, P_2, \eta_3) + O_{q \rightarrow \infty; d_1, d_2, d_3, k_3}(q^{2-1/2^{k_3-1}})$.

Let $f_i(x) = N(q, x; P_i)$. Then

$$N(q, c; P_1, P_2, P_3) = \sum_{x, y \in \mathbb{F}_q} f_1(x) f_2(c - x - P_3(y)) = q^2 \mathbb{E}_{x, y \in \mathbb{F}_q} f_1(x) f_2(c - x - P_3(y)).$$

Similarly,

$$N(q, c; P_1, P_2, \eta_3) = q^2 \mathbb{E}_{x, y \in \mathbb{F}_q} f_1(x) f_2(c - x - \eta_3(y)) = q^2 \mathbb{E}_{x \in \mathbb{F}_q} \mathbb{E}_{z \in H_{q,3}} f_1(x) f_2(c - x - z).$$

Hence, by the Cauchy-Schwarz inequality,

$$|N(q, c; P_1, P_2, P_3) - N(q, c; P_1, P_2, \eta_3)| \leq q^2 \|f_1\|_{L^2(\mathbb{F}_q)} \left\| \mathbb{E}_{y \in \mathbb{F}_q} f_2(c - x - P_3(y)) - \mathbb{E}_{z \in H_{q,3}} f_2(c - x - z) \right\|_{L^2(\mathbb{F}_q)}.$$

Now, by Theorem 1.10,

$$\left\| \mathbb{E}_{y \in \mathbb{F}_q} f_2(c - x - P_3(y)) - \mathbb{E}_{z \in H_{q,3}} f_2(c - x - z) \right\|_{L^2(\mathbb{F}_q)} \ll_{d_3, k_3} q^{-1/2^{k_3-1}} \|f_2\|_{L^2(\mathbb{F}_q)}.$$

Finally, for each $x \in \mathbb{F}_q$, the polynomial equation $P_i(u) = x$ has at most d_i solutions $u \in \mathbb{F}_q$, so $\|f_i\|_{L^2(\mathbb{F}_q)} \leq \|f_i\|_{L^\infty(\mathbb{F}_q)} \leq d_i$. Putting everything together,

$$N(q, c; P_1, P_2, P_3) = N(q, c; P_1, P_2, \eta_3) + O_{q \rightarrow \infty; d_1, d_2, d_3, k_3}(q^{2-1/2^{k_3-1}})$$

as claimed.

Applying the claim also to P_1 and P_2 , we obtain the estimate

$$N(q, c; P_1, P_2, P_3) = N(q, c; \eta_1, \eta_2, \eta_3) + O_{q \rightarrow \infty; d_1, d_2, d_3, k_1, k_2, k_3}(q^{2-\gamma}).$$

Let $\eta : \mathbb{F}_q^3 \rightarrow \mathbb{F}_q$, $\eta(x, y, z) = \eta_1(x) + \eta_2(y) + \eta_3(z)$. Then η is a group homomorphism with image H_q . Therefore, $N(q, c; \eta_1, \eta_2, \eta_3) = |\eta^{-1}(\{c\})|$ is constant in $c \in H_q$, so

$$N(q, c; \eta_1, \eta_2, \eta_3) = \frac{|\mathbb{F}_q^3|}{|H_q|} = \frac{q^3}{|H_q|}.$$

□

Problems in the vein of Proposition 6.1 counting solutions to polynomial equations over finite fields are well-studied. For instance, if the equation $P_1(x) + P_2(y) + P_3(z) = c$ defines a geometrically irreducible variety, then a theorem of Lang and Weil [LW54] states that the number of solutions is equal to $q^2 + O_{q \rightarrow \infty}(q^{3/2})$. The class of polynomials handled by Proposition 6.1 is very restricted, and the error term in Proposition 6.1 is in general much weaker than what can be obtained with the use of algebraic geometry. However, the elementary method of proof allows us to avoid any irreducibility assumption and is more flexible for combinatorial enhancements, such as the following Ramsey-theoretic result, restated from the introduction:

Theorem 1.19. *Let $P(x) \in \mathbb{F}_p[x]$ be a nonconstant polynomial, and let $Q(x) \in \mathbb{F}_p[x]$ be good for irrational equidistribution. For any $r \in \mathbb{N}$, there exists $K = K(P, Q, r) \in \mathbb{N}$ and $c = c(P, Q, r) > 0$ such that for any $k \geq K$ and any r -coloring $\mathbb{F}_{p^k} = \bigcup_{i=1}^r C_i$, there are at least $c(p^k)^2$ monochromatic solutions to the equation $P(x) - P(y) = Q(z)$. That is,*

$$\left| \left\{ (x, y, z) \in \mathbb{F}_{p^k}^3 : P(x) - P(y) = Q(z) \text{ and } \{x, y, z\} \subseteq C_i \text{ for some } i \in \{1, \dots, r\} \right\} \right| \geq c(p^k)^2.$$

Our proof of Theorem 1.19 combines Theorem 1.10 with tools from the theory of Loeb measures on ultraproduct spaces. We will need the following generalization of Theorem 1.10, which in fact comes as an easy consequence of the key estimate on character sums in Proposition 3.2:

Proposition 6.3. *Let $P(y) \in (\mathbb{F}[t])[y]$ be a nonconstant polynomial of degree d and derivational degree k . Write $P(y) = a_0 + \sum_{i=1}^n \eta_i(y^{r_i})$. Let $H_i = \eta_i(\mathbb{F}[t])$ and $H = \sum_{i=1}^n H_i$. Then for any $Q(t) \in \mathbb{F}[t]^+$, any $m \in \mathbb{N}$, and any $f : \mathbb{F}[t]_Q^m \rightarrow \mathbb{C}$,*

$$\begin{aligned} & \left\| \mathbb{E}_{y \in \mathbb{F}[t]_Q} f(x_1 + P(y), \dots, x_m + P(y)) - \mathbb{E}_{z \in H_Q} f(x_1 + a_0 + z, \dots, x_m + a_0 + z) \right\|_{L^2(\mathbb{F}[t]_Q^m)} \\ & \leq \left(p^{2 \lfloor \log_p d \rfloor} \frac{k-1}{\text{lpf}(Q)} \right)^{1/2^{k-1}} \|f\|_{L^2(\mathbb{F}[t]_Q^m)}, \end{aligned}$$

where $H_Q = \{z \pmod{Q} : z \in H\}$.

Proof. Expand f as a Fourier series:

$$f(x_1, \dots, x_m) = \sum_{s_1, \dots, s_m \in \mathbb{F}[t]_Q} \widehat{f}(s_1, \dots, s_m) e \left(\sum_{j=1}^m s_j x_j / Q \right).$$

Then for $\mathbf{x} = (x_1, \dots, x_m) \in \mathbb{F}[t]_Q^m$, we have

$$\begin{aligned} F(\mathbf{x}) &:= \mathbb{E}_{y \in \mathbb{F}[t]_Q} f(x_1 + P(y), \dots, x_m + P(y)) - \mathbb{E}_{z \in H_Q} f(x_1 + a_0 + z, \dots, x_m + a_0 + z) \\ &= \sum_{\mathbf{s} \in \mathbb{F}[t]_Q^m} \widehat{f}(\mathbf{s}) e(\mathbf{s} \cdot \mathbf{x} / Q) \left(\mathbb{E}_{y \in \mathbb{F}[t]_Q} e \left(\sum_{j=1}^m s_j P(y) / Q \right) - e \left(\sum_{j=1}^m s_j a_0 \right) \mathbb{1}_{H_Q^\perp} \left(\sum_{j=1}^m s_j \right) \right). \end{aligned}$$

For each $\mathbf{s} \in \mathbb{F}[t]_Q^m$, Proposition 3.2 gives the bound

$$\left| \mathbb{E}_{y \in \mathbb{F}[t]_Q} e \left(\sum_{j=1}^m s_j P(y) / Q \right) - e \left(\sum_{j=1}^m s_j a_0 \right) \mathbb{1}_{H_Q^\perp} \left(\sum_{j=1}^m s_j \right) \right| \leq \left(p^{2 \lfloor \log_p d \rfloor} \frac{k-1}{\text{lpf}(Q)} \right)^{1/2^{k-1}}.$$

Therefore, by Parseval's identity,

$$\|F\|_{L^2(\mathbb{F}[t]_Q^m)}^2 \leq \left(p^{2 \lfloor \log_p d \rfloor} \frac{k-1}{\text{lpf}(Q)} \right)^{1/2^{k-2}} \sum_{\mathbf{s} \in \mathbb{F}[t]_Q^m} |\widehat{f}(\mathbf{s})|^2 = \left(p^{2 \lfloor \log_p d \rfloor} \frac{k-1}{\text{lpf}(Q)} \right)^{1/2^{k-2}} \|f\|_{L^2(\mathbb{F}[t]_Q^m)}^2.$$

□

The relevant constructions for employing measure theory on ultraproducts are summarized as follows:

Definition 6.4.

- An *ultrafilter* on \mathbb{N} is a collection $\mathcal{U} \subseteq \mathcal{P}(\mathbb{N})$ of nonempty subsets of \mathbb{N} such that:
 - if $A, B \in \mathcal{U}$, then $A \cap B \in \mathcal{U}$;
 - for any $A \subseteq \mathbb{N}$, either $A \in \mathcal{U}$ or $\mathbb{N} \setminus A \in \mathcal{U}$.

The ultrafilter \mathcal{U} is *principal* if $\mathcal{U} = \{A \subseteq \mathbb{N} : n \in A\}$ for some $n \in \mathbb{N}$ and *non-principal* otherwise. The space of ultrafilters is denoted $\beta\mathbb{N}$.

- Given $\mathcal{U} \in \beta\mathbb{N}$ and a family of sets $(X_n)_{n \in \mathbb{N}}$, the *ultraproduct* is the set

$$\prod_{n \rightarrow \mathcal{U}} X_n = \left(\prod_{n \in \mathbb{N}} X_n \right) / \equiv_{\mathcal{U}},$$

where $\equiv_{\mathcal{U}}$ is the equivalence relation defined by $(x_n)_{n \in \mathbb{N}} \equiv_{\mathcal{U}} (y_n)_{n \in \mathbb{N}}$ if and only if $\{n \in \mathbb{N} : x_n = y_n\} \in \mathcal{U}$.

- Given $\mathcal{U} \in \beta\mathbb{N}$ and a sequence $(x_n)_{n \in \mathbb{N}}$ taking values in a compact Hausdorff space X , the *limit of $(x_n)_{n \in \mathbb{N}}$ along \mathcal{U}* is defined to be the unique point⁶ $x \in X$ such that for any neighborhood U of x , one has $\{n \in \mathbb{N} : x_n \in U\} \in \mathcal{U}$. The limit of $(x_n)_{n \in \mathbb{N}}$ along \mathcal{U} is denoted by $\lim_{n \rightarrow \mathcal{U}} x_n$.
- Let $\mathcal{U} \in \beta\mathbb{N}$, and let $(X_n, \mathcal{X}_n, \mu_n)_{n \in \mathbb{N}}$ be a family of probability spaces. Let $X = \prod_{n \rightarrow \mathcal{U}} X_n$.
 - An *internal set* is a set of the form $\prod_{n \rightarrow \mathcal{U}} A_n$ with $A_n \in \mathcal{X}_n$.
 - The *Loeb σ -algebra \mathcal{X}* is the σ -algebra on $\prod_{n \rightarrow \mathcal{U}} X_n$ generated by the algebra of internal sets.
 - The *Loeb measure μ* is the unique probability measure on \mathcal{X} with the property

$$\mu(A) = \lim_{n \rightarrow \mathcal{U}} \mu_n(A_n)$$

for any internal set $A = \prod_{n \rightarrow \mathcal{U}} A_n$.

The main property of the Loeb measure that we will use is the following version of Fubini's theorem:

Proposition 6.5 (cf. [K77], Theorem 1.12). *Let $(X_n)_{n \in \mathbb{N}}$ and $(Y_n)_{n \in \mathbb{N}}$ be sequences of finite sets. Let \mathcal{U} be a non-principal ultrafilter. Let $X = \prod_{n \rightarrow \mathcal{U}} X_n$ and $Y = \prod_{n \rightarrow \mathcal{U}} Y_n$. Let $f : X \times Y \rightarrow \mathbb{C}$ be a bounded Loeb-measurable function. Then*

- (1) *for any $x \in X$, the function $y \mapsto f(x, y)$ is Loeb-measurable on Y ;*
- (2) *the function $x \mapsto \int_Y f(x, y) d\mu_Y(y)$ is Loeb-measurable on X ; and*
- (3)

$$\int_{X \times Y} f d\mu_{X \times Y} = \int_X \left(\int_Y f(x, y) d\mu_Y(y) \right) d\mu_X(x).$$

Remark 6.6. Proposition 6.5 does not follow from standard version of Fubini's theorem. The subtlety lies in the structure of the Loeb σ -algebra on the product space $X \times Y$: there are internal subsets of $X \times Y$ that cannot be approximated by Boolean combinations of Cartesian products of internal subsets of X and Y (on the finitary level, this corresponds to approximating subsets of $X_n \times Y_n$ by products of boundedly many subsets of X_n and Y_n). Therefore, the function f need not be measurable with respect to the product of the Loeb σ -algebras on X and Y . Nevertheless, Proposition 6.5 shows that $\mu_{X \times Y}$ shares important features with the product measure $\mu_X \times \mu_Y$.

Proof of Theorem 1.19. Let $r \in \mathbb{N}$. Suppose for contradiction that there are r -colorings $\mathbb{F}_{p^{k_n}} = \bigcup_{i=1}^r C_{n,i}$ with $k_n \rightarrow \infty$ such that $|M_n| = o_{n \rightarrow \infty} \left((p^{k_n})^2 \right)$, where

$$M_n = \left\{ (x, y, z) \in \mathbb{F}_{p^{k_n}}^3 : P(x) - P(y) = Q(z) \text{ and } \{x, y, z\} \subseteq C_{n,i} \text{ for some } i \in \{1, \dots, r\} \right\}$$

is the collection of monochromatic solutions to the equation $P(x) - P(y) = Q(z)$.

Now we define a limit object associated with this sequence of colorings. Fix a non-principal ultrafilter \mathcal{U} on \mathbb{N} . Let \mathbb{F}_{∞} be the pseudo-finite field $\mathbb{F}_{\infty} = \prod_{n \rightarrow \mathcal{U}} \mathbb{F}_{p^{k_n}}$, let $C_i = \prod_{n \rightarrow \mathcal{U}} C_{n,i} \subseteq \mathbb{F}_{\infty}$, and let

⁶Such a point x exists by compactness and is unique by the Hausdorff property.

$M = \prod_{n \rightarrow \mathcal{U}} M_n$. Denote by μ the Loeb measure on \mathbb{F}_∞ obtained by equipping $\mathbb{F}_{p^{k_n}}$ with the normalized counting measure. For any $s \in \mathbb{N}$, we denote the Loeb measure on \mathbb{F}_∞^s by μ^s (not be confused with the product measure $\mu \times \cdots \times \mu$ on \mathbb{F}_∞^s). Let $V_n = \left\{ (x, y, z) \in \mathbb{F}_{p^{k_n}}^3 : P(x) - P(y) = Q(z) \right\}$ and $V = \prod_{n \rightarrow \mathcal{U}} V_n$. Finally, let μ_V be the Loeb measure on V obtained from the normalized counting measures on V_n .

Claim 1: $\mathbb{F}_\infty = \bigcup_{i=1}^r C_i$.

Let $x = (x_n)_{n \in \mathbb{N}} \in \mathbb{F}_\infty$. For $i \in \{1, \dots, r\}$, let $I_i = \{n \in \mathbb{N} : x_n \in C_{n,i}\}$. Then $\mathbb{N} = \bigcup_{i=1}^r I_i$, so $I_{i_0} \in \mathcal{U}$ for some $i_0 \in \{1, \dots, r\}$, since \mathcal{U} is an ultrafilter. By the definition of the sets C_1, \dots, C_r , it follows that $x \in C_{i_0}$. This proves the claim.

Arguing as in the proof of Claim 1 above, one can check that M is the set of monochromatic solutions $(x, y, z) \in \mathbb{F}_\infty^3$ to the equation $P(x) - P(y) = Q(z)$ with respect to the coloring $\mathbb{F}_\infty = \bigcup_{i=1}^r C_i$.

Claim 2: $\mu_V(M) = 0$.

We have constructed M as an internal set, so by the definition of the Loeb measure,

$$\mu_V(M) = \lim_{n \rightarrow \mathcal{U}} \frac{|M_n|}{|V_n|}.$$

Now, by Proposition 6.1, $|V_n| = (p^{k_n})^2 + O_{n \rightarrow \infty}((p^{k_n})^{2-\gamma})$ for some $\gamma > 0$. By assumption, $|M_n| = o_{n \rightarrow \infty}((p^{k_n})^2)$. Hence, $\frac{|M_n|}{|V_n|} = o_{n \rightarrow \infty}(1)$, so $\mu_V(M) = 0$, since \mathcal{U} is non-principal.

Let $A_i = P(C_i) = \prod_{n \rightarrow \mathcal{U}} P(C_{n,i})$. Note that

$$\frac{1}{d} \mu(C_i) \leq \mu(A_i) \leq \mu(C_i),$$

where $d = \deg P$. In particular, $\mu(A_i) = 0$ if and only if $\mu(C_i) = 0$.

Without loss of generality, we may assume that $\mu(C_i) > 0$ for $1 \leq i \leq s$ and $\mu(C_i) = 0$ for $s+1 \leq i \leq r$, for some $s \in \{1, \dots, r\}$. Let $A = A_1 \times \cdots \times A_s \subseteq \mathbb{F}_\infty^s$. Note that $\mu^s(A) = \prod_{i=1}^s \mu(A_i) > 0$. Let $T_z : \mathbb{F}_\infty^s \rightarrow \mathbb{F}_\infty^s$ be the map $T_z \mathbf{x} = (x_1 + z, \dots, x_s + z)$ for $z \in \mathbb{F}_\infty$, $\mathbf{x} = (x_1, \dots, x_s) \in \mathbb{F}_\infty^s$. For each $n \in \mathbb{N}$ and $i \in \{1, \dots, r\}$, let $A_{n,i} = P(C_{n,i})$, and let $A^{(n)} = A_{n,1} \times \cdots \times A_{n,s} \in \mathbb{F}_{p^{k_n}}^s$. Also let $T_z^{(n)} : \mathbb{F}_{p^{k_n}}^s \rightarrow \mathbb{F}_{p^{k_n}}^s$ be the map $T_z^{(n)} \mathbf{x} = (x_1 + z, \dots, x_s + z)$ for $z \in \mathbb{F}_{p^{k_n}}$ and $\mathbf{x} = (x_1, \dots, x_s) \in \mathbb{F}_{p^{k_n}}^s$. Now, since Q is good for irrational equidistribution, we have

$$(6.1) \quad \mathbb{E}_{\mathbf{x} \in \mathbb{F}_{p^{k_n}}^s} \mathbb{E}_{z \in \mathbb{F}_{p^{k_n}}} \mathbb{1}_{A^{(n)}}(\mathbf{x}) \mathbb{1}_{A^{(n)}}(T_{Q(z)}^{(n)} \mathbf{x}) = \mathbb{E}_{\mathbf{x} \in \mathbb{F}_{p^{k_n}}^s} \mathbb{E}_{y \in \mathbb{F}_{p^{k_n}}} \mathbb{1}_{A^{(n)}}(\mathbf{x}) \mathbb{1}_{A^{(n)}}(T_y^{(n)} \mathbf{x}) + o_{n \rightarrow \infty}(1)$$

by Proposition 6.3 and the Cauchy–Schwarz inequality. Hence,

$$\begin{aligned} \int_{\mathbb{F}_\infty} \mu^s(A \cap T_{Q(z)} A) d\mu(z) &\stackrel{(1)}{=} \int_{\mathbb{F}_\infty^{s+1}} \mathbb{1}_A(\mathbf{x}) \mathbb{1}_A(T_{Q(z)} \mathbf{x}) d\mu^{s+1}(\mathbf{x}, z) \\ &\stackrel{(2)}{=} \lim_{n \rightarrow \mathcal{U}} \mathbb{E}_{(\mathbf{x}, z) \in \mathbb{F}_{p^{k_n}}^{s+1}} \mathbb{1}_{A^{(n)}}(\mathbf{x}) \mathbb{1}_{A^{(n)}}(T_{Q(z)}^{(n)} \mathbf{x}) \\ &\stackrel{(3)}{=} \lim_{n \rightarrow \mathcal{U}} \mathbb{E}_{(\mathbf{x}, y) \in \mathbb{F}_{p^{k_n}}^{s+1}} \mathbb{1}_{A^{(n)}}(\mathbf{x}) \mathbb{1}_{A^{(n)}}(T_y^{(n)} \mathbf{x}) \\ &\stackrel{(4)}{\geq} \lim_{n \rightarrow \mathcal{U}} \left(\frac{|A^{(n)}|}{p^{sk_n}} \right)^2 \\ &\stackrel{(5)}{=} \mu^s(A)^2 > 0. \end{aligned}$$

The steps are justified as follows. Step (1) is a direct application of Proposition 6.5. The equality (2) comes from the definition of the Loeb measure μ^{s+1} . In step (3), we have taken the limit of both sides of (6.1) along \mathcal{U} . The inequality (4) holds for each $n \in \mathbb{N}$ by the Cauchy–Schwarz inequality:

$$\begin{aligned} \mathbb{E}_{(\mathbf{x}, y) \in \mathbb{F}_p^{s+1}} \mathbb{1}_{A^{(n)}}(\mathbf{x}) \mathbb{1}_{A^{(n)}}(T_y^{(n)} \mathbf{x}) &= \left\langle \mathbb{1}_{A^{(n)}}, \mathbb{E}_{y \in \mathbb{F}_p^{k_n}} T_y^{(n)} \mathbb{1}_{A^{(n)}} \right\rangle \\ &\geq \left\| \mathbb{E}_{y \in \mathbb{F}_p^{k_n}} T_y^{(n)} \mathbb{1}_{A^{(n)}} \right\|_{L^2(\mathbb{F}_p^{k_n})}^2 \\ &\geq \left(\frac{|A^{(n)}|}{p^{sk_n}} \right)^2. \end{aligned}$$

Finally, (5) follows from the definition of the Loeb measure μ^s .

Thus,

$$\mu(\{z \in \mathbb{F}_\infty : \mu^s(A \cap T_{Q(z)}A) > 0\}) > 0.$$

Let $G = \{z \in \bigcup_{i=1}^s C_i : \mu^s(A \cap T_{Q(z)}A) > 0\}$. Since the set $\bigcup_{i=s+1}^r C_i$ has Loeb measure zero, $\mu(G) > 0$.

For $z \in G$, let $i(z) \in \{1, \dots, s\}$ such that $z \in C_{i(z)}$. Noting that

$$\mu^s(A \cap T_{Q(z)}A) = \prod_{i=1}^s \mu(A_i \cap (A_i + Q(z))),$$

it follows that

$$\mu(A_{i(z)} \cap (A_{i(z)} + Q(z))) > 0.$$

The set $A_{i(z)}$ lies in the image of P by definition, so taking the inverse image under P ,

$$\mu(C_{i(z)} \cap P^{-1}(A_{i(z)} + Q(z))) > 0.$$

Therefore,

$$\begin{aligned} \mu_V(M) &= \lim_{n \rightarrow \mathcal{U}} \frac{|M_n|}{|V_n|} \\ &= \lim_{n \rightarrow \mathcal{U}} \frac{|M_n|}{(p^{k_n})^2} \\ &= \lim_{n \rightarrow \mathcal{U}} \frac{1}{(p^{k_n})^2} \sum_{i=1}^r \sum_{z \in \mathbb{F}_p^{k_n}} \mathbb{1}_{C_{n,i}}(z) |\{(x, y) \in C_{n,i}^2 : P(x) - P(y) = Q(z)\}| \\ &\geq \lim_{n \rightarrow \mathcal{U}} \sum_{i=1}^r \mathbb{E}_{z \in \mathbb{F}_p^{k_n}} \mathbb{1}_{C_{n,i}}(z) \frac{|C_{n,i} \cap P^{-1}(A_{n,i} + Q(z))|}{p^{k_n}} \\ &= \sum_{i=1}^r \int_{\mathbb{F}_\infty} \mathbb{1}_{C_i}(z) \mu(C_i \cap P^{-1}(A_i + Q(z))) \\ &\geq \int_G \mu(C_{i(z)} \cap P^{-1}(A_{i(z)} + Q(z))) d\mu(z) \\ &> 0. \end{aligned}$$

This final inequality contradicts Claim 2, so the theorem follows by *reductio ad absurdum*. \square

The proof of Theorem 1.19 is sufficiently flexible as to allow for many variations. We note that in the proof of Theorem 1.19, we do not use the full strength of the equidistribution assumption on Q . Carefully following each step in the proof reveals the following characterization of polynomials $Q(x) \in \mathbb{F}_p[x]$ for which the conclusion of Theorem 1.19 holds:

Corollary 6.7. *Let $Q(x) \in \mathbb{F}_p[x]$. Write $Q(x) = Q(0) + \sum_{i=1}^n \eta_i(x^{r_i})$ with r_1, \dots, r_n distinct with $p \nmid r_i$ and $\eta_1, \dots, \eta_n \in \mathbb{F}_p[x]$ additive polynomials. For $k \in \mathbb{N}$, let $H_i^{(k)} = \eta_i(\mathbb{F}_{p^k})$ and $H^{(k)} = \sum_{i=1}^n H_i^{(k)}$. The following are equivalent:*

- (i) *there exists $k_0 \in \mathbb{N}$ such that $Q(0) \in H^{(k)}$ for all $k \geq k_0$;*
- (ii) *for any nonconstant polynomial $P(x) \in \mathbb{F}_p[x]$ and any $r \in \mathbb{N}$, there exist $K = K(P, r) \in \mathbb{N}$ and $c(P, r) > 0$ such that for any $k \geq K$ and any r -coloring $\mathbb{F}_{p^k} = \bigcup_{i=1}^r C_i$, there are at least $c(p^k)^2$ monochromatic solutions to the equation $P(x) - P(y) = Q(z)$.*

Remark 6.8. Condition (i) in Corollary 6.7 holds for intersective Q ; see the first two lines in the proof of Corollary 1.12.

Proof. (i) \implies (ii). The proof of Theorem 1.19 goes through with only minor changes, which we will now describe. Equation (6.1) must be replaced with

$$\mathbb{E}_{\mathbf{x} \in \mathbb{F}_{p^{kn}}^s} \mathbb{E}_{z \in \mathbb{F}_{p^{kn}}} \mathbb{1}_{A^{(n)}}(\mathbf{x}) \mathbb{1}_{A^{(n)}}(T_{Q(z)}^{(n)} \mathbf{x}) = \mathbb{E}_{\mathbf{x} \in \mathbb{F}_{p^{kn}}^s} \mathbb{E}_{y \in H^{(kn)}} \mathbb{1}_{A^{(n)}}(\mathbf{x}) \mathbb{1}_{A^{(n)}}(T_y^{(n)} \mathbf{x}) + o_{n \rightarrow \infty}(1).$$

This follows from Proposition 6.3 together with the assumption (i) to eliminate the presence of the constant term $Q(0)$. We then note that the Cauchy–Schwarz inequality implies

$$\mathbb{E}_{\mathbf{x} \in \mathbb{F}_{p^{kn}}^s} \mathbb{E}_{y \in H^{(kn)}} \mathbb{1}_{A^{(n)}}(\mathbf{x}) \mathbb{1}_{A^{(n)}}(T_y^{(n)} \mathbf{x}) \geq \left(\frac{|A^{(n)}|}{p^{skn}} \right)^2$$

for each $n \in \mathbb{N}$. The steps (3) and (4) in the proof of Theorem 1.19 may be replaced by the above considerations.

Finally, by Proposition 6.1, the set $V_n = \left\{ (x, y, z) \in \mathbb{F}_{p^{kn}}^3 : P(x) - P(y) = Q(z) \right\}$ has cardinality $|V_n| = \frac{(p^{kn})^3}{|H_n|}$, where H_n is the subgroup generated by $H^{(kn)}$ and $\langle P(x) : x \in \mathbb{F}_{p^{kn}} \rangle$. By assumption, P is nonconstant, so $|H_n| \geq |P(\mathbb{F}_{p^{kn}})| \geq \frac{p^{kn}}{d}$, where $d = \deg P$. Hence, $(p^{kn})^2 \leq |V_n| \leq d(p^{kn})^2$. This upper and lower bound on the cardinality of V_n allow for the remainder of the argument to be carried out without difficulty.

(ii) \implies (i). We prove the contrapositive. Suppose $Q(0) \notin H^{(k)}$, and take $P = Q$. Then for any $x, y, z \in \mathbb{F}_{p^k}$, we have

$$\begin{aligned} P(x) - P(y) - Q(z) &= \left(Q(0) + \sum_{i=1}^n \eta_i(x^{r_i}) \right) - \left(Q(0) + \sum_{i=1}^n \eta_i(y^{r_i}) \right) - \left(Q(0) + \sum_{i=1}^n \eta_i(z^{r_i}) \right) \\ &\in -Q(0) + H^{(k)}, \end{aligned}$$

so $P(x) - P(y) = Q(z)$ does not have any solutions over \mathbb{F}_{p^k} , much less monochromatic solutions for an arbitrary coloring of \mathbb{F}_{p^k} . \square

Another feature of the proof of Theorem 1.19 is the following. Taking the ultraproduct of a sequence of finite fields \mathbb{F}_q with characteristic growing to infinity, the same method shows that for any nonconstant polynomials $P(x), Q(x) \in \mathbb{Z}[x]$, the equation $P(x) - P(y) = Q(z)$ is partition regular over all fields of sufficiently high characteristic. This follows by noting that P and Q will be nonconstant and separable (hence good for irrational equidistribution; see Example 1.16(1) above)

once the characteristic exceeds the degrees of P and Q and the size of some nonconstant coefficient.

In the special case $P = Q$, Theorem 1.19 can be seen as a polynomial version of Schur's theorem over finite fields. Indeed, the classical theorem of Schur asserts that the equation $x + y = z$ is partition regular over \mathbb{N} . We have just established partition regularity of the equation $P(x) + P(y) = P(z)$ over finite fields whenever P satisfies the equidistribution assumption in item (i) of Corollary 6.7. While the condition (i) in Corollary 6.7 depends on the characteristic p , it is automatically satisfied for polynomials with zero constant term. Hence, Corollary 1.20 holds.

The equation $P(x) - P(y) = Q(z)$ is often not partition regular (and may not even be solvable) over \mathbb{N} . A key fact leveraged in the proof of Theorem 1.19 is that polynomials take on a positive proportion of values in finite fields, something that is far from the case in \mathbb{N} . It remains an interesting and difficult open problem, asked by Erdős and Graham in [EG80], whether the Pythagorean equation $x^2 + y^2 = z^2$ is partition regular over \mathbb{N} . (This was settled with a computer-assisted proof in the case of 2-colorings in [HKM16] but is wide open for 3 or more colors.)

Some comments are in order on the use of ultraproducts in the proofs of the aforementioned partition regularity results. The basic strategy we have taken is to discard those colors that have zero Loeb measure in the ultraproduct and then to use recurrence along the polynomial Q to find the desired points x, y, z with $P(x) - P(y) = Q(z)$. One may be tempted to carry out this strategy in purely finitary terms, avoiding the use of ultraproducts and Loeb measure. Unfortunately, this does not work (at least in its most straightforward implementation). The following discussion illuminates the issues that arise. Fix a polynomial $Q(x) \in \mathbb{F}_p[x]$ that is good for irrational equidistribution. For simplicity, we will consider $P(x) = x$. Let $r \in \mathbb{N}$. Suppose $k \in \mathbb{N}$ is large and an r -coloring $\mathbb{F}_{p^k} = \bigcup_{i=1}^r C_i$ is given. We wish to use a function $\Phi : \mathbb{N} \rightarrow [0, \frac{1}{r}]$ as a cutoff for distinguishing "large" from "small" color classes. That is, we will consider a color class C_i large if $|C_i| \geq \Phi(k)p^k$ and small if $|C_i| < \Phi(k)p^k$. Without loss of generality, we may assume C_1, \dots, C_s are large and C_{s+1}, \dots, C_r are small for some $s \in \{1, \dots, r\}$. (The requirement that $\Phi(k) \leq \frac{1}{r}$ guarantees that at least one color class is large.) We now proceed as in the proof of Theorem 1.19, using "large" as a replacement for having positive Loeb measure. Let $A = C_1 \times \dots \times C_s$. Proposition 6.3 gives the bound

$$\mathbb{E}_{z \in \mathbb{F}_{p^k}} \frac{|A \cap (A + (Q(z), \dots, Q(z)))|}{p^{sk}} \geq \left(\frac{|A|}{p^{sk}} \right)^2 + O_{k \rightarrow \infty} (p^{-\gamma k})$$

where $\gamma = 2^{-(d - \deg Q - 1)}$. Since $|A \cap (A + (Q(z), \dots, Q(z)))| \leq |A|$ for each $z \in \mathbb{F}_{p^k}$, we deduce that

$$\begin{aligned} |\{z \in \mathbb{F}_{p^k} : A \cap (A + (Q(z), \dots, Q(z))) \neq \emptyset\}| &\geq \frac{p^{(s+1)k}}{|A|} \mathbb{E}_{z \in \mathbb{F}_{p^k}} \frac{|A \cap (A + (Q(z), \dots, Q(z)))|}{p^{sk}} \\ &\geq \frac{|A|}{p^{(s-1)k}} + O_{k \rightarrow \infty} \left(\frac{p^{(s+1)k}}{|A|} p^{-\gamma k} \right) \\ &\geq \Phi(k)^s p^k + O_{k \rightarrow \infty} \left(\Phi(k)^{-s} p^{(1-\gamma)k} \right), \end{aligned}$$

where in the last step we have used the bound $|A| \geq \Phi(k)^s p^{sk}$. In order to complete the argument, we want to find $z \in \bigcup_{i=1}^s C_i$ satisfying $A \cap (A + (Q(z), \dots, Q(z))) \neq \emptyset$. To that end, one would like to show

$$|\{z \in \mathbb{F}_{p^k} : A \cap (A + (Q(z), \dots, Q(z))) \neq \emptyset\}| > \left| \bigcup_{i=s+1}^r C_i \right|.$$

The total size of the small color classes is bounded by

$$\left| \bigcup_{i=s+1}^r C_i \right| \leq (r-1)\Phi(k)p^k.$$

The goal, then, is to choose the function $\Phi : \mathbb{N} \rightarrow [0, \frac{1}{r}]$ so that

$$\Phi(k)^s p^k > (r-1)\Phi(k)p^k + O_{k \rightarrow \infty} \left(\Phi(k)^{-s} p^{(1-\gamma)k} \right).$$

Dividing by p^k , this reduces to the inequality

$$\Phi(k)^s > (r-1)\Phi(k) + O_{k \rightarrow \infty} \left(\Phi(k)^{-s} p^{-\gamma k} \right).$$

But for $r \geq 2$, this requires $\Phi(k) > 1$, which violates the condition that $0 \leq \Phi(k) \leq \frac{1}{r}$.

Working with the ultraproduct allows us to replace “small” with measure zero. This is crucial, as we have just seen that “small” contributions in the finitary setting may accumulate and overtake individual “large” terms. In contrast, finite unions of measure zero sets remain of measure zero. However, our infinitary methods come at a cost: we are unable to provide any quantitative control on the values K and c appearing in the statement of Theorem 1.19 and related corollaries. It is therefore an interesting problem to obtain a purely finitary proof of Theorem 1.19 with effective bounds.

ACKNOWLEDGEMENTS

The first author was supported by the National Science Foundation under Grant No. DMS-1926686. We thank Peter Sarnak for pointing us to the work of Lang and Weil [LW54] and for insightful discussions that helped shape Section 6.

REFERENCES

- [BB23] Vitaly Bergelson and Andrew Best. The Furstenberg-Sárközy theorem and asymptotic total ergodicity phenomena in modular rings. *J. Number Theory* **243** (2023) 615–645.
- [BBI21] Vitaly Bergelson, Andrew Best, and Alex Iosevich. Sums of powers in large finite fields: a mix of methods. *Amer. Math. Monthly* **128** (2021) 701–718.
- [BL16] V. Bergelson and A. Leibman. A Weyl-type equidistribution theorem in finite characteristic. *Adv. Math.* **289** (2016) 928–950.
- [CGS12] Péter Csikvári, Katalin Gyarmati, and András Sárközy. Density and Ramsey type results on algebraic equations with restricted solution sets. *Combinatorica* **32** (2012) 425–449.
- [CLP17] Ernie Croot, Vsevolod F. Lev, and Péter Pál Pach. Progression-free sets in \mathbb{Z}_4^n are exponentially small. *Ann. of Math. (2)* **185** (2017) 331–337.
- [DLMS23] Sebastián Donoso, Anh N. Le, Joel Moreira, and Wenbo Sun. Additive averages of multiplicative correlation sequences and applications. *J. Analyse Math.* **149** (2023) 719–761.
- [EG80] P. Erdős and R. L. Graham. *Old and New Problems and Results in Combinatorial Number Theory*. L’Enseignement Mathématique **28** (Université de Genève, Geneva, 1980).
- [F77] Harry Furstenberg. Ergodic behavior of diagonal measures and a theorem of Szemerédi on arithmetic progressions. *J. Analyse Math.* **31** (1977) 204–256.
- [G17] Ben Green. Sárközy’s theorem in function fields. arXiv:1605.07263v4 (2017) 7 pp.
- [HKM16] Marijn J. H. Heule, Oliver Kullman, and Victor W. Marek. Solving and verifying the boolean Pythagorean triples problem via cube-and-conquer. In *Theory and Applications of Satisfiability Testing – SAT 2016: 19th International Conference, Bordeaux, France, July 5–8, 2016, Proceedings.*, Lecture Notes in Computer Science **9710** (Springer International Publishing, 2016) 228–245.
- [KM78] T. Kamae and M. Mendès France. Van der Corput’s difference theorem. *Israel J. Math.* **31** (1978) 335–342.
- [K77] H. Jerome Keisler. Hyperfinite model theory. In *Logic Colloquium 76*, Studies in Logic and Foundations of Mathematics **87** (North-Holland Publishing Company, Amsterdam, 1977) 5–110.
- [LW54] Serge Lang and André Weil. Number of points of varieties in finite fields. *Amer. J. Math.* **76** (1954) 819–827.
- [L98] Paul G. Larick. *Results in Polynomial Recurrence for Actions of Fields*. Ph.D. thesis (Ohio State University, 1998).

- [LS22] Anqi Li and Lisa Sauermann. Sárközy's theorem in various finite field settings. arXiv:2212.12754 (2022) 6 pp.
- [L18] Sofia Lindqvist. Partition regularity of generalised Fermat equations. *Combinatorica* **38** (2018) 1457–1483.
- [S78] A. Sárközy. On difference sets of sequences of integers. I. *Acta Math. Acad. Sci. Hungar.* **31** (1978) 125–149.
- [S16] J. Schur. Über die kongruenz $x^m + y^m \equiv z^m \pmod{p}$. *Jahresber. Deutschen Math. Verein.* **25** (1916) 114–117.

SCHOOL OF MATHEMATICS, INSTITUTE FOR ADVANCED STUDY, PRINCETON, NJ 08540
Email address: eackelsberg@ias.edu

DEPARTMENT OF MATHEMATICS, OHIO STATE UNIVERSITY, COLUMBUS, OH 43210
Email address: vitaly@math.ohio-state.edu