

FLAT BANDS AND HIGH CHERN NUMBERS IN TWISTED MULTILAYER GRAPHENE

MENGXUAN YANG

ABSTRACT. Motivated by recent Physical Review Letters of Wang–Liu [WL22] and Ledwith–Vishwanath–Khalaf [LVK22], we study Tarnopolsky–Kruchkov–Vishwanath [TKV19] chiral model of two sheets of n -layer Bernal stacked graphene twisted by a small angle using the framework developed by Becker–Embree–Wittsten–Zworski [BEWZ22]. We show that magic angles of this model are exactly the same as magic angles of chiral twisted bilayer graphene with multiplicity. For small inter-layer tunneling potentials, we compute the band separation at Dirac points as we turning on the tunneling parameter. Flat band eigenfunctions are also constructed using a new theta function argument and this yields a complex line bundle with the Chern number $-n$.

1. INTRODUCTION

When two or more sheets of graphene are stacked on top of each other and twisted, it has been observed that at certain angles, namely *magic angles*, the zero energy band becomes flat and the system becomes superconducting.

In this paper, we consider the chiral model [SJGG12] [TKV19] of two sheets of n -layer Bernal stacked graphene twisted by a small angle [WL22]. The twisted multilayer graphene (TMG) Hamiltonian is given by

$$H_n(\alpha; \mathbf{t}) := \begin{pmatrix} 0 & D_n^*(\alpha; \mathbf{t}) \\ D_n(\alpha; \mathbf{t}) & 0 \end{pmatrix} \quad (1.1)$$

where

$$D_n(\alpha; \mathbf{t}) := \begin{pmatrix} D(\alpha) & t_1 T_+ & & & & \\ t_1 T_- & D(0) & t_2 T_+ & & & \\ & t_2 T_- & D(0) & \ddots & & \\ & & \ddots & & & \\ & & & & t_{n-1} T_+ & \\ & & & & t_{n-1} T_- & D(0) \end{pmatrix}, \quad (1.2)$$

with $\mathbf{t} = (t_1, t_2, \dots, t_{n-1})$ and

$$D(\alpha) := \begin{pmatrix} 2D_{\bar{z}} & \alpha U(z) \\ \alpha U(-z) & 2D_{\bar{z}} \end{pmatrix}, \quad T_+ = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad T_- = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}. \quad (1.3)$$

In particular, $z = x_1 + ix_2$, $D_{\bar{z}} := \frac{1}{2i}(\partial_{x_1} + i\partial_{x_2})$ and

$$U(z) = U(z, \bar{z}) := \sum_{k=0}^2 \omega^k e^{\frac{1}{2}(z\bar{\omega}^k - \bar{z}\omega^k)}, \quad \omega := e^{2\pi i/3}. \quad (1.4)$$

In equation (1.2), $t_k T_+$ (resp. $t_k T_-$) denotes the tunneling between the top (resp. the bottom) k -th layer and $(k+1)$ -th layer. Without loss of generality we can assume that all t_i 's are non-zero, since otherwise we have direct sum decomposition of $D_n(\alpha; \mathbf{t})$ into smaller matrix blocks.

In the corresponding physics model, when two honeycomb lattices are twisted against each another, a periodic honeycomb superlattice, called the moiré lattice, is formed. Bistritzer–MacDonald [BM11] predicted that the symmetries of the periodic moiré lattice in twisted bilayer graphene (TBG) lead to dramatic flattening of the band spectrum. The chiral model of TBG (1.5) was obtained by Tarnopolsky–Kruchkov–Vishwanath [TKV19] by removing certain interaction terms from the operator constructed in [BM11]; it was recently studied in greater mathematical details by the work of Becker *et al.* [BEWZ22] and Becker–Humbert–Zworski [BHZ22a] [BHZ22b]. Recently, Wang–Liu [WL22] and Ledwith–Vishwanath–Khalaf [LVK22] generalized the chiral TBG model to chiral TMG models which also have the ideal quantum geometry and support high Chern number band,

Remark 1. For $n = 1$, the operator (1.1) reduces to the chiral twisted bilayer (TBG) model

$$H_n(\alpha; \mathbf{t}) = H(\alpha) = \begin{pmatrix} 0 & D^*(\alpha) \\ D(\alpha) & 0 \end{pmatrix} \quad (1.5)$$

independent of t , see [TKV19] [BEWZ22] [BHZ22a] [BHZ22b] for more details. The dimensionless parameter α in (1.1) and (1.5) is essentially the reciprocal of the twisting angle.

Note that the potential (1.4) satisfies the following properties:

$$\begin{aligned} U(z + \mathbf{a}) &= \bar{\omega}^{a_1 + a_2} U(z), \quad \mathbf{a} = \frac{4}{3}\pi i \omega (a_1 + \omega a_2), \quad a_j \in \mathbb{Z}, \\ U(\omega z) &= \omega U(z), \quad \overline{U(\bar{z})} = U(z). \end{aligned} \quad (1.6)$$

In particular, it is periodic with respect to the lattice 3Λ with

$$\Lambda := \left\{ \frac{4}{3}\pi i \omega (a_1 + \omega a_2) : a_1, a_2 \in \mathbb{Z} \right\}. \quad (1.7)$$

The dual lattice $\frac{1}{3}\Lambda^*$ of 3Λ consists of $\mathbf{k} \in \mathbb{C}$ satisfying

$$\langle \gamma, \mathbf{k} \rangle := \frac{1}{2}(\gamma \bar{\mathbf{k}} + \bar{\gamma} \mathbf{k}) \in 2\pi\mathbb{Z}, \quad \gamma \in 3\Lambda \implies \Lambda^* = \sqrt{3}\omega (\mathbb{Z} \oplus \omega\mathbb{Z}). \quad (1.8)$$

We consider a generalized Floquet condition under the modified translation operator $\mathcal{L}_{\mathbf{a}}$, with $\mathbf{a} \in \Lambda$ (defined in equation (2.3)), i.e., we study the spectrum of $H_n(\alpha; \mathbf{t})$

satisfying the following boundary conditions:

$$H_n(\alpha; \mathbf{t})\mathbf{u} = E(\alpha, \mathbf{k})\mathbf{u}, \quad \mathcal{L}_{\mathbf{a}}\mathbf{u}(z) = e^{i\langle \mathbf{a}, \mathbf{k} \rangle} \mathbf{u}(z), \quad \mathbf{k} \in \mathbb{C}. \quad (1.9)$$

The spectrum is symmetric with respect to the origin due to chiral symmetry and we index it as

$$\begin{aligned} \{E_j(\alpha, \mathbf{k})\}_{j \in \mathbb{Z}^*}, \quad E_j(\alpha, \mathbf{k}) = -E_{-j}(\alpha, \mathbf{k}), \quad \mathbb{Z}^* := \mathbb{Z} \setminus \{0\} \\ 0 \leq E_1(\alpha, \mathbf{k}) \leq E_2(\alpha, \mathbf{k}) \leq \cdots, \quad E_1(\alpha, \mathbf{0}) = E_1(\alpha, -\mathbf{i}) = 0, \end{aligned} \quad (1.10)$$

see §3.1 for details. The points $\mathbf{0}, -\mathbf{i}$ are called the *Dirac points* and are typically denoted by K and K' in the physics literature.

For $E(\alpha, \mathbf{k})$ satisfies equation (1.9) of $H_n(\alpha; \mathbf{t})$, we define the following set of *complex numbers*:

$$\mathcal{A}_n := \{\alpha \in \mathbb{C} : \forall \mathbf{k} \in \mathbb{C}, \quad E_1(\alpha, \mathbf{k}) \equiv 0\} \quad (1.11)$$

The *magic angles* are essentially the reciprocals of α 's. And $\alpha \in \mathcal{A}$ is call *simple* if $E_2(\alpha, \mathbf{k}) > 0$ and *multiplicity* of $\alpha \in \mathcal{A}$ is defined to be the number of j 's ($j > 0$) such that $E_j(\alpha, \mathbf{k}) \equiv 0$.

Our first theorem states that the magic angles of the TMG Hamiltonian $H_n(\alpha; \mathbf{t})$ coincides with magic angles of the TBG Hamiltonian $H(\alpha)$.

Theorem 1. *The set \mathcal{A}_n is independent of $n \in \mathbb{N}_+$. More precisely, the multiplicity of each $\alpha \in \mathcal{A}_n$ is independent of n .*

Therefore we can define $\mathcal{A} := \mathcal{A}_n$. This result is essentially contained in the paper [WL22]. A spectral theory characterization of the set \mathcal{A}_1 (for TBG) was given in [BEWZ22] using a Birman–Schwinger operator whose spectrum is given by $1/\mathcal{A}_1$.

Our second result studies the role of the tunneling parameter \mathbf{t} for $\alpha \in \mathcal{A}$. When $\mathbf{t} = \mathbf{0}$, the operator $D_n(\alpha; \mathbf{t})$ can be decomposed into direct sums of one operator $D(\alpha)$ and $(n - 1)$ operators $D(0)$, where the latter give rise to Dirac cones independent of $\alpha \in \mathbb{C}$. The Dirac cones are connected to the band $E_1(\alpha, \mathbf{k})$ for any $\alpha \in \mathbb{C}$. In particular, for $\alpha \in \mathcal{A}$, turning on the tunneling parameter \mathbf{t} such that $t_i > 0$ for $1 \leq i \leq n - 1$ results in these Dirac cones separate from the flat band and a band gap is formed. The band separation is studied for $H_2(\alpha; t)$:

Theorem 2. *Assume $\alpha \in \mathcal{A}$ is simple. For $t \ll 1$ and at $\mathbf{k} = K'$, the first two eigenvalues of $H_2(\alpha; t)$ are given by $E_1(\alpha, \mathbf{k}, \mathbf{t}) = 0$ and $E_2(\alpha, \mathbf{k}, \mathbf{t}) = Ct + \mathcal{O}(t^{3/2})$.*

Remark 2. *The constant C is computed in Proposition 5.3. The same result is expect to be true for $H_n(\alpha; \mathbf{t})$. We only present the case $n = 2$ here for notational simplicity purposes. This also gives an explicit result for the band separation mechanism discussed in [WL22].*

Using a new theta function argument and the resolvent formula for $(2D_{\bar{z}} - \mathbf{k})^{-1}$, for $\alpha \in \mathcal{A}$ simple, we construct flat band eigenfunctions of $H_n(\alpha; \mathbf{t})$. This gives rise to a holomorphic line bundle L (see discussions in §6.3 for more details).

Theorem 3. *The Chern number of the line bundle L is given by $c_1(L) = -n$.*

This justifies the high Chern number observation made in [WL22] and it relies on a numerical calculation of the integration of the flat band eigenfunctions (see Table 1). In [WKL22], the flat band eigenfunctions are considered as a section of rank n holomorphic vector bundle. Our new theta function argument gives an explicit analytic construction and answers a question proposed in [LB22].

We conclude our introduction by discussing its relation to physics:

Integer quantum Hall effect. For two dimensional electron gas in cold temperature, Hall conductance exhibits some plateaus at integer multiples of $e^2/2\pi\hbar$ as the magnetic field varies. It turns out that the Hall conductance at the plateaus is related (via the Berry curvature) to Chern number c_1 of the underlying Hermitian line bundles arising from ground state eigenfunctions. By Kubo's formula, the Hall conductance is given by

$$\sigma = -\frac{e^2}{2\pi\hbar}c_1$$

The higher Chern number in the TMG model can yield higher Hall conductance in this case.

Ideal Quantum Geometry and Fractional Chern insulators. Another observation (cf. [WL22]) is that flat band eigenfunctions constructed in (6.9) are holomorphic in $\mathbf{k} \in \mathbb{C}$. Such flat band is said to satisfy the *ideal quantum geometry* in physics literature (cf. [CLT⁺15][OM21][MO21] [WCM⁺21]), in the sense that the Berry curvature $\Omega(\mathbf{k})$ on the fundamental domain \mathbb{C}/Λ^* is non-vanishing and proportional to a Fubini–Study metric $g_{\mathbf{k}}$ on \mathbb{C}/Λ^* induced by the flat band eigenfunctions.

Such flat band was predicted and verified to have Fractional Chern Insulators (FCI) [RB11] [BL13], which are lattice generalizations of the conventional fractional quantum Hall effect in two-dimensional electron gases. For example, FCI can appear at band fillings given by

$$\nu = \frac{k}{(m-1)|c_1| + 1}$$

with $m, k \in \mathbb{N}$ such that $k \geq 1, m \geq 2$. Here $k = 1$ (resp. $k > 1$) corresponds to Abelian states (resp. non-Abelian states) and m is even (resp. odd) for bosons (resp. fermions). The stability of FCI are conjectured due to the ideal quantum geometry (cf. [TKV19] [WKL22] [LB22]).

Structure of the paper. In Section 2, we study symmetry groups and irreducible representations. This yields the so-called protected zero eigenstates for all $\alpha \in \mathbb{C}$. In Section 3, we recall a generalized Bloch–Floquet theory introduced in [BHZ22a] and give spectral characterizations of magic angles. In Section 4, we prove Theorem 1. Then in Section 5, we prove Theorem 2. In Section 6, we construct the flat band eigenfunctions assuming the simplicity of magic angles and prove Theorem 3.

Acknowledgement. The author is very grateful to Maciej Zworski for introducing him to the field of magic angles as well as for having many helpful discussions and comments on the early manuscript. The author would also like to thank Simon Becker for helpful discussions and providing various numerical supports, and Zhongkai Tao, Jie Wang and Jared Wunsch for helpful discussions related to this project and comments on the manuscript.

2. HAMILTONIANS OF TWISTED MULTILAYER GRAPHENE

In this section, we study symmetries of the Hamiltonian $H_n(\alpha; \mathbf{t})$. The symmetry group commutes with the Hamiltonian and split the functional space into irreducible representations. This gives rise to the so-called protected states. We also consider eigenstates with zero eigenvalue for $\mathbf{t} = 0$, for which some of them are unprotected in the sense that the corresponding eigenbranches separate from the flat band as we turning on the inter-layer parameter \mathbf{t} when $\alpha \in \mathcal{A}$.

2.1. Symmetries of the Hamiltonian.

2.1.1. *Translational symmetry.* The first identity in (1.6) shows that for $L_{\mathbf{a}}\mathbf{v}(z) := \mathbf{v}(z + \mathbf{a})$,

$$D_n(\alpha; \mathbf{t})L_{\mathbf{a}} = \text{diag}_n \left\{ \begin{pmatrix} \omega & 0 \\ 0 & 1 \end{pmatrix} \right\} L_{\mathbf{a}} D_n(\alpha; t) \text{diag}_n \left\{ \begin{pmatrix} \bar{\omega} & 0 \\ 0 & 1 \end{pmatrix} \right\}, \quad \mathbf{a} = \frac{4}{3}\pi i \omega^\ell, \quad \ell = 1, 2$$

where $\text{diag}_n\{(*)\}$ denotes block diagonal matrices with n blocks. Therefore,

$$\mathcal{L}_{\mathbf{a}} D(\alpha) = D(\alpha) \mathcal{L}_{\mathbf{a}}, \quad \mathcal{L}_{\mathbf{a}} := \text{diag}_n \left\{ \begin{pmatrix} \omega & 0 \\ 0 & 1 \end{pmatrix} \right\} L_{\mathbf{a}}, \quad \mathbf{a} = \frac{4}{3}\pi i \omega^\ell, \quad \ell = 1, 2. \quad (2.1)$$

Therefore for the lattice

$$\Lambda = \frac{4}{3}\pi i \omega (\mathbb{Z} \oplus \omega \mathbb{Z}) \quad (2.2)$$

and

$$\mathcal{L}_{\mathbf{a}} := \text{diag}_n \left\{ \begin{pmatrix} \omega^{a_1+a_2} & 0 \\ 0 & 1 \end{pmatrix} \right\} L_{\mathbf{a}}, \quad \mathbf{a} = \frac{4}{3}\pi i \omega (a_1 + \omega a_2) \in \Lambda, \quad (2.3)$$

we obtain a unitary action of Λ on $L^2(\mathbb{C}; \mathbb{C}^{2n})$, or more precisely on $L^2(\mathbb{C}/3\Lambda; \mathbb{C}^{2n})$, with $\mathbf{a} \mapsto \mathcal{L}_{\mathbf{a}}$. We can further extend the action of $\mathcal{L}_{\mathbf{a}}$ to $L^2(\mathbb{C}; \mathbb{C}^{4n})$ or $L^2(\mathbb{C}/3\Lambda; \mathbb{C}^{4n})$ block-diagonally and it yields

$$\mathcal{L}_{\mathbf{a}} H_n(\alpha) = H_n(\alpha) \mathcal{L}_{\mathbf{a}}, \quad \mathbf{a} \in \Lambda. \quad (2.4)$$

2.1.2. *Rotational symmetry.* The second identity in (1.6) shows that

$$[D(\alpha) \mathbf{u}(\omega \bullet)](z) = \bar{\omega} [D(\alpha) \mathbf{u}](\omega z).$$

Therefore, applying this to $D_n(\alpha, \mathbf{t})$ in equation (1.2), we have

$$[D_n(\alpha; \mathbf{t}) \mathbf{u}(\omega \bullet)](z) = \bar{\omega} [J D_n(\alpha; \mathbf{t}) J^* \mathbf{u}](\omega z), \quad (2.5)$$

$$J := \text{diag}(1, 1, \bar{\omega}, \omega, \bar{\omega}^2, \omega^2, \dots, \bar{\omega}^{n-1}, \omega^{n-1}). \quad (2.6)$$

Similarly for $D_n^*(\alpha; \mathbf{t})$, we have

$$[D_n^*(\alpha; \mathbf{t}) \mathbf{u}(\omega \bullet)](z) = \omega [J D_n^*(\alpha; \mathbf{t}) J^* \mathbf{u}](\omega z).$$

Hence, for $\mathbf{u} \in L^2(\mathbb{C}; \mathbb{C}^{4n})$,

$$\mathcal{C}' \begin{pmatrix} J & \\ & J \end{pmatrix} H_n(\alpha; \mathbf{t}) \begin{pmatrix} J^* & \\ & J^* \end{pmatrix} = H_n(\alpha; \mathbf{t}) \mathcal{C}', \quad (2.7)$$

$$\mathcal{C}' \mathbf{u}(z) := \begin{pmatrix} \text{Id}_{2n} & \\ 0 & \bar{\omega} \text{Id}_{2n} \end{pmatrix} \mathbf{u}(\omega z). \quad (2.8)$$

Therefore, we obtain

$$\mathcal{C} H_n(\alpha; \mathbf{t}) = H_n(\alpha; \mathbf{t}) \mathcal{C}, \quad \mathcal{C} := \mathcal{C}' \begin{pmatrix} J & \\ & J \end{pmatrix} \quad (2.9)$$

2.1.3. *Additional symmetries.* We record some additional actions and symmetries involving $H_n(\alpha; \mathbf{t})$.

$$H_n(\alpha; \mathbf{t}) = -\mathcal{W} H_n(\alpha; \mathbf{t}) \mathcal{W}^*, \quad \mathcal{W} := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \mathcal{W} \mathcal{C} = \mathcal{C} \mathcal{W}, \quad \mathcal{L}_{\mathbf{a}} \mathcal{W} = \mathcal{W} \mathcal{L}_{\mathbf{a}}, \quad (2.10)$$

This implies that the spectrum of $H_n(\alpha; \mathbf{t})$ is even.

Remark 3. *With the notation introduced in (3.6) we also have*

$$H_{n, \mathbf{k}}(\alpha; \mathbf{t}) = -\mathcal{W} H_{n, \mathbf{k}}(\alpha; \mathbf{t}) \mathcal{W}^*.$$

This implies that for each eigenvalue $E_j(\mathbf{k}) \geq 0$ of $H_{n, \mathbf{k}}(\alpha; \mathbf{t})$, there exists another eigenvalue $E_{-j}(\mathbf{k}) \leq 0$ such that $E_{-j}(\mathbf{k}) = -E_j(\mathbf{k})$.

2.1.4. *Group actions on functional spaces.* Following [BEWZ22], since $\mathcal{C}\mathcal{L}_{\mathbf{a}} = \mathcal{L}_{\bar{\omega}\mathbf{a}}\mathcal{C}$, we can combine the two actions into a unitary group action that commutes with $H_n(\alpha; \mathbf{t})$:

$$\begin{aligned} G := \Lambda \times \mathbb{Z}_3, \quad \mathbb{Z}_3 \ni k : \mathbf{a} \rightarrow \bar{\omega}^k \mathbf{a}, \quad (\mathbf{a}, k) \cdot (\mathbf{a}', \ell) &= (\mathbf{a} + \bar{\omega}^k \mathbf{a}', k + \ell), \\ (\mathbf{a}, \ell) \cdot \mathbf{u} &= \mathcal{L}_{\mathbf{a}} \mathcal{C}^\ell \mathbf{u}. \end{aligned} \quad (2.11)$$

Taking a quotient by 3Λ , we obtain a finite group acting unitarily on $L^2(\mathbb{C}/3\Lambda; \mathbb{C}^{4n})$ and commuting with $H_n(\alpha; \mathbf{t})$:

$$G_3 := G/3\Lambda = \Lambda/3\Lambda \times \mathbb{Z}_3 \simeq \mathbb{Z}_3^2 \times \mathbb{Z}_3. \quad (2.12)$$

Restricting to the components, G (resp. G_3) acts on $L^2(\mathbb{C}; \mathbb{C}^{2n})$ (resp. $L^2(\mathbb{C}/3\Lambda; \mathbb{C}^{2n})$) as well and we use the same notation for those actions. With the previous discussions, we have the following

Proposition 2.1. *The operator $H_n(\alpha; \mathbf{t}) : L^2(\mathbb{C}; \mathbb{C}^{4n}) \rightarrow L^2(\mathbb{C}; \mathbb{C}^{4n})$ is an unbounded self-adjoint operator with the domain given by $H^1(\mathbb{C}; \mathbb{C}^{4n})$. The operator $H_n(\alpha; \mathbf{t})$ commutes with the unitary action of the group G given by (2.11) and*

$$\text{Spec}_{L^2(\mathbb{C})} H_n(\alpha; \mathbf{t}) = -\text{Spec}_{L^2(\mathbb{C})} H_n(\alpha; \mathbf{t}).$$

The same conclusions are valid when $L^2(\mathbb{C})$ is replaced by $L^2(\mathbb{C}/3\Lambda)$ and G by G_3 given by (2.12). In that case, the spectrum is discrete.

2.2. **Protected states at zero.** To show the existence of protected zero eigenstates of the operator $H_n(\alpha; \mathbf{t})$, we start with considering all irreducible representations of G (resp. G_3) on $L^2(\mathbb{C})$ (resp. $L^2(\mathbb{C}/3\Lambda)$).

2.2.1. *Irreducible representations.* Recall as in [BEWZ22, Section 2.2], irreducible unitary representations of \mathbb{Z}_3^2 are one dimensional and are given by

$$\begin{aligned} \pi_{\mathbf{k}} : \mathbb{Z}_3^2 &\rightarrow \mathbf{U}(1), \quad \pi_{\mathbf{k}}(\mathbf{a}) = e^{i\langle \mathbf{a}, \mathbf{k} \rangle}, \\ \mathbf{a} &= \frac{4}{3}\pi i \omega(a_1 + a_2 \omega), \quad a_j \in \mathbb{Z}_3, \quad \mathbf{k} = \frac{1}{\sqrt{3}}\omega(\omega k_1 - k_2), \quad k_j \in \mathbb{Z}_3, \\ \langle \mathbf{a}, \mathbf{k} \rangle &= \frac{1}{2}(\mathbf{a}\bar{\mathbf{k}} + \bar{\mathbf{a}}\mathbf{k}) = \frac{2\pi}{3}(k_1 a_1 + k_2 a_2). \end{aligned} \quad (2.13)$$

Irreducible representations of G_3 are one dimensional for $\mathbf{k} \in \Delta$ (given by $\Delta(\mathbb{Z}_3) := \{(k, k), k \in \mathbb{Z}_3\}$ – note that $\langle \mathbf{k}, \omega \mathbf{a} \rangle = \langle \mathbf{k}, \mathbf{a} \rangle$, $\mathbf{a} \in \Lambda/3\Lambda$, if and only if $\mathbf{k} \in \Delta$),

$$\rho_{k,p}((\mathbf{a}, \ell)) = \bar{\omega}^{\ell p} \pi_{(k,k)}(\mathbf{a}),$$

or three dimensional, for $\mathbf{k} \notin \Delta$:

$$\rho_{\mathbf{k}}((\mathbf{a}, \ell)) = \begin{pmatrix} \omega^{\langle \mathbf{k}, \mathbf{a} \rangle} & 0 & 0 \\ 0 & \omega^{\langle \mathbf{k}, \omega \mathbf{a} \rangle} & 0 \\ 0 & 0 & \omega^{\langle \mathbf{k}, \omega^2 \mathbf{a} \rangle} \end{pmatrix} \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}^\ell \in \mathbf{U}(3).$$

The representations are equivalent for \mathbf{k} in the same orbit of the transpose of $\mathbf{a} \mapsto \omega\mathbf{a}$, and hence there are only two.

Defining the following functional space:

$$L_{k,p}^2 := \{\mathbf{u} \in L^2(\mathbb{C}/3\Lambda; \mathbb{C}^{4n}) : \mathcal{L}_{\mathbf{a}} \mathcal{C}^{\ell} \mathbf{u} = e^{i\langle \mathbf{a}, \mathbf{k} \rangle} \bar{\omega}^{\ell p} u\} \quad (2.14)$$

There are then 11 irreducible representations: 9 one dimensional and 2 three dimensional. We can decompose $L^2(\mathbb{C}/3\Lambda; \mathbb{C}^{4n})$ into 11 orthogonal subspaces:

$$L^2(\mathbb{C}/3\Lambda; \mathbb{C}^{4n}) = \left(\bigoplus_{k,p \in \mathbb{Z}_3} L_{k,p}^2 \right) \oplus L_{(1,0)}^2 \oplus L_{(2,0)}^2. \quad (2.15)$$

In view of Proposition 2.1 we define

$$H_{k,p}(\alpha; \mathbf{t}) := H_n(\alpha; \mathbf{t}) : L_{k,p}^2 \cap H^1(\mathbb{C}/3\Lambda; \mathbb{C}^{4n}) \longrightarrow L_{k,p}^2, \quad (2.16)$$

with $H_{(1,0)}$ and $H_{(0,1)}$ similarly defined. Here we omit the dimensional constant n when there is no ambiguity.

2.2.2. Protected zero eigenstates. We start with the case $\alpha = 0, \mathbf{t} \in (\mathbb{R} \setminus \{0\})^{n-1}$. The kernel of $H_n(\alpha; \mathbf{t})$ at $\alpha = 0$ is given by

$$\ker_{L^2(\mathbb{C}/3\Lambda; \mathbb{C}^{4n})} H_n(0; \mathbf{t}) = \{\mathbf{e}_1, \mathbf{e}_{2n}, \mathbf{e}_{2n+2}, \mathbf{e}_{4n-1}\},$$

where the set $\{\mathbf{e}_j\}_{j=1}^{4n}$ forms the standard basis of \mathbb{C}^{4n} . Now we consider the action of $G_3 = \mathbb{Z}_3^2 \rtimes \mathbb{Z}_3$ on $\ker_{L^2(\mathbb{C}/3\Lambda; \mathbb{C}^{4n})} H_n(0; \mathbf{t})$ with $\mathbf{a} = \frac{4}{3}\pi i \omega(a_1 + a_2 \omega)$, $(a_1, a_2) \in \mathbb{Z}_3^2$:

$$\begin{aligned} \mathcal{L}_{\mathbf{a}} \mathbf{e}_1 &= \omega^{a_1+a_2} \mathbf{e}_1, & \mathcal{L}_{\mathbf{a}} \mathbf{e}_{2n} &= \mathbf{e}_{2n}, & \mathcal{L}_{\mathbf{a}} \mathbf{e}_{2n+2} &= \mathbf{e}_{2n+2}, & \mathcal{L}_{\mathbf{a}} \mathbf{e}_{4n-1} &= \omega^{a_1+a_2} \mathbf{e}_{4n-1}, \\ \mathcal{C} \mathbf{e}_1 &= \mathbf{e}_1, & \mathcal{C} \mathbf{e}_{2n} &= \bar{\omega}^{1-n} \mathbf{e}_{2n}, & \mathcal{C} \mathbf{e}_{2n+2} &= \bar{\omega} \mathbf{e}_{2n+2}, & \mathcal{C} \mathbf{e}_{4n-1} &= \bar{\omega}^n \mathbf{e}_{4n-1}. \end{aligned}$$

If $n \neq 3k$ for $k \in \mathbb{N}$, then $\mathbf{e}_1, \mathbf{e}_{2n}, \mathbf{e}_{2n+2}, \mathbf{e}_{4n-1}$ are in different irreducible representations of $L^2(\mathbb{C}/3\Lambda; \mathbb{C}^{4n})$ under group action G_3 with

$$\mathbf{e}_1 \in L_{1,0}^2, \quad \mathbf{e}_{2n} \in L_{0,[1-n]}^2, \quad \mathbf{e}_{2n+2} \in L_{0,1}^2, \quad \mathbf{e}_{4n-1} \in L_{1,[n]}^2,$$

where $[k] \in \{0, 1, 2\}$ with $[k] \equiv k \pmod{3}$. Therefore if $n \neq 3k$ and $\alpha = 0$, each of these irreducible representation $H_{k,p}(0; \mathbf{t})$ has a single zero eigenvalue. Since \mathcal{W} (see (2.10)) commutes with the action of G_3 , the spectra of $H_{k,p}(\alpha; \mathbf{t})$ are symmetric with respect to 0 (see Proposition 2.1), it follows that for each k, p as above, $H_{k,p}(\alpha; \mathbf{t})$ has an eigenvalue at 0.

When $n = 3k$ for some $k \in \mathbb{N}$, we have

$$\mathbf{e}_1, \mathbf{e}_{4n-1} \in L_{1,0}^2, \quad \mathbf{e}_{2n}, \mathbf{e}_{2n+2} \in L_{0,1}^2.$$

since \mathbf{e}_{4n-1} and \mathbf{e}_{2n} are always eigenfunctions of $H_{1,0}(\alpha; \mathbf{t})$ and $H_{0,1}(\alpha; \mathbf{t})$ correspondingly, they are protected zero eigenstates naturally. The symmetry of the spectra of

$H_{1,0}$ (resp. $H_{0,1}$) implies that $H_{1,0}(\alpha; \mathbf{t})$ and $H_{0,1}(\alpha; \mathbf{t})$ both have two protected eigenvalues at 0. Since

$$\begin{aligned} \ker_{L^2(\mathbb{C}/3\Lambda; \mathbb{C}^{4n})} H_n(\alpha; \mathbf{t}) &= \ker_{L^2(\mathbb{C}/3\Lambda; \mathbb{C}^{2n})} D_n(\alpha; \mathbf{t}) \oplus \{0_{\mathbb{C}^{2n}}\} \\ &\quad + \{0_{\mathbb{C}^{2n}}\} \oplus \ker_{L^2(\mathbb{C}/3\Lambda; \mathbb{C}^{2n})} D_n^*(\alpha, \mathbf{t}), \end{aligned} \quad (2.17)$$

we obtained the following result about symmetric protected eigenstates at 0:

Proposition 2.2. *For all $\alpha \in \mathbb{C}$ and any $\mathbf{t} \in (\mathbb{R} \setminus \{0\})^{n-1}$*

$$\ker_{L_{1,0}^2} D_n(\alpha; \mathbf{t}) \neq \emptyset, \quad \ker_{L_{0,[1-n]}^2} D_n(\alpha; \mathbf{t}) = \{\mathbf{e}_{2n}\}.$$

Remark 4. *The protected states are also discussed in [WL22] using perturbation theory for \mathbf{t} infinitesimal. Here we provide a different approach to show the zero eigenstates are protected for all $\mathbf{t} \in \mathbb{R}^{n-1}$.*

2.2.3. Unprotected zero eigenstates and band separation. Now we consider the case $\mathbf{t} = 0$, i.e., no tunneling in the upper and lower n -layers. We show that in this case, there are additional $4n - 4$ zero eigenstates of $H_n(\alpha; 0)$ that is unprotected, in the sense that they result in band separations as we turning on the tunneling parameter \mathbf{t} (cf. Section 5.).

The operator $D_n(\alpha; 0)$ is decomposed into direct sums of one twisted bilayer operator $D(\alpha)$ (see equation (1.3)) and $D(0) = \text{diag}(2D_{\bar{z}}, 2D_{\bar{z}})$. The kernel of $H_n(\alpha; 0)$ is $4n$ -dimensional and given by direct sums of $\ker_{L^2(\mathbb{C}/3\Lambda; \mathbb{C}^2)} D(\alpha)$ and $\ker_{L^2(\mathbb{C}/3\Lambda; \mathbb{C}^2)} D^*(\alpha)$ and $(2n - 2)$ copies of $\ker_{L^2(\mathbb{C}/3\Lambda; \mathbb{C}^2)} D(0)$:

$$\ker_{L^2(\mathbb{C}/3\Lambda; \mathbb{C}^{4n})} H_n(\alpha; 0) = \{\Phi_i(\alpha)\}_{i=1,2,2n+1,2n+2} \cup \{\mathbf{e}_j\}_{j \neq 1,2,2n+1,2n+2}$$

where

$$\begin{aligned} \Phi_i(\alpha) &= (\varphi_i(\alpha), 0_{\mathbb{C}^{4n-2}}), \quad \Phi_{2n+i}(\alpha) = (0_{\mathbb{C}^{2n}}, \psi_i(\alpha), 0_{\mathbb{C}^{2n-2}}), \quad i = 1, 2; \\ D(\alpha)\varphi_i &= 0, \quad D^*(\alpha)\psi_i = 0, \quad i = 1, 2. \end{aligned}$$

We can again consider the action of $G_3 = \mathbb{Z}_3^2 \rtimes \mathbb{Z}_3$ on $\ker_{L^2(\mathbb{C}/3\Lambda; \mathbb{C}^{4n})} H_n(\alpha; 0)$ with $\mathbf{a} = \frac{4}{3}\pi i \omega(a_1 + a_2 \omega)$, $(a_1, a_2) \in \mathbb{Z}_3^2$.

For simplicity of presentation, we only discuss the case $n = 2$. This yields

$$\Phi_1 \in L_{1,0}^2, \quad \mathbf{e}_4 \in L_{0,2}^2, \quad \Phi_6 \in L_{0,1}^2, \quad \mathbf{e}_7 \in L_{1,2}^2, \quad (2.18)$$

$$\{\Phi_2, \mathbf{e}_8\} \subset L_{0,0}^2, \quad \{\mathbf{e}_3, \Phi_5\} \subset L_{1,1}^2. \quad (2.19)$$

Similar to previous discussion, for $t = 0$, each of these irreducible representation $H_{k,\ell}(\alpha; 0)$ ($k = 0, 1$; $\ell = 0, 2$) has a single zero eigenvalue, whereas $H_{\ell,\ell}(\alpha; 0)$ ($\ell = 0, 1$) has a double zero eigenvalue. The commutativity of \mathscr{W} with G_3 yields that $H_{k,\ell}(\alpha)$ each have an eigenvalue at 0. Whereas $\ker H_{\ell,\ell}(\alpha; t) = \emptyset$ for $\ell = 0, 1$ from the discussion in the previous subsection shows that the pairs of zero eigenfunctions $\{\Phi_2, \mathbf{e}_8\}$ and $\{\mathbf{e}_3, \Phi_5\}$ of $H_{\ell,\ell}(\alpha; 0)$ disappears as we turning on the tunneling parameter t . This, in

fact, is the origin of the Dirac band separation from the flatband, which is discussed in Section 5.

Same argument applies to any $n \in \mathbb{N}$, where all the irreducible representations $H_{k,\ell}(\alpha; 0)$ may have more than one zero eigenfunctions. The only protected zero eigenstates are eigenbranches corresponding to eigenfunctions $\{\Phi_1, \mathbf{e}_{2n}, \Phi_{2n+2}, \mathbf{e}_{4n-1}\}$.

3. BLOCH–FLOQUET THEORY

In this section we introduce a Bloch–Floquet theory of the Hamiltonian $H_n(\alpha; \mathbf{t})$ with respect to the translation operator $\mathcal{L}_{\mathbf{a}}$, $\mathbf{a} \in \Lambda$, which gives the band structure of the operator $H_{n,\mathbf{k}}(\alpha; \mathbf{t})$ (cf. equation (3.6)) for $\mathbf{k} \in \mathbb{C}/\Lambda^*$.

3.1. A generalized Floquet theory approach. We follow [BHZ22a] for the discussion of Floquet theory with respect to the operator $\mathcal{L}_{\mathbf{a}}$ with $\mathbf{a} \in \Lambda$, which has already implicitly appeared in the physics literature (cf. [TKV19]). One of the advantages of this type of Floquet theory is that the set $\ker_{L^2(\mathbb{C}/3\Lambda; \mathbb{C}^2)} D(\alpha)$ at the flat band, which is nine dimensional (cf. [BHZ22a, Section 3.3]), splits to elements of $\ker_{L^2_0(\mathbb{C}/\Lambda; \mathbb{C}^2)}(D(\alpha) + \mathbf{k})$ (see (3.8) below for the definition of L^2_0) for nine different $\mathbf{k} \in \mathbb{C}/\Lambda^*$. This generalized Floquet theory also simplifies the presentation of band separation mechanism (cf. Section 5).

We first define the inner product on \mathbb{C} by

$$\langle z, w \rangle := \frac{1}{2}(z\bar{w} + \bar{z}w), \quad z, w \in \mathbb{C}.$$

Note that if $z \in \Lambda$ and $w \in \Lambda^*$, then we have $\langle z, w \rangle = 2\pi\mathbb{Z}$.

The generalized Floquet condition is formulated for $\mathbf{w} \in H^1_{\text{loc}}(\mathbb{C}; \mathbb{C}^{4n})$ as follows:

$$\mathcal{L}_{\mathbf{a}}\mathbf{w}(z) := e^{i\langle \mathbf{a}, \mathbf{k} \rangle} \mathbf{w}(z), \quad \mathbf{a} \in \Lambda, \quad \mathbf{k} \in \mathbb{C}/\Lambda^*, \quad (3.1)$$

where

$$\begin{aligned} \langle \mathbf{a}, \mathbf{k} \rangle &= 2\pi(a_1k_1 + a_2k_2)/3, \quad \mathbf{a} = \frac{4}{3}\pi i\omega(a_1 + \omega a_2) \in \Lambda, \quad a_j \in \mathbb{Z}, \\ \mathbf{k} &= \frac{1}{\sqrt{3}}\omega(\omega k_1 - k_2) \in \mathbb{C}/\Lambda^*, \quad 0 \leq k_j < 3. \end{aligned} \quad (3.2)$$

We consider the Floquet spectrum (cf. Proposition 3.1 for the discreteness of the spectrum) with Floquet condition (3.1) under the twisted translation operator $\mathcal{L}_{\mathbf{a}}$:

$$H_n(\alpha; \mathbf{t})\mathbf{w}_j(\alpha, \mathbf{k}) = E_j(\alpha, \mathbf{k}; \mathbf{t})\mathbf{w}_j(\alpha, \mathbf{k}), \quad (3.3)$$

$$\mathcal{L}_{\mathbf{a}}\mathbf{w}_j(\alpha, \mathbf{k}) = e^{i\langle \mathbf{a}, \mathbf{k} \rangle} \mathbf{w}_j(\alpha, \mathbf{k}), \quad \mathbf{a} \in \Lambda, \quad \mathbf{k} \in \mathbb{C}/\Lambda^*. \quad (3.4)$$

We define

$$\mathbf{v}_j(\alpha, \mathbf{k}) := e^{-i\langle z, \mathbf{k} \rangle} \mathbf{w}_j(\alpha, \mathbf{k})(z), \quad \mathbf{a} \in \Lambda, \quad \mathbf{k} \in \mathbb{C}/\Lambda^*, \quad (3.5)$$

which yields $\mathcal{L}_{\mathbf{a}}\mathbf{v}_j(\alpha, \mathbf{k}) = \mathbf{v}_j(\alpha, \mathbf{k})$. We then consider the operator

$$H_{n,\mathbf{k}}(\alpha; \mathbf{t}) := e^{-i\langle z, \mathbf{k} \rangle} H_n(\alpha; \mathbf{t}) e^{i\langle z, \mathbf{k} \rangle} = \begin{pmatrix} 0 & D_n(\alpha; \mathbf{t})^* + \bar{\mathbf{k}} \\ D_n(\alpha; \mathbf{t}) + \mathbf{k} & 0 \end{pmatrix}. \quad (3.6)$$

with boundary condition $\mathcal{L}_{\mathbf{a}}\mathbf{v} = \mathbf{v}$ and that gives us *equivalent* Floquet spectrum:

$$H_{\mathbf{k},n}(\alpha; \mathbf{t})\mathbf{v}_j(\alpha, \mathbf{k}) = E_j(\alpha, \mathbf{k}, \mathbf{t})\mathbf{v}_j(\alpha, \mathbf{k}), \quad \mathcal{L}_{\mathbf{a}}\mathbf{v}_j(\alpha, \mathbf{k}) = \mathbf{v}_j(\alpha, \mathbf{k}), \quad \mathbf{a} \in \Lambda. \quad (3.7)$$

Note that $\mathbf{v}_j(\alpha, \mathbf{k})$ also depends on \mathbf{t} but we omit it here for simplicity. This suggests that we consider $H_{n,\mathbf{k}}(\alpha; \mathbf{t})$ as a self-adjoint operator on

$$L_0^2 = L_0^2(\mathbb{C}/3\Lambda; \mathbb{C}^{4n}) := \{\mathbf{v} \in L_{\text{loc}}^2(\mathbb{C}/3\Lambda, \mathbb{C}^{4n}) : \mathcal{L}_{\mathbf{a}}\mathbf{v} = \mathbf{v}, \mathbf{a} \in \Lambda\} \quad (3.8)$$

with the domain given by $H_0^1 := L_0^2 \cap H^1(\mathbb{C}/3\Lambda; \mathbb{C}^{4n})$. Note that we sometimes slightly abuse the notation of L_0^2 by restricting to the subspace \mathbb{C}^2 or \mathbb{C}^{2n} of \mathbb{C}^{4n} the operator $\mathcal{L}_{\mathbf{a}}$ is diagonal.

In other words, we can define a generalized Bloch transform (cf. [TZ23, Chapter 5]):

$$\mathcal{B}u(z, \mathbf{k}) := \sum_{\mathbf{a} \in \Lambda} e^{-i\langle z + \mathbf{a}, \mathbf{k} \rangle} \mathcal{L}_{\mathbf{a}}u(z) \quad (3.9)$$

which can be checked easily satisfies

$$\begin{aligned} \mathcal{B}u(z, \mathbf{k} + \mathbf{p}) &= e^{-i\langle z, \mathbf{p} \rangle} \mathcal{B}u(z, \mathbf{k}), \quad \mathbf{p} \in \Lambda^*, \\ \mathcal{L}_{\mathbf{a}'}\mathcal{B}u(\bullet, \mathbf{k}) &= \sum_{\mathbf{a} \in \Lambda} e^{-i\langle z + \mathbf{a} + \mathbf{a}', \mathbf{k} \rangle} \mathcal{L}_{\mathbf{a} + \mathbf{a}'}u(z) = \mathcal{B}u(\bullet, \mathbf{k}), \quad \mathbf{a}' \in \Lambda, \end{aligned}$$

so that $\mathcal{B}u(\bullet, \mathbf{k}) \in L_0^2$. By the commutativity of $H_n(\alpha; \mathbf{t})$ and $\mathcal{L}_{\mathbf{a}}$ we have

$$\mathcal{B}H_n(\alpha; \mathbf{t}) = H_{n,\mathbf{k}}(\alpha; \mathbf{t})\mathcal{B}.$$

For a fixed $\mathbf{k} \in \mathbb{C}/\Lambda^*$, $H_{n,\mathbf{k}} = \mathcal{B}H_n\mathcal{B}^*$ acts on L_0^2 as the operator in (3.6). The modified Bloch transform thus gives rise to bands of eigenfunctions $E_j(\alpha, \mathbf{k}, \mathbf{t})$, $j \in \mathbb{N}$ of $H_{n,\mathbf{k}}(\alpha; \mathbf{t})$ for $\mathbf{k} \in \Lambda^*$.

Remark 5. In \mathbb{C}/Λ^* , we denote two special points $\mathbf{k} = (0, 0) = \mathbf{0}$ by K and $\mathbf{k} = (1, 1) = \frac{1}{\sqrt{3}}\omega(\omega - 1) = -\mathbf{i}$ by K' as in physics literature.

3.2. Floquet theory and representations. We further define the spaces (cf. the decomposition (2.15)):

$$\begin{aligned} L_{\mathbf{p}}^2 &:= \{\mathbf{u} \in L^2(\mathbb{C}/3\Lambda; \mathbb{C}^{4n}) : \mathcal{L}_{\mathbf{a}}\mathbf{u} = e^{i\langle \mathbf{a}, \mathbf{p} \rangle} \mathbf{u}, \quad \forall \mathbf{a} \in \Lambda\}, \\ L^2(\mathbb{C}/3\Lambda; \mathbb{C}^{4n}) &= \bigoplus_{\mathbf{p} \in 3\Lambda^*/\Lambda^*} L_{\mathbf{p}}^2. \end{aligned} \quad (3.10)$$

Note that for

$$\mathbf{p} \in \Delta = \left\{ \frac{1}{\sqrt{3}}\omega(\omega p - p) : p \in \mathbb{Z}_3 \right\},$$

we have orthogonal decomposition

$$L_{\mathbf{p}}^2 = \bigoplus_{\ell \in \mathbb{Z}_3} L_{p,\ell}^2, \quad \mathbf{p} = \frac{1}{\sqrt{3}}\omega(\omega p - p).$$

Remark 6. Note that $u \in \ker_{L_0^2}(D_n(\alpha; \mathbf{t}) + \mathbf{p})$ if and only if for $v := e^{i\langle z, \mathbf{p} \rangle} u$, $v \in \ker_{L_{\mathbf{p}}^2} D_n(\alpha; \mathbf{t})$.

Comparing the remaining terms in the decomposition (2.15) and (3.10) we obtain

$$L_{(j,0)}^2 = \bigoplus_{\mathbf{p} \in \mathcal{O}_j} L_{\mathbf{p}}^2, \quad j = 1, 2,$$

where \mathcal{O}_j , $j = 1, 2$ being two orbits under action $\mathbf{p} \mapsto \bar{\omega}\mathbf{p}$ given by

$$\mathbb{Z}_3 \setminus \Delta = \mathcal{O}_1 \sqcup \mathcal{O}_2 = \{(1, 0), (0, 2), (2, 1)\} \sqcup \{(2, 0), (0, 1), (1, 2)\}. \quad (3.11)$$

3.3. Spectral characterization of magic angles. We start by introducing some basic properties of the operator $D_n(\alpha; \mathbf{t})$ with domain H_0^1 . We first observe that

$$\text{Spec}_{L_0^2} D_n(0; \mathbf{t}) = \Lambda^* + \{\mathbf{0}, \mathbf{i}\}, \quad (3.12)$$

since for $e_{\mathbf{k}}(z) := e^{-i\langle z, \mathbf{k} \rangle}$,

$$(D_n(0; \mathbf{t}) + \mathbf{k})e_{\mathbf{k}}\mathbf{e}_1 = 0, \quad \mathbf{k} \in \Lambda^* + \{-\mathbf{i}\}; \quad (D_n(0; \mathbf{t}) + \mathbf{k})e_{\mathbf{k}}\mathbf{e}_{2n} = 0, \quad \mathbf{k} \in \Lambda^* + \{\mathbf{0}\},$$

where the exponentials $e_{\mathbf{k}}/\text{vol}(\mathbb{C}/\Lambda)^{\frac{1}{2}}$, $\mathbf{k} \in \Lambda^*$ form an orthonormal basis of $L^2(\mathbb{C}/\Lambda)$ and $\{\mathbf{e}_j\}_{1 \leq j \leq 2n}$ is the standard basis of \mathbb{C}^{2n} . This leads to the following proposition:

Proposition 3.1. *The family $\mathbb{C} \ni \alpha \mapsto D_n(\alpha; \mathbf{t}) : H_0^1 \rightarrow L_0^2$ is a holomorphic family of elliptic Fredholm operators of index 0, and for all α , the spectrum of $D_n(\alpha; \mathbf{t})$ is Λ^* -periodic:*

$$\text{Spec}_{L_0^2} D_n(\alpha; \mathbf{t}) = \text{Spec}_{L_0^2} D_n(\alpha; \mathbf{t}) + \mathbf{k}, \quad \mathbf{k} \in \Lambda^*. \quad (3.13)$$

Proof. Since $D_{\bar{z}}$ is an elliptic operator in dimension 2, existence of parametrices (cf. [DZ19, Proposition E.32]) implies the Fredholm property (cf. [DZ19, §C.2] for Fredholm operators). In view of (3.12), $D_n(0; \mathbf{t}) + \mathbf{k}$ is invertible for $\mathbf{k} \notin \Lambda^* + \{\mathbf{0}, -\mathbf{i}\}$ and hence $D_n(0; \mathbf{t}) : H_0^1 \rightarrow L_0^2$ is an operator of index 0. The same is true for the Fredholm family $D_n(\alpha; \mathbf{t})$.

To see (3.13), note that if $(D_n(\alpha; \mathbf{t}) + \lambda)\mathbf{u} = 0$ with $\mathbf{u} \in H_0^1$, then $(D_n(\alpha; \mathbf{t}) + \lambda + \mathbf{k})(e_{\mathbf{k}}\mathbf{u}) = 0$ with $e_{\mathbf{k}}\mathbf{u} \in H_0^1$ for any $\mathbf{k} \in \Lambda^*$. \square

Therefore for each \mathbf{k} , the operator $H_{n,\mathbf{k}}(\alpha; \mathbf{t})$ is an elliptic differential system and hence it has a discrete spectrum that then describes the spectrum of $H_n(\alpha; \mathbf{t})$ on $L^2(\mathbb{C})$:

$$\text{Spec}_{L^2(\mathbb{C})} H_n(\alpha; \mathbf{t}) = \bigcup_{\mathbf{k} \in \mathbb{C}/\Lambda^*} \text{Spec}_{L_0^2} H_{n,\mathbf{k}}(\alpha; \mathbf{t}), \quad (3.14)$$

$$\text{Spec}_{L_0^2} H_{n,\mathbf{k}}(\alpha; \mathbf{t}) = \{\pm E_j(\mathbf{k}, \alpha, \mathbf{t})\}_{j=1}^{\infty}, \quad E_{j+1}(\mathbf{k}, \alpha, \mathbf{t}) \geq E_j(\mathbf{k}, \alpha, \mathbf{t}) \geq 0.$$

Note that sometimes we omit (α, \mathbf{t}) and simply write $H_{n,\mathbf{k}}, D_n$ and $E_j(\mathbf{k})$ when there is no ambiguity. To conclude the second identity in (3.14), note that

$$(\lambda - \mathcal{A})^{-1} = \begin{pmatrix} (\lambda^2 - A^*A)^{-1} & 0 \\ 0 & (\lambda^2 - AA^*)^{-1} \end{pmatrix} \begin{pmatrix} \lambda & A^* \\ A & \lambda \end{pmatrix}, \quad \mathcal{A} := \begin{pmatrix} 0 & A^* \\ A & 0 \end{pmatrix}. \quad (3.15)$$

Hence, non-zero eigenvalues of $H_{n,\mathbf{k}}$ on L_0^2 are given by \pm the non-zero singular values of $D_n + \mathbf{k}$, i.e., eigenvalues of the operator $[(D_n^* + \bar{\mathbf{k}})(D_n + \mathbf{k})]^{1/2}$, counted with their multiplicities.

The zero eigenvalue (if exists) of $(D_n^* + \bar{\mathbf{k}})(D_n + \mathbf{k})$ also has the same multiplicity as the zero eigenvalue of $(D_n + \mathbf{k})(D_n^* + \bar{\mathbf{k}})$, so that eigenvalues $E_j(\mathbf{k}) = 0$ are included exactly twice (for \pm), this follows from the identity

$$\dim \ker_{L_0^2}(D_n + \mathbf{k}) = \dim \ker_{L_0^2}(D_n^* + \bar{\mathbf{k}}), \quad (3.16)$$

which follows from Proposition 3.1: the operator $D_n + \mathbf{k}$ is a Fredholm operator of index 0.

Combining with the equation (3.6), The above discussion yields three equivalent characterizations of the existence of a *flat band* at energy zero:

Proposition 3.2. *The following statements are equivalent for $\alpha \in \mathbb{C}$:*

- (1) $0 \in \bigcap_{\mathbf{k} \in \mathbb{C}} \text{Spec}_{L_0^2} H_{n,\mathbf{k}}(\alpha; \mathbf{t})$
- (2) $E_0(\mathbf{k}, \alpha, \mathbf{t}) = 0$ for all $\mathbf{k} \in \mathbb{C}/\Lambda^*$.
- (3) $\text{Spec}_{L_0^2} D_n(\alpha; \mathbf{t}) = \mathbb{C}$.

4. MAGIC ANGLES OF THE TWISTED MULTILAYER HAMILTONIAN

The *magic angles* are defined as the angles $1/\alpha$'s at which

$$0 \in \bigcap_{\mathbf{k} \in \mathbb{C}} \text{Spec}_{L_0^2} H_{n,\mathbf{k}}(\alpha; \mathbf{t}) \quad (4.1)$$

As discussed in Section 3, the Hamiltonian $H_{n,\mathbf{k}}(\alpha; \mathbf{t})$ comes from the *Floquet theory* of $H_n(\alpha; \mathbf{t})$ and (4.1) means that $H_n(\alpha; \mathbf{t})$ has a *flat band* at 0 which is given by the equivalent condition:

$$0 \in \bigcap_{\mathbf{k} \in \mathbb{C}} \text{Spec}_{L_0^2} H_{n,\mathbf{k}}(\alpha; \mathbf{t}) \iff \text{Spec}_{L_0^2} D_n(\alpha; \mathbf{t}) = \mathbb{C}. \quad (4.2)$$

4.1. Magic angles of TBG. We briefly recall several results on flat bands and magic angles of TBG from [BHZ22a].

Let \mathcal{A} denote the sets of all magic angles of $D(\alpha)$ with $\alpha \in \mathbb{C}$, then the spectrum of $D(\alpha)$ on L_0^2 is given by

$$\text{Spec}_{L_0^2} D(\alpha) = \begin{cases} \Lambda^* + \{\mathbf{0}, \mathbf{i}\} & \alpha \notin \mathcal{A}, \\ \mathbb{C} & \alpha \in \mathcal{A}. \end{cases}$$

where $R_0(\mathbf{k}) := (D(0) - \mathbf{k})^{-1} : L_0^2 \rightarrow H_0^1$ for $\mathbf{k} \notin \Lambda^* + \{\mathbf{0}, \mathbf{i}\}$. We only need to show that for any $\alpha \in \mathbb{C}$ fixed $D_n(\alpha; \mathbf{t}) - \mathbf{k}$ is invertible if and only if $D(\alpha) - \mathbf{k}$ is invertible for $\mathbf{k} \notin \Lambda^* + \{\mathbf{0}, \mathbf{i}\}$. Following from a direct computation and the identity $T_+ R_0(\mathbf{k}) T_- = 0$, we have the following decomposition

$$D_n(\alpha; \mathbf{t}) - \mathbf{k} = (\text{Id} + J_{\mathbf{t},+}) \text{diag}(D(\alpha) - \mathbf{k}, D(0) - \mathbf{k}, \dots, D(0) - \mathbf{k}) (\text{Id} + J_{\mathbf{t},-}). \quad (4.5)$$

Since $\text{Spec}_{L_0^2} D(0) = \Lambda^* + \{\mathbf{0}, \mathbf{i}\}$, the operator $D_n(\alpha; \mathbf{t}) - \mathbf{k}$ is invertible if and only if $D(\alpha) - \mathbf{k}$ is invertible for $\mathbf{k} \notin \Lambda^* + \{\mathbf{0}, \mathbf{i}\}$. \square

Remark 7. *The decomposition (4.5) also concludes that the multiplicity of the flat band of $D_n(\alpha; \mathbf{t})$ is the same as the multiplicity of the flat band of $D(\alpha)$.*

5. BAND SEPARATION WITH INTER-LAYER TUNNELING

We discuss, using the perturbation theory, the band separation mechanism appeared in [WL22] once the tunneling \mathbf{t} being turned on. We set up a Grushin problem and compute the separation of the band explicitly. For a detailed introduction on the Grushin problem, we refer to [DZ19, Appendix C].

5.1. A Grushin Problem. Following the identity (3.15), eigenvalues of $H_{n,\mathbf{k}}(\alpha; \mathbf{t})$ are given by singular values of $D_n(\alpha; \mathbf{t}) + \mathbf{k}$, i.e., eigenvalues of $[(D_n^*(\alpha; \mathbf{t}) + \bar{\mathbf{k}})(D_n(\alpha; \mathbf{t}) + \mathbf{k})]^{1/2}$, counted with multiplicities. For simplicity purposes, we work with the case $n = 2$, the general case is essentially the same. We first introduce the Schur complement formula:

Proposition 5.1. *Suppose*

$$\begin{pmatrix} P & R_- \\ R_+ & R_{+-} \end{pmatrix} = \begin{pmatrix} E & E_+ \\ E_- & E_{-+} \end{pmatrix}^{-1} : X_1 \times X_- \rightarrow X_2 \times X_+ \quad (5.1)$$

are bounded operators on Banach spaces, then P is invertible if and only if E_{-+} is invertible. Moreover, in such case we have

$$P^{-1} = E - E_+ E_{-+}^{-1} E_-, \quad E_{-+}^{-1} = R_{+-} - R_+ P^{-1} R_-. \quad (5.2)$$

Proof. See [DZ19, Appendix C.1]. \square

Define $D_{\mathbf{k}}(\alpha) := D(\alpha) + \mathbf{k}$ and $D_{2,\mathbf{k}}(\alpha; t) := D_2(\alpha; t) + \mathbf{k}$. We have

$$D_{2,\mathbf{k}}(\alpha; t) = D_{2,\mathbf{k}}(\alpha; 0) + tT, \quad T = \begin{pmatrix} 0 & T_+ \\ T_- & 0 \end{pmatrix}.$$

We consider the operator $P_{\mathbf{k}}^t = D_{2,\mathbf{k}}^*(\alpha; t) D_{2,\mathbf{k}}(\alpha; t)$ given by

$$\begin{aligned} P_{\mathbf{k}}^t &= P_{\mathbf{k}}^0 + tQ + t^2 T^* T \\ &:= D_{2,\mathbf{k}}^*(\alpha; 0) D_{2,\mathbf{k}}(\alpha; 0) + t[T^* D_{2,\mathbf{k}}(\alpha; 0) + D_{2,\mathbf{k}}^*(\alpha; 0) T] + t^2 T^* T \end{aligned}$$




FIGURE 1. First three band structure $\mathbf{k} \mapsto E_j(\alpha, \mathbf{k}, t)$ of $H_2(\alpha; t)$ (cf. equation (3.7)) away from (left) and at the first magic angle of $H(\alpha)$ (right). Top row: $t = 0$; there exist two Dirac cones at points K, K' . Bottom row: $t = 0.5$; the Dirac bands separate from the flat band and a band gap is open.

We start by considering the zero spectrum of the operator $P_{\mathbf{k}}^0 = D_{2,\mathbf{k}}^*(\alpha; 0)D_{2,\mathbf{k}}(\alpha; 0)$. By equation (2.18) and (2.19) and the discussion of §3.2, $\ker D_2(\alpha; 0)$ are given by

$$\Phi_1, \mathbf{e}_3 \in L_{-\mathbf{i}}^2, \quad \Phi_2, \mathbf{e}_4 \in L_{\mathbf{0}}^2. \quad (5.3)$$

Here we slightly abuse notations by restricting $\ker H_2(\alpha; 0) = \{\Phi_1, \Phi_2, \mathbf{e}_3, \mathbf{e}_4\}$ to the first component $\ker D_2(\alpha; 0)$ because of the direct sum decomposition (2.17). Following the spirit of Remark 6, we define

$$\hat{\Phi}_1 := e^{i\langle z, \mathbf{i} \rangle} \Phi_1, \quad \hat{\mathbf{e}}_3 := e^{i\langle z, \mathbf{i} \rangle} \mathbf{e}_3, \quad \hat{\Phi}_2 := \Phi_2, \quad \hat{\mathbf{e}}_4 := \mathbf{e}_4 \in L_{\mathbf{0}}^2 \quad (5.4)$$

These are elements of $\ker_{L_{\mathbf{0}}^2} H_{2,\mathbf{k}}(\alpha; 0)$, or more precisely $\ker_{L_{\mathbf{0}}^2} P_{\mathbf{k}}^0$, for $\mathbf{k} = -\mathbf{i}, \mathbf{0}$ respectively.

5.2. Analysis at K' . To see how zero spectrum changes as we turning on $t > 0$, we consider the following Grushin problem for the perturbed operator $P_{\mathbf{k}}^t - z$ at the Dirac

point $\mathbf{k} = K' = -\mathbf{i}$:

$$\begin{pmatrix} P_{\mathbf{k}}^t - z & R_- \\ R_+ & 0 \end{pmatrix} : H_{\mathbf{0}}^1 \oplus \mathbb{C}^2 \longrightarrow L_{\mathbf{0}}^2 \oplus \mathbb{C}^2 \quad (5.5)$$

with

$$R_- : (u_-^{(1)}, u_-^{(2)})^T \mapsto u_-^{(1)} \hat{\Phi}_1 + u_-^{(2)} \hat{\mathbf{e}}_3, \quad R_+ : u \mapsto (\langle u, \hat{\Phi}_1 \rangle, \langle u, \hat{\mathbf{e}}_3 \rangle)^T.$$

We use \mathcal{I} to denote the set $\{\hat{\Phi}_1, \hat{\mathbf{e}}_3\}$. For $t = 0$, the Grushin problem (5.5) is invertible with the inverse given by

$$\mathcal{E} = \begin{pmatrix} E & E_+ \\ E_- & E_{-+} \end{pmatrix} : L_{\mathbf{0}}^2 \oplus \mathbb{C}^2 \longrightarrow H_{\mathbf{0}}^1 \oplus \mathbb{C}^2 \quad (5.6)$$

with

$$Ev = \sum_{u_j \notin \mathcal{I}} \frac{1}{z_j - z} \langle v, u_j \rangle u_j, \quad E_+ v_+ = R_- v_+, \\ E_- v = R_+ v, \quad E_{-+} = \begin{pmatrix} z \\ z \end{pmatrix}$$

where the set $\{u_j\} \subset L_{\mathbf{0}}^2$ is the set of all normalized eigenfunctions of $P_{\mathbf{k}}^0$ with eigenvalues $\{z_j\}$. Note that by definition of $D_n(\alpha; \mathbf{t})$ in equation (1.2), the eigenspace of $P_{\mathbf{k}}^0 = D_{2,\mathbf{k}}^*(\alpha; 0) D_{2,\mathbf{k}}(\alpha; 0)$ is given by the direct sum of eigenspaces of $D_{\mathbf{k}}^*(\alpha) D_{\mathbf{k}}(\alpha)$ and $D_{\mathbf{k}}^*(0) D_{\mathbf{k}}(0)$. Therefore we can take a complete basis of eigenfunctions $\{u_j\}$ with any element u_j is either of the form $(u_j^{(1)}, 0_{\mathbb{C}^2})^T$ or $(0_{\mathbb{C}^2}, u_j^{(2)})^T$, where $\{u_j^{(1)}\}$ are eigenfunctions of $D_{\mathbf{k}}^*(\alpha) D_{\mathbf{k}}(\alpha)$ and $\{u_j^{(2)}\}$ are eigenfunctions of $D_{\mathbf{k}}^*(0) D_{\mathbf{k}}(0)$. To work with the perturbed operator $P_{\mathbf{k}}^t$, we employ the following proposition.

Proposition 5.2. *For $|t| \ll 1$, the Grushin problem for $P_{\mathbf{k}}^t - z$:*

$$\mathcal{P}^t = \begin{pmatrix} P_{\mathbf{k}}^t - z & R_- \\ R_+ & 0 \end{pmatrix}$$

is invertible. If the inverse $(\mathcal{P}^t)^{-1}$ is given by $\begin{pmatrix} E^t & E_+^t \\ E_-^t & E_{-+}^t \end{pmatrix}$, then

$$E_{-+}^t = E_{-+} + \sum_{k=1}^{\infty} (-t)^k E_- (Q + tT) (E(Q + tT))^{k-1} E_+ \quad (5.7)$$

Proof. This follows directly from [TZ23, Proposition 2.12]. \square

Proposition 5.2 yields that

$$E_{-+}^t = E_{-+} - t E_- Q E_+ + t^2 E_- (Q E Q - T) E_+ + \mathcal{O}(t^3),$$

with

$$E_-QE_+ = \begin{pmatrix} \langle Q\hat{\Phi}_1, \hat{\Phi}_1 \rangle & \langle Q\hat{e}_3, \hat{\Phi}_1 \rangle \\ \langle Q\hat{\Phi}_1, \hat{e}_3 \rangle & \langle Q\hat{e}_3, \hat{e}_3 \rangle \end{pmatrix} \quad (5.8)$$

where the inner product is taken in the space $L_0^2(\mathbb{C}/3\Lambda; \mathbb{C}^4)$. The expression of Q and the fact that $\hat{\Phi}_1$ and \hat{e}_3 being zero eigenfunctions of $D_{2,\mathbf{k}}(\alpha; 0)$ at $\mathbf{k} = K'$ yields that equation (5.8) vanishes.

Now we compute the second order perturbation term in t . Writing $\hat{\Phi}_1 = (\varphi_1, 0_{\mathbb{C}^2})^T$ and $\hat{e}_3 = (0_{\mathbb{C}^2}, e_1)^T$ where $\varphi_1, e_1 \in L_0^2(\mathbb{C}/3\Lambda; \mathbb{C}^2)$, we have the following identities

$$T\hat{\Phi}_1 = (0_{\mathbb{C}^2}, T_-\varphi_1)^T, \quad T\hat{e}_3 = (T_+e_1, 0_{\mathbb{C}^2})^T.$$

The operator $E_-TE_+ : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ is then given by the matrix

$$\begin{pmatrix} \langle T\hat{\Phi}_1, T\hat{\Phi}_1 \rangle & \langle T\hat{\Phi}_1, T\hat{e}_3 \rangle \\ \langle T\hat{e}_3, T\hat{\Phi}_1 \rangle & \langle T\hat{e}_3, T\hat{e}_3 \rangle \end{pmatrix} = \begin{pmatrix} \|T_-\varphi_1\|^2 & 0 \\ 0 & \|T_+e_1\|^2 \end{pmatrix}, \quad (5.9)$$

while the operator $E_-QEQE_+ : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ is given by the matrix

$$\sum_{u_j \notin \mathcal{I}} \frac{1}{z_j - z} \begin{pmatrix} |\langle T\hat{\Phi}_1, D_{2,\mathbf{k}}(\alpha; 0)u_j \rangle|^2 & \langle T\hat{e}_3, D_{2,\mathbf{k}}(\alpha; 0)u_j \rangle \langle D_{2,\mathbf{k}}(\alpha; 0)u_j, T\hat{\Phi}_1 \rangle \\ \langle T\hat{\Phi}_1, D_{2,\mathbf{k}}(\alpha; 0)u_j \rangle \langle D_{2,\mathbf{k}}(\alpha; 0)u_j, T\hat{e}_3 \rangle & |\langle T\hat{e}_3, D_{2,\mathbf{k}}(\alpha; 0)u_j \rangle|^2 \end{pmatrix}.$$

Writing $u_j = (u_j^{(1)}, u_j^{(2)})^T$, the above matrix can be rewritten as

$$\sum_{u_j \notin \mathcal{I}} \frac{1}{z_j - z} \begin{pmatrix} |\langle T_-\varphi_1, D_{\mathbf{k}}(0)u_j^{(2)} \rangle|^2 & \langle T_+e_1, D_{\mathbf{k}}(\alpha)u_j^{(1)} \rangle \langle D_{\mathbf{k}}(0)u_j^{(2)}, T_-\varphi_1 \rangle \\ \langle D_{\mathbf{k}}(\alpha)u_j^{(1)}, T_+e_1 \rangle \langle T_-\varphi_1, D_{\mathbf{k}}(0)u_j^{(2)} \rangle & |\langle T_+e_1, D_{\mathbf{k}}(\alpha)u_j^{(1)} \rangle|^2 \end{pmatrix}.$$

By the direct sum decomposition of $P_{\mathbf{k}}^0$, any eigenfunctions $u_j = (u_j^{(1)}, u_j^{(2)})^T$ is of the form $(u_j^{(1)}, 0_{\mathbb{C}^2})^T$ or $(0_{\mathbb{C}^2}, u_j^{(2)})^T$, the off-diagonal terms in the above matrix vanishes. It thus reduces to a sum of diagonal matrices

$$\sum_j \frac{1}{z_j^{(2)} - z} \begin{pmatrix} |\langle T_-\varphi_1, D_{\mathbf{k}}(0)u_j^{(2)} \rangle|^2 & 0 \\ 0 & 0 \end{pmatrix} + \sum_i \frac{1}{z_i^{(1)} - z} \begin{pmatrix} 0 & 0 \\ 0 & |\langle T_+e_1, D_{\mathbf{k}}(\alpha)u_i^{(1)} \rangle|^2 \end{pmatrix} \quad (5.10)$$

in which $z_i^{(1)} \neq 0$ and $z_j^{(2)} \neq 0$ with

$$(D_{\mathbf{k}}^*(\alpha)D_{\mathbf{k}}(\alpha) - z_i^{(1)})u_i^{(1)} = 0, \quad (D_{\mathbf{k}}^*(0)D_{\mathbf{k}}(0) - z_j^{(2)})u_j^{(2)} = 0.$$

We consider the first term in the expression (5.10). Note that $\{u_j^{(2)}\}$ are joint eigenfunctions of $D_{\mathbf{k}}^*(0)D_{\mathbf{k}}(0)$ and $D_{\mathbf{k}}(0)$ with spectrum

$$\text{Spec}_{L_0^2} D_{\mathbf{k}}(0) = \Lambda^* + \{\mathbf{0}, -\mathbf{i}\} \text{ for } \mathbf{k} = -\mathbf{i}$$

Therefore,

$$|\langle T_-\varphi_1, D_{\mathbf{k}}(0)u_j^{(2)} \rangle|^2 = z_j^{(2)} |\langle T_-\varphi_1, u_j^{(2)} \rangle|^2.$$

Using the fact that $\langle T_- \varphi_1, e_1 \rangle = 0$ and Plancherel theorem, at $z = 0$ we obtain the equality

$$\sum_j \frac{z_j^{(2)}}{z_j^{(2)} - z} \begin{pmatrix} |\langle T_- \varphi_1, u_j^{(2)} \rangle|^2 & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} \|T_- \varphi_1\|^2 & 0 \\ 0 & 0 \end{pmatrix} \quad (5.11)$$

For $z = 0$, combining equations (5.9) (5.10) and (5.11), the second order perturbation is given by

$$E_-(QEQ - T)E_+ = \begin{pmatrix} 0 & 0 \\ 0 & -\|T_+ e_1\|^2 + \sum_i \frac{1}{z_i^{(1)}} |\langle T_+ e_1, D_{\mathbf{k}}(\alpha) u_i^{(1)} \rangle|^2 \end{pmatrix},$$

recall that $\{z_i^{(1)}\}$ is the set of non-zero eigenvalues of $D_{\mathbf{k}}^*(\alpha) D_{\mathbf{k}}(\alpha)$ with L_0^2 -eigenfunctions $\{u_i^{(1)}\}$. Therefore by Proposition 5.1, 5.2, zero is an eigenvalue for $P_{\mathbf{k}}^t$ closed to $\mathbf{k} = K'$. The zero eigenvalue corresponds to the flat band of $H_2(\alpha; t)$.

The eigenvalue of the operator $P_{\mathbf{k}}^t$ corresponding the Dirac cone of $P_{\mathbf{k}}^0$ is given by

$$\lambda_1(t) = t^2 \left[\|T_+ e_1\|_{L^2}^2 - \sum_i \frac{1}{z_i^{(1)}} |\langle T_+ e_1, D_{\mathbf{k}}(\alpha) u_i^{(1)} \rangle|^2 \right] + \mathcal{O}(t^3),$$

which corresponds to the separation of the Dirac band from the flat band. Since this eigenvalue is given by the solution to the equation

$$z - t^2 \left[\|T_+ e_1\|_{L^2}^2 - \sum_i \frac{1}{z_i^{(1)} - z} |\langle T_+ e_1, D_{\mathbf{k}}(\alpha) u_i^{(1)} \rangle|^2 \right] = 0,$$

a substitution $z = \lambda t^2$ and a geometric series argument yield the solution $z = \lambda_1(t)$.

Combining with the fact that there is a protected states at 0 for any $t \in \mathbb{R}$, this proves the following proposition:

Proposition 5.3. *For $t \ll 1$ and at $\mathbf{k} = K'$, the first two eigenvalues of $H_{2,\mathbf{k}}(\alpha; t)$ are given by $E_{1,\mathbf{k}}(t) = 0$ and $E_{2,\mathbf{k}}(t) = Ct + \mathcal{O}(t^{3/2})$, where the constant is given by*

$$C = \left(1 - \sum_i \frac{1}{z_i^{(1)}} |\langle T_+ e_1, D_{\mathbf{k}}(\alpha) u_i^{(1)} \rangle|^2 \right)^{1/2}.$$

6. FLATBAND EIGENFUNCTIONS AND CHERN NUMBERS

In the section, we construct flat band eigenfunctions of $H_{n,\mathbf{k}}(\alpha; \mathbf{t})$ at $E_1(\mathbf{k}, \alpha; \mathbf{t}) = 0$ for $\mathbf{k} \in \mathbb{C}/\Lambda^*$ using a theta function argument analogous to [BEWZ22] (see also [LTKV20]). The construction also gives an explicit method to compute the Chern number of the flat band of this twisted multilayer model. We start by reviewing some backgrounds in theta functions.

6.1. Review of theta functions. We recall that the theta function $\theta(\zeta) := -\theta_{\frac{1}{2}, \frac{1}{2}}(\zeta|\omega)$ is given by

$$\begin{aligned} \theta(\zeta) &= -\sum_{n \in \mathbb{Z}} \exp(\pi i(n + \frac{1}{2})^2 \omega + 2\pi i(n + \frac{1}{2})(\zeta + \frac{1}{2})), \quad \theta(-\zeta) = -\theta(\zeta), \\ \theta(\zeta + m) &= (-1)^m \theta(\zeta), \quad \theta(\zeta + n\omega) = (-1)^n e^{-\pi i n^2 \omega - 2\pi i \zeta n} \theta(\zeta), \end{aligned} \quad (6.1)$$

where

$$\theta_{a,b}(\zeta|\tau) := \sum_{n \in \mathbb{Z}} \exp(\pi i(a+n)^2 \tau + 2\pi i(n+a)(\zeta+b)), \quad \text{Im } \tau > 0.$$

The function $\theta(\zeta)$ has simple zeros at

$$\mathbb{Z}\omega + \mathbb{Z},$$

see [Mum83]. If we consider

$$g(\zeta) := e^{2\pi i \zeta a} \frac{\theta(\zeta + k)}{\theta(\zeta)},$$

we obtain a meromorphic function with simple poles at $\mathbb{Z} + \omega\mathbb{Z}$, simple zeros at $-k + \mathbb{Z} + \omega\mathbb{Z}$, and satisfying $g(\zeta + m + n\omega) = e^{2\pi i(am + (\omega a - k)n)} g(\zeta)$. Hence, by choosing $k = \omega a - b$, we define

$$g_k(\zeta) := e^{2\pi i \zeta(k - \bar{k})/\sqrt{3}} \frac{\theta(\zeta + k)}{\theta(z)}, \quad k = \omega a - b, \quad a, b \in \mathbb{R}. \quad (6.2)$$

It satisfies

$$\begin{aligned} g_k(\zeta + m + n\omega) &= e^{2\pi i(am + bn)} g_k(\zeta), \quad k = \omega a - b, \quad a, b \in \mathbb{R}, \\ am + nb &= \frac{i}{\sqrt{3}} (\zeta_0 \bar{k} - \bar{\zeta}_0 k), \quad \zeta_0 = m + n\omega, \quad m, n \in \mathbb{Z}, \\ g_{k+p\omega-q}(\zeta) &= (-1)^{p+q} e^{-2\pi i kp - \pi i p^2 \omega} g_k(\zeta), \quad p, q \in \mathbb{Z}. \end{aligned} \quad (6.3)$$

We then define a function periodic with respect to the lattice $\omega\mathbb{Z} + \mathbb{Z}$:

$$F_k(\zeta) := e^{2\pi(\zeta - \bar{\zeta})k/\sqrt{3}} \frac{\theta(\zeta + k)}{\theta(\zeta)}. \quad (6.4)$$

6.2. Construction of flat bands. We first transplant the functions constructed above to the lattice Λ and its dual Λ^* . Recall that in equation (2.2), they are given by

$$\Lambda = \frac{4}{3}\pi i \omega (\mathbb{Z} \oplus \omega\mathbb{Z}), \quad \Lambda^* = \sqrt{3}\omega (\omega\mathbb{Z} \oplus \mathbb{Z}).$$

We introduce change of variables

$$z = \frac{4}{3}\pi i \omega \zeta, \quad \mathbf{k} = \sqrt{3}\omega k$$

in the equation (6.2) (6.4), so that the new function is periodic with respect to the lattice Λ , i.e., we define

$$F_{\mathbf{k}}(z) := F_k\left(\frac{3z}{4\pi i \omega}\right), \quad F_{\mathbf{k}}(z + \mathbf{a}) = F_{\mathbf{k}}(z), \quad \mathbf{a} \in \Lambda, \quad (6.5)$$

Following from [BHZ22a, Lemma 3.1], $F_{\mathbf{k}}(z)$ is then the fundamental solution of $(2D_{\bar{z}} + \mathbf{k})$ in the sense that

$$(2D_{\bar{z}} + \mathbf{k})F_{\mathbf{k}}(z) = \alpha_{\mathbf{k}}\delta_0(z), \quad \alpha_{\mathbf{k}} \neq 0, \quad \mathbf{k} \notin \Lambda^*, \quad z \in \mathbb{C}/\Lambda. \quad (6.6)$$

Therefore the Schwartz kernel of

$$(2D_{\bar{z}} + \mathbf{k})^{-1} : L^2(\mathbb{C}/\Lambda) \rightarrow H^1(\mathbb{C}/\Lambda) \quad (6.7)$$

is given by $c_{\mathbf{k}}F_{\mathbf{k}}(z - z')$, with $c_{\mathbf{k}} = \theta'(0)/(\pi\theta(k)) = 1/\alpha_{\mathbf{k}}$.

Recall that in [BEWZ22, Proposition 3.4] for $\alpha \in \mathcal{A}$ simple, a solution to $(D(\alpha) + \mathbf{k})\mathbf{u}_{\mathbf{k}} = 0$ for any $\mathbf{k} \in \mathbb{C}$ is constructed using a theta function argument. In particular, for $\mathbf{k} \notin \Lambda^* + \{\mathbf{0}, -\mathbf{i}\}$ this implies, by Proposition 4.1, that $\alpha \in \mathbb{C}$ is a magic angle of $H(\alpha)$. Recall that the magic angles of $H_n(\alpha; \mathbf{t})$ coincides with the magic angles of $H(\alpha)$. We therefore have a similar construction of all flat band eigenfunctions of $H_{n,\mathbf{k}}(\alpha; \mathbf{t})$.

Recall that by Proposition 4.2, for $\alpha \in \mathcal{A}$ simple, i.e. flat band has multiplicity one, $\mathbf{u} \in \ker_{L_0^2} D(\alpha)$ vanishes at the point $-z_S + \Lambda^*$. Therefore we have

$$\mathbf{v}_{\mathbf{k}}(z) := F_{\mathbf{k}}(z + z_S)\mathbf{u}(z), \quad \mathbf{v}_{\mathbf{k}} \in L_0^2, \quad (D(\alpha) + \mathbf{k})\mathbf{v}_{\mathbf{k}} = 0, \quad (6.8)$$

Define the function

$$\Phi_{\mathbf{k}}(z) = (\phi_{1,\mathbf{k}}, \phi_{2,\mathbf{k}}, \dots, \phi_{2n,\mathbf{k}})^T \in L_0^2 \quad (6.9)$$

with

- (1) $\phi_{2j-1,\mathbf{k}} = 0$ for $j = 2, 3, \dots, n$;
- (2) $(\phi_{1,\mathbf{k}}, \phi_{2,\mathbf{k}})^T = \Theta_{n-1}(\mathbf{k})\mathbf{v}_{\mathbf{k}}(z)$;
- (3) $\phi_{2m+2,\mathbf{k}} = (\prod_{j=1}^m t_j) \cdot (2D_{\bar{z}} + \mathbf{k})^{-m}\phi_{2,\mathbf{k}}$, $m = 1, 2, \dots, n-1$,

where $(2D_{\bar{z}} + \mathbf{k})^{-m}$ denotes the m -th powers of the resolvent $(2D_{\bar{z}} + \mathbf{k})^{-1} : L_0^2 \rightarrow L_0^2$ and

$$\Theta_{n-1}(\mathbf{k}) := [\theta(\mathbf{k})]^{n-1} \quad (6.10)$$

It is easy to verify that $\Phi_{\mathbf{k}}(z)$ defined above solves the equation

$$(D_n(\alpha; \mathbf{t}) + \mathbf{k})\Phi_{\mathbf{k}}(z) = 0 \quad (6.11)$$

for $\mathbf{k} \notin \Lambda^*$, as the resolvent $(2D_{\bar{z}} + \mathbf{k})^{-1}$ is invertible in $\mathbb{C} \setminus \Lambda^*$.

For $\mathbf{k} \in \Lambda^*$, without loss of generality we consider $\mathbf{k} = \mathbf{0}$, otherwise multiplication by $e^{-i\langle z, \mathbf{k} \rangle} \in L^2(\mathbb{C}/\Lambda)$ reduces it to the zero case. Note that the (Schwartz kernel of) resolvent is meromorphic at $\mathbf{k} = \mathbf{0}$ with a simple pole, i.e.,

$$(2D_{\bar{z}} + \mathbf{k})^{-1} = \frac{1 \otimes 1}{\mathbf{k}} + f_{\mathbf{k}}(z, z'), \quad (6.12)$$

where $f_{\mathbf{k}}$ is holomorphic in \mathbf{k} . The function $\Theta_{n-1}(\mathbf{k})$ is holomorphic in \mathbf{k} and has a zero at $\mathbf{k} = \mathbf{0}$ with order $n-1$. Therefore, each component $\phi_{i,\mathbf{k}}$, $i = 1, \dots, 2n$ is

holomorphic in \mathbf{k} at $\mathbf{k} = \mathbf{0}$. In particular, the construction (3) above yields that at $\mathbf{k} = \mathbf{0}$

$$\phi_{2m+2,\mathbf{0}} = 0, \quad m = 1, 2, \dots, n-2; \quad \phi_{2n,\mathbf{0}} = \left(\prod_{j=1}^m t_j \right) \int_{C/\Lambda} u_2(z) dm(z), \quad (6.13)$$

where $\mathbf{v}_0 = \mathbf{u} = (u_1, u_2)$. This yields the flat band eigenfunction at $\mathbf{k} = \mathbf{0}$ provided that the integral appeared in the above equation is non-zero. This is verified numerically for the first five real magic angles, where the multiplicities of the flat band are one (see Table 1).

α	0.586	2.22	3.75	5.28	6.79	8.31
$I(u_2)$	0.2345	0.0542	0.0033	0.0022	0.0013	8.3963×10^{-4}

TABLE 1. Numerical results of the integral $I(u_2) = \int_{C/\Lambda} u_2(z) dm(z)$ in (6.13) of the flat band eigenfunction on one fundamental domain C/Λ

Remark 8. *This agrees with the protected eigenstates result shown using irreducible representations, where $\mathbf{e}_{2n} \in L_0^2$ is always a protected eigenstate with zero-energy.*

6.3. High Chern numbers of the flat band. In this section, assuming the flat band is simple, we show that the flat band eigenfunctions constructed above give rise to a holomorphic line bundle with Chern number $c_1(L) = -n$. For basics on holomorphic line bundles, Berry connections and Chern numbers, we refer to [TZ23].

We use the notations in §6.1 for simplicity. By equation (6.1) (6.3) and (6.7), we obtain

$$\Phi_{k+p} = [e_p^{(n)}(k)]^{-1} \tau(p) \Phi_k \quad (6.14)$$

where

$$[\tau(p)f](\zeta) = e^{-2\pi i(\zeta_1 p_1 + \zeta_2 p_2)} f(\zeta), \quad e_p^{(n)}(k) = [e^{\pi i p_1^2 \omega + 2\pi i k p_1} (-1)^{p_1 + p_2}]^n, \quad (6.15)$$

$$\zeta = \zeta_1 + \omega \zeta_2, \quad p = \omega p_1 - p_2, \quad p_j \in \mathbb{Z}, \quad k \in \mathbb{C}. \quad (6.16)$$

Note that

$$\tau(p) : \ker_{L_0^2}(D_n(\alpha; \mathbf{t}) + k) \rightarrow \ker_{L_0^2}(D_n(\alpha; \mathbf{t}) + k + p),$$

is a unitary transformation. Here we slightly abuse notations by identifying L_0^2 and $D_n(\alpha; \mathbf{t}) + k$ with corresponding parts in the rescaled lattice space. Following the standard construction we define

$$L := \left\{ [k, \Phi]_\tau \in (\mathbb{C} \times L_0^2) / \sim_\tau : \Phi \in \ker_{L_0^2}(D_n(\alpha; \mathbf{t}) + k) \right\}, \quad (6.17)$$

$$[k, \Phi]_\tau = [k', \Phi']_\tau \Leftrightarrow (k, \Phi) \sim_\tau (k', \Phi') \Leftrightarrow \exists p \in \Lambda \quad k' = k + p, \quad \Phi' = \tau(p)\Phi.$$

Therefore, we have the following lemma:

Lemma 6.1. *Definition (6.17) gives a holomorphic line bundle over \mathbb{C}/Λ^* :*

$$f : L \rightarrow \mathbb{C}/\Lambda^* \quad \text{with} \quad [k, \Phi]_\tau \mapsto [k] \in \mathbb{C}/\Lambda^*.$$

The corresponding family of multipliers is given by $k \mapsto e_p^{(n)}(k)$.

Proof. The action of the discrete group Λ^* , with $\Lambda^* \ni p : (k, \Phi) \mapsto (k + p, \tau(p)\Phi)$, on the trivial complex line bundle

$$\tilde{L} := \{(k, \kappa\Phi_k) : k \in \mathbb{C}, \kappa \in \mathbb{C}\} \simeq \mathbb{C}_k \times \mathbb{C}_\kappa, \quad (6.18)$$

(where Φ_k is defined in (6.9)) is free and proper, and the quotient map is given by $\pi_\tau(k, \kappa\Phi_k) = [k, \kappa\Phi_k]_\tau$. Hence its quotient by that action, L , is a smooth complex manifold of dimension 2.

The map $(p, k) \mapsto e_p^{(n)}(k)$ satisfies the cocycle conditions with the action φ_p on \tilde{L} :

$$\varphi_p(k, \kappa\Phi_k) := (k + p, e_p^{(n)}(k)\kappa\Phi_{k+p}), \quad e_{p+p'}^{(n)}(k) = e_{p'}^{(n)}(k + p)e_p^{(n)}(k), \quad p, p' \in \Lambda^*, \quad (6.19)$$

and for $p \in \Lambda^*$. We then have $\pi_\tau((k, \kappa\Phi_k)) = \pi_\tau(k + p, e_p^{(n)}(k)\kappa\Phi_{k+p})$ by equation (6.14) and this gives L the structure of a complex line bundle over \mathbb{C}/Λ^* . \square

The hermitian structure is inherited from $L^2(\mathbb{C}/\Lambda)$ and the resulting hermitian structure on \tilde{L} of (6.18). In coordinates (k, τ) on \tilde{L} , we get

$$h(k) = \|\Phi_k\|_{L^2(\mathbb{C}/3\Lambda)}^2,$$

and this gives a hermitian structure on L : from (6.14) we see that

$$h(k) = |e_p^{(n)}(k)|^2 h(k + p), \quad p \in \mathbb{Z} \oplus \omega\mathbb{Z} \cong \Lambda^*. \quad (6.20)$$

We then associate a Chern connection θ and a Berry curvature $\Omega = d\theta$ to the hermitian metric h . The Chern number of the line bundle L thus can be calculated using the method of multiplier as

$$\begin{aligned} c_1(L) &= \frac{i}{2\pi} \int_{\mathbb{C}/\Lambda^*} \Omega \\ &= \frac{i}{2\pi} \left(\log e_\omega^{(n)}(1) - \log e_\omega^{(n)}(0) - \log e_1^{(n)}(\omega) - \log e_1^{(n)}(0) \right) \\ &= -n. \end{aligned} \quad (6.21)$$

This concludes the proof to Theorem 3.

REFERENCES

- [ALB20] Ahmed Abouelkomsan, Zhao Liu, and Emil J Bergholtz. Particle-hole duality, emergent fermi liquids, and fractional chern insulators in moiré flatbands. *Physical Review Letters*, 124(10):106803, 2020.

- [BEWZ21] Simon Becker, Mark Embree, Jens Wittsten, and Maciej Zworski. Spectral characterization of magic angles in twisted bilayer graphene. *Physical Review B*, 103(16):165113, 2021.
- [BEWZ22] Simon Becker, Mark Embree, Jens Wittsten, and Maciej Zworski. Mathematics of magic angles in a model of twisted bilayer graphene. *Probability and Mathematical Physics*, 3(1):69–103, 2022.
- [BHZ22a] Simon Becker, Tristan Humbert, and Maciej Zworski. Fine structure of flat bands in a chiral model of magic angles. *arXiv preprint arXiv:2208.01628*, 2022.
- [BHZ22b] Simon Becker, Tristan Humbert, and Maciej Zworski. Integrability in the chiral model of magic angles. *arXiv preprint arXiv:2208.01620*, 2022.
- [BL13] Emil J Bergholtz and Zhao Liu. Topological flat band models and fractional chern insulators. *International Journal of Modern Physics B*, 27(24):1330017, 2013.
- [BM11] Rafi Bistritzer and Allan H MacDonald. Moiré bands in twisted double-layer graphene. *Proceedings of the National Academy of Sciences*, 108(30):12233–12237, 2011.
- [CLT⁺15] Martin Claassen, Ching Hua Lee, Ronny Thomale, Xiao-Liang Qi, and Thomas P Devereaux. Position-momentum duality and fractional quantum hall effect in chern insulators. *Physical Review Letters*, 114(23):236802, 2015.
- [DZ19] Semyon Dyatlov and Maciej Zworski. *Mathematical theory of scattering resonances*, volume 200. American Mathematical Soc., 2019.
- [LAB21] Zhao Liu, Ahmed Abouelkomsan, and Emil J Bergholtz. Gate-tunable fractional chern insulators in twisted double bilayer graphene. *Physical Review Letters*, 126(2):026801, 2021.
- [LB22] Zhao Liu and Emil J Bergholtz. Recent developments in fractional chern insulators. *arXiv preprint arXiv:2208.08449*, 2022.
- [LKL⁺19] Jong Yeon Lee, Eslam Khalaf, Shang Liu, Xiaomeng Liu, Zeyu Hao, Philip Kim, and Ashvin Vishwanath. Theory of correlated insulating behaviour and spin-triplet superconductivity in twisted double bilayer graphene. *Nature communications*, 10(1):5333, 2019.
- [LTKV20] Patrick J Ledwith, Grigory Tarnopolsky, Eslam Khalaf, and Ashvin Vishwanath. Fractional chern insulator states in twisted bilayer graphene: An analytical approach. *Physical Review Research*, 2(2):023237, 2020.
- [LVK22] Patrick J Ledwith, Ashvin Vishwanath, and Eslam Khalaf. Family of ideal chern flatbands with arbitrary chern number in chiral twisted graphene multilayers. *Physical Review Letters*, 128(17):176404, 2022.
- [MO21] Bruno Mera and Tomoki Ozawa. Kähler geometry and chern insulators: Relations between topology and the quantum metric. *Physical Review B*, 104(4):045104, 2021.
- [Mum83] D Mumford. *Tata Lectures on Theta, Vol. I*. Birkhäuser, Boston, 1983.
- [OM21] Tomoki Ozawa and Bruno Mera. Relations between topology and the quantum metric for chern insulators. *Physical Review B*, 104(4):045103, 2021.
- [RB11] N. Regnault and B. Andrei Bernevig. Fractional chern insulator. *Phys. Rev. X*, 1:021014, Dec 2011.
- [SJGG12] Pablo San-Jose, José González, and Francisco Guinea. Non-abelian gauge potentials in graphene bilayers. *Physical Review Letters*, 108(21):216802, 2012.
- [TKV19] Grigory Tarnopolsky, Alex Jura Kruchkov, and Ashvin Vishwanath. Origin of magic angles in twisted bilayer graphene. *Physical review letters*, 122(10):106405, 2019.
- [TZ23] Zhongkai Tao and Maciej Zworski. PDE methods in condensed matter physics. *Lecture Notes*, 2023.

- [WCM⁺21] Jie Wang, Jennifer Cano, Andrew J Millis, Zhao Liu, and Bo Yang. Exact landau level description of geometry and interaction in a flatband. *Physical Review Letters*, 127(24):246403, 2021.
- [WKL22] Jie Wang, Semyon Klevtsov, and Zhao Liu. Origin of model fractional chern insulators in all topological ideal flatbands: Explicit color-entangled wavefunction and exact density algebra. *arXiv preprint arXiv:2210.13487*, 2022.
- [WKML22] Alexander B Watson, Tianyu Kong, Allan H MacDonald, and Mitchell Luskin. Bistritzer-macdonald dynamics in twisted bilayer graphene. *arXiv preprint arXiv:2207.13767*, 2022.
- [WL21] Alexander B Watson and Mitchell Luskin. Existence of the first magic angle for the chiral model of bilayer graphene. *Journal of Mathematical Physics*, 62(9):091502, 2021.
- [WL22] Jie Wang and Zhao Liu. Hierarchy of ideal flatbands in chiral twisted multilayer graphene models. *Physical Review Letters*, 128(17):176403, 2022.
- [WZMC21] Jie Wang, Yunqin Zheng, Andrew J Millis, and Jennifer Cano. Chiral approximation to twisted bilayer graphene: Exact intravalley inversion symmetry, nodal structure, and implications for higher magic angles. *Physical Review Research*, 3(2):023155, 2021.
- [ZJF⁺11] Fan Zhang, Jeil Jung, Gregory A Fiete, Qian Niu, and Allan H MacDonald. Spontaneous quantum hall states in chirally stacked few-layer graphene systems. *Physical Review Letters*, 106(15):156801, 2011.

Email address: `mxyang@math.berkeley.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CA 94720, USA.