Comments on flavor symmetry breaking and three-dimensional superconformal index

Ilmar Gahramanov

¹ Department of Physics, Bogazici University, 34342 Bebek, Istanbul, Turkey

²Steklov Mathematical Institute of Russ. Acad. Sci., Gubkina str. 8, 119991 Moscow, Russia

³ Department of Mathematics, Khazar University, Mehseti St. 41, AZ1096, Baku, Azerbaijan

⁴ Institute of Radiation Problems ANAS, B. Vahabzade 9, AZ1143 Baku, Azerbaijan

Abstract

We investigate the superconformal index of a three-dimensional SU(2) $\mathcal{N} = 2$ supersymmetric QCD with $N_f = 4$ flavors. This theory confines with the breaking of a global symmetry. We obtain the index of this theory by integrating out a flavor from the theory with $N_f = 6$ flavors. The superconformal index of the theory vanishes for generic values of the flavor fugacities. However, the specific choice of fugacities allows us to describe the flavor symmetry-breaking phenomenon.

1 Introduction

Supersymmetric duality relates the infrared limits of different supersymmetric gauge theories. The basic example of such duality [1] is a four-dimensional $SU(N_c)$ "electric" gauge theory with N_f flavors of quarks in fundamental and antifundamental representations which possesses a dual description in terms of N_f "magnetic" flavors of quarks charged under $SU(N_f - N_c)$ gauge group. For $N_f = N_c$, the infrared theory is confined, and the classical constraint on the meson and baryon fields is modified by one instanton effect, leading to chiral symmetry breaking. In case $N_c = 2$ the global symmetry SU(4) is broken¹ to its maximal vector-like subgroup SP(4) by the chiral condensate [6]. The chiral symmetry breaking for four-dimensional $\mathcal{N} = 1$ supersymmetric gauge theories was considered on the level of the superconformal indices in [7] by Spiridonov and Vartanov.

Three-dimensional $\mathcal{N} = 2$ supersymmetric gauge theories have rich structures [8, 9, 10, 11, 12], such as duality, confinement, and chiral symmetry breaking, similar to four-dimensional $\mathcal{N} = 1$ supersymmetric gauge theories². In this work, we consider the flavor symmetry-breaking phenomenon for the three-dimensional SU(2) $\mathcal{N} = 2$ theory with $N_f = 4$ flavors by using the superconformal index technique. Similar to the case of the three-dimensional S^3 sphere partition function case [7] the superconformal index of such a theory involves the Dirac delta functions which reflect the presence of symmetry breaking

$$\sum_{m\in\mathbb{Z}} \oint \frac{(q^{1+(n_{\alpha}\pm n_{w}\pm m)/2}\alpha^{-1}w^{\mp}z^{\mp})_{\infty}}{(q^{n_{\alpha}\pm n_{w}\pm m}\alpha w^{\pm}z^{\pm})_{\infty}} \frac{(q^{1+(-n_{\alpha}\pm n_{y}\pm m)/2}\alpha y^{\mp}z^{\mp})_{\infty}}{(q^{-n_{\alpha}\pm n_{y}\pm m}\alpha^{-}y^{\pm}z^{\pm})_{\infty}} \frac{(1-q^{m}z^{\pm 2})}{q^{m}} \frac{dz}{4\pi i z}$$
$$= \frac{(q^{1+n_{\alpha}}\alpha^{\mp 2};q)_{\infty}}{(q^{n_{\alpha}}\alpha^{\pm 2};q)_{\infty}} \frac{(q^{1\pm n_{y}}w^{\mp 2};q)_{\infty}}{(q^{\pm n_{w}}w^{\pm 2};q)_{\infty}} \left(\delta(\theta+\chi)\delta_{n_{w}+n_{y},0} + \delta(\theta-\chi)\delta_{n_{w}-n_{y},0}\right) , \qquad (1)$$

where we use the usual q-Pochhammer symbol $(z;q)_{\infty} = \prod_{i=0}^{\infty}(1-zq^i)$ and shorthand notations $f(z^{\pm}) = f(z)f(\frac{1}{z})$. On the left-hand side of equality (1) we have the three-dimensional $\mathcal{N} = 2$ superconformal index of a theory with SU(2) gauge group and $N_f = 4$ chiral fields with the naive flavor group $SU(2) \times SU(2)$. However, the true flavor group is $(SU(2) \times SU(2))_{diag}$ and the index has non-zero support only on the corresponding subset of flavor fugacities. This is precisely the manifestation of chiral symmetry breaking in confining theories similar to the three-dimensional squashed-sphere partition functions case [7]. Note that the expression (1) plays a role of the local inversion relation in integrable lattice models [16, 17, 18].

One can consider the derivation of (1) as another algebraic approach to the inversion relation.

¹The same happens for the two colors non-supersymmetric QCD with the two flavors which enjoys the extended chiral symmetry SU(4) instead of $SU(2) \times SU(2)$, see e.g. [2, 3], see, e.g. [4, 5] for the three-dimensional case.

²The three-dimensional $\mathcal{N} = 2$ theory has the same number of supercharges as a four-dimensional $\mathcal{N} = 1$ theory. In fact, the three-dimensional $\mathcal{N} = 2$ theories discussed in this work are the dimensional reduction of the corresponding four-dimensional $\mathcal{N} = 1$ theories [7, 13, 14, 15].

1.1 Superconformal index

We consider supersymmetric index of a three-dimensional $\mathcal{N} = 2$ superconformal field theory³ which can be defined as a partition function on $S^2 \times S^1$ with the periodic boundary condition of the fields along S^1 [21, 22, 23, 24]

$$I(q, \{a_i\}) = \operatorname{Tr}\left[(-1)^{\mathrm{F}} e^{-\beta \{Q, Q^{\dagger}\}} q^{\frac{1}{2}(\Delta + j_3)} \prod_i a_i^{F_i} \right] , \qquad (2)$$

where the trace is taken over the Hilbert space of the theory on S^2 . The Q is a certain supersymmetric charge in three-dimensional $\mathcal{N} = 2$ superconformal algebra which satisfies the anti-commutation relation $\{Q, Q^{\dagger}\} = \Delta - R - j_3$, where Δ is the energy, j_3 is the third component of the angular momentum on S^2 , R is the R-charge. The operator F plays a role of the fermion number, F_i is the charge of global symmetry commuting with supercharge Q, and a_i is the corresponding fugacity.

The superconformal index can be computed exactly by the supersymmetric localization technique [25] and it takes the form of the following matrix integral⁴

$$I(q, \{a_j\}, \{n_j\}) = \sum_{m_i} \frac{1}{|W_m|} \oint \prod_{i=1}^{\operatorname{rank}G} \frac{dz_i}{2\pi i z_i} Z_{\operatorname{gauge}}(z_i, m_i; q) Z_{\operatorname{chiral}}(z_i, m_i; a_j, n_j; q) .$$
(3)

Here Z_{gauge} and Z_{chiral} stand for the contributions from the vector and chiral multiplets, respectively, $|W_m|$ is the dimension of the Weyl group, the sum is over monopole configurations, and the integral is over gauge fugacities. The form of the index is completely determined if one knows group-theoretical data, i.e. symmetries and corresponding representations of a theory. The reader interested in the precise formulation of the three-dimensional superconformal index can refer to the papers [20, 24].

1.2 Duality for 3d SU(2) $\mathcal{N} = 2$ SQCD with $N_f = 6$ flavor

As an example, we will be studying the global symmetry breaking in three-dimensional $\mathcal{N} = 2$ supersymmetric QCD with SU(2) gauge group and $N_f = 4$ flavors of quarks. To obtain such theory one can take first the *s*-confining theory⁵ with the gauge group SU(2) and $N_f = 6$ flavors. In the case of SU(2) gauge group quarks and antiquarks transform in the same way under the action of the gauge group, i.e. the fundamental and antifundamental representations are equivalent, and

³For more detailed introduction, see e.g. [16, 19, 20].

⁴We skip the contribution of the Chern–Simons term, since in this work we consider theories without this term.

⁵Three-dimensional s-confining theories were discussed in several works, e.g. [26, 27, 28, 29] and the superconformal indices for such theories were presented in [20, 30].

therefore the flavor group enhanced to SU(6) symmetry. The superconformal index of the theory can be expressed in the following form

$$I_E = \sum_{m=-\infty}^{\infty} \oint \prod_{i=1}^{6} \frac{(q^{1+(m+n_i)/2}/a_i z, q^{1+(n_i-m)/2} z/a_i; q)_{\infty}}{(q^{(m+n_i)/2} a_i z, q^{(n_i-m)/2} a_i/z; q)_{\infty}} \frac{(1-q^m z^2)(1-q^m z^{-2})}{q^m z^{6m}} \frac{dz}{4\pi i z}$$
(4)

with the balancing conditions

$$\prod_{i=1}^{6} a_i = q \text{ and } \sum_{i=1}^{6} n_i = 0.$$
(5)

The variables a_i and z are the fugacities associated with the SU(6) flavor group and SU(2) gauge group, respectively. In [31], it was shown that the superconformal index of this theory has an extended SO(12) flavor symmetry when coupled to four-dimensional multiplets with specific boundary conditions.

It is known that there is magnetic dual theory without gauge degrees of freedom and with fifteen chiral multiplets in the totally antisymmetric tensor representation of the same flavor group. The superconformal index of the magnetic theory is

$$I_M = \frac{1}{\prod_{i=1}^6 a_i^{n_i}} \prod_{1 \le i < j \le 6} \frac{(q^{1+(n_i+n_j)/2}/a_i a_j; q)_{\infty}}{(q^{(n_i+n_j)/2}a_i a_j; q)_{\infty}}.$$
 (6)

Since these theories are IR dual, the superconformal indices need to be in agreement [20, 32]. The result is given by the following q-hypergeometric identity

$$\sum_{m=-\infty}^{\infty} \oint \prod_{i=1}^{6} \frac{(q^{1+(m+n_i)/2}/a_i z, q^{1+(n_i-m)/2} z/a_i; q)_{\infty}}{(q^{(m+n_i)/2} a_i z, q^{(n_i-m)/2} a_i/z; q)_{\infty}} \frac{(1-q^m z^{\pm 2})}{q^m z^{6m}} \frac{dz}{4\pi i z}$$
$$= \frac{1}{\prod_{i=1}^{6} a_i^{n_i}} \prod_{1 \le i < j \le 6} \frac{(q^{1+(n_i+n_j)/2}/a_i a_j; q)_{\infty}}{(q^{(n_i+n_j)/2} a_i a_j; q)_{\infty}}$$
(7)

with the balancing conditions (5). This integral identity was proven in [20] and studied in [16, 20, 33].

The identity (7) can be written as a star-triangle relation (the non-operator form of the Yang-Baxter equation) [16, 18] which is a sufficient condition for integrability for the Ising-type lattice spin models. The Bailey pairs and the operator form of the Yang-Baxter equation related to this integral identity also were constructed in [16] (see, also [34]).

Note that the basic hypergeometric integral identity (7) is new in the literature and needs detailed study and classification [35]. Such identities (containing summation and integration) recently appear in supersymmetric gauge theory computations [14, 19, 20, 33, 36, 37, 38, 39, 40, 41] and integrable models of statistical mechanics [16, 34, 42, 43, 44, 45].

2 Breaking of flavor symmetry

Our goal will be to reduce the number of flavors from $N_f = 6$ to $N_f = 4$ case for the integral identity (7). One can perform the reduction of superconformal indices (7) by taking the following limit⁶

$$a_5 a_6 = q, \quad n_5 + n_6 = 0 \tag{8}$$

Generically this limit gives zero on both sides, however for special choices of fugacities a_i one can deal with such integrals. Let us denote

$$a_1 = \alpha w, \quad a_2 = \alpha w^{-1}, \quad a_3 = \beta y, \quad a_4 = \beta y^{-1},$$
(9)

and the corresponding discrete parameters

$$n_1 = n_\alpha + n_w, \quad n_2 = n_\alpha - n_w, \quad n_3 = n_\beta + n_y, \quad n_4 = n_\beta - n_y.$$
 (10)

In terms of new variables, the balancing conditions read as

$$a_5 a_6 \alpha^2 \beta^2 = q$$
, $2n_\alpha + 2n_\beta + n_5 + n_6 = 0$. (11)

To proceed further we need to consider the following integral: multiply the integral (4) by a holomorphic function near the unit circle f(y) and integrate over the variable y. Then using the integral identity (7) we obtain the following expression⁷

$$\int_{\mathbb{T}} f(y) I_E \frac{dy}{2\pi i y} = \alpha^{-2n_{\alpha}} w^{-2n_{\beta}} a_5^{-n_5} a_6^{-n_6} \frac{(q^{1+n_{\alpha}} \alpha^{-2}; q)_{\infty}}{(q^{n_{\alpha}} \alpha^2; q)_{\infty}} \frac{(q^{1+n_{\beta}} \beta^{-2}; q)_{\infty}}{(q^{n_{\beta}} \beta^2; q)_{\infty}} \\
\times \frac{(q^{1+\frac{n_5+n_6}{2}} (a_5 a_6)^{-1}; q)_{\infty}}{(q^{\frac{n_5+n_6}{2}} a_5 a_6; q)_{\infty}} \frac{(q^{1+\frac{n_{\alpha}\pm n_w+n_5}{2}} (\alpha w^{\pm 1} a_5)^{-1}; q)_{\infty}}{(q^{\frac{n_{\alpha}\pm n_w+n_5}} \alpha w^{\pm 1} a_5; q)_{\infty}} \frac{(q^{1+\frac{n_{\alpha}\pm n_w+n_6}{2}} (\alpha w^{\pm 1} a_6)^{-1}; q)_{\infty}}{(q^{\frac{n_{\alpha}\pm n_w+n_5}} \alpha w^{\pm 1} a_5; q)_{\infty}} \frac{(q^{1+\frac{n_{\alpha}\pm n_w+n_6}{2}} \alpha w^{\pm 1} a_6; q)_{\infty}}{(q^{\frac{n_{\alpha}\pm n_w+n_5}} \alpha w^{\pm 1} w^{\pm 1})^{-1}; q)_{\infty}} \\
\times \int_{\mathbb{T}} \frac{dy}{2\pi i y} f(y) y^{-2n_y} \frac{(q^{1+\frac{n_{\alpha}+n_{\beta}\pm n_w\pm n_y}{2}} (\alpha \beta w^{\pm 1} y^{\pm 1})^{-1}; q)_{\infty}}{(q^{\frac{n_{\alpha}+n_{\beta}\pm n_w\pm n_y}} \alpha \beta w^{\pm 1} y^{\pm 1}; q)_{\infty}} \prod_{j=5}^{6} \frac{(q^{1+\frac{n_{\beta}\pm n_y+n_j}{2}} (\beta y^{\pm 1} a_j)^{-1}; q)_{\infty}}{(q^{\frac{n_{\beta}\pm n_y+n_j}} \beta y^{\pm 1} a_j; q)_{\infty}}.$$
(12)

The first term in the integrand of (12) has the following sequences of poles

$$y_{in} = \alpha \beta w^{\pm 1} q^{\frac{n_\alpha + n_\beta \pm n_w - n_y}{2}} q^j, \tag{13}$$

$$y_{out} = \frac{1}{\alpha\beta} w^{\pm 1} q^{\frac{-n_{\alpha} - n_{\beta} \pm n_w - n_y}{2}} q^{-j}, \tag{14}$$

⁶Note that in work [30], the authors take another limit $a_5a_6 = q^{1/2}$ and obtain the s-confining theory without breaking SU(4) flavor symmetry. In [16], authors considered superconformal indices of dual theories with SU(4) flavor symmetry without superpotential term (without balancing condition).

⁷Here we assume that $\alpha^2 \neq 1$.

where $i, j \in \mathbb{Z}_{\geq 0}$ and the *in*-poles converge to zero y = 0 and *out*-poles go to infinity. In order to calculate the integral (12) we deform the counter \mathbb{T} such that only the residues of the in-poles are picked up and there are no singularities lying on the counter C. Then the residue calculation leads to the result

$$\int_{\mathbb{T}} f(y) I_E \frac{dy}{2\pi i y} = \alpha^{-2n_{\alpha}} w^{-2n_{w}} \beta^{-2n_{\beta}} a_5^{-n_5} a_6^{-n_6} \frac{(q^{1+n_{\alpha}} \alpha^{-2}; q)_{\infty}}{(q^{n_{\alpha}} \alpha^2; q)_{\infty}} \frac{(q^{1+n_{\beta}} \beta^{-2}; q)_{\infty}}{(q^{n_{\beta}} \beta^2; q)_{\infty}} \times \frac{(q^{1+\frac{n_5+n_6}{2}} (a_5 a_6)^{-1}; q)_{\infty}}{(q^{\frac{n_5+n_6}{2}} a_5 a_6; q)_{\infty}} \prod_{j=5}^{6} \frac{(q^{1+\frac{n_{\alpha}\pm n_{w}+n_j}{2}} (\alpha w^{\pm 1} a_j)^{-1}; q)_{\infty}}{(q^{\frac{n_{\alpha}\pm n_{w}+n_j}{2}} \alpha w^{\pm 1} a_j; q)_{\infty}} \left(I(w) + I(w^{-1}) + I(C)\right)$$
(15)

where

•

$$I(w) = f\left(\alpha\beta w q^{\frac{n\alpha+n_{\beta}+n_{w}-n_{y}}{2}}\right) \left(\alpha\beta w q^{\frac{n\alpha+n_{\beta}+n_{w}-n_{y}}{2}}\right)^{-2n_{y}} \frac{(q^{1+n_{y}} (\alpha\beta w)^{-2}; q)_{\infty}}{(q^{n\alpha+n_{\beta}+n_{w}} (\alpha\beta w)^{2}; q)_{\infty}} \\ \times \frac{(q^{1+n_{\alpha}+n_{\beta}+n_{w}-n_{y}}; q)_{\infty}}{(q; q)_{\infty}} \frac{(q^{1-n_{w}+n_{y}} (\alpha\beta)^{-2}; q)_{\infty}}{(q^{n_{\alpha}+n_{\beta}} (\alpha\beta)^{2}; q)_{\infty}} \frac{(q^{1+n_{\alpha}+n_{\beta}-n_{y}} w^{2}; q)_{\infty}}{(q^{-n_{w}} w^{-2}; q)_{\infty}} \\ \times \prod_{i=5}^{6} \frac{(q^{1+\frac{-n_{\alpha}+2n_{y}+n_{i}}{2}} (\alpha\beta^{2}wa_{i})^{-1}; q)_{\infty}}{(q^{\frac{2n_{\beta}+n_{\alpha}+n_{w}+n_{i}}{2}} \alpha\beta^{2}wa_{i}; q)_{\infty}} \frac{(q^{1+\frac{2n_{\beta}+n_{\alpha}+n_{i}+n_{w}-2n_{y}}{2}} (\alpha w^{-1}a_{i})^{-1}; q)_{\infty}}{(q^{\frac{-n_{w}-n_{\alpha}+n_{i}}{2}} \alpha w^{-1}a_{i}; q)_{\infty}}$$
(16)

$$I(w^{-1}) = f\left(\alpha\beta w^{-1}q^{\frac{n\alpha+n_{\beta}-nw-n_{y}}{2}}\right) (\alpha\beta w^{-1}q^{\frac{n\alpha+n_{\beta}-nw-n_{y}}{2}})^{-2n_{y}} \frac{(q^{1+n_{w}+n_{y}}(\alpha\beta)^{-2};q)_{\infty}}{(q^{n_{\alpha}+n_{\beta}}(\alpha\beta)^{2};q)_{\infty}} \\ \times \frac{(q^{1+n_{\alpha}+n_{\beta}-n_{y}}w^{-2};q)_{\infty}}{(q^{n_{w}}w^{2};q)_{\infty}} \frac{(q^{1+n_{y}}(\alpha\beta w^{-1})^{-2};q)_{\infty}}{(q^{n_{\alpha}+n_{\beta}-n_{w}}(\alpha\beta w^{-1})^{2};q)_{\infty}} \frac{(q^{1+n_{\alpha}+n_{\beta}-n_{w}-n_{y}};q)_{\infty}}{(q;q)_{\infty}} \\ \times \prod_{i=5}^{6} \frac{(q^{1+\frac{-n_{\alpha}+2n_{y}+n_{i}}{2}}(\alpha\beta^{2}w^{-1}a_{i})^{-1};q)_{\infty}}{(q^{\frac{2n_{\beta}+n_{\alpha}-n_{w}+n_{i}}{2}}\alpha\beta^{2}w^{-1}a_{i};q)_{\infty}} \frac{(q^{1+\frac{2n_{\beta}+n_{\alpha}+n_{i}}{2}}(\alphawa_{i})^{-1};q)_{\infty}}{(q^{\frac{nw-n_{\alpha}+n_{i}}{2}}\alphawa_{i};q)_{\infty}}$$
(17)

$$I(C) = \int_{\mathcal{C}} \frac{dy}{2\pi i y} f(y) y^{-2n_y} \frac{\left(q^{1+\frac{n_\alpha+n_\beta\pm n_w\pm n_y}{2}}(\alpha\beta w^{\pm 1}y^{\pm 1})^{-1};q\right)_{\infty}}{\left(q^{\frac{n_\alpha+n_\beta\pm n_w\pm n_y}{2}}\alpha\beta w^{\pm 1}y^{\pm 1};q\right)_{\infty}} \prod_{i=5}^{6} \frac{\left(q^{1+\frac{n_\beta\pm n_y+n_i}{2}}(\beta y^{\pm 1}a_i)^{-1};q\right)_{\infty}}{\left(q^{\frac{n_\beta\pm n_y+n_i}{2}}\beta y^{\pm 1}a_i;q\right)_{\infty}}$$
(18)

In the limit $\alpha\beta \to 1$ and $n_{\alpha} + n_{\beta} \to 0$, the poles at $\alpha\beta w^{\pm 1}q^{\pm n_w - n_y}$ and $w^{\pm 1}q^{\pm n_w - n_y}/\alpha\beta$ start to pinch \mathbb{T} . The integral (18) vanishes in this limit. As a result, we obtain⁸

$$\int_{\mathbb{T}} f(y) I_E \frac{dy}{2\pi i y} = \frac{(q^{1+n_\alpha} \alpha^{\mp 2}; q)_\infty}{(q^{n_\alpha} \alpha^{\pm 2}; q)_\infty} \frac{(q^{1\pm n_y} w^{\mp 2}; q)_\infty}{(q^{\pm n_w} w^{\pm 2}; q)_\infty} \times \left(f(wq^{(n_w - n_y)/2}) \delta_{n_w + n_y, 0} + f(w^{-1}q^{(-n_w - n_y)/2}) \delta_{n_w - n_y, 0} \right)$$
(20)

⁸In order to cancel out many terms we used the equality [16, 46]

$$\frac{(q^{1-m/2}z^{-1};q)_{\infty}}{(q^{-m/2}z;q)_{\infty}} = \frac{q^{m/2}}{(-z)^m} \frac{(q^{1+m/2}z^{-1};q)_{\infty}}{(q^{+m/2}z;q)_{\infty}}, \quad m \in \mathbb{Z}.$$
(19)

Here $\delta_{i,j}$ is the Kronecker delta. Now we denote $y = e^{2\pi i\theta}$, $w = e^{2\pi i\chi}$ and pass to the integration over real variable θ . Because of the arbitrariness of the function f(y), we can give to the function I_E a distributional sense and rewrite the expression (20) in the following form

$$\sum_{m\in\mathbb{Z}} \oint \frac{(q^{1+(n_{\alpha}\pm n_{w}\pm m)/2}\alpha^{-1}w^{\mp}z^{\mp})_{\infty}}{(q^{n_{\alpha}\pm n_{w}\pm m}\alpha w^{\pm}z^{\pm})_{\infty}} \frac{(q^{1+(-n_{\alpha}\pm n_{y}\pm m)/2}\alpha y^{\mp}z^{\mp})_{\infty}}{(q^{-n_{\alpha}\pm n_{y}\pm m}\alpha^{-}y^{\pm}z^{\pm})_{\infty}} \frac{(1-q^{m}z^{2})(1-q^{m}z^{-2})}{q^{m}} \frac{dz}{4\pi i z}$$
$$= \frac{(q^{1+n_{\alpha}}\alpha^{\mp 2};q)_{\infty}}{(q^{n_{\alpha}}\alpha^{\pm 2};q)_{\infty}} \frac{(q^{1\pm n_{y}}w^{\mp 2};q)_{\infty}}{(q^{\pm n_{w}}w^{\pm 2};q)_{\infty}} \left(\delta(\theta+\chi)\delta_{n_{w}+n_{y},0} + \delta(\theta-\chi)\delta_{n_{w}-n_{y},0}\right) \quad (21)$$

where $\delta(\theta)$ is the periodic Dirac delta function, $\delta(\phi + 1) = \delta(\phi)$. The appearance of the Dirac delta functions describes the symmetry breaking of the flavor group to the $(SU(2) \times SU(2))_{diag}$.

The equality (21) has an interpretation as the inversion relation⁹ for exactly solvable lattice spin model introduced in [16, 18]. In particular, this observation can be used [51] for a better understanding of the correspondence [42, 52, 53, 54] between supersymmetric gauge theories and integrable models of statistical mechanics.

3 Concluding remarks

We have shown that it is possible to obtain a three-dimensional $\mathcal{N} = 2$ s-confining theory with flavor symmetry breaking by analyzing the superconformal index of the theory. There are several potential directions to extend our work. It would be interesting to obtain the superconformal index with delta functions via localization technique if it is possible.

Though studied only SU(2) supersymmetric gauge theory with $N_f = 4$ flavors the approach discussed here also applies to gauge theories with other flavor groups. We would expect analogous behavior of the superconformal index for the general SU(N) theory, however, it is computationally complicated to work with the multivariate analog of the basic hypergeometric integral identity (7).

The four-dimensional s-confining theories are classified [55, 56, 57]. The chiral symmetry is always broken in such theories if one reduces flavor symmetries to $N_f = N_c$. The corresponding threedimensional analysis [26, 28, 58, 59] for such theories need to be performed.

Finally, there are many integrable lattice models [17, 60, 61, 62, 63, 64] obtained recently via exact results from supersymmetric gauge theories. A similar approach can be used for obtaining corresponding inversion relations. We hope to elaborate on this as well as on other topics elsewhere.

⁹Inversion relation is a simple relation as a star-triangle relation within the framework of integrable lattice spin models of statistical mechanics, see, e.g. [47, 48, 49, 50]. In the context of S-matrix for particle physics in 1 + 1 dimensions it has the meaning of the unitarity relation.

Acknowledgements

The work originated from stimulating discussions with Vyacheslav Spiridonov, I thank him for his help, suggestions, and contributions to the ideas presented here. I would like to thank Keita Nii, Mustafa Mullahasanoglu, Erdal Catak, Yasin Şale, Merve Kaftan, and Yağız Eren Kiremit for the discussions. This work has been supported by the Russian Science Foundation grant number 22-72-10122 2 (https://rscf.ru/en/project/22-72-10122). I am grateful to the Institut des Hautes Etudes Scientifiques, IHES (Bures-sur-Yvette, France), for the warm hospitality, where part of this work was carried out. The visit to IHES was supported by Nokia Fund in 2023.

References

- N. Seiberg, "Electric magnetic duality in supersymmetric nonAbelian gauge theories," Nucl. Phys. B435 (1995) 129-146, arXiv:hep-th/9411149 [hep-th].
- [2] A. V. Smilga and J. J. M. Verbaarschot, "Spectral sum rules and finite volume partition function in gauge theories with real and pseudoreal fermions," *Phys. Rev.* D51 (1995) 829-837, arXiv:hep-th/9404031 [hep-th].
- J. B. Kogut, M. A. Stephanov, and D. Toublan, "On two color QCD with baryon chemical potential," *Phys. Lett.* B464 (1999) 183-191, arXiv:hep-ph/9906346 [hep-ph].
- [4] U. Magnea, "The Orthogonal ensemble of random matrices and QCD in three-dimensions," *Phys. Rev. D* 61 (2000) 056005, arXiv:hep-th/9907096.
- [5] G. V. Dunne and S. M. Nishigaki, "Two color QCD in 3-D at finite baryon density," Nucl. Phys. B 654 (2003) 445-465, arXiv:hep-ph/0210219.
- [6] N. Seiberg, "Exact results on the space of vacua of four-dimensional SUSY gauge theories," *Phys. Rev. D* 49 (1994) 6857–6863, arXiv:hep-th/9402044.
- [7] V. P. Spiridonov and G. S. Vartanov, "Vanishing superconformal indices and the chiral symmetry breaking," JHEP 06 (2014) 062, arXiv:1402.2312 [hep-th].
- [8] K. A. Intriligator and N. Seiberg, "Mirror symmetry in three-dimensional gauge theories," *Phys. Lett.* B387 (1996) 513-519, arXiv:hep-th/9607207 [hep-th].
- [9] J. de Boer, K. Hori, H. Ooguri, and Y. Oz, "Mirror symmetry in three-dimensional gauge theories, quivers and D-branes," Nucl. Phys. B 493 (1997) 101-147, arXiv:hep-th/9611063.

- [10] O. Aharony, A. Hanany, K. A. Intriligator, N. Seiberg, and M. J. Strassler, "Aspects of N=2 supersymmetric gauge theories in three-dimensions," *Nucl. Phys.* B499 (1997) 67–99, arXiv:hep-th/9703110 [hep-th].
- [11] J. de Boer, K. Hori, and Y. Oz, "Dynamics of N=2 supersymmetric gauge theories in three-dimensions," Nucl. Phys. B 500 (1997) 163-191, arXiv:hep-th/9703100.
- [12] A. Amariti, D. Orlando, and S. Reffert, "String theory and the 4D/3D reduction of Seiberg duality. A review," *Phys. Rept.* **705-706** (2017) 1–53, arXiv:1611.04883 [hep-th].
- [13] F. A. H. Dolan, V. P. Spiridonov, and G. S. Vartanov, "From 4d superconformal indices to 3d partition functions," *Phys. Lett.* B704 (2011) 234-241, arXiv:1104.1787 [hep-th].
- [14] O. Aharony, S. S. Razamat, N. Seiberg, and B. Willett, "3d dualities from 4d dualities," JHEP 07 (2013) 149, arXiv:1305.3924 [hep-th].
- [15] A. Amariti, "4d/3d reduction of s-confining theories: the role of the "exotic" D instantons," JHEP 02 (2016) 139, arXiv:1507.05623 [hep-th].
- [16] I. Gahramanov and V. P. Spiridonov, "The star-triangle relation and 3d superconformal indices," JHEP 08 (2015) 040, arXiv:1505.00765 [hep-th].
- [17] I. Gahramanov and A. P. Kels, "The star-triangle relation, lens partition function, and hypergeometric sum/integrals," arXiv:1610.09229 [math-ph].
- [18] A. P. Kels, "New solutions of the star triangle relation with discrete and continuous spin variables," J. Phys. A48 no. 43, (2015) 435201, arXiv:1504.07074 [math-ph].
- [19] C. Krattenthaler, V. P. Spiridonov, and G. S. Vartanov, "Superconformal indices of three-dimensional theories related by mirror symmetry," *JHEP* 06 (2011) 008, arXiv:1103.4075 [hep-th].
- [20] I. Gahramanov and H. Rosengren, "Basic hypergeometry of supersymmetric dualities," Nucl. Phys. B913 (2016) 747-768, arXiv:1606.08185 [hep-th].
- [21] J. Bhattacharya, S. Bhattacharyya, S. Minwalla, and S. Raju, "Indices for Superconformal Field Theories in 3,5 and 6 Dimensions," JHEP 02 (2008) 064, arXiv:0801.1435 [hep-th].
- [22] J. Bhattacharya and S. Minwalla, "Superconformal Indices for N = 6 Chern Simons Theories," JHEP 01 (2009) 014, arXiv:0806.3251 [hep-th].

- [23] S. Kim, "The Complete superconformal index for N=6 Chern-Simons theory," Nucl. Phys. B 821 (2009) 241-284, arXiv:0903.4172 [hep-th]. [Erratum: Nucl.Phys.B 864, 884 (2012)].
- [24] Y. Imamura and S. Yokoyama, "Index for three dimensional superconformal field theories with general R-charge assignments," JHEP 04 (2011) 007, arXiv:1101.0557 [hep-th].
- [25] B. Willett, "Localization on three-dimensional manifolds,"
 J. Phys. A50 no. 44, (2017) 443006, arXiv:1608.02958 [hep-th].
- [26] C. Csáki, M. Martone, Y. Shirman, P. Tanedo, and J. Terning, "Dynamics of 3D SUSY Gauge Theories with Antisymmetric Matter," JHEP 08 (2014) 141, arXiv:1406.6684 [hep-th].
- [27] K. Nii, "3d s-confinement for three-index matters," JHEP 11 (2018) 099, arXiv:1805.06369 [hep-th].
- [28] K. Nii, "On s-confinement in 3d $\mathcal{N} = 2$ gauge theories with anti-symmetric tensors," arXiv:1906.03908 [hep-th].
- [29] A. Amariti and S. Rota, "3d N=2 SO/USp adjoint SQCD: s-confinement and exact identities," Nucl. Phys. B 987 (2023) 116068, arXiv:2202.06885 [hep-th].
- [30] I. Gahramanov and G. Vartanov, "Extended global symmetries for 4D N = 1 SQCD theories," J. Phys. A46 (2013) 285403, arXiv:1303.1443 [hep-th].
- [31] T. Dimofte and D. Gaiotto, "An E7 Surprise," JHEP 10 (2012) 129, arXiv:1209.1404 [hep-th].
- [32] V. Spiridonov and G. Vartanov, "Superconformal indices for N = 1 theories with multiple duals," Nucl. Phys. B 824 (2010) 192-216, arXiv:0811.1909 [hep-th].
- [33] H. Rosengren, "Rahman's biorthogonal functions and superconformal indices," arXiv:1612.05051 [math.CA].
- [34] E. Catak, "A remark on the q-hypergeometric integral Bailey pair and the solution to the star-triangle equation," arXiv:2212.13528 [math.CA].
- [35] M. J. Schlosser, "Hypergeometric and basic hypergeometric series and integrals associated with root systems," arXiv:1705.09221 [math.CA].
- [36] A. Kapustin and B. Willett, "Generalized Superconformal Index for Three Dimensional Field Theories," arXiv:1106.2484 [hep-th].

- [37] I. Gahramanov and H. Rosengren, "A new pentagon identity for the tetrahedron index," JHEP 11 (2013) 128, arXiv:1309.2195 [hep-th].
- [38] I. Gahramanov and H. Rosengren, "Integral pentagon relations for 3d superconformal indices," Proc. Symp. Pure Math. 93, 165 (2016), arXiv:1412.2926 [hep-th].
- [39] I. Gahramanov, "Mathematical structures behind supersymmetric dualities," Archivum Math. 51 (2015) 273-286, arXiv:1505.05656 [math-ph].
- [40] I. Gahramanov and O. E. Kaluc, "Bailey pairs for the q-hypergeometric integral pentagon identity," arXiv:2111.14793 [math-ph].
- [41] D. N. Bozkurt and I. Gahramanov, "Pentagon identities arising in supersymmetric gauge theory computations," *Teor. Mat. Fiz.* 198 no. 2, (2019) 215-224, arXiv:1803.00855 [math-ph].
- [42] I. Gahramanov and S. Jafarzade, "Integrable lattice spin models from supersymmetric dualities," *Phys. Part. Nucl. Lett.* **15** no. 6, (2018) 650–667, arXiv:1712.09651 [math-ph].
- [43] I. Gahramanov and S. Jafarzade, "Comments on the multi-spin solution to the Yang-Baxter equation and basic hypergeometric sum/integral identity," *Mod. Phys. Lett. A* 34 no. 18, (2019) 1950140, arXiv:1710.09106 [math-ph].
- [44] E. Eren, I. Gahramanov, S. Jafarzade, and G. Mogol, "Gamma function solutions to the star-triangle equation," *Nucl. Phys. B* 963 (2021) 115283, arXiv:1912.12271 [math-ph].
- [45] E. Catak, I. Gahramanov, and M. Mullahasanoglu, "Hyperbolic and trigonometric hypergeometric solutions to the star-star equation," *Eur. Phys. J. C* 82 no. 9, (2022) 789, arXiv:2107.06880 [hep-th].
- [46] T. Dimofte, D. Gaiotto, and S. Gukov, "3-Manifolds and 3d Indices," Adv. Theor. Math. Phys. 17 no. 5, (2013) 975-1076, arXiv:1112.5179 [hep-th].
- [47] Yu. G. Stroganov, "A new calculation method for partition functions in some lattice models," *Phys. Lett.* A74 (1979) 116–118.
- [48] R. J. Baxter, "The inversion relation method for some two-dimensional exactly solved models in lattice statistics," J. Statist. Phys. 28 (1982) 1–41.
- [49] M. T. Jaekel and J. M. Maillard, "INVERSE FUNCTIONAL RELATIONS FOR LATTICE MODELS," J. Phys. A 16 (1983) 1975.

- [50] J. H. H. Perk and F. Y. Wu, "Graphical Approach to the Nonintersecting String Model: Star Triangle Equation, Inversion Relation and Exact Solution," *Physica A* 138 (1986) 100–124.
- [51] I. Gahramanov, "Integrability from supersymmetric duality: a short review," arXiv:2201.00351 [hep-th].
- [52] V. P. Spiridonov, "Elliptic beta integrals and solvable models of statistical mechanics," Contemp. Math. 563 (2012) 181-211, arXiv:1011.3798 [hep-th].
- [53] M. Yamazaki, "Integrability As Duality: The Gauge/YBE Correspondence," *Phys. Rept.* 859 (2020) 1–20, arXiv:1808.04374 [hep-th].
- [54] J. Yagi, "Quiver gauge theories and integrable lattice models," JHEP 10 (2015) 065, arXiv:1504.04055 [hep-th].
- [55] C. Csaki, M. Schmaltz, and W. Skiba, "A Systematic approach to confinement in N=1 supersymmetric gauge theories," *Phys. Rev. Lett.* 78 (1997) 799-802, arXiv:hep-th/9610139.
- [56] C. Csaki, "The Confining N=1 supersymmetric gauge theories: A Review," in 3rd Workshop on Continuous Advances in QCD (QCD 98), pp. 489–499. 4, 1998. arXiv:hep-th/9807222.
- [57] C. Csaki and H. Murayama, "New confining N=1 supersymmetric gauge theories," *Phys. Rev. D* 59 (1999) 065001, arXiv:hep-th/9810014.
- [58] K. Nii, "Confinement in 3d $\mathcal{N} = 2$ Spin(N) gauge theories with vector and spinor matters," JHEP 03 (2019) 113, arXiv:1810.06618 [hep-th].
- [59] K. Nii, "Confinement in 3d $\mathcal{N} = 2$ exceptional gauge theories," arXiv:1906.10161 [hep-th].
- [60] A. P. Kels, "A new solution of the star-triangle relation," J. Phys. A47 (2014) 055203, arXiv:1302.3025 [math-ph].
- [61] A. P. Kels and M. Yamazaki, "Lens elliptic gamma function solution of the Yang-Baxter equation at roots of unity," J. Stat. Mech. 1802 no. 2, (2018) 023108, arXiv:1709.07148 [math-ph].
- [62] M. Mullahasanoglu and N. Tas, "Lens Partition Functions and Integrability Properties," arXiv:2112.15161 [hep-th].
- [63] J. de-la Cruz-Moreno and H. García-Compeán, "Star-triangle type relations from 2d $\mathcal{N} = (0,2) USp(2N)$ dualities," JHEP **01** (2021) 023, arXiv:2008.02419 [hep-th].
- [64] M. Dede, "A comment on the solutions of the generalized Faddeev-Volkov model," arXiv:2206.14271 [math-ph].