4D-EGB Black Holes in RPS Thermodynamics

Y. Ladghami^{*}, B. Asfour[†], A. Bouali[‡], A. Errahmani[§], and T. Ouali[¶]

Laboratory of Physics of Matter and Radiation, Mohammed I University, BP 717, Oujda, Morocco

March 22, 2023

Abstract

In this paper, we study thermodynamics of charged and uncharged 4-Dimension Einstein-Gauss-Bonnet (4D-EGB) black holes. The context of this study is the Visser's holographic thermodynamics with a fixed anti-de Sitter radius and a variable Newton constant known as restricted phase space thermodynamics (RPST). Our setup is constructed by using the AdS/CFT correspondence and by introducing a conjugate quantity of the Gauss-Bonnet parameter. By this ansatz, we conclude that the Gauss-Bonnet action multiplied by a temperature, behaves as a free energy. We derive the conjugate quantities corresponding to the first law in the RPST formalism. The study of the T-S processes and the effect of the Gauss-Bonnet constant, α , show that thermodynamic properties of charged black holes depend on the Gauss-Bonnet term and the charge of black holes. For an uncharged black holes, the effect of Gauss-Bonnet becomes crucial, as it behaves as a charged black hole with an effective charge. Finally, we find that the Hawking-Page phase transition occurs between a large black hole and a thermal AdS space.

1 Introduction

Black holes and their thermodynamics provide a fertile ground for research, especially after Hawking and Bekenstein's work on the temperature and the entropy of black holes [1,2], in addition to the four laws of black hole thermodynamics, through the thermodynamic analogy of the ordinary systems [3–5]. This thermodynamic is known as traditional black hole thermodynamics (TBHT), for which, the mass of the black hole is considered as its internal energy. The most important results of TBHT are the Hawking-Page phase transition for the space-time with a negative cosmological constant ($\Lambda < 0$) i.e an anti-de Sitter space-time (AdS) [6]. Another important formalism in the study of black holes is called extended phase space thermodynamics (EPST) started for the first time in [7]. This formalism is also called "Black hole chemistry" [8,9]. A number of works have been published in this regard [10–17]. In the EPST, the mass of black holes is interpreted as an enthalpy and thermodynamic parameters such as the pressure, P, and the volume, V, have been interpenetrated as thermodynamic variables, where the pressure is related to the cosmological constant via $P = -\Lambda/8\pi G$ [8]. AdS/CFT correspondence, speculated by Maldacena [18], has played an important role to understand the behavior of black hole and it is used as a framework to introduce a new thermodynamic approach. In this background, the AdS black hole is equivalent to a thermal state in the conformal field theories (CFT) [19]. Recently, within the framework of the AdS/CFT correspondence and based on the Visser's thermodynamics [20], a new thermodynamic called restricted phase space thermodynamics (RPST) was initiated by Gao et al in [21]. In RPST, black hole masses are reinterpreted as internal energy. However, instead of the pressure and the volume characterizing the EPST formalism, a new pair of conjugate thermodynamic variables characterizing the RPST formalism has been introduced. This new conjugates thermodynamic variables, are the chemical potential (color susceptibility) μ , and the central charge C which

^{*}yahya.ladghami@ump.ac.ma

[†]brahim.asfour@ump.ac.ma

[‡]a1.bouali@ump.ac.ma

[§]ahmederrahmani1@yahoo.fr

[¶]ouali1962@gmail.com

represents the number degrees of freedom (number of colors) in CFT [20,22]. The role of central charge C is similar to that of the number of particles in statistical physics of ordinary matter [23]. The central charge is related to the Newton's constant G and the AdS curvature radius l via $C = l^{d-2}/G$, where d is the dimension of the AdS space-time [21, 22]. Homogeneity of Smarr relation, thermodynamic processes and phase transitions, were studied in the RN-AdS, Kerr-AdS, BTZ and Taub-NUT black holes in the context of RPST [21, 23–26].

Weyl [27] and Eddington [28] first suggested modifications of General Relativity (GR) in 1918, and the idea has stuck around ever since. The current concept aims to develop GR alternatives for some cosmological issues, such as the cosmic acceleration in the early and late universe. One of these alternative theories is the Gauss-Bonnet gravity [29,30], in which an arbitrary function $f(\mathcal{G})$ is included in the gravitational action, where \mathcal{G} is the Gauss-Bonnet (GB) term. However, gravitational dynamics in 4-dimensional spacetime are unaffected by the GB term. For this reason, D. Glavan and C. Lin have recently proposed a novel 4-dimensional Einstein-Gauss-Bonnet (4D-EGB) gravity [31], which is founded on the idea of multiplying the GB term by the factor 1/(d-4) in order to circumvent the strict requirements of Lovelock's theorem [32] and avoid Ostrogradsky instability. Additionally, in recent years, this new approach has developed into a useful framework for researching physics and thermodynamics of black holes. For instance, vacuum solution for charged black holes, thermodynamic properties [33], extended thermodynamics of black holes [34, 35], a charged black hole with entangled particles and antiparticles [36], stable circular orbit and shadow of the black holes [37], power spectra of the Hawking radiation in de-Sitter spacetime [38], stability of black holes [39] and non-linear charged planar black holes [40, 41].

As far as we know, even with the number of applications of 4D-EGB, there is no study so far about the RPST formalism in this approach. This motivates us to consider the RPST formalism in the 4D-EGB gravity background. A second motivation is to study effects of the Gauss-Bonnet constant on thermodynamic behaviors of a charged and uncharged black holes in both approaches and to complete the study of black holes in the RPST formalism already made in [21, 23–26].

The paper is organised as follows: In Section 2, we present the 4D-EGB black holes solution in AdS space-time. In Section 3, we construct an holographic thermodynamics with a fixed AdS radius for 4D-EGB gravity. In section 4, the first law and thermodynamic variables are deduced from RPST. In section 5, we look at thermodynamic properties of the charged 4D-EGB black hole in the RPS thermodynamics by T-S processes. In section 6, we study uncharged 4D-EGB black holes. In section 7, we discuss our main conclusions. In this paper, we adopt the unit $\hbar = c = k_B = 1$.

2 4D-EGB Black Holes in AdS space-time

The action of the Gauss-Bonnet gravity in a d-dimensional space-time with a negative cosmological constant is given by [33]

$$I = \frac{1}{16\pi G} \int d^d x \left(R - 2\Lambda - F_{\mu\nu} F^{\mu\nu} + \frac{\alpha}{d-4} \mathcal{G} \right), \tag{1}$$

where the GB term writes

$$\mathcal{G} = R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} - 4R_{\mu\nu}R^{\mu\nu} + R^2, \qquad (2)$$

 α is Gauss-Bonnet coupling with dimension of $[length]^2$ and represents ultraviolet corrections to the Einstein theory, G is Newton's constant and the cosmological constant is given by

$$\Lambda = -\frac{(d-1)(d-2)}{2l^2}.$$
(3)

The AdS radius is given by l and $F_{\mu\nu} = \partial_{\mu}A_{\nu} - \partial_{\nu}A_{\mu}$ is the Maxwell strength field. The black hole metric, by adopting the limit $d \rightarrow 4$, is given by [31,42]

$$ds^{2} = -f(r)dt^{2} + \frac{1}{f(r)}dr^{2} + r^{2}\left(d\theta^{2} + \sin^{2}\theta d\phi^{2}\right),$$
(4)

where the metric function is given by

$$f(r) = 1 + \frac{r^2}{2\alpha} \left(1 - \sqrt{1 + 4\alpha \left(\frac{2MG}{r^3} - \frac{Q^2G}{r^4} - \frac{1}{l^2}\right)} \right),\tag{5}$$

where Q is the electric charge and M is the black hole's mass. Solving the equation f(r) = 0, we find two horizons: the event horizon r_+ and the Cauchy horizon r_- . Using Eq. (5) and the equation for the even horizon, $f(r_+) = 0$, the black hole's mass is given by

$$M = \frac{r_+}{2G} + \frac{\alpha}{2r_+G} + \frac{r_+^3}{2l^2G} + \frac{Q^2}{2r_+}.$$
(6)

The Hawking temperature of the 4D-EGB black hole is given by

$$T = \frac{f'(r_{+})}{4\pi} = \frac{3r^4l^{-2} + r^2 - GQ^2 - \alpha}{4\pi r^3 + 8\pi\alpha r},$$
(7)

and the electrical potential of the event horizon is defined by

$$\Phi = \frac{Q}{r_+}.\tag{8}$$

In the limit $\alpha \longrightarrow 0$, we recover the mass of Reissner–Nordström black hole in AdS space-time (RN-AdS) [21].

3 Holographic thermodynamics of black holes

In this section, we will construct holographic thermodynamics of 4D-EGB black hole based on AdS/CFT correspondence. In this construction, the function of a partition in the AdS space-time, Z_{AdS} , is equal to the one of the dual boundary theory i.e. Z_{CFT} [20,43]

$$Z_{AdS} = Z_{CFT}.$$
(9)

At finite temperature, the thermal partition function and the free energy W are related by

$$W = -T \ln Z_{CFT},\tag{10}$$

where T is the temperature. In the other hand, the grand canonical ensemble defines the free energy as

$$W = \mu C, \tag{11}$$

where the central charge C determines the number of degrees of microscopic freedom in CFT and its conjugate, μ , is considered as a chemical potential.

The partition function in the AdS space-time is given by the Euclidean action, I_E , of the gravitational theory [44]

$$I_{\rm E} = -\ln Z_{\rm AdS}.\tag{12}$$

In the 4D-EGB gravity, the Euclidean action I_E is given by Eq. (1) and can be decomposed as

$$I_E = I_{EM} + I_{GB},\tag{13}$$

where

$$I_{EM} = \frac{1}{16\pi G} \int d^d x \left(R - 2\Lambda - F_{\mu\nu} F^{\mu\nu} \right),$$
(14)

and

$$I_{GB} = \frac{1}{16\pi G} \int d^d x \frac{\alpha}{d-4} \mathcal{G},\tag{15}$$

are the Einstein-Maxwell (EM) action and the Gauss-Bonnet action, respectively. The EM action I_{EM} is given by [44]

$$TI_{EM} = M - TS_{BH} - \Phi Q - \Omega J, \tag{16}$$

where $S_{BH} = \pi r_+^2/G$ is the Bekenstein-Hawking entropy, T is its conjugate quantity, Φ is the electrical potential of the event horizon, Ω is the angular velocity and J the angular momentum. From now on, we will denote S_{BH} by S. By analogy to Eq. (16), we set the GB action I_{GB} as

$$TI_{GB} = -\alpha \mathcal{A},\tag{17}$$

where we have introduced a new quantity \mathcal{A} as the conjugate of the GB constant α . The expression of this new quantity will be given later in terms of thermodynamics and AdS/CFT quantities. The free energy W is expressed as the partition function Z such that

$$Z_{CFT} = e^{-\beta W},\tag{18}$$

where $\beta = 1/T$. From Eqs. (9)- (12), Eqs. (16) and (17), The classical gravitational action writes

$$Z_{AdS} = e^{-\beta \,\mu C} = e^{-\beta(W_{EM} + W_{GB})},\tag{19}$$

where

$$W_{EM} = M - TS - \Phi Q - \Omega J, \tag{20}$$

and

$$W_{GB} = -\alpha \mathcal{A}.$$
 (21)

are the Einstein-Maxwell and the Gauss-Bonnet free energies, respectively. Finally, the free energy, can be rewritten as

$$\mu C = M - TS - \Phi Q - \Omega J - \alpha \mathcal{A}.$$
(22)

4 Charged Black Holes

In this section, we study in the RPST formalism a charged 4D-EGB black hole i.e. $Q \neq 0$ and J = 0. As we indicated in the introduction, the pressure P and the volume V lose their meanings in the RPS thermodynamics standpoint by introducing the quantity C and its conjugate parameter μ [20,21]. The relationship between the central charge, the AdS radius and the Newton's constant are given by [21,23]

$$C = \frac{l^2}{G}.$$
(23)

The mass of black holes in RPST is interpreted as an internal energy [20, 21], and from Eq. (22), we find the Smarr relation of a charged 4D-EGB black hole in the RPS formalism as

$$M = TS + \tilde{\Phi}\tilde{Q} + \mu C + \mathcal{A}\alpha, \tag{24}$$

where the re-scaled electric potential, $\tilde{\Phi}$, and the re-scaled electric charge, \tilde{Q} , are defined by the dual CFT quantities as [21, 45]

 $\tilde{Q} = \frac{Ql}{\sqrt{G}},\tag{25}$

and

$$\tilde{\Phi} = \frac{\Phi\sqrt{G}}{l} = \frac{Q\sqrt{G}}{r_+l},\tag{26}$$

respectively. From Smarr's relation, we can say that the mass of a black hole is a function of extensive variables

$$M = M(S, Q, C, \alpha). \tag{27}$$

By differentiating Eq. (27), we can formulate the first law in the RPS thermodynamics for a charged 4D-EGB black hole as

$$dM = TdS + \Phi d\hat{Q} + \mu dC + \mathcal{A}d\alpha, \qquad (28)$$

with the corresponding conjugate quantities

$$T = \left(\frac{\partial M}{\partial S}\right)_{\tilde{Q},C,\alpha}, \qquad \tilde{\Phi} = \left(\frac{\partial M}{\partial \tilde{Q}}\right)_{S,C,\alpha}, \qquad \mu = \left(\frac{\partial M}{\partial C}\right)_{S,\tilde{Q},\alpha}, \quad \text{and} \quad \mathcal{A} = \left(\frac{\partial M}{\partial \alpha}\right)_{S,\tilde{Q},C}.$$
 (29)

We can also infer the Gibbs-Duheim relation

$$-SdT = \tilde{Q}d\Phi + Cd\mu + \alpha d\mathcal{A}.$$
(30)

Furthermore, the event horizon radius r_+ in terms of the Bekenstein-Hawking entropy and the AdS/CFT quantities writes

$$r_{+} = l \sqrt{\frac{S}{\pi C}}.$$
(31)

In order to compute thermodynamic variables in RPST, we rewrite the mass of the 4D-EGB black hole, Eq. (6), as

$$M = \frac{S^2 + \pi SC + \pi^2 \tilde{Q}^2 + \pi^2 \alpha C^2 l^{-2}}{2\pi^{3/2} l \sqrt{SC}}.$$
(32)

For $\alpha = 0$, we recover the mass of the RN-AdS black holes in the RPS formalism [21]. The RPST temperature of the 4D-EGB black hole is defined as a partial derivative of Eq. (32), i.e.

$$T = \frac{3S^2 + \pi SC - \pi^2 \tilde{Q}^2 - \alpha C^2 \pi^2 l^{-2}}{4\pi^{3/2} lS \sqrt{SC}}.$$
(33)

This temperature corresponds to the Hawking temperature, Eq. (7), for $A \gg 8\pi\alpha$, where A is the area of the event horizon .The chemical potential, which is the conjugate of the central charge, is given by Eq. (29)

$$\mu = -\frac{S^2 - \pi SC + \pi^2 \tilde{Q}^2 - 3\alpha C^2 \pi^2 l^{-2}}{4\pi^{3/2} l C \sqrt{SC}}.$$
(34)

Finally, the expression of the conjugate quantity of α or the GB potential is given by

$$\mathcal{A} = \frac{C^2 \sqrt{\pi}}{2 l^3 \sqrt{CS}}.$$
(35)

We notice that in the RPST formalism, Schwarzschild black holes i.e. $\tilde{Q} = 0$ are equivalent to RN black holes with an effective charge due to the 4D-EGB coupling. However, The Hawking-Page phase transition in Schwarzschild black holes may occurs at a entropy and temperature specific. Electrical and chemical potentials depend on the electric, the central charges, while the GB potential \mathcal{A} depends only on the central charge. Furthermore, the expression of the Gauss-Bonnet action I_{GB} in term of thermodynamics and AdS/CFT quantities by using Eqs. (17), (33) and (35) is given by

$$I_{GB} = -\frac{\alpha \pi^2 C^2 S}{3S^2 l^2 + \pi C S l^2 - \pi^2 \tilde{Q}^2 l^2 - \alpha C^2 \pi^2}.$$
(36)

This action observes a divergence for T = 0 as shown in Eq. (33). Such unphysical divergences were already found in the on-shell action and surface terms of the total action of the 4D Gauss-Bonnet theory [46]. We can check easily that the mass of the 4D-EGB black holes is a first-order homogeneity and T, Φ , μ and α are zeroth order homogeneity in S, C, μ and \mathcal{A} . Indeed, a re-scaling S, \tilde{Q} , C and \mathcal{A} as $S \to \lambda S$, $\tilde{Q} \to \lambda \tilde{Q}$, $C \to \lambda C$ and $\mathcal{A} \to \lambda \mathcal{A}$ the mass of the 4D-EGB black hole M, Eq.(32), is re-scaled as $M \to \lambda M$ while T, Φ , μ and α remain unchanged. This behavior is also observed from the Smarr relation Eq. (24).

5 Thermodynamic processes

This section is devoted to the study of the phase transition and thermodynamic processes of a charged 4D-EGB black hole by fixing the re-scaled electric charge, the Gauss-Bonnet constant and the central charge. Critical points of the T - S curve are obtained by inflection points of the temperature with respect to the entropy at fixed \tilde{Q} , α and C, i.e. by solving the system

$$\left(\frac{\partial T}{\partial S}\right)_{\tilde{Q},C,\alpha} = 0, \qquad \left(\frac{\partial^2 T}{\partial S^2}\right)_{\tilde{Q},C,\alpha} = 0. \tag{37}$$

Using Eqs. (33) and (37), we obtain the critical parameters as

$$S_c = \frac{\pi C}{6}$$
, and $\tilde{Q}_c = \pm \frac{C}{l} \sqrt{\frac{l^2}{36} - \alpha}$. (38)

Since the charge in the mass expression Eq. (32) is raised to the square root, we do not care about the sign of the charge. The critical charge imposes a bound on the parameter i.e $\alpha l^{-2} < \frac{1}{36}$ and for $\alpha l^{-2} = \frac{1}{36}$, there is no critical charge. By substituting these critical parameters in Eqs. (32) and (33), we obtain

$$M_c = \frac{1}{3}\sqrt{\frac{2}{3}}\frac{C}{l}$$
 and $T_c = \frac{\sqrt{6}}{3\pi l}$. (39)

The equation of state (EoS) in the RPST formalism, by setting $a = \alpha l^{-2}$, is given by

$$t = \frac{3s^2 + 6s - q^2 + 36a(q^2 - 1)}{8s^{3/2}},$$
(40)

where we have normalized the parameters as

$$t = \frac{T}{T_c}, \quad s = \frac{S}{S_c}, \qquad \text{and} \quad q = \frac{\tilde{Q}}{\tilde{Q}_c}.$$
 (41)

We can also introduce the Helmholtz free energy F, its the critical parameter and the normalized



Figure 1: T - S curve for different values of the GB constant and the electric charge.

quantity by

$$F = M - TS, (42)$$

$$F_c = \frac{C}{3\sqrt{6}l},\tag{43}$$

and

$$f = \frac{F}{F_c} = \frac{-4s^{3/2}t + q^2 + s^2 + 6s}{4\sqrt{s}},\tag{44}$$

respectively. We notice that, the EoS parameter and the expression of Helmholtz free energy, Eq. (44), are independent from the central charge C. This means that the central charge does not affect

the thermodynamic behavior of black holes. This is like "the law of corresponding states" in standard thermodynamics for ordinary matters in which the number has the same behavior.



Figure 2: F - T curve for different values of the GB constant and the electric charge.

Figs.1 and 2 show T - S and F - T curves corresponding to a fixed electric charge and the GB constant processes for different values of a and q. In Fig.1, we notice the same thermal behaviors for q = 1, i.e, $\tilde{Q} = \tilde{Q}_c$. This means that the Gauss-Bonnet coupling constant α has no effect on the thermal evolution of the 4D-EGB black holes and this case is accompanied with a second-order phase transition. This result can also be seen from Eq. (40), where RN-AdS black holes at the critical electric charge are recovered even for 4D-EGB black holes [21]. The same is true for the free energy F - T curves in Fig.2. For $0 < \tilde{Q} < \tilde{Q}_c$, the T - S curves are non-monotonic. This represents a first-order phase transition from small black holes to a large one passing through the medium sized black holes are stable. The F - T curves, in this case $(0 < \tilde{Q} < \tilde{Q}_c)$, contain a swallow tail with the root located at temperature $T > T_c$ (supercritical temperature). The Gauss-Bonnet constant affects the evolution of the T - S and the F - S curves for $0 < \tilde{Q} < \tilde{Q}_c$. Indeed, with the same value of q, transition temperatures T_{tran} , are different for different negative values of α . The transition temperature is, furthermore, more pronounced for a negative value of α compared to a vanishing or a positive value. For $\tilde{Q} > \tilde{Q}_c$ there are no phase transitions.

6 Uncharged 4D-EGB Black Holes

From Eqs. (33) and (40), we notice that for an uncharged 4D-EGB black hole, the GB constant behaves as an effective charge in the RPST formalism. This means that an uncharged 4D-EGB black holes behaves like a RN-AdS black holes in AdS/CFT. In this section, we will study an uncharged 4D-EGB black holes in restricted phase space thermodynamics. The first law and the Smarr relation of an uncharged 4D-EGB black holes write as

$$dM = TdS + \mu dC + \mathcal{A}d\alpha,\tag{45}$$

$$M = TS + \mu C + \mathcal{A}\alpha,\tag{46}$$

respectively. The expression of the mass of the uncharged 4D-EGB black hole becomes from Eq. (32)

$$M = \frac{S^2 + \pi SC + \pi^2 \alpha C^2 l^{-2}}{2\pi^{3/2} l \sqrt{SC}}.$$
(47)

The conjugate thermodynamics variables are as follows

$$T = \frac{3S^2 + \pi SC - \alpha C^2 \pi^2 l^{-2}}{4\pi^{3/2} l S \sqrt{SC}},\tag{48}$$

$$\mu = -\frac{S^2 - \pi SC - 3\alpha C^2 \pi^2 l^{-2}}{4\pi^{3/2} l C \sqrt{SC}},\tag{49}$$

$$\mathcal{A} = \frac{C^2 \sqrt{\pi}}{2l^3 \sqrt{CS}}.$$
(50)

Given the conjugate thermodynamics variable in the RPST formalism, we proceed by studying the phase transition and thermodynamic processes (T - S processes) of an uncharged 4D-EGB black holes with a fixed Gauss-Bonnet constant and a central charge. The critical points in the T - S curve at fixed α and C, are

$$S_c = \frac{\pi C}{6}, \quad \alpha_c = \frac{l^2}{36}, \quad T_c = \frac{\sqrt{6}}{3\pi l}, \quad \text{and} \quad F_c = \frac{\sqrt{6}C}{18l}.$$
 (51)

The equation of state and the Helmhotz free energy in RPST formalism for the uncharged 4D-EGB black holes are

$$t = \frac{3s^2 + 6s - b}{8s^{3/2}},\tag{52}$$

and

$$f = \frac{b + s^2 + 6s - 4t \, s^{3/2}}{4s^{1/2}},\tag{53}$$

where we have defined the relative parameter b as $\frac{\alpha}{\alpha_c}$.



Figure 3: T - S curve for different values of the GB constant.

In Figs. 3 and 4, the T-S and F-T curves correspond to the iso- α processes. Although the black hole is not electrically charged, it behaves like a RN-AdS [21]. This means that, in the RPST formalism, the Gauss-Bonnet constant causes an uncharged black hole to behave like a charged one. That is, α may be related to an effective electric charge. Indeed, for $\alpha < \alpha_c$, we have a first-order phase transition, for $\alpha = \alpha_c$ we have a second-order phase transition, and for $\alpha > \alpha_c$ there is no phase transition.



Figure 4: F - T curve for different values of the GB constant.

To find out this relationship, we consider the mass of RN-AdS black holes and that of an uncharged 4D-EGB black hole in the RPST formalism. The mass of an uncharged 4D-EGB black hole is

$$M_{4D-EGB} = \frac{S^2 + \pi SC + \pi^2 \alpha C^2 l^{-2}}{2\pi^{3/2} l \sqrt{SC}},$$
(54)

and the one of RN-AdS black holes is

$$M_{RN-AdS} = \frac{S^2 + \pi SC + \pi^2 \tilde{Q}^2}{2\pi^{3/2} l \sqrt{SC}}.$$
(55)

By comparing the expression of these two masses, a similitude can be made by introducing a re-scaled effective Gauss-Bonnet charge \tilde{Q}_{GB} as

$$\tilde{Q}_{GB} = \frac{C\sqrt{\alpha}}{l}.$$
(56)

In the RPST formalism, Schwarzschild black holes in 4D-EGB are similar to those of RN-AdS one.

One of the well known transition between the Schwarzschild black hole and the thermal AdS space is the Hawking-Page phase transition. These transitions may exist with respect to the conservation of charge and angular momentum. In our setup, i.e. the RPST formalism for 4D-EGB gravity, the Schwarzschild black hole becomes charged at thermodynamic properties level. This effective charge has a topological not electromagnetic properties. Thus, we can still consider the Hawking-Page phase transition between the 4D-EGB black holes and the thermal AdS space in the RPST formalism without violating the conservation of charge and angular momentum. The Gibbs free energy G of the thermal AdS space is zero. The Hawking-Page phase transition requires a null Gibbs free energy.

$$G = M - TS + PV. (57)$$

However, as the pressure and the volume are not thermodynamic variables in the RPST formalism, the Gibbs free energy is none other than the Helmholtz free energy F. Putting G = 0, we get the Hawking-Page entropy and temperature from Eqs. (47) and (48)

$$S_{HP} = \frac{\pi C}{2} \left(\sqrt{\frac{b}{3} + 1} + 1 \right),\tag{58}$$

and

$$T_{HP} = \frac{\sqrt{2} \left(\frac{b}{9} + \sqrt{\frac{b}{3} + 1} + 1\right)}{\pi l \left(\sqrt{\frac{b}{3} + 1} + 1\right)^{3/2}},\tag{59}$$

respectively. From Figs. 3 and 4 and Eqs. (58)-(59), we infer that $S_{HP}/S_c \leq 6$ and T_{HP}/T_c is approximately equal to 1.25, i.e, the Hawking-Page phase transition is between a large black hole and thermal AdS space.

7 Discussion and Conclusion

In this paper, we have studied the 4D-EGB black hole in the context of restricted phase space thermodynamics. We have started by solving the space-time black hole with a negative cosmological constant, with respect to 4D-EGB gravity. From the metric, we deduced the expression of the mass of the black hole in terms of the parameters of the black hole, the geometry, and the GB constant.

We constructed a holographic thermodynamic of the 4D-EGB black hole, based on the AdS/CFT correspondence by equating the partition function of AdS to that of CFT ($Z_{AdS} = Z_{CFT}$) and the free energy $W = \mu C$, to the Euclidean action $I_E = -\ln Z_{CFT}$. We were able to formulate Smarr relation, the first law and the Gibbs-Duhem relation for a charged and an uncharged 4D-EGB black holes. We show that Smarr relation is homogeneous at a first order. In addition, we have derived thermodynamic variables in the RPST formalism.

We have also studied thermodynamic properties of a charged black hole, and we found that the equation of state does not depend on the central charge. This result is similar to that in the law of corresponding states in standard thermodynamics for ordinary matter. We show that when the charge of black hole is greater than the critical one, there is no phase transition, but when $Q = Q_c$ we find a phase transition of the second order. For $0 < Q < Q_c$, we find a first-order phase transition. A remarkable result that not only the charge that influences thermodynamic properties but even the GB constant, precisely when the charge is sub-critical.

For the study of the uncharged black hole, we find that its thermodynamic properties depend on the value of the GB constant. Indeed, for $\alpha = \alpha_c$, it corresponds to a second-order phase transition and $0 < \alpha < \alpha_c$, it corresponds to a first-order phase transition. But, for $\alpha > \alpha_c$, there is no phase transition. Although it is not charged, the black hole in 4D-EGB behaves as if it is charged one with an effective charge that depends on the GB constant and the central charge.

Finally, the Hawking-Page phase transition may occurs between a large black hole and a thermal AdS space.

References

- S. W. Hawking. Particle Creation by Black Holes. Commun. Math. Phys., 43:199–220, 1975.
 [Erratum: Commun.Math.Phys. 46, 206 (1976)].
- [2] Jacob D. Bekenstein. Black Holes and Entropy. Phys. Rev. D, 7:2333–2346, 1973.
- [3] J. D. Bekenstein. Black holes and the second law. Lettere al Nuovo Cimento (1971-1985), 4(15):737-740, 1972.
- [4] J. M. Bardeen, B. Carter, and S. W. Hawking. The four laws of black hole mechanics. Communications in Mathematical Physics, 31(2):161–170, 1973.
- [5] Stephen W Hawking. Black holes and thermodynamics. Physical Review D, 13(2):191, 1976.
- [6] S. W. Hawking and Don N. Page. Thermodynamics of black holes in anti-de Sitter space. Communications in Mathematical Physics, 87(4):577–588, 1983.
- [7] David Kastor, Sourya Ray, and Jennie Traschen. Enthalpy and the mechanics of AdS black holes. Classical and Quantum Gravity, 26(19):195011, 2009.
- [8] David Kubizňák and Robert B Mann. Black hole chemistry. Canadian Journal of Physics, 93(9):999–1002, 2015.
- [9] David Kubizňák, Robert B Mann, and Mae Teo. Black hole chemistry: thermodynamics with Lambda. Classical and Quantum Gravity, 34(6):063001, 2017.
- [10] Brian P. Dolan. Compressibility of rotating black holes. *Physical Review D*, 84(12), 2011.
- [11] Rong-Gen Cai, Li-Ming Cao, Li Li, and Run-Qiu Yang. P-V criticality in the extended phase space of Gauss-Bonnet black holes in AdS space. *Journal of High Energy Physics*, 2013(9), 2013.
- [12] David Kubizňák and Robert B. Mann. P V criticality of charged AdS black holes. Journal of High Energy Physics, 2012(7), 2012.
- [13] Brian P Dolan. Pressure and volume in the first law of black hole thermodynamics. *Classical and Quantum Gravity*, 28(23):235017, 2011.
- [14] Krishnakanta Bhattacharya. Extended phase space thermodynamics of black holes: A study in einstein's gravity and beyond. Nuclear Physics B, page 116130, 2023.
- [15] Peng Wang, Houwen Wu, Haitang Yang, and Feiyu Yao. Extended phase space thermodynamics for black holes in a cavity. *Journal of High Energy Physics*, 2020(9):1–19, 2020.
- [16] SH Hendi and MH Vahidinia. Extended phase space thermodynamics and P- V criticality of black holes with a nonlinear source. *Physical Review D*, 88(8):084045, 2013.
- [17] Milko Estrada and Rodrigo Aros. Thermodynamic extended phase space and PV criticality of black holes at Pure Lovelock gravity. The European Physical Journal C, 80(5):1–16, 2020.
- [18] Juan Maldacena. The Large-N Limit of Superconformal Field Theories and Supergravity. International Journal of Theoretical Physics, 38(4):1113–1133, 1999.
- [19] Koji Hashimoto, Shunichiro Kinoshita, and Keiju Murata. Imaging black holes through the AdS/CFT correspondence. *Physical Review D*, 101(6), 2020.
- [20] Manus R. Visser. Holographic thermodynamics requires a chemical potential for color. *Physical Review D*, 105(10), 2022.
- [21] Zeyuan Gao and Liu Zhao. Restricted phase space thermodynamics for AdS black holes via holography. *Classical and Quantum Gravity*, 39(7):075019, 2022.
- [22] Andreas Karch and Brandon Robinson. Holographic black hole chemistry. Journal of High Energy Physics, 2015(12):1–15, 2015.

- [23] Zeyuan Gao, Xiangqing Kong, and Liu Zhao. Thermodynamics of Kerr-AdS black holes in the restricted phase space. The European Physical Journal C, 82(2), 2022.
- [24] Jafar Sadeghi, Mehdi Shokri, Saeed Noori Gashti, and Mohammad Reza Alipour. RPS thermodynamics of Taub–NUT AdS black holes in the presence of central charge and the weak gravity conjecture. *General Relativity and Gravitation*, 54(10):129, 2022.
- [25] Yan-Ying Bai, Xuan-Rui Chen, Zhen-Ming Xu, and Bin Wu. Revisit on thermodynamics of BTZ black hole with variable Newton constant. arXiv preprint arXiv:2208.11859, 2022.
- [26] Xiangqing Kong, Tao Wang, Zeyuan Gao, and Liu Zhao. Restricted phased space thermodynamics for black holes in higher dimensions and higher curvature gravities. *Entropy*, 24(8):1131, 2022.
- [27] H Weyl. Sitzungsber. Preuss. Akad. Wiss, Berlin, 465,(1918). Annalen der Physik (Lpz), 59:101, 1919.
- [28] Arthur Stanley Eddington. A generalisation of Weyl's theory of the electromagnetic and gravitational fields. Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character, 99(697):104–122, 1921.
- [29] Baojiu Li, John D Barrow, and David F Mota. Cosmology of modified Gauss-Bonnet gravity. *Physical Review D*, 76(4):044027, 2007.
- [30] Guido Cognola, Emilio Elizalde, Shin'ichi Nojiri, Sergei D Odintsov, and Sergio Zerbini. Dark energy in modified Gauss-Bonnet gravity: Late-time acceleration and the hierarchy problem. *Physical Review D*, 73(8):084007, 2006.
- [31] Draž en Glavan and Chunshan Lin. Einstein-Gauss-Bonnet Gravity in Four-Dimensional Spacetime. *Physical Review Letters*, 124(8), 2020.
- [32] David Lovelock. The Einstein tensor and its generalizations. Journal of Mathematical Physics, 12(3):498–501, 1971.
- [33] Pedro G.S. Fernandes. Charged black holes in AdS spaces in 4D Einstein Gauss-Bonnet gravity. *Physics Letters B*, 805:135468, 2020.
- [34] Shao-Wen Wei and Yu-Xiao Liu. Extended thermodynamics and microstructures of fourdimensional charged Gauss-Bonnet black hole in AdS space. *Physical Review D*, 101(10):104018, 2020.
- [35] Seyed Ali Hosseini Mansoori. Thermodynamic geometry of the novel 4-D Gauss–Bonnet AdS black hole. *Physics of the Dark Universe*, 31:100776, 2021.
- [36] M Bousder and M Bennai. Particle-antiparticle in 4D charged Einstein-Gauss-Bonnet black hole. *Physics Letters B*, 817:136343, 2021.
- [37] M Guo and PC Li. The innermost stable circular orbit and shadow in the novel 4D. Einstein-Gauss-Bonnet gravity, 2003.
- [38] Cheng-Yong Zhang, Peng-Cheng Li, and Minyong Guo. Greybody factor and power spectra of the Hawking radiation in the 4 D Einstein–Gauss–Bonnet de-Sitter gravity. *The European Physical Journal C*, 80(9):874, 2020.
- [39] Cheng-Yong Zhang, Shao-Jun Zhang, Peng-Cheng Li, and Minyong Guo. Superradiance and stability of the novel 4D charged Einstein-Gauss-Bonnet black hole. arXiv preprint arXiv:2004.03141, 2020.
- [40] Moisés Bravo-Gaete, Luis Guajardo, and Julio Oliva. Nonlinear charged planar black holes in four-dimensional scalar-Gauss-Bonnet theories. *Physical Review D*, 106(2):024017, 2022.
- [41] Ke Yang, Bao-Min Gu, Shao-Wen Wei, and Yu-Xiao Liu. Born–Infeld black holes in 4D Einstein– Gauss–Bonnet gravity. The European Physical Journal C, 80(7):1–10, 2020.

- [42] Pedro G S Fernandes, Pedro Carrilho, Timothy Clifton, and David J Mulryne. The 4D Einstein–Gauss–Bonnet theory of gravity: a review. *Classical and Quantum Gravity*, 39(6):063001, 2022.
- [43] Edward Witten. Anti De Sitter Space And Holography. 1998.
- [44] G. W. Gibbons and S. W. Hawking. Action integrals and partition functions in quantum gravity. *Phys. Rev. D*, 15:2752–2756, 1977.
- [45] Wan Cong and David Kubizň ák and Robert B. Mann. Thermodynamics of AdS Black Holes: Critical Behavior of the Central Charge. *Physical Review Letters*, 127(9), 2021.
- [46] Subhash Mahapatra. A note on the total action of 4 D Gauss–Bonnet theory. *The European Physical Journal C*, 80, 2020.