


Hayden-Preskill Recovery in Hamiltonian Systems

Yoshifumi Nakata¹ * and Masaki Tezuka² †

¹*Yukawa Institute for Theoretical Physics, Kyoto University,
Kitashirakawa, Sakyo-ku, Kyoto 606-8502, Japan, and*

²*Department of Physics, Kyoto University, Kitashirakawa, Sakyo-ku, Kyoto 606-8502, Japan.*

(Dated: March 24, 2023)

The key to understanding complex quantum systems is information scrambling originally proposed in relation to the Hayden-Preskill recovery, which refers to localized information being spread over the whole system and being accessible from small subsystems. This phenomenon is well-understood in random unitary models but is hardly explored in Hamiltonian systems. In this Letter, we investigate the information recovery for various time-independent Hamiltonians, including Sachdev-Ye-Kitaev (SYK) models and chaotic spin chains. Despite the fact that the Hayden-Preskill recovery typically relies on the assumption never satisfied in Hamiltonian systems, we show that information recovery is possible in certain, but not all, chaotic models. This highlights the difference between information scrambling and quantum chaos and offers the information recovery as a fully operational probe for the manifestation of novel complex quantum dynamics.

A central challenge in modern physics is to characterize the dynamics in far-from-equilibrium quantum many-body systems. The Hayden-Preskill protocol [1] offers an operational approach toward this goal and is at the heart of the interdisciplinary researches [2–26], including proposals of experiments of quantum gravity [27–29]. The protocol addresses if the information initially localized in a small subsystem can be recovered from other subsystems after unitary time evolution. It turned out that, if the unitary dynamics is sufficiently random, the information rapidly spreads over the system, and information can be recovered from any small subsystems. This is closely related to quantum random encoding [30] and is known as the *Hayden-Preskill recovery*. The Hayden-Preskill recovery is of use for operationally characterizing complex quantum many-body dynamics.

The discovery of the Hayden-Preskill recovery substantially accelerated the interaction over theoretical physics: it is inspired by the information paradox of quantum black holes [31–33], is formulated in the language of quantum information, and is investigated by the technique of random matrix theory (RMT) [34, 35]. The unitary dynamics leading to the Hayden-Preskill recovery is referred to as *information scrambling* [1]. A huge number of studies aiming to characterize information scrambling were performed from different perspectives, such as entanglement generation [2, 4], operator mutual information (OMI) [19, 36–38], and out-of-time-ordered correlators (OTOCs) [26, 39, 40].

A canonical model is the SYK model [11, 12, 41, 42] as it is expected to be a holographic dual to quantum gravity [13, 14, 43–45]. The dynamics of SYK model turns out to rapidly saturate OTOCs [11, 12] and to take the

maximum possible value of the quantum Lyapunov exponent [46]. The static properties of the SYK model have also been studied and were shown to have RMT-like energy statistics, implying that it is quantum chaotic in a conventional notion based on the Bohigas-Giannoni-Schmit (BGS) conjecture [47]. To better understand the model, various variants have been proposed [48], and transitions from quantum chaos to a many-body localization (MBL) have been studied in detail [49, 50]. A family of SYK models is, hence, of significant interest in high energy and strongly-correlated physics.

Despite these progresses, the Hayden-Preskill recovery in a family of SYK models has been rarely explored so far [51]. The Hayden-Preskill recovery is widely believed to be achieved in quantum chaotic systems, but this is yet to be confirmed: while the Hayden-Preskill recovery strongly relies on the assumption that the unitary time evolution is sufficiently random, often called as Haar random, the dynamics in time-independent Hamiltonian systems is never as random as Haar even approximately [8]. Furthermore, quantum chaos is conventionally characterized by eigenenergy statistics, as in the BGS conjecture, which is a *static* property, but the Hayden-Preskill recovery is about the *dynamical* properties. Thus, the relation between quantum chaos and the Hayden-Preskill recovery is not a priori trivial.

In this Letter, we investigate information scrambling in the original sense of the Hayden-Preskill recovery in various quantum Hamiltonian systems. We first quantitatively confirm the absence and presence of the Hayden-Preskill recovery in commuting and the original SYK Hamiltonians, respectively, and then show that, unlike the common belief, quantum chaotic Hamiltonian does not always achieve the Hayden-Preskill recovery. We also show that chaotic-to-integrable transitions in a family of SYK models are well-captured by the information recovery, reflecting the drastic changes of quan-

* yoshifumi.nakata@yukawa.kyoto-u.ac.jp

† tezuka@scphys.kyoto-u.ac.jp




FIG. 1. A diagram of the Hayden-Preskill protocol. Time flows from bottom to top. Horizontal lines imply that the qubits connected by the line may be entangled. The initial states on AR and BB' are given by a maximally entangled state of k ebits, i.e., k EPR pairs, that keeps track of quantum information in A , and a thermofield double state $|\xi(\beta)\rangle$ at the inverse temperature β in B , respectively. The system $S := AB$ undergoes Hamiltonian dynamics by \hat{H}_S , and then, is split into two random subsystems C of ℓ qubits and D of $N - \ell$ qubits. By applying a quantum channel \mathcal{D} onto $B'C$, one aims to decode the quantum information in A , that is, to recover the k EPR pairs between \hat{A} and R . This protocol has a natural interpretation in the context of information paradox [1]. See also a tutorial [52].

tum information-theoretic properties of the system. The Hayden-Preskill protocol in Hamiltonian systems, thus, offers an information-theoretic probe of manifestation of complex quantum dynamics.

The Hayden-Preskill recovery.— Given quantum many-body system S of N qubits, we encode quantum information into a localized subsystem $A \subset S$ of k qubits ($k \ll N$). We then let the system S undergo the Hamiltonian time-evolution $U_{\hat{H}}(t) := e^{-i\hat{H}t}$ for some time t , where \hat{H} is the Hamiltonian in S . After the time-evolution, the information is tried to be recovered from an ℓ -qubit subsystem $C \subset S$. Throughout our analysis, we assume $C \subset B = S \setminus A$ as far as $\ell \leq N - k$. The question is how large ℓ should be for a successful recovery.

The answer highly depends on the initial state in B as well as available resources in the recovery process. Here, we assume that the initial state in B is a thermal state $\xi^B(\beta)$ at the inverse temperature β , which is purified to a thermofield double state $|\xi(\beta)\rangle^{BB'}$ by a system B' . We also assume that the purifying system B' can be made use of in the recovery process, which has a natural interpretation in the information paradox of black holes [1]. See Fig. 1.

The recovery of quantum information is defined using k Einstein-Podolsky-Rosen (EPR) pairs $|\Phi\rangle^{AR}$ between A and R , where the reference system R is virtually introduced for analyses. If we can reproduce k EPR pairs in AR from the state in $B'CR$ by a quantum operation \mathcal{D} on $B'C$ within some error, then it implies that infor-

mation in A is recoverable within that error. The precise definition is provided in Supplemental Material S1.

For given \hat{H} , t , and β , it is in general difficult to exactly compute the recovery error $\Delta_{\hat{H}}(t, \beta)$, which we normalize such that $\Delta_{\hat{H}} \in [0, 1]$. Nevertheless, its upper bound can be computed based on the decoupling approach [53–55]. To this end, we introduce

$$\Psi_{\text{fin}}^{SB'R}(t, \beta) = U_{\hat{H}}^S(t)(\Phi^{AR} \otimes \xi^{BB'})U_{\hat{H}}^S(t)^\dagger. \quad (1)$$

Here, we indicate the system over which the partial trace is taken by the superscript, such as $\Psi_{\text{fin}}^{DR}(t, \beta)$, where $D = S \setminus C$, is the partial trace of $\Psi_{\text{fin}}^{SB'R}$ over CB' . Using this notation, a calculable and typically good upper bound on the recovery error $\Delta_{\hat{H}}(t, \beta)$ is obtained (see Supplemental Material S1)

$$\Delta_{\hat{H}}(t, \beta) \leq \bar{\Delta}_{\hat{H}}(t, \beta) := \min\{1, \sqrt{2\Sigma_{\hat{H}}(t, \beta)}\}, \quad (2)$$

where $\Sigma_{\hat{H}}(t, \beta) := \|\Psi_{\text{fin}}^{DR}(t, \beta) - \Psi_{\text{fin}}^D(t, \beta) \otimes \pi^R\|_1/2$, $\|\cdot\|_1$ is the trace norm, and $\pi^R = I^R/2^k$ is the completely mixed state in R .

Note that $\Sigma_{\hat{H}}(t, \beta)$ can be characterized by the mutual information $I(R : D)_{\Psi_{\text{fin}}(t, \beta)} = 2k - I(R : B'C)_{\Psi_{\text{fin}}(t, \beta)}$ as $2\Sigma_{\hat{H}}^2/\ln 2 \leq I(R : D)_{\Psi_{\text{fin}}(t, \beta)} \leq 2k\Sigma_{\hat{H}} + (1 + \Sigma_{\hat{H}})h(\Sigma_{\hat{H}}/(1 + \Sigma_{\hat{H}}))$, where $h(x)$ is the binary entropy [30]. Since $I(R : D)_{\Psi_{\text{fin}}(t, \beta)}$ have been studied in the context of the OMI [19] especially when $\beta = 0$, the recovery error can be computed from the OMI (see Supplemental Material S1), which, however, results in less tight bound. Hence, we focus on $\bar{\Delta}_{\hat{H}}(t, \beta)$.

Information scrambling and quantum chaos.— The recovery error has been easily computed in a random unitary model, in which the time evolution $U_{\hat{H}}^S(t)$ is replaced with a Haar random unitary, or equivalently a circular unitary ensemble (CUE). Denoting by $\Delta_{\text{Haar}}(\beta)$ the recovery error in this case, it holds with high probability that [1, 24, 55]

$$\Delta_{\text{Haar}}(\beta) \leq \bar{\Delta}_{\text{Haar}}(\beta) := \min\{1, 2^{\frac{1}{2}(\ell_{\text{Haar,th}}(\beta) - \ell)}\}. \quad (3)$$

see Supplemental Material S2. Note that there is no concept of time t in a random unitary model. Here, $\ell_{\text{Haar,th}}(\beta) := \frac{1}{2}(N + k - H(\beta))$, and $H(\beta)$ is the Renyi-2 entropy of $\xi^B(\beta)$, defined by $2^{-H(\beta)} = \text{Tr}[(\xi^B(\beta))^2]$. The bound (Eq. (3)) can also be obtained from the property of entanglement generated by a Haar random unitary [24].

It is clear from Eq. (3) that $\Delta_{\text{Haar}}(\beta) \ll 1$ if $\ell \gg \ell_{\text{Haar,th}}(\beta)$. In particular, $\ell_{\text{Haar,th}}(0) = k$. Hence, if the system is initially at infinite temperature, the k -qubit quantum information in A is recoverable from any subsystem of the size that is independent of N . This is the Hayden-Preskill recovery. Note that the Hayden-Preskill recovery is fully operational since all the processes, i.e., encoding and recovery of information, are operative and can be in principle realized in practical experiments.

Following the original proposal [1], we refer to the dynamics achieving the Hayden-Preskill recovery as information scrambling. Although it is common to interchangeably use information scrambling and quantum chaos, we clearly distinguish them: quantum chaos is based on a RMT-like eigenenergy spectrum.

Commuting Hamiltonians.— We start with the simplest instance of commuting Hamiltonians in the form of $\hat{H} = \hat{H}^A + \hat{H}^B + \hat{H}^{A:B}$, where \hat{H}^A and \hat{H}^B are Hamiltonians in A and B , and $\hat{H}^{A:B}$ is an interaction Hamiltonian over A and B and commutes with both \hat{H}^A and \hat{H}^B . In Supplemental Material S3, we show that this system is never information scrambling, while it can show chaotic features (see e.g. [56] and Appendix D in [57]).

The absence of information scrambling is due to a lack of information spreading. Note that a spread of information is necessary for the information recovery. We can explicitly show that the dynamics by \hat{H} spreads information only to the neighbours in terms of $\hat{H}^{A:B}$. This leads to $\Delta_{\hat{H}}(t, \beta) \geq 1 - 2^{-(\kappa-k-1)/2}$ for any t and β whenever $A \subseteq D$, where κ is the number of qubits in A that directly interact with B . Hence, when $\kappa \leq k - 1$, the error remains constant and the Hayden-Preskill recovery is not achieved. The bound can be improved for concrete commuting Hamiltonians.

A family of SYK Hamiltonians.— We next investigate the Hayden-Preskill recovery for the SYK model, SYK₄, consisting of $2N_q$ Majorana fermions. The Hamiltonian is

$$\hat{H}_{\text{SYK}_4} = \sum_{1 \leq a_1 < a_2 < a_3 < a_4 \leq 2N_q} J_{a_1 a_2 a_3 a_4} \hat{\psi}_{a_1} \hat{\psi}_{a_2} \hat{\psi}_{a_3} \hat{\psi}_{a_4}, \quad (4)$$

in which $\hat{\psi}_j$ is Majorana fermion operators. The real couplings $J_{a_1 a_2 a_3 a_4}$ are independently chosen at random from the Gaussian distribution with average zero and variance $\sigma^2 = \binom{2N_q}{4}^{-1}$. To avoid the effect of parity symmetry of SYK₄, which induces significant deviations to the Hayden-Preskill recovery [9, 10, 24, 58], we focus on the even-parity sector and set $N = N_q - 1$. A Haar random unitary in the corresponding situation leads to the recovery error bounded from above with high probability by $\Delta'_{\text{Haar}}(\beta) = \min\{1, 2^{(\ell_{\text{Haar, th}}(\beta) - \ell) - \frac{1}{4}}\}$. Here, the initial thermal state $\xi(\beta)^B$ is set to the even parity. See Supplemental Material S4 for details. We have also checked that the effect by the periodicity, characterized by $N_q \bmod 4$, is negligible.

In Fig. 2, we numerically plot the upper bound on the recovery error at infinite temperature, $\bar{\Delta}_{\text{SYK}_4}(t, \beta = 0)$, against time t for various ℓ . It clearly shows that $\bar{\Delta}_{\text{SYK}_4}$ quickly approaches $\bar{\Delta}'_{\text{Haar}}$. This is also the case for finite temperature case. The time scale needed for $\bar{\Delta}_{\text{SYK}_4}$ converging to $\bar{\Delta}'_{\text{Haar}}$ is evaluated to be $O(\sqrt{N_q})$, which corresponds to $O(1)$ in the normalization of the Hamiltonian commonly used in the context of holographic duality. This supports that the fast scrambling conjecture [2–4], stating that information scrambling occurs in time $O(\ln N_q)$, is likely to hold in SYK₄, although our result is




FIG. 2. Semilogarithmic plot of the value of $\bar{\Delta}_{\hat{H}_{\text{SYK}_4}}(t, \beta = 0)$ against t for $k = 1, N_q = 13$, and $2 \leq \ell < N_q - k$. Average over 64 samples is taken. For $\ell = 1$, $\bar{\Delta}_{\hat{H}_{\text{SYK}_4}}(t, \beta = 0)$ is ~ 1 for all t . The dashed lines represent $\bar{\Delta}'_{\text{Haar}}(\beta = 0) = 2^{\frac{1}{2}(1-\ell)}$ given in Eq. (3) for $\ell = 2, 3, \dots, N_q - 2$. For smaller values of N_q and finite β , see respectively Fig. S1 and Fig. S2 in the Supplemental Material S4.

not sufficiently precise to distinguish $O(1)$ and $O(\ln N_q)$.

These results imply that the SYK₄ dynamics, while differs from Haar random, quickly achieves the Hayden-Preskill recovery. Thus, SYK₄ is both quantum chaotic and fast information scrambling, highlighting an excellent agreement of the model with the prediction by RMT both in terms of static and dynamic properties.

In practice, however, the SYK₄ model is too complex to be realized due to its all-to-all interaction. For this reason, a simplified models have been studied. A common simplification is a sparse SYK model (spSYK₄), in which the number of non-zero random coupling constant is fixed to K_{cpl} . It recovers SYK₄ when $K_{\text{cpl}} = \binom{N_q}{4}$, but $K_{\text{cpl}} = O(N_q)$ is known to suffice to have chaotic features and to reproduce holographic properties [59, 60]. This leads to a natural question about whether the information scrambling remains by this simplification.

In Fig. 3, we plot the upper bound on the recovery error for a further simplified sparse SYK model ($\pm\text{spSYK}_4$), in which a half of the non-zero couplings is set to $1/\sqrt{K_{\text{cpl}}}$ and the other half to $-1/\sqrt{K_{\text{cpl}}}$ [61]. We clearly observe that the value is nearly identical to the Haar value for $K_{\text{cpl}} \gtrsim 30$, which is of order N_q . This implies that $\pm\text{spSYK}_4$ with $K_{\text{cpl}} = O(N_q)$, which is substantially smaller than $K_{\text{cpl}} = O(N_q^4)$ of SYK₄, suffices also to reproduce information-theoretic properties of SYK₄. See Supplemental Material S5 for details. This would help experimental realizations of the model.

Yet another SYK model attracting much attention is the SYK₄₊₂ model [62]. The Hamiltonian is

$$\hat{H}_{\text{SYK}_{4+2}}(\theta) = \cos \theta \hat{H}_{\text{SYK}_4} + \sin \theta \hat{H}_{\text{SYK}_2}, \quad (5)$$

where $\hat{H}_{\text{SYK}_2} = i \sum_{1 \leq b_1 < b_2 \leq 2N_q} K_{b_1 b_2} \hat{\psi}_{b_1} \hat{\psi}_{b_2}$, and $\theta \in$




FIG. 3. The late-time value of $\bar{\Delta}_{\pm\text{spSYK}_4}(t, \beta = 0)$ against ℓ is plotted for $k = 1$ and various numbers K_{cpl} of non-zero coupling constant. We set N_q to 13, and the number of samples is 64. The average for $t = (1, 2, \dots, 10) \times 10^6$ is plotted as the late-time value in all figures.

$[0, \pi/2]$ is a mixing parameter. The coupling constants $\{K_{b_1 b_2}\}$ satisfy $K_{b_2 b_1} = -K_{b_1 b_2}$ and are normalized for the variance of eigenenergies of $\hat{H}_{\text{SYK}_{4+2}}(\theta)$ to be unity.

The SYK₄₊₂ model has been studied since it has a peculiar energy-shell structure in the sense of the local density of states in Fock space. The structure is known to be drastically changed by varying θ . Accordingly, the range of $\theta \in [0, \pi/2]$ is divided into four regimes I, II, III, and IV [49, 50]. In I, there is only one energy-shell as large as the whole Hilbert space, and it is quantum chaotic. In II, the size of the energy-shell is diminished, and $\mathcal{O}(\text{poly}(N_q))$ energy-shells appear. This diminishing of each energy-shell further proceeds in III. The energy statistics remains RMT-like in these two regimes, and characterizing physics in II and III has been under intense investigations [63]. In IV, the number of energy-shell approaches $\mathcal{O}(\exp(N_q))$, and Fock-space localization is observed.

By investigating The Hayden-Preskill recovery in SYK₄₊₂, we can operationally characterize each regime. In Fig. 4, we plot the late time values of the upper bound $\bar{\Delta}_{\text{SYK}_{4+2}}(t, \beta = 0)$ of the recovery error against $\tan \theta$. Two characteristic values of θ , $\tan \theta_1 \approx 0.5$ and $\tan \theta_2 \approx 20$, are observed, by which three plateaus appear. In the first plateau ($\theta \in [0, \theta_1)$) corresponding to the regime I, $\bar{\Delta}_{\text{SYK}_{4+2}} \approx \bar{\Delta}'_{\text{Haar}}$ and the system is information scrambling. The second is for $\theta \in (\theta_1, \theta_2]$, corresponding to II and III, where $\bar{\Delta}_{\text{SYK}_{4+2}}$ is substantially larger than $\bar{\Delta}'_{\text{Haar}}$. The difference between them seems to remain constant even if N_q increases (see Supplemental Material S6). The third plateau, $\theta \in (\theta_2, \pi/2]$, corresponds to IV, where the system is almost SYK₂.

For sufficiently small and large θ , the behaviour of $\bar{\Delta}_{\text{SYK}_{4+2}}$ can be naturally understood. For small θ , the model is approximately SYK₄. Since the dynamics of SYK₄ quickly achieves the Hayden-Preskill recovery, so




FIG. 4. The late-time value of $\bar{\Delta}_{\text{SYK}_{4+2}}(t, \beta = 0)$ plotted for $k = 1$ and various ℓ against the value of δ . N_q is set to 13. 64 samples are used. The lines connecting the data points are guide to the eye. The horizontal dashed lines indicate $\bar{\Delta}'_{\text{Haar}}$ for various ℓ .

does that in the regime I. In contrast, for sufficiently large θ , the model is almost SYK₂ and the Fock-space localization occurs. Hence, the recovery error should remain a large constant in the regime IV. Despite these natural behaviours in the regimes I and IV, $\bar{\Delta}_{\text{SYK}_{4+2}}$ is smoothly changing for the intermediate values of θ , which is in tension with the division of the regimes II and III in terms of the energy-shell structure.

To understand the intermediate plateau, we shall recall that, in II and III, transitions from one energy-shell to the other are strongly suppressed, which effectively results in the division of the whole Hilbert space into $\mathcal{O}(\text{poly}(N_q))$ energy-shells [49]. Additionally, it is known that the dynamics in each energy-shell seems to be approximately Haar random *within* the subspace [50]. The intermediate plateau of $\bar{\Delta}_{\text{SYK}_{4+2}}$ can be explained from these common features in II and III. Since the whole Hilbert space is effectively divided into smaller ones, within which the dynamics remains still Haar random, the unitary dynamics in II and III induces *partial decoupling* [64] rather than decoupling. While decoupling results in the Hayden-Preskill recovery (Eq. (3)), partial decoupling leads to the recovery error into the form of $2^{\ell'_{\text{th}} - \ell} + \Delta_{\text{rem}}$ [24]. Here, $\ell'_{\text{th}} \approx \ell_{\text{Haar, th}} + \mathcal{O}(\sqrt{k})$ and Δ_{rem} , which is inverse proportional to the standard deviation of energy in the subsystem D , represents the amount of information that cannot be recovered even when ℓ is large. While ℓ'_{th} is hardly observed in our analysis, as we set $k = 1$, we can clearly observe Δ_{rem} as an intermediate plateau. Since the standard deviation of energy in B shall be $\mathcal{O}(\sqrt{N_q})$, that in the subsystem D is at least $\mathcal{O}(\sqrt{N_q})$. This provides a qualitative estimation, $\Delta_{\text{rem}} = \mathcal{O}(1/\sqrt{N_q})$, of the recovery error in the plateau even for large ℓ .

From this perspective, the two transitions can be understood as consequences of drastic changes of the de-

coupling properties induced by the dynamics. In I, the combined regime over II and III, and IV, the SYK₄₊₂ dynamics leads to full, partial, and no decoupling, respectively. Accordingly, each regime has qualitatively different information-theoretic properties and results in different behaviours of information recovery.

The emerging difference between II and III should be an artifact due to the fact that the energy shell is viewed in the Fock basis. The Fock basis shall be appropriate in III, as it is nearly SYK₂, but not in II, as it should inherit the basis of SYK₄ and is more random. Since the choice of basis is not intrinsic, the difference between II and III is not observed in terms of the information recovery.

It is worth noting that the information recovery provides an operational probe for characterizing SYK₄₊₂ in the experimentally-realizable manner. This is in contrast to the eigenenergy and eigenstates statistics, which are in general difficult to experimentally access.

Heisenberg and transversal Ising Hamiltonians with magnetic field.— We finally investigate information recovery in spin-1/2 chains, particularly, the Heisenberg with random magnetic field $\hat{H}_{\text{XXZ}} = \sum_{j=1}^{N-1} (S_j^x S_{j+1}^x + S_j^y S_{j+1}^y + J_z S_j^z S_{j+1}^z) + \sum_{j=1}^N h_j S_j^z$, where h_j are independently sampled from a uniform distribution in $[-W, W]$, and the transversal Ising with constant magnetic field, $\hat{H}_{\text{Ising}} = -\sum_{j=1}^{N-1} (S_j^z S_{j+1}^z) - g \sum_{j=1}^L S_j^x - h \sum_{j=1}^L S_j^z$.

Both Hamiltonians show integrable–chaotic transitions by varying W or (g, h) . However, our analysis reveals that, for any value of the parameters, the recovery errors remain a high value for any t , clearly implying an absence of the the Hayden-Preskill recovery in those systems. See Supplemental Materials S7 and S8 for the details. This should be compared to the facts that they are quantum chaotic for certain parameters and that OTOCs for local observables saturate at late time [65–70]. Thus, neither quantum chaotic features nor the saturation of OTOCs for local observables implies the Hayden-Preskill recovery.

These highlight that information scrambling, in the sense of the Hayden-Preskill recovery, is a manifestation of complex dynamics that captures an aspect different from quantum chaos and OTOCs.

Summary and discussions.— In this Letter, we have studied information scrambling in the sense of the Hayden-Preskill recovery. We have first confirmed the absence and the presence of information scrambling in commuting and SYK₄ Hamiltonians, respectively. The latter remains even if SYK₄ is substantially simplified to $\pm\text{spSYK}_4$. The information scrambling was also shown to information-theoretically characterize the transitions in SYK₄₊₂. Finally, no information scrambling occurs in chaotic spin chains despite the saturation of OTOCs at late time.

Our analyses reveal that the Hayden-Preskill recovery is potentially useful for investigating the manifestation of novel complex quantum dynamics from the information-theoretic viewpoint, which cannot be captured by conventional concepts, such as quantum chaos and OTOCs for local observables. It will be of crucial importance to further clarify the relation over the Hayden-Preskill recovery, quantum chaos, and the saturation of OTOCs in realistic situations.

The work was supported by Grants-in-Aid for Transformative Research Areas (A) No. JP21H05182, No. JP21H05183 and No. JP21H05185 from MEXT of Japan. The work of M.T. was partly supported by Grants-in-Aid No. JP20H05270 and No. JP20K03787 from MEXT of Japan. Y.N. was supported by JST, PRESTO Grant Number JPMJPR1865, Japan, and by JSPS KAKENHI Grant Number JP22K03464, Japan. Part of the computation in this paper was conducted using the Supercomputing Facilities of the Institute for Solid State Physics, University of Tokyo. The authors thank Satyam S. Jha for collaboration in the early stage of the work. M.T. also thanks Masanori Hanada, Chisa Hotta, and Norihiro Iizuka for valuable discussions.

-
- [1] P. Hayden and J. Preskill, Black holes as mirrors: quantum information in random subsystems, *J. High Energy Phys.* **0709**, 120 (2007), [arXiv:0708.4025 \[hep-th\]](#).
- [2] Y. Sekino and L. Susskind, Fast scramblers, *J. High Energy Phys.* **0810**, 065 (2008), [arXiv:0808.2096 \[hep-th\]](#).
- [3] L. Susskind, Addendum to Fast Scramblers (2011), [arXiv:1101.6048 \[hep-th\]](#).
- [4] N. Lashkari, D. Stanford, M. Hastings, T. Osborne, and P. Hayden, Towards the fast scrambling conjecture, *J. High Energy Phys.* **1304**, 022 (2013), [arXiv:1111.6580 \[hep-th\]](#).
- [5] S. H. Shenker and D. Stanford, Black holes and the butterfly effect, *J. High Energy Phys.* **1403**, 067 (2014), [arXiv:1306.0622 \[hep-th\]](#).
- [6] S. H. Shenker and D. Stanford, Stringy effects in scrambling, *J. High Energy Phys.* **1505**, 132 (2015), [arXiv:1412.6087 \[hep-th\]](#).
- [7] D. A. Roberts and D. Stanford, Diagnosing Chaos Using Four-Point Functions in Two-Dimensional Conformal Field Theory, *Phys. Rev. Lett.* **115**, 131603 (2015), [arXiv:1412.5123 \[hep-th\]](#).
- [8] D. A. Roberts and B. Yoshida, Chaos and complexity by design, *J. High Energy Phys.* **1704**, 121 (2017), [arXiv:1610.04903 \[quant-ph\]](#).
- [9] B. Yoshida, Soft mode and interior operator in the Hayden-Preskill thought experiment, *Phys. Rev. D* **100**, 086001 (2019), [arXiv:1812.07353 \[hep-th\]](#).
- [10] J. Liu, Scrambling and decoding the charged quantum information, *Phys. Rev. Res.* **2**, 043164 (2020), [arXiv:2003.11425 \[quant-ph\]](#).
- [11] A. Kitaev, “Hidden correlations in the Hawking radiation and thermal noise.”, talk at KITP (2015), <http://online.kitp.ucsb.edu/online/joint98/kitaev/>.
- [12] A. Kitaev, “A simple model of quan-

- tum holography.”, talks at KITP (2015), <http://online.kitp.ucsb.edu/online/entangled15/kitaev/> and <http://online.kitp.ucsb.edu/online/entangled15/kitaev2/>.
- [13] K. Jensen, Chaos in AdS₂ Holography, *Phys. Rev. Lett.* **117**, 111601 (2016), [arXiv:1605.06098](https://arxiv.org/abs/1605.06098) [hep-th].
- [14] J. Maldacena and D. Stanford, Remarks on the Sachdev-Ye-Kitaev model, *Phys. Rev. D* **94**, 106002 (2016), [arXiv:1604.07818](https://arxiv.org/abs/1604.07818) [hep-th].
- [15] S. Sachdev, Bekenstein-Hawking Entropy and Strange Metals, *Phys. Rev. X* **5**, 041025 (2015), [arXiv:1506.05111](https://arxiv.org/abs/1506.05111) [hep-th].
- [16] M. Blake, Universal Charge Diffusion and the Butterfly Effect in Holographic Theories, *Phys. Rev. Lett.* **117**, 091601 (2016), [arXiv:1603.08510](https://arxiv.org/abs/1603.08510) [hep-th].
- [17] C. W. von Keyserlingk, T. Rakovszky, F. Pollmann, and S. L. Sondhi, Operator Hydrodynamics, OTOCs, and Entanglement Growth in Systems without Conservation Laws, *Phys. Rev. X* **8**, 021013 (2018), [arXiv:1705.08910](https://arxiv.org/abs/1705.08910) [cond-mat.str-el].
- [18] V. Khemani, A. Vishwanath, and D. A. Huse, Operator Spreading and the Emergence of Dissipative Hydrodynamics under Unitary Evolution with Conservation Laws, *Phys. Rev. X* **8**, 031057 (2018), [arXiv:1710.09835](https://arxiv.org/abs/1710.09835) [cond-mat.stat-mech].
- [19] P. Hosur, X.-L. Qi, D. A. Roberts, and B. Yoshida, Chaos in quantum channels, *J. High Ener. Phys.* **1602**, 004 (2016), [arXiv:1511.04021](https://arxiv.org/abs/1511.04021) [hep-th].
- [20] F. Pastawski, B. Yoshida, D. Harlow, and J. Preskill, Holographic quantum error-correcting codes: toy models for the bulk/boundary correspondence, *J. High Energy Phys.* **1506**, 149 (2015), [arXiv:1503.06237](https://arxiv.org/abs/1503.06237) [hep-th].
- [21] F. Pastawski, J. Eisert, and H. Wilming, Towards Holography via Quantum Source-Channel Codes, *Phys. Rev. Lett.* **119**, 020501 (2017), [arXiv:1611.07528](https://arxiv.org/abs/1611.07528) [quant-ph].
- [22] T. Kohler and T. Cubitt, Toy models of holographic duality between local Hamiltonians, *J. High Energy Phys.* **1908**, 017 (2019), [arXiv:1810.08992](https://arxiv.org/abs/1810.08992) [hep-th].
- [23] P. Hayden and G. Penington, Learning the Alpha-bits of black holes, *J. High Energy Phys.* **1912**, 007 (2019), [arXiv:1807.06041](https://arxiv.org/abs/1807.06041) [hep-th].
- [24] Y. Nakata, E. Wakakuwa, and M. Koashi, Black holes as clouded mirrors: the Hayden-Preskill protocol with symmetry, *Quantum* **7**, 928 (2023), [arXiv:2007.00895](https://arxiv.org/abs/2007.00895) [quant-ph].
- [25] Y. Nakata, T. Matsuura, and M. Koashi, Constructing quantum decoders based on complementarity principle (2022), [arXiv:2210.06661](https://arxiv.org/abs/2210.06661) [quant-ph].
- [26] Y. Cheng, C. Liu, J. Guo, Y. Chen, P. Zhang, and H. Zhai, Realizing the Hayden-Preskill protocol with coupled Dicke models, *Phys. Rev. Res.* **2**, 043024 (2020), [arXiv:1909.12568](https://arxiv.org/abs/1909.12568) [cond-mat.quant-gas].
- [27] K. A. Landsman, C. Figgatt, T. Schuster, N. M. Linke, B. Yoshida, N. Y. Yao, and C. Monroe, Verified quantum information scrambling, *Nature* **567**, 61 (2019), [arXiv:1806.02807](https://arxiv.org/abs/1806.02807) [quant-ph].
- [28] A. R. Brown, H. Gharibyan, S. Leichenauer, H. W. Lin, S. Nezami, G. Salton, L. Susskind, B. Swingle, and M. Walter, Quantum Gravity in the Lab: Teleportation by Size and Traversable Wormholes (2019), [arXiv:1911.06314](https://arxiv.org/abs/1911.06314) [quant-ph].
- [29] S. Nezami, H. W. Lin, A. R. Brown, H. Gharibyan, S. Leichenauer, G. Salton, L. Susskind, B. Swingle, and M. Walter, Quantum Gravity in the Lab: Teleportation by Size and Traversable Wormholes, Part II (2021), [arXiv:2102.01064](https://arxiv.org/abs/2102.01064) [quant-ph].
- [30] M. M. Wilde, *Quantum Information Theory* (Cambridge University Press, 2013).
- [31] S. W. Hawking, Black hole explosions?, *Nature* **248**, 30 (1974).
- [32] S. W. Hawking, Particle creation by black holes, *Commun. Math. Phys.* **43**, 199 (1975).
- [33] S. W. Hawking, Breakdown of predictability in gravitational collapse, *Phys. Rev. D* **14**, 2460 (1976).
- [34] F. Haake, *Quantum Signatures of Chaos* (Springer, Berlin, Heidelberg, 2001).
- [35] M. Mehta, *Random Matrices* (Elsevier Science, 2004).
- [36] P. Caputa, J. Simón, A. Štikonas, T. Takayanagi, and K. Watanabe, Scrambling time from local perturbations of the eternal BTZ black hole, *J. High Energy Phys.* **1508** (8), 011, [arXiv:1503.08161](https://arxiv.org/abs/1503.08161) [hep-th].
- [37] L. Nie, M. Nozaki, S. Ryu, and M. T. Tan, Signature of quantum chaos in operator entanglement in 2d CFTs, *J. Stat. Mech.* **2019**, 093107 (2019), [arXiv:1812.00013](https://arxiv.org/abs/1812.00013) [hep-th].
- [38] K. Goto, M. Nozaki, S. Ryu, K. Tamaoka, and M. T. Tan, Scrambling and Recovery of Quantum Information in Inhomogeneous Quenches in Two-dimensional Conformal Field Theories (2023), [arXiv:2302.08009](https://arxiv.org/abs/2302.08009) [hep-th].
- [39] A. I. Larkin and Y. N. Ovchinnikov, Quasiclassical Method in the Theory of Superconductivity, *Sov. Phys. JETP* **28**, 1200 (1969).
- [40] H. Shen, P. Zhang, R. Fan, and H. Zhai, Out-of-Time-Order Correlation at a Quantum Phase Transition, *Phys. Rev. B* **96**, 054503 (2017), [arXiv:1608.02438](https://arxiv.org/abs/1608.02438) [cond-mat.quant-gas].
- [41] S. Sachdev and J. Ye, Gapless spin-fluid ground state in a random quantum Heisenberg magnet, *Phys. Rev. Lett.* **70**, 3339 (1993), [arXiv:cond-mat/9212030](https://arxiv.org/abs/cond-mat/9212030).
- [42] S. Sachdev, Holographic Metals and the Fractionalized Fermi Liquid, *Phys. Rev. Lett.* **105**, 151602 (2010), [arXiv:1006.3794](https://arxiv.org/abs/1006.3794) [hep-th].
- [43] J. S. Cotler, G. Gur-Ari, M. Hanada, J. Polchinski, P. Saad, S. H. Shenker, D. Stanford, A. Streicher, and M. Tezuka, Black Holes and Random Matrices, *J. High Energy Phys.* **1705**, 118 (2017), [arXiv:1611.04650](https://arxiv.org/abs/1611.04650) [hep-th].
- [44] G. Sarosi, AdS₂ holography and the SYK model, in *Proceedings of XIII Modave Summer School in Mathematical Physics — PoS(Modave2017)* (Sissa Medialab, 2018) [arXiv:1711.08482](https://arxiv.org/abs/1711.08482) [hep-th].
- [45] D. A. Trunin, Pedagogical introduction to the Sachdev–Ye–Kitaev model and two-dimensional dilaton gravity, *Physics-Uspekhi* **64**, 219 (2021), [arXiv:2002.12187](https://arxiv.org/abs/2002.12187) [hep-th].
- [46] J. Maldacena, S. H. Shenker, and D. Stanford, A bound on chaos, *J. High Ener. Phys.* **1608**, 106 (2016), [arXiv:1503.01409](https://arxiv.org/abs/1503.01409) [hep-th].
- [47] O. Bohigas, M. J. Giannoni, and C. Schmit, Characterization of chaotic quantum spectra and universality of level fluctuation laws, *Phys. Rev. Lett.* **52**, 1 (1984).
- [48] S. Sachdev, Statistical mechanics of strange metals and black holes (2022), [arXiv:2205.02285](https://arxiv.org/abs/2205.02285) [hep-th].
- [49] F. Monteiro, T. Micklitz, M. Tezuka, and A. Altland, Minimal model of many-body localization, *Phys. Rev. Res.* **3**, 013023 (2021), [arXiv:2005.12809](https://arxiv.org/abs/2005.12809) [cond-mat.str-el].
- [50] F. Monteiro, M. Tezuka, A. Altland, D. A. Huse, and T. Micklitz, Quantum Ergodicity in the Many-Body Lo-

- calization Problem, *Phys. Rev. Lett.* **127**, 030601 (2021), [arXiv:2012.07884 \[cond-mat.dis-nn\]](#).
- [51] V. Balasubramanian, A. Kar, C. Li, O. Parrikar, and H. Rajgadia, *Quantum Error Correction from Complexity in Brownian SYK* (2023), [arXiv:2301.07108 \[hep-th\]](#).
- [52] S. Xu and B. Swingle, *Scrambling Dynamics and Out-of-Time Ordered Correlators in Quantum Many-Body Systems: a Tutorial* (2022), [arXiv:2202.07060 \[quant-ph\]](#).
- [53] P. Hayden, M. Horodecki, A. Winter, and J. Yard, A Decoupling Approach to the Quantum Capacity, *Open Syst. Inf. Dyn.* **15**, 7 (2008), [arXiv:quant-ph/0702005](#).
- [54] F. Dupuis, M. Berta, J. Wullschleger, and R. Renner, One-Shot Decoupling, *Commun. Math. Phys.* **328**, 251 (2014), [arXiv:1012.6044 \[quant-ph\]](#).
- [55] F. Dupuis, *The decoupling approach to quantum information theory*, Ph.D. thesis, Université de Montréal (2009), [arXiv:1004.1641 \[quant-ph\]](#).
- [56] H. G. Katzgraber and F. Krzakala, Temperature and Disorder Chaos in Three-Dimensional Ising Spin Glasses, *Phys. Rev. Lett.* **98**, 017201 (2007), [arXiv:cond-mat/0606180 \[cond-mat.dis-nn\]](#).
- [57] G. Gur-Ari, R. Mahajan, and A. Vaezi, Does the SYK model have a spin glass phase?, *J. High Energ. Phys.* **1811**, 070 (2018).
- [58] H. Tajima and K. Saito, *Universal limitation of quantum information recovery: symmetry versus coherence* (2021).
- [59] S. Xu, L. Susskind, Y. Su, and B. Swingle, A Sparse Model of Quantum Holography (2020), [arXiv:2008.02303 \[cond-mat.str-el\]](#).
- [60] A. M. García-García, Y. Jia, D. Rosa, and J. J. M. Verbaarschot, Sparse Sachdev-Ye-Kitaev model, quantum chaos, and gravity duals, *Phys. Rev. D* **103**, 106002 (2021), [arXiv:2007.13837 \[hep-th\]](#).
- [61] M. Tezuka, O. Oktay, E. Rinaldi, M. Hanada, and F. Nori, Binary-coupling sparse Sachdev-Ye-Kitaev model: an improved model of quantum chaos and holography, *Phys. Rev. B* **107**, L081103 (2023), [arXiv:2208.12098 \[quant-ph\]](#).
- [62] A. M. García-García, B. Loureiro, A. Romero-Bermúdez, and M. Tezuka, Chaotic-Integrable Transition in the Sachdev-Ye-Kitaev Model, *Phys. Rev. Lett.* **120**, 241603 (2018), [arXiv:1707.02197 \[hep-th\]](#).
- [63] D. K. Nandy, T. Čadež, B. Dietz, A. Andreanov, and D. Rosa, Delayed thermalization in the mass-deformed Sachdev-Ye-Kitaev model, *Phys. Rev. B* **106**, 245147 (2022), [arXiv:2206.08599 \[cond-mat.str-el\]](#).
- [64] E. Wakakuwa and Y. Nakata, One-Shot Randomized and Nonrandomized Partial Decoupling, *Commun. Math. Phys.* **386**, 589 (2021), [arXiv:1903.05796 \[quant-ph\]](#).
- [65] J. Li, R. Fan, H. Wang, B. Ye, B. Zeng, H. Zhai, X. Peng, and J. Du, Measuring Out-of-Time-Order Correlators on a Nuclear Magnetic Resonance Quantum Simulator, *Phys. Rev. X* **7**, 031011 (2017), [arXiv:1609.01246 \[cond-mat.str-el\]](#).
- [66] R. Fan, P. Zhang, H. Shen, and H. Zhai, Out-of-time-order correlation for many-body localization, *Sci. Bull.* **62**, 707 (2017), [arXiv:1608.01914 \[cond-mat.quant-gas\]](#).
- [67] J. Riddell and E. S. Sørensen, Out-of-time ordered correlators and entanglement growth in the random-field XX spin chain, *Phys. Rev. B* **99**, 054205 (2019), [arXiv:1810.00038 \[cond-mat.stat-mech\]](#).
- [68] J. Lee, D. Kim, and D.-H. Kim, Typical growth behavior of the out-of-time-ordered commutator in many-body localized systems, *Phys. Rev. B* **99**, 184202 (2019), [arXiv:1812.00357 \[cond-mat.str-el\]](#).
- [69] S. Xu and B. Swingle, Accessing scrambling using matrix product operators, *Nature Physics* **16**, 199 (2020), [arXiv:1802.00801 \[quant-ph\]](#).
- [70] R. K. Shukla, A. Lakshminarayan, and S. K. Mishra, Out-of-time-order correlators of nonlocal block-spin and random observables in integrable and nonintegrable spin chains, *Phys. Rev. B* **105**, 224307 (2022), [arXiv:2203.05494 \[quant-ph\]](#).
- [71] I. Dumitriu and A. Edelman, Matrix models for beta ensembles, *J. Math. Phys.* **43**, 5830 (2002), [arXiv:math-ph/0206043](#).
- [72] L. F. Santos, Integrability of a disordered Heisenberg spin-1/2 chain, *J. Phys. A Math. Theor.* **37**, 4723 (2004), [arXiv:cond-mat/0310035](#).
- [73] K. Kudo and T. Deguchi, Level statistics of XXZ spin chains with a random magnetic field, *Phys. Rev. B* **69**, 132404 (2004), [arXiv:cond-mat/0310752 \[cond-mat.stat-mech\]](#).
- [74] L. Viola and W. G. Brown, Generalized entanglement as a framework for complex quantum systems: purity versus delocalization measures, *J. Phys. A Math. Theor.* **40**, 8109 (2007), [arXiv:quant-ph/0702014 \[quant-ph\]](#).
- [75] M. Žnidarič, T. Prosen, and P. Prelovšek, Many-body localization in the Heisenberg XXZ magnet in a random field, *Phys. Rev. B* **77**, 064426 (2008), [arXiv:0706.2539 \[quant-ph\]](#).
- [76] A. Pal and D. A. Huse, Many-body localization phase transition, *Phys. Rev. B* **82**, 174411 (2010), [arXiv:1010.1992 \[cond-mat.dis-nn\]](#).
- [77] A. D. Luca and A. Scardicchio, Ergodicity breaking in a model showing many-body localization, *EPL (Europhysics Letters)* **101**, 37003 (2013), [arXiv:1206.2342 \[cond-mat.str-el\]](#).
- [78] F. Alet and N. Laflorencie, Many-body localization: An introduction and selected topics, *C. R. Phys.* **19**, 498 (2018), [arXiv:1711.03145 \[cond-mat.str-el\]](#).
- [79] D. A. Abanin, E. Altman, I. Bloch, and M. Serbyn, Colloquium: Many-body localization, thermalization, and entanglement, *Rev. Mod. Phys.* **91**, 021001 (2019), [arXiv:1804.11065 \[cond-mat.dis-nn\]](#).
- [80] D. J. Luitz, N. Laflorencie, and F. Alet, Many-body localization edge in the random-field Heisenberg chain, *Phys. Rev. B* **91**, 081103 (2015), [arXiv:1411.0660 \[cond-mat.dis-nn\]](#).
- [81] A. Morningstar and D. A. Huse, Renormalization-group study of the many-body localization transition in one dimension, *Phys. Rev. B* **99**, 224205 (2019), [arXiv:1903.02001 \[cond-mat.stat-mech\]](#).
- [82] J. Šuntajs, J. Bonča, T. Prosen, and L. Vidmar, Quantum chaos challenges many-body localization, *Phys. Rev. E* **102**, 062144 (2020), [arXiv:1905.06345 \[cond-mat.str-el\]](#).
- [83] P. Sierant, D. Delande, and J. Zakrzewski, Thouless Time Analysis of Anderson and Many-Body Localization Transitions, *Phys. Rev. Lett.* **124**, 186601 (2020), [arXiv:1911.06221 \[cond-mat.dis-nn\]](#).
- [84] M. Kiefer-Emmanouilidis, R. Unanyan, M. Fleischhauer, and J. Sirker, Evidence for Unbounded Growth of the Number Entropy in Many-Body Localized Phases, *Phys. Rev. Lett.* **124**, 243601 (2020), [arXiv:2003.04849 \[cond-mat.dis-nn\]](#).
- [85] T. Chanda, P. Sierant, and J. Zakrzewski, Many-body localization transition in large quantum spin chains:

- The mobility edge, *Phys. Rev. Res.* **2**, 032045 (2020), [arXiv:2006.02860 \[cond-mat.dis-nn\]](#).
- [86] D. Sels and A. Polkovnikov, Dynamical obstruction to localization in a disordered spin chain, *Phys. Rev. E* **104**, 054105 (2021), [arXiv:2009.04501 \[quant-ph\]](#).
- [87] M. Kiefer-Emmanouilidis, R. Unanyan, M. Fleischhauer, and J. Sirker, Slow delocalization of particles in many-body localized phases, *Phys. Rev. B* **103**, 024203 (2021), [arXiv:2010.00565 \[cond-mat.dis-nn\]](#).
- [88] A. Morningstar, L. Colmenarez, V. Khemani, D. J. Luitz, and D. A. Huse, Avalanches and many-body resonances in many-body localized systems, *Phys. Rev. B* **105**, 174205 (2022), [arXiv:2107.05642 \[cond-mat.dis-nn\]](#).
- [89] D. Sels, Bath-induced delocalization in interacting disordered spin chains, *Phys. Rev. B* **106**, L020202 (2022), [arXiv:2108.10796 \[cond-mat.dis-nn\]](#).
- [90] P. Sierant and J. Zakrzewski, Challenges to observation of many-body localization, *Phys. Rev. B* **105**, 224203 (2022), [arXiv:2109.13608 \[cond-mat.dis-nn\]](#).
- [91] R. Ghosh and M. Žnidarič, Resonance-induced growth of number entropy in strongly disordered systems, *Phys. Rev. B* **105**, 144203 (2022), [arXiv:2112.12987 \[cond-mat.dis-nn\]](#).
- [92] M. C. Bañuls, J. I. Cirac, and M. B. Hastings, Strong and Weak Thermalization of Infinite Nonintegrable Quantum Systems, *Phys. Rev. Lett.* **106**, 050405 (2011), [arXiv:1007.3957 \[quant-ph\]](#).

Supplemental Materials: Hayden-Preskill Recovery in Hamiltonian Systems

S1. UPPER AND LOWER BOUNDS ON THE RECOVERY ERROR

In this section, we derive an upper bound on the recovery error in the Hayden-Preskill protocol based on the decoupling approach [53–55]. The recovery error in a Hamiltonian system is defined by

$$\Delta_{\hat{H}}(t, \beta) := \frac{1}{2} \min_{\mathcal{D}} \left\| \Phi^{AR} - \mathcal{D}^{CB' \rightarrow A}(\Psi_{\text{fin}}^{CB'R}(t)) \right\|_1, \quad (\text{S1})$$

where the minimum is taken over all CPTP maps \mathcal{D} from CB' to A , $\Psi_{\text{fin}}^{CB'R}(t, \beta) = \text{Tr}_D[\Psi_{\text{fin}}^{SB'R}(t, \beta)]$, and

$$|\Psi_{\text{fin}}(t, \beta)\rangle^{SB'R} = e^{-i\hat{H}_S t} (|\Phi\rangle^{AR} \otimes |\xi(\beta)\rangle^{BB'}). \quad (\text{S2})$$

Here, \hat{H}_S being the Hamiltonian in the system $S = AB = CD$ of N qubits. $|\xi(\beta)\rangle^{BB'}$ is a thermofield-double state in BB' of a thermal state $\xi(\beta)^B$ at the inverse temperature β .

Below, we show

$$1 - \sqrt{1 - (\Sigma_{\hat{H}}^{\text{opt}}(t, \beta))^2} \leq \Delta_{\hat{H}}(t, \beta) \leq \min\{1, \sqrt{2\Sigma_{\hat{H}}^{\text{opt}}(t, \beta)}\}. \quad (\text{S3})$$

Here, $\Sigma_{\hat{H}}^{\text{opt}}(t, \beta)$ is the degree of decoupling between D and R , defined by

$$\Sigma_{\hat{H}}^{\text{opt}}(t, \beta) := \frac{1}{2} \min_{\sigma} \left\| \Psi_{\text{fin}}^{DR}(t, \beta) - \sigma^D \otimes \pi^R \right\|_1, \quad (\text{S4})$$

where the minimization is taken over all quantum states on D , and $\Psi_{\text{fin}}^{DR}(t, \beta) = \text{Tr}_C[\Psi_{\text{fin}}^{CB'R}(t, \beta)]$. This implies that $\Sigma_{\hat{H}}^{\text{opt}}(t, \beta) \ll 1$ is essentially necessary and sufficient for the recovery error to be small. This is a well-known relation, but we show it for the sake of completeness. In the derivation, we do not explicitly write (t, β) .

We use the trace distance between ρ and σ by $T(\rho, \sigma)$, i.e.,

$$T(\rho, \sigma) := \frac{1}{2} \|\rho - \sigma\|_1. \quad (\text{S5})$$

For pure states $|\phi\rangle$ and $|\psi\rangle$, we denote the trace distance by $T(|\phi\rangle, |\psi\rangle) = \frac{1}{2} \|\phi\rangle\langle\phi| - |\psi\rangle\langle\psi|\|_1$ for simplicity. To obtain the lower bound, we use the relation between the trace distance and the fidelity:

$$1 - \sqrt{F(\rho, \sigma)} \leq T(\rho, \sigma) \leq \sqrt{1 - F(\rho, \sigma)}, \quad (\text{S6})$$

where $F(\rho, \sigma) := \|\sqrt{\rho}\sqrt{\sigma}\|_1^2$ is the fidelity. We also use the Uhlmann's theorem:

$$F(\rho^A, \sigma^A) = \max_V |\langle \rho|^{AB} V^{C \rightarrow B} |\sigma\rangle^{AC}|^2, \quad (\text{S7})$$

where the maximization is over all isometries $V^{C \rightarrow B}$ from C to B ($\dim \mathcal{H}_C \leq \dim \mathcal{H}^B$), and $|\rho\rangle^{AB}$ and $|\sigma\rangle^{AC}$ are purifications of ρ^A and σ^A , respectively.

The upper bound directly follows from the Uhlmann's theorem in terms of the trace norm (see, e.g., [55]). It states that, if there are two states ρ^A and σ^A such that $T(\rho, \sigma) \leq \epsilon$, then there exists systems B (C) that purifies ρ^A (σ^A) to a pure state $|\rho\rangle^{AB}$ ($|\sigma\rangle^{AC}$) and a partial isometry $V^{B \rightarrow C}$ that satisfies $T(V^{B \rightarrow C} |\rho\rangle^{AB}, |\sigma\rangle^{AC}) \leq \sqrt{2\epsilon}$.

Recall that a purification of $\Psi_{\text{fin}}^{DR}(t, \beta)$ is $|\Psi_{\text{fin}}(t, \beta)\rangle^{SB'R}$ and that that of π^R is $|\Phi\rangle^{\hat{A}R}$. Hence, denoting by $|\sigma\rangle^{DE}$ a purification of σ^D in Eq. (S4) by some system E , there is a partial isometry $V^{DB' \rightarrow \hat{A}E}$ such that

$$T(V^{DB' \rightarrow \hat{A}E} |\Psi_{\text{fin}}(t, \beta)\rangle^{SB'R}, |\sigma\rangle^{DE} \otimes |\Phi\rangle^{\hat{A}R}) \leq \sqrt{2\Sigma_{\hat{H}}^{\text{opt}}}. \quad (\text{S8})$$

By tracing out DE , we obtain

$$T(\mathcal{D}^{DB' \rightarrow \hat{A}} |\Psi_{\text{fin}}(t, \beta)\rangle^{CB'R}, |\Phi\rangle^{\hat{A}R}) \leq \sqrt{2\Sigma_{\hat{H}}^{\text{opt}}}, \quad (\text{S9})$$

where \mathcal{D} is a CPTP map obtained from the isometry V and the partial trace. As the left-hand side is exactly $\Delta_{\hat{H}}$, this provides an upper bound of the recovery error.

The lower bound in Eq. (S3) is obtained using Eqs. (S6) and (S7):

$$\Delta_{\hat{H}} := \min_{\mathcal{D}} T(\Phi^{AR}, \mathcal{D}^{CB' \rightarrow A}(\Psi_{\text{fin}}^{CB'R})) \quad (\text{S10})$$

$$\geq \min_{\mathcal{D}} \{1 - \sqrt{F(\Phi^{AR}, \mathcal{D}^{CB' \rightarrow A}(\Psi_{\text{fin}}^{CB'R}))}\} \quad (\text{S11})$$

$$= \min_{V, \sigma} \{1 - |\langle \Psi_{\text{fin}} |^{SB'R} V^{AS' \rightarrow B'C} (|\Phi\rangle^{AR} \otimes |\sigma\rangle^{DS'})|\} \quad (\text{S12})$$

$$\geq \min_{\sigma} \{1 - \sqrt{F(\Psi_{\text{fin}}^{DR}, \pi^R \otimes \sigma^D)}\} \quad (\text{S13})$$

$$\geq \min_{\sigma} \{1 - \sqrt{1 - (T(\Psi_{\text{fin}}^{DR}, \pi^R \otimes \sigma^D))^2}\} \quad (\text{S14})$$

$$\geq 1 - \sqrt{1 - (\Sigma_{\hat{H}}^{\text{opt}})^2}. \quad (\text{S15})$$

Here, the second and the second last lines follow from Eq. (S6), the third line from the Uhlmann's theorem, and the fourth line from the monotonicity of the fidelity under the partial trace.

The degree of decoupling $\Sigma_{\hat{H}}^{\text{opt}}$ is still computationally intractable in general due to the minimization over all states in D . However, it typically suffices to set the state σ in Eq. (S4) to Ψ_{fin}^D . Thus, we define

$$\Sigma_{\hat{H}}(t, \beta) := \frac{1}{2} \|\Psi_{\text{fin}}^{DR}(t, \beta) - \Psi_{\text{fin}}^D(t, \beta) \otimes \pi^R\|_1, \quad (\text{S16})$$

and investigate $\Sigma_{\hat{H}}(t, \beta)$ in our analysis. The corresponding upper bound is

$$\Delta_{\hat{H}}(t, \beta) \leq \bar{\Delta}_{\hat{H}}(t, \beta) := \min\{1, \sqrt{2\Sigma_{\hat{H}}(t, \beta)}\}. \quad (\text{S17})$$

The degree of decoupling $\Sigma_{\hat{H}}(t, \beta)$ can be characterized by the mutual information between D and R . To this end, we start with

$$\frac{2}{\ln 2} \delta^2 \leq I(X : Y)_\rho \leq 2\delta \log[\min\{d_X, d_Y\}] + (1 + \delta)h\left(\frac{\delta}{1 + \delta}\right), \quad (\text{S18})$$

where $\delta = \frac{1}{2} \|\rho^{XY} - \rho^X \otimes \rho^Y\|_1$, d_W is the dimension of the Hilbert space of W ($W = X, Y$), and $h(x) = -x \log x - (1 - x) \log(1 - x)$ is the binary entropy. See, e.g., [30]. Setting $X = D$ and $Y = R$, we have

$$\frac{2}{\ln 2} \Sigma_{\hat{H}}^2 \leq I(R : D)_{\Psi_{\text{fin}}(t, \beta)} \leq 2k\Sigma_{\hat{H}} + (1 + \Sigma_{\hat{H}})h\left(\frac{\Sigma_{\hat{H}}}{1 + \Sigma_{\hat{H}}}\right), \quad (\text{S19})$$

when $k \leq N - \ell$. This might be of independent importance since $I(D : R)_{\Psi_{\text{fin}}}$ has been studied as the OMI, especially for the infinite temperature ($\beta = 0$) [19, 36–38]. Hence, it shall be possible to quantitatively connect a number of results on the OMI with the recovery error in the Hayden-Preskill protocol. Note that, in relation to the Hayden-Preskill protocol, R is the system carrying quantum information and D corresponds to the subsystem that is traced out, which corresponds to the remaining black hole in the context of the information paradox. Hence, the size of the subsystems D and R should be k and $N - \ell$. This implies that the parameters of the OMI relevant to the Hayden-Preskill recovery is $k = |R| \ll |D| = N - \ell$, whereas it is sometimes set to $|R| \approx |D|$ in the study of the OMI.

From Eq. (S19), we also have

$$\Sigma_{\hat{H}}(t, \beta)^2 \leq \frac{\ln 2}{2} I(D : R)_{\Psi_{\text{fin}}} \quad (\text{S20})$$

$$= \frac{\ln 2}{2} (S(D)_{\Psi_{\text{fin}}} + S(R)_{\Psi_{\text{fin}}} - S(DR)_{\Psi_{\text{fin}}}) \quad (\text{S21})$$

$$= \frac{\ln 2}{2} (S(D)_{\Psi_{\text{fin}}} + k - S(CB')_{\Psi_{\text{fin}}}), \quad (\text{S22})$$

where we used in the last line the fact that $\Psi_{\text{fin}}^{SC'R}$ is a pure state. Hence, we have another upper bound on the recovery error in terms of the entropies:

$$\Delta_{\hat{H}}(t, \beta) \leq \min\left\{1, \left(2 \ln 2 (S(D)_{\Psi_{\text{fin}}} + k - S(CB')_{\Psi_{\text{fin}}})\right)^{1/4}\right\}. \quad (\text{S23})$$

S2. HAYDEN-PRESKILL RECOVERY BY A HAAR RANDOM DYNAMICS

To compute the recovery error when the unitary is given by a Haar random unitary, we consider the degree of decoupling $\Sigma_{\text{Haar}}(\beta)$ defined by

$$\Sigma_{\text{Haar}}(\beta) := \frac{1}{2} \|\Psi_{\text{fin}}^{DR}(t, \beta) - \Psi_{\text{fin}}^D(t, \beta) \otimes \pi^R\|_1, \quad (\text{S24})$$

where the unitary is Haar random. It is well known that this can be easily computed on average. The result in general cases is summarized to the so-called one-shot decoupling theorem [54], which in our case reads

$$\log[2\mathbb{E}_U[\Sigma_{\text{Haar}}(\beta)]] \leq -\frac{1}{2} \left(H_2(S|R)_{\xi \otimes \Phi} + H_2(\tilde{S}|D)_\tau \right), \quad (\text{S25})$$

where \mathbb{E}_U is the average over a Haar random unitary, and $H_2(X|Y)_\rho$ is the conditional collision entropy for the state ρ . Here and in the following, \log denotes logarithm to base 2 unless stated otherwise. The state $\tau^{\tilde{S}D}$ is defined by introducing a virtual system \tilde{S} that has the same structure of S . That is, \tilde{S} consists of N qubits and can be decomposed as $\tilde{C}\tilde{D}$. The state is given by

$$\tau^{S\tilde{D}} = \text{Tr}_{\tilde{C}}[|\Phi\rangle\langle\Phi|^{S\tilde{S}}] = \frac{I^C}{2^\ell} \otimes |\Phi\rangle\langle\Phi|^{D\tilde{D}}, \quad (\text{S26})$$

where $\Phi^{S\tilde{S}}$ is the maximally entangled state between S and \tilde{S} . The conditional collision entropy differs from the conditional von Neumann entropy in general, but when $\beta = 0, \infty$, they take the same values. Below, we denote it simply by $H(X|Y)_\rho$.

It is also known that a Haar random unitary shows the concentration of measure phenomena in the sense that, if a unitary is randomly drawn from the Haar measure, sufficiently continuous functions of the unitary take the values close to their average with high probability. We thus obtain the statement that, when U is a Haar random unitary,

$$\log[2\Sigma_{\text{Haar}}(\beta)] \leq -\frac{1}{2} \left(H(S|R)_{\xi \otimes \Phi} + H(S|D)_\tau \right), \quad (\text{S27})$$

with high probability. See, e.g., [55] for a more quantitative analysis.

The entropies are also easily computed in our case. Recalling that the conditional collision entropy satisfies the additivity regarding the tensor product and that $S = BA$, we have

$$H(S|R)_{\xi \otimes \Phi} = H(B)_\xi + H(A|R)_\Phi \quad (\text{S28})$$

$$= H(\beta) - k, \quad (\text{S29})$$

where we have used the notation $H(B)_\xi = H(\beta)$, as $\xi(\beta)$ is a thermal state. We have also used the fact that $H(A|R)_\Phi = -k$. On the other hand, using Eq. (S26), we have

$$H_2(S|\tilde{D})_\tau = \ell - (N - \ell) = 2\ell - N. \quad (\text{S30})$$

Note that C and D contain ℓ and $N - \ell$ qubits, respectively.

Combining these, we arrive at $\Sigma_{\text{Haar}}(\beta) \leq 2^{\ell_{\text{Haar,th}} - \ell}$ with high probability, where $\ell_{\text{Haar,th}} = \frac{1}{2}(N + k - H(\beta))$. Substituting this into Eq. (S17), we obtain

$$\Delta_{\text{Haar}}(\beta) \leq \bar{\Delta}_{\text{Haar}}(\beta) = 2^{\frac{1}{2}(\ell_{\text{Haar,th}} - \ell)}. \quad (\text{S31})$$

S3. HAYDEN-PRESKILL RECOVERY IN COMMUTING HAMILTONIAN MODELS

We provide a detailed analysis of the Hayden-Preskill protocol for commuting Hamiltonians. A commuting Hamiltonian is the one in the form of

$$\hat{H}_S = \hat{H}_A + \hat{H}_B + \hat{H}_{A:B}, \quad (\text{S32})$$

where \hat{H}_A and \hat{H}_B are Hamiltonians on A and B , respectively, and $\hat{H}_{A:B}$ is an interacting Hamiltonian between A and B that commutes with the other two. Below, we assume that $A \subset D$, which implies $C \subset B$ as $S = AB = CD$.

Due to the commuting condition, the time-evolving operator generated by \hat{H}_S is decomposed to $e^{-i\hat{H}_S t} = e^{-i\hat{H}_A t} e^{-i\hat{H}_B t} e^{-i\hat{H}_{A:B} t}$. Since a thermal state $\xi(\beta)^B \propto e^{-\beta\hat{H}_B}$ is invariant under the time-evolution by \hat{H}_B , we have $\Psi_{\text{fin}}^{DR}(t, \beta) = \text{Tr}_D [e^{-i\hat{H}_A t} e^{-i\hat{H}_{A:B} t} (\Phi^{AR} \otimes \xi(\beta)^B) e^{i\hat{H}_{A:B} t} e^{i\hat{H}_A t}]$. Since we have assumed $A \subset D$, it further reduces to $\Psi_{\text{fin}}^{DR}(t, \beta) = \text{Tr}_D [e^{-i\hat{H}_{A:B} t} (\Phi^{AR} \otimes \xi(\beta)^B) e^{i\hat{H}_{A:B} t}]$. Hence, we have

$$\Sigma_{\hat{H}}^{\text{opt}}(t, \beta) = \frac{1}{2} \min_{\sigma} \left\| \text{Tr}_C [e^{-i\hat{H}_{A:B} t} (\Phi^{AR} \otimes \xi(\beta)^B) e^{i\hat{H}_{A:B} t}] - \sigma^D \otimes \pi^R \right\|_1, \quad (\text{S33})$$

for commuting Hamiltonians.

Denote by \bar{B} an extended region of B in terms of $\hat{H}_{A:B}$, that is, a union of the system B and the set of qubits on which $\hat{H}_{A:B}$ acts non-trivially. We also denote the boundary of B in terms of $\hat{H}_{A:B}$ by $\partial\bar{B} := \bar{B} \setminus B$. By taking the trace over $\bar{B} \setminus C$ in the right-hand side Eq. (S33) and using the monotonicity of the trace norm under the partial trace, we have

$$\begin{aligned} \Sigma_{\hat{H}}^{\text{opt}}(t, \beta) &\geq \frac{1}{2} \min_{\sigma} \left\| \text{Tr}_{\bar{B}} [e^{-i\hat{H}_{A:B} t} (\Phi^{AR} \otimes \xi(\beta)^B) e^{i\hat{H}_{A:B} t}] - \text{Tr}_{D \cap \bar{B}} [\sigma^D] \otimes \pi^R \right\|_1 \\ &= \frac{1}{2} \min_{\sigma} \left\| \text{Tr}_{\partial\bar{B}} [\Phi^{AR}] - \sigma^{A \setminus \partial\bar{B}} \otimes \pi^R \right\|_1, \end{aligned} \quad (\text{S34})$$

where the first equality holds since the interaction Hamiltonian $\hat{H}_{A:B}$ nontrivially acts only within \bar{B} .

Since $\partial\bar{B} \subset A$, we have $\text{Tr}_{\partial\bar{B}} [\Phi^{AR}] = \Phi^{A-R_-} \otimes \pi^{R_+}$, where R is divided into $R_- R_+$, Φ^{A-R_-} is a maximally entangled state between $A_- := A \setminus \partial\bar{B}$ and R_- , and π^{R_+} is the completely mixed state in R_+ . Using κ , defined by the number of qubits in A that directly interact with B by $\hat{H}_{A:B}$, Φ^{A-R_-} is simply $(k-\kappa)$ EPR pairs. As $\pi^R = \pi^{R_-} \otimes \pi^{R_+}$, it follows that

$$\begin{aligned} \Sigma_{\hat{H}}^{\text{opt}}(t, \beta) &\geq \frac{1}{2} \min_{\sigma} \left\| (\Phi^{A-R_-} - \sigma^{A_-} \otimes \pi^{R_-}) \otimes \pi^{R_+} \right\|_1, \\ &= \frac{1}{2} \min_{\sigma} \left\| \Phi^{A-R_-} - \sigma^{A_-} \otimes \pi^{R_-} \right\|_1. \end{aligned} \quad (\text{S35})$$

Using Eq. (S6) and the fact that $F(\Phi^{A-R_-}, \sigma^{A_-} \otimes \pi^{R_-}) = 1/2^{2(k-\kappa)}$, we arrive at $\Sigma_{\hat{H}}^{\text{opt}}(t, \beta) \geq 1 - 2^{k-\kappa}$. Substituting this into the lower bound given in Eq. (S3), we have a lower bound on the recovery error as

$$\Delta_{\hat{H}} \geq 1 - 2^{(\kappa-k-1)/2}. \quad (\text{S36})$$

As $\kappa \leq k$, this implies that the recovery error $\Delta_{\hat{H}}$ is bounded from below by a constant.

We can improve the bound for specific Hamiltonians. Let us particularly consider the Sherrington-Kirkpatrick (SK) Hamiltonian given by $\hat{H}_{\text{SK}} := -\sum_{n < m} J_{nm} Z_n \otimes Z_m$, where J_{nm} is chosen from a given distribution. Similarly to Eq. (S32), we first divide the Hamiltonian into three:

$$\hat{H}_{\text{SK}} = \hat{H}_A + \hat{H}_B + \hat{H}_{A:B}, \quad (\text{S37})$$

where \hat{H}_A and \hat{H}_B nontrivially act only on A and B , respectively.

Using the fact that ξ^B is a thermal state of the Hamiltonian \hat{H}_B , the calculation same as Eq. (S33) leads to

$$\Sigma_{\hat{H}}^{\text{opt}}(t, \beta) = \frac{1}{2} \min_{\sigma} \left\| \text{Tr}_C [e^{-i\hat{H}_{A:B} t} (\Phi^{AR} \otimes \xi(\beta)^B) e^{i\hat{H}_{A:B} t}] - \sigma^D \otimes \pi^R \right\|_1. \quad (\text{S38})$$

We further decompose $\hat{H}_{A:B}$ in terms of the division of D and C as

$$\hat{H}_{A:B} = \hat{H}_{(A:B) \cap D} + \hat{H}_{A:C}, \quad (\text{S39})$$

where the former nontrivially acts only on D , while the latter acts on A and C as well as A and B . Again using the unitary invariance of the trace norm, it follows that

$$\Sigma_{\hat{H}}^{\text{opt}}(t, \beta) = \frac{1}{2} \min_{\sigma} \left\| \text{Tr}_C [e^{-i\hat{H}_{A:C} t} (\Phi^{AR} \otimes \xi(\beta)^B) e^{i\hat{H}_{A:C} t}] - \sigma^D \otimes \pi^R \right\|_1. \quad (\text{S40})$$

By applying a CPTP map onto R that maps ρ^R to $\sum_j \langle j | \rho^R | j \rangle |j\rangle\langle j|^R$, where $\{|j\rangle\}_j$ ($j = 0, \dots, 2^k - 1$) is the Pauli- Z basis in R , we have

$$\begin{aligned} \Sigma_{\hat{H}}^{\text{opt}}(t, \beta) &\frac{1}{2} \geq \min_{\sigma} \left\| \text{Tr}_C [e^{-i\hat{H}_{A:C} t} (\Omega^{AR} \otimes \xi(\beta)^B) e^{i\hat{H}_{A:C} t}] - \sigma^D \otimes \pi^R \right\|_1 \\ &\geq \frac{1}{2} \min_{\sigma} \left\| \text{Tr}_B [e^{-i\hat{H}_{A:C} t} (\Omega^{AR} \otimes \xi(\beta)^B) e^{i\hat{H}_{A:C} t}] - \sigma^A \otimes \pi^R \right\|_1, \end{aligned} \quad (\text{S41})$$

where $\Omega^{AR} := 2^{-k} \sum_j |j\rangle\langle j|^A \otimes |j\rangle\langle j|^R$. Note that π^R is invariant under the above CPTP map, and that the second inequality follows from $B \subset C$. We now use the relation that, for $n \in A$ and $m \in C$,

$$e^{iJ_{nm}Z_n \otimes Z_m t} (|j\rangle\langle j|^A \otimes \xi(\beta)^B) e^{-iJ_{nm}Z_n \otimes Z_m t} = |j\rangle\langle j|^A \otimes e^{(-1)^{j_n} iJ_{nm}Z_m} \xi(\beta)^B e^{-(-1)^{j_n} iJ_{nm}Z_m}, \quad (\text{S42})$$

where we express j in binary as $j_1 \dots j_k$ ($j_n = 0, 1$). The terms such as $e^{(-1)^{j_n} iJ_{nm}Z_m}$ then disappears when Tr_B is taken due to the cyclic property of the trace. Hence, it follows that

$$\text{Tr}_B [e^{-i\hat{H}_{A:C}t} (\Omega^{AR} \otimes \xi(\beta)^B) e^{i\hat{H}_{A:C}t}] = \Omega^{AR}. \quad (\text{S43})$$

Hence, we have

$$\Sigma_{\hat{H}}^{\text{opt}}(t, \beta) \geq \frac{1}{2} \min_{\sigma^A} \|\Omega^{AR} - \sigma^A \otimes \pi^R\|_1 \quad (\text{S44})$$

$$\geq \min_{\sigma^A} F(\Omega^{AR}, \sigma^A \otimes \pi^R) \quad (\text{S45})$$

$$\geq 1 - 2^{-k}, \quad (\text{S46})$$

where we have used the lower bound given in Eq. (S6) and the last line follows from the direct calculation. This leads to $\Delta_{\hat{H}_{\text{SK}}} \geq 1 - 2^{(1-k)/2}$.

S4. HAYDEN-PRESKILL RECOVERY IN PURE SYK₄

Here, we study the Hayden-Preskill recovery in SYK₄ in detail. The original SYK model with all-to-all four fermion interactions among $2N_q$ Majorana fermions is henceforth denoted as the SYK₄ model. The Hamiltonian is

$$\hat{H}_{\text{SYK}_4} = \sum_{1 \leq a_1 < a_2 < a_3 < a_4 \leq 2N_q} J_{a_1 a_2 a_3 a_4} \hat{\psi}_{a_1} \hat{\psi}_{a_2} \hat{\psi}_{a_3} \hat{\psi}_{a_4}, \quad (4)$$

in which the $2N_q$ Majorana fermions $\{\hat{\psi}_j\}_{j=1}^{2N_q}$ satisfy the anti-commutation relation $\{\hat{\psi}_j, \hat{\psi}_k\} = \hat{\psi}_j \hat{\psi}_k + \hat{\psi}_k \hat{\psi}_j = 2\delta_{jk}$, in which δ_{ij} is the Kronecker delta. The time-independent real couplings $J_{a_1 a_2 a_3 a_4}$ obey the Gaussian distribution $P(\{J_{a_1 a_2 a_3 a_4}\}) = \exp[-J_{a_1 a_2 a_3 a_4}^2 / 2\sigma^2] / \sqrt{2\pi\sigma^2}$ with the variance σ^2 . As $\text{Tr}[\hat{H}_{\text{SYK}_4}^2] = \sum_{1 \leq a_1 < a_2 < a_3 < a_4 \leq 2N} J_{a_1 a_2 a_3 a_4}^2$, we choose

$$\sigma^2 = \binom{2N_q}{4}^{-1} = \frac{12}{N(2N_q - 1)(2N_q - 2)(2N_q - 3)}, \quad (\text{S47})$$

so as to set the variance of the many-body energy eigenvalue to unity. The corresponding recovery error is denoted by $\bar{\Delta}_{\text{SYK}_4}(t, \beta)$.

A. Effectively canceling the parity symmetry

The SYK₄ model has symmetry, which divides the system into odd and even parity sectors. Accordingly, the time evolution operator is decomposed as

$$\hat{U}_{\text{SYK}_4}(t) = \hat{U}_{\text{SYK}_4, \text{odd}}(t) \oplus \hat{U}_{\text{SYK}_4, \text{even}}(t), \quad (\text{S48})$$

where $\hat{U}_{\text{SYK}_4, p}(t)$ ($p = \text{even}, \text{odd}$) is the unitary acting only on the sector with parity p . Since the presence of symmetry is known to induce drastic changes in the Hayden-Preskill recovery [9, 10, 24, 58], which we would like to ignore in this analysis, we provide a slight modification that allows us to effectively cancel the effect of symmetry.

The modification is to embed the system A of k qubits into an even-parity sector of a larger system A' , whose Hilbert space is $\mathcal{H}^{A'} := \mathcal{H}^a \otimes \mathcal{H}^A$ with \mathcal{H}^a being a two-dimensional space $\text{span}\{|o\rangle, |e\rangle\}$ that labels the parity in A' . For instance, when $k = 1$,

$$|e\rangle^a \otimes |0\rangle^A = |00\rangle^{A'}, |e\rangle^a \otimes |1\rangle^A = |11\rangle^{A'}, |o\rangle^a \otimes |0\rangle^A = |01\rangle^{A'}, \text{ and } |o\rangle^a \otimes |1\rangle^A = |10\rangle^{A'}. \quad (\text{S49})$$

The maximally entangled state between A and R is embedded to $|\Phi\rangle^{A'R} := |e\rangle^a \otimes |\Phi\rangle^{AR}$, which is a pure state in the 2^{2k+1} -dimensional system A' with Schmidt rank 2^k .

Embedding A into A' enlarges the whole system S to $S' := A'B$ of $N_q = N+1$ qubits, which we similarly decompose as $S' = sS$, where s is the two-dimensional space for labeling the parity of S . The time-evolution operator in S' by $\hat{U}_{\text{SYK}_4}^{S'}(t)$ is then given by

$$\hat{U}_{\text{SYK}_4}^{S'}(t) = |o\rangle\langle o|^s \otimes \hat{U}_{\text{SYK}_4, \text{odd}}^S(t) + |e\rangle\langle e|^s \otimes \hat{U}_{\text{SYK}_4, \text{even}}^S(t). \quad (\text{S50})$$

Assuming that the initial thermal state $\xi(\beta)^B$ is in the even-parity sector, which is 2^{N-k-1} dimensional, the state after the time evolution is

$$\begin{aligned} \Psi^{S'R}(t, \beta) &:= \hat{U}_{\text{SYK}_4}^{S'}(t)(\Phi^{A'R} \otimes \xi(\beta)^B)\hat{U}_{\text{SYK}_4}^{S'\dagger}(t) \\ &= |e\rangle\langle e|^s \otimes \hat{U}_{\text{SYK}_4, \text{even}}^S(t)(\Phi^{AR} \otimes \xi(\beta)^B)\hat{U}_{\text{SYK}_4, \text{even}}^{S\dagger}(t). \end{aligned} \quad (\text{S51})$$

Assuming that the information that the support of the state $\Psi^{S'R}(t, \beta)$ is the even-parity sector of $\mathcal{H}^{S'}$ is available in the decoding process, the state relevant to the degree of decoupling is the one in which s and C are traced over:

$$\begin{aligned} \Psi^{DR}(t, \beta) &= \text{Tr}_{s,C}[\Psi^{S'R}(t, \beta)] \\ &= \text{Tr}_C[\hat{U}_{\text{SYK}_4, \text{even}}^S(t)(\Phi^{AR} \otimes \xi(\beta)^B)\hat{U}_{\text{SYK}_4, \text{even}}^{S\dagger}(t)]. \end{aligned} \quad (\text{S52})$$

In this way, the effect of symmetry of the SYK₄ model is effectively canceled.

It should be emphasized that this modification leads to a slight change in the degree of decoupling, resulting in a slight change of the upper bound $\bar{\Delta}$ on the recovery error. To see this, we consider the degree of decoupling when $\hat{U}_{\text{SYK}_4, \text{even}}^S(t)$ in Eq. (S52) is replaced with the Haar random unitary U_{Haar}^S acting on the even-parity sector, whose dimension is 2^{N-k-1} . We denote it by Σ'_{Haar} . Since the system s , storing the parity information of S' , is traced out, the state τ in Eq. (S26) changes into the state in the system $S'\bar{D}$ such as

$$\tau^{S'\bar{D}} = \text{Tr}_{\bar{s}, \bar{C}}[|\Phi\rangle\langle\Phi|^{S'\bar{S}'}] = \frac{I^s}{2} \otimes \frac{I^C}{2^\ell} \otimes |\Phi\rangle\langle\Phi|^{D\bar{D}}. \quad (\text{S53})$$

By computing the conditional collision entropy of this state, we have

$$\log[2\Sigma'_{\text{Haar}}(\beta)] \leq \frac{N+k-H(\beta)-1}{2} - \ell. \quad (\text{S54})$$

This leads to

$$\Delta'_{\text{Haar}}(\beta) \leq 2^{\frac{1}{2}\left(\frac{N+k-H(\beta)-1}{2}-\ell\right)} = 2^{\frac{1}{2}\left(\frac{N+k-H(\beta)}{2}-\ell\right)-\frac{1}{4}}. \quad (\text{S55})$$

This differs from Eq. (3) by $2^{-\frac{1}{4}}$, which arises from the fact that the information about the parity sector is available in the decoding process. Note also that, since we have restricted the initial thermal state $\xi(\beta)^B$ on the even-parity sector, its entropy also changes. For instance, its maximum value is $N-k-1$ rather than $N-k$.

B. Recovery errors in the SYK₄ model

The collision entropy of $\xi_B(\beta)$ is

$$H(\beta) = -\log \text{Tr}[\rho(\beta)^2] = -\log \text{Tr} \frac{[e^{-2\beta H}]}{Z(\beta)^2} = -\log \frac{Z(2\beta)}{Z(\beta)^2} = 2 \log Z(\beta) - \log Z(2\beta) = \frac{1}{\ln 2}(F(2\beta) - 2F(\beta)), \quad (\text{S56})$$

in which $F(\beta) = -\ln Z(\beta)$ is the Helmholtz free energy. For $\beta = 0$, $Z(\beta = 0)$ is simply the dimension of the Hilbert space of B , and we have $H(\beta = 0) = \log Z(\beta = 0) = N-k-1$. Therefore, the right-hand side of (S55) is

$$2^{\frac{1}{2}\left(\frac{N+k-(N-k-1)}{2}-\ell\right)-\frac{1}{4}} = 2^{\frac{k-\ell}{2}}; \quad \bar{\Delta}'_{\text{Haar}}(\beta = 0) = \min\{1, 2^{\frac{k-\ell}{2}}\}. \quad (\text{S57})$$

In Fig. S1, we plot upper bound $\bar{\Delta}_{\text{SYK}_4}$ on the recovery error against time t for various ℓ , for $9 \leq N_q \leq 12$ and infinite temperature $\beta = 0$. Since the SYK₄ model has randomness of choosing the coupling constant $J_{a_1 a_2 a_3 a_4}$, we




FIG. S1. Semilogarithmic plot of the value of $\bar{\Delta}_{\text{SYK}_4}(t, \beta = 0)$ against t for $k = 1, N_q = 9, 10, 11, 12$ and $2 \leq \ell < N_q - k$. Average over 2^{19-N_q} samples is taken. For $\ell = 1$, $\bar{\Delta}_{\text{SYK}_4}(t, \beta = 0)$ is ~ 1 for all t . The dashed lines represent $\bar{\Delta}_{\text{Haar}}(\beta = 0) = 2^{\frac{1}{2}(1-\ell)}$ given in Eq. (3) for $\ell = 2, 3, \dots, N_q - 2$. See Fig. 2 for the plot for $N_q = 13$.




FIG. S2. Left: Semilogarithmic plot of the value of $\bar{\Delta}_{\text{SYK}_4}(t, \beta = 1)$ against t for $k = 1, N_q = 13$. For $\beta = 0$, see Fig. 2 in the main text. Center: Late-time value of $\bar{\Delta}_{\text{SYK}_4}(t, \beta)$ averaged over 64 samples plotted against ℓ for $N_q = 13$. Right: Late-time value $\bar{\Delta}_{\text{SYK}_4}(t, \beta)$ averaged over 64 samples plotted against β for $N_q = 13$. Dashed lines represent $\bar{\Delta}'_{\text{Haar}}(\beta)$.

take an average over many samples. We clearly observe that all curves rapidly decay and approach the bound $\bar{\Delta}_{\text{Haar}}$ for the Haar random dynamics, indicating that the Hayden-Preskill recovery is quickly achieved by the dynamics of SYK₄. This is also the case for finite temperature, as observed in Fig. S2. The late-time value of $\bar{\Delta}_{\text{SYK}_4}$ is in proportion to $2^{-H(\beta)/4}$ as expected from (S55) for $\beta \lesssim 10$, beyond which small deviations occur presumably due to the small number of eigenenergies within $1/\beta$ of the ground state.

Note that the SYK₄ model has a $N_q \bmod 4$ periodicity, depending on which the eigenenergy statistics resembles




FIG. S3. $\bar{\Delta}_{\hat{H}_{\text{SYK}_4}}(t, \beta = 0)$ plotted against $t/\sqrt{N_q}$, for $\ell = 4, 6, 8, 10$ from top to bottom. Average over 2^{19-N_q} samples is taken. The curves show better agreement as ℓ is increased, which is not observed for other choices of the exponent of N_q for scaling.

the Gaussian Unitary, Orthogonal, and Symplectic Ensembles (GUE, GOE, and GSE). We have checked that this periodicity does not affect much on the Hayden-Preskill recovery, implying that the Hayden-Preskill recovery is achieved by those ensembles of random Hermitian matrices. This might be of interest since the dynamics generated by GUE, GOE, and GSE is not Haar random, but achieves the Hayden-Preskill recovery that was shown based on a Haar random unitary.

C. Convergence time

To check the time scale needed for $\bar{\Delta}_{\text{SYK}_4}$ converging to $\bar{\Delta}_{\text{Haar}}$, we plot in Fig. S3 $\bar{\Delta}(t, \beta = 0)$ for $\ell = 4, 6, 8, 10$ against re-scaled time $t/\sqrt{N_q}$. The plot indicates that the convergence time is likely to be $\sqrt{N_q}$.

S5. HAYDEN-PRESKILL RECOVERY IN SPARSE SYK₄

We consider the sparse SYK₄ model [59–61], referred to as spSYK₄. The model has a parameter K_{cpl} that counts the number of non-zero coupling constant $J_{a_1 a_2 a_3 a_4}$ in Eq. (4). Other than that, the Hamiltonian of spSYK₄ is defined similarly to that of SYK₄. When $K_{\text{cpl}} = \binom{N_q}{4}$, then spSYK₄ reduces to SYK₄. Note that this approach to reducing the number of non-zero parameter is drastically different from the approach in [71], where tridiagonal matrix models generalizing the Gaussian and Wishart dense matrix models were introduced. The SYK₄ Hamiltonian is still sparse when presented as a matrix in the many-body Hilbert space.

For simplicity, we further introduce a constraint that a half of the non-zero couplings is set to $1/\sqrt{K_{\text{cpl}}}$ and the other half is $-1/\sqrt{K_{\text{cpl}}}$, which we call $\pm\text{spSYK}_4$ model. The corresponding upper bound of the recovery error is denoted by $\bar{\Delta}_{\pm\text{spSYK}_4}(t, \beta)$ for time t and the inverse temperature β .

The sparse SYK₄ model is known to have a transition from quantum chaos to integrability by varying the number K_{cpl} of non-zero coupling constant. For $K_{\text{cpl}} \ll 2N_q$, we typically have extra degeneracy in the spectrum than expected from the symmetry, and the spectral statistics of the distinct eigenvalues is not random-matrix like. For $K_{\text{cpl}} \gtrsim 2N_q$, such extra degeneracy disappears for practically all samples, and the spectral statistics strongly resembles that of the Gaussian random matrix ensemble with the corresponding symmetry [59–61]. It is surprising to some extent that the $\pm\text{spSYK}_4$ recovers important properties of SYK₄ when $K_{\text{cpl}} = O(N)$, which is by far smaller than $K_{\text{cpl}} = \binom{N_q}{4}$, which is needed for \hat{H}_{spSYK_4} reducing to \hat{H}_{SYK_4} .

In Fig. S4(a–b), we plot $\bar{\Delta}_{\pm\text{spSYK}_4}(t, \beta)$ for $\ell = 4, 10$ against t for various values of K_{cpl} . At an earlier time, $\bar{\Delta}_{\pm\text{spSYK}_4}(t, \beta)$ starts decreasing for any $K_{\text{cpl}}(> 0)$ as time t increases, which is similar to the SYK₄ model, but it soon witnesses the difference from SYK₄ when K_{cpl} is small. For $K_{\text{cpl}} \lesssim 16$, $\bar{\Delta}_{\pm\text{spSYK}_4}$ is unlikely to converge to the Haar value $\bar{\Delta}_{\text{Haar,th}}$ even in the large time limit, while for large K_{cpl} , $\bar{\Delta}_{\pm\text{spSYK}_4}$ seems to eventually converge to $\bar{\Delta}_{\text{Haar,th}}$ as t increases. Hence, we observe from these plots that at least two differences come up in the $\pm\text{spSYK}_4$




FIG. S4. The value of $\bar{\Delta}_{\pm\text{spSYK}_4}(t, \beta = 0)$ plotted against t for $k = 1$, $N_q = 13$ and $\ell = 4$ (left), $\ell = 10$ (center). 64 samples are used. The result for the SYK₄ model as well as the Haar value, $\bar{\Delta}_{\text{Haar}}(\beta = 0) = 2^{\frac{1-\ell}{2}}$, are also shown. (right) The late-time value of $\bar{\Delta}_{\pm\text{spSYK}_4}(t, \beta = 0)$ plotted for $N_q = 13$, $k = 1$, and various ℓ against the number K_{cpl} of nonzero couplings. 64 samples are used.

model in comparison with the SYK₄ model. One is the converging value of $\bar{\Delta}_{\pm\text{spSYK}_4}(t = \infty, \beta)$, and the other is the time scale for convergence.

To closely investigate the converging value, we plot in Fig. S4(c) the late-time value of $\bar{\Delta}_{\pm\text{spSYK}_4}$ for various ℓ against K_{cpl} for $2N_q = 26$ Majorana fermions. It is clear that the value is nearly identical to the Haar value for $K_{\text{cpl}} \gtrsim 30$. This convergence to the dense SYK limit is consistent with the eigenenergy statistics.[61] For the density of states plotted in the left panel of Fig. S5, we observe that while the overall shape stays similar, the fluctuation disappears as we increase K_{cpl} . The fluctuation observed for smaller K_{cpl} originates from the decrease in the number of distinct eigenvalues due to the additional degeneracy when emergent conserved quantities appear.[60, 61]

S6. HAYDEN-PRESKILL RECOVERY IN SYK₄₊₂

Because the trace of the product of six Majorana fermions is zero when four of them differ with each other, for the SYK₄₊₂ model defined in eq. (5), we have

$$\text{Tr} \hat{H}_{\text{SYK}_{4+2}}(\theta)^2 = (\cos^2 \theta) \text{Tr} \hat{H}_{\text{SYK}_4}^2 + (\sin^2 \theta) \text{Tr} \hat{H}_{\text{SYK}_2}^2 = 2^{N_q}. \quad (\text{S58})$$

Note that the normalization in [49] is so that $\hat{H}_{\text{SYK}_{4+2}}(\theta) = \hat{H}_{\text{SYK}_4} + \delta \hat{H}_{\text{SYK}_2}$ with the variance of the many-body eigenvalues for \hat{H}_{SYK_4} is $24(2N_q)^{-4} \binom{2N_q}{4} \sim 1$, and the variance of the *single-particle* eigenvalues for \hat{H}_{SYK_2} is $\delta^2(2N_q)^{-3}(2N_q - 2)(2N_q - 3) \sim \delta^2/(2N_q)$.

A. The density of states for SYK₄₊₂

We plot in the right panel of Fig. S5 the density of states for various values of $\tan \theta$. The variance of the eigenstate energy is fixed at unity. As $\tan \theta$ is increased, the peak becomes higher and the tails become thicker, changing from a RMT-like spectrum, which is a sign for quantum chaos and is the case for SYK₄, to a Gaussian-shape spectrum, which differs from SYK₄.

B. Numerical results for SYK₄₊₂

We here provide numerical results of the Hayden-Preskill recovery in the SYK₄₊₂ model. The Hamiltonian is given by

$$\begin{aligned} \hat{H}_{\text{SYK}_{4+2}}(\theta) &= (\cos \theta) \hat{H}_{\text{SYK}_4} + (\sin \theta) \hat{H}_{\text{SYK}_2} \\ &= (\cos \theta) \left(\hat{H}_{\text{SYK}_4} + (\tan \theta) \hat{H}_{\text{SYK}_2} \right), \end{aligned} \quad (\text{5})$$

which maps to the normalization in [49] by $\delta = \tan \theta$. The SYK₄₊₂ model trivially reduces to SYK₄ when $\delta = 0$. Below, we denote by $\bar{\Delta}_{\text{SYK}_{4+2}}(t, \beta)$ the corresponding upper bound on the recovery error and investigate it.




FIG. S5. The normalized density of states for (left) $\hat{H}_{\pm\text{spSYK}_4}$ plotted for $N_q = 13$ and 2048 samples, and for (right) $\hat{H}_{\text{SYK}_{4+2}}$ plotted for $N_q = 13$ and 64 samples.




FIG. S6. The value of $\bar{\Delta}_{\text{SYK}_{4+2}}(t, \beta = 0)$ plotted for $k = 1$ and various ℓ against the value of δ for $N_q = 9, 10, 11, 12$. $2^{19-N_q} = 2^{10}, 2^9, 2^8, 2^7$ samples are used and average over the values for $t = (1, 2, \dots, 10) \times 10^6$ is taken. The lines connecting the data points are guide to the eye.

By varying the strength δ of the SYK₂ term, this Hamiltonian shows a transition from quantum chaos, in the sense of eigenenergy statistics, to Fock space many-body localization (F-MBL) at $\delta_c \simeq N_q^2 \ln N_q$ [49], which corresponds to $\tan \theta = \mathcal{O}(10^2)$ for $9 \leq N_q \leq 13$ that we numerically study in this work.

In Fig. S6 we plot the late time behaviour of $\bar{\Delta}_{\text{SYK}_{4+2}}(t, \beta)$ by setting $t = 10^3$ and $\beta = 0$ as a function of the SYK₂ coupling strength $\delta = \tan \theta$. There are two characteristic values of $\delta = \tan \theta$, that is, $\delta_0 = \tan \theta_0 \approx 0.1$ and $\delta_1 = \tan \theta_1 \approx 3$, yielding three plateau-like shapes of $\bar{\Delta}_{\text{SYK}_{4+2}}$. When $0 \leq \theta \leq \theta_0$, $\bar{\Delta}_{\text{SYK}_{4+2}} \approx \bar{\Delta}_{\text{Haar}}$. This implies that the Hayden-Preskill recovery in the SYK₄ model is stable against small perturbations by the SYK₂ terms. When $\theta_0 \leq \theta \leq \theta_1$, $\bar{\Delta}_{\text{SYK}_{4+2}}$ starts deviating from the Haar value $\bar{\Delta}_{\text{Haar}}(\beta = 0)$ and increases linearly in terms of δ until it reaches the second plateau. The second plateau ends at $\theta = \theta_1$ and, for $\theta \geq \theta_1$, $\bar{\Delta}_{\text{SYK}_{4+2}}$ restarts increasing until the




FIG. S7. The value of $\bar{\Delta}$ for the XY model with $(L, \beta) =$ (a) $(8, 0)$, (b) $(10, 0)$, (c) $(12, 0)$, and (d) $(12, 1)$.

third plateau.

S7. HAYDEN-PRESKILL RECOVERY IN THE HEISENBERG MODEL WITH RANDOM MAGNETIC FIELD

We consider a one-dimensional quantum spin chain with site-dependent random magnetic field to the z direction,

$$\hat{H}_{\text{XXZ}} = \sum_{j=1}^{L-1} (S_j^x S_{j+1}^x + S_j^y S_{j+1}^y + J_z S_j^z S_{j+1}^z) + \sum_{j=1}^L h_j S_j^z, \quad (\text{S59})$$

in which L is the number of $S = 1/2$ spins, h_j are independently sampled from a uniform distribution in $[-W, W]$, J_z is the ratio of the coupling in the z direction to that in the xy plane.

A. Integrable case

For $J_z = 0$ and $h_j = 0$ ($j = 1, 2, \dots, L$), the model is integrable. In Fig. S7, we plot the time dependence of $\bar{\Delta}$ for the XY model ($J = 0$) without random magnetic field. The plots at $\beta = 0$ for different $L = 8, 10, 12$ are qualitatively similar to each other, and they do not converge to a constant because we are here working with a single, integrable Hamiltonian for each L . Introducing finite temperature, $\beta = 1$, increases the value of $\bar{\Delta}$, reflecting the decrease of the effective dimension of the Hilbert space.

B. Chaotic and many-body localized cases

For $J_z = 1$ and $W > 0$, the model has been extensively studied as a prototypical model of many-body localization in one spatial dimension [72–79]. For system sizes accessible by numerical diagonalization, various measures of localization point to the MBL transition at finite critical W [80]. We note that the location of the transition to the genuine MBL phase in the thermodynamic limit have been heavily debated in more recent studies.[81–91]

In Fig. S8, we plot the time dependence of $\bar{\Delta}$ for the XXZ model with $J = 1$. For $W = 0$, this is the Heisenberg model, which is integrable. Again, the plots are not converging. For $W = 1$, the average over 128 samples is plotted. The late-time behavior of the averaged value of $\bar{\Delta}$ is smoother compared to the $W = 0$ case. The value decreases as l is increased. (For $l = 9 = L - 1$, we always obtain $\bar{\Delta} = 0$, because the system is in the $S^z = 0$ subspace.)

In Fig. S9, we plot the sample average of $\bar{\Delta}$ for various values of l and W . $\bar{\Delta}$ monotonically increases and approaches unity as W is increased.

We numerically observe that as W is increased, $\bar{\Delta}$ monotonically increases. Even though a small W introduces a chaotic behavior to the integrable spin chain at $W = 0$, we conclude this does not lead to successful quantum error correction.




FIG. S8. The value of $\bar{\Delta}$ for the XXZ model with $J_z = 1$, $L = 10$ and $(W, \beta) =$ (a) $(0, 0)$, (b) $(0, 1)$, (c) $(1, 0)$, and (d) $(1, 1)$. For (c) and (d), averages over 16 samples are plotted for $W > 0$.




FIG. S9. The value of $\bar{\Delta}$ for the XXZ model with $J_z = 1$, $L = 12$, $\beta = 0$ for various values of W and (a) $\ell = 1$, (b) $\ell = 4$, (c) $\ell = 7$, and (d) $\ell = 10$. Averages over 16 samples are plotted for $W > 0$.

S8. ISING MODEL WITH CHAOTIC SPIN CHAIN

A translationally invariant spin chain with nearest-neighbor Ising-type interaction and magnetic field,

$$\hat{H}_{\text{Ising}} = - \sum_{j=1}^{L-1} (S_j^z S_{j+1}^z) - g \sum_{j=1}^L S_j^x - h \sum_{j=1}^L S_j^z, \quad (\text{S60})$$

is exactly solvable if $g = 0$ or $h = 0$. For other choices of (g, h) , the model is non-integrable, and the case with $(g, h) = (-1.05, 0.5)$ [92] has been often studied as a prototypical model of chaotic spin chain. In Fig. S10 we plot the value of Δ for $\beta = 0$ and various ℓ . We observe that while Δ decreases as ℓ is increased, the decrease is very slow. For finite β , the value of Δ is generally larger, indicating less efficient error correction.




FIG. S10. Ising spin chain, open boundary condition, $L = 12$, $(g, h) = (-1.05, 0.5)$.