

# ADAPTIVE INTERVENTIONS FOR GLOBAL HEALTH: A CASE STUDY OF MALARIA

**África Perriñez**

benshi.ai  
Barcelona, Spain  
africa@benshi.ai

**Andrew Trister**

Bill and Melinda Gates Foundation  
Seattle, USA  
andrew.trister@gatesfoundation.org

**Madhav Nekkar & Ana Fernández del Río**

benshi.ai  
Barcelona, Spain  
{madhav, ana}@benshi.ai

**Pedro L. Alonso**

Faculty of Medicine, Universidad de Barcelona  
Barcelona, Spain  
alonso@ub.edu

## ABSTRACT

Malaria can be prevented, diagnosed, and treated; however, every year, there are more than 200 million cases and 200,000 preventable deaths. Malaria remains a pressing public health concern in low- and middle-income countries, especially in sub-Saharan Africa. We describe how by means of mobile health applications, machine-learning-based adaptive interventions can strengthen malaria surveillance and treatment adherence, increase testing, measure provider skills and quality of care, improve public health by supporting front-line workers and patients (e.g., by capacity building and encouraging behavioral changes, like using bed nets), reduce test stockouts in pharmacies and clinics and informing public health for policy intervention.

## 1 INTRODUCTION

Not everyone benefits equally from the same treatment, has the same environment, or can receive therapy at the same time. Adaptive interventions consider human differences as treatment begins and during its course to ensure each person gets the treatment that continuously works for them. Interventions are adaptive in that they adapt to the individual (providing personalization) and their evolving context and needs (i.e., adjusting dynamically).

Both clinical and patient actions are essential to high-quality, low-cost, effective healthcare. Human behavior constitutes the primary mode for activating health improvements, including decisions made by clinicians and their patients. In medical practice, clinicians can be gently nudged to improve decision-making by providing a decision architecture in which optimal default clinical actions are suggested. Outside controlled clinical environments (e.g., intensive care units), patient actions (e.g., lifestyle choices and treatment adherence) primarily determine health outcomes. Mobile health is key to promoting continuous and proactive healthcare monitoring. It can, for example, mitigate the lack of medication adherence, which can be low and challenging to address, as patient actions control healthcare delivery outside medical environments.

The data generated by the users of digital applications are instrumental in determining past and current behaviors and predicting future conduct. This information can be used to deliver personalized mobile-mediated interventions. Such interventions aim to provide the appropriate type or degree of support by adapting to an individual's changing internal and contextual states (Nahum-Shani et al., 2018; Menictas et al., 2019; Carpenter et al., 2020; Bidargaddi.N amd Schrader et al., 2020; Coppersmith et al., 2021).

Mobile health applications also serve as a direct channel of communication with their users, from front-line healthcare workers to patients and the general public. Interventions and incentives can be delivered directly to the users through their phones. Machine learning (ML) can help generate predictions regarding the behavior of app users, health outcomes, and their contexts. These predictions

can be integrated with real-time information pertaining to the users’ choices and circumstances to determine the individuals or groups needing additional support. Moreover, these predictions can be used to personalize the timing of the interventions delivered to each user. Similarly, reliable predictions regarding the evolution or variations in the demand for different medical supplies can help establish reminder and suggestion systems that can help ensure that all the supply chain actors (e.g., pharmacists) maintain adequate stocks of essential supplies at all times. Motivational prompts, personalized reminders, incentives, engaging elements (such as game-designed elements), and other nudges can be used to boost and reinforce good practices (Hrnjic & Tomczak, 2019).

Adaptive interventions in healthcare have primarily been deployed in resource-rich countries (Nahum-Shani et al., 2018; Hardeman et al., 2019). Given the immense disease burdens borne by low- and middle-income countries and the increasing smartphone penetration in these regions, the use of adaptive interventions to improve health outcomes may be highly beneficial to global health. In this research, we explore this potential based on a case study of malaria. A discussion of the ML methods can be found in the appendix A.

## 2 MALARIA: A CASE STUDY

Malaria, caused by the parasite *Plasmodium falciparum*, remains a deadly illness across much of the Global South.

Malaria in pregnancy and infancy is a major public health concern and a key driver of maternal and newborn mortality (Tarning, 2016). In 2020, nearly 630,000 deaths worldwide were caused by malaria, a disproportionate share of which (96%) occurred in Africa (Organization, 2022). In Sub-Saharan Africa, malaria is responsible for 12% of all child fatalities (Roser & Ritchie, 2019), and children under five account for approximately 80% of all malaria deaths (Organization, 2022).

Although global malaria incidence and mortality have substantially decreased in the recent two decades, progress stalled since 2015 (Noor & Alonso, 2022), and researchers worry that the COVID-19 pandemic has disrupted malaria intervention coverage, reversing these gains (Weiss et al., 2021).

### 2.1 SURVEILLANCE: PROGRESS TOWARDS ELIMINATION

Even with the World Health Organization’s (WHO) historic approval of a malaria vaccine (Li et al., 2022), senior leaders caution that a broader approach will be needed for malaria eradication, including improved collection and usage of high-quality data—from health-management information systems and electronic databases to geospatial models (Alonso, 2021)—and flexible evaluation and implementation of interventions by local decision-makers (Alonso & O’Brien, 2022). According to a global landscape review, malaria surveillance systems in 2015—2016 were “insufficient to support the planning and implementing of targeted interventions and measure progress toward malaria elimination” (Lourenço et al., 2019). Such elimination efforts require the accurate notification of individual cases within 24 h of diagnosis to provide timely and targeted responses, which essentially represent adaptive interventions (WHO, 2015).

Notably, such malaria surveillance frameworks must be integrated, data-driven, tailored, and based on mobile platforms. Mobile health for malaria surveillance, that uses a combination of message- and application-based reporting, can support health workers and clinicians in recording malaria case information (Githinji et al., 2014; Baliga et al., 2019; RTI, 2020; Oo et al., 2021; Bhowmick et al., 2021), decreasing delays in case reporting to health officials, and improving the quality of data collection. Data fragmentation remains a barrier to a cohesive malaria response, and previously siloed data from diverse sources (private/public healthcare providers, government agencies, etc.) must be integrated to create a real-time, case-based malaria surveillance system (Rahi & Sharma, 2020). Automation is essential, as analytics related to threat monitoring, requirement identification, and system performance must be readily available for decision-makers (Ohrt et al., 2015).

A comprehensive malaria surveillance system can inform policy interventions, e.g., the allocation of mosquito nets, tests, and antimalarials can be targeted to favor individuals and communities in need. By understanding the changes in people’s mobility and clustering, the impact of non-pharmaceutical interventions can be evaluated, and geographic areas in which additional actions may be helpful can be identified (Grantz et al., 2020). Several studies revealed that the characterization of travel patterns

through geolocation data and their combination with contextual information on malaria incidence can inform strategies to target travelers and reduce transmission (Milusheva, 2020).

## 2.2 PREVENTION

For pregnant women visiting institutions for antenatal care, the WHO recommends the administration of intermittent preventive treatment with sulfadoxine–pyrimethamine and distribution of long-lasting insecticide-treated mosquito nets (LLINs) (Salomão et al., 2017).

Unfortunately, many pregnant women in low- and middle-income countries do not receive either intervention owing to stock-outs in data-fragmented health systems (Salomão et al., 2017). Accurate demand forecasts that can integrate real-time contextual information are needed to ensure equitable and efficient allocation of important preventive goods. For example, by building geospatial models and combining data from net manufacturers, national programs, and cross-sectional household surveys, researchers can develop detailed maps of LLIN access and usage (Bertozzi-Villa et al., 2021). These analyses can also be automated to identify geographic areas in which additional supplies or different actions may be required.

Patient behavior also determines the efficacy of the interventions, as many pregnant women fail to take their pills (Mubyazi et al., 2005) or do not use bed nets for various practical reasons (Manu et al., 2017; Gultie et al., 2020). These aspects highlight the importance of communication to encourage behavioral changes (Ricotta et al., 2014).

For children aged 3–59 months, WHO recommends seasonal malaria chemoprevention (SMC) during the months of peak malaria transmission. Although SMC effectively controls malaria and reduces hospitalizations (Diawara et al., 2017; Baba et al., 2020; Issiaka et al., 2020; Cairns et al., 2021), clinical data and pharmacokinetic analyses reveal that complete adherence to treatment is observed in fewer than 20% of children outside the study setting (Ding et al., 2020). Targeted interventions for behavioral changes are required to improve the real-world effectiveness of SMC.

Caregiver-targeted message-based interventions to increase preventive health behaviors (such as sleeping under a net) have been noted to be successful in decreasing the malaria prevalence in children under the age of five (Mohammed et al., 2019). Despite the vast potential of mobile health solutions, one-size-fits-all interventions are typically implemented in which a standard message is sent to all participants. There exists an enormous opportunity for the delivery of personalized interventions that can appropriately incentivize a given user and guide targeted public health campaigns, which may include gamification and leveraging of individual social networks (Ernst et al., 2017). As a potential example, researchers recently tested an ML model in conjunction with an accelerometer-based approach for measuring a range of LLIN use behaviors: Although these technologies represent a proof of concept at present, they can support the implementation of financial incentives based on granular LLIN-use monitoring over longer time periods (Koudou et al., 2022).

The only way to understand which interventions work best and which incentives drive behavioral change is by running experiments on the ground (Banerjee et al., 2010; Bates et al., 2012; Zhou et al., 2020a; 2021).

## 2.3 QUALITY OF CARE

### 2.3.1 DIAGNOSIS

Malaria is one of the most under-diagnosed and over-treated diseases. The potential for severe outcomes means that any patient with a fever (especially a child) may be administered treatment for malaria, often without having received a diagnostic test (Amankwa et al., 2019; Ajibaye et al., 2019; Boadu et al., 2016; Beisel et al., 2016). The administration of drugs without a conclusive test may accelerate antimalarial resistance, an increasingly worrying problem according to WHO malaria experts (Rasmussen et al., 2022).

Although microscopy is the gold-standard for malaria diagnosis, its implementation remains infeasible in many resource-constrained settings (Beisel et al., 2016). Consequently, many stakeholders have turned to rapid diagnostic tests (RDTs). However, their access remains limited, with frequent stock-outs (Boadu et al., 2016; Blanas et al., 2013) that may be related to inaccurate record-keeping.

Digital tools and demand forecasting algorithms can alleviate this problem, as has been proved in India with an app that included a supply chain management component (Rajvanshi et al., 2021).

Furthermore, many lay community health workers struggle to appropriately perform RDTs (Boadu et al., 2016; Blanas et al., 2013; Beisel et al., 2016), and thus, capacity-building efforts to increase their skills are urgently required. Digital monitoring of the health workers' performance can serve as an effective quality control strategy and a mode of delivering feedback (Laktabai et al., 2018). Personalized digital nudges can direct the health workers who need additional support to specific online learning resources or encourage them to sign up for an in-person training session.

Patient-facing steering may also be required, as sick patients who receive a negative test may still expect treatment. Patient and provider education around antimicrobial resistance may help, in addition to increasing awareness and clinical decision support for the management of other febrile illnesses.

### 2.3.2 ARTEMISININ COMBINATION THERAPY (ACT)

ACT is the first-line malaria treatment throughout most of the malaria-endemic world. Notable issues include stock-outs (Blanas et al., 2013; Rowe et al., 2009; O'Connell et al., 2011) and the distribution of low-quality antimalarials, especially in urban areas Newton et al. (2017). Most antimalarials are distributed through the private sector, where non-artemisinin therapies are prevalent and 5–24 times less expensive than quality-assured ACT (O'Connell et al., 2011). In this context, it is important to strengthen the supply chain with a data-driven approach, potentially by using demand forecasting algorithms to send appropriately timed reminders to pharmacists to ensure that they restock necessary supplies.

The quality of care and clinical management of malaria remains widely varying and substandard. In Sub-Saharan Africa, less than one-third of the children diagnosed with malaria receive both a blood test diagnosis and appropriate antimalarial treatment (Cohen et al., 2020). Evidence for gaps in the community health workers' and drug dispensers' ability to appropriately manage malaria has been found in several countries (Rowe et al., 2009; Blanas et al., 2013; Chowdhury et al., 2020; Buabeng, 2010; Kamuhabwa & Silumbe, 2013). Capacity-building efforts are clearly necessary, and adaptive interventions can facilitate the assignment of appropriate content to each worker (Katsaris et al., 2021). AI-based user segmentation and behavioral phenotyping can yield highly granular cohorts to be focused on. For example, we research the capacity development of midwives by predicting their demand for specific contents (Guitart et al., 2021a) and building recommendation systems for an e-learning app by predicting the chance that a given user clicks on a certain item (Guitart et al., 2021b).

Lastly, patient adherence to antimalarial regimes is a notable issue, with estimates suggesting that it ranges from 40% to 65% (Amponsah et al., 2015; Mace et al., 2011; Onyango et al., 2012). Trials of message-based-reminders to healthcare workers (Zurovac et al., 2011; Kaunda-Khangamwa et al., 2018) and patients (Macías Saint-Gerons et al., 2022) have shown promise while highlighting the necessity of personalized interventions involving relevant and actionable messages (Buabeng, 2010). In the case of tuberculosis, where adherence is also a problem, researchers have noted that an ML-driven approach to predict TB adherence risk in conjunction with adaptive interventions dynamically increases the odds ratio of next-day treatment adherence verification by 35% (Boutillier et al., 2021).

### 2.3.3 PRE-REFERRAL TREATMENT

Another crucial area where malaria care delivery can be strengthened is pre-referral treatment, usually rectal artesunate (RAS), recommended by WHO for young children with suspected severe malaria when injections are not available (WHO, 2022). Although the effectiveness of RAS has been demonstrated in placebo-controlled trials (Gomes et al., 2009), more recent studies have indicated that its introduction in real-world conditions may increase the malaria case-fatality rates (WHO, 2022). This phenomenon is attributable to several reasons, such as low patient adherence to referral guidelines (Simba et al., 2010) and inadequate ability of health workers to deliver pre-referral care (Amboko et al., 2022) appropriately. Patient- and provider-facing steering and capacity building are needed to strengthen health systems and ensure that guidelines are followed.

### 3 CONCLUSIONS

Combating the devastating toll of malaria requires multifaceted and innovative strategies. Technology alone is not a panacea, and no simple solution exists. However, ensuring high-quality data collection from diverse sources, such as mobile phones, supply chains, public surveys, and electronic health records, can play a crucial step. These data can be integrated to obtain intrinsic and contextual information that can drive personalized and adaptive interventions.

The use cases are vast: Reminding antenatal care clinics to stock up on LLINs; providing capacity-building resources to struggling community health workers; supporting a district health officer in responding to a new malaria outbreak; incentivizing pharmacists to administer diagnostic tests before prescribing treatments; or encouraging patients to adhere to a treatment regime.

#### ACKNOWLEDGMENTS

This work was supported, in whole or in part, by the Bill & Melinda Gates Foundation INV-022480. Under the grant conditions of the Foundation, a Creative Commons Attribution 4.0 Generic License has been assigned to the Author Accepted Manuscript version that might arise from this submission.

#### REFERENCES

- Olusola Ajibaye, Emmanuel O Balogun, Yetunde A Olukosi, Bassey A Orok, Kolapo M Oyebola, Bamidele A Iwalokun, Olugbenga O Aina, Olalere Shittu, Adeniyi K Adeneye, Oyesola O Ojewunmi, K Kita, and Samson T Awolola. Impact of training of mothers, drug shop attendants and voluntary health workers on effective diagnosis and treatment of malaria in lagos, nigeria. *Tropical Parasitology*, 9(1):36–44, 2019. doi: 10.4103/tp.TP\_36\_1810.1186/s12936-016-1174-0.
- Pedro L Alonso. Malaria: a problem to be solved and a time to be bold. *Nature Medicine*, 27(9): 1506–1509, 2021. doi: 10.1038/s41591-021-01492-6.
- Pedro L Alonso and Katherine L O’Brien. A malaria vaccine for africa—an important step in a century-long quest. *New England Journal of Medicine*, 386(11):1005–1007, 2022. doi: 10.1056/NEJMp2116591.
- Charles Enyaah Amankwa, Harriet Affran Bonful, Kofi Agyabeng, and Priscillia A. Nortey. Dispensing practices for anti-malarials in the la nkwantanang-madina municipality, greater accra, ghana: a cross-sectional study. *Malaria Journal*, 18:260, 2019. doi: 10.1186/s12936-019-2897-5.
- Beatrice Amboko, Beatrice Machini, George Githuka, Philip Bejon, Dejan Zurovac, and Robert W Snow. Readiness of the kenyan public health sector to provide pre-referral care for severe paediatric malaria. *Tropical Medicine & International Health*, 27(3):330–336, 2022. doi: 10.1111/tmi.13728.
- Alexandria O. Amponsah, Helen Vosper, and Afia F. A. Marfo. Patient related factors affecting adherence to antimalarial medication in an urban estate in ghana. *Malaria Research and Treatment*, 2015:1–8, 2015. doi: 10.1155/2015/452539.
- Ebenezer Baba, Prudence Hamade, Harriet Kivumbi, Maddy Marasciulo, Kolawole Maxwell, Diego Moroso, Arantxa Roca-Feltrer, Adama Sanogo, Joanna Stenstrom Johansson, James Tibenderana, et al. Effectiveness of seasonal malaria chemoprevention at scale in west and central africa: an observational study. *The Lancet*, 396(10265):1829–1840, 2020. doi: 10.1016/S0140-6736(20)32227-3.
- B Shantharam Baliga, Animesh Jain, Naren Koduvattat, BG Kumar, Manu Kumar, Arun Kumar, and Susanta K Ghosh. Indigenously developed digital handheld android-based geographic information system (gis)-tagged tablets (tabs) in malaria elimination programme in mangaluru city, karnataka, india. *Malaria journal*, 18:444, 2019. doi: 10.1186/s12936-019-3080-8.
- Abhijit Vinayak Banerjee, Esther Duflo, Rachel Glennerster, and Dhruva Kothari. Improving immunisation coverage in rural india: clustered randomised controlled evaluation of immunisation campaigns with and without incentives. *BMJ*, 340:c2220, 2010. doi: 10.1136/bmj.c2220.

- Andrea Barraza-Urbina and Dorota Glowacka. Introduction to bandits in recommender systems. In *Fourteenth ACM Conference on Recommender Systems, RecSys '20*, pp. 748–750, New York, 2020. Association for Computing Machinery. ISBN 9781450375832. doi: 10.1145/3383313.3411547. URL <https://doi.org/10.1145/3383313.3411547>.
- Mary Ann Bates, Rachel Glennerster, Kamilla Gumedde, and Esther Dufflo. The price is wrong. *Field Actions Science Reports*, Special Issue 4, 2012. URL <https://journals.openedition.org/factsreports/1554>.
- Uli Beisel, René Umlauf, Eleanor Hutchinson, and Clare I. R. Chandler. The complexities of simple technologies: re-imagining the role of rapid diagnostic tests in malaria control efforts. *Malaria Journal*, 15:64, 2016. doi: 10.1186/s12936-016-1083-2.
- Konstantinos Benidis, Syama Sundar Rangapuram, Valentin Flunkert, Bernie Wang, Danielle Maddix, Caner Turkmen, Jan Gasthaus, Michael Bohlke-Schneider, David Salinas, Lorenzo Stella, Laurent Callot, and Tim Januschowski. Neural forecasting: Introduction and literature overview, 2020.
- Amelia Bertozzi-Villa, Caitlin A Bever, Hannah Koenker, Daniel J Weiss, Camilo Vargas-Ruiz, Anita K Nandi, Harry S Gibson, Joseph Harris, Katherine E Battle, Susan F Rumisha, et al. Maps and metrics of insecticide-treated net access, use, and nets-per-capita in africa from 2000-2020. *Nature communications*, 12:3589, 2021. doi: 10.1038/s41467-021-23707-7.
- Ipsita Pal Bhowmick, Dibyajyoti Chutia, Avinash Chouhan, Nilay Nishant, PLN Raju, Kanwar Narain, Harpreet Kaur, Rocky Pebam, Jayanta Debnath, Rabindra Tripura, et al. Validation of a mobile health technology platform (fevertracker) for malaria surveillance in india: Development and usability study. *JMIR Formative Research*, 5(11):e28951, 2021. doi: 10.2196/28951.
- G Bidargaddi.N amd Schrader, P Klasnja, J Licinio, and S Murphy. Designing m-health interventions for precision mental health support. *Translational Psychiatry*, 10(222), 2020. doi: 10.1038/s41398-020-00895-2.
- Demetri A Blanas, Youssoupha Ndiaye, Kim Nichols, Andrew Jensen, Ammar Siddiqui, and Nils Hennig. Barriers to community case management of malaria in saraya, senegal: training, and supply-chains. *Malaria Journal*, 12:95, 2013. doi: 10.1186/1475-2875-12-95.
- Nana Yaa Boadu, John Amuasi, Daniel Ansong, Edna Einsiedel, Devidas Menon, and Stephanie K. Yanow. Challenges with implementing malaria rapid diagnostic tests at primary care facilities in a ghanaian district: a qualitative study. *Malaria Journal*, 15:126, 2016. doi: 10.1186/s12936-016-1174-0.
- Justin J Boutilier, Jonas Oddur Jonasson, and Erez Yoeli. Improving tuberculosis treatment adherence support: The case for targeted behavioral interventions. *Manufacturing & Service Operations Management*, 2021. doi: 10.1287/msom.2021.1046.
- Kwame Ohene Buabeng. *The role of the pharmaceutical sector in malaria control in Ghana*. PhD thesis, University of Eastern Finland, 2010. URL [https://erepo.uef.fi/bitstream/handle/123456789/9639/urn\\_isbn\\_978-952-61-0015-9.pdf](https://erepo.uef.fi/bitstream/handle/123456789/9639/urn_isbn_978-952-61-0015-9.pdf)
- Giuseppe Burtini, Jason Loepky, and Ramon Lawrence. A survey of online experiment design with the stochastic multi-armed bandit, 2015.
- Matthew Cairns, Serign Jawo Ceesay, Issaka Sagara, Issaka Zongo, Hamit Kessely, Kadidja Gamougam, Abdoulaye Diallo, Johnbull Sonny Ogbai, Diego Moroso, Suzanne Van Hulle, et al. Effectiveness of seasonal malaria chemoprevention (smc) treatments when smc is implemented at scale: Case-control studies in 5 countries. *PLoS medicine*, 18(9):e1003727, 2021. doi: 10.1371/journal.pmed.1003727.
- Stephanie M Carpenter, Marianne Menictas, Inbal Nahum-Shani, David W Wetter, and Susan A Murphy. Developments in mobile health just-in-time adaptive interventions for addiction science. *Current addiction reports*, 7(3):280–290, 2020. doi: 10.1007/s40429-020-00322-y.
- Minmin Chen, Alex Beutel, Paul Covington, Sagar Jain, Francois Belletti, and Ed Chi. Top-k off-policy correction for a reinforce recommender system, 2020.

- Purvita Chowdhury, Subrata Baidya, Debosmita Paul, Biraj Kalita, Gunendra Saikia, Sanjoy Karmakar, and Sanghita Paul. A comparative study on knowledge and practice against malaria among accredited social health activists (ASHAs) of low and high endemic regions of tripura, northeast india. *Journal of Family Medicine and Primary Care*, 9(5):2420, 2020. doi: 10.4103/jfmpe.jfmpe.1169.19.
- Jessica L Cohen, Hannah H Leslie, Indrani Saran, and Günther Fink. Quality of clinical management of children diagnosed with malaria: A cross-sectional assessment in 9 sub-saharan african countries between 2007–2018. *PLoS medicine*, 17(9):e1003254, 2020. doi: 10.1371/journal.pmed.1003254.
- Daniel DL Coppersmith, Walter Dempsey, Evan Kleiman, Kate Bentley, Susan Murphy, and Matthew Nock. Just-in-time adaptive interventions for suicide prevention: Promise, challenges, and future directions, 2021.
- Fatou Diawara, Laura C Steinhardt, Almahamoudou Mahamar, Tiangoua Traore, Daouda T Kone, Halimatou Diawara, Beh Kamate, Diakalia Kone, Mouctar Diallo, Aboubacar Sadou, et al. Measuring the impact of seasonal malaria chemoprevention as part of routine malaria control in kita, mali. *Malaria journal*, 16:325, 2017. doi: 10.1186/s12936-017-1974-x.
- Junjie Ding, Matthew E Coldiron, Bachir Assao, Ousmane Guindo, Daniel Blessborn, Markus Winterberg, Rebecca F Grais, Alena Koscalova, Celine Langendorf, and Joel Tarning. Adherence and population pharmacokinetic properties of amodiaquine when used for seasonal malaria chemoprevention in african children. *Clinical Pharmacology & Therapeutics*, 107(5):1179–1188, 2020. doi: 10.1002/cpt.1707.
- Raaz Dwivedi, Susan Murphy, and Devavrat Shah. Counterfactual inference for sequential experimental design. 2022.
- Kacey C Ernst, Steven Erly, Charity Adusei, Melanie L Bell, David Komla Kessie, Alberta Biritwum-Nyarko, and John Ehiri. Reported bed net ownership and use in social contacts is associated with uptake of bed nets for malaria prevention in pregnant women in ghana. *Malaria journal*, 16:13, 2017. doi: 10.1186/s12936-016-1660-4.
- Wei Fu and Jeffrey S. Simonoff. Survival trees for left-truncated and right-censored data, with application to time-varying covariate data. *Biostatistics*, 18(2):352–369, 2016. ISSN 1468–4357. doi: 10.1093/biostatistics/kxw047.
- Sophie Githinji, Samwel Kigen, Dorothy Memusi, Andrew Nyandigisi, Andrew Wamari, Alex Muturi, George Jagoe, René Ziegler, Robert W Snow, and Dejan Zurovac. Using mobile phone text messaging for malaria surveillance in rural kenya. *Malaria journal*, 13:107, 2014. doi: 10.1186/1475-2875-13-107.
- MF Gomes, MA Faiz, JO Gyapong, M Warsame, T Agbenyega, A Babiker, F Baiden, EB Yunus, F Binka, C Clerk, et al. Pre-referral rectal artesunate to prevent death and disability in severe malaria: a placebo-controlled trial. *The Lancet*, 373(9663):557–566, 2009. doi: 10.1016/S0140-6736(08)61734-1.
- Kyra H Grantz, Hannah R Meredith, Derek AT Cummings, C Jessica E Metcalf, Bryan T Grenfell, John R Giles, Shruti Mehta, Sunil Solomon, Alain Labrique, Nishant Kishore, et al. The use of mobile phone data to inform analysis of covid-19 pandemic epidemiology. *Nature communications*, 11:4961, 2020. doi: 10.1038/s41467-020-18190-5.
- Anna Guitart, Ana Fernández del Río, Africa Perriñez, and Lauren Bellhouse. Midwifery learning and forecasting: Predicting content demand with user-generated logs. In *Proceedings of 2021 KDD Workshop on Applied Data Science for Healthcare (DSHealth 2021)*. ACM, 2021a. URL <https://arxiv.org/abs/2107.02480>.
- Anna Guitart, Afsaneh Heydari, Eniola Olaleye, et al. A recommendation system to enhance midwives’ capacities in low-income countries, 2021b.

- Teklemariam Gultie, Gistane Ayele, Befikadu Tariku, Mekdes Kondale, Zerihun Zerdo, Behailu Merdekiyos, Tsegaye Tsalla, Mesfin Kote, Alemayehu Bekele, Mulugeta Shigaz, Gebrekiros Gebremichael, Feleke Gebremeskel, and Alazar Baharu. Trend of declining bed net utilization among pregnant women in ethiopia: new data from the arba minch health and demographic surveillance system, 2010–2016. *Malaria Journal*, 19:142, 2020. doi: 10.1186/s12936-020-03211-x.
- Huifeng Guo, Ruiming Tang, Yunming Ye, Zhenguo Li, and Xiuqiang He. Deepfm: A factorization-machine based neural network for ctr prediction, 2017.
- Wendy Hardeman, Julie Houghton, Kathleen Lane, Andy Jones, and Felix Naughton. A systematic review of just-in-time adaptive interventions (jitais) to promote physical activity. *International Journal of Behavioral Nutrition and Physical Activity*, 16:31, 2019. doi: 10.1186/s12966-019-0792-7.
- Emir Hrnjic and Nikodem Tomczak. Machine learning and behavioral economics for personalized choice architecture, 2019.
- Eugene Ie, Vihan Jain, Jing Wang, Sanmit Narvekar, Ritesh Agarwal, Rui Wu, Heng-Tze Cheng, Morgane Lustman, Vince Gatto, Paul Covington, Jim McFadden, Tushar Chandra, and Craig Boutilier. Reinforcement learning for slate-based recommender systems: A tractable decomposition and practical methodology, 2019.
- Djibrilla Issiaka, Amadou Barry, Tiangoua Traore, Boubacar Diarra, David Cook, Mohamed Keita, Issaka Sagara, Patrick Duffy, Michal Fried, and Alassane Dicko. Impact of seasonal malaria chemoprevention on hospital admissions and mortality in children under 5 years of age in oulessebouyou, mali. *Malaria Journal*, 19:103, 2020. doi: 10.1186/s12936-020-03175-y.
- Appolinary AR Kamuhabwa and Richard Silumbe. Knowledge among drug dispensers and anti-malarial drug prescribing practices in public health facilities in dar es salaam. *Drug, Healthcare and Patient Safety*, 5:181–189, 2013. doi: 10.2147/DHPS.S50071.
- Iraklis Katsaris, , and Nikolas Vidakis and. Adaptive e-learning systems through learning styles: A review of the literature. *Advances in Mobile Learning Educational Research*, 1(2):124–145, 2021. doi: 10.25082/amler.2021.02.007.
- Blessings N Kaunda-Khangamwa, Laura C Steinhardt, Alexander K Rowe, Austin Gumbo, Dubulao Moyo, Humphreys Nsona, Peter Troell, Dejan Zurovac, and Don Mathanga. The effect of mobile phone text message reminders on health workers’ adherence to case management guidelines for malaria and other diseases in malawi: lessons from qualitative data from a cluster-randomized trial. *Malaria journal*, 17:481, 2018. doi: 10.1186/s12936-018-2629-2.
- Guibehi B Koudou, April Monroe, Seth R Irish, Michael Humes, Joseph D Krezanoski, Hannah Koenker, David Malone, Janet Hemingway, and Paul J Krezanoski. Evaluation of an accelerometer-based monitor for detecting bed net use and human entry/exit using a machine learning algorithm. *Malaria journal*, 21:85, 2022. doi: 10.1186/s12936-022-04102-z.
- Jeremiah Laktabai, Alyssa Platt, Diana Menya, Elizabeth L Turner, Daniel Aswa, Stephen Kinoti, and Wendy Prudhomme O’Meara. A mobile health technology platform for quality assurance and quality improvement of malaria diagnosis by community health workers. *PLoS One*, 13(2): e0191968, 2018. doi: 10.1371/journal.pone.0191968.
- Changhee Lee, Jinsung Yoon, and Mihaela van der Schaar. Dynamic-DeepHit: A deep learning approach for dynamic survival analysis with competing risks based on longitudinal data. *IEEE Transactions on Biomedical Engineering*, 67(1):122–133, 2020. doi: 10.1109/tbme.2019.2909027.
- Xiaohong Li, Youyou Tu, Linhua Tang, Qi Gao, and Pedro L. Alonso. The role of research in china’s successful elimination of malaria. *Nature Medicine*, 2022. doi: <https://doi.org/10.1038/s41591-022-01824-0>.



- Jianxun Lian, Xiaohuan Zhou, Fuzheng Zhang, Zhongxia Chen, Xing Xie, and Guangzhong Sun. xdeepfm: Combining explicit and implicit feature interactions for recommender systems. In *Proceedings of the 24th ACM SIGKDD International Conference on Knowledge Discovery & Data Mining*, pp. 1754–1763. ACM, 2018. doi: 10.1145/3219819.3220023.
- Timothy P. Lillicrap, Jonathan J. Hunt, Alexander Pritzel, Nicolas Heess, Tom Erez, Yuval Tassa, David Silver, and Daan Wierstra. Continuous control with deep reinforcement learning, 2019.
- Bryan Lim, Sercan O. Arik, Nicolas Loeff, and Tomas Pfister. Temporal fusion transformers for interpretable multi-horizon time series forecasting, 2019.
- Feng Liu, Ruiming Tang, Xutao Li, Weinan Zhang, Yunming Ye, Haokun Chen, Huifeng Guo, and Yuzhou Zhang. Deep reinforcement learning based recommendation with explicit user-item interactions modeling, 2019.
- Christopher Lourenço, Andrew J Tatem, Peter M Atkinson, Justin M Cohen, Deepa Pindolia, Darlene Bhavnani, and Arnaud Le Menach. Strengthening surveillance systems for malaria elimination: a global landscaping of system performance, 2015–2017. *Malaria journal*, 18:315, 2019. doi: 10.1186/s12936-019-2960-2.
- Wantong Lu, Yantao Yu, Yongzhe Chang, Zhen Wang, Chenhui Li, and Bo Yuan. A dual input-aware factorization machine for ctr prediction. In *IJCAI’20: Proceedings of the Twenty-Ninth International Joint Conference on Artificial Intelligence*, pp. 3139–3145, 2020. URL <https://www.ijcai.org/proceedings/2020/0434.pdf>.
- K. E. Mace, D. Mwandama, J. Jafali, M. Luka, S. J. Filler, J. Sande, D. Ali, S. P. Kachur, D. P. Mathanga, and J. Skarbinski. Adherence to treatment with artemether-lumefantrine for uncomplicated malaria in rural malawi. *Clinical Infectious Diseases*, 53(8):772–779, 2011. doi: 10.1093/cid/cir498.
- Diego Macías Saint-Gerons, Sheila Rodovalho, Ádila Liliâne Barros Dias, André Lacerda Ulysses de Carvalho, Andrea Beratarrechea, Wuelton Marcelo Monteiro, Myrna Barata Machado, Cristiano Fernandes da Costa, Marcelo Yoshito Wada, Márcia Helena Maximiano Faria de Almeida, et al. Strengthening therapeutic adherence and pharmacovigilance to antimalarial treatment in manaus, brazil: a multicomponent strategy using mhealth. *Malaria journal*, 21:28, 2022. doi: 10.1186/s12936-022-04047-3.
- Grace Manu, Ellen Abrafi Boamah-Kaali, Lawrence Gyabaa Febir, Emmanuel Ayipah, Seth Owusu-Agyei, and Kwaku Poku Asante. Low utilization of insecticide-treated bed net among pregnant women in the middle belt of ghana. *Malaria Research and Treatment*, 2017:1–7, 2017. doi: 10.1155/2017/7481210.
- Aditya Mate, Lovish Madaan, Aparna Taneja, Neha Madhiwalla, Shresth Verma, Gargi Singh, Aparna Hegde, Pradeep Varakantham, and Milind Tambe. Field study in deploying restless multi-armed bandits: Assisting non-profits in improving maternal and child health. In *AAAI Conference on Artificial Intelligence*, Vancouver, Canada, 2022.
- Marianne Menictas, Mashfiqui Rabbi, Predrag Klasnja, and Susan Murphy. Artificial intelligence decision-making in mobile health. *The Biochemist*, 41(5):20–24, 2019. doi: 10.1042/BIO04105020.
- Sveta Milusheva. Managing the spread of disease with mobile phone data. *Journal of Development Economics*, 147:102559, 2020. doi: 10.1016/j.jdevco.2020.102559.
- Aliyu Mohammed, Princess Ruhama Acheampong, Easmon Otupiri, Francis Adjei Osei, Roderick Larson-Reindorf, and Ellis Owusu-Dabo. Mobile phone short message service (SMS) as a malaria control tool: a quasi-experimental study. *BMC Public Health*, 19(1):1193, 2019. doi: 10.1186/s12889-019-7336-6.
- Godfrey Mubyazi, Paul Bloch, Mathias Kamugisha, Andrew Kitua, and Jasper Ijumba. Intermittent preventive treatment of malaria during pregnancy: a qualitative study of knowledge, attitudes and practices of district health managers, antenatal care staff and pregnant women in korogwe district, north-eastern tanzania. *Malaria Journal*, 4:31, 2005. doi: 10.1186/1475-2875-4-31.

- Inbal Nahum-Shani, Shawna N Smith, Bonnie J Spring, Linda M Collins, Katie Witkiewitz, Ambuj Tewari, and Susan A Murphy. Just-in-time adaptive interventions (jitais) in mobile health: key components and design principles for ongoing health behavior support. *Annals of Behavioral Medicine*, 52(6):446–462, 2018. doi: 10.1007/s12160-016-9830-8.
- Paul N. Newton, , Kara Hanson, and Catherine Goodman. Do anti-malarials in africa meet quality standards? the market penetration of non quality-assured artemisinin combination therapy in eight african countries. *Malaria Journal*, 16:204, 2017. doi: 10.1186/s12936-017-1818-8.
- Van Bach Nguyen, Belaid Mohamed Karim, Bao Long Vu, Jörg Schlötterer, and Michael Granitzer. Policy learning for malaria control. *ArXiv*, abs/1910.08926, 2019.
- Abdisalan M Noor and Pedro L Alonso. The message on malaria is clear: progress has stalled. *The Lancet*, 399:1777, 2022. doi: 10.1016/S0140-6736(22)00732-2.
- Kathryn A O’Connell, Hellen Gatakaa, Stephen Poyer, Julius Njogu, Illah Evance, Erik Munroe, Tsione Solomon, Catherine Goodman, Kara Hanson, Cyprien Zinsou, et al. Got acts? availability, price, market share and provider knowledge of anti-malarial medicines in public and private sector outlets in six malaria-endemic countries. *Malaria journal*, 10:326, 2011. doi: 10.1186/1475-2875-10-326.
- Colin Ohrt, Kathryn W Roberts, Hugh JW Sturrock, Jennifer Wegbreit, Bruce Y Lee, and Roly D Gosling. Information systems to support surveillance for malaria elimination. *The American journal of tropical medicine and hygiene*, 93(1):145–152, 2015. doi: 10.4269/ajtmh.14-0257.
- Elizabeth O Onyango, George Ayodo, Carren A Watsierah, Tom Were, Wilson Okumu, Samuel B Anyona, Evans Raballah, John M Okoth, Sussy Gumo, George O Orinda, and Collins Ouma. Factors associated with non-adherence to artemisinin-based combination therapy (ACT) to malaria in a rural population from holoendemic region of western kenya. *BMC Infectious Diseases*, 12(1), 2012. doi: 10.1186/1471-2334-12-143.
- Win Han Oo, Win Htike, Julia C Cutts, Kyawt Mon Win, Kaung Myat Thu, May Chan Oo, Galau Naw Hkawng, Paul A Agius, Pyae Phyo Htoon, Nick Scott, et al. A mobile phone application for malaria case-based reporting to advance malaria surveillance in myanmar: a mixed methods evaluation. *Malaria Journal*, 20:167, 2021. doi: 10.1186/s12936-021-03701-6.
- World Health Organization. Fact sheet about malaria, 2022. URL <https://www.who.int/news-room/fact-sheets/detail/malaria>.
- Yanru Qu, Han Cai, Kan Ren, Weinan Zhang, Yong Yu, Ying Wen, and Jun Wang. Product-based neural networks for user response prediction. In *2016 IEEE 16th International Conference on Data Mining (ICDM)*, pp. 1149–1154. IEEE, 2016. doi: 10.1109/ICDM.2016.0151.
- Manju Rahi and Amit Sharma. For malaria elimination india needs a platform for data integration. *BMJ Global Health*, 5(12):e004198, 2020. doi: 10.1136/bmjgh-2020-004198.
- Harsh Rajvanshi, Yashpal Jain, Nidhi Kaintura, Chaitanya Soni, Raja Chandramohan, Ramanathan Srinivasan, Vinay Telasey, Praveen K. Bharti, Deepak Jain, Mangeshi Surve, Sachin Saxena, Vilas Gangamwar, M. S. Anand, and Altaf A. Lal. A comprehensive mobile application tool for disease surveillance, workforce management and supply chain management for malaria elimination demonstration project. *Malaria Journal*, 20:91, 2021. doi: 10.1186/s12936-021-03623-3.
- Charlotte Rasmussen, Pedro Alonso, and Pascal Ringwald. Current and emerging strategies to combat antimalarial resistance. *Expert Review of Anti-infective Therapy*, 20(3):353–372, 2022. doi: 10.1080/14787210.2021.1962291.
- Emily Ricotta, Hannah Koenker, Albert Kilian, and Matthew Lynch. Are pregnant women prioritized for bed nets? an assessment using survey data from 10 african countries. *Global Health: Science and Practice*, 2(2):165–172, 2014. doi: 10.9745/GHSP-D-14-00021.
- Max Roser and Hannah Ritchie. Malaria, 2019. URL <https://ourworldindata.org/malaria>.

- Alexander K Rowe, Gabriel F Ponce de León, Jules Mihigo, Ana Carolina FS Santelli, Nathan P Miller, and Pedro Van-Dúnem. Quality of malaria case management at outpatient health facilities in angola. *Malaria Journal*, 8:275, 2009. doi: 10.1186/1475-2875-8-275.
- RTI. Coconut surveillance: Open-source mobile tool designed to take on malaria, other infectious diseases, 2020. Retrieved from <https://www.rti.org/impact/coconut-surveillance>.
- David Salinas, Michael Bohlke-Schneider, Laurent Callot, Roberto Medico, and Jan Gasthaus. High-dimensional multivariate forecasting with low-rank gaussian copula processes. arXiv, 2019.
- David Salinas, Valentin Flunkert, Jan Gasthaus, and Tim Januschowski. Deepar: Probabilistic forecasting with autoregressive recurrent networks. *International Journal of Forecasting*, 36(3): 1181–1191, 2020. ISSN 0169-2070. doi: <https://doi.org/10.1016/j.ijforecast.2019.07.001>. URL <https://www.sciencedirect.com/science/article/pii/S0169207019301888>.
- Cristolde Salomão, Jahit Sacarlal, and Eduardo Samo Gudo. Assessment of coverage of preventive treatment and insecticide-treated mosquito nets in pregnant women attending antenatal care services in 11 districts in mozambique in 2011: the critical role of supply chain. *Malaria journal*, 16:223, 2017. doi: 10.1186/s12936-017-1872-2.
- Matthias Seeger, Syama Rangapuram, Yuyang Wang, David Salinas, Jan Gasthaus, Tim Januschowski, and Valentin Flunkert. Approximate bayesian inference in linear state space models for intermittent demand forecasting at scale, 2017.
- Daudi O Simba, Deodatus C Kakoko, Marian Warsame, Zul Premji, Melba F Gomes, Goran Tomson, and Eva Johansson. Understanding caretakers’ dilemma in deciding whether or not to adhere with referral advice after pre-referral treatment with rectal artesunate. *Malaria journal*, 9:123, 2010. doi: 10.1186/1475-2875-9-123.
- Dexian Tang, Guillem Francès, and África Perriñez. A data-centric behavioral machine learning platform to reduce health inequalities. In *35th Conference on Neural Information Processing Systems (NeurIPS 2021)*, 2021. URL [https://datacentricai.org/neurips21/papers/166\\_CameraReady\\_main.pdf](https://datacentricai.org/neurips21/papers/166_CameraReady_main.pdf).
- Joel Tarning. Treatment of malaria in pregnancy. *New England Journal of Medicine*, 374(10): 981–982, 2016. doi: 10.1056/nejme1601193.
- Daniel J Weiss, Amelia Bertozzi-Villa, Susan F Rumisha, Punam Amratia, Rohan Arambepola, Katherine E Battle, Ewan Cameron, Elisabeth Chestnutt, Harry S Gibson, Joseph Harris, et al. Indirect effects of the covid-19 pandemic on malaria intervention coverage, morbidity, and mortality in africa: a geospatial modelling analysis. *The Lancet Infectious Diseases*, 21(1):59–69, 2021. doi: 10.1016/S1473-3099(20)30700-3.
- WHO. Global technical strategy for malaria 2016-2030, 2015.
- WHO. The use of rectal artesunate as a pre-referral treatment for severe *P. falciparum* malaria, 2022. Retrieved from <https://apps.who.int/iris/bitstream/handle/10665/351187/9789240042513-eng.pdf>.
- Marvin N. Wright, Theresa Dankowski, and Andreas Ziegler. Unbiased split variable selection for random survival forests using maximally selected rank statistics. *Statistics in Medicine*, 36(8): 1272–1284, 2017. doi: <https://doi.org/10.1002/sim.7212>.
- Jiayu Yao, Emma Brunskill, Weiwei Pan, Susan Murphy, and Finale Doshi-Velez. Power constrained bandits. In Ken Jung, Serena Yeung, Mark Sendak, Michael Sjoding, and Rajesh Ranganath (eds.), *Proceedings of the 6th Machine Learning for Healthcare Conference*, volume 149 of *Proceedings of Machine Learning Research*, pp. 209–259. PMLR, 2021. URL <https://proceedings.mlr.press/v149/yao21a.html>.
- Kelly W Zhang, Lucas Janson, and Susan A Murphy. Statistical inference after adaptive sampling in non-markovian environments, 2022.

- Guofa Zhou, Ming chieh Lee, Harrysone E. Atieli, John I. Githure, Andrew K. Githeko, James W. Kazura, and Guiyun Yan. Adaptive interventions for optimizing malaria control: an implementation study protocol for a block-cluster randomized, sequential multiple assignment trial. *Trials*, 21(1), jul 2020a. doi: 10.1186/s13063-020-04573-y.
- Guofa Zhou, Ming chieh Lee, Xiaoming Wang, Daibin Zhong, Elizabeth Hemming-Schroeder, and Guiyun Yan. An adaptive intervention trial design for finding the optimal integrated strategies for malaria control and elimination in africa: A model simulation study. *The American Journal of Tropical Medicine and Hygiene*, 105(5):1173–1183, nov 2021. doi: 10.4269/ajtmh.20-0254.
- Wenjun Zhou, Taposh Dutta Roy, and Iryna Skrypnik. The kdd cup 2019 report. *SIGKDD Explor. Newsl.*, 22(1):8–17, 2020b. ISSN 1931-0145. doi: 10.1145/3400051.3400056. URL <https://doi.org/10.1145/3400051.3400056>.
- Lixin Zou. Data-efficient reinforcement learning for malaria control. In Zhi-Hua Zhou (ed.), *Proceedings of the Thirtieth International Joint Conference on Artificial Intelligence, IJCAI-21*, pp. 507–513. International Joint Conferences on Artificial Intelligence Organization, 2021. doi: 10.24963/ijcai.2021/71. URL <https://doi.org/10.24963/ijcai.2021/71>. Main Track.
- Dejan Zurovac, Raymond K Sudoi, Willis S Akhwale, Moses Ndiritu, Davidson H Hamer, Alexander K Rowe, and Robert W Snow. The effect of mobile phone text-message reminders on kenyan health workers' adherence to malaria treatment guidelines: a cluster randomised trial. *The Lancet*, 378(9793):795–803, 2011. doi: 10.1016/s0140-6736(11)60783-6.

## A APPENDIX: TOOLS AND METHODS

This appendix describes the proposed conceptual and methodological framework: a data-centric behavioral ML platform (Tang et al., 2021) that leverages logs from different types of mobile health solutions—together with contextual information—to deliver adaptive interventions to healthcare workers and their patients directly through their phones.

### A.1 DATA-CENTRIC PLATFORM

Our platform is data-centric in that data tracking, and labeling lies at its core. Integration with the different apps is achieved through our *Software Developer Kit* (HealthKit SDK), which provides specifications on what information should be logged (and how) and the messaging service that delivers the interventions. Integration through this SDK ensures that various user metrics and traits characterizing engagement and behavior are readily available, which can help clarify the individuals to be targeted by specific interventions and the time for delivering the interventions.

### A.2 MACHINE LEARNING

The application of data science and ML methodologies to extract and predict valuable information to inform intervention design lies at the core of the software we build. As is generally the case, no single model is best across all datasets and use cases. Use-case-specific data pipelines transform incoming information through the HealthKit SDK into metrics ready to be consumed by the statistical and ML models, which in turn produce additional metrics that can be used in the intervention definition.

#### A.2.1 REINFORCEMENT LEARNING

Reinforcement learning (RL) is the ideal paradigm for sequential decision-making in dynamically evolving contexts that respond to those decisions. It allows us to continually improve how we make choices for a patient at any given moment, maximizing the potential for positive outcomes while minimizing undesired side effects. A competition on policy learning for malaria control using RL was for example included as part of the KDD Cup Challenge 2019 (Zhou et al., 2020b; Nguyen et al., 2019; Zou, 2021).

Contextual are a formulation of the RL problem with limited state representation, and dynamics (Burtini et al., 2015; Yao et al., 2021; Dwivedi et al., 2022; Zhang et al., 2022). Similarly, restless bandits can be used for resource allocation (Mate et al., 2022). They are significantly less data-intensive than other RL approaches while being robust and tractable. This framework is instrumental in the context of adaptive intervention delivery. At their core lies the exploration-exploitation trade-off, i.e., the compromise between clinical research (to gather knowledge about treatments) and clinical practice (to benefit the patients) by assigning the best intervention possible based on all available information at that point.

#### A.2.2 TIME-VARYING AND DYNAMIC PREDICTION MODELING

Time series datasets such as clinical records represent valuable sources of information sometimes spanning a patient’s entire lifetime of care. The approach to decision-making described in this paper is dynamic (i.e., sequential and adaptive). The analytic and predictive modeling to support this needs to be similarly dynamic, with the time-dependent evolution of the systems of interest and their characteristics at its core. Supervised and unsupervised ML with time-varying data (such as the survival analysis described in the next section), time series modeling, and longitudinal data processing should be part of the toolbox. We can use them to understand and predict how the systems of interest behave and evolve to inform our decision-making.

For behavioral nudging, besides the individual predictions generated using survival analysis described below, multivariate time series forecasting can be critical to optimizing specific interventions, such as reminders to prevent stockouts of medical supplies based on demand prediction. Furthermore, considering the time-varying nature of data and interventions within an RL framework allows us to understand and optimize a patient’s treatment as different sequential interventions in time instead of as a single-point decision.

Predictive modeling can help define the target individuals for the intervention and appropriate delivery time. As discussed, accurate demand prediction is key to optimizing the supply chain and inventory. Forecasting methods that can learn simultaneously from multiple time series, often combining deep and state space modeling elements, have been noted to be effective (Seeger et al., 2017; Salinas et al., 2019; Lim et al., 2019; Salinas et al., 2020; Benidis et al., 2020).

#### A.2.3 DEEP AND ENSEMBLE SURVIVAL ANALYSIS

With time dependence at its core, survival analysis refers to a collection of algorithms used to predict time to an event of interest (which was traditionally death or organ failure) (Wright et al., 2017; Fu & Simonoff, 2016; Lee et al., 2020). These methodologies can be used to predict behavior and health outcomes at the individual level, as they are well suited to extract information from censored data (i.e., models that can learn from subjects who have yet to experience the event of interest besides those that have). Their output is a survival curve for each subject, indicating the probability of not having experienced the event of interest depending on the time. By carefully selecting the events we predict and how we measure time, we can profile individuals using multiple predictions and risk scores (of clinical complications or of lack of adherence to treatment, for example).

#### A.2.4 RECOMMENDATION SYSTEMS

Recommendation algorithms can be used to decide the techniques to target health workers, patients, and the general public. Different methodologies are expected to be suitable for different apps and use cases, from deep learning-based click-through-rate predictions (Qu et al., 2016; Guo et al., 2017; Lian et al., 2018; Lu et al., 2020; Guitart et al., 2021b) to collaborative, interactive recommendation systems Ie et al. (2019); Liu et al. (2019); Lillicrap et al. (2019); Chen et al. (2020). We foresee a growing shift toward reinforcement-learning-based methods beyond the (relatively widespread) use of contextual bandits (Barraza-Urbina & Glowacka, 2020). This approach is particularly useful for optimizing a *sequence of interventions* while reconciling short- and long-term goals in the desired outcome.