Asymptotic theory for extreme value generalized additive models

Takuma Yoshida¹

¹Kagoshima University, Kagoshima 890-8580, Japan E-mail: yoshida@sci.kagoshima-u.ac.jp

Abstract

The classical approach to analyzing extreme value data is the generalized Pareto distribution (GPD). When the GPD is used to explain a target variable with the large dimension of covariates, the shape and scale function of covariates included in GPD are sometimes modeled using the generalized additive models (GAM). In contrast to many results of application, there are no theoretical results on the hybrid technique of GAM and GPD, which motivates us to develop its asymptotic theory. We provide the rate of convergence of the estimator of shape and scale functions, as well as its local asymptotic normality.

keywords: Extreme value theory; Generalized additive model; Generalized Pareto distribution; Peak over threshold; Penalized spline.

MSC Classification 62G08, 62G20, 62G32.

1 Introduction

Generalized Pareto distribution (GPD) is a typical model to express the tail probability of the tail event. The extreme value theory (EVT) explains that the GPD can fit the data exceeding some high threshold value. The GPD contains two parameters, one characterizes the shape of distribution whereas other can be regarded as scale. Smith (1987) and Drees et al. (2004) developed the maximum likelihood method for the shape and scale parameter of GPD and established an asymptotic theory of the maximum likelihood estimator.

To predict tail probability of a target variable associated with covariate information, we often consider the GPD with shape and scale function of the covariate using GPD regression. Hall and Tajvidi (2000), Ramesh and Davison (2002), and Beirlant et al. (2004) investigated GPD regression in the context of nonparametric smoothing with kernel methods. However, the ordinary nonparametric estimator would be drastically incorrect when the number of covariates is large, the so-called curse of dimensionality. In the regression with a large number of covariates, some specific modeling would be better instead of fully nonparametric methods. To avoid the curse of dimensionality, one of the efficient approaches is the generalized additive model (GAM). Buja et al. (1989) proposed the GAM, which has been developed by many authors in several regression models, summarized by Hastie and Tibshirani (1990) and Wood (2017). Chavez-Demoulin and Davison (2005) provided the GPD regression with shape and scale functions assumed to be GAM. Yee and Stephenson (2007), Chavez-Demoulin (2015), Vatter and Chavez-Demoulin (2015), Mhalla et al. (2019), and Youngman (2019) contributed the additive modeling for extreme value analysis. Since the GPD regression has two target functions for shape and scale, its additive modeling is sometimes denoted by vector generalized additive models (see, Yee 2015). The important work on GAM in GPD regression is Youngman (2022), who have published the Rpackage evgam, and hence, everyone can easily use GAM in GPD in extreme value data analysis.

On the other hand, to the best of our knowledge, there is a lack of theoretical investigation of GPD regression with GAM. This motivates us to establish its asymptotic theory. The estimator of shape and (logarithm of) scale function under GAM is constructed via penalized maximum-likelihood method and the technique of spline-type smoothing method (see, Wood 2017). Such method is also the standard approach for the estimator obtained by evgam. As the spline smoothing method, we consider the normalized *B*-spline model (see, Wang and Yang (2007), Liu et al. (2011), and Liu et al. (2013)). The advantage of using the normalized *B*-spline method is that, unlike other spline bases, those of normalized *B*-spline model are linear independent (see, Marx and Eilers (1998), Wang and Yang (2007)), and hence, the non-singularity of the Fisher information matrix of log-likelihood of GPD can be guaranteed. This is the key property to evaluate the asymptotic variance of the estimator. Meanwhile, asymptotic bias of the estimator can be analyzed using the derivative of normalized *B*-spline model (see, Xiao (2019)) and the second-order condition of EVT (see, de Haan and Ferreira (2006)). Consequently, we derive the optimal rate of convergence of the estimators of shape and scale functions. The local asymptotic normality of the estimator is also presented.

The rest of this paper is organized as follows. Basic conditions of EVT are summarized in section 2. Section 3 provides the estimator of shape and scale function under GAM in the GPD regression. Main results are presented in section 4. We first introduce the mathematical conditions to obtain the asymptotic theory for the estimator in section 4.1. Section 4.2 presents the L_2 and L_{∞} -rate of convergence of the estimator as well as local asymptotic normality. In section 5, we establish the asymptotic theory for the additive estimator with orthogonalreparametrized GPD model proposed by Chavez-Demoulin and Davison (2005). Future work is discussed in section 6. All proofs of theorems and related lemmas are described in the Appendix (see, Supplementary file). This study focuses on mathematical result of the estimator under GAM in GPD regression. Its numerical performance can be easily confirmed via the evgam, and hence, we omit the numerical study.

2 Preriminalies

2.1 Extreme value theory

We first review the EVT for univariate random variable. Let $Y \in \mathbb{R}_+$ be the random variable with distribution function F, denoted by F(y) = P(Y < y) for $y \in \mathbb{R}_+$, where $\mathbb{R}_+ = (0, \infty)$. For some threshold value $w \in \mathbb{R}_+$, we define

$$F_w(y) = \frac{F(w+y) - F(w)}{1 - F(w)}.$$

Meanwhile, we define the GPD function as

$$H(y|\gamma) = 1 - (1 + \gamma y)^{-1/\gamma}$$

with $1 + \gamma y > 0$ and parameter $\gamma \in \mathbb{R}$. If $\gamma = 0$, we set $H(y|0) = \lim_{\gamma \to 0} H(y|\gamma) = 1 - e^{-y}$, which is the distribution function of standard exponential distribution. Then, it is well known that there exists a constant $\sigma_w \in \mathbb{R}_+$ such that for any $y \in (0, y^* - w)$,

$$\lim_{w \to y^*} \left| F_w(y) - H\left(\left. \frac{y}{\sigma_w} \right| \gamma \right) \right| \to 0, \tag{1}$$

where $y^* = \sup\{t : F(t) < 1\}$ if and only if F belongs to domain of attraction of distribution function $G(y|\gamma) = \exp[-(1+\gamma y)^{-1/\gamma}], 1+\gamma y > 0$ with $\gamma \in \mathbb{R}$, denoted by $F \in \mathcal{D}(G(\cdot|\gamma))$ (e.g.

Theorem 1.2.5 in de Haan and Ferreira (2006)). From the above, we see that if $\gamma > 0$, $y^* = \infty$ and y^* is finite for $\gamma < 0$. For $\gamma = 0$, y^* can obtain both finite or infinite. In this paper, we only consider the case that $y^* = \infty$ if $\gamma = 0$. In addition, σ_w can be taken as

$$\begin{cases} \lim_{w \to \infty} \frac{\sigma_w}{w_{\sigma_w}} = \gamma, & \gamma > 0, \\ \lim_{w \to y^*} \frac{\sigma_w}{y^* - w} = -\gamma, & \gamma < 0, \\ \lim_{w \to \infty} \sigma_w = \sigma, & \gamma = 0, \end{cases}$$
(2)

where $\sigma > 0$ is some constant (see, Theorem 1.2.5 in de Haan and Ferreira 2006).

To predict the probability using the GPD model, the estimation of (γ, σ_w) is important. Smith (1987) and Drees et al. (2004) investigated the maximum-likelihood estimator of (γ, σ_w) and its asymptotic result. To establish the asymptotic theory of the estimator of (γ, σ_w) , (1) should be modified in the context of second-order condition of EVT.

Let $F^{-1}(x) = \inf\{y : F(y) \ge x\}$ and let $U(t) = F^{-1}(1 - 1/t)$. Then, $U(t) \to y^*$ as $t \to \infty$. According to Theorem 1.1.6 in de Haan and Ferreira (2006), $F \in \mathcal{D}(G(\cdot|\gamma))$ if and only if function $a : \mathbb{R}_+ \to \mathbb{R}_+$ exists such that $\{U(tx) - U(t)\}/a(t) \to (x^{\gamma} - 1)/\gamma$ as $t \to \infty$. Note that $a(t) = \sigma_w$ with w = U(t). Let us assume that function $A : \mathbb{R}_+ \to \mathbb{R}_+$ and $\tilde{Q} : \mathbb{R}_+ \to \mathbb{R}_+$ exists such that

$$\lim_{t \to \infty} \left| \frac{\frac{U(tx) - U(t)}{a(t)} - \frac{x^{\gamma} - 1}{\gamma}}{A(t)} - \tilde{Q}(x|\gamma, \rho) \right| = 0,$$
(3)

where $\rho \leq 0$ is the so-called second-order parameter,

$$\tilde{Q}(x|\gamma,\rho) = \frac{1}{\rho} \left(\frac{x^{\gamma+\rho-1}-1}{\gamma+\rho} - \frac{x^{\gamma}-1}{\gamma} \right),$$

and $A(t) \to 0$ as $t \to \infty$. Then, Theorem 2.3.3 in de Haan and Ferreira (2006) demonstrates that A is ρ -regularly varying function at $t \to \infty$, that is, for all $x \in \mathbb{R}_+$, $A(xt)/A(t) \to x^{\rho}$ as $t \to \infty$. If $\gamma = 0$ or $\rho = 0$, we get $\tilde{Q}(x|\gamma, \rho)$ as $\gamma \to 0$ or $\rho \to 0$. By Theorem 2.3.8 in de Haan and Ferreira (2006), under (3), we obtain

$$\lim_{w \to y^*} \left| \frac{F_w(y) - H(y/\sigma_w|\gamma)}{\alpha(w)} - Q(y/\sigma_w|\gamma, \rho) \right| = 0$$
(4)

where $Q(y|\gamma,\rho) = \bar{H}(y|\gamma)^{1+\gamma}\tilde{Q}(\bar{H}^{-1}(y|\gamma)|\gamma,\rho)$, $\bar{H}(y|\gamma) = 1 - H(y|\gamma)$ and $\alpha(w) = A(1/(1 - F(w)))$. When (4) is satisfied, F belongs to domain of attraction of $G(\cdot|\gamma)$ with second-order parameter ρ . The asymptotic behavior of the estimator of (γ, σ_w) is dependent not only (γ, σ_w, w) but also ρ . Therefore, the second-order condition of EVT is an important assumption to examine the asymptotic theory for the estimator of (γ, σ_w) .

Remark 1. In (2), the ordinary property of scale parameter for $\gamma = 0$ is $d\sigma_w/dw \to 0$. That is, for $\gamma = 0$, $\sigma_w = \sigma \log w$ with some constant $\sigma > 0$ and $\sigma_w \to 0$ are also allowed. However if we consider such general condition, the discussion for $\gamma = 0$ becomes more complicated (see, Zhou 2009). Therefore, in this paper, we only focus on the simple case that σ_w converges to constant, that is, $H(y/\sigma_w|0) \approx 1 - e^{-y/\sigma}$.

2.2 Extreme value theory in regression

We now extend the univariate EVT discussed in previous section to conditional EVT. Let (Y^*, X, Z) be triplet random variables with response $Y^* \in \mathbb{R}_+$ and covariates $X = (X^{(1)}, \dots, X^{(p)})^T \in \mathbb{R}_+$

 $\mathcal{X} \subset \mathbb{R}^p$ and $Z = (Z^{(1)}, \ldots, Z^{(d)})^T \in \mathcal{Z} \subset \mathbb{R}^d$. Here, \mathcal{X} and \mathcal{Z} are assumed to be a compact set. Let $F(y|x,z) = P(Y^* < y|X = x, Z = z)$ be conditional distribution function of Y^* given $(X^*, Z^*) = (x, z) = (x^{(1)}, \ldots, x^{(p)}, z^{(1)}, \ldots, z^{(d)})^T \in \mathcal{X} \times \mathcal{Z}$. The covariate dependent threshold function is denoted by $\tau(w|x, z)$, where w is some sequence, which controls the level of threshold function. We choose τ so that for any $(x, z) \in \mathcal{X} \times \mathcal{Z}$, $\tau(w|x, z) \to y^*(x, z)$ as $w \to \infty$, where $y^*(x, z) = \sup\{t : F(t|x, z) < 1\}$. Let $F_{w,\tau}(y|x, z) = \{F(\tau(w|x, z) + y|x, z) - F(\tau(w|x, z)|x, z)\}/\{1 - F(\tau(w|x, z)|x, z)\}$. For simplicity, we write $F_{w,\tau}(y|x, z) = F_w(y|x, z)$. We see that $F_w(y|x, z) = P(Y^* < \tau(w|x, z) + y|Y^* > \tau(w|x, z), x, z)$.

As an extension of (1) to regression version, we assume that for any $(x, z) \in \mathcal{X} \times \mathcal{Z}, F(\cdot|x, z) \in \mathcal{D}(G(\cdot|\gamma_0(x, z)))$ with $\gamma_0 : \mathcal{X} \times \mathcal{Z} \to \mathbb{R}$ and there exists some function $\sigma_{w,\tau}^{\dagger} : \mathcal{X} \times \mathcal{Z} \to \mathbb{R}_+$ such that

$$\lim_{w \to \infty} \sup_{(x,z) \in \mathcal{X} \times \mathcal{Z}} \left| F_w(y|x,z) - H\left(\frac{y}{\sigma_{w,\tau}^{\dagger}(x,z)} | \gamma_0(x,z)\right) \right| = 0.$$

For simplicity, $\sigma_{w,\tau}^{\dagger}$ is denoted by σ_{w}^{\dagger} below. Similar to the previous section, we see that $y^{*}(x,z) = \infty$ if $\gamma_{0}(x,z) > 0$ whereas $y^{*}(x,z)$ is finite when $\gamma_{0}(x,z) < 0$. For $\gamma_{0}(x,z) = 0$, we assume that $y^{*}(x,z) = \infty$. Furthermore, we can obtain that for any $(x,z) \in \mathcal{X} \times \mathcal{Z}$,

$$\begin{cases} \lim_{w \to \infty} \frac{\sigma_w^{\dagger}(x,z)}{\tau(w|x,z)} = \gamma_0(x,z), & \gamma_0(x,z) > 0, \\ \lim_{w \to \infty} \frac{\sigma_w^{\dagger}(x,z)}{y^*(x,z) - \tau(w|x,z)} = -\gamma_0(x,z), & \gamma_0(x,z) < 0, \\ \lim_{w \to \infty} \sigma_w^{\dagger}(x,z) = \sigma^{\dagger}(x,z), & \gamma_0(x,z) = 0 \end{cases}$$
(5)

for some function $\sigma^{\dagger} : \mathcal{X} \times \mathcal{Z} \to \mathbb{R}_+$ independent from w. As the second-order condition of conditional EVT, we assume that

$$\lim_{w \to \infty} \sup_{(x,z) \in \mathcal{X} \times \mathcal{Z}} \left| \frac{F_w(y|x,z) - H\left(\frac{y}{\sigma_w^{\dagger}(x,z)} | \gamma_0(x,z)\right)}{\alpha(\tau(w|x,z)|x,z)} - Q\left(\frac{y}{\sigma_w^{\dagger}(x,z)} | \gamma_0(x,z), \rho(x,z)\right) \right| = 0 \quad (6)$$

for some function $\rho(x,z) \leq 0$ and $\alpha(\tau(w|x,z)|x,z)$ satisfying $\alpha(\tau(w|x,z)|x,z) \to 0$ as $w \to \infty$. The random variable $Y = Y^* - \tau(w|x,z)$ is approximately distributed as $H(y/\sigma_w^{\dagger}(x,z)|\gamma_0(x,z))$ under Y > 0.

Remark 2. Determining the threshold function τ should be carefully considered in practice. One typical choice of τ is $\tau(w|x, z) = \min\{w, y^*(x, z)\}$, which is intrinsically independent from covariates (x, z). This simple choice would be useful if the dimension of covariates p+d is large. The classical approach for the covariate dependent threshold function is that the conditional quantile of Y^* given (X, Z) = (x, z), denoted by q(a|x, z) with quantile level $a \in (0, 1)$ (see, Koenker (2005), Daouia et al. (2013)). Beirlant et al. (2006) also recommend quantile regression as τ in GPD regression. In this case, we can put $\tau(w|x, z) = q(1 - 1/w|x, z)$.

3 Extreme value generalized additive models

In this section, we provide the estimation method of shape and scale functions.

3.1 Peak over threshold

Let $\{(Y_i^*, X_i, Z_i) : i = 1, ..., N\}$ be *i.i.d.* random sample from same distribution as (Y^*, X, Z) defined in the previous section, where $Y_i^* \in \mathbb{R}_+$, $X_i = (X_i^{(1)}, ..., X_i^{(p)})^T \in \mathcal{X}$ and $Z_i = (Z_i^{(1)}, ..., Z_i^{(d)})^T \in \mathcal{Z}$. For a given threshold function $\tau(w|x, z)$, we let $Y_i = \max\{Y_i^* - \tau(w|X_i, Z_i), 0\}$. We then define $n = \sum_{i=1}^N I(Y_i > 0)$, which is the sample size of the data exceeding the threshold function. Note that n/N converges to $P(Y_i > 0)$ in probability. The estimation method of unknown objects using the data exceeding threshold is the so-called peak over threshold. The conditional distribution of Y_i given $Y_i > 0$ and $(X_i, Z_i) = (x, z)$ is $F_w(y|x, z) = P(Y_i < y|Y_i > 0, x, z)$. Our aim is then to estimate $(\gamma_0, \sigma_w^{\dagger})$ using (6), that is $F_w(y|x, z) \approx H(y/\sigma_w^{\dagger}(x, z))\gamma_0(x, z))$.

Here, we provide the likelihood-based estimation method. The density function associated with H with arbitrary shape and scale function (γ, σ) is obtained as

$$h(y|\gamma(x,z),\sigma(x,z)) := \frac{d}{dy} H\left(\frac{y}{\sigma(x,z)}|\gamma(x,z)\right) = \frac{1}{\sigma(x,z)} \left(1 + \gamma(x,z)\frac{y}{\sigma(x,z)}\right)^{-1/\gamma(x,z)-1}.$$

When $\gamma(x, z) = 0$, $h(y|0, \sigma(x, z)) = \lim_{t\to 0} h(y|t, \sigma(x, z)) = \{1/\sigma(x, z)\} \exp[-y/\sigma(x, z)]$. The log-likelihood for (γ, σ) is

$$\ell(\gamma, \sigma) = \sum_{i=1}^{N} \log h(Y_i | \gamma(X_i, Z_i), \sigma(X_i, Z_i)) I(Y_i > 0).$$

Beirlant and Goegebeur (2004) examined the nonparametric estimation of (γ, σ) using the kernelweighted log-likelihood approach. However, if the dimension of covariate is large, fully nonparametric estimation faces the curse of dimensionality, which leads poor estimation. To avoid this, we introduce the GAM and penalized log-likelihood estimation.

3.2 generalized additive model

For fixed point $x = (x^{(1)}, \ldots, x^{(p)})^T \in \mathcal{X}$ and $z = (z^{(1)}, \ldots, z^{(d)})^T \in \mathcal{Z}$, γ and σ are modeled using the additive model:

$$\gamma(x,z) = \sum_{j=1}^{p} \beta_j x^{(j)} + \sum_{j=1}^{d} g_j(z^{(j)}) = \beta^T x + \sum_{j=1}^{d} g_j(z^{(j)})$$

and

$$\log \sigma(x,z) = \sum_{j=1}^{p} u_j x^{(j)} + \sum_{j=1}^{d} s_j(z^{(j)}) = u^T x + \sum_{j=1}^{d} s_j(z^{(j)}),$$

where $\beta = (\beta_1, \ldots, \beta_p)^T \in \mathbb{R}^p$ and $u = (u_1, \ldots, u_p)^T \in \mathbb{R}^p$ are unknown parameter vectors, and $g_j, s_j : \mathbb{R} \to \mathbb{R}$ are unknown univariate nonparametric functions. For convenience, we assume that $X^{(1)} \equiv 1$ and $X_i^{(1)} \equiv 1(i = 1, \ldots, n)$. Thus, β_1 and u_1 are intercept parameters for γ and $\log \sigma$, respectively. To obtain the identifiability of nonparametric function, we assume that $E[g_j(Z_i^{(j)})] = E[s_j(Z_i^{(j)})] = 0$ for $j = 1, \ldots, d$. The above model in GPD regression is the so-called extreme value generalized additive models (see, Chavez-Demoulin (2005), Youngman (2019)).

We provide the estimation method of $(\beta, u, g_1, \ldots, g_d, s_1, \ldots, s_d)$. In the following, for simplicity, the support of Z is set as $\mathcal{Z} = [0, 1]^d$, that is, $Z^{(j)} \in [0, 1]$ and $Z_i^{(j)} \in [0, 1]$ for all

 $j = 1, \ldots, d$. The nonparametric functions g_j, s_j 's are approximated by the *B*-spline model. Let $0 = \kappa_0 < \kappa_1 < \cdots < \kappa_{K+1} = 1$ be the sequence of knots. In addition, for some $\xi > 0$, we define another 2ξ knots as $\kappa_{-\xi} = \cdots = \kappa_{-1} = \kappa_0$ and $\kappa_{K+1} = \kappa_{K+2} \cdots = \kappa_{K+\xi+1}$. For simplicity, we assume that the location of knots is equidistant, that is, $\kappa_j - \kappa_{j-1} = 1/(K+1)$. Then, let $\{\psi_0^{[\xi]}(\cdot), \ldots, \psi_{K+\xi}^{[\xi]}(\cdot)\}$ be ξ th degree or $(\xi + 1)$ th order *B*-spline bases, where $\psi_k^{[\xi]} : [0, 1] \to \mathbb{R}_+$. The definition and some basic properties of *B*-spline bases are clarified in de Boor (2001). We next transform the ordinary *B*-spline bases $\{\psi_0^{[\xi]}(\cdot), \ldots, \psi_{K+\xi}^{[\xi]}(\cdot)\}$ to the normalized *B*-spline bases (see, Liu et al. 2011). For $j = 1, \ldots, d$ and $k = 1, \ldots, K + \xi - 1$, we define

$$B_{j,k}^{[\xi]}(\tilde{z}) = \frac{\bar{\psi}_{j,k}^{[\xi]}(\tilde{z})}{||\bar{\psi}_{j,k}^{[\xi]}||}, \quad k = 1, \dots, K + \xi,$$

where $\bar{\psi}_{j,k}^{[\xi]}(\tilde{z}) = \psi_k^{[\xi]}(\tilde{z}) - (\phi_{j,k}/\phi_{j,k-1})\psi_{k-1}^{[\xi]}(\tilde{z})$ and $\phi_{j,k} = E[\psi_k^{[\xi]}(Z_i^{(j)})]$. We note that the definition of ordinary *B*-spline bases $\psi_k^{[\xi]}$'s are common for all *j*, but this is not true for the normalized *B*-spline bases $B_{jk}^{[\xi]}$'s since $\phi_{j,k}$ depends on distribution of $Z^{(j)}$. For $\tilde{z} \in [0, 1]$, the normalized *B*-spline model is defined as

$$\bar{g}_j(\tilde{z}) = \sum_{k=1}^{K+\xi} B_{j,k}^{[\xi]}(\tilde{z}) b_{j,k} = B_j(\tilde{z})^T b_j$$

and

$$\bar{s}_j(\tilde{z}) = \sum_{k=1}^{K+\xi} B_{j,k}^{[\xi]}(\tilde{z}) c_{j,k} = B_j(\tilde{z})^T c_j,$$

where $B_j(\tilde{z}) = (B_{j,1}^{[\xi]}(\tilde{z}), \ldots, B_{j,K+\xi}^{[\xi]}(\tilde{z}))^T$, and $b_j = (b_{j,1}, \ldots, b_{j,K+\xi})^T$ and $c_j = (c_{j,1}, \ldots, c_{j,K+\xi})^T$ are $(K + \xi)$ -unknown parameter vectors. By the definition of $B_{j,k}^{[\xi]}$, we can easily confirm that $E[\bar{g}_j(Z^{(j)})] = E[\bar{s}_j(Z^{(j)})] = 0$. We then consider that the additive functions g_j, s_j are approximated by the normalized *B*-spline model: $g_j \approx \bar{g}_j$ and $s_j \approx \bar{s}_j$ for $j = 1, \ldots, d$. Then, for fixed point $x \in \mathbb{R}^p$ and $z = (z^{(1)}, \ldots, z^{(d)})^T \in [0, 1]^d$, γ and σ are approximated by

$$\bar{\gamma}(x,z) = x^T \beta + \sum_{j=1}^d \bar{g}_j(z^{(j)}) = x^T \beta + \sum_{j=1}^d B_j(z^{(j)})^T b_j$$

and

$$\log \bar{\sigma}(x,z) = x^T u + \sum_{j=1}^d \bar{s}_j(z^{(j)}) = x^T u + \sum_{j=1}^d B_j(z^{(j)})^T c_j$$

Accordingly, our purpose is to estimate the parameter vector (β, u, b, c) , where $b = (b_1^T, \dots, b_d^T)^T \in \mathbb{R}^{p+K+\xi}$ and $c = (c_1^T, \dots, c_d^T)^T \in \mathbb{R}^{p+K+\xi}$.

The minus log-likelihood of (β, u, b, c) is

$$\ell(\beta, u, b, c) = -\sum_{i=1}^{N} \log h(Y_i | \bar{\gamma}(X_i, Z_i), \bar{\sigma}(X_i, Z_i)) I(Y_i > 0)$$

Then, (β, u, b, c) is estimated by minimizing the penalized log-likelihood

$$\ell_{pen}(\beta, u, b, c) = \ell(\beta, u, b, c) + \sum_{j=1}^{d} \left\{ \lambda_j \int_0^1 \{\bar{g}_j^{(m)}(z)\}^2 dz + \nu_j \int_0^1 \{\bar{s}_j^{(m)}(z)\}^2 dz \right\},\tag{7}$$

where λ_j 's and ν_j 's are smoothing parameters and m is some integer smaller than ξ , where ξ is the degree of *B*-spline function. In fact, if $m > \xi$, $\bar{\gamma}_j^{(m)} = \bar{s}_j^{(m)} \equiv 0$. The estimator obtained from (7) is denoted by $(\hat{\beta}, \hat{u}, \hat{b}, \hat{c})$, where $\hat{b} = (\hat{b}_1^T, \dots, \hat{b}_d^T)^T$ and $\hat{c} = (\hat{c}_1^T, \dots, \hat{c}_d^T)^T$. From these estimators, we construct

$$\hat{\gamma}(x,z) = x^T \hat{\beta} + \sum_{j=1}^d \hat{g}_j(z^{(j)})$$

and

$$\hat{\sigma}(x,z) = \exp\left[x^T \hat{u} + \sum_{j=1}^d \hat{s}_j(z^{(j)})\right],\,$$

where $\hat{g}_j(z^{(j)}) = B^{[\xi]}(z^{(j)})^T \hat{b}_j$ and $\hat{s}_j(z^{(j)}) = B^{[\xi]}(z^{(j)})^T \hat{c}_j$. The above estimator can be implemented via the R-package evgam proposed by Youngman (2022). Thus, numerical performance of $\hat{\gamma}$ and $\hat{\sigma}$ is already guaranteed by evgam. Meanwhile, there is no result of the theoretical evidence of $\hat{\gamma}$ and $\hat{\sigma}$ thus far. We establish the asymptotic theory for $\hat{\gamma}$ and $\hat{\sigma}$ in the next section. For simplicity, the tuning parameters for each shape and scale function are assumed to be common among all covariates, denoted by $\lambda = \lambda_1 = \cdots = \lambda_d > 0$ and $\nu = \nu_1 = \cdots = \nu_d > 0$.

4 Asymptotic theory

4.1 General condition

We first define the functional space of the additive model. Let

$$\mathcal{A}_j = \{ \alpha : [0,1] \to \mathbb{R} | E[\alpha(Z_i^{(j)})] = 0, V[\alpha(Z_i^{(j)})] < \infty \}$$

We further let

$$\mathcal{T}_A = \left\{ f : \mathcal{X} \times \mathcal{Z} \to \mathbb{R} | f(x, z) = x^T \beta + \sum_{j=1}^d q_j(z^{(j)}), q_j \in \mathcal{A}_j, \beta \in \mathbb{R}^p \right\}.$$

We then assume that for any $(x, z) \in \mathcal{X} \times \mathcal{Z}$, $F(\cdot|x, z) \in \mathcal{D}(G(\cdot|\gamma_0(x, z)))$ with $\gamma_0 \in \mathcal{T}_A$. However, although $\hat{\sigma}(x, z)$ is constructed via the additive model, we cannot know whether $\log \sigma_w^{\dagger} \in \mathcal{T}_A$. Therefore, we use the alternative target scale function to investigate the asymptotic behavior of $\hat{\sigma}$. Define the target functions as

$$(\gamma_0, \sigma_{w0}) = \operatorname*{argmin}_{(\gamma, \log \sigma) \in \mathcal{T}_A \times \mathcal{T}_A} - E[\log h(Y|\gamma(X, Z), \sigma(X, Z))].$$
(8)

Furthermore, we let

$$\varepsilon_w(x,z) = \frac{\sigma_{w0}(x,z)}{\sigma_w^{\dagger}(x,z)} - 1.$$

We note that $\varepsilon_w(x, z)$ depends on the signature of γ_0 . Then, $\varepsilon_w(x, z)$ can be seen as model bias between true function σ_w^{\dagger} and the best approximation of the additive model σ_{w0} . We assume that for all $(x, z) \in \mathcal{X} \times \mathcal{Z}$, $|\varepsilon_w(x, z)|$ is quite small so that σ_{w0} can be identified with σ_w^{\dagger} . For simplicity, we assume that $\varepsilon_w(x, z) = 0$ for all $(x, z) \in \mathcal{X} \times \mathcal{Z}$ below. Actually, $\varepsilon_w(x, z)$ cannot be evaluated or improved without any prior information. Using this and (5), σ_{w0} holds

$$\lim_{w \to \infty} \frac{\sigma_{w0}(x,z)}{\tau(w|x,z)} = O(1), \qquad \gamma_0(x,z) > 0, \\
\lim_{w \to \infty} \frac{\sigma_{w0}(x,z)}{y^*(x,z) - \tau(w|x,z)} = O(1) \quad \gamma_0(x,z) < 0, \\
\lim_{w \to \infty} \sigma_{w0}(x,z) = \sigma_0(x,z), \qquad \gamma_0(x,z) = 0,$$
(9)

where $\log \sigma_0 \in \mathcal{T}_A$.

We continue about the discussion on the behavior of σ_{w0} . Since σ_{w0} depends on w, the true coefficients of the parametric part and each additive function of the nonparametric part in $\log \sigma_{w0}$ may also vary with w. Thus, for γ_0 and σ_{w0} , there exist $\beta_0 = (\beta_{01}, \ldots, \beta_{0p})^T, u_{w0} = (u_{w,01}, \ldots, u_{w,0p})^T \in \mathbb{R}^p$ and $g_{0j}, s_{w,0j} \in \mathcal{A}_j (j = 1, \ldots, d)$ such that for any $x = (x^{(1)}, \ldots, x^{(p)}) \in \mathcal{X}$ with $x^{(1)} = 1$ and $z = (z^{(1)}, \ldots, z^{(d)}) \in \mathcal{Z}$,

$$\gamma_0(x,z) = x^T \beta_0 + \sum_{j=1}^d g_{0j}(z^{(j)})$$
(10)

and

$$\log \sigma_{w0}(x,z) = x^T u_{w0} + \sum_{j=1}^d s_{w,0j}(z^{(j)}).$$
(11)

For (11), we assume that coefficients except for intercept and each additive function are independent from w, denoted by $u_{w,0j} = u_{0j}$ for j = 2, ..., p and $s_{w,0k} = s_{0k}$ for k = 1, ..., d. That is, (11) can be written as

$$\log \sigma_{w0}(x,z) = u_{w,01} + \sum_{j=2}^{p} u_{0j} x^{(j)} + \sum_{j=1}^{d} s_{0j}(z^{(j)}).$$
(12)

For $\gamma_0(x,z) > 0$, if we choose τ as $\tau(w|x,z) = a_1(w)\tau_1(x,z)$ for some functions $a_1 : \mathbb{R}_+ \to \mathbb{R}_+$ and $\tau_1 : \mathcal{X} \times \mathcal{Z} \to \mathbb{R}_+$ with $a_1(w) \to \infty(w \to \infty)$, (12) holds with $u_{w,01} = \log a_1(w) + u_0, u_0 \in \mathbb{R}$. A simple but important example is $\tau(w|x,z) = w$. For $\gamma_0(x,z) < 0$, we obtain (12) when τ can be written as $y^*(x,z) - \tau(w|x,z) = a_2(w)\tau_2(x,z)$ for some functions $a_2 : \mathbb{R}_+ \to \mathbb{R}_+$ and $\tau_2 : \mathcal{X} \times \mathcal{Z} \to \mathbb{R}_+$ with $a_2(w) \to 0(w \to \infty)$. Since $y^*(x,z)$ is unknown, identifying such a_2 and τ_2 is difficult in practice.

Remark 3. We provide additional explanation of the assumption of (12). For simplicity, we assume that $\gamma_0(z) > 0$ for all $z = (z^{(1)}, z^{(2)}) \in [0, 1]^2$ and (11) as $\sigma_{w0}(z) = \exp[u_{w0} + s_{w,01}(z^{(1)}) + s_{w,02}(z^{(2)})]$. Then, from (9), roughly speaking, $s_{w,01}$ may tend to infinity since $\tau(w|z) \to \infty$ and $\sigma_{w0} \to \infty$ as $w \to \infty$. However, from the condition $E[s_{w,01}(Z^{(1)})] = 0$ for identifiability of function, if $s_{w,01}(z^*) \to \infty$ for a point $z^* \in [0, 1]$, another point $z^{**} \in [0, 1]$ exists such that $s_{w,01}(z^{**}) \to -\infty$. Nevertheless, for all $z \in [0, 1]^2$, $\sigma_{w0}(z)$ needs to diverge to infinity as $w \to \infty$. This yields that $s_{w,01}(z^{**})/u_{w0} \to 0$ as $w \to \infty$. Thus, u_{w0} should dominate other components as the sequence of w. Furthermore, $s_{w,0j} = O(1)$ or $s_{w,0j} = o(1)$ can also be allowed as long as $\sigma_{w0} \to \infty$. In fact, if $\tau(w|z) = w$, we can obtain that $u_{w0} = \log w + u_0$ with $u_0 \in \mathbb{R}$ and $s_{w,0j} = O(1)$ from (9). Consequently, the construction of $s_{w,0j}$ as the sequence of w becomes quite complicated in (11), and (12) is simplest model with no contradiction as (9).

The behavior of σ_{w0} depends on the signature of γ_0 . From (9), if $\gamma_0(x,z) > 0$, $\sigma_{w0}(x,z) \to \infty$, and if $\gamma_0(x,z) < 0$, we have $\sigma_{w0}(x,z) \to 0$. Therefore, if $\gamma_0(x_0,z_0) = 0$ at the point $(x_0,z_0) \in \mathcal{X} \times \mathcal{Z}$ and $\gamma_0(x,z)$ can take both positive and negative at the neighbourhood of (x_0,z_0) , $\sigma_{w0}(x,z)$ is not continuous at (x_0,z_0) although $\log \sigma$ is modeled by continuous additive functions. Thus, when we consider the additive model for scale function, the signature of shape function is expected to be fixed even if its signature is unknown. We consider the following three cases separately: (i) $\inf_{(x,z)\in\mathcal{X}\times\mathcal{Z}}\gamma_0(x,z) > 0$, (ii) $\sup_{(x,z)\in\mathcal{X}\times\mathcal{Z}}\gamma_0(x,z) < 0$ and (iii) $\gamma_0(x,z) \equiv 0$ for all $(x,z) \in \mathcal{X} \times \mathcal{Z}$. One may consider case (iii) to be unrealistic, but if $F(\cdot|x,z)$ is the Gaussian distribution, it belongs to $\mathcal{D}(G(\cdot|\gamma_0(x,z)))$ with $\gamma_0(x,z) = 0$. If the signature of γ_0 varies among $(x,z) \in \mathcal{X} \times \mathcal{Z}$, the additive modeling for σ_{w0} would not be performed better. Such case is beyond the scope of this study. Thus, we first state following conditions concerned with positive, negative, and zero shape function.

- (S1) Constant $\gamma_{min} > 0$ exists such that $\gamma_{min} < \gamma_0(x, z)$ for all $(x, z) \in \mathcal{X} \times \mathcal{Z}$.
- (S2) For some $\delta > 0$, $-1/(2+\delta) < \gamma_0(x,z) < 0$ for all $(x,z) \in \mathcal{X} \times \mathcal{Z}$.
- (S3) For all $(x, z) \in \mathcal{X} \times \mathcal{Z}, \gamma_0(x, z) = 0.$

We establish the asymptotic theory of $(\hat{\gamma}, \hat{\sigma})$ under each condition among (S1), (S2), or (S3).

As the true setting, we assume that (6), (8), (9), and (12). In addition, we state the technical conditions to investigate the asymptotic property of the estimator for all cases (S1)–(S3). Let $B(Z) = (B_1(Z^{(1)})^T, \ldots, B_d(Z^{(d)})^T)^T$ be $d(K + \xi)$ -vector of *B*-spline bases. For a > 0, let C^a be the class of functions with *a*th continuously differentiable on [0,1]. We note that te sample size $n = \sum_{i=1}^{N} I(Y_i > \tau(w|X_i, Z_i))$ is the random variable. For easy interpretation, the following condition is given by *n*. However, *n* can be replaced with $NP(Y_i > 0) = NP(Y_i^* > \tau(w|X_i, Z_i))$ below.

- (C1) Without loss of generality, for j = 1, ..., p, $E[X^{(j)}] = 0$. Furthermore, the matrix $E[(X, B(Z))(X^T B(Z)^T)^T]$ is positive definite.
- (C2) Let $\zeta > m$, where *m* is appeared in the penalty term of (7). Then, $g_{0k} \in \mathcal{A}_k \cap \mathcal{C}^{\zeta}$ and $s_{0k} \in \mathcal{A}_k \cap \mathcal{C}^{\zeta}$ for $k = 1, \ldots, d$.
- (C3) As $N \to \infty$, $n \to \infty$ and $n/N \to 0$.
- (C4) In (6), there exists a continuous and bounded function $\alpha_1(x,z)$ such that $\alpha(\tau(w_N \mid x, z) \mid x, z) = \alpha_1(x,z)P(Y_i > 0)^{-\rho(x,z)}(1 + o(1))$ as $N \to \infty$. In addition, there exists a constant $\rho < 0$ such that $\sup_{(x,z) \in \mathcal{X} \times \mathcal{Z}} \rho(x, z) \leq \rho < 0$.
- (C5) The number of knots $K = K_N$ and smoothing parameters $\lambda = \lambda_N$ and $\nu = \nu_N$ satisfy $K \to \infty$, $K \log(N)/n \to 0$, $\lambda K^{2m} = O(1)$ and $\nu K^{2m} = O(1)$ as $N \to \infty$.

(C1) is the standard condition for covariates in the regression. Lemma 2 in Appendix B shows that $E[B(Z)B(Z)^T]$ is positive definite. Therefore, the last assumption of (C1) is natural. (C2) defines the smoothness of true additive function. Since we use the *m*th derivative of spline in (7), the order of smoothness of the true function is needed to be larger than *m*. (C3) explains the condition of sample size of data exceeding threshold, is quite popular in EVT (see, Drees et al. (2004)). (C3) can be reexpressed by that $w = w_N$ satisfies $NP(Y_i^* > \tau(w_N | X_i, Z_i)) \to 0$ and $P(Y_i^* > \tau(w_N | X_i, Z_i)) \to 0$ as $N \to \infty$. More detailed discussion about (C3) is in 4 (C4) controls the asymptotic order of bias of the estimator. We provide some justification of first condition of (C4) in Remark 5. The second condition of (C4) means that the case $\rho(x, z) = 0$ is removed. If $\rho(x, z) = 0$, this corresponds to $\alpha(\tau(w_N \mid x, z) \mid x, z) = O(\log P(Y_i > 0))$ for example. This rate is too slow to explain the efficiency of the estimator. (C5) examines the rates of tuning parameters to obtain the optimal rate of convergence of the estimator. Xiao (2019) uses similar conditions of the number of knots and smoothing parameters.

Remark 4. For condition (C3), we have $P(Y_i > 0) = E_{X,Z}[P(Y_i^* > \tau(w_N|x,z)|x,z)|X = x, Z = z] = E_{X,Z}[1 - F(\tau(w_N|X,Z)|X,Z)]$, where $E_{X,Z}$ is the expectation operator with the distribution function of (X,Z). According to Theorem 1.2.1 of de Haan and Ferreira (2006), we can assume that there exists a function $c_1(x,z) > 0$ such that $1 - F(\tau(w_N|x,z)|x,z) \approx c_1(x,z)\tau(w_N|x,z)^{-1/\gamma_0(x,z)}$ under (S1) or (S2). This suggests that

$$P(Y_i > 0) \approx E_{X,Z}[c_1(X,Z)\tau(w_N|X,Z)^{-1/\gamma_0(X,Z)}].$$

Thus, $P(Y_i > 0) \approx n/N$ depends on γ_0 , and hence the distribution of n is also unknown. Therefore, (C3) seems to be somewhat strong. However, in the context of data analysis, (C3) is natural since the sample size of extreme value data tends to be relatively small compared to the total number of data N. See also Theorem 1 of Wang and Tsai (2009).

Remark 5. We note in (C4). For t > 0, let $U(t \mid x, z) = F^{-1}(1 - 1/t \mid x, z)$. Then, the equivalence condition to (6) is that there exists a function A such that

$$\lim_{t \to \infty} \sup_{(x,z) \in \mathcal{X} \times \mathcal{Z}} \left| \frac{\frac{U(t\eta|x,z) - U(t|x,z)}{a_t(x,z)} - \frac{(\eta^{-\gamma_0(x,z)} - 1)}{\gamma_0(x,z)}}{A(t \mid x, z)} - \tilde{Q}(\eta \mid \gamma_0(x,z), \rho(x,z)) \right| \to 0$$

for all $\eta > 0$, where $a_t(x, z) = \sigma_w^{\dagger}(x, z)$ with $\tau(w \mid x, z) = U(t \mid x, z)$ and the function $A(t \mid x, z)$ satisfying $A(\eta t \mid x, z)/A(t \mid x, z) \to \eta^{\rho(x,z)}$ as $t \to \infty$. Such a result is given by Theorem 2.3.3 ans 2.3.8 of de Haan and Ferreira (2006). Thus, we can assume that $A(t \mid x, z) = \alpha_1(x, z)t^{\rho(x,z)}(1 + o(1))$ as $t \to \infty$. Although we can consider another choice of A such as $A(t \mid x, z) = \alpha_1(x, z)t^{\rho(x,z)}\log t$, this case is omitted in this paper. From Theorem 2.3.8 of de Haan and Ferreira (2006), we obtain $\alpha(t \mid x, z) = A(1/\{1 - F(t \mid x, z)\} \mid x, z)$. This implies that $\alpha(\tau(w \mid x, z) \mid x, z) = A(1/\{1 - F(\tau(w \mid x, z))\} \mid x, z) = A(1/P(Y_i > 0) \mid x, z)$. Therefore, $\alpha(\tau(w \mid x, z) \mid x, z) = \alpha_1(x, z)P(Y_i > 0)^{-\rho(x,z)}(1 + o(1))$ is one of natural assumption. Since $P(Y_i > 0) \sim n/N$, it can be interpreted by $\alpha(\tau(w \mid x, z) \mid x, z) = \alpha_1(x, z)(N/n)^{\rho(x,z)}(1 + o(1))$.

4.2 Main result

For any function $r: \mathcal{X} \times \mathcal{Z} \to \mathbb{R}$, we define L_2 -norm of r as $||r||_{L_2} = \sqrt{E[r(\tilde{X}, \tilde{Z})^2]}$, where (\tilde{X}, \tilde{Z}) is the random variable having the same distribution as (X, Z) independently. If r depends on sample $\{(Y_i, X_i, Z_i) : i = 1, ..., n\}$, E takes the expectation for not only (\tilde{X}, \tilde{Z}) but also (Y_i, X_i, Z_i) . The L_{∞} -norm of r is defined as $||r||_{\infty} = \sup_{(x,z)\in\mathcal{X}\times\mathcal{Z}} |r(x,z)|$. We first describe the L_2 -rate of convergence of the estimator.

Theorem 1. Suppose that (C1)–(C5). In each scenario (S1), (S2) or (S3), as $N \to \infty$,

$$\begin{aligned} ||\hat{\gamma} - \gamma_0||_{L_2} &\leq O\left(\sqrt{\frac{K}{n}}\right) + O(\lambda K^m) + O(\nu K^m) + O\left(\left(\frac{N}{n}\right)^{\rho}\right), \\ ||\log\hat{\sigma} - \log\sigma_{w0}||_{L_2} &\leq O\left(\sqrt{\frac{K}{n}}\right) + O(\lambda K^m) + O(\nu K^m) + O\left(\left(\frac{N}{n}\right)^{\rho}\right). \end{aligned}$$

Under the optimal rate of number of knots $K = O(n^{1/(2m+1)})$,

$$||\hat{\gamma} - \gamma_0||_{L_2} \leq O\left(n^{-m/(2m+1)}\right) + O\left(\left(\frac{N}{n}\right)^{\rho}\right),$$

$$||\log\hat{\sigma} - \log\sigma_{w0}||_{L_2} \leq O\left(n^{-m/(2m+1)}\right) + O\left(\left(\frac{N}{n}\right)^{\rho}\right).$$

As the threshold $w = w_N$ is selected so that n takes the optimal order: $n = O(N^{-\rho(2m+1)/\{m-\rho(2m+1)\}}),$

$$\begin{aligned} ||\hat{\gamma} - \gamma_0||_{L_2} &\leq O\left(N^{\frac{\rho}{1-2\rho+1/m}}\right), \\ |\log\hat{\sigma} - \log\sigma_{w0}||_{L_2} &\leq O\left(N^{\frac{\rho}{1-2\rho+1/m}}\right). \end{aligned}$$

Next, we investigate L_{∞} -convergence of the estimator.

Theorem 2. Suppose that (C1)–(C5). Under (S2), suppose that $\{K \log N\}^{1+\delta/2}/n^{\delta/2} \to 0$ as $N \to \infty$. Then, in each scenario (S1), (S2) or (S3), as $N \to \infty$,

$$\begin{aligned} ||\hat{\gamma} - \gamma_0||_{L_{\infty}} &\leq O\left(\sqrt{\frac{K\log N}{n}}\right) + O(\lambda K^m) + O(\nu K^m) + O\left(\left(\frac{N}{n}\right)^{\rho}\right), \\ ||\log \hat{\sigma} - \log \sigma_{w0}||_{L_{\infty}} &\leq O\left(\sqrt{\frac{K\log N}{n}}\right) + O(\lambda K^m) + O(\nu K^m) + O\left(\left(\frac{N}{n}\right)^{\rho}\right). \end{aligned}$$

Under the optimal rate of number of knots $K = O((n/\log N)^{1/(2m+1)})$,

$$||\hat{\gamma} - \gamma_0||_{L_{\infty}} \leq O\left((n/\log N)^{-m/(2m+1)}\right) + O\left(\left(\frac{N}{n}\right)^{\rho}\right),$$

$$||\log\hat{\sigma} - \log\sigma_{w0}||_{L_{\infty}} \leq O\left((n/\log N)^{-m/(2m+1)}\right) + O\left(\left(\frac{N}{n}\right)^{\rho}\right).$$

As the threshold $w = w_N$ is selected so that n takes the optimal order: $n = O((N/\log N)^{-\rho(2m+1)/\{m-\rho(2m+1)\}}),$

$$\begin{aligned} ||\hat{\gamma} - \gamma_0||_{L_{\infty}} &\leq O\left(\left(\frac{N}{\log N}\right)^{\frac{\rho}{1-2\rho+1/m}}\right),\\ ||\log\hat{\sigma} - \log\sigma_{w0}||_{L_{\infty}} &\leq O\left(\left(\frac{N}{\log N}\right)^{\frac{\rho}{1-2\rho+1/m}}\right). \end{aligned}$$

Remark 6. In Theorem 2. under (S2), we need the extra condition: $\{K \log N\}^{1+\delta/2}/n^{\delta/2} \to 0$. This condition is for the technical reason to prove L_{∞} -convergence of the estimator (see Appendix C). When $K = O((n/\log N)^{1/(2m+1)})$ is used, we have

$$\{K \log N\}^{1+\delta/2} = O\left(n^{\frac{2+\delta}{4m+2}} (\log N)^{\frac{2m(2+\delta)}{4m+2}}\right).$$

Then, $\{K \log N\}^{1+\delta/2}/n^{\delta/2} \to 0$ holds if $\delta > 1/m$. This implies that (S2) is replaced with $\gamma_0(x,z) > -m/(2m+1)$ for some $m \in \mathbb{N}$. In the spline method, we often use m = 2, which indicates that (S2) becomes $0 > \gamma_0(x,z) > -2/5$. Thus, when γ_0 is estimated via the penalized maximum-likelihood method, the bound condition of the negative shape function is somewhat stronger than that for the non-regression case: $\gamma_0 > -1/2$ (see, Drees et al. (2004)).

Next, the local asymptotic normality of the estimator is analyzed. For $(x, z) \in \mathcal{X} \times \mathcal{Z}$, let $(p + K + \xi)$ -vector $A(x, z) = (x^T, B(z)^T)^T$ and let $2(p + K + \xi) \times 2$ matrix

$$D(x,z) = \begin{bmatrix} (x^T, B(z)^T)^T & 0_{p+K+\xi}^T \\ 0_{p+K+\xi}^T & (x^T, B(z)^T)^T \end{bmatrix}^T,$$

where $0_{p+K+\xi}$ is the $(p+K+\xi)$ -zero vector. We further let

$$\Sigma = E \left[\frac{1}{2\gamma_0(X,Z) + 1} \left(\begin{array}{c} \frac{2}{\gamma_0(X,Z) + 1} A(X,Z) A(X,Z)^T & \frac{1}{\gamma_0(X,Z) + 1} A(X,Z) A(X,Z)^T \\ \frac{1}{\gamma_0(X,Z) + 1} A(X,Z) A(X,Z)^T & A(X,Z) A(X,Z)^T \end{array} \right) \right] + \Omega_{\gamma,\sigma},$$

where $\Omega_{\gamma,\sigma}$ is defined in the proof of Lemma 6. Since each element of B(z) has an order $O(\sqrt{K})$ and the element of Σ^{-1} has O(1), we obtain $D(x,z)^T \Sigma^{-1} D(x,z)/K = O(1)$.

Theorem 3. Suppose that (C1)–(C5). Furthermore, assume that (K, w_N) satisfies

$$K = O\left(\left(\frac{n}{(\log n)^{2m}}\right)^{1/(2m+1)}\right)$$

and $(n/\log n)^{m/(2m+1)}(N/n)^{\rho} \to 0$ as $N \to \infty$. Then, in each scenario (S1), (S2) or (S3), for any $(x, z) \in \mathcal{X} \times \mathcal{Z}$,

$$\left(\frac{n}{\log n}\right)^{m/(2m+1)} \begin{bmatrix} \hat{\gamma}(x,z) - \gamma_0(x,z) \\ \frac{\hat{\sigma}(x,z)}{\sigma_{w0}(x,z)} - 1 \end{bmatrix} \xrightarrow{D} N_2\left(0, \lim_{N \to \infty} D(x,z)^T \Sigma^{-1} D(x,z) / K\right),$$

as $N \to \infty$, where N_2 means the bivariate normal distribution.

In Theorem 3, the bias disappears because of the choice of (K, w_N) . In fact, the order of the bias of the estimator is $O(K^{-m}) + O((N/n)^{\rho})$ whereas the standard deviation has $O(\sqrt{K/n})$. From Theorem 1, $K = O(n^{1/(2m+1)})$ balances $O(\sqrt{K/n}) + O(K^{-m})$. The choice of $K = O(n^{1/(2m+1)}/(\log n)^{2m/(2m+1)})$ is slightly larger than $O(n^{1/(2m+1)})$, and hence, the term $O(\sqrt{K/n}) = O((n/\log n)^{-m/(2m+1)})$ dominates the bias $O(K^{-m})$. On the other hand, if $(n/\log n)^{m/(2m+1)}(N/n)^{\rho} \to 0$, the term $O((N/n)^{\rho})$ disappears compared with the order of the standard deviation $O((n/\log n)^{-m/(2m+1)})$. Thus, the condition of (K, w_N) implies that the asymptotic order of the standard deviation of the estimator is slightly larger than that of the bias. Since the form of bias is complicated, deriving its consistent estimator is quite challenging. Therefore, we establish the variance dominated asymptotic normality of the estimator $(\hat{\gamma}, \hat{\sigma})$. Drees et al. (2004) proved that the estimator has an order $O(n^{-1/2})$ in non-regression case of GPD model. Compared with their result, the rate $O((n/\log n)^{-m/(2m+1)})$ is that of the parametric estimator.

Let 2p-square matrix

$$\Sigma_{\beta,u} = E \left[\frac{1}{2\gamma_0(X,Z) + 1} \left(\begin{array}{cc} \frac{2}{\gamma_0(X,Z) + 1} X X^T & \frac{1}{\gamma_0(X,Z) + 1} X X^T \\ \frac{1}{\gamma_0(X,Z) + 1} X X^T & X X^T \end{array} \right) \right].$$

If one wants to focus the linear part of the estimator, the following result would be helpful.

Theorem 4. Suppose that (C1)–(C5). Furthermore, we choose (K, w_N) such that $K \ge O(n^{1/2m} \log n)$ as $N \to \infty$. Then, in each scenario (S1), (S2) or (S3), if $n^{1/2}(N/n)^{\rho} = O(1)$, as $N \to \infty$,

$$\begin{aligned} ||\hat{\beta} - \beta_0|| &\leq O(n^{-1/2}) + O((N/n)^{\rho}) = O(N^{\rho/(1-2\rho)}), \\ ||\hat{u} - u_{w,0}|| &\leq O(n^{-1/2}) + O((N/n)^{\rho}) = O(N^{\rho/(1-2\rho)}). \end{aligned}$$

Furthermore, for each (S1), (S2) or (S3), if $n^{1/2}(N/n)^{\rho} \to 0$, as $N \to \infty$,

$$\sqrt{n} \left[\begin{array}{c} \hat{\beta} - \beta_0 \\ \hat{u} - u_{w,0} \end{array} \right] \xrightarrow{D} N_{2p}(0, \Sigma_{\beta, u}^{-1}).$$

Theorem 4 is the optimal result for the parametric part of the estimator for the both shape and scale function. We note that the condition $K > O(n^{1/2m} \log n)$ is not optimal choice for the nonparametric part, which indicates that the bias is so small to be negligible. Other studies have reported such technique as the important property of the partially linear additive models (e.g. Theorem 4 in Liu et al. (2017)). We can interpret that Theorem 4 is the regression coefficient version of the result in Corollary 2.1 in Drees et al. (2004). The condition $n^{1/2}(N/n)^{\rho} \rightarrow 0$ means that the standard deviation of the estimator of parametric part dominates the bias appeared from second-order condition of EVT. If we choose large w_N , such situation can be obtained. Even when the optimal w_N is chosen, deriving the explicit form of the bias is difficult and has quite complicated expression since the bias from second-order condition of EVT contains not only the parametric part but also nonparametric part. Thus, in the second assertion of Theorem 4, we established the simple but convenient result as a no-biased version.

Remark 7. If $\gamma_0(x, z) = \gamma_0$ and we use non-covariates in analysis, Theorems 1 and 4 can reduce to the result in Smith (1987). In this case, our result is also related to Drees et al. (2004), however, they utilized the threshold value as random variable unlike deterministic threshold used in our study. Therefore, the situation in our study is somewhat different from Drees et al. (2004). Meanwhile, if we know that $\gamma(x, z)$ is positive, GPD can be reformulated as Pareto-type distribution. Therefore, our model with a non-nonparametric part connects to the result of Wang and Tsai (2009) whereas Li et al. (2022) is related to our model with d = 1. Thus, the results in this section can be widely applied to related models.

Remark 8. In Theorem 1, the asymptotic biases of the estimator occurring from spline approximation and penalisation have order $O(K^{-\zeta})$ and $O((\lambda + \nu)K^m)$ although the term $O(K^{-\zeta})$ is negligible under (C2) and (C5). According to Claeskens et al. (2009), such situation can be explained in that the rates of convergence of the penalized B-spline estimator are equivalent to that of the smoothing spline estimator (see, Green and Silverman (1994)). Claeskens et al. (2009) and Xiao (2019) refer to the above setting as a large K scenario. Meanwhile, we can establish the asymptotic result of the estimator with another setting as $\max\{\lambda,\nu\} \leq o(K^{-m-\zeta}) = o(K^{-2m})$. Then, the order of spline approximation $O(K^{-\zeta})$ dominates that of penalty $O((\lambda + \nu)K^m)$. This situation matches the result for unpenalized B-spline regression (see, Zhou et al. (1998)). Then, the optimal L₂-rate of the estimator becomes $O(n^{-\zeta/(2\zeta+1)}) + O((N/n)^{\rho})$ and $O(N^{\rho/(1-\rho+1/\zeta)})$ under $K = O(n^{1/(2\zeta+1)})$. Since $\zeta > m$ and $K = O(n^{1/(2\zeta+1)}) < O(n^{1/(2m+1)})$, this situation means that the number of knots cannot be set as large. From this, Claeskens et al. (2009) and Xiao (2019) refer to this condition as the small K scenario. The estimator obtained via evgam is based on the penalized spline method including smoothing spline, and hence, we mainly adopt the large K scenario. However, we can easily use the unpenalized spline method via evgam (see also R-package mgcv, Wood 2017), the result for small K scenario would also be important.

5 Reparametrization

As shown in Chavez-Demoulin and Davison (2005), we provide the reparametrization of $(\gamma_0, \sigma_w^{\dagger})$ in (6) to $(\gamma_0, \varsigma_w^{\dagger})$, where $\varsigma_w^{\dagger}(x, z) = \sigma_w^{\dagger}(x, z)(\gamma_0(x, z) + 1)$. If $\gamma_0(x, z)$ is greater than -1 for all $(x, z) \in \mathcal{X} \times \mathcal{Z}, \varsigma_w^{\dagger}(x, z) > 0$ can be guaranteed. We note that this transformation is identity under (S3). The equivalent condition as (6) is that $(\gamma_0, \varsigma_w^{\dagger})$ satisfies

$$\lim_{w \to \infty} \sup_{(x,z) \in \mathcal{X} \times \mathcal{Z}} \left| \frac{F_w(y|x,z) - H\left(\frac{(\gamma_0(x,z)+1)y}{\varsigma_w^{\dagger}(x,z)} | \gamma_0(x,z)\right)}{\alpha(\tau(w|x,z)|x,z)} - Q\left(\frac{(\gamma_0(x,z)+1)y}{\varsigma_w^{\dagger}(x,z)} | \gamma_0(x,z), \rho(x,z)\right) \right| \to 0.(13)$$

Furthermore, (9) implies that

$$\begin{cases} \lim_{w \to \infty} \frac{\zeta_w^{\dagger}(x,z)}{\tau(w|x,z)} = \gamma_0(x,z)(\gamma_0(x,z)+1), & \gamma_0(x,z) > 0, \\ \lim_{w \to \infty} \frac{\zeta_w^{\dagger}(x,z)}{y^* - \tau(w|x,z)} = -\gamma_0(x,z)(\gamma_0(x,z)+1), & \gamma_0(x,z) < 0, \\ \lim_{w \to \infty} \zeta_w^{\dagger}(x,z) = \sigma^{\dagger}(x,z), & \gamma_0(x,z) = 0. \end{cases}$$
(14)

In this section, γ_0 and ς_w^{\dagger} can be approximated by the additive model. Let

$$(\gamma_0,\varsigma_{w,0}) = \operatorname*{argmin}_{(\gamma,\log\varsigma)\in\mathcal{T}_{\mathcal{A}}\times\mathcal{T}_{\mathcal{A}}} - E[h^{\S}(Y_i|\gamma(X_i,Z_i),\varsigma(X_i,Z_i))],$$

where h^{\S} is the density function corresponding to distribution function $H((\gamma(x, z)+1)y/\varsigma(x, z)|\gamma(x, z))$:

$$h^{\S}(y|\gamma(x,z),\varsigma(x,z)) = \frac{\gamma(x,z)+1}{\varsigma(x,z)} \left(1 + \frac{\gamma(x,z)(\gamma(x,z)+1)y}{\varsigma(x,z)}\right)^{-1/\gamma(x,z)-1}$$

Then, define

$$\varepsilon_w^{\S}(x,z) = rac{\varsigma_{w0}(x,z)}{\varsigma_w^{\dagger}(x,z)} - 1.$$

Again, ε_w^{\S} cannot be improved without some prior information; it must be assumed to be very small to connect between EVT and additive modeling of scale function. From this point forwards, we assume that $\varepsilon_w^{\S}(x,z) = 0$ for all $(x,z) \in \mathcal{X} \times \mathcal{Z}$. Similar to (9), we obtain

$$\begin{cases} \lim_{w \to \infty} \frac{\varsigma_{w0}(x,z)}{\tau(w|x,z)} = O(1), & \gamma_0(x,z) > 0, \\ \lim_{w \to \infty} \frac{\varsigma_{w0}(x,z)}{y^*(x,z) - \tau(w|x,z)} = O(1), & \gamma_0(x,z) < 0, \\ \lim_{w \to \infty} \varsigma_{w0}(x,z) = \sigma_0(x,z), & \gamma_0(x,z) = 0. \end{cases}$$
(15)

By the definition of $(\gamma_0, \varsigma_{w0})$, γ_0 has the form (10) and ς_{w0} is assumed to be expressed as

$$\log \varsigma_{w0}(x,z) = x^T u_{w,0}^{\S} + \sum_{k=1}^d s_{0k}^{\S}(z^{(k)}),$$

where $u_{w,0}^{\S} = (u_{w,01}^{\S}, u_{02}^{\S}, \cdots, u_{0p}^{\S})^T \in \mathbb{R}^p$ and $s_{0k}^{\S} \in \mathcal{A}_k(k = 1, \ldots, d)$. Similar to (12), the intercept parameter $u_{w,01}^{\S}$ only depends on w while other slopes for parametric components and each additive components are independent from w.

We construct the penalized spline estimator of $(\gamma_0, \varsigma_{w,0})$. Let $\log \bar{\varsigma}(x, z) = x^T u^{\S} + B(z)^T c^{\S}$, where $u^{\S} \in \mathbb{R}^p$ and $c^{\S} = ((c_1^{\S})^T, \dots, (c_d^{\S})^T)^T \in \mathbb{R}^{d(K+\xi)}, c_j^{\S} \in \mathbb{R}^{K+\xi}$ are unknown parameter vectors. The *B*-spline model for γ_0 is already defined in section 2. Then, the estimator of $(\beta, b, u^{\S}, c^{\S})$ is defined as

$$(\hat{eta}^{\S}, \hat{b}, \hat{u}^{\S}, \hat{c}^{\S}) = \operatorname*{argmin}_{(eta, b, u^{\S}, c^{\S})} \ell_{pen}^{\S}(eta, b, u^{\S}, c^{\S}),$$

where

$$\ell_{pen}^{\S}(\beta, b, u^{\S}, c^{\S}) = -\sum_{i=1}^{N} \log h^{\S}(Y_i | \bar{\gamma}(X_i, Z_i), \bar{\varsigma}(X_i, Z_i)) I(Y_i > 0) + \lambda \sum_{j=1}^{d} \int_0^1 \{ d^m B_j(z)^T b_j / dz^m \}^2 dz + \nu \sum_{j=1}^{d} \int_0^1 \{ d^m B_j(z)^T c_j^{\S} / dz^m \}^2 dz.$$

For simplicity, we use same λ and ν as those given in section 2. We then construct $\hat{\gamma}(x, z) = x^T \hat{\beta} + B(z)^T \hat{b}$ and $\hat{\varsigma}(x, z) = \exp[x^T \hat{u}^{\S} + B(z)^T \hat{c}^{\S}]$.

We establish some asymptotic properties of $(\hat{\gamma}, \hat{\varsigma})$.

Theorem 5. Suppose that (C1), (C2) with replaced s_{0k} by s_{0k}^{\S} for j = 1, ..., d and (C3)–(C5). In each scenario (S1), (S2) or (S3), as $N \to \infty$,

$$\begin{aligned} ||\hat{\gamma} - \gamma_0||_{L_2} &\leq O\left(\sqrt{\frac{K}{n}}\right) + O(\lambda K^m) + O\left(\left(\frac{N}{n}\right)^{\rho}\right), \\ |\log\hat{\varsigma} - \log\varsigma_{w0}||_{L_2} &\leq O\left(\sqrt{\frac{K}{n}}\right) + O(\nu K^m) + O\left(\left(\frac{N}{n}\right)^{\rho}\right). \end{aligned}$$

Under (S2), we further assume that $\{K \log N\}^{1+\delta/2}/n^{\delta/2} \to 0$ as $N \to \infty$. In each (S1), (S2) or (S3), as $N \to \infty$,

$$\begin{aligned} ||\hat{\gamma} - \gamma_0||_{L_{\infty}} &\leq O\left(\sqrt{\frac{K\log N}{n}}\right) + O(\lambda K^m) + +O\left(\left(\frac{N}{n}\right)^{\rho}\right),\\ |\log\hat{\varsigma} - \log\varsigma_{w0}||_{L_{\infty}} &\leq O\left(\sqrt{\frac{K\log N}{n}}\right) + O(\nu K^m) + O\left(\left(\frac{N}{n}\right)^{\rho}\right). \end{aligned}$$

The optimal rates for L_2 and L_{∞} convergence of $(\hat{\gamma}, \hat{\varsigma})$ are similar to those given in Theorems 1 and 2.

Here, $||\hat{\gamma} - \gamma_0||_{L_2}$ in Theorem 5 does not depend on $O(\nu K^m)$ unlike Theorem 1. Similarly, $||\hat{\varsigma} - \varsigma||_{L_2}$ is not affected by $O(\lambda K^m)$. We also obtained the same conclusion for L_{∞} -convergence. Before describing this reason, we state local asymptotic normality of $(\hat{\gamma}, \hat{\varsigma})$. Let

$$\Sigma^{\S} = \begin{bmatrix} E \left[\frac{1}{(\gamma_0(X,Z)+1)^2} A(X,Z) A(X,Z)^T \right] + \lambda K^{2m} \Omega & O \\ O & E \left[\frac{1}{2\gamma_0(X,Z)+1} A(X,Z) A(X,Z)^T \right] + \nu K^{2m} \Omega \end{bmatrix}$$

,

where O is $(p + K + \xi)$ -square zero matrix, Ω is that defined in the proof of Lemma 6. We then obtain following theorem.

Theorem 6. Suppose that (C1), (C2) with replaced s_{0k} by s_{0k}^{\S} for $j = 1, \ldots, d$ and (C3)–(C5). Furthermore, assume that (K, w_N) satisfies $K = O(n^{1/(2m+1)}/(\log n)^{2m/(2m+1)})$ and $n^{m/(2m+1)}(N/n)^{\rho} \to 0$ as $N \to \infty$. Then, in each scenario (S1), (S2) or (S3), for any $(x, z) \in \mathcal{X} \times \mathcal{Z}$,

$$\left(\frac{n}{\log n}\right)^{m/(2m+1)} \left[\begin{array}{c} \hat{\gamma}(x,z) - \gamma_0(x,z) \\ \frac{\hat{\zeta}(x,z)}{\zeta_{w0}(x,z)} - 1 \end{array}\right] \xrightarrow{D} N_2(0,\lim_{N \to \infty} D(x,z)^T (\Sigma^{\S})^{-1} D(x,z)/K)$$

as $N \to \infty$.

In Theorem 6, since the asymptotic covariance matrix of $(\hat{\gamma}, \hat{\varsigma})$ is block diagonal, $\hat{\gamma}$ and $\hat{\varsigma}$ are asymptotically independent. This orthogonal property of $\hat{\gamma}$ and $\hat{\varsigma}$ is derived from reparametrisation of the scale function, which achieves the computationally efficiency to construct the estimator (see, Chavez-Demoulin and Davison (2005)). We note that when we use the reparameterization of scale function, the estimator of (logarithm of) original scale function $\hat{\sigma} = \hat{\varsigma}/(1+\hat{\gamma})$ cannot retain the additive structure. However, the rate of convergence of $\hat{\sigma} = \hat{\varsigma}/(1+\hat{\gamma})$ is similar to $\hat{\varsigma}$ and hence, it is faster than the fully nonparametric estimator and $\hat{\sigma}$ can avoid the curse of dimensionality. Thus, the reparametrisation is very useful if we do not stick with the additive structure of original scale function in data analysis.

Lastly, we introduce the optimal rate for the parametric part. Define 2p-square matrix

$$\Sigma_{\beta,u}^{\S} = \begin{bmatrix} E \begin{bmatrix} \frac{1}{(\gamma_0(X,Z)+1)^2} X X^T \end{bmatrix} & O \\ O & E \begin{bmatrix} \frac{1}{2\gamma_0(X,Z)+1} X X^T \end{bmatrix} \end{bmatrix},$$

where O is p-square zero matrix.

Theorem 7. Suppose that (C1), (C2) with replaced s_{0k} by s_{0k}^{\S} for j = 1, ..., d and (C3)–(C5). Furthermore, assume that $K \ge O(n^{1/2m} \log n)$ as $N \to \infty$. Then, in each scenario (S1), (S2) or (S3), if $n^{1/2}(N/n)^{\rho} = O(1)$, as $N \to \infty$,

$$\begin{aligned} ||\hat{\beta} - \beta_0|| &\leq O(n^{-1/2}) + O((N/n)^{\rho}) = O(N^{\rho/(1-2\rho)}), \\ ||\hat{u}^{\S} - u_{w,0}^{\S}|| &\leq O(n^{-1/2}) + O((N/n)^{\rho}) = O(N^{\rho/(1-2\rho)}). \end{aligned}$$

Furthermore, under $n^{1/2}(N/n)^{\rho} \to 0$,

$$\sqrt{n} \left[\begin{array}{c} \hat{\beta} - \beta_0 \\ \hat{u}^{\S} - u^{\S}_{w,0} \end{array} \right] \xrightarrow{D} N_{2p}(0, (\Sigma^{\S}_{\beta,u})^{-1}).$$

The proof of Theorem 7 is omitted since it is quite similar to Theorem 4. In Theorem 7, we also see that $\hat{\beta}$ and \hat{u}^{\S} are asymptotically independent from the structure of $\Sigma_{\beta,u}^{\S}$.

6 Conclusion

We developed the asymptotic theory for the regression with GPD. The shape and scale parameters (functions of covariates) included in GPD are modeled by the GAM. The estimator of each component is constructed by applying the normalized *B*-spline method to the extreme value data, which are picked using the peak over threshold method. By using the property of *B*-spline basis, some concentration inequality, and asymptotic theory, we demonstrate the L_2

and L_{∞} rate of convergences of the estimators of shape and scale functions. We also present the asymptotic normality of the estimator as the local asymptotics of the additive estimator.

The motivation of this study came from the previous study that the GPD regression with GAM is available in the R-package evgam provided by Youngman (2022). In this package, shape and (logarithm of) scale function are modeled using GAM. Indeed, such modeling is reasonable in numerical study and data analysis. However, from the theoretical perspective, the additive modeling of scale function needed somewhat strong conditions. First, since the scale function depends on shape and threshold function in EVT, we cannot know whether the scale function can be written as the form of GAM. In particular, the threshold function is predetermined before considering GPD. We assumed that the true scale function and its best approximation of GAM are nearly equal. However, this bias cannot generally be improved. Second, in the scale function with GAM, we assumed that only the intercept parameter depends on the threshold level unlike other components, which is detailed in Remark 2. Lastly, we assumed that the signature of the shape function is fixed for all data points of covariates. If this condition fails, scale function is discontinuous, and the GAM would not be performed well. We believe that relaxing these conditions is quite challenging. However, further discussion about three above conditions is important to elucidate the mathematical property of GAM in GPD or evgam.

In R-package evgam, we can also use GAM for the generalized extreme value (GEV) distribution instead of GPD. The developments in theoretical results for GAM of GEV regression would be an important topic for further research. Such topic can be regarded as the extension of Bücher and Segers (2017), which developed the asymptotic theory for maximum-likelihood estimator of GEV with a non-regression case.

Appendix A: Outline of proof of theorems

Our purpose is to evaluate the asymptotic behavior of the estimator $\hat{\gamma}(x, z)$ and $\hat{\sigma}(x, z)$ of the true additive model $\gamma_0(x, z)$ and $\sigma_{w0}(x, z)$. Let

$$L(\theta) = -E[\log h(Y|\bar{\gamma}(X,Z),\bar{\sigma}(X,Z))]$$

and let $\theta_0 = (\beta_0^T, b_0^T, u_{w,0}^T, c_0^T)^T$ be the best spline approximation parameters that

$$\theta_0 = \operatorname{argmin}_{(\beta, b, u, c)} L(\theta).$$

We write $b_0 = (b_{01}^T, \dots, b_{0d}^T)^T$ and $b_{0j} = (b_{0,j,1}, \dots, b_{0,j,K+\xi})^T \in \mathbb{R}^{K+\xi}$. Similarly, $c_0 = (c_{01}^T, \dots, c_{0d}^T)^T$ and $c_{0j} = (c_{0,j,1}, \dots, b_{c,j,K+\xi})^T \in \mathbb{R}^{K+\xi}$. Then, for $(x, z) \in \mathcal{X} \times \mathcal{Z}$ with $z = (z^{(1)}, \dots, z^{(d)})^T$, let $\bar{g}_{0j}(z^{(j)}) = B_j(z^{(j)})^T b_{0j}$ and $\bar{s}_0(z^{(j)}) = B_j(z^{(j)})^T c_{0j}$ for $j = 1, \dots, d$. Therefore, the best spline approximation of $(\gamma_0, \log \sigma_{w0})$ can be written as

$$\bar{\gamma}_0(x,z) = x^T \beta_0 + \sum_{j=1}^d \bar{g}_{0j}(z^{(j)}),$$

and

$$\log \bar{\sigma}_{w0}(x,z) = x^T u_{w,0} + \sum_{j=1}^d \bar{s}_0(z^{(j)}),$$

respectively. From Lemma 3 in Appendix B, we see that $\bar{g}_{0j}(z^{(j)}) - g_{0j}(z^{(j)}) = O(K^{-\zeta})$ and $\bar{s}_{0j}(z^{(j)}) - s_{0j}(z^{(j)}) = O(K^{-\zeta})$. Thus, we obtain

$$\hat{g}_j(z^{(j)}) - g_{0j}(z^{(j)}) = \hat{g}_j(z^{(j)}) - \bar{g}_{0j}(z^{(j)}) + \bar{g}_{0j}(z^{(j)}) - g_{0j}(z^{(j)}) = B_j(z^{(j)})^T(\hat{b} - b_0) + O(K^{-\zeta})$$

and

$$\hat{s}_j(z^{(j)}) - s_{0j}(z^{(j)}) = \hat{s}_j(z^{(j)}) - \bar{s}_{0j}(z^{(j)}) + \bar{s}_{0j}(z^{(j)}) - s_{0j}(z^{(j)}) = B_j(z^{(j)})^T (\hat{c} - c_0) + O(K^{-\zeta}).$$

Accordingly, we have

$$\hat{\gamma}(x,z) - \gamma_0(x,z) = \hat{\gamma}(x,z) - \bar{\gamma}_0(x,z) + \bar{\gamma}_0(x,z) - \gamma_0(x,z) = x^T (\hat{\beta} - \beta_0) + B(z)^T (\hat{b} - b_0) + O(K^{-\zeta})$$

and

$$\log \hat{\sigma}(x, z) - \log \sigma_{w0}(x, z) = \log \hat{\sigma}(x, z) - \log \bar{\sigma}_{w0}(x, z) + \log \bar{\sigma}_{w0}(x, z) - \log \sigma_{w0}(x, z) = x^T (\hat{u} - u_{w0}) + B(z)^T (\hat{c} - c_0) + O(K^{-\zeta}).$$

Thus, our purpose is to analyze the asymptotic behavior of $\hat{\theta} - \theta_0$. We write $\ell_{pen}(\theta) = \ell_{pen}(\beta, b, u, c)$. The first key result is the consistency of $||\hat{\theta} - \theta_0||$, which is described in Lemma 8. The consistency of $||\hat{\theta} - \theta_0||$ guarantees to use the Taylor expansion to $\ell_{pen}(\theta)$ as

$$\left(\frac{\partial^{2}\ell_{pen}(\theta_{0})}{\partial\theta\partial\theta^{T}}\right)(\hat{\theta}-\theta_{0}) = \left\{\frac{\partial\ell_{pen}(\theta_{0})}{\partial\theta} - E\left[\frac{\partial\ell_{pen}(\theta_{0})}{\partial\theta}\right]\right\}(1+o_{P}(1)) + E\left[\frac{\partial\ell_{pen}(\theta_{0})}{\partial\theta}\right](1+o(1)).$$
(16)

Lemmas 9 and 10 indicate the rate of convergence of $||\hat{\theta} - \theta_0||$ and asymptotic normality of $\hat{\theta} - \theta_0$ using (16). Lemmas 1–7 are used to demonstrate Lemmas 8–10. The preliminary results are Lemmas 1–4. As our result, the asymptotic order of first and second term of the right-hand side of (16) is presented in Lemmas 5 and 6 whereas Lemma 7 proves the boundness of $(\partial^2 \ell_{pen}(\theta_0)/\partial\theta \partial\theta^T)$. From these lemmas, theorems in section 3 can be shown in Appendix C. Proofs of theorems in section 4 are mentioned in Appendix D.

We note about how to prove theorems in this paper below. Let $\Lambda : \mathbb{R} \times \mathcal{X} \times \mathcal{Z}$ be any function. In te following, Λ is assumed to be penalized log-likelihood function, its derivative or other functions. Then, for example, since $P(Y_i > 0) \approx n/N$, we have

$$E\left[\sum_{i=1}^{N} \Lambda(Y_i, X_i, Z_i) I(Y_i > 0)\right] = NE[\Lambda(Y_i, X_i, Z_i) I(Y_i > 0)]$$

$$= NP(Y_i > 0)E[\Lambda(Y_i, X_i, Z_i)|Y_i > 0]$$

$$\approx nE[\Lambda(Y_i, X_i, Z_i)|Y_i > 0],$$

which is asymptotically equivalent to

$$E\left[\sum_{i=1}^{n} \Lambda(Y_i, X_i, Z_i)\right]$$

conditioning on $Y_1, \ldots, Y_n > 0$. The latter case corresponds to the situation that the samples (Y_i^*, X_i, Z_i) are retained for *i* such that $Y_i^* \ge \tau(w|X_i, Z_i)$ and removing the sample (Y_i^*, X_i, Z_i) with $Y_i^* < \tau(w|X_i, Z_i)$ in advance. In the following, for simplicity, we assume such a case, that is, dataset is remarked as $\{(Y_i, X_i, Z_i) : i = 1, \ldots, n\}$, where $Y_i > 0$. Then, the log-likelihood is redefined by

$$\ell(\beta, u, b, c) = -\sum_{i=1}^{n} \log h(Y_i | \bar{\gamma}(X_i, Z_i), \bar{\sigma}(X_i, Z_i))$$

conditioning on $Y_1, \ldots, Y_n > 0$.

Appendix B: Lemmas

Lemma 1 (Bernstein's inequality). Let W_1, \ldots, W_n be univariate i.i.d. random variable having $E[W_i] = 0$, $V[W_i] = \sigma^2 \in (0, \infty)$ and $|W_i| < M$ almost surely for some constant M > 0. Then, for any $\varepsilon > 0$,

$$P\left(\sum_{i=1}^{n} W_i > \varepsilon\right) \le \exp\left[\frac{-2^{-1}\varepsilon^2}{\sum_{i=1}^{n} \sigma_i^2 + 3^{-1}\varepsilon M}\right].$$

Lemma 1 is the famous result in probability theory. Its proof is clarified in van der Geer (2000).

Lemma 2. The matrix $E[B(Z)B(Z)^T]$ is positive definite, that is, for any nonzero-vector $v \in \mathbb{R}^{d(K+\xi)}$, there exist constants $M_{min}, M_{max} > 0$ such that

$$M_{min} \le \frac{v^T E[B(Z)B(Z)^T]v}{||v||} \le M_{max}.$$

The proof of Lemma 2 is shown in Lemma A.2 of Liu et al. (2011).

Lemma 3. Suppose that (C2). Then, as $K \to \infty$,

$$\sup_{z^{(j)} \in [0,1]} \left| \sum_{k=1}^{K+\xi} B_k^{[\xi]}(z^{(j)}) b_{0,j,k} - g_{0j}(z^{(j)}) \right| = O(K^{-\zeta}), \quad j = 1, \dots, d,$$

and

$$\sup_{z^{(j)} \in [0,1]} \left| \sum_{k=1}^{K+\xi} B_k^{[\xi]}(z^{(j)}) c_{0,j,k} - s_{0j}(z^{(j)}) \right| = O(K^{-\zeta}), \quad j = 1, \dots, d.$$

Consequently,

$$\sup_{(x,z)\in\mathcal{X}\times\mathcal{Z}}|\bar{\gamma}(x,z)-\gamma_0(x,z)|=O(K^{-\zeta})$$

and

$$\sup_{(x,z)\in\mathcal{X}\times\mathcal{Z}} |\log \bar{\sigma}(x,z) - \log \sigma_{w0}(x,z)| = O(K^{-\zeta})$$

Proof of Lemma 3. From de Boor (2001), for $g_{0j} \in \mathcal{C}^{(\ell)}[0,1]$, there exists $b_j^* = (b_{j,0}^*, \dots, b_{j,K+\xi}^*)^T \in \mathbb{R}^{K+\xi+1}$ such that

$$\sup_{z \in [0,1]} \left| \sum_{k=0}^{K+\xi} \psi_k^{[\xi]}(z) b_{j,k}^* - g_{0j}(z) \right| = O(K^{-\zeta})$$

When putting $b_{0,j,1} = -(\phi_{j,1}/\phi_{j,0})^{-1} ||\bar{\psi}_{j,0}^{[\xi]}||b_{j,0}^*$ and

$$b_{0,j,k+1} = ||\bar{\psi}_{j,k+1}||\frac{\phi_{j,k}}{\phi_{j,k+1}} \left\{\frac{b_{0,j,k}}{||\bar{\psi}_{j,k}||} - b_{j,k}^*\right\}, \quad k = 1, \dots, K + \xi - 1,$$

we have that for $z \in [0, 1]$,

$$\sum_{k=1}^{K+\xi} B_k^{[\xi]}(z) b_{0,j,k} = \sum_{k=0}^{K+\xi} \psi_k^{[\xi]}(z) b_{j,k}^*,$$

which yields that

$$\sup_{z \in [0,1]} \left| \sum_{k=1}^{K+\xi} B_k^{[\xi]}(z) b_{0,j,k} - g_{0j}(z) \right| = O(K^{-\zeta}).$$

Other assertions of this lemma can be shown as the same mannar. This completes the proof. \Box

We next show the quadratic form and asymptotic order of penalty term. Let $\Psi_j = (\Psi_{j,i,k})_{ik}$ be $(K + \xi + 1) \times (K + \xi)$ matrix with $\Psi_{j,i,i} = 1/||\bar{\psi}_{j,i}^{[\xi]}||_2$, $\Psi_{j,i+1,i} = -(\phi_{j,i}/\phi_{j,i-1})/||\bar{\psi}_{j,i}^{[\xi]}||_2$ for $i = 1, \ldots, K + \xi$ and $\Psi_{j,i,k} = 0$ for $|i - k| \ge 2$. Then, we can write $\bar{g}_j(z_j) = B(z_j)^T b_j = \psi^{[\xi]}(z_j)^T \Psi_j b_j$, where $\psi^{[\xi]}(z_j) = (\psi_0^{[\xi]}(z_j), \ldots, \psi_{K+\xi}^{[\xi]}(z_j))^T$ is the vector of original *B*-spline bases. Let $D_1 = (D_{1,i,j})_{ij}$ be $(K + \xi) \times (K + \xi + 1)$ matrix with $D_{1,i,i} = 1$ and $D_{1,i,i+1} = -1$ for $i = 1, \ldots, K + \xi$, and $D_{1,i,j} = 0$ for |i - j| > 2. Then, D_1 is the so called first order difference matrix (see, Xiao 2019). For $q \ge 2$, let D_q be $(K + \xi + 1 - q) \times (K + \xi + 1)$ matrix satisfying $D_q = D_{q-1} \times D_1$ recursively. Then, D_q is denoted by qth order difference matrix. Lastly, we define $K + \xi + 1 - m$ matrix $R_m = (R_{i,k})_{ik}$ with $R_{i,k} = \int_0^1 \psi_{i-1}^{[\xi-m]}(z)\psi_{k-1}^{[\xi-m]}(z)dz$ for $i, k = 1, \ldots, K + \xi + 1 - m$. By using the derivative of $\psi_k^{[\xi]}(z_j)$, we can evaluate the quadratic form of the penalty term.

Lemma 4. For any $v \in \mathbb{R}^{K+\xi}$,

$$\frac{\partial^2}{\partial v \partial v^T} \int_0^1 \left\{ \frac{d^m B_j(z)^T v}{dz^m} \right\}^2 dz = K^{2m} (m!)^2 \Psi_j^T D_m^T R_m D_m \Psi_j = O(K^{2m}), j = 1, \dots, d.$$

Proof of Lemma 4. By the definition of normalized B-spline, we obtain $B_j(z)^T v = \psi^{[\xi]}(z)^T \Psi_j v$ for $z \in [0, 1]$. Furthermore, from the property of mth derivative of B-spline function (see, de Boor 2001, Xiao 2019), we obtain that for $z \in [0, 1]$,

$$\frac{d^m}{dz^m}\psi^{[\xi]}(z)^T\Psi_j v = K^m m! \psi^{[\xi-m]}(z)^T D_m \Psi_j v,$$

where $\psi^{[\xi-m]}(z) = (\psi_0^{[\xi-m]}(z_j), \dots, \psi_{K+\xi-m}^{[\xi-m]}(z_j))^T$. Accordingly,

$$\int_{0}^{1} \left\{ \frac{d^{m} B_{j}(z)^{T} v}{dz^{m}} \right\}^{2} dz = K^{2m} (m!)^{2} v^{T} \Psi_{j}^{T} D_{m}^{T} R_{m} D_{m} \Psi_{j} v$$

Here, for $R_m = (R_{i,k})_{ik}$, we have $R_{i,k} = O(K^{-1})$ for $|i - k| \leq \xi - m$ and $R_{i,k} = 0$ otherwise, that is, R_m is the band matrix. Furthermore, for $\Psi_j = (\Psi_{j,i,k})_{ik}$, $\Psi_{j,i,k} = O(K^{1/2})$ for k = i + 1 and $\Psi_{j,i,k} = 0$ otherwise. Thus, Ψ_j is also band matrix. This implies that $K^{2m}(m!)^2 \Psi_j^T D_m^T R_m D_m \Psi_j = O(K^{2m})$.

Define $\theta = (\beta^T, b^T, u^T, c^T)^T$ and we write $\ell_{pen}(\theta) = \ell_{pen}(\beta, b, u, c)$. Furthermore, let $\hat{\theta} = (\hat{\beta}, \hat{b}, \hat{u}, \hat{c})$ be the estimator, which is obtained by solving

$$\frac{\partial}{\partial \theta} \ell_{pen}(\theta) = \underline{0},$$

where $\underline{0}$ is the zero-vector with dimesion $2(p+K+\xi)$. Define $\theta_{\gamma} = (\beta^T, b^T)^T$ and $\theta_{\sigma} = (u^T, c^T)^T$. From these symbols, we have $\bar{\gamma}(X, Z) = A(X, Z)^T \theta_{\gamma}$ and $\bar{\sigma}(X, Z) = \exp[A(X, Z)^T \theta_{\sigma}]$. We further define some symbols. Let

$$\begin{aligned} \ell_{\gamma}(y|\gamma(x,z),\sigma(x,z)) &= \left. \frac{\partial}{\partial a} \left\{ -\log h(y|a,b) \right\} \right|_{a=\gamma(x,z),b=\sigma(x,z)} \\ &= \left. (\gamma(x,z)^{-1}+1) \frac{y/\sigma(x,z)}{1+y\gamma(x,z)/\sigma(x,z)} - \gamma(x,z)^{-2} \log \left(1 + \frac{y\gamma(x,z)}{\sigma(x,z)} \right) \right. \end{aligned}$$

and

$$\ell_{\sigma}(y|\gamma(x,z),\sigma(x,z)) = \frac{\partial}{\partial \log b} \left\{ -\log h(y|a,b) \right\} \Big|_{a=\gamma(x,z),b=\sigma(x,z)}$$
$$= 1 - (\gamma(x,z)^{-1} + 1) \frac{y\gamma(x,z)/\sigma(x,z)}{1 + y\gamma(x,z)/\sigma(x,z)}.$$

If $\gamma(x, z) = 0$,

$$\ell_{\gamma}(y|0,\sigma(x,z)) = \frac{y}{\sigma(x,z)} - \frac{1}{2}\frac{y^2}{\sigma(x,z)}$$

and $\ell_{\sigma}(y|0,\sigma(x,z)) = 1 - y/\sigma(x,z)$. We can show that

$$\int \ell_{\gamma}(y|\gamma_0(x,z),\sigma_{w0}(x,z))dH(y|\gamma_0(x,z),\sigma_{w0}(x,z))$$
$$= \int \ell_{\sigma}(y|\gamma_0(x,z),\sigma_{w0}(x,z))dH(y|\gamma_0(x,z),\sigma_{w0}(x,z))$$
$$= 0$$

from the property of gradient of log-likelihood function of GPD. Furthermore, the hessian matrix of log-likelihood of GPD involves

$$\int \ell_{\gamma}(y|\gamma_{0}(x,z),\sigma_{w0}(x,z))^{2} dH(y|\gamma_{0}(x,z),\sigma_{w0}(x,z)) = \frac{2}{(2\gamma_{0}(x,z)+1)(\gamma_{0}(x,z)+1)},$$

$$\int \ell_{\sigma}(y|\gamma_{0}(x,z),\sigma_{w0}(x,z))^{2} dH(y|\gamma_{0}(x,z),\sigma_{w0}(x,z)) = \frac{1}{2\gamma_{0}(x,z)+1}$$

and

$$\int \ell_{\gamma}(y|\gamma_{0}(x,z),\sigma_{w0}(x,z))\ell_{\sigma}(y|\gamma_{0}(x,z),\sigma_{w0}(x,z))dH(y|\gamma_{0}(x,z),\sigma_{w0}(x,z))$$

= $\frac{1}{(2\gamma_{0}(x,z)+1)(\gamma_{0}(x,z)+1)}.$

Lemma 5. Suppose that (C1)–(C5). In each scenario (S1), (S2) or (S3), as $N \to \infty$,

$$E\left[\left|\left|\frac{\partial}{\partial\theta}\ell_{pen}(\theta_0) - E\left[\frac{\partial}{\partial\theta}\ell_{pen}(\theta_0)\right]\right|\right|^2\right] = O(K/n)$$

Proof of Lemma 5. Since the penalty term of ℓ_{pen} does not contain the stochastic part, we have

$$\left\| \frac{\partial}{\partial \theta} \ell_{pen}(\theta_0) - E\left[\frac{\partial}{\partial \theta} \ell_{pen}(\theta_0) \right] \right\|^2 = \left\| \frac{\partial}{\partial \theta} \ell(\theta_0) - E\left[\frac{\partial}{\partial \theta} \ell(\theta_0) \right] \right\|^2.$$

We let $\bar{\ell}_A = \ell_A(Y_i|\bar{\gamma}_0(X_i, Z_i), \bar{\sigma}_{w0}(X_i, Z_i))$ for $A = \{\gamma, \sigma\}$ and let I be $(p + K + \xi)$ -identity

matrix. Then, from Lemma 3, for *i.i.d.* sample (Y_i, X_i, Z_i) ,

$$\begin{split} & E\left[\left|\left|\frac{\partial}{\partial\theta}\ell(\theta_{0})-E\left[\frac{\partial}{\partial\theta}\ell(\theta_{0})\right]\right|\right|^{2}\right] \\ &\leq \frac{1}{n}E\left[\left(X^{T},B(Z)^{T},X^{T},B(Z)^{T}\right)\left(\begin{array}{cc} \overline{\ell}_{\gamma}^{2}I & \overline{\ell}_{\gamma}\overline{\ell}_{\sigma}I\\ \overline{\ell}_{\sigma}\overline{\ell}_{\gamma}I & \overline{\ell}_{\sigma}^{2}I\end{array}\right)\left(\begin{array}{c} X\\ B(Z)\\ X\\ B(Z)\end{array}\right)\right] \\ &= \frac{1}{n}E\left[\left(X^{T},B(Z)^{T},X^{T},B(Z)^{T}\right)\left(\begin{array}{c} \frac{2}{(2\gamma(X,Z)+1)(\gamma(X,Z)+1)}I & \frac{1}{(2\gamma(X,Z)+1)(\gamma(X,Z)+1)}I\\ (2\gamma(X,Z)+1)(\gamma(X,Z)+1)I\end{array}\right)\left(\begin{array}{c} X\\ B(Z)\\ X\\ B(Z)\end{array}\right)\right] \\ &= \frac{1}{n}E\left[\frac{\gamma(X,Z)+5}{(2\gamma(X,Z)+1)(\gamma(X,Z)+1)}(||X||^{2}+||B(Z)||^{2})\right] \\ &\leq C\frac{d(K+\xi)}{n} \end{split}$$

for some constant C > 0. The last inequality can be shown by the property of normalized *B*-spline: $E[||B(Z)||^2] = d(K + \xi)$.

By Lemma 4, the penalty term of ℓ_{pen} can be written as quadratic form. Thus, there exists a $2(p + K + \xi)$ -square matrix $\Omega_{\gamma,\sigma}$ such that

$$\sum_{j=1}^{d} \left\{ \lambda \int_{0}^{1} \{ \bar{g}_{j}^{(m)}(z) \}^{2} dz + \nu \int_{0}^{1} \{ \bar{s}_{j}^{(m)}(z) \}^{2} dz \right\} = \theta_{0}^{T} \Omega_{\gamma,\sigma} \theta_{0}.$$

Using above, we obtain next Lemma.

Lemma 6. Suppose that (C1)–(C5). Then, in each scenario (S1), (S2) or (S3), as $N \to \infty$,

$$\left| \left| E\left[\frac{\partial}{\partial \theta} \ell_{pen}(\theta_0)\right] \right| \right| \le O\left(\left(\frac{N}{n}\right)^{\rho}\right) + O(\lambda K^m) + O(\nu K^m).$$

Proof of Lemma 6. By the Cauchy–Schwarz inequality, we have

$$\left\| E\left[\frac{\partial}{\partial\theta}\ell_{pen}(\theta_0)\right] \right\| \leq \left\| E\left[\frac{\partial}{\partial\theta}\ell(\theta_0)\right] \right\| + \left\| \frac{\partial}{\partial\theta}\theta_0^T\Omega_{\gamma,\sigma}\theta_0 \right\|.$$

We first consider the part of log-likelihood. Again, we use the symbola $\bar{\ell}_A, A \in \{\gamma, \sigma\}$ defined in the proof of Lemma 5. Then, we have

$$E\left[\frac{\partial}{\partial\theta}\ell(\theta_{0})\right] = E\left[\begin{array}{c} \bar{\ell}_{\gamma}\frac{\partial\bar{\gamma}}{\partial\theta_{\gamma}}\\ \bar{\ell}_{\sigma}\frac{\partial\log\bar{\sigma}}{\partial\theta_{\sigma}}\end{array}\right] \\ = E\left[\begin{array}{c} \bar{\ell}_{\gamma}X\\ \bar{\ell}_{\gamma}B(Z)\\ \bar{\ell}_{\sigma}X\\ \bar{\ell}_{\sigma}B(Z)\end{array}\right].$$

Here, for simplicity, we write $\alpha(w|x,z)$ as $\alpha(\tau(w|x,z)|x,z)$. For any integrable function q(Y), we have that for any $(x,z) \in \mathcal{X} \times \mathcal{Z}$,

$$E_{Y|x,z}[q(Y)] = \int q(y)dF(y|x,z) = \int q(y)dH(y|x,z) + \alpha(w|x,z) \int q(y)Q'(y/\sigma_{w0}(x,z)|x,z)dy(1+o(1)),$$

where, $E_{Y|x,z}$ is the expectation by the conditional distribution of Y_i given $(X_i, Z_i) = (x, z)$ and $H(y|x, z) = H(y|\gamma_0(x, z), \sigma_{w0}(x, z))$. If $q(y) = \bar{\ell}_A, A \in \{\gamma, \sigma\}$, we have $\int q(y)dH(y|x, z) = 0$. Since $\bar{\gamma}_0(x, z) - \gamma_0(x, z) = O(K^{-\zeta})$ and $\bar{\sigma}_{w0}(x, z) - \sigma_{w0}(x, z) = O(K^{-\zeta})$ from Lemma 3, we obtain

$$\begin{split} &E_{Y|x,z}\left[\bar{\ell}_{\gamma}\right] \\ &= E_{Y|x,z}\left[\ell_{\gamma}(y|\bar{\gamma}_{0}(x,z),\bar{\sigma}_{w0}(x,z))\right] \\ &= \alpha(\tau(w|x,z)|x,z)\int \ell_{\gamma}(y|\gamma_{0}(x,z),\sigma_{w0}(x,z))Q'(y/\sigma_{w0}(x,z)|x,z)dy(1+o(1)) + O(K^{-\zeta}) \\ &\equiv \alpha(\tau(w|x,z)|x,z)q_{\gamma}(x,z) + O(K^{-\zeta}). \end{split}$$

By the definition of Q, we can find that $|q_{\gamma}(x,z)| < \infty$ for all $(x,z) \in \mathcal{X} \times \mathcal{Z}$ under each case (S1)–(S3). Similarly, we have

$$E_{Y|x,z}\left[\bar{\ell}_{\sigma}\right] = \alpha(\tau(w|x,z)|x,z)q_{\sigma}(x,z)(1+o(1)) + O(K^{-\zeta}),$$

where

$$q_{\sigma}(x,z) = \int \ell_{\sigma}(y|\gamma_0(x,z),\sigma_{w0}(x,z))Q'(y/\sigma_{w0}(x,z)|x,z)dy$$

and $|q_{\sigma}(x,z)| < \infty$ for all $(x,z) \in \mathcal{X} \times \mathcal{Z}$ from the form of Q. Thus,

$$E\left[\frac{\partial}{\partial\theta}\ell(\theta_0)\right] = E_{X,Z} \begin{bmatrix} \alpha(\tau(w|x,z)|X,Z)q_{\gamma}(X,Z)\{1+O(K^{-\zeta})\}X\\ \alpha(\tau(w|x,z)|X,Z)q_{\gamma}(X,Z)\{1+O(K^{-\zeta})\}B(Z)\\ \alpha(\tau(w|x,z)|X,Z)q_{\sigma}(X,Z)\{1+O(K^{-\zeta})\}X\\ \alpha(\tau(w|x,z)|X,Z)q_{\sigma}(X,Z)\{1+O(K^{-\zeta})\}B(Z) \end{bmatrix}.$$

From the property of normalized *B*-spline, for some bounded function $q : \mathbb{R} \to \mathbb{R}$, each element of E[q(Z)B(Z)] has an order $O(1/\sqrt{K})$. Since B(Z) is $d(K+\xi)$ -matrix, we have ||E[q(Z)B(Z)]|| = O(1). Thus, from conditions (C1), (C3), (C4) and the property of *B*-spline, we can evaluate

$$\left\| E\left[\frac{\partial}{\partial\theta}\ell(\theta_0)\right] \right\| \le O\left(\left(\frac{N}{n}\right)^{\rho}\right).$$

We next derive the asymptotic order of the penalty term:

$$\frac{\partial}{\partial\theta_0} \sum_{j=1}^d \left\{ \lambda \int_0^1 \{ \bar{g}_j^{(m)}(z) \}^2 dz + \nu \int_0^1 \{ \bar{s}_j^{(m)}(z) \}^2 dz \right\}.$$

From Lemma 4, we have

$$\sum_{j=1}^{d} \lambda \int_{0}^{1} \{\bar{g}_{j}^{(m)}(z)\}^{2} dz = \frac{\lambda K^{2m}}{2} b_{0}^{T} \Omega_{\gamma} b_{0},$$

where Ω is the $d(K + \xi)$ square matrix with

$$\Omega = (m!)^2 \begin{bmatrix} \Psi_1^T D_m^T R_m D_m \Psi_1 & & \\ & \ddots & \\ & & \Psi_d^T D_m^T R_m D_m \Psi_d \end{bmatrix}$$

Here, all elements in the off diagonal block (the part of blank) of Ω are zero. Similarly, we can write

$$\sum_{j=1}^{d} \nu \int_{0}^{1} \{\bar{s}_{j}^{(m)}(z)\}^{2} dz = \frac{\nu K^{2m}}{2} c_{0}^{T} \Omega c_{0}.$$

Therefore, using

$$\Omega_{\gamma,\sigma} = \begin{bmatrix} O & & & \\ & \lambda K^{2m}\Omega & & \\ & & O & \\ & & & \nu K^{2m}\Omega \end{bmatrix},$$

where the all elements of O and the off-diagonal block are zero, we obtain

$$\frac{\partial}{\partial\theta_0} \sum_{j=1}^d \left\{ \lambda \int_0^1 \{\bar{g}_j^{(m)}(z)\}^2 dz + \nu \int_0^1 \{\bar{s}_j^{(m)}(z)\}^2 dz \right\} = \Omega_{\gamma,\sigma} \boldsymbol{\theta}_0$$

Here, from the definition of R_m , D_m , Ψ_j , b_{0j} and Lemma 4, we obtain

$$\begin{split} \lambda K^{2m}(m!)^2 \Psi_j^T D_m^T R_m D_m \Psi_j b_{0j} \\ &= \lambda K^m m! \Psi_j^T D_m^T \begin{bmatrix} \int_0^1 \psi_1^{[\xi-m]}(z) g_{0j}^{(m)}(z) dz \\ \vdots \\ \int_0^1 \psi_{K+\xi-m}^{[\xi-m]}(z) g_{0j}^{(m)}(z) dz \end{bmatrix} (1+o(1)) \\ &= O(\lambda K^{m-1/2}) \end{split}$$

and

$$\nu K^{2m}(m!)^2 \Psi_j^T D_m^T R_m D_m \Psi_j c_{0j} = O(\nu K^{m-1/2}).$$

Therefore, we obtain $\lambda K^{2m}\Omega b_0 = O(\lambda K^{m-1/2})$ and $\nu K^{2m}\Omega c_0 = O(\nu K^{m-1/2})$. Since b_0 and c_0 are $(K + \xi)$ -matrix, we obtain $||\lambda K^{2m}\Omega b_0|| = O(\lambda K^m)$ and $||\nu K^{2m}\Omega c_0|| = O(\nu K^m)$.

Lemma 7. Suppose that (C1)–(C5). In each scenario (S1), (S2) or (S3), for any vector $v \in \mathbb{R}^{2(p+K+\xi)} - \{0\}$ with $||v|| \leq 1$, there exist positive constants C_m and C_M such that

$$C_m < v^T E\left[\frac{\partial^2}{\partial \theta \partial \theta^T} \ell_{pen}(\theta_0)\right] v < C_M.$$

Proof of Lemma 7. We first obtain

$$\frac{\partial^2}{\partial\theta\partial\theta^T}\ell_{pen}(\theta) = \frac{\partial^2}{\partial\theta\partial\theta^T}\ell(\theta) + \Omega_{\gamma,\sigma},$$

where $\Omega_{\gamma,\sigma}$ is that given in the proof of Lemma 6. By the straightforward calculation of hessian matrix of log-likelihood of GPD, we have

$$E_{Y|x,z} \left[\frac{\partial^2}{\partial \theta \partial \theta^T} \ell(\theta) \right] = \frac{1}{2\gamma_0(x,z)+1} \left[\begin{array}{cc} \frac{2}{\gamma_0(x,z)+1} A(x,z) A(x,z)^T & \frac{1}{\gamma_0(x,z)+1} A(x,z) A(x,z)^T \\ \frac{1}{\gamma_0(x,z)+1} A(x,z) A(x,z)^T & A(x,z) A(x,z)^T \end{array} \right] (1+o_P(1)) + O_P(1) \left[\begin{array}{c} \frac{2}{\gamma_0(x,z)+1} A(x,z) A(x,z)^T & \frac{1}{\gamma_0(x,z)+1} A(x,z) A(x,z)^T \\ \frac{1}{\gamma_0(x,z)+1} A(x,z) A(x,z)^T & A(x,z) A(x,z)^T \end{array} \right] (1+o_P(1)) + O_P(1) \left[\begin{array}{c} \frac{2}{\gamma_0(x,z)+1} A(x,z) A(x,z)^T & \frac{1}{\gamma_0(x,z)+1} A(x,z) A(x,z)^T \\ \frac{1}{\gamma_0(x,z)+1} A(x,z) A(x,z)^T & A(x,z) A(x,z)^T \end{array} \right] (1+o_P(1)) + O_P(1) \left[\begin{array}{c} \frac{2}{\gamma_0(x,z)+1} A(x,z) A(x,z)^T & \frac{1}{\gamma_0(x,z)+1} A(x,z) A(x,z)^T \\ \frac{1}{\gamma_0(x,z)+1} A(x,z) A(x,z)^T & A(x,z) A(x,z)^T \end{array} \right] (1+o_P(1)) + O_P(1) \left[\begin{array}{c} \frac{2}{\gamma_0(x,z)+1} A(x,z) A(x,z)^T & \frac{1}{\gamma_0(x,z)+1} A(x,z) A(x,z)^T \\ \frac{1}{\gamma_0(x,z)+1} A(x,z) A(x,z)^T & A(x,z) A(x,z)^T \end{array} \right] (1+o_P(1)) + O_P(1) \left[\begin{array}{c} \frac{2}{\gamma_0(x,z)+1} A(x,z) A(x,z)^T & \frac{1}{\gamma_0(x,z)+1} A(x,z) A(x,z)^T \\ \frac{1}{\gamma_0(x,z)+1} A(x,z) A(x,z)^T & A(x,z) A(x,z)^T \end{array} \right] (1+o_P(1)) + O_P(1) \left[\begin{array}{c} \frac{2}{\gamma_0(x,z)+1} A(x,z) A(x,z) A(x,z)^T \\ \frac{1}{\gamma_0(x,z)+1} A(x,z) A(x,z) A(x,z)^T \\ \frac{1}{\gamma_0(x,z)+1} A(x,z) A(x,z) A(x,z)^T \end{array} \right] (1+o_P(1)) + O_P(1) \left[\begin{array}{c} \frac{2}{\gamma_0(x,z)+1} A(x,z) A(x$$

From (C5) and Lemma 4, $\Omega_{\gamma,\sigma}$ is non-negative definite matrix. Therefore, the proof is completed if we can show that $E\left[\frac{\partial^2}{\partial\theta\partial\theta^T}\ell(\theta_0)\right]$ is positive definite. From Lemma 2, $E[q(X,Z)B(Z)B(Z)^T]$ is the positive definite for all continuous and bounded function $q: \mathcal{X} \times \mathcal{Z} \to \mathbb{R}$. Condition (C1) implies that $E[q(X,Z)A(X,Z)A(X,Z)^T]$ is also positive definite. Lastly, for any $v = \mathbb{R}^{2(p+K+\xi)}$ with ||v|| = 1, which can be devided as $v = (v_{\gamma}^T, v_{\sigma}^T)^T$ with $v_{\gamma}, v_{\sigma} \in \mathbb{R}^{p+K+\xi}$ and $||v_{\gamma}|| \leq 1, ||v_{\sigma}|| \leq 1$. Then,

$$v^{T}E\left[\frac{\partial^{2}}{\partial\theta\partial\theta^{T}}\ell(\theta_{0})\right]v = 2v_{\gamma}^{T}E[q_{1}(X,Z)A(X,Z)A(X,Z)^{T}]v\gamma + 2v_{\gamma}^{T}E[q_{1}(X,Z)A(X,Z)A(X,Z)^{T}]v_{\sigma} + v_{\sigma}^{T}E[q_{2}(X,Z)A(X,Z)A(X,Z)^{T}]v_{\sigma},$$

where $q_1(x, z) = 2/(2\gamma_0(x, z) + 1)(\gamma_0(x, z) + 1)$ and $q_2(x, z) = 1/(2\gamma_0(x, z) + 1)$. Consequently, Cauchy-Schwarz integrality yeids that the maximum (minimum) eigen value of $v^T E\left[\frac{\partial^2}{\partial\theta\partial\theta^T}\ell(\theta_0)\right]v$ is bounded from above (below) by some positive constant, which completes the proof.

We next show the consistency of $||\hat{\theta} - \theta_0||$. However, the purpose of Lemma 8 is not deriving optimal rate of convergence, which is shown in Lemma 9. Since Lemma 8 is used to prove Lemma 9, we show Lemma 8 in advance. Throughout the proof of Lemma 8, we use C, C_1, C_2 and C^* to denote universal constants and these values may change from line to line.

Lemma 8. Suppose that (C1)–(C5). Define the positive sequence $\{\varepsilon_N\}$, which satisfies $\varepsilon_N \to 0$ and

$$\frac{\max\{K^{-m}, (N/n)^{\rho}, n^{-1/4}\}}{\varepsilon_N} \to 0.$$

Then, in each scenario (S1), (S2) or (S3), as $N \to \infty$,

$$P\left(||\hat{\theta} - \theta_0|| > \varepsilon_N\right) \to 0.$$

Proof of Lemma 8. We consider the probability of the event that $||\hat{\theta} - \theta_0|| > \varepsilon_N$. However, the important things is only rate of ε_N , and the constant part is not important. Therefore, for some constant C > 0, $C\varepsilon_N$ is replaced with ε_N below for simple notation.

From Hijort and Pollard (2011), we see that for $\varepsilon_N > 0$,

$$P\left(||\hat{\theta} - \theta_0|| > \varepsilon_N\right) \le P\left(\sup_{||\theta - \theta_0|| \le \varepsilon_N} |\ell_{pen}(\theta) - L(\theta)| \ge 2^{-1} \inf_{||\theta - \theta_0|| = \varepsilon_N} |L(\theta) - L(\theta_0)|\right).$$

Thus, our purpose is to show as $N \to \infty$,

$$P\left(\sup_{||\theta-\theta_0||\leq\varepsilon_N} |\ell_{pen}(\theta) - L(\theta)| \geq 2^{-1} \inf_{||\theta-\theta_0||=\varepsilon_N} |L(\theta) - L(\theta_0)|\right) \to 0.$$

Under $||\theta - \theta_0|| = \varepsilon_N$, we can write $\theta = \theta_0 + \varepsilon_N \eta$, where η is $2(p + K + \xi)$ -vector satisfying $||\eta|| = 1$. Then, we have

$$L(\theta) - L(\theta_0) = L'(\theta_0)(\theta - \theta_0) + \frac{1}{2}(\theta - \theta_0)^T L''(\theta^*)(\theta - \theta_0)$$
$$= \varepsilon_N L'(\theta_0)\eta + \frac{\varepsilon_N^2}{2}\eta^T L''(\theta^*)\eta,$$

where $L'(\theta) = \partial L(\theta)/\partial \theta$, $L''(\theta) = \partial^2 L(\theta)/\partial \theta \partial \theta^T$ and θ^* is the $2(p + K + \xi)$ -vector satisfying $||\theta^* - \theta_0|| < ||\hat{\theta} - \theta_0||$. Similar to the proof of Lemma 6, we have

$$L'(\theta_0)$$

$$= E_{X,Z} \left[E_{Y|X,Z} \left[\begin{array}{c} \ell_{\gamma}(Y|\bar{\gamma}_0(X,Z),\bar{\sigma}_{w0}(X,Z))A(X,Z) \\ \ell_{\sigma}(Y|\bar{\gamma}_0(X,Z),\bar{\sigma}_{w0}(X,Z))A(X,Z) \end{array} \right] \right]$$

$$= E_{X,Z} \left[\begin{array}{c} O(K^{-\zeta} + \alpha(w|X,Z))A(X,Z) \\ O(K^{-\zeta} + \alpha(w|X,Z))A(X,Z) \end{array} \right].$$

Next, as $\varepsilon_N \to 0$, we have

$$\begin{split} L''(\theta_0) \\ &= E_{X,Z} \begin{bmatrix} \frac{2}{(2\gamma_0(X,Z)+1)(\gamma_0(X,Z)+1)} A(X,Z) A(X,Z)^T & \frac{1}{(2\gamma_0(X,Z)+1)(\gamma_0(X,Z)+1)} A(X,Z) A(X,Z)^T \\ \frac{1}{(2\gamma_0(X,Z)+1)(\gamma_0(X,Z)+1)} A(X,Z) A(X,Z)^T & \frac{1}{2\gamma_0(X,Z)+1} A(X,Z) A(X,Z)^T \\ \times (1+o(1)), \end{split}$$

which is obtained by the straightforward calculation of expectation of Hessian matrix of loglikelihood of GPD. Here, we can obtain that $||L'(\theta_0)|| < O(K^{-\zeta}) + O((N/n)^{\rho}) = o(\varepsilon_N)$. Since $L''(\theta_0)$ is positive definite, there exist constants $C, C_1 > 0$ such that

$$L(\theta) - L(\theta_0) = \varepsilon_N L'(\theta_0) \eta + \frac{\varepsilon_N^2}{2} \eta^T L''(\theta^*) \eta$$

= $o(\varepsilon_N^2) + C_1 \varepsilon_N^2$
 $\geq C \varepsilon_N^2.$

This implies that

$$P\left(\sup_{||\theta-\theta_0||\leq\varepsilon_N} |\ell(\theta) - L(\theta)| \geq 2^{-1} \inf_{||\theta-\theta_0||=\varepsilon_N} |L(\theta) - L(\theta_0)|\right)$$
$$\leq P\left(\sup_{||\theta-\theta_0||\leq\varepsilon_N} |\ell(\theta) - L(\theta)| \geq \varepsilon_N^2\right)$$

Next, since

$$\sup_{\substack{||\theta-\theta_0|| \le \varepsilon_N}} |\ell_{pen}(\theta) - L(\theta)|$$

$$\leq |\ell_{pen}(\theta_0) - L(\theta_0)| + \sup_{\substack{||\theta-\theta_0|| \le \varepsilon_N}} |\ell_{pen}(\theta) - \ell_{pen}(\theta_0) - L(\theta) + L(\theta_0)|,$$

we have

$$P\left(\sup_{||\theta-\theta_0||\leq\varepsilon_N} |\ell_{pen}(\theta) - L(\theta)| \geq \varepsilon_N^2\right)$$

$$\leq P\left(|\ell_{pen}(\theta_0) - L(\theta_0)| \geq 2^{-1}\varepsilon_N^2\right) + P\left(\sup_{||\theta-\theta_0||\leq\varepsilon_N} |\ell_{pen}(\theta) - \ell_{pen}(\theta_0) - L(\theta) + L(\theta_0)| \geq 2^{-1}\varepsilon_N^2\right)$$

$$\equiv \mathcal{J}_1 + \mathcal{J}_2.$$

We first show $\mathcal{J}_1 \to 0$. From de Boor (2001), since $m < \zeta$, we have that for $z^{(j)} \in [0,1]$, $|d^m(\bar{g}_{0j}(z^{(j)}) - g_{0j}(z^{(j)}))/dz^m| = O(K^{-\zeta+m}) = o(1)$ and $|(d^m\bar{s}_{0j}(z^{(j)}) - s_{0j}(z^{(j)}))/dz^m| = O(K^{-\zeta+m}) = o(1)$ for j = 1, ..., d, we obtain

$$\int_0^1 \{\bar{g}_j^{(m)}(z)\}^2 dz = \int_0^1 \{g_j^{(m)}(z)\}^2 dz = O(1)$$

and

$$\int_{0}^{1} \{\bar{s}_{j}^{(m)}(z)\}^{2} dz = \int_{0}^{1} \{s_{j}^{(m)}(z)\}^{2} dz = O(1).$$

Since $(\lambda + \nu)/\varepsilon_N^2 = O((K^{-m}/\varepsilon_N)^2) \to 0$, we obtain

$$\frac{1}{\varepsilon_N^2} \left\{ \sum_{j=1}^d \lambda \int_0^1 \{ \bar{g}_j^{(m)}(z) \}^2 dz + \nu \int_0^1 \{ \bar{s}_j^{(m)}(z) \}^2 dz \right\} \to 0.$$

Therefore, to show $\mathcal{J}_1 \to 0$, we next prove the probability that

$$\left|\frac{1}{n}\sum_{i=1}^{n} -\log h(Y_i|\bar{\gamma}_0(X_i, Z_i), \bar{\sigma}_{w0}(X_i, Z_i)) - E[-\log h(Y_i|\bar{\gamma}_0(X_i, Z_i), \bar{\sigma}_{w0}(X_i, Z_i))]\right|$$

is greater than ε_N^2 converges to zero. Let $E_i = -\log h(Y_i | \bar{\gamma}_0(X_i, Z_i), \bar{\sigma}_{w0}(X_i, Z_i))$ and define the event

$$\mathcal{M} = \left\{ \max_{1 \le i \le n} E_i < M \right\}$$

with M > 0. Then, $P(\mathcal{M}^c) \leq ne^{-C_2 M}$. This implies that

$$|E_i I(\mathcal{M}^c) - E[E_i I(\mathcal{M}^c)]| \le o_P(e^{-M}) = o(\varepsilon_N^2)$$

if $M = 1/\varepsilon_n^2$. We next show that

$$P\left(\left|\frac{1}{n}\sum_{i=1}^{n}E_{i}I(\mathcal{M})-E[E_{i}I(\mathcal{M})]\right|>\varepsilon_{N}^{2}\right)\to0.$$

It is easy to show that $V[(1/n)E_i] < n^{-2}C_1$ for some constant $C_1 > 0$ and for some constant M > 0, we obtain $|n^{-1}E_i| < M/n$ for some constant $C_2 > 0$ on the event \mathcal{M} . Therefore, from Lemma 1, we have

$$P\left(\left|\frac{1}{n}\sum_{i=1}^{n}E_{i}I(\mathcal{M})-E[E_{i}I(\mathcal{M})]\right|>\varepsilon_{N}^{2}\right)\leq 2\exp\left[-\frac{2^{-1}C^{2}\varepsilon_{N}^{4}}{C_{1}/n+3^{-1}C\varepsilon_{N}^{2}M/n}\right].$$

Putting $M = 1/\varepsilon_N$, we obtain as $N \to \infty$,

$$P\left(\left|\frac{1}{n}\sum_{i=1}^{n}E_{i}I(\mathcal{M})-E[E_{i}I(\mathcal{M})]\right|>\varepsilon_{N}^{2}\right)\leq2\exp\left[-C_{3}n\varepsilon_{N}^{4}\right]\rightarrow0$$

since $n^{1/4}\varepsilon_N \to \infty$, where C_3 is some positive constant. Thus, we obtain

$$\mathcal{J}_1 \to 0 \ (N \to \infty).$$

Next, we show $\mathcal{J}_2 \to 0$ as $n \to \infty$. For $\{\theta : ||\theta - \theta_0|| \leq \varepsilon_N\}$, we write $\theta = \theta_0 + \varepsilon_N \eta$ with $||\eta|| \leq 1$. From Taylor expansion, we have

$$\ell_{pen}(\theta) - \ell_{pen}(\theta_0) = \varepsilon_N \frac{\partial}{\partial \theta} \ell(\theta_0) \eta + \frac{\varepsilon_N^2}{2} \eta^T \frac{\partial^2 \ell(\theta_0 + \delta_{1n}\eta)}{\partial \theta \partial \theta^T} \eta + \sum_{j=1}^d \lambda \left[\int_0^1 \left\{ \frac{d^m}{dx^m} B^{[\xi]}(x)^T (b_{0j} + \varepsilon_N \eta_{\gamma,j}) \right\}^2 dx - \int_0^1 \left\{ \frac{d^m}{dx^m} B^{[\xi]}(x)^T b_{0j} \right\}^2 dx \right] + \sum_{j=1}^d \nu \left[\int_0^1 \left\{ \frac{d^m}{dx^m} B^{[\xi]}(x)^T (c_{0j} + \varepsilon_N \eta_{\sigma j}) \right\}^2 dx - \int_0^1 \left\{ \frac{d^m}{dx^m} B^{[\xi]}(x)^T c_{0j} \right\}^2 dx \right],$$

where $\delta_{1n} = \varepsilon_N \operatorname{diag}(q_1, \ldots, q_{2(p+K+\xi)})$ with $q_i \in (0, 1)$, $\eta_{\gamma,j}$ and $\eta_{\sigma,j}$ are the $(K + \xi)$ -subvector of η corresponding to b_j and c_j in θ . Similarly, we have

$$L(\theta) - L(\theta_0) = \varepsilon_N \frac{\partial}{\partial \theta} L(\theta_0) \eta + \frac{\varepsilon_N^2}{2} \eta^T \frac{\partial^2 L(\theta_0 + \delta_{2n} \eta)}{\partial \theta \partial \theta^T} \eta_2$$

where $\delta_{2n} = \varepsilon_N \operatorname{diag}(q_{2,1}, \dots, q_{2,2(p+K+\xi)})$ with $q_{2,i} \in (0,1)$.

From the property of *B*-spline (see, de Boor 2001), we have $\int ||d^m B^{[\xi]}(x)/dx^m||dx = O(K^{m-1/2})$. Since $||\eta_{\gamma,j}|| < 1$ and $||\eta_{\sigma,j}|| < 1$, we have

$$\sum_{j=1}^{d} \lambda \left[\int_0^1 \left\{ \frac{d^m}{dx^m} B^{[\xi]}(x)^T (b_{0j} + \varepsilon_N \eta_{\gamma,j}) \right\}^2 dx - \int_0^1 \left\{ \frac{d^m}{dx^m} B^{[\xi]}(x)^T b_{0j} \right\}^2 dx \right] = O(\lambda K^m \varepsilon_N)$$

and

$$\sum_{j=1}^{d} \nu \left[\int_{0}^{1} \left\{ \frac{d^{m}}{dx^{m}} B^{[\xi]}(x)^{T} (c_{0j} + \varepsilon_{N} \eta_{\sigma j}) \right\}^{2} dx - \int_{0}^{1} \left\{ \frac{d^{m}}{dx^{m}} B^{[\xi]}(x)^{T} c_{0j} \right\}^{2} dx \right] = O(\nu K^{m} \varepsilon_{N}).$$

From the condition of ε_N , we have $(\lambda + \nu)K^m \varepsilon_N = o(\varepsilon_N^2)$. Next, $\partial^2 \ell(\theta) / \partial \theta \partial \theta^T$ is Hessian matrix of log-likelihood of θ and is continuous for θ , it converges to its expectation. Thus,

$$\frac{\partial^2 \ell(\theta_0 + \delta_{1n}\eta)}{\partial \theta \partial \theta^T} \xrightarrow{P} \frac{\partial^2 L(\theta_0)}{\partial \theta \partial \theta^T}, \quad \frac{\partial^2 L(\theta_0 + \delta_{2n}\eta)}{\partial \theta \partial \theta^T} \rightarrow \frac{\partial^2 L(\theta_0)}{\partial \theta \partial \theta^T}$$

and

$$\eta^T \left(\frac{\partial^2 \ell(\theta_0 + \delta_{1n} \eta)}{\partial \theta \partial \theta^T} - \frac{\partial^2 L(\theta_0 + \delta_{2n} \eta)}{\partial \theta \partial \theta^T} \right) \eta^T \xrightarrow{P} 0$$

as $n \to \infty$. That is, we have

$$\ell_{pen}(\theta) - \ell_{pen}(\theta_0) - L(\theta) + L(\theta_0) = \varepsilon_N \left(\frac{\partial \ell(\theta_0)}{\partial \theta} - \frac{\partial L(\theta_0)}{\partial \theta}\right) \eta + o(\varepsilon_N^2).$$

This implies that it only needs to show

$$P\left(\sup_{||\eta||\leq 1} \left| \varepsilon_N\left\{\frac{\partial}{\partial\theta}\ell(\theta_0) - E\left[\frac{\partial}{\partial\theta}\ell(\theta_0)\right]\right\}\eta \right| > \varepsilon_N^2\right) \to 0$$

to prove $\mathcal{J}_2 \to 0$. Define

$$\ell_1(\eta) = \left\{ \frac{\partial}{\partial \theta} \ell(\theta_0) - E\left[\frac{\partial}{\partial \theta} \ell(\theta_0) \right] \right\} \eta.$$

Let $\mathcal{E} = \{\eta \in \mathbb{R}^{2(p+K+\xi)} : ||\eta|| \leq 1\}$ and let $\mathcal{E}_1, \ldots, \mathcal{E}_N$ be a covering of \mathcal{E} with a diameter $R_n = c^* \varepsilon_N$ for some constant $c^* > 0$ such that $\mathcal{E} \subseteq \bigcup_{i=1}^{N} \mathcal{E}_i$. Thanks to Lemma 2.5 of van de Geer (2000), we can set $\mathcal{N} \leq c^* (1/\varepsilon_N)^{2(p+K+\xi)}$. Let fixed point $\eta^{(i)} \in \mathcal{E}_i, i = 1, \ldots, \mathcal{N}$ and then, $\mathcal{E}_i = \{\eta \in \mathbb{R}^{2(p+K+\xi)} || |\eta - \eta^{(i)} || < R_n\}$. We further define the event

$$\mathcal{M}_1 = \left\{ \max_{1 \le i \le n} -\log h(Y_i | \gamma_0(X_i, Z_i), \sigma_{w0}(X_i, Z_i)) < M_1 \right\}$$

for some constant $M_1 > 0$. Then,

$$\ell_1(\eta) = \ell_1(\eta)I(\mathcal{M}_1) + \ell_1(\eta)I(\mathcal{M}_1^c).$$

From $P(\mathcal{M}_1^c) = ne^{-M_1}(1 + o(1))$, it is easy to show that

$$|\ell_1(\eta)I(\mathcal{M}_1^c)| \le o_P(\varepsilon_N)$$

if M_1 is appropriately chozen (see, below). Next, we consider

$$P\left(\sup_{||\eta|| \le 1} |\ell_1(\eta)I(\mathcal{M}_1)| \middle| > \varepsilon_N\right) = P\left(\max_{1 \le j \le \mathcal{N}} \sup_{\eta \in \mathcal{E}_j} |\ell_1(\eta)I(\mathcal{M}_1)| \middle| > \varepsilon_N\right)$$

$$\leq \sum_{j=1}^{\mathcal{N}} P\left(\sup_{\eta \in \mathcal{E}_j} |\ell_1(\eta)I(\mathcal{M}_1)| \middle| > \varepsilon_N\right)$$

$$\leq \sum_{j=1}^{\mathcal{N}} P\left(\left|\ell_1(\eta^{(j)})I(\mathcal{M}_1)\right| + \sup_{\eta \in \mathcal{E}_j} \left|\{\ell_1(\eta) - \ell_1(\eta^{(j)})\}I(\mathcal{M}_1)\right| \middle| > \varepsilon_N\right).$$

It is easy to show from Lemma 5 that

$$\sup_{\eta \in \mathcal{E}_j} \left| \{\ell_1(\eta) - \ell_1(\eta^{(j)})\} I(\mathcal{M}_1) \right| \le \left| \left| \frac{\partial}{\partial \theta} \ell(\theta_0) - E\left[\frac{\partial}{\partial \theta} \ell(\theta_0) \right] \right| \left| \sup_{\eta \in \mathcal{E}_j} ||\eta - \eta^{(j)}|| = O_P\left(\sqrt{\frac{K}{n}} R_n\right) = o_P\left(\varepsilon_N\right) \right|$$

Latstly, we show

$$\sum_{j=1}^{\mathcal{N}} P\left(\left|\ell_1(\eta^{(j)})I(\mathcal{M}_1)\right| > \varepsilon_N \middle| \mathcal{M}_1\right) \to 0.$$
(17)

We show that (17) converges to zero for each case of (S1)–(S3). Here, by the definition of $\partial \ell(\theta_0)/\partial \theta$, we obtain

$$\ell_1(\eta^{(j)}) = \frac{1}{n} \sum_{i=1}^n \left[\begin{array}{c} \ell_{\gamma}(Y_i | \bar{\gamma}_0(X_i, Z_i), \bar{\sigma}_{w0}(X_i, Z_i)) A(X_i, Z_i) \\ \ell_{\sigma}(Y_i | \bar{\gamma}_0(X_i, Z_i), \bar{\sigma}_{w0}(X_i, Z_i)) A(X_i, Z_i) \end{array} \right] \eta^{(j)} + o(1) \equiv \sum_{i=1}^n \frac{1}{n} \ell_{i,1}(Y_i, X_i, Z_i) + o(1)$$

From Lemma 7 and the fact that $||\eta^{(j)}|| \leq 1$, we see that $V[\ell_{i,1}(Y_i, X_i, Z_i)/n] < C_1 n^{-2}$ for some constant $C_1 > 0$. We first consider case (S1). We put $M_1 = 1/\varepsilon_N$. Then, the straightforward calculation yields that $|\ell_{\gamma}(Y_i|\bar{\gamma}_0(X_i, Z_i), \bar{\sigma}_{w0}(X_i, Z_i))| \leq M_1/\gamma_{min}$ and $|\ell_{\sigma}(Y_i|\bar{\gamma}_0(X_i, Z_i), \bar{\sigma}_{w0}(X_i, Z_i))| \leq$ 1 on the event \mathcal{M}_1 . In addition, we have $\max_i |\ell_{i,1}(Y_i, X_i, Z_i)/n| < C_2 M_1/n$ for some constant $C_2 > 0$. From Bernstein's inequality (see, Lemma 1), we obtain

$$P\left(|\ell_1(\eta^{(j)})I(\mathcal{M}_1)| > \varepsilon_N\right) \le 2\exp\left[-\frac{2^{-1}\varepsilon_N^2}{C_1/n + C_2\varepsilon_N M_1/n}\right] \le 2\exp\left[-C^*n\varepsilon_N^2\right].$$

for some constant $C^* > 0$. From the condition of ε_N that $n\varepsilon_N^2/\{K\log(N)\} \to \infty$, we have $\mathcal{N}\exp[-C^*n\varepsilon_N^2] \to 0$, which implies (17).

We next assume (S2). Then, on the event \mathcal{M}_1 , we have $|\ell_{\gamma}(Y_i|\bar{\gamma}_0(X_i, Z_i), \bar{\sigma}_{w0}(X_i, Z_i))| \leq Ce^{M_1/(2+\delta)}$ and $|\ell_{\sigma}(Y_i|\bar{\gamma}_0(X_i, Z_i), \bar{\sigma}_{w0}(X_i, Z_i))| \leq Ce^{M_1/(2+\delta)}$ for some constant C > 0. These imply that $\max_i |\ell_{i,1}(Y_i, X_i, Z_i)/n| < C_2 e^{M_1/(2+\delta)}/n$. In this case, we put $M_1 = \log(n \log N)$. Then, Lemma 1 and the condition $n^{(1+\delta)/(2+\delta)} \varepsilon_N \to \infty$ yield that

$$P\left(|\ell_1(\eta^{(j)})| > \varepsilon_N\right) \le 2\exp\left[-\frac{2^{-1}\varepsilon_N^2}{C_1/n + C_2\varepsilon_N e^{M_1/(2+\delta)}/n}\right] \le \exp\left[-C^* n^{(1+\delta)/(2+\delta)}\varepsilon_N\right]$$

for some constant $C^* > 0$. Since $\varepsilon_N > O(n^{-1/4})$, for any $\delta > 0$, we have $n^{(1+\delta)/(2+\delta)}\varepsilon_N \to \infty$. Thus, (17) can be shown.

In the case (S3), we have $|\ell_{\gamma}(Y_i|\bar{\gamma}_0(X_i,Z_i),\bar{\sigma}_{w0}(X_i,Z_i))| \leq CM_1^2$ and

$$|\ell_{\sigma}(Y_i|\bar{\gamma}_0(X_i, Z_i), \bar{\sigma}_{w0}(X_i, Z_i))| \le CM_1$$

for some constant C > 0. When we put $M_1 = 1/\sqrt{\varepsilon_N}$, (17) can be proven similar to the case (S1). Thus, the proof was completed.

Lemma 9. Suppose that (C1)–(C5). In each scenario (S1), (S2) or (S3), as $N \to \infty$,

$$E[||\hat{\theta} - \theta_0||^2] \le O_P\left(\frac{K}{n}\right) + O(\lambda^2 K^{2m}) + O(\nu^2 K^{2m}) + O\left(\left(\frac{N}{n}\right)^{2\rho}\right).$$

Proof of Lemma 9. From Lemma 8, we can use the Taylor expansion to $\partial \ell_{pen}(\theta)/\partial \theta$ around $\hat{\theta} = \theta_0$. This indicates that

$$\left(\frac{\partial^2 \ell_{pen}(\theta^*)}{\partial \theta \partial \theta^T}\right)(\hat{\theta} - \theta_0) = \frac{\partial}{\partial \theta} \ell_{pen}(\theta_0),$$

where θ^* is the parameter satisfying $||\theta^* - \theta_0|| < ||\hat{\theta} - \theta_0||$. From Lemma 7 and Cauchy–Schwarz inequality, we obtain

$$\left| \left| \left(\frac{\partial^2 \ell_{pen}(\theta^*)}{\partial \theta \partial \theta^T} \right) (\hat{\theta} - \theta_0) \right| \right|^2 \ge C_m^2 ||\hat{\theta} - \theta_0||^2.$$

Together with Lemmas 5 and 6, we obtain

$$E\left[\left\|\left|\frac{\partial}{\partial\theta}\ell_{pen}(\theta_0)\right|\right\|^2\right] \le O_P\left(\frac{K}{n}\right) + O(\lambda^2 K^{2m}) + O(\nu^2 K^{2m}) + O\left(\left(\frac{N}{n}\right)^{2\rho}\right).$$

For two random variable A_n and B, $A_n \stackrel{a.s.}{\sim} B$ means that A_n and B have same distribution as $n \to \infty$.

Lemma 10. Suppose that (C1)–(C5). In each scenario (S1), (S2) or (S3), as $N \to \infty$,

$$\sqrt{n} \left\{ \frac{\partial \ell(\theta_0)}{\partial \theta} - E\left[\frac{\partial \ell(\theta_0)}{\partial \theta} \right] \right\} \stackrel{a.s}{\sim} N_{2(p+K+\xi)}(0,\Sigma).$$

Proof of Lemma 10. In this proof, we denote $\ell_{\gamma}(Y_i|X_i, Z_i) = \ell_{\gamma}(Y_i|\bar{\gamma}_0(X_i, Z_i), \bar{\sigma}_{w0}(X_i, Z_i))$ and $\ell_{\sigma}(Y_i|X_i, Z_i) = \ell_{\sigma}(Y_i|\bar{\gamma}_0(X_i, Z_i), \bar{\sigma}_{w0}(X_i, Z_i))$. Similar to the proof of Lemma 6), we have

$$\frac{\partial \ell(\theta_0)}{\partial \theta} = \frac{1}{n} \sum_{i=1}^n \left[\begin{array}{c} \ell_{\gamma}(Y_i | X_i, Z_i) A(X_i, Z_i) \\ \ell_{\sigma}(Y_i | X_i, Z_i) A(X_i, Z_i) \end{array} \right].$$

To show the asymptotic normality of derivative of log-likelihood, we use Cramér-Wold theorem. For any vector $r = (r_{\gamma}^T, r_{\sigma}^T)^T \in \mathbb{R}^{2(p+K+\xi)}$ with $r_{\gamma} \in \mathbb{R}^{p+K+\xi}, r_{\sigma} \in \mathbb{R}^{p+K+\xi}$, we consider

$$W = \sqrt{n}r^T \left\{ \frac{\partial \ell(\theta_0)}{\partial \theta} - E\left[\frac{\partial \ell(\theta_0)}{\partial \theta}\right] \right\} = \sum_{i=1}^n W_i,$$

where

$$W_{i} = \frac{1}{\sqrt{n}} \left[r_{\gamma}^{T} \left\{ \ell_{\gamma}(Y_{i}|X_{i}, Z_{i}) A(X_{i}, Z_{i}) - E\left[\ell_{\gamma}(Y|X, Z) A(X, Z) \right] \right\} + r_{\sigma}^{T} \left\{ \ell_{\gamma}(Y_{i}|X_{i}, Z_{i}) A(X_{i}, Z_{i}) - E\left[\ell_{\gamma}(Y|X, Z) A(X, Z) \right] \right\} \right].$$

Our aim is to check whether $\sum_{i=1}^{n} W_i$ satisfies the Lyapnov's condition of central limit theorem (CLT). It is easy to find that $E[W_i] = 0$. Next, from the proof of Lemma 6, we have $E_{Y|x,z}[\ell_{\gamma}(Y|x,z)] = o(1)$ and $E_{Y|x,z}[\ell_{\sigma}(Y|x,z)] = o(1)$. Thus, $V[W_i] = E[W_i^2](1+o(1))$. In addition, the property of derivative of log-likelihood, Lemma 3 and proof of Lemma 7, we obtain

$$E[\{r_{\gamma}^{T}\{\ell_{\gamma}(Y_{i}|X_{i},Z_{i})A(X_{i},Z_{i})\}^{2}] = r_{\gamma}^{T}E\left[\frac{2}{(2\gamma_{0}(X,Z)+1)(\gamma_{0}(X,Z)+1)}A(X,Z)A(X,Z)^{T}\right]r_{\gamma}(1+o(1))$$

$$= O(K),$$

$$E[\{r_{\sigma}^{T}\{\ell_{\sigma}(Y_{i}|X_{i},Z_{i})^{2}X_{i}B(Z_{i})^{T}] = r_{\sigma}^{T}E\left[\frac{1}{2\gamma_{0}(X,Z)+1}A(X,Z)A(X,Z)^{T}\right]r_{\sigma}(1+o(1))$$

$$= O(K)$$

and

$$E[\{r_{\gamma}^{T}\{\ell_{\gamma}(Y_{i}|X_{i},Z_{i})A(X_{i},Z_{i})\}\{r_{\sigma}^{T}\{\ell_{\sigma}(Y_{i}|X_{i},Z_{i})A(X_{i},Z_{i})\}]$$

= $r_{\gamma}^{T}E\left[\frac{1}{(2\gamma_{0}(X,Z)+1)(\gamma_{0}(X,Z)+1)}A(X,Z)A(X,Z)^{T}\right]r_{\sigma}(1+o(1))$
= $O(K).$

Thus, $E[W_i^2] = O(K/n)$. Lastly, we need to show that

$$\frac{1}{(\sum_{i=1}^{n} V[W_i])^{(2+\varepsilon)/2}} \sum_{i=1}^{n} E[|W_i|^{2+\varepsilon}] \to 0$$

for some $\varepsilon > 0$. Here, we put $\varepsilon = \delta$, where δ is given in (S2). Note that we can use similar $\varepsilon = \delta$ even for (S1) and (S3). To evaluate $E[|W_i|^{2+\delta}]$, we calculate

$$E_{Y|x,z}[|\ell_{\gamma}(Y|x,z)|^{2+\delta}] = \int_{0}^{y^{*}(x,z)} |\ell_{\gamma}(y|x,z)|^{2+\delta} \frac{1}{\sigma(x,z)} \left(1 + \frac{\gamma_{0}(x,z)y}{\sigma_{w0}(x,z)}\right)^{-1/\gamma_{0}(x,z)-1} dy + o(1).$$

The part of o(1) is followed by Conditions (C2), (C4) and the definition of Q. Under (S1), that is, when $\gamma_0(x,z) > 0$, it is easy to show that $E_{Y|x,z}[|\ell_{\gamma}(Y|x,z)|^{2+\delta}] < \infty$ since $||\ell_{\gamma}(Y|x,z)|^{2+\delta}| < C \log(1 + y\gamma_0(x,z)/\sigma_{w0}(x,z))$ for some constant C > 0. For (S2), we need to carefully calculate $E_{Y|x,z}[|\ell_{\gamma}(Y|x,z)|^{2+\delta}]$. Since $y^*(x,z) = -\sigma_{w0}(x,z)/\gamma_0(x,z)$, the straightforward calculation yields that

$$E_{Y|x,z}[|\ell_{\gamma}(Y|x,z)|^{2+\delta}] < \frac{C}{(2+\delta)\gamma_0(x,z)+1}(1+o(1))$$

for some constant C > 0. Thus, from the condition of $(S2) : -1/(2+\delta) < \gamma_0(x,z) < 0$, we have $E_{Y|x,z}[|\ell_{\gamma}(Y|x,z)|^{2+\delta}] < \infty$. For case (S3), $E_{Y|x,z}[|\ell_{\gamma}(Y|x,z)|^{2+\delta}] < \infty$ can easily be shown since the distribution of Y is approximated to exponential distribution. Similarly, we can evaluate $E_{Y|x,z}[|\ell_{\sigma}(Y|x,z)|^{2+\delta}] < \infty$ under each (S1), (S2) or (S3).

Next, from the property of B-spline function, we have

$$E[|r_g^T B(Z) + r_s^T B(Z)|^{2+\delta}] = O(K^{(2+\delta)/2} K^{-1} K) = O(K^{(2+\delta)/2}).$$

Accordingly, we obtain $E[|W_i|^{2+\delta}] = O(K^{(2+\delta)/2})$ and

$$\sum_{i=1}^{n} E[|W_i|^{2+\delta}] = O\left(\frac{K^{(2+\delta)/2}}{n^{\delta/2}}\right).$$

Consequently, we obtain

$$\frac{1}{\left(\sum_{i=1}^{n} V[W_i]\right)^{(2+\delta)/2}} \sum_{i=1}^{n} E[|W_i|^{2+\delta}] = O(n^{-\delta/2}) = o(1)$$

as $n \to \infty$. Thus, Lyapunov's condition of CLT is satisfied. From the proof of Lemma 5, we have $V[W] = r^T \Sigma r$. Consequently,

$$W = \sqrt{n}r^T \left\{ \frac{\partial \ell(\theta_0)}{\partial \theta} - E\left[\frac{\partial \ell(\theta_0)}{\partial \theta}\right] \right\} \xrightarrow{D} N(0, r^T \Sigma r)$$

can be obtained, which leads the assetion of this lemma.

Appendix C: Proof of Theorems

Proof of Theorem 1. Let $(\tilde{X}, \tilde{Z}) \in \mathcal{X} \times \mathcal{Z}$ be random variable with same distribution as (X_i, Z_i) independently. According to Appendix A, the difference between $\hat{\gamma}$ and γ_0 can be shown as

$$\hat{\gamma}(\tilde{X},\tilde{Z}) - \gamma_0(\tilde{X},\tilde{Z}) = \tilde{X}^T(\hat{\beta} - \beta_0) + B(\tilde{Z})^T(\hat{b} - b_0) + O(K^{-\zeta})$$

The condition (C1) implies that $E[|\tilde{X}(\hat{\beta} - \beta_0)^T|||^2] \leq CE[||\hat{\beta} - \beta_0||^2]$ for some constant C > 0. Lemma 2 and Cauchy-Schwarz inequality yield that

$$E||B(\tilde{Z})(\hat{b}-b_0)||^2] \le M_{\max}E[||\hat{b}-b_0||^2].$$

Thus, the asymptotic rate of $||\hat{\gamma} - \gamma_0||_{L_2}$ is dominated by that of $||\hat{\theta} - \theta_0||$. From Lemma 9, we obtain

$$||\hat{\gamma} - \gamma_0||_{L_2} \le O\left(\sqrt{\frac{K}{n}}\right) + O(\lambda K^m) + O(\nu K^m) + O\left(\left(\frac{N}{n}\right)^{\rho}\right) + O(K^{-\zeta}).$$

Here, we note that the term $O(K^{-\zeta})$ is negrible order compared with $O(\lambda K^m)$. Similarly, we have

$$\left|\left|\log\hat{\sigma} - \log\sigma_{w0}\right|\right|_{L_2} \le O\left(\sqrt{\frac{K}{n}}\right) + O(\lambda K^m) + O(\nu K^m) + O\left(\left(\frac{N}{n}\right)^{\rho}\right) + O(K^{-\zeta}).$$

Thus, the first assertion of Theorem 1 is shown. The remaining two assetions are obtained directly from straightforward calculation. $\hfill \Box$

Proof of Theorem 2. First, from Lemma 7, Σ is positive definite matrix. Accordingly, we write

$$\Sigma^{-1} = \left[\begin{array}{cc} S_{11} & S_{12} \\ S_{21} & S_{22} \end{array} \right],$$

where each $S_{ij}(i, j = 1, 2)$ is $(p + K + \xi)$ -square matrix. Similar to the proof of Lemma 9, we have that for $(x, z) \in \mathcal{X} \times \mathcal{Z}$,

$$\hat{\gamma}(x,z) - \gamma_0(x,z) = A(x,z)^T \begin{bmatrix} \hat{\beta} - \beta_0 \\ \hat{b} - b_0 \end{bmatrix} = A(x,z)^T \left\{ S_{11} \begin{bmatrix} \frac{\partial \ell_{pen}(\theta_0)}{\partial \beta} \\ \frac{\partial \ell_{pen}(\theta_0)}{\partial b} \end{bmatrix} + S_{12} \begin{bmatrix} \frac{\partial \ell_{pen}(\theta_0)}{\partial u} \\ \frac{\partial \ell_{pen}(\theta_0)}{\partial c} \end{bmatrix} \right\} + O(K^{-\zeta}).$$

From the proof of Lemma 10, we obtain

$$\begin{aligned} \hat{\gamma}(x,z) - \gamma(x,z) &= \frac{1}{n} \sum_{i=1}^{n} \left\{ \ell_{\gamma}(Y_{i}|X_{i},Z_{i})A(x,z)^{T}S_{11}A(X_{i},Z_{i}) + \ell_{\sigma}(Y_{i}|X_{i},Z_{i})A(x,z)^{T}S_{12}A(X_{i},Z_{i}) \right\} \\ &+ A(x,z)^{T}S_{11} \left[\begin{array}{c} \underline{0}_{p} \\ \lambda K^{2m}\Omega b_{0} \end{array} \right] + A(x,z)^{T}S_{12} \left[\begin{array}{c} \underline{0}_{p} \\ \nu K^{2m}\Omega c_{0} \end{array} \right] + O(K^{-\zeta}), \end{aligned}$$

where $\underline{0}_p$ is *p*-zero vector, $\ell_{\gamma}(y|x,z) = \ell_{\gamma}(y|\bar{\gamma}_0(x,z), \bar{\sigma}_{w0}(x,z)), \ \ell_{\sigma}(y|x,z) = \ell_{\sigma}(y|\bar{\gamma}_0(x,z), \bar{\sigma}_{w0}(x,z)),$ and Ω is that appeared in Lemma 6. From definition of normalized *B*-spline basis, $\sup_{(x,z)\in\mathcal{X}\times\mathcal{Z}} ||A(x,z)|| = O(\sqrt{K})$. Therefore, Lemma 6 implies that

$$A(x,z)^T S_{11} \begin{bmatrix} \underline{0}_p \\ \lambda K^{2m} \Omega b_0 \end{bmatrix} + A(x,z)^T S_{12} \begin{bmatrix} \underline{0}_p \\ \nu K^{2m} \Omega c_0 \end{bmatrix} = O(\lambda K^m) + O(\nu K^m).$$

Define

$$G_i(x,z) = \ell_{\gamma}(Y_i|X_i, Z_i)A(x,z)^T S_{11}A(X_i, Z_i) + \ell_{\sigma}(Y_i|X_i, Z_i)A(x,z)^T S_{12}A(X_i, Z_i).$$

and $G(x,z) = E[G_i(x,z)]$. Lemma 6 implies that $\sup_{(x,z)\in\mathcal{X}\times\mathcal{Z}} |G(x,z)| < O((N/n)^{\rho})$.

The remain of proof is to show

$$\sup_{(x,z)\in\mathcal{X}\times\mathcal{Z}} \left| \frac{1}{n} \sum_{i=1}^{n} G_i(x,z) - G(x,z) \right| = O(\sqrt{K \log N/n}).$$

Let $E_i = \gamma_0(X_i, Z_i)^{-1} \log(1 + Y_i \gamma_0(X_i, Z_i) / \sigma_{w0}(X_i, Z_i))$ for the cases (S1) and (S2). For the case (S3): $\gamma_0(x, z) = 0$, we set $E_i = Y_i / \sigma_{w0}(X_i, Z_i)$. Then, for constant M > 0, we have $P(E_i > M) = e^{-M}(1 + o(1))$. Define the event $\mathcal{E} = \{\max_i E_i < M\}$ and the sequence $\varepsilon_n = C_{\varepsilon} \sqrt{K \log N/n}$, where $C_{\varepsilon} > 0$ is the constant defined below. Then, we have $P(\mathcal{E}^c) = ne^{-M}(1 + o(1))$. We then have

$$\sup_{(x,z)\in\mathcal{X}\times\mathcal{Z}} \left| \frac{1}{n} \sum_{i=1}^{n} G_i(x,z) - G(x,z) \right| \leq \sup_{(x,z)\in\mathcal{X}\times\mathcal{Z}} \left| \frac{1}{n} \sum_{i=1}^{n} G_i(x,z)I(\mathcal{E}) - G(x,z)I(\mathcal{E}) \right| + \sup_{(x,z)\in\mathcal{X}\times\mathcal{Z}} \left| \frac{1}{n} \sum_{i=1}^{n} G_i(x,z)I(\mathcal{E}^c) - G(x,z)I(\mathcal{E}^c) \right|.$$

Since $\sup_{(x,z)\in\mathcal{X}\times\mathcal{Z}} ||A(x,z)|| = O(\sqrt{K})$, we have $\sup_{(x,z)\in\mathcal{X}\times\mathcal{Z}} V[G_i(x,z)] \leq O(K/n) = o(\varepsilon_n^2)$. Therefore, if we set $M > \log(n\log(N))$ defined as below, we have

$$\sup_{(x,z)\in\mathcal{X}\times\mathcal{Z}} \left| \frac{1}{n} \sum_{i=1}^{n} G_i(x,z) I(\mathcal{E}^c) - G(x,z) I(\mathcal{E}^c) \right| \le O(\varepsilon_n n e^{-M}) = o(\varepsilon_n).$$

Thus, all that remains is to show

$$\sup_{(x,z)\in\mathcal{X}\times\mathcal{Z}} \left| \frac{1}{n} \sum_{i=1}^{n} G_i(x,z) I(\mathcal{E}) - G(x,z) I(\mathcal{E}) \right| = O(\varepsilon_n).$$

We first obtain

$$P\left(\sup_{(x,z)\in\mathcal{X}\times\mathcal{Z}}\left|\frac{1}{n}\sum_{i=1}^{n}G_{i}(x,z)I(\mathcal{E})-G(x,z)I(\mathcal{E})\right|>\varepsilon_{n}\right)$$

$$\leq P\left(\sup_{(x,z)\in\mathcal{X}\times\mathcal{Z}}\left|\frac{1}{n}\sum_{i=1}^{n}G_{i}(x,z)I(\mathcal{E})-E[G_{i}(x,z)I(\mathcal{E})]\right|>\varepsilon_{n}(1+o(1))\right).$$

We take *J*-fixed points $(x_j^*, z_j^*) \in \mathcal{X} \times \mathcal{Z}(j = 1, ..., J)$ and define the set $\mathcal{X}_j \times \mathcal{Z}_j = \{(x, z) || | (x, z) - (x_j^*, z_j^*) || < n^{-\eta} \}$ with $\eta > 0$ for j = 1, ..., J. The number *J* is quite large in order to satisfy $\mathcal{X} \times \mathcal{Z} \subset \bigcup_{j=1}^J \mathcal{X}_j \times \mathcal{Z}_j$. From Lemma 2.5 of van der Geer (2000), at least, for some constant $C_{\eta} > 0, J \leq C_{\eta} n^{\eta(p+d)}$ holds. Then,

$$\begin{split} \sup_{(x,z)\in\mathcal{X}\times\mathcal{Z}} \left| \frac{1}{n} \sum_{i=1}^{n} \{G_i(x,z)I(\mathcal{E}) - E[G_i(x,z)I(\mathcal{E})\}] \right| \\ &= \max_{j} \sup_{(x,z)\in\mathcal{X}_j\times\mathcal{Z}_j} \left| \frac{1}{n} \sum_{i=1}^{n} \{G_i(x,z)I(\mathcal{E}) - E[G_i(x,z)I(\mathcal{E})]\} - \{G_i(x_j^*,z_j^*)I(\mathcal{E}) - E[G_i(x_j^*,z_j^*)I(\mathcal{E})]\} \\ &+ \max_{j} \left| \frac{1}{n} \sum_{i=1}^{n} G_i(x_j^*,z_j^*)I(\mathcal{E}) - E[G_i(x_j^*,z_j^*)I(\mathcal{E})] \right|. \end{split}$$

Here, we choose η such that $n^{-\eta} = O(K^{-2})$. Then, from the Lipschitz continuity of *B*-spline basis (see, de Boor 2001) yields that $B(z) - B(z_j^*)$ has an order $O(K^{3/2}n^{-\eta}) = O(K^{-1/2})$ for only few element of and zero for other many elements. Therefore, under \mathcal{E} , we have from the

proof of Lemma 5 that

$$\max_{j} \sup_{(x,z)\in\mathcal{X}_{j}\times\mathcal{Z}_{j}} \left| \frac{1}{n} \sum_{i=1}^{n} \{G_{i}(x,z)I(\mathcal{E}) - E[G_{i}(x,z)I(\mathcal{E})]\} - \{G_{i}(x_{j}^{*},z_{j}^{*})I(\mathcal{E}) - E[G_{i}(x_{j}^{*},z_{j}^{*})I(\mathcal{E})]\} \right|$$

$$\leq O_{P}(Kn^{-1}) = o_{P}(\varepsilon_{n}).$$

Lastly, we show that

$$P\left(\max_{j}\left|\frac{1}{n}\sum_{i=1}^{n}G_{i}(x_{j}^{*},z_{j}^{*})I(\mathcal{E})-E[G_{i}(x_{j}^{*},z_{j}^{*})I(\mathcal{E})]\right|>\varepsilon_{n}\right)\to0$$
(18)

by using Lemma 1.

It can easily be described that for any $(x, z) \in \mathcal{X} \times \mathcal{Z}$, $V[n^{-1}G_i(x, z)] < C_1 n^{-2}K$ for some constant $C_1 > 0$ for all (S1)–(S3). We now consider the case (S1). Then, under \mathcal{E} , we have $|\ell_{\gamma}(Y_i|Z_i, X_i)| \leq M/\gamma_{min}$ and $|\ell_{\sigma}(Y_i|X_i, Z_i)| < 1$. Together with $A(x, z) = O(\sqrt{K})$, we see that $|n^{-1}\{G_i(x, z)I(\mathcal{E}) - E[G_i(x, z)I(\mathcal{E})]\}| \leq C_2 K n^{-1} M = C_2 K n^{-1} M$. Putting $M = 1/\varepsilon_n$, $P(\mathcal{E}^c) = ne^{-M}(1 + o(1)) = o(1)$. In addition, from Lemma 1, we obtain

$$P\left(\left|\frac{1}{n}\sum_{i=1}^{n}G_{i}(x_{j}^{*},z_{j}^{*})I(\mathcal{E})-E[G_{i}(x_{j}^{*},z_{j}^{*})I(\mathcal{E})]\right| > \varepsilon_{n}\right) \leq 2\exp\left[\frac{2^{-1}\varepsilon_{n}^{2}}{C_{1}K/n+3^{-1}C_{2}\varepsilon_{n}MK/n}\right]$$
$$\leq 2\exp\left[-C^{*}C_{\varepsilon}\log N\right]$$

for some constant $C^* > 0$. Since $J = O(n^{\eta(p+d)}) = O(K^{2(p+d)})$, if we choose C_{ε} such that $K^{2(p+d)}/N^{C^*C_{\varepsilon}} \to 0$, we have

$$P\left(\max_{j=1,\dots,J}\left|\frac{1}{n}\sum_{i=1}^{n}G_{i}(x_{j}^{*},z_{j}^{*})I(\mathcal{E})-E[G_{i}(x_{j}^{*},z_{j}^{*})I(\mathcal{E})]\right|>\varepsilon_{n}\right)\leq 2\mathcal{J}\exp\left[-C^{*}C_{\varepsilon}\log N\right]\to 0.$$

This implies (18).

We next focus on the case (S2): $-(2+\delta)^{-1} < \gamma_0(x,z) < 0$. Under \mathcal{E} , we obtain $|\ell_{\gamma}(Y_i|X_i, Z_i)| < Ce^{M/(2+\delta)}$ and $|\ell_{\sigma}(Y_i|X_i, Z_i)| < Ce^{M/(2+\delta)}$ for some constants C > 0. Thus, for any fixed point $(x,z) \in \mathcal{X} \times \mathcal{Z}$,

$$\frac{1}{n}\sum_{i=1}^{n}G_{i}(x,z)I(\mathcal{E}) - E[G_{i}(x,z)I(\mathcal{E})] \leq C_{2}e^{M/(2+\delta)}Kn^{-1}$$

for some constant $C_2 > 0$. Therefore, we put $M = \log(n \log N)$. This implies $P(\mathcal{E}^c) \to 0$ and under the condition that $\{K \log N\}^{1+\delta/2}/n^{\delta/2} \to 0$, $e^{M/(2+\delta)} = n^{1/(2+\delta)} \{\log N\}^{1/(2+\delta)} \le O(\varepsilon_n^{-1})$. Consequently, Lemma 1 shows that

$$P\left(\left|\frac{1}{n}\sum_{i=1}^{n}G_{i}(x_{j}^{*},z_{j}^{*})I(\mathcal{E})-E[G_{i}(x_{j}^{*},z_{j}^{*})I(\mathcal{E})]\right| > \varepsilon_{n}\right) \leq 2\exp\left[\frac{2^{-1}\varepsilon_{n}^{2}}{C_{1}K/n+3^{-1}C_{2}\varepsilon_{n}e^{M/(2+\delta)}K/n}\right]$$
$$\leq 2\exp\left[-C^{*}C_{\varepsilon}\log N\right].$$

Similar to the case (S1), we obtain (18).

For the case (iii): $\gamma_0(x, z) = 0$, we obtain $|\ell_{\gamma}(Y_i|X_i, Z_i)| < CM^2$ and $|\ell_{\sigma}(Y_i|X_i, Z_i)| < CM$ for some constants C > 0. When putting $M = M_n = 1/\sqrt{\varepsilon_n}$, $P(\mathcal{E}^c) \to 0$ and (18) can be shown as the same mannar as case (S1). Thus, in each (S1), (S2) or (S3), we obtain

$$||\hat{\gamma} - \gamma_0||_{L_{\infty}} \le O\left(\sqrt{\frac{K\log N}{n}}\right) + O(\lambda K^m) + O(\nu K^m) + O((N/n)^{\rho})$$

Similarly, the rate of convergence of $||\hat{\gamma} - \gamma_0||_{L_{\infty}}$ is shown. The optimal rate of $||\hat{\gamma} - \gamma_0||_{L_{\infty}}$ and $||\hat{\gamma} - \gamma_0||_{L_{\infty}}$ are proven directly. Thus, the proof is completed.

Proof of Theorem 3. From Lemma 3, we have

$$\begin{bmatrix} \hat{\gamma}(x,z) - \gamma_0(x,z) \\ \log \hat{\sigma}(x,z) - \log \sigma_{w0}(x,z) \end{bmatrix} = \begin{bmatrix} A(x,z)^T & 0_{p+K+\xi}^T \\ 0_{p+K+\xi}^T & A(x,z)^T \end{bmatrix} (\hat{\theta} - \theta_0) + o(\sqrt{K/N}).$$

Similar to proof of Lemma 9, Taylor expansion yields that

$$\hat{\theta} - \theta_0 = \Sigma^{-1} \frac{\partial \ell_{pen}(\theta_0)}{\partial \theta} (1 + o_P(1)) \\ = \Sigma^{-1} \left[\left\{ \frac{\partial \ell(\theta_0)}{\partial \theta} - E \left[\frac{\partial \ell(\theta_0)}{\partial \theta} \right] \right\} + E \left[\frac{\partial \ell(\theta_0)}{\partial \theta} \right] + \Omega_{\gamma,\sigma} \theta_0 \right] (1 + o_P(1)).$$

Here, we note that $A(x,z) = O(\sqrt{K})$. From the proof of Lemma 6 and condition that $\sqrt{K/n} = O((N/\log N)^{\rho/(1-2\rho+1/m)})$, we obtain

$$\begin{split} &\sqrt{\frac{n}{K}} \begin{bmatrix} A(x,z)^T & 0_{p+K+\xi}^T \\ 0_{p+K+\xi}^T & A(x,z)^T \end{bmatrix} \Sigma^{-1} \left\{ E \begin{bmatrix} \frac{\partial \ell(\theta_0)}{\partial \theta} \end{bmatrix} + \Omega_{\gamma,\sigma} \theta_0 \right\} \\ &\leq O \left(\sqrt{\frac{n}{K}} K^{1/2} \left\{ K^{-1/2} (N/n)^\rho + \lambda K^{m-1/2} + \nu K^{m-1/2} \right\} \right) \\ &= O(1/\log N) \\ &= o(1). \end{split}$$

Next, Lemma 10 yields that

$$\sqrt{\frac{n}{K}} \begin{bmatrix} A(x,z)^T & 0_{p+K+\xi}^T \\ 0_{p+K+\xi}^T & A(x,z)^T \end{bmatrix} \Sigma^{-1} \left\{ \frac{\partial \ell(\theta_0)}{\partial \theta} - E\left[\frac{\partial \ell(\theta_0)}{\partial \theta}\right] \right\} \xrightarrow{\mathcal{D}} N(0, \lim_{N \to \infty} D(x,z)^T \Sigma^{-1} D(x,z)/K).$$

Consequently, we obtain

$$\sqrt{\frac{n}{K}} \begin{bmatrix} \hat{\gamma}(x,z) - \gamma_0(x,z) \\ \log \hat{\sigma}(x,z) - \log \sigma_{w0}(x,z) \end{bmatrix} \xrightarrow{\mathcal{D}} N(0, \lim_{N \to \infty} D(x,z)^T \Sigma^{-1} D(x,z)/K).$$

Define $g(a) = e^a$ for any $a \in \mathbb{R}$. We then apply the delta method to $g(\log \hat{\sigma}(x, z)) - g(\log \sigma_{w0}(x, z))$, we obtain

$$\sqrt{\frac{n}{K}} \begin{bmatrix} \hat{\gamma}(x,z) - \gamma_0(x,z) \\ \frac{\hat{\sigma}(x,z)}{\sigma_{w0}(x,z)} - 1 \end{bmatrix} \xrightarrow{\mathcal{D}} N(0, \lim_{N \to \infty} D(x,z)^T \Sigma^{-1} D(x,z)/K),$$

which completes the proof.

Proof of Theorem 4. Let $\theta_P = (\beta^T, u^T)^T \in \mathbb{R}^{2p}$. When we use the Taylor expansion to $\partial \ell_{pen}(\beta, \hat{b}, u, \hat{c}) / \partial \theta_P$ around $(\hat{\beta}^T, \hat{u}^T)^T = (\beta_0^T, u_{w,0}^T)^T$, we have

$$0 = \frac{\partial \ell_{pen}(\hat{\beta}, \hat{b}, \hat{u}, \hat{c})}{\partial \theta_P} = \frac{\partial \ell_{pen}(\beta_0, \hat{b}, u_0, \hat{c})}{\partial \theta_P} + \left(\frac{\partial^2 \ell_{pen}(\beta^*, \hat{b}, u^*, \hat{c})}{\partial \theta_P \partial \theta_P^T}\right) \begin{bmatrix} \hat{\beta} - \beta_0 \\ \hat{u} - u_{w,0} \end{bmatrix}$$

where $((\beta^*)^T, (u^*)^T)^T$ is 2*p*-vector satisfying $||\beta^* - \beta_0||^2 + ||u^* - u_{w,0}||^2 < ||\hat{\beta} - \beta_0||^2 + ||\hat{u} - u_{w,0}||^2$. From Lemma 9, we have $||\hat{b} - b_0|| + ||\hat{c} - c_0|| \xrightarrow{P} 0$. Thefore, Lemma 7 implies that

$$\frac{\partial^2 \ell_{pen}(\beta^*, b, u^*, \hat{c})}{\partial \theta_P \partial \theta_P^T} = \Sigma_{\beta, u} (1 + o_P(1)).$$

Next, we evaluate

$$\frac{\partial \ell_{pen}(\beta_0, \hat{b}, u_{w,0}, \hat{c})}{\partial \theta_P} = \frac{\partial \ell_{pen}(\beta_0, b_0, u_{w,0}, c_0)}{\partial \theta_P} + \frac{\partial \ell_{pen}(\beta_0, \hat{b}, u_{w,0}, \hat{c})}{\partial \theta_P} - \frac{\partial \ell_{pen}(\beta_0, b_0, u_{w,0}, c_0)}{\partial \theta_P}$$

Similar to the proof of Lemmas 10, 6 and Theorem 3,

$$\sqrt{n} \left\{ \frac{\partial \ell_{pen}(\beta_0, b_0, u_{w,0}, c_0)}{\partial \theta_P} - E \left[\frac{\partial \ell_{pen}(\beta_0, b_0, u_{w,0}, c_0)}{\partial \theta_P} \right] \right\} \xrightarrow{D} N(0, \Sigma_{\beta, u})$$

and $E[\partial \ell_{pen}(\beta_0, b_0, u_{w,0}, c_0)/\partial \theta_P] = O((N/n)^{\rho})$. We next show that

$$R_{n}(\hat{b},\hat{c}) = \frac{\partial \ell_{pen}(\beta_{0},b,u_{w,0},\hat{c})}{\partial \theta_{p}} - \frac{\partial \ell_{pen}(\beta_{0},b_{0},u_{w,0},c_{0})}{\partial \theta_{p}} = O((N/n)^{\rho}) + o_{P}(n^{-1/2}).$$

Let $\gamma(x, z|b) = x^T \beta_0 + B(z)^T b$ for any $b \in \mathbb{R}^{d(K+\xi)}$ and $\sigma(x, z|c) = \exp[x^T u_{w,0} + B(z)^T c]$ for any $c \in \mathbb{R}^{d(K+\xi)}$. Note that $\gamma_0(x, z) = \gamma(x, z|b_0)$ and $\sigma_{w0}(x, z) = \sigma(x, z|c_0)$. We further let

$$\ell_{\gamma}(y|x,z,\hat{b},\hat{c}) = (\gamma(x,z|\hat{b})^{-1} + 1) \frac{y/\sigma(x,z|\hat{c})}{1 + y\gamma(x,z|\hat{b})/\sigma(x,z|\hat{c})} - \gamma(x,z|\hat{b})^{-2} \log\left(1 + \frac{y\gamma(x,z|\hat{b})}{\sigma(x,z|\hat{c})}\right)$$

and

$$\ell_{\sigma}(y|x, z, \hat{b}, \hat{c}) = 1 - (\gamma(x, z|\hat{b})^{-1} + 1) \frac{y\gamma(x, z|b)/\sigma(x, z|\hat{c})}{1 + y\gamma(x, z|\hat{b})/\sigma(x, z|\hat{c})}$$

Then, we have

$$R_n(\hat{b},\hat{c}) = \frac{1}{n} \sum_{i=1}^n \left[\begin{array}{c} \{\ell_\gamma(Y_i|X_i, Z_i, \hat{b}, \hat{c}) - \ell_\gamma(Y_i|X_i, Z_i, b_0, c_0)\}X_i \\ \{\ell_\sigma(Y_i|X_i, Z_i, \hat{b}, \hat{c}) - \ell_\sigma(Y_i|X_i, Z_i, b_0, c_0)\}X_i \end{array} \right].$$

Since $||\hat{b} - b_0|| + ||\hat{c} - c_0|| \xrightarrow{P} 0$ and $R_n(b,c)$ is continuous with respect to (b,c), the standard deviation of $R_n(\hat{b},\hat{c})$ is $o(n^{-1/2})$. The remain of proof is to show $E[R_n(\hat{b},\hat{c})] = O((N/n)^{\rho})$. We now focus on deriving

$$E[\{\ell_{\gamma}(Y_i|X_i, Z_i, \hat{b}, \hat{c}) - \ell_{\gamma}(Y_i|X_i, Z_i, b_0, c_0)\}X_i] = O((N/n)^{\rho}) + o(n^{-1/2}).$$

The Taylor expansion implies that

$$\ell_{\gamma}(Y_i|X_i, Z_i, \hat{b}, \hat{c}) - \ell_{\gamma}(Y_i|X_i, Z_i, b_0, c_0) = \frac{\partial \ell_{\gamma}(Y_i|X_i, Z_i, b^*, c^*)}{\partial b}(\hat{b} - b_0) + \frac{\partial \ell_{\gamma}(Y_i|X_i, Z_i, b^*, c^*)}{\partial c}(\hat{c} - c_0)$$

where $b^*, c^* \in \mathbb{R}^{d(K+\xi)}$ such that $||b^* - b_0|| + ||c^* - c_0|| \le ||\hat{b} - b_0|| + ||\hat{c} - c_0||$. Here, we obtain

$$\frac{\partial \ell_{\gamma}(Y_i|X_i, Z_i, b^*, c^*)}{\partial b} = \ell_{\gamma\gamma}(Y_i|X_i, Z_i, b, c^*)B(Z_i),$$

where

$$\ell_{\gamma\gamma}(y|x,z,b,c) = -2\gamma(x,z|b)^{-2} \frac{y/\sigma(x,z|c)}{1+y\gamma(x,z|b)/\sigma(x,z|c)} - (\gamma(x,z|b)^{-1}+1) \frac{-y^2/\sigma^2(x,z|c)}{\{1+y\gamma(x,z|b)/\sigma(x,z|c)\}^2} + 2\gamma(x,z|b)^{-3} \log\left(1+\frac{y\gamma(x,z|b)}{\sigma(x,z|c)}\right).$$

We note that $E[\ell_{\gamma\gamma}(Y_i|x, z, b_0, c_0)] = 2(2\gamma_0(x, z) + 1)^{-1}(\gamma_0(x, z) + 1)^{-1}(1 + o(1))$, which is related to hessian of $-\log h(Y_i|\gamma_0(x, z), \sigma_0(x, z))$ (see, the proof of Lemma 5). Similarly, we have

$$\frac{\partial \ell_{\gamma}(Y_i|X_i, Z_i, b^*, c^*)}{\partial c} = \ell_{\gamma\sigma}(Y_i|X_i, Z_i, b^*, c^*)B(Z_i),$$

where

$$\ell_{\gamma\sigma}(Y_i|X_i, Z_i, b, c) = \gamma(x, z|b)^{-2} \frac{y\gamma(x, z|b)/\sigma(x, z|c)}{1 + y\gamma(x, z|b)/\sigma(x, z|c)} - (\gamma(x, z|b)^{-1} + 1) \left\{ \frac{y/\sigma_0(x, z|c)}{1 + y\gamma(x, z|b)/\sigma(x, z|c)^2} - \frac{y^2\gamma(x, z|b)/\sigma(x, z|c)^2}{\{1 + y\gamma(x, z|b)/\sigma(x, z|c)^2\}^2} \right\}$$

and $E[\ell_{\gamma\sigma}(Y_i|x, z, b_0, c_0)] = (2\gamma_0(x, z) + 1)^{-1}(1 + o(1))$. We write

$$\Sigma^{-1} = \begin{bmatrix} \bar{\Sigma}_{\beta\beta} & \bar{\Sigma}_{\beta b} & \bar{\Sigma}_{\beta u} & \bar{\Sigma}_{\beta c} \\ \bar{\Sigma}_{b\beta} & \bar{\Sigma}_{bb} & \bar{\Sigma}_{bu} & \bar{\Sigma}_{bc} \\ \bar{\Sigma}_{u\beta} & \bar{\Sigma}_{ub} & \bar{\Sigma}_{uu} & \bar{\Sigma}_{uc} \\ \bar{\Sigma}_{c\beta} & \bar{\Sigma}_{cb} & \bar{\Sigma}_{cu} & \bar{\Sigma}_{cc} \end{bmatrix},$$

where the size of each block of Σ^{-1} are similar to the length of vector appeared in index: $\beta, u \in \mathbb{R}^p, b, c \in \mathbb{R}^{d(K+\xi)}$. Since

$$\begin{bmatrix} \hat{\beta} - \beta_0 \\ \hat{b} - b_0 \\ \hat{u} - u_0 \\ \hat{c} - c_0 \end{bmatrix} = \Sigma^{-1} \begin{bmatrix} \frac{\partial}{\partial \beta} \ell_{pen}(\theta_0) \\ \frac{\partial}{\partial b} \ell_{pen}(\theta_0) \\ \frac{\partial}{\partial u} \ell_{pen}(\theta_0) \\ \frac{\partial}{\partial c} \ell_{pen}(\theta_0) \end{bmatrix} (1 + o_P(1)),$$

we obtain

$$\begin{split} \hat{b} &- b_0 \\ &= \left[\bar{\Sigma}_{b\beta} \frac{\partial}{\partial \beta} \ell_{pen}(\theta_0) + \bar{\Sigma}_{bb} \frac{\partial}{\partial b} \ell_{pen}(\theta_0) + \bar{\Sigma}_{bu} \frac{\partial}{\partial u} \ell_{pen}(\theta_0) + \bar{\Sigma}_{bc} \frac{\partial}{\partial c} \ell_{pen}(\theta_0) \right] (1 + o_P(1)) \\ &= \frac{1}{n} \sum_{i=1}^n \left[\ell_{\gamma}(Y_i | X_i, Z_i) \{ \bar{\Sigma}_{b\beta} X_i + \bar{\Sigma}_{bb} B(Z_i) \} + \ell_{\sigma}(Y_i | X_i, Z_i) \{ \bar{\Sigma}_{bu} X_i + \bar{\Sigma}_{bc} B(Z_i) \} \right. \\ &\quad + \lambda K^{2m} \bar{\Sigma}_{bb} \Omega b_0 + \nu K^{2m} \bar{\Sigma}_{bc} \Omega c_0 \right] (1 + o_P(1)) \\ &= \frac{1}{n} \sum_{i=1}^n L_b(Y_i | X_i, Z_i) (1 + o_P(1)), \end{split}$$

where $\ell_{\gamma}(Y_i|X_i, Z_i) = \ell_{\gamma}(Y_i|X_i, Z_i, b_0, c_0)$ and $\ell_{\sigma}(Y_i|X_i, Z_i) = \ell_{\sigma}(Y_i|X_i, Z_i, b_0, c_0)$. The asymptotic property of L_b is obtained from property of the normalized *B*-splines, Lemmas 4 and 6. Although we omit the detail, simple calculation yields that

$$\begin{split} E\left[\frac{1}{n}\sum_{i=1}^{n} \{\ell_{\gamma}(Y_{i}|X_{i},Z_{i},\hat{b},\hat{c}) - \ell_{\gamma}(Y_{i}|X_{i},Z_{i},b_{0},c_{0})\}X_{i}\right] \\ &= \frac{1}{n^{2}}\sum_{i,j=1}^{n} E[\ell_{\gamma\gamma}(Y_{i}|X_{i},Z_{i},b^{*},c^{*})B(Z_{i})^{T}L_{b}(Y_{j}|X_{j},Z_{j})X_{i}] \\ &= \frac{1}{n}E[\ell_{\gamma\gamma}(Y_{i}|X_{i},Z_{i},b^{*},c^{*})B(Z_{i})^{T}L_{b}(Y_{i}|X_{i},Z_{i})X_{i}] \\ &\quad + \frac{1}{n^{2}}\sum_{i,j=1,i\neq j}^{n} E[\ell_{\gamma\gamma}(Y_{i}|X_{i},Z_{i},b^{*},c^{*})B(Z_{i})^{T}L_{b}(Y_{j}|X_{j},Z_{j})X_{i}] \\ &\leq O(n^{-1}) + O(K^{-m} + (N/n)^{\rho}) \\ &= O(K^{-m} + (N/n)^{\rho}). \end{split}$$

Similarly, we have

$$E\left[\frac{1}{n}\sum_{i=1}^{n} \{\ell_{\sigma}(Y_{i}|X_{i}, Z_{i}, \hat{b}, \hat{c}) - \ell_{\sigma}(Y_{i}|X_{i}, Z_{i}, b_{0}, c_{0})\}X_{i}\right] \le O(K^{-m} + (N/n)^{\rho}).$$

Under the condition that $K \ge O(n^{1/2m} \log n)$, (K^{-m}) is negrigible and we obtain $E[R(\hat{b}, \hat{c})] = O((N/n)^{\rho}) + o(n^{-1/2})$, which completes the proof of first assertion of Theorem 4. Second assertion can also be obtained under $\sqrt{n}(N/n)^{\rho} \to 0$.

Appendix D: Proof of theorem in Section 4

We describe the proofs of Theorems 5 and 6. However, these proofs is quite similar to those of Theorems 1, 2 and 3. Therefore, we only illustrate the outline of the proofs and the details are omitted.

Proof of Theorems 5 and 6. Let $\vartheta_0 = (\beta_0^T, b_0^T, (u_{w,0}^{\S})^T, (c_0^{\S})^T)^T$, where

$$\vartheta_0 = \underset{(\beta,b,u^{\S},c^{\S})}{\operatorname{argmin}} - E[h^{\S}(Y|\bar{\gamma}(X_i,Z_i),\bar{\varsigma}(X_i,Z_i))].$$

We further let $\hat{\vartheta} = (\hat{\beta}^T, \hat{b}^T, (\hat{u}^{\S})^T, (\hat{c}^{\S})^T)^T$. Similar to Lemma 8, we have $||\hat{\vartheta} - \vartheta_0|| \xrightarrow{P} 0$ as $N \to \infty$. The Taylor expansion implies that

$$\left(\frac{\partial^2 \ell_{pen}^{\S}(\vartheta^*)}{\partial \vartheta \partial \vartheta^T}\right) (\hat{\vartheta} - \vartheta_0) = \frac{\partial \ell_{pen}^{\S}(\vartheta_0)}{\partial \vartheta},$$

where ϑ^* is $2(p + K + \xi)$ -vector satisfying $||\vartheta^* - \vartheta_0|| \le ||\hat{\vartheta} - \vartheta_0||$. Here, for the random variable Y_* having the distribution function $H((\gamma + 1)y/\varsigma|\gamma)$, the straightforward calculation yields that

$$E\left[\begin{array}{cc} \frac{\partial^2 -\log h^{\S}(Y_*|\gamma,\varsigma)}{\partial\gamma^2} & \frac{\partial^2 -\log h^{\S}(Y_*|\gamma,\varsigma)}{\partial\gamma\partial\log\varsigma} \\ \frac{\partial^2 -\log h^{\S}(Y_*|\gamma,\varsigma)}{\partial\log\varsigma\partial\gamma} & \frac{\partial^2 -\log h^{\S}(Y_*|\gamma,\varsigma)}{\partial(\log\varsigma)^2} \end{array}\right] = \left[\begin{array}{cc} (\gamma+1)^{-1} & 0 \\ 0 & (2\gamma+1)^{-1} \end{array}\right].$$

This implies that

$$\left(\frac{\partial^2 \ell_{pen}^{\S}(\vartheta^*)}{\partial \vartheta \partial \vartheta^T}\right) = \Sigma^{\S}(1 + o_P(1)).$$

Thus, we have from the definition and non-singularity of Σ^{\S} that

$$\hat{\gamma}(x,z) - \gamma_0(x,z) = A(x,z)^T \Sigma_{\gamma}^{-1} \left[\begin{array}{c} \frac{\partial \ell_{pen}^{\S}(\vartheta_0)}{\partial \beta} \\ \frac{\partial \ell_{pen}^{\S}(\vartheta_0)}{\partial b} \end{array} \right] (1 + o_P(1)) + O(K^{-\zeta})$$

and

$$\log \hat{\varsigma}(x,z) - \log \varsigma_0(x,z) = A(x,z)^T \Sigma_{\varsigma}^{-1} \left[\begin{array}{c} \frac{\partial \ell_{pen}^{\S}(\vartheta_0)}{\partial u^{\S}} \\ \frac{\partial \ell_{pen}^{\S}(\vartheta_0)}{\partial c^{\S}} \end{array} \right] (1 + o_P(1)) + O(K^{-\zeta}),$$

where $\Sigma_{\gamma} = E[(\gamma_0(X,Z)+1)^{-2}A(X,Z)A(X,Z)^T] + \lambda K^{2m}\Omega$ and $\Sigma_{\varsigma} = E[(2\gamma_0(X,Z)+1)^{-1}A(X,Z)A(X,Z)^T] + \nu K^{2m}\Omega$. Thus, we can find that $\hat{\gamma}(x,z) - \gamma_0(x,z)$ does not depend on the term of penalty for reparametrized scale function and hence, the term including ν does not appeared. Similarly, λ does not affect to $\log \hat{\varsigma}(x,z) - \log \varsigma_0(x,z)$. The remains of the proof can be derived in the same mannar as the proofs of Theorems 1, 2, 3 and related Lemmas.

Acknowledgements

This research was financially supported by JSPS KAKENHI (Grant No. 22K11935, 23H03353).

References

- Beirlant, J. and Goegebeur, Y. (2004). Local polynomial maximum likelihood estimation for Pareto-type distributions. *Journal of Multivariate Analysis*. 89 97–118.
- [2] Beirlant, J., Goegebeur, Y., Segers, J., and Teugels, J. (2004). *Statistics of extremes: Theory and applications*. John Wiley & Sons.
- [3] Buja, A., Hastie, T. and Tibshirani, R. (1989). Linear smoothers and additive models. Annals of Statistics. 17 453–510.
- [4] Chavez-Demoulin,V., Embrechts,P. and Hofert,M. (2015). An extreme vakue approach for modeling operational risk losses depending on covariates. *Journal of Risk Insuarance*. 83 735—776.
- [5] Chavez-Demoulin, V. and Davison, A.C. (2005). Generalized additive modeling of sample extremes. *Journal of the Royal Statistical Society Series C.* 54 207–222.
- [6] Claeskens, G., Krivobokova, T. and Opsomer, J.D. (2009). Asymptotic properties of penalized spline estimators. *Biometrika*. 96 529–544.
- [7] Daouia, A., Gardes, L., and Girard, S. (2013). On kernel smoothing for extremal quantile regression. *Bernoulli* 19 2557–2589.
- [8] de Boor, C. (2001). A practical guide to splines. Springer. New York.

- [9] de Haan, L. and Ferreira, A. (2006). *Extreme value theory: An introduction*. New York: Springer-Verlag.
- [10] Drees, H., Ferreira, A. and de Haan, L. On maximum likelihood estimation of the extreme value index. Annals of Applied Probability. 14 1179–1201.
- [11] Green, P.J. and Silverman, B.W. (1994). Nonparametric Regression and Generalized Linear Models: A roughness penalty approach. Chapman & Hall/CRC. New York.
- [12] Hastie, T. and Tibshirani, R. (1990). Generalized additive models. Chapman & Hall/CRC. London.
- [13] Hijort,N.L. and Pollard,D. (2011). Asymptotics for minimisers of convex processes. arXiv::1107.3806
- [14] Lin,R., Leng,C. and You,J. (2022). Semiparametric tail index regression. Journal of Business & Economic Statistics. 40 82–95.
- [15] Liu,X., Wang,L. and Liang,H. (2011). Estimation and variable selection for semiparametric additive partial linear models. *Statistica Sinica*. **21** 1225–1248.
- [16] Liu,R., Yang,L. and Härdle,W.K. (2013). Oracally efficient two-step estimation of generalized additive model. *Journal of American Statistical Association*. 108 619–631.
- [17] Liu,R., Härdle,W.K., and Zhang,G. (2017). Statistical inference for generalized additive partially linear models. *Jornal od Multicariate Analysis.* 162 1–15.
- [18] Marx,B.D. and Eilers,P.H.C. (1998). Direct generalized additive modeling with penalized likelihood. *Computational Statistics & Data Analysis.* 28 193–209.
- [19] Mhalla,L., de Carvalho,M. and Chavez-Demoulin,V. (2019). Regression-type models for extreme dependence. Scandinavian Journal of Statistics. 46 1141–1167.
- [20] Mhalla,L., Opitz,T. and Chavez-Demoulin,V. (2019). Exceedance-based nonlinear regression of tail dependence. *Extremes.* 22 523--552.
- [21] Reiss, R.D. and Thomas, M. (2007). Statistical analysis of extreme values: with applications to insurance, finance, hydrology and other fields. Birkhaeuser
- [22] Smith, R.L. (1987). Estimating tails of probability distributions. The Annals of Statistics. 15 1174—1207.
- [23] van de Geer,S. (2000). Empirical Processes in M-Estimation. Cambridge University Press, Cambridge.
- [24] Vatter, T. and Chavez-Demoulin, V. (2015). Generalized additive models for conditional dependence structures. *Journal of Multivariate Analysis.* 141 147–167.
- [25] Wang,H., and Tsai, C. L. (2009). Tail Index Regression. Journal of the American Statistical Association. 104 1233–1240.
- [26] Wang,L. and Yang,L. (2007). Spline-backfitted kernel smoothing of nonlinear additive autoregression model. Annals of Statistics. 35 2474–2503.

- [27] Wood,S.N. (2017). Generalized additive models: An Introduction with R, Second Edition. Chapman & Hall/CRC. London.
- [28] Xiao,L. (2019). Asymptotic theory of penalized spline. Electronic Journal of Statistics. 13 747–794.
- [29] Yee, T.W. (2015). Vector generalized linear and additive models: With an implementation in R. Springer-Verlag. New-York.
- [30] Yee, T.W. and Stephenson, A.G. (2007). Vector generalized linear and additive extreme value models. *Extremes.* 10 1––19.
- [31] Youngman,B.D. (2019). Generalized additive models for exceedances of high threshilds with an application to return level estimation for U.S. wing gusts. *Journal of the American Statistical Association.* **114** 1865–1879.
- [32] Youngman,B.D. (2022). evgam: An R package for generalized additive extreme value models. Journal of Statistical Software. 103 1–26.
- [33] Zhou, C. (2009). Existence and consistency of the maximum likelihood estimator for the extreme value index. *Journal of Multivariate Analysis.* **100** 794–815.
- [34] Zhou,S., Shen,X., and Wolfe,D.A. (1998). Local asymptotics for regression splines and confidence regions. Annals of Statistics 26 1760–1782.