# Testing equality between two-sample dependence structure using Bernstein polynomials

Guanjie Lyu

Department of Mathematics and Statistics, University of Windsor, Canada

Mohamed Belalia\*

Department of Mathematics and Statistics, University of Windsor, Canada

December 19, 2023

#### Abstract

Tests for the equality of copulas between two samples, utilizing the empirical Bernstein copula process, are introduced and studied. Three statistics are proposed, and their asymptotic properties are established. Furthermore, the empirical Bernstein copula process is investigated in association with subsampling methods and the multiplier bootstrap. Through a simulation study, it is demonstrated that the Bernstein tests significantly outperform the tests based on the empirical copula.

Keywords: Empirical Bernstein copula; Empirical process; Multiplier bootstrap; Subsampling method

# 1 Introduction

Copulas play a crucial role in characterizing the complete dependence structure between random variables, making them indispensable in various fields such as statistics, finance, and actuarial science. They have gained widespread recognition and are considered as a fundamental tool. One noteworthy related topic is the testing of copula function equality between different groups, which has seen growing interest in recent years due to its practical implications, see Rémillard and Scaillet (2009), Bouzebda et al. (2011), Bouzebda and El Faouzi (2012), and Seo (2021). In particular, the identification of similarities in dependence structures holds significant relevance across diverse domains. For instance, Dupuis et al. (2009) have demonstrated that the dependence structures between credit default swaps and equity returns can differ significantly. Additionally, Szolgay et al. (2016) conducted an exploration into the homogeneity of dependence structures pertaining to flood peaks and their corresponding volumes, both within different flood types and across various catchments.

<sup>\*</sup>Corresponding author: Mohamed.Belalia @ uwindsor.ca

However, the underlying copula function is often unknown, necessitating the adoption of an appropriate parametric model or, more commonly, a nonparametric approach. Among nonparametric estimators, the empirical copula, developed by Rüschendorf (1976), stands as the most popular. A multitude of nonparametric tests, including the equality tests mentioned in the previous paragraph, have been built upon this approach in the literature. One major drawback of the empirical copula estimator is its discreteness, which violates the continuity of the copula function when the marginals are continuous. This limitation becomes more pronounced when the sample size is relatively small. To address this issue, several smooth estimators have been introduced. For instance, Hall and Neumeyer (2006) and Morettin et al. (2010) explored wavelet-smoothed empirical copula, while Fermanian et al. (2004), Chen and Huang (2007) and Omelka et al. (2009) investigated kernel-smoothed empirical copula. For a comprehensive overview of nonparametric estimation of copula and copula density, refer to Charpentier et al. (2007).

We employ the empirical Bernstein copula, a trending smooth estimator for the copula function, as our choice for conducting the equality test. This selection is motivated by two main factors. Firstly, estimation based on Bernstein polynomials is known to be asymptotically bias-free at boundary points, as compared to kernel-based methods that often suffer from excess bias near the boundaries (see, Sancetta and Satchell (2004), Janssen et al. (2012), Leblanc (2012), Belalia (2016), and Belalia et al. (2017)). A comprehensive discussion on boundary bias for kernel-based methods can be found in Chen and Huang (2007). Secondly, the empirical Bernstein copula is a polynomial and therefore possesses all partial derivatives, which are crucial for constructing our two resampling methods. Furthermore, as noted by Neumann et al. (2019), "in practice, it may not be most important which smoothing method to choose, while it is recommendable to smooth at all." The empirical Bernstein copula offers a favourable balance between simplicity and effectiveness when compared to other nonparametric estimators.

In the context of hypothesis testing relying on nonparametric estimators, standard Monte Carlo procedures often encounter challenges, as elaborated in Section 3. To ensure the generation of valid p-values, the widely accepted approach is the application of the multiplier bootstrap method, known for its effectiveness and promising performance. Expanding on this understanding, we introduce a multivariate Bernstein version of the multiplier bootstrap, building upon the foundational work of Lyu and Belalia (2023). However, it is worth noting that the Bernstein multiplier bootstrap, while advantageous in various aspects, can be computationally intensive and does not readily accommodate the popular empirical beta copula.

Recent studies by Beare and Seo (2020) and Seo (2021) have explored a randomization test strategy to obtain feasible p-values. Nevertheless, it is essential to recognize that this approach introduces a further computational burden compared to the multiplier bootstrap method. In response to these computational challenges, we propose an additional subsampling method for the empirical Bernstein copula process, with the potential to facilitate the generation of bootstraps for the empirical beta copula. Our simulation results indicate that these two procedures offer complementary advantages in addressing both computational and statistical aspects.

The remainder of the paper is organized as follows. Section 2 presents the proposed test statistics and examines their asymptotic behaviours. Section 3 and 4 focus on the deployment of the multiplier bootstrap and subsampling method for the empirical Bernstein copula process, respectively. In Section 5, a simulation study is conducted, implementing the multiplier bootstrap and subsampling method. The proofs are provided in the Appendix.

# 2 Testing Procedure

In this section, a testing procedure is presented both for independent and paired samples. Heuristically, it can be shown that the two cases can be treated similarly for testing two-sample homogeneity of the dependence structure.

## 2.1 Independent samples

Consider two independent samples of  $\mathbb{R}^d$ -valued (*d* is much less than the sample size) *i.i.d.* vectors, that is,  $(X_{11}, \ldots, X_{1d}), \ldots, (X_{n_11}, \ldots, X_{n_1d})$  with distribution function *F* associated with continuous margins  $F_1, \ldots, F_d$  and  $(Y_{11}, \ldots, Y_{1d}), \ldots, (Y_{n_21}, \ldots, Y_{n_2d})$  with distribution function *G* associated with continuous margins  $G_1, \ldots, G_d$ . Let *C*, *D* denote the underlying copula function of the two samples, by Sklar (1959), for any  $\boldsymbol{u} = (u_1, \ldots, u_d) \in [0, 1]^d$ ,

$$C(\boldsymbol{u}) = F(F_1^{-1}(u_1), \dots, F_d^{-1}(u_d)), \quad D(\boldsymbol{u}) = G(G_1^{-1}(u_1), \dots, G_d^{-1}(u_d)),$$

where  $F_{\ell}^{-1}(u_{\ell}) = \inf\{t \in \mathbb{R} : F_{\ell}(t) \ge u_{\ell}\}, \ell = 1, \dots, d$ . To determine whether the dependence structure of the two samples is identical, the corresponding hypothesis test is

$$\begin{cases} \mathscr{H}_{0}: C(\boldsymbol{u}) = D(\boldsymbol{u}) \quad \forall \boldsymbol{u} \in [0, 1]^{d} \\ & \text{versus} \\ \mathscr{H}_{1}: C(\boldsymbol{u}) \neq D(\boldsymbol{u}) \quad \exists \boldsymbol{u} \in [0, 1]^{d}. \end{cases}$$
(1)

Based on the two samples, the empirical copulas are defined as

$$C_{n_1}(\boldsymbol{u}) = \frac{1}{n_1} \sum_{i=1}^{n_1} \prod_{\ell=1}^d \mathbb{I}\left(\widehat{U}_{i\ell} \le u_\ell\right), \quad D_{n_2}(\boldsymbol{u}) = \frac{1}{n_2} \sum_{i=1}^{n_2} \prod_{\ell=1}^d \mathbb{I}\left(\widehat{V}_{i\ell} \le u_\ell\right),$$

where  $\mathbb{I}(\cdot)$  denotes the indicator function and  $\hat{U}_{i\ell}, \hat{V}_{i\ell}$  are the pseudo-observations with

$$\widehat{U}_{i\ell} = \frac{1}{n_1} \sum_{j=1}^{n_1} \mathbb{I}(X_{j\ell} \le X_{i\ell}), \quad \widehat{V}_{i\ell} = \frac{1}{n_2} \sum_{j=1}^{n_2} \mathbb{I}(Y_{j\ell} \le Y_{i\ell})$$

One can smooth empirical copulas by Bernstein polynomials, that is,

$$C_{n_1,m_1}(\boldsymbol{u}) = \frac{1}{n_1} \sum_{i=1}^{n_1} \left[ \sum_{k_1=0}^{m_1} \cdots \sum_{k_d=0}^{m_1} \prod_{\ell=1}^d \mathbb{I}\left(\hat{U}_{i\ell} \le k_\ell/m_1\right) P_{k_\ell,m_1}(u_\ell) \right],$$

$$D_{n_2,m_2}(\boldsymbol{u}) = \frac{1}{n_2} \sum_{i=1}^{n_2} \left[ \sum_{k_1=0}^{m_2} \cdots \sum_{k_d=0}^{m_2} \prod_{\ell=1}^d \mathbb{I}\left(\hat{V}_{i\ell} \le k_\ell/m_2\right) P_{k_\ell,m_2}(u_\ell) \right],$$
(2)

where for simplicity, considering the same Bernstein order (which is assumed to be dependent on the sample size) for smoothing each dimension. Note that, when the Bernstein order equals the sample size, the empirical Bernstein copula reduces to the empirical beta copula, see Segers et al. (2017) and Kiriliouk et al. (2021). Three statistics are proposed relying on Equation (2). Let  $\boldsymbol{m} = (m_1, m_2), \boldsymbol{n} = (n_1, n_2), \lambda_{\boldsymbol{n}} = n_1/(n_1 + n_2)$ . Define

$$R_{n}^{m} = n_{2}\lambda_{n}\int_{[0,1]^{d}} \left\{ C_{n_{1},m_{1}}(\boldsymbol{u}) - D_{n_{2},m_{2}}(\boldsymbol{u}) \right\}^{2} \mathrm{d}\boldsymbol{u},$$

$$S_{n}^{m} = n_{2}\lambda_{n}\int_{[0,1]^{d}} \left\{ C_{n_{1},m_{1}}(\boldsymbol{u}) - D_{n_{2},m_{2}}(\boldsymbol{u}) \right\}^{2} \mathrm{d}C_{n_{1},m_{1}}(\boldsymbol{u}),$$

$$T_{n}^{m} = \sqrt{n_{2}\lambda_{n}} \sup_{\boldsymbol{u}\in[0,1]^{d}} |C_{n_{1},m_{1}}(\boldsymbol{u}) - D_{n_{2},m_{2}}(\boldsymbol{u})|.$$
(3)

For the establishment of asymptotic behaviours of the proposed statistics, the following assumptions (Segers, 2012) are needed.

Assumption 1. Assume every partial derivative  $\dot{C}_{\ell}(\boldsymbol{u}) = \partial C(\boldsymbol{u}) / \partial u_{\ell}, \ell \in \{1, \ldots, d\}$  exists and is continuous on the set  $V_{d,\ell} = \{\boldsymbol{u} \in [0,1]^d : 0 < u_{\ell} < 1\}.$ 

Assumption 2. For every  $j, \ell \in \{1, ..., d\}$ , the second-order partial derivative  $\ddot{C}_{j\ell} = \partial^2 C(\boldsymbol{u})/\partial u_j u_\ell$  is defined and continuous, on the sets  $V_{d,j} \cap V_{d,\ell}$ , and there exists a constant K > 0 such that

$$\left|\ddot{C}_{j\ell}(\boldsymbol{u})\right| \leq K \min\left(\frac{1}{u_j(1-u_j)}, \frac{1}{u_\ell(1-u_\ell)}\right), \quad \boldsymbol{u} \in V_{d,j} \cap V_{d,\ell}.$$

Let  $\mathbb{C}_{n_1} = \sqrt{n_1} \{ C_{n_1}(\boldsymbol{u}) - C(\boldsymbol{u}) \}, \mathbb{D}_{n_2} = \sqrt{n_2} \{ D_{n_2}(\boldsymbol{u}) - D(\boldsymbol{u}) \}$  be the empirical copula processes of the two samples. Under Assumption 1 and  $m_1 = cn_1^{\alpha}$  with  $c > 0, \alpha \ge 1$ , the empirical Bernstein copula process (Segers et al., 2017) converges to a *d*-dimensional Brownian pillow, *i.e.*,

$$\mathbb{C}_{n_1,m_1}(\boldsymbol{u}) = \sqrt{n_1} \{ C_{n_1,m_1}(\boldsymbol{u}) - C(\boldsymbol{u}) \} \rightsquigarrow \mathbb{C}(\boldsymbol{u}) = \mathbb{B}_C(\boldsymbol{u}) - \sum_{\ell=1}^d \mathbb{B}_C(\boldsymbol{u}^\ell) \dot{C}_\ell(\boldsymbol{u}),$$
(4)

where  $u^{\ell} = (1, \ldots, 1, u_{\ell}, 1, \ldots, 1)$ , as all elements are replaced by 1 except the  $\ell$ -th component and  $\mathbb{B}_C$  is a Brownian bridge with covariance function

$$\mathbb{E}\left[\mathbb{B}_C(\boldsymbol{u})\mathbb{B}_C(\boldsymbol{v})\right] = C(\boldsymbol{u}\wedge\boldsymbol{v}) - C(\boldsymbol{u})C(\boldsymbol{v}), \qquad \boldsymbol{u}, \boldsymbol{v}\in[0,1]^d.$$

Similarly, under Assumption 1 and  $m_2 = cn_2^{\alpha}$  with  $c > 0, \alpha \ge 1$ ,

$$\mathbb{D}_{n_2,m_2}(\boldsymbol{u}) = \sqrt{n_2} \{ D_{n_2,m_2}(\boldsymbol{u}) - D(\boldsymbol{u}) \} \rightsquigarrow \mathbb{D}(\boldsymbol{u}) = \mathbb{B}_D(\boldsymbol{u}) - \sum_{\ell=1}^d \mathbb{B}_D(\boldsymbol{u}^\ell) \dot{D}_\ell(\boldsymbol{u}).$$

Because of the weak convergence of the empirical Bernstein copula process, the following lemma holds.

**Lemma 1.** Suppose that  $\lambda_n \to \lambda$  and  $m_r = cn_r^{\alpha}$  with  $c > 0, \alpha \ge 1$  for r = 1, 2. If Assumption 1 is satisfied, then, as  $\min(n_1, n_2) \to \infty$ ,

$$\mathbb{F}_{n}^{m} = \sqrt{1 - \lambda_{n}} \mathbb{C}_{n_{1}, m_{1}} - \sqrt{\lambda_{n}} \mathbb{D}_{n_{2}, m_{2}} \rightsquigarrow \mathbb{F} = \sqrt{1 - \lambda} \mathbb{C} - \sqrt{\lambda} \mathbb{D}.$$
(5)

*Proof.* Since the two samples are independent, clearly,

$$\left(\sqrt{1-\lambda_n}, \mathbb{C}_{n_1,m_1}, \sqrt{\lambda_n}, \mathbb{D}_{n_2,m_2}\right) \rightsquigarrow \left(\sqrt{1-\lambda}, \mathbb{C}, \sqrt{\lambda}, \mathbb{D}\right).$$

Applying the continuous mapping theorem, this yields the desired result.

With the help of Lemma 1, weak convergence of the proposed statistics can be established under the same conditions.

**Theorem 1.** Suppose that assumptions in Lemma 1 are satisfied. Then under the null hypothesis, as  $\min(n_1, n_2) \to \infty$ ,

$$R_{n}^{m} = \int_{[0,1]^{d}} \left\{ \mathbb{F}_{n}^{m}(\boldsymbol{u}) \right\}^{2} \mathrm{d}\boldsymbol{u} \rightsquigarrow R_{\mathbb{F}} = \int_{[0,1]^{d}} \{\mathbb{F}(\boldsymbol{u})\}^{2} \mathrm{d}\boldsymbol{u},$$

$$S_{n}^{m} = \int_{[0,1]^{d}} \left\{ \mathbb{F}_{n}^{m}(\boldsymbol{u}) \right\}^{2} \mathrm{d}C_{n_{1},m_{1}}(\boldsymbol{u}) \rightsquigarrow S_{\mathbb{F}} = \int_{[0,1]^{d}} \{\mathbb{F}(\boldsymbol{u})\}^{2} \mathrm{d}C(\boldsymbol{u}),$$

$$T_{n}^{m} = \sup_{\boldsymbol{u} \in [0,1]^{d}} |\mathbb{F}_{n}^{m}(\boldsymbol{u})| \rightsquigarrow T_{\mathbb{F}} = \sup_{\boldsymbol{u} \in [0,1]^{d}} |\mathbb{F}(\boldsymbol{u})|.$$
(6)

*Proof.* The proof is postponed to the Appendix.

More generally, the asymptotic behaviour of the test statistics under the alternative hypothesis are given as follows.

**Theorem 2.** Suppose that assumptions in Lemma 1 are satisfied. Then, as  $\min(n_1, n_2) \to \infty$ ,

$$n_2^{-1} R_n^m \xrightarrow{a.s.} R(C, D) = \lambda \int_{[0,1]^d} \left\{ C(\boldsymbol{u}) - D(\boldsymbol{u}) \right\}^2 d\boldsymbol{u}$$
$$n_2^{-1} S_n^m \xrightarrow{a.s.} S(C, D) = \lambda \int_{[0,1]^d} \left\{ C(\boldsymbol{u}) - D(\boldsymbol{u}) \right\}^2 dC(\boldsymbol{u}),$$
$$\sqrt{n_2^{-1}} T_n^m \xrightarrow{a.s.} T(C, D) = \sqrt{\lambda} \sup_{\boldsymbol{u} \in [0,1]^d} |C(\boldsymbol{u}) - D(\boldsymbol{u})|.$$

Specifically, under the alternative hypothesis, as  $\min(n_1, n_2) \to \infty$ ,

$$R_n^m \xrightarrow{a.s.} \infty, \quad S_n^m \xrightarrow{a.s.} \infty, \quad T_n^m \xrightarrow{a.s.} \infty,$$

which provides a guarantee of the consistency of the proposed test statistics.

**Remark 1.** Note that, as a consequence of the following Lemma 2, the convergences in Theorem 1-2 also hold under Assumptions 1-2 with a weaker condition on Bernstein order, that is,  $m_r = cn_r^{\alpha}$  with  $c > 0, \alpha > 3/4$  for r = 1, 2.

**Lemma 2.** Suppose that C satisfies Assumptions 1-2, additionally, if  $m_r = cn_r^{\alpha}$  with  $c > 0, \alpha > 3/4$  for r = 1, 2, then, in  $\ell^{\infty}([0, 1]^d)$ , Equation (4) holds.

*Proof.* The proof is postponed to the Appendix.

Comparing with Theorem 1, imposing more conditions, a strong approximation of the proposed statistics follows.

**Theorem 3.** Suppose that  $\lambda_n \to \lambda$  and  $m_r = cn_r^{\alpha}$  with  $c > 0, \alpha \ge 1$  for r = 1, 2 as  $\min(n_1, n_2) \to \infty$ . Additionally, Assumptions 1-2 are satisfied. On a suitable probability space, it is possible to define  $\mathbb{F}_n^m(\boldsymbol{u}), \boldsymbol{u} \in [0,1]^d$ , jointly with Gaussian processes  $\{\mathbb{F}_n^*(\boldsymbol{u})\}_{\boldsymbol{n}\in\mathbb{N}^+\times\mathbb{N}^+}$ , in such a way that, under the null hypothesis, as  $\min(n_1, n_2) \to \infty$ , we have

$$\begin{vmatrix} R_{n}^{m} - \int_{[0,1]^{d}} \{\mathbb{F}_{n}^{*}(\boldsymbol{u})\}^{2} \, \mathrm{d}\boldsymbol{u} \end{vmatrix} \stackrel{a.s.}{=} O\left( \max\left( (\log \log n_{1})^{1/2}, (\log \log n_{2})^{1/2} \right) \Psi(n_{1}, n_{2}) \right), \\ S_{n}^{m} - \int_{[0,1]^{d}} \{\mathbb{F}_{n}^{*}(\boldsymbol{u})\}^{2} \, \mathrm{d}C(\boldsymbol{u}) \end{vmatrix} \stackrel{a.s.}{=} O\left( \max\left( (\log \log n_{1})^{1/2}, (\log \log n_{2})^{1/2} \right) \Psi(n_{1}, n_{2}) \right), \\ \left| T_{n}^{m} - \sup_{\boldsymbol{u} \in [0,1]^{d}} |\mathbb{F}_{n}^{*}(\boldsymbol{u})| \right| \stackrel{a.s.}{=} O\left( \Psi(n_{1}, n_{2}) \right), \end{aligned}$$

where

$$\Psi(n_1, n_2) = \max\left(n_1^{-1/(2(2d-1))} \log n_1, n_2^{-1/(2(2d-1))} \log n_2\right).$$

and

$$\mathbb{F}_{\boldsymbol{n}}^{*}(\boldsymbol{u}) = \sqrt{1 - \lambda_{\boldsymbol{n}}} \mathbb{C}^{(n_{1})} + \sqrt{\lambda_{\boldsymbol{n}}} \mathbb{D}^{(n_{2})}$$

where  $\{\mathbb{C}^{(n_1)}\}_{n_1\in\mathbb{N}^+}$  (resp.  $\{\mathbb{D}^{(n_2)}\}_{n_2\in\mathbb{N}^+}$ ) are independent copies of  $\mathbb{C}$  (resp.  $\mathbb{D}$ ). Additionally,  $\{\mathbb{C}^{(n_1)}\}_{n_1\in\mathbb{N}^+}$  are also independent of  $\{\mathbb{D}^{(n_2)}\}_{n_2\in\mathbb{N}^+}$ .

*Proof.* The proof is postponed to the Appendix.

**Remark 2.** For the precise meaning of "suitable probability space", we refer to Bouzebda et al. (2011, Remark 2.1).

## 2.2 Paired samples

Sometimes, one may encounter paired samples. This subsection addresses a testing procedure in this scenario. Consider independent vectors  $\mathbf{Z}_1 = (X_{11}, \ldots, X_{d1}, Y_{11}, \ldots, Y_{d1}), \ldots, \mathbf{Z}_n = (X_{1n}, \ldots, X_{dn}, Y_{1n}, \ldots, Y_{dn})$ with copula function  $\mathcal{C}$  on  $[0, 1]^{2d}$ , satisfying the property that for any  $\mathbf{u}, \mathbf{v} \in [0, 1]^d$ ,

$$C(\boldsymbol{u}, \underbrace{1, \ldots, 1}_{d}) = C(\boldsymbol{u}), \quad C(\underbrace{1, \ldots, 1}_{d}, \boldsymbol{v}) = D(\boldsymbol{v}),$$

where C, D are copulas for X and Y, respectively. The objective is to investigate if C and D are equal. Define the empirical Bernstein copula for sample Z as follows. For  $\boldsymbol{u}, \boldsymbol{v} \in [0, 1]^d$ ,

$$\mathcal{C}_{n,m}(\boldsymbol{u},\boldsymbol{v}) = \frac{1}{n} \sum_{i=1}^{n} \left[ \sum_{k_1=0}^{m} \cdots \sum_{k_d=0}^{m} \sum_{\kappa_1=0}^{m} \cdots \sum_{\kappa_d=0}^{m} \prod_{\ell=1}^{d} \prod_{j=1}^{d} \mathbb{I}\left(\widehat{U}_{i\ell} \le k_\ell/m\right) \mathbb{I}\left(\widehat{V}_{ij} \le \kappa_j/m\right) P_{k_\ell,m}(u_\ell) P_{\kappa_j,m}(v_j) \right].$$

Further, under Assumption 1 and as  $n/m \to c > 0$ , let  $\boldsymbol{z} \in [0, 1]^{2d}$ . As  $n \to \infty$ ,

$$\mathfrak{C}_{n,m}(\boldsymbol{z}) = \sqrt{n} \{ \mathcal{C}_{n,m}(\boldsymbol{z}) - \mathcal{C}(\boldsymbol{z}) \} \rightsquigarrow \mathfrak{C}(\boldsymbol{z}) = \mathbb{B}_{\mathcal{C}}(\boldsymbol{z}) - \sum_{\ell=1}^{2d} \mathbb{B}_{\mathcal{C}}(\boldsymbol{z}^{\ell}) \dot{\mathcal{C}}_{\ell}(\boldsymbol{z}).$$

It is readily observed that

$$\mathcal{C}_{n,m}(\boldsymbol{u},1,\ldots,1) = C_{n,m}(\boldsymbol{u}), \quad \mathcal{C}_{n,m}(1,\ldots,1,\boldsymbol{v}) = D_{n,m}(\boldsymbol{v}), \tag{7}$$

where  $C_{n,m}(\boldsymbol{u}), D_{n,m}(\boldsymbol{v})$  are defined in Equation (2), *i.e.*, the empirical Bernstein copulas for X and Y, respectively. It immediately follows that, as  $n \to \infty$ ,

$$\mathbb{C}_{n,m}(\boldsymbol{u}) = \sqrt{n} \{ C_{n,m}(\boldsymbol{u}) - C(\boldsymbol{u}) \} \rightsquigarrow \mathbb{C}(\boldsymbol{u}), \quad \mathbb{D}_{n,m}(\boldsymbol{v}) = \sqrt{n} \{ D_{n,m}(\boldsymbol{u}) - D(\boldsymbol{u}) \} \rightsquigarrow \mathbb{D}(\boldsymbol{v}).$$

Hence, the paired samples case can be treated the same as the independent samples case by separating X and Y as in Equation (7), and all the procedures and results for independent samples are valid for paired samples as well. For this reason, the remainder of this article focuses on the independent samples case.

## 3 Multiplier bootstrap

The conventional Monte Carlo procedure is unsuitable for this scenario due to its reliance on the underlying copula C and D, thereby rendering it inadequate for accurately evaluating the limits of proposed test statistics. To address this limitation, a widely acknowledged approach in the literature is the multiplier bootstrap method, originally introduced by Scaillet (2005) and further advanced by Rémillard and Scaillet (2009). Prominently, this method has garnered significant attention and adoption by numerous researchers (Kojadinovic and Yan, 2011; Genest et al., 2012; Harder and Stadtmüller, 2017; Bahraoui et al., 2018) across diverse research contexts. The multiplier bootstrap technique offers notable benefits, including the provision of valid p-values and exceptional efficiency, particularly when applied to goodness-of-fit tests. Recently, Lyu and Belalia (2023) constructed multiplier bootstraps for the empirical Bernstein copula process dealing with bivariate copulas. Here, we will extend these results to the multivariate case. Let  $H \in \mathbb{N}^+$  and for each  $h \in \{1, \ldots, H\}$ , let  $\boldsymbol{\xi}_{n_1+n_2}^{(h)} \coloneqq \left(\boldsymbol{\xi}_1^{(h)}, \ldots, \boldsymbol{\xi}_{n_1+n_2}^{(h)}\right)$  be a vector of independent random variables with unit mean and unit variance (taking  $\boldsymbol{\xi}_i^{(h)} \sim \operatorname{Exp}(1), i = 1, \ldots, n_1 + n_2$ ). Denote the sample mean of the first  $n_1$  observations by  $\overline{\boldsymbol{\xi}}_{n_1}^{(h)}$ , and the sample mean of the remaining  $n_2$  observations by  $\overline{\boldsymbol{\xi}}_{n_2}^{(h)}$ . Set

$$G_{n_1,m_1}(\boldsymbol{u}) = \frac{1}{n_1} \sum_{i=1}^{n_1} \left[ \sum_{k_1=0}^{m_1} \cdots \sum_{k_d=0}^{m_1} \prod_{\ell=1}^d \mathbb{I} \left( U_{i\ell} \le k_\ell / m_1 \right) P_{k_\ell,m_1}(u_\ell) \right]$$
$$= \sum_{k_1=0}^{m_1} \cdots \sum_{k_d=0}^{m_1} G_{n_1} \left( \frac{k_1}{m_1}, \dots, \frac{k_d}{m_1} \right) \prod_{\ell=1}^d P_{k_\ell,m_1}(u_\ell),$$

where  $U_{i\ell} = F_{\ell}(X_i), \ell = 1, ..., d$ . For  $h \in \{1, ..., H\}$ , define

$$\mathbb{G}_{n_1,m_1}(\boldsymbol{u}) = \sqrt{n_1} \left\{ G_{n_1,m_1}(\boldsymbol{u}) - C(\boldsymbol{u}) \right\} = \frac{1}{\sqrt{n_1}} \sum_{i=1}^{n_1} \left[ \sum_{k_1=0}^{m_1} \cdots \sum_{k_d=0}^{m_1} \prod_{\ell=1}^d \mathbb{I} \left( U_{i\ell} \le k_\ell/m_1 \right) P_{k_\ell,m_1}(u_\ell) - C(\boldsymbol{u}) \right],$$

and

$$\mathbb{G}_{n_1,m_1}^{(h)}(\boldsymbol{u}) = \frac{1}{\sqrt{n_1}} \sum_{i=1}^{n_1} \left( \xi_i^{(h)} - \overline{\xi}_{n_1}^{(h)} \right) \left[ \sum_{k_1=0}^{m_1} \cdots \sum_{k_d=0}^{m_1} \prod_{\ell=1}^d \mathbb{I}\left( \widehat{U}_{i\ell} \le k_\ell/m_1 \right) P_{k_\ell,m_1}(u_\ell) \right].$$

Note that,  $\mathbb{G}_{n_1,m_1}$  is the empirical copula process when we have information about margins. Define  $G_{n_2}(\boldsymbol{u}), G_{n_2,m_2}(\boldsymbol{u}), \mathbb{G}_{n_2,m_2}(\boldsymbol{u}), \mathbb{G}_{n_2,m_2}(\boldsymbol{u})$  similarly. The following proposition states that  $\mathbb{G}_{n_1,m_1}(\boldsymbol{u}), \mathbb{G}_{n_2,m_2}(\boldsymbol{u})$  converge weakly to the Brownian bridge mentioned in preceding section. Moreover, the replicates are valid.

**Proposition 1.** Suppose that  $\lambda_n \to \lambda$  and  $m_r = cn_r^{\alpha}$  with  $c > 0, \alpha > 3/4$  for r = 1, 2 as  $\min(n_1, n_2) \to \infty$ . Additionally, Assumptions 1-2 are satisfied. Then for  $\boldsymbol{u} \in [0, 1]^d$ ,

1.

$$\mathbb{G}_{n_1,m_1}(\boldsymbol{u}) \rightsquigarrow \mathbb{B}_C(\boldsymbol{u}), \quad \mathbb{G}_{n_2,m_2}(\boldsymbol{u}) \rightsquigarrow \mathbb{B}_D(\boldsymbol{u}),$$

2. For each  $h \in \{1, ..., H\}$ ,

$$\mathbb{G}_{n_1,m_1}^{(h)}(oldsymbol{u}) \rightsquigarrow \mathbb{B}_C(oldsymbol{u}), \quad \mathbb{G}_{n_2,m_2}^{(h)}(oldsymbol{u}) \rightsquigarrow \mathbb{B}_D(oldsymbol{u}).$$

*Proof.* The proof is postponed to the Appendix.

Therefore, bootstrap replicates of  $\mathbb{C}_{n_1,m_1}$  for  $h \in \{1,\ldots,H\}$  can be defined as

$$\mathbb{C}_{n_1,m_1}^{(h)}(\boldsymbol{u}) = \mathbb{G}_{n_1,m_1}^{(h)}(\boldsymbol{u}) - \sum_{\ell=1}^d \mathbb{G}_{n_1,m_1}^{(h)}(\boldsymbol{u}^\ell) \, \partial C_{n_1,m_1}(\boldsymbol{u}) / \partial u_\ell,$$

where

$$\partial C_{n_1,m_1}(\boldsymbol{u})/\partial u_{\ell} = m_1 \sum_{k_1=0}^{m_1} \cdots \sum_{k_{\ell}=0}^{m_1-1} \cdots \sum_{k_d=0}^{m_1} \left\{ C_{n_1}\left(\frac{k_1}{m_1}, \dots, \frac{k_{\ell}+1}{m_1}, \dots, \frac{k_d}{m_1}\right) \right\}$$

$$-C_{n_1}\left(\frac{k_1}{m_1},\ldots,\frac{k_{\ell}}{m_1},\ldots,\frac{k_d}{m_1}\right)\right\}P_{k_1,m_1}(u_1)\cdots P_{k_{\ell},m_1-1}(u_{\ell})\cdots P_{k_d,m_1}(u_d).$$

Similarly, the replicates of  $\mathbb{D}_{n_2,m_2}$  are defined as

$$\mathbb{D}_{n_2,m_2}^{(h)}(\boldsymbol{u}) = \mathbb{G}_{n_2,m_2}^{(h)}(\boldsymbol{u}) - \sum_{\ell=1}^d \mathbb{G}_{n_2,m_2}^{(h)}(\boldsymbol{u}^\ell) \,\partial D_{n_2,m_2}(\boldsymbol{u}) / \partial u_\ell.$$

The partial derivatives of the empirical Bernstein copula for bivariate case were studied in Janssen et al. (2016). Unlike the empirical copula, these empirical Bernstein copula partial derivatives can be calculated directly without any further estimation. The following proposition shows the uniform consistency of these partial derivatives.

**Proposition 2.** Suppose that  $\lambda_n \to \lambda$  and  $m_r = cn_r^{\alpha}$  with  $c > 0, \alpha > 3/4$  for r = 1, 2 as  $\min(n_1, n_2) \to \infty$ . Additionally, Assumptions 1-2 are satisfied. Then for any  $\ell \in \{1, \ldots, d\}$  and  $b_{n_r} = \kappa n_r^{-\beta}$  with  $\kappa > 0, \beta < \alpha/2$ , as  $\min(n_1, n_2) \to \infty$ ,

$$\sup_{\substack{\{\boldsymbol{u}\in[0,1]^d:u_\ell\in[b_{n_1},1-b_{n_1}]\}}} \left|\partial C_{n_1,m_1}(\boldsymbol{u})/\partial u_\ell - \dot{C}_\ell(\boldsymbol{u})\right| \stackrel{a.s.}{=} O\left(m_1^{1/2}n_1^{-1/2}(\log\log n_1)^{1/2}\right),$$
$$\sup_{\substack{\{\boldsymbol{u}\in[0,1]^d:u_\ell\in[b_{n_2},1-b_{n_2}]\}}} \left|\partial D_{n_2,m_2}(\boldsymbol{u})/\partial u_\ell - \dot{D}_\ell(\boldsymbol{u})\right| \stackrel{a.s.}{=} O\left(m_2^{1/2}n_2^{-1/2}(\log\log n_2)^{1/2}\right).$$

*Proof.* The proof is postponed to the Appendix.

Combining Propositions 1-2, for  $\boldsymbol{u} \in (0,1)^d$ ,  $\mathbb{C}_{n_1,m_1}^{(h)}(\boldsymbol{u})$ ,  $\mathbb{D}_{n_2,m_2}^{(h)}(\boldsymbol{u})$  are valid replicates for  $\mathbb{C}_{n_1,m_1}(\boldsymbol{u})$ ,  $\mathbb{D}_{n_2,m_2}(\boldsymbol{u})$ . Further, plugging  $\mathbb{C}_{n_1,m_1}^{(h)}(\boldsymbol{u})$ ,  $\mathbb{D}_{n_2,m_2}^{(h)}(\boldsymbol{u})$  into (5), under some conditions (see conditions in Theorem 4), one has

$$\left(\mathbb{F}_{n}^{m},\mathbb{F}_{n}^{m,(1)},\ldots,\mathbb{F}_{n}^{m,(H)}\right) \rightsquigarrow \left(\mathbb{F},\mathbb{F}^{(1)},\ldots,\mathbb{F}^{(H)}\right),$$

where  $\mathbb{F}^{(1)}, \ldots, \mathbb{F}^{(H)}$  are independent copies of  $\mathbb{F}$ . Therefore, applying Theorem 1, the asymptotic properties of replicates of proposed statistics are established in the following theorem.

**Theorem 4.** Suppose that  $\lambda_n \to \lambda$  and  $m_r = cn_r^{\alpha}$  with  $c > 0, 3/4 < \alpha < 1$  for r = 1, 2 as  $\min(n_1, n_2) \to \infty$ .  $\infty$ . Additionally, Assumptions 1-2 are satisfied. Then under the null hypothesis, as  $\min(n_1, n_2) \to \infty$ , for all  $H \in \mathbb{N}^+$ ,

$$\begin{pmatrix} R_n^m, R_n^{m,(1)}, \dots, R_n^{m,(H)} \end{pmatrix} \rightsquigarrow \begin{pmatrix} \mathbb{F}_R, \mathbb{F}_R^{(1)}, \dots, \mathbb{F}_R^{(H)} \end{pmatrix}, \begin{pmatrix} S_n^m, S_n^{m,(1)}, \dots, S_n^{m,(H)} \end{pmatrix} \rightsquigarrow \begin{pmatrix} \mathbb{F}_S, \mathbb{F}_S^{(1)}, \dots, \mathbb{F}_S^{(H)} \end{pmatrix}, \begin{pmatrix} T_n^m, T_n^{m,(1)}, \dots, T_n^{m,(H)} \end{pmatrix} \rightsquigarrow \begin{pmatrix} \mathbb{F}_T, \mathbb{F}_T^{(1)}, \dots, \mathbb{F}_T^{(H)} \end{pmatrix}.$$

Clearly, p-values of the statistics can be computed as

$$\frac{1}{H}\sum_{h=1}^{H}\mathbb{I}\left(R_{n}^{\boldsymbol{m},(h)}\right), \quad \frac{1}{H}\sum_{h=1}^{H}\mathbb{I}\left(S_{n}^{\boldsymbol{m},(h)}\right), \quad \frac{1}{H}\sum_{h=1}^{H}\mathbb{I}\left(T_{n}^{\boldsymbol{m},(h)}\right)$$

It is worth noting that the conditions imposed on the Bernstein order, namely  $m_r = cn_r^{\alpha}$  with  $c > 0, 3/4 < \alpha < 1$  for r = 1, 2 as  $\min(n_1, n_2) \to \infty$ . This particular condition renders the application of the proposed multiplier bootstrap method infeasible for empirical beta copula for which  $m_r = n_r$ . In the following section, an alternative resampling technique will be introduced that empowers interested readers to explore its potential application with empirical beta copula.

# 4 Subsampling method

An alternative to the bootstrap multiplier method, which is easier to implement, is the subsampling technique developed by Kojadinovic and Stemikovskaya (2019). This approach can be flexibly adapted to various smooth and weighted empirical copula processes, as demonstrated in the original paper. Unlike the multiplier method, it does not require the estimation of partial derivatives. Therefore, this method is valid for the empirical beta copula. In the subsequent section, we demonstrate the feasibility of applying this subsampling method to the empirical Bernstein copula process. Notably, this procedure offers comparable performance to the multiplier method while demanding significantly fewer computational resources and relying on fewer theoretical assumptions.

Sample  $b_1 < n_1, b_2 < n_2$  observations from X and Y without replacement, respectively. In this way, the number of possible subsamples are  $N_{b_1,n_1} = \binom{n_1}{b_1}, N_{b_2,n_2} = \binom{n_2}{b_2}$ . Let  $I_h \coloneqq (h_1, h_2)$  with  $h_1 \in \{1, \ldots, N_{b_1,n_1}\}, h_2 \in \{1, \ldots, N_{b_2,n_2}\}$ . If  $m_{(r)}, r = 1, 2$  are the Bernstein orders for subsamples of X and Y respectively, then define

$$C_{b_{1},m_{(1)}}^{(I_{h})}(\boldsymbol{u}) = \frac{1}{b_{1}} \sum_{i=1}^{b_{1}} \left[ \sum_{k_{1}=0}^{m_{(1)}} \cdots \sum_{k_{d}=0}^{m_{(1)}} \prod_{\ell=1}^{d} \mathbb{I}\left(\widehat{U}_{i\ell} \le k_{\ell}/m_{(1)}\right) P_{k_{\ell},m_{(1)}}(u_{\ell}) \right],$$
$$D_{b_{2},m_{(2)}}^{(I_{h})}(\boldsymbol{u}) = \frac{1}{b_{2}} \sum_{i=1}^{b_{2}} \left[ \sum_{k_{1}=0}^{m_{(2)}} \cdots \sum_{k_{d}=0}^{m_{(2)}} \prod_{\ell=1}^{d} \mathbb{I}\left(\widehat{V}_{i\ell} \le k_{\ell}/m_{(2)}\right) P_{k_{\ell},m_{(2)}}(u_{\ell}) \right],$$

as the empirical Bernstein copulas for the subsample h. The corrected subsample replicates of the empirical Bernstein copula process can be defined according to these subsample replicates,

$$\mathbb{C}_{b_1,m_{(1)}}^{(I_h)}(\boldsymbol{u}) = \sqrt{\frac{b_1}{1 - b_1/n_1}} \{ C_{b_1,m_{(1)}}^{(I_h)}(\boldsymbol{u}) - C_{n_1,m_1}(\boldsymbol{u}) \}, \quad \mathbb{D}_{b_2,m_{(2)}}^{(I_h)}(\boldsymbol{u}) = \sqrt{\frac{b_2}{1 - b_2/n_2}} \{ D_{b_2,m_{(2)}}^{(I_h)}(\boldsymbol{u}) - D_{n_2,m_2}(\boldsymbol{u}) \}.$$

The validity of these replicates is provided in the following theorem.

**Theorem 5.** Suppose that assumptions in Lemma 1 are satisfied, and that  $b_1, b_2$  go to infinity as  $\min(n_1, n_2) \rightarrow \infty$ . Let  $I_1, \ldots, I_H$  be independent bivariate random vectors with associated component being independent of X, Y and uniformly distributed on the set  $\{1, \ldots, N_{b_1, n_1}\}, \{1, \ldots, N_{b_2, n_2}\}$ . Then

$$\left( \mathbb{C}_{n_1,m_1}, \mathbb{C}_{b_1,m_{(1)}}^{(I_1)}, \dots, \mathbb{C}_{b_1,m_{(1)}}^{(I_H)} \right) \rightsquigarrow \left( \mathbb{C}, \mathbb{C}^{(1)}, \dots, \mathbb{C}^{(H)} \right),$$
$$\left( \mathbb{D}_{n_2,m_2}, \mathbb{D}_{b_2,m_{(2)}}^{(I_1)}, \dots, \mathbb{D}_{b_2,m_{(2)}}^{(I_H)} \right) \rightsquigarrow \left( \mathbb{D}, \mathbb{D}^{(1)}, \dots, \mathbb{D}^{(H)} \right).$$

*Proof.* The proof is postponed to the Appendix.

Plug 
$$\mathbb{C}_{b_1,m_{(1)}}^{(I_h)}(\boldsymbol{u}), \mathbb{D}_{b_2,m_{(2)}}^{(I_h)}(\boldsymbol{u})$$
 into (5), and let  $\tilde{m} = (m_{(1)}, m_{(2)})$ . Then  
 $\left(\mathbb{F}_{b_1,b_2}^{\tilde{m}}, \mathbb{F}_{b_1,b_2}^{\tilde{m},(I_1)}, \dots, \mathbb{F}_{b_1,b_2}^{\tilde{m},(I_H)}\right) \rightsquigarrow \left(\mathbb{F}, \mathbb{F}^{(I_1)}, \dots, \mathbb{F}^{(I_H)}\right),$ 

where  $\mathbb{F}^{(I_1)}, \ldots, \mathbb{F}^{(I_H)}$  are independent copies of  $\mathbb{F}$ . Clearly, using these replicates one can readily follow the strategies in Section 3 to construct valid replicates of the three proposed statistics. We skip them to reduce repetition.

## 5 Simulation study

Finite sample performance is investigated in this section. All tests were conducted with 500 repetitions under 5% nominal level using H = 200 multiplier replicates. Also given that the statistics shown in (3) involve integration, a discrete approximation is applied with a grid spanning a range of  $20 \times \cdots \times 20$  points on  $[0, 1]^d$ . To show the improvement of empirical Bernstein copula tests, empirical level and power of the proposed tests were compared with tests in Rémillard and Scaillet (2009).

The selection of an optimal Bernstein order is important and left for future study. According to the recommendation of Segers et al. (2017), we set  $m_r = \lfloor n_r/5 \rfloor$ , r = 1, 2, where  $\lfloor a \rfloor$  denote the largest integer less than or equal to a. And for the subsample sizes of the subsampling method, we follow the recommendation in Kojadinovic and Stemikovskaya (2019) with  $b_r = \lfloor 0.28 \times n_r \rfloor$  and associated Bernstein order  $m_{(r)} = b_r$  for convenience.

In Table 1 and 2, it is evident that all the tests effectively maintain their nominal level. Notably, the proposed Bernstein test is particularly preferable, given that the Crámer-von Mises type statistics consistently exhibit the highest power (indicated by bold font). When considering the Clayton copula from the Archimedean family,  $S_{n,mul}^m$  outperforms other statistics across most cases. Conversely, for the Gaussian copula from the elliptical family,  $R_{n,mul}^m$  generally outperforms the other statistics. In the case of 3-dimensional scenarios, the superiority of the Bernstein tests slightly diminishes, but they still demonstrate an advantage over empirical copula-based tests, especially for the Gaussian copula models.

When comparing the two resampling methods, it is important to consider their respective trade-offs. The subsampling method exhibits slightly inferior performance compared to the multiplier method but offers the advantage of reduced computational burden. This finding is consistent with the conclusions drawn by Kojadinovic and Stemikovskaya (2019). In summary, for small or moderate sample sizes, we recommend utilizing the multiplier method due to its superior performance. However, for large sample sizes, the subsampling method serves as a suitable alternative, striking a balance between computational difficulty and performance.

| Model                     | τ   | $R_{n,mul}$ | $R^{m}_{n,mul}$ | $R^{m}_{n,sub}$ | $S_{\boldsymbol{n},mul}$ | $S^{m}_{n,mul}$ | $S^{m{m}}_{m{n},sub}$ | $T_{\boldsymbol{n},mul}$ | $T^{\boldsymbol{m}}_{\boldsymbol{n},mul}$ | $T^{\boldsymbol{m}}_{\boldsymbol{n},sub}$ |
|---------------------------|-----|-------------|-----------------|-----------------|--------------------------|-----------------|-----------------------|--------------------------|---|---|
| $(n_1, n_2) = (50, 50)$   | 0.2 |             |                 |                 |                          |                 |                       |                          |   |   |
|                           | 0.2 | 2.6         | 5.6             | 3.4             | 2.4                      | 5.6             | 3.4                   | 1.0                      | 5.0                                       | 3.2                                       |
|                           | 0.3 | 3.4         | 8.8             | 6.4             | 2.2                      | 9.0             | 5.8                   | 1.6                      | 6.8                                       | 4.8                                       |
|                           | 0.4 | 16.0        | 26.2            | 20.4            | 9.8                      | 27.8            | 20.4                  | 7.6                      | 21.0                                      | 12.4                                      |
| Clayton                   | 0.5 | 40.8        | 60.6            | 51.2            | 32.0                     | 61.2            | 51.0                  | 24.6                     | 51.2                                      | 43.0                                      |
|                           | 0.6 | 72.0        | 86.6            | 81.4            | 63.4                     | 87.2            | 80.8                  | 43.8                     | 77.8                                      | 69.4                                      |
|                           | 0.7 | 88.6        | 97.6            | 95.4            | 83.6                     | 97.4            | 95.0                  | 67.4                     | 92.8                                      | 89.0                                      |
|                           | 0.8 | 99.0        | 99.8            | 99.4            | 97.6                     | 99.8            | 99.8                  | 90.6                     | 99.2                                      | 98.8                                      |
| $(n_1, n_2) = (100, 50)$  | 0.2 |             |                 |                 |                          |                 |                       |                          |   |   |
|                           | 0.2 | 0.8         | 5.0             | 2.0             | 1.2                      | 4.6             | 1.8                   | 0.8                      | 3.2                                       | 0.8                                       |
|                           | 0.3 | 5.0         | 8.2             | 6.0             | 4.0                      | 8.0             | 5.2                   | 4.0                      | 6.6                                       | 4.6                                       |
|                           | 0.4 | 22.2        | 26.4            | 24.0            | 19.8                     | 27.2            | 24.2                  | 11.4                     | 20.8                                      | 16.0                                      |
| Clayton                   | 0.5 | 56.6        | 60.0            | 56.8            | 51.4                     | 62.2            | 55.2                  | 33.8                     | 46.8                                      | 41.8                                      |
|                           | 0.6 | 84.4        | 89.0            | 85.8            | 83.6                     | 89.6            | 86.2                  | 66.2                     | 82.0                                      | 76.8                                      |
|                           | 0.7 | 98.4        | 99.2            | 99.2            | 98.2                     | 99.4            | 99.2                  | 91.4                     | 97.0                                      | 96.4                                      |
|                           | 0.8 | 99.8        | 100.0           | 100.0           | 99.8                     | 100.0           | 100.0                 | 99.0                     | 100.0                                     | 99.6                                      |
| $(n_1, n_2) = (150, 100)$ | 0.2 |             |                 |                 |                          |                 |                       |                          |   |   |
|                           | 0.2 | 2.0         | 2.8             | 2.6             | 1.8                      | 3.4             | 2.2                   | 1.2                      | 2.6                                       | 1.2                                       |
|                           | 0.3 | 15.2        | 18.0            | 15.2            | 11.8                     | 17.0            | 14.2                  | 9.2                      | 13.6                                      | 11.0                                      |
|                           | 0.4 | 52.6        | 57.4            | 54.2            | 48.6                     | <b>59.4</b>     | 54.8                  | 33.0                     | 45.4                                      | 39.6                                      |
| Clayton                   | 0.5 | 92.4        | 94.2            | 92.4            | 89.8                     | 94.6            | 93.2                  | 72.2                     | 88.0                                      | 84.2                                      |
|                           | 0.6 | 99.2        | 99.8            | 99.4            | 98.6                     | 99.8            | 99.4                  | 94.8                     | 98.8                                      | 98.0                                      |
|                           | 0.7 | 100.0       | 100.0           | 100.0           | 100.0                    | 100.0           | 100.0                 | 99.8                     | 100.0                                     | 100.0                                     |
|                           | 0.8 | 100.0       | 100.0           | 100.0           | 100.0                    | 100.0           | 100.0                 | 100.0                    | 100.0                                     | 100.0                                     |

Table 1: Empirical level (%) and power (%) of the tests are estimated from 500 samples from 2-dimensional Clayton copula given various Kendall's tau with different sample sizes.

| Model                     | au  | $R_{n,mul}$ | $R^{\boldsymbol{m}}_{\boldsymbol{n},mul}$ | $R^{m}_{n,sub}$ | $S_{\boldsymbol{n},mul}$ | $S^{\boldsymbol{m}}_{\boldsymbol{n},mul}$ | $S^{m}_{n,sub}$ | $T_{\boldsymbol{n},mul}$ | $T^{\boldsymbol{m}}_{\boldsymbol{n},mul}$ | $T^{\boldsymbol{m}}_{\boldsymbol{n},sub}$ |
|---------------------------|-----|-------------|---|-----------------|--------------------------|---|-----------------|--------------------------|---|---|
| $(n_1, n_2) = (50, 50)$   | 0.2 |             |   |                 |                          |   |                 |                          |   |   |
|                           | 0.2 | 1.8         | 4.6                                       | 3.4             | 2.6                      | 4.4                                       | 3.2             | 1.2                      | 4.2                                       | 1.8                                       |
|                           | 0.3 | 3.8         | 9.8                                       | 6.2             | 3.0                      | 10.2                                      | 5.8             | 1.6                      | 7.0                                       | 3.6                                       |
|                           | 0.4 | 14.2        | <b>26.6</b>                               | 17.4            | 8.4                      | 25.8                                      | 17.8            | 5.0                      | 18.2                                      | 11.6                                      |
| Gaussian                  | 0.5 | 40.4        | <b>60.4</b>                               | 52.4            | 30.0                     | 60.2                                      | 50.8            | 17.4                     | 45.4                                      | 35.6                                      |
|                           | 0.6 | 69.2        | 86.8                                      | 81.0            | 59.2                     | 86.2                                      | 81.0            | 41.6                     | 74.0                                      | 67.0                                      |
|                           | 0.7 | 93.4        | 99.0                                      | 97.6            | 90.2                     | 99.0                                      | 98.4            | 68.0                     | 95.0                                      | 91.6                                      |
|                           | 0.8 | 98.4        | 100.0                                     | 100.0           | 98.2                     | 100.0                                     | 99.8            | 87.2                     | 99.8                                      | 99.4                                      |
| $(n_1, n_2) = (100, 50)$  | 0.2 |             |   |                 |                          |   |                 |                          |   |   |
|                           | 0.2 | 2.0         | 5.6                                       | 3.2             | 2.2                      | 5.0                                       | 4.2             | 0.8                      | 3.4                                       | 1.4                                       |
|                           | 0.3 | 5.6         | 8.4                                       | 5.8             | 4.4                      | 7.8                                       | 5.4             | 3.0                      | 5.8                                       | 2.8                                       |
|                           | 0.4 | 23.6        | <b>30.4</b>                               | 24.4            | 21.6                     | 28.6                                      | 23.2            | 11.2                     | 19.8                                      | 12.8                                      |
| Gaussian                  | 0.5 | 60.4        | 66.4                                      | 61.6            | 56.0                     | 68.0                                      | 61.2            | 29.8                     | 50.4                                      | 43.4                                      |
|                           | 0.6 | 89.6        | 92.4                                      | 89.8            | 87.0                     | <b>92.4</b>                               | 89.6            | 62.2                     | 82.0                                      | 76.6                                      |
|                           | 0.7 | 97.4        | 99.6                                      | 99.8            | 96.8                     | 99.8                                      | 99.0            | 87.2                     | 96.6                                      | 94.4                                      |
|                           | 0.8 | 100.0       | 100.0                                     | 100.0           | 99.8                     | 100.0                                     | 100.0           | 97.8                     | 99.6                                      | 99.8                                      |
| $(n_1, n_2) = (150, 100)$ | 0.2 |             |   |                 |                          |   |                 |                          |   |   |
|                           | 0.2 | 2.4         | 5.2                                       | 3.8             | 3.2                      | 5.6                                       | 3.6             | 1.4                      | 3.8                                       | 2.6                                       |
|                           | 0.3 | 13.8        | 18.0                                      | 14.2            | 12.8                     | 17.8                                      | 15.2            | 7.6                      | 11.6                                      | 10.4                                      |
|                           | 0.4 | 50.6        | 57.4                                      | 53.6            | 48.2                     | 58.4                                      | 54.2            | 29.6                     | 44.2                                      | 39.0                                      |
| Gaussian                  | 0.5 | 89.4        | 92.6                                      | 91.0            | 85.8                     | 92.0                                      | 91.2            | 66.4                     | 81.4                                      | 77.6                                      |
|                           | 0.6 | 99.4        | 99.8                                      | 99.6            | 99.6                     | 99.6                                      | 99.6            | 92.8                     | 98.6                                      | 98.0                                      |
|                           | 0.7 | 100.0       | 100.0                                     | 100.0           | 100.0                    | 100.0                                     | 100.0           | 99.8                     | 100.0                                     | 100.0                                     |
|                           | 0.8 | 100.0       | 100.0                                     | 100.0           | 100.0                    | 100.0                                     | 100.0           | 100.0                    | 100.0                                     | 100.0                                     |

Table 2: Empirical level (%) and power (%) of the tests are estimated from 500 samples of 2-dimensional Gaussian copula given various Kendall's tau with different sample sizes.

| Model                   | θ    | $R_{n,mul}$ | $R^{\boldsymbol{m}}_{\boldsymbol{n},mul}$ | $R^{m}_{n,sub}$ | $S_{\boldsymbol{n},mul}$ | $S^{m}_{n,mul}$ | $S^{\boldsymbol{m}}_{\boldsymbol{n},sub}$ | $T_{\boldsymbol{n},mul}$ | $T^{\boldsymbol{m}}_{\boldsymbol{n},mul}$ | $T^{\boldsymbol{m}}_{\boldsymbol{n},sub}$ |
|-------------------------|------|-------------|---|-----------------|--------------------------|-----------------|---|--------------------------|---|---|
| $(n_1, n_2) = (50, 50)$ | 1.00 |             |   |                 |                          |                 |   |                          |   |   |
|                         | 1.00 | 3.6         | 2.6                                       | 1.8             | 2.6                      | 2.8             | 1.6                                       | 2.2                      | 2.0                                       | 0.4                                       |
|                         | 1.25 | 5.8         | 6.2                                       | 4.6             | 4.6                      | 6.8             | 5.2                                       | 3.6                      | 4.0                                       | 2.4                                       |
|                         | 1.50 | 11.4        | 13.4                                      | 10.2            | 7.6                      | 12.6            | 9.8                                       | 7.6                      | 9.0                                       | 5.6                                       |
| Clayton                 | 1.75 | 21.6        | 21.2                                      | 18.0            | 13.2                     | <b>22.8</b>     | 18.2                                      | 10.8                     | 15.2                                      | 9.6                                       |
|                         | 2.00 | <b>28.8</b> | 27.8                                      | 23.6            | 17.6                     | 27.2            | 24.2                                      | 13.4                     | 17.0                                      | 12.0                                      |
|                         | 2.25 | 44.6        | 42.0                                      | 38.4            | 30.6                     | 42.8            | 40.0                                      | 21.8                     | 29.6                                      | 21.0                                      |
|                         | 2.50 | 50.0        | 51.6                                      | 42.0            | 40.4                     | 47.6            | 43.2                                      | 27.6                     | 33.2                                      | 28.4                                      |
| $(n_1, n_2) = (50, 50)$ | 0.00 |             |   |                 |                          |                 |   |                          |   |   |
|                         | 0.00 | 2.6         | 4.0                                       | 2.4             | 3.0                      | 4.2             | 2.2                                       | 1.4                      | 3.0                                       | 0.8                                       |
|                         | 0.05 | 4.4         | 7.4                                       | 3.2             | 3.4                      | 6.4             | 4.0                                       | 2.8                      | 3.6                                       | 1.4                                       |
|                         | 0.10 | 5.8         | 11.4                                      | 6.6             | 3.0                      | 11.4            | 6.0                                       | 4.0                      | 6.8                                       | 2.2                                       |
| Gaussian                | 0.15 | 13.2        | 18.2                                      | 9.4             | 4.6                      | 17.8            | 9.2                                       | 5.6                      | 11.2                                      | 5.2                                       |
|                         | 0.20 | 19.6        | 26.2                                      | 18.0            | 9.6                      | 25.2            | 17.4                                      | 9.8                      | 15.0                                      | 7.6                                       |
|                         | 0.25 | 30.4        | 37.8                                      | 26.0            | 17.8                     | 36.0            | 28.6                                      | 17.4                     | 25.2                                      | 13.4                                      |
|                         | 0.30 | 47.4        | <b>52.8</b>                               | 43.2            | 28.2                     | 52.6            | 42.0                                      | 24.2                     | 37.0                                      | 22.6                                      |

Table 3: Empirical level (%) and power (%) of the tests are estimated from 500 samples from 3-dimensional Clayton and Gaussian copula given various parameters 50 sample size.

# 6 Concluding remarks

New tests for equality of multivariate copulas, based on Bernstein polynomials, have been proposed and studied. These tests improve upon the empirical tests introduced by Rémillard and Scaillet (2009) and demonstrate better performance in simulation studies. The limiting distribution of the proposed test statistics has been thoroughly investigated, and resampling methods using Bernstein polynomials have been developed and implemented to accurately simulate p-values.

However, the performance of these tests deteriorates as the dimensionality increases. It is essential to explore methods to mitigate the impact of dimensionality in nonparametric copula tests. Moreover, it would be intriguing to extend our results to k-sample equality tests of copulas (as seen in Bouzebda, Keziou et al. (2011) Bakam and Pommeret (2022) and Derumigny et al. (2022)) and to equality tests of copulas involving a large number of populations (as explored in Zhan and Hart (2014), Cousido-Rocha et al. (2019) and Jiménez-Gamero et al. (2022)).

# Appendix: Proofs of the results

**Lemma 3.** Let C be a copula function with continuous margins, then

$$\sup_{\boldsymbol{u}\in[0,1]^d} \left| C_{n,m}(\boldsymbol{u}) - C(\boldsymbol{u}) \right| \stackrel{a.s.}{=} O\left(\sqrt{\frac{\log\log n}{n}}\right) + O\left(m^{-1/2}\right)$$

as  $n \to \infty$ .

*Proof.* From Deheuvels (1981) and reference therein, a multivariate version of Glivenko-Cantelli theorem holds uniformly, *i.e.*,

$$\sup_{\boldsymbol{u}\in[0,1]^d} \left| C_n(\boldsymbol{u}) - C(\boldsymbol{u}) \right| \stackrel{a.s.}{=} O\left( \sqrt{\frac{\log\log n}{n}} \right).$$

The following steps are adapted from Janssen et al. (2012, Proof of Theorem 1) for the multivariate case. Let  $B_m$  denote the Bernstein copula, that is, the margins are known. Then by triangle inequality,

$$\sup_{\boldsymbol{u}\in[0,1]^d} |C_{n,m}(\boldsymbol{u}) - C(\boldsymbol{u})| \le \sup_{\boldsymbol{u}\in[0,1]^d} |C_{n,m}(\boldsymbol{u}) - B_m(\boldsymbol{u})| + \sup_{\boldsymbol{u}\in[0,1]^d} |B_m(\boldsymbol{u}) - C(\boldsymbol{u})|.$$

Further, let  $\boldsymbol{k} = (k_1, \ldots, k_d),$ 

$$\sup_{\boldsymbol{u}\in[0,1]^d} |C_{n,m}(\boldsymbol{u}) - B_m(\boldsymbol{u})| = \sup_{\boldsymbol{u}\in[0,1]^d} \left| \sum_{k_1=0}^m \cdots \sum_{k_d=0}^m \left[ C_n\left(\boldsymbol{k}/m\right) - C(\boldsymbol{k}/m) \right] \prod_{\ell=1}^d P_{k_\ell,m}(u_\ell) \right| \\ \leq \max_{0 \le k_1, \dots, k_d \le m} |C_n(\boldsymbol{k}/m) - C(\boldsymbol{k}/m)| \\ \leq \sup_{\boldsymbol{u}\in[0,1]^d} |C_n - C| = O\left(\sqrt{\frac{\log\log n}{n}}\right).$$

And using Lipschitz property of copulas,

$$\begin{split} \sup_{\boldsymbol{u}\in[0,1]^d} |B_m(\boldsymbol{u}) - C(\boldsymbol{u})| &= \sup_{\boldsymbol{u}\in[0,1]^d} \left| \sum_{k_1=0}^m \cdots \sum_{k_d=0}^m \left[ C(\boldsymbol{k}/m) - C(\boldsymbol{u}) \right] \prod_{\ell=1}^d P_{k_\ell,m}(u_\ell) \right| \\ &\leq \sup_{\boldsymbol{u}\in[0,1]^d} \left| \sum_{k_1=0}^m \cdots \sum_{k_d=0}^m |k_1/m - u_1| \prod_{\ell=1}^d P_{k_\ell,m}(u_\ell) + \cdots \right| \\ &+ \sum_{k_1=0}^m \cdots \sum_{k_d=0}^m |k_d/m - u_d| \prod_{\ell=1}^d P_{k_\ell,m}(u_\ell) \right|. \end{split}$$

Let  $\{B_{\ell} : \ell = 1, ..., d\}$  denote binomial random variables with parameter m and  $u_{\ell}$ . Then by the Cauchy-Schwarz inequality,

$$\sup_{\boldsymbol{u}\in[0,1]^d} |B_m(\boldsymbol{u}) - C(\boldsymbol{u})| \le \sum_{\ell=1}^d \sup_{u_\ell\in[0,1]} \left| \sqrt{m^{-2} \mathbb{V}\mathrm{ar}\left(B_\ell\right)} \right|$$
$$= \sum_{\ell=1}^d \sup_{u_\ell\in[0,1]} \sqrt{m^{-1} u_\ell(1-u_\ell)}$$
$$= \frac{d}{2} m^{-1/2}.$$

This completes the proof.

## A.1 Proof of Theorem 1

*Proof.* This proof is adapted from Harder and Stadtmüller (2017, Proof of Theorem 3.4). The weak convergence of  $R_n^m, T_n^m$  follows directly from the continuous mapping theorem and Lemma 1. For the convergence of  $S_n^m$ , one need to apply Harder and Stadtmüller (2017, Lemma A.1). Let  $C[0,1]^d$  denote the space of continuous functions on  $[0,1]^d$ ,  $D[0,1]^d$  denote the space of functions with continuity from upper right quadrant and limits from other quadrants on  $[0,1]^d$ , equipped with sup-norm. Further, denote  $BV_1[0,1]^d$  by the subspace of  $D[0,1]^d$  where functions have total variation bounded by 1. By the continuous mapping theorem,

$$\left( \{\mathbb{F}_n^m\}^2, \mathbb{C}_{n_1, m_1} \right) \rightsquigarrow \left(\mathbb{F}^2, \mathbb{C}\right)$$

in the space  $\ell^{\infty}[0,1]^d \times \ell^{\infty}[0,1]^d$ . Rewrite this as

$$\left(\{\mathbb{F}_{n}^{m}\}^{2},\mathbb{C}_{n_{1},m_{1}}\right)=\sqrt{n_{1}}\{(A_{n}^{m},C_{n_{1},m_{1}})-(A,C)\},\$$

where  $A \equiv 0$  and  $A_n^m := \frac{\sqrt{n_1 n_2}}{n_1 + n_2} \{C_{n_1, m_1} - D_{n_2, m_2}\}^2$ . Then, consider the map  $\phi : \ell^{\infty}[0, 1]^d \times BV_1[0, 1]^d \to \mathbb{R}$  defined by

$$\phi(a,b) = \int_{(0,1]^d} a \,\mathrm{d}b$$

Clearly,

$$S_{n}^{m} = \phi\left(\{\mathbb{F}_{n}^{m}\}^{2}, \mathbb{C}_{n_{1}, m_{1}}\right) = \sqrt{n_{1}}\{\phi(A_{n}^{m}, C_{n_{1}, m_{1}}) - \phi(A, C)\}.$$

To conclude the proof, by Harder and Stadtmüller (2017, Lemma A.1),  $\phi$  is Hadamard differentiable tangentially to  $C[0,1]^d \times D[0,1]^d$  at each  $(\alpha,\beta)$  in  $\ell^{\infty}[0,1]^d \times BV_1[0,1]^d$  such that  $\int |d\alpha| < \infty$  with derivative

$$\phi'_{(\alpha,\beta)}(a,b) = \int \alpha \, \mathrm{d}b + \int a \, \mathrm{d}\beta.$$

Then applying the Functional Delta Method,  $S_n^m \rightsquigarrow \phi'_{(A,C)} \left( \mathbb{F}^2, \mathbb{C} \right)$ , where

$$\phi'_{(A,C)}\left(\mathbb{F}^2,\mathbb{C}\right) = \int_{(0,1]^d} A \, \mathrm{d}\mathbb{C} + \int_{(0,1]^d} \mathbb{F}^2 \, \mathrm{d}C = \int_{(0,1]^d} \mathbb{F}^2 \, \mathrm{d}C.$$

This yields the desired result.

## A.2 Proof of Theorem 2

*Proof.* This proof is inspired by Genest et al. (2012, Proposition 4). The strongly uniform consistency of the *d*-variate empirical Bernstein copula is provided in Lemma 3. Then by the continuous mapping theorem and independence between the two samples, it follows immediately that  $R_n^m$  and  $T_n^m$  converge to R(C, D) and T(C, D) almost surely, respectively. Further, to prove the convergence of  $S_n^m$ , write

$$|S_{n}^{m} - S(C, D)| \le n_{2}\lambda_{n}\{|\gamma_{n}^{m}| + |\zeta_{n_{1}, m_{1}}|\},\$$

where

$$\gamma_{\boldsymbol{n}}^{\boldsymbol{m}} = \int_{[0,1]^d} \left\{ C_{n_1,m_1}(\boldsymbol{u}) - D_{n_2,m_2}(\boldsymbol{u}) \right\}^2 \mathrm{d}C_{n_1,m_1}(\boldsymbol{u}) - \int_{[0,1]^d} \left\{ C(\boldsymbol{u}) - D(\boldsymbol{u}) \right\}^2 \mathrm{d}C_{n_1,m_1}(\boldsymbol{u}),$$

and

$$\zeta_{n_1,m_1} = \int_{[0,1]^d} \left\{ C(\boldsymbol{u}) - D(\boldsymbol{u}) \right\}^2 \mathrm{d}C_{n_1,m_1}(\boldsymbol{u}) - \int_{[0,1]^d} \left\{ C(\boldsymbol{u}) - D(\boldsymbol{u}) \right\}^2 \mathrm{d}C(\boldsymbol{u})$$

Since

$$\begin{aligned} \left| \{C_{n_1,m_1}(\boldsymbol{u}) - D_{n_2,m_2}(\boldsymbol{u})\}^2 - \{C(\boldsymbol{u}) - D(\boldsymbol{u})\}^2 \right| &= \left| [C_{n_1,m_1}(\boldsymbol{u}) + C(\boldsymbol{u})] - [D_{n_2,m_2}(\boldsymbol{u}) + D(\boldsymbol{u})] \right| \\ &\times \left| [C_{n_1,m_1}(\boldsymbol{u}) - C(\boldsymbol{u})] - [D_{n_2,m_2}(\boldsymbol{u}) - D(\boldsymbol{u})] \right| \\ &\leq \left[ |C_{n_1,m_1}(\boldsymbol{u}) + C(\boldsymbol{u})| + |D_{n_2,m_2}(\boldsymbol{u}) + D(\boldsymbol{u})| \right] \\ &\times \left[ |C_{n_1,m_1}(\boldsymbol{u}) - C(\boldsymbol{u})| + |D_{n_2,m_2}(\boldsymbol{u}) - D(\boldsymbol{u})| \right] \\ &\leq 4 \sup_{\boldsymbol{u} \in [0,1]^d} |C_{n_1,m_1}(\boldsymbol{u}) - C(\boldsymbol{u})| \\ &+ 4 \sup_{\boldsymbol{u} \in [0,1]^d} |D_{n_2,m_2}(\boldsymbol{u}) - D(\boldsymbol{u})| \,, \end{aligned}$$

using Lemma 3,

$$|\gamma_{n_1,n_2}^{\boldsymbol{m}}| \le 4 \left\{ \sup_{\boldsymbol{u} \in [0,1]^d} \left| C_{n_1,m_1}(\boldsymbol{u}) - C(\boldsymbol{u}) \right| + \sup_{\boldsymbol{u} \in [0,1]^d} \left| D_{n_2,m_2}(\boldsymbol{u}) - D(\boldsymbol{u}) \right| \right\} \xrightarrow[\min(n_1,n_2) \to \infty]{a.s.} 0.$$

As for  $\zeta_{n_1,m_1}$ , note that,  $C_{n_1,m_1} : \Omega \times [0,1]^d \to [0,1]$  is a random mappings for fixed  $\omega \in \Omega$ , and by Lemma 3,  $C_{n_1,m_1}$  converges pointwise to C in an almost sure sense. Let  $\tilde{U}_{n_1,m_1} \sim C_{n_1,m_1}$  and  $U \sim C$ . Hence

$$\widetilde{U}_{n_1,m_1} \xrightarrow[n_1 \to \infty]{d} U.$$

In light of Genest et al. (1995, Proposition A.1 (i)) and Harder and Stadtmüller (2017, Proof of Lemma 3.1), let

$$\psi: [0,1]^d \to \mathbb{R}, \quad \psi(\boldsymbol{u}) := \{C(\boldsymbol{u}) - D(\boldsymbol{u})\}^2.$$

Then by the continuous mapping theorem,

$$V_{n_1,m_1} \coloneqq \psi\left(\widetilde{U}_{n_1,m_1}\right) \xrightarrow[n_1 \to \infty]{d} \psi(U) \eqqcolon V.$$

If  $V_{n_1,m_1}$  is asymptotically uniformly integrable, then applying van der Vaart and Wellner (1996, Theorem 1.11.3), one has

$$\int_{[0,1]^d} \left\{ C(\boldsymbol{u}) - D(\boldsymbol{u}) \right\}^2 \mathrm{d}C_{n_1,m_1}(\boldsymbol{u}) \xrightarrow[n_1 \to \infty]{} \int_{[0,1]^d} \left\{ C(\boldsymbol{u}) - D(\boldsymbol{u}) \right\}^2 \mathrm{d}C(\boldsymbol{u}).$$

Then  $\zeta_{n_1,m_1} \xrightarrow[n_1 \to \infty]{} 0$ . To this end, it will be shown that there exists  $\varepsilon > 0$  such that  $\mathbb{E}\left[|V_{n_1,m_1}|^{1+\varepsilon}\right]$  is bounded. Indeed, since  $\{C(\boldsymbol{u}) - D(\boldsymbol{u})\}^2 \leq 1$ ,

$$\mathbb{E}\left[|V_{n_1,m_1}|^{1+\varepsilon}\right] = \int_{[0,1]^d} |\psi(\boldsymbol{u})|^{1+\varepsilon} \, \mathrm{d}C_{n_1,m_1}(\boldsymbol{u}) \leq 1.$$

Combining  $\gamma_{n_1,n_2}^m$  and  $\zeta_{n_1,m_1}$ ,  $S_n^m$  converges to S(C,D) almost surely.

## A.3 Proof of Lemma 2

*Proof.* Under these assumptions, one need to use the framework in Segers et al. (2017). Specifically, let  $\mu_{m_1,u}$  be the law of random vector  $(B_1/m_1, \ldots, B_d/m_1)$ , where  $\{B_\ell : \ell = 1, \ldots, d\}$  follow  $\mathsf{Binomial}(m_1, u_\ell)$ . The empirical Bernstein copula in Equation (2) can be rewritten as

$$C_{n_1,m_1}(\boldsymbol{u}) = \int_{[0,1]^d} C_{n_1}(\boldsymbol{w}) \,\mathrm{d}\mu_{m,\boldsymbol{u}}(\boldsymbol{w}), \qquad \boldsymbol{u} \in [0,1]^d.$$

Moreover, write w(t) = u + t(w - u) with  $t \in [0, 1]$ . Then, the empirical Bernstein copula process is

$$\mathbb{C}_{n_{1},m_{1}}(\boldsymbol{u}) = \sqrt{n_{1}} \{C_{n_{1},m_{1}}(\boldsymbol{u}) - C(\boldsymbol{u})\} \\
= \sqrt{n_{1}} \left\{ C_{n_{1},m_{1}}(\boldsymbol{u}) - \int_{[0,1]^{d}} C(\boldsymbol{w}) \,\mathrm{d}\mu_{m_{1},\boldsymbol{u}}(\boldsymbol{w}) + \int_{[0,1]^{d}} C(\boldsymbol{w}) \,\mathrm{d}\mu_{m_{1},\boldsymbol{u}}(\boldsymbol{w}) - C(\boldsymbol{u}) \right\} \\
= \int_{[0,1]^{d}} \sqrt{n_{1}} \{C_{n_{1}}(\boldsymbol{w}) - C(\boldsymbol{w})\} \,\mathrm{d}\mu_{m_{1},\boldsymbol{u}}(\boldsymbol{w}) + \sqrt{n_{1}} \left\{ \int_{[0,1]^{d}} C(\boldsymbol{w}) \,\mathrm{d}\mu_{m_{1},\boldsymbol{u}}(\boldsymbol{w}) - C(\boldsymbol{u}) \right\} \\
= T_{1} + T_{2}.$$
(8)

The two terms are dealt with separately.

• For the term  $T_1$ , according to Segers et al. (2017, Proposition 3.1), one has

$$\sup_{\boldsymbol{u}\in[0,1]^d} \left| \int_{[0,1]^d} \sqrt{n_1} \left\{ C_{n_1}(\boldsymbol{w}) - C(\boldsymbol{w}) \right\} \, \mathrm{d}\mu_{m_1,\boldsymbol{u}}(\boldsymbol{w}) - \sqrt{n_1} \left\{ C_{n_1}(\boldsymbol{u}) - C(\boldsymbol{u}) \right\} \right| = o_p(1).$$

And note that,  $\sqrt{n_1} \{C_{n_1}(\boldsymbol{u}) - C(\boldsymbol{u})\} \rightsquigarrow \mathbb{B}_C(\boldsymbol{u})$  in  $\ell^{\infty}([0,1]^d)$  under the Assumption 1 (see Segers (2012)). Therefore,  $T_1 \rightsquigarrow \mathbb{B}_C(\boldsymbol{u})$  as n goes to infinity.

• For the term  $T_2$ , Let  $m_1 = cn_1^{\alpha}$  for some c > 0, by Kojadinovic (2022, Lemma 3.1), under Assumption 1-2, one has

$$\sup_{\boldsymbol{u}\in[0,1]^d} \sqrt{n_1} \left| \int_{[0,1]^d} C(\boldsymbol{w}) \, \mathrm{d}\mu_{m_1,\boldsymbol{u}}(\boldsymbol{w}) - C(\boldsymbol{u}) \right| = O\left( n_1^{(3-4\alpha)/6} \right)$$

almost surely. Therefore, if  $\alpha > 3/4$ ,  $T_2$  goes to zero as n goes to infinity.

Combining above results completes the proof.

#### A.4Proof of Theorem 3

*Proof.* One can verify an intermediate result that as  $\min(n_1, n_2) \to \infty$ 

$$\sup_{\boldsymbol{u}\in[0,1]^d} \left| \mathbb{F}_{\boldsymbol{n}}^{\boldsymbol{m}}(\boldsymbol{u}) - \mathbb{F}_{\boldsymbol{n}}^*(\boldsymbol{u}) \right| \stackrel{a.s.}{=} O\left( \Psi(n_1, n_2) \right).$$
(9)

ı.

Indeed, let  $\mathbb{F}_n(u) := \sqrt{1 - \lambda_n} \mathbb{C}_{n_1} - \sqrt{\lambda_n} \mathbb{D}_{n_2}$ , in the light of Bouzebda et al. (2011, Proof of Theorem 2.1). By Borisov (1982), for a sequence of Brownian bridge  $\{\mathbb{B}_{n_1,C}\}_{n_1\in\mathbb{N}^+}$ , which are independent copies of  $\mathbb{B}_C$ , one has as  $n_1 \to \infty$ ,

$$\sup_{\boldsymbol{u}\in[0,1]^d} \left| \mathbb{G}_{n_1}(\boldsymbol{u}) - \mathbb{B}_{n_1,C}(\boldsymbol{u}) \right| \stackrel{a.s.}{=} O\left( n_1^{-1/(2(2d-1))} \log n_1 \right),$$

where

$$\mathbb{G}_{n_1}(\boldsymbol{u}) = \sqrt{n} \left\{ G_{n_1}(\boldsymbol{u}) - C(\boldsymbol{u}) \right\} = \frac{1}{\sqrt{n_1}} \sum_{i=1}^{n_1} \left[ \prod_{\ell=1}^d \mathbb{I} \left( U_{i\ell} \le u_\ell \right) - C(\boldsymbol{u}) \right].$$

Then, under our assumptions, by Segers (2012), as  $n_1 \to \infty$ ,

$$\sup_{\boldsymbol{u}\in[0,1]^d} \left| \mathbb{C}_{n_1}(\boldsymbol{u}) - \mathbb{G}_{n_1}(\boldsymbol{u}) + \sum_{\ell=1}^d \mathbb{G}_{n_1}(\boldsymbol{u}^\ell) \dot{C}_\ell(\boldsymbol{u}) \right| \stackrel{a.s.}{=} O\left( n_1^{-1/4} (\log n_1)^{1/2} (\log \log n_1)^{1/4} \right).$$

Т

Since  $\dot{C}_{\ell}(\boldsymbol{u}), \ell = 1, \dots, d$  are bounded,

$$\begin{split} \sup_{\boldsymbol{u}\in[0,1]^d} \left| \mathbb{C}_{n_1}(\boldsymbol{u}) - \mathbb{C}^{(n_1)}(\boldsymbol{u}) \right| &\leq \sup_{\boldsymbol{u}\in[0,1]^d} \left| \mathbb{C}_{n_1}(\boldsymbol{u}) - \mathbb{G}_{n_1}(\boldsymbol{u}) + \sum_{\ell=1}^d \mathbb{G}_{n_1}(\boldsymbol{u}^\ell) \dot{C}_\ell(\boldsymbol{u}) \right| \\ &+ \sup_{\boldsymbol{u}\in[0,1]^d} \left| \mathbb{G}_{n_1}(\boldsymbol{u}) - \sum_{\ell=1}^d \mathbb{G}_{n_1}(\boldsymbol{u}^\ell) \dot{C}_\ell(\boldsymbol{u}) - \mathbb{C}^{(n_1)}(\boldsymbol{u}) \right| \\ &= O\left( n_1^{-1/4} (\log n_1)^{1/2} (\log \log n_1)^{1/4} \right) + O\left( n_1^{-1/(2(2d-1))} \log n_1 \right) \\ &= O\left( n_1^{-1/(2(2d-1))} \log n_1 \right) \end{split}$$

almost surely as  $n_1 \to \infty$ . Similarly, one can obtain

$$\sup_{\boldsymbol{u}\in[0,1]^d} \left| \mathbb{D}_{n_2}(\boldsymbol{u}) - \mathbb{D}^{(n_2)}(\boldsymbol{u}) \right| = O\left( n_2^{-1/(2(2d-1))} \log n_2 \right)$$

almost surely as  $n_2 \rightarrow \infty$ . Therefore, one has almost surely

$$\sup_{\boldsymbol{u}\in[0,1]^d} \left| \mathbb{F}_{\boldsymbol{n}}(\boldsymbol{u}) - \mathbb{F}_{\boldsymbol{n}}^*(\boldsymbol{u}) \right| = O\left( \Psi(n_1, n_2) \right).$$

Therefore,

$$\begin{split} \sup_{\bm{u}\in[0,1]^d} |\mathbb{F}_{\bm{n}}^{\bm{m}}(\bm{u}) - \mathbb{F}_{\bm{n}}^*(\bm{u})| &\leq \sup_{\bm{u}\in[0,1]^d} |\mathbb{F}_{\bm{n}}^{\bm{m}}(\bm{u}) - \mathbb{F}_{\bm{n}}(\bm{u})| + \sup_{\bm{u}\in[0,1]^d} |\mathbb{F}_{\bm{n}}(\bm{u}) - \mathbb{F}_{\bm{n}}^*(\bm{u})| \\ &\leq n_2\lambda_{\bm{n}} \left[ \sup_{\bm{u}\in[0,1]^d} |C_{n_1,m_1}(\bm{u}) - C_{n_1}(\bm{u})| + \sup_{\bm{u}\in[0,1]^d} |D_{n_2,m_2}(\bm{u}) - D_{n_2}(\bm{u})| \right] \end{split}$$

$$+ O\left(\Psi(n_{1}, n_{2})\right)$$
  
=  $(1 - \lambda_{n}) \sup_{\boldsymbol{u} \in [0, 1]^{d}} |\mathbb{C}_{n_{1}, m_{1}}(\boldsymbol{u}) - \mathbb{C}_{n_{1}}(\boldsymbol{u})| + \lambda_{n} \sup_{\boldsymbol{u} \in [0, 1]^{d}} |\mathbb{D}_{n_{2}, m_{2}}(\boldsymbol{u}) - \mathbb{D}_{n_{2}}(\boldsymbol{u})|$   
+  $O\left(\Psi(n_{1}, n_{2})\right).$  (10)

Further, adapted from Segers et al. (2017), under our assumptions, as  $\min(n_1, n_2) \to \infty$ ,

$$\sup_{\boldsymbol{u}\in[0,1]^d} \left| \mathbb{C}_{n_1,m_1}(\boldsymbol{u}) - \mathbb{C}_{n_1}(\boldsymbol{u}) \right| = O\left(n_1^{-1/2}\right),$$
  
$$\sup_{\boldsymbol{u}\in[0,1]^d} \left| \mathbb{D}_{n_2,m_2}(\boldsymbol{u}) - \mathbb{D}_{n_2}(\boldsymbol{u}) \right| = O\left(n_2^{-1/2}\right).$$
(11)

One has

$$(10) = O\left(n_1^{-1/2}\right) + O\left(n_2^{-1/2}\right) + O\left(\Psi(n_1, n_2)\right) = O\left(\Psi(n_1, n_2)\right).$$

Based on the intermediate result Equation (9), one can show strong approximation of these statistics.

• For  $R_n^m$ ,

$$\begin{split} \left| R_n^m - \int_{[0,1]^d} \{ \mathbb{F}_n^*(u) \}^2 \, \mathrm{d}u \right| &= \left| \int_{[0,1]^d} \{ \mathbb{F}_n^m(u) \}^2 - \{ \mathbb{F}_n^*(u) \}^2 \, \mathrm{d}u \right| \\ &\leq \left( \sup_{u \in [0,1]^d} \left| \mathbb{F}_n^m(u) - \mathbb{F}_n^*(u) \right| \right) \left( \sup_{u \in [0,1]^d} \left| \mathbb{F}_n^m(u) + \mathbb{F}_n^*(u) \right| \right) \\ &\leq \left( \sup_{u \in [0,1]^d} \left| \mathbb{F}_n^m(u) - \mathbb{F}_n^*(u) \right| \right) \\ &\times \left( \sup_{u \in [0,1]^d} \left| \mathbb{F}_n^m(u) - \mathbb{F}_n^*(u) \right| + 2 \sup_{u \in [0,1]^d} \left| \mathbb{F}_n^m(u) \right| \right) \\ &= O\left( \max\left( (\log \log n_1)^{1/2}, (\log \log n_2)^{1/2} \right) \Psi(n_1, n_2) \right), \end{split}$$

where the last equality follows by Janssen et al. (2012, Theorem 1).

• For  $S_n^m$ ,

$$\begin{aligned} \left| S_{n}^{m} - \int_{[0,1]^{d}} \{\mathbb{F}_{n}^{*}(\boldsymbol{u})\}^{2} \, \mathrm{d}C(\boldsymbol{u}) \right| &\leq \left| S_{n}^{m} - \int_{[0,1]^{d}} \{\mathbb{F}_{n}^{*}(\boldsymbol{u})\}^{2} \, \mathrm{d}C_{n_{1},m_{1}}(\boldsymbol{u}) \right| \\ &+ \left| \int_{[0,1]^{d}} \{\mathbb{F}_{n}^{*}(\boldsymbol{u})\}^{2} \, \mathrm{d}C_{n_{1},m_{1}}(\boldsymbol{u}) - \int_{[0,1]^{d}} \{\mathbb{F}_{n}^{*}(\boldsymbol{u})\}^{2} \, \mathrm{d}C(\boldsymbol{u}) \right| \\ &= O\left( \max\left( (\log \log n_{1})^{1/2}, (\log \log n_{2})^{1/2} \right) \Psi(n_{1},n_{2}) \right) \\ &+ \left| \int_{[0,1]^{d}} \{\mathbb{F}_{n}^{*}(\boldsymbol{u})\}^{2} \, \mathrm{d}\{C_{n_{1},m_{1}}(\boldsymbol{u}) - C(\boldsymbol{u})\} \right| \\ &\leq O\left( \max\left( (\log \log n_{1})^{1/2}, (\log \log n_{2})^{1/2} \right) \Psi(n_{1},n_{2}) \right) \end{aligned}$$

$$+ \sup_{\boldsymbol{u} \in [0,1]^d} \left| \{ \mathbb{F}_{\boldsymbol{n}}^*(\boldsymbol{u}) \}^2 \right| \sup_{\boldsymbol{u} \in [0,1]^d} \left| \{ C_{n_1,m_1}(\boldsymbol{u}) - C(\boldsymbol{u}) \} \right|$$
  
=  $O\left( \max\left( (\log \log n_1)^{1/2}, (\log \log n_2)^{1/2} \right) \Psi(n_1,n_2) \right)$   
+  $O(1)O\left( n_1^{-1/2} (\log \log n_1)^{1/2} \right)$   
=  $O\left( \max\left( (\log \log n_1)^{1/2}, (\log \log n_2)^{1/2} \right) \Psi(n_1,n_2) \right),$ 

where the O(1) comes form the fact that  $\mathbb{F}_n^*$  is a centred Gaussian process.

• For  $T_n^m$ , the result immediately follows.

This completes the proof.

## A.5 Proof of Proposition 1

*Proof.* We only prove the results for a sample X with size  $n_1$ . According to Rémillard and Scaillet (2009), one has for  $\boldsymbol{u} \in [0, 1]^d$  and  $h \in \{1, \ldots, H\}$ ,

$$\mathbb{G}_{n_1}(\boldsymbol{u}) = \sqrt{n} \left\{ G_{n_1}(\boldsymbol{u}) - C(\boldsymbol{u}) \right\} = \frac{1}{\sqrt{n_1}} \sum_{i=1}^{n_1} \left[ \prod_{\ell=1}^d \mathbb{I} \left( U_{i\ell} \le u_\ell \right) - C(\boldsymbol{u}) \right] \rightsquigarrow \mathbb{B}_C(\boldsymbol{u}),$$

and

$$\mathbb{G}_{n_1}^{(h)}(\boldsymbol{u}) = \frac{1}{\sqrt{n_1}} \sum_{i=1}^{n_1} \left( \xi_i^{(h)} - \overline{\xi}_{n_1}^{(h)} \right) \left[ \prod_{\ell=1}^d \mathbb{I}\left( \widehat{U}_{i\ell} \le k_\ell / m_1 \right) \right] \rightsquigarrow \mathbb{B}_C(\boldsymbol{u}).$$

To complete the proof, one needs to show that the difference between  $\left(\mathbb{G}_{n_1}, \mathbb{G}_{n_1}^{(h)}\right)$  and  $\left(\mathbb{G}_{n_1,m_1}, \mathbb{G}_{n_1,m_1}^{(h)}\right)$  is asymptotically negligible.

By Lemma 2, under the assumptions, one has

$$\mathbb{C}_{n_1,m_1}(\boldsymbol{u}) = \sqrt{n_1} \{ C_{n_1,m_1}(\boldsymbol{u}) - C(\boldsymbol{u}) \} \rightsquigarrow \mathbb{C}(\boldsymbol{u}) = \mathbb{B}_C(\boldsymbol{u}) - \sum_{\ell=1}^d \mathbb{B}_C(\boldsymbol{u}^\ell) \dot{C}_\ell(\boldsymbol{u})$$

which implies that

$$\mathbb{G}_{n_1,m_1}(\boldsymbol{u}) \rightsquigarrow \mathbb{B}_C(\boldsymbol{u}).$$

Furthermore, given that  $\mathbb{C}_{n_1,m_1}(\boldsymbol{u})$  and  $\mathbb{C}_{n_1}(\boldsymbol{u})$  converge to the same limit, one has that  $\mathbb{G}_{n_1,m_1}^{(h)}(\boldsymbol{u})$  and  $\mathbb{G}_{n_1}^{(h)}(\boldsymbol{u})$  will also converge to the same limit. This leads to the conclusion that  $\mathbb{G}_{n_1,m_1}^{(h)}(\boldsymbol{u}) \rightsquigarrow \mathbb{B}_C(\boldsymbol{u})$ .  $\Box$ 

## A.6 Proof of Proposition 2

*Proof.* We only show the uniform consistency for

$$\partial C_{n_1,m_1}(\boldsymbol{u})/\partial u_1 = m_1 \sum_{k_1=0}^{m_1-1} \cdots \sum_{k_d=0}^{m_1} \left\{ C_{n_1}\left(\frac{k_1+1}{m_1}, \dots, \frac{k_d}{m_1}\right) - C_{n_1}\left(\frac{k_1}{m_1}, \dots, \frac{k_d}{m_1}\right) \right\}$$

$$\times P_{k_1,m_1-1}(u_1)\cdots P_{k_d,m_1}(u_d).$$

The result for other partial derivatives can be obtained similarly. For any  $\{u \in [0,1]^d : u_1 \in [b_{n_1}, 1-b_{n_1}]\}$ , one has

$$\begin{aligned} \left| \frac{\partial C_{n_1,m_1}(\boldsymbol{u})}{\partial u_1} - \dot{C}_1(\boldsymbol{u}) \right| \\ &\leq \left| m_1 \sum_{k_1=0}^{m_1-1} \cdots \sum_{k_d=0}^{m_1} \left\{ C_{n_1} \left( \frac{k_1+1}{m_1}, \dots, \frac{k_d}{m_1} \right) - C_{n_1} \left( \frac{k_1}{m_1}, \dots, \frac{k_d}{m_1} \right) - C \left( \frac{k_1+1}{m_1}, \dots, \frac{k_d}{m_1} \right) \right. \\ &+ C \left( \frac{k_1}{m_1}, \dots, \frac{k_d}{m_1} \right) \right\} P_{k_1,m_1-1}(u_1) \cdots P_{k_d,m_1}(u_d) \left| + \left| m_1 \sum_{k_1=0}^{m_1-1} \cdots \sum_{k_d=0}^{m_1} \left\{ C \left( \frac{k_1+1}{m_1}, \dots, \frac{k_d}{m_1} \right) \right. \right. \\ &\left. - C \left( \frac{k_1}{m_1}, \dots, \frac{k_d}{m_1} \right) \right\} P_{k_1,m_1-1}(u_1) \cdots P_{k_d,m_1}(u_d) - \dot{C}_1(\boldsymbol{u}) \right| \\ &= A_1 + A_2. \end{aligned}$$

Further, let  $P'_{k_1,m_1}(u_1)$  be the derivative of  $P_{k_1,m_1}(u_1)$ , then

$$A_{1} \leq \sum_{k_{1}=0}^{m_{1}} \cdots \sum_{k_{d}=0}^{m_{1}} \left| C_{n_{1}} \left( \frac{k_{1}}{m_{1}}, \dots, \frac{k_{d}}{m_{1}} \right) - C \left( \frac{k_{1}}{m_{1}}, \dots, \frac{k_{d}}{m_{1}} \right) \right| \left| P_{k_{1},m_{1}}'(u_{1}) \right| \cdots P_{k_{d},m_{1}}(u_{d})$$

$$\leq \sup_{u \in [0,1]^{d}} \left| C_{n_{1}}(u) - C(u) \right| \cdot \sum_{k_{1}=0}^{m_{1}} \left| P_{k_{1},m_{1}}'(u_{1}) \right|$$

$$= O \left( m_{1}^{1/2} n_{1}^{-1/2} (\log \log n_{1})^{1/2} \right)$$

almost surely as  $n_1 \to \infty$  and where  $\sum_{k_1=0}^{m_1} |P'_{k_1,m_1}(u_1)| = O\left(m_1^{1/2}\right)$  by Janssen et al. (2014, Lemma 1). For dealing with  $A_2$ , let  $\nu_{m_1,u}$  be the law of random vector  $(S_1/(m_1-1), B_2/m_1, \dots, B_d/m_1)$ , where

For dealing with  $A_2$ , let  $\nu_{m_1,u}$  be the law of random vector  $(S_1/(m_1-1), B_2/m_1, \ldots, B_d/m_1)$ , where  $S_1$  follows Binomial $(m_1 - 1, u_1)$  and  $\{B_\ell : \ell = 2, \ldots, d\}$  follow Binomial $(m_1, u_\ell)$ . One has

$$\sum_{k_1=0}^{m_1-1} \cdots \sum_{k_d=0}^{m_1} \left\{ C\left(\frac{k_1+1}{m_1}, \dots, \frac{k_d}{m_1}\right) - C\left(\frac{k_1}{m_1}, \dots, \frac{k_d}{m_1}\right) \right\} P_{k_1,m_1-1}(u_1) \cdots P_{k_d,m_1}(u_d)$$
$$= \int_{[0,1]^d} C\left(\left(w_1 + \frac{1}{m_1-1}\right) \frac{m_1-1}{m_1}, w_2, \dots, w_d\right) d\nu_{m_1,u}(\boldsymbol{w})$$
$$- \int_{[0,1]^d} C\left(w_1 \frac{m_1-1}{m_1}, w_2, \dots, w_d\right) d\nu_{m_1,u}(\boldsymbol{w}).$$

Using the representation in Segers et al. (2017, Proof of Proposition 3.4), for 0 < t < 1, one has

$$A_{2} = \left| m_{1} \int_{[0,1]^{d}} \left[ C\left( \left( w_{1} + \frac{1}{m_{1} - 1} \right) \frac{m_{1} - 1}{m_{1}}, w_{2}, \dots, w_{d} \right) - C\left( w_{1} \frac{m_{1} - 1}{m_{1}}, w_{2}, \dots, w_{d} \right) \right] d\nu_{m_{1}, \boldsymbol{u}}(\boldsymbol{w}) - \dot{C}_{1}(\boldsymbol{u}) \right] d\nu_{m_{1}, \boldsymbol{u}}(\boldsymbol{w}) dt \right|.$$

$$= \left| \int_{0}^{1} \int_{[0,1]^{d}} \left[ \dot{C}_{1}\left( \left( \frac{m_{1} - 1}{m_{1}} w_{1} + \frac{1 + t}{m_{1}} \right), w_{2}, \dots, w_{d} \right) - \dot{C}_{1}(\boldsymbol{u}) \right] d\nu_{m_{1}, \boldsymbol{u}}(\boldsymbol{w}) dt \right|.$$
(12)

Let 
$$w_1'(w_1, t) := \frac{m_1 - 1}{m_1} w_1 + \frac{1 + t}{m_1}$$
 and  $\varepsilon_{n_1} = b_{n_1}/2$ , then  

$$(12) \leq \left| \int_0^1 \int_{[0,1]^d} \left[ \dot{C}_1\left(w_1', w_2, \dots, w_d\right) - \dot{C}_1(u) \right] \mathbb{I}\left( \max(|w_1' - u_1|, |w_2 - u_2|, \dots, |w_d - u_d|) \leq \varepsilon_{n_1} \right) d\nu_{m_1, u}(w) dt \right|$$

$$+ \left| \int_0^1 \int_{[0,1]^d} \left[ \dot{C}_1\left(w_1', w_2, \dots, w_d\right) - \dot{C}_1(u) \right] \mathbb{I}\left( \max(|w_1' - u_1|, |w_2 - u_2|, \dots, |w_d - u_d|) > \varepsilon_{n_1} \right) d\nu_{m_1, u}(w) dt \right|$$

$$= A_{21} + A_{22}.$$

Further, the two terms are dealt with separately using the strategy in Kojadinovic (2022, Proof of Lemma 3.1),

• For term  $A_{21}$ , under Assumption 1-2, by Segers (2012, Lemma 4.3), for a constant L > 0, one has

$$\left| \dot{C}_1 \left( w'_1, w_2, \dots, w_d \right) - \dot{C}(\boldsymbol{u}) \right| \mathbb{I} \left( \max(|w'_1 - u_1|, |w_2 - u_2|, \dots, |w_d - u_d|) \le \varepsilon_{n_1} \right)$$
$$\le L b_{n_1}^{-1} \left[ |w'_1 - u_1| + |w_2 - u_2| + \dots + |w_d - u_d| \right].$$

Further,

$$A_{21} \leq Lb_{n_1}^{-1} \int_{[0,1]^d} \int_0^1 \left[ |w_1 - u_1| + |w_2 - u_2| + \dots + |w_d - u_d| + |w_1' - w_1| \right] dt \, d\nu_{m_1, \boldsymbol{u}}(\boldsymbol{w})$$
  

$$\leq Lb_{n_1}^{-1} \int_{[0,1]^d} \int_0^1 \left[ |w_1 - u_1| + |w_2 - u_2| + \dots + |w_d - u_d| + \left| \frac{1+t}{m_1} \right| + \left| \frac{w_1}{m_1} \right| \right] dt \, d\nu_{m_1, \boldsymbol{u}}(\boldsymbol{w})$$
  

$$= Lb_{n_1}^{-1} \int_{[0,1]^d} \left[ |w_1 - u_1| + |w_2 - u_2| + \dots + |w_d - u_d| + \frac{3}{2m_1} + \frac{w_1}{m_1} \right] d\nu_{m_1, \boldsymbol{u}}(\boldsymbol{w})$$
(13)

By Cauchy-Schwarz inequality,

$$(13) \le Lb_{n_1}^{-1} \left[ O\left(m_1^{-1/2}\right) + O\left(m_1^{-1}\right) \right]$$
$$= O\left(b_{n_1}^{-1}m_1^{-1/2}\right).$$

• For term  $A_{22}$ , since  $\dot{C}_1 \in [0,1]$  and using the result in the proof of  $A_{21}$ ,

$$A_{22} \left| \int_{0}^{1} \int_{[0,1]^{d}} \frac{2}{\varepsilon_{n_{1}}} \max(|w_{1}' - u_{1}|, |w_{2} - u_{2}|, \dots, |w_{d} - u_{d}|) \,\mathrm{d}\nu_{m_{1},\boldsymbol{u}}(\boldsymbol{w}) \,\mathrm{d}t \right|$$
(14)

$$\leq \left| \int_{0}^{1} \int_{[0,1]^{d}} \frac{2}{\varepsilon_{n_{1}}} \max(|w_{1}'-u_{1}|+|w_{2}-u_{2}|+\dots+|w_{d}-u_{d}|) \,\mathrm{d}\nu_{m_{1},\boldsymbol{u}}(\boldsymbol{w}) \,\mathrm{d}t \right| \tag{15}$$

$$\leq \varepsilon_{n_1}^{-1} \left[ O\left( m_1^{-1/2} \right) + O(m_1^{-1}) \right] \tag{16}$$

$$= O\left(b_{n_1}^{-1}m_1^{-1/2}\right).$$
(17)

Therefore,

$$\sup_{\{\boldsymbol{u}\in[0,1]^d:u_1\in[b_{n_1},1-b_{n_1}]\}} \left| \frac{\partial C_{n_1,m_1}(\boldsymbol{u})}{\partial u_1} - \dot{C}_1(\boldsymbol{u}) \right| = O\left(m_1^{1/2}n_1^{-1/2}(\log\log n_1)^{1/2}\right) + O\left(b_{n_1}^{-1}m_1^{-1/2}\right)$$

almost surely as  $n_1 \to \infty$ , which completes the proof.

## A.7 Proof of Theorem 5

*Proof.* We only show the result for  $\mathbb{C}$ . The  $\mathbb{D}$  process can be treated similarly. The empirical copula for subsample h is defined as

$$C_{b_1}^{(I_h)}(\boldsymbol{u}) = \frac{1}{b_1} \sum_{i=1}^{b_1} \prod_{\ell=1}^d \mathbb{I}\left(\widehat{U}_{i\ell} \le u_\ell\right), \qquad h \in \{1, \dots, H\}.$$

The corresponding corrected subsample empirical copula process is

$$\mathbb{C}_{b_1}^{(I_h)}(\boldsymbol{u}) = \sqrt{\frac{b_1}{1 - b_1/n_1}} \{ C_{b_1}^{(I_h)}(\boldsymbol{u}) - C_{n_1}(\boldsymbol{u}) \}.$$

By Kojadinovic and Stemikovskaya (2019, Theorem 1), under our assumptions,

$$\left(\mathbb{C}_{n_1},\mathbb{C}_{b_1}^{(I_1)},\ldots,\mathbb{C}_{b_1}^{(I_H)}\right)\rightsquigarrow \left(\mathbb{C},\mathbb{C}^{(1)},\ldots,\mathbb{C}^{(H)}\right).$$

Then one can show that  $\mathbb{C}_{b_1,m_{(1)}}^{(I_h)}$  and  $\mathbb{C}_{b_1}^{(I_h)}$  are asymptotically equivalent. Indeed, applying Segers et al. (2017, Theorem 3.6), as  $n_1 \to \infty$ ,

$$\begin{split} &\sqrt{\frac{b_1}{1-b_1/n_1}}\{C_{b_1,m_{(1)}}^{(I_h)}(\boldsymbol{u}) - C(\boldsymbol{u})\} = \sqrt{\frac{b_1}{1-b_1/n_1}}\{C_{b_1}^{(I_h)}(\boldsymbol{u}) - C(\boldsymbol{u})\} + o_p(1) \\ &\sqrt{\frac{b_1}{1-b_1/n_1}}\{C_{n_1,m_1}(\boldsymbol{u}) - C(\boldsymbol{u})\} = \sqrt{\frac{b_1}{1-b_1/n_1}}\{C_{n_1}(\boldsymbol{u}) - C(\boldsymbol{u})\} + o_p(1). \end{split}$$

Therefore,

$$\mathbb{C}_{b_1,m_{(1)}}^{(I_h)} = \mathbb{C}_{b_1}^{(I_h)} + o_p(1), \quad n_1 \to \infty,$$

which completes the proof.

# References

- Bahraoui, T., T. Bouezmarni, and J.-F. Quessy (2018). Testing the symmetry of a dependence structure with a characteristic function. *Dependence Modeling* 6(1), 331–355.
- Bakam, Y. I. N. and D. Pommeret (2022). K-sample test for equality of copulas. arXiv preprint arXiv:2112.05623.
- Beare, B. K. and J. Seo (2020). Randomization tests of copula symmetry. *Econometric Theory* 36(6), 1025–1063.
- Belalia, M. (2016). On the asymptotic properties of the Bernstein estimator of the multivariate distribution function. Statistics and Probability Letters 110, 249–256.
- Belalia, M., T. Bouezmarni, F. C. Lemyre, and A. Taamouti (2017). Testing independence based on Bernstein empirical copula and copula density. *Journal of Nonparametric Statistics* 29(2), 346–380.

| n |  |  |
|---|--|--|
|   |  |  |
| L |  |  |
|   |  |  |
|   |  |  |

- Borisov, I. (1982). Approximation of empirical fields that are constructed from vector observations with dependent coordinates. *Sibirskii Matematicheskii Zhurnal* 23(5), 31–41.
- Bouzebda, S. and N.-E. El Faouzi (2012). New two-sample tests based on the integrated empirical copula processes. *Statistics* 46(3), 313–324.
- Bouzebda, S., N.-E. El Faouzi, and T. Zari (2011). On the multivariate two-sample problem using strong approximations of empirical copula processes. *Communications in Statistics—Theory and Methods* 40(8), 1490–1509.
- Bouzebda, S., A. Keziou, and T. Zari (2011). K-sample problem using strong approximations of empirical copula processes. *Mathematical Methods of Statistics 20*, 14–29.
- Charpentier, A., J.-D. Fermanian, and O. Scaillet (2007). The estimation of copulas : theory and practice. In J. Rank (Ed.), *Copulas: from theory to application in finance*, Chapter 2, pp. 35–64. London: Risk Books.
- Chen, S. X. and T.-M. Huang (2007). Nonparametric estimation of copula functions for dependence modelling. *Canadian Journal of Statistics* 35(2), 265–282.
- Cousido-Rocha, M., J. de Uña-Alvarez, and J. D. Hart (2019). Testing equality of a large number of densities under mixing conditions. *Test 28*, 1203–1228.
- Deheuvels, P. (1981). An asymptotic decomposition for multivariate distribution-free tests of independence. Journal of Multivariate Analysis 11(1), 102–113.
- Derumigny, A., J.-D. Fermanian, and A. Min (2022). Testing for equality between conditional copulas given discretized conditioning events. *Canadian Journal of Statistics*. In press.
- Dupuis, D., E. Jacquier, N. Papageorgiou, and B. Rémillard (2009). Empirical evidence on the dependence of credit default swaps and equity prices. Journal of Futures Markets: Futures, Options, and Other Derivative Products 29(8), 695–712.
- Fermanian, J.-D., D. Radulović, and M. Wegkamp (2004). Weak convergence of empirical copula processes. Bernoulli 10(5), 847–860.
- Genest, C., K. Ghoudi, and L.-P. Rivest (1995). A semiparametric estimation procedure of dependence parameters in multivariate families of distributions. *Biometrika* 82(3), 543–552.
- Genest, C., J. Nešlehová, and J.-F. Quessy (2012). Tests of symmetry for bivariate copulas. Annals of the Institute of Statistical Mathematics 64(4), 811–834.

- Hall, P. and N. Neumeyer (2006). Estimating a bivariate density when there are extra data on one or both components. *Biometrika* 93(2), 439–450.
- Harder, M. and U. Stadtmüller (2017). Testing exchangeability of copulas in arbitrary dimension. *Journal* of Nonparametric Statistics 29(1), 40–60.
- Janssen, P., J. Swanepoel, and N. Veraverbeke (2012). Large sample behavior of the Bernstein copula estimator. *Journal of Statistical Planning and Inference* 142(5), 1189–1197.
- Janssen, P., J. Swanepoel, and N. Veraverbeke (2014). A note on the asymptotic behavior of the Bernstein estimator of the copula density. *Journal of Multivariate Analysis* 124, 480–487.
- Janssen, P., J. Swanepoel, and N. Veraverbeke (2016). Bernstein estimation for a copula derivative with application to conditional distribution and regression functionals. *Test* 25(2), 351–374.
- Jiménez-Gamero, M., M. Cousido-Rocha, M. Alba-Fernández, F. Jiménez-Jiménez, et al. (2022). Testing the equality of a large number of populations. *Test 31*, 1–21.
- Kiriliouk, A., J. Segers, and H. Tsukahara (2021). Resampling procedures with empirical beta copulas. In Pioneering Works on Extreme Value Theory: In Honor of Masaaki Sibuya, pp. 27–53. Springer.
- Kojadinovic, I. (2022). On Stute's representation for a class of smooth, possibly data-adaptive empirical copula processes. arXiv preprint arXiv:2204.11240.
- Kojadinovic, I. and K. Stemikovskaya (2019). Subsampling (weighted smooth) empirical copula processes. Journal of Multivariate Analysis 173, 704–723.
- Kojadinovic, I. and J. Yan (2011). A goodness-of-fit test for multivariate multiparameter copulas based on multiplier central limit theorems. *Statistics and Computing* 21, 17–30.
- Leblanc, A. (2012). On the boundary properties of Bernstein polynomial estimators of density and distribution functions. *Journal of Statistical Planning and Inference* 142(10), 2762–2778.
- Lyu, G. and M. Belalia (2023). Testing symmetry for bivariate copulas using Bernstein polynomials. arXiv preprint arXiv:2202.12787.
- Morettin, P. A., C. M. C. Toloi, C. Chiann, and J. C. S. de Miranda (2010). Wavelet-smoothed empirical copula estimators. *Revista Brasileira de Finanças* 8(3), 263–281.
- Neumann, A., T. Bodnar, D. Pfeifer, and T. Dickhaus (2019). Multivariate multiple test procedures based on nonparametric copula estimation. *Biometrical Journal* 61(1), 40–61.

- Omelka, M., I. Gijbels, and N. Veraverbeke (2009). Improved kernel estimation of copulas: Weak convergence and goodness-of-fit testing. *The Annals of Statistics* 37(5B), 3023 3058.
- Rémillard, B. and O. Scaillet (2009). Testing for equality between two copulas. Journal of Multivariate Analysis 100(3), 377–386.
- Rüschendorf, L. (1976). Asymptotic distributions of multivariate rank order statistics. The Annals of Statistics 4(5), 912–923.
- Sancetta, A. and S. Satchell (2004). The Bernstein copula and its applications to modeling and approximations of multivariate distributions. *Econometric Theory* 20(3), 535–562.
- Scaillet, O. (2005). A Kolmogorov-Smirnov type test for positive quadrant dependence. *Canadian Journal* of Statistics 33(3), 415–427.
- Segers, J. (2012). Asymptotics of empirical copula processes under non-restrictive smoothness assumptions. Bernoulli 18(3), 764–782.
- Segers, J., M. Sibuya, and H. Tsukahara (2017). The empirical beta copula. Journal of Multivariate Analysis 155, 35–51.
- Seo, J. (2021). Randomization tests for equality in dependence structure. Journal of Business & Economic Statistics 39(4), 1026–1037.
- Sklar, A. (1959). Fonctions de répartition à n dimensions et leurs marges. Publications de l'Institut de Statistique de l'Université de Paris 8, 229–231.
- Szolgay, J., L. Gaál, T. Bacigál, S. Kohnová, K. Hlavčová, R. Výleta, J. Parajka, and G. Blöschl (2016). A regional comparative analysis of empirical and theoretical flood peak-volume relationships. J. Hydrol. Hydromech 64(4), 367–381.
- van der Vaart, A. W. and J. A. Wellner (1996). Weak Convergence and Empirical Processes: With Applications to Statistics. New York, NY, USA: Springer.
- Zhan, D. and J. Hart (2014). Testing equality of a large number of densities. *Biometrika* 101(2), 449–464.