Sum rules for light-dressed matter

Bing $\rm{Gu}^{1,2,*}$

¹Department of Chemistry, School of Science, Westlake University, 600 Dunyu Road, Hangzhou 310030, Zhejiang Province, China ²Institute of Natural Sciences, Westlake Institute for Advanced Study, 18 Shilongshan Road, Hangzhou 310024, Zhejiang Province, China

(Dated: March 7, 2023)

Light-driven matter can exhibit qualitatively distinct electronic and optical properties from those observed at equilibrium. We introduce generalized sum rules for the optical properties of driven systems by both quantum and classical light. For classical light, it shows that the sum of all Fourier components, indexed by n , of the time-dependent dipole matrix elements between dressed states weighted by the corresponding quasienergy difference in the first Floquet Brillouin zone plus n driving frequency is a constant, determined by the number of electrons. An analogous sum rule for quantum light-dressing is also derived. These developments provide guidance for the control of effective optical properties of matter by light fields.

I. INTRODUCTION

Matter ranging from atoms, through molecules to solid state materials, driven out of equilibrium by external stimulus, can have emergent electronic and optical properties that are qualitatively different from their equilibrium counterparts [\[1](#page-4-0)[–5](#page-4-1)]. For example, laser-dressed atoms exhibit Autler-Townes effects and Mollow triplets [\[6](#page-4-2)], dressed diatomic molecules contain light-induced conical intersections [\[7\]](#page-4-3). There are a wealth of light-induced phenomena for solid state materials [\[8,](#page-4-4) [9](#page-4-5)] such as Thouless pumping [\[10\]](#page-4-6), modulated optical susceptibility [\[11](#page-4-7), [12](#page-4-8)], and tunable band topology [\[13](#page-4-9)[–15](#page-4-10)].

Optical driving provide a means to tune the

optical properties of matter in an ultrafast femtosecond timescale. A particular interest is to employ far off-resonant drive, that can alleviate the heating induced by resonant pulses [\[11](#page-4-7)]. It has been experimentally demonstrated that ZnO driven by an infrared light shows below gap absorption [\[12](#page-4-8)]. A giant modulation of the secondorder nonlinearity has been observed for a layered magnetic insulator, manganese phosphorus trisulfide $(MnPS₃)$ under strong far off-resonant driving [\[16](#page-5-0)].

Understanding the emergent optical properties of driven matter poses challenges to the current theories that are developed under equilibrium conditions. A general endeavor is to extend the concepts and theories that underlie our standing of equilibrium matter to non-

[∗] gubing@westlake.edu.cn

tical sum rules such as Thomas–Reiche–Kuhn electronic Hamiltonian (TRK) and Bethe sum rules [\[17](#page-5-1)] play significant roles in our understanding of the optical properties of matter owning to its generality to all kinds of systems including atoms, molecules, and strongly interacting systems [\[18,](#page-5-2) [19](#page-5-3)]. The TRK sum rule is also important in the study of superradiant phase transition [\[20\]](#page-5-4) However, no such general rules have been found for dressed materials.

Here we generalize the TRK sum rule to systems driven by external light. Both classical and quantum light are considered. The sum rule for classical light applies to any electronic material under periodic drive, and thus can provide useful insights into designing driven materials with desired effective optical properties. The sum rule for quantum light resembles closer the original TRK sum rule with the joint light-matter states replacing the electronic states. These sum rules provide guidance to the manipulation of effective optical properties of electronic materials by light fields, e.g., in Floquet engineering and cavity quantum electrodynamics [\[21](#page-5-5), [22\]](#page-5-6).

Atomic units $\hbar = e = m_e = 1$ are used throughout.

II. GENERALIZED TRK SUM RULE FOR FLOQUET MATTER

The general Hamiltonian for electronic materials interacting with a classical light field is

equilibrium driven systems. Particularly, op- given by $H(t) = H_M + H_M(t)$, where H_M is the

$$
H_{\rm M} = \sum_{i} \frac{\mathbf{p}_i^2}{2} + V(\mathbf{r}_i) + \sum_{i \neq j} V_{\rm ee}(\mathbf{r}_i, \mathbf{r}_j), \qquad (1)
$$

 \mathbf{p}_i is the momentum operator for *i*-th electron, $V(\mathbf{r}_i)$ the electron-nuclear interaction, and $V_{\text{ee}}(\mathbf{r}_i, \mathbf{r}_j)$ the electron-electron Coulomb interaction, and $H_{LM}(t)$ the light-matter interaction. In the electric-dipole gauge

$$
H_{\text{LM}}(t) = -\mathbf{d} \cdot \mathbf{E}(t),\tag{2}
$$

where $\mathbf{d} = -\sum_i \mathbf{r}_i$ is the dipole operator and $\mathbf{E}(t)$ is the total electric field of the dressing light. The driving can be strong to defy perturbative treatment of the light-matter interaction.

A general framework to understand strongly driven systems is provided by the Floquet theory. The Floquet theory provides not only an intuitive picture to understand driven matter but also an efficient computational method. For periodic Hamiltonian $H(t) = H(t + T)$, according to the Floquet theorem [\[11](#page-4-7), [23](#page-5-7), [24](#page-5-8)], there exits so-called Floquet states

$$
\psi_{\lambda}(\mathbf{r},t) = e^{-i\epsilon_{\lambda}t} \phi_{\lambda}(\mathbf{r},t)
$$
\n(3)

that satisfy the time dependent Schrödinger equation . Here $|\phi_{\lambda}(t)\rangle = |\phi_{\lambda}(t+T)\rangle$ are the many-body Floquet modes, $T = 2\pi/\Omega$ is the driving period with Ω the driving frequency, and ϵ_{λ} are the quasienergies. The Floquet theorem is the analog of the Bloch theorem in the time domain. It applies to a system driven by a laser

with carrier frequency Ω together with any com- which follows directly from the commutation rebination of its harmonics. Inserting Eq. [\(3\)](#page-1-0) into the time dependent Schrödinger equation yields

$$
\Gamma|\phi_{\lambda}\rangle\!\rangle = \epsilon_{\lambda}|\phi_{\lambda}\rangle\!\rangle \tag{4}
$$

where $\Gamma = H(t) - i \frac{\partial}{\partial t}$ is the Floquet Hamiltonian. Equation [\(4\)](#page-2-0) is a generalized eigenvalue equation defined in an extended Hilbert space, a tensor product of the physical Hilbert space and the temporal space consisting of T-periodic functions. In this so-called Sambe space, Γ is Hermitian and can be represented in a basis set $\{ | \chi_{\alpha}\rangle e^{in\Omega t} \}$ for numerical computations, where α runs over electronic states and $n = 0, \pm 1, \cdots$. We use $\ket{\cdot \cdot}$ to denote states in the Sambe space. The electronic operator can be promoted to the extended Hilbert space by $A(\mathbf{r}, \mathbf{p}) \rightarrow A \otimes I$ where I is the identity in time space.

An important relation between the Floquet modes is that if $|\phi_{\lambda}\rangle$ is a Floquet mode with quasienergy ϵ_{λ} , then $e^{-in\Omega t}\phi_{\lambda}(\mathbf{r},t)$ will also be an eigenstate of the Floquet Hamiltonian with quasienergy $\epsilon_{\lambda} + n\hbar\Omega$, i.e., . This can be verified by inserting $e^{-in\Omega t}\phi_{\lambda}(\mathbf{r},t)$ into Eq. [\(4\)](#page-2-0). Therefore, the full spectrum can be understood as the quasienergies uniquely defined in the first Floquet Brillouin zone (FFBZ) $[-\hbar\Omega/2,\hbar\Omega/2]$ and their replicas with energies shifted by an integer number of driving frequencies.

To derive the sum rule, we start from the equality (assume the polarization in the \hat{z} direction)

$$
\[z_i, \left[\Gamma, z_j\right] \] = \delta_{ij} \tag{5}
$$

lation $[z_i, p_z^j] = i\delta_{ij}$. Multiplying both sides of Eq. [\(5\)](#page-2-1) by e^2 and sum over i, j yields

$$
[d_z, [\Gamma, d_z]] = N_e \tag{6}
$$

where $d_z = \sum_i -e z_i$ is the dipole operator in zdirection, $N_{\rm e}$ the number of electrons.

Inserting resolution of identity $I =$ $\sum_{\beta} |\phi_{\beta}\rangle\langle |\phi_{\beta}|$ of the extended space in Eq. [\(6\)](#page-2-2) and taking expectation value of state $|\alpha\rangle$ yields

$$
2\sum_{\beta} \left(\epsilon_{\beta} - \epsilon_{\alpha}\right) |\langle \phi_{\alpha} | d_z | \phi_{\beta} \rangle \rangle|^2 = N_e \tag{7}
$$

where $\epsilon_{\beta\alpha} = \epsilon_{\beta} - \epsilon_{\alpha}$. Equation [\(7\)](#page-2-3) is a sum rule that involves all Floquet states in the Sambe space. However, the number of Floquet states is infinity as $n \in (-\infty, \infty)$. It is more useful to focus on the Floquet states in the FFBZ. To reduce the summation in Eq. [\(7\)](#page-2-3) to states only in the FFBZ, we use the relation

$$
\sum_{\beta} \epsilon_{\beta} = \sum_{\lambda \in \text{ FFBZ}} \sum_{n = -\infty}^{\infty} (\epsilon_{\lambda} + n\hbar \Omega) \tag{8}
$$

Let $\alpha = \lambda'$ be another Floquet mode in the FFBZ, the transition dipole matrix elements reads

$$
\langle \langle \phi_{\lambda'} | d_z | \phi_{\lambda} \rangle \rangle = \sum_n d_{\lambda' \lambda}^{(n)} \tag{9}
$$

where $d_{\lambda'\lambda}^{(n)} = T^{-1} \int_0^T$ $\int_0^T dt \langle\!\langle \phi_{\lambda'}(t)|d_z|\phi_{\lambda}(t)\rangle\!\rangle e^{-in\Omega t}$ is the n-th Fourier component of the timedependent transition dipole between the Floquet modes. Such expansion is allowed due to the periodicity of the Floquet modes.

Using eqs. (8) and (9) into Eq. (7) yields

$$
2\sum_{\lambda \in \text{FFBZ}} \sum_{n=-\infty}^{\infty} \left(\epsilon_{\lambda \lambda'} + n \hbar \Omega\right) \left| d_{\lambda' \lambda}^{(n)} \right|^2 = N_{\text{e}} \qquad (10)
$$

It can also be written as

$$
\int_{-\infty}^{\infty} \frac{d\omega}{2\pi} \omega S(\omega) = N_{\rm e} \tag{11}
$$

where $S(\omega) = 2\pi \sum_{n=-\infty}^{\infty} \delta(\omega - \omega_{\lambda\lambda'} - n\Omega) |d_{\lambda'\lambda}^{(n)}|^2$.

Equation [\(10\)](#page-3-0) is the sum rule for periodically driven materials. It is valid for strongly correlated materials and for strongly driven matter. It reduces to the TKS sum rule (i.e. $\sum_{\beta} |d_{\alpha\beta}|^2 2\epsilon_{\beta\alpha} = N$) in the equilibrium (nondriven) limit. For instance, consider a highfrequency driving where all the pristine eigenstates are contained in the FFBZ. The dressed states become the eigenstates of H_M and the inter-FBZ transitions are forbidden as the transition dipole between eigenstates are constants such that $n = 0$ is the only non-vanishing Fourier component.

The sum rule provides a guideline for the manipulation of the optical absorption and emission properties of matter by lasers. An intuitive way to understand Eq. [\(10\)](#page-3-0) is that any two Floquet modes in the FFBZ can be associated with a set of transitions indicated by an integer number n . While it is tempting to consider an transition with $n \neq 0$ as photon-assisted transitions, this is only meaningful under a high-frequency driving while all pristine eigenstates reside in the FFBZ. the characteristic energy gap of the pristine ma-brid light-matter states replacing bare electronic

terial, even the bare transitions will be associated with a large n as all pristine states folds into a narrow spectral range. We can envision that the bare absorption spectrum of a material be broadened due to the replica structure of the quasienergies.

We now consider the case where the materials is driven by quantum light. Quantum light refers to electromagnetic fields with manifest quantum mechanical properties including entanglement as in entangled photon pairs generated through parametric down-conversion. Coherent laser light is considered to be classical. Quantum light cannot be described by the semiclassical theory but requires a full quantum electrodynamical approach. The full Hamiltonian then reads

$$
\mathcal{H} = \mathcal{H}_0 - \mathbf{d} \cdot \hat{\mathbf{E}} \tag{12}
$$

where E is the electric field operator at the molecule and \mathcal{H}_0 is the noninteracting Hamiltonian of matter and quantized field. Proceeding along the same lines as above, a similar sum rule is obtained

$$
2\sum_{\beta} \epsilon_{\beta \alpha} |\langle \Phi_{\alpha} | d_z | \Phi_{\beta} \rangle|^2 = N_e \tag{13}
$$

In a low frequency driving $\Omega \ll E_g$ where E_g is to the non-driven TRK sum rule with the hywhere $|\Phi_{\alpha}\rangle$, eigenstate of H with eigenenergy ε_{α} , is a quantum light-dressed state, e.g., polariton state under strong light-matter coupling. Here $\langle \cdots \rangle$ refers to the joint light-matter space and β runs over all eigenstates of the joint light-matter Hamiltonian, i.e., \mathcal{H}_0 . Eq. [\(13\)](#page-3-1) is analogous states. It does not have the Floquet quasienergy ing the electronic states with coupled electron-

To summarize, we have derived generalized TRK sum rules for driven matter, valid for strongly driven systems. While we have assumed a classical nuclei in the Hamiltonian, it can be easily generalized to quantized nuclei by replac-

structure as its spectrum is bounded from below. nuclear states. This is due to that the nuclear kinetic energy operator does not change the commutation relation of electronic operators eq. [\(5\)](#page-2-1). Moreover, there can be variants of its form if a different gauge is used and generalizations if a general operator is considered [\[18](#page-5-2)].

- [1] U. D. Giovannini and H. Huebener, Floquet analysis of excitations in materials, J. Phys. Mater. [10.1088/2515-7639/ab387b](https://doi.org/10.1088/2515-7639/ab387b) (2019).
- [2] T. Oka and S. Kitamura, Floquet Engineering of Quantum Materials, [Annu. Rev. Condens. Matter Phys.](https://doi.org/10.1146/annurev-conmatphys-031218-013423) 10, 387 (2019).
- [3] M. Grifoni and P. Hänggi, Driven quantum tunneling, Phys. Rep. 304[, 229 \(1998\).](https://doi.org/10.1016/S0370-1573(98)00022-2)
- [4] C. Bao, P. Tang, D. Sun, and S. Zhou, Light-induced emergent phenomena in 2D materials and topological materials, [Nat Rev Phys](https://doi.org/10.1038/s42254-021-00388-1) 4, 33 (2021).
- [5] S.-I. Chu and D. A. Telnov, Beyond the Floquet theorem: Generalized Floquet formalisms and quasienergy methods for atomic and molecular multiphoton processes in intense laser fields, [Phys. Rep.](https://doi.org/10.1016/j.physrep.2003.10.001) 390, 1 (2004).
- [6] B. R. Mollow, Power Spectrum of Light Scattered by Two-Level Systems, Phys. Rev. 188[, 1969 \(1969\).](https://doi.org/10.1103/PhysRev.188.1969)
- [7] A. Natan, M. R. Ware, V. S. Prabhudesai, U. Lev, B. D. Bruner, O. Heber, and P. H. Bucksbaum, Observation of Quantum Interferences via Light-Induced Conical Intersections in Diatomic Molecules, [Phys. Rev. Lett.](https://doi.org/10.1103/PhysRevLett.116.143004) 116, 143004 (2016).
- [8] K. I. Seetharam, Controlled Population

of Floquet-Bloch States via Coupling to Bose and Fermi Baths, Phys. Rev. X 5, [10.1103/PhysRevX.5.041050](https://doi.org/10.1103/PhysRevX.5.041050) (2015).

- [9] A. Zenesini, H. Lignier, D. Ciampini, O. Morsch, and E. Arimondo, Coherent Control of Dressed Matter Waves, [Phys. Rev. Lett.](https://doi.org/10.1103/PhysRevLett.102.100403) 102, 100403 (2009).
- [10] D. J. Thouless, Quantization of particle transport, [Phys. Rev. B](https://doi.org/10.1103/PhysRevB.27.6083) 27, 6083 (1983).
- [11] B. Gu and I. Franco, Optical absorption properties of laser-driven matter, Phys. Rev. A 98[, 063412 \(2018\).](https://doi.org/10.1103/PhysRevA.98.063412)
- [12] S. Ghimire, A. D. DiChiara, E. Sistrunk, P. Agostini, L. F. DiMauro, and D. A. Reis, Observation of high-order harmonic generation in a bulk crystal, Nat. Phys. 7[, 138 \(2011\).](https://doi.org/10.1038/nphys1847)
- [13] Y. H. Wang, H. Steinberg, P. Jarillo-Herrero, and N. Gedik, Observation of Floquet-Bloch States on the Surface of a Topological Insulator, Science 342[, 453 \(2013\).](https://doi.org/10.1126/science.1239834)
- [14] K. Schwennicke and J. Yuen-Zhou, Optical Activity from the Exciton Aharonov–Bohm Effect: A Floquet Engineering Approach, [J. Phys. Chem. C](https://doi.org/10.1021/acs.jpcc.9b10030) 124, 4206 (2020).
- [15] K. Wintersperger, C. Braun, F. N. Unal, A. Eckardt, M. D. Liberto, N. Goldman, I. Bloch, and M. Aidelsburger, Realization of an

anomalous Floquet topological system with ultracold atoms, Nat. Phys. 16[, 1058 \(2020\).](https://doi.org/10.1038/s41567-020-0949-y)

- [16] J.-Y. Shan, M. Ye, H. Chu, S. Lee, J.-G. Park, L. Balents, and D. Hsieh, Giant modulation of optical nonlinearity by Floquet engineering, Nature 600[, 235 \(2021\).](https://doi.org/10.1038/s41586-021-04051-8)
- [17] H. Bethe, Zur Theorie des Durchgangs schneller Korpuskularstrahlen durch Materie, [Annalen der Physik](https://doi.org/10.1002/andp.19303970303) 397, 325 (1930).
- [18] S. Wang, Generalization of the Thomas-Reiche-Kuhn and the Bethe sum rules, [Phys. Rev. A](https://doi.org/10.1103/PhysRevA.60.262) 60, 262 (1999).
- [19] V. Chernyak and S. Mukamel, Generalized sum rules for optical nonlinearities of many-electron systems, [The Journal of Chemical Physics](https://doi.org/10.1063/1.470283) 103, 7640 (1995).
- [20] K. Rza˙zewski, K. W´odkiewicz, and W. Zakowicz, Phase Transitions, Two-

Level Atoms, and the ${\frac{1}{4}A}{\hat{2}}$ Term, [Phys. Rev. Lett.](https://doi.org/10.1103/PhysRevLett.35.432) 35, 432 (1975).

- [21] K. Wang, M. Seidel, K. Nagarajan, T. Chervy, C. Genet, and T. W. Ebbesen, Large optical nonlinearity enhancement under electronic strong coupling, arXiv:2005.13325 [physics] (2020), [arXiv:2005.13325 \[physics\].](https://arxiv.org/abs/2005.13325)
- [22] F. J. Garcia-Vidal, C. Ciuti, and T. W. Ebbesen, Manipulating matter by strong coupling to vacuum fields, Science 373, [10.1126/science.abd0336](https://doi.org/10.1126/science.abd0336) (2021).
- [23] G. Floquet, Sur les équations différentielles linéaires à coefficients périodiques, Ann. Sci. Éc. Norm. Supér 12, 47 (1883).
- [24] H. Sambe, Steady States and Quasienergies of a Quantum-Mechanical System in an Oscillating Field, [Phys. Rev. A](https://doi.org/10.1103/PhysRevA.7.2203) 7, 2203 (1973).