

Empirical partially Bayes multiple testing and compound χ^2 decisions

Nikolaos Ignatiadis
n.ignatiadis@columbia.edu

Bodhisattva Sen
bodhi@stat.columbia.edu

Draft manuscript: March, 2023

Abstract

We study multiple testing in the normal means problem with estimated variances that are shrunk through empirical Bayes methods. The situation is asymmetric in that a prior is posited for the nuisance parameters (variances) but not the primary parameters (means). If the prior were known, one could proceed by computing p-values conditional on sample variances; a strategy called partially Bayes inference by Sir David Cox. These conditional p-values satisfy a Tweedie-type formula and are approximated at nearly-parametric rates when the prior is estimated by nonparametric maximum likelihood. If the variances are in fact fixed, the approach retains type-I error guarantees.

Keywords: partially Bayes inference, conditionality principle, nonparametric maximum likelihood, Tweedie’s formula, Benjamini-Hochberg, average significance control

1 Introduction

We study an empirical Bayes method for testing n null hypotheses, $H_i : \mu_i = 0$, in the normal means problem with estimated variances. To be concrete, we assume that for the i -th hypothesis we observe a Gaussian measurement $Z_i \sim \mathcal{N}(\mu_i, \sigma_i^2)$ centered around μ_i with variance σ_i^2 , as well as an independent unbiased measurement S_i^2 of σ_i^2 whose law is the scaled χ^2 distribution with $\nu \geq 1$ degrees of freedom:

$$(Z_i, S_i^2) \stackrel{\text{ind}}{\sim} \mathcal{N}(\mu_i, \sigma_i^2) \otimes \frac{\sigma_i^2}{\nu} \chi_\nu^2 \quad \text{for } i = 1, \dots, n. \quad (1)$$

Here $\sigma_i^2 > 0$ and $\mu_i \in \mathbb{R}$ are unknown. Z_i and S_i^2 arise from data summarization in Gaussian linear models. As one concrete example, suppose that for the i -th hypothesis we observe $Y_{ij} \mid \mu_i, \tilde{\sigma}_i^2 \stackrel{\text{iid}}{\sim} \mathcal{N}(\mu_i, \tilde{\sigma}_i^2)$ for $j = 1, \dots, K$. Then $Z_i := \sum_{j=1}^K Y_{ij}/K$ and $S_i^2 := \sum_{j=1}^K (Y_{ij} - Z_i)^2/(K-1)$ satisfy (1) with $\mu_i = \mu_i, \sigma_i^2 = \tilde{\sigma}_i^2/K$ and $\nu = K-1$. Data of the form (1) are ubiquitous and arise naturally in high-throughput biological studies, for example in transcriptomics [Lönstedt, 2005, Smyth, 2004, Ritchie et al., 2015], proteomics [Kammers et al., 2015, Terkelsen et al., 2021, Zheng et al., 2021], and methylomics [Zhang et al., 2013, Maksimovic et al., 2017] (cf. Sections 1.1, 1.2, and 6).

The crux of a standard empirical Bayesian [Robbins, 1956, Efron, 2010] analysis with data from (1) is to further posit that (μ_i, σ_i^2) are i.i.d. draws from an unknown distribution Π and

to then make statistical decisions by imitating the actions of an oracle Bayesian who knows Π . The imitation step can be accomplished by estimating Π based on $(Z_1, S_1^2), \dots, (Z_n, S_n^2)$ as $\hat{\Pi}$ and then pretending that $(\mu_i, \sigma_i^2) \stackrel{\text{iid}}{\sim} \hat{\Pi}$.

This is *not* the empirical Bayes approach we pursue in this paper. Instead, we imitate an oracle Bayesian that is a “*partial Bayesian*” [Cox, 1975, McCullagh, 1990] who posits a distribution G for the nuisance parameters σ_i^2 ,

$$\sigma_i^2 \stackrel{\text{iid}}{\sim} G, \text{ for } i = 1, \dots, n, \quad (2)$$

but treats the parameters of primary interest, μ_i , as fixed and unknown. What’s more, for the partial Bayesian that we seek to imitate the *conditionality principle* is in full force (see explanation in Section 2): their inferences for μ_i would be based on the conditional distribution of $Z_i \mid S_i^2$, for example, a two-sided p-value for testing $H_i : \mu_i = 0$ would be given by $P_i = P_G(Z_i, S_i^2)$, where

$$P_G(z, s^2) := \mathbb{P}_G [|Z^{H_0}| \geq |z| \mid S^2 = s] = \mathbb{E}_G [2(1 - \Phi(|z|/\sigma)) \mid S^2 = s^2]. \quad (3)$$

Above Φ is the standard normal distribution function and we write (σ^2, Z^{H_0}, S^2) for the triple generated as $\sigma^2 \sim G$ (2) and (Z^{H_0}, S^2) generated as in (1) conditional on σ^2 and $\mu = 0$. In a purely frequentist analysis with σ_i^2 treated as fixed, the distribution of Z^{H_0} conditional on S^2 is *not pivotal*, since $Z^{H_0} \mid S^2, \sigma^2 \sim \mathcal{N}(0, \sigma^2)$. However, the partial Bayesian can marginalize over $\sigma^2 \mid S^2$, and so the distribution of $Z^{H_0} \mid S^2$ becomes pivotal. The conditional p-value $P_G(Z, S^2)$ has the following elegant interpretation in view of the right hand side of (3): it is a weighted average of standard z-test based p-values in the case of known variance σ^2 with weighting determined by the posterior of σ^2 conditional on S^2 .

Our empirical Bayes proposal is as follows: in accordance to the empirical Bayes principle, we estimate G by \hat{G} based on S_1^2, \dots, S_n^2 . Then we conduct partially Bayes inference *conditional* on S_1^2, \dots, S_n^2 with the prior \hat{G} . That is, we form the conditional p-values $P_{\hat{G}}(Z_i, S_i^2)$ (3) for $i = 1, \dots, n$. Finally, we apply the method of Benjamini and Hochberg [1995] on these p-values to control the false discovery rate. We call the resulting approach to testing “*empirical partially Bayes multiple hypothesis testing*”— such empirical Bayes developments had been explicitly anticipated by Sir David Cox [1975] (more on this historical connection in Section 2). Algorithm 1 provides a high-level description of our approach.

In this paper, we demonstrate, that the empirical partially Bayes multiple testing approach has substantial upshots compared to alternative approaches for the same problem.

Comparison to standard t-tests: A standard solution to the multiple testing task with data from (1) is to compute p-values P_i^{ttest} based on the t-statistics $T_i = Z_i/S_i$. The latter are pivotal under the null, that is, $T_i \sim t_\nu$ for $\mu = 0$, where t_ν is the t-distribution with ν degrees of freedom. When the degrees of freedom ν are small, as is common in applications (we give concrete examples later), the t-test may have almost no power (due to the heavy tails of the t-distribution with few degrees of freedom). In such cases, the empirical partially Bayes p-values P_i often lead to substantially more powerful tests. For example, if G in (2) is very concentrated around a fixed $\bar{\sigma}^2$, the P_i will be approximately equal to the p-value of the standard z-test with *known* variance (as formalized in Theorem 11 and Example 12 below). Furthermore, in Proposition 4 we show that the conditional type-I error of the standard t-test for small observed value of S_i^2 can be arbitrarily inflated. Such a property is undesirable in view of the conditionality principle; we *do not* want to prioritize a hypothesis that had a small sample variance by chance. The empirical partially Bayes approach avoids the aforementioned caveat since the p-values are computed *conditional* on S_i^2 (3).

Algorithm 1: Empirical partially Bayes multiple hypothesis testing.**Input:** Pairs (Z_i, S_i^2) , $i = 1, \dots, n$.Nominal false discovery rate control level $\alpha \in (0, 1)$.

- 1 Let \widehat{G} be an estimate of the distribution of σ_i^2 based on (S_1^2, \dots, S_n^2) only, e.g., (9).
- 2 Let $P_i = P_{\widehat{G}}(Z_i, S_i^2)$, where $P_G(z, s^2) = \mathbb{E}_G [2(1 - \Phi(|z|/\sigma)) \mid S^2 = s^2]$ is defined in (3) and Φ is the standard normal distribution function.
- 3 Apply the Benjamini-Hochberg procedure with p-values P_1, \dots, P_n at level α as in (14).

Comparison to a full empirical Bayes treatment: As mentioned in the beginning of this paper, an alternative approach for multiple testing with data from (1) treats the problem in a fully empirical Bayes fashion by positing $(\mu_i, \sigma_i^2) \stackrel{\text{iid}}{\sim} \Pi$ and estimating $\widehat{\Pi}$ [Lönstedt and Speed, 2002, Ploner et al., 2006, Hwang and Liu, 2010, Lu and Stephens, 2019, Zheng et al., 2021]. Asymptotically optimal testing procedures [Sun and Cai, 2007] can then be derived by thresholding the (estimated) local false discovery rate $\mathbb{P}_{\widehat{\Pi}}[\mu_i = 0 \mid Z_i, S_i^2]$. The approach we propose has three main advantages: first, it is computationally streamlined—the most intensive step is the solution of a nonparametric maximum likelihood task for a one-dimensional distribution; there are established software solutions for this task [Koenker and Gu, 2017, Kim et al., 2020]. Given such software solutions, Algorithm 1 can be implemented in a few lines of code. In contrast, solving a nonparametric maximum likelihood problem over two dimensions is substantially more difficult [Gu and Koenker, 2017b]—both computationally and statistically—and may require the practitioner to take a stance e.g., on possible independence of σ_i^2 and μ_i (e.g., Zheng et al. [2021], also see Section 5), or to make a semiparametric assumption [Lu and Stephens, 2019]. Second, in Theorem 14* we prove that Algorithm 1 asymptotically controls the false discovery rate even if all the parameters μ_1, \dots, μ_n and $\sigma_1^2, \dots, \sigma_n^2$ are in fact fixed and not random. We are not aware of an analogous result for empirical Bayes approaches that threshold $\mathbb{P}_{\widehat{\Pi}}[\mu_i = 0 \mid Z_i, S_i^2]$.¹ Finally, the empirical partially Bayes approach leads to an intermediate computation of p-values. This can be an advantage for practitioners who may be more comfortable reporting results that are accompanied by p-values [Jeffery et al., 2006].

1.1 Limma: Parametric empirical partially Bayes multiple testing

The empirical partially Bayes multiple testing approach is not new—a parametric instantiation of the framework has found broad usage and success in genomics applications. Lönstedt [2005] and Lönstedt and Speed [2002] proposed a model for detecting differentially expressed genes with replicated microarray data, which was generalized and implemented as the R package `limma` by Smyth [2004]. Limma, in our terminology, is a parametric empirical partially Bayes multiple testing approach that has become the de-facto testing approach for the analysis of microarray data (Smyth [2004] has over 12,000 citations to date), and is also commonly used for the analysis of RNA-Seq data with over 19,000 citations in Ritchie et al. [2015]. Limma has also been successfully applied in many other areas, e.g., for the analysis of methylation data [Maksimovic et al., 2017], in proteomics [Kammers et al., 2015,

¹In recent breakthrough work, Castillo and Roquain [2020] derive such guarantees when σ_i^2 is known to the analyst and equal to 1 for all i . It is plausible—but far from obvious—that such results could be extended to the setting we consider wherein the variances are heteroscedastic and unknown.

Terkelsen et al., 2021], and genetic interaction screening [Laufer et al., 2013].

The model underlying limma specifies G in (2) parametrically as

$$\frac{1}{\sigma_i^2} \stackrel{\text{iid}}{\sim} \frac{1}{\nu_0 s_0^2} \chi_{\nu_0}^2, \quad \text{for } i = 1, \dots, n, \quad (4)$$

where s_0^2, ν_0 are estimated in an empirical Bayes fashion as $\hat{s}_0^2, \hat{\nu}_0$, e.g., by the method of moments [Smyth, 2004] or maximum marginal likelihood [Wright and Simon, 2003, de Menezes et al., 2004]. The p-value for the i -th hypothesis is defined as $P_i = P_{\hat{\nu}_0, \hat{s}_0^2}^{\text{limma}}(Z_i, S_i)$, where

$$P_{\nu_0, s_0^2}^{\text{limma}}(z, s^2) := 2\bar{F}_{t, \nu_0 + \nu}(|\tilde{t}|), \quad \tilde{t} = \frac{z}{\tilde{s}}, \quad \tilde{s}^2 = \frac{\nu_0 s_0^2 + \nu s^2}{\nu_0 + \nu}, \quad (5)$$

and $\bar{F}_{t, \nu_0 + \nu}$ is the survival function of the t -distribution with $\nu_0 + \nu$ degrees of freedom. The formal argument justifying the limma p-values is the following: when (Z_i, S_i^2) are sampled according to (1) with $\mu_i = 0$ and we marginalize over $\sigma_i^2 \sim G$ (4), then \tilde{T}_i , i.e., the random variable with realization \tilde{t} in (5), follows the t -distribution with $\nu_0 + \nu$ degrees of freedom (see Proposition 2 below).

In Section 2 we will see that limma is a special case of the general approach set forth in Algorithm 1. The test statistic underlying limma, \tilde{T}_i , is called a “moderated” t -statistic. While the standard t -statistic $T_i = Z_i/S_i$ is studentized by S_i^2 , the moderated t -statistic is studentized by \tilde{S}_i^2 , a convex combination of S_i^2 and s_0^2 . At a heuristic level, the moderated t -statistic leads to an increase in power compared to the standard t -statistic because p-values are computed with respect to the t -distribution with $\nu + \nu_0$ degrees of freedom (rather than ν , as in the standard t -test). The increase in degrees of freedom may be attributed to the additional information we retrieve for each hypothesis by “borrowing” information across $i = 1, \dots, n$. Furthermore, excessively small values of S_i^2 are pulled upward toward s_0^2 , and so a hypothesis with very small S_i^2 is less likely to be rejected (compared to the standard t -test).

In this paper we go beyond the parametric assumption for the distribution of σ_i^2 in (4) and instead propose to estimate G (recall (2)) by nonparametric maximum likelihood (NPMLE) [Kiefer and Wolfowitz, 1956] (see Section 3 and (9)). Methodologically we thus provide an alternative in cases wherein the parametric assumption is unlikely to hold. From a theoretical perspective, in Theorem 7 we study the statistical properties of the NPMLE with scaled χ^2 -data and derive parametric rates (up to logarithmic factors) of Hellinger consistency for the marginal density that are analogous to related results for the NPMLE with Gaussian data [Ghosal and van der Vaart, 2001, Zhang, 2009, Soloff et al., 2021], or Poisson data [Jana et al., 2022]. Furthermore, our results put forth a theoretical foundation for understanding empirical partially Bayes multiple testing methods and their success in applications.

In Proposition 9 we demonstrate that the oracle partially Bayes p-values satisfy a Tweedie-type formula [Efron, 2011]. The Tweedie-type formula along with Theorem 7 in turn enable us to prove (Theorem 11) that the oracle partially Bayes p-values can be approximated via empirical Bayes at parametric rates up to logarithmic factors, even when G in (2) is specified fully nonparametrically. In Theorem 14*, we prove the remarkable property that our proposed method can asymptotically control the false discovery rate even when the σ_i^2 are in fact fixed (and not random). Our formal results are corroborated by the data analyses in Sections 1.2 and 6, as well as the simulation study in Section 7.

1.2 A demonstration with high-throughput methylation data

Before delving into a more formal treatment, we provide a high-level overview of the methodological developments in this paper through a concrete data analysis. Zhang et al. [2013] compared two types of immune cells, naïve T-cells and antigen-activated naïve T-cells, in terms of DNA methylation, a heritable epigenetic mark. We refer to Zhang et al. [2013] and Maksimovic et al. [2017] for details on the dataset and the analysis; what is important for our analysis is the following: after data preprocessing and summarization, the statistical task amounts to a large scale multiple testing problem with 439,918 hypotheses (one for each methylation probe) based on data that may be assumed to be drawn from (1) with $\nu = 4$ degrees of freedom.²

Fig. 1a) shows the histogram of the sample variances S_i^2 , as well as the implied marginal densities of S_i^2 (marginalizing over (2)) for two choices of $G = \hat{G}$: the nonparametric maximum likelihood estimate (NPMLE; more of which in Section 3.1), and limma’s parametric prior ($\hat{\nu}_0 \approx 3.96$ and $\hat{s}_0^2 \approx 0.037$ in (4)). The NPMLE leads to a slightly better fit at the mode of the histogram. Fig. 1b) shows the two estimated priors \hat{G} ; the NPMLE is discrete. We note that the two priors are qualitatively quite different, yet the implied marginal densities for S_i^2 in Fig. 1a) are quite similar.

Finally, Fig. 1c) allows us to demonstrate the heart of the matter: it shows a 2d histogram of all the (Z_i, S_i^2) pairs as well as the rejection thresholds of different methods. There are three dashed lines; one corresponding to standard t-test p-values, and the other two to the NPMLE and parametric (limma) partially Bayes p-values. For each value of S_i^2 , the dashed rejection threshold demarcates the value of z which would be such that the corresponding p-value would be equal to 0.05; in each case, hypotheses with $Z_i > 0$ (resp. $Z_i < 0$) above (resp. below) the corresponding rejection threshold are rejected. The solid lines demarcate the rejection thresholds of the different methods after adjusting for multiple testing with the Benjamini-Hochberg procedure with a nominal false discovery rate control of $\alpha = 0.05$.

With the Benjamini-Hochberg correction, the NPMLE-based method leads to 986 discoveries, compared to 627 with the parametric limma prior, and to only 1 discovery with the standard t-test. In this case the empirical partially Bayes methods lead to a substantial increase in discoveries. Fig. 1c) provides some insight into the mechanism. The (solid) rejection threshold corresponding to the t-test is a lot more liberal for small values of S_i^2 (that is: closer to 0) compared to the empirical partially Bayes methods, but quickly becomes a lot more conservative as S_i^2 increases.³ The single discovery made with the t-test has sample variance $S_i^2 = 4 \cdot 10^{-5}$, which is the second smallest sample variance in the whole dataset, and this hypothesis is *not* rejected by the empirical partially Bayes methods. At a heuristic level, the empirical Bayes approach shrinks this outlying sample variance upwards, so that this hypothesis is not rejected. On the other hand, for the bulk of the hypotheses, the empirical Bayes approach leads to more accurate estimate of the sample variance, which in turn leads to a more liberal rejection threshold and thus to more power overall. In this problem the NPMLE makes more discoveries than limma. In general this will not be true,

²Indeed, the analysis in Zhang et al. [2013] and Maksimovic et al. [2017] uses limma, and posits the sampling model (1), as well as the parametric assumption (4) on the distribution of σ_i^2 .

³The reader may at this point wonder why the difference is less striking for the dashed thresholds that correspond to p-values ≤ 0.05 compared to the solid thresholds (Benjamini-Hochberg rejection thresholds). One reason is that the Benjamini-Hochberg procedure adapts to the signal in the p-values, and so, the implied rejection threshold in terms of p-values varies for the different methods. To be concrete, the data-driven p-value thresholds (for the different methods) determined by the Benjamini-Hochberg procedure are equal to: $P_i^{\text{NPMLE}} \lesssim 10^{-4}$, $P_i^{\text{limma}} \lesssim 0.7 \cdot 10^{-4}$, and $P_i^{\text{ttest}} \lesssim 10^{-7}$.




Figure 1: **Partially empirical Bayes analysis of methylation data of Zhang et al. [2013]**: We are testing $n = 439,918$ hypotheses $H_i : \mu_i = 0$ based on summary data as in (1) with $\nu = 4$ degrees of freedom. **a)** Histogram of the sample variances S_i^2 , as well as the implied marginal densities of S_i^2 (marginalizing over (2)) for two choices of $G = \widehat{G}$: the nonparametric maximum likelihood estimate (NPMLE), and limma’s parametric prior. **b)** The two estimated priors \widehat{G} . For limma we plot the Lebesgue density of the prior. The NPMLE is a discrete measure; we have rescaled the mass of the support points by a multiplicative constant for visualization purposes. **c)** Two dimensional histogram of (S_i^2, Z_i) pairs with the x -axis logarithmically scaled. The number of hypotheses in each histogram bin is indicated by color (color legend on the right) wherein the denser regions have darker green color. The range of histogram counts is wide and any non-empty bin (even with a single hypothesis) is colored in green. The plot also shows the rejection thresholds of 3×2 different methods; in each case any hypothesis with Z_i above the upper line or below the lower line will be rejected. The different methods correspond to three constructions of p-values (empirical partially Bayes p-values with NPMLE, with limma, and standard t-test p-values) combined with two different multiple testing procedures: reject any hypothesis with p-value ≤ 0.05 (dashed lines, unadjusted for multiple testing) and reject any hypothesis with Benjamini-Hochberg (BH) adjusted p-value ≤ 0.05 (solid lines).




Figure 2: **P-value histograms in the analysis methylation data of Zhang et al. [2013]**: The first row shows t-test p-values, while the second rows shows the empirical partially Bayes p-values based on the NPMLE. The columns correspond to three different subsets of hypotheses: the first column shows all hypotheses, the second column only retains hypotheses with sample variance S_i^2 in the bottom 10% percentile, and the third column retains hypotheses with sample variance S_i^2 in the top 10% percentile.

and the main benefit of the NPMLE is that it has theoretical type-I error guarantees also when the parametric model for G in (4) does not hold.

We now discuss the conditional properties of procedures that we alluded to earlier. Fig. 2 shows the histogram of the t-test p-values, as well as the NPMLE-based partially empirical Bayes p-values. Three histograms are shown for each method: a histogram of all hypotheses, a histogram of hypotheses with sample variance S_i^2 in the bottom 10% percentile, as well as a histogram of hypotheses with sample variance S_i^2 in the top 10% percentile. We observe that for large sample variances, the t-test leads to a histogram that is superuniform with almost no p-values ≤ 0.05 . The t-test p-values are enriched for smaller p-values when only retaining hypotheses with small sample variance. In contrast, the empirical partially Bayes p-values appear to have better conditional properties. We explain these empirical findings theoretically in Proposition 4 which shows that the type-I error of the t-test may be arbitrarily inflated when the sample variance is small, while Proposition 13 demonstrates that the NPMLE p-values are asymptotically uniform conditional on S_i^2 .

1.3 Related work

Our work builds on a rich literature on multiple testing that was motivated by the microarray technology [Tusher et al., 2001, Efron et al., 2001, Baldi and Long, 2001, Newton et al., 2001, Cui and Churchill, 2003, Wright and Simon, 2003, de Menezes et al., 2004, Newton et al., 2004, Delmar et al., 2005, Cui et al., 2005, Ploner et al., 2006, Witten and Tibshirani, 2007]. As mentioned earlier, the specific model we elaborate upon was originally developed by Lönnstedt [2005], Lönnstedt and Speed [2002] and turned into a widely used “empirical

partially Bayes” testing methodology by [Smyth \[2004\]](#). [Ignatiadis et al. \[2022\]](#) consider testing in the limma model of [\[Smyth, 2004\]](#) with e-values instead of p-values.

Closest to our paper, [Lu and Stephens \[2016\]](#) consider a nonparametric generalization of the limma method based on a class of unimodal priors. We consider an even broader nonparametric assumption and lay out the theoretical foundations and framework that justify our proposal.

When testing for $\mu_i = 0$, uncertainty in σ_i is often ignored by assuming that σ_i is known exactly [\[Sun and McLain, 2012, Stephens, 2016, Gu and Shen, 2018, Fu et al., 2022\]](#). This can be a good approximation in some cases, e.g., when σ_i^2 is estimated by S_i^2 and the degrees of freedom ν are large. In this paper we are mostly interested in the situation wherein the degrees of freedom ν are small; such settings benefit the most from the empirical partially Bayes methodology that we outline in this paper.

Going beyond the goal of multiple testing, several authors have studied the empirical Bayes problem with sample variances S_i^2 or joint draws of (Z_i, S_i^2) as in (1). [Yu and Hoff \[2018\]](#) develop confidence intervals for μ_i with finite-sample frequentist coverage that borrow strength from $(Z_j, S_j^2)_{j \neq i}$. A further strand of the literature considers shrinkage estimation of e.g., μ_i , σ_i^2 , or $1/\sigma_i^2$ [\[Robbins, 1982, Muralidharan, 2012, Gu and Koenker, 2017a,b, Banerjee, Fu, James, and Sun, 2021, Kwon and Zhao, 2022\]](#). When $\nu = 2$, the mixture model we consider for S_i^2 is a mixture of exponential distributions; see e.g., [Jewell \[1982\]](#) and [Polyanskiy and Wu \[2020\]](#) for some results for exponential mixtures.

Finally, the empirical partially Bayes approach was introduced in the setting wherein there is a single primary parameter of interest and many nuisance parameters [\[Lindsay, 1985, Zhao et al., 2019\]](#); a prior is posited (and estimated by empirical Bayes) only for the nuisance parameters but not the primary parameter. Here we use the same terminology and study the empirical partially Bayes approach when we are facing a simultaneous inference task with many parameters of primary interest, as well as many nuisance parameters.

2 Oracle partially Bayes hypothesis testing

“It would be possible to develop this investigation in much more detail, but in the absence of a specific application this will not be done”

— Sir David Roxbee Cox, [1975](#)

We first consider the setting in which the random variance assumption (2) holds for a known prior G . To be concrete, we assume that we observe (Z_i, S_i^2) drawn from the following model,

$$\sigma_i^2 \stackrel{\text{iid}}{\sim} G, \quad (Z_i, S_i^2) \mid \sigma_i^2 \stackrel{\text{ind}}{\sim} \mathcal{N}(\mu_i, \sigma_i^2) \otimes \frac{\sigma_i^2}{\nu} \chi_\nu^2 \quad \text{for } i = 1, \dots, n, \quad (6)$$

where G is known and μ_i is unknown. Later, we will drop the assumptions that G is known (Section 3) and that σ_i^2 are random (Section 4). We study (6) to answer the question: *how* would an oracle analyst that knows (6) test the hypothesis $H_i : \mu_i = 0$? Our nonparametric empirical Bayes approach will then seek to mimic that oracle analyst. In the rest of this section we drop the subscript i since we study only a single draw from (6).

We start with the following characterization of (Z, S^2) in (6), see Supplement C.1 for a proof of the result.

Proposition 1. Suppose that G in (6) is not degenerate, that is, the support of G consists of 2 points or more and $G((0, \infty)) = 1$. Then (Z, S^2) is the minimal sufficient statistic⁴ for μ in (6). Furthermore, S^2 is ancillary for μ , that is, its distribution does not depend on μ .

The implication of the proposition is the following: when conducting inference for μ , we need to use both Z and S^2 even though S^2 does not contain any information about μ . The conditionality principle is in full force [Cox and Hinkley, 1974, Chapter 2.3] and any inference for μ should be based on the conditional distribution of Z given S^2 .⁵ Hence, an oracle analyst that knows (6) may use the conditional p-value $P = P_G(Z, S^2)$ defined in (3) to test $H : \mu = 0$.

Now, what if G has the parametric form (4)? It turns out that then the limma p-value $P_{\nu_0, s_0^2}^{\text{limma}}(z, s^2)$ is identical to the conditional p-value (3). This is a consequence of the following result (proved in Supplement C.2).

Proposition 2. Let (Z, S^2) be drawn from (6) with G equal to the distribution of σ^2 in (4), i.e., the inverse scaled chi-square distribution with parameters ν_0, s_0^2 . Then:

$$Z \mid (S^2 = s^2) \stackrel{\mathcal{D}}{=} (\mu + \tilde{s}T_{\nu_0+\nu}), \quad (7)$$

where $T_{\nu_0+\nu}$ is a random variable following the t-distribution with $\nu_0 + \nu$ degrees of freedom and \tilde{s} is as in (5). It follows that for such G , $P_G(z, s^2) = P_{\nu_0, s_0^2}^{\text{limma}}(z, s^2)$.

The above result gives an alternative interpretation of the limma p-value showing that it is exactly the same as the conditional p-value, which is the right thing to look at given the conditionality principle.

The claim of Proposition 2 is known, e.g., it is mentioned in Lu and Stephens [2016], yet its implications are far-reaching and of historical importance: Sir David Cox [1975] considered model (6) with G as an inverse scaled chi-square distribution (4) and concluded that estimation/inference for μ should be based on (7), that is, the conditional distribution of Z given S^2 . He called the resulting inference “*partially Bayes*”; the nuisance parameter is randomly distributed according to a prior, while the primary parameter of interest is fixed. What’s more, Sir David Cox anticipated that the above model would be very fruitful in an empirical Bayes analysis [Cox, 1975, “If the parameters of the prior distribution are estimated from a large amount of data”]. We do not believe that this contribution of Sir David Cox is well-known.⁶ We find it a source of great inspiration that yet another idea of Sir David Cox had ground-breaking impact in applied statistical work, in this case through the idea’s wide adoption in high-throughput biology via the limma software [Smyth, 2004].

The following proposition states three properties of the conditional p-values $P_G(Z, S^2)$; see Supplement C.3 for a proof.

Proposition 3. Let (Z, S^2) be drawn from model (6) with $\mu = 0$. Then, $P = P_G(Z, S^2)$ has the following properties.

- a. Conditional uniformity: $\mathbb{P}_G [P \leq t \mid S^2] = t$ for all $t \in [0, 1]$ almost surely.

⁴However, (Z, S^2) is *not* complete for μ .

⁵Suppose G is degenerate, i.e., $G(\{\bar{\sigma}\}) = 1$ for some $\bar{\sigma} > 0$. Then the conclusion of Proposition 1 does not hold and Z (rather than (Z, S^2)) is minimal sufficient for μ . In that case Z is independent of S^2 , and so inference based on the conditional distribution of Z given S^2 is identical to inference based on Z .

⁶Cox [1975] has less than 40 citations to date. But this is not for lack of importance: this work is included in the collection of selected statistical papers of Sir David Cox [Cox, 2005] and it appears to have been a source of inspiration in the development of predictive p-values by Meng [1994].

- b. Unconditional uniformity: $\mathbb{P}_G [P \leq t] = t$ for all $t \in [0, 1]$.
- c. Monotonicity: $P_G(z, s^2)$ is non-increasing in $|z|$ for any fixed value of $s^2 > 0$, and non-decreasing in $s^2 > 0$ for any fixed value of $|z|$.

The first property follows from the definition in (3) and the probability integral transform applied conditional on S^2 . Such conditional validity is desirable (cf. Cox and Hinkley [1974, Chapter 2.3] and Lehmann and Romano [2005, Chapter 10]), and may not be taken for granted. For example, this property does not hold for the standard t-test as we will prove formally in Proposition 4 below; we also refer the reader to the data analysis of Section 1.2 (in particular, Figure 2) that showcases empirical ramifications of the standard t-test’s lack of conditional type-I error control. The second property (unconditional uniformity) follows from the conditional uniformity result (and iterated expectation). Finally, the monotonicity posits a “natural” requirement for a testing procedure based on (Z, S^2) : the larger the value of $|Z|$ is, the stronger the evidence, and for a fixed value of $|Z|$ the evidence is stronger when S^2 is smaller. Empirical Bayes procedure often satisfy such natural monotonicity requirements, e.g., van Houwelingen [1976], Koenker and Mizera [2014], but not always [Gu and Koenker, 2023].

We contrast the results from Proposition 3 with the following result for the standard t-test (see Supplement C.4 for a proof):

Proposition 4. Consider the t-test p-value $P = P^{\text{ttest}}(Z, S^2)$ with $P^{\text{ttest}}(z, s^2) = 2\bar{F}_{t,\nu}(|z|/s)$ and $\bar{F}_{t,\nu}$ is the survival function of the t-distribution with ν -degrees of freedom. Let (Z, S^2) be drawn from model (6) with $\mu = 0$. Then, P is unconditionally uniform and satisfies the monotonicity property in Proposition 3. On the other hand, if $\nu \geq 2$ and $\mathbb{E}_G [\sigma^{-\nu}] < \infty$, then:

$$\lim_{\delta \searrow 0} \mathbb{P}_G [P^{\text{ttest}}(Z, S^2) \leq t \mid S^2 \leq \delta] = 1 \text{ for all } t \in (0, 1].$$

Hence the standard t-test p-value does not satisfy the conditional uniformity property.⁷ If we observe S^2 that is unusually small, then Proposition 4 implies that the conditional level of the t-test will be inflated and the resulting inference untrustworthy. The concern about poor conditional behavior, however, is not actionable with a single hypothesis and without prior information for σ^2 : without prior knowledge about plausible values of σ^2 , it is impossible to know whether S^2 is unusually small. The partially Bayes model (6) provides knowledge of plausible values of σ^2 , since $\sigma^2 \sim G$. Perhaps more interestingly, such knowledge is also provided in the empirical Bayes setting of the following section wherein we have access to measurements S_1^2, \dots, S_n^2 of many related variances $\sigma_1^2, \dots, \sigma_n^2$. Hence the setting we consider provides a further demonstration of the value of large scale data [Efron, 2010] with new opportunities that were not available previously.

3 Empirical partially Bayes multiple hypothesis testing

“This formula for the mean error depends only on the observed distribution”
— Sir Frank Dyson, 1926

We now turn to the practically relevant case wherein we have n independent draws of (Z_i, S_i^2) from (6) with G and μ_i unknown. Our goal is to mimic the oracle analyst who

⁷In contrast, by Proposition 3, the oracle partially Bayes p-value P satisfies $\mathbb{P}_G [P \leq t \mid S^2 \leq \delta] = t$ for all $t \in [0, 1]$ and $\delta > 0$.

would test $H_i : \mu_i = 0$ with the conditional p-value (3). The strategy we pursue for this task is a nonparametric generalization of the limma approach. First, we estimate \widehat{G} with the nonparametric maximum likelihood estimator (NPMLE) of Robbins [1950] and Kiefer and Wolfowitz [1956] based on (S_1^2, \dots, S_n^2) ,

$$\sigma_i^2 \stackrel{\text{iid}}{\sim} G, \quad S_i^2 \mid \sigma_i^2 \stackrel{\text{ind}}{\sim} \frac{\sigma_i^2}{\nu} \chi_\nu^2 \quad \text{for } i = 1, \dots, n, \quad (8)$$

and then we use the plug-in p-values $P_{\widehat{G}}(Z_i, S_i^2)$ in place of $P_G(Z_i, S_i^2)$. We call this approach *empirical partially Bayes hypothesis testing*; a similar nonparametric extension of limma was also considered by Lu and Stephens [2016] based on a nonparametric class of unimodal priors.

3.1 Nonparametric maximum likelihood estimation for χ^2 data

Our proposed estimator of G based on (S_1^2, \dots, S_n^2) from (8) is defined as a solution to the following optimization problem:

$$\widehat{G} \in \operatorname{argmax} \left\{ \sum_{i=1}^n \log(f_G(S_i^2)) : G \text{ distribution supported on } (0, +\infty) \right\}. \quad (9)$$

In (9), we use the notation $f_G(\cdot)$ to denote the marginal density of S_i^2 when $\sigma_i^2 \sim G$ (for any distribution G supported on $(0, \infty)$), and $S_i^2 \mid \sigma_i^2$ follows (8). More formally, for $s^2 \geq 0$:

$$f_G(s^2) \equiv f_G(s^2; \nu) := \int_0^\infty p(s^2 \mid \sigma^2, \nu) dG(\sigma^2), \quad (10)$$

where $p(s^2 \mid \sigma^2, \nu)$ is the (scaled) χ_ν^2 -density which is equal to,

$$p(s^2 \mid \sigma^2) \equiv p(s^2 \mid \sigma^2, \nu) := \frac{\nu^{\nu/2}}{(\sigma^2)^{\nu/2} 2^{\nu/2} \Gamma(\nu/2)} (s^2)^{\nu/2-1} \exp\left(-\frac{\nu s^2}{2\sigma^2}\right). \quad (11)$$

For both (10) and (11) we typically suppress the dependence on ν . Thus (9) seeks the maximizer of the log marginal likelihood over all possible choices of the prior G .

The following proposition establishes a few formal properties of the optimization problem (9) by following the arguments of e.g., Lindsay [1983] and Jewell [1982]; see Supplement D.1 for a proof.

Proposition 5 (Properties of NPMLE optimization). Suppose that $S_i^2 \in (0, \infty)$ for all i , $S_i^2 \neq S_j^2$ for all $i \neq j$ and that $\nu \geq 2$. Then there exists a unique maximizer \widehat{G} of (9). Further, \widehat{G} is a discrete probability measure with at most n points of support, all lying in the interval $[\min_i \{S_i^2\}, \max_i \{S_i^2\}]$.

As a consequence of the proposition, the NPMLE may be written as $\widehat{G} = \sum_{j=1}^{\widehat{K}} \delta_{\hat{\sigma}_j^2}$, where δ_{σ^2} is a Dirac point mass at σ^2 . The number of components \widehat{K} and the locations $\hat{\sigma}_j^2$ are determined in a fully data-driven way. It is worthwhile to compare this result to the method of Delmar et al. [2005] who assumed a clustering structure for the variances σ_i^2 and posited a prior of the form $G = \sum_{j=1}^K \delta_{\sigma_j^2}$ for a fixed value of K . For the NPMLE, the number of components is automatically learned, and furthermore, as we will see below, the NPMLE performs well for the downstream task of empirical partially Bayes testing even if no such clustering structure is present.

Remark 6 (Computation). To solve (9) we use the interior point convex programming solver MOSEK [MOSEK ApS, 2020] after discretization of (9) as proposed by Koenker and Mizera [2014]. To be concrete, we discretize as follows: we let a , resp. b , be the 1% quantile,⁸ resp. maximum, of $\{S_1^2, \dots, S_n^2\}$ and then we consider a logarithmically equispaced grid between a and b with 300 points. We maximize the marginal likelihood over all distributions supported on the aforementioned grid. Our theoretical results below do not take into account the discretization; we refer to Dicker and Zhao [2016] and Soloff et al. [2021] for analyses of the NPMLE that also consider the discretization error.

As is common in the literature [Ghosal and van der Vaart, 2001, Zhang, 2009], we study the statistical properties of \widehat{G} in terms of the Hellinger distance between f_G and $f_{\widehat{G}}$. The squared Hellinger distance between two densities f and h on $(0, \infty)$ is defined as follows:

$$\mathfrak{H}^2(f, h) = \frac{1}{2} \int_0^\infty \left(\sqrt{f(t)} - \sqrt{h(t)} \right)^2 dt. \quad (12)$$

In Supplement D.2 we prove the following result.

Theorem 7 (Convergence in Hellinger distance). Let $0 < \underline{L} \leq L \leq U \leq \bar{U} < \infty$. Suppose $\nu \geq 2$, $\sigma_i^2 \in [L, U]$ almost surely and that we compute the NPMLE \widehat{G} (9) under the additional constraint $G([L, \bar{U}]) = 1$. Then, there exist constants $C = C(\nu, L, \bar{U}) > 0$ and $c = c(\nu, L, \bar{U}) > 0$ such that:

$$\mathbb{P}_G \left[\mathfrak{H}(f_{\widehat{G}}, f_G) \geq C \frac{\log n}{\sqrt{n}} \right] \leq \exp(-c(\log n)^2) \text{ for all } n \in \mathbb{N}_{\geq 1}.$$

We comment on the requirement of compact support bounded away from 0 and ∞ . As we explained in Footnote 8, we view the lower bound (L, \underline{L}) as a substantive requirement: it is a conservative choice that protects us from spuriously small p-values for hypotheses with sample variance $S_i^2 \approx 0$. The upper bound (U, \bar{U}) is a technical assumption that streamlines our proofs; it could be relaxed to a light tail assumption with a more careful analysis as in Zhang [2009].

According to Theorem 7, estimation of f_G in Hellinger distance is a relatively easy statistical task that may be achieved at the parametric rate $1/\sqrt{n}$ up to a logarithmic factor.

With Theorem 7 in hand and identifiability results of Teicher [1961], we also prove the following in Supplement D.4.

Corollary 8. In the setting of Theorem 7, it holds with probability 1 that:

$$\widehat{G} \overset{d}{\rightsquigarrow} G \text{ as } n \rightarrow \infty.$$

Here $\overset{d}{\rightsquigarrow}$ denotes weak convergence.

⁸ The choice of the 1% quantile for the lower end of the support of the estimated prior is motivated by the following considerations: on one hand, it is a conservative choice that protects us from spuriously small p-values. The choice implies that the conditional p-values we compute in (3) will be at least as large as $2(1 - \Phi(|Z|/S_{0.01}))$, where $S_{0.01}^2$ is the 1% quantile of the sample variances. Second, Lönnerstedt and Speed [2002] consider filtering out hypotheses with variances in the bottom 1% as a simple solution to avoid an excessive number of rejections of hypotheses with small sample variance.

We do not pursue the rate of the above convergence for two reasons: first, as in related deconvolution problems, e.g., with additive Gaussian noise, minimax convergence rates can be very slow, that is, polynomial in $1/\log n$ [Zhang, 1990, Fan, 1991]. Belomestny and Goldenshluger [2020, paragraph after Remark 5] suggest that such slow rates also hold in the deconvolution problem with *multiplicative* noise (8) that we are studying. Second and more importantly, in what follows we will establish the (perhaps surprising) result that the statistical difficulty of the empirical partially Bayes hypothesis testing problem follows from the fast rates in Theorem 7 and is not impacted by slow rate in the convergence of \widehat{G} to G .

3.2 A Tweedie-type formula for conditional p-values

As mentioned above, in empirical Bayes problems it is common that the prior G can only be estimated at very slow rates, while the marginal density f_G can be estimated at faster rates. The statistical difficulty of an empirical Bayes analysis, that is, the difficulty in imitating the oracle Bayesian, can vary substantially across possible models and statistical tasks, see e.g., Efron [2014], Ignatiadis and Wager [2022]. In the worst case, the empirical Bayes problem may be as difficult as the task of estimating G .

However, for certain empirical Bayes problems, the action of the oracle Bayesian may be described as a benign functional of $f_G(\cdot)$; in that case the task of the empirical Bayesian is easier. A concrete example of such a situation occurs when the empirical Bayesian seeks mean squared error optimal estimation of the natural parameter in an exponential family. For example, the distribution of the sample variances S_i^2 in (8) forms an exponential family with natural parameter $1/\sigma_i^2$. When $\sigma_i^2 \sim G$ (2), the optimal estimator of $1/\sigma_i^2$ under squared error loss, is the posterior mean $\mathbb{E}_G [1/\sigma_i^2 | S_i^2]$ which may be expressed in terms of the marginal density f_G of S_i^2 as follows:

$$\mathbb{E}_G \left[\frac{1}{\sigma_i^2} \mid S_i^2 \right] = \frac{\nu - 2}{\nu} \frac{1}{S_i^2} - \frac{2}{\nu} \frac{f'_G(S_i^2)}{f_G(S_i^2)}. \quad (13)$$

The above formula is called Tweedie’s formula [Efron, 2011] and extends to other exponential families, e.g., in the homoskedastic Gaussian case an analogous formula was known by Dyson [1926].

There are further settings where analogous “ F -formulas” hold, see e.g., Robbins [1982], Cressie [1982], Du and Hu [2022], Kwon and Zhao [2022]. Our next result demonstrates that such a formula also exists for the conditional p-values (3); see Supplement E.1 for a proof.

Proposition 9. The conditional p-value (3) may be expressed as

$$P_G(z, s^2) = C(\nu) \frac{(s^2)^{\nu/2-1}}{f_G(s^2; \nu)} \int_0^\infty \mathbf{1} \left(t^2 \geq \frac{\nu s^2 + z^2}{\nu + 1} \right) \frac{(t^2)^{-(\nu-1)/2}}{\sqrt{(\nu+1)t^2 - \nu s^2}} f_G(t^2; \nu+1) d(t^2),$$

where $C(\nu) := \{(1 + 1/\nu)^{-\nu/2} \Gamma((\nu + 1)/2)\} / \{\sqrt{\pi}(\nu + 1)^{-1/2} \Gamma(\nu/2)\}$.

In words, the p-value function $P_G(\cdot)$ only depends on the unknown G through $f_G(\cdot; \nu)$ and $f_G(\cdot; \nu + 1)$. $f_G(\cdot; \nu)$ is easily handled, since we have access to direct measurements from it. As shown in Theorem 7, $f_G(\cdot; \nu)$ may be estimated at a rate that is parametric (up to logarithmic factors) with the NPML under nonparametric specification of G .

The term $f_G(\cdot; \nu + 1)$ is slightly unusual since it corresponds to the density of sample variances with one additional degree of freedom compared to our actual observations ($\nu + 1$

vs. ν). Hence we do not have access to direct measurements from $f_G(\cdot; \nu + 1)$. Nevertheless, the following lemma establishes that if we have \widehat{G} such that $f_G(\cdot; \nu) \approx f_{\widehat{G}}(\cdot; \nu)$, then also $f_G(\cdot; \nu + 1) \approx f_{\widehat{G}}(\cdot; \nu + 1)$. The key technical argument for the proof (in Supplement E.2) hinges on properties of the Mellin transform [Epstein, 1948, Butzer and Stefan, 1999, Brenner Miguel et al., 2021].

Lemma 10. Consider two distributions G, G' as in (2) with $\mathbb{E}_G[\sigma^{-1}], \mathbb{E}_{G'}[\sigma^{-1}] < M$, for a $M < \infty$. Suppose that $\nu \geq 2$. Then there exists a positive constant $C = C(\nu, M)$ that only depends on ν, M such that:

$$\|f_G(\cdot; \nu + 1) - f_{G'}(\cdot; \nu + 1)\|_{L^2}^2 \leq C \left(1 + \left| \log \left(\|f_G(\cdot; \nu) - f_{G'}(\cdot; \nu)\|_{L^2}^2 \right) \right| \right) \|f_G(\cdot; \nu) - f_{G'}(\cdot; \nu)\|_{L^2}^2,$$

where $\|f\|_{L^2}^2 = \int_{[0, \infty)} f^2(t) dt$.

By combining Theorem 7, Proposition 9, and Lemma 10, we prove in Supplement E.3 that the empirical Bayes p-values $P_{\widehat{G}}(Z_i, S_i^2)$ converge to the oracle p-values $P_G(Z_i, S_i^2)$ at almost parametric rates.

Theorem 11. Suppose the assumptions of Theorem 7 hold and that $\nu \geq 3$. Then, for any $\zeta \in (1/2, 1)$, there exists a constant $C = C(\nu, \underline{L}, \bar{U}, \zeta)$ that depends only on $\nu, \underline{L}, \bar{U}, \zeta$ such that:

$$\max_{1 \leq i \leq n} \mathbb{E}_G \left[\left| P_G(Z_i, S_i^2) \wedge \zeta - P_{\widehat{G}}(Z_i, S_i^2) \wedge \zeta \right| \right] \leq C \frac{(\log n)^{5/2}}{\sqrt{n}} \text{ for all } n \in \mathbb{N}_{\geq 3}.$$

Above we write $a \wedge b := \min\{a, b\}$ for two numbers $a, b \in \mathbb{R}$.

The truncation of the p-values to $\zeta \in (1/2, 1)$ is helpful in controlling the tail behavior of $P_G(z, s^2)$ when $z \approx 0$. In the next section we will study the Benjamini-Hochberg procedure; the above guarantee suffices as long as $\alpha < \zeta$, where α is the nominal level at which we seek to control the false discovery rate.

Example 12 (Identical variances). Suppose that all the variances σ_i^2 are almost surely equal to $\bar{\sigma}^2 > 0$, that is, $G = \delta_{\bar{\sigma}^2}$, where δ_u denotes the point mass at u . In this case, the oracle procedure for testing $H_i : \mu_i = 0$ reduces to the usual z-test with p-value $2(1 - \Phi(|Z_i|/\bar{\sigma}))$. Theorem 11 then implies that the empirical partially Bayes p-values using the NPMLE in (9) automatically adapt and are close to the oracle z-test p-values.

The assumption that G is a point mass at $\bar{\sigma}$, may appear to be contrived. In the statistical literature, related assumptions are often considered [Robbins, 1956, Ignatiadis et al., 2021, Soloff et al., 2021] to demonstrate the adaptation of nonparametric empirical Bayes procedures. In the present setting, the assumption is substantive: the assumption that all σ_i are equal appears regularly in scientific publications [Kerr et al., 2000, Houde et al., 2011, Weis, 2019]. The motivation is to stabilize the variance estimate when the individual sample variances are very noisy, (e.g., (8) with small degrees of freedom ν). Our nonparametric strategy will automatically adapt to the equal variance assumption, if it is indeed warranted, and can thus match the power of the oracle z-test with known variance. The gains in that case, e.g., compared to the t-test, can be substantial, see the classical treatment of Walsh [1949] and the simulation study in Section 7.

We conclude with an important consequence of Theorem 11: the empirical partially Bayes p-values $P_{\widehat{G}}(Z_i, S_i^2)$ asymptotically satisfy the uniformity properties as the oracle p-values $P_G(Z_i, S_i^2)$ that we laid out in Proposition 3; see the proof in Supplement E.4.

Proposition 13. Let $\mathcal{H}_0 := \{i \in \{1, \dots, n\} : \mu_i = 0\}$ be the subset of null hypotheses. Then in the setting of Theorem 11 the following hold for $P_i = P_{\hat{G}}(Z_i, S_i^2)$:

1. Asymptotic conditional uniformity:

$$\max_{i \in \mathcal{H}_0} \mathbb{E}_G \left[\sup_{t \in [0,1]} |\mathbb{P}_G [P_i \leq t \mid S_1^2, \dots, S_n^2] - t| \right] \rightarrow 0 \text{ as } n \rightarrow \infty.$$

2. Asymptotic uniformity:

$$\max_{i \in \mathcal{H}_0} \sup_{t \in [0,1]} |\mathbb{P}_G [P_i \leq t] - t| \rightarrow 0 \text{ as } n \rightarrow \infty.$$

In the statement about conditional uniformity we condition on all of S_1^2, \dots, S_n^2 ⁹ to make explicit the following: the empirical partially Bayes approach proceeds in two steps: first learn information from all the sample variances S_1^2, \dots, S_n^2 to construct test statistics with desirable properties, and then use the randomness remaining (i.e., the randomness in (Z_1, \dots, Z_n) conditional on S_1^2, \dots, S_n^2) for multiple testing.

3.3 Benjamini-Hochberg with empirical partially Bayes p-values

In this subsection we describe the last step of Algorithm 1 in more detail. Suppose we are testing n hypotheses H_1, \dots, H_n based on p-values P_1, \dots, P_n . Given $\alpha \in (0, 1)$, the [Benjamini and Hochberg \[1995\]](#) (BH) procedure produces a subset \mathcal{D} of $\{1, \dots, n\}$ representing the indices of rejected hypotheses through the following rule:

$$\text{Reject } H_k \text{ if } P_k \leq P_{(k^*)}, \text{ where } k^* := \max \left\{ \ell \in \{0, \dots, n\} : P_{(\ell)} \leq \frac{\alpha \cdot \ell}{n} \right\}, \quad (14)$$

and $P_{(1)} \leq P_{(2)} \leq \dots \leq P_{(n)}$ are the order statistics of P_1, \dots, P_n and $P_{(0)} = 0$.

The most basic type-I error result for the BH procedure is as follows. Let $V_n = \#\mathcal{D} \cap \mathcal{H}_0$ be the total number of false discoveries of BH, $R_n = \#\mathcal{D}$ the total number of discoveries, and define the false discovery rate as the expectation of the false discovery proportion, that is, $\text{FDR}_n = \mathbb{E} [V_n / (R_n \vee 1)]$, where we write $a \vee b = \max \{a, b\}$ for $a, b \in \mathbb{R}$. Then, if all p-values are independent and $P_i \sim U[0, 1]$ for $i \in \mathcal{H}_0$, the BH procedure satisfies $\text{FDR}_n = \alpha n_0 / n$ where $n_0 = \#\mathcal{H}_0$ is the number of null hypotheses. In words: the BH procedure controls the false discovery rate at the slightly conservative level $\alpha n_0 / n$.¹⁰

In our setting, we only have approximate p-values $P_i = P_{\hat{G}}(Z_i, S_i^2)$, which in general will not be exactly uniformly distributed for $i \in \mathcal{H}_0$ and finite n . Our next results builds on Theorem 11 and demonstrates that BH applied to $P_i = P_{\hat{G}}(Z_i, S_i^2)$ (Algorithm 1) asymptotically controls the false discovery rate at level $\alpha n_0 / n$ as $n \rightarrow \infty$; see Supplement F.2 for a proof.

⁹For the oracle p-values it automatically holds that $\mathbb{P}_G [P_G(Z_i, S_i^2) \leq t \mid S_1^2, \dots, S_n^2] = \mathbb{P}_G [P_G(Z_i, S_i^2) \leq t \mid S_i^2]$, since we assume independence across i .

¹⁰The conservative FDR control of the BH procedure has motivated the development of null proportion adaptive methods, e.g., Storey's procedure [[Storey et al., 2004](#)]. At a high level, null proportion adaptive methods proceed by estimating n_0/n as $\widehat{\pi}_0$ and then applying the BH procedure at level $\alpha/\widehat{\pi}_0$. It is possible to prove that if we were to use Storey's procedure [[Storey et al., 2004](#)] instead of BH in Algorithm 1, then the false discovery rate would be asymptotically controlled in the setting of Theorem 14.

Theorem 14. Suppose $\liminf_{n \rightarrow \infty} \inf_{t \in [t_1^*, t_2^*]} \left\{ \sum_{i=1}^n \mathbb{P}_{G, \mu_i} [P_G(Z_i, S_i^2) \leq t] / (nt) \right\} \geq 1/\alpha'$ for $t_1^*, t_2^*, \alpha' \in (0, \alpha)$ with $\alpha \in (0, 1/2)$ fixed. Then, under the conditions of Theorem 7 and for $\nu \geq 3$, Algorithm 1 asymptotically controls the false discovery rate, that is:

$$\limsup_{n \rightarrow \infty} \left\{ \frac{n}{n_0} \text{FDR}_n \right\} = \limsup_{n \rightarrow \infty} \left\{ \frac{n}{n_0} \mathbb{E}_{G, \mu} \left[\frac{V_n}{R_n \vee 1} \right] \right\} \leq \alpha.$$

The assumption we make on the signal strength $\sum_{i=1}^n \mathbb{P}_{G, \mu_i} [P_G(Z_i, S_i^2) \leq t^*]$ is common in the analysis of the asymptotics of BH and is similar to e.g., the assumption of Storey et al. [2004, Theorem 4]. It requires that there is an interval close to the origin in which the averaged distribution of oracle p-values is stochastically smaller than the uniform distribution by a margin.

4 Oracle hypothesis testing in the compound setting

“Let us use a mixed model, even if it might not be appropriate”

— Hans C. van Houwelingen, 2014

In this section we study our procedure in the purely frequentist compound setting wherein σ_i^2 are fixed. More formally, we observe independent (but not identically distributed) draws of (Z_i, S_i^2) from (1) with $\boldsymbol{\sigma} = \boldsymbol{\sigma}_n = (\sigma_1, \dots, \sigma_n)$, $\boldsymbol{\mu} = \boldsymbol{\mu}_n = (\mu_1, \dots, \mu_n)$ deterministic but unknown. We nevertheless proceed as before and pretend that the data generating process is given by (6). Our main result in this section is that the conclusion of Theorem 14 on the asymptotic false discovery rate control continues to hold in the compound setting.

It is instructive to consider a single hypothesis, say $n = i = 1$, as in Section 2. Suppose we posit model (6) for a pre-specified G and compute $P_i = P_G(Z_i, S_i^2)$ as in (3), but seek inferential validity under (1), that is, for fixed σ_1 . Then, we may interpret P_i as a conditional predictive p-value as defined by Bayarri and Berger [2000]—they introduce conditional predictive p-value as an inferential device for dealing with nuisance parameters through marginalization (over G). Let σ_1 be fixed, $\mu_1 = 0$ and take $\nu \rightarrow \infty$. Then (under regularity assumptions on G) the results of Robins, van der Vaart, and Ventura [2000] imply that P_1 is asymptotically uniform.

In contrast, our asymptotic results are attained as $n \rightarrow \infty$ and ν is *fixed*. For example, Theorem 14* below establishes asymptotic control of the false discovery rate with Algorithm 1 as $n \rightarrow \infty$ and ν is fixed. Hence, we find this conclusion striking. For example, if $\mu_1 = 0$, then $P_G(Z_1, S_1^2)$ will not be uniform for *any* choice of G , unless $G = \delta_{\sigma_1}$. This is in sharp contrast to e.g., the result in Proposition 3, wherein $P_G(Z_1, S_1^2)$ is uniform conditional on S_1^2 . In other words, in the compound setting $P_{\hat{G}}(Z_i, S_i^2)$ are not asymptotically “valid” p-values. Nevertheless, the asymptotic false discovery rate guarantee holds (Theorem 14*). The reason is that while type-I error for individual hypotheses is not correctly controlled, the *average* type-I error across hypotheses *is controlled*—such notion of average significance control was introduced by Armstrong [2022]. We make this argument rigorous below.

To explain the core ideas, we will first provide a summary of compound decision theory in the Gaussian sequence model in Section 4.1 and then explain how compound decision theory applies to our setting in Section 4.2.

4.1 A modicum of compound decision theory

We provide a bird’s eye overview of compound decision theory in the Gaussian sequence model, and refer the reader for more details to [Robbins \[1951\]](#), [Copas \[1969\]](#), [Jiang and Zhang \[2009\]](#), [Weinstein \[2021\]](#) and references therein. Consider the following two Gaussian location sequence models (where M is a distribution on \mathbb{R}):

$$\boldsymbol{\mu} = (\mu_1, \dots, \mu_n) \text{ fixed,} \quad Z_i \stackrel{\text{iid}}{\sim} \mathcal{N}(\mu_i, 1) \text{ for } i = 1, \dots, n \quad (15a)$$

$$\mu_i \stackrel{\text{iid}}{\sim} M \text{ for } i = 1, \dots, n, \quad Z_i | \mu_i \stackrel{\text{iid}}{\sim} \mathcal{N}(\mu_i, 1) \text{ for } i = 1, \dots, n. \quad (15b)$$

In the first model, the parameters of interest μ_i are fixed, while in the second model they are randomly distributed according to M . The crux of compound decision theory is that for certain statistical tasks, models (15a) and (15b) behave very “similarly” under the choice of prior $M = M(\boldsymbol{\mu}) = \frac{1}{n} \sum_{i=1}^n \delta_{\mu_i}$, that is, when M is equal to the empirical measure of μ_1, \dots, μ_n .

Perhaps the most famous result in compound decision theory is the fundamental theorem of compound decisions. Suppose we seek to estimate μ_1, \dots, μ_n in either of the models in (15) with $\hat{\mu}_i = \eta(Z_i)$ where $\eta : \mathbb{R} \rightarrow \mathbb{R}$ is a fixed function. Then:

$$\mathbb{E}_{\boldsymbol{\mu}} \left[\frac{1}{n} \sum_{i=1}^n (\eta(Z_i) - \mu_i)^2 \right] = \frac{1}{n} \sum_{i=1}^n \mathbb{E}_{\mu_i} \left[(\eta(Z_i) - \mu_i)^2 \right] = \mathbb{E}_{\mu \sim M(\boldsymbol{\mu})} \left[(\eta(Z) - \mu)^2 \right]. \quad (16)$$

The critical observation is that the LHS is the mean squared error in estimating μ_i with $\hat{\mu}_i$ under the compound model (15a) while the RHS is the mean squared under the hierarchical model (15b). Under (15b), (μ_i, Z_i) are i.i.d. and so we drop the subscript i in the RHS of (16). The equality in (16) is merely formal; however a deep (and challenging to prove) consequence is that an empirical Bayes approach that estimates M as \widehat{M} based on Z_1, \dots, Z_n and then uses $\hat{\eta}(z) = \mathbb{E}_{\widehat{M}}[\mu | Z_i = z]$ will perform well in mean squared error under both models in (15) [[Jiang and Zhang, 2009](#)].

4.2 Average significance controlling tests

We are ready to study our empirical partially Bayes multiple testing method in the compound setting. In Section 3 we treated $\boldsymbol{\mu} = (\mu_1, \dots, \mu_n)$ as fixed, but assumed that $\sigma_i^2 \sim G$ (2). Now we also treat $\boldsymbol{\sigma} = (\sigma_1, \dots, \sigma_n)$ as fixed. Analogously to our recap of compound decision theory in Section 4.1, the crux of our argument will be that the setting with $\boldsymbol{\sigma}$ fixed is quantitatively similar to the hierarchical model (6) with a specific choice of prior G for σ_i^2 , namely the empirical distribution of the σ_i^2 :

$$G \equiv G(\boldsymbol{\sigma}) := \frac{1}{n} \sum_{i=1}^n \delta_{\sigma_i^2}. \quad (17)$$

The compound decision argument suggests considering the following compound “p-value” in lieu of the conditional p-values (3):

$$P_{\boldsymbol{\sigma}}(z, s^2) := P_{G(\boldsymbol{\sigma})}(z, s^2) = \frac{\sum_{j=1}^n 2(1 - \Phi(|z|/\sigma_j))p(s^2 | \sigma_j^2)}{\sum_{j=1}^n p(s^2 | \sigma_j^2)}. \quad (18)$$

Suppose we knew $\boldsymbol{\sigma}$ and could use $P_{\boldsymbol{\sigma}}(Z_i, S_i^2)$, $i = 1, \dots, n$, to test the null hypotheses $H_i : \mu_i = 0$. Would we have any frequentist guarantees? The following theorem proves that

this is indeed the case and the $P_{\boldsymbol{\sigma}}(Z_i, S_i^2)$ correspond to average significance controlling tests introduced by [Armstrong \[2022\]](#).

Theorem 15. Consider n independent draws from model (1) with $\boldsymbol{\mu}$ and $\boldsymbol{\sigma}$ fixed. Denote the null indices by $\mathcal{H}_0 := \{1 \leq i \leq n : \mu_i = 0\}$. The oracle compound p-values $P_{\boldsymbol{\sigma}}(Z_i, S_i^2)$ (18) satisfy the following guarantee:

$$\frac{1}{n} \sum_{i \in \mathcal{H}_0} \mathbb{P}_{\boldsymbol{\sigma}} [P_{\boldsymbol{\sigma}}(Z_i, S_i^2) \leq t] = \frac{1}{n} \sum_{i \in \mathcal{H}_0} \mathbb{P}_{\sigma_i} [P_{\boldsymbol{\sigma}}(Z_i, S_i^2) \leq t] \leq t \text{ for all } t \in [0, 1]. \quad (19)$$

Proof. Let $Z'_i = Z_i - \mu_i$ so that $Z'_i = Z_i$ for $i \in \mathcal{H}_0$. By definition, $P_{\boldsymbol{\sigma}}(Z_i, S_i^2) = P_{G(\boldsymbol{\sigma})}(Z_i, S_i^2)$ with $G(\boldsymbol{\sigma})$ defined in (17). Then:

$$\frac{1}{n} \sum_{i \in \mathcal{H}_0} \mathbb{P}_{\sigma_i} [P_{G(\boldsymbol{\sigma})}(Z_i, S_i^2) \leq t] \leq \frac{1}{n} \sum_{i=1}^n \mathbb{P}_{\sigma_i} [P_{G(\boldsymbol{\sigma})}(Z'_i, S_i^2) \leq t] = \mathbb{P}_{G(\boldsymbol{\sigma})} [P_{G(\boldsymbol{\sigma})}(Z', S^2) \leq t].$$

The last equality is only formal (cf. Section 4.1). Nevertheless, this formal equality enables us to conclude, since by Proposition 3 it holds that $\mathbb{P}_{G(\boldsymbol{\sigma})} [P_{G(\boldsymbol{\sigma})}(Z', S^2) \leq t] = t$ for all $t \in [0, 1]$. \square

We comment on the average significance controlling property in (19) as far as it pertains to empirical partially Bayes multiple hypothesis testing and refer the reader to [Armstrong \[2022\]](#) and [Armstrong, Kolesár, and Plagborg-Møller \[2022\]](#) for elaboration in other settings. (19) has two weaknesses compared to Proposition 3b. First, control only holds on *average* (rather than holding for each individual null hypothesis), and second the guarantee is inflated by n/n_0 , where $n_0 = \#\mathcal{H}_0$.¹¹ The first weakness seems unavoidable in view of the nature of the compound setting. The second weakness is inherent to the empirical partially Bayes approach we have described (and which is predominant in practice): by applying the NPMLE (9) to S_i^2 , $i = 1, \dots, n$, we estimate the empirical distribution of all the σ_i^2 , that is, $G(\boldsymbol{\sigma})$ (17). However, the correct reference distribution is the empirical distribution of the null σ_i^2 , that is, $G((\sigma_i)_{i \in \mathcal{H}_0})$.¹²

To make things concrete, suppose for simplicity that $n = 2$ and $i = 1$ is a null hypothesis, while $i = 2$ an alternative hypothesis. Take $\nu = 3$ degrees of freedom and suppose that $\sigma_1^2 = 1, \sigma_2^2 = 0.5$, that is, the alternative has a smaller variance than the null. Then we can numerically compute for $t = 0.01$, that $\mathbb{P}_{\sigma_1} [P_{(\sigma_1, \sigma_2)}(Z_1, S_1^2) \leq 0.01] \approx 0.0199$, i.e., the upper bound $n/n_0 \cdot t = 0.02$ in (19) is nearly tight. The intuitive reason that type-I error control is lost by a factor of nearly 2 is that the reference distribution for the null variances, that is $G = \delta_1$, is pushed stochastically downwards to $G = (\delta_1 + \delta_{0.5})/2$ due to the presence of a smaller variance under the alternative.

In view of the preceding example, we find it surprising that even under adversarial arrangement of the variances σ_i^2 , the oracle compound p-values have the type-I error guarantee in (19) which is only inflated by a factor n/n_0 . In many applications there will only be few alternatives, so that $n/n_0 \approx 1$. A further silver lining is that the average significance control guarantee in (19) suffices to control the false discovery rate asymptotically when applying

¹¹In the setting of Proposition 3b: $\sum_{i \in \mathcal{H}_0} \mathbb{P}_G [P_G(Z_i, S_i^2) \leq t] = tn_0$.

¹²To estimate $G((\sigma_i)_{i \in \mathcal{H}_0})$, we would also need to use information about the μ_i , which is contained in the Z_i , but not the S_i^2 . This hints at possible benefits of joint empirical Bayesian modeling of μ_i and σ_i^2 compared to the partially empirical Bayes approach. We refer to the Introduction (Section 1) for further discussion of this point.

the Benjamini-Hochberg procedure [Armstrong, 2022]. Hence, below (Theorem 14*) we establish a false discovery control guarantee for Algorithm 1 in the compound setting that is analogous to the result in Theorem 14.

4.3 Asymptotic results in the compound setting

In this section we show that most of the asymptotic results of Section 3 have an analogue in the compound setting wherein $\sigma = (\sigma_1, \dots, \sigma_n)$ is fixed. There is one notable exception of a result without a matching analogue in the compound setting: as we explained in the beginning of this section, the asymptotic conditional and unconditional uniformity of null p-values that we demonstrated in Proposition 13 typically would not hold in the compound setting.

To state our results, it will be helpful to introduce the averaged density f_σ of the S_i^2 :

$$f_\sigma(s^2) \equiv f_\sigma(s^2; \nu) := \frac{1}{n} \sum_{i=1}^n p(s^2 \mid \sigma_i^2, \nu). \quad (20)$$

Note that, as in the general compound argument, $f_\sigma(\cdot)$ is formally equal to the marginal density $f_{G(\sigma)}(\cdot)$ in (10) with the prior $G(\sigma)$ (17). The probabilistic interpretation of these two objects however is distinct.

First we state the matching result to Theorem 7:

Theorem 7*. Let $0 < \underline{L} \leq L \leq U \leq \bar{U} < \infty$. Suppose $\nu \geq 2$, $\sigma_i^2 \in [L, U]$ for all i and that we compute the NPML \widehat{G} (9) under the additional constraint $G([\underline{L}, \bar{U}]) = 1$. Then, there exist constants $C = C(\nu, \underline{L}, \bar{U}) > 0$ and $c = c(\nu, L, \bar{U}) > 0$ such that:

$$\mathbb{P}_\sigma [\mathfrak{H}(f_{\widehat{G}}, f_\sigma) \geq C \log n / \sqrt{n}] \leq \exp(-c(\log n)^2) \text{ for all } n \in \mathbb{N}_{\geq 1}.$$

Compared to Theorem 7, Theorem 7* only has the following differences: first, the probability in the statement is taken with respect to the n independent draws of S_i^2 in (8) with σ fixed (rather than $\sigma_i^2 \stackrel{\text{iid}}{\sim} G$), and second, the Hellinger distance bound pertains to the average density f_σ (rather than f_G). The argument for these two theorems are almost identical (cf. Zhang [2009] in the Gaussian location setting) and so we also prove both in the Supplement concurrently.¹³

The following theorem, which parallels Theorem 11, also follows immediately:

Theorem 11*. Suppose the assumptions of Theorem 7* hold and that $\nu \geq 3$. Then, for any $\zeta \in (1/2, 1)$, there exists a constant $C = C(\nu, \underline{L}, \bar{U}, \zeta)$ that depends only on $\nu, \underline{L}, \bar{U}, \zeta$ such that:

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E}_\sigma [|P_\sigma(Z_i, S_i^2) \wedge \zeta - P_{\widehat{G}}(Z_i, S_i^2) \wedge \zeta|] \leq C \frac{(\log n)^{5/2}}{\sqrt{n}} \text{ for all } n \in \mathbb{N}_{\geq 3}.$$

The difference between the conclusions of the above result with that of Theorem 11 is that we only bound the expectation averaged over all $i = 1, \dots, n$.

¹³While we do not explicitly state it here, Corollary 8 also has an analogue in the compound setting: for any bounded and continuous function $\psi : [0, \infty) \rightarrow \mathbb{R}$ it holds that $\int \psi(u)(d\widehat{G}(u) - dG(\sigma)(u))$ converges to 0 in probability. We refer the reader to Greenshtein and Ritov [2022, Theorem 4] for the rationale and rigorous justification of such a result.

More interestingly, as already announced, the asymptotic false discovery rate control guarantee of Theorem 14 also continues to hold in the compound setting. Unlike the two preceding theorems, the following result does not follow immediately and the proof relies on the machinery developed in Section 4.2 on average significance controlling tests. Asymptotic false discovery rate control with p-values satisfying the average significance control property (19) was already shown by Armstrong [2022]. Here our argument also accounts for the fact that the average significance control property is only approximately satisfied since the BH procedure uses $P_i = P_{\widehat{G}}(Z_i, S_i^2)$ rather than $P_{\sigma}(Z_i, S_i^2)$.

Theorem 14*. Suppose $\liminf_{n \rightarrow \infty} \inf_{t \in [t_1^*, t_2^*]} \{ \sum_{i=1}^n \mathbb{P}_{\sigma_i, \mu_i} [P_G(Z_i, S_i^2) \leq t] / (nt) \} \geq 1/\alpha'$ for $t_1^*, t_2^*, \alpha' \in (0, \alpha)$ with $\alpha \in (0, 1/2)$ fixed. Then, under the conditions of Theorem 7* and for $\nu \geq 3$, Algorithm 1 asymptotically controls the false discovery rate, that is:

$$\limsup_{n \rightarrow \infty} \{\text{FDR}_n\} = \limsup_{n \rightarrow \infty} \left\{ \mathbb{E}_{\sigma, \mu} \left[\frac{V_n}{R_n \vee 1} \right] \right\} \leq \alpha.$$

It is worth pointing out a difference in the statement of Theorem 14* compared to Theorem 14. In the latter, the FDR is controlled at level $n_0/n \cdot \alpha$, while here it is controlled only at level α : BH is conservative by a factor n_0/n [Benjamini and Hochberg, 1995, equation (2)], the oracle compound p-values may be anticonservative by a factor of at most n/n_0 (Theorem 15), and these two behaviors cancel out. A note of caution: our conclusion in Footnote 10 no longer holds and null-proportion adaptive methods such as Storey's procedure may not control the FDR in the compound setting (see results of simulation study in Section 7).

5 Results under a joint hierarchical model

So far, we have treated the following two cases: $\boldsymbol{\mu} = (\mu_1, \dots, \mu_n)$ is fixed and $\sigma_i^2 \stackrel{\text{iid}}{\sim} G$ (Section 3) and both $\boldsymbol{\mu}$ and $\boldsymbol{\sigma}^2 = (\sigma_1^2, \dots, \sigma_n^2)$ are fixed (Section 4). It is instructive to also study Algorithm 1 when *both* μ_i and σ_i^2 are random; the following discussion will further illuminate the distinction between the formal results in Sections 3 and 4. To be concrete, we suppose that there exists an unknown distribution Π supported on $\mathbb{R} \times \mathbb{R}_{>0}$ such that

$$(\mu_i, \sigma_i^2) \stackrel{\text{iid}}{\sim} \Pi \text{ for } i = 1, \dots, n. \quad (21)$$

The distribution $G(\cdot; \Pi)$ of σ_i^2 is given by marginalizing Π over μ_i , that is, G is defined as follows for any Borel set $A \subset [0, \infty)$:

$$G(A; \Pi) := \Pi(\mathbb{R} \times A). \quad (22)$$

We then have that (2) holds, that is, $\sigma_i^2 \stackrel{\text{iid}}{\sim} G(\cdot; \Pi)$.

We may directly apply the results of Section 4 by conditioning on the random vectors $\boldsymbol{\mu}$ and $\boldsymbol{\sigma}$, in which case we are back to draws from model (1). The more interesting question is the following: when are the results of Section 3 applicable? For example, when can we argue that the empirical partially Bayes p-values for null hypotheses are approximately uniform conditional on (S_1^2, \dots, S_n^2) (Proposition 13 in Section 3)? In the following we demonstrate that it suffices that $\mathbb{1}(\mu_i = 0)$ is independent of σ_i^2 under Π (21), and we also contextualize this requirement.

As our initial attempt to applying the results of Section 3, we condition on $\boldsymbol{\mu}$ (not on $\boldsymbol{\sigma}$). Then, (6) specifies the following (where we make the conditioning on $\boldsymbol{\mu}$ explicit in the notation): there exists a distribution \tilde{G} such that¹⁴

$$\sigma_i^2 \mid \mu_i \stackrel{\text{iid}}{\sim} \tilde{G} \text{ for } i = 1, \dots, n.$$

For the above to hold for almost all $\boldsymbol{\mu}$ from (21), we require that the marginals of Π factorize, that is, we require that σ_i^2 is *independent* of μ_i in which case $\tilde{G} = G(\cdot; \Pi)$ (22). Hence, in the above case, we may apply the results from Section 3 by conditioning on $\boldsymbol{\mu}$. The assumption of independence is relatively common in the literature, see e.g., Ghosal and van der Vaart [2001, equation (2.2)], Jiang [2020], Zheng et al. [2021], and Soloff et al. [2021]. This assumption can be reasonable in some situations, namely when one believes that all primary parameters μ_i are of similar magnitude for all i [Efron and Morris, 1973, Section 8]. On the other hand, it is also well recognized that the assumption may be violated in problems of practical interest [Stephens, 2016, Gu and Koenker, 2017b, Weinstein et al., 2018, Chen, 2022].

It turns out, however, that independence of μ_i and σ_i^2 is not necessary to e.g., prove an analogous result to Proposition 13 in Section 3: in the empirical partially Bayes approach we model and estimate the prior distribution of the nuisance parameters σ_i^2 based on (S_1^2, \dots, S_n^2) only, i.e., a vector of statistics that are ancillary for μ_i (Proposition 1). Hence most of the technical results in Section 3, e.g., Theorem 7 and Corollary 8 only rely on the marginal distribution $G(\cdot; \Pi)$ of the σ_i^2 ; and not the joint distribution Π (21). The next proposition provides a sufficient condition, substantially more general than independence of μ_i and σ_i^2 , such that the oracle conditional “p-values” $P_{G(\cdot; \Pi)}(Z_i, S_i)$, i.e., (3) for the prior $G(\cdot; \Pi)$, are bona-fide conditional p-values that satisfy the properties in Proposition 3 (see Supplement G for a proof).

Proposition 16. Suppose $(\mu_i, \sigma_i) \sim \Pi$ (21) and (Z_i, S_i^2) are generated as in (1) conditional on μ_i, σ_i . Suppose further that,

$$\mathbf{1}(\mu_i = 0) \text{ is independent of } \sigma_i^2, \tag{23}$$

then it holds that:

$$\mathbb{P}_\Pi [P_{G(\cdot; \Pi)}(Z_i, S_i^2) \leq t \mid S_i^2, \mu_i = 0] = t \text{ for all } t \in [0, 1].$$

Assumption (23) seems unavoidable for the empirical partially Bayes methodology—it is the price we pay for using all of (S_1^2, \dots, S_n^2) to get a reference *null* distribution of Z_i conditional on S_i^2 .¹⁵ At the same time, (23) encapsulates a requirement that is substantially weaker than independence of μ_i and σ_i^2 . For example, Stephens [2016] considers the following modeling assumption: there exists $a \geq 0$ such that μ_i/σ_i^a is independent of σ_i^2 under Π .¹⁶ This assumption implies (23).¹⁷ From a practical perspective, we anticipate that (23) will approximately hold in many situations (and if it does not; then our methods still have the guarantees of Section 4).

¹⁴Here we are implicitly using the i.i.d. structure in (21) which implies that the conditional distribution of σ_i^2 given $\boldsymbol{\mu}$ is equal to the conditional distribution of σ_i^2 given μ_i .

¹⁵In the simulation study of Section 7 we will see that when (16) is violated, then the conclusion of Theorem 14 may not hold.

¹⁶By taking $a = 0$, this assumption posits independence of μ_i and σ_i^2 .

¹⁷The reason is that $\mathbf{1}(\mu_i = 0) = \mathbf{1}(\mu_i/\sigma_i^a = 0)$.

6 Applications

As we mentioned in the Introduction, the framework we consider is widely applicable. Here we complement the differential methylation analysis from Section 1.2 by presenting two additional use-cases for our framework: detecting differentially expressed genes with microarray data and detecting differentially abundant proteins with mass spectrometry data. We further choose these datasets so as to get insights into empirical partially Bayes analyses with different choices of the degrees of freedom ν : $\nu = 4$ for the methylation dataset, $\nu = 11$ for the microarray dataset and $\nu = 28$ for the proteomics dataset.

6.1 Differentially expressed genes after treatment with Ibrutinib

Here we consider a typical data analysis with microarray data. As explained in the Introduction (also see Efron [2010] for an eloquent exposition), microarrays were the driving force of several developments in multiple testing and empirical Bayes methodology (e.g., limma) in the beginning of the century. Nowadays the microarray technology has been largely superseded by RNA-Seq, but nevertheless it finds use, e.g., for clinical applications [Klaus and Reisenauer, 2018].

The concrete application we consider is the following: Dietrich et al. [2017] collected samples from 12 chronic lymphocytic leukemia (CLL) patients. Cells were incubated for 12 hours with the drug Ibrutinib (which is used to treat CLL) and compared against negative controls. Gene expression profiles were measured with Illumina (incorporation, San Diego, California, USA) microarrays. After performing a paired analysis for each probe (pairing each patient’s treated sample vs. the negative control), the data may be summarized in the form (1) with $n = 48,107$ probes and $\nu = 12 - 1 = 11$ degrees of freedom.

Applying Benjamini-Hochberg to control the FDR at $\alpha = 0.05$, we get 92 discoveries with the NPMLE-based p-values, 107 discoveries with limma (the estimated prior (4) has parameters $\hat{s}_0^2 \approx 0.007$, $\hat{\nu}_0 \approx 10.4$), and 14 discoveries with the standard t-test. Thus also in this problem, the empirical partially Bayes approaches showcase substantial increase in discoveries compared to the t-test. The results are visualized in Fig. 3. We note (Fig. 3a) that the NPMLE leads to a slightly better fit to the histogram of sample variances S_i^2 compared to limma’s inverse gamma model. The latter slightly overshoots for $S_i^2 \geq 0.02$, and this in turn leads to less conservative rejection regions (Fig. 3c) for large values of S_i^2 (which in turns explains the slight increase of discoveries with limma compared to the NPMLE).

6.2 Differentially abundant proteins in breast cancer subtypes

Our next application pertains to a proteomics dataset. Terkelsen et al. [2021] conducted a liquid chromatography tandem mass spectrometry analysis of 34 tumor interstitial fluid samples of three breast cancer subtypes: luminal, Her2, and triple negative that were processed in four experimental batches. To describe the application of our methods to this dataset, we first need to take a detour to explain how the framework enables handling of different experimental designs.

Remark 17 (Testing for contrasts in linear regression with repeated design). Here we demonstrate how (Z_i, S_i^2) as in (1) can arise from data summarization in Gaussian linear models when the goal is to test for contrasts in linear regressions with repeated design. The




Figure 3: **Partially empirical Bayes analysis with microarray data of Dietrich et al. [2017]**: We are testing $n = 48,107$ hypotheses $H_i : \mu_i = 0$ based on summary data as in (1) with $\nu = 11$ degrees of freedom. The three panels are analogous to the four panels of Fig. 1.

results herein are explicated by Smyth [2004] and lie at the core of the wide applicability of the empirical partially Bayes approach.

Suppose our data for the i -th hypothesis is modelled by a linear regression with K samples and p covariates, i.e.,

$$\mathbf{Y}_i \mid \mathbf{X} \sim \mathcal{N}(\mathbf{X}\boldsymbol{\beta}_i, \tilde{\sigma}_i^2 \mathbf{I}_p), \quad (24)$$

where $\mathbf{Y}_i \in \mathbb{R}^K$, $\mathbf{X} \in \mathbb{R}^{K \times p}$. Here \mathbf{I}_p is the $p \times p$ identity matrix and $\boldsymbol{\beta}_i \in \mathbb{R}^p$, $\tilde{\sigma}_i^2 > 0$ are the unknown parameters. Furthermore, suppose the parameter of interest is $\mu_i = \mathbf{c}^\top \boldsymbol{\beta}_i$, where $\mathbf{c} \in \mathbb{R}^p$ is a known contrast. Let $\hat{\boldsymbol{\beta}}_i := (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{Y}_i$ be the ordinary least squares coefficient estimate, and let

$$Z_i := \mathbf{c}^\top \hat{\boldsymbol{\beta}}_i, \quad S_i^2 := \mathbf{c}^\top (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{c} \cdot \left\| \mathbf{X} \hat{\boldsymbol{\beta}}_i - \mathbf{Y}_i \right\|_2^2 / (K - p).$$

Then (Z_i, S_i^2) satisfy (1) with $\mu_i = \mathbf{c}^\top \boldsymbol{\beta}_i$, $\sigma_i^2 = \tilde{\sigma}_i^2 \mathbf{c}^\top (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{c}$, and $\nu = K - p$.

We are ready to continue with the analysis of Terkelsen et al. [2021]: for each of $n = 6,763$ proteins, the investigators fit a linear model with design matrix consisting of the indicators




Figure 4: **Partially empirical Bayes analysis with proteomics data of [Terkelsen et al. \[2021\]](#)**: We are testing $n = 6,763$ hypotheses $H_i : \mu_i = 0$ based on summary data as in (1) with $\nu = 28$ degrees of freedom. The three panels are analogous to the four panels of Fig. 1.

for the 4 experimental batches and the 3 cancer subtypes. Hence the variance is estimated with $\nu = 34 - 1 - (4 - 1) - (3 - 1) = 28$ degrees of freedom. Here we seek to test for each protein whether the contrast for the difference of the coefficients for the luminal subtype minus the coefficient for the Her2 subtype is equal to 0. This problem can be cast into the framework of this paper with $\nu = 28$ degrees of freedom (Remark 17).

The Benjamini-Hochberg procedure (applied to control the FDR at $\alpha = 0.05$) yields 93 discoveries with the NPMLE-based p-values, 102 discoveries with limma (the estimated prior (4) has parameters $\hat{s}_0^2 \approx 0.077$, $\hat{\nu}_0 \approx 5.2$), and 74 discoveries with a standard t-test. As in our other applications, the empirical partially Bayes methods lead to more discoveries compared to the standard t-test, however the difference is less pronounced in this case. The reason is that the degrees of freedom are larger in this problem, so that there is less benefit in an empirical Bayes treatment of the sample variances S_i^2 , which are already reasonably accurate estimates of σ_i^2 .¹⁸ We visualize the results in Fig. 4. In Fig. 4c) we see that indeed the rejection thresholds of the t-test are more closely aligned with the empirical partially Bayes testing thresholds compared to Fig. 1c) and Fig. 3c). Nevertheless, it still holds true

¹⁸Indeed, all three p-values are asymptotically equivalent as $\nu \rightarrow \infty$.

that the t-test is more liberal for small values of S_i^2 , and more conservative for larger values of S_i^2 .

7 Simulations

In this section we conduct a simulation study to accompany our theoretical findings and the data analyses above. Throughout our simulations, we generate data from model (6) with $n = 10,000$. We vary the degrees of freedom $\nu \in \{2, 4, 8, 16, 32, 64\}$ and consider three choices of prior G for σ_i^2 :

1. A point mass at $\sigma_i^2 = 1$, i.e., $G = \delta_1$ as considered, e.g., in Example 12.
2. The scaled inverse χ^2 prior in (4) with $\nu = 6$ and $s_0^2 = 1$.
3. The two point prior that assigns equal mass to $\sigma^2 = 10$ and $\sigma^2 = 1$, that is, $G = (\delta_{10} + \delta_1)/2$.

The μ_i are generated as follows: first, we let $\mu_i = 0$ for 9000 of hypotheses (that is, the null proportion is equal to 0.9) with the null indices picked at random. For the alternative hypotheses, we let $\mu_i \sim \mathcal{N}(0, 16\sigma_i^2)$.

We compute p-values with the following four methods:

1. Standard **t-test**.
2. **Limma**.
3. **NPMLE**-based empirical partially Bayes.
4. **Oracle** partially Bayes that is provided with access to the ground truth G .

We correct for multiple testing with the Benjamini-Hochberg (**BH**) procedure applied at a nominal false discovery rate level $\alpha = 0.1$. For the oracle partially Bayes p-values, we also apply an alternative multiple testing strategy based on Storey’s null proportion adaptive procedure [Storey et al., 2004].¹⁹ In total this leads to 5 methods under evaluation—t-test (BH), limma (BH), NPMLE (BH), oracle (BH), and oracle (Storey). Finally, we evaluate methods in terms of their false discovery rate (FDR), and power, which we define as the expected proportion of true discoveries among all alternatives. Both the FDR and the power are computed by averaging over 3,000 Monte Carlo replicates of each simulation setting.

The results of the simulation are shown in Fig. 5. The first row shows the results for the simulation with the Dirac prior. All methods control the false discovery rate at $\alpha = 0.1$.²⁰ Limma and NPMLE are well-specified here²¹ and have power similar to the oracle test (which here is just a z-test with known variance $\sigma^2 = 1$, cf. Example 12), which is substantially greater than the power of the t-test. For example, for $\nu = 2$, the t-test has power almost equal to 0, while the other methods have power ≈ 0.5 . The t-test becomes more competitive as ν increases, while the performance of the other methods remains almost constant as ν increases (since here the empirical partially Bayes p-values are dominated by the point mass prior and the χ^2 -likelihood plays a minimal role). The second row of Fig. 5 shows the scaled inverse χ^2 simulation. Here the conclusions are similar to the Dirac prior. However, the power of the (empirical) partially Bayes methods is not as large as in the Dirac case

¹⁹The rationale for considering Storey’s procedure is to demonstrate that null proportion adaptive procedures may not control the FDR in settings wherein BH does; see the discussion after Theorem 14*.

²⁰In fact, we expect them to control the false discovery rate at level $0.09 = n_0/n \cdot \alpha$, see e.g., Theorem 14 and the discussion preceding it.

²¹For limma this is not immediately obvious. Note that the distribution of σ_i^2 in (4) converges weakly to δ_1 as $\nu_0 \rightarrow \infty$ and $s_0^2 = 1$.




Figure 5: **Simulation results:** We compare 5 methods—t-test (BH), limma (BH), NPMLE (BH), oracle (BH), and oracle (Storey)—across simulation settings that span three different choice of prior G for the variances σ_i^2 (rows of the figure), as well as varying degrees of freedom $\nu \in \{2, 4, 8, 16, 32, 64\}$ (x -axis of each plot). The panels of the first column show the false discovery rate of the different methods, while the panels of the second column show their power.

and it increases with ν ; the reason is that in this setting there is unexplained heterogeneity in the variances. The results for the two point prior are shown in the last row of Fig. 5. In contrast to the previous two simulation settings, here the limma model is misspecified. We observe that this misspecification can lead to both conservative results (that is, type-I error substantially below nominal), as occurs when $\nu = 2$, but also to anticonservative results (type-I error above nominal) as occurs when $\nu = 4$. Finally, we note that throughout these simulations, Storey’s multiple testing correction with oracle p-values leads to FDR control (as follows from Proposition 3b and Storey et al. [2004]) and slightly higher power than its BH counterpart.

Adversarial simulation: The results of the preceding simulations are in accordance to the theoretical results in Sections 2, 3, and 5. We next consider a challenging setting for the empirical partially Bayes methods. We repeat the simulations from above with the following tweak: we do not choose the null indices at random, but instead we sort the variances σ_i^2 and let $\mu_i = 0$ for the 9,000 hypotheses with the largest σ_i^2 . Hence we are violating (23) and the




Figure 6: **Adversarial simulation results:** The panels in this figure are analogous to the panels of the last two rows of Fig. 5. In contrast to Fig. 5, wherein the indices of null hypotheses were chosen at random, here null hypotheses are enriched for small variances σ_i^2 .

results of Section 3 are not applicable. Furthermore, as explained at the end of Section 4.2, the constellation in which the alternative hypotheses have smaller variances than the nulls is adversarial/challenging for the empirical partially Bayes methods. On the other hand, as explained in Section 5, the theoretical results of the compound setting (Section 4) are applicable.

Results of this simulation study are shown in Fig. 6.²² We make the following observations: first, as already explained in Section 4, we observe that Storey’s procedure with the oracle partially Bayes p-values no longer controls the FDR in contrast to the BH procedure. Second, we note that under such adversarial arrangement of null hypotheses, in some cases the t -test can have more power than the empirical partially Bayes based approaches. For example, for the scaled inverse χ^2 prior, the t -test has more power than the empirical partially Bayes approaches for $\nu \geq 8$ degrees of freedom, but less power for $\nu \leq 4$ degrees of freedom. An explanation for this phenomenon is that under the adversarial constellation we consider, partially Bayes p-values for null hypotheses are smaller than they should be (so that Storey’s procedure no longer controls FDR), but larger than they should be for alternative hypotheses (that is, the smaller sample variances of the alternatives are shrunk upwards towards the bulk of larger sample variances of the nulls). When the degrees of freedom are small, shrinkage (even in the wrong direction) is sufficiently beneficial to lead to an increase in power compared to the t -test, while for larger degrees of freedom the t -test outperforms the partially Bayes p-values. All in all, we consider the results of this simulation to be encouraging: even under this adversarial setting (that would be unlikely to occur in many real applications), Algorithm 1 with the NPMLE performs well, and in all cases

²²Fig. 6 does not show the Dirac prior because in that case the adversarial and non-adversarial simulations are identical.

closely tracks the performance of the oracle partially Bayes p-values.

8 Discussion

In this paper, we have made progress towards theoretically understanding the problem of empirical partially Bayes (multiple) hypothesis testing. The basic idea is to apply empirical Bayes techniques to “shrink” the nuisance parameters, while treating the primary parameter of interest in standard frequentist fashion. As we have mentioned throughout this work, the limma approach and model is ubiquitous in applications of high-throughput biology. The more general idea of empirical partially Bayes testing comes up in even more applications (albeit not under this name), e.g., in some of the most popular approaches for analyzing RNA-Seq data [Robinson et al., 2010, Love et al., 2014] based on models for count data. We hope that our work will spur further research into empirical partially Bayes testing and illuminate *why* this approach has been so successful in practice.

We have also made methodological progress by deriving a fully nonparametric generalization of the limma approach and by showing that even under nonparametric assumptions it is possible to approximate the “oracle” p-values at a rate that is parametric (that is, of order $1/\sqrt{n}$) up to a logarithmic factor. This suggests that our approach may have substantial benefits in practice and at little cost when the parametric model is justified. Further work could also consider finite-sample issues in more detail and modify the procedure to be more conservative, e.g., based on the ideas in Phipson et al. [2016], Stephens [2016], Ignatiadis and Wager [2022], and Xie and Stephens [2022].

Finally, our results also have implications for practitioners who use the parametric limma approach. They suggest the following:

1. Sample variance histogram as a diagnostic: The inverse chi-squared assumption for G in (4) underlying the limma method may appear to be overly strong in applications. Nevertheless, the results of Proposition 9 and Lemma 10 imply that limma can perform well even if \hat{G}^{limma} is a bad approximation of the “true” G as long as the parametric model provides a good fit for the marginal density of the sample variances, that is, $f_{\hat{G}^{\text{limma}}} \approx f_G$. The latter may be visually assessed by overlaying the implied marginal density with a histogram of the sample variances as e.g., in Fig. 1a) in this manuscript, as well as Wright and Simon [2003, Fig. 3 and 4] and Kammers et al. [2015, Supplementary Fig. 1]. Hence we recommend accompanying a limma analysis, when appropriate, with this diagnostic plot.

2. Benjamini-Hochberg compared to Storey’s adjustment: A well-known disadvantage of the Benjamini-Hochberg (BH) method is that it is not null proportion adaptive, that is, it typically controls the false discovery rate at level $\alpha \cdot \#\mathcal{H}_0/n$ when applied at level α and so can be slightly conservative. Adaptive methods such as Storey’s procedure [Storey et al., 2004] address this issue by estimating the null proportion $\#\mathcal{H}_0/n$ and then appropriately inflating the level at which the BH procedure is applied. Our results in Sections 4, 5, as well as the simulation study in Section 7 suggest that one should be cautious when applying adaptive multiple testing procedures in the context of limma (or more generally Algorithm 1): adaptive procedures can fail to control the false discovery rate, e.g., when (23) is violated, while the BH procedure can still work in such settings. Hence BH is to be preferred; we note that the software underlying limma indeed uses BH by default.

Reproducibility

All numerical results of this paper are fully third-party reproducible, and we provide the code and data in Github under the following link:

<https://github.com/nignatiadis/empirical-partially-bayes-paper>.

Acknowledgments

This research was supported by NSF grant DMS-2015376.

References

- J. Aitchison and I. R. Dunsmore. *Statistical Prediction Analysis*. Cambridge University Press, Cambridge, 1975.
- G. E. Andrews, R. Askey, and R. Roy. *Special Functions*. Cambridge University Press, Cambridge, first edition, 1999.
- T. B. Armstrong. False discovery rate adjustments for average significance level controlling tests. 2022. URL <http://arxiv.org/abs/2209.13686>.
- T. B. Armstrong, M. Kolesár, and M. Plagborg-Møller. Robust empirical Bayes confidence intervals. *Econometrica*, 90(6):2567–2602, 2022.
- P. Baldi and A. D. Long. A Bayesian framework for the analysis of microarray expression data: Regularized t-test and statistical inferences of gene changes. *Bioinformatics*, 17(6): 509–519, 2001.
- T. Banerjee, L. J. Fu, G. M. James, and W. Sun. Nonparametric Empirical Bayes Estimation on Heterogeneous Data. 2021. URL <http://arxiv.org/abs/2002.12586>.
- M. J. Bayarri and J. O. Berger. p -values for composite null models. *Journal of the American Statistical Association*, 95(452):1127–1142, 2000.
- D. Belomestny and A. Goldenshluger. Nonparametric density estimation from observations with multiplicative measurement errors. *Annales de l'Institut Henri Poincaré, Probabilités et Statistiques*, 56(1):36–67, 2020.
- Y. Benjamini and Y. Hochberg. Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B*, 57(1): 289–300, 1995.
- P. Billingsley. *Probability and Measure*. Wiley Series in Probability and Statistics. Wiley, New York, NY, third edition, 1995.
- L. Breiman. *Probability*. SIAM - Society for Industrial and Applied Mathematics, classic edition, 1992.
- S. Brenner Miguel, F. Comte, and J. Johannes. Spectral cut-off regularisation for density estimation under multiplicative measurement errors. *Electronic Journal of Statistics*, 15(1):3551–3573, 2021.

- P. L. Butzer and J. Stefan. A self-contained approach to Mellin transform analysis for square integrable functions; applications. *Integral Transforms and Special Functions*, 8 (3-4):175–198, 1999.
- I. Castillo and É. Roquain. On spike and slab empirical Bayes multiple testing. *The Annals of Statistics*, 48(5), 2020.
- J. Chen. Gaussian heteroskedastic empirical Bayes without independence, 2022. URL <http://arxiv.org/abs/2212.14444>.
- J. B. Copas. Compound decisions and empirical Bayes (with discussion). *Journal of the Royal Statistical Society: Series B*, 31(3):397–417, 1969.
- D. R. Cox. A note on partially Bayes inference and the linear model. *Biometrika*, 62(3): 651–654, 1975.
- D. R. Cox. Selected statistical papers of Sir David Cox. Volume 2. Foundations of Statistical Inference, Theoretical Statistics, Time Series and Stochastic Processes. volume 2. Cambridge University Press, Cambridge, 2005. Edited by D. J. Hand and A. M. Herzberg.
- D. R. Cox and D. V. Hinkley. *Theoretical Statistics*. Chapman and Hall, London, first edition, 1974.
- N. Cressie. A useful empirical Bayes identity. *The Annals of Statistics*, 10(2):625–629, 1982.
- X. Cui and G. A. Churchill. Statistical tests for differential expression in cDNA microarray experiments. *Genome Biology*, 4(4):210, 2003.
- X. Cui, J. T. G. Hwang, J. Qiu, N. J. Blades, and G. A. Churchill. Improved statistical tests for differential gene expression by shrinking variance components estimates. *Biostatistics*, 6(1):59–75, 2005.
- R. X. de Menezes, J. M. Boer, and H. C. van Houwelingen. Microarray data analysis: A hierarchical t-test to handle heteroscedasticity. *Applied Bioinformatics*, 3(4):229–235, 2004.
- P. Delmar, S. Robin, D. Tronik-Le Roux, and J. J. Daudin. Mixture model on the variance for the differential analysis of gene expression data. *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, 54(1):31–50, 2005.
- L. H. Dicker and S. D. Zhao. High-dimensional classification via nonparametric empirical Bayes and maximum likelihood inference. *Biometrika*, 103(1):21–34, 2016.
- S. Dietrich, M. Oleś, J. Lu, L. Sellner, S. Anders, B. Velten, B. Wu, J. Hüllein, M. da Silva Liberio, T. Walther, L. Wagner, S. Rabe, S. Ghidelli-Disse, M. Bantscheff, A. K. Oleś, M. Słabicki, A. Mock, C. C. Oakes, S. Wang, S. Oppermann, M. Lukas, V. Kim, M. Sill, A. Benner, A. Jauch, L. A. Sutton, E. Young, R. Rosenquist, X. Liu, A. Jethwa, K. S. Lee, J. Lewis, K. Putzker, C. Lutz, D. Rossi, A. Mokhir, T. Oellerich, K. Zirlik, M. Herling, F. Nguyen-Khac, C. Plass, E. Andersson, S. Mustjoki, C. von Kalle, A. D. Ho, M. Hensel, J. Dürig, I. Ringshausen, M. Zapatka, W. Huber, and T. Zenz. Drug-perturbation-based stratification of blood cancer. *Journal of Clinical Investigation*, 128(1):427–445, 2017.

- L. Du and I. Hu. An empirical Bayes method for chi-squared data. *Journal of the American Statistical Association*, 117(537):334–347, 2022.
- F. Dyson. A method for correcting series of parallax observations. *Monthly Notices of the Royal Astronomical Society*, 86:686, 1926.
- B. Efron. *Large-Scale Inference: Empirical Bayes Methods for Estimation, Testing, and Prediction*. Institute of Mathematical Statistics Monographs. Cambridge University Press, Cambridge, 2010.
- B. Efron. Tweedie’s formula and selection bias. *Journal of the American Statistical Association*, 106(496):1602–1614, 2011.
- B. Efron. Two modeling strategies for empirical Bayes estimation. *Statistical Science*, 29(2):285–301, 2014.
- B. Efron and C. Morris. Stein’s estimation rule and its competitors—an empirical Bayes approach. *Journal of the American Statistical Association*, 68(341):117–130, 1973.
- B. Efron, R. Tibshirani, J. D. Storey, and V. Tusher. Empirical Bayes analysis of a microarray experiment. *Journal of the American Statistical Association*, 96(456):1151–1160, 2001.
- B. Epstein. Some applications of the Mellin transform in statistics. *The Annals of Mathematical Statistics*, 19(3):370–379, 1948.
- J. Fan. On the optimal rates of convergence for nonparametric deconvolution problems. *The Annals of Statistics*, 19(3):1257–1272, 1991.
- A. M. Fink and M. Jodeit, Jr. On Chebyshev’s other inequality. In *Institute of Mathematical Statistics Lecture Notes - Monograph Series*, pages 115–120. Institute of Mathematical Statistics, Hayward, CA, 1984.
- L. Fu, B. Gang, G. M. James, and W. Sun. Heteroscedasticity-adjusted ranking and thresholding for large-scale multiple testing. *Journal of the American Statistical Association*, 117(538):1028–1040, 2022.
- W. Gabcke. *Neue Herleitung und explizite Restabschätzung der Riemann-Siegel-Formel*. PhD thesis, Georg-August-Universität zu Göttingen, Göttingen, 1979.
- S. Ghosal and A. W. van der Vaart. Entropies and rates of convergence for maximum likelihood and Bayes estimation for mixtures of normal densities. *The Annals of Statistics*, 29(5):1233–1263, 2001.
- E. Greenshtein and Y. Ritov. Generalized maximum likelihood estimation of the mean of parameters of mixtures, with applications to sampling, 2022. URL <http://arxiv.org/abs/2107.09296>.
- J. Gu and R. Koenker. Empirical Bayesball remixed: Empirical Bayes methods for longitudinal data. *Journal of Applied Econometrics*, 32(3):575–599, 2017a.
- J. Gu and R. Koenker. Unobserved heterogeneity in income dynamics: An empirical Bayes perspective. *Journal of Business & Economic Statistics*, 35(1):1–16, 2017b.

- J. Gu and R. Koenker. Invidious comparisons: Ranking and selection as compound decisions. *Econometrica*, 91(1):1–41, 2023.
- J. Gu and S. Shen. Oracle and adaptive false discovery rate controlling methods for one-sided testing: Theory and application in treatment effect evaluation. *The Econometrics Journal*, 21(1):11–35, 2018.
- D. Houde, S. A. Berkowitz, and J. R. Engen. The utility of Hydrogen/Deuterium exchange mass spectrometry in biopharmaceutical comparability studies. *Journal of Pharmaceutical Sciences*, 100(6):2071–2086, 2011.
- J. G. Hwang and P. Liu. Optimal tests shrinking both means and variances applicable to microarray data analysis. *Statistical Applications in Genetics and Molecular Biology*, 9(1), 2010.
- N. Ignatiadis and S. Wager. Confidence intervals for nonparametric empirical Bayes analysis. *Journal of the American Statistical Association*, 117(539):1149–1166, 2022.
- N. Ignatiadis, S. Saha, D. L. Sun, and O. Muralidharan. Empirical Bayes mean estimation with nonparametric errors via order statistic regression on replicated data. *Journal of the American Statistical Association*, (forthcoming), 2021.
- N. Ignatiadis, R. Wang, and A. Ramdas. E-values as unnormalized weights in multiple testing. 2022. URL <http://arxiv.org/abs/2204.12447>.
- S. Jana, Y. Polyanskiy, and Y. Wu. Optimal empirical Bayes estimation for the Poisson model via minimum-distance methods, 2022. URL <http://arxiv.org/abs/2209.01328>.
- I. B. Jeffery, D. G. Higgins, and A. C. Culhane. Comparison and evaluation of methods for generating differentially expressed gene lists from microarray data. *BMC Bioinformatics*, 7(1):359, 2006.
- N. P. Jewell. Mixtures of exponential distributions. *The Annals of Statistics*, pages 479–484, 1982.
- W. Jiang. On general maximum likelihood empirical Bayes estimation of heteroscedastic IID normal means. *Electronic Journal of Statistics*, 14(1):2272–2297, 2020.
- W. Jiang and C.-H. Zhang. General maximum likelihood empirical Bayes estimation of normal means. *The Annals of Statistics*, 37(4):1647–1684, 2009.
- K. Kammers, R. N. Cole, C. Tiengwe, and I. Ruczinski. Detecting significant changes in protein abundance. *EuPA Open Proteomics*, 7:11–19, 2015.
- R. W. Keener. *Theoretical Statistics*. Springer Texts in Statistics. Springer New York, New York, NY, 2010.
- M. K. Kerr, M. Martin, and G. A. Churchill. Analysis of variance for gene expression microarray data. *Journal of Computational Biology*, 7(6):819–837, 2000.
- J. Kiefer and J. Wolfowitz. Consistency of the maximum likelihood estimator in the presence of infinitely many incidental parameters. *The Annals of Mathematical Statistics*, 27(4): 887–906, 1956.

- Y. Kim, P. Carbonetto, M. Stephens, and M. Anitescu. A fast algorithm for maximum likelihood estimation of mixture proportions using sequential quadratic programming. *Journal of Computational and Graphical Statistics*, 29(2):261–273, 2020.
- B. Klaus and S. Reisenauer. An end to end workflow for differential gene expression using Affymetrix microarrays. *F1000Research*, 5:1384, 2018.
- R. Koenker and J. Gu. REBayes : An R Package for Empirical Bayes Mixture Methods. *Journal of Statistical Software*, 82(8), 2017.
- R. Koenker and I. Mizera. Convex optimization, shape constraints, compound decisions, and empirical Bayes rules. *Journal of the American Statistical Association*, 109(506): 674–685, 2014.
- Y. Kwon and Z. Zhao. On F-modelling-based empirical Bayes estimation of variances. *Biometrika*, page asac019, 2022.
- C. Laufer, B. Fischer, M. Billmann, W. Huber, and M. Boutros. Mapping genetic interactions in human cancer cells with RNAi and multiparametric phenotyping. *Nature Methods*, 10(5):427–431, 2013.
- B. Laurent and P. Massart. Adaptive estimation of a quadratic functional by model selection. *The Annals of Statistics*, 28(5):1302–1338, 2000.
- E. L. Lehmann and J. P. Romano. *Testing Statistical Hypotheses*. Springer Texts in Statistics. Springer New York, New York, NY, 2005.
- B. G. Lindsay. The geometry of mixture likelihoods, part II: The exponential family. *The Annals of Statistics*, 11(3):783–792, 1983.
- B. G. Lindsay. Using empirical partially Bayes inference for increased efficiency. *The Annals of Statistics*, 13(3):914–931, 1985.
- I. Lönnstedt. *Empirical Bayes Methods for DNA Microarray Data*. PhD thesis, Matematiska institutionen / Uppsala University, Department of Mathematics / Uppsala University, Department of Mathematics, 2005.
- I. Lönnstedt and T. Speed. Replicated microarray data. *Statistica sinica*, pages 31–46, 2002.
- M. I. Love, W. Huber, and S. Anders. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology*, 15(12):550, 2014.
- M. Lu and M. Stephens. Variance adaptive shrinkage (*vash*): Flexible empirical Bayes estimation of variances. *Bioinformatics*, page btw483, 2016.
- M. Lu and M. Stephens. Empirical Bayes estimation of normal means, accounting for uncertainty in estimated standard errors, 2019. URL <http://arxiv.org/abs/1901.10679>.
- J. Maksimovic, B. Phipson, and A. Oshlack. A cross-package Bioconductor workflow for analysing methylation array data. *F1000Research*, 5(1281), 2017.
- P. McCullagh. A note on partially Bayes inference for generalized linear models. Technical Report 284, Department of Statistics, University of Chicago, 1990.

- X.-L. Meng. Posterior predictive p -values. *The Annals of Statistics*, 22(3):1142–1160, 1994.
- MOSEK ApS. *The MOSEK Optimization Suite Manual, Version 9.2*, 2020. URL <https://www.mosek.com/>.
- O. Muralidharan. High dimensional exponential family estimation via empirical Bayes. *Statistica Sinica*, pages 1217–1232, 2012.
- M. A. Newton, C. M. Kendzierski, C. S. Richmond, F. R. Blattner, and K. W. Tsui. On differential variability of expression ratios: Improving statistical inference about gene expression changes from microarray data. *Journal of Computational Biology*, 8(1):37–52, 2001.
- M. A. Newton, A. Noueir, D. Sarkar, and P. Ahlquist. Detecting differential gene expression with a semiparametric hierarchical mixture method. *Biostatistics*, 5(2):155–176, 2004.
- B. Phipson, S. Lee, I. J. Majewski, W. S. Alexander, and G. K. Smyth. Robust hyperparameter estimation protects against hypervariable genes and improves power to detect differential expression. *The Annals of Applied Statistics*, 10(2):946–963, 2016.
- A. Ploner, S. Calza, A. Gusnanto, and Y. Pawitan. Multidimensional local false discovery rate for microarray studies. *Bioinformatics*, 22(5):556–565, 2006.
- G. Pólya and G. Szegő. *Aufgaben Und Lehrsätze Aus Der Analysis. Zweiter Band: Funktionentheorie · Nullstellen Polynome · Determinanten · Zahlentheorie*. Springer Berlin Heidelberg, Berlin, Heidelberg, 1925.
- Y. Polyanskiy and Y. Wu. Self-regularizing property of nonparametric maximum likelihood estimator in mixture models, 2020. URL <http://arxiv.org/abs/2008.08244>.
- M. E. Ritchie, B. Phipson, D. Wu, Y. Hu, C. W. Law, W. Shi, and G. K. Smyth. Limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Research*, 43(7):e47–e47, 2015.
- H. Robbins. A generalization of the method of maximum likelihood: Estimating a mixing distribution (abstract). *The Annals of Mathematical Statistics*, 21:314–315, 1950.
- H. Robbins. Asymptotically subminimax solutions of compound statistical decision problems. In *Proceedings of the Second Berkeley Symposium on Mathematical Statistics and Probability*, volume 2, pages 131–149. University of California Press, 1951.
- H. Robbins. An empirical Bayes approach to statistics. In *Proceedings of the Third Berkeley Symposium on Mathematical Statistics and Probability, Volume 1: Contributions to the Theory of Statistics*, pages 157–163. The Regents of the University of California, 1956.
- H. Robbins. Estimating many variances. In *Statistical Decision Theory and Related Topics III*, pages 251–261. Elsevier, 1982.
- J. M. Robins, A. van der Vaart, and V. Ventura. Asymptotic distribution of p values in composite null models. *Journal of the American Statistical Association*, 95(452):1143–1156, 2000.

- M. D. Robinson, D. J. McCarthy, and G. K. Smyth. edgeR: A Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics*, 26(1): 139–140, 2010.
- G. K. Smyth. Linear models and empirical bayes methods for assessing differential expression in microarray experiments. *Statistical applications in genetics and molecular biology*, 3(1), 2004.
- J. A. Soloff, A. Guntuboyina, and B. Sen. Multivariate, Heteroscedastic Empirical Bayes via Nonparametric Maximum Likelihood, 2021. URL <http://arxiv.org/abs/2109.03466>.
- M. Stephens. False discovery rates: A new deal. *Biostatistics*, 18(2):275–294, 2016.
- J. D. Storey, J. E. Taylor, and D. Siegmund. Strong control, conservative point estimation and simultaneous conservative consistency of false discovery rates: A unified approach. *Journal of the Royal Statistical Society: Series B*, 66(1):187–205, 2004.
- W. Sun and T. T. Cai. Oracle and adaptive compound decision rules for false discovery rate control. *Journal of the American Statistical Association*, 102(479):901–912, 2007.
- W. Sun and A. C. McLain. Multiple testing of composite null hypotheses in heteroscedastic models. *Journal of the American Statistical Association*, 107(498):673–687, 2012.
- H. Teicher. Identifiability of mixtures. *The Annals of Mathematical Statistics*, 32(1):244–248, 1961.
- T. Terkelsen, M. Pernemalm, P. Gromov, A.-L. Børresen-Dale, A. Krogh, V. D. Haakensen, J. Lethiö, E. Papaleo, and I. Gromova. High-throughput proteomics of breast cancer interstitial fluid: Identification of tumor subtype-specific serologically relevant biomarkers. *Molecular Oncology*, 15(2):429–461, 2021.
- V. G. Tusher, R. Tibshirani, and G. Chu. Significance analysis of microarrays applied to the ionizing radiation response. *Proceedings of the National Academy of Sciences*, 98(9): 5116–5121, 2001.
- H. C. van Houwelingen. The role of empirical Bayes methodology as a leading principle in modern medical statistics. *Biometrical Journal*, 56(6):919–932, 2014.
- J. C. van Houwelingen. Monotone empirical Bayes tests for the continuous one-parameter exponential family. *The Annals of Statistics*, 4(5):981–989, 1976.
- J. E. Walsh. On the “Information” lost by using a t-test when the population variance is known. *Journal of the American Statistical Association*, 44(245):122–125, 1949.
- A. Weinstein. On permutation invariant problems in large-scale inference, 2021. URL <http://arxiv.org/abs/2110.06250>.
- A. Weinstein, Z. Ma, L. D. Brown, and C.-H. Zhang. Group-linear empirical Bayes estimates for a heteroscedastic normal mean. *Journal of the American Statistical Association*, 113(522):698–710, 2018.

- D. D. Weis. Comment on Houde, D.; Berkowitz, S. A.; Engen, J. R., The utility of Hydrogen/Deuterium exchange mass spectrometry in biopharmaceutical comparability studies. *J. Pharm. Sci.* 2011, 100, 2071-2086. *Journal of Pharmaceutical Sciences*, 108(2):807–810, 2019.
- D. M. Witten and R. Tibshirani. A comparison of fold-change and the t-statistic for microarray data analysis. Technical report, Stanford University, 2007.
- G. W. Wright and R. M. Simon. A random variance model for detection of differential gene expression in small microarray experiments. *Bioinformatics*, 19(18):2448–2455, 2003.
- D. Xie and M. Stephens. Discussion of “Confidence intervals for nonparametric empirical Bayes analysis”. *Journal of the American Statistical Association*, 117(539):1186–1191, 2022.
- C. Yu and P. D. Hoff. Adaptive multigroup confidence intervals with constant coverage. *Biometrika*, 105(2):319–335, 2018.
- C.-H. Zhang. Fourier methods for estimating mixing densities and distributions. *The Annals of Statistics*, 18(2):806–831, 1990.
- C.-H. Zhang. Generalized maximum likelihood estimation of normal mixture densities. *Statistica Sinica*, pages 1297–1318, 2009.
- Y. Zhang, J. Maksimovic, G. Naselli, J. Qian, M. Chopin, M. E. Blewitt, A. Oshlack, and L. C. Harrison. Genome-wide DNA methylation analysis identifies hypomethylated genes regulated by FOXP3 in human regulatory T cells. *Blood*, 122(16):2823–2836, 2013.
- Q. Zhao, Y. Chen, J. Wang, and D. S. Small. Powerful three-sample genome-wide design and robust statistical inference in summary-data Mendelian randomization. *International Journal of Epidemiology*, 48(5):1478–1492, 2019.
- Z. Zheng, A. M. Mergaert, I. M. Ong, M. A. Shelef, and M. A. Newton. MixTwice: Large-scale hypothesis testing for peptide arrays by variance mixing. *Bioinformatics*, 37(17):2637–2643, 2021.

A General remarks on proofs and notation

In the main text for many of the theoretical statements we provided two versions: one that applies to the hierarchical setting with $\sigma_i^2 \stackrel{\text{iid}}{\sim} G$ (2), and a second starred version that applies to the compound setting in which $\boldsymbol{\sigma} = (\sigma_1, \dots, \sigma_n)$ is fixed. For example, we have Theorem 7 and Theorem 7*. Furthermore, in the main text we clarified whether an expectation or probability is computed with respect to the hierarchical or compound setting by using the subscript G ($\mathbb{E}_G[\cdot]$, $\mathbb{P}_G[\cdot]$), resp. $\boldsymbol{\sigma}$ ($\mathbb{E}_{\boldsymbol{\sigma}}[\cdot]$, $\mathbb{P}_{\boldsymbol{\sigma}}[\cdot]$).

Throughout the supplement we often omit the subscript G and $\boldsymbol{\sigma}$, in which case an argument is to be understood as going through in both the hierarchical and compound settings. This allows us to provide unified proofs for the two versions of our theoretical results. We clarify within the proofs whenever a step requires different arguments for the two cases. For the unified treatment, we also introduce the following notation: we write $P_i^* = P_*(z, s^2)$ for the oracle p-values, where $P_*(z, s^2) = P_G(z, s^2)$ (3) when $\sigma_i^2 \sim G$ (2), and $P_*(z, s^2) = P_{\boldsymbol{\sigma}}(z, s^2)$ (18) in the compound setting. Analogously, we write $f_* = f_G$ (10) when $\sigma_i^2 \sim G$, and $f_* = f_{\boldsymbol{\sigma}}$ (20) in the compound setting.

Further notation: Throughout the proofs we write $A \lesssim_a B$ to denote that there exists a constant $C = C(a)$ that depends only on a such that $A \leq C \cdot B$. Analogously we also write $A \gtrsim_a B$.

B Properties of χ^2 -likelihood and precision parametrization

Often in the proofs it will be convenient to reparameterize the likelihood by the precisions $\tau_i^2 = 1/\sigma_i^2$:

$$p(s^2 | \tau^2) \equiv p(s^2 | \tau^2, \nu) := \frac{\nu^{\nu/2} (\tau^2)^{\nu/2}}{2^{\nu/2} \Gamma(\nu/2)} (s^2)^{\nu/2-1} \exp\left(-\frac{\nu \tau^2 s^2}{2}\right). \quad (25)$$

Lemma 18. The χ^2 -likelihood $p(s^2 | \tau^2) = p(s^2 | \tau^2, \nu)$ has the following properties ($\nu \geq 2$).

- A. $\sup \{p(s^2 | \tau^2) : s^2 > 0, 0 < \tau^2 < L^{-1}\} \lesssim_{\nu} L^{-1} \lesssim_{\nu, L} 1$.
- B. $\tau^2 \mapsto p(s^2 | \tau^2)$ is decreasing in the interval $[1/s^2, \infty)$ and increasing in $(0, 1/s^2)$.
- C. It holds that:

$$p(s^2 | \tau^2) \leq p(s^2 | 1/U) \lesssim_{\nu, U} \exp(-\nu s^2 / (2U)) \text{ for } s^2 \geq U \geq 1/\tau^2.$$

In particular, suppose $s^2 \geq 2(\kappa/\nu)U$ with $\kappa \geq \nu/2$ and that $\tau^2 \geq 1/U$, then:

$$p(s^2 | \tau^2) \leq p(s^2 | 1/U) \lesssim_{\nu, U} \exp(-\kappa).$$

- D. $\sup \left\{ \frac{\partial p(s^2 | \tau^2)}{\partial \tau^2} : \tau^2 > 0, s^2 > 0 \right\} \lesssim_{\nu} 1$.

Proof of Lemma 18. The following inequality will be useful:

$$x^k \exp(-x) \leq k^k \exp(-k) \text{ for all } x \geq 0, k \geq 0. \quad (26)$$

The inequality follows by observing that the LHS is maximized when $x = k$.

Let us now start with the claim in A. Note that,

$$p(s^2 | \tau^2) \lesssim_{\nu} (\tau^2)^{\nu/2} \left(\frac{\nu s^2}{2}\right)^{\nu/2-1} \exp\left(-\frac{\nu \tau^2 s^2}{2}\right) = \tau^2 \cdot \left\{ \left(\frac{\nu \tau^2 s^2}{2}\right)^{\nu/2-1} \exp\left(-\frac{\nu \tau^2 s^2}{2}\right) \right\}.$$

Apply inequality (26) with $k = \nu/2 - 1$ and conclude that A holds by using the fact that we seek to take the supremum over $\tau^2 \leq L^{-1}$.

We next take the derivative of (25) with respect to τ^2 :

$$\frac{\partial p(s^2 | \tau^2)}{\partial \tau^2} = \frac{\nu^{\nu/2} (s^2)^{\nu/2-1}}{2^{\nu/2} \Gamma(\nu/2)} \frac{\nu}{2} (\tau^2)^{\nu/2-1} (1 - \tau^2 s^2) \exp\left(-\frac{\nu \tau^2 s^2}{2}\right).$$

From here we can read off the monotonicity claimed in B. The result in C. also follows by the monotonicity property. Finally, for part D. we apply inequality (26) twice: once with $k = \nu/2$ and once with $k = \nu/2 - 1$. \square

C Proofs for Section 2

C.1 Proof of Proposition 1

Proof of Proposition 1. Ancillarity of S^2 follows since under (6), the distribution of S^2 does not depend on μ : S^2 is distributed as the product $\sigma^2 \cdot \mathcal{X}$ for independently drawn $\sigma^2 \sim G$, and $\nu \mathcal{X} \sim \chi_{\nu}^2$ (where neither G , nor χ_{ν}^2 depend on μ).

We continue with minimal sufficiency of (Z, S^2) . Sufficiency of (Z, S^2) is immediate in this setting (it is the full data generated from model (6)). Hence it remains to prove minimality; we will use the characterization of minimal sufficiency in terms of likelihood shapes [Keener, 2010, Theorem 3.11]. Write $p(z, s^2 | \mu)$ for the joint density of (z, s^2) on $\mathbb{R} \times \mathbb{R}_+$ under (6) with respect to the Lebesgue measure, where we make the dependence on μ explicit. Next, fix $x_1 = (z_1, s_1^2)$, $x_2 = (z_2, s_2^2)$ with $s_1^2, s_2^2 > 0$ and suppose that:

$$p(z_1, s_1^2 | \mu) = C(x_1, x_2) p(z_2, s_2^2 | \mu) \text{ for all } \mu, \quad (27)$$

with $C(\cdot)$ being a function of x_1, x_2 only (and not of μ). To prove minimality, it suffices to prove that (27) implies that $(z_1, s_1^2) = (z_2, s_2^2)$.

We will start by showing $z_1 = z_2$. Suppose otherwise, i.e., $z_1 \neq z_2$. Using the precision parametrization described in Supplement B, (27) is equivalent to:

$$\begin{aligned} & \int (s_1^2)^{\nu/2-1} (\tau^2)^{(\nu+1)/2} \exp(-\tau^2(z_1 - \mu)^2/2) \exp(-\nu \tau^2 s_1^2/2) dH(\tau^2) \\ &= C(x_1, x_2) \int (s_2^2)^{\nu/2-1} (\tau^2)^{(\nu+1)/2} \exp(-\tau^2(z_2 - \mu)^2/2) \exp(-\nu \tau^2 s_2^2/2) dH(\tau^2). \end{aligned}$$

Now pick μ such that: $(z_2 - \mu)^2 - (z_1 - \mu)^2 = (s_1^2 - s_2^2)\nu$. This is possible because we assumed $z_1 \neq z_2$. Then, the terms within $\exp(\cdot)$ on the LHS and RHS become identical, i.e., for some $\ell = \ell(x_1, x_2) > 0$:

$$\int (s_1^2)^{\nu/2-1} (\tau^2)^{(\nu+1)/2} \exp(-\ell \tau^2) dH(\tau^2) = C(x_1, x_2) \int (s_2^2)^{\nu/2-1} (\tau^2)^{(\nu+1)/2} \exp(-\ell \tau^2) dH(\tau^2).$$

This in turn implies that $C(x_1, x_2) = (s_1^2)^{\nu/2-1}/(s_2^2)^{\nu/2-1}$ and so:

$$\begin{aligned} & \int (\tau^2)^{(\nu+1)/2} \exp(-\tau^2/2 \{(z_1 - \mu)^2 + \nu s_1^2\}) dH(\tau^2) \\ &= \int (\tau^2)^{(\nu+1)/2} \exp(-\tau^2/2 \{(z_2 - \mu)^2 + \nu s_2^2\}) dH(\tau^2) \text{ for all } \mu \in \mathbb{R}. \end{aligned}$$

However, the above leads to a contradiction: suppose without loss of generality that $s_1^2 > s_2^2$ and pick $\mu = z_2$. Then $(z_1 - \mu)^2 + \nu s_1^2 > \nu s_2^2 = (z_2 - \mu)^2 + \nu s_2^2$, so that the above equality could not hold. Thus $z_1 = z_2$.

It remains to prove that also $s_1^2 = s_2^2$. We have already shown that $z_1 = z_2$. By translation, we may assume without loss of generality that in fact $z_1 = z_2 = 0$. Further let $u = \mu^2/2$ and $\tilde{C}(x_1, x_2) = C(x_1, x_2)((s_2/s_1)^2)^{\nu/2-1}$. Then:

$$\begin{aligned} & \int \exp(-u\tau^2)(\tau^2)^{(\nu+1)/2} \exp(-\tau^2\nu s_1^2/2) dH(\tau^2) \\ &= \tilde{C}(x_1, x_2) \int \exp(-u\tau^2)(\tau^2)^{(\nu+1)/2} \exp(-\tau^2\nu s_2^2/2) dH(\tau^2) \text{ for all } u \geq 0. \end{aligned}$$

On the LHS and RHS we have two finite measures supported on $(0, \infty)$ that have identical Laplace transforms. Hence by uniqueness of the Laplace transform (e.g., Billingsley [1995, Theorem 22.2]) these two measures must be identical, which is to say:

$$(\tau^2)^{(\nu+1)/2} \exp(-\tau^2\nu s_1^2/2) = \tilde{C}(x_1, x_2)(\tau^2)^{(\nu+1)/2} \exp(-\tau^2\nu s_2^2/2) \quad H\text{-almost surely.}$$

Since G , and so H is not degenerate, the above must hold for at least two values of $\tau \in (0, \infty)$, which implies that $s_1^2 = s_2^2$. \square

C.2 Proof of Proposition 2

Proof of Proposition 2. It will be convenient to parameterize the problem in terms of the precision $\tau^2 = 1/\sigma^2$ as in Supplement B. In what follows we may assume without loss of generality that $\mu = 0$; otherwise one can consider the translated family $Z \mapsto Z - \mu$.

The joint density of Z, S^2, τ^2 with respect to the Lebesgue measure on $\mathbb{R} \times \mathbb{R}_{>0} \times \mathbb{R}_{>0}$ is given by (up to a constant):

$$\begin{aligned} & p(z, s^2, \tau^2) \\ &= C (\tau^2)^{1/2} \exp\left(-\frac{z^2\tau^2}{2}\right) \cdot (s^2)^{\nu/2-1} (\tau^2)^{\nu/2} \exp\left(-\frac{\nu s^2\tau^2}{2}\right) \cdot (\tau^2)^{\nu_0/2-1} \exp\left(-\frac{\nu_0 s_0^2\tau^2}{2}\right) \\ &= C (s^2)^{\nu/2-1} (\tau^2)^{(\nu_0+\nu+1)/2-1} \exp\left(-\frac{(z^2 + \nu s^2 + \nu_0 s_0^2)\tau^2}{2}\right). \end{aligned}$$

Hence marginalizing over τ^2 :

$$\begin{aligned} p(z, s^2) &= \int_0^\infty p(z, s^2, \tau^2) d(\tau^2) \\ &= C (s^2)^{\nu/2-1} \int_0^\infty (\tau^2)^{(\nu_0+\nu+1)/2-1} \exp\left(-\frac{(z^2 + \nu s^2 + \nu_0 s_0^2)\tau^2}{2}\right) d(\tau^2) \\ &= C (s^2)^{\nu/2-1} \left(\frac{z^2 + \nu s^2 + \nu_0 s_0^2}{2}\right)^{-(\nu_0+\nu+1)/2} \Gamma(\nu_0/2 + \nu/2) \\ &= C' (s^2)^{\nu/2-1} (z^2 + \nu s^2 + \nu_0 s_0^2)^{-(\nu_0+\nu+1)/2}. \end{aligned}$$

We make the following parenthetical remarks at this point: first, by also marginalizing out over S^2 , we find that Z is marginally distributed according to the t -distribution with ν_0 degrees of freedom scaled by s_0 , that is,

$$Z \stackrel{\mathcal{D}}{=} s_0 \cdot T_{\nu_0}.$$

Second, the joint marginal distribution of (Z, S^2) is called the Student-Siegel distribution in [Aitchison and Dunsmore \[1975, Table 2.2., page 25\]](#).

Continuing with our proof, let $\tilde{t} = z/\sqrt{(\nu_0 s_0^2 + \nu s^2)/(\nu_0 + \nu)}$ and consider the mapping $\psi(z, s^2) = (\tilde{t}, s^2)$. Then,

$$|\det(\nabla\psi(z, s^2))| = |\partial\psi(z, s^2)/\partial z| = 1/\sqrt{(\nu_0 s_0^2 + \nu s^2)/(\nu_0 + \nu)}.$$

Hence:

$$\begin{aligned} p(\tilde{t}, s^2) &= p(\psi^{-1}(\tilde{t}, s^2))/|\det(\nabla\psi(z, s^2))| \\ &= C' (s^2)^{\nu/2-1} \sqrt{(\nu_0 s_0^2 + \nu s^2)/(\nu_0 + \nu)} \left(\tilde{t}^2 \frac{\nu_0 s_0^2 + \nu s^2}{\nu_0 + \nu} + \nu s^2 + \nu_0 s_0^2 \right)^{-(\nu_0 + \nu + 1)/2} \\ &= C'' (\tilde{t}^2 + \nu_0 + \nu)^{-(\nu_0 + \nu + 1)/2} \cdot (s^2)^{\nu/2-1} (\nu_0 s_0^2 + \nu s^2)^{-(\nu_0 + \nu)/2}. \end{aligned}$$

From the above result we can immediately read off the following two distribution statements: \tilde{T} and S^2 are independent (with respect to their marginal distribution—they are *not* independent conditional on τ^2). Second, $\tilde{T} \sim t_{\nu_0 + \nu}$. We also refer the reader to [Smyth \[2004, Section 4\]](#) for further discussion of the above distributional conclusions.

It remains to note the following two distributional equalities to conclude the proof of the first part of the proposition. First,

$$Z \mid (S^2 = s^2) \stackrel{\mathcal{D}}{=} \tilde{S} \cdot \tilde{T} \mid (S^2 = s^2),$$

since $Z = \tilde{T}\tilde{S}$. Second,

$$\tilde{S} \cdot \tilde{T} \mid (S^2 = s^2) \stackrel{\mathcal{D}}{=} \tilde{s} \cdot T_{\nu_0 + \nu},$$

since \tilde{S} is a deterministic function of S^2 , S^2 and \tilde{T} are independent, and $\tilde{T} \sim t_{\nu_0 + \nu}$.

For the second part of the proposition, fix $z \in \mathbb{R}$. Also write \tilde{s} , $\tilde{t} = z/\tilde{s}$ for the observed values of \tilde{S} and \tilde{T} . It holds that:

$$\begin{aligned} P_G(z, s^2) &\stackrel{(i)}{=} \mathbb{P}_G [|Z^{H_0}| \geq |z| \mid S^2 = s^2] \\ &\stackrel{(ii)}{=} \mathbb{P}_G [\tilde{s} \cdot T_{\nu_0 + \nu} \geq |z|] \\ &= \mathbb{P}_G [T_{\nu_0 + \nu} \geq |z|/\tilde{s}] \\ &= \mathbb{P}_G [T_{\nu_0 + \nu} \geq |\tilde{t}|] \\ &\stackrel{(iii)}{=} P_{\nu_0, s_0^2}^{\text{limma}}(z, s^2). \end{aligned}$$

In (i) we used the definition of the conditional p-values in (3). In (ii) we used the distributional result of the first part of the proposition. Finally, in (iii) we used the definition of the limma p-value in (5).

Finally, for completeness we also prove the RHS equality of (3). Note that,

$$\begin{aligned}\mathbb{P}_G [|Z^{H_0}| \geq |z| \mid S^2 = s^2] &= \mathbb{E}_G [\mathbb{P}_G [|Z| \geq |z| \mid \sigma^2, S^2 = s^2] \mid S^2 = s^2] \\ &= \mathbb{E}_G [\mathbb{P}_G [|Z^{H_0}| \geq |z| \mid \sigma^2] \mid S^2 = s^2] \\ &= \mathbb{E}_G [2(1 - \Phi(|z|/\sigma)) \mid S^2 = s^2].\end{aligned}$$

□

C.3 Proof of Proposition 3

Proof. As mentioned directly after the proposition, the first result is an application of the probability integral transform conditional on S^2 . The second result follows by iterated expectation:

$$\mathbb{P}_G [P_G(Z, S^2) \leq t] = \mathbb{E}_G [\mathbb{P}_G [P_G(Z, S^2) \leq t \mid S^2]] = \mathbb{E}_G [t] = t \text{ for all } t \in [0, 1].$$

It remains to prove the third result on the monotonicity. We use the precision parameterization of Supplement B wherein $\tau^2 \sim H$. We also let $\bar{\Phi} = 1 - \Phi$. By definition, $P_G(z, s^2)/2 = \mathbb{E}_G [\bar{\Phi}(|z|\tau) \mid S^2 = s^2]$. The monotonicity in terms of $|z|$ directly follows because $\bar{\Phi}(|z|\tau) \leq \bar{\Phi}(|z'|\tau)$ for any $|z'| \leq |z|$.

To show monotonicity with respect to s^2 , we first note that (as $S^2 \mid \tau^2 \sim \chi_\nu^2/(\nu\tau^2)$ and $\tau^2 \sim H$):

$$\frac{1}{2} P_G(z, s^2) = \frac{\int \bar{\Phi}(|z|\tau) (\tau^2)^{\nu/2} \exp\left(-\frac{\nu\tau^2 s^2}{2}\right) dH(\tau^2)}{\int (\tau^2)^{\nu/2} \exp\left(-\frac{\nu\tau^2 s^2}{2}\right) dH(\tau^2)}.$$

Let A_s be the probability measure that is absolutely continuous with respect to H with Radon-Nikodym derivative:

$$\frac{dA_s(\tau^2)}{dH} = (\tau^2)^{\nu/2} \exp\left(-\frac{\nu\tau^2 s^2}{2}\right) \Big/ \int (\tau^2)^{\nu/2} \exp\left(-\frac{\nu\tau^2 s^2}{2}\right) dH(\tau^2).$$

Then, taking the derivative of $P_G(z, s^2)$ with respect to s^2 , we find that:

$$\frac{\partial P_G(z, s^2)}{\partial s^2} = -\frac{\nu}{2} \left(\int \bar{\Phi}(|z|\tau) \tau^2 dA_s(\tau^2) - \int \bar{\Phi}(|z|\tau) dA_s(\tau^2) \int \tau^2 dA_s(\tau^2) \right).$$

To show the monotonicity claim, it suffices to show that $\partial P_G(z, s^2)/\partial s^2 \geq 0$, i.e., that $\text{Cov}_{A_s} [\bar{\Phi}(|z|\tau), \tau^2] \leq 0$ when $\tau^2 \sim A_s$. But note that $\tau^2 \mapsto \tau^2$ is non-decreasing and that $\tau^2 \mapsto \bar{\Phi}(|z|\tau)$ is non-increasing. Hence the conclusion follows by Chebyshev's "other" inequality (e.g., Fink and Jodeit [1984]). □

C.4 Proof of Proposition 4

Lemma 19. Let χ^2 be a random variable distributed according to the χ^2 -distribution with $\nu \geq 2$ degrees of freedom. Let F_ν be the distribution function of χ^2 . Then:

$$F_\nu(u)u^{-\nu/2} \rightarrow c(\nu) := \frac{1}{2^{\nu/2}\Gamma(\nu/2)} > 0 \text{ as } u \searrow 0.$$

Furthermore, there exists $C(\nu)$ such that:

$$F_\nu(u) \leq C(\nu)u^{\nu/2} \text{ for all } u > 0.$$

Proof. Let us write $f_\nu = F'_\nu$ for the density of χ^2 . Then, we have:

$$f_\nu(s) = \frac{1}{2^{\nu/2}\Gamma(\nu/2)} (s)^{\nu/2-1} \exp(-s/2) = \frac{1}{2^{\nu/2}\Gamma(\nu/2)} \sum_{k=0}^{\infty} \frac{(-1)^k}{2^k k!} s^{k+\nu/2-1}.$$

Integrating the above term by term (and noting that all series defined are absolutely convergent), we see that:

$$F_\nu(s) = \frac{1}{2^{\nu/2}\Gamma(\nu/2)} \sum_{k=0}^{\infty} \frac{(-1)^k}{2^k k!(k+\nu/2)} s^{k+\nu/2}.$$

From here we can immediately read off the first claim about the limit of $F_\nu(u)u^{-\nu/2}$ as $u \rightarrow 0$.

For the second claim, let us define the function $h(u) := C(\nu)u^{\nu/2} - F_\nu(u)$ (where we will pick $C(\nu)$ below). Note that $h(0) = 0$ and

$$h'(u) = (\nu/2)C(\nu)u^{\nu/2-1} - \frac{1}{2^{\nu/2}\Gamma(\nu/2)} u^{\nu/2-1} \exp(-u/2),$$

and so this can be made ≥ 0 for all $u > 0$ by choosing $C(\nu)$ large enough. \square

Proof of Proposition 4. The uniformity of the t-test p-values and their monotonicity are elementary. It remains to prove the failure of conditional uniformity, which is equivalent to the following statement:

$$\lim_{\delta \searrow 0} \mathbb{P}_G [P^{\text{ttest}}(Z, S^2) > t \mid S^2 \leq \delta] = 0 \text{ for all } t \in (0, 1].$$

We start by observing that the event $\{P^{\text{ttest}}(Z, S^2) > t\}$ is identical to the event $\{|Z| < qS\}$, where $q > 0$ is the $1 - t/2$ quantile of the t -distribution with ν -degrees of freedom. Hence:

$$\begin{aligned} \mathbb{P}_G [P^{\text{ttest}}(Z, S^2) > t \mid S^2 \leq \delta] &= \mathbb{P}_G [|Z| < qS \mid S^2 \leq \delta] \\ &= \mathbb{E}_G [\mathbb{P}_G [|Z| < qS \mid \sigma^2, S^2] \mid S^2 \leq \delta] \\ &= \mathbb{E}_G [2\Phi(qS/\sigma) \mid S^2 \leq \delta] \\ &= \frac{\mathbb{E}_G [(2\Phi(qS/\sigma) - 1) \mathbf{1}(S^2 \leq \delta)]}{\mathbb{P}_G [S^2 \leq \delta]}. \end{aligned}$$

As in Lemma 19, we next write χ^2 for a random variable distributed according to the χ^2 -distribution with ν degrees of freedom. We also write F_ν for its distribution function. Then:

$$\begin{aligned} \mathbb{P}_G [S^2 \leq \delta] &= \mathbb{P}_G [\chi^2 \leq \nu\delta/\sigma^2] \\ &= \mathbb{E}_G [\mathbb{P}_G [\chi^2 \leq \nu\delta/\sigma^2 \mid \sigma^2]] \\ &= \mathbb{E}_G [F_\nu(\nu\delta/\sigma^2)]. \end{aligned}$$

By Lemma 19, $F_\nu(u)u^{-\nu/2} \rightarrow c(\nu) > 0$ as $u \rightarrow 0$. Hence:

$$\begin{aligned} &\liminf_{\delta \rightarrow 0} \left\{ \mathbb{P}_G [S^2 \leq \delta] / \delta^{\nu/2} \right\} \\ &= \liminf_{\delta \rightarrow 0} \left\{ \mathbb{E}_G \left[(\sigma^2/\nu)^{-\nu/2} \cdot F_\nu(\nu\delta/\sigma^2) \cdot (\nu\delta/\sigma^2)^{-\nu/2} \right] \right\} \\ &\stackrel{(*)}{\geq} \mathbb{E}_G \left[\liminf_{\delta \rightarrow 0} \left\{ (\sigma^2/\nu)^{-\nu/2} \cdot F_\nu(\nu\delta/\sigma^2) \cdot (\nu\delta/\sigma^2)^{-\nu/2} \right\} \right] \\ &= c(\nu)\nu^{\nu/2} \mathbb{E}_G [\sigma^{-\nu}] > 0. \end{aligned}$$

In (*) we used Fatou's lemma. Next, we recall also by Lemma 19 that there exists $C(\nu)$ such that:

$$F_\nu(u) \leq C(\nu)u^{\nu/2} \text{ for all } u > 0.$$

Hence, we also get that:

$$\begin{aligned} & \limsup_{\delta \rightarrow 0} \left\{ \mathbb{E}_G \left[(2\Phi(qS/\sigma) - 1) \mathbf{1}(S^2 \leq \delta) \right] / \delta^{\nu/2} \right\} \\ & \leq \limsup_{\delta \rightarrow 0} \left\{ \mathbb{E}_G \left[\left(2\Phi(q\sqrt{\delta}/\sigma) - 1 \right) \mathbf{1}(S^2 \leq \delta) \right] / \delta^{\nu/2} \right\} \\ & = \limsup_{\delta \rightarrow 0} \left\{ \mathbb{E}_G \left[\left(2\Phi(q\sqrt{\delta}/\sigma) - 1 \right) F_\nu(\nu\delta/\sigma^2) / \delta^{\nu/2} \right] \right\} \\ & \lesssim_\nu \limsup_{\delta \rightarrow 0} \left\{ \mathbb{E}_G \left[\left(2\Phi(q\sqrt{\delta}/\sigma) - 1 \right) \sigma^{-\nu} \right] \right\} \\ & = 0, \end{aligned}$$

where in the last step we used dominated convergence. To conclude, we write:

$$\mathbb{P}_G [P^{\text{ttest}}(Z, S^2) > t \mid S^2 \leq \delta] = \frac{\mathbb{E}_G \left[(2\Phi(qS/\sigma) - 1) \mathbf{1}(S^2 \leq \delta) \right] / \delta^{\nu/2}}{\mathbb{P}_G [S^2 \leq \delta] / \delta^{\nu/2}},$$

and take limits as $\delta \rightarrow 0$. □

D Proofs for Section 3.1

D.1 Proof of Proposition 5

We will need the following result:

Lemma 20 (Pólya and Szegő [1925], Page 48). Let $P_1(x), \dots, P_n(x)$ be polynomials $\neq 0$ of degrees $m_1 - 1, \dots, m_n - 1$, where $m_i \geq 2$ for all i . Also let $a_1, \dots, a_n \in \mathbb{R}$. Then, the function

$$g(x) = \sum_{i=1}^n P_i(x) \exp(a_i x),$$

has at most $(\sum_{i=1}^n m_i) - 1$ zeros in \mathbb{R} (counting multiplicity).

Proof of Proposition 5. We prove these results when the problem is parametrized in terms of precisions τ^2 as in Section B. The results for this parametrization will then imply the statement of the proposition via the pushforward $\tau^2 \mapsto \frac{1}{\tau^2} = \sigma^2$.

A NPMLE with respect to the precision parametrization is defined as follows:

$$\hat{H} \in \operatorname{argmax} \left\{ \sum_{i=1}^n \log(f_H(S_i^2)) : H \text{ distribution supported on } (0, \infty) \right\}, \quad (28)$$

Let us start by defining \mathcal{H} as the class of all measures with mass ≤ 1 supported on $(0, \infty)$ and $\mathcal{H}_1 \subset \mathcal{H}$ the subclass of distributions.

We define the map,

$$\psi : \mathcal{H} \rightarrow \mathbb{R}^n, \text{ where } \psi(H) = (\psi_1(H), \dots, \psi_n(H)) \text{ with } \psi_i(H) = \int p(S_i^2 \mid \tau^2) dH(\tau^2).$$

The image $\psi(\mathcal{H}) \subset \mathbb{R}^n$ is convex. It is also compact by the Helly Bray selection theorem [Breiman, 1992, Theorem 6.8] and the fact that $\tau^2 \mapsto p(S_i^2 | \tau^2)$ is bounded and continuous (and also converges to 0 as $\tau \searrow 0$ or $\tau \nearrow +\infty$). Furthermore, there must exist unique $\hat{\beta} \in \psi(\mathcal{H})$ such that:

$$\hat{\beta} \in \operatorname{argmax} \{h(\beta) : \beta \in \psi(\mathcal{H})\}, \quad h(\beta) := \sum_{i=1}^n \log(\beta_i). \quad (29)$$

This follows by strict concavity of $h(\beta)$ as well as the fact that $h(\beta) \rightarrow -\infty$ as any $\beta_i \rightarrow 0$. In particular there also must exist $\hat{H} \in \mathcal{H}$ such that $\beta = \psi(\hat{H})$. Our next goal is to prove that \hat{H} is in fact a probability measure. Suppose otherwise, i.e., $\hat{H}((0, \infty)) = 1 - \eta$ for some $\eta > 0$. Let t be any point in the interval $(0, \infty)$ and consider the measure $\tilde{H} = \hat{H} + \eta\delta_t$. Then $\tilde{H} \in \mathcal{H}$ and $h(\psi(\tilde{H})) > h(\psi(\hat{H}))$ which is a contradiction. Thus $\hat{H} \in \mathcal{H}_1$.

For any two distributions $H_1, H_2 \in \mathcal{H}_1$ we define the directional derivative from H_2 to H_1 as:

$$D(H_1, H_2) = \lim_{\lambda \searrow 0} \frac{h(\psi(\lambda H_1 + (1-\lambda)H_2)) - h(\psi(H_2))}{\lambda} = \sum_{i=1}^n \frac{\psi_i(H_1) - \psi_i(H_2)}{\psi_i(H_2)}.$$

We next define the directional derivative from the measure \hat{H} to H . Since \hat{H} maximizes the log marginal likelihood, it must hold that $D(H_1, \hat{H}) \leq 0$ for any $H_1 \in \mathcal{H}_1$. In particular it must also hold for any Dirac measure $H_1 = \delta_{\tau^2}$ with $\tau^2 \in (0, \infty)$. Hence for such τ^2 we get by rearranging the inequality $D(\delta_{\tau^2}, \hat{H}) \leq 0$ that:

$$\xi(\tau^2) := \sum_{i=1}^n \frac{p(S_i^2 | \tau^2)}{f_{\hat{H}}(S_i^2)} \leq n \text{ for all } \tau^2 \in (0, \infty), \quad (30)$$

where above we introduced the function $\xi(\cdot)$. The integral of the LHS (i.e., of $\xi(\cdot)$) of the inequality with respect to \hat{H} is equal to $\sum_{i=1}^n \int 1 = n$, and so we get an equality of the integrated LHS and RHS of (30) w.r.t. \hat{H} . This implies that equality must hold \hat{H} almost surely, i.e.,

$$\xi(\tau^2) = \sum_{i=1}^n \frac{p(S_i^2 | \tau^2)}{f_{\hat{H}}(S_i^2)} = n \quad \hat{H}\text{-almost surely.} \quad (31)$$

Absorbing constants with respect to τ^2 for the i -th summand into a constant c_i , we may write $\xi(\cdot)$ as:

$$\xi(\tau^2) = \sum_{i=1}^n c_i (\tau^2)^{\nu/2} \exp\left(-\frac{\nu\tau^2 S_i^2}{2}\right).$$

At this point we note the following: the function $\tau^2 \mapsto \xi(\tau^2) - n$ is an exponential polynomial of the form considered in Lemma 20. Hence its number of zeros has to be finite, i.e., the support of \hat{H} must be finite (we will make this more precise below). Here we also used the fact that $\xi(\tau^2)$ cannot be constant, for example, $\xi(\tau^2) \rightarrow 0$ as $\tau^2 \rightarrow 0$.

Next, the derivative of ξ with respect to τ^2 is equal to:

$$\xi'(\tau^2) = \frac{d\xi(\tau^2)}{d(\tau^2)} = \sum_{i=1}^n c_i \cdot \frac{\nu}{2} (\tau^2)^{\nu/2-1} \exp\left(-\frac{\nu\tau^2 S_i^2}{2}\right) (1 - \tau^2 S_i^2). \quad (32)$$

From here we can derive our conclusion about the support of \widehat{H} . First we will prove that \widehat{H} has zero mass in the interval $(0, \min_i \{1/S_i^2\})$. By (32) it holds that $\xi'(\tau^2) > 0$ in the interval $(0, \min \{1/S_i^2\})$, that is, $\xi(\cdot)$ is strictly increasing. Now suppose there is $\tau^2 \in (0, \min_i \{1/S_i^2\})$ in the support of \widehat{H} . Then $\xi(\tau^2) = n$ by (31) and $\xi(\tau^2 + \varepsilon) > n$ for sufficiently small ε , in contradiction to (31). Thus we conclude that \widehat{H} assigns zero mass to $(0, \min_i \{1/S_i^2\})$. Analogously we can argue that \widehat{H} has zero mass in the interval $(\max_i \{1/S_i^2\}, +\infty)$ since on the latter interval it holds that $\xi'(\tau^2) < 0$, i.e., $\xi(\cdot)$ is strictly decreasing.

It remains to accomplish the following tasks: to provide an upper bound on the cardinality of the support of \widehat{H} and to prove its uniqueness.

Suppose the support of \widehat{H} consists of S points. Then for each support point the function $\xi(\tau^2)$ has a local maximum due to (30) and (31). Furthermore, between any consecutive support points, there must be a local minimum. Hence ξ' must have at least $2S - 1$ zeros. On the other hand, by (32) we see that $\xi'(\tau^2) = 0$ implies (for $\tau^2 > 0$) that:

$$\zeta(\tau^2) := \sum_{i=1}^n c_i \exp\left(-\frac{\nu\tau^2 S_i^2}{2}\right) (1 - \tau^2 S_i^2) = 0.$$

But ζ is of the form considered in Lemma 20 with $m_i = 2$ and hence can have at most $\sum_{i=1}^n m_i - 1 = 2n - 1$ zeros. Hence $2n - 1 \geq 2S - 1$, i.e., $S \leq n$.

It remains to prove the uniqueness of \widehat{H} . Suppose there is another maximizer \widetilde{H} of the log marginal likelihood. From the beginning of our argument, namely (29), we then know that it must be the case that

$$f_{\widetilde{H}}(S_i^2) = f_{\widehat{H}}(S_i^2) = \hat{\beta}_i \text{ for } i = 1, \dots, n. \quad (33)$$

Hence (31) is identical for both \widetilde{H} and \widehat{H} , and it must hold that $\widetilde{H}(S) = \widehat{H}(S) = 1$, where S is the set of roots in (31). Thus we can write:

$$\widehat{H} = \sum_{j=1}^S \widehat{\pi}_j \delta_{\tau_j^2}, \quad \widetilde{H} = \sum_{j=1}^S \widetilde{\pi}_j \delta_{\tau_j^2},$$

where we indexed the elements of S as $\{\tau_j^2\}$. By (33) we get:

$$\begin{aligned} & \sum_{j=1}^S (\widehat{\pi}_j - \widetilde{\pi}_j) p(S_i^2 | \tau_j^2) = 0 \text{ for } i = 1, \dots, n, \\ \implies & \sum_{j=1}^S (\widehat{\pi}_j - \widetilde{\pi}_j) (\tau_j^2)^{\nu/2} \exp\left(-\frac{\nu\tau_j^2 S_i^2}{2}\right) = 0 \text{ for } i = 1, \dots, n. \end{aligned}$$

Hence the function $s \mapsto \sum_{j=1}^S (\widehat{\pi}_j - \widetilde{\pi}_j) (\tau_j^2)^{\nu/2} \exp\left(-\frac{\nu\tau_j^2 s}{2}\right)$ has at least n zeros. On the other hand, by Lemma 20 with $m_i = 1$, it can have at most $S - 1$ zeros; otherwise it must be identically equal to 0. Suppose it is not identically to zero. Then $S - 1 \geq n$. But we already proved that $S \leq n$, which leads to a contradiction. Hence the above function is identically equal to zero, which means that $\widehat{\pi}_j = \widetilde{\pi}_j$ for all j , that is, $\widehat{G} = \widetilde{G}$. \square

D.2 Proof of Theorems 7 and 7*

We first define the class of marginal densities of S^2 when $\sigma^2 \in [L, U]$ almost surely with $0 < L < U < \infty$:

$$\mathcal{F} \equiv \mathcal{F}(L, U, \nu) := \{f_G(\cdot) = f_G(\cdot; \nu) : G([L, U]) = 1\}. \quad (34)$$

For the proof of the theorem we will use the following lemmata (which we prove later).

Lemma 21. For all $\varepsilon > 0$ sufficiently small (that is, with $\varepsilon \leq c \equiv c(\nu, L, U)$), it holds that:

$$\log(N(\varepsilon, \mathcal{F}(L, U, \nu), \|\cdot\|_\infty)) \lesssim_{\nu, L, U} |\log(\varepsilon)|^2.$$

Lemma 22. Suppose $\sigma_i^2 \leq U$ almost surely for all i . Then, for any $C > 0, \gamma > 0, n \in \mathbb{N}_{\geq 1}$ and any $B \geq 4U$, it holds that:

$$\mathbb{P} \left[\prod_{i: S_i^2 > B} \frac{C(S_i^2)^2}{\eta B^2} \geq \exp(2\gamma) \right] \leq \exp \left(-2\gamma + \frac{nC(B^2 + 8UB + 32U^2)}{\eta B^2} \exp(-B/(4U)) \right).$$

Proof of Theorems 7 and 7.* Recall from Supplement A that we write $f_* = f_G$, resp. $f_* = f_\sigma$ depending on whether we are studying the hierarchical or compound setting. Our goal is to obtain high-probability bounds on

$$\mathbb{P}[A_n], \text{ where } A_n := \{\mathfrak{H}(f_{\hat{G}}, f_*) \geq \varepsilon_n\}. \quad (35)$$

We will figure out later what the choice of ε_n should be.

Our strategy is as follows. Consider the following class of marginal densities:

$$\mathcal{F}(\varepsilon_n, L, \bar{U}, \nu) = \left\{ f_{\tilde{G}}(\cdot) = f_{\tilde{G}}(\cdot; \nu) : \tilde{G}([L, \bar{U}]) = 1, \mathfrak{H}(f_{\tilde{G}}, f_*) \geq \varepsilon_n \right\}. \quad (36)$$

This is essentially the class of mixture densities in (34) subject to the additional constraint that their Hellinger distance to f_* is sufficiently large. Furthermore, let $\mathcal{S} = \{f_j : j \in \mathcal{J}\}$, $\mathcal{J} = \{1, \dots, J\}$, $J = \#\mathcal{S}$, be a *proper* $(\|\cdot\|_\infty, \eta)$ -cover of (36).²³ Lemma 21 already provides a cover of a larger class of functions. However, the latter cover is not a proper cover for (36), that is the cover will contain elements that do not necessarily lie in the class (36). However, by a standard argument we can turn an improper cover to a proper cover at the cost of doubling the radius (i.e., an $\eta/2$ -cover in Lemma 21 yields a η -cover for (36)). Hence we get that:

$$\log J \lesssim_{\nu, L, \bar{U}} |\log(\eta)|^2.$$

Hence, on the event A_n (35), there must exist $\hat{j} \in \mathcal{J}$ such that $\|f_{\hat{j}} - f_{\hat{G}}\|_\infty \leq \eta$. This further implies that on A_n :

$$f_{\hat{G}}(s^2) \leq f_{\hat{j}}(s^2) + \eta \leq \max_{j \in \mathcal{J}} f_j(s^2) + \eta \text{ for all } s^2 > 0.$$

From here we proceed as follows. We introduce the following function parameterized by B :

$$\eta(z) := \eta \mathbf{1}\{|z| \leq B\} + \eta \frac{B^2}{z^2} \mathbf{1}\{|z| > B\}, \text{ for } z > 0.$$

²³By “proper cover”, we mean that the centers of the cover are themselves elements of (36).

Notice that by construction

$$\int_0^\infty \eta(z) dz = \eta B + \eta B = 2\eta B,$$

and also

$$f_{\hat{G}}(s^2) \leq \max_{j \in \mathcal{J}} \{f_j(s^2) + \eta(s^2)\} \text{ if } s^2 \leq B, \text{ and } f_{\hat{G}}(s^2) \leq C(\nu, \underline{L}) \text{ otherwise.}$$

The boundedness required for the ‘‘otherwise’’ part of the branch above follows by Lemma 18A with $C(\nu, \underline{L}) > 0$ being a constant that only depends on ν and \underline{L} . For any f , write:

$$L_n(f, f_*) = \prod_{i=1}^n \frac{f(S_i^2)}{f_*(S_i^2)}.$$

Since $f_{\hat{G}}$ is the NPMLE, it must hold that $L_n(f_{\hat{G}}, f_*) \geq 1$. Next, on the event A_n :

$$\begin{aligned} L_n(f_{\hat{G}}, f_*) &\leq \prod_{i: S_i^2 \leq B} \frac{f_{\hat{j}}(S_i^2) + \eta(S_i^2)}{f_*(S_i^2)} \prod_{i: S_i^2 > B} \frac{C(\nu, \underline{L})}{f_*(S_i^2)} \\ &= \prod_{i=1}^n \frac{f_{\hat{j}}(S_i^2) + \eta(S_i^2)}{f_*(S_i^2)} \prod_{i: S_i^2 > B} \frac{C(\nu, \underline{L})}{f_{\hat{j}}(S_i^2) + \eta(S_i^2)} \\ &\leq \prod_{i=1}^n \frac{f_{\hat{j}}(S_i^2) + \eta(S_i^2)}{f_*(S_i^2)} \prod_{i: S_i^2 > B} \frac{C(\nu, \underline{L})}{\eta(S_i^2)}. \end{aligned}$$

Pick any γ . We get:

$$\begin{aligned} \mathbb{P} [\mathfrak{H}(f_{\hat{G}}, f_*) \geq \varepsilon_n] &= \mathbb{P} \left[\mathfrak{H}(f_{\hat{G}}, f_*) \geq \varepsilon_n, L_n(f_{\hat{G}}, f_*) \geq 1 \right] \\ &\leq \mathbb{P} \left[\left\{ \left(\sup_{j \in \mathcal{J}} \prod_{i=1}^n \frac{f_j(S_i^2) + \eta(S_i^2)}{f_*(S_i^2)} \right) \cdot \left(\prod_{i: S_i^2 > B} \frac{C(\nu, \underline{L})}{\eta(S_i^2)} \right) \right\} \geq 1 \right] \\ &\leq \mathbb{P} \left[\left(\sup_{j \in \mathcal{J}} \prod_{i=1}^n \frac{f_j(S_i^2) + \eta(S_i^2)}{f_*(S_i^2)} \right) \geq \exp(-2\gamma) \right] + \mathbb{P} \left[\prod_{i: S_i^2 > B} \frac{C(\nu, \underline{L})}{\eta(S_i^2)} \geq \exp(2\gamma) \right]. \end{aligned}$$

In Lemma 22 we already bounded the second probability. Hence it remains to bound the

first term. Observe that,

$$\begin{aligned}
& \mathbb{P} \left[\left(\sup_{j \in \mathcal{J}} \prod_{i=1}^n \frac{f_j(S_i^2) + \eta(S_i^2)}{f_*(S_i^2)} \right) \geq \exp(-2\gamma) \right] \\
& \leq J \sup_{j \in \mathcal{J}} \mathbb{P} \left[\prod_{i=1}^n \frac{f_j(S_i^2) + \eta(S_i^2)}{f_*(S_i^2)} \geq \exp(-2\gamma) \right] \\
& = J \sup_{j \in \mathcal{J}} \mathbb{P} \left[\prod_{i=1}^n \sqrt{\frac{f_j(S_i^2) + \eta(S_i^2)}{f_*(S_i^2)}} \geq \exp(-\gamma) \right] \\
& \leq J \sup_{j \in \mathcal{J}} \left\{ \exp(\gamma) \prod_{i=1}^n \mathbb{E} \left[\sqrt{\frac{f_j(S_i^2) + \eta(S_i^2)}{f_*(S_i^2)}} \right] \right\} \\
& \leq J \sup_{j \in \mathcal{J}} \left\{ \exp(\gamma) \exp \left(\sum_{i=1}^n \left\{ \mathbb{E} \left[\sqrt{\frac{f_j(S_i^2) + \eta(S_i^2)}{f_*(S_i^2)}} \right] - 1 \right\} \right) \right\} \\
& \stackrel{(*)}{=} J \sup_{j \in \mathcal{J}} \left\{ \exp(\gamma) \exp \left(n \left\{ \int_0^\infty \sqrt{f_j(z) + \eta(z)} \sqrt{f_*(z)} dz - 1 \right\} \right) \right\} \\
& \stackrel{(**)}{\leq} J \exp(\gamma) \exp \left(-n\mathfrak{H}^2(f_j, f_*) + n\sqrt{2\eta B} \right) \\
& \leq \exp \left\{ -n\varepsilon_n^2 + n\sqrt{2\eta B} + \gamma + \log J \right\}.
\end{aligned}$$

In (*) we have used the following argument, which is slightly different in the hierarchical setting compared to the compound setting. In the hierarchical setting:

$$\sum_{i=1}^n \mathbb{E}_G \left[\sqrt{\frac{f_j(S_i^2) + \eta(S_i^2)}{f_G(S_i^2)}} \right] = \sum_{i=1}^n \int_0^\infty \sqrt{\frac{f_j(z) + \eta(z)}{f_G(z)}} f_G(z) dz = n \int_0^\infty \sqrt{f_j(z) + \eta(z)} \sqrt{f_G(z)} dz.$$

In the compound setting:

$$\begin{aligned}
\sum_{i=1}^n \mathbb{E}_{\sigma_i} \left[\sqrt{\frac{f_j(S_i^2) + \eta(S_i^2)}{f_\sigma(S_i^2)}} \right] &= \sum_{i=1}^n \int_0^\infty \sqrt{\frac{f_j(z) + \eta(z)}{f_\sigma(z)}} p(z | \sigma_i^2) dz \\
&= n \int_0^\infty \sqrt{\frac{f_j(z) + \eta(z)}{f_\sigma(z)}} \left(\frac{1}{n} \sum_{i=1}^n p(z | \sigma_i^2) \right) dz \\
&= n \int_0^\infty \sqrt{f_j(z) + \eta(z)} \sqrt{f_\sigma(z)} dz.
\end{aligned}$$

In (***) we used the following argument:

$$\begin{aligned}
\int_0^\infty \sqrt{f_j(z) + \eta(z)} \sqrt{f_*(z)} dz - 1 &\leq \int_0^\infty \left(\sqrt{f_j(z)f_*(z)} + \sqrt{\eta(z)f_*(z)} \right) dz - 1 \\
&\leq -\mathfrak{H}^2(f_j, f_*) + \left(\int_0^\infty \eta(z) dz \right)^{1/2} \left(\int_0^\infty f_*(z) dz \right)^{1/2} \\
&= -\mathfrak{H}^2(f_j, f_*) + \sqrt{2\eta B}.
\end{aligned}$$

We are now ready to pick all parameters. First we pick $\gamma = (\log n)^2$ and $\eta = 1/n^2$. Then $\log J \lesssim_{\nu, L, \bar{U}} (\log n)^2$. Also let $B = 8\bar{U}(1 \vee \log n)$. Finally, we pick:

$$\varepsilon_n^2 = C \frac{(\log n)^2}{n},$$

where we pick C sufficiently large so that $n\varepsilon_n^2$ dominates all other terms that are of order $(\log n)^2$. \square

D.3 Proof of Lemmata 21 and 22

Lemma 23. Take any G supported on $[0, U]$ and suppose $\nu \geq 2$. Then, for any $\varepsilon \in (0, 1/e)$ it holds that:

$$\mathbb{P}_G [S^2 \geq t(\varepsilon)] \leq \varepsilon, \text{ where } t(\varepsilon) = 4U |\log \varepsilon|.$$

Proof of Lemma 23. Let $t' \geq 1$ and define $t = 4Ut'$. Since $\nu \geq 2$, it follows that

$$\nu + 2\sqrt{\nu t'} + 2t' \leq (3\nu + 2)t' \leq 4\nu t' = (\nu/U)t.$$

Write $S^2 = \chi_\nu^2 \sigma^2 / \nu$ where χ_ν^2 follows the χ^2 -distribution with ν degrees of freedom. Then, as $\sigma^2 \leq U$ with probability 1,

$$\mathbb{P}_G [S^2 \geq t] = \mathbb{P}_G \left[\frac{\sigma^2}{\nu} \chi_\nu^2 \geq t \right] \leq \mathbb{P}_G \left[\chi_\nu^2 \geq \frac{\nu}{U} t \right] \leq \mathbb{P}_G \left[\chi_\nu^2 \geq \nu + 2\sqrt{\nu t'} + 2t' \right] \stackrel{(*)}{\leq} \exp(-t').$$

(*) follows by standard concentration results for χ^2 -random variables (e.g., [Laurent and Massart \[2000, Lemma 1\]](#)). Hence, for $\varepsilon \in (0, 1/e)$ and $t' = |\log \varepsilon| \geq 1$, i.e., for $t(\varepsilon) = 4U |\log \varepsilon|$, it follows that $\mathbb{P}_G [S^2 \geq t(\varepsilon)] \leq \varepsilon$. \square

Proof of Lemma 21. We will use the precision parametrization (Supplement B) of the priors for most of the proof. Throughout we fix H supported on $[U^{-1}, L^{-1}]$.

Step 1: Let $\varepsilon > 0$ be given. In subsequent steps we will only seek to control the marginal densities for $s^2 \in (0, B]$. We choose this B in such a way that:

$$f_H(s^2) \lesssim_{\nu, L, U} \varepsilon \text{ for all } s^2 \geq B.$$

By Lemma 18C., we have that for $\kappa \geq \nu/2$ and $B = 2(\kappa\nu)U$, then $p(s^2 | \tau^2) \lesssim_{\nu, U} \exp(-\kappa)$. Hence we can pick $\kappa = \max\{\nu/2, \log(1/\varepsilon)\}$ and

$$B = (2/\nu)U \max\{\nu/2, \log(1/\varepsilon)\}.$$

Step 2: In addition to H , take another distribution H' supported on $[U^{-1}, L^{-1}]$. Then, for $s^2 \leq B$ and any $K \in \mathbb{N}_{\geq 1}$ it holds that:

$$\begin{aligned} & f_H(s^2) - f_{H'}(s^2) \\ &= C(\nu)(s^2)^{\nu/2-1} \int (\tau^2)^{\nu/2} \exp\left(-\frac{\nu\tau^2 s^2}{2}\right) (dH(\tau^2) - dH'(\tau^2)) \\ &\stackrel{(*)}{=} C(\nu)(s^2)^{\nu/2-1} \int (\tau^2)^{\nu/2} \left(R_K(\tau, s) + \sum_{k=0}^K \frac{(-1)^k \nu^k (\tau^2)^k (s^2)^k}{k! 2^k} \right) (dH(\tau^2) - dH'(\tau^2)) \\ &= C(\nu)(s^2)^{\nu/2-1} \int \left(R_K(\tau, s) (\tau^2)^{\nu/2} + \sum_{k=0}^K \frac{(-1)^k \nu^k (\tau^2)^{k+\nu/2} (s^2)^k}{k! 2^k} \right) (dH(\tau^2) - dH'(\tau^2)). \end{aligned}$$

In (*) we applied Taylor's theorem. The remainder term R_K therein satisfies:

$$|R_K(\tau, s)| \leq \frac{(\nu\tau^2 s^2/2)^{K+1}}{(K+1)!} \leq \left(\frac{e\nu\tau^2 s^2/2}{K+1}\right)^{K+1} \leq \left(\frac{2\nu L^{-1}B}{K+1}\right)^{K+1},$$

where we used that $(K+1)! \geq ((K+1)/e)^{K+1}$ from Stirling's approximation. We now build upon a moment matching argument. Suppose H' matches the following (fractional) moments:

$$\int (\tau^2)^{k+\nu/2} dH(\tau^2) = \int (\tau^2)^{k+\nu/2} dH'(\tau^2) \text{ for } k = 0, \dots, K. \quad (37)$$

Then:

$$\begin{aligned} |f_H(s^2) - f_{H'}(s^2)| &= C(\nu)(s^2)^{\nu/2-1} \int R_K(\tau, s) (\tau^2)^{\nu/2} (dH(\tau^2) - dH'(\tau^2)) \\ &\lesssim_\nu (B)^{\nu/2-1} L^{-\nu/2} \left(\frac{2\nu L^{-1}B}{K+1}\right)^{K+1}. \end{aligned}$$

Now let $K \geq \max\{2^{\nu/2}e\nu L^{-1}B, \log(1/\varepsilon)\}$. Then noting that $K^{\nu/2-1} \leq (2^{\nu/2-1})^{K+1}$, we get:

$$\begin{aligned} &(B)^{\nu/2-1} L^{-\nu/2} \left(\frac{2\nu L^{-1}B}{K+1}\right)^{K+1} \\ &\lesssim_{\nu, L} K^{\nu/2-1} \left(\frac{2\nu L^{-1}B}{K+1}\right)^{K+1} \leq \left(\frac{2^{\nu/2}\nu L^{-1}B}{K+1}\right)^{K+1} \leq e^{-(K+1)} \leq \varepsilon. \end{aligned}$$

Step 3: By Caratheodory's theorem, there exists a discrete distribution H' supported on $[U^{-1}, L^{-1}]$ with at most $K' = K + 2$ points of support that matches the above moments of H .

Step 4: In Step 3 we constructed H' of the following form:

$$H' = \sum_{j=1}^{K'} \pi_j \delta_{x_j}, \quad (38)$$

where $U^{-1} \leq x_1 \leq \dots \leq x_{K'} \leq L^{-1}$ and $\pi \in \text{Simplex}(K')$, the probability simplex embedded in $\mathbb{R}^{K'}$. Now take another measure of the form:

$$H'' = \sum_{k=1}^{K'} p_k \delta_{y_k},$$

where $U^{-1} \leq y_1 \leq \dots \leq y_{K'} \leq L^{-1}$ and $p \in \text{Simplex}(K')$. Then:

$$\begin{aligned} |f_{H'}(s^2) - f_{H''}(s^2)| &= \left| \sum_{j=1}^{K'} \pi_j p(s^2 | x_j) - \sum_{j=1}^{K'} p_j p(s^2 | y_j) \right| \\ &= \left| \sum_{j=1}^{K'} \pi_j (p(s^2 | x_j) - p(s^2 | y_j)) + \sum_{j=1}^{K'} (\pi_j - p_j) p(s^2 | y_j) \right| \\ &\lesssim_{\nu, L, U} \max_{j=1, \dots, K'} \{|x_j - y_j|\} + \|\pi - p\|_1. \end{aligned}$$

In the last step we used Lemma 18, parts A. and D.

Step 5 (putting everything together): The remaining crux of the argument is an explicit construction of the cover we seek.

To this end, first let Ξ be an $(\varepsilon, |\cdot|)$ -cover of $[U^{-1}, L^{-1}]$. By picking an equidistant grid we see that $|\Xi| \lesssim_{L,U} 1/\varepsilon$.

Second, let $\mathcal{S}(\text{Simplex}(K'))$ be an $(\varepsilon, \|\cdot\|_1)$ cover of $\text{Simplex}(K')$. A volume argument yields $|\mathcal{S}(\text{Simplex}(K'))| \leq (5/\varepsilon)^{K'}$ for $\varepsilon \leq 1$.

Combining the above, we let:

$$\mathcal{H} := \left\{ H'' = \sum_{j=1}^{K'} p_j \delta_{y_j} : p \in \mathcal{S}(\text{Simplex}(K')), y_j \in \Xi \right\}.$$

Combining the previous steps, we see that for any $f = f_H \in \mathcal{F}$, there exists $H'' \in \mathcal{H}$ such that:

$$\|f_H - f_{H''}\|_\infty \lesssim_{\nu,L,U} \varepsilon,$$

and so our cover of (34) consists of all elements of the form $f_H(\cdot) = \int p(\cdot | \tau^2) dH''(\tau^2)$ with $H'' \in \mathcal{H}$. It only remains to count $|\mathcal{H}|$. We have that:

$$|\mathcal{H}| \leq \binom{\Xi}{K'} \cdot |\mathcal{S}(\text{Simplex}(K'))| \leq \left(\frac{C(U,L)}{\varepsilon} \right)^{K'} \cdot \left(\frac{5}{\varepsilon} \right)^{K'}.$$

Hence:

$$\log(|\mathcal{H}|) \lesssim_{L,U} K' \log(1/\varepsilon).$$

The above is a $C(\nu, L, U) \cdot \varepsilon$ cover for a constant $C(\nu, L, U)$, and so:

$$\log(N(\varepsilon, \mathcal{F}(L, U, \nu), \|\cdot\|_\infty)) \lesssim_{\nu,L,U} K' \log(C(\nu, L, U)/\varepsilon) \lesssim_{\nu,L,U} K' \log(1/\varepsilon).$$

Observe that our construction allows to choose $K' \lesssim_{\nu,L,U} \log(1/\varepsilon)$ to conclude. \square

Proof of Lemma 22.

$$\begin{aligned} & \mathbb{P} \left[\prod_{i: S_i^2 > B} \frac{C(S_i^2)^2}{\eta B^2} \geq \exp(2\gamma) \right] \\ & \leq \exp(-2\gamma) \prod_{i=1}^n \mathbb{E} \left[\left(\frac{C(S_i^2)^2}{\eta B^2} \right)^{\mathbf{1}(S_i^2 > B)} \right] \\ & \leq \exp(-2\gamma) \prod_{i=1}^n \mathbb{E} \left[\left(1 + \mathbf{1}(S_i^2 > B) \frac{C(S_i^2)^2}{\eta B^2} \right) \right] \\ & \leq \exp(-2\gamma) \exp \left(\frac{C}{\eta B^2} \sum_{i=1}^n \mathbb{E} \left[\mathbf{1}(S_i^2 > B) (S_i^2)^2 \right] \right). \end{aligned}$$

Hence it suffices to bound the inner expectation. We have that:

$$\begin{aligned}
\mathbb{E} \left[\mathbf{1}(S_i^2 > B) (S_i^2)^2 \right] &= 2 \int_0^\infty t \mathbb{P} [S_i^2 \mathbf{1}(S_i^2 > B) \geq t] dt \\
&= 2 \left(\int_0^B t \mathbb{P} [S_i^2 \geq B] dt + \int_B^\infty t \mathbb{P} [S_i^2 \geq t] dt \right) \\
&\stackrel{(*)}{\leq} B^2 \exp(-B/(4U)) + 2 \int_B^\infty t \exp(-t/(4U)) dt \\
&= (B^2 + 8UB + 32U^2) \exp(-B/(4U)).
\end{aligned}$$

The tail bound (*) follows precisely as in the proof of Lemma 23 and we use the fact here that $B \geq 4U$. \square

D.4 Proof of Corollary 8

Proof. Our proof operates on the following event

$$A := \bigcup_{k=1}^\infty \bigcap_{n=k}^\infty A_n, \text{ where } A_n := \left\{ \mathfrak{H}(f_{\widehat{G}_n}, f_*) < \frac{C \log n}{\sqrt{n}} \right\},$$

with the constant C in the statement of Theorem 7. We make the dependence of \widehat{G} on n explicit by writing $\widehat{G}_n \equiv \widehat{G}$. By Theorem 7 we also have that $\mathbb{P}_G [A_n^c] \leq \exp(-c(\log n)^2)$, where A_n^c denotes the complement of A_n , and so:

$$\sum_{n=1}^\infty \mathbb{P}_G [A_n^c] < \infty.$$

Thus, by the first Borel-Cantelli Lemma (e.g., [Billingsley, 1995, Theorem 4.3]), it follows that $\mathbb{P}_G [A] = 1$. We will prove that $\widehat{G}_n \xrightarrow{d} G$ on the event A .

First by construction it holds that $(\widehat{G}_n)_{n \geq 1}$ is a tight (since \widehat{G} is supported on the compact set $[\underline{L}, \bar{U}]$). Next take any subsequence \widehat{G}_{n_k} such that $\widehat{G}_{n_k} \xrightarrow{d} \widetilde{G}$ as $k \rightarrow \infty$ for some probability measure \widetilde{G} . By Billingsley [1995, Chapter 25, corollary on page 337], the weak convergence $\widehat{G}_n \xrightarrow{d} G$ will follow if we can show that $\widetilde{G} = G$.

To see this, first note that for any $u > 0$:

$$f_{\widehat{G}_{n_k}}(u) = \int p(u | \sigma^2) d\widehat{G}_{n_k}(\sigma^2) \rightarrow \int p(u | \sigma^2) d\widetilde{G}(\sigma^2) = f_{\widetilde{G}}(u) \text{ as } k \rightarrow \infty.$$

Here we used the definition of weak convergence along with the facts that $\sigma^2 \mapsto p(u | \sigma^2)$ is continuous and bounded on $[\underline{L}, \bar{U}]$. Hence by Scheffe's theorem [Billingsley, 1995, Theorem 16.12.] applied to the Lebesgue measure on $(0, \infty)$, we get convergence in total variation distance (TV),

$$\text{TV}(f_{\widehat{G}_{n_k}}, f_{\widetilde{G}}) := \frac{1}{2} \int \left| f_{\widehat{G}_{n_k}}(u) - f_{\widetilde{G}}(u) \right| du \rightarrow 0 \text{ as } k \rightarrow \infty.$$

On the other hand:

$$\text{TV}(f_{\widehat{G}_{n_k}}, f_G) \stackrel{(*)}{\leq} \sqrt{2} \mathfrak{H}(f_{\widehat{G}_{n_k}}, f_G) \stackrel{(**)}{\rightarrow} 0 \text{ as } k \rightarrow \infty.$$

(**) follows because on the event A it holds that:

$$\limsup_{k \rightarrow \infty} \mathfrak{H}(f_{\tilde{G}^{n_k}}, f_G) \leq \limsup_{k \rightarrow \infty} \frac{C \log n_k}{\sqrt{n_k}} \rightarrow 0, \text{ as } k \rightarrow \infty.$$

(*) is a standard result that follows e.g., by the Cauchy-Schwarz inequality. Hence by the triangle inequality, we find that $\text{TV}(f_{\tilde{G}}, f_G) = 0$, which implies that $f_G(u) = f_{\tilde{G}}(u)$ for almost all $u > 0$, and so by continuity, for all $u > 0$. Let H and \tilde{H} be the push-forwards of G and \tilde{G} under the map $\sigma^2 \mapsto 1/\sigma^2$. In other words, we are switching to the precision parametrization of Supplement B. Then it follows by Teicher [1961, Section 4, Example 1] that $H = \tilde{H}$. This in turn implies that $G = \tilde{G}$, and so we conclude with the proof of this corollary. \square

E Proofs for Section 3.2

E.1 Proof of Proposition 9

Proof. To simplify notation suppose that $z \geq 0$. Then:

$$\begin{aligned} & 2\mathbb{E} [(1 - \Phi(z/\sigma)) | S^2 = s^2] \\ &= 2 \frac{1}{f_G(s^2; \nu)} \int_0^\infty (1 - \Phi(z/\sigma)) p(s^2 | \sigma^2, \nu) dG(\sigma^2) \\ &= 2 \frac{1}{f_G(s^2; \nu)} \int_0^\infty \left(\int_z^\infty \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{u^2}{2\sigma^2}\right) du \right) p(s^2 | \sigma^2) dG(\sigma^2) \\ &= \sqrt{\frac{2}{\pi}} \frac{(\nu/2)^{\nu/2}}{\Gamma(\nu/2)} \frac{(s^2)^{\nu/2-1}}{f_G(s^2; \nu)} \int_z^\infty \int_0^\infty (\sigma^2)^{-1/2} (\sigma^2)^{-\nu/2} \exp\left(-\frac{\nu s^2 + u^2}{2\sigma^2}\right) dG(\sigma^2) du \\ &= \sqrt{\frac{2}{\pi}} \frac{(\nu/2)^{\nu/2}}{\Gamma(\nu/2)} \frac{(s^2)^{\nu/2-1}}{f_G(s^2; \nu)} \int_z^\infty \int_0^\infty (\sigma^2)^{-\frac{\nu+1}{2}} \exp\left(-\frac{(\nu+1)\{(\nu s^2 + u^2)/(\nu+1)\}}{2\sigma^2}\right) dG(\sigma^2) du \\ &= \sqrt{\frac{2}{\pi}} \frac{(\nu/2)^{\nu/2}}{\Gamma(\nu/2)} \frac{(s^2)^{\nu/2-1}}{f_G(s^2; \nu)} \int_z^\infty \Gamma\left(\frac{\nu+1}{2}\right) \left(\frac{2}{\nu+1}\right)^{\frac{\nu+1}{2}} \left(\frac{\nu s^2 + u^2}{\nu+1}\right)^{-\frac{\nu-1}{2}} f_G\left(\frac{\nu s^2 + u^2}{\nu+1}; \nu+1\right) du \\ &= \sqrt{\frac{2}{\pi}} \frac{\sqrt{2}(1+1/\nu)^{-\nu/2} \Gamma((\nu+1)/2)}{(\nu+1)^{1/2} \Gamma(\nu/2)} \frac{(s^2)^{\nu/2-1}}{f_G(s^2; \nu)} \int_z^\infty \left(\frac{\nu s^2 + u^2}{\nu+1}\right)^{1/2-\nu/2} f_G\left(\frac{\nu s^2 + u^2}{\nu+1}; \nu+1\right) du \\ &= \frac{(1+1/\nu)^{-\nu/2} \Gamma((\nu+1)/2)}{\sqrt{\pi}(\nu+1)^{-1/2} \Gamma(\nu/2)} \frac{(s^2)^{\nu/2-1}}{f_G(s^2; \nu)} \int_0^\infty \frac{(t^2)^{-\frac{\nu-1}{2}}}{\sqrt{(\nu+1)t^2 - \nu s^2}} f_G(t^2; \nu+1) \mathbb{1}\left(t^2 \geq \frac{\nu s^2 + z^2}{\nu+1}\right) d(t^2). \end{aligned}$$

\square

E.2 Proof of Lemma 10

As mentioned in the main text, our proof of Lemma 10 builds upon Mellin transform analysis. For conciseness we do not provide the necessary background and definitions here and instead refer to Brenner Miguel et al. [2021]; our notation in the proof matches the notation therein. We do however provide references for any mathematical property of the Mellin transform that we invoke.

Our proof also requires the following preliminary lemmata.

Lemma 24 (Theorem 4.4.3. in [Gabcke \[1979\]](#)). Let $\kappa \geq 1$. Then, for any $T \geq \kappa$ it holds that:

$$\int_T^\infty t^{\kappa-1} \exp(-t) dt \leq \kappa T^{\kappa-1} \exp(-T).$$

Proof. The proof appears in [Gabcke \[1979\]](#). Since the latter is written in German, we provide the argument here as well. In the case $\kappa = 1$, the above in fact holds with equality (by integrating the exponential function). Henceforth we assume that $\kappa > 1$. By partial integration:

$$\int_T^\infty t^{\kappa-1} \exp(-t) dt = T^{\kappa-1} \exp(-T) + (\kappa - 1) \int_T^\infty t^{\kappa-2} \exp(-t) dt.$$

Hence we now look at the second summand of the RHS. Consider the transformation $x = 1/t$, then:

$$\int_T^\infty t^{\kappa-2} \exp(-t) dt = \int_0^{1/T} x^{-\kappa} \exp(-1/x) dx \stackrel{(*)}{\leq} \frac{1}{T} T^\kappa \exp(-T) = T^{\kappa-1} \exp(-T).$$

Combining with the preceding display yields the result of the lemma.

It remains to justify (*). Define $h(x) = x^{-\kappa} \exp(-1/x)$. This function has derivative $h'(x) = \exp(-1/x) (-\kappa x^{-\kappa-1} + x^{-\kappa-2})$. Hence $h'(x) > 0$ for all $x \in (0, 1/T)$, that is, h is non-decreasing in the interval $(0, 1/T)$. \square

Lemma 25 (Corollary 1.4.4 in [Andrews et al. \[1999\]](#)). Fix $a_1 \leq a_2$. Then, for any $a \in [a_1, a_2]$, it holds that as $|b| \rightarrow \infty$:

$$|\Gamma(a + ib)| = \sqrt{2\pi} |b|^{a-1/2} \exp(-\pi |b|/2) \{1 + O(1/|b|)\},$$

and the constant in $O(\cdot)$ only depends on a_1 and a_2 .

Proof of Lemma 10. Below we carry out a slightly more general argument which allows to bound the L^2 distance weighted by the function $x \mapsto x^{2c-1}$. We fix c for now, and suppose that it satisfies the following inequality:²⁴

$$c > 1 - \nu/2. \tag{39}$$

For the results in this slightly more general setting, we also consider the more general form of our assumption on G, G' , namely that:

$$\mathbb{E}_G \left[(\sigma^2)^{c-1} \right] \leq M, \quad \mathbb{E}_{G'} \left[(\sigma^2)^{c-1} \right] \leq M, \tag{40}$$

which again reduces to the assumption of the lemma when $c = 1/2$.

Let $X_\nu \sim \chi_\nu^2/\nu$. Then, S^2 in (8) is equal in distribution to $\sigma^2 X_\nu$. We write $p_\nu(\cdot)$ for the Lebesgue density of X_ν , that is

$$p_\nu(x) = \frac{\nu^{\nu/2}}{2^{\nu/2} \Gamma(\nu/2)} x^{\nu/2-1} \exp\left(-\frac{\nu x}{2}\right), \quad \text{for } x > 0.$$

²⁴The result stated in the lemma corresponds to the choice $c = 1/2$. For $c = 1/2$, the inequality (39) is satisfied, since we only consider $\nu \geq 2$.

We first verify that the following integrals are finite:

$$\int_0^\infty x^{c-1} p_\nu(x) dx < \infty, \quad \int_0^\infty x^{2c-1} p_\nu^2(x) dx < \infty.$$

For the first integral:

$$\int_0^\infty x^{c-1} p_\nu(x) dx = \frac{\nu^{\nu/2}}{2^{\nu/2} \Gamma(\nu/2)} \int_0^\infty x^{(c+\nu/2-1)-1} \exp\left(-\frac{\nu x}{2}\right) dx = \left(\frac{2}{\nu}\right)^{c-1} \frac{\Gamma(c+\nu/2-1)}{\Gamma(\nu/2)} < \infty.$$

Above we used the fact that $c + \nu/2 - 1 > 0$ by (39). For the second integral:

$$\int_0^\infty x^{2c-1} p_\nu^2(x) dx \lesssim_\nu \int_0^\infty x^{2c-1} \left(x^{\nu/2-1} \exp\left(-\frac{\nu x}{2}\right)\right)^2 dx = \int_0^\infty x^{(2c+\nu-2)-1} \exp(-\nu x) dx,$$

hence this integral is also finite since $2c + \nu - 2 > 0$ by (39). Using the above result, (40) and Lemma 2.2a) in Butzer and Stefan [1999], we also get that:

$$\int_0^\infty x^{2c-1} f_G^2(x; \nu) ds \leq M^2 \int_0^\infty x^{2c-1} p_\nu^2(x) dx < \infty.$$

Furthermore:

$$\int_0^\infty x^{c-1} f_G(x; \nu) dx = \mathbb{E}_G [(S^2)^{c-1}] = \mathbb{E}_G \left[(\sigma^2)^{c-1} X_\nu^{c-1} \right] = \mathbb{E}_G \left[(\sigma^2)^{c-1} \right] \mathbb{E} [X_\nu^{c-1}] < \infty.$$

The above steps establish integrability conditions that simplify the application of the Mellin transform theory.

The next calculation computes the Mellin transform of p_ν for c that satisfies (39) and arbitrary $t \in \mathbb{R}$ (its definition being shown in the first line below). Define

$$\begin{aligned} \mathcal{M}_c[p_\nu](t) &:= \mathbb{E}_{X \sim p_\nu} [X^{c-1+it}] \\ &= \int_0^\infty x^{c+\nu/2-1+it-1} \frac{\nu^{\nu/2}}{2^{\nu/2} \Gamma(\nu/2)} \exp\left(-\frac{\nu x}{2}\right) dx \\ &= \int_0^\infty \left(\frac{2}{\nu}\right)^{c-1+it} u^{c+\nu/2-1+it-1} \frac{1}{\Gamma(\nu/2)} \exp(-u) du \\ &= \left(\frac{2}{\nu}\right)^{c-1+it} \frac{\Gamma(c+\nu/2-1+it)}{\Gamma(\nu/2)}. \end{aligned}$$

Write $r(t) := \mathcal{M}_c[p_{\nu+1}](t)/\mathcal{M}_c[p_\nu](t)$, which by the preceding calculation has absolute value equal to

$$|r(t)| = \left(\frac{\nu}{\nu+1}\right)^{c-1} \frac{\Gamma(\nu/2)}{\Gamma((\nu+1)/2)} \frac{|\Gamma(c+(\nu+1)/2-1+it)|}{|\Gamma(c+\nu/2-1+it)|}.$$

Hence, using Lemma 25, we see that there exist positive constants $k = k(\nu, c)$ and $K = K(\nu, c)$ that depend only on ν and c such that:

$$|r(t)| \leq K(\nu, c) |T|^{1/2} \quad \text{for all } |t| \leq |T|, \text{ and all } |T| \geq k(\nu, c). \quad (41)$$

Let us also compute the Mellin transform of $f_G(\cdot; \nu)$:

$$\mathcal{M}_c[f_G(\cdot; \nu)](t) = \mathbb{E}_G [(S^2)^{c-1+it}] = \mathbb{E}_G \left[(\sigma^2)^{c-1+it} X_\nu^{c-1+it} \right] = \mathbb{E}_G \left[(\sigma^2)^{c-1+it} \right] \mathcal{M}_c[p_\nu](t).$$

This computation, (40), and Lemma 25 furnish the existence of positive constants $\ell = \ell(\nu, c)$ and $L = L(\nu, c, M)$ such that:

$$|\mathcal{M}_c[f_G(\cdot; \nu + 1)](t)| \leq L(\nu, c, M) |t|^{c+\nu/2-1} \exp(-\pi |t|/2) \text{ for all } |t| \geq \ell(\nu, c). \quad (42)$$

Before the main argument, we still need the following result that relates the Mellin transform of $f_G(\cdot; \nu)$ and $f_G(\cdot; \nu + 1)$ via the ratio $r(\cdot)$ defined above.

$$\begin{aligned} \mathcal{M}_c[f_G(\cdot; \nu + 1)](t) &= \mathbb{E}_G \left[(\sigma^2)^{c-1+it} \right] \mathcal{M}_c[p_{\nu+1}](t) \\ &= \mathbb{E}_G \left[(\sigma^2)^{c-1+it} \right] \mathcal{M}_c[p_\nu](t) r(t) \\ &= \mathcal{M}_c[f_G(\cdot; \nu)](t) \cdot r(t). \end{aligned}$$

We are ready to proceed with our main argument. In (*) below we apply the Plancherel isometry for the Mellin transform, see e.g., Lemma 2.3. in Butzer and Stefan [1999].

$$\begin{aligned} & \int_0^\infty x^{2c-1} (f_G(x; \nu + 1) - f_{G'}(x; \nu + 1))^2 dx \\ \stackrel{(*)}{=} & \frac{1}{2\pi} \int_{\mathbb{R}} |\mathcal{M}_c[f_G(\cdot; \nu + 1)](t) - \mathcal{M}_c[f_{G'}(\cdot; \nu + 1)](t)|^2 dt \\ = & \frac{1}{2\pi} \int_{\mathbb{R}} |r(t)|^2 |\mathcal{M}_c[f_G(\cdot; \nu)](t) - \mathcal{M}_c[f_{G'}(\cdot; \nu)](t)|^2 dt \\ = & \frac{1}{2\pi} \left(\int_{|t| \geq T} |r(t)|^2 |\mathcal{M}_c[f_G(\cdot; \nu)](t) - \mathcal{M}_c[f_{G'}(\cdot; \nu)](t)|^2 dt + \int_{|t| < T} \dots dt \right) \\ = & \text{I} + \text{II}, \end{aligned}$$

where I, resp. II refer to the two integrals and we will pick $T \geq \ell(\nu, c) \vee k(\nu, c) \vee (2c + \nu)$ later. We analyze the two integrals in turn. By (42),

$$\begin{aligned} \text{I} &= \frac{1}{2\pi} \int_{|t| \geq T} |\mathcal{M}_c[f_G(\cdot; \nu + 1)](t) - \mathcal{M}_c[f_{G'}(\cdot; \nu + 1)](t)|^2 dt \\ &\leq 2L(\nu, c, M) \int_{|t| \geq T} |t|^{2c+\nu-1} \exp(-\pi |t|) dt \\ &\stackrel{(**)}{\lesssim_{\nu, c, M}} T^{2c+\nu-1} \exp(-\pi T). \end{aligned}$$

For (**) we used Lemma 24 also noting that $2c + \nu > 2 \geq 1$ by (39). For the second integral, by (41),

$$\begin{aligned} \text{II} &= \frac{1}{2\pi} \int_{|t| < T} |r(t)|^2 |\mathcal{M}_c[f_G(\cdot; \nu)](t) - \mathcal{M}_c[f_{G'}(\cdot; \nu)](t)|^2 dt \\ &\leq \frac{1}{2\pi} K(\nu, c)^2 T \int_{\mathbb{R}} |\mathcal{M}_c[f_G(\cdot; \nu)](t) - \mathcal{M}_c[f_{G'}(\cdot; \nu)](t)|^2 dt \\ &= K(\nu, c)^2 \cdot T \cdot \int_0^\infty x^{2c-1} (f_G(x; \nu) - f_{G'}(x; \nu))^2 dx. \end{aligned}$$

In the second line we used (41) and for the last step we used the Plancherel isometry for the Mellin transform. Now let us call:

$$\rho(\nu, c) := \int_0^\infty x^{2c-1} (f_G(x; \nu) - f_{G'}(x; \nu))^2 dx.$$

We have proved that:

$$\rho(\nu + 1, c) \lesssim_{\nu, c, M} T^{2c+(\nu-1)} \exp(-\pi T) + T\rho(\nu, c).$$

The above bound holds for any c that satisfies (39). We now specialize to the case discussed in the lemma, namely $c = 1/2$. Then, the above general inequality reduces to:

$$\rho(\nu + 1, 1/2) \lesssim_{\nu, M} T^\nu \exp(-\pi T) + T\rho(\nu, 1/2).$$

Using e.g., (26), we get $T^\nu \exp(-\pi T) \lesssim_\nu \exp(-T)$, so that:

$$\rho(\nu + 1, 1/2) \lesssim_{\nu, M} \exp(-T) + T\rho(\nu, 1/2).$$

Let us pick

$$T = |\log(\rho(\nu, 1/2))| \vee \ell(\nu, 1/2) \vee k(\nu, 1/2) \vee (1 + \nu).$$

Notice that

$$T \lesssim_\nu 1 + |\log(\rho(\nu, 1/2))|,$$

and also that

$$\exp(-T) \leq \exp(\log(\rho(\nu, 1/2))) = \rho(\nu, 1/2).$$

Hence,

$$\rho(\nu + 1, 1/2) \lesssim_{\nu, M} (1 + |\log(\rho(\nu, 1/2))|)\rho(\nu, 1/2).$$

The above is the claim made in the lemma, since by definition:

$$\rho(\nu, 1/2) = \|f_G(\cdot; \nu) - f_{G'}(\cdot; \nu)\|_{L^2}^2, \quad \rho(\nu + 1, 1/2) = \|f_G(\cdot; \nu + 1) - f_{G'}(\cdot; \nu + 1)\|_{L^2}^2.$$

□

E.3 Proof of Theorems 11 and 11*

Proof of Theorems 11 and 11.* We start by the following observation. For any distribution \tilde{G} such that $\sigma_i^2 \geq \underline{L}$, it holds that $P_{\tilde{G}}(z, s^2) \geq 2\bar{\Phi}(|z|/\underline{L}^{1/2})$. Let $z_{1-\zeta/2}$ be the $(1 - \zeta/2)$ -quantile of the standard normal distribution. Then for $|z| \leq \underline{z} := \underline{L}^{1/2}z_{1-\zeta/2}$, it holds that:

$$P_{\tilde{G}}(z, s^2) \geq 2\bar{\Phi}(z_{1-\zeta/2}) = \zeta.$$

Henceforth (as mentioned in Supplement A) we write $P_*(z, s^2)$ to denote either $P_G(z, s^2)$ (hierarchical setting) or $P_\sigma(z, s^2) = P_{G(\sigma)}(z, s^2)$ (compound setting).

By the above considerations we can conclude that $P_*(z, s^2) \wedge \zeta - P_{\tilde{G}}(Z_i, S_i^2) \wedge \zeta = 0$ for $|z| \leq \underline{L}^{1/2}z_{1-\zeta/2}$. On the other hand, for all i , it holds that

$$|P_*(Z_i, S_i^2) \wedge \zeta - P_{\tilde{G}}(Z_i, S_i^2) \wedge \zeta| \leq |P_*(Z_i, S_i^2) - P_{\tilde{G}}(Z_i, S_i^2)|.$$

This holds since the projection of a number in $[0, 1]$ to the set $[0, \zeta]$ is a contraction. Combining the above two observations and letting $\tilde{Z}_i := \text{sign}(Z_i) \cdot (|Z_i| \vee \underline{L}^{1/2}z_{1-\zeta/2})$, we find the following:

$$\begin{aligned} |P_*(Z_i, S_i^2) \wedge \zeta - P_{\tilde{G}}(Z_i, S_i^2) \wedge \zeta| &\leq |P_*(\tilde{Z}_i, S_i^2) - P_{\tilde{G}}(\tilde{Z}_i, S_i^2)| \\ &\leq \sup_{z: |z| \geq \underline{z}} |P_*(z, S_i^2) - P_{\tilde{G}}(z, S_i^2)|. \end{aligned} \quad (43)$$

We now need one argument that is slightly different for Theorem 11 versus Theorem 11*. To bound the LHS in the inequality stated in Theorem 11* we build on the inequality

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E}_{\sigma} [|P_*(Z_i, S_i^2) \wedge \zeta - P_{\widehat{G}}(Z_i, S_i^2) \wedge \zeta|] \leq \frac{1}{n} \sum_{i=1}^n \mathbb{E}_{\sigma} \left[\sup_{z:|z| \geq z} |P_*(z, S_i^2) - P_{\widehat{G}}(z, S_i^2)| \right],$$

which follows by averaging (43) over $i = 1, \dots, n$. To bound the LHS of the inequality in the hierarchical version of Theorem 11, let us first fix $j \in \{1, \dots, n\}$. We have that:

$$\begin{aligned} \mathbb{E}_G [|P_*(Z_j, S_j^2) \wedge \zeta - P_{\widehat{G}}(Z_j, S_j^2) \wedge \zeta|] &\leq \mathbb{E}_G \left[\sup_{z:|z| \geq z} |P_*(z, S_j^2) - P_{\widehat{G}}(z, S_j^2)| \right] \\ &= \frac{1}{n} \sum_{i=1}^n \mathbb{E}_G \left[\sup_{z:|z| \geq z} |P_*(z, S_i^2) - P_{\widehat{G}}(z, S_i^2)| \right]. \end{aligned}$$

For the last equality we used the fact that under (6), (S_1^2, \dots, S_n^2) are all exchangeable. Thus:

$$\max_{i=1, \dots, n} \mathbb{E}_G [|P_*(Z_i, S_i^2) \wedge \zeta - P_{\widehat{G}}(Z_i, S_i^2) \wedge \zeta|] \leq \frac{1}{n} \sum_{i=1}^n \mathbb{E}_G \left[\sup_{z:|z| \geq z} |P_*(z, S_i^2) - P_{\widehat{G}}(z, S_i^2)| \right].$$

In either case, the preceding arguments establish that it suffices to bound

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[\sup_{z:|z| \geq z} |P_*(z, S_i^2) - P_{\widehat{G}}(z, S_i^2)| \right],$$

and we devote the rest of this proof to this task. Let A be the event:

$$A := \{\mathfrak{H}(f_{\widehat{G}}, f_*) < C \log n / \sqrt{n}\}, \quad (44)$$

with the constant C in the statement of Theorem 7. We have that:

$$\mathbb{E} [|P_*(Z_i, S_i^2) \wedge \zeta - P_{\widehat{G}}(Z_i, S_i^2) \wedge \zeta|] \leq \mathbb{E} \left[|P_*(\tilde{Z}_i, S_i^2) - P_{\widehat{G}}(\tilde{Z}_i, S_i^2)| \mathbf{1}(A) \right] + \mathbb{P}[A^c].$$

Above we used the fact that $P_*(\tilde{Z}_i, S_i^2), P_{\widehat{G}}(\tilde{Z}_i, S_i^2) \in [0, 1]$.

Next, for any distribution \tilde{G} supported on $(0, \infty)$, and any $z, z_i \in \mathbb{R}, s^2 > 0$, let us write:

$$\begin{aligned} N_{\tilde{G}}(z, s^2) &:= P_{\tilde{G}}(z, s^2) \cdot f_{\tilde{G}}(s^2), \\ N_i(z_i, \tilde{G}) &:= N_{\tilde{G}}(z_i, S_i^2), \\ D_i(\tilde{G}) &:= f_{\tilde{G}}(S_i^2), \text{ and} \\ P_{\tilde{G}}(z_i, S_i^2) &:= N_i(z_i, \tilde{G}) / D_i(\tilde{G}). \end{aligned}$$

Throughout the rest of the proof we also write G_* for G in the hierarchical setting or for

$G(\boldsymbol{\sigma})$ in the compound setting and we also let $\widehat{G}_* = (\widehat{G} + G_*)/2$. Then:

$$\begin{aligned}
& |P_*(z_i, S_i^2) - P_{\widehat{G}}(z_i, S_i^2)| \\
&= \left| \frac{N_i(z_i, G_*)}{D_i(G_*)} - \frac{N_i(z_i, \widehat{G})}{D_i(\widehat{G})} \right| \\
&= \left| \frac{N_i(z_i, G_*)}{D_i(G_*)} - \frac{N_i(z_i, G_*)}{D_i(\widehat{G}_*)} + \frac{N_i(z_i, G_*)}{D_i(\widehat{G}_*)} - \frac{N_i(z_i, \widehat{G})}{D_i(\widehat{G}_*)} + \frac{N_i(z_i, \widehat{G})}{D_i(\widehat{G}_*)} - \frac{N_i(z_i, \widehat{G})}{D_i(\widehat{G})} \right| \\
&\leq \frac{N_i(z_i, G_*)}{D_i(G_*)} \left| \frac{D_i(G_*) - D_i(\widehat{G}_*)}{D_i(\widehat{G}_*)} \right| + \frac{|N_i(z_i, G_*) - N_i(z_i, \widehat{G})|}{D_i(\widehat{G}_*)} + \frac{N_i(z_i, \widehat{G})}{D_i(\widehat{G})} \left| \frac{D_i(\widehat{G}_*) - D_i(\widehat{G})}{D_i(\widehat{G}_*)} \right| \\
&\leq \frac{|N_i(z_i, G_*) - N_i(z_i, \widehat{G})|}{D_i(\widehat{G}_*)} + \frac{|D_i(G_*) - D_i(\widehat{G})|}{D_i(\widehat{G}_*)}.
\end{aligned}$$

In the last step we used two facts: first, it holds that $N_i(z_i, \widetilde{G})/D_i(\widetilde{G}) \in [0, 1]$ for all \widetilde{G} (since they correspond to conditional p-values), and second, the map $\widetilde{G} \mapsto D_i(\widetilde{G})$ is linear, which implies that:

$$D_i(G_*) - D_i(\widehat{G}_*) = (D_i(G_*) - D_i(\widehat{G}))/2, \quad D_i(\widehat{G}_*) - D_i(\widehat{G}) = (D_i(G_*) - D_i(\widehat{G}))/2.$$

Combining all of the above results, we get:

$$\begin{aligned}
& \frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[\sup_{z: |z| \geq z} |P_*(z, S_i^2) - P_{\widehat{G}}(z, S_i^2)| \right] \\
&\leq \mathbb{P}[A^c] + \frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[\sup_{z: |z| \geq z} \left| \frac{N_i(z, G_*) - N_i(z, \widehat{G})}{D_i(\widehat{G}_*)} \right| \mathbf{1}(A) \right] + \frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[\left| \frac{D_i(G_*) - D_i(\widehat{G})}{D_i(\widehat{G}_*)} \right| \mathbf{1}(A) \right] \\
&= \mathbb{P}[A^c] + \text{I} + \text{II}.
\end{aligned}$$

By Theorem 7, $\mathbb{P}_G[A^c] \leq \exp(-c(\log n)^2)$. Furthermore, in two upcoming Lemmata 26 and 27 we bound terms I and II above and so conclude. \square

Lemma 26. It holds that:

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[\sup_{z: |z| \geq z} \left| \frac{N_i(z, G_*) - N_i(z, \widehat{G})}{D_i(\widehat{G}_*)} \right| \mathbf{1}(A) \right] \lesssim_{L, \bar{U}, \nu, \zeta} \frac{(\log n)^{5/2}}{\sqrt{n}}.$$

Proof. Below we will prove that there exists non-random $\zeta_n > 0$ such that for all i :

$$\sup_{z: |z| \geq z} |N_i(z, G_*) - N_i(z, \widehat{G})| \mathbf{1}(A) \leq \zeta_n.$$

Let $B_n > 0$ be another non-random sequence that we will pick later. Also recall that $D_i(\widehat{G}_*) = D_i(G_*)/2 + D_i(\widehat{G})/2 \geq D_i(G_*)/2 = f_*(S_i^2)/2$. We get:

$$\left| \frac{N_i(z, G_*) - N_i(z, \widehat{G})}{D_i(\widehat{G}_*)} \right| \mathbf{1}(A) \leq 4\mathbf{1}(S_i^2 > B_n) + 2\zeta_n \frac{1}{f_*(S_i^2)} \mathbf{1}(S_i^2 \leq B_n),$$

where we also used the fact that $N_i(z, \tilde{G})/D_i(\tilde{G}) \in [0, 1]$ for any distribution \tilde{G} .

Taking $B_n = 4\bar{U} \log n$ for $n \geq 3$, we get via Lemma 23 that $\mathbb{P}[S_i^2 > B_n] \leq 1/n$. Hence:

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[\sup_{z: |z| \geq z} \left| \frac{N_i(z, G_*) - N_i(z, \hat{G})}{D_i(\hat{G}_*)} \right| \mathbf{1}(A) \right] \leq \frac{4}{n} + 2\zeta_n \frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[\frac{\mathbf{1}(S_i^2 \leq B_n)}{f_*(S_i^2)} \right] \stackrel{(*)}{\leq} \frac{4}{n} + 2\zeta_n B_n.$$

(*) requires a slightly different argument in the hierarchical vs. compound setting. In the hierarchical setting it suffices to note that:

$$\mathbb{E}_G \left[\frac{\mathbf{1}(S_i^2 \leq B_n)}{f_G(S_i^2)} \right] = \int_0^{B_n} \frac{1}{f_G(t)} f_G(t) dt = B_n. \quad (45)$$

In the compound setting:

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^n \mathbb{E}_{\sigma_i} \left[\frac{\mathbf{1}(S_i^2 \leq B_n)}{f_{\sigma}(S_i^2)} \right] &= \frac{1}{n} \sum_{i=1}^n \int_0^{B_n} \frac{1}{f_{\sigma}(t)} p(t | \sigma_i^2) dt \\ &= \int_0^{B_n} \frac{1}{f_{\sigma}(t)} \left(\frac{1}{n} \sum_{i=1}^n p(t | \sigma_i^2) \right) dt \\ &= \int_0^{B_n} \frac{1}{f_{\sigma}(t)} f_{\sigma}(t) dt = B_n. \end{aligned}$$

It remains to determine ζ_n . Recall that we had defined (in the proof of Theorem 11) $N_i(z, \tilde{G}) = N_{\tilde{G}}(z, S_i^2)$ for any distribution \tilde{G} . For any $z \geq \underline{z}$ (we take z to be > 0 without loss of generality) and $s^2 > 0$, it holds that:

$$\begin{aligned} &\{N_{G_*}(z, s^2) - N_{\hat{G}}(z, s^2)\} / \{C(\nu) (s^2)^{\nu/2-1}\} \\ &= \int_0^{\infty} \mathbf{1} \left(t^2 \geq \frac{\nu s^2 + z^2}{\nu + 1} \right) \frac{(t^2)^{-(\nu-1)/2}}{\sqrt{(\nu+1)t^2 - \nu s^2}} (f_{G_*}(t^2; \nu+1) - f_{\hat{G}}(t^2; \nu+1)) d(t^2) \\ &\leq \left\{ \int_{\frac{\nu s^2 + z^2}{\nu+1}}^{\infty} \frac{(t^2)^{-(\nu-1)}}{(\nu+1)t^2 - \nu s^2} d(t^2) \right\}^{1/2} \left\{ \int_0^{\infty} (f_{G_*}(t^2; \nu+1) - f_{\hat{G}}(t^2; \nu+1))^2 d(t^2) \right\}^{1/2}. \end{aligned}$$

We bound these terms in turn. Since $\nu \geq 3$:

$$\int_{\frac{\nu s^2 + z^2}{\nu+1}}^{\infty} \frac{(t^2)^{-(\nu-1)}}{(\nu+1)t^2 - \nu s^2} d(t^2) \leq \frac{1}{(\nu-2)z^2} \left[-(t^2)^{-\nu+2} \right]_{\frac{\nu s^2 + z^2}{\nu+1}}^{\infty} = \frac{(\nu+1)^{\nu-2}}{(\nu-2)} \frac{1}{z^2 (\nu s^2 + z^2)^{\nu-2}}.$$

For the second term, it will be helpful to first note the following:

$$\begin{aligned} &\int_0^{\infty} (f_{G_*}(u) - f_{\hat{G}}(u))^2 d(u) \\ &= \int_0^{\infty} \left(\sqrt{f_{G_*}(u)} + \sqrt{f_{\hat{G}}(u)} \right)^2 \left(\sqrt{f_{G_*}(u)} - \sqrt{f_{\hat{G}}(u)} \right)^2 d(u) \\ &\lesssim_{L, \nu} \frac{1}{2} \int_0^{\infty} \left(\sqrt{f_{G_*}(u)} - \sqrt{f_{\hat{G}}(u)} \right)^2 d(u) \\ &= \mathfrak{H}^2(f_{G_*}, f_{\hat{G}}). \end{aligned}$$

For the step with the inequality $\lesssim_{L,\nu}$ we used Lemma 18A. Next we will use Lemma 10, which in combination with the above display yields:

$$\int_0^\infty (f_{G_*}(t^2; \nu + 1) - f_{\widehat{G}}(t^2; \nu + 1))^2 d(t^2) \lesssim_{L,\bar{U},\nu} |\log(\mathfrak{H}^2(f_{G_*}, f_{\widehat{G}}))| \mathfrak{H}^2(f_{G_*}, f_{\widehat{G}}).$$

On the event A (see (44)), the RHS above is $\lesssim_{L,\bar{U},\nu} (\log n)^3/n$. Combining the above results, we find that on the event A and for any z such that $|z| \geq z > 0$:

$$|N_{G_*}(z, s^2) - N_{\widehat{G}}(z, s^2)| \lesssim_{L,\bar{U},\nu} \frac{(s^2)^{\nu/2-1}}{z(\nu s^2 + z^2)^{\nu/2-1}} \cdot \frac{(\log n)^{3/2}}{\sqrt{n}} \lesssim_{z,L,\bar{U},\nu} \frac{(\log n)^{3/2}}{\sqrt{n}}.$$

Thus we may take:

$$\zeta_n \lesssim_{L,\bar{U},\nu} \frac{(\log n)^{3/2}}{\sqrt{n}}.$$

□

Lemma 27.

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[\left| \frac{D_i(G_*) - D_i(\widehat{G})}{D_i(\widehat{G}_*)} \right| \mathbf{1}(A) \right] \lesssim_{\nu,L,\bar{U}} \frac{\log n}{\sqrt{n}}.$$

Proof. Let $\eta = 1/n$. The starting point of the proof will be similar to the start of the proof of Theorem 7 and will consist of picking a suitable proper $(\eta, \|\cdot\|_\infty)$ -cover of the following class:

$$\mathcal{F}^-(L, \bar{U}, \nu) := \left\{ f_{\widehat{G}}(\cdot) = f_{\widehat{G}}(\cdot; \nu) : \tilde{G}([L, \bar{U}]) = 1, \mathfrak{H}(f_{\widehat{G}}, f_*) < \frac{C \log n}{\sqrt{n}} \right\}. \quad (46)$$

The constant $C > 0$ is as in Theorem 7. This definition is similar to the one in (36) except we now only consider densities close to f_* (rather than far).

Let us call the η -cover $\mathcal{S} = \{f_j : j \in \mathcal{J}\}$, $\mathcal{J} = \{1, \dots, J\}$, $J = \#\mathcal{S}$. Furthermore, since this a proper cover, for each j there exists a distribution G_j such that $f_j = f_{G_j}$. With an analogous argument as in the beginning of the proof of Theorem 7, we have that:

$$\log J \lesssim_{\nu,L,\bar{U}} |\log(\eta)|^2 = (\log n)^2.$$

We are ready to start with the main argument of the lemma.

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[\left| \frac{D_i(G_*) - D_i(\widehat{G})}{D_i(\widehat{G}_*)} \right| \mathbf{1}(A) \right] \\ &= \frac{2}{n} \sum_{i=1}^n \mathbb{E} \left[\left| \frac{D_i(G_*) - D_i(\widehat{G})}{D_i(G_*) + D_i(\widehat{G})} \right| \mathbf{1}(A) \right] \\ &\stackrel{(*)}{\leq} \frac{2}{n} + 8\eta\bar{U} \log n + \mathbb{E} \left[\sup_{j \in \mathcal{J}} \left\{ \frac{2}{n} \sum_{i=1}^n \left| \frac{D_i(G_*) - D_i(G_j)}{D_i(G_*) + D_i(G_j)} \right| \right\} \right] \\ &\leq \frac{2}{n} + 8\eta\bar{U} \log n + \mathbb{E} \left[\sup_{j \in \mathcal{J}} \left\{ \frac{2}{n} \sum_{i=1}^n \left(\left| \frac{D_i(G_*) - D_i(G_j)}{D_i(G_*) + D_i(G_j)} \right| - \mathbb{E} \left[\left| \frac{D_i(G_*) - D_i(G_j)}{D_i(G_*) + D_i(G_j)} \right| \right] \right) \right\} \right] \\ &\quad + \sup_{j \in \mathcal{J}} \left\{ \mathbb{E} \left[\frac{2}{n} \sum_{i=1}^n \left| \frac{D_i(G_*) - D_i(G_j)}{D_i(G_*) + D_i(G_j)} \right| \right] \right\} \\ &= \frac{2}{n} + 8\eta\bar{U} \log n + \text{I} + \text{II}. \end{aligned} \quad (47)$$

We first justify (*). On the event A (see (44)), there must exist a (random) index \hat{j} such that $\|f_{G_{\hat{j}}} - f_{\hat{G}}\|_{\infty} \leq \eta$. Let $B_n := 4\bar{U} \log n$, so that $\mathbb{P}[S_i^2 \geq B_n] \leq 1/n$ ($n \geq 3$, Lemma 23). Then:

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[\left| \frac{D_i(G_*) - D_i(\hat{G})}{D_i(G_*) + D_i(\hat{G})} \right| \mathbf{1}(A) \right] \\ & \leq \frac{1}{n} \sum_{i=1}^n \left\{ \mathbb{E} \left[\left| \frac{D_i(G_*) - D_i(\hat{G})}{D_i(G_*) + D_i(\hat{G})} \right| \mathbf{1}(A) \mathbf{1}(S_i^2 \leq B_n) \right] + \mathbb{P}[S_i^2 > B_n] \right\} \\ & \leq \frac{1}{n} + \frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[\left| \frac{D_i(G_*) - D_i(G_{\hat{j}})}{D_i(G_*) + D_i(G_{\hat{j}})} \right| \mathbf{1}(A) \mathbf{1}(S_i^2 \leq B_n) \right] \\ & \quad + \frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[\left| \frac{D_i(G_*) - D_i(\hat{G})}{D_i(G_*) + D_i(\hat{G})} - \frac{D_i(G_*) - D_i(G_{\hat{j}})}{D_i(G_*) + D_i(G_{\hat{j}})} \right| \mathbf{1}(A) \mathbf{1}(S_i^2 \leq B_n) \right]. \end{aligned}$$

We may bound the second summand in the last display by noting that $\hat{j} \in \mathcal{J}$ on the event A , and then taking the supremum over all $j \in \mathcal{J}$. On the other hand, we can show that the last summand is at most $2\eta B_n$:

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[\left| \frac{D_i(G_*) - D_i(\hat{G})}{D_i(G_*) + D_i(\hat{G})} - \frac{D_i(G_*) - D_i(G_{\hat{j}})}{D_i(G_*) + D_i(G_{\hat{j}})} \right| \mathbf{1}(A) \mathbf{1}(S_i^2 \leq B_n) \right] \\ & \leq \frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[\left| \frac{2D_i(G_*)(D_i(G_{\hat{j}}) - D_i(\hat{G}))}{(D_i(G_*) + D_i(G_{\hat{j}}))(D_i(G_*) + D_i(\hat{G}))} \right| \mathbf{1}(A) \mathbf{1}(S_i^2 \leq B_n) \right] \\ & \leq 2\eta \frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[\frac{\mathbf{1}(S_i^2 \leq B_n)}{D_i(G_*)} \right] \\ & = 2\eta B_n. \end{aligned}$$

The argument for the penultimate line is identical to the argument in equation (45) (as well as just below that equation for the compound setting).

We still need to bound I and II in (47). We start with II. Fix any $j \in \mathcal{J}$. Then,

$$\begin{aligned} \mathbb{E} \left[\frac{2}{n} \sum_{i=1}^n \left| \frac{D_i(G_*) - D_i(G_j)}{D_i(G_*) + D_i(G_j)} \right| \right] & \leq \frac{2}{n} \sum_{i=1}^n \mathbb{E} \left[\left(\frac{D_i(G_*) - D_i(G_j)}{D_i(G_*) + D_i(G_j)} \right)^2 \right]^{1/2} \\ & \leq 2 \left\{ \frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[\left(\frac{D_i(G_*) - D_i(G_j)}{D_i(G_*) + D_i(G_j)} \right)^2 \right] \right\}^{1/2}. \end{aligned}$$

For both inequality steps above we used Jensen's inequality. Continuing:

$$\begin{aligned}
& \frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[\left(\frac{D_i(G_*) - D_i(G_j)}{D_i(G_*) + D_i(G_j)} \right)^2 \right] \\
&= \frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[\left(\frac{f_*(S_i^2) - f_{G_j}(S_i^2)}{f_*(S_i^2) + f_{G_j}(S_i^2)} \right)^2 \right] \\
&= \int \left(\frac{f_*(u) - f_{G_j}(u)}{f_*(u) + f_{G_j}(u)} \right)^2 f_*(u) du \\
&= \int \left(\sqrt{f_*(u)} - \sqrt{f_{G_j}(u)} \right)^2 \frac{\left(\sqrt{f_*(u)} + \sqrt{f_{G_j}(u)} \right)^2}{f_*(u) + f_{G_j}(u)} \frac{f_*(u)}{f_*(u) + f_{G_j}(u)} du \\
&\leq 2 \int \left(\sqrt{f_*(u)} - \sqrt{f_{G_j}(u)} \right)^2 du \\
&= 4\mathfrak{J}^2(f_*, f_{G_j}) \leq 4C \frac{(\log n)^2}{n}.
\end{aligned}$$

At last, let us turn our attention to I. We introduce the notation:

$$U_{ij} = \left| \frac{D_i(G_*) - D_i(G_j)}{D_i(G_*) + D_i(G_j)} \right|, \text{ for } i \in \{1, \dots, n\}, j \in \mathcal{J},$$

and note that for any j , U_{ij} is a function of S_i^2 only. Furthermore, U_{ij} is bounded and takes values in the interval $[0, 1]$. Hence, by Hoeffding's inequality (for fixed j) and any $t \geq 0$:

$$\mathbb{P} \left[\left| \frac{1}{n} \sum_{i=1}^n (U_{ij} - \mathbb{E}[U_{ij}]) \right| \geq t \right] \leq \exp(-2nt^2).$$

Hence by applying the union bound over $j \in \mathcal{J}$:

$$\mathbb{P} \left[\sup_{j \in \mathcal{J}} \left\{ \left| \frac{1}{n} \sum_{i=1}^n (U_{ij} - \mathbb{E}[U_{ij}]) \right| \geq t \right\} \right] \leq J \exp(-2nt^2).$$

Hence:

$$\begin{aligned}
& \mathbb{E} \left[\sup_{j \in \mathcal{J}} \left\{ \left(\frac{1}{n} \sum_{i=1}^n (U_{ij} - \mathbb{E}[U_{ij}]) \right) \right\}^2 \right] \\
&\leq \mathbb{E} \left[\sup_{j \in \mathcal{J}} \left\{ \left(\frac{1}{n} \sum_{i=1}^n (U_{ij} - \mathbb{E}[U_{ij}]) \right)^2 \right\} \right] \\
&= \int_0^\infty \mathbb{P} \left[\sup_{j \in \mathcal{J}} \left| \frac{1}{n} \sum_{i=1}^n (U_{ij} - \mathbb{E}[U_{ij}]) \right| \geq \sqrt{t} \right] dt \\
&\leq \int_0^\infty \min\{1, J \exp(-2nt)\} dt \\
&= \frac{\log J}{2n} + J \left[-\frac{1}{2n} \exp(-2nt) \right]_{u^*}^\infty \\
&= \frac{1 + \log J}{2n} \lesssim_{\nu, L, \bar{U}} \frac{(\log n)^2}{n}.
\end{aligned}$$

Above u^* was chosen such that, $2nu^* = \log J$, i.e., $u^* = \log J/(2n)$.

Let us put everything together. In (47), we showed that:

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[\left| \frac{D_i(G_*) - D_i(\widehat{G})}{D_i(\widehat{G}_*)} \right| \mathbf{1}(A) \right] \leq \frac{2}{n} + 8\eta\bar{U} \log n + \text{I} + \text{II},$$

with $\eta = 1/n$. We also showed that:

$$\text{I} \lesssim_{\nu, L, \bar{U}} \frac{\log n}{\sqrt{n}},$$

and that,

$$\text{II} \leq 4\sqrt{C} \frac{\log n}{\sqrt{n}}.$$

Hence:

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[\left| \frac{D_i(G_*) - D_i(\widehat{G})}{D_i(\widehat{G}_*)} \right| \mathbf{1}(A) \right] \lesssim_{\nu, L, \bar{U}} \frac{\log n}{\sqrt{n}}.$$

□

E.4 Proof of Proposition 13

Proof. Let us fix $i \in \mathcal{H}_0$. Throughout this proof we write $P_i = P_{\widehat{G}}(Z_i, S_i^2)$ and $P_i^* = P_G(Z_i, S_i^2)$. Fix any $\zeta \in (0, 1)$ and $\delta \in (0, \zeta)$. Then by Lemma 29 below, it holds for $t \leq \zeta - \delta$ that:

$$\mathbf{1}(P_i \leq t) \leq \mathbf{1}(P_i^* \leq t + \delta) + \frac{1}{\delta} |P_i \wedge \zeta - P_i^* \wedge \zeta|.$$

Notice that $\mathbb{P}_G [P_i^* \leq t' \mid S_1^2, \dots, S_n^2] = t'$ for any $t' \in [0, 1]$. Hence taking the conditional expectation of the above inequality:

$$\mathbb{P}_G [P_i \leq t \mid S_1^2, \dots, S_n^2] \leq t + \delta + \frac{1}{\delta} \mathbb{E}_G [|P_i \wedge \zeta - P_i^* \wedge \zeta| \mid S_1^2, \dots, S_n^2].$$

Rearranging:

$$\mathbb{P}_G [P_i \leq t \mid S_1^2, \dots, S_n^2] - t \leq \delta + \frac{1}{\delta} \mathbb{E}_G [|P_i \wedge \zeta - P_i^* \wedge \zeta| \mid S_1^2, \dots, S_n^2].$$

Let us now take $t \in (\zeta - \delta, 1]$. Then $\mathbb{P}_G [P_i \leq t \mid S_1^2, \dots, S_n^2] \leq 1$ so that:

$$\mathbb{P}_G [P_i \leq t \mid S_1^2, \dots, S_n^2] - t \leq 1 - t \leq 1 - \zeta + \delta.$$

We next write $(a)_+ = a \vee 0$ for the positive part of a number $a \in \mathbb{R}$. Then the above inequalities have established that:

$$\sup_{t \in [0, 1]} (\mathbb{P}_G [P_i \leq t \mid S_1^2, \dots, S_n^2] - t)_+ \leq 1 - \zeta + \delta + \frac{1}{\delta} \mathbb{E}_G [|P_i \wedge \zeta - P_i^* \wedge \zeta| \mid S_1^2, \dots, S_n^2].$$

By iterated expectation:

$$\mathbb{E}_G \left[\sup_{t \in [0, 1]} (\mathbb{P}_G [P_i \leq t \mid S_1^2, \dots, S_n^2] - t)_+ \right] \leq 1 - \zeta + \delta + \frac{1}{\delta} \mathbb{E}_G [|P_i \wedge \zeta - P_i^* \wedge \zeta|].$$

The inequality above is valid for any $i \in \mathcal{H}_0$, so:

$$\max_{i \in \mathcal{H}_0} \mathbb{E}_G \left[\sup_{t \in [0,1]} (\mathbb{P}_G [P_i \leq t \mid S_1^2, \dots, S_n^2] - t)_+ \right] \leq 1 - \zeta + \delta + \frac{1}{\delta} \max_{i \in \mathcal{H}_0} \mathbb{E}_G [|P_i \wedge \zeta - P_i^* \wedge \zeta|].$$

By Theorem 11 and taking $n \rightarrow \infty$ we thus find that:

$$\limsup_{n \rightarrow \infty} \max_{i \in \mathcal{H}_0} \mathbb{E}_G \left[\sup_{t \in [0,1]} (\mathbb{P}_G [P_i \leq t \mid S_1^2, \dots, S_n^2] - t)_+ \right] \leq 1 - \zeta + \delta.$$

We now first take $\delta \rightarrow 0$ and then $\zeta \rightarrow 1$ to conclude that:

$$\limsup_{n \rightarrow \infty} \max_{i \in \mathcal{H}_0} \mathbb{E}_G \left[\sup_{t \in [0,1]} (\mathbb{P}_G [P_i \leq t \mid S_1^2, \dots, S_n^2] - t)_+ \right] = 0.$$

Now write $(a)_- = (-a) \vee 0$ for the negative part of a . Then we can analogously establish that:

$$\limsup_{n \rightarrow \infty} \max_{i \in \mathcal{H}_0} \mathbb{E}_G \left[\sup_{t \in [0,1]} (\mathbb{P}_G [P_i \leq t \mid S_1^2, \dots, S_n^2] - t)_- \right] = 0.$$

We conclude by noting the fact that $|a| \leq (a)_+ + (a)_-$ so that:

$$\limsup_{n \rightarrow \infty} \max_{i \in \mathcal{H}_0} \mathbb{E}_G \left[\sup_{t \in [0,1]} |\mathbb{P}_G [P_i \leq t \mid S_1^2, \dots, S_n^2] - t| \right] = 0.$$

Asymptotic uniformity follows from conditional uniformity shown above as follows:

$$\begin{aligned} & \max_{i \in \mathcal{H}_0} \sup_{t \in [0,1]} |\mathbb{P}_G [P_i \leq t] - t| \\ &= \max_{i \in \mathcal{H}_0} \sup_{t \in [0,1]} |\mathbb{E} [\mathbb{P}_G [P_i \leq t \mid S_1^2, \dots, S_n^2] - t]| \\ &\leq \max_{i \in \mathcal{H}_0} \mathbb{E} \left[\sup_{t \in [0,1]} |\mathbb{P}_G [P_i \leq t \mid S_1^2, \dots, S_n^2] - t| \right] \rightarrow 0 \text{ as } n \rightarrow \infty. \end{aligned}$$

□

F Proofs for Section 3.3

F.1 Further background on the Benjamini-Hochberg procedure

For the proofs it will be helpful to recall the equivalent characterization of the Benjamini-Hochberg procedure (14) through an empirical process viewpoint [Storey et al., 2004]. Let

$$R_n(t) := \sum_{i=1}^n \mathbf{1}(P_i \leq t). \quad (48)$$

Also let:

$$\hat{t} := \sup \left\{ t \in [0, \alpha] : \frac{nt}{R_n(t) \vee 1} \leq \alpha \right\}. \quad (49)$$

Then the following holds for the Benjamini-Hochberg procedure applied at level α :

$$H_i \text{ rejected if and only if } P_i \leq \hat{t}. \quad (50)$$

In particular, the total number of rejections of BH, R_n is equal to $R_n(\hat{t})$ (where we slightly abuse notation). In analogy to the empirical process of total discoveries (48), we may also define the empirical process of null discoveries:

$$V_n(t) := \sum_{i \in \mathcal{H}_0} \mathbf{1}(P_i \leq t),$$

so that the number of false discoveries of BH is equal to $V_n(\hat{t})$. Finally, with the above notation, the false discovery rate is equal to $\mathbb{E} [V_n(\hat{t}) / (R_n(\hat{t}) \vee 1)]$.

F.2 Proof of Theorems 14 and 14*

Proof. Let us write $\pi_* = 1$ in the compound setting and $\pi_* = n_0/n$ when $\sigma_i^2 \stackrel{\text{iid}}{\sim} G$. Using the empirical process characterization of the BH procedure from Section F.1, we may write the following:

$$\pi_*^{-1} \frac{V_n(\hat{t})}{R_n(\hat{t}) \vee 1} = \underbrace{\frac{n\hat{t}}{R_n(\hat{t}) \vee 1}}_{\leq \alpha} + \pi_*^{-1} \underbrace{\frac{V_n(\hat{t}) - n\pi_*\hat{t}}{R_n(\hat{t}) \vee 1}}_{\leq 0 + o_{\mathbb{P}}(1)}.$$

The expectation of the LHS is the false discovery rate multiplied by π_*^{-1} . Hence, once we justify the two steps in the underbraces, the conclusion will follow by the dominated convergence theorem since $V_n(\hat{t}) / (R_n(\hat{t}) \vee 1) \in [0, 1]$.

The argument for the first underbrace follows from the definition of \hat{t} in (49). For the second underbrace we need to argue a bit more. In particular we will also need to account for the fact that we applied the BH procedure to the ‘‘approximate’’ p-values P_i .

Let us define the event $A_n = \{\hat{t} \geq t_2^*\}$, where t_2^* was defined in the statement of the theorem. In Lemma 31 we will prove that $\mathbb{P}[A_n] \rightarrow 1$ as $n \rightarrow \infty$ and that $R_n(\hat{t})/n \geq t_2^*/\alpha'$ on the event A_n . The above implies that $(R_n(\hat{t}) \vee 1)/n$ asymptotically remains bounded away from 0.

In Lemma 30 we also show that:

$$\sup_{t \in [0, \alpha]} \left\{ \frac{V_n(t)}{n} - \pi_* t \right\} \leq 0 + o_{\mathbb{P}}(1).$$

Thus we have that

$$\frac{V_n(\hat{t})}{n} - \pi_* \hat{t} \leq 0 + o_{\mathbb{P}}(1).$$

Since A_n has asymptotic probability 1, we conclude that:

$$\frac{V_n(\hat{t}) - n\pi_*\hat{t}}{R_n(\hat{t}) \vee 1} = \left(\frac{V_n(\hat{t})}{n} - \pi_*\hat{t} \right) \bigg/ \frac{R_n(\hat{t}) \vee 1}{n} \leq 0 + o_{\mathbb{P}}(1).$$

□

First we state a Glivenko-Cantelli-type theorem. Since the oracle p-values change as n changes (since $\sigma = (\sigma_1, \dots, \sigma_n)$ is being updated) we require a result under triangular array asymptotics.

Lemma 28 (Glivenko-Cantelli). Let $U_{i,n} \in [0, 1]$, $i = 1, \dots, n$ be independent (but not necessarily identically distributed) random variables. Then under triangular array asymptotics:

$$\sup_{t \in [0,1]} \left| \frac{1}{n} \sum_{i=1}^n (\mathbf{1}(U_{i,n} \leq t) - \mathbb{P}[U_{i,n} \leq t]) \right| \rightarrow 0 \text{ almost surely as } n \rightarrow \infty.$$

Furthermore, if $\mathcal{H}_0 \subset \{1, \dots, n\}$ is a deterministic subset with $n_0 = \#\mathcal{H}_0 \rightarrow \infty$ as $n \rightarrow \infty$, then also:

$$\sup_{t \in [0,1]} \left| \frac{1}{n_0} \sum_{i \in \mathcal{H}_0} (\mathbf{1}(U_{i,n} \leq t) - \mathbb{P}[U_{i,n} \leq t]) \right| \rightarrow 0 \text{ almost surely as } n \rightarrow \infty.$$

Proof. The above follows from a standard bracketing argument. \square

For all results that follow, for compactness of notation, we write $P_i = P_{\hat{G}}(Z_i, S_i^2)$ (as we have done throughout this section), and also use the notation P_i^* for the oracle p-values, that is $P_i^* = P_G(Z_i, S_i^2)$ when $\sigma^2 \sim G$ (2), and $P_i^* = P_{\sigma}(Z_i, S_i^2)$ in the compound setting (recall Supplement A).

Lemma 29. Let $\zeta \in (0, 1)$, $\delta \in (0, \zeta)$ and $t \in [0, \zeta - \delta]$. Then:

$$\mathbf{1}(P_i \leq t) - \mathbf{1}(P_i^* \leq t + \delta) \leq \frac{1}{\delta} |P_i \wedge \zeta - P_i^* \wedge \zeta|.$$

Proof.

$$\begin{aligned} & \mathbf{1}(P_i \leq t) - \mathbf{1}(P_i^* \leq t + \delta) \\ &= \mathbf{1}(P_i \wedge \zeta \leq t) - \mathbf{1}(P_i^* \wedge \zeta \leq t + \delta) \leq \mathbf{1}(P_i \wedge \zeta \leq t, P_i^* \wedge \zeta > t + \delta) \leq \frac{1}{\delta} |P_i \wedge \zeta - P_i^* \wedge \zeta|. \end{aligned}$$

\square

Lemma 30. It holds that:

$$\sup_{t \in [0, \alpha]} \left\{ \frac{1}{n} \sum_{i \in \mathcal{H}_0} \mathbf{1}(P_i \leq t) - \pi_* t \right\} \leq 0 + o_{\mathbb{P}}(1).$$

Proof. Take any $\zeta \in (\alpha, 1)$. Then, for any $\delta \in (0, \zeta - \alpha)$ and $t \in [0, \alpha]$ by Lemma 29 we have the following:

$$\mathbf{1}(P_i \leq t) - \mathbf{1}(P_i^* \leq t + \delta) \leq \frac{1}{\delta} |P_i \wedge \zeta - P_i^* \wedge \zeta|.$$

From the above we get:

$$\begin{aligned} \frac{1}{n} \sum_{i \in \mathcal{H}_0} \mathbf{1}(P_i \leq t) &= \frac{1}{n} \sum_{i \in \mathcal{H}_0} \mathbf{1}(P_i^* \leq t + \delta) + \frac{1}{n} \sum_{i \in \mathcal{H}_0} [\mathbf{1}(P_i \leq t) - \mathbf{1}(P_i^* \leq t + \delta)] \\ &\leq \frac{1}{n} \sum_{i \in \mathcal{H}_0} \mathbf{1}(P_i^* \leq t + \delta) + \frac{1}{\delta} \frac{1}{n} \sum_{i \in \mathcal{H}_0} |P_i \wedge \zeta - P_i^* \wedge \zeta| \\ &\stackrel{(*)}{=} \frac{1}{n} \sum_{i \in \mathcal{H}_0} \mathbb{P}[P_i^* \leq t + \delta] + o_{\mathbb{P}}(1) + \frac{1}{\delta} o_{\mathbb{P}}(1) \\ &\stackrel{(**)}{\leq} \pi_*(t + \delta) + o_{\mathbb{P}}(1) + \frac{1}{\delta} o_{\mathbb{P}}(1). \end{aligned}$$

In (*), both $o_{\mathbb{P}}(1)$ terms are uniform in t . This follows by Lemma 28 for the first term and by Theorem 11 (along with Markov's inequality) for the second term. (**) follows by Proposition 3 when $\sigma_i^2 \stackrel{\text{iid}}{\sim} G$. In the compound case, (**) follows by the average significance control property of the oracle compound p-values which we proved in Theorem 15.

To conclude, take $\delta \searrow 0$ sufficiently slowly. Then:

$$\sup_{t \in [0, \alpha]} \left\{ \frac{1}{n} \sum_{i \in \mathcal{H}_0} \mathbb{1}(P_i \leq t) - \pi_* t \right\} \leq 0 + o_{\mathbb{P}}(1).$$

□

Lemma 31. Define the event $A_n := \{\hat{t} \geq t_2^*\}$, where t_2^* was defined in the statement of Theorem 14. Then: $\mathbb{P}[A_n] \rightarrow 1$ as $n \rightarrow \infty$ and furthermore on the event A_n it holds that $R_n(\hat{t})/n \geq t_2^*/\alpha$.

Proof. We start by arguing analogously to Lemma 30, but in the opposite direction. Take any $t \in [t_1^*, \alpha]$, $\zeta \in (\alpha, 1)$ and $\delta \in (0, t_1^* \wedge (t_2^* - t_1^*))$. Then:

$$\mathbb{1}(P_i \leq t) - \mathbb{1}(P_i^* \leq t - \delta) \geq -\mathbb{1}(P_i > t, P_i^* \leq t - \delta) \geq -\frac{1}{\delta} |P_i \wedge \zeta - P_i^* \wedge \zeta|.$$

Then:

$$\begin{aligned} \frac{R_n(t)}{n} &= \frac{1}{n} \sum_{i=1}^n \mathbb{1}(P_i \leq t) \\ &= \frac{1}{n} \sum_{i=1}^n \mathbb{1}(P_i^* \leq t - \delta) + \frac{1}{n} \sum_{i=1}^n [\mathbb{1}(P_i \leq t) - \mathbb{1}(P_i^* \leq t - \delta)] \\ &\geq \frac{1}{n} \sum_{i=1}^n \mathbb{1}(P_i^* \leq t - \delta) - \frac{1}{\delta} \frac{1}{n} \sum_{i=1}^n |P_i \wedge \zeta - P_i^* \wedge \zeta| \\ &= \frac{1}{n} \sum_{i=1}^n \mathbb{P}[P_i^* \leq t - \delta] - o_{\mathbb{P}}(1) - \frac{1}{\delta} o_{\mathbb{P}}(1). \end{aligned}$$

As in the proof of Lemma 30, the $o_{\mathbb{P}}(1)$ terms are uniform in t . Now take $t = t_2^*$ in the above display to get:

$$R_n(t_2^*)/n \geq (t_2^* - \delta)/\alpha' - o_{\mathbb{P}}(1) - \frac{1}{\delta} o_{\mathbb{P}}(1).$$

By taking $\delta \searrow 0$ sufficiently slowly, the above yields that:

$$R_n(t_2^*)/n \geq t_2^*/\alpha' - o_{\mathbb{P}}(1).$$

Now let A'_n be the event that $\{R_n(t_2^*)/n \geq t_2^*/\alpha'\}$. Then, since $\alpha > \alpha'$, it follows that $\mathbb{P}[A'_n] \rightarrow 1$ as $n \rightarrow \infty$. To conclude we note that by (49) it follows that on A'_n it holds that $\hat{t} \geq t_2^*$, i.e., $A_n \supset A'_n$ and $\mathbb{P}[A_n] \rightarrow 1$ as $n \rightarrow \infty$. Finally, since $R_n(\cdot)$ is non-decreasing in t , we have that $R_n(\hat{t})/n \geq R_n(t_2^*)/n \geq t_2^*/\alpha$ on the event A_n . □

G Proof of Proposition 16 in Section 5

Proof. Let $(\mu_i, \sigma_i^2, Z_i, S_i^2)$ be generated as in the statement of the proposition, that is, suppose $(\mu_i, \sigma_i) \sim \Pi$ (21) and (Z_i, S_i^2) are generated as in (1) conditional on μ_i, σ_i . It will be helpful to also generate (compare to the notation in (3)):

$$Z_i^{H_0} \mid \mu_i, \sigma_i^2, Z_i, S_i^2 \sim \mathcal{N}(0, \sigma_i^2).$$

Further note that the distribution of $(\sigma_i^2, Z_i^{H_0}, S_i^2)$ may be factorized as follows:

$$\sigma_i^2 \sim G(\cdot; \Pi), \quad (Z_i^{H_0}, S_i^2) \mid \sigma_i^2 \sim \mathcal{N}(0, \sigma_i^2) \otimes \frac{\sigma_i^2}{\nu} \chi_\nu^2,$$

where $G(\cdot; \Pi)$ has been defined in (22). The distribution of $(\sigma_i^2, Z_i^{H_0}, S_i^2)$ depends on Π only through $G(\cdot; \Pi)$.

By (23), it holds that $\sigma_i^2 \mid (\mu_i = 0) \sim G(\cdot; \Pi)$. Since the distribution of S_i^2 conditional on σ_i^2, μ_i only depends on σ_i^2 , it also holds that:

$$(\sigma_i^2, S_i^2) \mid (\mu_i = 0) \stackrel{\mathcal{D}}{=} (\sigma_i^2, S_i^2).$$

This also implies that:

$$(Z_i, \sigma_i^2, S_i^2) \mid (\mu_i = 0) \stackrel{\mathcal{D}}{=} (Z_i^{H_0}, \sigma_i^2, S_i^2).$$

Thus, S_i^2 -almost surely,

$$Z_i \mid S_i^2, \mu_i = 0 \stackrel{\mathcal{D}}{=} Z_i^{H_0} \mid S_i^2. \quad (51)$$

We are ready to conclude. For any $t \in [0, 1]$, it almost surely holds that:

$$\begin{aligned} \mathbb{P}_\Pi [P_{G(\cdot; \Pi)}(Z_i, S_i^2) \leq t \mid S_i^2, \mu_i = 0] &\stackrel{(*)}{=} \mathbb{P}_\Pi [P_{G(\cdot; \Pi)}(Z_i^{H_0}, S_i^2) \leq t \mid S_i^2] \\ &\stackrel{(**)}{=} \mathbb{P}_{G(\cdot; \Pi)} [P_{G(\cdot; \Pi)}(Z_i^{H_0}, S_i^2) \leq t \mid S_i^2] \stackrel{(***)}{=} t. \end{aligned}$$

In (*) we used (51), in (**) we recalled the observation that the distribution of $(Z_i^{H_0}, S_i^2)$ only depends on $G(\cdot; \Pi)$, and in (***) we used Proposition 3. \square