

Large p SYK from chord diagrams

Baur Mukhametzhanov

Department of Physics, Cornell University
Ithaca, NY 14853, USA

Abstract

The p -body SYK model at finite temperature exhibits submaximal chaos and contains stringy-like corrections to the dual JT gravity. It can be solved exactly in two different limits: “large p ” SYK $1 \ll p \ll N$ and “double-scaled” SYK $N, p \rightarrow \infty$ with $\lambda = 2p^2/N$ fixed. We clarify the relation between the two. Starting from the exact results in the double-scaled limit, we derive several observables in the large p limit. In particular, we find that the density of states takes a very simple form up to non-perturbative corrections in λ . We compute euclidean $2n$ -point correlators and out-of-time-order four-point function at long lorentzian times. To compute the correlators we find the relevant asymptotics of the $\mathcal{U}_q(\mathfrak{su}(1,1))$ 6j-symbol.

Contents

| | | |
|----------|--|-----------|
| 1 | Introduction | 1 |
| 2 | Double-scaled SYK review | 2 |
| 2.1 | Density of states | 2 |
| 2.2 | Correlation functions | 4 |
| 3 | Partition function | 5 |
| 3.1 | Density of states | 5 |
| 3.2 | Partition function | 6 |
| 4 | Two-point function | 7 |
| 5 | Four-point function | 9 |
| 5.1 | 6j-symbol | 9 |
| 5.2 | OTOC at long times | 11 |
| 5.3 | Euclidean four-point function | 13 |
| 5.4 | Euclidean $2n$ -point function | 14 |
| A | Asymptotics of the 6j-symbol | 15 |

1 Introduction

SYK model [1–3] has played a significant role in recent years as a solvable example of holographic duality [4–9]. At low temperatures it is approximately described by JT gravity in a nearly AdS_2 spacetime, e.g. see [10] for a review and a more extensive list of references. In this regime the SYK model exhibits maximal chaos related to the presence of the black hole horizon [11].

At finite temperatures there are two additional limits of parameters where the SYK model is exactly solvable. One is the “large p” SYK¹ [4] in the limit $1 \ll p \ll N$ for a p -body SYK hamiltonian. The other solvable limit is the “double-scaled” SYK (DSSYK) [12, 13], in the limit $N, p \rightarrow \infty$ while keeping the ratio $\lambda = 2p^2/N$ finite.

Our interest in these solvable limits is two-fold. One is that the SYK model at finite temperature exhibits submaximal chaos. Second, from the perspective of the holographic gravity dual, large p/DSSYK could be thought of as including stringy corrections to JT gravity. The two are in fact related [14]. Having a solvable model for both of these phenomena is valuable.

In the present paper we derive various observables in large p SYK as the limit $\lambda = 2p^2/N \rightarrow 0$ of the exact results in DSSYK. We consider the partition function, euclidean correlators and OTOC at long lorentzian times. This was partially studied in [12]. Our contribution is to clarify the relation and see that it agrees with existing results in the literature derived using different methods [4, 15, 16].

One of our findings is that the density of states in large p SYK takes a simple form²

$$\rho_q(s) \approx \frac{1}{(2\pi)^2} \frac{2 \sin(\lambda s)}{\lambda} 2 \sinh(2\pi s) e^{-2\lambda s^2} + O(e^{4\pi(s-\frac{\pi}{\lambda})}), \quad E(s) = -\frac{2}{\lambda} \cos(\lambda s). \quad (1.1)$$

The second equation defines $s \in [0, \pi/\lambda]$ that parametrizes the energy. The energies relevant for large p SYK at finite temperature turn out to be

$$\lambda \rightarrow 0, \quad \lambda s \text{ - finite.} \quad (1.2)$$

In this regime the main difference of (1.1) from the Schwarzian density of states is the factor $e^{-2\lambda s^2}$, which changes the saddle in the partition function and correlators. At $\lambda \rightarrow 0$ and finite s (1.1) reduces to the Schwarzian density of states.

Another finding is that the Lyapunov exponent is given by

$$\lambda_L = \frac{2\pi}{\beta} \left(1 - \frac{2}{\pi} \lambda s \right), \quad (1.3)$$

where s is the energy parameter determined by the saddle in the partition function, see section 3. The Lyapunov exponent in large p SYK was first derived in [4]. In (1.3) we relate it to the average energy in the system. In particular, the bound on chaos [17] is satisfied $\lambda_L \leq 2\pi/\beta$ since $s \geq 0$.

The paper is organized as follows. In section 2, we review the exact results in DSSYK [12] following the conventions of section 8 in [18]. In section 3, we derive the density of states and

¹In the literature this is often referred to as “large q” SYK, but in the present paper we reserve “q” for a different quantity.

²Up to an overall s -independent factor, see eq. (3.6).

the partition function in large p SYK. In section 4 we derive the euclidean two-point function. In section 5 we derive the lorentzian out-of-time-order (OTO) four-point function and euclidean $2n$ -point functions.

Note: After this work had been completed and was in preparation for publication, we learned that partially intersecting results were independently obtained in [19]. The current work was presented by the author at the “Abu Dhabi Meeting on Theoretical Physics” on January 10, 2023 [20].

2 Double-scaled SYK review

We start with a brief review of correlation functions in DSSYK [12]. Our conventions for the normalization of the density of states and correlators will follow the section 8 in [18], that are more convenient for matching with JT gravity in the low temperature limit. The SYK model has a hamiltonian

$$H = i^{p/2} \sum_{i_1 < \dots < i_p} J_{i_1 \dots i_p} \psi_{i_1} \dots \psi_{i_p}, \quad \langle J_{i_1 \dots i_p}^2 \rangle = \frac{\mathcal{J}^2}{\lambda \binom{N}{p}}, \quad (2.1)$$

where $\psi_i; i = 1, \dots, N$ are Majorana fermions $\{\psi_i, \psi_j\} = 2\delta_{ij}$. Our normalization for \mathcal{J} is different from [12] by a factor of $\lambda = \frac{2p^2}{N}$ and instead follows [4]. This will be more appropriate to compare with large p SYK results in the literature. Below we set $\mathcal{J} = 1$.

There are two interesting limits where exact correlators can be computed at finite temperatures

$$\text{“Double-scaled SYK”}: \quad N, p \rightarrow \infty, \quad \lambda = \frac{2p^2}{N} - \text{fixed}. \quad (2.2)$$

$$\text{“Large } p \text{ SYK”}: \quad 1 \ll p \ll N. \quad (2.3)$$

In the second case, the limit can be taken in various ways. For example, one can take $N \rightarrow \infty$ with fixed p first, and then take $p \rightarrow \infty$. This order was adopted in [4]. Alternatively, one can take the double-scaling limit (2.2) first, and then take $\lambda \rightarrow 0$. In this paper we will take the latter approach.

2.1 Density of states

The density of states is given by³

$$E(s) = -\frac{2}{\lambda} \cos(\lambda s), \quad s \in [0, \pi/\lambda], \quad (2.4)$$

$$\rho_q(s) = \frac{1}{2\pi \Gamma_q(\pm 2is)}. \quad (2.5)$$

The first equation is the definition of s that parametrizes the energy. The energy spectrum has a finite support $E(s) \in [-2/\lambda, 2/\lambda]$. We also introduced an important parameter

$$q = e^{-\lambda} \quad (2.6)$$

³More precisely, $E(s) = -\frac{2}{\sqrt{\lambda(1-e^{-\lambda})}} \cos(\lambda s)$. Our main interest will be the limit $\lambda \rightarrow 0$, where this makes no difference.

that will appear throughout the paper. The density of states is expressed in terms of the q -gamma function $\Gamma_q(x)$ formally defined as

$$\Gamma_q(x) = (1-q)^{1-x} \frac{(q; q)_\infty}{(q^x; q)_\infty}, \quad (a; q)_\infty = \prod_{n=0}^{\infty} (1-aq^n). \quad (2.7)$$

To understand this definition one first considers q -deformation of integers (“ q -numbers”)

$$[n]_q = 1 + q + \dots + q^{n-1} = \frac{1-q^n}{1-q} \xrightarrow{q \rightarrow 1} n. \quad (2.8)$$

Then for integer n the q -gamma function is defined as the q -factorial

$$\Gamma_q(n) = [n-1]_q! \quad (2.9)$$

$$= [1]_q [2]_q \dots [n-1]_q. \quad (2.10)$$

For integer x in (2.7) most factors in the two Pochhammer symbols cancel out and we are left with (2.9). Therefore (2.7) is the analytic continuation of (2.9) to complex x .

Clearly, in the limit $q \rightarrow 1$ we get the usual gamma-function $\lim_{q \rightarrow 1} \Gamma_q(x) = \Gamma(x)$. In particular, the density of states reduces to the Schwarzian density of states⁴

$$\rho_q(s) \approx \frac{1}{2\pi\Gamma(\pm 2is)} \quad (2.11)$$

$$= \frac{2s}{(2\pi)^2} 2 \sinh(2\pi s), \quad \lambda \rightarrow 0. \quad (2.12)$$

While the energy near the edge of the spectrum $E_0 = -2/\lambda$ is

$$E(s) - E_0 = \frac{2}{\lambda} (1 - \cos \lambda s) \quad (2.13)$$

$$\approx \lambda s^2, \quad \lambda \rightarrow 0. \quad (2.14)$$

In the second line we took the limit $\lambda \rightarrow 0$ with s fixed. The overall λ signals that we work in the low energy limit. Respectively, we work at low temperatures $\beta = \frac{\tilde{\beta}}{\lambda}$, such that $\beta E = \tilde{\beta} s^2$ stays finite.⁵

The partition function of DSSYK is of course

$$Z(\beta) = \int_0^{\pi/\lambda} ds \rho_q(s) e^{-\beta E(s)}. \quad (2.15)$$

And our density of states (2.5) is normalized by⁶

$$Z(\beta = 0) = \int_0^{\pi/\lambda} ds \rho_q(s) = \frac{1}{\lambda(1-q)^2(q; q)_\infty^3} \equiv N_q. \quad (2.16)$$

⁴Our conventions are related to [21] by $\rho(s) = 2\rho_{\text{there}}(s)$ that could be achieved by rescaling S_0 .

⁵The relation to JT gravity is $\beta\mathcal{J} = \frac{\tilde{\beta}J\mathcal{T}}{\epsilon}$, $\lambda = 4G_N\epsilon/\tilde{\phi}_r$, where we restored \mathcal{J} .

⁶Our density of states is related to [12] by $\frac{1}{N_q}\rho_q(s) ds = \frac{(q; q)_\infty}{2\pi} (e^{\pm 2i\theta}; q)_\infty d\theta$, where $\theta = \lambda s$.

2.2 Correlation functions

In [12] the authors also considered operators

$$\mathcal{O} = i^{p'/2} \sum_{i_1 < \dots < i_{p'}} J_{i_1 \dots i_{p'}}^{(\mathcal{O})} \psi_{i_1} \dots \psi_{i_{p'}} , \quad \left\langle \left(J_{i_1 \dots i_{p'}}^{(\mathcal{O})} \right)^2 \right\rangle = \binom{N}{p'}^{-1} . \quad (2.17)$$

These are similar to the hamiltonian (2.1), but contain a different number of fermions p' , as well as independent gaussian random couplings $J_{i_1 \dots i_{p'}}^{(\mathcal{O})}$. In DSSYK we work in the limit (2.2) with

$$p' = p\Delta , \quad (2.18)$$

where Δ is interpreted as the scaling dimension of the operator \mathcal{O} . **This makes sense at low temperatures where we have conformal symmetry and the hamiltonian H has dimension 1.**

One might ask why we are interested in random operators (2.17). A technical reason is that their correlation functions can be computed exactly in the double-scaling limit. Another reason is that in large p SYK the correlators of \mathcal{O} will have a very similar form to those of single fermion operators, as we will see in subsequent sections.

Correlation functions of \mathcal{O} take the form

$$\langle \text{Tr} e^{-\beta_1 H} \mathcal{O} e^{-\beta_2 H} \mathcal{O} \dots e^{-\beta_n H} \mathcal{O} \rangle = \frac{1}{Z} \int_0^{\pi/\lambda} \prod_{j=1}^n (ds_j \rho_q(s_j) e^{-\beta_j E_j(s_j)}) \langle \mathcal{O}_{E_1 E_2} \dots \mathcal{O}_{E_n E_1} \rangle . \quad (2.19)$$

The correlators in the energy basis are computed as a sum over all ‘‘chord diagrams’’

$$\langle \mathcal{O}_{E_1 E_2} \dots \mathcal{O}_{E_n E_1} \rangle = \begin{array}{c} \mathcal{O}_{E_1 E_2} \qquad \mathcal{O}_{E_2 E_3} \\ \circlearrowleft \\ \begin{array}{c} s_3 \\ s_2 \\ s_4 \\ \dots \\ s_1 \end{array} \end{array} + \dots \quad (2.20)$$

In each diagram operators are connected pairwise. The chords split the diagram into regions, to each of which we assign an energy parameter s . Then for each chord diagram we write a formula as follows:

- For each matrix element $\mathcal{O}_{E_1 E_2}$ we write a factor of⁷

$$\Gamma_{12}^{1/2} \equiv \left(\frac{\Gamma_q(\Delta \pm is_1 \pm is_2)}{\Gamma_q(2\Delta)} \right)^{1/2} , \quad (2.21)$$

- For each intersection of chords there is a factor of the 6j-symbol of the quantum group $\mathcal{U}_{q^{1/2}}(su(1,1))$

$$\begin{array}{c} \Delta \qquad \Delta \\ \diagdown \quad \diagup \\ s_2 \\ \diagup \quad \diagdown \\ s_1 \quad s_3 \\ \diagdown \quad \diagup \\ s_4 \end{array} = \left\{ \begin{array}{ccc} \Delta & s_1 & s_2 \\ \Delta & s_3 & s_4 \end{array} \right\}_q . \quad (2.22)$$

⁷We adopt a common convention that ‘‘ \pm ’’ denotes a product for all choices of signs: $\Gamma_q(\Delta \pm is_1 \pm is_2) = \Gamma_q(\Delta + is_1 + is_2)\Gamma_q(\Delta + is_1 - is_2)\Gamma_q(\Delta - is_1 + is_2)\Gamma_q(\Delta - is_1 - is_2)$

For now, this is just some special function with group-theoretic origin. We will discuss it in more detail in section 5.

Let's consider some examples. The two-point function has only one diagram computed by

$$\langle \mathcal{O}_{E_1 E_2} \mathcal{O}_{E_2 E_1} \rangle = \begin{array}{c} \text{---} s_1 \text{---} \\ | \\ \text{---} s_2 \text{---} \\ \text{---} \end{array} = \frac{\Gamma_q(\Delta \pm i s_1 \pm i s_2)}{\Gamma_q(2\Delta)}. \quad (2.23)$$

The four-point function has three diagrams

$$\begin{aligned} \langle \mathcal{O}_{E_1 E_2} \mathcal{O}_{E_2 E_3} \mathcal{O}_{E_3 E_4} \mathcal{O}_{E_4 E_1} \rangle &= \begin{array}{c} \text{---} s_2 \text{---} \\ | \\ \text{---} s_3 \text{---} \\ | \\ \text{---} s_4 \text{---} \\ | \\ \text{---} s_1 \text{---} \end{array} + \begin{array}{c} \text{---} s_2 \text{---} \\ | \\ \text{---} s_3 \text{---} \\ | \\ \text{---} s_4 \text{---} \\ | \\ \text{---} s_1 \text{---} \end{array} + \begin{array}{c} \text{---} s_2 \text{---} \\ | \\ \text{---} s_3 \text{---} \\ | \\ \text{---} s_4 \text{---} \\ | \\ \text{---} s_1 \text{---} \end{array} \quad (2.24) \\ &= (\Gamma_{12} \Gamma_{23} \Gamma_{34} \Gamma_{41})^{1/2} \left(\frac{\delta(s_1 - s_3)}{\rho_q(s_1)} + \frac{\delta(s_2 - s_4)}{\rho_q(s_2)} + \left\{ \begin{array}{c} \Delta \quad s_1 \quad s_2 \\ \Delta \quad s_3 \quad s_4 \end{array} \right\}_q \right). \quad (2.25) \end{aligned}$$

Each region must have only one integration parameter s . In (2.19) we introduced integrals over s_1, \dots, s_n for the n -point function. Therefore, in the first two diagrams in (2.24) we compensate redundant integrations by writing appropriate delta-functions.

The six-point function has $\binom{6}{2} = 15$ chord diagrams. Out of those, 11 are disconnected in the energy basis, i.e. products of two- and four-point functions. There are 4 connected diagrams

$$\langle \mathcal{O}_{E_1 E_2} \mathcal{O}_{E_2 E_3} \mathcal{O}_{E_3 E_4} \mathcal{O}_{E_4 E_5} \mathcal{O}_{E_5 E_6} \mathcal{O}_{E_6 E_1} \rangle_{conn} = \begin{array}{c} \text{---} \text{---} \\ | \quad | \\ \text{---} \text{---} \\ | \quad | \\ \text{---} \text{---} \end{array} + \begin{array}{c} \text{---} \text{---} \\ | \quad | \\ \text{---} \text{---} \\ | \quad | \\ \text{---} \text{---} \end{array} + \begin{array}{c} \text{---} \text{---} \\ | \quad | \\ \text{---} \text{---} \\ | \quad | \\ \text{---} \text{---} \end{array} + \begin{array}{c} \text{---} \text{---} \\ | \quad | \\ \text{---} \text{---} \\ | \quad | \\ \text{---} \text{---} \end{array}. \quad (2.26)$$

3 Partition function

We now turn to the computation of the partition function and correlators in large p SYK $1 \ll p \ll N$. As was discussed in the previous section, this can be achieved by taking the limit $\lambda \rightarrow 0$ of DSSYK results. We are interested in keeping the temperature finite. The relevant limit for the energies will turn out to be

$$\lambda \rightarrow 0, \quad \lambda s \text{ - fixed}. \quad (3.1)$$

3.1 Density of states

We start with computing the relevant limit of the density of states (2.5). Note that we can express it as a Jacobi function⁸

$$\rho_q(s) = N_q \frac{2\lambda \sin(\lambda s) e^{\lambda/8}}{2\pi} \vartheta_{11} \left(\frac{\lambda s}{\pi}, \frac{i\lambda}{2\pi} \right), \quad (3.2)$$

⁸We use conventions of [22].

where⁹

$$\vartheta_{11}(\nu, \tau) = -\frac{e^{\pi i \tau/4}}{2 \sin(\pi \nu)} (q; q)_{\infty} (z^{\pm 1}; q)_{\infty}, \quad q = e^{2\pi i \tau}, \quad z = e^{2\pi i \nu}. \quad (3.3)$$

Using a modular transformation of the Jacobi function

$$\vartheta_{11}(\nu, \tau) = \frac{i}{\sqrt{-i\tau}} e^{-\pi i \nu^2/\tau} \vartheta_{11}(\nu/\tau, -1/\tau) \quad (3.4)$$

we find

$$\rho_q(s) = \frac{1}{(2\pi)^2} \frac{\lambda^2}{(1-q)^2} \frac{2 \sin(\lambda s)}{\lambda} 2 \sinh(2\pi s) e^{-2\lambda s^2} \frac{\prod_{n=1}^{\infty} (1 - e^{\pm 4\pi s} q'^n)}{(q'; q')_{\infty}^2}, \quad (3.5)$$

where $q' = e^{-4\pi^2/\lambda}$. Up to non-perturbative corrections in λ and if we stay away from the right edge $s = \pi/\lambda$, the density of states takes a very simple form

$$\rho_q(s) = \frac{1}{(2\pi)^2} \frac{\lambda^2}{(1-q)^2} \frac{2 \sin(\lambda s)}{\lambda} 2 \sinh(2\pi s) e^{-2\lambda s^2} + O(e^{4\pi(s-\frac{\pi}{\lambda})}). \quad (3.6)$$

If we were interested in the low temperature limit, we would take $\lambda \rightarrow 0$ with s fixed. Then the above formula reduces to the Schwarzian density (2.12): $\frac{\sin(\lambda s)}{\lambda} e^{-2\lambda s^2} \approx s$. At finite temperatures we instead keep λs fixed and find

$$\rho_q(s) \approx \frac{1}{(2\pi)^2} \frac{2 \sin(\lambda s)}{\lambda} 2 \sinh(2\pi s) e^{-2\lambda s^2}. \quad (3.7)$$

It is important to retain the factors $\sin(\lambda s) e^{-2\lambda s^2}$ that will play a crucial role in computations at finite temperature and lead to a new saddle point for the average energy. Similar approximations were discussed in spin glass models [23] where the exact density of states turns out to have the same form (2.5).

3.2 Partition function

Now we compute the partition function in the limit $\lambda \rightarrow 0$ with β fixed

$$Z(\beta) = \int_0^{\pi/\lambda} ds \rho_q(s) e^{-\beta E(s)} \quad (3.8)$$

$$\approx \int_0^{\infty} ds \frac{1}{(2\pi)^2} \frac{2 \sin(\lambda s)}{\lambda} e^{2\pi s - 2\lambda s^2 + \beta \frac{2}{\lambda} \cos \lambda s}. \quad (3.9)$$

Note that if $s \sim \frac{1}{\lambda}$ then all three terms in the exponential are large and of order $\frac{1}{\lambda}$. We therefore compute in the saddle approximation

$$\partial_s \left(2\pi s - 2\lambda s^2 + \beta \frac{2}{\lambda} \cos \lambda s \right) = 0 \quad \Rightarrow \quad 2 \sin(\lambda s) = \frac{2\pi}{\beta} \left(1 - \frac{2\lambda s}{\pi} \right). \quad (3.10)$$

⁹ $(z^{\pm 1}; q)_{\infty} \equiv (z; q)_{\infty} (z^{-1}; q)_{\infty}$

Equivalently, the saddle-point equation is

$$\frac{v}{2 \cos \frac{\pi}{2} v} = \frac{\beta}{2\pi}, \quad v \equiv 1 - \frac{2}{\pi} \lambda s. \quad (3.11)$$

We recognize the equation derived by Maldacena and Stanford [4], where we also related their function $v(\beta)$ to the average energy.

The leading asymptotics of $v(\beta)$ are easy to understand from the second relation in (3.11) $v = 1 - \frac{2}{\pi} \lambda s$. Recall that $E(s) = -\frac{2}{\lambda} \cos(\lambda s)$, $s \in [0, \pi/\lambda]$. At low temperatures the energies are near the ground state $E_0 = -2/\lambda$ and $s \approx 0$. This corresponds to $v \approx 1$. On the other hand, at high temperatures we expect to be at the maximum entropy point. This corresponds to the middle of the spectrum $s = \frac{\pi}{2\lambda}$ and therefore $v \approx 0$. More precisely we find from solving the saddle equation

$$v = \begin{cases} 1 - \frac{2}{\beta} + \frac{4}{\beta^2} \dots, & \beta \gg 1, \\ \frac{\beta}{\pi} + \dots, & \beta \ll 1. \end{cases} \quad (3.12)$$

Also including gaussian fluctuations, we compute the partition function

$$\frac{1}{(2\pi)^{3/2} N_q} Z(\beta) \approx \frac{1}{2\sqrt{\pi}\beta^{3/2}} \frac{v}{\sqrt{\sin \frac{\pi v}{2} + \frac{2}{\beta}}} \exp \left\{ \frac{2\pi v}{\lambda} \left(\tan \frac{\pi v}{2} - \frac{\pi v}{4} \right) \right\}. \quad (3.13)$$

The exponential precisely matches the free energy derived in [4]. Here, we also computed the 1-loop correction that interpolates between low and high temperatures. At low temperatures $\beta \gg 1$

$$\frac{1}{(2\pi)^{3/2} N_q} Z(\beta) \approx \left(1 - \frac{3}{\beta} + \dots \right) \frac{1}{2\sqrt{\pi}\beta^{3/2}} \exp \left\{ \frac{1}{\lambda} \left[2\beta - \frac{\pi^2}{2} + \frac{\pi^2}{\beta} - \frac{2\pi^2}{\beta^2} + \dots \right] \right\}. \quad (3.14)$$

In the exponent, the linear in β term corresponds to the ground state energy $-\beta E_0$. The second term is a correction to ground state entropy. The third term is the Schwarzian action on-shell. The fourth term is the first correction beyond the Schwarzian. At high temperatures $\beta \ll 1$

$$\frac{1}{N_q} Z(\beta) \approx (\beta^{3/2} + \dots) \frac{1}{\beta^{3/2}} \exp \left\{ \frac{1}{\lambda} \left(\frac{\beta^2}{2} + \dots \right) \right\}. \quad (3.15)$$

In particular, the temperature dependence cancels out in the 1-loop prefactor. We recover the correct normalization at infinite temperature $\frac{1}{N_q} Z \approx 1$ as $\beta \rightarrow 0$.

The saddle equation (3.10) and the partition function (3.13) were independently obtained in [19].

4 Two-point function

Now we compute the 2pt function in the limit $\lambda \rightarrow 0$ with β fixed. The exact result is

$$\langle \text{Tr} e^{-\beta H} \mathcal{O}(\tau) \mathcal{O}(0) \rangle = \frac{1}{Z} \int_0^{\pi/\lambda} ds_1 ds_2 \rho_q(s_1) \rho_q(s_2) e^{-\beta_1 E_1 - \beta_2 E_2} \frac{\Gamma_q(\Delta \pm i s_1 \pm i s_2)}{\Gamma_q(2\Delta)}, \quad (4.1)$$

where $\beta_1 = \beta - \tau, \beta_2 = \tau$. To compute this and also other correlation functions later on, it is convenient to change variables to the average energy and energy differences. The saddle in the average energy will turn out to be the same as in the computation of the partition function (3.11). While the integrals over energy differences will turn out to be essentially the same as in JT gravity, except for the fact that the relation between average energy and temperature is different and determined by (3.11). For the case of the two-point function we introduce new variables

$$s = \frac{1}{2}(s_1 + s_2), \quad \omega = s_1 - s_2, \quad (4.2)$$

$$E = \frac{1}{2}(E_1 + E_2), \quad \Omega = E_1 - E_2. \quad (4.3)$$

The relevant limit is $\lambda \rightarrow 0$, while keeping λs fixed. The differences of momenta are finite $\omega \sim 1$. In this limit the energies and densities of states are

$$E_j = -\frac{2}{\lambda} \cos \lambda s_j, \quad (j = 1, 2), \quad (4.4)$$

$$E \approx -\frac{2}{\lambda} \cos \lambda s, \quad \Omega \approx \partial_s E \omega = 2 \sin(\lambda s) \omega, \quad (4.5)$$

$$\rho_q(s_1) \rho_q(s_2) \approx \rho_q(s)^2. \quad (4.6)$$

We also have

$$\beta_1 E_1 + \beta_2 E_2 = \beta E + \left(\frac{\beta}{2} - \tau \right) \Omega \quad (4.7)$$

$$\approx \beta E + \left(\frac{\beta}{2} - \tau \right) \partial_s E \omega. \quad (4.8)$$

Further, we can approximate the q-gamma functions as follows

$$\Gamma_q(\Delta \pm i s_1 \pm i s_2) = \Gamma_q(\Delta \pm 2i s) \Gamma_q(\Delta \pm i \omega) \quad (4.9)$$

$$= \frac{1}{2\pi \rho_q(s)} \frac{\Gamma_q(\Delta \pm 2i s)}{\Gamma_q(\pm 2i s)} \Gamma_q(\Delta \pm i \omega) \quad (4.10)$$

$$\approx \frac{1}{2\pi \rho_q(s)} \left(\frac{2 \sin(\lambda s)}{\lambda} \right)^{2\Delta} \Gamma(\Delta \pm i \omega). \quad (4.11)$$

In the second line we multiplied and divided by $\Gamma_q(\pm 2i s) = \frac{1}{2\pi \rho_q(s)}$. To compute the ratio $\frac{\Gamma_q(\Delta \pm 2i s)}{\Gamma_q(\pm 2i s)}$ it is convenient to use the definition as the infinite product (2.7). We can imagine first that Δ is an integer. Then most factors in the infinite products cancel out between $\Gamma_q(\Delta \pm 2i s)$ and $\Gamma_q(\pm 2i s)$. The remainder can be shown to be $\left(\frac{2 \sin(\lambda s)}{\lambda} \right)^{2\Delta}$ in the limit $\lambda \rightarrow 0, s \sim \frac{1}{\lambda}$. This can of course be continued to non-integer Δ . Equivalently, we use (A.5). This is an analog of the formula $\frac{\Gamma(\Delta \pm 2i s)}{\Gamma(\pm 2i s)} \approx (2s)^{2\Delta}, s \rightarrow \infty$ for the ordinary gamma function.

Putting it all together, we find

$$\frac{1}{Z} \int_0^\infty ds \rho_q(s) e^{-\beta E(s)} \left(\frac{2 \sin(\lambda s)}{\lambda} \right)^{2\Delta} \int_{-\infty}^\infty \frac{d\omega}{2\pi} e^{\omega \partial_s E(\tau - \frac{\beta}{2})} \frac{\Gamma(\Delta \pm i \omega)}{\Gamma(2\Delta)}. \quad (4.12)$$

Here, the role of the integral over s is to determine the relation between the average energy and temperature. This is in fact the same saddle as in the computation of the partition function (3.11). In the rest of the integral we just set the average energy to take the saddle value. In particular, we have a factor $e^{\omega \partial_s E(\tau - \frac{\beta}{2})}$. The saddle equation (3.11) determines $\partial_s E = \frac{2\pi v}{\beta}$. Except for this factor $\partial_s E$, the integral is identical to the one arising in the semi-classical computation of JT gravity two-point function, e.g. see section 3 in [24]. We have

$$\left(\frac{2 \cos \frac{\pi v}{2}}{\lambda}\right)^{2\Delta} \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} e^{\omega \frac{2\pi v}{\beta}(\tau - \frac{\beta}{2})} \frac{\Gamma(\Delta \pm i\omega)}{\Gamma(2\Delta)} = \lambda^{-2\Delta} \left(\frac{\cos \frac{\pi v}{2}}{\cos \frac{\pi v}{\beta}(\frac{\beta}{2} - \tau)}\right)^{2\Delta}. \quad (4.13)$$

The factor $\lambda^{-2\Delta}$ can be absorbed by rescaling the operators \mathcal{O} . The rest agrees with the two-point function in [4].

5 Four-point function

Now we turn to the computation of the four-point function in large p SYK. In euclidean signature, we will show that it factorizes into Wick contractions with the two-point function (4.13). In fact, we will show that in euclidean signature all higher correlators factorize into products of two-point functions (4.13). This is of course just the large N factorization and is not surprising. Our contribution is to see how this arises from chord diagrams.

More interestingly, we will compute the out-of-time-order four-point correlator (OTOC) in large p SYK. We will consider the long lorentzian time limit $u \rightarrow \infty$, $\lambda \rightarrow 0$ while keeping $\lambda e^{\lambda_L u}$ fixed. The Lyapunov exponent λ_L will be given below.

To carry out these computations, the main object of interest is the 6j-symbol of the quantum group $\mathcal{U}_{q^{1/2}}(su(1,1))$. Many useful properties of this 6j-symbol and the related Askey-Wilson function are collected in appendix A of [18]. Other useful references include [25, 26]. We start with writing down some explicit formulas for this object.

5.1 6j-symbol

In [12] it was computed that the crossed digram in (2.24) is given in terms of a certain “basic (q -deformed) hypergeometric series” ${}_8\phi_7$, that we will define shortly. In [25] it was shown that this hypergeometric series is in fact a 6j-symbol of the quantum group $\mathcal{U}_{q^{1/2}}(su(1,1))$. It obeys numerous identities coming from group-theoretic considerations, such as orthogonality and Yang-Baxter relations. For our purposes it is more convenient to write an equivalent representation as a sum of two ${}_4\phi_3$ series¹⁰

$$\left\{ \begin{matrix} \Delta & s_1 & s_2 \\ \Delta & s_3 & s_4 \end{matrix} \right\}_q = \frac{\lambda}{1-q} (\gamma_{12}\gamma_{23}\gamma_{34}\gamma_{41})^{1/2} W_{q^{is_4}}(q^{is_2}; q^{\Delta \pm is_1}, q^{\Delta \pm is_3} | q), \quad \gamma_{12} \equiv \Gamma_q(\Delta \pm is_1 \pm is_2). \quad (5.1)$$

¹⁰See formula 8.15 in [25]. Our definition of Askey-Wilson function differs from [25] by $d \rightarrow 1 - d$ and a factor that depends only of q . The former is to make our function symmetric in a, b, c, d . The latter is to make the limit $q \rightarrow 1$ transparent.

where the Askey-Wilson function is defined by

$$W_{q^y}(q^x; q^a, q^b, q^c, q^d|q) = \frac{\Gamma_q(d-a)}{\Gamma_q(a+b)\Gamma_q(a+c)\Gamma_q(d\pm x)\Gamma_q(\tilde{d}\pm y)} \quad (5.2)$$

$${}_4\phi_3 \left(\begin{matrix} q^{a\pm x}, q^{\tilde{a}\pm y} \\ q^{a+b}, q^{a+c}, q^{a-d+1} \end{matrix}; q, q \right) + (a \leftrightarrow d), \quad (5.3)$$

where the “dual” parameters are $\tilde{a} = \frac{a+b+c-d}{2}$, $\tilde{d} = \frac{-a+b+c+d}{2}$. In (5.1), (5.3) “ \pm ” in the arguments of W and ${}_4\phi_3$ means that they depend on both parameters and the order is not important because these functions are symmetric. The basic (q-deformed) hypergeometric series is defined as

$${}_4\phi_3 \left(\begin{matrix} A_1, A_2, A_3, A_4 \\ B_1, B_2, B_3 \end{matrix}; q, x \right) = \sum_{n=0}^{\infty} \frac{(A_1, A_2, A_3, A_4; q)_n}{(B_1, B_2, B_3; q)_n} \frac{x^n}{(q; q)_n}, \quad (5.4)$$

where $(A_1, \dots, A_r; q)_n = \prod_{j=1}^r (A_j; q)_n$.

The Askey-Wilson function (5.3) is symmetric in a, b, c, d . This implies that the 6j-symbol (5.1) is symmetric in s_1, s_3 . For the particular choice of parameters in (5.1) the Askey-Wilson function, and therefore the 6j-symbol, has two additional symmetries. One is the symmetry in s_2, s_4 and follows from the so called “duality” property of the Askey-Wilson function [25]. The other is the symmetry under the permutation of columns $s_1 \leftrightarrow s_2$, $s_3 \leftrightarrow s_4$ and follows from the representation as ${}_8\phi_7$, see eq. 8.14 in [25].

In the limit $q \rightarrow 1$ the basic hypergeometric series reduces to the hypergeometric series

$$\lim_{q \rightarrow 1} {}_4\phi_3 \left(\begin{matrix} q^{a_1}, q^{a_2}, q^{a_3}, q^{a_4} \\ q^{b_1}, q^{b_2}, q^{b_3} \end{matrix}; q, x \right) = {}_4F_3 \left(\begin{matrix} a_1, a_2, a_3, a_4 \\ b_1, b_2, b_3 \end{matrix}; x \right). \quad (5.5)$$

Moreover, in the limit $q \rightarrow 1$ the q-gamma functions become gamma functions. Then the Askey-Wilson function (5.3) becomes the Wilson function [25, 27], while the quantum 6j-symbol (5.1) becomes the classical 6j-symbol of $sl(2, \mathbb{R})$. In this way we recover the four-point function in JT gravity [28].

Now we compute the asymptotics of the 6j-symbol. It depends on four energy parameters s_j . To compute the corresponding energy integrals in the four-point function (2.19), we will be interested in the limit when the average $s = \frac{1}{4} \sum_{j=1}^4 s_j$ is large, but the differences $s_i - s_j$ are finite. More precisely, we take

$$\lambda \rightarrow 0, \quad \lambda s \text{ - fixed}, \quad (s_i - s_j) \text{ - fixed}. \quad (5.6)$$

In this limit we find (see appendix A for details)

$$\left\{ \begin{matrix} \Delta & s_1 & s_2 \\ \Delta & s_3 & s_4 \end{matrix} \right\}_q \approx \frac{1}{2\pi\rho_q(s)} \left(\frac{2 \sin \lambda s}{\lambda} \right)^{i(\nu_2 + \nu_4)} \Gamma(i(\nu_2 + \nu_4)) \prod_{n=1}^4 \frac{\Gamma(\Delta - (-1)^n i\nu_n)^{1/2}}{\Gamma(\Delta + (-1)^n i\nu_n)^{1/2}} + \text{c.c.} \quad (5.7)$$

where we defined

$$s = \frac{1}{4} \sum_{j=1}^4 s_j, \quad \nu_1 = s_1 - s_2, \quad \nu_2 = s_2 - s_3, \quad \nu_3 = s_3 - s_4, \quad \nu_4 = s_4 - s_1. \quad (5.8)$$

A few comments about the approximation (5.7) are in order. Importantly, (5.7) has large oscillations $\lambda^{-i(\nu_2+\nu_4)}$. At long lorentzian times these oscillations will cancel with the oscillations from the long time evolution e^{-itE_j} , giving a finite result.

At finite euclidean times there will be no other oscillating factors in the four-point function that can cancel $\lambda^{-i(\nu_2+\nu_4)}$. Therefore this oscillating factor effectively sets $\nu_2 + \nu_4 = \nu_1 + \nu_3 = 0$. In this regime gamma functions in numerator and denominator cancel out. We also have a pole from $\Gamma(i(\nu_2 + \nu_4)) \approx \frac{1}{i(\nu_2+\nu_4)}$ and find in the limit $\lambda \rightarrow 0$

$$\left\{ \begin{array}{ccc} \Delta & s_1 & s_2 \\ \Delta & s_3 & s_4 \end{array} \right\}_q \approx \frac{1}{2\pi\rho_q(s)} \left(\frac{1}{i(\nu_2 + \nu_4)} \left(\frac{2 \sin \lambda s}{\lambda} \right)^{i(\nu_2+\nu_4)} + \text{c.c.} \right) \quad (5.9)$$

$$\approx \frac{\delta(s_1 + s_3 - s_2 - s_4)}{\rho_q(s)}. \quad (5.10)$$

In the second line we obtained a delta-function from $\lim_{\Lambda \rightarrow \infty} \frac{2 \sin(\Lambda x)}{x} = 2\pi\delta(x)$. In our case $\Lambda = \frac{2 \sin(\lambda s)}{\lambda} \gg 1$.

Now we use the approximations (5.7) and (5.10) to compute the lorentzian OTOC and euclidean four-point functions respectively.

5.2 OTOC at long times

Now we compute the OTOC four-point function at long times. This comes from the crossed diagram in (2.24) where we use the approximation (5.7) for the 6j-symbol. We have the lorentzian correlator

$$\langle \text{Tr} e^{-\beta H} \mathcal{O}(u_1) \dots \mathcal{O}(u_4) \rangle = \frac{1}{Z} \int_0^{\pi/\lambda} \prod_{j=1}^4 (ds_j \rho_q(s_j) e^{-\beta_j E_j}) (\Gamma_{12} \Gamma_{23} \Gamma_{34} \Gamma_{41})^{1/2} \left\{ \begin{array}{ccc} \Delta & s_1 & s_2 \\ \Delta & s_3 & s_4 \end{array} \right\}_q \quad (5.11)$$

where we analytically continue to lorentzian times u_j ($u_{ij} = u_i - u_j$)

$$\beta_1 = \beta - iu_{14}, \quad \beta_2 = iu_{12}, \quad \beta_3 = iu_{23}, \quad \beta_4 = iu_{34}. \quad (5.12)$$

We are interested in the OTOC regime when $u_2, u_4 \gg u_1, u_3$. It is convenient to change variables s_1, \dots, s_4 to the average energy and energy differences s, ν_1, \dots, ν_4 (5.8)

$$ds_1 ds_2 ds_3 ds_4 = ds d\nu_1 d\nu_2 d\nu_3 \quad (5.13)$$

$$= ds d\nu_1 d\nu_2 d\nu_3 d\nu_4 \delta(\nu_1 + \nu_2 + \nu_3 + \nu_4). \quad (5.14)$$

In the second line we introduced an extra variable to make permutation symmetries manifest at the cost of a delta-function. The inverse transformation can be written as

$$s_1 = s + \frac{2\nu_1 + \nu_2 - \nu_4}{4}, \quad s_2 = s + \frac{2\nu_2 + \nu_3 - \nu_1}{4}, \quad (5.15)$$

$$s_3 = s + \frac{2\nu_3 + \nu_4 - \nu_2}{4}, \quad s_4 = s + \frac{2\nu_4 + \nu_1 - \nu_3}{4}. \quad (5.16)$$

The q-gamma functions in the limit $s \sim 1/\lambda$ with ν_j finite are approximated similarly to (4.9) - (4.11)

$$(\Gamma_{12}\Gamma_{23}\Gamma_{34}\Gamma_{41})^{1/2} \approx \frac{1}{[2\pi\rho_q(s)]^2} \left(\frac{2\sin\lambda s}{\lambda}\right)^{4\Delta} \prod_{n=1}^4 \left(\frac{\Gamma(\Delta \pm i\nu_n)}{\Gamma(2\Delta)}\right)^{1/2}. \quad (5.17)$$

We also use that $\prod_{j=1}^4 \rho_q(s_j) \approx \rho_q(s)^4$. Putting it all together the four-point function takes the form

$$\frac{1}{\Gamma(2\Delta)^2} \frac{1}{Z} \int_0^\infty ds \rho_q(s) e^{-\beta E(s)} \left(\frac{2\sin\lambda s}{\lambda}\right)^{4\Delta} \int_{-\infty}^\infty \frac{\prod_{n=1}^4 d\nu_n}{(2\pi)^3} \delta\left(\sum_{n=1}^4 \nu_n\right) e^{-\sum_{n=1}^4 \beta_n (E_n - E)} \quad (5.18)$$

$$\left(\frac{2\sin\lambda s}{\lambda}\right)^{i(\nu_2 + \nu_4)} \Gamma(i(\nu_2 + \nu_4)) \prod_{n=1}^4 \Gamma(\Delta - (-1)^n i\nu_n). \quad (5.19)$$

We dropped the second complex conjugate term in (5.7) because it gives an exponentially decaying in time, instead of growing, contribution. The energy differences can be expressed as $E_n - E \approx (s_n - s)\partial_s E$ and then using (5.15), (5.16) for $s_n - s$.

Let us make a few comments about (5.18), (5.19). Similarly to the two-point function, the integral over the average energy just sets the relation between average energy and temperature according to the saddle equation (3.11). In the rest of the integral the average energy is set to its saddle value. In fact, the rest of the integral has the same form as in JT gravity [29], except for $\partial_s E = \frac{2\pi v}{\beta}$ in the Boltzmann factors. So the only difference with the computation in the Schwarzian limit, is the relation between the average energy and temperature, which is determined by the saddle of the average energy s integral. The integrals over ν_j can be computed and give¹¹

$$\frac{\langle \mathcal{O}_1 \mathcal{O}_2 \mathcal{O}_3 \mathcal{O}_4 \rangle_\beta}{\langle \mathcal{O}_1 \mathcal{O}_3 \rangle_\beta \langle \mathcal{O}_2 \mathcal{O}_4 \rangle_\beta} = z^{-2\Delta} U(2\Delta, 1; 1/z), \quad z = \frac{\lambda}{4\lambda_L} \frac{e^{\frac{\lambda_L}{2}(-i\frac{\beta}{2} + u_2 + u_4 - u_1 - u_3)}}{\cosh \frac{\lambda_L}{2}(i\frac{\beta}{2} + u_{13}) \cosh \frac{\lambda_L}{2}(i\frac{\beta}{2} + u_{24})}. \quad (5.20)$$

where $U(a, 1, x) = \int_0^\infty \frac{dy}{y} e^{-xy} \frac{y^a}{(1+y)^a}$ is the confluent hypergeometric function. The Lyapunov exponent is

$$\lambda_L = \frac{2\pi v}{\beta} = \partial_s E = \frac{2\pi}{\beta} \left(1 - \frac{2}{\pi} \lambda s\right). \quad (5.21)$$

For example, we can choose a symmetric configuration of operators

$$iu_1 = \frac{3}{4}\beta - i\frac{u}{2}, \quad iu_2 = \frac{1}{2}\beta + i\frac{u}{2}, \quad iu_3 = \frac{1}{4}\beta - i\frac{u}{2}, \quad iu_4 = i\frac{u}{2}, \quad z = \frac{\lambda}{4\lambda_L} e^{\lambda_L u}. \quad (5.22)$$

The result (5.20) is valid in the long time limit

$$\lambda \rightarrow 0, \quad z - \text{fixed}. \quad (5.23)$$

¹¹In this computation, it is convenient to use the integral identity (4.13) several times.

It resums an infinite power series in z , interpolating between early time Lyapunov growth and late time quasinormal mode decay

$$z^{-2\Delta}U(2\Delta, 1; 1/z) \approx \begin{cases} 1 - 4\Delta^2 z, & z \ll 1, \\ \frac{\log z}{\Gamma(2\Delta)z^{2\Delta}}, & z \gg 1. \end{cases} \quad (5.24)$$

Note that both the early time growth and the late time decay are slower than in JT gravity.

It is also interesting to note that the last expression in (5.21) for the Lyapunov exponent makes it clear that the bound on chaos [17] is satisfied

$$\lambda_L = \frac{2\pi}{\beta} \left(1 - \frac{2}{\pi} \lambda s \right) \leq \frac{2\pi}{\beta} \quad (5.25)$$

since $s \geq 0$. The deviation from maximality is akin to a stringy correction [14].

Our result (5.20) agrees with [15] where it was derived by summing multi-ladder diagrams and with [16] where it was derived from the $G\Sigma$ effective action. The leading linear in z term was previously computed in [30, 31]. The form of OTOC (5.20) is similar to JT gravity [6], except that z now has more non-trivial dependence on the temperature through the Lyapunov exponent $\lambda_L(\beta)$.

5.3 Euclidean four-point function

We can also compute the euclidean four-point function in large p SYK as a limit of the DSSYK correlator (2.19), (2.24). Of course, the answer we expect is that at leading order it factorizes into Wick contractions with the two-point function given in (4.13). Nevertheless, it is interesting to see how this arises from the chord diagrams. In fact, this is similar for all higher point correlators. The result is that the approximation of the 6j-symbol as a delta-function (5.10) leads to factorized n -point correlators.

We compute the four-point function with euclidean time separations in large p SYK, i.e. in the limit $\lambda \rightarrow 0$ and β_j finite. We can approximate the 6j-symbol by the delta-function (5.10) and the four-point function (2.19), (2.24) takes the form

$$\langle \text{Tr} e^{-\beta H} \mathcal{O}(\tau_1) \dots \mathcal{O}(\tau_4) \rangle = \frac{1}{Z} \int_0^{\pi/\lambda} \prod_{j=1}^4 (ds_j \rho_q(s_j) e^{-\beta_j E_j(s_j)}) (\Gamma_{12}\Gamma_{23}\Gamma_{34}\Gamma_{41})^{1/2} \quad (5.26)$$

$$\left(\frac{\delta(s_1 - s_3)}{\rho_q(s_1)} + \frac{\delta(s_2 - s_4)}{\rho_q(s_2)} + \frac{\delta(s_1 + s_3 - s_2 - s_4)}{\rho_q(s)} \right), \quad (5.27)$$

where $(\tau_{ij} = \tau_i - \tau_j)$

$$\beta_1 = \beta - \tau_{14}, \quad \beta_2 = \tau_{12}, \quad \beta_3 = \tau_{23}, \quad \beta_4 = \tau_{34}, \quad \beta > \tau_1 > \tau_2 > \tau_3 > \tau_4 > 0. \quad (5.28)$$

We will focus on computing the last term coming from the 6j-symbol. The other two terms can be computed in a similar manner. This computation is essentially identical to the semi-classical limit of the four-point function in JT gravity [24].

We proceed similarly to the computation of OTOC in the previous subsection. We change variables to the average energy s and energy differences ν_1, \dots, ν_4 and make the same approximations for the q-gamma functions. We find for the last term in (5.27)

$$\frac{1}{Z} \int_0^\infty ds \rho_q(s) e^{-\beta E(s)} \left(\frac{2 \sin \lambda s}{\lambda} \right)^{4\Delta} \quad (5.29)$$

$$\int_{-\infty}^\infty \frac{d\nu_1 d\nu_2 d\nu_3 d\nu_4}{(2\pi)^2} \exp \left\{ -\nu_1 \partial_s E \frac{\beta_{1-2-3+4}}{2} - \nu_2 \partial_s E \frac{\beta_{1+2-3-4}}{2} \right\} \prod_{n=1}^4 \left(\frac{\Gamma(\Delta \pm i\nu_n)}{\Gamma(2\Delta)} \right)^{1/2} \delta(\nu_1 + \nu_3) \delta(\nu_2 + \nu_4), \quad (5.30)$$

where $\beta = \sum_{j=1}^4 \beta_j$ and $\beta_{1-2+3-4} = \beta_1 - \beta_2 + \beta_3 - \beta_4$. The integral over the average energy s has the saddle (3.11). In the rest of the integral we set the average energy to take the saddle value, e.g. $\lambda_L = \partial_s E = \frac{2\pi v}{\beta}$. Two integrals, e.g. over ν_3, ν_4 , are straightforward using the delta-functions. The remaining two integrals give

$$\int_{-\infty}^\infty d\nu_1 d\nu_2 \exp \left\{ -\nu_1 \partial_s E \left(\frac{\beta}{2} - \tau_{13} \right) - \nu_2 \partial_s E \left(\frac{\beta}{2} - \tau_{24} \right) \right\} \frac{\Gamma(\Delta \pm i\nu_1) \Gamma(\Delta \pm i\nu_2)}{\Gamma(2\Delta)^2} \quad (5.31)$$

$$= \lambda^{-4\Delta} \left(\frac{\cos \frac{\pi v}{2}}{\cos \frac{\pi v}{2} \left(\frac{\beta}{2} - \tau_{13} \right)} \frac{\cos \frac{\pi v}{2}}{\cos \frac{\pi v}{2} \left(\frac{\beta}{2} - \tau_{24} \right)} \right)^{2\Delta}. \quad (5.32)$$

This is just a product of two-point functions (4.13). A similar computation for the first two terms in (5.27) gives the other two Wick contractions. Altogether we have a factorized answer

$$\langle \mathcal{O}_1 \mathcal{O}_2 \mathcal{O}_3 \mathcal{O}_4 \rangle_\beta = \langle \mathcal{O}_1 \mathcal{O}_2 \rangle \langle \mathcal{O}_3 \mathcal{O}_4 \rangle_\beta + \langle \mathcal{O}_1 \mathcal{O}_4 \rangle \langle \mathcal{O}_2 \mathcal{O}_3 \rangle_\beta + \langle \mathcal{O}_1 \mathcal{O}_3 \rangle \langle \mathcal{O}_2 \mathcal{O}_4 \rangle_\beta \quad (5.33)$$

with the two-point functions given in (4.13).

5.4 Euclidean $2n$ -point function

The computation for the euclidean four-point function above is in fact straightforward to generalize to any $2n$ -point function and the corresponding chord diagrams (2.20). In each chord diagram we substitute all 6j-symbols by delta-functions according to (5.10). Then the energy integrals can be computed and one obtains that the $2n$ -point function is given by Wick contractions with the two-point function (4.13). This is similar to how the correlators in JT gravity in the semi-classical limit reduce to the generalized free field [24, 28]. Corrections in λ can also be computed if desired.

Acknowledgements

I am grateful to Tom Hartman, Henry Lin, Juan Maldacena, Joaquin Turiaci, Jinzhao Wang for useful discussions and Daniel Jafferis, David Kolchmeyer, Julian Sonner for collaboration on related topics. I am supported by NSF grant PHY-2014071. A part of this work was completed at KITP, Santa Barbara and supported by the National Science Foundation under Grant No. NSF PHY-1748958.

A Asymptotics of the 6j-symbol

To derive (5.7) we do the following. The parameters of the Askey-Wilson function (5.3) corresponding to (5.1) are

$$a = d^* = \Delta + is_1, \quad b = c^* = \Delta + is_3. \quad (\text{A.1})$$

The basic hypergeometric function in (5.1), (5.3) can be reduced to the ordinary hypergeometric function

$${}_4\phi_3 \left(\begin{matrix} q^{\Delta+i(s_1+s_2)}, q^{\Delta+i\nu_1}, q^{\Delta+i(s_1+s_4)}, q^{\Delta-i\nu_4} \\ q^{2\Delta+i(s_1+s_3)}, q^{2\Delta+i(\nu_1+\nu_2)}, q^{2is_1+1} \end{matrix} ; q, q \right) \approx {}_2\phi_1 \left(\begin{matrix} q^{\Delta+i\nu_1}, q^{\Delta-i\nu_4} \\ q^{2\Delta+i(\nu_1+\nu_2)} \end{matrix} ; q, q \right) \quad (\text{A.2})$$

$$\approx {}_2F_1 \left(\begin{matrix} \Delta + i\nu_1, \Delta - i\nu_4 \\ 2\Delta + i(\nu_1 + \nu_2) \end{matrix} ; 1 \right) \quad (\text{A.3})$$

$$= \frac{\Gamma(2\Delta + i(\nu_1 + \nu_2))\Gamma(i(\nu_2 + \nu_4))}{\Gamma(\Delta + i\nu_2)\Gamma(\Delta - i\nu_3)}. \quad (\text{A.4})$$

A useful formula for computing q-gamma function asymptotics is

$$\frac{\Gamma_q(a + is)}{\Gamma_q(b + is)} \approx \left(\frac{1 - e^{-i\lambda s}}{\lambda} \right)^{a-b}, \quad \lambda \rightarrow 0, \lambda s - \text{fixed}. \quad (\text{A.5})$$

Then the q-gamma functions in the prefactor in (5.2) are estimated as

$$\frac{\Gamma_q(d - a)}{\Gamma_q(a + b)\Gamma_q(a + c)\Gamma_q(d \pm x)\Gamma_q(\tilde{d} \pm y)} \quad (\text{A.6})$$

$$= 2\pi\rho_q(s) \frac{\Gamma_q(2is)\Gamma_q(-2is)\Gamma_q(-2is_1)}{\Gamma_q(2\Delta + is_{1+3})\Gamma_q(\Delta - is_{1+2})\Gamma_q(\Delta - is_{1+4})} \frac{1}{\Gamma_q(2\Delta + i\nu_{1+2})\Gamma_q(\Delta - i\nu_1)\Gamma_q(\Delta + i\nu_4)} \quad (\text{A.7})$$

$$\approx 2\pi\rho_q(s) \left(\frac{2\sin(\lambda s)}{\lambda} \right)^{-4\Delta+i(\nu_2+\nu_4)} \frac{1}{\Gamma_q(2\Delta + i\nu_{1+2})\Gamma_q(\Delta - i\nu_1)\Gamma_q(\Delta + i\nu_4)}, \quad (\text{A.8})$$

where we used short notation e.g. $s_{1+2} = s_1 + s_2$. In the second line we multiplied and divided by $2\pi\rho_q(s) = \frac{1}{\Gamma_q(\pm 2is)}$. In the third line we used (A.5). We have from (A.4) and (A.8)

$$W_{q^{is_4}}(q^{is_2}; q^{\Delta \pm is_1}, q^{\Delta \pm is_3} | q) \approx 2\pi\rho_q(s) \left(\frac{2\sin(\lambda s)}{\lambda} \right)^{-4\Delta+i(\nu_2+\nu_4)} \frac{\Gamma(i(\nu_2 + \nu_4))}{\prod_{n=1}^4 \Gamma_q(\Delta + (-1)^n i\nu_n)}. \quad (\text{A.9})$$

Similarly we find

$$(\gamma_{12}\gamma_{23}\gamma_{34}\gamma_{41})^{1/2} \approx \frac{1}{[2\pi\rho_q(s)]^2} \left(\frac{2\sin \lambda s}{\lambda} \right)^{4\Delta} \prod_{n=1}^4 \Gamma(\Delta \pm i\nu_n)^{1/2}. \quad (\text{A.10})$$

Putting everything together we derive (5.7).

References

- [1] S. Sachdev and J. Ye, *Gapless spin-fluid ground state in a random quantum heisenberg magnet*, *Physical Review Letters* **70** (1993) 3339–3342.
- [2] A. Kitaev, *A simple model of quantum holography 1*, Talk at KITP, April 7, 2015, <http://online.kitp.ucsb.edu/online/entangled15/kitaev/>.
- [3] A. Kitaev, *A simple model of quantum holography 2*, Talk at KITP, May 27, 2015, <http://online.kitp.ucsb.edu/online/entangled15/kitaev2/>.
- [4] J. Maldacena and D. Stanford, *Remarks on the Sachdev-Ye-Kitaev model*, *Phys. Rev. D* **94** (2016) 106002 [[1604.07818](#)].
- [5] K. Jensen, *Chaos in AdS₂ Holography*, *Phys. Rev. Lett.* **117** (2016) 111601 [[1605.06098](#)].
- [6] J. Maldacena, D. Stanford and Z. Yang, *Conformal symmetry and its breaking in two dimensional Nearly Anti-de-Sitter space*, *PTEP* **2016** (2016) 12C104 [[1606.01857](#)].
- [7] J. Engelsöy, T.G. Mertens and H. Verlinde, *An investigation of AdS₂ backreaction and holography*, *JHEP* **07** (2016) 139 [[1606.03438](#)].
- [8] A. Kitaev and S.J. Suh, *The soft mode in the Sachdev-Ye-Kitaev model and its gravity dual*, *JHEP* **05** (2018) 183 [[1711.08467](#)].
- [9] A. Kitaev and S.J. Suh, *Statistical mechanics of a two-dimensional black hole*, *JHEP* **05** (2019) 198 [[1808.07032](#)].
- [10] T.G. Mertens and G.J. Turiaci, *Solvable Models of Quantum Black Holes: A Review on Jackiw-Teitelboim Gravity*, [[2210.10846](#)].
- [11] S.H. Shenker and D. Stanford, *Black holes and the butterfly effect*, *JHEP* **03** (2014) 067 [[1306.0622](#)].
- [12] M. Berkooz, M. Isachenkov, V. Narovlansky and G. Torrents, *Towards a full solution of the large N double-scaled SYK model*, *JHEP* **03** (2019) 079 [[1811.02584](#)].
- [13] H.W. Lin, *The bulk Hilbert space of double scaled SYK*, *JHEP* **11** (2022) 060 [[2208.07032](#)].
- [14] S.H. Shenker and D. Stanford, *Stringy effects in scrambling*, *JHEP* **05** (2015) 132 [[1412.6087](#)].
- [15] Y. Gu, A. Kitaev and P. Zhang, *A two-way approach to out-of-time-order correlators*, *JHEP* **03** (2022) 133 [[2111.12007](#)].
- [16] C. Choi, F.M. Haehl, M. Mezei and G. Sárosi, *Effective description of sub-maximal chaos: stringy effects for SYK scrambling*, [[2301.05698](#)].
- [17] J. Maldacena, S.H. Shenker and D. Stanford, *A bound on chaos*, *JHEP* **08** (2016) 106 [[1503.01409](#)].
- [18] D.L. Jafferis, D.K. Kolchmeyer, B. Mukhametzhanov and J. Sonner, *JT gravity with matter, generalized ETH, and Random Matrices*, [[2209.02131](#)].

- [19] A. Goel, V. Narovlansky and H. Verlinde, *Semiclassical geometry in double-scaled SYK*, [[2301.05732](#)].
- [20] B. Mukhametzhanov, *Comments on Double-Scaled SYK*, Abu Dhabi Meeting on Theoretical Physics, January 10, 2023, <https://nyuad.nyu.edu/en/events/2023/january/abu-dhabi-meeting-on-theoretical-physics/agenda.html>.
- [21] P. Saad, S.H. Shenker and D. Stanford, *JT gravity as a matrix integral*, [[1903.11115](#)].
- [22] J. Polchinski, *String Theory*, vol. 1 of *Cambridge Monographs on Mathematical Physics*, Cambridge University Press (1998), [10.1017/CBO9780511816079](#).
- [23] M. Berkooz, P. Narayan and J. Simon, *Chord diagrams, exact correlators in spin glasses and black hole bulk reconstruction*, *JHEP* **08** (2018) 192 [[1806.04380](#)].
- [24] D.L. Jafferis, D.K. Kolchmeyer, B. Mukhametzhanov and J. Sonner, *Matrix models for eigenstate thermalization*, [[2209.02130](#)].
- [25] W. Groenevelt, *Wilson function transforms related to Racah coefficients*, *Acta Appl. Math.* **91** (2006) 133 [[math/0501511](#)].
- [26] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, Encyclopedia of Mathematics and its Applications, Cambridge University Press, 2 ed. (2004), [10.1017/CBO9780511526251](#).
- [27] W. Groenevelt, *The Wilson function transform*, *Int. Math. Res. Not.* (2003) 2779 [[math/0306424](#)].
- [28] T.G. Mertens, G.J. Turiaci and H.L. Verlinde, *Solving the Schwarzian via the Conformal Bootstrap*, *JHEP* **08** (2017) 136 [[1705.08408](#)].
- [29] H.T. Lam, T.G. Mertens, G.J. Turiaci and H. Verlinde, *Shockwave S-matrix from Schwarzian Quantum Mechanics*, *JHEP* **11** (2018) 182 [[1804.09834](#)].
- [30] A. Streicher, *SYK Correlators for All Energies*, *JHEP* **02** (2020) 048 [[1911.10171](#)].
- [31] C. Choi, M. Mezei and G. Sárosi, *Exact four point function for large q SYK from Regge theory*, *JHEP* **05** (2021) 166 [[1912.00004](#)].