

Properly outer and strictly outer actions of finite groups on prime C*-algebras

Costel Peligrad

Department of Mathematical Sciences, University of Cincinnati, PO Box 210025, Cincinnati, OH 45221-0025, USA. E-mail address: costel.peligrad@uc.edu

Key words and phrases. dynamical system, algebra of local multipliers, strictly outer actions.

2020 Mathematics Subject Classification. Primary 46L55, 46L40; Secondary 20F29.

ABSTRACT. An action of a compact, in particular finite group on a C*-algebra is called properly outer if no automorphism of the group that is distinct from identity is implemented by a unitary element of the algebra of local multipliers of the C*-algebra. In this paper I define the notion of strictly outer action (similar to the definition for von Neumann factors in [11]) and prove that for finite groups and prime C*-algebras, it is equivalent to the proper outerness of the action. For finite abelian groups this is equivalent to other relevant properties of the action.

1 Introduction

An action α of a locally compact group or a quantum group, G , on a von Neumann factor M is said to be strictly outer when the relative commutant of the factor in the crossed product is trivial. The action is called minimal if the inclusion $M^\alpha \subset M$ is irreducible, that is the relative commutant $(M^\alpha)' \cap M$ of the fixed point algebra M^α in M is trivial. For integrable actions of quantum groups on factors an action is minimal if and only if it is strictly outer [11]. It can be shown that, in particular, for finite groups, an action α on a factor M is strictly outer if and only if it is properly outer (that is, no $\alpha_g, g \neq e$ is implemented by a unitary element of M). In the next Section, I define strictly outer actions of finite groups on C*-algebras and, in Section 3.1., prove that an action of a finite group, G , on a prime C*-algebra, A , is strictly outer if and only if it is properly outer (that is, no $\alpha_g, g \neq e$, is implemented by a unitary element in the algebra of local multipliers of A). In Section 3.2. I study actions of finite abelian groups on prime C*-algebras and state some equivalent conditions to proper and therefore strict outerness. These results extend and complement some known results in the literature ([4], [5], [7]). In [2] similar results were

obtained for compact abelian group actions on separable prime C*-algebras with an additional condition. I do not require separability of C*-algebras.

2 Definitions and preliminary results

If A is a C*-algebra, denote by \mathcal{I} the set of all closed essential, two sided ideals of A . A two sided ideal $I \subset A$ is called essential if $\{a \in A : aI = (0)\} = (0)$. If $I \subset J$ are two such ideals then according to [7, Proposition 3.12.8.] we have the inclusion $M(J) \subset M(I)$ where $M(I)$, respectively $M(J)$ are the multiplier algebras of I , respectively J . The inductive limit of the system $\{M(I) : I \in \mathcal{I}\}$ is called the algebra of local multipliers of A and denoted $M_{loc}(A)$ [1]. Originally, this algebra was denoted $M^\infty(A)$ and called the algebra of essential multipliers of A [6]. Recall that a C*-algebra, A , is called prime if every (two sided) ideal of A is essential.

In the next lemma I collect some known results

Lemma 1 *Let A be a C*-algebra. The following properties are equivalent*

- a) A is prime.
- b) The center of $M_{loc}(A)$ consists of scalars.
- c) If B is a C*-subalgebra of $M_{loc}(A)$ such that $A \subset B \subset M_{loc}(A)$, then the center of B consists of scalars.

Proof. The equivalence of a) and b) follows from [2, Proposition 3.1.] (see also [6, page 303, statements 6), 4) and 5])). The implication b) \Rightarrow c) follows from [1, Lemma 3.2.2.] and the implication c) \Rightarrow b) is obvious. ■

In the rest of this paper we will consider a C*-dynamical system (A, G, α) where A is a C*-algebra, G a finite (or compact) group and α a faithful action of G on A , that is for each $g \in G$, α_g is an automorphism of A such that $\alpha_{g_1 g_2} = \alpha_{g_1} \alpha_{g_2}$ for all $g_1, g_2 \in G$. We will denote by $A^\alpha = \{a \in A : \alpha_g(a) = a, g \in G\}$ the fixed point algebra of the action. The action α is called faithful if $\alpha_g \neq id$ for $g \neq e$ where e is the unit of G . If the group G is finite, the action α of G on A has a natural extension to $M_{loc}(A)$.

Lemma 2 *Suppose G is a compact group. Then,*

- a) Every approximate identity of A^α is an approximate identity of A ,
- b) $M(A)^\alpha = M(A^\alpha)$.

Proof. a) follows from the more general case of compact quantum group actions [3, Lemma 2.7.]. To prove b), notice first that $M(A)^\alpha \subset M(A^\alpha)$, then, using a) it follows that $M(A^\alpha) \subset M(A)$, so $M(A^\alpha) \subset M(A)^\alpha$. ■

Lemma 3 *Let (A, G, α) be a C^* -dynamical system with G finite and A prime. Then,*

- a) $M_{loc}(A)^\alpha \subset M_{loc}(A^\alpha)$.
- b) *If $x \in M_{loc}(A)$ commutes with A^α then x commutes with $M_{loc}(A)^\alpha$.*
- c) *If A^α is prime, then the center of $M_{loc}(A)^\alpha$ consists of scalars.*
- d) $A' \cap M_{loc}(A)$ consists of scalars.

Proof. a) Since G is finite, every essential ideal $J \subset A$ contains the α -invariant essential ideal $I = \bigcap_{g \in G} \alpha_g(J)$. Therefore, $M_{loc}(A)$ is the inductive limit of $\{M(I) : I \subset A \text{ } \alpha\text{-invariant essential ideal}\}$. If $I \subset A$ is a closed two sided α -invariant essential ideal of A , it is obvious that I^α is an essential ideal of A^α . From Lemma 2 b) applied to I instead of A it follows that $M(I)^\alpha = M(I^\alpha) \subset M_{loc}(A^\alpha)$ and, therefore, $M_{loc}(A)^\alpha \subset M_{loc}(A^\alpha)$.

To prove b), let $I \subset A$ be an α -invariant essential ideal. Then I^α is ideal of A^α so x commutes with I^α . If $m \in M(I^\alpha)$ and $a \in I^\alpha$ we have

$$xam = amx.$$

and

$$xam = axm.$$

Therefore

$$a(xm - mx) = 0.$$

so $I^\alpha(xm - mx) = \{0\}$. Applying Lemma 2 a) to I instead of A , it follows that $I(xm - mx) = \{0\}$. From [1, Lemma 2.3.3.ii)] we get $xm - mx = 0$. Applying Lemma 2 b) to I , we have $M(I^\alpha) = M(I)^\alpha$, so x commutes with $M(I)^\alpha$ for every α -invariant essential ideal of A . Therefore, x commutes with $\lim_I M(I)^\alpha = M_{loc}(A)^\alpha$.

c) If A^α is prime, from Lemma 1 b) applied to A^α instead of A it follows that the center of $M_{loc}(A^\alpha)$ consists of scalars. Applying a) it follows that $M_{loc}(A)^\alpha \subset M_{loc}(A^\alpha)$. From Lemma 1 c) applied to A^α instead of A it follows that the center of $M_{loc}(A)^\alpha$ consists of scalars.

d) Follows from part b) for trivial α . ■

The cross product $A \times_\alpha G$ of A by a finite group, G , is defined as the C^* -completion of the algebra of all functions $f : G \rightarrow A$ with the following operations:

$$(f \cdot h)(g) = \sum_{p \in G} f(p) \alpha_p(h(p^{-1}g)).$$

$$f^*(g) = \alpha_g(f(g^{-1}))^*.$$

As G is a finite group, each $f \in A \times_\alpha G$ can be written $f = \sum a_g \delta_g$ where $a_g \in A$ and δ_g is the function on G with values in the multiplier algebra $M(A)$ of A such that $\delta_g(g) = 1$ and $\delta_g(p) = 0$ for $p \neq g$.

If A is a C^* -algebra then, $A \subset A'' \subset B(H_u)$ where A'' is enveloping von Neumann algebra of A in $B(H_u)$ and H_u is the Hilbert space of the universal

representation of A [7, Section 3.7.]. Denote by id the identity representation of A in $B(H_u)$. If G is a finite group, denote by λ the left regular representation of G on $l^2(G)$. Denote by $(\tilde{id}, \lambda, l^2(G, H_u))$, the covariant representation of (A, G, α) on $l^2(G, H_u)$, where

$$\tilde{id}(a)(h) = \alpha_{h^{-1}}(a), \lambda_g(\xi)(h) = \xi(g^{-1}h).$$

Consider the regular representation, $\tilde{id} \times \lambda$, of $A \times_\alpha G$ induced by $(\tilde{id}, \lambda, l^2(G, H_u))$:

$$(\tilde{id} \times \lambda)(f)(p) = \sum_{g \in G} \alpha_{p^{-1}}(f(g)) \xi(g^{-1}p).$$

Then, $\tilde{id} \times \lambda$ is a faithful non degenerate representation of $A \times_\alpha G$ on $l^2(G, H_u)$ [7, Section 7.7.].

Remark 4 $A \times_\alpha G \subset M(A) \times_\alpha G \subset A'' \times_\alpha G$ where $A'' \times_\alpha G$ is the W^* -cross product of A'' by the double dual action α'' of α .

I will also use the following

Lemma 5 *Suppose G is finite. Then, $M_{loc}(A) \subset M_{loc}(A \times_\alpha G)$.*

Proof. According to the proof of Lemma 3. a), $M_{loc}(A)$ is the inductive limit of $\{M(I) : I \subset A \text{ } \alpha\text{-invariant essential ideal}\}$. A standard argument, using that G is finite, shows that for every α -invariant essential ideal $I \subset A$, $I \times_\alpha G$ is an essential ideal of $A \times_\alpha G$. Since $M(I) \subset M(I \times_\alpha G) \subset M_{loc}(A \times_\alpha G)$, it follows that $M_{loc}(A) \subset M_{loc}(A \times_\alpha G)$. ■

In the following definitions we assume that the group G is finite.

Definition 6 a) *If (A, G, α) is a C^* -dynamical system with A prime, the action α is called properly outer if no $\alpha_g, g \neq e$ is implemented by a unitary element in $M_{loc}(A)$.*

b) *The action is called strictly outer if $A' \cap M_{loc}(A \times_\alpha G) = \mathbb{C}I$, where A' is the commutant of A .*

In [2, Proposition 3.2.] the authors prove that two other statements are equivalent with the above Definition 6. part a) for separable C^* -algebras. For not necessarily separable C^* -algebras, I use the statement given in the above Definition 6. a). Actually, as proven in [2], the other two statements imply the Definition 6. a) for not necessarily separable C^* -algebras.

Now, let (A, G, α) be a C^* -dynamical system with G compact abelian. Denote by \widehat{G} the dual of G . If $\gamma \in \widehat{G}$ let $A_\gamma = \{a \in A : \alpha_g(a) = \langle g, \gamma \rangle a, g \in G\}$ be the corresponding spectral subspace of A . Denote by $sp(\alpha)$ the Arveson spectrum of α and by $\Gamma(\alpha)$ the Connes spectrum of α :

$$sp(\alpha) = \left\{ \gamma \in \widehat{G} : A_\gamma \neq \{0\} \right\}.$$

and, if \mathcal{H} is the set of all non zero α -invariant hereditary subalgebras of A

$$\Gamma(\alpha) = \cap \{sp(\alpha|_B : B \in \mathcal{H})\}.$$

I will list some standard facts about the Arveson spectrum of actions of compact abelian groups (in particular finite abelian groups) that will be used in Section 3.2.:

- 1) The linear span of $\cup \{A_\gamma : \gamma \in Sp(\alpha)\}$ is dense in A . Therefore, if $\langle t, \gamma \rangle = 1$ for every $\gamma \in Sp(\alpha)$, then α_t is the identity automorphism.
- 2) If $\gamma_1, \gamma_2 \in Sp(\alpha)$, then $A_{\gamma_1} A_{\gamma_2} \subset A_{\gamma_1 + \gamma_2}$. Here, $A_{\gamma_1} A_{\gamma_2}$ denotes the linear span of $\{ab : a \in A_{\gamma_1}, b \in A_{\gamma_2}\}$.
- 3) If $\gamma \in Sp(\alpha)$, $A_\gamma A_\gamma^*$ and $A_\gamma^* A_\gamma$ are ideals of A^α . Here $A_\gamma A_\gamma^*$ denotes the linear span of $\{ab^* : a, b \in A_\gamma\}$.
- 4) If A^α is prime, $Sp(\alpha)$ is a subgroup of \widehat{G} . This follows from 2) and 3).
- 5) If A^α is prime and the action α is faithful (that is $\alpha_t \neq id$ for $t \neq 0$), then $Sp(\alpha) = \widehat{G}$. This follows from 4) and 1).

The following result was proven by Olesen for simple C^* -algebras, A and actions for which $sp(\alpha)/\Gamma(\alpha)$ is compact in $\widehat{G}/\Gamma(\alpha)$ in [4, Theorem 4.2.] (see also [7, Theorem 8.9.7.]). If (A, G, α) is a C^* -dynamical system, A is called G -prime if every non zero α -invariant ideal of A is an essential ideal.

Proposition 7 *Let (A, G, α) be a C^* -dynamical system with A G -prime and G compact abelian such that $\widehat{G}/\Gamma(\alpha)$ is finite (in particular G finite group). If $t_0 \in G$ is such that $\langle t_0, \gamma \rangle = 1$ for every $\gamma \in \Gamma(\alpha)$ then there exists an α -invariant (essential) ideal $J \subset A$ and a unitary operator $u \in M(J)^\alpha$ such that $\alpha_{t_0} = Adu$.*

Proof. Applying [7. Proposition 8.8.7.] for A G -prime and G compact with

$\widehat{G}/\Gamma(\alpha)$ finite, it follows that there exists a non zero α -invariant hereditary C^* -subalgebra, $C \subset A$ such that $sp(\alpha|_C) = \Gamma(\alpha)$. Therefore, $\alpha_{t_0}|_C = id$. Let $J = \overline{ACA}$ be the closed two sided ideal of A generated by C . Since C is α -invariant and A is G -prime, it follows that J is an essential α -invariant ideal of A . Using an approximate identity of A it follows that $C \subset J \subset J''$.

Let p be the open projection corresponding to C in A'' . Then $p \in J''$. Denote by 1_J the unit of J'' (1_J is the open projection in A'' corresponding to J). Notice that the central support $c(p)$ of p in J'' equals 1_J . Indeed if $q \geq p$ is a central projection in J'' , then qJ'' is an ideal of J'' that contains C , so $q = 1_J$.

Applying [7, Lemma 8.9.1.] to $M = J''$ and $w = p$, it follows that there exists a unique unitary $u \in J''$ such that $up = p$ and $\alpha''_{t_0}|_{J''} = Adu|_{J''}$. Since $\alpha''_{t_0}(p) = p = upu^*$ we have $up = pu = p$. Since $\alpha''_{t_0}|_{J''} = Adu|_{J''}$ and $M(J) \subset J''$, it follows that $\alpha_{t_0}|_{M(J)} = Adu|_{M(J)}$. Also, since G is abelian, $\alpha_t(p) = p$, for every $t \in G$, and u is unique with the listed properties, it follows that, $\alpha_t(u) = u$ for every $t \in G$.

We will prove next that $u \in M(J)$. Let $J_0 = \{x \in J : ux \in J\}$. Clearly J_0 is a closed right ideal of J . Now, if $y \in J$, and $x \in J_0$ we have $uyx = uy u^* ux = \alpha_{t_0}(y)ux \in J$, so J_0 is also a left ideal. Notice that $C \subset J_0$. Indeed, if $x \in C$, we have $ux = upx = px = x \in C \subset J$, so $x \in J_0$. Therefore, $C \subset J_0 \subset J$. Since J is the (two sided) ideal generated by C it follows that $J_0 = J$, so u is a left multiplier of J . To prove that u is also a right multiplier of J , notice that if $x \in J$ we have $xu = uu^*xu = u\alpha_{t_0}^{-1}(x) \in J$. Since $M(A) \subset M(J)$ (in $M_{loc}(A)$), it follows that $\alpha_{t_0} = Adu$. ■

Corollary 8 *Let (A, G, α) be a C^* -dynamical system with A G -prime and G finite abelian. If $t_0 \in G$ is such that $\langle t_0, \gamma \rangle = 1$ for every $\gamma \in \Gamma(\alpha)$ then α_{t_0} is inner in $M_{loc}(A)^\alpha$.*

Proof. Proposition 7 shows that $\alpha_{t_0} = Adu, u \in M(J)^\alpha \subset M_{loc}(J)$. By [6, page 303, statement 3)], since J is an essential ideal of A , $M_{loc}(J) = M_{loc}(A)$, so α_{t_0} is inner in $M_{loc}(A)^\alpha$. ■

I will close this section with the following

Theorem 9 *Let (A, G, α) be a C^* -dynamical system with G finite abelian, α faithful and A G -prime. Then the following statements are equivalent:*

- a) $Sp(\alpha) = \Gamma(\alpha)$.
- b) A^α is prime.
- c) The center of $M_{loc}(A)^\alpha$ consists of scalars.
- d) No $\alpha_t \neq id$ is implemented by a unitary in $M_{loc}(A)^\alpha$.

Proof. a) \Leftrightarrow b) This is [7 Theorem 8.10.4].

b) \Rightarrow c) Since A^α is prime, from Lemma 1, a) \Leftrightarrow b), with A^α instead of A it follows that the center of $M_{loc}(A^\alpha)$ is $\mathbb{C}I$. Applying Lemma 1, b) \Leftrightarrow c) to A^α instead of A , it follows that the center of $M_{loc}(A)^\alpha \subset M_{loc}(A^\alpha)$ is $\mathbb{C}I$.

c) \Rightarrow d) If for some $t_0 \in G$, α_{t_0} is implemented by a unitary $u \in M_{loc}(A)^\alpha$, then u commutes with A^α . From Lemma 3 b) it follows that u commutes with $M_{loc}(A)^\alpha$, so u belongs to the center of $M_{loc}(A)^\alpha$, hence u is a scalar and $\alpha_{t_0} = id$.

d) \Rightarrow a) Suppose $Sp(\alpha) \neq \Gamma(\alpha)$. Since $\Gamma(\alpha)$ is a subgroup of \widehat{G} , there exists $t_0 \in G$ such that $\langle t_0, \gamma \rangle = 1$ for every $\gamma \in \Gamma(\alpha)$ and $\langle t_0, \gamma \rangle \neq 1$ for some $\gamma \in Sp(\alpha)$. From Corollary 8 it follows that $\alpha_{t_0} \neq id$ and α_{t_0} is implemented by a unitary in $M_{loc}(A)^\alpha$, contradiction. ■

3 Outer and strictly outer actions of finite groups

3.1 Finite non abelian groups

Lemma 10 *Let (A, G, α) be a C^* -dynamical system with G finite.*

a) *Then $M(A \times_\alpha G) = M(A) \times_\alpha G$.*

b) *If A is prime and α is properly outer, then $M_{loc}(A \times_\alpha G)$ is isomorphic to $M_{loc}(A) \times_\alpha G$.*

Proof. a) It is obvious that $(\tilde{id} \times \lambda)(A \times_\alpha G)$ is included in the W^* -crossed product $(\tilde{id} \times \lambda)(A'' \times_{\alpha''} G)$. Since $\tilde{id} \times \lambda$ is a faithful and non degenerate representation of $A \times_\alpha G$, the multiplier algebra $M(A \times_\alpha G)$ is isometrically isomorphic with the algebra of multipliers of $(\tilde{id} \times \lambda)(A \times_\alpha G)$ in $((\tilde{id} \times \lambda)(A'' \times_{\alpha''} G)) \subset B(l^2(G, H_u))$ [7, Prop. 3.12.3]. Let $b = \sum_{g \in G} b_g \delta_g \in A'' \times_{\alpha''} G$ be a multiplier of $A \times_\alpha G$. If $a \in A$ is arbitrary, $\tilde{a} = a \delta_e \in A \times_\alpha G$, where e is the neutral element of G and \cdot is the multiplication of $A'' \times_{\alpha''} G$, we have

$$(b \cdot \tilde{a})_g = \sum_{p \in G} b_p \alpha_p(\tilde{a}_{p^{-1}g}) = b_g \alpha_g(a) \in A.$$

$$(\tilde{a} \cdot b)_g = \sum_{p \in G} \tilde{a}_p \alpha_p(b_{p^{-1}g}) = ab_g \in A.$$

Therefore, $b_g \in M(A)$ for every $g \in G$, so $b \in M(A) \times_\alpha G$.

b) Let I be an α -invariant essential ideal of A . Then, a simple computation shows that $I \times_\alpha G$ is an essential ideal of $A \times_\alpha G$ [10, Proposition 2.2.]. By a), $M(I \times_\alpha G) = M(I) \times_\alpha G$. If J is an essential ideal of $A \times_\alpha G$ then there exists an α -invariant essential ideal I of A such that $I \times_\alpha G \subset J$, so $M(J) \subset M(I \times_\alpha G) = M(I) \times_\alpha G$. To prove that such an ideal I exists, notice that, since A is prime and α properly outer, the action α is, in particular, "purely outer" in the sense used by Rieffel in [10]. From [10 Theorem 1.1.] it follows that $I_0 = J \cap A \neq \{0\}$. Since $J \subset A \times_\alpha G$ is essential it follows that I_0 is essential in A . Since G is finite, it follows that $I = \bigcap_{g \in G} \alpha_g(I_0) \subset I_0$ is an essential α -invariant ideal of A and $I \times_\alpha G \subset J$. Therefore, $M(J) \subset M(I \times_\alpha G)$. According to a), $M(I \times_\alpha G) = M(I) \times_\alpha G$. Therefore

$$M(J) \subset M(I) \times_\alpha G \subset M_{loc}(A) \times_\alpha G.$$

Hence

$$M_{loc}(A \times_\alpha G) \subset M_{loc}(A) \times_\alpha G.$$

On the other hand, by Lemma 5, $M_{loc}(A) \subset M_{loc}(A \times_\alpha G)$, so

$$M_{loc}(A) \times_\alpha G \subset M_{loc}(A \times_\alpha G).$$

■

Theorem 11 *Let (A, G, α) be a C^* -dynamical system with G finite, A prime and α faithful. Then α is properly outer if and only if it is strictly outer.*

Proof. Suppose α is properly outer and let $v = \sum_{h \in G} v(h)\delta_h \in A' \cap M_{loc}(A \times_\alpha G)$ ($= M_{loc}(A) \times_\alpha G$ by Lemma 10) be a unitary element. Then, for $a \in A$ denote $\tilde{a} = a\delta_e \in A \times_\alpha G \subset M_{loc}(A) \times_\alpha G$ ($= M_{loc}(A \times_\alpha G)$ by Lemma 10.). We have

$$v\tilde{a} = \tilde{a}v \text{ in } M_{loc}(A) \times_\alpha G \text{ for all } a \in A.$$

Therefore

$$\sum_h v(h)\alpha_h(\tilde{a}(h^{-1}g)) = \sum_h \tilde{a}(h)\alpha_h(v(h^{-1}g)), g \in G.$$

So

$$v(g)\alpha_g(a) = av(g) \text{ for all } g \in G, a \in A. \quad (*)$$

We will prove next that if $v(g) \neq 0$ then $v(g)$ is a scalar multiple of a unitary element in $M_{loc}(A)$. If we denote

$$\tilde{v}_{g_0} = v(g_0)\delta_{g_0} \in M_{loc}(A) \times_\alpha G.$$

then, since $v(g)\alpha_g(a) = av(g), g \in G$, we have $\tilde{v}_{g_0}\tilde{a} = \tilde{a}\tilde{v}_{g_0}$ and therefore $\tilde{v}_{g_0}^*\tilde{a} = \tilde{a}\tilde{v}_{g_0}^*$ for all $g_0 \in G, a \in A$. Then,

$$\tilde{v}_{g_0}\tilde{v}_{g_0}^*, \tilde{v}_{g_0}^*\tilde{v}_{g_0} \in A' \cap (M_{loc}(A) \times_\alpha G).$$

Standard calculations in $M_{loc}(A) \times_\alpha G$ show that $(\tilde{v}_{g_0}\tilde{v}_{g_0}^*) = v(g_0)^*v(g_0)\delta_e$, and $(\tilde{v}_{g_0}^*\tilde{v}_{g_0}) = \alpha_{g_0^{-1}}(v^*(g_0)v(g_0))\delta_e$ so,

$$\tilde{v}_{g_0}\tilde{v}_{g_0}^*, \tilde{v}_{g_0}^*\tilde{v}_{g_0} \in A' \cap M_{loc}(A).$$

Since A is prime, by Lemma 3 d) we have that $v(g_0)v(g_0)^* \in \mathbb{C}1$ and $v(g_0)^*v(g_0) \in \mathbb{C}1$. Therefore, there exists $\omega \in \mathbb{C}, \omega \neq 0$ such that $\omega v(g_0)$ is unitary. We can assume that $\omega = 1$. By relation (*) above we have

$$\alpha_{g_0}(a) = v^*(g_0)av(g_0), a \in A.$$

Hence α_{g_0} is inner in $M_{loc}(A)$, so $g_0 = e$. Therefore $v \in \mathbb{C}1$ in $M_{loc}(A) \times_\alpha G = M_{loc}(A \times_\alpha G)$ and we are done.

Now, suppose that α is strictly outer, i.e. $(A')' \cap M_{loc}(A \times_\alpha G) = \mathbb{C}I$. Let $g_0 \in G$ with $\alpha_{g_0}(a) = v_0av_0^*$ for a unitary element $v_0 \in M_{loc}(A)$. I will show that $g_0 = e$ where e is the unit of G . If we denote $\hat{v}_0 = v_0\delta_e \in M_{loc}(A) \times_\alpha G$ ($= M_{loc}(A \times_\alpha G)$ according to Lemma 10 b)) and $\tilde{b} = b\delta_e \in A \times_\alpha G \subset M_{loc}(A) \times_\alpha G$ for $b \in A$, then

$$\widetilde{\alpha_{g_0}(a)} = \hat{v}_0\tilde{a}\hat{v}_0^*, a \in A.$$

On the other hand, a standard calculation in $M_{loc}(A) \times_\alpha G$ gives

$$\widetilde{\alpha_{g_0}(a)} = \delta_{g_0}\tilde{a}\delta_{g_0^{-1}}.$$

Therefore $\delta_{g_0^{-1}} \widehat{v}_0 \in (A)' \cap M_{loc}(A \times_\alpha G) = \mathbb{C}1$, so $\widehat{v}_0 = \mu \delta_{g_0}$ for some scalar μ . Since the support of \widehat{v}_0 as a function on G is $\{e\}$ and the support of δ_{g_0} is $\{g_0\}$ it follows that $g_0 = e$. Hence every α_{g_0} with $g_0 \neq e$ is properly outer. ■

Remark 12 a) If $(A^\alpha)' \cap M_{loc}(A) = \mathbb{C}1$, then every automorphism $\alpha_t, t \neq e$ is properly outer and therefore, the action is strictly outer by the above theorem. Indeed if some α_{g_0} is implemented by a unitary $u \in M_{loc}(A)$ then, clearly $u \in (A^\alpha)' \cap M_{loc}(A)$, so $g_0 = e$.

b) Suppose A and $A \times_\alpha G$ are prime, G compact (in particular finite) and α faithful. If for every automorphism, β of A whose restriction to A^α is the identity, there exists $g \in G$ such that $\beta = \alpha_g$, then $(A^\alpha)' \cap M_{loc}(A) = \mathbb{C}1$. Indeed, if A and $A \times_\alpha G$ are prime and α is faithful, by [8, Corollary 3.12] and [9, Lemma 2.2.] the hypotheses of [9, Theorem 2.3.] are satisfied and the conclusion follows from that result.

In the next Section I will prove that, for finite abelian groups, the condition $(A^\alpha)' \cap M_{loc}(A) = \mathbb{C}1$ is equivalent with the action being properly outer, and therefore strictly outer.

3.2 Finite abelian groups

The proof of the following result is inspired by the proof of [5, Theorem 4] for simple C*-algebras (see also [7, Proposition 8.10.13.]). The hypothesis that A is simple is essential in their result.

Proposition 13 Let (A, G, α) be a C*-dynamical system with G finite abelian, α faithful and A prime. Then $(A^\alpha)' \cap M_{loc}(A) = \mathbb{C}1$ if and only if for every $t \neq 0$, α_t is a properly outer automorphism.

Proof. Suppose $(A^\alpha)' \cap M_{loc}(A) = \mathbb{C}1$. If α_{t_0} is implemented by a unitary u_{t_0} in $M_{loc}(A)$, then, clearly, $u_{t_0} \in (A^\alpha)' \cap M_{loc}(A) = \mathbb{C}1$. Conversely suppose that α_t is properly outer for every $t \in G, t \neq 0$. According to Theorem 9, the center of $M_{loc}(A)^\alpha$ is trivial.

Let $B = (A^\alpha)' \cap M_{loc}(A)$. Since the center of $M_{loc}(A)^\alpha$ is trivial, from Lemma 3 b) it follows that the fixed point algebra of the action α restricted to the α -invariant subalgebra B is trivial. Suppose that $B \neq \mathbb{C}1$. Then, there exists $\gamma_0 \in \widehat{G}$ such that $B_{\gamma_0} \neq \{0\}$. If $x, y \in B_{\gamma_0}$, then $x^*y \in \mathbb{C}1$ (the fixed point algebra of the restriction of α to B). Therefore there exists a unitary, $u \in B_{\gamma_0}$, such that $B_{\gamma_0} = \mathbb{C}u$. Since u is unitary, it follows that $n\gamma_0 \in Sp(\alpha|_B)$ for every $n \in \mathbb{Z}$, so $B_{n\gamma_0} = \mathbb{C}u^n$ for every $n \in \mathbb{Z}$.

Let $H = \{t \in G : \langle t, \gamma_0 \rangle = 1\}$. Then, since $\alpha_t(u) = \langle t, \gamma_0 \rangle u$ for every $t \in G$, it follows that $\alpha_t(u) = u$ for all $t \in H$. Notice that the annihilator, H^\perp , of H in \widehat{G} equals $\mathbb{Z}\gamma_0$. Now, consider the C*-subalgebra

$$M_{loc}(A)^H = \{x \in M_{loc}(A) : \alpha_t(x) = x \text{ for all } t \in H\}.$$

By [7, 8.10.3.], if $\tilde{\alpha}$ is the action of G/H on $M_{loc}(A)^H$, we have $Sp(\tilde{\alpha}) \subset \mathbb{Z}\gamma_0$. Therefore, the spectral subspaces $(M_{loc}(A)^H)_{n\gamma_0}$ have dense span in $M_{loc}(A)^H$. Next, we will prove that u commutes with $(M_{loc}(A)^H)_{n\gamma_0}$, $n \in \mathbb{Z}$ and thus u belongs to the center of $M_{loc}(A)^H$. Let $x \in (M_{loc}(A)^H)_{n\gamma_0} \subset (M_{loc}(A))_{n\gamma_0}$. Then

$$x = x(u^*)^n u^n \in (M_{loc}(A))^\alpha u^n.$$

On the other hand, since $u \in B = (A^\alpha)' \cap M_{loc}(A)$, in particular $u \in (A^\alpha)'$. Applying Lemma 3 b) again, it follows that u commutes with $M_{loc}(A)^\alpha$, so u commutes with x . Hence u belongs to the center of $M_{loc}(A)^H$. Thus the center of $M_{loc}(A)^H$ contains a non scalar element. Since, by hypothesis, A is prime, in particular H -prime, applying Theorem 9 for $G = H$, it follows that there exists $t_0 \in H$ such that α_{t_0} is implemented by a unitary in $M_{loc}(A)^H$, thus α_{t_0} is not properly outer, which is a contradiction. ■

As stated before, all the results obtained in this paper, hold for not necessarily separable C^* -algebras. The next theorem was considered in [2] for compact groups and separable C^* -algebras. In addition to separability, in the equivalence of a) and b) below the authors make the assumption that A^α is prime.

Theorem 14 *Let (A, G, α) be a C^* -dynamical system with A prime, G finite abelian and α faithful. The following conditions are equivalent*

- a) α_g is properly outer for every $g \in G, g \neq 0$.
- b) $(A^\alpha)' \cap M_{loc}(A) = \mathbb{C}I$.
- c) $(A)' \cap M_{loc}(A \times_\alpha G) = \mathbb{C}I$.
- d) A^α is prime and $\hat{\alpha}_\gamma$ is properly outer (in $M_{loc}(A \times_\alpha G)$) for every $\gamma \in \hat{G}, \gamma \neq 0$.

Proof. The equivalence of a) and b) follows from Proposition 13. The

equivalence of a) and c) follows from Theorem 11 in the particular case of abelian groups.

c) \Rightarrow d), Notice first that from c) it follows that the center of $M_{loc}(A \times_\alpha G)$ consists of scalars. Therefore, by Lemma 1, $A \times_\alpha G$ is prime, so, by [8, Corollary 3.13], A^α is prime. If for some $\gamma \in \hat{G}, \hat{\alpha}_\gamma$ is implemented by a unitary u in $M_{loc}(A \times_\alpha G)$, then, as $(A \times_\alpha G)^{\hat{\alpha}} = A$, it follows that $u \in (A)' \cap M_{loc}(A \times_\alpha G) = \mathbb{C}I$. Hence $\gamma = 0$.

d) \Rightarrow c) If A^α is prime and α is faithful, from property 5) in the list of properties of $Sp(\alpha)$ discussed after Definition 6, Section 2, it follows that $Sp(\alpha) = \hat{G}$. Applying [8 Cor 3.13.] it follows that $A \times_\alpha G$ is prime. Therefore $A \times_\alpha G$ is prime and $\hat{\alpha}_\gamma$ is properly outer (in $M_{loc}(A \times_\alpha G)$) for every $\gamma \in \hat{G}, \gamma \neq 0$. Since $(A \times_\alpha G)^{\hat{\alpha}} = A$, from Proposition 13 applied to the C^* -dynamical system $(A \times_\alpha G, \hat{G}, \hat{\alpha})$ (instead of (A, G, α)) it follows that $(A)' \cap M_{loc}(A \times_\alpha G) = \mathbb{C}I$. ■

References

- [1] P. Ara and M. Mathieu, *Local Multipliers of C^* -Algebras*, Springer, 2003.
- [2] O. Bratteli, G. A. Elliott, D. E. Evans and A. Kishimoto, *Quasi-Product Actions of a Compact Abelian Group on a C^* -Algebra*, Tohoku J. of Math., 41 (1989), 133-161.
- [3] R. Dumitru and C. Peligrad, *Compact Quantum Actions on C^* -Algebras and Invariant Derivations*, Proceedings of the Amer. Math. Soc., vol. 135, Nr.12 (2007), 3977-3984.
- [4] D. Olesen, *Inner $*$ -Automorphisms of Simple C^* -Algebras*, Commun. math. Phys. 44, (1975), 175—190.
- [5] D. Olesen, G. K. Pedersen and E. Stormer, *Compact Abelian Groups of Automorphisms of Simple C^* -Algebras*, Inventiones Math. 39 (1977), 55–64.
- [6] G. K. Pedersen, *Approximating Derivations on Ideals of C^* -Algebras*, Inventiones Math. 45 (1978), 299-305.
- [7] G. K. Pedersen, *C^* -Algebras and Their Automorphism Groups*, Second edition, Academic Press, 2018.
- [8] C. Peligrad, *Locally Compact Group Actions on C^* -Algebras and Compact Subgroups*, J. Functional Anal. 76, Nr. 1 (1988), 126-139.
- [9] C. Peligrad, *Duality for Compact Group Actions on Operator Algebras and Applications: Irreducible Inclusions and Galois Correspondence*, Internat. J. Math. 31, Nr. 9 (2020), 12 pp.
- [10] M. Rieffel, *Actions of Finite Groups on C^* -Algebras*, Math. Scand., 47 (1980), 157-176.
- [11] S. Vaes, *The Unitary Implementation of a Locally Compact Group Action*, J. Functional Anal. 180 (2001), 426-480.