

Asymptotic Quantum Many-Body Scars

Lorenzo Gotta,^{1,2,*} Sanjay Moudgalya,^{3,4,†} and Leonardo Mazza^{2,‡}

¹*Department of Quantum Matter Physics, University of Geneva,
24 Quai Ernest-Ansermet, 1211 Geneva, Switzerland*

²*Université Paris-Saclay, CNRS, LPTMS, 91405, Orsay, France*

³*Department of Physics and Institute for Quantum Information and Matter,
California Institute of Technology, Pasadena, California 91125, USA*

⁴*Walter Burke Institute for Theoretical Physics, California Institute of Technology, Pasadena, California 91125, USA*
(Dated: March 10, 2023)

We consider a quantum lattice spin model featuring exact quasiparticle towers of eigenstates with low entanglement at finite size, known as quantum many-body scars (QMBS). We show that the states in the neighboring part of the energy spectrum can be superposed to construct entire families of low-entanglement states whose energy variance decreases asymptotically to zero as the lattice size is increased. As a consequence, they have a relaxation time that diverges in the thermodynamic limit, and therefore exhibit the typical behavior of exact QMBS although they are not exact eigenstates of the Hamiltonian for any finite size. We refer to such states as *asymptotic* QMBS. These states are orthogonal to any exact QMBS at any finite size, and their existence shows that the presence of an exact QMBS leaves important signatures of non-thermalness in the rest of the spectrum; therefore, QMBS-like phenomena can hide in what is typically considered the thermal part of the spectrum. We support our study using numerical simulations in the spin-1 XY model, a paradigmatic model for QMBS, and we conclude by presenting a weak perturbation of the model that destroys the exact QMBS while keeping the asymptotic QMBS.

Introduction — Quantum Many-Body Scars (QMBS) [1–4] in non-integrable quantum lattice models of any dimension have recently emerged as one of the paradigms for the weak violation of the Eigenstate Thermalization Hypothesis (ETH) [5, 6], according to which all local properties of energy eigenstates in the middle of the spectra of non-integrable models coincide with those of a thermal Gibbs density matrix at a suitable temperature [7–10]. QMBS are isolated energy eigenstates that are outliers in many respects, e.g., in the expectation value of a local observable or in the entanglement entropy. Numerous instances of lattice models featuring exact QMBS at finite size have been discovered, including towers of QMBS in well-known models such as the spin-1 AKLT model [11–13], the spin-1 XY model [13–15], and the Hubbard model [16–23]. Most of these results have also been understood via unified frameworks or systematic construction recipes [3, 13, 17, 20, 24–30].

While most of the literature focuses on the exact QMBS states, a question that has been less explored is whether the presence of a finite-size QMBS affects the properties of the rest of the spectrum. An exception is Ref. [31], which pointed out the existence of low-entanglement states in the PXP model which exhibit slow relaxation even though they are orthogonal to the known exact QMBS [32]. In particular, they showed that the energy variance of such states is independent of system size and thus that their fidelity relaxation time does not decrease with system size. This is a remarkable phenomenology to be contrasted with that of typical product or short-range correlated states, whose energy variance grows with system size [33], whereas the fidelity relax-

ation time decreases [34].

A natural question that follows is whether there are even more drastic examples of slowly relaxing states, for instance with an energy variance decreasing with system size, which would lead to a relaxation time that *diverges* in the thermodynamic limit (TL). Two settings are typically possible. First, slow relaxation of hydrodynamic origin is ubiquitous in systems with continuous symmetries, where it is said to occur at a diverging timescale known as the *Thouless time* [35–38], and is related to phenomena such as diffusion and subdiffusion [39–44]. The interpretation of QMBS as an unconventional non-local symmetry [29] motivates the search for such slow relaxation. A different type of slow relaxation usually occurs due to the presence of long-lived quasiparticles in the system, e.g. the phonons of a superfluid with Beliaev decay [45]. QMBS are associated to quasiparticles with specific momenta and infinite lifetime [4], hence it is natural to look for long-lived quasiparticles at neighboring momenta.

In this article we address these questions by considering the spin-1 XY model featuring exact QMBS at any finite size [14] and show that it is possible to construct slowly-relaxing low-entanglement initial states that exhibit QMBS-like features, but nevertheless are orthogonal to the exact QMBS. They have an energy variance that goes to zero in the TL and asymptotically display the typical dynamical phenomenology of a QMBS, i.e. the lack of thermalization; hence we refer to such initial states as *asymptotic* QMBS. Our work widens the range of initial states that qualitatively exhibit a non-thermalizing phenomenology and motivates the search for non-thermal features also in regions of the spectrum

where entanglement signatures do not make them evident.

The model and the exact QMBS — We consider a one-dimensional spin-1 chain of length L , which we take to be even for simplicity, and consider a spin-1 XY model with external magnetic field and axial anisotropy:

$$H = J \sum_j (S_j^x S_{j+1}^x + S_j^y S_{j+1}^y) + h \sum_j S_j^z + D \sum_j (S_j^z)^2 + J_3 \sum_j (S_j^x S_{j+3}^x + S_j^y S_{j+3}^y), \quad (1)$$

where S_j^α , with $\alpha = x, y, z$, are the spin-1 operators on site j . We use open boundary conditions (OBC) for the numerical simulations and periodic boundary conditions (PBC) for some of the analytical results. This model with OBC has been numerically shown to be non-integrable; nonetheless, we have added the last term to break a hidden non-local symmetry [46] without affecting the QMBS physics [4, 14].

The Hamiltonian in Eq. (1) exhibits QMBS for any finite value of L , as it was first identified in [14]. In order to see that, we define the fully-polarised state $|\Downarrow\rangle = |-\dots -\rangle$ with all spins in the eigenstate of S_j^z with eigenvalue -1 , and the operator

$$J_k^+ = \frac{1}{2} \sum_{j=1}^L e^{ikj} (S_j^+)^2, \quad (2)$$

The scar states read:

$$|n, \pi\rangle = \frac{1}{\sqrt{N_{n,\pi}}} (J_\pi^+)^n |\Downarrow\rangle, \quad (3)$$

where $N_{n,\pi}$ is a normalisation constant. The state satisfies the stationary Schrödinger eigenvalue equation $H|n, \pi\rangle = (-Lh + 2nh + LD)|n, \pi\rangle$ and for generic values of h and D it lies in the middle of the spectrum of the Hamiltonian.

Moreover, it is possible to consider the reduced density matrix $\rho_{A,n,\pi}$ of $|n, \pi\rangle$ defined on half the system (conventionally, the region A is $1 \leq j < L/2$), and to compute its entanglement entropy, $S_{n,\pi} = -\text{tr}[\rho_{A,n,\pi} \log \rho_{A,n,\pi}]$. The explicit calculation has been done in Ref. [14], and it shows that it scales as $\log L$, displaying a mild logarithmic violation of an entanglement area law, see Supplementary Materials (SM) [47] for details. QMBS are easily found numerically by plotting the entanglement entropy S_{E_i} of ρ_{A,E_i} , the reduced density matrix of the eigenstate $|E_i\rangle$, as a function of energy. Indeed, almost all the eigenstates appear to satisfy the ETH and are characterised by an S_{E_i} that is only a function of the energy E_i ; they have a higher amount of entanglement than the QMBS states, which indeed violate ETH.

A family of states obtained by deforming the exact QMBS — We now consider other initial states

for the dynamics of the model in Eq. (1); they read as follows:

$$|n, k\rangle = \frac{1}{\sqrt{N_{n,k}}} J_k^+ (J_\pi^+)^{n-1} |\Downarrow\rangle, \quad (4)$$

where $N_{n,k}$ is a normalisation constant, which coincide with the exact QMBS in Eq. (3). Remarkably, when $k \neq \pi$ and is an integer multiple of $\frac{2\pi}{L}$, they are *orthogonal* to the exact QMBS: the relation $\langle n, k | n', \pi \rangle = \delta_{n,n'} \delta_{k,\pi}$ for any $1 \leq n, n' \leq L-1$ is proved in the SM [47]. Models where such classes of multimagnon states are exact eigenstates have been studied in [48], however for $k \neq \pi$ these are *not* eigenstates of the spin-1 XY model. It is easy to show that the average energy of these states does not depend on k and reads $\langle n, k | H | n, k \rangle = -Lh + 2nh + LD$ [47].

Furthermore, the entanglement of the states in Eq. (4) scales with system size as a sub-volume law. For a quick proof, since $|n, k\rangle \propto J_k^+ |n-1, \pi\rangle$, we note that J_k^+ can be straightforwardly expressed as a Matrix Product Operator (MPO) of bond dimension $\chi = 2$ [3, 49, 50], hence the half-subsystem entanglement entropies of $|n-1, \pi\rangle$ and $|n, k\rangle$ can differ at most of an additive term $\log 2$. In other words, since the operator J_k^+ can be split in two terms, one acting on $j < L/2$ and one on $j \geq L/2$, it is possible to show (see the SM [47] for details) that the total number of Schmidt states in $|n, k\rangle$ is at most twice than that in $|n-1, \pi\rangle$. Since we know that $|n-1, \pi\rangle$ has an entanglement that scales with the logarithm of the system size [3, 12, 14, 51], we are led to conclude that $|n, k\rangle$ has the same scaling for extensive n .

To further characterise the states in Eq. (4), we compute the variance of the energy ΔH^2 under the Hamiltonian H in PBC, and as we show in the SM [47], we obtain:

$$\Delta H^2 = 4 \left[J^2 \cos^2 \left(\frac{k}{2} \right) + J_3^2 \cos^2 \left(\frac{3k}{2} \right) \right]. \quad (5)$$

Note that among the states defined in Eq. (4), the $|n, \pi\rangle$ are the only eigenstates of the Hamiltonian, because $\Delta H^2 = 0$ only for $k = \pi$. When $k \neq \pi$, $|n, k\rangle$ must be a linear superposition of the energy eigenstates of H , which are mostly in a window centered around the same energy of $|n, \pi\rangle$ and in a width of about ΔH . Remarkably, when $k \neq \pi$ is chosen to be an integer multiple of $\frac{2\pi}{L}$, $|n, \pi\rangle$ is not part of this set of states due to orthogonality. Since $|n, \pi\rangle$ numerically appear to be the only exact QMBS of H [14], we conclude that such states $|n, k\rangle$ must be a linear superposition of “thermal” eigenstates, i.e., those that are typically said to satisfy ETH, having an entanglement entropy and expectation values of local observables that are smooth functions of energy.

We have numerically verified this statement using the python-based package QuSpin [52]: we diagonalize the Hamiltonian (1) and compute the bipartition entanglement entropy S_{E_i} and the average square magnetisation




FIG. 1. Top: Squared overlap of $|n, k\rangle$ for $n = L/2$ and $k = \pi - \frac{2\pi}{L}$ with the eigenstates $|E_i\rangle$ of Hamiltonian (1) with zero magnetisation, $S_z = 0$; the parameters of the simulation are $\{J, \hbar, D, J_3\} = \{1, 0, 0.1, 0.1\}$ and $L = 10$. The information on $|\langle E_i | n, k \rangle|^2$ is also encoded in the color code of the marker of all panels using a logarithmic scale, see colorbar. Middle and bottom: We plot the data of the top panel in a diagram with the energy E on the abscissa and the bipartition entanglement entropy S_E or the average square magnetisation $S_E^2(E)$ of the eigenstate on the ordinate, respectively. The state $|n, k\rangle$ has overlap only with states whose S_{E_i} or $S_E^2(E_i)$ lies on the continuous “thermal” curve. The red circle and the blue square highlight the regions of the plots where the QMBS $|n = L/2, \pi\rangle$ appear: the absence of any grey mark means that the scalar product is compatible with the numerical zero.

$S_E^2(E_i) = \frac{1}{L} \sum_j \langle (S_j^z)^2 \rangle$ of all eigenstates. Subsequently, we compute the scalar product of the state $|n, k\rangle$ with all eigenstates for $n = L/2$ and $k = \pi - \frac{2\pi}{L}$ and look at the properties of the eigenstates with whom the overlap is not zero. The results are reported in Fig. 1, and support our thesis.

Dynamics and asymptotic QMBS — The dynamical properties of the states $|n, k\rangle$ for large system sizes depend on how we approach $L \rightarrow \infty$. If the limit is taken while the momentum k is held fixed, then the variance is finite in the TL. Loosely speaking, we can invoke the well-known energy-time uncertainty relation, linking the typical timescale of the dynamics τ of a quantum state to the standard deviation of the energy:

$$\tau \geq \frac{\hbar}{2\Delta H} \quad (6)$$

to claim that for these states the dynamics is frozen up to a given time-scale τ that is independent of L . That is, we expect that for $t \lesssim \tau$, the state behaves essentially as an eigenstate; for $t \gtrsim \tau$, the dynamics sets in, and evolution towards thermal equilibration values is expected (see SM [47]). We should caution the reader that the loss of fidelity does not by itself imply relaxation of expectation




FIG. 2. The properties of the state $e^{-iHt} |n, k\rangle$ for $n = L/2$ and $k = \pi - 2\pi/L$ as a function of time for various system sizes L . Left: time evolution of the squared magnetisation $S^2(t)$. Right: time evolution of the fidelity with the initial state $F(t)$.

values of local observables, and the latter timescale can in principle be much longer. Nevertheless, we should think of the fidelity relaxation time, determined purely by the variance of the initial state [53], to be a lower bound for the relaxation time of local observables [10, 54]. Note that an initial state with similar finite-variance properties has been discussed in Ref. [31] in the context of exact QMBS in the PXP model [32], and the authors reach similar conclusions.

Another class of states can be obtained by approaching the TL while letting k flow to π . This can be done by setting $k = \pi + \frac{2\pi}{L}m$, with the coefficient $m \in \mathbb{Z}$ kept constant while $L \rightarrow \infty$. In this case the energy variance scales as $\Delta H^2 \sim (J^2 + 9J_3^2)(k - \pi)^2$ and tends to zero as $1/L^2$. We refer to this second class of states as *asymptotic QMBS* of the model, since according to (6), the typical relaxation timescale of their dynamics scales as $\tau \sim L$, i.e., the system is frozen for timescales that *increase* polynomially with the system size. On the contrary, low entanglement states, by virtue of their diverging variance [33], are typically expected to lose fidelity on timescales that decrease with system size, and the expectation values of typical observables relax in timescales that do not change drastically with system size [10, 55–61]. Hence the dynamics of this class of states asymptotically approaches QMBS-like behavior even though they are *not* exact QMBS of the system at finite size, and moreover they are *orthogonal* to all the exact QMBS $|n, \pi\rangle$. To the best of our knowledge, this phenomenology has never been discussed before.

We support the previous statements with a numerical simulation of the dynamics of the states $|n, k\rangle$ under the action of H using a time-evolving block decomposition (TEBD) code based on a Matrix-Product-State (MPS) representation of the state obtained via the ITen-

sor library [62, 63]. We consider in particular the state $|n = L/2, k = \pi - 2\pi/L\rangle$ for several system sizes up to $L = 60$ and truncation error 10^{-12} . We then compute the observable $S^{z2}(t) = \frac{1}{L} \sum_j \langle (S_j^z)^2 \rangle_t$ and the fidelity of the time-evolved state with the initial state $F(t) = |\langle n, k | e^{-iHt} | n, k \rangle|^2$. The results, reported in Fig. 2, show in both cases an important slow-down of the dynamics as the size increases. In the SM we show that the data concerning the fidelity can be collapsed via a rescaling of time by a factor of L [47], which suggests the divergence of the relaxation time in the TL. The result on the fidelity $F(t)$ is particularly important as it shows undoubtedly that the time-evolved state maintains an overlap with the initial state that increases with L and it implies the freezing of the state. In the SM we complement this analysis by contrasting it with the typical dynamics of other states [47].

Slow relaxation and non-thermalness in the middle of the energy spectrum — At this point, it is important to highlight the two properties that make the asymptotic QMBS particularly interesting: (a) they have a limited amount of entanglement, i.e., a sub-volume law, but an extensive amount of energy; (b) they have an energy variance ΔH^2 that drops fast enough to zero in the TL. Any state that satisfies these conditions is guaranteed to have a long relaxation time, both in the fidelity and in the observables, while having an average energy that lies in the middle of the Hamiltonian spectrum. Note that both (a) and (b) are necessary features that make the behavior of asymptotic QMBS atypical. While any linear superposition of thermal eigenstates with small energy variance relaxes slowly, one typically cannot construct a low-entanglement state if its energy variance decreases fast enough with system size [64]. On the other hand, a typical low-entanglement state has an energy variance that increases with system size [33], which usually implies a short relaxation time.

It is tempting to think that the existence of asymptotic QMBS should imply some kind of “non-thermalness” [31] or ETH-violation in the “thermal” states orthogonal to the exact QMBS, even at finite system size. Note that ETH consists of two parts [6, 9, 65], pertaining to diagonal and off-diagonal matrix elements of a local operator in the energy eigenbasis. The diagonal matrix elements control the late-time expectation values of observables, and we do not expect the violation of diagonal ETH since the asymptotic QMBS eventually thermalize. On the other hand, the timescale of relaxation is controlled by both the energy variance of the initial state and the off-diagonal matrix elements, although it is challenging to make a precise connection [60]. Nevertheless, given that asymptotic QMBS cannot be supported by any ETH-satisfying generic local Hamiltonian [33], it is plausible that our result entails a violation of off-diagonal ETH at least in a part of the Hamiltonian spectrum. In any case, our work shows that the exact tower of QMBS leaves some marks




FIG. 3. Properties of the eigenstates of Hamiltonian H' in the zero magnetization sector $S_z = 0$; the parameters of the simulation are $\{J, h, D, J_3, J_z\} = \{1, 0, 0.1, 0.1, 1\}$ and $L = 10$. Top and bottom: We plot the bipartition entanglement entropy S_{E_i} and the average square magnetisation $S^{z2}(E_i)$ as a function of the energy of the eigenstate E_i . All eigenstates collapse on a continuous curve; no exact QMBS outlier is observed.

in the neighboring part of the energy spectrum.

Asymptotic QMBS without exact QMBS —

Our definition of asymptotic QMBS is based on a deformation of the tower of exact QMBS supported at finite size; it seems likely that our construction can be adapted also to other models with that property [2–4]. However, it is not clear whether asymptotic QMBS can exist in models without any exact QMBS or if they can exist at energies distant from those where the exact QMBS.

Nevertheless, we now show that it is possible to weakly perturb the Hamiltonian H in a way that destroys all exact QMBS, but such that the exact QMBS become asymptotic QMBS of the perturbed model. As an example, we consider $H' = H + V$ with $V = (J_z/L) \sum_j S_j^z S_{j+1}^z$, which is still a non-trivial local perturbation since its spectral norm $\|V\|_\infty$ corresponding to its largest singular value is subextensive and scales as $O(1)$. Using the python-based QuSpin package [52], we numerically diagonalize H' and compute the entanglement entropy S_{E_i} and the average square magnetisation $S^{z2}(E_i)$ for all eigenstates. The plots, shown in Fig. 3, do not indicate the presence of any exact QMBS.

We now consider the state $|n, \pi\rangle$ of Eq. (3), which is an exact QMBS of H but *not* an eigenstate of H' . Using the ITensor library [62, 63], we compute $S^{z2}(t)$ and the fidelity $F(t)$ for the time-evolved state $|\Psi(t)\rangle = e^{-iH't} |n, \pi\rangle$; the results are in Fig. 4. The plots display the phenomenology of an asymptotic QMBS in a Hamiltonian that does not show any exact QMBS at finite size, and the $F(t)$ curves exhibit a collapse when time is rescaled by a factor $\sqrt{L}$ [47], indicating a diverging re-




FIG. 4. The properties of the state $e^{-iH't} |n, \pi\rangle$ for $n = L/2$ as a function of time; left: time evolution of the squared magnetisation $S^{z2}(t)$; right: time evolution of the fidelity with the initial state $F(t)$. The inset shows the scaling as a function of size of the values of $F(t = 3/J)$; we find a scaling to 1 as $1/L$.

laxation time. This behavior can be directly attributed to the fact that the variance of the state $|n, \pi\rangle$ under the Hamiltonian H' scales as $\sim 1/L$ when n is a finite fraction of L , as it is proven in the SM [47][66].

Conclusions — In this letter we have revisited the paradigmatic one-dimensional spin-1 XY model that supports exact QMBS at finite size, and we explored the properties of the rest of the spectrum. We showed that it is possible to construct other states, dubbed asymptotic QMBS, with little entanglement and whose relaxation time diverges polynomially in the thermodynamic limit. From the late-time hydrodynamics point of view, these asymptotic QMBS indicate the existence of slowly relaxing modes in systems with exact QMBS. On the other hand, they indicate the existence of long-lived quasiparticle bands on top of the QMBS deep in the spectrum.

Remarkably, asymptotic QMBS are linear combinations of “thermal” eigenstates whose entanglement entropy and average squared magnetization are a “smooth” function of energy; therefore, their existence hints at a possible violation of ETH in its conventional form, and in particular of the off-diagonal ETH. We leave for future work the direct and careful study of the eigenstates to identify such violation using standard methods developed for such purposes [67–73].

Asymptotic QMBS with similar properties can also be constructed in higher dimensional spin-1 XY models [47], but extending our results to even more general settings would be interesting. First, the asymptotic QMBS might be a generic feature of the exhaustive algebra of local Hamiltonians that have the same exact QMBS $|n, \pi\rangle$ [29]. Second, they likely can always be constructed in Hamiltonians with simple quasiparticle towers of exact QMBS [11–13, 17–19, 25, 74]. Third, there are many different types of exact QMBS [3], e.g., the “quasi-

particles” could be non-local [13, 27, 75], or the states could be isolated [24, 32], or they could appear in gauge theories [76, 77] or Floquet systems [28, 78–80]: are there asymptotic QMBS in these models? Finally, one could also consider deformations of Hamiltonians with exact QMBS (a problem that we partially addressed in the final part of this letter for weak perturbations), and ask the following intriguing question: what are the conditions for a Hamiltonian to display an asymptotic QMBS without any exact QMBS?

Acknowledgements — We warmly acknowledge enlightening discussions with Saverio Bocini, Xiangyu Cao, Maurizio Fagotti, David Huse, Michael Knap, Lesik Motrunich and Nicolas Regnault. We also thank Lesik Motrunich for useful comments on a draft. L.G. and L.M. also thank Guillaume Roux and Pascal Simon for discussions on previous shared projects. This work is supported by the Walter Burke Institute for Theoretical Physics at Caltech and the Institute for Quantum Information and Matter, by LabEx PALM (ANR-10-LABX-0039-PALM) in Orsay, by Region Ile-de-France in the framework of the DIM Sirteq and by the Swiss National Science Foundation under Division II. S.M. also acknowledges the hospitality of the Laboratoire de Physique Théorique et Modèles Statistiques (LPTMS) in Orsay, where this collaboration was initiated, and the Physik-Institut of the University of Zurich, where some of this work was performed.

* lorenzo.gotta@unige.ch

† sanjaym@caltech.edu

‡ leonardo.mazza@universite-paris-saclay.fr

- [1] M. Serbyn, D. A. Abanin, and Z. Papić, *Nature Physics* **17**, 675 (2021).
- [2] Z. Papić, (2021), arXiv:2108.03460 [cond-mat.quant-gas].
- [3] S. Moudgalya, B. A. Bernevig, and N. Regnault, *Reports on Progress in Physics* **85**, 086501 (2022).
- [4] A. Chandran, T. Iadecola, V. Khemani, and R. Moessner, arXiv e-prints (2022), arXiv:2206.11528 [cond-mat.str-el].
- [5] J. M. Deutsch, *Physical Review A* **43**, 2046 (1991).
- [6] M. Srednicki, *Physical Review E* **50**, 888 (1994).
- [7] M. Rigol, V. Dunjko, and M. Olshanii, *Nature* **452**, 854 (2008).
- [8] A. Polkovnikov, K. Sengupta, A. Silva, and M. Vengalattore, *Reviews of Modern Physics* **83**, 863 (2011).
- [9] L. D’Alessio, Y. Kafri, A. Polkovnikov, and M. Rigol, *Advances in Physics* **65**, 239 (2016).
- [10] T. Mori, T. N. Ikeda, E. Kaminishi, and M. Ueda, *Journal of Physics B: Atomic, Molecular and Optical Physics* **51**, 112001 (2018).
- [11] S. Moudgalya, S. Rachel, B. A. Bernevig, and N. Regnault, *Physical Review B* **98**, 235155 (2018).
- [12] S. Moudgalya, N. Regnault, and B. A. Bernevig, *Phys. Rev. B* **98**, 235156 (2018).

- [13] D. K. Mark, C.-J. Lin, and O. I. Motrunich, Phys. Rev. B **101**, 195131 (2020).
- [14] M. Schecter and T. Iadecola, Phys. Rev. Lett. **123**, 147201 (2019).
- [15] J. Wildeboer, C. M. Langlett, Z.-C. Yang, A. V. Gorshkov, T. Iadecola, and S. Xu, Phys. Rev. B **106**, 205142 (2022).
- [16] C. N. Yang, Physical Review letters **63**, 2144 (1989).
- [17] S. Moudgalya, N. Regnault, and B. A. Bernevig, Phys. Rev. B **102**, 085140 (2020).
- [18] D. K. Mark and O. I. Motrunich, Phys. Rev. B **102**, 075132 (2020).
- [19] K. Pakrouski, P. N. Pallegar, F. K. Popov, and I. R. Klebanov, Phys. Rev. Lett. **125**, 230602 (2020).
- [20] K. Pakrouski, P. N. Pallegar, F. K. Popov, and I. R. Klebanov, Phys. Rev. Research **3**, 043156 (2021).
- [21] H. Yoshida and H. Katsura, Phys. Rev. B **105**, 024520 (2022).
- [22] L. Gotta, L. Mazza, P. Simon, and G. Roux, Phys. Rev. B **106**, 235147 (2022).
- [23] M. Nakagawa, H. Katsura, and M. Ueda, arXiv e-prints, arXiv:2205.07235 (2022), arXiv:2205.07235 [cond-mat.str-el].
- [24] N. Shiraishi and T. Mori, Phys. Rev. Lett. **119**, 030601 (2017).
- [25] S. Moudgalya, E. O'Brien, B. A. Bernevig, P. Fendley, and N. Regnault, Phys. Rev. B **102**, 085120 (2020).
- [26] J. Ren, C. Liang, and C. Fang, Phys. Rev. Lett. **126**, 120604 (2021).
- [27] N. O'Dea, F. Burnell, A. Chandran, and V. Khemani, Phys. Rev. Research **2**, 043305 (2020).
- [28] P.-G. Rozon, M. J. Gullans, and K. Agarwal, Phys. Rev. B **106**, 184304 (2022).
- [29] S. Moudgalya and O. I. Motrunich, arXiv e-prints (2022), arXiv:2209.03377 [cond-mat.str-el].
- [30] P.-G. Rozon and K. Agarwal, arXiv e-prints, arXiv:2302.04885 (2023).
- [31] C.-J. Lin, A. Chandran, and O. I. Motrunich, Phys. Rev. Res. **2**, 033044 (2020).
- [32] C.-J. Lin and O. I. Motrunich, Phys. Rev. Lett. **122**, 173401 (2019).
- [33] M. C. Bañuls, D. A. Huse, and J. I. Cirac, Phys. Rev. B **101**, 144305 (2020).
- [34] Note that the precise scaling of the relaxation time depends on the definition. The fidelity of an initial state at short times decays as $F(t) \sim \exp(-\Delta H^2 t^2)$ [53], where ΔH^2 is the variance, this gives a timescale $\tau \sim 1/\Delta H$. On the other hand, one can define the fidelity relaxation time as the timescale at which fidelity decays to the the typical fidelity between two many body states, which scales as $\exp(-L)$ [53], this adds an extra factor of $\sqrt{L}$. In this work, we use the former definition, and are mostly interesting in the relative decay timescales between initial states of different variances.
- [35] D. J. Thouless, Phys. Rev. Lett. **39**, 1167 (1977).
- [36] A. Chan, A. De Luca, and J. T. Chalker, Phys. Rev. Lett. **121**, 060601 (2018).
- [37] M. Schiulaz, E. J. Torres-Herrera, and L. F. Santos, Phys. Rev. B **99**, 174313 (2019).
- [38] A. Dymarsky, Phys. Rev. Lett. **128**, 190601 (2022).
- [39] P. M. Chaikin, T. C. Lubensky, and T. A. Witten, Principles of condensed matter physics, Vol. 10 (Cambridge university press Cambridge, 1995).
- [40] S. Mukerjee, V. Oganesyan, and D. Huse, Phys. Rev. B **73**, 035113 (2006).
- [41] J. Lux, J. Müller, A. Mitra, and A. Rosch, Phys. Rev. A **89**, 053608 (2014).
- [42] A. Gromov, A. Lucas, and R. M. Nandkishore, Phys. Rev. Res. **2**, 033124 (2020).
- [43] J. Feldmeier, P. Sala, G. De Tomasi, F. Pollmann, and M. Knap, Phys. Rev. Lett. **125**, 245303 (2020).
- [44] S. Moudgalya, A. Prem, D. A. Huse, and A. Chan, Phys. Rev. Res. **3**, 023176 (2021).
- [45] L. Pitaevskii and S. Stringari, Bose-Einstein Condensation (Oxford University Press, 2016).
- [46] A. Kitazawa, K. Hijii, and K. Nomura, Journal of Physics A: Mathematical and General **36**, L351 (2003).
- [47] "See the supplemental material,".
- [48] L.-H. Tang, N. O'Dea, and A. Chandran, Phys. Rev. Res. **4**, 043006 (2022).
- [49] G. M. Crosswhite and D. Bacon, Phys. Rev. A **78**, 012356 (2008).
- [50] J. Motruk, M. P. Zaletel, R. S. K. Mong, and F. Pollmann, Phys. Rev. B **93**, 155139 (2016).
- [51] O. Vafek, N. Regnault, and B. A. Bernevig, SciPost Phys. **3**, 043 (2017).
- [52] P. Weinberg and M. Bukov, SciPost Phys. **2**, 003 (2017).
- [53] L. Campos Venuti and P. Zanardi, Phys. Rev. A **81**, 022113 (2010).
- [54] T. Mori and N. Shiraishi, Phys. Rev. E **96**, 022153 (2017).
- [55] S. Goldstein, T. Hara, and H. Tasaki, Phys. Rev. Lett. **111**, 140401 (2013).
- [56] A. S. L. Malabarba, L. P. García-Pintos, N. Linden, T. C. Farrelly, and A. J. Short, Phys. Rev. E **90**, 012121 (2014).
- [57] S. Goldstein, T. Hara, and H. Tasaki, New Journal of Physics **17**, 045002 (2015).
- [58] P. Reimann, Nature Communications **7**, 10821 (2016).
- [59] L. P. García-Pintos, N. Linden, A. S. L. Malabarba, A. J. Short, and A. Winter, Phys. Rev. X **7**, 031027 (2017).
- [60] H. Wilming, T. R. de Oliveira, A. J. Short, and J. Eisert, "Equilibration times in closed quantum many-body systems," in Thermodynamics in the Quantum Regime: Fundamental Aspects and edited by F. Binder, L. A. Correa, C. Gogolin, J. Anders, and G. Adesso (Springer International Publishing, Cham, 2018) pp. 435–455.
- [61] J. Riddell, L. P. García-Pintos, and Á. M. Alhambra, arXiv e-prints, arXiv:2112.09475 (2021).
- [62] M. Fishman, S. R. White, and E. M. Stoudenmire, SciPost Phys. Codebases, **4** (2022).
- [63] M. Fishman, S. R. White, and E. M. Stoudenmire, SciPost Phys. Codebases, **4** (2022).
- [64] The results of [33] show that Matrix Product States (MPS) with polynomially large bond dimension and variance that decreases logarithmically with system size can always be constructed.
- [65] N. Shiraishi and T. Mori, Phys. Rev. Lett. **121**, 038902 (2018).
- [66] Note that even this timescale matches the rigorous lower bounds on relaxation times for weak perturbations of models with exact QMBS [31] by setting the perturbation strength $\lambda = 1/L$, the latter is the observable relaxation time, which we generically expect to be different from the fidelity relaxation time we discuss.

- [67] E. Khatami, G. Pupillo, M. Srednicki, and M. Rigol, Phys. Rev. Lett. **111**, 050403 (2013).
- [68] R. Steinigeweg, J. Herbrych, and P. Prelovšek, Phys. Rev. E **87**, 012118 (2013).
- [69] W. Beugeling, R. Moessner, and M. Haque, Phys. Rev. E **91**, 012144 (2015).
- [70] J. Richter, A. Dymarsky, R. Steinigeweg, and J. Gemmer, Phys. Rev. E **102**, 042127 (2020).
- [71] J. Wurtz and A. Polkovnikov, Phys. Rev. B **101**, 195138 (2020).
- [72] S. Sugiura, P. W. Claeys, A. Dymarsky, and A. Polkovnikov, Phys. Rev. Res. **3**, 013102 (2021).
- [73] F. M. Surace, M. Votto, E. G. Lazo, A. Silva, M. Dalmonte, and G. Giudici, Phys. Rev. B **103**, 104302 (2021).
- [74] T. Iadecola and M. Schecter, Phys. Rev. B **101**, 024306 (2020).
- [75] C. M. Langlett, Z.-C. Yang, J. Wildeboer, A. V. Gorshkov, T. Iadecola, and S. Xu, Phys. Rev. B **105**, L060301 (2022).
- [76] S. Biswas, D. Banerjee, and A. Sen, SciPost Phys. **12**, 148 (2022).
- [77] D. Banerjee and A. Sen, Phys. Rev. Lett. **126**, 220601 (2021).
- [78] K. Mizuta, K. Takasan, and N. Kawakami, Phys. Rev. Research **2**, 033284 (2020).
- [79] S. Sugiura, T. Kuwahara, and K. Saito, Phys. Rev. Research **3**, L012010 (2021).
- [80] T. Iadecola and S. Vijay, Phys. Rev. B **102**, 180302 (2020).

Online supplementary material for: Asymptotic Quantum Many-Body Scars

Lorenzo Gotta^{1,2}, Sanjay Moudgalya^{3,4}, Leonardo Mazza²

¹ *Department of Quantum Matter Physics, University of Geneva, 24 Quai Ernest-Ansermet, 1211 Geneva, Switzerland*

² *Université Paris-Saclay, CNRS, LPTMS, 91405, Orsay, France*

³ *Department of Physics and Institute for Quantum Information and Matter,
California Institute of Technology, Pasadena, California 91125, USA*

⁴ *Walter Burke Institute for Theoretical Physics, California Institute of Technology, Pasadena, California 91125, USA*

March 10, 2023

In this Supplementary Material we present the explicit calculations of the main relevant properties of the asymptotic QMBS presented in the main text:

- S1. Orthogonality of the asymptotic QMBS with the exact QMBS
- S2. Average energy and energy variance for the asymptotic QMBS
- S3. Entanglement entropy of the exact and asymptotic QMBS
- S4. Variance of the exact QMBS for the perturbed Hamiltonian
- S5. Dynamics of initial states that are not asymptotic QMBS
- S6. Spectral properties of the Hamiltonian H'
- S7. Universal rescaling of fidelities
- S8. Higher dimensional generalisations of asymptotic QMBS

S1. ORTHOGONALITY OF THE ASYMPTOTIC QMBS WITH THE EXACT QMBS

In this section, we demonstrate the orthogonality of the states $\{|n, k\rangle\}$, defined in Eq. (4). First, we note that $|n, k\rangle$ is orthogonal to $|n', k'\rangle$ when $n \neq n'$ because they have a different magnetisation $S^z = \sum_j S_j^z$, which is a simple function of n : $S^z = -L + 2n$. We now consider states with the same n and take the system size L to be even and k to be an integer multiple of $\frac{2\pi}{L}$ for simplicity. We then observe that $\langle n, k | n, k' \rangle \propto \langle n-1, \pi | J_k^- J_{k'}^+ | n-1, \pi \rangle$ for $n \geq 1$. By definition of the operators $J_k^\pm$ in Eq. (2) we have:

$$\begin{aligned}
 \langle n-1, \pi | J_k^- J_{k'}^+ | n-1, \pi \rangle &= \frac{1}{4} \sum_{j, j'=1}^L e^{-i(kj-k'j')} \langle n-1, \pi | (S_j^-)^2 (S_{j'}^+)^2 | n-1, \pi \rangle \\
 &= \frac{1}{4} \sum_{j=1}^L e^{-i(k-k')j} \langle n-1, \pi | (S_j^-)^2 (S_j^+)^2 | n-1, \pi \rangle + \frac{1}{4} \sum_{j \neq j'} e^{-i(kj-k'j')} \langle n-1, \pi | (S_j^-)^2 (S_{j'}^+)^2 | n-1, \pi \rangle \\
 &= \frac{1}{4} \alpha \sum_{j=1}^L e^{-i(k-k')j} + \frac{1}{4} \beta \sum_{j=1}^L e^{i(\pi-k)j} \sum_{j' \neq j} e^{i(k'-\pi)j'} = \frac{1}{4} L(\alpha - \beta) \delta_{k, k'} + \frac{1}{4} \beta L^2 \delta_{k, \pi} \delta_{k', \pi},
 \end{aligned} \tag{S7}$$

where $\alpha = 4 \frac{\binom{L-1}{n-1}}{\binom{L}{n}} = 4 \frac{L+1-n}{L}$ and $\beta = 4 \frac{\binom{L-2}{n-2}}{\binom{L}{n}} = 4 \frac{(n-1)(L-n+1)}{L(L-1)}$, and we have used the fact that k and k' are integer multiples of $\frac{2\pi}{L}$. This calculation is done directly by using the expression of $|n, \pi\rangle$ as an equal amplitude superposition of “fully-magnetised” product states

$$|n, \pi\rangle = \sqrt{\frac{1}{2^{2n} \binom{L}{n}}} \sum_{1 \leq j_1 < j_2 < \dots < j_n \leq L} e^{i\pi \sum_{i=1}^n j_i} (S_{j_1}^+)^2 (S_{j_2}^+)^2 \dots (S_{j_n}^+)^2 |\Downarrow\rangle, \tag{S8}$$

and studying the action of the sandwiched operator on the basis states separately when $j = j'$ and when $j \neq j'$, and carefully accounting for the phase factors and normalization factors. It is important to visualise the combinatorial

nature of this state, expanded on a basis of states where the bimagnons created by $(S_j^+)^2$ are equally distributed everywhere. When $j = j'$, we obtain that α in Eq. (S7) is simply related to the number of fully-magnetised product states that do not have a bimagnon at site j , or else the action of $(S_j^-)^2(S_j^+)^2$ vanishes on such a basis state. This number is $\binom{L-1}{n-1}$; if we consider the normalisation factor and the specific matrix elements of $(S_j^-)^2(S_j^+)^2$, we obtain its expression, given after Eq. (S7). Similarly, when $j \neq j'$ and $n > 1$, we obtain that β in Eq. (S7) is related to the number of fully-magnetised product states that have one bimagnon at site j , and no bimagnon at j' , which is $\binom{L-2}{n-2}$. Its expression, given after Eq. (S7), then follows directly after taking into account the normalization factors and matrix elements. Hence using Eq. (S7) for any $k \neq k'$ it is clear that we obtain $\langle n, k' | n, k \rangle = 0$. Given that we work with normalised states, we can combine the arguments above to conclude that $\langle n, k | n', \pi \rangle = \delta_{n,n'} \delta_{k,\pi}$ whenever k is an integer multiple of $\frac{2\pi}{L}$ and L is even.

S2. AVERAGE ENERGY AND ENERGY VARIANCE FOR THE ASYMPTOTIC QMBS

In this section, we compute the average energy and variance of the asymptotic QMBS states $\{|n, k\rangle\}$ defined in Eq. (4).

A. Rewriting the asymptotic QMBS

For the convenience of explicit calculations, we propose the following rewriting of the asymptotic QMBS:

$$|n, k\rangle = \frac{1}{\sqrt{\mathcal{M}_{n,k}}} J_k^+ |n-1, \pi\rangle. \quad (\text{S9})$$

where the states $|n-1, \pi\rangle$ and $|n, k\rangle$ are normalised. As a first step, we compute the normalization factor coefficient $\mathcal{M}_{n,k}$, which can be directly deduced from Eq. (S7). That is, its expression reads

$$\mathcal{M}_{n,k} = \langle n-1, \pi | J_k^- J_k^+ | n-1, \pi \rangle = \frac{(L-n+1)(L-n)}{L-1} + \frac{L(L-n+1)(n-1)}{L-1} \delta_{k,\pi} \quad (\text{S10})$$

B. Average energy

To compute the average energy of the state $|n, k\rangle$, we first rewrite the OBC spin-1 XY Hamiltonian, along with the symmetry breaking perturbation [see discussion below Eq. (1)], as:

$$H = \frac{J}{2} \sum_{j=1}^{L-1} (S_j^+ S_{j+1}^- + S_j^- S_{j+1}^+) + h \sum_{j=1}^L S_j^z + D \sum_j (S_j^z)^2 + \frac{J_3}{2} \sum_{j=1}^{L-3} (S_j^+ S_{j+3}^- + S_j^- S_{j+3}^+). \quad (\text{S11})$$

In order to compute the average energy, we need to study the action of $[S_j^+ S_{j+1}^- + h.c.]$ onto the state $|n, k\rangle$, and for this it is convenient to consider the decomposition of $|n, k\rangle$ over sites j and $j+1$. For example, we can rewrite $|n, \pi\rangle$ as

$$|n, \pi\rangle = \alpha_{n,\pi} |+\rangle_j |+\rangle_{j+1} |\psi_{n,\pi,1}\rangle + \beta_{n,\pi} |-\rangle_j |-\rangle_{j+1} |\psi_{n,\pi,2}\rangle + \gamma_{n,\pi} \left(\frac{|+\rangle_j |-\rangle_{j+1} - |-\rangle_j |+\rangle_{j+1}}{\sqrt{2}} \right) |\psi_{n,\pi,3}\rangle; \quad (\text{S12})$$

where $\alpha_{n,\pi}$, $\beta_{n,\pi}$, and $\gamma_{n,\pi}$ are numbers with $|\alpha_{n,\pi}|^2 + |\beta_{n,\pi}|^2 + |\gamma_{n,\pi}|^2 = 1$, and $\{|\psi_{n,\pi,\ell}\rangle\}$ for $1 \leq \ell \leq 3$ are some states with support on sites other than j and $j+1$, and we have denoted the three spin-1 states on a site j by $|+\rangle_j$, $|-\rangle_j$, and $|0\rangle_j$. One can similarly rewrite the $|n, k\rangle$ as:

$$|n, k\rangle = \alpha_{n,k} |+\rangle_j |+\rangle_{j+1} |\psi_{n,k,1}\rangle + \beta_{n,k} |-\rangle_j |-\rangle_{j+1} |\psi_{n,k,2}\rangle + \gamma_{n,k} \left(\frac{|+\rangle_j |-\rangle_{j+1} - |-\rangle_j |+\rangle_{j+1}}{\sqrt{2}} \right) |\psi_{n,k,3}\rangle + v_{n,k} \left(\frac{|+\rangle_j |-\rangle_{j+1} + e^{ik} |-\rangle_j |+\rangle_{j+1}}{\sqrt{2}} \right) |\psi_{n,k,4}\rangle, \quad (\text{S13})$$

where $\alpha_{n,k}$, $\beta_{n,k}$, $\gamma_{n,k}$, and $v_{n,k}$ are numbers such that $|n, k\rangle$ is normalized and $\{|\psi_{n,\pi,\ell}\rangle\}$ for $1 \leq \ell \leq 4$ are some states without support on j and $j+1$. The action of the term $[S_j^+ S_{j+1}^- + h.c.]$ can then be directly computed to be:

$$(S_j^+ S_{j+1}^- + h.c.) |n, k\rangle = \sqrt{2} v_{n,k} (1 + e^{ik}) |0\rangle_j |0\rangle_{j+1} |\psi_{n,k,4}\rangle. \quad (\text{S14})$$

Using Eq. (S13), it then directly follows that $\langle n, k | (S_j^+ S_{j+1}^- + h.c.) |n, k\rangle = 0$. A similar reasoning can be carried out for the interaction term proportional to J_3 to show that $\langle n, k | (S_j^+ S_{j+3}^- + h.c.) |n, k\rangle = 0$, hence in all we obtain $\langle n, k | H |n, k\rangle = h(-L + 2n) + DL$ for all k . We conclude by noticing that the same result holds in PBC as well.

C. Energy variance

To compute the energy variance in any state, it is easy to see that the contribution of the terms in the Hamiltonian for which the state is an eigenstate simply vanishes. Hence, in the computation of the variance of $|n, k\rangle$, we can simply ignore the magnetic field and anisotropy terms in H of Eq. (S11), i.e., those that are proportional to h and D , since $|n, k\rangle$ are their eigenstates. For simplicity, we refer to the terms in H proportional to J and J_3 as H_1 and H_3 , respectively, and work with PBC. As we showed in the previous section, $\langle n, k | H_1 |n, k\rangle = \langle n, k | H_3 |n, k\rangle = 0$, and using similar ideas one can also show that $\langle n, k | H_1 H_3 |n, k\rangle = \langle n, k | H_3 H_1 |n, k\rangle = 0$. Hence the expression of the variance of $|n, k\rangle$ in H reduces to $\Delta H^2 = \langle n, k | (H_1 + H_3)^2 |n, k\rangle = \langle n, k | (H_1^2 + H_3^2) |n, k\rangle$. We now propose a rewriting of each term:

$$\begin{aligned} \langle n, k | H_\ell^2 |n, k\rangle &= \frac{1}{\mathcal{M}_{n,k}} \langle n-1, \pi | J_k^- H_\ell^2 J_k^+ |n-1, \pi\rangle = \frac{1}{\mathcal{M}_{n,k}} \langle n-1, \pi | [J_k^-, H_\ell] [H_\ell, J_k^+] |n-1, \pi\rangle = \\ &= \frac{1}{\mathcal{M}_{n,k}} \langle n-1, \pi | [H_\ell, J_k^+]^\dagger [H_\ell, J_k^+] |n-1, \pi\rangle, \end{aligned} \quad (\text{S15})$$

where $\ell = 1, 3$, and we have exploited the fact that $H_\ell |n-1, \pi\rangle = 0$. We a few straightforward algebraic passages, it is possible to show that:

$$\begin{aligned} [H_1, J_k^+] &= \frac{J}{2} \sum_{j=1}^L e^{ikj} [S_j^+ S_{j+1}^- + S_j^- S_{j+1}^+, \frac{1}{2}(S_j^+)^2 + \frac{e^{ik}}{2}(S_{j+1}^+)^2] \\ &= -\frac{J}{2} \sum_{j=1}^L e^{ikj} [\{S_j^z, S_j^+\} S_{j+1}^+ + e^{ik} S_j^+ \{S_{j+1}^z, S_{j+1}^+\}], \end{aligned} \quad (\text{S16})$$

where $\{\cdot, \cdot\}$ denotes the anti-commutator and we have used the identity $[S_m^+ S_n^-, (S_n^+)^2] = -2S_m^+ \{S_n^z, S_n^+\}$. The calculation proceeds by substituting Eq. (S16) into Eq. (S15) and it is greatly simplified by the fact that $S_j^z S_j^+ |n-1, \pi\rangle = S_j^z S_j^- |n-1, \pi\rangle = 0$. First, using this identity simplifies the action of $[H_1, J_k^+]$ on $|n-1, \pi\rangle$ to

$$[H_1, J_k^+] |n-1, \pi\rangle = -\frac{J}{2} \sum_{j=1}^L e^{ikj} [S_j^+ S_j^z S_{j+1}^+ + e^{ik} S_j^+ S_{j+1}^+ S_{j+1}^z] |n-1, \pi\rangle, \quad (\text{S17})$$

and Eq. (S15) then reads

$$\langle n, k | H_1^2 |n, k\rangle = \frac{J^2}{4\mathcal{M}_{n,k}} \sum_{j,j'=1}^L e^{ik(j-j')} \langle n-1, \pi | [S_j^z, S_{j'}^- S_{j'+1}^- + e^{-ik} S_{j'}^- S_{j'+1}^z S_{j'+1}^-] [S_j^+ S_j^z S_{j+1}^+ + e^{ik} S_j^+ S_{j+1}^+ S_{j+1}^z] |n-1, \pi\rangle. \quad (\text{S18})$$

We then notice that in Eq. (S18), all the terms with $j \neq j'$ in the sum vanish since the action of the sandwiched operator on $|n-1, \pi\rangle$ in such cases leads to inevitable appearance of spins with states $|0\rangle_m$ on certain sites m , which

in turn have a vanishing overlap with $\langle n-1, \pi |$. Hence, we can simplify Eq. (S18) to

$$\begin{aligned} \langle n, k | H_1^2 | n, k \rangle &= \frac{J^2}{4\mathcal{M}_{n,k}} \sum_{j=1}^L \langle n-1, \pi | [S_j^z S_j^- S_{j+1}^- + e^{-ik} S_j^- S_{j+1}^z S_{j+1}^-] [S_j^+ S_j^z S_{j+1}^+ + e^{ik} S_j^+ S_{j+1}^z S_{j+1}^+] | n-1, \pi \rangle \\ &= \frac{J^2}{4\mathcal{M}_{n,k}} \langle n-1, \pi | \sum_{j=1}^L \left[e^{-ik} S_j^- S_j^+ S_j^z S_{j+1}^z S_{j+1}^- S_{j+1}^+ S_{j+1}^- + S_j^- S_j^+ S_{j+1}^z S_{j+1}^- S_{j+1}^+ S_{j+1}^z \right. \\ &\quad \left. + S_j^z S_j^- S_j^+ S_{j+1}^- S_{j+1}^+ + e^{ik} S_j^z S_j^- S_j^+ S_{j+1}^- S_{j+1}^+ S_{j+1}^z \right] | n-1, \pi \rangle, \end{aligned} \quad (\text{S19})$$

Now we consider the expansion of $|n-1, \pi\rangle$ in the product state basis, as shown in Eq. (S8) and note that each of the terms in Eq. (S19) vanish on the basis states unless there is no bimagnon on both sites j and $j+1$. Hence we can simply count the number of such states and incorporate the normalization factor to obtain:

$$\langle n, k | H_1^2 | n, k \rangle = \frac{J^2}{\mathcal{M}_{n,k}} \sum_{j=1}^L \frac{\binom{L-2}{n-1}}{\binom{L}{n-1}} [e^{-ik} + 1 + 1 + e^{ik}] = \frac{4J^2 \cos^2\left(\frac{k}{2}\right)}{1 + \delta_{k,\pi} \frac{L(n-1)}{L-n}} = 4J^2 \cos^2\left(\frac{k}{2}\right), \quad (\text{S20})$$

where in the last step we have used the fact that the numerator anyway vanishes for $k = \pi$. The same calculation can be carried out in OBC and amounts to a multiplication of the result in Eq. (S20) by a factor $1 - \frac{1}{L}$, which does not change the PBC result in the thermodynamic limit. With similar arguments one can prove that:

$$\langle n, k | H_3^2 | n, k \rangle = 4J_3^2 \cos^2\left(\frac{3k}{2}\right), \quad (\text{S21})$$

thus recovering the result in Eq. (5). Once again, the choice of OBC amounts to a correction factor $1 - \frac{3}{L}$, which is irrelevant in the thermodynamic limit.

S3. ENTANGLEMENT ENTROPY OF THE EXACT AND ASYMPTOTIC QMBS

In this section we review the calculation of the entanglement entropy for the states $|n, k\rangle$, which proceeds along the lines of calculations performed in [14, 51]. We first divide the lattice into two parts, A and B . Typically, one considers A as the set of lattice sites with $j \leq L/2$ and B the rest, but this is not necessary. The key observation is that it is always possible to split the J_k^+ operators as a sum of an operator acting on A and of an operator acting on B :

$$J_k^+ = J_{k,A}^+ + J_{k,B}^+ = \frac{1}{2} \sum_{j \in A} e^{ikj} (S_j^+)^2 + \frac{1}{2} \sum_{j \in B} e^{ikj} (S_j^+)^2. \quad (\text{S22})$$

The state $|\Downarrow\rangle$ is a product state: $|\Downarrow\rangle_A \otimes |\Downarrow\rangle_B$. Hence, for $|n, \pi\rangle$, we obtain [14, 51]

$$|n, \pi\rangle = \frac{1}{\sqrt{N_{n,\pi}}} \left(J_{\pi,A}^+ + J_{\pi,B}^+ \right)^n |\Downarrow\rangle_A \otimes |\Downarrow\rangle_B = \frac{1}{\sqrt{N_{n,\pi}}} \sum_{m=0}^n \binom{n}{m} \left(J_{\pi,A}^+ \right)^m |\Downarrow\rangle_A \otimes \left(J_{\pi,B}^+ \right)^{n-m} |\Downarrow\rangle_B, \quad (\text{S23})$$

where $N_{n,\pi}$ is the normalization factor for the state $|n, \pi\rangle$, given by $\binom{L}{n}$. Additional care must be used in truncating the sum in the proper way: if A is composed of L_A lattice sites, it is not possible to apply the $J_{k,A}^+$ operator more than L_A times; similarly for L_B . Hence for simplicity, here we assume that $n < L_A, L_B$. Therefore, the expansion in Eq. (S23) gives the Schmidt decomposition of the state, which is composed of the $n+1$ orthogonal states $\{J_{k,\ell}^m |\Downarrow\rangle\}_{m=0}^n$ for the $\ell \in \{A, B\}$ part. In the presence of $n+1$ orthogonal states, the highest entropy state is the maximally mixed one, where they all have the same Schmidt coefficients; in that case $S_A = \log(n+1)$. If we consider a lattice of length L and the bipartition with $L_A = L_B = L/2$, the states with an extensive number of bimagnons are those such that $n = \alpha L$, with $0 < \alpha < 1$, and thus these states satisfy the following $S_A \sim \log L + \log \alpha$. As it is well-known, the quantum many-body scars have an entropy scaling with the logarithm of the volume.

Let us now consider the asymptotic QMBS states, $|n, k\rangle$. In this case, we use Eq. (S9) to obtain

$$|n, k\rangle = \frac{1}{\sqrt{\mathcal{M}_{n,k}}} J_k^+ |n-1, \pi\rangle = \frac{1}{\sqrt{\mathcal{M}_{n,k} N_{n-1, \pi}}} \sum_{m=0}^{n-1} \binom{n-1}{m} J_{k,A}^+ (J_{\pi,A}^+)^m |\Downarrow\rangle_A \otimes (J_{\pi,B}^+)^{n-1-m} |\Downarrow\rangle_B \\ + \frac{1}{\sqrt{\mathcal{M}_{n,k} N_{n-1, \pi}}} \sum_{m=0}^{n-1} \binom{n-1}{m} (J_{\pi,A}^+)^m |\Downarrow\rangle_A \otimes J_{k,B}^+ (J_{\pi,B}^+)^{n-1-m} |\Downarrow\rangle_B. \quad (\text{S24})$$

Note that unlike for the $|n, \pi\rangle$, Eq. (S24) is in general is not the Schmidt decomposition of the state. Yet, if we consider one subsystem, say A, the Schmidt states of a fixed magnetisation $-L_A + 2m$ are in the two-dimensional subspace spanned by the following linearly independent states:

$$\left(J_{\pi,A}^+\right)^m |\Downarrow\rangle_A, \quad J_{k,A}^+ \left(J_{\pi,A}^+\right)^{m-1} |\Downarrow\rangle_A. \quad (\text{S25})$$

Hence we can conclude that the total number of Schmidt states is at most $2n$, and for an extensive number of bimagnons $n = \alpha L$, we obtain that in the highest entropy situation $S_A \sim \log 2 + \log \alpha + \log L$. Thus, with respect to the exact QMBS $|n-1, \pi\rangle$, the asymptotic QMBS $|n, k\rangle$ has at most an additive correction of $\log 2$.

S4. VARIANCE OF THE EXACT QMBS FOR THE PERTURBED HAMILTONIAN

We consider the perturbed Hamiltonian $H' = H + V$, where H is the Hamiltonian (S11) with exact scars at finite size with PBC and $V = \frac{J_z}{L} \sum_j S_j^z S_{j+1}^z$. Since $|n, \pi\rangle$ is an eigenstate of H , the variance can be computed focusing only on V :

$$\Delta H'^2 = \Delta V^2 = \langle n, \pi | V^2 | n, \pi \rangle - \langle n, \pi | V | n, \pi \rangle^2 \\ = \frac{J_z^2}{L^2} \sum_{j,j'=1}^L (\langle n, \pi | S_j^z S_{j+1}^z S_{j'}^z S_{j'+1}^z | n, \pi \rangle - \langle n, \pi | S_j^z S_{j+1}^z | n, \pi \rangle \langle n, \pi | S_{j'}^z S_{j'+1}^z | n, \pi \rangle). \quad (\text{S26})$$

We can then use the structure of $|n, \pi\rangle$ to compute various correlation functions that appear in Eq. (S26). We first compute the two point correlation function to be

$$\langle n, \pi | S_j^z S_{j+1}^z | n, \pi \rangle = \frac{\binom{L-2}{n-2} + \binom{L-2}{n} - 2\binom{L-2}{n-1}}{\binom{L}{n}} \equiv F_2, \quad (\text{S27})$$

where we have used the action of $S_j^z S_{j+1}^z$ on the product basis states that compose $|n, \pi\rangle$, i.e., Eq. (S8), and noting that it takes the value of $+1$ if there are zero or two bimagnons on sites j and $j+1$, and -1 if there is one bimagnon. Using similar ideas, we obtain that when $j' \neq j-1, j, j+1$, the four point correlation function reads

$$\langle n, \pi | S_j^z S_{j+1}^z S_{j'}^z S_{j'+1}^z | n, \pi \rangle = \frac{\binom{L-4}{n} - 4\binom{L-4}{n-1} + 6\binom{L-4}{n-2} - 4\binom{L-4}{n-3} + \binom{L-4}{n-4}}{\binom{L}{n}} \equiv F_4 \quad (\text{S28})$$

Note that F_2 and F_4 in Eqs. (S27) and (S28) are numbers that only depend on L and n , and are independent of j ; and we have assumed that $n \geq 4$ and PBC. When $j' = j-1, j, j+1$, we obtain the following expressions for the “four point” correlation functions

$$\langle n, \pi | S_{j-1}^z (S_j^z)^2 S_{j+1}^z | n, \pi \rangle = \langle n, \pi | S_j^z (S_{j+1}^z)^2 S_{j+2}^z | n, \pi \rangle = \langle n, \pi | S_j^z S_{j+1}^z | n, \pi \rangle = F_2, \quad \langle n, \pi | (S_j^z)^2 (S_{j+1}^z)^2 | n, \pi \rangle = 1. \quad (\text{S29})$$

Combining Eqs. (S26)-(S29), and using translation invariance, we obtain that

$$\Delta H'^2 = \frac{J_z^2}{L^2} \left[\sum_j \sum_{j' \neq j-1, j, j+1} (F_4 - F_2^2) + \sum_j (1 - F_2^2) + 2 \sum_j (F_2 - F_2^2) \right] = J_z^2 \left[F_4 \left(1 - \frac{3}{L} \right) - F_2^2 + \frac{2}{L} F_2 + \frac{1}{L} \right] \quad (\text{S30})$$

Using Eq. (S30), we find that when $n/L = \nu$, where ν is a constant, $\Delta H'^2$ asymptotically scales as $\sim \frac{16\nu^2(1-\nu)^2}{L}$. On the other hand, when n is kept finite, $\Delta H'^2$ asymptotically scales as $\sim \frac{16n(n-1)}{L^3}$.




FIG. S1. First and second panel: The properties of the state $e^{-iHt} |n, k\rangle$ for $n = L/2$ and $k = 0$ as a function of time. First panel: time evolution of the squared magnetization $S^{z2}(t)$. Second panel: time evolution of the fidelity with the initial state $F(t)$. Third and fourth panel: The properties of the state $e^{-iHt} |+-+ \dots\rangle$ as a function of time. Third panel: time evolution of the squared magnetization. Fourth panel: time evolution of the fidelity with the initial state $F(t)$; in the inset we show the bare data, whereas in the main plot we rescale time by a factor $\sqrt{L}$ to display a clear collapse.

S5. DYNAMICS OF INITIAL STATES THAT ARE NOT ASYMPTOTIC QMBS

In this section, we study the dynamics of certain initial states, that are not asymptotic QMBS, under the Hamiltonian H of Eq. (1). We present this study in order to further support our claim that the dynamics of $|n, k = \pi - 2\pi/L\rangle$ is special.

A. Initial state with finite energy variance

First, we consider the states $|n, k = 0\rangle$, which are in the family of states of Eq. (4), but are not asymptotic QMBS since they have a finite energy variance in the thermodynamic limit, as evident from Eq. (5). Note that a state with finite energy variance was already discussed in Ref. [31], reaching similar conclusions. In Fig. S1 we study the dynamics of $|n, k = 0\rangle$ by presenting similar numerical results for the time-evolution of the latter state. The dynamics of the observable $S^{z2}(t)$ is “activated” on a short time-scale of order J^{-1} that does not depend on L (see the first panel of Fig. S1). The dynamics reaches a “pre-thermal” plateau [54] that increases to the initial value for $L \rightarrow \infty$. Note that this result does not contradict the fact that at finite size and in the long-time limit, observables should relax to their thermal value predicted by the diagonal ensemble. However, the thermalization timescale is much longer than the typical times that we can probe numerically using MPS-based methods. We have performed long-time simulations using exact diagonalization on small system sizes, and verified that this is indeed the case. Although the apparently long thermalization time may lead one to consider these states as asymptotic QMBS, the study of the fidelity with the initial state $F(t)$ is qualitatively very different. This is shown in the second panel of Fig. S1: on the same time-scale J^{-1} the state becomes essentially orthogonal to the initial one, and the data for different sizes are basically indistinguishable. The data on the fidelity relaxation time can be understood as a consequence of the finite energy-variance of the state $|n = L/2, k = 0\rangle$.

B. Initial Product State

It is also interesting to contrast the dynamics of the asymptotic QMBS with that of an uncorrelated product state; we consider here the staggered state $|\dots + - + - + - \dots\rangle$ which has the same zero magnetisation as the states considered in the main text and the same average squared magnetisation as the asymptotic QMBS, equal to one. The data on the dynamics of $S^{z2}(t)$ collapse on the same curve for all L considered (third panel of Fig. S1); the fidelity relaxation time instead becomes shorter with increasing L (fourth panel of Fig. S1). The behaviour is consistent with expectations for the time evolution of generic product states [53, 60], and is radically different from that of the asymptotic QMBS.




FIG. S2. Top: Squared overlap of $|n, \pi\rangle$ for $n = L/2$ with the eigenstates $|E_i\rangle$ of Hamiltonian H' with zero magnetisation, $S_z = 0$; the parameters of the simulation are $\{J, h, D, J_3, J_z\} = \{1, 0, 0.1, 0.1, 1.0\}$ and $L = 10$. The information on $|\langle E_i | n, \pi \rangle|^2$ is also encoded in the color code of the marker of all panels using a logarithmic scale, see colorbar. Middle and bottom: We plot the data of the top panel in a diagram with the energy E on the abscissa and the bipartition entanglement entropy S_E or the average square magnetisation $S^{z2}(E)$ of the eigenstate on the ordinate, respectively. The state $|n, \pi\rangle$ has overlap only with states whose S_{E_i} or $S^{z2}(E_i)$ lies on a continuous curve.

S6. SPECTRAL PROPERTIES OF THE HAMILTONIAN H'

In this section, we analyze the spectrum of the Hamiltonian H' discussed in the main text, which is a weak perturbation of the spin-1 XY Hamiltonian H exhibiting exact QMBS. We have numerically diagonalized the Hamiltonian H' and computed the overlap of the state $|n, \pi\rangle$ with all eigenvectors of the Hamiltonian. The results are reported in Fig. S2, they show clearly that the state has overlap only on “thermal” states whose bipartition entanglement entropy and average square magnetisation is determined by only their energy.

S7. UNIVERSAL RESCALING OF FIDELITIES

In this section, we present the data collapse of the fidelities for the asymptotic QMBS for various system sizes presented in the main text. Such a data collapse occurs at short times, once the time is rescaled by a factor that depends on the size of the system, as shown in Fig. S3. In the left panel, we present data for the asymptotic QMBS $|n, k = \pi - 2\pi/L\rangle$ time-evolved with the spin-1 XY Hamiltonian H of Eq. (1), which includes the term proportional to J_3 , and the collapse is obtained by rescaling the time as $\tau = t/L$. In the right panel, we present data for the state $|n, \pi\rangle$ time-evolved with the Hamiltonian $H' = H + V$; the collapse is obtained by rescaling the time as $\tau = t/L^{1/2}$.

It is interesting to link these results to the energy-time uncertainty relation of Eq. (6), whose proof is presented in many quantum mechanics textbooks and will not be reviewed here. The overlap of the time-evolved state with the initial one is related to the expectation value of the Hamiltonian and of its powers as [53]

$$\langle \Psi | e^{-iHt} | \Psi \rangle \approx 1 - it \langle \Psi | H | \Psi \rangle - \frac{1}{2}t^2 \langle \Psi | H^2 | \Psi \rangle + \frac{i}{6}t^3 \langle \Psi | H^3 | \Psi \rangle + \dots \quad (\text{S31})$$

and thus we can express the fidelity as

$$F(t) = |\langle \Psi | e^{-iHt} | \Psi \rangle|^2 \approx 1 - t^2 \left(\langle \Psi | H^2 | \Psi \rangle - \langle \Psi | H | \Psi \rangle^2 \right) + \dots \quad (\text{S32})$$

The short-time fidelity dynamics is thus completely dictated by the energy-variance of the initial state with respect to the Hamiltonian of the dynamics.

It is interesting to study a state with a Gaussian energy spread, for which the calculation of the time-dynamics of the fidelity is exactly possible. In fact here it is possible to show that it minimizes the inequality and has a fidelity $F(t)$




FIG. S3. Rescaling of the fidelities $F(t)$ plotted in Fig. 2 and Fig. 4 of the main text.

whose dynamics happens on the shortest possible timescale. Consider indeed an initial state that is a Gaussian linear superposition of energy eigenstates with average energy E_0 and energy variance σ^2 (we introduce also a normalisation prefactor $\alpha \in \mathbb{C}$): Assuming that the density of states in the energy window $[E_0 - \sigma, E_0 + \sigma]$ is approximately constant and takes the value $\rho(E_0)$, the scalar product between the time-evolved state and the initial one is given by:

$$\langle \Psi_0 | e^{-iHt} | \Psi_0 \rangle \approx \int |\alpha|^2 e^{-\frac{(E-E_0)^2}{2\sigma^2}} e^{-iEt} \rho(E) dE = |\alpha|^2 \sqrt{2\pi\sigma^2} e^{-iE_0 t} e^{-\frac{\sigma^2 t^2}{2}} \rho(E_0). \quad (\text{S33})$$

The normalisation of the state, computed for $t = 0$, requires that $|\alpha|^2 \sqrt{2\pi\sigma^2} \rho(E_0) = 1$. The fidelity $F(t)$ is the squared modulus of this scalar product and hence $F(t) = \exp[-\sigma^2 t^2]$; we can define the typical time scale of the fidelity dynamics as $\tau = 1/(2\sigma)$, and the energy-time inequality is satisfied and minimised.

In general terms, we thus expect that the dynamics of the fidelity at short times takes place on time-scales that are the shortest possible and minimize the energy-time inequality. This short-time behaviour is indeed verified by the numerics plotted in Fig. S3. In the left panel we have $\tau \sim L$ and $\sigma \sim 1/L$; in the right panel we have $\tau \sim \sqrt{L}$ and $\sigma \sim 1/\sqrt{L}$. However, it is important to keep in mind that the numerics has been performed only at short times and that long-time behaviours would need further investigation.

S8. HIGHER DIMENSIONAL GENERALISATIONS OF ASYMPTOTIC QMBS

Finally, we show that the existence of the asymptotic QMBS is not limited to one-dimensional systems, but can be easily generalised to higher-dimensional lattices. As an example, we consider a simple cubic Bravais lattice in $d > 1$ dimensions with primitive vectors $\mathbf{t}_i$ and $i = 1, \dots, d$; the vectors are adimensional and orthonormal: $\mathbf{t}_i \cdot \mathbf{t}_j = \delta_{ij}$. The lattice has linear dimension L and is composed of L^d sites; periodic boundary conditions (PBC) are applied. On each site of the lattice there is a spin-1 degree of freedom and we define the spin-1 operators $S_{\mathbf{r}}^\alpha$, with $\alpha = x, y, z$. We then consider a nearest-neighbor XY model with external magnetic field:

$$H = J \sum_{\mathbf{r}} \sum_{i=1}^d (S_{\mathbf{r}}^x S_{\mathbf{r}+\mathbf{t}_i}^x + S_{\mathbf{r}}^y S_{\mathbf{r}+\mathbf{t}_i}^y) + h \sum_{\mathbf{r}} S_{\mathbf{r}}^z. \quad (\text{S34})$$

As discussed in [14, 18], this model in Eq. (S34) exhibits exact QMBS for any finite value of L and for any dimension d . Note that when $d > 1$, the model of Eq. (S34) is non-integrable, and unlike in the one-dimensional case in Eq. (1), we

need not add the anisotropy term proportional to D or the longer range term proportional to J_3 to break integrability or unusual symmetries. Starting from the fully-polarised state $|\Downarrow\rangle$, we define the quasiparticle creation operator $J_{\mathbf{k}}^+ = \frac{1}{2} \sum_{\mathbf{r}} e^{i\mathbf{k}\cdot\mathbf{r}} (S_{\mathbf{r}}^+)^2$.

The exact QMBS states then read $|n, \boldsymbol{\pi}\rangle = \frac{1}{\sqrt{N_{n, \boldsymbol{\pi}}}} (J_{\boldsymbol{\pi}}^+)^n |\Downarrow\rangle$ where $\boldsymbol{\pi}$ is the vector with all d components equal to π . It is easy to show that $H|n, \boldsymbol{\pi}\rangle = h(-L^d + 2n)|n, \boldsymbol{\pi}\rangle$, hence the state is an exact QMBS in the middle of the spectrum of the Hamiltonian [14, 15, 18]. The states that we are interested in are:

$$|n, \mathbf{k}\rangle = \frac{1}{\sqrt{N_{n, \mathbf{k}}}} J_{\mathbf{k}}^+ (J_{\boldsymbol{\pi}}^+)^{n-1} |\Downarrow\rangle, \quad (\text{S35})$$

where $\mathbf{k}$ is any vector of the reciprocal space confined to the first Brillouin zone (1BZ). Similar to the one-dimensional case, it is possible to show that as long as the momentum $\mathbf{k}$ is chosen compatible with PBC in all directions, we can show that $\langle n, \mathbf{k} | n', \mathbf{k}' \rangle = \delta_{n, n'} \delta_{\mathbf{k}, \mathbf{k}'}$. With these states, we can directly repeat the proof in Sec. S2 *mutatis mutandis*. We find that the average energy is given by $\langle n, \mathbf{k} | H | n, \mathbf{k} \rangle = h(-L^d + 2n)$, and the energy variance is given by

$$\Delta H^2 = \frac{4J^2 \sum_{i=1}^d \cos^2\left(\frac{k_i}{2}\right)}{1 + \frac{(n-1)L^d}{L^d - n} \delta_{\vec{k}, \vec{\pi}}} = 4J^2 \sum_{i=1}^d \cos^2\left(\frac{k_i}{2}\right). \quad (\text{S36})$$

Thus, if we consider $\mathbf{k}$ with components $k_i = \pi + \frac{2\pi}{L} m_i$ and keep the $m_i \in \mathbb{Z}$ fixed while $L \rightarrow \infty$, the variance reduces to zero while being orthogonal to the exact QMBS. For such states, we expect the same phenomenology of asymptotic QMBS discussed for the one-dimensional case.