V. R. Shaginyan,^{1, 2, [∗](#page-0-0)} A. Z. Msezane,² and G. S. Japaridze²

 1 Petersburg Nuclear Physics Institute of NRC "Kurchatov Institute", Gatchina, 188300, Russia

 2 Clark Atlanta University, Atlanta, GA 30314, USA

The authors of Ref. [\[1\]](#page-0-1) consider the superconductivity in a twisted bilayer graphene and explore the profound effect of vanishingly small velocity in a superconducting Dirac flat band system. They claim: "In a flat band superconductor, the charge carriers' group velocity v_F is extremely slow. Superconductivity therein is particularly intriguing, being related to the long-standing mysteries of high-temperature superconductors and heavy-fermion systems". They continue: "In conclusion, twisted bilayer graphene (tBLG) requires us to face the challenge of ultra-strong coupling superconductivity in flat band Dirac systems, where, naively, there can be no transport or superconductivity".

The authors "naively" suggest that the superconductivity based on flat bands has not been thoroughly studied and analyzed. The emergence of flat bands has been predicted about thirty years ago which was accompanied by the astonishing claim that the superconducting gap $\Delta \propto g$, with g being the superconducting coupling constant [\[2](#page-0-2)[–7](#page-0-3)]. Simultaneously Fermi systems with flat band represent both the new liquid and the new state of matter $[3, 5, 6]$ $[3, 5, 6]$ $[3, 5, 6]$ $[3, 5, 6]$ $[3, 5, 6]$. It was also shown that an absolutely flat band supports the superconducting state, with the finite superconducting order parameter and the zero gap $[5, 6, 8]$ $[5, 6, 8]$ $[5, 6, 8]$ $[5, 6, 8]$ $[5, 6, 8]$. If the coupling constant g is finite, then the superconducting phase transition temperature T_c becomes finite and the flat band vanishes, since flat bands should be tuned with the superconducting state. As a result, the flat band becomes slightly tilted and rounded off at its end points $[5, 9-11]$ $[5, 9-11]$ $[5, 9-11]$, as predicted $[10]$.

Measurements of the Fermi velocity v_F in magic-angle twisted bilayer graphene as a function of T_c revealed that $T_c \propto v_F \propto 1/N_s$, where N_s is the density of states at the Fermi level. It was also shown that the high- T_c compounds $Bi_2Sr_2CaCu_2O_{8+x}$ exhibit similar behavior. This observation challenges theories of high- T_c superconductivity, since v_F normally correlates negatively with T_c : $T_c \propto 1/v_F \propto N_s$, see [\[9,](#page-0-8) [11\]](#page-0-9) and references therein. Note that our topical reviews $[5, 11]$ $[5, 11]$ and the monograph $[6]$ describe the physics of heavy fermion metals and high- T_c superconductors. There the topological fermion condensation quantum phase transition, leading to flat bands is considered and the special behavior of the strongly correlated Fermi systems is elucidated, a feature not exhibited by common Fermi liquids described by the Landau Fermiliquid theory. Topical reviews [\[5,](#page-0-5) [11\]](#page-0-9) and monograph [\[6](#page-0-6)] combine theoretical explanations and experimental data on the thermodynamic, transport and relaxation properties collected on heavy fermion metals, high- T_c superconductors and strongly correlated Fermi systems.

In summary, we have demonstrated that an absolutely flat band retains the superconducting state at $T_c \rightarrow 0$ with $v_F \rightarrow 0$. When T_c becomes finite, the flat band disappears, since it must be modified by the superconducting state. The authors [\[1\]](#page-0-1) have to take into account that at $T_c > 0$ the flat band distorts, becoming tilted, and $T_c \propto v_F$ [\[9](#page-0-8), [11\]](#page-0-9). Thus, the statement "the charge carriers' group velocity v_F is extremely slow" is incorrect and leads the authors [\[1](#page-0-1)] to the conceptional misunderstanding, confusing the reader.

[∗] Electronic address: vrshag@thd.pnpi.spb.ru

- [1] Tian H.; Gao X.; Zhang Y.; Che S.; et. al. Evidence for Dirac flat band superconductivity enabled by quantum geometry. Nature 2023, 614, 440.
- [2] Khodel, A.V.; Shaginyan, V.R. Superfluidity in systems with fermion condensate. JETP Lett. 1990, 51, 553.
- [3] Volovik, G.E. A new class of normal Fermi liquids. JETP Lett. 1991, 53, 222.
- [4] Khodel, V.A.; Shaginyan, V.R.; Khodel, V.V. New approach in the microscopic Fermi systems theory. Phys. Rep. 1994, 249, 1.
- [5] Shaginyan, V.R.; Amusia, M.Y.; Msezane, A.Z.; Popov, K.G. Scaling behavior of heavy fermion metals. Phys. Rep. 2010, 492, 31.
- [6] Amusia, M.Y.; Shaginyan, V.R. Strongly Correlated Fermi Systems: A New State of Matter. In Springer Tracts in Modern Physics; Springer Nature: Cham, Switzerland, 2020; Volume 283.
- [7] Törmä, P.; Peotta, S.; Bernevig, B.A. Superconductivity, superfluidity and quantum geometry in twisted multilayer systems. Nat. Rev. Phys. 2022, 4, 528.
- [8] Shaginyan, V.R.; Msezane, A.Z.; Stephanovich, V.A.; Kirichenko, E.V. Quasiparticles and quantum phase transition in universal low-temperature properties of heavy-fermion metals. Europhys. Lett. 2006, 76, 898.
- [9] Shaginyan, V.R.; Msezane, A.Z.; Amusia, M.Y.; Japaridze, G.S. Effect of superconductivity on the shape of flat bands. Europhys. Lett. 2022, 138, 16004.
- [10] Amusia, M.Y.; Shaginyan, V.R. Quasiparticle dispersion and lineshape in a strongly correlated liquid with the fermion condensate, Phys. Lett. 2000, A 275, 124.
- [11] Shaginyan, V.R.; Msezane, A.Z.; Japaridze, G.S. Peculiar Physics of Heavy-Fermion Metals: Theory versus Experiment, Atoms 2022, 10, 67.