

# Random Walk in the Boundary and Slow Roll in the Bulk

Yoshihisa KITAZAWA<sup>2),3)</sup> \*

<sup>2)</sup> *KEK Theory Center, Tsukuba, Ibaraki 305-0801, Japan*

<sup>3)</sup> *Department of Particle and Nuclear Physics*

*The Graduate University for Advanced Studies (Sokendai)  
Tsukuba, Ibaraki 305-0801, Japan*

## Abstract

The slow rolling inflation is dual to the random walk of conformal zero-mode. The  $O(N)$  enhancements of the two point functions : $N$  is the e-folding number, suppress the slow roll parameters by the same magnitude. The distribution function of conformal mode  $\rho_t(\omega)$  satisfies the Fokker Planck equation just like the Brownian particles. We identify the quantum corrections to de Sitter entropy with the von Neumann (entangled) entropy of the conformal zero-mode. They sum up as the running gravitational coupling:  $1/g(t) = \log N(t)/2$ . Under the Gaussian approximation, the distribution function obeys Fokker-Planck equation. FP equation boil down to a solvable first order partial differential equation (GFP). The identical equation is derived by the thermodynamic arguments. We study two types of the solutions of GFP:(1) UV complete spacetime and (2) inflationary spacetime with power potentials. The concavity of entangled entropy dictates the potential of inflaton is also concave. The maximum entropy principle favors the scenario: The universe is (a) born with small  $\epsilon$  and (b) grows large by inflation in the concave potentials.  $1 - n_s, r$  are  $O(1/N)$ . We predict  $1 - n_s \geq 0.02(0.016)$  and  $r \leq 0.08(0.066)$  for  $N = 50(60)$  at the pivot angle  $0.002 Mpc^{-1}$ . We propose a scenario to produce the curvature perturbation in the right ball park.

---

\*E-mail address: kitazawa@post.kek.jp

# 1 Introduction

The important questions on quantum gravity are specified with the Hubble scales. [1]

We may raise the cosmological constant problem, inflation [2–5], dark energy [6, 7] and so on [8].

It is persuasive to relate them to the infra-red behavior of quantum gravity just like hadron physics with the infra-red behaviors of QCD. With the accumulating observational information, the time is ripe to make a substantial progress on quantum gravity. Consistent quantum gravity such as string theory and matrix models are manifestly UV finite. What is difficult is to understand their infra-red behavior.

In the inflation theory, the universe underwent the accelerated expansion. Quantum fluctuations are redshifted to exit the cosmological horizon one after another. From the static observer's point of view, nothing goes out of the horizon but the conformal zero mode accumulates at the horizon. We build on the stochastic picture of infrared (IR) fluctuations [9], [10].

It is the diffusion process of the conformal zero-mode like that of the Brownian particles. The resummation of IR logarithmic effects in quantum gravity, which are  $O(\log^n a)$  in terms of the scale factor  $a$  of the universe play an essential role in elucidating cosmic random walk. [11, 12].

In our study of IR effects in quantum gravity, Einstein gravity plays the central role as the mass-less sector of string theory contains Einstein gravity. Some modifications of Einstein gravity are proposed in connection with inflation and dark energy. Phenomenologically it is necessary to introduce a single scalar field (inflaton) which couples minimally to gravity with a potential. By satisfying a slow roll condition, inflaton can explain the essence of inflation. Although it is a semi-classical modification of the Einstein gravity, the quantum effect exists at the heart of the theory, i.e. the vacuum fluctuation is stretched beyond Hubble scale by inflation. It should be possible to construct fully quantum theory by superseding these semi-classical theories. We argue de Sitter duality is the key to unify stochastic (quantum fluctuation) and dissipative (entropy generation) aspects.

We have computed the one loop quantum correction to Einstein gravity [24]. The bare Einstein gravity is inflation theory since it requires the inflaton field to construct the counter terms.: Einstein gravity is dual to inflation theory in this sense. The opposite is true as we obtain the former by integrating out the inflaton from the latter.

In section 2, our key observation is explained. The duality between the quantum effects of Einstein gravity and slowly rolling inflaton holds as the enhancement of the two point function leads to the suppression of the slow roll parameters.

In section 3, we argue the random walk of conformal zero mode is dual to slow roll inflation: This intuition has been our main impetus to pursue dS duality. In section 4, we derive the GFP from classical thermodynamics for inflation theory in a dual picture to the original quantum stochastic picture. It is based on the relation  $S_A = S_B$ . The bulk and boundary entropy are the same. It justifies de Sitter duality at the non-perturbative level. In section 5, we conclude this paper by listing our successes so far and tasks remaining ahead. Duality between random walk and slow roll defines the universal class in inflation theories. The slow roll parameters are  $O(1/N)$  and entropy favors the linear potential. If the de Sitter entropy is quantum :EE, the inflaton potential is concave. We construct the composite spacetime in which the curvature perturbation comes out naturally of right magnitude.

## 2 de Sitter Duality

The gravity has always been investigated from geometric point of view following Einstein. Inflation theory is no exception. On the other hand, the inflation theory leaves too many unexplained freedom. What is the inflaton? Why they perform slow roll on the potential? What determines the potential? and so forth. There is an alternative (dual) approach, i.e. stochastic point of view. This approach relies on a different point of view from classical geometric picture. The stochastic equations on the boundary play the major role like the Einstein's field equation. This dual approach is capable to explain the essence of inflaton. Namely, the mysterious slow roll can be interpreted as the random walk of conformal zero mode in a dual picture. This idea enables us to predict that the magnitude of slow roll parameters as  $\epsilon, \sim 1/N$ . This universal prediction is explained by the Langevin equation as we recall the nutshell of the argument as follows [15].

At large times, we assume the distribution should be canonical.

$$\dot{x} = b(t). \tag{2.1}$$

In Einstein gravity, there is only one massless scalar field. So this assumption holds. Let us consider

$$B(t)^\delta = \int_t^{t+\delta} dt' b(t'). \tag{2.2}$$

We assume  $\delta$  can be taken arbitrary small. In the low energy approximation, we can ignore the drift term.

Under such circumstances,

$$\overline{b(t_1)b(t_2)} = \delta(t_1 - t_2)H\tilde{g}. \tag{2.3}$$

$\tilde{g} = (3/4)g$  is the coupling projected in  $X$  field. The variance is computed as

$$\overline{(x(t) - x(0))^2} = \int_0^t dt_1 \int_0^t dt_2 \overline{b(t_1)b(t_2)} = \int dN \tilde{g}. \tag{2.4}$$

the slow roll conditions are satisfied, it is clear that the tensor to scalar ratio  $r$  is  $g^*/\int^* dN\tilde{g} \sim 1/N$ . The scalar spectral index is  $\tilde{g}^*/\int^* dN\tilde{g} \sim 1/N$ .

We assume  $N$  is large enough such that  $\tilde{g}, H$  are well approximated to be constant. More generic case will be discussed in the next section.

The probability distribution of  $x(t) - x(0)$  is

$$P(x, 0) = \sqrt{\frac{1}{2\pi\tilde{g}N}} \exp\left[-\frac{(x - x(0))^2}{2\tilde{g}N}\right]. \quad (2.5)$$

We simultaneously derived the kernel of Fokker Planck equation (3) as the major apparatus to explore the universe. We have thus far formulated inflationary universe in terms of Langevin and FP equation. The logic behind it is the duality between Random walk and slow roll inflation. The two point functions displays the identical scaling law for large  $N$  which indicates they belong to the same universal class. Alternatively we could resum IR logs to all orders. In fact the  $n$  point functions contain logarithmic contributions  $\log(a) \sim N$ . In order to determine the cosmological constant operator, we need to resum them.

We obtain FP equation from the renormalization group. It can determine the evolution of the entropy of the universe. So each solution of FP equation corresponds to a history of the universe.

In de Sitter space, the scale invariance of the infra-red quantum fluctuations may give rise to time dependent effects such as shielding the cosmological constant. While the inflaton generates entropy by dissipation, it represents the quantum effect in de Sitter dual picture in accordance with the fluctuation-dissipation theorem.  $(Ht)^n \sim (\log a)^n$  grow during the long cosmic history, we need to resum IR logarithms to all orders in order to realize such a mechanism. The renormalization group is one of the most powerful tool to accomplish it. In the critical phenomena, the divergence of the correlation length is caused by IR degrees of freedom. The important point is that there is only one massless scalar mode plays the significant role as explained in Appendix A [16]. We call it  $X$  field. It is nothing but the curvature perturbation  $\zeta$ .

The remarkable relation is  $S_A = S_B$ . The entropy of bulk and boundary is the same. The FP equation possesses the solutions in the both UV and IR finite. This fact offers a strong evidence for de Sitter duality.

Its solution with the ultraviolet fixed point indicates that the Universe starts with the de Sitter expansion near the Planck scale with  $\beta \sim \epsilon = 0$ .

In order to elucidate the IR logarithmic effects non-perturbatively, we formulate a Fokker Planck equation for the conformal zero modes of the metric. We obtain the  $\beta$  function for dimension-less gravitational coupling  $g = G_N H^2/\pi$  in a Gaussian approximation.  $G_N$  is the Newton's coupling and  $H$  is the Hubble parameter. It is valid for a small coupling  $g$  in quantum gravity except at the beginning of the Universe. We first sum up the  $O(\log^n a)$

terms to all orders to identify the one-loop running coupling  $g = 2/\log N$ . We next resum the  $1/(\log^n N)$  type corrections to take account of the quantum back-reaction on  $g$ . Since the  $\beta$  function turns out to be negative, it implies asymptotic freedom for  $g$  toward the future [17,18]. Furthermore, the  $\beta$  function possesses the ultraviolet (UV) fixed point in the past with the critical coupling  $g = 1/2$ . This fact indicates that our Universe begun with the dS expansion near the Planck scale with a minimal entropy.

Since  $\epsilon \propto \beta$ , the smallness of  $\epsilon$  in the beginning of the Universe is naturally explained. Strictly speaking, our approximation is justifiable when the coupling is small. Nevertheless,  $g = 1/2$  might be small enough such that our picture qualitatively holds. In our approach, the quantum correction to de Sitter entropy is given by the Gibbs entropy of the zero mode. We may decompose the total Hilbert space  $H = H_A \otimes H_B$  into  $H_A$  and  $H_B$ . Quantum definition of  $\rho_B$  involves the integration on A:  $\rho_B = \text{Tr}_A \rho^{\text{tot}}$ . This operation corresponds to the renormalization of  $\rho$  at the horizon scale. Then  $S_B = -\text{Tr}_B \rho_B \log(\rho_B)$ . We have counted the entropy of the mass less sector in the super-horizon. At the quantum level, the de Sitter entropy is von Neumann entropy, i.e. entangled entropy

### 3 Conformal mode in the leading log

The solution of FP equation implies the conformal mode behaves as the random walk. Since the fractal dimension of random walk  $D_f = 2$  is universal, the 2 point function is  $O(N) \sim 1/\epsilon$ . We can sum up the IR logarithms  $\log^n k$  by this equation to find a running coupling  $g(t)$ . We first work out the leading log solution which is valid in the large N limit. The FP (diffusion) equation shows that the solution is the Gaussian distribution with the standard deviation  $O(1/N(t))$  as shown in section 2 [15]. We first check whether we are on the right track. The distribution entropy of the conformal mode increases logarithmically,  $\frac{1}{2} \log \frac{1}{\xi} \sim \frac{1}{2} \log N(t)$ . In de Sitter space, entropy may be identified with the effective action with the opposite sign.  $-\Gamma = S$  as  $E = 0$ . We maximize the entropy in order to minimize the effective action.

Identifying the EE of conformal zero mode with the quantum correction to dS entropy, we obtain the bare action with the counter term

$$\frac{1}{g_B} = \frac{1}{g(N)} - \frac{1}{2} \log(N). \quad (3.1)$$

By requiring the bare action is independent of the renormalization scale: namely  $N$ , as we have just done to derive FP equation, we obtain the one loop  $\beta$  function.

$$\beta = \frac{\partial}{\partial \log(N)} g(N) = -\frac{1}{2} g(N)^2. \quad (3.2)$$

We find the running gravitational coupling in the leading log approximation:  $g(N) = 2/\log(N)$ . Semiclassical de Sitter entropy is in agreement with this estimate  $S(N) = 1/g(N) = \log(N)/2$ . We have collected the convincing support to our postulate. The

semiclassical evidence is that the EE is given by the gravitational coupling  $1/g = (\log N)/2$ . We have shown the renormalized distribution function obeys gravitational Fokker-Planck equation (GFP). That is equivalent to tune  $\xi$  self-consistently. and Langevin equations . Under the Gaussian approximation, they boil down to a solvable first order partial differential equation. The identical equation are derived by the thermodynamic arguments in the inflationary space-time. GFP determines the evolution of de Sitter entropy of the universe. It coincides with  $\beta$  function of  $g$ .

The holographic investigation at the boundary shows that  $g$  is asymptotically free toward the future. The renormalization group trajectory must reach Einstein gravity in the weak coupling limit for the consistency with general covariance [26]. We find that it approaches a flat space-time in agreement with this requirement.

We have evaluated the time evolution of EE entropy to the leading log order in [24]. In order to take account of the higher loop corrections in  $g$ , the FP equation should be precisely formulated. It turns out to be just necessary to make the equation covariant and local:  $\frac{\partial}{\partial N}\rho(N) - \frac{3g(N)}{4} \cdot \frac{1}{2} \frac{\partial^2}{\partial \omega^2}\rho(N) = 0$ . The covariance and locality ensure that the all the  $n$ -th loop divergences are local if all the sub divergence are subtracted. Furthermore the bare operator does not depend on the renormalization point (time).

In what follows, we consider the following Gaussian solutions of FP equation.

$$\rho = \sqrt{\frac{4\xi(t)}{\pi g(t)}} \exp\left(-\frac{4\xi(t)}{g(t)}\omega^2\right). \quad (3.3)$$

The standard deviatio  $\xi \propto 1/N$  decreases as we pointed out in the preceding section.

This distribution function determines the two point function of the conformal zero mode.  $\int d\omega \omega^2 \rho(\omega) = g(t)/8\xi(t)$ . We put the Gaussian ansatz into the FP equation (3) and find the following condition for the background to satisfy.  $\partial/\partial N (g/8\xi) = 3g/4$ .

We thus obtain a simple equation in terms of e-foldings.

$$\frac{\partial}{\partial N} \log \frac{g(t)}{\xi} = 6\xi. \quad (3.4)$$

This is the equation to investigate in what follows . Since the EE is  $(1/2) \log \frac{g(t)}{\xi}$ , this equation determines the evolution of entropy of a universe. This equation is derived from FP equation with the Gaussian approximation. The same equation holds in the inflation theory as explained later. In this way (3.4) supports de Sitter duality between Stochastic and thermodynamic approaches.

We have looked at FP equations from many view points. We have first adopted the microscopic approach. We derive FP equation from renormalization group. We then derive

it from macroscopic thermodynamics. We have also checked the consistency with Langevin formalism.

We first investigate a two point function of the conformal mode in the coincident limit  $\langle \omega^2 \rangle$  as it is a subtle step.

We recall the following identity holds at the horizon exit  $t = t^*$  [14].

$$\dot{N}^* \exp(N^*) = k. \quad (3.5)$$

where  $*$  denotes the exit time. It is the reason why the renormalization scales are related  $\log k = Ht^*$  in (3.5). It is the first come, the first served system. The softer the plain wave, the earlier its exit time.

We need to use time independent UV cut-off like dimensional regularization to identify time dependent quantum corrections. This is nothing but the second term of the Schwinger-de Witt expansion of the propagator. It is logarithmically divergent near 4 dimensions as  $1/\delta \sim 1/4 - d$  pole indicates

$$\langle \omega^2 \rangle = -\frac{3}{16\pi} \frac{G_{NR}}{6} \left( \frac{2}{\delta} - 2\log(k_*) \right)$$

Although the leading term is quadratically divergent, it vanishes in the dimensional regularization. As we emphasize, there is no time dependent UV contributions. We focus on the Hubble scale physics where  $a(t) = 1/(-\tau H) = \exp(Ht)$  and  $|k\tau| \sim 1$ . Our renormalization scale is  $\log k \sim -\log(-\tau) \sim Ht = N$ . Note the metric is negative, but  $k$  is the exiting momenta at the horizon and it increases with exiting time  $t^*$ . The renormalized 2 point function is

$$\langle \omega^2 \rangle_R = (3/4)g^* \log(k_*) = (3/4)g^* N^*$$

The bare distribution function is

$$\rho_B = \exp(\tilde{g}^*(1/\delta - \log(k_*))\partial^2/\partial\omega^2)\rho(\omega)$$

The FP equation is obtained as  $\rho_B$  is independent of the renormalization time.

$$\dot{\rho} = \tilde{g}\partial^2/\partial\omega^2\rho(\omega)$$

While the wave functions of the bulk modes oscillate with respect to  $\tau$ , the boundary mode is constant. That is why we call it the zero mode. Our strategy is to integrate out oscillating modes first.

We thus construct low energy effective theory around the Hubble scale by renormalizing the distribution function. This is the one of the fundamental results of our work [16]. We investigate it in the both its foundation and implications in this work. Our theory is identical

to that of Brownian particles in 1 + 1 dimensions. It is holographic in comparison to  $dS_4$ . In our theory conformal zero-mode random walk on the boundary (horizon).

As we have emphasized, Gaussian approximation must be valid unless we probe Planck scale. (3.4) determines the evolution of EE  $S = 1/2 \log g/\xi$  with respect to  $N$ . This formula confirms the validity of our postulate that distribution entropy of conformal zero mode constitutes the quantum correction to de Sitter entropy [20].

Secondly, the leading log approximation is valid in the large  $N$  limit since  $g \sim O(\log N)$  can be regarded as a constant. We obtain the equation which is the large  $N$  limit of (3.4). Equivalently we put it as follows.

$$\partial/\partial N \log \xi = -6\xi.$$

The solution is

$$\xi = \frac{1}{6N}. \quad (3.6)$$

Here we have neglected time dependence of  $1/g \sim \log N$  in comparison to  $1/\xi \sim N$ . In such a limit, (3.4) turns into (3). For finite  $N$ ,  $\xi = 1/6N$  will acquire finite  $N$  corrections like (3.7).

As it happens  $\epsilon = 1/4nN$  for the power potential, the result is in agreement with the slow roll picture  $\langle \zeta^2 \rangle = g/\epsilon$ . In this estimate, we have used Gaussian ansatz and large  $N$  expansion. It appears that only concave potential is allowed:  $n > 1$  if the de Sitter entropy is EE.

The solutions with more generic potentials are considered in the next section from de Sitter duality point of view. Our basic conjecture is that the mysterious slow roll of inflaton represents the Brownian motion of the conformal zero mode. This idea explains generic features and may develop further to demystify de Sitter duality. The attractive point of our theory is its simplicity. The inflaton is replaced by the random walk of the conformal zero mode. The universal enhancement of scalar to tensor ratio by  $O(N)$  is explained by its fractal dimension of the scalar modes. However the precise coefficient depends on the details.

There is a UV fixed point in our renormalization group. FP equation (3.4) enables us to evaluate higher order corrections to the  $\beta$  function. The expansion parameter is  $1/\log N$ . We can confirm that the following  $g_f$  and  $\xi_f$  satisfies (3.4),

$$g_f = \frac{2}{\log N} \left(1 - \frac{1}{\log N}\right), \quad \xi_f = \frac{1}{6N} \left(1 - \frac{1}{\log N}\right). \quad (3.7)$$

The leading log results are (3.2) and (3.6) respectively. Thus, the  $\beta$  function,  $\epsilon$  and the semi-classical entropy generation rate are given by  $\beta = \partial/\partial \log N g_f = -2/\log^2 N + 4/\log(N)^3$ .



$$\epsilon_f = -1/2 \partial/\partial N \log(g_f) = -1/2 g_f N \beta_f.$$

$$\frac{\partial}{\partial N} S_{sc} = \frac{\partial}{\partial N} \frac{1}{g_f} = -\frac{1}{N g_f^2} \beta_f \quad (3.8)$$

A remarkable feature is that the coupling has the maximum value  $g = 1/2$  at the beginning. It steadily decreases toward the future as the  $\beta$  function is negative in the whole region of time flow. It has two fixed points at the beginning and at the future of the Universe. The existence of the UV fixed point may indicate the consistency of quantum gravity. The single stone solves the  $\epsilon$  problem [25] as well since it vanishes at the fixed point. The  $\beta$  function describes a scenario that our Universe started the dS expansion with a minimal entropy  $S = 2$  while it has  $S = 10^{120}$  now.

Since we work with the Gaussian approximation, our results on the UV fixed points are not water tight as the coupling is not weak. Nevertheless we find it remarkable that they support the idea that quantum gravity has a UV fixed point with a finite coupling. In fact 4 dimensional de Sitter space is constructed in the target space at the UV fixed point of  $2 + \epsilon$  dimensional quantum gravity [26]. 4 dimensional de Sitter space also appears at the UV fixed point of the exact renormalization group [27] [28].

Such a theory might be a strongly interacting conformal field theory. However, it is not an ordinary field theory as the Hubble scale is Planck scale. Our dynamical  $\beta$  function is closely related to the cosmological horizon and physics around it. The existence of the UV fixed point could solve the trans-Planckian physics problem. A consistent quantum gravity theory can be constructed under the assumption that there are no degrees of freedom at trans-Planckian physics [29]. In this sense, it is consistent with string theory and matrix models. The Universe might be governed by (3.7) in the beginning as it might be indispensable to construct the UV finite solutions of the FP equation.

## 4 Inflationary Universe

The inflaton may be identified with the stochastic variable  $f$  whose correlators show characteristic features of Brownian motion.  $\langle f^2 \rangle = \tilde{N}$  and  $g \propto \tilde{N}^{m/2} = \langle f^m \rangle$ . The increase of the entropy  $S = 1/g$  can be evaluated by the first law  $T \Delta S = \Delta E$  where  $\Delta E$  is the incoming energy flux of the inflaton [30]. Obviously we are estimating the entropy of super-horizon.

Since these solutions are scale invariant, Langevin equation determines the scaling behavior precisely in contrast to the logarithmic correlators.

In this way, the one of the Einstein's equation is obtained:

$$\dot{H}(t) = -4\pi G_N \dot{f}^2. \quad (4.1)$$

This formula can be re-expressed in terms of the slow roll parameter:  $\epsilon = \dot{f}^2/2H^2$

$2\epsilon = -\frac{\partial}{\partial N} \log g(t)$ . The tilt of the gravitational wave spectrum is checked:  $n_t = -2\epsilon$ . We add  $O(1/\tilde{N})$  quantity to the both sides of the equation.  $2\epsilon + \frac{1}{\tilde{N}} = -\frac{\partial}{\partial N} \log(g\tilde{N})$ .

We find the relation between  $\xi$  and  $\epsilon$  in (4.4) :  $6\xi = 2\epsilon + \frac{1}{\tilde{N}} = \frac{(m+2)}{2\tilde{N}}$  is reproduced. The tilt of the conformal mode is reproduced. We have thus shown that curvature and conformal perturbation are identical up to the constant factor (4.2). For power potential inflationary universe, (4.7) is rewritten as

$$6\xi = \frac{\partial}{\partial N} \log \frac{g}{\xi} = \frac{\partial}{\partial N} \log \frac{g}{\epsilon}.$$

It is consistent with our GFP (3.4). The second equality arises since  $\xi/\epsilon$  is independent of  $N$  as shown in (4.4). This ambiguity corresponds to a constant  $c$  in (4.2). In other words,  $\xi = \frac{m+2}{12\tilde{N}}, \epsilon = \frac{m}{4\tilde{N}}$  are  $O(1/\tilde{N})$ . The both can balance the equation.

In this form, it is evident that  $6\xi$  is the scalar spectral index in agreement with slow roll inflation theory. Tensor to scalar ratio is  $16\epsilon$ . We have shown GFP can be derived by macroscopic arguments also. From quantum gravity point of view, this is an important non-perturbative evidence for de Sitter duality: quantum/geometric duality.

EE must be concave from strong sub-additivity [21] [22]. Inflationary universe with a slow roll parameter may be well approximated as de Sitter space locally

The concave power solutions can be obtained from convex solutions by formally replacing  $m$  by  $1/n$  where  $n > 1$  has to be a real number .

$$\frac{\partial^2}{\partial N^2} \left( \frac{1}{g} \right) = \frac{(1/n)(1/n-1)}{N^2} \left( \frac{1}{g} \right), \quad \frac{\partial^2}{\partial N^2} S < 0$$

We conclude that potentials for consistent inflation theory are concave. In fact convex potentials are disfavored by observation.

The conformal mode of the inflationary universe is performing the Brownian motion. Its trajectories constitute 2 dimensional fractal. The concave potentials are promising avenue to explore right now. The potential of inflation theory may be identified with the entangled entropy which is always concave.

As it turns out, the equation (3.4) has another class of solutions: the inflationary universe with the power potentials.

$$g = c\tilde{N}^{\frac{m}{2}}, \quad \xi = \frac{m+2}{12\tilde{N}}. \quad (4.2)$$

Since these are solutions of FP equation, they have many common features. The enhancement of scalar 2 point function by  $O(N)$  is such an example.

Here  $c$  is an integration constant.  $m$  denotes the power of the potential:  $f^m$ . We have changed the variables by replacing  $N$  by  $\tilde{N}$  where  $\tilde{N} = N_e - N$ .  $N_e$  denotes the e-folding number at the end of inflation. To be precise, they are the solutions of the following equation:

$$-\frac{\partial}{\partial N} \log \frac{g(\tilde{N})}{\xi(\tilde{N})} = 6\xi(\tilde{N}). \quad (4.3)$$

(4.2) shows  $\epsilon$  and  $\xi$  are related as

$$\epsilon = \frac{m}{4\tilde{N}} = \frac{3m}{m+2}\xi. \quad (4.4)$$

In this way, the one of the Einstein's equation is obtained:

$$\dot{H}(t) = -4\pi G_N \dot{f}^2. \quad (4.5)$$

This formula can be re-expressed in terms of the slow roll parameter:  $\epsilon = \dot{f}^2/2H^2$

$$2\epsilon = -\frac{\partial}{\partial N} \log g(t). \quad (4.6)$$

The tilt of the gravitational wave spectrum is  $n_t = -2\epsilon$ . We add  $O(1/\tilde{N})$  quantity to the both sides of the equation.

$$2\epsilon + \frac{1}{\tilde{N}} = -\frac{\partial}{\partial N} \log(g\tilde{N}). \quad (4.7)$$

We find the relation between  $\xi$  and  $\epsilon$  in (4.4) :

$6\xi = 2\epsilon + \frac{1}{\tilde{N}} = \frac{(m+2)}{2\tilde{N}}$  is reproduced. We have thus shown that curvature and conformal perturbation are identical up to the constant factor (4.2).

For power potential inflationary universe, (4.7) is rewritten as

$$6\xi = \frac{\partial}{\partial N} \log \frac{g}{\xi} = \frac{\partial}{\partial N} \log \frac{g}{\epsilon}. \quad (4.8)$$

It is consistent with our GFP (3.4). The second equality arises since  $\xi/\epsilon$  is independent of  $N$  as shown in (4.4). This ambiguity corresponds to a constant  $c$  in (4.2). In other words,  $\xi = m + 2/12\tilde{N}, \epsilon = m/4\tilde{N}$  are  $O(1/\tilde{N})$ . The both can balance the equation.

In this form, it is evident that  $6\xi$  is the scalar spectral index in agreement with slow roll inflation theory. Tensor to scalar ratio is  $16\epsilon$ . We have shown GFP can be derived by macroscopic arguments also. From quantum gravity point of view, this is an important non-perturbative evidence for de Sitter duality: quantum/geometric duality.

	$n = 1$	$n = 2$	$n = 3$
$\epsilon$	$\frac{1}{4\tilde{N}}$	$\frac{1}{8\tilde{N}}$	$\frac{1}{12\tilde{N}}$
$n_s$	$1 - \frac{3}{2\tilde{N}}$	$1 - \frac{5}{4\tilde{N}}$	$1 - \frac{7}{6\tilde{N}}$
$\frac{\partial}{\partial \log k} n_s$	$-\frac{3}{2\tilde{N}^2}$	$-\frac{5}{4\tilde{N}^2}$	$-\frac{7}{6\tilde{N}^2}$

Table 1:

$\tilde{N} = 50$	$n = 1$	$n = 2$	$n = 3$
$r$	0.08	0.04	0.027
$n_s$	0.97	0.975	0.977
$\frac{\partial}{\partial \log k} n_s$	-0.0006	-0.0005	-0.00046

Table 2: N=50

$\tilde{N} = 60$	$n = 1$	$n = 2$	$n = 3$
$r$	0.066	0.033	0.022
$n_s$	0.975	0.97911	0.981
$\frac{\partial}{\partial \log k} n_s$	-0.0006	-0.0005	-0.00046

Table 3: ,N=60

Planck 2018
$r_{0.002} < 0.065$
$n_{s,0.002} = 0.979 \pm 0.0041$
$dn_s/d \log k = -0.006 \pm 0.013$

Table 4: , Planck 2018

In Table 1, we list expected  $\epsilon$ ,  $1 - n_s$ ,  $\frac{\partial n_s}{\partial \log k}$  in the power potential model with  $f^{1/n}$ ,  $n = 1, 2, 3$ .  $n > 1$  for the concave potential. We note  $1 - n_s$  is bounded from below by  $1/\tilde{N}$  while  $r = 16\epsilon$  is not. Table II lists our numerical estimates by putting  $\tilde{N} = 60$  in Table I.

They are consistent with the current observations [33].

We adopt the same pivot scale  $0.002Mpc^{-1}$  for  $r, n_s$  to respect self-consistency. It is important to establish the bound on  $n$ . We recall here the curvature perturbation:

$$P \sim 2.2 \times 10^{-9}. \quad (4.9)$$

It is bounded from below  $g > 10^{-11}$  if  $\epsilon > 1/200$ .

The entropy grows logarithmically  $1/g = (\log N)/2$  as  $N$  grows after the Universe emerges out of the UV fixed point. In the inflationary phase, the entropy grows power like  $1/g \sim 1/\tilde{N}^{1/2n}$ . The entropy increases slowly in the pre-inflation era while it grows much faster in the inflation era. Inflation phase inevitably takes over the pre-inflation phase as the entropy of the former dominates in time.

The simplest scenario assumes that the inflation starts sometime after the birth of the Universe. We hope to explain the observed magnitude of the curvature perturbation  $P \sim 10^{-10}$ . This is a fundamental quantity of the Universe.

Since the inflation does not stop unless  $\epsilon$  grows up to 1. It is necessary to cross over to the slow roll inflation phase.

We find it attractive to consider the composite solutions, which consist of the logarithmic solution (3.7) and the the power-law solution (4.2). They may describe the whole history of the Universe from its birth at the UV fixed point to the completion of inflation. Let us investigate a pre-inflation scenario. The entropy grows logarithmically  $1/g = (\log N)/2$  as  $N$  grows after the Universe emerges out of the UV fixed point. In the inflationary phase, the entropy grows power like  $1/g \sim 1/\tilde{N}^{1/2n}$ . The entropy increases slowly in the pre-inflation era while it grows much faster in the inflation era.

## 5 Conclusions and Discussions

Although we have argued the necessity of the composite universe, it is another matter to understand the mechanism and the consistency of the transition. We have explored composite solutions which realize the above transition scenario [32]. The system consists of the logarithmic solution (3.7) and the the power-law solution (4.2). The entropy  $1/g$  always increase with cosmic evolution. They may describe the whole history of the universe from its birth at the UV fixed point until the finale of the inflation.

Let us investigate pre-inflation scenario. The entropy grows logarithmically  $1/g = (\log N)/2$  as  $N$  grows after the Universe emerges out of the UV fixed point. In the inflationary phase, the entropy grows power like  $1/g \sim 1/\tilde{N}^{1/2n}$ . The entropy increases slowly in the pre-inflation era while it grows much faster in the inflation era. Inflation phase inevitably takes over the pre-inflation phase as the entropy of the former dominates in time.

The simplest scenario assumes that our universe is the 50 : 50 composite of the UV complete and inflationary universes. We recall  $\epsilon = -(1/2)\frac{\partial}{\partial N} \log H^2$ . We may consider composite universe with the geometric averaged Hubble parameter:  $H^2 = H_1 H_2$ . The linear average  $\epsilon$  is  $\bar{\epsilon} = (\epsilon_1 + \epsilon_2)/2$ .

$\epsilon_1 \sim 1/2N \log(N)$ ,  $\epsilon_2 \sim 1/4n\tilde{N}$ . Note the former decreases while the latter increases with  $N$ .

Therefore the inflation starts when  $|\epsilon_1| \sim |\epsilon_2|$ . A more explicit estimate with  $n = 2$  is  $N \sim \tilde{N} \sim 10^{11}$ .

After this epoch, the universe is dominated by the inflation theory with  $\tilde{N} < 10^{11}$ . It is remarkable that the curvature perturbation  $P \sim 10^{-11}$  comes out of the right magnitude without fine tuning.

Finally we reproduce  $\epsilon_* = 1/4n\tilde{N}_*$  at the horizon exit. Since  $\epsilon \sim 1$  at the end of inflation, the whole period takes much longer time than the minimal inflation [31]. From holographic perspective, we have formulated the FP equation for the conformal mode at the boundary, i.e., the horizon. The GFP is derived under the Gaussian approximation, which may be justifiable for small  $g = G_N H^2/\pi$ . The von Neumann entropy of conformal zero mode gives the quantum correction to the dS entropy  $S = 1/g$ .

Our transition scenario is supported by the expectation that the solution with dominant entropy ( $\epsilon$ ) is chosen. Since the entropy  $1/g$  for (3.7) increases logarithmically and that for (4.2) increases in a power-law, the former is chosen initially and the latter is chosen if we wait long enough.

Holography is a specific feature of quantum gravity. In this paper we have derived the Fokker-Planck equation on the boundary from the renormalization group equation in the bulk. We have stressed the de Sitter duality between the bulk geometry and the boundary random walk at the horizon. de Sitter duality is deeply connected with the von Neumann or entangled entropy of the conformal zero mode.

Suppose the total Hilbert space is factorized into the bulk states and boundary states. The entropy of the bulk and the boundary are the same. This explains why de Sitter duality is important. It is not expected that  $\beta$  function at the UV fixed point can be accessible from the FP equation at the boundary. Nevertheless the dual pair have the identical entropy which coincide with the effective action. This set-up allows us to evaluate EE in the both quantum procedure and in the classical thermodynamics.

Furthermore we can even pull out the promising candidate for the curvature perturbation. Its an encouraging sign for our research achievements in quantum gravity on de Sitter space.

## acknowledgment

This work is supported by Grant-in-Aid for Scientific Research (C) No. 16K05336. We thank Hiroyuki Kitamoto, Masashi Hazumi, Satoshi Iso, Hikaru Kawai, Kozunori Kohri, Takahiko Matsubara, Jun Nishimura, Hirotaka Sugawara, and Takao Suyama for discussions.

## References

- [1] A. Polyakov, arXiv:0709.2899 [hep-th].
- [2] A. A. Starobinsky, JETP Lett. **30**, 682 (1979) [Pisma Zh. Eksp. Teor. Fiz. **30**, 719 (1979)].
- [3] A. H. Guth, *The Inflationary Universe* (Perseus Books, New York, 1997)
- [4] A. D. Linde, Phys. Lett. **108B**, 389 (1982) [Adv. Ser. Astrophys. Cosmol. **3**, 149 (1987)].
- [5] A. Albrecht and P. J. Steinhardt, Phys. Rev. Lett. **48**, 1220 (1982) [Adv. Ser. Astrophys. Cosmol. **3**, 158 (1987)].
- [6] B. Ratra and P. J. E. Peebles, Phys. Rev. D **37**, 3406 (1988).
- [7] R. R. Caldwell, R. Dave and P. J. Steinhardt, Phys. Rev. Lett. **80**, 1582 (1998) [astro-ph/9708069].
- [8] Brandenberger. Robert , Comeau. Vincent, Fossati. Leonardo and Lavini, Heisenberg arXiv: 2201.08824[hep-th].
- [9] A. A. Starobinsky, Lect. Notes Phys. **246**, 107 (1986); A. A. Starobinsky and J. Yokoyama, Phys. Rev. D **50**, 6357 (1994).
- [10] N. C. Tsamis and R. P. Woodard, Nucl. Phys. B **724**, 295 (2005).
- [11] N. C. Tsamis and R. P. Woodard, Commun. Math. Phys. **162**, 217 (1994).
- [12] H. Kitamoto and Y. Kitazawa, Phys. Rev. D **87**, 124007 (2013).
- [13] H. Kitamoto and Y. Kitazawa, Int. J. Mod. Phys. A **29**, 1430016 (2014).
- [14] J. M. Maldacena, JHEP **0305**, 013 (2003) [astro-ph/0210603].
- [15] G. Parisi, Statistical Field Theory, Frontiers in Physics **66**, (1987).

- [16] Yoshihisa Kitazawa, J.Phys.Soc.Jap. 91(2022)7,074003 2011.14640[hep-th].
  - [17] D. J. Gross and F. Wilczek, Phys. Rev. Lett. **30**, 1343 (1973).
  - [18] H. D. Politzer, Phys. Rev. Lett. **30**, 1346 (1973).
  - [19] G. W. Gibbons and S. W. Hawking, Phys. Rev. D **15**, 2752 (1977).
  - [20] Leonard Susskind and John Uglum, Phys.Rev.D**50**, 2700-2711 (1994) [hep-th/9401070].
  - [21] T. Nishioka, S. Ryu, T. Takayanagi, arXiv:0905.0932, J.Phys.A42:504008,2009.
  - [22] Calabrese-Cardy, arXiv:0905.4013, J.Phys.A42:504005,2009.
- 7d
- [23] H. Kitamoto and Y. Kitazawa, Phys. Rev. D **99**, 085015 (2019).
  - [24] H. Kitamoto, Y. Kitazawa and T. Matsubara, Phys. Rev. D **101**, 023504 .
  - [25] R. Penrose, Annals N. Y. Acad. Sci. **571**, 249 (1989).
  - [26] H. Kawai, Y. Kitazawa and M. Ninomiya, Nucl. Phys. B **404**, 684 (1993).
  - [27] M. Reuter, Phys. Rev. D**57** (1998) 971.
  - [28] W. Souma PTP **102**, 181 (1999).
  - [29] A. Bedroya and C. Vafa, arXiv:1909.11063 [hep-th].
  - [30] A. V. Frolov and L. Kofman, JCAP **0305**, 009 (2003).
  - [31] H. Kitamoto, Y. Kitazawa, R. Kojima, Phys. Rev. D**96**, 023535 (2017).
  - [32] J. K. Jain, *Composite Fermions*, (Cambridge University Press, New York, 2007).
  - [33] N. Aghanim *et al.* [Planck Collaboration], arXiv:1807.06209 [astro-ph.CO].