

Electron wave functions in beta-decay formulas revisited (II): Completion including recoil-order and induced currents

Wataru Horiuchi^{1,2,3,4}, Toru Sato⁵, Yuichi Uesaka⁶, and Kenichi Yoshida^{5,3}

¹*Department of Physics, Osaka Metropolitan University, Osaka 558-8585, Japan*

²*Nambu Yoichiro Institute of Theoretical and Experimental Physics (NITEP), Osaka Metropolitan University, Osaka 558-8585, Japan*

³*RIKEN Nishina Center, Wako, Saitama 351-0198, Japan*

⁴*Department of Physics, Hokkaido University, Sapporo 060-0810, Japan*

⁵*Research Center for Nuclear Physics, Osaka University, Ibaraki, Osaka 567-0047, Japan*

⁶*Faculty of Science and Engineering, Kyushu Sangyo University, Fukuoka 813-8503, Japan*

.....
 We present complete formulas of the allowed and first-forbidden transitions of the nuclear beta decay taking into account the recoil-order and induced currents up to the next-to-leading order (NLO). The longitudinal part of the vector current is cleared away by the use of the conservation of the vector current for the multipole operators of the natural-parity transitions, which makes the effect of the meson exchange current for the vector current as small as possible. The formula is transparent enough to be applied to various beta-decay processes. As a numerical demonstration, we apply our formulas to the beta decay of a neutron-rich nucleus ^{160}Sn . We find that the NLO corrections amount to 10–20% of the total decay rate, whereas the induced currents alter the rate at most 5%.

1 Introduction

Nuclear beta decay has attracted renewed interest. The astrophysical rapid-neutron-capture process, the r-process nucleosynthesis, is believed to produce more than half of heavier elements than iron. The beta decay is a predominant process in the freeze-out phase [1], and thus plays a decisive role in forming the abundance pattern. The detection of gravitational waves from a binary neutron-star merger and associated electromagnetic waves from a kilonova/macronova has provided an evidence of the r-process site [2, 3]. In such an exotic environment, heavy neutron-rich nuclei even close to the dripline are involved in the r-process.

An interest in the nuclear beta decay with a precise description also lies in neutrino physics, such as the reactor neutrino anomaly [4, 5] and the search for the beyond-standard-model (BSM) physics [6]. The long-standing issue of nuclear physics, the effective (weak) axial vector coupling constant (g_A^{eff}), might be resolved in the electron spectrum of the higher-order-forbidden transitions [7].

In our previous paper [8] [referred to as paper (I)], formulas of the beta-decay rate in a convenient and practical form were proposed. The nuclear beta decay was formulated using the effective multipole operators [9, 10]. The analytic forms of the electron wave function in the next-to-leading order (NLO) of the integral equation of the electron Dirac equation and the formulas of the effective nuclear multipole operators were presented. It was demonstrated that the NLO formula reproduces well the ‘exact’ decay rate of the allowed Gamow–Teller and first-forbidden spin-dipole transitions for heavy nuclei with large Z , where the frequently used formula[11] corresponding to a leading order formula of the systematic expansion in terms of nuclear radius, the Coulomb potential, the energy and the mass of electron[12], possibly overestimates the decay rates.

The formula in paper (I) is expressed in terms of the nuclear current density that is general enough to incorporate any models of the nuclear current such as many-body nuclear current in addition to the one-body current, non-relativistic and relativistic description of nuclear current. However, the explicit formula in the impulse approximation is given only for the leading $\sigma\tau$ operator of the axial vector current. It is certainly necessary and valuable to update the formula of paper (I) by including the velocity-dependent terms such as $A_0 \sim \boldsymbol{\sigma} \cdot \mathbf{p}/M\tau$, $\mathbf{V} \sim \mathbf{p}/M\tau$ and the induced currents.

In this paper, we give the formulas of the nuclear multipole operators using the non-relativistic impulse approximation of the nuclear current. The velocity-dependent terms of the time component of the axial-vector current, the space component of the vector current and the induced terms, i.e., the weak-magnetism term and the induced pseudo-scalar term,

are included. These nuclear currents play an important role in the first-forbidden transitions. For the 0^- transition, for example, the time component of the axial-vector current competes with the spin-dipole operator [13]. The induced pseudo-scalar term would play a minor role in the nuclear beta decay. However, we keep it because the formulas can be used for the neutrino reactions including muon and tau leptons, while the so-called second class currents, the G-parity irregular terms [14, 15], are not included.

In neutrino reactions and beta decay in the plane-wave approximation of lepton wave functions, a longitudinal part of the vector current is eliminated using the conservation of current, and further the LO term of the transverse electric current can be written in terms of the Coulomb multipole operator [16–18]. In the beta decay or muon capture, because of the complication due to the Coulomb effect, this is done in the leading order of the long-wave-length approximation. In the muon capture, this was discussed in connection with the extended Siegert’s theorem [19]. In this paper, we rewrite the formula of paper (I) using the Fourier–Bessel transformation and the current conservation, which makes the effect of the meson-exchange current for the vector current as small as possible for the allowed and first-forbidden transitions. This work provides complete formulas of the allowed and first-forbidden transitions of the nuclear beta decay up to NLO within the non-relativistic impulse approximation of the nuclear current.

In section 2, the multipole operators for the parity $P = (-1)^J$ and $(-1)^{J+1}$ are given. The longitudinal part of the vector current is rewritten in terms of the time component of the vector current with the help of the current conservation. In section 3, a formula of the multipole operators in the impulse approximation is given. As a numerical demonstration of the usefulness of our method, the formula is applied to the beta decay of ^{160}Sn in section 4 together with the formula written in terms of nuclear transition densities. The transition densities are obtained by a nuclear energy-density-functional method. A summary is given in section 5.

2 Multipole expansion of the effective Hamiltonian

The effective Hamiltonian for the beta (β^\mp) decay is given by

$$H_{\text{eff}} = \frac{G_F V_{ud}}{\sqrt{2}} \int d\mathbf{x} [\bar{e}(\mathbf{x}) \gamma^\mu (1 - \gamma_5) \nu_e(\mathbf{x}) J_\mu(\mathbf{x}) + \text{H.c.}], \quad (1)$$

where $G_F = 1.166 \times 10^{-5} \text{ GeV}^{-2}$ is the Fermi coupling constant and $V_{ud} = 0.9737$ is the up-down element of the Cabibbo–Kobayashi–Maskawa matrix. This Hamiltonian consists of the

electron field e , the neutrino field ν , and the hadronic current

$$J^\mu(\mathbf{x}) = \mathcal{V}^\mu(\mathbf{x}) - \mathcal{A}^\mu(\mathbf{x}), \quad (2)$$

with the vector $[\mathcal{V}^\mu = (V^0, \mathbf{V})]$ and axial-vector $[\mathcal{A}^\mu = (A^0, \mathbf{A})]$ currents. Using the multipole operator $\Xi_{JL}(\kappa_e, \kappa_\nu)$ constructed from partial waves of electron and neutrino specified in terms of κ_e and κ_ν [8], one obtains the rate of the β^\mp decay from the initial (i) to final (f) nuclear states, $i \rightarrow f + e^\mp + \bar{\nu}_e(\nu_e)$, as

$$\Gamma = \frac{(G_F V_{ud})^2}{\pi^2} \int_{m_e}^{E_0} dE_e p_e E_e (E_0 - E_e)^2 \sum_{J, \kappa_e, \kappa_\nu} \frac{1}{2J_i + 1} \left| \sum_L \langle f || \Xi_{JL}(\kappa_e, \kappa_\nu) || i \rangle \right|^2, \quad (3)$$

where J_i is the angular momentum of the initial nuclear state and E_0 is the maximum energy of the electron. The nonzero integers κ_e and κ_ν represent the lepton angular momentum for the electron and neutrino, respectively, with

$$j_\kappa = |\kappa| - 1/2, \quad l_\kappa = \begin{cases} \kappa & (\kappa > 0) \\ -\kappa - 1 & (\kappa < 0) \end{cases}. \quad (4)$$

The bracket $\langle f || \Xi_{JL} || i \rangle$ denotes the reduced matrix element (RDME) [20] of the multipole operator Ξ_{JL} , which will be detailed in the next subsection.

2.1 Multipole operator

As shown in Eq. (12) of Ref. [8], the multipole operator Ξ_{JL} is written down as

$$\begin{aligned} \Xi_{JLM}(\kappa_e, \kappa_\nu) = S_{\kappa_e} \int d\mathbf{r} \\ \times \{ \mp Y_{JM}(\hat{r}) V_0(\mathbf{r}) \delta_{L,J} \Phi_{0JJ}(\kappa_e, \kappa_\nu; r) \pm i [Y_L(\hat{r}) \otimes \mathbf{V}(\mathbf{r})]_{JM} \phi_{1LJ}(\kappa_e, \kappa_\nu; r) \\ + i Y_{JM}(\hat{r}) A_0(\mathbf{r}) \delta_{L,J} \phi_{0JJ}(\kappa_e, \kappa_\nu; r) - [Y_L(\hat{r}) \otimes \mathbf{A}(\mathbf{r})]_{JM} \Phi_{1LJ}(\kappa_e, \kappa_\nu; r) \}, \end{aligned} \quad (5)$$

where the plus-minus signs correspond to the β^\mp decay. Here, $S_\kappa = \text{sgn}(\kappa)$ is the sign of κ and

$$\Phi_{KLJ}(\kappa_e, \kappa_\nu; r) = G_{\kappa_e}(r) g_{\kappa_\nu}(r) S_{KLJ}(\kappa_e, \kappa_\nu) + F_{\kappa_e}(r) f_{\kappa_\nu}(r) S_{KLJ}(-\kappa_e, -\kappa_\nu), \quad (6)$$

$$\phi_{KLJ}(\kappa_e, \kappa_\nu; r) = G_{\kappa_e}(r) f_{\kappa_\nu}(r) S_{KLJ}(\kappa_e, -\kappa_\nu) - F_{\kappa_e}(r) g_{\kappa_\nu}(r) S_{KLJ}(-\kappa_e, \kappa_\nu) \quad (7)$$

with

$$\begin{aligned} S_{KLJ}(\kappa', \kappa) = \sqrt{2(2j_\kappa + 1)(2j_{\kappa'} + 1)(2l_\kappa + 1)(2l_{\kappa'} + 1)(2K + 1)} \\ \times (l_\kappa, 0, l_{\kappa'}, 0 | L, 0) \begin{Bmatrix} l_{\kappa'} & 1/2 & j_{\kappa'} \\ l_\kappa & 1/2 & j_\kappa \\ L & K & J \end{Bmatrix}. \end{aligned} \quad (8)$$

Here $(j_1, m_1, j_2, m_2 | J, M)$ is the Clebsch–Gordan coefficient and the curly brackets denote the 9-j symbol [21, 22]. The radial wave functions of the emitted electron or the positron, $G_\kappa(r)$ and $F_\kappa(r)$, are obtained by solving the radial Dirac equation with the Coulomb potential $V_C(r)$ as

$$\frac{dG_\kappa(r)}{dr} + \frac{1 + \kappa}{r} G_\kappa(r) - (m_e + E_e - V_C(r)) F_\kappa(r) = 0, \quad (9)$$

$$\frac{dF_\kappa(r)}{dr} + \frac{1 - \kappa}{r} F_\kappa(r) - (m_e - E_e + V_C(r)) G_\kappa(r) = 0. \quad (10)$$

On the other hand, the neutrino is treated as a plane wave with the momentum $p_\nu = E_0 - E_e$, and its radial wave functions are given by $g_\kappa(r) = j_{l_\kappa}(p_\nu r)$ and $f_\kappa(r) = S_\kappa j_{l-\kappa}(p_\nu r)$, where $j_l(x)$ is the l th-order spherical Bessel function.

For a case of the angular momentum J with the parity $P = (-1)^{J+1}$ such as $J^P = 0^-, 1^+,$ and 2^- , the multipole operator is written as

$$\begin{aligned} \sum_L \Xi_{JLM}(\kappa_e, \kappa_\nu) = S_{\kappa_e} \int d\mathbf{r} \left\{ \pm i [Y_J(\hat{r}) \otimes \mathbf{V}(\mathbf{r})]_{JM} \phi_{1JJ}(\kappa_e, \kappa_\nu; r) \right. \\ \left. + i Y_{JM}(\hat{r}) A_0(\mathbf{r}) \phi_{0JJ}(\kappa_e, \kappa_\nu; r) \right. \\ \left. - \sum_{L=J\pm 1} [Y_L(\hat{r}) \otimes \mathbf{A}(\mathbf{r})]_{JM} \Phi_{1LJ}(\kappa_e, \kappa_\nu; r) \right\}. \quad (11) \end{aligned}$$

The spatial component of the vector current (1st term) is the ‘magnetic’ operator, while that of the axial-vector current (3rd term) is the ‘electric’ operator. The time component (2nd term) and the spatial component (3rd term) of the axial vector current contribute to the 0^- operator.

For a case of the angular momentum J with parity $P = (-1)^J$ such as $J^P = 0^+, 1^-,$ and 2^+ , the multipole operator is written as

$$\begin{aligned} \sum_L \Xi_{JLM}(\kappa_e, \kappa_\nu) = S_{\kappa_e} \int d\mathbf{r} \left\{ \mp Y_{JM}(\hat{r}) V_0(\mathbf{r}) \Phi_{0JJ}(\kappa_e, \kappa_\nu; r) \right. \\ \left. \pm i \sum_{L=J\pm 1} [Y_L(\hat{r}) \otimes \mathbf{V}(\mathbf{r})]_{JM} \phi_{1LJ}(\kappa_e, \kappa_\nu; r) \right. \\ \left. - [Y_J(\hat{r}) \otimes \mathbf{A}(\mathbf{r})]_{JM} \Phi_{1JJ}(\kappa_e, \kappa_\nu; r) \right\}. \quad (12) \end{aligned}$$

In this case, the spatial component of the vector current (2nd term) is the ‘electric’ operator, while the spatial component of the axial-vector current (3rd term) is the ‘magnetic’ operator. For the 0^+ operator, the time component (1st term) and the spatial component (2nd term) of the vector current contribute to the transition.

2.2 Longitudinal component of the vector current

The meson-exchange current (MEC) for the spatial component of the isovector vector current plays an important role, while the impulse approximation works well for the time component of the vector current in most cases. Since it is rather hard to construct many-body nuclear currents that satisfy the current conservation relation consistent with the nuclear wave function obtained from the effective nuclear potential, it is effective to rewrite a part of the spatial component of the vector current in terms of the time component of the vector current and use the impulse approximation. Here we focus on the second term of Eq. (12) for the ‘electric’ transition. The first term of Eq. (11) has nothing to do with the current conservation law.

For the nuclear beta decay, due to the Coulomb distortion of the electron wave function, it is not straightforward to implement the current conservation relation in contrast to the plane-wave formula for neutrino reaction and beta decay rate [16, 17]. Here we use the Bessel–Fourier transformation to take advantage of the momentum space representation. The relevant terms of the multipole operator are written as

$$\sum_{L=J\pm 1} [Y_L(\hat{r}) \otimes \mathbf{V}(\mathbf{r})]_{JM} \phi_{1LJ}(\kappa_e, \kappa_\nu; r) = \frac{2}{\pi} \int_0^\infty dq q^2 \mathcal{O}^J(\kappa_e, \kappa_\nu; q), \quad (13)$$

where

$$\mathcal{O}^J(\kappa_e, \kappa_\nu; q) = T^{JJ-1M}(q) \tilde{\phi}_{1J-1J}(\kappa_e, \kappa_\nu; q) + T^{JJ+1M}(q) \tilde{\phi}_{1J+1J}(\kappa_e, \kappa_\nu; q). \quad (14)$$

Here we used the identity,

$$\frac{2}{\pi} \int_0^\infty dq q^2 j_L(qr) j_L(qr') = \frac{\delta(r-r')}{r^2}. \quad (15)$$

The momentum-space multipole operator and lepton wave function are respectively defined as

$$T^{JLM}(q) = \int d\mathbf{r} j_L(qr) [Y_L(\hat{r}) \otimes \mathbf{V}(\mathbf{r})]_{JM}, \quad (16)$$

$$\tilde{\phi}_{1LJ}(\kappa_e, \kappa_\nu; q) = \int_0^\infty dr r^2 j_L(qr) \phi_{1LJ}(\kappa_e, \kappa_\nu; r). \quad (17)$$

Hereafter, $\tilde{\phi}_{1LJ}(\kappa_e, \kappa_\nu; q)$ is simplified as $\tilde{\phi}_L(q)$.

The electric T_E^{JM} and longitudinal T_L^{JM} multipole operators are defined by

$$\begin{aligned} T_E^{JM}(q) &= \frac{1}{q} \int d\mathbf{r} \nabla \times [j_J(qr) \mathbf{Y}_{JJM}(\hat{r})] \cdot \mathbf{V}(\mathbf{r}) \\ &= -i\sqrt{\frac{J}{2J+1}} T^{JJ+1M}(q) + i\sqrt{\frac{J+1}{2J+1}} T^{JJ-1M}(q), \end{aligned} \quad (18)$$

$$\begin{aligned} T_L^{JM}(q) &= \frac{i}{q} \int d\mathbf{r} \nabla [j_J(qr) Y_{JM}(\hat{r})] \cdot \mathbf{V}(\mathbf{r}) \\ &= i\sqrt{\frac{J+1}{2J+1}} T^{JJ+1M}(q) + i\sqrt{\frac{J}{2J+1}} T^{JJ-1M}(q). \end{aligned} \quad (19)$$

Note that in Ref. [16] T_E^{JM} and T_L^{JM} are called T_{JM}^{el} and L_{JM} , respectively. Using these longitudinal and electric multipole operators, Eq. (14) can be written as

$$\begin{aligned} i\mathcal{O}^J(\kappa_e, \kappa_\nu; q) &= T_E^{JM}(q) \left(\sqrt{\frac{J+1}{2J+1}} \tilde{\phi}_{J-1}(q) - \sqrt{\frac{J}{2J+1}} \tilde{\phi}_{J+1}(q) \right) \\ &\quad + T_L^{JM}(q) \left(\sqrt{\frac{J}{2J+1}} \tilde{\phi}_{J-1}(q) + \sqrt{\frac{J+1}{2J+1}} \tilde{\phi}_{J+1}(q) \right). \end{aligned} \quad (20)$$

We incorporate the current conservation relation in Eq. (14). The current conservation relation of the vector current of the charged current is expressed by

$$\nabla \cdot \mathbf{V} + i[H, V_0] = 0, \quad (21)$$

where the nuclear Hamiltonian H is assumed to be isospin invariant. To include the Coulomb interaction for protons and mass difference between proton and neutron, the extended Siegert's theorem has to be used. In the momentum space, the relation is expressed in terms of the longitudinal multipole operator $T_L^{JM}(q)$ and the Coulomb multipole operator $T_C^{JM}(q)$ as

$$qT_L^{JM}(q) + [H, T_C^{JM}(q)] = 0. \quad (22)$$

Here,

$$T_C^{JM}(q) = \int d\mathbf{r} j_J(qr) Y_{JM}(\hat{r}) V_0(\mathbf{r}), \quad (23)$$

which is called M_{JM} in Ref. [16]. When we take the nuclear matrix element of the operator, the commutator with the Hamiltonian is reduced to the energy difference between the final and initial nuclear states $\omega = E_f - E_i$, which corresponds to the maximum energy of electron

$E_0 = -\omega$. Then we obtain

$$T_L^{JM}(q) = -\frac{\omega}{q} T_C^{JM}(q). \quad (24)$$

Furthermore, the term $T^{JJ-1M}(q)$, which is the leading term of the electric multipole operator $T_E^{JM}(q)$ in the long-wave-length limit, can be written as

$$T^{JJ-1M}(q) = i\sqrt{\frac{2J+1}{J}} \frac{\omega}{q} T_C^{JM}(q) - \sqrt{\frac{J+1}{J}} T^{JJ+1M}(q). \quad (25)$$

Inserting Eq. (25) to Eq. (14), we get

$$\begin{aligned} i\mathcal{O}^J(\kappa_e, \kappa_\nu; q) &= -\sqrt{\frac{2J+1}{J}} \frac{\omega}{q} T_C^{JM}(q) \tilde{\phi}_{J-1}(q) \\ &\quad + iT^{JJ+1M}(q) \left(\tilde{\phi}_{J+1}(q) - \sqrt{\frac{J+1}{J}} \tilde{\phi}_{J-1}(q) \right), \end{aligned} \quad (26)$$

for $J \geq 1$.

For $J = 0$, \mathcal{O}^J can be expressed by the Coulomb form factor as

$$i\mathcal{O}^0(\kappa_e, \kappa_\nu; q) = -\frac{\omega}{q} T_C^{00}(q) \tilde{\phi}_1(q). \quad (27)$$

Finally, we transform the momentum space expression back to the coordinate space, using Eq. (15) and the following well-known identities:

$$\int_0^\infty dq q j_J(qr) j_{J-1}(qr') = \frac{\pi}{2} \frac{r'^{J-1}}{r^{J+1}} \theta(r-r'), \quad (28)$$

$$\int_0^\infty dq q^2 j_{J+1}(qr) j_{J-1}(qr') = \frac{\pi}{2} \left[(2J+1) \frac{r'^{J-1}}{r^{J+2}} \theta(r-r') - \frac{\delta(r-r')}{r^2} \right]. \quad (29)$$

The multipole operator for $P = (-1)^J$ ($J \geq 1$) is given as

$$\begin{aligned} \sum_L \Xi_{JLM}(\kappa_e, \kappa_\nu) &= S_{\kappa_e} \int d\mathbf{r} \left\{ \mp Y_{JM}(\hat{r}) V_0(\mathbf{r}) \left(\Phi_{0JJ}(\kappa_e, \kappa_\nu; r) + \omega \sqrt{\frac{2J+1}{J}} \frac{\varphi_J(r)}{r^{J+1}} \right) \right. \\ &\quad \pm i[Y_{J+1}(\hat{r}) \otimes \mathbf{V}(\mathbf{r})]_{JM} \left[\phi_{1J+1J}(\kappa_e, \kappa_\nu; r) + \sqrt{\frac{J+1}{J}} \left(\phi_{1J-1J}(\kappa_e, \kappa_\nu; r) - (2J+1) \frac{\varphi_J(r)}{r^{J+2}} \right) \right] \\ &\quad \left. - [Y_J(\hat{r}) \otimes \mathbf{A}(\mathbf{r})]_{JM} \Phi_{1JJ}(\kappa_e, \kappa_\nu; r) \right\}, \end{aligned} \quad (30)$$

where

$$\varphi_J(r) = \int_0^r dr' (r')^{J+1} \phi_{1J-1J}(\kappa_e, \kappa_\nu; r'). \quad (31)$$

The magnetization current, $\mathbf{V} \sim \nabla \times \boldsymbol{\mu}$, is purely transverse, which gives $T_L^{JM}(q) = 0$ and $T^{JJ-1M}(q) = -\sqrt{(J+1)/J}T^{JJ+1M}(q)$. The relation shows that the expression for the magnetization current in Eq. (12) is equivalent to that of Eq. (30).

For $J = 0$, the multipole operator is written as

$$\Xi_{000}(\kappa_e, \kappa_\nu) = \mp S_{\kappa_e} \int d\mathbf{r} Y_{00}(\hat{r}) V_0(\mathbf{r}) \varphi_0(r) \quad (32)$$

with

$$\varphi_0(r) = \Phi_{000}(\kappa_e, \kappa_\nu; r) + \omega \int_r^\infty dr' \phi_{110}(\kappa_e, \kappa_\nu; r'). \quad (33)$$

The second term of Eq. (33) is the contribution of the longitudinal term of the spatial component of the vector current.

For the transition to the isobaric analog state (IAS), $\omega = 0$ assuming the isospin invariance of the nuclear Hamiltonian and we obtain

$$\varphi_0(r) = \Phi_{000}(\kappa_e, \kappa_\nu; r). \quad (34)$$

The formula shows that the decay rate is given by the time component of the vector current alone. There is no need to include the correction originating from the spatial component of the vector current.

The 0^+ transition to the non-IAS states is weak compared to the transition to the IAS because of the orthogonality relation $\langle f | \int d\mathbf{r} V^0 | i \rangle = \langle f | T^\pm | i \rangle = 0$. For practical computations, it is convenient to rewrite Eq. (33) using this orthogonality relation. For the transition to the non-IAS, we can use $\varphi_0(r)$ subtracted by a constant value as

$$\varphi_0(r) \rightarrow \varphi_0(r) - \varphi_0(0). \quad (35)$$

Thus, we obtain

$$\varphi_0(r) = \Phi_{000}(\kappa_e, \kappa_\nu; r) - \Phi_{000}(\kappa_e, \kappa_\nu; 0) - \omega \int_0^r dr' \phi_{110}(\kappa_e, \kappa_\nu; r'). \quad (36)$$

An advantage of Eq. (36) is that $\varphi_0(r)$ is an operator of the form of ‘second-forbidden transition’ starting from $\mathcal{O}(r^2)$ and the integration of r to $r \rightarrow \infty$ of the ω term is avoided.

3 Impulse approximation

In this section, we present explicit expressions of the multipole operator in the impulse approximation of the nuclear current. For the axial-vector current \mathcal{A} , the space and time

components of the one-body charged-current operators in non-relativistic approximation are respectively given as

$$\mathbf{A}(\mathbf{r}) = \sum_{j=1}^A \delta(\mathbf{r} - \mathbf{r}_j) \left[g_A(q^2) \boldsymbol{\sigma}_j - g_P(q^2) \frac{\mathbf{q}(\boldsymbol{\sigma}_j \cdot \mathbf{q})}{2M} \right] \tau_j^\pm, \quad (37)$$

$$A_0(\mathbf{r}) = \sum_{j=1}^A \delta(\mathbf{r} - \mathbf{r}_j) \left[g_A(q^2) \frac{\boldsymbol{\sigma}_j \cdot (\mathbf{p}'_j + \mathbf{p}_j)}{2M} - g_P(q^2) \frac{\omega(\boldsymbol{\sigma}_j \cdot \mathbf{q})}{2M} \right] \tau_j^\pm, \quad (38)$$

where \mathbf{p}'_j and \mathbf{p}_j are the momentum operators of the j th nucleon for the final and initial states, respectively, and \mathbf{q} is the momentum transfer from the lepton, $\mathbf{q} = \mathbf{p}'_j - \mathbf{p}_j$. Here the nucleon form factors $g_A(q^2)$ and $g_P(q^2)$ are functions of momentum transfer q^2 to the nucleon. In the nuclear beta decay, the use of the form factors at $q^2 = 0$ is a good approximation: $g_A(0) = 1.27$ and $g_P(0) = g_A(0)2M/m_\pi^2$ assuming the pion-pole dominance [23].

For the vector current \mathcal{V} , the operators are given as

$$\mathbf{V}(\mathbf{r}) = \sum_{j=1}^A \delta(\mathbf{r} - \mathbf{r}_j) \left[g_V(q^2) \frac{(\mathbf{p}'_j + \mathbf{p}_j)}{2M} + \frac{g_V(q^2) + g_M(q^2)}{2M} i \boldsymbol{\sigma}_j \times \mathbf{q} \right] \tau_j^\pm, \quad (39)$$

$$V_0(\mathbf{r}) = \sum_{j=1}^A \delta(\mathbf{r} - \mathbf{r}_j) g_V(q^2) \tau_j^\pm, \quad (40)$$

where $g_V(0) = 1$, $g_V(0) + g_M(0) = \mu_p - \mu_n = 4.7$. Hereafter, we simply write the form factors at $q^2 = 0$ as g_A, g_V, g_M and g_P .

The $g_A \boldsymbol{\sigma} \tau^\pm$ term of the spatial component of the axial-vector current and $g_V \tau^\pm$ term of the time component of the vector current are main contributors remaining in the non-relativistic limit of the nucleon current. The $g_A \boldsymbol{\sigma} \cdot (\mathbf{p} + \mathbf{p}')/2M$ term of the time component of the axial-vector current and the $g_V(\mathbf{p} + \mathbf{p}')/2M$ term of the spatial component of the vector current (convection current), are velocity-dependent (v/c) correction terms. The $(g_V + g_M)/(2M)$ term is the induced weak-magnetism term.

3.1 Case of $P = (-1)^{J+1}$

In the impulse approximation, the multipole operator for $P = (-1)^{J+1}$ is given as

$$\begin{aligned} \sum_L \Xi_{JLM}(\kappa_e, \kappa_\nu) &= S_{\kappa_e} \int d\mathbf{r} \\ &\times \left[\sum_{j=1}^A \tau_j^\pm \delta(\mathbf{r} - \mathbf{r}_j) \left\{ [Y_{J-1}(\hat{r}) \otimes \boldsymbol{\sigma}_j]_{JM} \phi_a(r) + [Y_{J+1}(\hat{r}) \otimes \boldsymbol{\sigma}_j]_{JM} \phi_b(r) \right\} \right] \end{aligned}$$

$$+ Y_{JM} \boldsymbol{\sigma}_j \cdot \nabla_j \phi_c(r) + [Y_J(\hat{r}) \otimes \nabla_j]_{JM} \phi_d(r) \Big\} , \quad (41)$$

with

$$\begin{aligned} \phi_a(r) &= -g_A \Phi_{1J-1J}(r) + \frac{1}{2M} \sqrt{\frac{J}{2J+1}} D_-^J [g_A \phi_{0JJ}(r) + g_P m_e \bar{\phi}_{0JJ}(r)] \\ &\mp \frac{g_V + g_M}{2M} \sqrt{\frac{J+1}{2J+1}} D_-^J \phi_{1JJ}(r), \end{aligned} \quad (42)$$

$$\begin{aligned} \phi_b(r) &= -g_A \Phi_{1J+1J}(r) - \frac{1}{2M} \sqrt{\frac{J+1}{2J+1}} D_+^J [g_A \phi_{0JJ}(r) + g_P m_e \bar{\phi}_{0JJ}(r)] \\ &\mp \frac{g_V + g_M}{2M} \sqrt{\frac{J}{2J+1}} D_+^J \phi_{1JJ}(r), \end{aligned} \quad (43)$$

$$\phi_c(r) = \frac{g_A}{M} \phi_{0JJ}, \quad (44)$$

$$\phi_d(r) = \pm \frac{g_V}{M} \phi_{1JJ}(r), \quad (45)$$

where

$$D_+^J = \frac{d}{dr} - \frac{J}{r}, \quad (46)$$

$$D_-^J = \frac{d}{dr} + \frac{J+1}{r}, \quad (47)$$

$$\bar{\phi}_{0JJ}(r) = G_{\kappa_e}(r) f_{\kappa_\nu}(r) S_{0JJ}(\kappa_e, -\kappa_\nu) + F_{\kappa_e}(r) g_{\kappa_\nu}(r) S_{0JJ}(-\kappa_e, \kappa_\nu). \quad (48)$$

The simplified notations $\Phi_{KLS}(r)$ and $\phi_{KLS}(r)$ denoting $\Phi_{KLS}(\kappa_e, \kappa_\nu; r)$ and $\phi_{KLS}(\kappa_e, \kappa_\nu; r)$ respectively are used.

The first terms of ϕ_a (Eq. (42)) and ϕ_b (Eq. (43)) are contributions of the spatial component of the axial-vector current, while the g_A/M terms of ϕ_a , ϕ_b and ϕ_c (Eq. (44)) are contributions of the time component of the axial vector current. The last terms, $(g_V + g_M)/(2M)$, of ϕ_a and ϕ_b are the weak-magnetism terms. The ϕ_d (Eq. (45)) is the contribution of the convection current.

The Dirac equation, Eqs. (9) and (10), is used to derive the g_P term in Eqs. (42) and (43). Here only m_e remains and the Coulomb potential $V_C(r)$ disappears due to the gauge invariance [24]. The g_P term is usually neglected for the nuclear beta decay [11] because the contributions of the induced pseudo-scalar term g_P are about $\mathcal{O}(m_e E_0/m_\pi^2)$, typically $\sim 10^{-3}$, smaller than those of the leading g_A terms of ϕ_a and ϕ_b .

3.2 Case of $P = (-1)^J$

In the impulse approximation, the multipole operator for $P = (-1)^J$ is given as

$$\begin{aligned} \sum_L \Xi_{JLM}(\kappa_e, \kappa_\nu) &= S_{\kappa_e} \int d\mathbf{r} \\ &\times \left[\sum_{j=1}^A \tau_j^\pm \delta(\mathbf{r} - \mathbf{r}_j) \left\{ Y_{JM}(\hat{r}) \phi_A(r) + [Y_J(\hat{r}) \otimes \boldsymbol{\sigma}_j]_{JM} \phi_B(r) \right. \right. \\ &\left. \left. + [Y_{J+1}(\hat{r}) \otimes \boldsymbol{\nabla}_j]_{JM} \phi_C(r) \right\} \right] \end{aligned} \quad (49)$$

with

$$\phi_A(r) = \mp g_V \left(\Phi_{0JJ}(r) + \omega \sqrt{\frac{2J+1}{J}} \frac{\varphi_J(r)}{r^{J+1}} - \frac{1}{2M} \sqrt{\frac{J+1}{J}} \psi_J(r) \right), \quad (50)$$

$$\phi_B(r) = -g_A \Phi_{1JJ}(r) \mp \frac{g_V + g_M}{2M} \psi_J(r), \quad (51)$$

$$\phi_C(r) = \pm \frac{g_V}{M} \left\{ \phi_{1J+1J}(r) + \sqrt{\frac{J+1}{J}} \left[\phi_{1J-1J}(r) - (2J+1) \frac{\varphi_J(r)}{r^{J+2}} \right] \right\}, \quad (52)$$

and

$$\psi_J(r) = \sqrt{\frac{J+1}{2J+1}} D_+^{J-1} \phi_{1J-1J}(r) + \sqrt{\frac{J}{2J+1}} D_-^{J+1} \phi_{1J+1J}(r). \quad (53)$$

The first terms of ϕ_A (Eq. (50)) and ϕ_B (Eq. (51)) are the contribution of the time component of the vector current and the space component of the axial-vector current, respectively. The second term of ϕ_B is the weak-magnetism term. The second and third terms of ϕ_A and the ϕ_C (Eq. (52)) are contributions of the convection current. The second term of ϕ_A , which is proportional to ω , is usually taken into account in the leading order approximation of the φ_J .

4 Explicit formulas and applications

In this section, we give explicit formulas for the allowed $J^P = 1^+$ and the first-forbidden $J^P = 0^-, 1^-,$ and 2^- transitions useful in practical nuclear-structure calculations. We include all the recoil order currents $\mathcal{O}(p/M)$ that are considered as higher-order corrections to the Fermi and Gamow–Teller operators. We show that these formulas are reduced into the widely-used leading-order formula by Behrens–Bühring (LOB)[12] with the approximation of the electron and neutrino wave functions described in Eqs. (60)–(65) of paper (I). We then

apply our formulas to the beta-decay rate of ^{160}Sn , where the first-forbidden transitions, in particular the $J^P = 1^-$ transition, are expected to give a sizable contribution to the total beta-decay rate [25].

The RDME of the multipole operator is given by the radial integral of the transition densities together with the lepton wave functions. For $P = (-1)^{J+1}$, the RDME is given as

$$\begin{aligned} \langle f || \sum_L \Xi_{JL}(\kappa_e, \kappa_\nu) || i \rangle &= S_{\kappa_e} \int_0^\infty dr r^2 \\ &\times \left[\rho_{J-1J}^\sigma(r) \phi_a(r) + \rho_{J+1J}^\sigma(r) \phi_b(r) + \rho_J^{\sigma\nabla}(r) \phi_c(r) + \rho_{JJ}^{\nabla}(r) \phi_d(r) \right], \end{aligned} \quad (54)$$

and for $P = (-1)^J$

$$\langle f || \sum_L \Xi_{JL}(\kappa_e, \kappa_\nu) || i \rangle = S_{\kappa_e} \int_0^\infty dr r^2 \left[\rho_J(r) \phi_A(r) + \rho_{JJ}^\sigma(r) \phi_B(r) + \rho_{J+1J}^{\nabla}(r) \phi_C(r) \right], \quad (55)$$

where the transition densities used in the impulse approximation are defined as

$$\rho_J(r) = \langle f || \sum_{j=1}^A \int d\Omega_r \delta(\mathbf{r} - \mathbf{r}_j) \tau_j^\pm Y_J(\hat{r}) || i \rangle, \quad (56)$$

$$\rho_{LJ}^\sigma(r) = \langle f || \sum_{j=1}^A \int d\Omega_r \delta(\mathbf{r} - \mathbf{r}_j) \tau_j^\pm [Y_L(\hat{r}) \otimes \boldsymbol{\sigma}_j]_J || i \rangle, \quad (57)$$

$$\rho_{LJ}^{\nabla}(r) = \langle f || \sum_{j=1}^A \int d\Omega_r \delta(\mathbf{r} - \mathbf{r}_j) \tau_j^\pm [Y_L(\hat{r}) \otimes \boldsymbol{\nabla}_j]_J || i \rangle, \quad (58)$$

$$\rho_J^{\sigma\nabla}(r) = \langle f || \sum_{j=1}^A \int d\Omega_r \delta(\mathbf{r} - \mathbf{r}_j) \tau_j^\pm Y_J(\hat{r}) \boldsymbol{\sigma}_j \cdot \boldsymbol{\nabla}_j || i \rangle. \quad (59)$$

Here, $\int d\Omega_r$ denotes the integration over solid angles. We employ a nuclear EDF method to generate these transition densities. As the details can be found in Ref. [26], here we recapitulate the numerical procedures relevant to the present analysis. In the framework of the nuclear EDF method, the ground state of a mother nucleus is described by solving the Kohn–Sham–Bogoliubov (KSB) equation [27]. We solve the KSB equation in cylindrical coordinates assuming the axial symmetry [28]. Then, the transitions to a daughter nucleus are described by the proton–neutron quasiparticle-random-phase approximation (pnQRPA).

The residual interactions entering into the pnQRPA equation are given by the EDF self-consistently. The EDF consists of the Skyrme SLy4 functional [29] and the Yamagami–Shimizu–Nakatsukasa functional [30] for the particle-hole and pairing energy, respectively. Since we assume the axial symmetry, the z -component of the total angular momentum $J_z = K$ is a good quantum number.

In the actual computations, we evaluate the following transition densities:

$$\delta\rho^1(\mathbf{r}) := \langle f | \sum_{\sigma, \sigma', \tau, \tau'} \psi^\dagger(\mathbf{r}\sigma'\tau') \langle \sigma' | 1 | \sigma \rangle \langle \tau' | \tau^\pm | \tau \rangle \psi(\mathbf{r}\sigma\tau) | i \rangle, \quad (60)$$

$$\delta\rho_{\mu\sigma}^\sigma(\mathbf{r}) := \langle f | \sum_{\sigma, \sigma', \tau, \tau'} \psi^\dagger(\mathbf{r}\sigma'\tau') \langle \sigma' | \sigma_{\mu\sigma} | \sigma \rangle \langle \tau' | \tau^\pm | \tau \rangle \psi(\mathbf{r}\sigma\tau) | i \rangle, \quad (61)$$

$$\delta\rho_{\mu r}^\nabla(\mathbf{r}) := \langle f | \sum_{\sigma, \sigma', \tau, \tau'} \psi^\dagger(\mathbf{r}\sigma'\tau') \langle \sigma' | 1 | \sigma \rangle \langle \tau' | \tau^\pm | \tau \rangle \nabla_{\mu r} \psi(\mathbf{r}\sigma\tau) | i \rangle, \quad (62)$$

$$\begin{aligned} \delta\rho^{\sigma\nabla}(\mathbf{r}) := & \langle f | \sum_{\sigma, \sigma', \tau, \tau'} \sum_{\mu\sigma, \mu r} \psi^\dagger(\mathbf{r}\sigma'\tau') \langle \sigma' | \sigma_{\mu\sigma} | \sigma \rangle \langle \tau' | \tau^\pm | \tau \rangle \nabla_{\mu r} \psi(\mathbf{r}\sigma\tau) \\ & \times (1, \mu\sigma, 1, \mu r | 0, 0) | i \rangle, \end{aligned} \quad (63)$$

where $\sigma_{\mu\sigma}$ and $\nabla_{\mu r}$ are spherical components of the Pauli matrix $\boldsymbol{\sigma}$ and the operator ∇ , respectively. With these, we can evaluate the transition densities appearing in the RDMs as

$$\rho_{JK}(r) = \int d\Omega_r \delta\rho^1(\mathbf{r}) Y_{JK}(\hat{r}), \quad (64)$$

$$\rho_{LJK}^\sigma(r) = \int d\Omega_r \sum_{\mu\sigma, M} \delta\rho_{\mu\sigma}^\sigma(\mathbf{r}) Y_{LM}(\hat{r}) (L, M, 1, \mu\sigma | J, K), \quad (65)$$

$$\rho_{LJK}^\nabla(r) = \int d\Omega_r \sum_{\mu r, M} \delta\rho_{\mu r}^\nabla(\mathbf{r}) Y_{LM}(\hat{r}) (L, M, 1, \mu r | J, K), \quad (66)$$

$$\rho_{JK}^{\sigma\nabla}(r) = \int d\Omega_r \delta\rho^{\sigma\nabla}(\mathbf{r}) Y_{JK}(\hat{r}). \quad (67)$$

For spherical nuclei, we only need to evaluate the transition densities for $K = 0$. Then, one obtains the transition densities as

$$\rho_J(r) = \sqrt{2J+1} \rho_{J0}(r), \quad (68)$$

$$\rho_{LJ}^\sigma(r) = \sqrt{2J+1} \rho_{LJ0}^\sigma(r), \quad (69)$$

$$\rho_{LJ}^\nabla(r) = \sqrt{2J+1} \rho_{LJ0}^\nabla(r), \quad (70)$$

$$\rho_J^{\sigma\nabla}(r) = \sqrt{2J+1} \rho_{J0}^{\sigma\nabla}(r) \quad (71)$$

for any L and J .

4.1 Case for $J^P = 1^+$

An explicit formula of the multipole operator for the allowed Gamow–Teller transition in the impulse approximation is given as

$$\langle f || \sum_L \Xi_{1L}(\kappa_e, \kappa_\nu) || i \rangle = S_{\kappa_e} \int_0^\infty dr r^2 \times \left[\rho_{01}^\sigma(r) \phi_a(r) + \rho_{21}^\sigma(r) \phi_b(r) + \rho_1^{\sigma\nabla}(r) \phi_c(r) + \rho_{11}^{\nabla}(r) \phi_d(r) \right], \quad (72)$$

where

$$\begin{aligned} \phi_a(r) &= -g_A \Phi_{101}(r) + \frac{1}{2M} \sqrt{\frac{1}{3}} \left(\frac{d}{dr} + \frac{2}{r} \right) [g_A \phi_{011}(r) + g_P m_e \bar{\phi}_{011}(r)] \\ &\mp \frac{g_V + g_M}{2M} \sqrt{\frac{2}{3}} \left(\frac{d}{dr} + \frac{2}{r} \right) \phi_{111}(r), \end{aligned} \quad (73)$$

$$\begin{aligned} \phi_b(r) &= -g_A \Phi_{121}(r) - \frac{1}{2M} \sqrt{\frac{2}{3}} \left(\frac{d}{dr} - \frac{1}{r} \right) [g_A \phi_{011}(r) + g_P m_e \bar{\phi}_{011}(r)] \\ &\mp \frac{g_V + g_M}{2M} \sqrt{\frac{1}{3}} \left(\frac{d}{dr} - \frac{1}{r} \right) \phi_{111}(r), \end{aligned} \quad (74)$$

$$\phi_c(r) = \frac{g_A}{M} \phi_{011}(r), \quad (75)$$

$$\phi_d(r) = \pm \frac{g_V}{M} \phi_{111}(r). \quad (76)$$

In LOB, ϕ_a for $(\kappa_e, \kappa_\nu) = (-1, -1)$ and $(1, 1)$ is non zero and is given by

$$\phi_a(r) = -\sqrt{2} g_A \alpha_{\kappa_e}. \quad (77)$$

The spectrum shape $C(E_e)$ is given as

$$(2J_i + 1) F(Z, E_e) C(E_e) = (\alpha_{-1}^2 + \alpha_1^2) |\langle f || \sum_{j=1}^A g_A \tau_j^\pm \sigma_j || i \rangle|^2 \quad (78)$$

$$= 4\pi (\alpha_{-1}^2 + \alpha_1^2) \left| g_A \int_0^\infty dr r^2 \rho_{01}^\sigma(r) \right|^2. \quad (79)$$

Note that $\alpha_{-1}^2 + \alpha_1^2$ is often replaced with the Fermi function evaluated at the nuclear surface [31].

4.2 Case for $J^P = 0^-$

For the $J = 0$ with the parity change $P = -1$ transition (0^-), only the axial-vector current can contribute. In the impulse approximation, the RDME of the multipole operators is given by the transition densities ρ_{01}^σ and $\rho_0^{\sigma\nabla}$, which respectively come from the

space $g_A \boldsymbol{\sigma} \cdot \mathbf{r}$ and time $g_A \boldsymbol{\sigma} \cdot \nabla / M$ components of the axial-vector current. The latter is velocity dependent. Their contributions will be equally important as they are estimated by $\mathcal{O}(\langle p_N \rangle / M) \approx 200 \text{ MeV} / 1 \text{ GeV}$ and $\mathcal{O}(E_e R_A)$, respectively.

The multipole operator for $J^P = 0^-$ is given as

$$\langle f || \sum_L \Xi_{0L}(\kappa_e, \kappa_\nu) || i \rangle = S_{\kappa_e} \int_0^\infty dr r^2 \left[\rho_{10}^\sigma(r) \phi_b(r) + \rho_0^{\sigma\nabla}(r) \phi_c(r) \right], \quad (80)$$

where ϕ_b and ϕ_c of Eq. (41) are given as

$$\phi_b(r) = -g_A \Phi_{110}(r) - \frac{1}{2M} \frac{d}{dr} \left[g_A \phi_{000}(r) + g_P m_e \bar{\phi}_{000}(r) \right], \quad (81)$$

$$\phi_c(r) = \frac{g_A}{M} \phi_{000}(r). \quad (82)$$

In LOB, two partial waves with $(\kappa_e, \kappa_\nu) = (-1, 1), (1, -1)$ can contribute. Then ϕ_b and ϕ_c are expressed as

$$\phi_b(r) = -\sqrt{2} S_{\kappa_e} \alpha_{\kappa_e} \frac{g_A}{3} \left(E_0 + \frac{3V_{D1}(r)}{r} + S_{\kappa_e} m_e \right) r, \quad (83)$$

$$\phi_c(r) = -\sqrt{2} S_{\kappa_e} \alpha_{\kappa_e} \frac{g_A}{M}, \quad (84)$$

with $V_{D1}(r)$ defined in Eq. (42) of Ref. [8],

$$V_{D1}(r) = - \int_0^r \left(\frac{r'}{r} \right)^{2k} V_C(r') dr'. \quad (85)$$

In order to compare with LOB for the first-forbidden transition, we rewrite the RDMEs using the parametrization of Ref. [11] summarized in the appendix. The RDME in LOB is written as

$$\sqrt{2\pi} \langle f || \sum_L \Xi_{0L}(\kappa_e, \kappa_\nu) || i \rangle = -\alpha_{\kappa_e} \left(\zeta_0 + S_{\kappa_e} \frac{m_e w}{3} \right). \quad (86)$$

Here ζ_0 and w are the nuclear matrix elements defined in the appendix. The spectrum shape is given as

$$(2J_i + 1) F(Z, E_e) C(E_e) = \alpha_{-1}^2 \left(\zeta_0 - \frac{m_e w}{3} \right)^2 + \alpha_1^2 \left(\zeta_0 + \frac{m_e w}{3} \right)^2, \quad (87)$$

which is in agreement with Ref. [11]. We note that this formula is often used with the following approximations

$$\alpha_{-1}^2 + \alpha_1^2 \sim F(Z, E_e), \quad (88)$$

$$\lambda_2, \mu_1 \sim 1. \quad (89)$$

The accuracy of the above treatment depends on Z and E_e . Deviation from the approximations of Eqs. (88) and (89) can be very large for large Z [8, 32].

4.3 Case for $J^P = 1^-$

The operators for the $J = 1$ and parity $P = -1$ transition include the contribution of V_0 , \mathbf{V} , and \mathbf{A} . The multipole operator in the impulse approximation is given by

$$\langle f || \sum_L \Xi_{1L}(\kappa_e, \kappa_\nu) || i \rangle = S_{\kappa_e} \int_0^\infty dr r^2 \left[\rho_1(r) \phi_A(r) + \rho_{11}^\sigma(r) \phi_B(r) + \rho_{21}^\nabla(r) \phi_C(r) \right] \quad (90)$$

with

$$\phi_A(r) = \mp g_V \left[\Phi_{011}(r) + \sqrt{3}\omega \frac{\varphi_1(r)}{r^2} - \frac{\sqrt{2}}{2M} \psi_1(r) \right], \quad (91)$$

$$\phi_B(r) = -g_A \Phi_{111}(r) \mp \frac{g_V + g_M}{2M} \psi_1(r), \quad (92)$$

$$\phi_C(r) = \pm \frac{g_V}{M} \left\{ \phi_{121}(r) + \sqrt{2} \left[\phi_{101}(r) - 3 \frac{\varphi_1(r)}{r^3} \right] \right\}, \quad (93)$$

where

$$\psi_1(r) = \sqrt{\frac{2}{3}} \frac{d}{dr} \phi_{101}(r) + \sqrt{\frac{1}{3}} \left(\frac{d}{dr} + \frac{3}{r} \right) \phi_{121}(r). \quad (94)$$

When we evaluate the convection current without using the current conservation relation, the multipole operator Ξ is given as

$$\langle f || \sum_L \Xi_{1L}(\kappa_e, \kappa_\nu) || i \rangle = S_{\kappa_e} \int_0^\infty dr r^2 \times \left[\rho_1(r) \phi_{A'}(r) + \rho_{11}^\sigma(r) \phi_B(r) + \rho_{21}^\nabla(r) \phi_{C'}(r) + \rho_{01}^\nabla(r) \phi_D(r) \right], \quad (95)$$

where the leading-order convection current term ρ_{01}^∇ is given together with the lepton wave function

$$\phi_D(r) = \pm \frac{g_V}{M} \phi_{101}(r), \quad (96)$$

and $\phi_{A'}$ and $\phi_{C'}$ are:

$$\phi_{A'}(r) = \mp g_V \left[\Phi_{011}(r) - \frac{1}{2M} \psi_1'(r) \right], \quad (97)$$

$$\phi_{C'}(r) = \pm \frac{g_V}{M} \phi_{121}(r), \quad (98)$$

where

$$\psi_1'(r) = \sqrt{\frac{1}{3}} \frac{d}{dr} \phi_{101}(r) - \sqrt{\frac{2}{3}} \left(\frac{d}{dr} + \frac{3}{r} \right) \phi_{121}(r). \quad (99)$$

In LOB, lepton wave functions are approximated as explained in I and the contributions of $\mathcal{O}(1/M)$ terms are neglected. The RDME is given by the dipole ρ_1 and spin-dipole ρ_{11}^σ

terms as

$$\langle f || \sum_L \Xi_{1L}(\kappa_e, \kappa_\nu) || i \rangle = S_{\kappa_e} \int_0^\infty dr r^2 [\rho_1(r)\phi_A(r) + \rho_{11}^\sigma(r)\phi_B(r)]. \quad (100)$$

Here, $\rho_1\phi_A$ and $\rho_{11}^\sigma\phi_B$ terms correspond to $g_V\mathbf{r}$ and $g_A\mathbf{r} \times \boldsymbol{\sigma}$, respectively. Six partial waves with $(\kappa_e, \kappa_\nu) = (\mp 1, \pm 1), (\mp 1, \mp 2), (\mp 2, \mp 1)$ contribute. For the partial waves with $(\kappa_e, \kappa_\nu) = (-1, -2), (1, 2), (-2, -1)$, and $(2, 1)$, ϕ_A and ϕ_B are given by

$$\phi_A(r) = \mp g_V \alpha_{\kappa_e} \frac{2\sqrt{3}}{9} (p_\nu \delta_{|\kappa_e|,1} + p_e \delta_{|\kappa_e|,2}) r, \quad (101)$$

$$\phi_B(r) = g_A \alpha_{\kappa_e} \frac{\sqrt{6}}{9} (p_\nu \delta_{|\kappa_e|,1} - p_e \delta_{|\kappa_e|,2}) r. \quad (102)$$

The RDME can be written as

$$\sqrt{2\pi} \langle f || \sum_L \Xi_{1L}(\kappa_e, \kappa_\nu) || i \rangle = S_{\kappa_e} \alpha_{\kappa_e} \frac{1}{3\sqrt{2}} [(2x+u)p_\nu \delta_{|\kappa_e|,1} + (2x-u)p_e \delta_{|\kappa_e|,2}]. \quad (103)$$

Here, x and u are defined in the appendix.

For the partial waves with $(\kappa_e, \kappa_\nu) = (-1, 1)$ and $(1, -1)$, ϕ_A and ϕ_B are given as

$$\phi_A(r) = \pm g_V S_{\kappa_e} \alpha_{\kappa_e} \frac{\sqrt{6}}{9} \left(E_0 + \frac{3V_{D1}}{r} + 3\omega + S_{\kappa_e} m_e \right) r, \quad (104)$$

$$\phi_B(r) = g_A S_{\kappa_e} \alpha_{\kappa_e} \frac{2\sqrt{3}}{9} \left(E_0 - 2E_e - \frac{3V_{D1}}{r} - S_{\kappa_e} m_e \right) r, \quad (105)$$

and

$$\sqrt{2\pi} \langle f || \sum_L \Xi_{1L}(\kappa_e, \kappa_\nu) || i \rangle = \alpha_{\kappa_e} \left[\zeta_1 - \frac{2}{3} u E_e - S_{\kappa_e} \frac{m_e}{3} (x+u) \right]. \quad (106)$$

From the Siegert's theorem, the RDME of the convection current is written in terms of that of the dipole operator; the following relation is used in the above formula

$$\xi' y \rightarrow -\omega x. \quad (107)$$

Here ζ_1 and $\xi' y$ are defined in the appendix.

Finally, the spectrum shape is given as

$$(2J_i + 1)F(E_e, Z)C(E) = (\alpha_{-1}^2 + \alpha_1^2) \left[\frac{(2x+u)^2 p_\nu^2}{18} + \lambda_2 \frac{(2x-u)^2 p_e^2}{18} \right] + \sum_{\kappa_e = \mp 1} \alpha_{\kappa_e}^2 \left[\zeta_1 - \frac{2}{3} u E_e - S_{\kappa_e} \frac{m_e}{3} (x+u) \right]^2. \quad (108)$$

The above expression agrees with that given in Ref. [11].

4.4 Case for $J^P = 2^-$

The 2^- transition offers a unique first-forbidden transition, where only the single spin-dipole operator $[Y_1 \otimes \mathbf{A}]_2(\rho_{12}^\sigma)$ contributes in LOB. Due to this property, the 2^- transition has been used for a precise evaluation of the beta-ray spectrum in BSM searches. However, by including NLO, we have an additional contribution of the weak magnetism, $[Y_3 \otimes \mathbf{A}]_2(\rho_{32}^\sigma)$, the time component of the axial-vector current $Y_2 A_0(\rho_2^{\sigma\nabla})$, and the vector current $[Y_2 \otimes \mathbf{V}]_2(\rho_{22}^{\nabla})$. In the impulse approximation, the multipole operator is given by

$$\begin{aligned} \langle f || \sum_L \Xi_{2L}(\kappa_e, \kappa_\nu) || i \rangle &= S_{\kappa_e} \int_0^\infty dr r^2 \\ &\times \left[\rho_{12}^\sigma(r) \phi_a(r) + \rho_{32}^\sigma(r) \phi_b(r) + \rho_2^{\sigma\nabla}(r) \phi_c(r) + \rho_{22}^{\nabla}(r) \phi_d(r) \right] \end{aligned} \quad (109)$$

with

$$\begin{aligned} \phi_a(r) &= -g_A \Phi_{112}(r) + \frac{1}{2M} \sqrt{\frac{2}{5}} D_-^2 (g_A \phi_{022}(r) + g_P m_e \bar{\phi}_{022}(r)) \\ &\mp \frac{g_V + g_M}{2M} \sqrt{\frac{3}{5}} D_-^2 \phi_{122}(r), \end{aligned} \quad (110)$$

$$\begin{aligned} \phi_b(r) &= -g_A \Phi_{132}(r) - \frac{1}{2M} \sqrt{\frac{3}{5}} D_+^2 (g_A \phi_{022}(r) + g_P m_e \bar{\phi}_{022}(r)) \\ &\mp \frac{g_V + g_M}{2M} \sqrt{\frac{2}{5}} D_+^2 \phi_{122}(r), \end{aligned} \quad (111)$$

$$\phi_c(r) = \frac{g_A}{M} \phi_{022}(r), \quad (112)$$

$$\phi_d(r) = \pm \frac{g_V}{M} \phi_{122}(r). \quad (113)$$

For $|\kappa_e|, |\kappa_\nu| \leq 2$, six partial waves with $(\kappa_e, \kappa_\nu) = (\pm 1, \pm 2), (\pm 2, \pm 1), (\pm 2, \mp 2)$ contribute. In LOB, where only the operator $[Y_1 \otimes \boldsymbol{\sigma}]_2$ contributes, the RDME for four partial waves with $(\kappa_e, \kappa_\nu) = (\pm 1, \pm 2), (\pm 2, \pm 1)$ is given as

$$\langle f || \sum_L \Xi_{2L}(\kappa_e, \kappa_\nu) || i \rangle = S_{\kappa_e} \int_0^\infty dr r^2 \rho_{12}^\sigma(r) \phi_a(r) \quad (114)$$

with

$$\phi_a(r) = -g_A \alpha_{\kappa_e} \frac{\sqrt{2}}{3} (p_\nu \delta_{|\kappa_e|,1} + p_e \delta_{|\kappa_e|,2}) r. \quad (115)$$

By using the parametrization of the matrix elements given in the appendix, the RDME reads

$$\sqrt{2\pi} \langle f || \sum_L \Xi_{JL}(\kappa_e, \kappa_\nu) || i \rangle = S_{\kappa_e} \alpha_{\kappa_e} \frac{z}{2\sqrt{3}} (p_\nu \delta_{|\kappa_e|,1} + p_e \delta_{|\kappa_e|,2}), \quad (116)$$

and the spectrum shape is given by

$$(2J_i + 1)F(E_e, Z)C(E_e) = (\alpha_{-1}^2 + \alpha_1^2) \frac{z^2}{12} (p_\nu^2 + \lambda_2 p_e^2). \quad (117)$$

The expression agrees with that obtained in Ref. [11].

4.5 Beta-decay rate of ^{160}Sn

Table 1 Beta-decay rate in sec^{-1} . Electron wave functions obtained by solving the Dirac equation are employed for ‘Exact’, while the approximated electron wave functions described in paper (I) are used for NLO and LOB.

J^P		\mathbf{A}	$+V_0$	$+A_0$	$+\mathbf{V}(\text{WM})$	$+\mathbf{V}(\text{Conv.})$	Total
1^+	Exact	280.	-	285.	296.	296.	296.
	NLO	279.	-	285.	295.	295.	295.
	LOB	320.	-	-	-	-	320.
0^-	Exact	51.4	-	14.1	-	-	14.1
	NLO	51.2	-	14.1	-	-	14.1
	LOB	60.3	-	14.1	-	-	14.1
1^-	Exact	40.1	75.8	-	78.3	90.0	90.0
	NLO	40.0	75.5	-	78.2	90.3	90.3
	LOB	45.5	87.5	-	-	93.4	93.4
2^-	Exact	24.3	-	25.1	26.2	26.6	26.6
	NLO	24.2	-	25.0	26.1	26.5	26.5
	LOB	28.7	-	-	-	-	28.7

The formulas of the beta-decay rate in the impulse approximation developed in the previous subsection are applied to ^{160}Sn . The total decay rate is obtained by the sum of the partial decay rates of about 1400 states obtained from the pnQRPA approach. The allowed transition to the 1^+ states and the first-forbidden transitions to the 0^- , 1^- , and 2^- states are examined. We take $g_A = 1$ and include partial waves of the lepton wave function up to $|\kappa_e|, |\kappa_\nu| \leq 2$.

The results are summarized in Tab. 1. By ‘Exact’, we use the electron wave function obtained from the numerical solution of the Dirac equation and the plane-wave neutrino wave function. An approximate electron wave function of NLO is described in paper (I). The decay rate calculated from the LO terms of the nucleon current in the non-relativistic approximation is given in the third (the space component of the axial-vector current, \mathbf{A})

and the fourth column (adding the time component of the vector current, V_0). The 5th, 6th, and 7th columns show the decay rates calculated from the $\mathbf{A} + V_0 + A_0$, $\mathbf{A} + V_0 + A_0 + \mathbf{V}(\text{WM})$, and $\mathbf{A} + V_0 + A_0 + \mathbf{V}(\text{WM}) + \mathbf{V}(\text{Conv.})$ terms, respectively. The decay rates including all the components of the nucleon current are given in the 8th column (Total). Here the ‘Convection current’ (Conv.) includes the term obtained using the current conservation relation and all the terms associated with the convection current as explained in the previous subsection. In all the cases, NLO reproduces the ‘Exact’ results with 1% accuracy.

For the allowed transition 1^+ and the unique first-forbidden transition 2^- , the \mathbf{A} term gives a dominant contribution. The A_0 , $\mathbf{V}(\text{WM})$, and $\mathbf{V}(\text{Conv.})$ terms respectively contribute to the decay rate approximately by 2%, 3% and -0.1% for 1^+ and 3%, 4% and 2% for 2^- . LOB overestimates the decay rate by 14% for 1^+ and 12% for 2^- when ‘Exact’ is calculated using the \mathbf{A} term alone, while it reduces to 8% for 1^+ and 7% for 2^- when the A_0 and $\mathbf{V}(\text{WM})$ terms are included in ‘Exact’. The contributions of higher-order terms $Y_2\boldsymbol{\sigma} \cdot \nabla$ and $[Y_2 \otimes \nabla]_2$ are both less than 1% and their contributions are negligible.

The $\mathcal{O}(p/M)$ currents play an important role in the 0^- and 1^- transitions. For the transition to the 0^- states, the velocity-dependent current A_0 is as important as the spin-dipole operator in the \mathbf{A} term. Tab. 2 shows the decay rate calculated using the \mathbf{A} , V_0 , A_0 , and $\mathbf{V}(\text{Conv.})$ terms for 0^- and 1^- . A strong cancellation between two components takes place. This can be seen from the transition densities shown in Fig. 1 for the state with the largest partial decay width. Although the effect of improving the treatment of lepton wave function from LOB to ‘Exact’ is about 17% for the \mathbf{A} term and 9% for the A_0 term, the effect becomes small for the total decay rate in this case. Since the pion-exchange current due to the soft-pion production is known to contribute as largely as the impulse current [33], it would be important to include the meson-exchange current for the realistic estimation of the decay rate of 0^- [13, 33].

Table 2 Calculated beta-decay rates in sec^{-1} for the \mathbf{A} , V_0 , A_0 and $\mathbf{V}(\text{Conv.})$ terms.

J^P	\mathbf{A}	V_0	A_0	$\mathbf{V}(\text{Conv.})$
0^-	51.4	-	110.	-
1^-	40.1	64.9	-	114.

For the transition to the 1^- states, the dipole operator from the V_0 and $\mathbf{V}(\text{Conv.})$ terms and the spin-dipole operator from the \mathbf{A} term are equally important. Their contributions are shown in Tab. 2. If we do not use the current conservation relation for the convection current but use Eq. (95), the calculated decay rate becomes 140 sec^{-1} . The use of the




Fig. 1 Transition densities for the 0^- state at $E_0 = 13.94$ MeV of ^{160}Sn . The solid (blue) and dashed (red) curves show $\rho_{10}^\sigma r^3 E_0$ and $(\rho_0^{\sigma\nabla}/M)r^2$, respectively.




Fig. 2 Transition densities for the 1^- state at $E_0 = 15.63$ MeV of ^{160}Sn . The solid (blue) and dashed (red) curves show $\rho_1 r^3 E_0$ and $\rho_{11}^\sigma r^3 E_0$, respectively.

current conservation relation for the electric multipole of the vector current reduces the total decay rate by about 45%, while LOB overestimates the decay rate by about 5%. The weak magnetism alters the decay rate by about 5%, which is sizable, while the contribution of the ρ_{21}^∇ term is less than 1%. The transition densities for the dipole and spin-dipole operators are shown in Fig. 2. One sees coherent contributions of the V_0 and \mathbf{A} terms.

5 Summary

The formulas of the beta-decay rate including the velocity-dependent weak currents as well as the induced currents have been presented for the allowed and first-forbidden transitions. The longitudinal part of the vector current is eliminated by using the conservation of the vector current for the multipole operator of $P = (-1)^J$ transition.

The allowed and first-forbidden transitions of the beta decay of ^{160}Sn have been studied as an application of the present formulas. The transition densities are obtained by the EDF method. The results obtained with the electron wave functions in LO and NLO are compared with the ‘exact’ ones. It is shown that the conventional LO formula misses the transition rate by about 10 to 20 %. Including the NLO term reproduces the ‘exact’ with high accuracy. The induced weak-magnetism term typically contributes by about a few percent to the decay rate through the interference term of the leading-order amplitude. Reorganizing the formula using the current conservation is important for the 1^- transition operator of the vector current as known well. We reconfirmed the velocity-dependent term is as equally important as the spin-dipole term for the 0^- transition. The induced pseudo-scalar term can be neglected for the nuclear beta decay.

The formulas offer semi-analytic expressions of the transition matrix element of the nuclear beta decay, which are transparent and allow us to apply not only to various beta-decay processes but also the shape and angular correlations of the beta decay and neutrino reactions. The precise formulas are also useful to search signals of BSM in nuclear beta decay.

Acknowledgment

We would like to thank Prof. K. Koshigiri for the useful discussions. This work was in part supported by the JSPS KAKENHI Grants Nos. JP18H01210, JP18H04569, JP18K03635, JP19H05140, JP19K03824, JP21H00081, JP22H01237, JP22K03602, and JP22H01214, the Collaborative Research Program 2022, Information Initiative Center, Hokkaido University, and the JSPS/NRF/NSFC A3 Foresight Program “Nuclear Physics in the 21st Century.” The nuclear EDF calculation was performed on Yukawa-21 at the Yukawa Institute for Theoretical Physics, Kyoto University.

A Nuclear matrix element

Nuclear matrix elements w, x, u, z and those with Coulomb potential and velocity-dependent operators that are used in the literature are expressed in terms of the transition

densities introduced in section 4:

$$w = g_A \sqrt{4\pi} \int_0^\infty dr r^3 \rho_{10}^\sigma(r), \quad (\text{A1})$$

$$x = \mp g_V \sqrt{\frac{4\pi}{3}} \int_0^\infty dr r^3 \rho_1(r), \quad (\text{A2})$$

$$u = g_A \sqrt{\frac{8\pi}{3}} \int_0^\infty dr r^3 \rho_{11}^\sigma(r), \quad (\text{A3})$$

$$z = -g_A \sqrt{\frac{16\pi}{3}} \int_0^\infty dr r^3 \rho_{12}^\sigma(r), \quad (\text{A4})$$

$$\xi w' = g_A \sqrt{4\pi} \int_0^\infty dr r^2 \rho_{10}^\sigma(r) V_{D1}(r), \quad (\text{A5})$$

$$\xi x' = \mp g_V \sqrt{\frac{4\pi}{3}} \int_0^\infty dr r^2 \rho_1(r) V_{D1}(r), \quad (\text{A6})$$

$$\xi u' = g_A \sqrt{\frac{8\pi}{3}} \int_0^\infty dr r^2 \rho_{11}^\sigma(r) V_{D1}(r), \quad (\text{A7})$$

$$\xi' v = \sqrt{4\pi} \frac{g_A}{M} \int_0^\infty dr r^2 \rho_0^{\sigma\nabla}(r), \quad (\text{A8})$$

$$\xi' y = \mp \sqrt{4\pi} \frac{g_V}{M} \int_0^\infty dr r^2 \rho_{01}^{\sigma\nabla}(r), \quad (\text{A9})$$

$$\zeta_0 = \xi w' + \xi' v + \frac{E_0 w}{3}, \quad (\text{A10})$$

$$\zeta_1 = \xi' y - \xi u' - \xi x' + \frac{(u-x)E_0}{3} \quad (\text{A11})$$

for β^\mp decay and

$$\lambda_2 = \frac{\alpha_{-2}^2 + \alpha_2^2}{\alpha_{-1}^2 + \alpha_1^2}, \quad (\text{A12})$$

$$\frac{m_e}{E_e} \gamma_1 \mu_1 = \frac{\alpha_{-1}^2 - \alpha_1^2}{\alpha_{-1}^2 + \alpha_1^2}. \quad (\text{A13})$$

References

- [1] M. R. Mumpower, R. Surman, G. C. McLaughlin, and A. Aprahamian, *Prog. Part. Nucl. Phys.*, **86**, 86–126, [Erratum: *Prog.Part.Nucl.Phys.* 87, 116–116 (2016)] (2016), arXiv:1508.07352.
- [2] B. P. Abbott et al., *Phys. Rev. Lett.*, **119**, 161101 (2017), arXiv:1710.05832.
- [3] B. P. Abbott et al., *Astrophys. J. Lett.*, **848**, L12 (2017), arXiv:1710.05833.
- [4] G. Mention, M. Fechner, Th. Lasserre, Th. A. Mueller, D. Lhuillier, M. Cribier, and A. Letourneau, *Phys. Rev. D*, **83**, 073006 (Apr 2011).
- [5] A. C. Hayes, J. L. Friar, G. T. Garvey, Gerard Jungman, and G. Jonkmans, *Phys. Rev. Lett.*, **112**, 202501 (May 2014).
- [6] A. Glick-Magid and D. Gazit, *Jou. of Phys. G*, **49**(10), 105105 (sep 2022).

- [7] J. Kostensalo, M. Haaranen, and J. Suhonen, *Phys. Rev. C*, **95**, 044313 (Apr 2017).
- [8] W. Horiuchi, T. Sato, Y. Uesaka, and K. Yoshida, *Prog. Theor. Exp. Phys.*, **2021**(10), 103D03 (2021).
- [9] K. Koshigiri, M. Nishimura, H. Ohtsubo, and M. Morita, *Nucl. Phys. A*, **319**, 301, [Erratum: *Nucl.Phys.A* 340, 482 (1980)] (1979).
- [10] M. Morita and A. Fujii, *Phys. Rev.*, **118**, 606 (1960).
- [11] H. F. Schopper, *Weak Interactions and Nuclear Beta Decay*, (North-Holland Publishing Company, 1966).
- [12] H. Behrens and W. Bühring, *Nucl. Phys. A*, **162**, 111–144 (1971).
- [13] I. S. Towner, *Ann. Rev. Nucl. Part. Sci.*, **36**(1), 115–136 (1986).
- [14] S. Weinberg, *Phys. Rev.*, **112**, 1375–1379 (Nov 1958).
- [15] D. H. Wilkinson, *Physics Letters B*, **66**(2), 105–108 (1977).
- [16] J. D. Walecka, *Semileptonic weak interactions in nuclei in Muon physics II*, (Academic Press, Inc., 1975), Edited by V. W. Hughes and C. S. Wu.
- [17] S. Nakamura, T. Sato, Vladimir P. Gudkov, and K. Kubodera, *Phys. Rev. C*, **63**, 034617, [Erratum: *Phys.Rev.C* 73, 049904 (2006)] (2001), nucl-th/0009012.
- [18] A. C. Hayes and J. L. Friar, *Phys. Rev. C*, **98**, 065505 (Dec 2018).
- [19] A. Fujii, J.-I. Fujita, and M. Morita, *Prog. Theor. Phys.*, **32**(3), 438–449 (1964).
- [20] J.J. Sakurai and S.F. Tuan, *Modern Quantum Mechanics*, (Benjamin/Cummings Pub., 1985).
- [21] M. E. Rose, *Elementary theory of angular momentum*, (John Wiley & Sons, Inc., New York, 1957).
- [22] A.R. Edmonds, *Angular Momentum in Quantum Mechanics*, Investigations in Physics Series. (Princeton University Press, 1996).
- [23] T. Gorringe and H. W. Fearing, *Rev. Mod. Phys.*, **76**, 31–91 (Dec 2003).
- [24] L.D. Blokhintsev and E.I. Dolinsky, *Nucl. Phys.*, **34**(2), 498–499 (1962).
- [25] M. T. Mustonen and J. Engel, *Phys. Rev. C*, **93**, 014304 (2016), arXiv:1510.02136.
- [26] K. Yoshida, *Prog. Theor. Exp. Phys.*, **2013**, 113D02, [Erratum: *Prog. Theor. Exp. Phys.*, 2021, 019201 (2021)] (2013), arXiv:1308.0424.
- [27] J. Dobaczewski, H. Flocard, and J. Treiner, *Nucl. Phys. A*, **422**, 103 – 139 (1984).
- [28] H. Kasuya and K. Yoshida, *Prog. Theor. Exp. Phys.*, **2021**, 013D01 (2021), arXiv:2005.03276.
- [29] E. Chabanat, P. Bonche, P. Haensel, J. Meyer, and R. Schaeffer, *Nucl. Phys. A*, **635**, 231–256, [Erratum: *Nucl.Phys.A* 643, 441–441 (1998)] (1998).
- [30] M. Yamagami, Y. R. Shimizu, and T. Nakatsukasa, *Phys. Rev. C*, **80**, 064301 (2009), arXiv:0812.3197.
- [31] M. Morita, *Beta decay and muon capture*, (Benjamin, 1973).
- [32] H. Behrens, J. Jänecke, and H. Schopper, *Numerical tables for beta-decay and electron capture*, (Springer-Verlag Berlin, 1969).
- [33] K. Kubodera, J. Delorme, and M. Rho, *Phys. Rev. Lett.*, **40**, 755–758 (Mar 1978).