Low-frequency quantum oscillations in LaRhIn₅: Dirac point or nodal line?^a

G. P. Mikitik $^{1,\,\dagger}\,$ and Yu. V. Sharlai $^{1,\,2}$

¹B. Verkin Institute for Low Temperature Physics and Engineering of National Academy of Sciences of Ukraine, Kharkiv 61103, Ukraine ²Institute of Low Temperature and Structure Research, Polish Academy of Sciences, 50-422 Wrocław, Poland

^a ARISING FROM C. Guo et al. Nature Communications https://doi.org/10.1038/s41467-021-26450-1 (2021)

In the recent paper [1], a new method based on measuring a temperature correction to a quantum-oscillation frequency was proposed to study an energy-band dispersion of charge carriers in small Fermi surface (FS) pockets of crystals. To illustrate their approach, Guo et al. [1] applied it to a number of materials and, in particular, to the multiband metal LaRhIn₅ which, apart from high-frequency oscillations associated with a large FS, also exhibits the oscillations with the low frequency $F \approx 7$ T. Although the method of Ref. [1] really detects charge carriers with a linear dispersion, it does not distinguish between the carriers near a Dirac point and near a nodal line, since all such quasiparticles disperse linearly. Here we ask what is the nature of the carriers associated with the frequency F in LaRhIn₅ and call attention to the puzzling origin of this frequency.

Many years ago [2], we argued that the 7 T frequency is due to the minimal cross section of a FS surrounding a nodal line in LaRhIn₅, whereas Guo et al. [1] now relate this frequency with a cross section of a FS pocket enclosing a Dirac point. Below we show that the main experimental result of Ref. [1] does not contradict our assumption of the nodal line in LaRhIn₅. The degeneracy of two bands $\varepsilon_c(\mathbf{p})$ and $\varepsilon_v(\mathbf{p})$ along a nodal line, strictly speaking, occurs in LaRhIn₅ only when neglecting a weak spin-orbit interaction. Consider now these bands in the vicinity of some point \mathbf{p}_0 of the line, taking into account this interaction [3, 4] (Fig. 1),

$$\varepsilon_{\mathrm{c},\mathrm{v}}(\mathbf{p}) = \varepsilon_{\mathrm{d}} + \mathbf{a}\mathbf{p} + bp_{z}^{2} \pm \sqrt{\Delta^{2} + (v_{x}p_{x})^{2} + (v_{y}p_{y})^{2}}$$
(1)

where $\Delta \equiv \Delta(\mathbf{p}_0)$ is half of the spin-orbit gap at the point \mathbf{p}_0 , ε_d is the band-degeneracy energy at this point in absence of the spin-orbit coupling (i.e., when $\Delta = 0$), v_x , v_y , $\mathbf{a} = (a_x, a_y, a_z)$, and b are constant parameters, the quasimomentum \mathbf{p} is measured from \mathbf{p}_0 , the p_z axis coincides with the tangent to the band-contact line at this point, and the p_x , p_y axes are chosen in such a way that the quadratic form under the square root is diagonal. Let the magnetic field be directed along p_z . The cross section of a FS surrounding the nodal line at the plane $p_z = \text{constant}$ is a closed curve (an ellipse) only if $\tilde{a}_{\perp}^2 \equiv (a_x/v_x)^2 + (a_y/v_y)^2 < 1$. The parameter \tilde{a}_{\perp} characterizes the tilt of the spectrum at constant p_z , and $\tilde{a}_{\perp} \neq 0$ for all real situations. If the cross-sectional area at $p_z = 0$ is extremal with respect to p_z , then $a_z = 0$, and the term bp_z^2 is taken into account in Eq. (1).

The temperature dependence of the quantum-oscillation frequency F looks as follows [1]:

$$F(E_{\rm F}) = F_0 - \theta \frac{(\pi k_{\rm B} T)^2}{F_0 \beta^2}$$
(2)

where $E_{\rm F}$ is the Fermi energy, F_0 is the frequency of these oscillations at zero temperature, $\beta = e\hbar/2m_{\rm c}$, $m_{\rm c}$ is the cyclotron mass, and $\theta = 1/16$ for a band with a linear dispersion. With Eq. (1)



FIG. 1. The energy bands $\varepsilon_c(\mathbf{p})$ and $\varepsilon_v(\mathbf{p})$, Eq. (1), in the vicinity of their nodal line in the plane $p_z = 0$ perpendicular to the line. The red solid and black dashed lines show the bands with and without the spin-orbit interaction, respectively. The red circles mark the minimum of $\varepsilon_c(\mathbf{p})$ and the maximum of $\varepsilon_v(\mathbf{p})$ in the plane. The minimal indirect gap $2\Delta_{\min} = 2\Delta(1 - \tilde{a}_{\perp}^2)^{1/2}$ determined by these two points is less than 2Δ , the spin-orbit gap at $\mathbf{p} = 0$. Here $p_1 \equiv (a_x p_x + a_y p_y)/(\tilde{a}_{\perp}\Delta)$ is the dimensionless quasimomentum measured along the vector $(\tilde{a}_x, \tilde{a}_y)$ in the plane with the coordinates $p_x v_x/\Delta$ and $p_y v_y/\Delta$; $\tilde{a}_i \equiv a_i/v_i$, and $\tilde{a}_{\perp} \equiv (\tilde{a}_x^2 + \tilde{a}_y^2)^{1/2}$. The dotted line indicates the Fermi level E_F . Upper inset: The Fermi surface enclosing the nodal line (the dash-dotted line) at $(E_F - \varepsilon_d)b < 0$. Lower inset: The cross section (ellipse) of the Fermi surface on the plane $p_z = 0$. The black dashed line marks the direction along which the bands are shown in the main panel. The black asterisk and green cross mark the point $\mathbf{p} = 0$ and the center of the ellipse, respectively.

and formulas of Guo et al. [1] for θ , we arrive at

$$\theta = \frac{1}{16} \left(1 - \frac{\Delta_{\min}^2}{(E_{\rm F} - \varepsilon_{\rm d})^2} \right) \tag{3}$$

where $\Delta_{\min} = \Delta (1 - \tilde{a}_{\perp}^2)^{1/2}$ is the minimal indirect half-gap in the plane $p_z = 0$ (Fig. 1). In Ref. [1], the simplified spectrum with $a_x = a_y = \tilde{a}_{\perp} = 0$ was implied, and it was concluded that the value $\theta = 1/16$, which was experimentally obtained for LaRhIn₅, can occur only if the direct spin-orbit gap is perturbatively small, $\Delta^2/(E_{\rm F} - \varepsilon_{\rm d})^2 \ll 1$. On the other hand, the band-structure calculations [1] revealed that this ratio, in general, is not very small for LaRhIn₅, and Guo et al. ascribed the



FIG. 2. Magnetization of LaRhIn₅. The dots are the experimental data [5], the blue line shows the magnetization [2] produced by the nodal line in Fig. 1, whereas the red line is the magnetization of the Dirac pocket, with the background term $\chi_0 B = -0.7Cg(1)B$ being added to Eq. (4). This term is determined by the charge carriers that are far away from the point $\mathbf{p} = 0$ [2]. The inset is a zoom into the region $2 \le B \le 4$ T.

frequency *F* to a cross section passing through a Dirac point (when $\Delta \equiv 0$), excluding the case of the nodal line from their consideration. However, Eq. (3) demonstrates that the perturbatively small Δ is not necessary to obtain $\theta \approx 1/16$. It is sufficient if only the indirect spin-orbit gap in the plane of the extremal cross section of the FS is small, $\Delta_{\min}^2/(E_F - \varepsilon_d)^2 \ll 1$, and so a nodal line can lead to $\theta \approx 1/16$ even though the spin-orbit coupling is not perturbatively weak in LnRhIn₅.

It was shown earlier [2] that the experimental dependence of the longitudinal magnetization of LaRhIn₅ on the magnetic induction *B* [5] can be explained if a nodal line penetrates the minimal cross section of the FS in this material (Fig. 2). Let us now discuss the case of a FS pocket enclosing the Dirac point assumed by Guo et al. [1]. A formula for the magnetization of such a pocket with a linear dispersion of its charge carriers was derived many years ago [6], and a convenient representation [4, 7] of this formula reads:

$$M = CFg(u) \tag{4}$$

where the positive coefficient *C* depends on Dirac-spectrum parameters, $u \equiv F/B$, and g(u) is a universal function independent of any parameters. The magnetization calculated with Eq. (4) at

F = 7 T is shown in Fig. 2. The value of *C* is chosen in such a way that the calculated amplitude of the oscillations agrees with the experimental data. Figure 2 reveals a qualitative disagreement between this theoretical curve and the data. The theoretical curve (which is the same for electron and hole Dirac pockets) exhibits sharp peaks, whereas the experimental data reveal sharp troughs. Moreover, the behavior of the magnetization at B > F essentially deviates from the experimental dependence M(B). In other words, the assumption that the low-frequency oscillations are determined by a Dirac pocket is incompatible with the data [5] on the magnetization of LaRhIn₅. On the other hand, Supplementary Fig. 4 of Guo et al. [1] shows that the frequency F = 7 T of the Shubnikov–de Haas oscillations is practically independent of the direction of the magnetic field. This result is inconsistent with the nodal-line assumption, which leads to $F(\psi) \sim 1/\cos \psi$ where ψ is the angle between *B* and the line, and so Guo et al. [1] assumed the existence of the Dirac point. Thus, at present there is no self-consistent explanation of the 7 T oscillations in LaRhIn₅.

The Fermi surface near the nodal line (upper inset in Fig. 1) and the appropriate M(B) (blue line in Fig. 2) are shown for the case $(E_F - \varepsilon_d)b < 0$. If the small difference $E_F - \varepsilon_d$ changes its sign, the anisotropy of the frequency $F(\psi)$ noticeably decreases, and M(B) resembles the red curve in Fig. 2 [2]. Thus, the previously published data [1, 5] look as if they were obtained on crystals with slightly different E_F but with practically equal $|E_F - \varepsilon_d|$. This hypothesis can be verified, measuring both the Shubnikov–de Haas oscillations and the longitudinal magnetization M(B) in one and the same sample. Then, in a sample with M(B) like in Goodrich et al. [5], the dependence $F(\psi)$ has to be strongly anisotropic, whereas in a sample with a weak dependence $F(\psi)$, M(B) cannot exhibit the sharp troughs visible in the oscillations in Fig. 2.

Acknowledgements. Yu.V.S. acknowledges the Poland NCN program for scientists from Ukraine [R-2022/01/3/ST3/00083].

[†] Corresponding author: mikitik@ilt.kharkov.ua

- Guo, C. et al. Temperature dependence of quantum oscillations from non-parabolic dispersion. *Nat. Commun.* 12, 6213 (2021).
- [2] Mikitik, G. P. & Sharlai, Yu. V. Berry phase and de Haas van Alphen effect in LaRhIn₅. *Phys. Rev. Lett.* 93, 106403 (2004).
- [3] Mikitik, G.P. & Sharlai Yu.V. Semiclassical energy levels of electrons in metals with band degeneracy

lines. JETP 87, 747-755 (1998).

- [4] Mikitik, G. P. & Sharlai, Yu. V. Magnetic Susceptibility of Topological Semimetals. *J. Low Temp. Phys.* 197, 272-309 (2019).
- [5] Goodrich, R.G. et al. Magnetization in the ultraquantum limit. Phys. Rev. Lett. 89, 026401 (2002).
- [6] Mikitik, G.P., Sharlai, Yu.V. Field dependences of magnetic susceptibility of crystals under conditions of degeneracy of their electron energy bands. *Low Temp. Phys.* **22**, 585-592 (1996).
- [7] Mikitik, G. P. & Sharlai, Yu. V. Magnetic susceptibility of topological nodal semimetals. *Phys. Rev. B* 94, 195123 (2016).