NMR in ³He-B

V. V. Zavjalov[∗]

March 17, 2023

Abstract

This text contains a collection of equations useful for understanding Nuclear Magnetic Resonance (NMR) experiments in superfluid ³He-B. This is a part of my notebook where I try to describe some parts of this sophisticated system.

Introduction

Nuclear magnetic resonance (NMR) in superfluid ³He is a very powerful tool used for studying this system since its discovery in 1972. Superfluidity in fermionic ³He is formed via Cooper pairing of atoms, with pairs having spin 1 and orbital momentum 1. As a result the order parameter of the system includes both spin and orbital degrees of freedom and can be written as 3x3 complex matrix. A few different superfluid phases with different broken symmetries are possible. For the B-phase degenerate space of the order parameter includes an arbitrary 3D rotation matrix which describes fixed mutual orientation of spin and orbital spaces. In NMR experiments we observe motion of this matrix, which means three degrees of freedom and three spin-wave modes. This is different from many other magnetic materials where we study motion of magnetization vector with only two degrees of freedom. Another important feature of this system is possibility of non-uniform spatial distribution and topological defects in the orderparameter field (so-called textures). This can be clearly observed in linear NMR where small oscillations of the order parameter are happening around the equilibrium texture. One more feature of this system is spin-orbit interaction. It introduces an additional non-linear force acting on the order parameter and leads to important effects such as longitudinal NMR or Homogeneously Precessing Domain (HPD).

This text contains two almost independent parts. The first one is about linear NMR. It was written during my work in Vladimir Eltsov's group in Aalto University (Finland) in 2012-2015. We had a really nice experimental system for measuring optical magnons trapped in a harmonic potential formed by the order parameter texture and magnetic field. It was possible to control shape of the trap and population of individual levels in it, to observe Bose condensation of magnons, parametric excitation of other spinwave modes, interaction of the magnon condensate with texture, quantized vortices, and free surface of helium. I would like to thank Petri Heikkinen, Samuli Autti, and Jere Mäkinen who also worked on this project.

The second part contains Leggett equations for non-linear NMR and many things related to Homogeneously Precessing Domain (HPD), a unique coherent state which is also a very useful tool for various ³He studies. I worked with HPD in Vladimir Dmitriev's group in Kapitza Institute (Russia) in 2000-2006, and in Pertti Hakonen's group in Aalto University in 2016-2019.

The text contains many references to Vollhardt and Wölfle book $[1]$, usually written as $(WV \leq \text{equa--}$ tion number>). Many derivations require straightforward but huge tensor manipulations. A useful tool for this kind of calculations is cadabra $[2]$. Some of ³He-related parameters and equations can be found in my he3lib library [\[3\]](#page-18-2).

I hope this work will be useful for understanding NMR experiments in $^3{\rm He}\text{-B}.$

[∗]e-mail: v.zavjalov@lancaster.ac.uk

Part 1. Linear spin waves in ³He-B

At energy scales much less then energy gap Δ the state of superfluid ³He-B is described by the order parameter:

$$
A_{aj} = \frac{1}{\sqrt{3}} \Delta e^{i\varphi} R_{aj} \tag{1}
$$

where φ is phase, and R_{aj} is a rotation matrix which can be written in terms of rotation axis **n** and rotation angle ϑ as

$$
R_{aj} = \cos\vartheta \, \delta_{aj} + (1 - \cos\vartheta) \, n_a n_j - \sin\vartheta \, e_{ajk} n_k. \tag{2}
$$

In small magnetic fields, $h \gamma H \ll \Delta$ the gap Δ and magnetic susceptibility χ_B are isotropic. There are four degrees of freedom in the system, the phase φ , the unit vector **n** and the angle ϑ . Oscillations of the phase is sound (here we do not consider normal component and thus have only one sound mode), and oscillations of the $R_{ai}(\mathbf{n},\vartheta)$ matrix are spin waves. In the following discussion we are interested only in these three spin-wave modes. Hamiltonian of the system can be written as a combination of three components: magnetic energy, energy of spin-orbit interaction and gradient energy (see VW 6.103, VW 7.17).

$$
\mathcal{H} = F_M + F_{SO} + F_{\nabla},\tag{3}
$$

$$
F_M = -(\mathbf{S} \cdot \gamma \mathbf{H}) + \frac{\gamma^2}{2\chi_B} \mathbf{S}^2,\tag{4}
$$

$$
F_{SO} = g_D \Delta^2 \Big[R_{jj} R_{kk} + R_{jk} R_{kj} - \frac{2}{3} R_{jk} R_{jk} \Big] = g_D \Delta^2 \Big[R_{jj} R_{kk} + R_{jk} R_{kj} \Big] + \text{const.},
$$
 (5)

$$
F_{\nabla} = \frac{1}{2} \Delta^2 \Big[K_1(\nabla_j R_{ak}) (\nabla_j R_{ak}) + K_2(\nabla_j R_{ak}) (\nabla_k R_{aj}) + K_3(\nabla_j R_{aj}) (\nabla_k R_{ak}) \Big],\tag{6}
$$

where **S** is spin and **H** is magnetic field.

We are going to write equations of motion for this system. Spin is a not a canonical variable, its components do not commute with each other and thus we have to use equations with Poisson brackets. In this approach evolution of any parameter a can be written as $\dot{a} = \{H, a\}$. Poisson brackets can be found from microscopic considerations, from commutation rules in quantum mechanics, or from symmetry $[4, 4]$ $[4, 4]$ [5\]](#page-18-4).

The matrix R_{aj} for any value of index j can be treated as a vector in spin space. For such a vector commutation rules can be written as:

$$
\{S_a, S_b\} = -e_{abc}S_c, \quad \{R_{aj}, S_b\} = \{S_a, R_{bj}\} = -e_{abc}R_{cj}, \quad \{R_{aj}, R_{bk}\} = 0.
$$
 (7)

It is possible to write equations of motion for the spin **S** and matrix R_{aj} (see later, in the second part of this text), but here we will use small rotations θ^s of the spin space as coordinates.

Using small rotations θ_a^s as coordinates

We follow derivation of Theodorakis and Fetter $[6]$, see also VW 8.4.6. Consider some equilibrium distribution of the order parameter matrix R^0 . We are going to study small oscillations around the equilibrium. Arbitrary change of the rotation matrix can be represented as an additional rotation:

$$
R_{aj} = R_{ab}(\theta^s) R_{bj}^0.
$$
\n⁽⁸⁾

Here θ^s is a vector and $R_{ab}(\theta^s)$ is a rotation matrix around the direction of θ^s by angle $|\theta^s|$. We will use small rotations, $|\theta^s| \ll 1$. Using formula [\(2\)](#page-1-0) for $R_{ab}(\theta^s)$ one can write up to second order terms in $|\theta^s|$.

$$
R_{aj} = \left(\delta_{ab} - e_{abc}\theta_c^s + \frac{1}{2}\theta_a^s \theta_b^s - \frac{1}{2}\delta_{ab}\theta_c^s \theta_c^s + o(\theta^{s2})\right) R_{bj}^0
$$
\n
$$
(9)
$$

The Hamiltonian can be written as a function of the spin and angles θ^s . Poisson brackets in the equations of motion can be expanded as

$$
\dot{S}_a = \{ \mathcal{H}, S_a \} = \frac{\delta \mathcal{H}}{\delta S_b} \{ S_b, S_a \} + \frac{\delta \mathcal{H}}{\delta \theta_b^s} \{ \theta_b^s, S_a \}
$$
(10)

$$
\dot{\theta}_a^s = \{ \mathcal{H}, \theta_a^s \} = \frac{\delta \mathcal{H}}{\delta S_b} \{ S_b, \theta_a^s \} + \frac{\delta \mathcal{H}}{\delta \theta_b^s} \{ \theta_b^s, \theta_a^s \}
$$
(11)

We are going to obtain linear equations for small θ^s . Derivatives of the Hamiltonian are zero in the equilibrium (energy is in the minimum at $\theta^s = 0$) and thus we have only first-order terms there. Thus in the Poisson brackets we need only zero-order terms. As the matrix R_{aj} is a function of θ^s one can write

$$
\{R_{aj}, S_b\} = \frac{dR_{aj}}{d\theta_c^s} \{\theta_c^s, S_b\}.
$$
\n(12)

Using (7) and (9) one can find commutation rules for our set of coordinates:

$$
\{S_a, S_b\} = -e_{abc}S_c, \quad \{\theta_a^s, S_b\} = -\{S_a, \theta_b^s\} = -\delta_{ab}, \quad \{\theta_a^s, \theta_b^s\} = 0.
$$
 (13)

It is easy to find the derivative $\delta \mathcal{H}/\delta S_h$. Using it and the commutation rules we have:

$$
\dot{\mathbf{S}} = [\mathbf{S} \times \gamma \mathbf{H}] - \frac{\delta \mathcal{H}}{\delta \theta^{\mathbf{s}}}
$$
(14)

$$
\dot{\theta}^s = \gamma \left(\frac{\gamma}{\chi_B} \mathbf{S} - \mathbf{H} \right) = \frac{\gamma^2}{\chi_B} \delta \mathbf{S}
$$
 (15)

where we introduced δS , deviation of the spin from its equilibrium value $S^0 = \chi_B H/\gamma$. These equations describe a simple Larmor precession with the additional term $-\delta \mathcal{H}/\delta \theta^s$ — a torque acting on the spin because of gradient and spin-orbit interactions. This derivative is not trivial, and will be calculated below.

Variational derivative for small rotations

Since the energy depends on both angles θ_a^s and their gradients $\nabla_j \theta_a^s$ we have to use a so-called variational derivative $\delta \mathcal{H}/\delta \theta_a^s$ [\[7\]](#page-18-6). An additional difficulty appears because rotations θ^s do not commute. Let us take this into account. Using formula [\(9\)](#page-1-2) one can check that two small successive rotations, θ_A and then θ_B are equivalent to the rotation by the angle $\theta_A + \theta_B + \frac{1}{2}\theta_A \times \theta_B$ up to second order terms in the angles:

$$
R_{ac}(\theta_B)R_{cb}(\theta_A) = R_{ab}\left(\theta_A + \theta_B + \frac{1}{2}[\theta_A \times \theta_B] + o(\theta_A^2, \theta_B^2)\right).
$$
 (16)

Two rotations done in a different order produce a difference $[\theta_A \times \theta_B]$. This can be written as a commutation rule for differentials (see VW 9.14):

$$
\delta \nabla \theta^s - \nabla \delta \theta^s = [\delta \theta^s \times \nabla \theta^s]. \tag{17}
$$

Consider a small non-uniform rotation, $\delta\theta^s(\mathbf{r})$, with boundary condition $\delta\theta^s = 0$ which changes gradients by $\delta \nabla_j \theta^s$. The functional derivative $\delta \mathcal{H}/\delta \theta^s_a$ is given by

$$
\int_{V} \frac{\delta \mathcal{H}}{\delta \theta_{a}^{s}} \delta \theta_{a}^{s} dr = \int_{V} \left(\frac{\partial F}{\partial \theta_{a}^{s}} \delta \theta_{a}^{s} + \frac{\partial F}{\partial \nabla_{j} \theta_{a}^{s}} \delta \nabla_{j} \theta_{a}^{s} \right) dr
$$
\n(18)

From mechanical point of view this means that work produced against a torque $T_a = -\delta \mathcal{H}/\delta \theta_a^s$ equals to the total change of energy. Now we swap differentials $\delta \nabla \theta^s$ using the formula [\(17\)](#page-2-0) and integrate this term by parts using the zero boundary condition.

$$
\int_{V} \frac{\delta \mathcal{H}}{\delta \theta_{a}^{s}} \delta \theta_{a}^{s} dr = \int_{V} \left(\frac{\partial F}{\partial \theta_{a}^{s}} \delta \theta_{a}^{s} - \nabla_{j} \frac{\partial F}{\partial \nabla_{j} \theta_{a}^{s}} \delta \theta_{a}^{s} + \frac{\partial F}{\partial \nabla_{j} \theta_{a}^{s}} e_{abc} \delta \theta_{b}^{s} \nabla_{j} \theta_{c}^{s} \right) dr \tag{19}
$$

This equation is true for any possible variations $\delta\theta^s$. This means that the integrands are equal at any point and the derivative is:

$$
\frac{\delta \mathcal{H}}{\delta \theta_a^s} = \frac{\partial F}{\partial \theta_a^s} - \nabla_j \frac{\partial F}{\partial \nabla_j \theta_a^s} + \frac{\partial F}{\partial \nabla_j \theta_c^s} e_{abc} \nabla_j \theta_b^s \tag{20}
$$

Gradient energy

The gradient energy in 3 He-B is given by [\(6\)](#page-1-3). Now put the expression for the distorted matrix [\(9\)](#page-1-2) into it. Both original matrix R^0 and rotation angles θ^s can be non-uniform here:

$$
\frac{2}{\Delta^{2}}(F_{\nabla} - F_{\nabla}^{0}) = (2K_{1} + K_{2} + K_{3}) \left(\nabla_{j} \theta_{a}^{s} \right) \left(\nabla_{j} \theta_{a}^{s} \right) - \left[K_{2} R_{aj}^{0} R_{bk}^{0} + K_{3} R_{ak}^{0} R_{bj}^{0} \right] \left(\nabla_{j} \theta_{a}^{s} \right) \left(\nabla_{k} \theta_{b}^{s} \right) \n+ \left[K_{1} R_{ak}^{0} (\nabla_{j} R_{bk}^{0}) + K_{2} R_{ak}^{0} (\nabla_{k} R_{bj}^{0}) + K_{3} R_{aj}^{0} (\nabla_{k} R_{bk}^{0}) \right] \left(2e_{abc} + \delta_{ac} \theta_{b}^{s} - \delta_{bc} \theta_{a}^{s} \right) \left(\nabla_{j} \theta_{c}^{s} \right)
$$

where F_{∇}^{0} is an energy calculated for the undistorted matrix R^{0} . Substituting this into [\(20\)](#page-2-1) we have the torque $T_c^{\nabla} = -\delta F_{\nabla}/\delta \theta^s$:

$$
\frac{T_c^{\nabla}}{\Delta^2} = (2K_1 + K_2 + K_3) (\nabla_j \nabla_j \theta_c^s) - \nabla_j \left[(K_2 R_{ak}^0 R_{cj}^0 + K_3 R_{aj}^0 R_{ck}^0) (\nabla_k \theta_a^s) \right]
$$
\n
$$
+ \frac{1}{2} R_{ak}^0 \left[K_1 (\nabla_j \nabla_j R_{bk}^0) + (K_2 + K_3) (\nabla_j \nabla_k R_{bj}^0) \right] (2e_{abc} + \delta_{ac} \theta_b^s - \delta_{bc} \theta_a^s)
$$
\n(22)

If gradients of equilibrium texture R^0 can be neglected ("uniform texture") then

$$
T_c^{\nabla} = \Delta^2 \left[K \ \delta_{ac} \ \nabla_j \nabla_j - K' \ R^0_{ak} R^0_{cj} \nabla_j \nabla_k \right] \theta^s_a \tag{23}
$$

where $K = 2K_1 + K_2 + K_3$ and $K' = K_2 + K_3$.

Dipolar energy

Dipolar energy is given by (6) . Substituting the small rotation (9) we get:

$$
R_{jj}R_{kk} + R_{jk}R_{kj} = [R_{jj}^{0}R_{kk}^{0} + R_{jk}^{0}R_{kj}^{0}] (1 - |\theta^{s}|^{2})
$$

+
$$
[R_{aj}^{0}R_{kk}^{0} + R_{ak}^{0}R_{kj}^{0}] (\theta_{j}^{s}\theta_{a}^{s} - 2e_{jab} \theta_{b}^{s})
$$

+
$$
[R_{aj}^{0}R_{a'k}^{0} + R_{ak}^{0}R_{a'j}^{0}] e_{jab} \theta_{b}^{s} e_{ka'b'} \theta_{b'}^{s}
$$
 (24)

Or in terms of axis **n** and rotation angle ϑ of matrix R^0 :

$$
R_{jj}R_{kk} + R_{jk}R_{kj} = \frac{1}{2}(4\cos\vartheta + 1)^2 - \frac{1}{2}
$$

\n
$$
- 4\sin\vartheta(4\cos\vartheta + 1)(\mathbf{n} \cdot \theta^s)
$$

\n
$$
- (4\cos\vartheta + 1)(\cos\vartheta + 1) |\theta^s|^2
$$

\n
$$
+ (9 + 3\cos\vartheta - 12\cos^2\vartheta)(\mathbf{n} \cdot \theta^s)^2
$$

The torque $T_c^D = -\delta F_D/\delta\theta^s$ is

$$
\frac{\mathbf{T}^D}{\Delta^2} = \frac{2g_D(4\cos\vartheta + 1)[2\sin\vartheta \mathbf{n} + (\cos\vartheta + 1)\theta^{\mathbf{s}}]}{-2g_D(9 + 3\cos\vartheta - 12\cos^2\vartheta)(\mathbf{n} \cdot \theta^{\mathbf{s}})\mathbf{n}}
$$
\n(26)

Equilibrium texture stays in the minimum of the dipolar energy where $\cos \theta = -1/4$. In this case the dipolar torque is

$$
\mathbf{T}^{D} = -15g_{D}\Delta^{2} \ (\mathbf{n} \cdot \theta^{\mathbf{s}})\mathbf{n}
$$
 (27)

Spin waves in a uniform texture

Put the dipolar torque [\(27\)](#page-3-0) and the gradient torque [\(23\)](#page-3-1) into the equation [\(14\)](#page-2-2), differentiate it over time and exclude θ^s using the equation [\(15\)](#page-2-2). Here we work only with first order terms in θ^s and thus do not care about commutation of derivatives [\(17\)](#page-2-0).

$$
\delta \ddot{\mathbf{S}} = [\delta \dot{\mathbf{S}} \times \gamma \mathbf{H}] + \hat{\mathbf{\Lambda}} \delta \mathbf{S},
$$

where $\Lambda_{ab} = \frac{\Delta^2 \gamma^2}{\chi_B} \left[K \delta_{ab} \delta_{kj} - K' R_{ak}^0 R_{bj}^0 \right] \nabla_j \nabla_k - \Omega_B^2 n_a n_b.$ (28)

Here we introduced Leggett frequency:

$$
\Omega_B^2 = \frac{\gamma^2 \Delta^2}{\chi_B} 15g_D. \tag{29}
$$

For a flat wave $\delta S = s \exp(i\omega t + i kx)$ with frequency ω and wave vector k in the magnetic field directed along z axis

$$
\left[\left(\begin{array}{ccc} \omega^2 & -i\omega\omega_L & 0 \\ i\omega\omega_L & \omega^2 & 0 \\ 0 & 0 & \omega^2 \end{array} \right) + \hat{\mathbf{\Lambda}} \right] \left(\begin{array}{c} s_x \\ s_y \\ s_z \end{array} \right) = 0, \tag{30}
$$

$$
\Lambda_{ab} = \frac{\Delta^2 \gamma^2}{\chi_B} \left[K \ \delta_{ab} \mathbf{k}^2 - K' \ R_{ak}^0 k_k R_{bj}^0 k_j \right] - \Omega_B^2 \ n_a n_b. \tag{31}
$$

where $\omega_L = \gamma H$. Solution corresponds to zero determinant of the matrix. Using the fact that $\hat{\Lambda}$ is symmetric this can be written as:

$$
\begin{aligned}\n\left[(\Lambda_{xx} + \omega^2)(\Lambda_{yy} + \omega^2) - \Lambda_{xy}^2 - \omega^2 \omega_L^2 \right] (\Lambda_{zz} + \omega^2) &+ \\
2\Lambda_{xy}\Lambda_{yz}\Lambda_{xz} - (\Lambda_{xx} + \omega^2)\Lambda_{yz}^2 - (\Lambda_{yy} + \omega^2)\Lambda_{xz}^2 &= 0,\n\end{aligned}
$$
\n(32)

or

$$
\omega^6 + \omega^4 [\Lambda_{xx} + \Lambda_{yy} + \Lambda_{zz} - \omega_L^2]
$$

+
$$
\omega^2 [\Lambda_{xx} \Lambda_{yy} + \Lambda_{yy} \Lambda_{zz} + \Lambda_{zz} \Lambda_{xx} - \Lambda_{xy}^2 - \Lambda_{yz}^2 - \Lambda_{xz}^2 - \omega_L^2 \Lambda_{zz}]
$$

+
$$
\Lambda_{xx} \Lambda_{yy} \Lambda_{zz} + 2\Lambda_{xy} \Lambda_{yz} \Lambda_{xz} - \Lambda_{xx} \Lambda_{yz}^2 - \Lambda_{yy} \Lambda_{zx}^2 - \Lambda_{zz} \Lambda_{xy}^2 = 0,
$$
 (33)

We have a third-order equation for ω^2 . It gives three doubly degenerate spin-wave modes with positive and negative frequencies. The equation can be analytically solved for arbitrary wave-vector **k**, **n** and ω_L in the uniform texture ($\nabla R^0 \ll k$). Calculation is implemented in he3lib [\[3\]](#page-18-2).

Uniform precession

If we neglect all gradient terms then $\Lambda_{ab} = -\Omega_B^2 n_a n_b$ and equation [\(28\)](#page-3-2) is (compare with [\[8\]](#page-18-7)):

$$
\omega^6 - \omega^4 \left(\omega_L^2 + \Omega_B^2\right) + \omega^2 \omega_L^2 \Omega_B^2 n_z^2 = 0.
$$
\n(34)

NMR experiments are often done at a fixed frequency ω . We can solve the equation for $\gamma H = \omega_L$ and find magnetic field where the resonance is observed:

$$
\gamma H = \omega \sqrt{\frac{\omega^2 - \Omega_B^2}{\omega^2 - \Omega_B^2 n_z^2}}.
$$
\n(35)

At a fixed magnetic field we solve the equation for ω and find three spin-wave modes:

$$
\omega^2 = \frac{1}{2}(\omega_L^2 + \Omega_B^2) \pm \sqrt{\frac{1}{4}(\omega_L^2 + \Omega_B^2)^2 - \omega_L^2 \Omega_B^2 n_z^2}, \qquad \omega^2 = 0
$$
\n(36)

For $\gamma H \gg \Omega_B$ we have a well-known expressions for transverse and longitudinal NMR frequency:

$$
\omega = \gamma H + \frac{\Omega_B^2}{2\gamma H} (1 - n_z^2), \qquad \omega = \Omega_B n_z, \qquad \omega = 0,
$$
\n(37)

Here meaning of Ω_B becomes clear, it is frequency of the longitudinal NMR in a texture with n||H.

Separating equations for transverse and longitudinal magnons

Let's use complex *spherical coordinates*. For an arbitrary vector A:

$$
A_{\pm} = \frac{1}{\sqrt{2}} (A_x \pm iA_y), \quad A_0 = A_z, \qquad A_i B_i = A_q B_q^*, \quad \text{where } q = 0, +, -
$$
 (38)

In terms of polar and azimuthal angles α and β :

$$
A_{\pm} = \frac{|A|}{\sqrt{2}} \sin \beta \exp(\pm i\alpha), \qquad A_0 = |A| \cos \beta \tag{39}
$$

If field **H** is directed alone the z axis, then $[A \times H]_q = -qiA_qH$. Now one can write the equation in spherical coordinates for a harmonic oscillation $\delta S_q = s_q e^{i\omega t}$.

$$
\left[\frac{\gamma^2 \Delta^2}{\chi_B} \left[-K \ \delta_{qp} \nabla_j \nabla_j + K' \ R_{qj}^0 R_{p^*k}^0 \nabla_j \nabla_k\right] + \Omega_B^2 \ n_q n_p^* \right] s_p = \omega(\omega - q \gamma H) s_q \tag{40}
$$

Now let's remove non-diagonal coupling between different modes. According to [\[6\]](#page-18-5) this coupling is small in the high-frequency limit $\omega \gg \Omega_B$ (as Ω_B^2/ω^2). Then we can write (without summation of the q index):

$$
\left[\frac{\gamma^2 \Delta^2}{\chi_B} \left[-K \nabla_j \nabla_j + K' R^0_{qj} R^0_{q^*k} \nabla_j \nabla_k\right] + \Omega_B^2 |n_q|^2\right] s_q = \omega(\omega - q\gamma H)s_q \tag{41}
$$

Now we can write separate equations for transverse $(q = +, -)$ and longitudinal $(q = 0)$ modes. We use the fact that

$$
R_{+j}^{0}R_{-k}^{0} = \frac{1}{2}(R_{xj}^{0}R_{xk} + R_{yj}^{0}R_{yk}) = \frac{1}{2}(\delta_{jk} - R_{zj}^{0}R_{zk}),
$$
\n(42)

and R_{zj}^0 is an equilibrium orbital anisotropy axis: $\hat{L}_j^0 = R_{aj}^0 \hat{S}_a^0 = R_{zj}^0$.

$$
\left[-c_{\perp}^{2} \nabla^{2} - (c_{\parallel}^{2} - c_{\perp}^{2}) \hat{L}_{j}^{0} \hat{L}_{k}^{0} \nabla_{j} \nabla_{k} + \frac{1}{2} \Omega_{B}^{2} \sin^{2} \beta_{N} \right] s_{+} = \omega(\omega - \gamma H) s_{+}
$$
\n(43)

$$
\[-C_{\perp}^{2} \nabla^{2} - (C_{\parallel}^{2} - C_{\perp}^{2}) \hat{L}_{j}^{0} \hat{L}_{k}^{0} \nabla_{j} \nabla_{k} + \Omega_{B}^{2} \cos^{2} \beta_{N} \]_{s_{0}} = \omega^{2} s_{0}
$$
\n(44)

here we introduce parameters:

$$
c_{\parallel}^{2} = C_{\perp}^{2} = \frac{\gamma^{2} \Delta^{2}}{\chi_{B}} K, \quad c_{\perp}^{2} = \frac{\gamma^{2} \Delta^{2}}{\chi_{B}} (K - K'/2), \quad C_{\parallel}^{2} = \frac{\gamma^{2} \Delta^{2}}{\chi_{B}} (K - K'), \tag{45}
$$

We have skipped the equation for s_. It can be obtained from one for s₊ by changing ω by $-\omega$ and thus have same solutions but with an opposite sign. As expected we have three doubly degenerate modes with positive and negative ω . Note that these equations have been obtained in the assumption $\omega \gg \Omega_B$.

Spin-wave velocities in a weak coupling approximation

Gradient energy coefficients have been obtained in [\[9\]](#page-18-8) (see also VW 7.25).

$$
2K_1 + K_2 + K_3 = -\frac{2}{\Delta^2} \left(\frac{\hbar}{2m}\right)^2 (4+\delta)c, \qquad K_2 = -\frac{2}{\Delta^2} \left(\frac{\hbar}{2m}\right)^2 c, \qquad K_3 = -\frac{2}{\Delta^2} \left(\frac{\hbar}{2m}\right)^2 (1+\delta)c \tag{46}
$$

where

$$
c = -\frac{\rho_s}{10} \frac{3 + F_1^a}{3 + F_1^s} \frac{1}{1 + F_1^a (5 - 3\rho_s/\rho)/15}, \qquad \delta = \frac{F_1^a \rho_s/\rho}{3 + F_1^s (1 - \rho_s/\rho)}
$$
(47)

Then

$$
c_{\parallel}^{2} = C_{\perp}^{2} = -\frac{2\gamma^{2}}{\chi_{B}} \left(\frac{\hbar}{2m}\right)^{2} (4+\delta)c, \quad c_{\perp}^{2} = -\frac{2\gamma^{2}}{\chi_{B}} \left(\frac{\hbar}{2m}\right)^{2} (3+\delta/2)c, \quad C_{\parallel}^{2} = -\frac{2\gamma^{2}}{\chi_{B}} \left(\frac{\hbar}{2m}\right)^{2} 2c \quad (48)
$$

Without Fermi liquid corrections $K_1 = K_2 = K_3$ and

$$
c_{\perp}/c_{\parallel} = \sqrt{3/4}, \qquad C_{\perp}/C_{\parallel} = \sqrt{2}.
$$
 (49)

In many papers one of these two sets of spin-wave velocities is usually used. For example, in early Leggett's papers it is C_{\perp}/C_{\parallel} [\[10\]](#page-18-9), in Fomin's papers it is c_{\perp}/c_{\parallel} [\[11\]](#page-18-10).

Motion of L, n and ϑ in the spin wave

A common question is how **L**, **n** and ϑ move in the transverse NMR. Here we calculate this explicitly. First let us find deviations of rotation angle and axis of the matrix R_{aj} , δ n and $\delta \vartheta$ caused by rotation θ^s . In a linear approximation:

$$
R_{aj} = R_{aj}^0 + \frac{\partial R_{aj}^0}{\partial \vartheta} \delta \vartheta + \frac{\partial R_{aj}^0}{\partial n_k} \delta n_k = (\delta_{ab} - e_{abc} \theta_c^s) R_{bj}^0
$$
\n
$$
\tag{50}
$$

Using expression [\(2\)](#page-1-0) for R_{aj}^0 and finding convolutions with δ_{aj} , n_a and n_j one can find:

$$
\delta\vartheta = (\mathbf{n} \cdot \theta^{\mathbf{s}}), \qquad \delta\mathbf{n} = -\frac{1}{2} [\mathbf{n} \times \theta^{\mathbf{s}}] + \frac{\theta^{\mathbf{s}} - (\theta^{\mathbf{s}} \cdot \mathbf{n}) \mathbf{n}}{2 \tan \vartheta/2}
$$
(51)

Orbital and spin anisotropy axes are connected by the order parameter: $L_j = S_a R_{aj}$. Using [\(9\)](#page-1-2) and (15) one can write up to the first order of θ^s :

$$
L_j = \frac{\chi_B}{\gamma^2} (\gamma \mathbf{H} + \dot{\theta}^s - [\gamma \mathbf{H} \times \theta^s])_a R_{aj}^0
$$
 (52)

Consider a harmonic transverse spin wave with frequency ω in the field **H** \parallel \hat{z} . Then

$$
\dot{\theta}^s = \frac{\gamma^2}{\chi_B} \delta \mathbf{S}, \qquad \theta^s = -\frac{\gamma^2}{\omega \chi_B} [\hat{z} \times \delta \mathbf{S}] \tag{53}
$$

and substituting this to [\(51\)](#page-6-0) and [\(52\)](#page-6-1) we write $\delta \vartheta$ and **L**:

$$
\delta\vartheta = -\frac{\gamma^2}{\omega\chi_B}[\mathbf{n}\times\delta\mathbf{S}]_z = \frac{\gamma^2}{\omega\chi_B}(n_x\delta S_y - n_y\delta S_x)
$$
\n(54)

$$
L_j = \frac{\chi_B H}{\gamma} R_{zj}^0 + \frac{\omega - \gamma H}{\omega} \delta S_a R_{aj}^0
$$
\n
$$
(55)
$$

One can see, that the motion of the orbital anisotropy axis \bf{L} is small if the precession frequency is close to γH , and motion of ϑ is is small if the texture is close to the Leggett configuration $(\mathbf{n} \parallel \hat{\mathbf{z}})$.

In the equilibrium dipolar energy stays at the minimum, where $\cos \theta = -1/4$ (so-called Leggett angle). In the spin wave it goes out of the minimum and thus some additional energy appear. It depends on the orientation of n and thus can modify equilibrium texture. This effect causes self-localization of magnons $[12]$. From the expression (25) for the dipolar energy:

$$
F_D - F_D^0 = \frac{15}{2} g_D \Delta^2 \left(\mathbf{n} \cdot \theta^{\mathbf{s}} \right)^2 = \frac{15}{2} g_D \Delta^2 \delta \vartheta^2 \tag{56}
$$

Average change in the energy, caused by the transverse spin wave can be calculated using [\(54\)](#page-6-2):

$$
\langle F_D - F_D^0 \rangle = \frac{15}{4} g_D \Delta^2 \left(\frac{\gamma H}{\omega} \right)^2 \sin^2 \beta_N \beta_M^2 \tag{57}
$$

Another useful formula is a relation between equilibrium textural angles β_N and β_L . In the equilibrium $\hat{\mathbf{S}}^0 = \hat{\mathbf{z}}$ and

$$
\cos \beta_L = \hat{L}_z^0 = R_{zz}^0 = \cos \vartheta + (1 - \cos \vartheta) \cos^2 \beta_N. \tag{58}
$$

In the minimum of the dipolar energy

$$
\sin^2 \frac{\beta_L}{2} = \frac{5}{8} \sin^2 \beta_N. \tag{59}
$$

Quasiclassical equation for magnons

Consider case of short spin waves, in which spin changes on much smaller distances then the texture. We can represent such a wave as a set of flat waves $s_q \propto \cos(\mathbf{k} \cdot \mathbf{r})$ where wave vector k also changes slowly. Substituting the flat wave into (43) and (44) we get:

$$
c_{\perp}^2 \mathbf{k}^2 + (c_{\parallel}^2 - c_{\perp}^2)(\mathbf{k} \cdot \hat{\mathbf{L}}^0)^2 + \frac{1}{2}\Omega_B^2 \sin^2 \beta_N = \omega(\omega - \gamma H) \tag{60}
$$

$$
C_{\perp}^{2} \mathbf{k}^{2} + (C_{\parallel}^{2} - C_{\perp}^{2}) (\mathbf{k} \cdot \hat{\mathbf{L}}^{0})^{2} + \Omega_{B}^{2} \cos^{2} \beta_{N} = \omega^{2}
$$
 (61)

Here meaning of $c_{\perp,\parallel}$ and $C_{\perp,\parallel}$ becomes clear. In a short-wave limit where magnetic field and spinorbit interaction are not important $(\omega \gg \Omega_B, \gamma H)$ we have a linear dispersion laws where $c_{\perp,\parallel}$ are velocities of transverse waves, propagating perpendicular and parallel to $\hat{\mathbf{L}}^0$ direction; $C_{\perp,\parallel}$ are same velocities for longitudinal waves.

The first equation describes transverse spin waves, which are similar to that in other magnetic systems. In the presence of magnetic field it has two solutions $\omega(k)$, which are called acoustic (low ω) and optical (high ω) magnons. The second equation for longitudinal waves is unique for ³He.

Schrödinger equation for magnons

Consider a long-wave optical magnons with frequencies $\omega \approx \gamma H$ in the texture where **n** is close to vertical. In this case we can write equation for transverse spin waves (43) in a form of a Schrödinger equation for magnon quasiparticles, where complex transverse spin s_{+} plays role of a wave function and precessing frequency ω plays role of an energy. Effect of texture on the gradient terms is neglected because it adds only a small correction to the total gradient energy. The dipolar energy, which also depends on texture, can be of the same order as the total gradient energy.

$$
\left[-\frac{c_{\perp}^2}{\gamma H} \left(\nabla_x^2 + \nabla_y^2 \right) - \frac{c_{\parallel}^2}{\gamma H} \nabla_z^2 + \frac{\Omega_B^2}{2\gamma H} \sin^2 \beta_N + \gamma H \right] s_+ = \omega s_+ \tag{62}
$$

$$
s_{+} = \frac{1}{\sqrt{2}} \frac{\chi_B H}{\gamma} \sin \beta_M \ e^{i(\alpha_M + \omega t)} \tag{63}
$$

Both texture (spatial distribution of β_N) and magnetic field form a potential for magnons. It our experiment we used a combined effect of a flare-out texture in the cylindrical cell and non-uniform field of a small longitudinal coil to create a harmonic energy trap for magnons.

Magnons in a harmonic trap

Schrödinger equation in a cylindrical harmonic potential can be solved analytically. Let us write it in the form: \lceil

$$
-\frac{\hbar^2}{2m_{\perp}}\left(\nabla_x^2 + \nabla_y^2\right) - \frac{\hbar^2}{2m_{\parallel}}\nabla_z^2 + \hbar\omega_0 + \frac{m_{\perp}\omega_r^2}{2}r^2 + \frac{m_{\parallel}\omega_z^2}{2}z^2\right]s_+ = \hbar\omega s_+\tag{64}
$$

where parameters ω_0 , ω_r and ω_z describes the potential and masses $m_{\perp,\parallel}$ in the case of magnons are:

$$
m_{\perp, \parallel} = \frac{\hbar}{2c_{\perp, \parallel}^2} \gamma H. \tag{65}
$$

Then eigenvalues are

$$
\omega = \omega_0 + \omega_r (2n_r + |n_\phi| + 1) + \omega_z (n_z + 1/2), \tag{66}
$$

for $n_r, n_z = 0, 1, 2, \ldots, n_\phi = 0, \pm 1, \pm 2, \ldots$, and normalized solutions are:

$$
s_{+} = s_{+}^{r}(r) s_{+}^{\phi}(\phi) s_{+}^{z}(z), \qquad (67)
$$

where

$$
s_{+}^{r}(r) = \frac{1}{a_{r}} \sqrt{\frac{2 n_{r}!}{(n_{r} + |n_{\phi}|)!}} \left(\frac{r}{a_{r}}\right)^{|n_{\phi}|} \exp\left(-\frac{r^{2}}{2a_{r}^{2}}\right) L_{n_{r}}^{|n_{\phi}|}\left(\frac{r^{2}}{a_{r}^{2}}\right)
$$
(68)

$$
s_{+}^{\phi}(\phi) = \frac{1}{\sqrt{2\pi}} \exp(in_{\phi}\phi)
$$

\n
$$
s_{+}^{z}(z) = \sqrt{\frac{1}{a_{z}\sqrt{\pi}} \frac{1}{2^{n_{z}} n_{z}!}} \exp\left(-\frac{z^{2}}{2a_{z}^{2}}\right) H_{n_{z}}\left(\frac{z}{a_{z}}\right),
$$
\n(69)

 $L_n^m(x)$ and $H_n(x)$ are Laguerre and Hermite polynomials:

$$
L_n^m(x) = \frac{1}{n!} x^{-m} e^x \frac{d^n}{dx^n} (x^{n+m} e^{-x}), \qquad H_n(x) = (-1)^n e^{x^2} \frac{d^n}{dx^n} e^{-x^2}, \tag{70}
$$

and a_r and a_z are sizes of the wave:

$$
a_r = \sqrt{\frac{\hbar}{m_\perp \omega_r}} = c_\perp \sqrt{\frac{2}{\omega_r \gamma H}}, \qquad a_z = \sqrt{\frac{\hbar}{m_\parallel \omega_z}} = c_\parallel \sqrt{\frac{2}{\omega_z \gamma H}}.\tag{71}
$$

Usually we are interested only in non-antisymmetric states which have non-zero integral transverse magnetization and can be excited and observed in NMR experiments. For these states $n_{\phi} = 0$, $n_z =$ $0, 2, 4...,$ and

$$
s_{+}(r,z) = \left(\pi^{3/2} a_r^2 a_z \ 2^{n_z} \ n_z!\right)^{-1/2} \ \exp\left(-\frac{r^2}{2a_r^2} - \frac{z^2}{2a_z^2}\right) L_{n_r}^0\left(\frac{r^2}{a_r^2}\right) H_{n_z}\left(\frac{z}{a_z}\right),\tag{72}
$$

$$
\omega = \omega_0 + \omega_r (2n_r + 1) + \omega_z (n_z + 1/2), \qquad n_r = 0, 1, 2, \dots, n_z = 0, 2, 4 \dots \tag{73}
$$

Part 2. Non-linear spin dynamics

Leggett equations in R_{aj} coordinates

In this part non-linear Leggett equations of spin dynamics in ³He-B [\[10\]](#page-18-9) are derived. We start from same Hamiltonian as in the first part and use components of matrix R_{aj} as coordinates.

Order parameter of ³He-B is

$$
A_{aj} = \frac{1}{\sqrt{3}} \Delta e^{i\varphi} R_{aj}, \qquad (74)
$$

where φ is the phase, and R_{aj} is a rotation matrix with axis **n** and rotation angle ϑ :

$$
R_{aj}^{0} = \cos\vartheta \, \delta_{aj} + (1 - \cos\vartheta) \, n_a n_j - \sin\vartheta \, e_{ajk} n_k. \tag{75}
$$

Hamiltonian of this system is:

$$
\mathcal{H} = F_M + F_{SO} + F_{\nabla},\tag{76}
$$

$$
F_M = -(\mathbf{S} \cdot \gamma \mathbf{H}) + \frac{\gamma^2}{2\chi_B} \mathbf{S}^2,\tag{77}
$$

$$
F_{SO} = g_D \Delta^2 \Big[R_{jj} R_{kk} + R_{jk} R_{kj} - \frac{2}{3} R_{jk} R_{jk} \Big] = g_D \Delta^2 \Big[R_{jj} R_{kk} + R_{jk} R_{kj} \Big] + \text{const.}, \tag{78}
$$

$$
F_{\nabla} = \frac{1}{2} \Delta^2 \Big[K_1(\nabla_j R_{ak}) (\nabla_j R_{ak}) + K_2(\nabla_j R_{ak}) (\nabla_k R_{aj}) + K_3(\nabla_j R_{aj}) (\nabla_k R_{ak}) \Big]. \tag{79}
$$

Equations of motion are

$$
\dot{S}_a = \{ \mathcal{H}, S_a \} = \frac{\delta \mathcal{H}}{\delta S_b} \{ S_b, S_a \} + \frac{\delta \mathcal{H}}{\delta R_{bj}} \{ R_{bj}, S_a \} = \frac{\delta \mathcal{H}}{\delta S_b} e_{abc} S_c + \frac{\delta \mathcal{H}}{\delta R_{bj}} e_{abc} R_{cj}, \quad (80)
$$

$$
\dot{R}_{aj} = \{ \mathcal{H}, R_{aj} \} = \frac{\delta \mathcal{H}}{\delta S_b} \{ S_b, R_{aj} \} + \frac{\delta \mathcal{H}}{\delta R_{bk}} \{ R_{bk}, R_{aj} \} = \frac{\delta \mathcal{H}}{\delta S_b} e_{abc} R_{cj}.
$$
\n(81)

The derivatives of the Hamiltonian are much simpler then in the case of θ^s angles. There are no commutation problem in the variational derivative. The result is:

$$
\frac{\delta \mathcal{H}}{\delta \mathbf{S}} = -\gamma \mathbf{H} + \frac{\gamma^2}{\chi_B} \mathbf{S},\tag{82}
$$

$$
\frac{\delta \mathcal{H}}{\delta R_{bj}} = \Delta^2 \left[2g_D(\delta_{bj} R_{kk} + R_{jb} - 2/3R_{bj}) - K_1(\nabla_k \nabla_k R_{bj}) - (K_2 + K_3)(\nabla_k \nabla_j R_{bk}) \right].
$$
 (83)

Substituting this into equations of motion $(80,81)$ we obtain Leggett equations:

$$
\dot{S}_a = [\mathbf{S} \times \gamma \mathbf{H}]_a + T_a^D + T_a^{\nabla}, \n\dot{R}_{aj} = e_{abc} R_{cj} \left(\frac{\gamma^2}{\chi_B} \mathbf{S} - \gamma \mathbf{H} \right)_b, \tag{84}
$$

with dipolar and gradient torques:

$$
T_a^D = e_{abc} \frac{\delta F_D}{\delta R_{bj}} R_{cj} = 2\Delta^2 g_D \ e_{abc} R_{cj} (\delta_{bj} R_{kk} + R_{jb}),
$$

\n
$$
T_a^{\nabla} = e_{abc} \frac{\delta F_{\nabla}}{\delta R_{bj}} R_{cj} = -\Delta^2 e_{abc} R_{cj} \left[K_1 (\nabla_k \nabla_k R_{bj}) + (K_2 + K_3)(\nabla_k \nabla_j R_{bk}) \right].
$$
\n(85)

The gradient torque can be rewritten in terms of spin current J_{ak} which carries a component of the spin in the direction k :

$$
T_a^{\nabla} = -\nabla_k J_{ak}, \qquad J_{ak} = \Delta^2 e_{abc} R_{cj} \left[K_1 (\nabla_k R_{bj}) + (K_2 + K_3)(\nabla_j R_{bk}) \right]. \tag{86}
$$

Here we used equality $e_{abc}\nabla_k R_{bj}\nabla_k R_{cj} = e_{abc}\nabla_k R_{bj}\nabla_j R_{ck} = 0.$ It is also interesting to study motion of orbital anisotropy vector $L_j = S_a R_{aj}$:

$$
\dot{L}_k = \{ \mathcal{H}, S_a R_{ak} \} = \{ \mathcal{H}, S_a \} R_{ak} + \{ \mathcal{H}, R_{ak} \} S_a = \dot{S}_a R_{aj} + \dot{R}_{aj} S_a.
$$
 (87)

Substituting equations for \dot{S}_a and \dot{R}_{aj} we have:

$$
\dot{L}_k = (T_a^D + T_a^{\nabla}) R_{ak}.
$$
\n(88)

Leggett equations in n and ϑ coordinates

Let's rewrite Leggett equations in **n** and ϑ coordinates (axis and angle or rotation of the matrix R_{ai}). Time derivative of matrix $R(\mathbf{n}, \vartheta)$ is:

$$
\dot{R}_{aj} = \left[\sin \vartheta \left(n_a n_j - \delta_{aj} \right) - \cos \vartheta \left(e_{ajk} n_k \right) \dot{\vartheta} + (1 - \cos \vartheta) [n_j \dot{n}_a + n_a \dot{n}_j] - \sin \vartheta \left(e_{ajk} \dot{n}_k \right) \right] \tag{89}
$$

Let's calculate following combinations:

$$
n_a \dot{R}_{aj} + n_a \dot{R}_{ja} = 2(1 - \cos \vartheta) \dot{n}_j,
$$

\n
$$
R_{ja} \dot{R}_{aj} = -4 \cos \vartheta \sin \vartheta \dot{\vartheta}.
$$
\n(90)

By substituting R from [\(75\)](#page-9-1) and \dot{R} from [\(84\)](#page-9-2) we can write Leggett equations in \dot{n} and $\dot{\theta}$ coordinates:

$$
\dot{\mathbf{S}} = \mathbf{S} \times \gamma \mathbf{H} + \mathbf{T}^{D} - \nabla_{k} J_{ak},
$$
\n
$$
\dot{\mathbf{n}} = -\frac{1}{2} \mathbf{n} \times \left[\left(\frac{\gamma^{2}}{\chi_{B}} \mathbf{S} - \gamma \mathbf{H} \right) + \frac{\sin \vartheta}{1 - \cos \vartheta} \mathbf{n} \times \left(\frac{\gamma^{2}}{\chi_{B}} \mathbf{S} - \gamma \mathbf{H} \right) \right],
$$
\n
$$
\dot{\vartheta} = \mathbf{n} \cdot \left(\frac{\gamma^{2}}{\chi_{B}} \mathbf{S} - \gamma \mathbf{H} \right),
$$
\n(91)

Expression for dipolar torque via **n** and ϑ :

$$
\mathbf{T}^{D} = 4\Delta^{2} g_{D} \sin \vartheta (4 \cos \vartheta + 1) \mathbf{n}.
$$
 (92)

Spin current:

$$
J_{ak} = 2\Delta^2 K 1 \left[-n_a \nabla_k \vartheta - \sin \vartheta \nabla_k n_a + (1 - \cos \vartheta) e_{abc} n_c \nabla_k n_b \right]
$$
(93)
+ $\Delta^2 (K_2 + K_3) \left[-n_a \nabla_k \vartheta + \cos \vartheta \delta_{ak} n_j \nabla_j \vartheta + (1 - \cos \vartheta) n_k n_j n_a \nabla_j \vartheta + \cos \vartheta \sin \vartheta e_{abj} n_b n_k \nabla_j \vartheta + (1 - \cos \vartheta) \sin \vartheta e_{abk} n_b n_j \nabla_j \vartheta - \cos \vartheta \sin \vartheta e_{kbj} n_b n_a \nabla_j \vartheta - \cos \vartheta \sin \vartheta e_{akj} \nabla_j \vartheta + \sin \vartheta \cos \vartheta \delta_{ak} \nabla_j n_j + (1 - \cos \vartheta) \sin \vartheta n_a n_k \nabla_j n_j + (1 - \cos \vartheta) \sin \vartheta n_a n_j \nabla_j n_k - (1 - \cos \vartheta) \sin \vartheta n_k n_j \nabla_j n_a - \sin^2 \vartheta \delta_{ak} e_{cjb} n_b \nabla_j n_c + (1 - \cos \vartheta)^2 e_{abc} n_c n_j n_k \nabla_j n_b + (1 - \cos \vartheta) \cos \vartheta e_{abj} n_k \nabla_j n_b + (1 - \cos \vartheta) \cos \vartheta e_{abj} n_b \nabla_j n_k - \cos \vartheta \sin \vartheta \nabla_k n_a - (1 - \cos \vartheta) \sin \vartheta \nabla_a n_k + \sin^2 \vartheta e_{kj} n_b \nabla_a n_j \right].$ (93)

Expression for gradient torque can be obtain by finding spin current derivative [\(86\)](#page-9-3).

Spin created by motion of n and ϑ

It could be useful to solve second and third Leggett equations [\(91\)](#page-10-0) for spin. We use temporary notation

$$
\mathbf{s} = \frac{\gamma^2}{\chi_B} \mathbf{S} - \gamma \mathbf{H}, \qquad C = \frac{\sin \vartheta}{1 - \cos \vartheta}.
$$
 (94)

Then the equations are:

$$
\dot{\mathbf{n}} = -\frac{1}{2} \mathbf{n} \times [\mathbf{s} + C \mathbf{n} \times \mathbf{s}], \qquad \dot{\mathcal{V}} = \mathbf{n} \cdot \mathbf{s}
$$
 (95)

Let's solve them for s. Solution can be written as $s = n\dot{\theta} + n \times A$ with $A \cdot n = 0$. This satisfies the equation for $\dot{\theta}$ and splits vector s into two parts, parallel and perpendicular to **n**. Substitute s into the equation for $\dot{\mathbf{n}}$:

$$
\mathbf{A} + C\mathbf{n} \times \mathbf{A} = 2\dot{\mathbf{n}},\tag{96}
$$

find vector product of this equation with n , multiply it by C and subtract from original equation. From this we can find $\mathbf{A} = (1 - \cos \vartheta) \mathbf{\dot{n}} - \sin \vartheta \mathbf{n} \times \mathbf{\dot{n}}$ and get the result:

$$
\frac{\gamma^2}{\chi_B} \mathbf{S} = \gamma \mathbf{H} + \mathbf{n} \dot{\vartheta} + (1 - \cos \vartheta) \mathbf{n} \times \dot{\mathbf{n}} + \sin \vartheta \dot{\mathbf{n}}.
$$
 (97)

Here one can see how spin is created by magnetic field and motion of **n** and ϑ .

There is an interesting observation: if we have a soliton moving with a constant velocity, we can have a uniform spin distribution in the moving frame, but non-uniform in the static one.

Leggett equations for n and ϑ in the rotating frame

For numerical integration of Leggett equations [\(91\)](#page-10-0) it is convenient to switch to a "rotating frame" by doing substitution

$$
S_{a'}^r = U_{a'a} S_a, \quad T_{a'}^r = U_{a'a} T_a, \quad R_{a'k'}^r = U_{k'k} U_{a'a} R_{ak}, \quad \dots \tag{98}
$$

with rotation matrix $U_{jk} = R_{jk}(\omega t, \hat{\mathbf{z}}) = \cos \omega t \, \delta_{jk} + (1 - \cos \omega t) \, \hat{\mathbf{z}}_j \hat{\mathbf{z}}_k - \sin \omega t \, e_{jkl} \hat{\mathbf{z}}_l$.

For me it is still not clear is this "rotating frame" has a physical meaning (for example in 3D space we do not specify any axis of rotation). Anyway, we can always do the substitution, solve equations and return back to original vectors if needed.

By making the substitution in [\(84\)](#page-9-2) and using facts that $U\mathbf{a} \cdot U\mathbf{b} = \mathbf{a} \cdot \mathbf{b}$, $[U\mathbf{a} \times U\mathbf{b}] = U[\mathbf{a} \times \mathbf{b}]$ and $U\dot{U}^{-1}\mathbf{a} = \omega[\mathbf{a} \times \hat{\mathbf{z}}]$ we have equations in the rotating frame:

$$
\dot{\mathbf{S}}^{r} + \omega[\mathbf{S}^{r} \times \hat{\mathbf{z}}] = [\mathbf{S}^{r} \times \gamma \mathbf{H}^{r}] + \mathbf{T}^{Dr} + \mathbf{T}^{\nabla r}
$$
\n
$$
\dot{\mathbf{n}}^{r} + \omega[\mathbf{n}^{r} \times \hat{\mathbf{z}}] = -\frac{1}{2} \mathbf{n}^{r} \times \left[\left(\frac{\gamma^{2}}{\chi_{B}} \mathbf{S}^{r} - \gamma \mathbf{H}^{r} \right) + \frac{\sin \vartheta}{1 - \cos \vartheta} \mathbf{n}^{r} \times \left(\frac{\gamma^{2}}{\chi_{B}} \mathbf{S}^{r} - \gamma \mathbf{H}^{r} \right) \right]
$$
\n
$$
\dot{\vartheta} = \mathbf{n}^{r} \cdot \left(\frac{\gamma^{2}}{\chi_{B}} \mathbf{S}^{r} - \gamma \mathbf{H}^{r} \right).
$$
\n(99)

with dipolar torque

$$
\mathbf{T}^{Dr} = 4\Delta^2 g_D \sin \vartheta (4\cos \vartheta + 1) \mathbf{n}^r \tag{100}
$$

Writing the gradient torque in the rotating frame is not trivial, because matrix U will act on gradient operators. To avoid this difficulty we will be solving only 1D problems with all gradients directed along \hat{z} axis. Then spin current is not affected by rotation and we can simply write:

$$
\mathbf{T}^{\nabla r} = -\nabla_3 J_{a3} = -\mathbf{J}', \qquad J_a = \Delta^2 e_{abc} R_{cj} \left[K_1 R'_{bj} + (K_2 + K_3) R'_{b3} \delta_{j3} \right], \tag{101}
$$

using primes for ∇_3 . Usually we assume that magnetic field in the rotating frame \mathbf{H}^r is constant and choose axis \hat{x} in such a way that field is in $x - z$ plane:

$$
\gamma \mathbf{H}^{\mathbf{r}} = \hat{\mathbf{z}} \,\omega_z + \hat{\mathbf{x}} \,\omega_x \tag{102}
$$

This is a good description of a usual continuous-wave NMR experiments with static field $\gamma H_z = \omega_z$ and rotating radio-frequency field $\gamma H_x = \omega_x$.

Spin in the rotating frame

In the following calculations we will work only with vectors in the rotating frame and omit indices "r". It's convenient to introduce a dimensionless vector w:

$$
\mathbf{w} = \frac{1}{\omega} \left(\frac{\gamma^2}{\chi_B} \mathbf{S} - \gamma \mathbf{H} \right) + \hat{\mathbf{z}}
$$
(103)

Then Leggett equations will be written as

$$
\dot{\mathbf{w}} = \mathbf{w} \times (\hat{\mathbf{z}}(\omega_z - \omega) + \hat{\mathbf{x}} \omega_x) + \frac{\gamma^2}{\omega \chi_B} (\mathbf{T}^D + \mathbf{T}^\nabla)
$$
\n
$$
\dot{\mathbf{n}}/\omega = -\frac{1}{2} \mathbf{n} \times [(\mathbf{w} + \hat{\mathbf{z}}) + \frac{\sin \vartheta}{1 - \cos \vartheta} \mathbf{n} \times (\mathbf{w} - \hat{\mathbf{z}})]
$$
\n
$$
\dot{\vartheta}/\omega = \mathbf{n} \cdot (\mathbf{w} - \hat{\mathbf{z}}).
$$
\n(104)

We can solve second and third equations for w in the same way as (97) :

$$
\mathbf{w} = \cos \vartheta \, \hat{\mathbf{z}} + (1 - \cos \vartheta) \, \mathbf{n} \, n_z + \sin \vartheta \, \mathbf{n} \times \hat{\mathbf{z}} \n+ \frac{1}{\omega} \Big[\mathbf{n} \, \dot{\vartheta} + (1 - \cos \vartheta) \, \mathbf{n} \times \dot{\mathbf{n}} + \sin \vartheta \, \dot{\mathbf{n}} \Big]
$$
\n(105)

This can be also written as:

$$
\mathbf{w} = R\hat{\mathbf{z}} + \frac{1}{\omega} \left[\mathbf{n} \dot{\vartheta} + \frac{1 - \cos \vartheta}{\sin \vartheta} \left(R\dot{\mathbf{n}} + \dot{\mathbf{n}} \right) \right]
$$
(106)

Or in original notation:

$$
\frac{\gamma^2}{\chi_B} \mathbf{S} = \gamma \mathbf{H} + \mathbf{n} \dot{\vartheta} + (1 - \cos \vartheta) \mathbf{n} \times [\dot{\mathbf{n}} + \omega \mathbf{n} \times \hat{\mathbf{z}}] + \sin \vartheta [\dot{\mathbf{n}} + \omega \mathbf{n} \times \hat{\mathbf{z}}]
$$
(107)

$$
\frac{\gamma^2}{\chi_B} \mathbf{S} = \gamma \mathbf{H} + \omega \left(R \hat{\mathbf{z}} - \hat{\mathbf{z}} \right) + \mathbf{n} \dot{\vartheta} + \frac{1 - \cos \vartheta}{\sin \vartheta} \left(R \hat{\mathbf{n}} + \hat{\mathbf{n}} \right)
$$
(108)

Non-uniform equilibrium in the rotating frame

Equilibrium state can be found by setting all time derivatives to zero. From [\(106\)](#page-12-0) we have

$$
w_a = R_{a3},\tag{109}
$$

it means that in the equilibrium $|\mathbf{w}| = 1$.

Using this equation we can exclude w from the first equation and write equilibrium problem using only R_{aj} (or **n** and ϑ) coordinates:

$$
e_{abc}R_{bj}\left[\frac{\omega\chi_B}{\gamma^2\Delta^2}\left(\delta_{c3}\left(\omega_z-\omega\right)+\delta_{c1}\,\omega_x\right)\delta_{j3}-2g_D\,\left(\delta_{cj}R_{kk}+R_{jc}\right)+K_1R_{cj}''+\left(K_2+K_3\right)\delta_{j3}R_{c3}''\right]=0.\tag{110}
$$

Here we use expressions (85) for dipolar and gradient torques and keep only gradients in \hat{z} direction, represented by primes.

This can be written in term of Leggett frequency and spin-wave velocities:

$$
e_{abc}R_{bj}\left[\omega((\omega_z-\omega)\,\delta_{c3}+\omega_x\,\delta_{c1})\,\delta_{j3}-\frac{2}{15}\Omega_B^2\,\left(\delta_{cj}R_{kk}+R_{jc}\right)+\frac{1}{2}C_{\parallel}^2R_{cj}''+\left(C_{\perp}^2-C_{\parallel}^2\right)\delta_{j3}R_{c3}''\right]=0.\tag{111}
$$

This is a set of three equations for three parameters (**n** and ϑ). It describes non-uniform texture of a precessing state and can be used for calculating HPD boundaries or solitons.

Uniform equilibrium

Let's remove gradient terms in the first equation in (104) and use expression (100) for dipolar torque:

$$
\mathbf{w} \times (\hat{\mathbf{z}}(\omega_z - \omega) + \hat{\mathbf{x}} \omega_x) = \frac{4}{15} \frac{\Omega_B^2}{\omega} \sin \vartheta (4 \cos \vartheta + 1) \mathbf{n}, \tag{112}
$$

By multiplying it by w we have left-hand side equals zero and three cases: $\mathbf{n} \cdot \mathbf{w} = 0$, or $\cos \vartheta = -1/4$, or $\sin \theta = 0$. The last one is unstable equilibrium, it corresponds to maximum of dipolar energy. Two other cases are called HPD and NPD, now we can find them.

HPD (Homogeneously-precessing domain):

We already showed in [\(109\)](#page-12-1) that $\mathbf{w} = R\hat{\mathbf{z}}$. Using this, [\(112\)](#page-12-2), and HPD condition $\mathbf{n} \cdot \mathbf{w} = 0$ we can find

$$
\mathbf{n} = \pm \hat{\mathbf{y}}, \qquad \mathbf{w} = \cos \vartheta \; \hat{\mathbf{z}} + n_y \sin \vartheta \; \hat{\mathbf{x}}.
$$
 (113)

$$
\cos \vartheta = -\frac{1}{4} + \frac{15}{16} \frac{\omega}{\Omega_B^2} \left((\omega - \omega_z) + n_y \frac{\cos \vartheta}{\sin \vartheta} \omega_x \right). \tag{114}
$$

We can use only positive sign of ϑ because matrix $R(\mathbf{n}, \vartheta)$ is same as $R(-\mathbf{n}, -\vartheta)$. Later it will be shown that state with $\vartheta > 0$ and $\mathbf{n} = -\hat{\mathbf{y}}$ is unstable. To check it we will keep parameter $n_y = \pm 1$, which can be always set to 1 for HPD equilibrium state.

Usually RF-pumping ω_x is much smaller then frequency shift $\omega - \omega_z$, cos ϑ is close to -1/4 and we can use $\cos \vartheta / \sin \vartheta = -1/\sqrt{15}$ in the right part of the last equation:

$$
\cos \vartheta = -\frac{1}{4} + \frac{15}{16} \frac{\omega}{\Omega_B^2} \left(\omega - \omega_z - \frac{n_y}{\sqrt{15}} \omega_x \right). \tag{115}
$$

In arbitrary case to find ϑ we can do substitutions

$$
t = \tan \vartheta/2, \qquad A = \frac{15}{16} \frac{\omega(\omega - \omega_z)}{\Omega_B^2}, \qquad B = \frac{15}{16} \frac{\omega}{\Omega_B^2} n_y \omega_x,\tag{116}
$$

and obtaining 4-th order equation for t:

$$
Bt^4 - \frac{1}{2}(3+4A)t^3 + \frac{1}{2}(5-4A)t - B = 0.
$$
 (117)

NPD (Non-precessing domain), Brinkman-Smith mode:

Using the NPD condition $\cos \theta = -1/4$ we have:

$$
w_1 = \frac{-\omega_x}{\sqrt{\omega_x^2 + (\omega_z - \omega)^2}}, \qquad w_2 = 0, \qquad w_3 = \frac{-(\omega_z - \omega)}{\sqrt{\omega_x^2 + (\omega_z - \omega)^2}}, \tag{118}
$$

$$
n_1 = \frac{w_1 n_3}{1 + w_3}, \qquad n_2 = \frac{w_1 \sqrt{3/5}}{1 + w_3}, \qquad n_3 = \pm (4w_3 + 1)/5. \tag{119}
$$

(There is also solution with opposite signs of w_1 and w_3 which corresponds to unstable equilibrium).

Here n is precessing around \hat{z} axis. This is case of a "vertical" texture. In real NMR experiments if RF pumping is small enough then texture is determined by boundary conditions in the experimental cell and possible topological defects such as n-solitons. It can have any equilibrium orientation of n vector and small oscillations around this equilibrium. These solutions can not be found as an equilibrium in our rotating frame.

Small oscillations of HPD in the rotating frame

This section follows calculations in my PhD thesis (2012, available only in Russian).

Let's again consider a uniform case with zero gradient terms. Introduce dimensionless parameters:

$$
d = \frac{\omega - \omega_z}{\omega}, \qquad h = \frac{\omega_x}{\omega}, \qquad b = \frac{15\gamma^2 \Delta^2 g_D}{\chi_B \omega^2} = \frac{\Omega_B^2}{\omega^2}
$$
(120)

Then equations [\(104\)](#page-11-0) will be

$$
\dot{\mathbf{w}}/\omega = \mathbf{w} \times (h\,\hat{\mathbf{x}} - d\,\hat{\mathbf{z}}) + \frac{4b}{15} \sin \vartheta (4\cos \vartheta + 1) \mathbf{n}
$$
\n
$$
\dot{\mathbf{n}}/\omega = -\frac{1}{2} \mathbf{n} \times (\mathbf{w} + \hat{\mathbf{z}}) - \frac{1}{2} \frac{\sin \vartheta}{1 - \cos \vartheta} \left[\mathbf{n} \left(\mathbf{n} \cdot (\mathbf{w} - \hat{\mathbf{z}}) \right) - (\mathbf{w} - \hat{\mathbf{z}}) \right],
$$
\n
$$
\dot{\vartheta}/\omega = \mathbf{n} \cdot (\mathbf{w} - \hat{\mathbf{z}}),
$$
\n(121)

(In my PhD thesis slightly different definition of dimensionless parameters were used. Also there was an error in equilibrium magnetization values, but it does not change the answer for small oscillation frequencies.)

The equilibrium which corresponds to HPD [\(113,](#page-12-3)[115\)](#page-12-4) is

$$
\mathbf{n}^0 = \pm \hat{\mathbf{y}}, \qquad \mathbf{w}^0 = \cos \vartheta \ \hat{\mathbf{z}} + n_y \sin \vartheta \ \hat{\mathbf{x}} \qquad \cos \vartheta^0 = -\frac{1}{4} - \frac{15}{16} \left(d - \frac{h \, n_y}{\sqrt{15}} \right) \frac{1}{b} \tag{122}
$$

Let's rewrite equation (121) in coordinates:

$$
\begin{array}{rcl}\n\dot{w}_x/\omega & = & Fb \; n_x - w_y d \\
\dot{w}_y/\omega & = & Fb \; n_y + w_z h + w_x d \\
\dot{w}_z/\omega & = & Fb \; n_z - w_y h \\
2\dot{n}_x/\omega & = & -n_y(w_z + 1) + n_z w_y - C \; n_x \left(\mathbf{n} \cdot (\mathbf{w} - \hat{\mathbf{z}}) \right) + C \; w_x \\
2\dot{n}_y/\omega & = & -n_z w_x + n_x(w_z + \omega) - C \; n_y \left(\mathbf{n} \cdot (\mathbf{w} - \hat{\mathbf{z}}) \right) + C \; w_y \\
2\dot{n}_z/\omega & = & -n_x w_y + n_y w_x - C \; n_z \left(\mathbf{n} \cdot (\mathbf{w} - \hat{\mathbf{z}}) \right) + C \left(w_z - 1 \right) \\
\dot{\vartheta}/\omega & = & \mathbf{n} \cdot (\mathbf{w} - \hat{\mathbf{z}}).\n\end{array} \tag{123}
$$

where

$$
C = \frac{\sin \vartheta}{1 - \cos \vartheta}, \qquad F = \frac{4}{15} \sin \vartheta (4 \cos \vartheta + 1). \tag{124}
$$

Three equations for \dot{n} , are dependent because $|\mathbf{n}| = 1$. This can be checked by calculating $n_x \dot{n}_x +$ $n_y \dot{n}_y + n_z \dot{n}_z.$

Lets write the equations for small deviations $d\mathbf{n}$, $d\mathbf{w}$, $d\vartheta$ from the equilibrium. Here we keep only first-order terms with the deviations. Also we put $dn_y = 0$ and omit equation for n_y because **n** is moving near $\hat{\mathbf{y}}$:

$$
d\dot{w}_x/\omega = F^0 b \, dn_x - d \, dw_y,
$$

\n
$$
d\dot{w}_y/\omega = dF b \, n_y^0 + h \, dw_z + d \, dw_x,
$$

\n
$$
d\dot{w}_z/\omega = F^0 b \, dn_z - h \, dw_y,
$$

\n
$$
2d\dot{n}_x/\omega = -n_y^0 \, dw_z + C^0 \, dw_x + w_x^0 \, dC,
$$

\n
$$
2d\dot{n}_z/\omega = n_y^0 \, dw_x + C^0 \, dw_z + (w_z - 1) \, dC,
$$

\n
$$
d\dot{v}/\omega = w_x^0 \, dn_x + (w_z^0 - 1) \, dn_z + n_y^0 \, dw_y.
$$
\n(125)

For periodic motion with dimensionless frequency $x = \Omega/\omega$ this gives an equation

$$
\begin{vmatrix}\n-ix & -d & 0 & bF^0 & 0 & 0 \\
d & -ix & h & 0 & 0 & n_y^0 b dF/d\vartheta \\
0 & -h & -ix & 0 & bF^0 & 0 \\
C & 0 & -n_y & -2ix & 0 & n_y^0 \sin \vartheta dC/d\vartheta \\
n_y & 0 & C & 0 & -2ix & (\cos \vartheta^0 - 1) dC/d\vartheta \\
0 & n_y^0 & 0 & w_x^0 & (\cos \vartheta^0 - 1) & -ix\n\end{vmatrix} = 0
$$
\n(126)

If $d, h \ll b \ll 1$, then the equation is

 $\bigg\}$ $\bigg\}$ $\overline{}$ $\overline{}$ $\overline{}$ $\overline{}$ $\overline{}$ $\overline{}$ $\overline{}$ $\overline{}$ \vert

$$
-x^{6} + (1+b)x^{4} - \frac{1}{\sqrt{15}}\left(\frac{3}{8}(\sqrt{15}d - n_{y}h) + b(\sqrt{15}d + 3n_{y}h)\right)x^{2} + \frac{1}{10}(\sqrt{15}d - n_{y}h) \,bn_{y}h = 0; \quad (127)
$$

where $n_y^0 = \pm 1$ and ϑ^0 is positive (changing sign of ϑ is equivalent to changing sign of **n**). Solutions are

$$
x_1^2 = 1 + b,
$$

\n
$$
x_2^2 = \frac{4}{\sqrt{15}} \frac{n_y^0 h b}{1 + 8/3 b},
$$

\n
$$
x_3^2 = \frac{\sqrt{15} d - n_y^0 h}{\sqrt{15}} \frac{3/8 + b}{1 + b}.
$$
\n(128)

One can see that $n_y^0 = -1$ corresponds to unstable state $(x_2^2 < 0)$ and we can use $n_y^0 = +1$. In original notations

$$
\Omega_1^2 = \omega^2 + \Omega_B^2,
$$
\n
$$
\Omega_2^2 = \frac{4}{\sqrt{15}} \frac{\omega_x \omega \Omega_B^2}{\omega^2 + 8/3 \Omega_B^2},
$$
\n
$$
\Omega_3^2 = \frac{\sqrt{15} (\omega - \omega_z) \omega - \omega_x \omega}{\sqrt{15}} \frac{3/8 \omega^2 + \Omega_B^2}{\omega^2 + \Omega_B^2}.
$$
\n(129)

The second mode is a useful tool for calibrating RF pumping ω_x or measuring Leggett frequency Ω_B . The third mode as fas as I know have not been observed in experiments.

Leggett-Takagi relaxation

Leggett-Takagi equations [\[13\]](#page-18-12) can be made from Leggett equations [\(91\)](#page-10-0) by adding a relaxation term with parameter τ to the equation for $\dot{\vartheta}$.

$$
\dot{\vartheta} = \mathbf{n} \cdot \left(\frac{\gamma^2}{\chi_B} \mathbf{S} - \gamma \mathbf{H}\right) + \frac{4}{15} \sin \vartheta (4 \cos \vartheta + 1) \frac{1}{\tau}.
$$
 (130)

Switching to the rotating frame and using dimensionless parameters \mathbf{w}, d, h, b [\(103,](#page-11-2) [120\)](#page-13-1) we can write equations of motion [\(121\)](#page-13-0) with the relaxation term:

$$
\dot{\mathbf{w}}/\omega = \mathbf{w} \times (-d\hat{\mathbf{z}} + h\hat{\mathbf{x}}) + \frac{4}{15}b \sin \vartheta (4 \cos \vartheta + 1) \mathbf{n}
$$
(131)
\n
$$
\dot{\mathbf{n}}/\omega = -\frac{1}{2} \mathbf{n} \times (\mathbf{w} + \hat{\mathbf{z}}) - \frac{1}{2} \frac{\sin \vartheta}{1 - \cos \vartheta} \left[\mathbf{n} (\mathbf{n} \cdot (\mathbf{w} - \hat{\mathbf{z}})) - (\mathbf{w} - \hat{\mathbf{z}}) \right]
$$

\n
$$
\dot{\vartheta}/\omega = \mathbf{n} \cdot (\mathbf{w} - \hat{\mathbf{z}}) + \frac{4}{15} \sin \vartheta (4 \cos \vartheta + 1) \frac{1}{\omega \tau},
$$

or in coordinates with C and F defined in (124) :

$$
\dot{w}_x/\omega = -d w_y + bF n_x
$$
\n
$$
\dot{w}_y/\omega = h w_z + d w_x + bF n_y
$$
\n
$$
\dot{w}_z/\omega = -h w_y + bF n_z
$$
\n
$$
2\dot{n}_x/\omega = -n_y(w_z + 1) + n_z w_y - C n_x (\mathbf{n} \cdot (\mathbf{w} - \hat{\mathbf{z}})) + C w_x
$$
\n
$$
2\dot{n}_y/\omega = -n_z w_x + n_x(w_z + 1) - C n_y (\mathbf{n} \cdot (\mathbf{w} - \hat{\mathbf{z}})) + C w_y
$$
\n
$$
2\dot{n}_z/\omega = -n_x w_y + n_y w_x - C n_z (\mathbf{n} \cdot (\mathbf{w} - \hat{\mathbf{z}})) + C(w_z - 1)
$$
\n
$$
\dot{\vartheta}/\omega = \mathbf{n} \cdot (\mathbf{w} - \hat{\mathbf{z}}) + \frac{F}{\omega \tau}.
$$
\n(132)

For the equilibrium state we have:

$$
\mathbf{n}^{0} \cdot \mathbf{w}^{0} = 0, \qquad n_{z}^{0} = \frac{F^{0}}{\omega \tau}, \qquad n_{x}^{0} = \frac{d}{h} \frac{F^{0}}{\omega \tau}, \qquad w_{y}^{0} = \frac{b}{h} \frac{(F^{0})^{2}}{\omega \tau}.
$$
 (133)

One can see that equilibrium position of vector **n** is rotated from \hat{y} direction by Leggett-Takagi relaxation, and vector \mathbf{w}^0 stays perpendicular to \mathbf{n}^0 . Note that even at low relaxation n_x^0 can be large because of d/h factor. This probably determines maximum frequency shift where HPD can exist for a given RF pumping and Leggett-Takagi relaxation. It is very interesting to measure this effect in experiment.

θ -solitons

Minimum of spin-orbit interaction corresponds to $\cos \theta = -1/4$. It is possible to have a so-called theta-soliton [\[14\]](#page-18-13) there ϑ changes between two minima, $\vartheta_L = \cos^{-1}(-1/4)$ and $2\pi - \vartheta_L$, as a solution of equilibrium Leggett equations.

Let's assume $\mathbf{n} = \hat{\mathbf{z}}, \, \mathbf{H} = H_z \hat{\mathbf{z}}$, and $\mathbf{S} = \chi_B \mathbf{H}/\gamma$. Only ϑ is changing along $\hat{\mathbf{z}}$ axis.

Using usual expression $R_{aj} = \cos \vartheta \, \delta_{aj} + (1 - \cos \vartheta) \, n_a n_j - \sin \vartheta \, e_{ajk} n_k$, and $n_j = \delta_{j3}$ one can find that only J_{33} component of spin current is non-zero:

$$
J_{33} = -2\Delta^2 K_1 \vartheta' \tag{134}
$$

Second and third of Leggett equations are satisfied by chosen values of **n** and **S**. The first equation gives non-trivial expression for distribution of ϑ :

$$
T_3^D = J'_{33} \tag{135}
$$

By substituting spin current and dipolar torque we obtain a second-order differential equation for ϑ :

$$
\vartheta'' = -\frac{2g_D}{K1} \sin \vartheta (4\cos \vartheta + 1) \tag{136}
$$

or with usual definition of dipolar length $\xi_D^2 = \frac{13}{24} \frac{K_1}{g_D}$.

$$
\frac{12}{13}\xi_D^2 \ \vartheta'' = -\sin\vartheta (4\cos\vartheta + 1) \tag{137}
$$

To solve this equation we multiply both sides by ϑ' and integrate using boundary conditions at $z = \pm \infty$: cos $\vartheta = -1/4$, $\vartheta' = 0$:

$$
\sqrt{\frac{12}{13}}\xi_D \,\vartheta' = \pm 2(\cos\vartheta + 1/4) \tag{138}
$$

Second integration will give us formula of the ϑ -soliton. Integration constant determines position of the soliton, we choose it to localize the soliton at $z = 0$. There are two solutions, one goes from $-\theta_L$ to $+\theta_L$, another from $+\theta_L$ to $2\pi - \theta_L$. To find the second solution one can make substitution $\bar{\theta} = \theta - \pi$ and do similar integration. Result is:

$$
\vartheta = 2 \tan^{-1} \left[\sqrt{\frac{5}{3}} \tanh \left(\sqrt{\frac{65}{64}} \frac{\pm z}{\xi_D} \right) \right], \qquad \vartheta = \pi + 2 \tan^{-1} \left[\sqrt{\frac{3}{5}} \tanh \left(\sqrt{\frac{65}{64}} \frac{\pm z}{\xi_D} \right) \right].
$$
 (139)

The first solution is not topologically protected because it is possible to move from $-\theta_L$ to $+\theta_L$ by rotating **vector, always staying in the minimum of spin-orbit interaction. If vector** $**n**$ **is oriented along** \hat{z} for example by magneto-dipolar energy (not discussed in this text yet) or by spin precession, it will form a so-called n-soliton which has much smaller energy and bigger size. The second solution is the ϑ -soliton.

 ϑ -solitons can exist in HPD state. In this case they have small core where ϑ is changing and large tails where n vector is rotated. This have been discussed in [\[15\]](#page-18-14) and numerically calculated in [\[16\]](#page-18-15).

Motion of arbitrary soliton (in the rotating frame)

Consider a soliton with some distribution $n(z)$ and $\vartheta(z)$ which are equilibrium solution of Leggett equations.

Consider a soliton which has the same distribution of n and ϑ and moves with velocity v. For the moving soliton we can write $n(z, dt) = n(z + v dt)$ and $\dot{n}(z) = n' v$, and same for theta. Now put this into [\(108\)](#page-12-5) and find spin S distribution of the moving soliton:

$$
\frac{\gamma^2}{\chi_B} \mathbf{S}(v) = \left[\mathbf{n} \; \vartheta' + (1 - \cos \vartheta) \; \mathbf{n} \times \mathbf{n}' + \sin \vartheta \; \mathbf{n}' \right] v + \gamma \mathbf{H} + \omega \; (R\hat{\mathbf{z}} - \hat{\mathbf{z}}) \tag{140}
$$

Calculate kinetic energy of the soliton using [\(77\)](#page-9-5), as an energy difference between moving and static soliton. Note that expression for energy is same in the rotating frame:

$$
E_K = \int_{-\infty}^{\infty} \left[-(\mathbf{S}(v) \cdot \gamma \mathbf{H}) + \frac{\gamma^2}{2\chi_B} \mathbf{S}(v)^2 \right] dz - \int_{-\infty}^{\infty} \left[-(\mathbf{S}(0) \cdot \gamma \mathbf{H}) + \frac{\gamma^2}{2\chi_B} \mathbf{S}(0)^2 \right] dz \tag{141}
$$

$$
= -\frac{\chi_B}{\gamma^2} \omega v \int_{-\infty}^{\infty} 2(1 - \cos \vartheta) [\mathbf{n} \times \mathbf{n}']_z dz + \frac{\chi_B}{\gamma^2} \frac{v^2}{2} \int_{-\infty}^{\infty} [\mathbf{n} \; \vartheta' + (1 - \cos \vartheta) \; \mathbf{n} \times \mathbf{n}' + \sin \vartheta \; \mathbf{n}']^2 dz
$$

The first term is proportional to ωv , which is interesting: it says that if the soliton in the rotating frame has a helical structure where \bf{n} rotates around $\hat{\bf{z}}$ axis then energy minimum corresponds to a non-zero velocity. The second term is proportional to v^2 , from this term we can get effective mass of the soliton:

$$
m = \frac{\chi_B}{\gamma^2} \int_z \left[(\vartheta')^2 + 2(1 - \cos \vartheta) (\mathbf{n}')^2 \right] dz \tag{142}
$$

This calculation is valid for small velocities. Arbitrary motion of ϑ -solitons have been calculated in [\[17\]](#page-18-16).

Motion of ϑ -solitons

Let's look at a 3D case: a soliton perpendicular to \hat{z} direction in an experimental cell. It can be considered as a membrane attached to walls because of some pinning effects. Let's calculate effective mass and tension of this membrane using profile of the soliton $\vartheta(z)$ from [\(139\)](#page-16-0).

Tension of the soliton membrane is its total potential energy (spin-orbit + gradient) per unit area. If n is uniform the gradient energy [\(79\)](#page-9-5) is

$$
E_{\nabla} = \frac{1}{2} \Delta^2 \left[(2K_1 + K_2 + K_3)(\nabla \vartheta)^2 - (K_2 + K_3)(\mathbf{n} \cdot \nabla \vartheta)^2 \right].
$$
 (143)

For the soliton [\(136](#page-15-0)[-139\)](#page-16-0) it is

$$
F_{\nabla} = \Delta^2 K_1 (\vartheta')^2 = 8\Delta^2 g_D (\cos \vartheta + 1/4)^2
$$
\n(144)

Energy of spin-orbit interaction [\(78\)](#page-9-5) is also

$$
F_{SO} = 8\Delta^2 g_D(\cos\vartheta + 1/4)^2\tag{145}
$$

And tension is integral of total energy density over z :

$$
T = 16\Delta^2 g_D \int_{-\infty}^{+\infty} (\cos \vartheta(z) + 1/4)^2 dz
$$
 (146)

Mass of the soliton per unit area can be found by integrating [\(142\)](#page-16-1):

$$
M = 8\frac{\chi_B}{\gamma^2} \frac{g_D}{K_1} \int_{-\infty}^{+\infty} (\cos\vartheta + 1/4)^2 dz \tag{147}
$$

Speed of sound is square root of tension divided by mass:

$$
C^{2} = 2\frac{\Delta^{2}\gamma^{2}}{\chi_{B}}K_{1} = 2c_{\perp}^{2} - c_{\parallel}^{2}
$$
\n(148)

The first oscillation mode of a circular membrane with radius R is $2.405 C/R$.

References

- 1. Vollhardt, D. & Wölfle, P. The superfluid phases of helium 3 (Taylor and Francis, London, 1990).
- 2. <https://cadabra.science/>.
- 3. <https://github.com/slazav/he3lib>.
- 4. Dzyaloshinskii, I. & Volovick, G. Poisson brackets in condensed matter physics. Annals of Physics 125, 67 –97. [https://doi.org/10.1016/0003-4916\(80\)90119-0](https://doi.org/10.1016/0003-4916(80)90119-0) (1980).
- 5. Bunkov, Y. M. & Golo, V. L. Spin-Orbital Dynamics in the B-Phase of Superfluid Helium-3. Journal of Low Temperature Physics 137. <https://doi.org/10.1007/s10909-004-0897-4> (5-6 2004).
- 6. Theodorakis, S. & Fetter, A. L. Vortices and NMR in Rotating ³He-B.
- 7. Haar, D. T. Elements of Hamiltonian mechanics (Pergamon Press, 1971).
- 8. Bunkov, Y. M., L'vov, V. S. & Volovik, G. E. Solution of the problem of catastrophic relaxation of homogeneous spin precession in superfluid ³He-B. JETP Letters 83. [https://doi.org/10.1134/](https://doi.org/10.1134/s0021364006120022) [s0021364006120022](https://doi.org/10.1134/s0021364006120022) (12 2006).
- 9. Cross, M. A generalized Ginzburg-Landau approach to the superfluidity of helium 3. English. Journal of Low Temperature Physics 21, 525-534. <https://doi.org/10.1007/BF01141607> (1975).
- 10. Leggett, A. J. A theoretical description of the new phases of liquid ³He. Rev. Mod. Phys. 47, 331– 414. <https://doi.org/10.1103/RevModPhys.47.331> (2 1975).
- 11. Fomin, I. A. Large-amplitude spin waves and magnetic relaxation in the superfluid phases of ³He. JETP 51, 1203–1212 (6 1980).
- 12. Autti, S. et al. Self-Trapping of Magnon Bose-Einstein Condensates in the Ground State and on Excited Levels: From Harmonic to Box Confinement. Phys. Rev. Lett. 108, 145303. [https://doi.](https://doi.org/10.1103/PhysRevLett.108.145303) [org/10.1103/PhysRevLett.108.145303](https://doi.org/10.1103/PhysRevLett.108.145303) (14 2012).
- 13. Leggett, A. & Takagi, S. Orientational dynamics of superfluid 3He: A two-fluid model. I. Spin dynamics with relaxation. Annals of Physics 106, 79-133. [https://doi.org/10.1016/0003-](https://doi.org/10.1016/0003-4916%2877%2990005-7) [4916%2877%2990005-7](https://doi.org/10.1016/0003-4916%2877%2990005-7) (1 1977).
- 14. Maki, K. & Kumar, P. Magnetic solitons in superfluid ³He. Phys. Rev. B 14, 118–127. [https:](https://doi.org/10.1103/PhysRevB.14.118) [//doi.org/10.1103/PhysRevB.14.118](https://doi.org/10.1103/PhysRevB.14.118) (1 1976).
- 15. Misirpashaev, T. S. & Volovik, G. Topology of coherent precession in superfluid ³He-B. JETP 75, 650–665. http://inis.iaea.org/search/search.aspx?orig_q=RN:24044181 (4 1992).
- 16. Zavjalov, V. V. Dynamics of ϑ -solitons in the HPD state of superfluid ${}^{3}He-B$ 2022. https:// arxiv.org/abs/2212.12232.
- 17. Rozhkov, S. S. Dynamics of the order parameter of superfluid phases of helium-3. Soviet Physics Uspekhi 29, 186–198. <https://doi.org/10.1070/pu1986v029n02abeh003163> (1986).