Properties of 2D anyon gas

Douglas Lundholm¹

¹Department of Mathematics, Uppsala University, Box 480, SE-751 06, Uppsala, Sweden

An overview is given of the 2D many-anyon gas, including its definition (both for ideal and certain less-than-ideal particles, as well as for abelian and nonabelian braid group representations), its corresponding known properties starting out from the intricate relationship between exchange and exclusion, as well as its emergence from bosons and/or fermions in 3D. For the Encyclopedia of Condensed Matter Physics, 2e. Date: June, 2023

PACS numbers: 05.30.Pr, 03.75.Lm, 71.15.Mb, 73.43.-f Keywords: quantum statistics, anyons, intermediate and fractional statistics, braid group representations, statistics transmutation, exclusion principle, density functional theory, Thomas-Fermi approximation

Key points/objectives

• How can the 2D anyon gas be defined physically and modelled mathematically?

• What are the essential properties of the anyon gas? Does it admit some notion of exclusion statistics intermediate to bosons and fermions?

• How can an anyon gas emerge in a realistic physical system consisting of only bosons and/or fermions?

I. INTRODUCTION

Quantum statistics refers to the symmetry and organizing principle assumed by the components of a quantum system. Primarily we think of particles, described by a joint state or wave function, which can fall into certain symmetry classes due to the fundamental constraints on the information that can be extracted from the system, such as those imposed by the uncertainty principle. Bosons and fermions are ensembles of identical such particle subsystems characterized by symmetry, respectively antisymmetry, with respect to particle exchange. In their ideal limits they manifest the statistical state distributions defined by Bose and Einstein, respectively Fermi and Dirac, with the latter subject to Pauli's exclusion principle. In a sense, whereas bosons unify, fermions individualize. Anyons are defined as identical quantum particles with an intermediate exchange symmetry characterized by representations of the braid group, which is the relevant group of continuous exchanges of particle configurations in 2D (two spatial dimensions). An anyon gas is a quantum system consisting of a large number of such particles subject to a fixed class of exchange symmetries, which is dependent on the specific underlying kinematics and dynamics of the quantum systems from which it emerged. The study of the behavior of the proper anyon gas presents a significant challenge to mathematics and physics. Our aim with this overview is to make these statements precise and to give an updated status report on some of the many facets of the topic.



FIG. 1. Three different perspectives on anyons.

A. Brief historic overview

For completeness and to set our reference point for this overview, we give a very brief account on the origins of the anyon gas, referring to [Biedenharn et al. 1990, Fröhlich 1990, Goldin 2022, Leinaas and Myrheim 2022] for further background.

1. Three perspectives on anyons

The discovery of anyons and intermediate quantum statistics in two spatial dimensions may be attributed to three independent research groups, working from three distinctly different perspectives¹. Chronologically, these are (cf. Fig. 1):

1. Geometric: In 1977, Leinaas and Myrheim used geometric methods from gauge theory, fiber bundles, and extensions of the kinetic energy operator via boundary conditions (also known as "Schrödinger quantization") to arrive at the possibility for what they termed "intermediate statistics" in 2D (as well as in 1D), and worked out some of its detailed consequences for two particles [Leinaas and Myrheim 1977]. They further pointed to the role of the fundamental group of the configuration space for arbitrary numbers of identical particles and the possibility for higher-dimensional fibers (necessary

¹ These three groups were initially unaware of each other's earlier contributions and their initial works therefore lack the respective citations. There have subsequently been attempts to remedy this in the literature [Biedenharn et al. 1990].

for nonabelian representations, but they did not explicitly mention the braid group or its representation theory in their work).

2. Algebraic: In 1980-'81, starting out from the representation theory of the Lie algebra of local currents on \mathbb{R}^d and the group of diffeomorphisms (akin to "Heisenberg quantization" and Dirac's canonical quantization scheme [Dirac 1967]). Goldin, Menikoff and Sharp found a kinematical derivation of the possibilities for quantum statistics with an unexpected dependence on the dimension d [Goldin et al. 1981, 1980]. In [Goldin et al. 1981] they also explored connections to the Aharonov-Bohm effect (in 3D), while subsequently in 1983 they highlighted the role of the one-dimensional unitary representations of the braid group [Goldin et al. 1983], and in 1985 pointed out that higher-dimensional braid group representations induce inequivalent representations of the current algebra [Goldin et al. 1985]. Such identical particles were later termed "plektons" (see, e.g., [Dell'Antonio et al. 1997, Korff et al. 1999, Mund and Schrader 1995]) or simply "nonabelian anyons" (nowadays the more common term; cf. [Nayak et al. 2008]). It was also pointed out in [Goldin et al. 1985] that nontrivial exchange phases in 2D do not require indistinguishability; the *colored* braid group serves as the fundamental group for configurations of *distinguishable* particles. The general algebraic (and potentially nonabelian) approach to quantum statistics was further developed in the field theory direction by Tsuchiya and Kanie [Tsuchiya and Kanie 1987, 1988], Kohno [Kohno 1987], as well as Fröhlich, Gabbiani, Kerler and Marchetti [Fröhlich and Gabbiani 1990, Fröhlich and Marchetti 1988, 1989, Fröhlich 1988, 1990, Fröhlich and Kerler 1993], and Fredenhagen, Rehren and Schroer [Fredenhagen et al. 1989].

3. Magnetic: In 1982, motivated by topological gauge theory examples in higher dimensions (" θ -vacua" and "dyons"), Wilczek considered a composite of a 2D magnetic flux and a charged particle, coined for it the name "anyon", as well as predicted its fractional spin and worked out the basic quantum mechanics for two such composite particles [Wilczek 1982a,b]. The same line of investigation was continued by Wu in 1984, who connected it to path integrals, extended it to arbitrary numbers of particles, and independently and explicitly invoked the braid group and its one-dimensional representations [Wu 1984a,b]. Further, Arovas, Schrieffer, Wilczek and Zee considered in 1985 the statistical mechanics of a gas of such abelian anyons by means of its lowest-order virial expansion [Arovas et al. 1985] (see also [Dowker 1985]).

The first concrete physical application of intermediate/fractional statistics and anyons came from the magnetic perspective and concerned the **fractional quantum Hall effect** (FQHE) [Laughlin 1999, Stormer 1999, Tsui 1999], in which the emergent fractionally charged quasiparticles/holes introduced by Laughlin were proposed by Halperin to have such properties [Halperin 1984]. Arovas, Schrieffer and Wilczek subsequently verified this [Arovas et al. 1984] (though under tacit assumptions of adiabaticity [Forte 1991]) by computing the corresponding Berry phase for quasiholes, thereby converging the geometric and magnetic (and eventually algebraic) perspectives. Further, the latent power of the algebraic perspective was brought to light in the new millennium as Kitaev proposed the usefulness of nonabelian anyons in topological quantum computation [Freedman et al. 2003, Kitaev 2003, 2006]. Most of the development of the field throughout this time span has been covered in the books and reviews [Canright and Johnson 1994, Date et al. 2003, Forte 1992, Fröhlich 1990, Iengo and Lechner 1992, Jackiw 1990, Khare 2005, Lerda 1992, Myrheim 1999, Nayak et al. 2008, Ouvry 2007, Stern 2008, Wilczek 1990].

2. Precursor ideas

As is evident from the above, anyons and intermediate ("fractional") quantum statistics are the convergence of many central ideas and concepts in mathematical physics. The former notion (anyon) is more closely tied into the concepts of exchange phases and symmetry classes of wave functions, while the latter (fractional statistics) to that of the **exclusion principle** and permissible probability distributions for particles. It has been—and still is—a nontrivial task to rigorously connect these two notions in the strictly intermediate case (see, e.g., the review [Canright and Johnson 1994]). Exploration of intermediate statistics from the latter approach of exclusion with a finite occupation number of one-body states was considered by Gentile already in 1940-'42 [Gentile 1940, 1942], and—inspired by anyonsanother, dimension-independent, approach allowing for fractional exclusion statistics was suggested by Haldane in 1991 [Haldane 1991].

Ideas of charged particles encircling regions of magnetic flux were developed by Ehrenberg and Siday [Ehrenberg and Siday 1949], Aharonov and Bohm [Aharonov and Bohm 1959], and the more general concept of geometric phases by Pancharatnam [Pancharatnam 1956], Berry [Berry 1984] and Simon [Simon 1983]. Methods of geometric quantization were developed by Souriau around 1967-'70, and he pointed out that if the configuration space is not simply connected then more than two different group characters could arise and this "would lead to prequantizations of a new type" [Souriau 1970]. Around the same time, homotopy classes of Feynman paths brought another topological perspective on quantum statistics [Dowker 1972, Laidlaw and DeWitt 1971, Schulman 1968] (cf., e.g., [Mouchet 2021]). For example, Laidlaw and DeWitt remarked that possibilities in two space dimensions are not limited to bosons and fermions, but they did not elaborate further [Laidlaw and DeWitt 1971].

Foundations for nonabelian anyons were laid much ear-

lier in the theory of parastatistics (nonabelian representations of the permutation group) [Green 1953, Messiah and Greenberg 1964]. Further, the central idea employed by Leinaas and Myrheim that equivalent configurations of indistinguishable particles need to be identified goes back all the way to Gibbs (cf. [Fröhlich 1990, Leinaas and Myrheim 1977]).

The above shows that the essential ideas that were necessary for the discoveries of anyons were floating around at the time, and it is also of interest to remark that, while it took about half a century from the discovery of bosons and fermions to that of anyons, it took another half a century to see their experimental confirmation.

3. Some remarks concerning one-dimensional anyons

Although this article concerns anyons in the 2D setting, it is unavoidable to note the influence of ideas from generalized notions of quantum statistics in 1D (i.e. 1+1 spacetime dimensions). One of the first works in which generalized commutation relations for field operators can be found is Klaiber in 1968 [Klaiber 1968], followed by Streater and Wilde [Streater and Wilde 1970], and Fröhlich [Fröhlich 1976]. Transmutation of 1D hard-core bosons (studied classically by Tonks [Tonks 1936) into fermions and vice versa was considered by Girardeau [Girardeau 1960], who also noted topological and dimensional consequences for the configuration space [Girardeau 1965]. The Lieb–Liniger model [Lieb and Liniger 1963] for point-interacting bosons in 1D constitutes the intermediate statistics found by Leinaas and Myrheim in 1D, however exchange phases (thus "anyonic Lieb-Liniger") may also be added to that model [Kundu 1999]. Leinaas and Myrheim later considered a different approach to statistics in 1D (and in higher dimensions) akin to "Heisenberg quantization" [Leinaas and Myrheim 1993], and found a relationship to the Calogero–Sutherland model [Calogero 1969a,b, Polychronakos 1989, Sutherland 1971]. 2D anyons that are forced into the lowest Landau level of a strong magnetic field may be understood using such 1D concepts [Ouvry 2007].

B. Key points/objectives

We will in this overview focus on providing answers to the following key questions:

- 1. How can the 2D anyon gas be defined physically and modelled mathematically?
- 2. What are the essential properties of the anyon gas? Does it admit some notion of exclusion statistics intermediate to bosons and fermions?
- 3. How can an anyon gas emerge in a realistic physical system consisting of only bosons and/or fermions?



FIG. 2. The circle of abelian anyons, parametrized by the statistics parameter α . Note that any choice of proper anyons (i.e. neither bosons nor fermions) marks an additional point in this circle and thus breaks its orientation symmetry (mirror symmetry / complex conjugation).

The first question is addressed in the first part of Section II for the case of the ideal abelian anyon gas, and in the first part of Section III for the nonideal abelian gas. Section IV introduces the nonabelian gas in brief. The second question is the subject of the later parts of Sections II, III and IV. The third and final question is addressed in Section V. Conclusions and an outlook are given in Section VI.

II. THE IDEAL ANYON GAS

A. Quantum statistics

The possibility for anyons in 2D boils down to the peculiar geometry and topology of the plane as compared to spaces of higher dimensions. Consider for simplicity two particles at positions \mathbf{x}_1 and \mathbf{x}_2 in the Euclidean plane \mathbb{R}^2 , with a wave function Ψ encoding the probability density $|\Psi(\mathbf{x}_1, \mathbf{x}_2)|^2$ of observing the particles at these locations. An exchange of the two positions admits then a change in phase:

$$\Psi(\mathbf{x}_2, \mathbf{x}_1) = e^{\pm i\theta} \Psi(\mathbf{x}_1, \mathbf{x}_2),$$

where $\theta \in \mathbb{R}$. The sign is included above to specify the way in which the exchange is made, and if there is **orien-tation symmetry** then we cannot distinguish between clockwise and counterclockwise exchanges, so that

$$e^{i\theta} = e^{-i\theta} \quad \Leftrightarrow \quad e^{2i\theta} = 1$$

so $\theta/\pi \in \mathbb{Z}$ is either an even integer (bosons) or an odd integer (fermions). However, if we relax the assumption on orientation symmetry and consider a multivalued function for which we keep track of the number and orientation of elementary exchanges (the winding number), then we can allow *any* phase θ (anyons) to appear here. Conventionally, we write $\theta = \alpha \pi$ where α is known as the **statistics parameter** and can be either $\alpha \in \mathbb{R}$ (defined modulo periodicity 2) or in a suitable interval $\alpha \in (-1, 1]$ or $\alpha \in [0, 2)$; cf. Fig. 2.

More generally, we may consider a wave function Ψ for N distinct particles in d-dimensional Euclidean space \mathbb{R}^d and its probability density:

$$\Psi(\mathbf{x})|^2, \qquad \mathbf{x} = (\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N) \in \mathbb{R}^{dN} \setminus \mathbb{A}_N,$$

where, in order to indeed make them distinct we have removed the (fat) **diagonal set** where any two or more particles overlap:

$$\Delta_N := \{ \mathbf{x} \in \mathbb{R}^{dN} : \mathbf{x}_j = \mathbf{x}_k \text{ for some } j \neq k \}.$$

Our reason for considering distinct particles is that we, for simplicity at this stage, fix the particle number to Nand assume that number to be certain and conserved (we shall return to possibilities of collisions on Δ_N later).

In the case that the particles are **identical** and thus **indistinguishable**, then we should reduce the configuration space further and ignore the (now artificial) labels:

$$X = \{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N\} \in \mathcal{C}^N := (\mathbb{R}^{dN} \setminus \Delta_N) / S_N,$$

where the permutation group S_N acts on the labels jof $\mathbf{x} = (\mathbf{x}_j)_{j=1}^N$ while leaving the equivalence class (set) X subject to this action fixed. Indeed, our N-particle configuration space is identical to the manifold

$$\mathcal{C}^N = \{ X = \{ \mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N \} \subset \mathbb{R}^d : |X| = N \}$$

of all N-point subsets of \mathbb{R}^d . If we now consider exchanges of particles then we can no longer act by permutations since it has lost meaning on configurations X of identical particles, however what we can consider instead is the *continuous* exchange of positions in \mathcal{C}^N . Namely, among the continuous paths from a point $X \in \mathcal{C}^N$ to another point $Y \in \mathcal{C}^N$ there are the *loops*, starting and ending at the same point X = Y, which we hereby take to represent continuous physical exchanges of identical particles [Leinaas and Myrheim 1977]. Given such exchange loops σ (continuous functions $[0,1] \to \mathcal{C}^N$ of a parameter, s.t. $\sigma(0) = \sigma(1)$), we may then consider lifting any wave function Ψ which has been assigned a value on a configuration $X \in \mathcal{C}^N$ to a multivalued function² upon which the loops can act. Let us denote the action of a loop σ on Ψ at the point X by $\Psi(\sigma X)$. Then we demand the exchange conditions

$$\Psi(\sigma X) = \rho(\sigma)\Psi(X), \tag{1}$$

where $|\rho(\sigma)| = 1$, so that

$$|\Psi(\sigma X)|^{2} = |\Psi(X)|^{2}$$
(2)

unambiguously defines a probability density at $X \in C^N$. Further, $\rho(1) = 1$ for the trivial (constant) loop $\sigma = 1$, and if we compose two loops σ_1 and σ_2 into a new loop $\sigma_1 \sigma_2$ then we demand the compatibility of the action,

$$\rho(\sigma_1 \sigma_2) = \rho(\sigma_1)\rho(\sigma_2)$$

In other words, ρ is a group homomorphism and thus a **representation** of the group of exchange loops at X with composition.

In comparison, if we consider a single particle, $C^{N=1} = \mathbb{R}^d$, charged and subject to magnetism, then the action of a path should be understood as the (parallel) transport of the charge along the path, and different loops σ may then give rise to different **holonomies** or phases given by the flux enclosed by the loop:

$$\rho(\sigma) = e^{i\Phi_{\sigma}}, \qquad \Phi_{\sigma} = \oint_{\sigma} \mathbf{A} \cdot d\mathbf{r} = \int_{\text{interior of } \sigma} B(\mathbf{x}) \, d\mathbf{x},$$

where curl $\mathbf{A} = B$ is the magnetic field (viewed as a (pseudo)scalar field in d = 2 and appropriately projected in $d \geq 3$). However, for $N \geq 1$ and in the case that there is no external field so that $\rho(\sigma)$ should depend only on the topology of the loop σ , i.e. if we require that $\rho(\sigma) = \rho(\sigma')$ for any two topologically equivalent loops $\sigma \sim \sigma'$ (connected by some continuous deformation), then we consider loops only up to homotopy and the relevant group of (equivalence classes of) loops is the **fundamental group** of the configuration space [Goldin et al. 1983, 1985, Wu 1984a]:

$$\sigma \in \pi_1(\mathcal{C}^N) = \begin{cases} \{1\}, & d = 1, \\ B_N, & d = 2, \\ S_N, & d \ge 3, \end{cases}$$

where B_N denotes the **braid group**, defined below. The group is trivial³ in one dimension because particles then cannot exchange continuously without colliding, i.e. hitting the diagonal set Δ_N .

Now, if Ψ takes values in \mathbb{C} , then $\rho(\sigma)$ is a phase — the **exchange phase** corresponding to the exchange loop σ — and we may identify different **exchange quantum** statistics as symmetry classes of Ψ defined by ρ in (1):

Bosons: $\rho(\sigma) = +1$, the trivial representation. Wave functions Ψ extend symmetrically from \mathcal{C}^N to \mathbb{R}^{dN} ,⁴

$$L^2_{\mathrm{sym}}(\mathbb{R}^{dN}) := \{ \Psi \in L^2 : \Psi(\sigma.\mathbf{x}) = \Psi(\mathbf{x}), \ \sigma \in S_N \}$$

(the usual action of permutations, and with any missing data on Δ_N to be filled in, depending on our choice of Hamiltonian). For example, particles may be independent and identically distributed: $\Psi_0 = \otimes^N \psi_0 \in L^2_{\text{sym}}$,

$$\Psi_0(\mathbf{x}) = (\otimes^N \psi_0)(\mathbf{x}) = \prod_{j=1}^N \psi_0(\mathbf{x}_j)$$

corresponding to Bose–Einstein condensation in the single one-body state $\psi_0 \in L^2(\mathbb{R}^d)$.

 $^{^2}$ More concretely, a section of a bundle with base space $\mathcal{C}^N,$ or a ρ -equivariant function on the covering space of $\mathcal{C}^N;$ see e.g. [Dell'Antonio et al. 1997, Dowker 1985, Fröhlich 1990, Lundholm 2019, Lundholm and Qvarfordt 2020, Mund and Schrader 1995, Myrheim 1999]. The range/fiber is initially $\mathbb{C}.$

 $^{^3}$ Interesting statistics can still emerge as a choice of boundary condition on \triangle_N [Leinaas and Myrheim 1977].

⁴ L² denotes the Hilbert space of square-integrable functions with inner product $\langle \Psi, \Phi \rangle = \int \overline{\Psi(\mathbf{x})} \Phi(\mathbf{x}) d\mathbf{x}$.

Fermions: $\rho(\sigma) = \operatorname{sign}(\sigma) \in \{-1, +1\}$, the signed representation on S_N (signature of permutation). In this case wave functions Ψ extend *antisymmetrically from* \mathcal{C}^N to \mathbb{R}^{dN} ,

$$L^{2}_{\text{asym}}(\mathbb{R}^{dN}) := \{ \Psi \in L^{2} : \Psi(\sigma.\mathbf{x}) = \text{sign}(\sigma)\Psi(\mathbf{x}), \, \sigma \in S_{N} \}$$

(in this case it is natural to require that Ψ vanishes on Δ_N). We apparently obtain determinantal correlations and the **Pauli principle**, since functions in L^2_{asym} are spanned by **Slater determinants**, such as $\Psi_0 = \psi_0 \wedge \psi_1 \wedge \ldots \wedge \psi_{N-1}$, where

$$\bigwedge_{k=0}^{N-1} \psi_k(\mathbf{x}_1,\ldots,\mathbf{x}_N) := \det \begin{bmatrix} \psi_0(\mathbf{x}_1) & \cdots & \psi_0(\mathbf{x}_N) \\ \vdots & & \vdots \\ \psi_{N-1}(\mathbf{x}_1) & \cdots & \psi_{N-1}(\mathbf{x}_N) \end{bmatrix}.$$

Anyons: in d = 2 the most general ρ is a unitary representation of B_N . Proper anyons then correspond to those ρ which do not factor through S_N , i.e. for which orientation symmetry is broken. Let us denote by L_{ρ}^2 the Hilbert space of square-integrable multivalued wave functions Ψ based on the configuration space C^N and subject to the exchange conditions (1). A central question is, whether we can anticipate **intermediate or fractional exclu**sion statistics in L_{ρ}^2 ? Actually, it turns out that the information at hand is not sufficient to decide this, and even the above conclusions for bosons and fermions can be misleading, as we will see below.

B. The braid group

For d = 2, the group of continuous loops in \mathcal{C}^N modulo homotopy is conveniently visualized as a group of Nparticle world lines that evolve in 2+1-dimensional spacetime, subject to composition via some arbitrary but fixed reference configuration $X_0 \in \mathcal{C}^N$. The resulting group, denoted B_N , is the **braid group on** N **strands** [Artin 1947, 1925, Birman 1974]:

$$B_N = \left\langle \sigma_1, \dots, \sigma_{N-1} : \sigma_j \sigma_{j+1} \sigma_j = \sigma_{j+1} \sigma_j \sigma_{j+1}, \\ \sigma_j \sigma_k = \sigma_k \sigma_j \right\rangle_{\substack{j,k=1\dots N-1\\k\neq j\pm 1}}^{j,k=1\dots N-1}$$

That is, the group generated by N-1 elementary braids

$$\sigma_j: \quad \left| \begin{array}{c} | \\ 1 \end{array} \right| \quad \left| \begin{array}{c} \searrow \\ j \end{array} \right| \quad \left| \\ j \end{array} \right| \qquad \sigma_j^{-1}: \quad \left| \begin{array}{c} | \\ j \end{array} \right| \quad \left| \begin{array}{c} \bigotimes \\ j \end{array} \right| \quad \left| \\ j \end{array} \right|$$

which are composed by stacking one on top of another. These elementary braids are subject to two sets of topological relations; one of which are known as the **Yang-Baxter relations**, and the other implement the commutativity of independent braids. In B_4 we have, e.g.,





FIG. 3. Exchange loops for a single resp. a pair of particles with p other particles enclosed, and corresponding braid diagrams and phases. From [Lundholm and Solovej 2013a].

If we add a third set of relations $\sigma_j^2 = 1 \forall j$, or equivalently, $\sigma_j = \sigma_j^{-1}$, i.e. clockwise and counterclockwise exchanges cannot be distinguished, then we obtain the **permutation group** S_N . This will be the case for $d \geq 3$ since an elementary exchange may be rotated out of the plane (continuously and homotopically). From now on we shall stick to the anyonic case d = 2.

If one looks for irreducible abelian unitary representations, i.e. **exchange phases**

$$\rho \colon B_N \to \mathrm{U}(1) = \{ z \in \mathbb{C} : |z| = 1 \},\$$

then the Yang-Baxter relations single out a unique phase:

$$\rho(\sigma_j) = e^{i\alpha\pi}, \quad \text{for all } j = 1, 2, \dots, N-1; \quad (3)$$

cf. Figs. 2-3. The only representations that factor through S_N require either $\alpha = 0$ ($\alpha \in 2\mathbb{Z}$) or $\alpha = 1$ ($\alpha \in 2\mathbb{Z} + 1$), thus corresponding to bosons or fermions. We will return to the nonabelian case, and representations with higher rank than 1, in Section IV.

C. Kinetic energy

Above we considered wave functions $\Psi \in L^2_{\rho}$ that implement an exchange symmetry (1) by choice of a representation ρ of B_N . However, it turns out that this is only one part of the data that specifies an anyon model. The other, equally essential, part is the choice of a Hamiltonian or kinetic energy operator. Namely, in order to be able to consider continuous exchanges of particles we also need to resolve space and time at the appropriate energy scale.

Our starting point is the nonrelativistic⁵ free kinetic energy for N particles with mass m:

$$T = \frac{1}{2m} \sum_{j=1}^{N} \mathbf{p}_j^2, \qquad (4)$$

⁵ We remain nonrelativistic throughout this overview and likewise we will ignore spin; see however [Fröhlich 2009, Jackiw 1990, Mund 2009].

where $\mathbf{p}_j \in \mathbb{R}^2$ is the momentum of the particle with position $\mathbf{x}_j \in \mathbb{R}^2$. If the particles are all distinguishable then the canonical quantization of this expression is obtained with momentum operators $\hat{\mathbf{p}}_j = -i\hbar\nabla_{\mathbf{x}_j}$ canonically conjugate to the position operators $\hat{\mathbf{x}}_{ij}$, i.e.

$$\hat{T}_{\text{dist}} := \frac{\hbar^2}{2m} \sum_{j=1}^N (-i\nabla_{\mathbf{x}_j})^2, \qquad (5)$$

acting on (suitably regular) $\Psi \in L^2(\mathbb{R}^{2N})$.

In our case of identical anyons we may consider the particles to be *locally* distinguishable. Namely, at any fixed point $X \in \mathcal{C}^N$ we may attach arbitrary labels to them:

$$\mathcal{C}^N \ni X = \{\mathbf{x}_1, \dots, \mathbf{x}_N\} \iff (\mathbf{x}_1, \dots, \mathbf{x}_N) \in \mathbb{R}^{2N} \setminus \Delta_N,$$

and we can keep track of these labels along any paths from X in \mathcal{C}^N as long as they cannot exchange nontrivially, i.e. as long as any two possible paths do not compose to a topologically nontrivial loop. So in order to define free anyons, the idea is then that we should keep the kinetic energy operator as above whenever we consider wave functions Ψ supported on topologically trivial subsets of \mathcal{C}^N (i.e. not encircling any of the holes in the manifold), which then can anyway be treated as subsets of the configuration space $\mathbb{R}^{2N} \setminus \Delta_N$ of distinguishable and distinct particles. Additional information must then be added to \hat{T}_{dist} whenever we pass between different such subsets of \mathcal{C}^N so as to complete topologically nontrivial loops $\sigma \in \pi_1(\mathcal{C}^N) = B_N, \ \sigma \neq 1$. One way to implement this concretely is to treat the exchange condition (1) as a topological boundary condition⁶ that Ψ must satisfy in order to be in the domain of the operator, which we denote by

$$\hat{T}_{\rho} := \frac{\hbar^2}{2m} \sum_{j=1}^{N} (-i\nabla_{\mathbf{x}_j}^{(\rho)})^2, \qquad (6)$$

or simply by \hat{T}_{α} if $\rho(\sigma_j) = e^{i\alpha\pi}$. For convenience we also denote the Hilbert space L^2_{ρ} by L^2_{α} in this case. Again, we stress that \hat{T}_{α} acts *locally* as the standard Laplacian $\hat{T}_{\text{dist}} = \frac{\hbar^2}{2m} \sum_{j=1}^{N} (-\Delta_{\mathbf{x}_j})$ on (suitably regular) $\Psi \in L^2_{\alpha}$, thus *independently* of α , but *globally* we have also the exchange conditions to account for, such as **pair exchange**

$$\Psi(\mathbf{x}_1, \dots, \mathbf{x}_k, \dots, \mathbf{x}_j, \dots, \mathbf{x}_N) = e^{i(2p+1)\alpha\pi} \Psi(\mathbf{x}_1, \dots, \mathbf{x}_j, \dots, \mathbf{x}_k, \dots, \mathbf{x}_N), \quad (7)$$

where p is the number of other particles enclosed in the exchange loop as \mathbf{x}_j and \mathbf{x}_k are exchanged once in a counterclockwise manner (as well as other conditions; cf. Fig. 3).

As an example, consider the two-particle case N = 2 in center-of-mass and relative coordinates $\mathbf{X} = \frac{1}{2}(\mathbf{x}_1 + \mathbf{x}_2)$, $\mathbf{r} = \mathbf{x}_1 - \mathbf{x}_2$. Then the kinetic energy separates,

$$\hat{T}_{\alpha} = \frac{\hbar^2}{4m} (-\Delta_{\mathbf{X}}) + \frac{\hbar^2}{m} (-\Delta_{\mathbf{r}}), \qquad (8)$$

and further, expressed in polar coordinates $(r = |\mathbf{r}|, \varphi)$:

$$-\Delta_{\mathbf{r}} = -\partial_r^2 - \frac{1}{r}\partial_r - \frac{1}{r^2}\partial_{\varphi}^2.$$
 (9)

These operators act on functions $\Psi(\mathbf{X}, r, \varphi)$ (here with its proper multivaluedness on \mathcal{C}^2 incorporated in the winding of $\varphi \in \mathbb{R}$ over multiples of π) such that

$$\Psi(\mathbf{X}, r, \varphi + \pi) = e^{i\alpha\pi}\Psi(\mathbf{X}, r, \varphi), \quad r > 0,$$
(10)

corresponding to the exchange condition (7) (all braids in B_2 are generated by this simple exchange σ_1). The manifold C^2 may be parametrized by the set in \mathbb{R}^4 given by the half-space

$$\Omega = \{ \mathbf{X} \in \mathbb{R}^2, \ r > 0, \ 0 \le \varphi < \pi \},\$$

and its boundary

$$\partial \Omega = \{ \mathbf{X} \in \mathbb{R}^2, \ r = 0 \text{ or } \varphi = 0 \text{ or } \varphi = \pi \}$$

relates the two points $(r > 0, \varphi = 0)$ and $(r > 0, \varphi = \pi)$ topologically to the same point in C^2 . An equivalent description is therefore (8)-(9) defined on $L^2(\Omega)$ and subject to the boundary condition

$$\Psi(\mathbf{X}, r, \varphi = \pi) = e^{i\alpha\pi}\Psi(\mathbf{X}, r, \varphi = 0), \quad r > 0.$$

The operator \hat{T}_{ρ} as defined above is formally hermitian but not self-adjoint (and not even necessarily *essentially* self-adjoint) since we have still not supplied any information at the diagonal set Δ_N , which cuts out holes of codimension 2 in the manifold \mathcal{C}^N (for N = 2 this set is $\Delta_2 = \{\mathbf{X} \in \mathbb{R}^2, r = 0\}$). There is however a canonical choice of a self-adjoint realization (known in the mathematical literature as the **Friedrichs extension**), obtained by considering the expectation value of the kinetic energy in its formal quadratic form expression⁷

$$\langle \Psi, \hat{T}_{\rho} \Psi \rangle = \frac{\hbar^2}{2m} \int_{\mathbb{R}^{2N}} \sum_{j=1}^N |\nabla_{\mathbf{x}_j}^{(\rho)} \Psi|^2 d\mathbf{x} \ge 0$$
(11)

initially on (multivalued) wave functions Ψ that are smooth and compactly supported away from \mathbb{A}_N and

⁶ Technically, Ψ is defined as a complex-valued function on the covering space of \mathcal{C}^N subject to the ρ -equivariance condition (1), which is equivalent to regarding Ψ as a section of a complex line bundle over \mathcal{C}^N with its geometry specified by $\rho: B_N \to U(1)$ (locally flat, but globally curved by the holonomies specified by $\rho(\sigma)$ for $\sigma \in B_N$); see, e.g., [Dowker 1985, Lundholm and Qvarfordt 2020, Mund and Schrader 1995].

⁷ This is more appropriately defined on C^N and observed to extend unambiguously to the covering space [Lundholm and Qvarfordt 2020].

subject to the exchange conditions (7) resp. (1).⁸ For N = 2 the energy form (11) is

$$\langle \Psi, \hat{T}_{\alpha} \Psi \rangle = \frac{\hbar^2}{m} \int_{\mathbf{X} \in \mathbb{R}^2} \int_{r=0}^{\infty} \int_{\varphi=0}^{2\pi} \left(\frac{1}{4} |\nabla_{\mathbf{X}} \Psi|^2 + |\partial_r \Psi|^2 + \frac{1}{r^2} |\partial_{\varphi} \Psi|^2 \right) d\varphi \, r dr \, d\mathbf{X}, \quad (12)$$

again subject to the exchange condition (10). In general then, this specific choice of a self-adjoint operator \hat{T}_{ρ} resp. \hat{T}_{α} is what we take to mean precisely by **free** (noninteracting, as compared to point-interacting; see below) **ideal anyons**, by analogy with bosons and fermions.

D. Statistics transmutation in 2D

So far we have worked exclusively in the geometric perspective of anyons, but will now be able to make the connection to the magnetic perspective.

Since we have fixed our attention on the plane \mathbb{R}^2 , it is occasionally convenient to switch to complex notation:

$$\mathbb{R}^{2N} \ni \mathbf{x} = (\mathbf{x}_1, \dots, \mathbf{x}_N) \iff \mathbf{z} = (z_1, z_2, \dots, z_N) \in \mathbb{C}^N,$$

where the real (imaginary) part in \mathbb{C} is canonically identified with the first (second) coordinate in \mathbb{R}^2 .

Now, consider either a bosonic or a fermionic wave function $\Psi \in L^2_{\text{sym/asym}}(\mathbb{C}^N)$ restricted to the subset of distinct coordinates $z \in \mathbb{C}^N \setminus \Delta_N$ and make there the gauge transformation $\Psi = U\tilde{\Psi}$, where

$$U(z) := \prod_{j < k} \frac{z_j - z_k}{|z_j - z_k|} = \exp\left(i\sum_{j < k} \arg(z_j - z_k)\right), \quad (13)$$

the product of all relative phases of pairs of particles. One may verify that this expression is well defined and single valued on $\mathbb{C}^N \setminus \Delta_N$, unitary $|U(\mathbf{z})| = 1$, and antisymmetric w.r.t. permutations of labels in $\mathbf{z} = (z_j)_{j=1}^N$, so that if Ψ is symmetric then $\tilde{\Psi} = U^{-1}\Psi$ is antisymmetric, and vice versa. Thus, modulo the smaller set of diagonals Δ_N where it is not defined, the unitary transformation by U achieves a **statistics transmutation** $L^2_{\text{sym}} \leftrightarrow L^2_{\text{asym}}$ between bosons and fermions in 2D. This perhaps seems confusing, however, recall that the kinetic energy is equally important in defining the exchange statistics. Indeed, if we consider the momentum then this transmutation comes at the cost of a **gauge potential**:

$$-i\nabla_{\mathbf{x}_j}\Psi = U(-i\nabla_{\mathbf{x}_j} + \mathbf{A}_j)\Psi,\tag{14}$$

where we have defined the vector potentials

$$\mathbf{A}_{j}(\mathbf{x}) := -iU^{-1}\nabla_{\mathbf{x}_{j}}U = \sum_{k \neq j} \frac{(\mathbf{x}_{j} - \mathbf{x}_{k})^{\perp}}{|\mathbf{x}_{j} - \mathbf{x}_{k}|^{2}}, \qquad (15)$$

for j = 1, ..., N, and we denoted a $\pi/2$ rotation in \mathbb{R}^2

$$\mathbf{x} = (x, y) \quad \mapsto \quad \mathbf{x}^{\perp} = (-y, x)$$

(or $z \mapsto iz$). In other words, if we had initially

$$\hat{T}_{\rm asym} = \frac{\hbar^2}{2m} \sum_{j=1}^{N} (-i\nabla_{\mathbf{x}_j})^2 \quad \text{acting on } L^2_{\rm asym}$$

then we now obtain

$$\hat{T}_{\text{sym}\to\text{asym}} := \frac{\hbar^2}{2m} \sum_{j=1}^{N} \left(-i\nabla_{\mathbf{x}_j} + \mathbf{A}_j \right)^2 \quad \text{acting on } L^2_{\text{sym}},$$

so that indeed fermions may be represented as bosons, although with an additional magnetic interaction. The magnetic field seen by particle j,

$$\operatorname{curl}_{\mathbf{x}_j} \mathbf{A}_j = 2\pi \sum_{k \neq j} \delta(\mathbf{x}_j - \mathbf{x}_k), \qquad (16)$$

corresponds to the attachment of **Aharonov–Bohm point fluxes** of magnitude 2π to each of the other particles. The set where this field is singular is precisely Δ_N .

Similarly, if we fix a statistics parameter $\alpha \in \mathbb{R}$ and consider $\Psi = U^{\alpha} \tilde{\Psi}$ (now typically a singular gauge transformation involving multivalued functions over $\mathbb{C}^N \setminus \Delta_N$) then

$$-i\nabla_{\mathbf{x}_j}\Psi = U^{\alpha}(-i\nabla_{\mathbf{x}_j} + \alpha \mathbf{A}_j)\Psi, \qquad (17)$$

and thus if we originally had the anyonic kinetic energy

$$\hat{T}_{\alpha} = \frac{\hbar^2}{2m} \sum_{j=1}^{N} (-i\nabla_{\mathbf{x}_j})^2 \quad \text{acting on } \Psi \in L^2_{\alpha}$$

then we obtain an equivalent description in terms of bosons:

$$\hat{T}_{\mathrm{sym}\to\alpha} := \frac{\hbar^2}{2m} \sum_{j=1}^N \left(-i\nabla_{\mathbf{x}_j} + \alpha \mathbf{A}_j \right)^2, \text{ acting on } \tilde{\Psi} \in L^2_{\mathrm{sym}}$$

 $^{^{8}}$ There is a natural extension (closure) of the energy form (11) to the largest subspace of the Hilbert space L^2_{ρ} upon which the energy is defined and finite, which is also the largest domain of the momentum operator $-i\nabla^{(\rho)}$. Denoting its adjoint $(-i\nabla^{(\rho)})^*$, one then defines $\hat{T}_{\rho} := \frac{\hbar^2}{2m} (-i\nabla^{(\rho)})^* (-i\nabla^{(\rho)})$, canonically self-adjoint on a corresponding smaller subspace of L^2_{ρ} . One might object that this approach seems to assume a hard-core condition on the anyons since functions are initially taken to vanish close to Δ_N , however for bosons and fermions this eventually yields exactly the standard free kinetic energy $\hat{T}_{\rm asym/sym}$ (with the same expression as \hat{T}_{dist}) on the (Sobolev) spaces of anti/symmetric square-integrable functions with both one and two partial derivatives, that are also square-integrable. It is in fact a peculiarity of dimensions $d \ge 2$ that any extra conditions at the diagonals disappear in this procedure, so that our assumption on distinct particles is justified a posteriori; cf [Bourdeau and Sorkin 1992, Lundholm and Solovej 2014]. This can also be understood by means of a more precise version of the Hardy inequality (34) discussed below [Lundholm and Qvarfordt 2020].

If we instead consider the transformation $\Psi = U^{\alpha-1}\tilde{\Psi}$ where $\tilde{\Psi} \in L^2_{asym}$, then we arrive at a description in terms of fermions:

$$\hat{T}_{\mathrm{asym} \to \alpha} := \frac{\hbar^2}{2m} \sum_{j=1}^N \left(-i \nabla_{\mathbf{x}_j} + (\alpha - 1) \mathbf{A}_j \right)^2,$$

acting on L^2_{asym} . The magnetic field of $\alpha \mathbf{A}_j$ corresponds to the attachment of Aharonov–Bohm fluxes of magnitude $2\pi\alpha$ to each of the other particles. In this way we have the freedom to treat anyons as either bosons or fermions with topological magnetic interactions. This tool can certainly be beneficial as we can trade a simple operator (the free kinetic energy) but on a complicated geometry (multivalued functions with appropriate exchange conditions), for a complicated operator (magnetic interactions) but on a simple geometry (symmetric or antisymmetric functions on \mathbb{R}^{2N}). The former choice is usually referred to as the **anyon gauge** while the latter is the **magnetic gauge**, and this connection also brings us from an idealized world of quantum statistics of point particles and closer to the various possibilities for emergent statistics transmutation phenomena.

The procedure we referred to above to ensure the selfadjointness of the kinetic energy operator by extending the energy form (Friedrichs extension) also applies in this magnetic gauge formulation of the problem, so that indeed the operators $\hat{T}_{\text{sym/asym}} \rightarrow \alpha$ are self-adjoint on appropriate domains in $L^2_{\text{sym/asym}}$ where the energy is finite [Lundholm and Solovej 2014]. Further, we remark that multiples of the unitary multiplication operator

$$U^2 \colon L^2_{\text{sym/asym}/\alpha} \to L^2_{\text{sym/asym}/(\alpha+2)}$$

may be used, together with complex conjugation symmetry, to obtain that the free ideal anyonic kinetic energies are unitarily equivalent w.r.t. the transformations

$$\alpha \mapsto \alpha + 2n, \ n \in \mathbb{Z} \quad \text{and} \quad \alpha \mapsto -\alpha.$$
 (18)

E. Exchange vs. exclusion

Now that we have managed to define more or less concrete but still hypothetical models for ideal abelian anyons, we come to the next question: How do such anyons actually behave? If we are ever to observe them in nature then we need to know their behavior. So, are they more like bosons or more like fermions, or something in between, like the terms "intermediate statistics" or "fractional statistics" suggest? In other words, can we in any way relate the intermediate (braid) **exchange statistics** encoded into anyons to some intermediate **exclusion statistics** concerning the occupation numbers of various states as seen from a one-body perspective. Of course, since the magnetic gauge picture clarifies that anyons are (possibly) at least as complicated as strongly interacting systems of bosons or fermions, it is a priori



FIG. 4. Energy spectra for N = 2 (upper) and N = 3 (lower) anyons in a harmonic trap (minus the center of mass energy 1, and in suitable units). From [Yakaboylu et al. 2020]; cf. also [Leinaas and Myrheim 1977, Murthy et al. 1991, Sporre et al. 1991].

not at all obvious whether this is possible. The issue of exchange vs. exclusion was referred to as " α to β " by Canright and Johnson in their review [Canright and Johnson 1994] on the matter.

Only in cases of very few particles, N = 2, 3, 4, are energy spectra for ideal anyons well understood, because they have then been computed analytically (for N = 2[Arovas et al. 1985, Leinaas and Myrheim 1977, Wilczek 1982b], also extending to a collective [Date and Murthy 1993, Date et al. 2003] part of the spectrum for all N[Chou 1991, Wu 1984b]) or numerically (the remaining part of the spectrum [Murthy et al. 1991, Sporre et al. 1991, 1992]), in a few situations.

In order to lift the degeneracy of the spectrum and facilitate comparison, it is convenient to confine the particles in a harmonic trapping potential $V(\mathbf{x}) = \frac{1}{2}m\omega^2 |\mathbf{x}|^2$. For N anyons in the magnetic gauge description w.r.t. bosons we thus consider the Hamiltonian operator

$$\hat{H}_N = \hat{T}_{\text{sym}\to\alpha} + \hat{V} = \sum_{j=1}^N \left[\frac{\hbar^2}{2m} \left(-i\nabla_{\mathbf{x}_j} + \alpha \mathbf{A}_j \right)^2 + V(\mathbf{x}_j) \right],$$



FIG. 5. Schematic energy spectrum for $N \gg 1$ (left) and estimates for the g.s.e. near bosons and fermions (right) for anyons with statistics $\alpha = \theta/\pi$ in a harmonic trap. From [Chitra and Sen 1992].

and its corresponding ground-state energy (g.s.e.)

$$E_N := \inf \operatorname{spec} \hat{H}_N = \inf_{0 \neq \Psi \in L^2_{\operatorname{sym}}} \langle \hat{H}_N \rangle_{\Psi}, \qquad (19)$$

where $\langle \hat{A} \rangle_{\Psi} := \langle \Psi, \hat{A}\Psi \rangle / \langle \Psi, \Psi \rangle$ denotes the expectation of any operator \hat{A} in a state $\Psi \neq 0$. The spectrum of the 2D harmonic oscillator $\hat{H}_{N=1}$ is well known:

spec
$$\hat{H}_1 = \{\hbar \omega n\}_{n=1,2,3,\dots}$$
 (each with multiplicity n),

and for bosons one thus obtains N times the lowest eigenvalue, $E_N(\alpha = 0) = \hbar \omega N$, while for (spinless) fermions, due to the Pauli principle, we must sum over the N first eigenvalues according to their multiplicity: $E_N(\alpha = 1) \sim \frac{\sqrt{8}}{3} \hbar \omega N^{3/2}$ as $N \to \infty$. Further, because of rotation symmetry, the Hamiltonian commutes with the total angular momentum operator

$$\hat{L}/\hbar = -i\sum_{j=1}^{N} \left[x_j \partial_{y_j} - y_j \partial_{x_j} \right] = \sum_{j=1}^{N} \left[z_j \frac{\partial}{\partial z_j} - \bar{z}_j \frac{\partial}{\partial \bar{z}_j} \right],$$

and its eigenvalues $L \in \mathbb{Z}$ can therefore be used to separate the energy spectrum further.

The spectrum of H_N for N = 2 and N = 3 is shown in Fig. 4, and while in the former case it manifests a strictly linear interpolation between bosons and fermions, [Leinaas and Myrheim 1977]

$$E_2(\alpha) = \hbar\omega \left(2 + \min_{q \in \mathbb{Z}} |2q + \alpha|\right), \qquad (20)$$

in the latter case there are also nonlinearly interpolating states at various L, and such a state with $L \neq 0$ comes down from an excited level at $\alpha = 0$ to reach the Fermi energy at $\alpha = 1$. Similar features are also seen in the numerical spectrum for N = 4 [Sporre et al. 1992] and expected to occur for all $N \geq 3$ [Chitra and Sen 1992].

9

tained simply by considering trial states which localize the particles on separate domains in \mathbb{R}^2 , so that their exchange statistics is not seen (in the magnetic gauge picture one may then gauge away all magnetic interactions, while in the anyon gauge picture one then stays within a domain of distinguishability). By balancing the localization energy and the spread into the potential one then finds that $\widetilde{E}_N(\alpha) \lesssim \hbar \omega N^{3/2}$, i.e. the energy is at most at the order of fermions. Note however that the fermionic energy does not provide a strict upper bound, as seen for N = 3 where $E_3(\alpha \approx 0.7) > E_3(\alpha = 1)$.

On the other hand, it is also useful to note that the bosonic energy, for which the g.s. Ψ_0 is a Gaussian at the origin x = 0, always gives a rigorous lower bound:

$$E_N(\alpha) \ge E_N(0) = \hbar \omega N, \tag{21}$$

since the kinetic energy satisfies a diamagnetic inequality [Lundholm and Solovej 2014]

$$\langle \hat{T}_{\alpha} \rangle_{\Psi} \ge \langle \hat{T}_{\text{sym}} \rangle_{|\Psi|}.$$
 (22)

This is observed by factoring Ψ into its modulus and phase, and dropping the phase contribution to the momentum $-i\nabla\Psi$.

Chitra and Sen [Chitra and Sen 1992] provided a more interesting lower bound to the spectrum for any N, extending (20): for an arbitrary state Ψ with angular momentum L, it holds

$$\langle \hat{H}_N \rangle_{\Psi} \geq \hbar \omega \left(N + \left| L + \alpha \frac{N(N-1)}{2} \right| \right).$$
 (23)

Combined with our a priori upper bound, this implies that in the ground state the angular momentum needs to be close to

$$L \approx -\alpha \binom{N}{2}$$
, since $\left| L + \alpha \binom{N}{2} \right| \lesssim O(N^{3/2})$, (24)

so that on the average there needs to be an angular momentum $-\alpha$ for every pair of particles in order to cancel the shifted inherent/anyonic angular momentum (the number of attached units of magnetic flux per particle). This also means that as N grows large there will be a large number $O(N^{1/2})$ of level crossings in the g.s. between different integer angular momenta, because any fixed angular momentum L can only be good on a scale $\Delta \alpha = O(N^{-1/2})$. The situation is sketched in Fig. 5, where at fixed $N \gg 1$ one expects to find curves corresponding to families of states at fixed L, that are smoothly varying with α , to eventually come down to an energy $O(N^{3/2})$ on a $O(N^{1/2})$ -neighborhood around a suitable minimum $\alpha \approx \alpha_0(L)$, and then to again shoot upwards in energy with a slope $\gtrsim O(N^2)$.

Chitra and Sen went on to estimate (again, see Fig. 5) the enveloping g.s.e. curve close to bosons (suggesting $E_N/(\hbar\omega N^{3/2}) \approx \frac{7\sqrt{3}}{9}\sqrt{\alpha}$ and to fermions (suggesting $E_N/(\hbar\omega N^{3/2}) \approx \frac{\sqrt{8}}{3}$, but left it as an open problem whether there are nonanalyticities in the limiting function

$$h(\alpha) := \liminf_{N \to \infty} E_N(\alpha) / (\hbar \omega N^{3/2}), \qquad (25)$$

concluding that "This is an interesting but difficult question to answer."

F. Towards density functionals for anyons

The few-particle spectra led to some initial results on the thermodynamics of the anyon gas, by considering the high-temperature, low-density virial expansions [Arovas et al. 1985, Dowker 1985] (see [Khare 2005, Mancarella et al. 2013a, Myrheim 1999] for review). In the lowtemperature regime, in order to make the many-anyon ground state problem more manageable, we could wish for a limiting effective description with a relatively limited number of degrees of freedom that are sufficient to describe the collective state of the system at $N \to \infty$. Such a state might converge towards the minimizer of a suitable functional, i.e. (taking some suitable scale a > 0for normalization):

$$E_N/N^a \xrightarrow{N \to \infty}$$
 minimum of $\mathcal{E}[\psi]$ or $\mathcal{E}[\varrho]$, and

some marginal of g.s. Ψ_0 of $\hat{H}_N \xrightarrow{N \to \infty}$ minimizer of \mathcal{E} ,

where \mathcal{E} is typically an energy functional of either onebody states $\psi \in L^2(\mathbb{R}^2)$ or of corresponding probability densities $\varrho = |\psi|^2 \in L^1(\mathbb{R}^2)$ (with mass 1 or N).

For example, for 2D fermions in a trapping potential V, one has the **Thomas–Fermi approximation** [Fermi 1927, Thomas 1927]:

$$E_N = \inf_{0 \neq \Psi \in L^2_{\text{asym}}} \langle \hat{T}_{\text{asym}} + \hat{V} \rangle_{\Psi} \approx \inf_{\varrho \ge 0: \int_{\mathbb{R}^2} \varrho = N} \mathcal{E}^{\text{TF}}[\varrho],$$
(26)

where the density $\rho = \rho^{\text{TF}} \approx \rho_{\Psi_0}$ (one-body marginal of g.s.) minimizes the **Thomas–Fermi (TF) functional**

$$\mathcal{E}^{\mathrm{TF}}[\varrho] := \int_{\mathbb{R}^2} \left[2\pi \varrho(\mathbf{x})^2 + V(\mathbf{x})\varrho(\mathbf{x}) \right] d\mathbf{x}.$$
(27)

(Note: To simplify our discussion, we shall now and in the remainder of this article assume that the physical constant $\hbar^2/(2m) = 1$, i.e. $T = \mathbf{p}^2$, by an appropriate choice of units $\hbar = 2m = c = e = 1$.) The kinetic energy density is here given by the energy per unit area of the **homogeneous 2D Fermi gas** at density $\varrho(\mathbf{x})$, obtained e.g. by confining N fermions in a box of area $L \times L$, summing the one-body eigenvalues, and taking the thermodynamic limit:

$$E_N/L^2 \xrightarrow{N,L \to \infty} 2\pi \bar{\varrho}^2$$
, at fixed $\bar{\varrho} = N/L^2$

On the other hand, for bosons with sufficiently weak *scalar* interactions one has the **Gross–Pitaevskii (GP)**

functional [Gross 1961, Pitaevskii 1961]:

$$\mathcal{E}^{\rm GP}[\psi] := \int_{\mathbb{R}^2} \left[\left| (-i\nabla + \mathbf{A}_{\rm ext})\psi \right|^2 + V|\psi|^2 + g|\psi|^4 \right], \quad (28)$$

describing approximate condensation into the state $\psi \in L^2(\mathbb{R}^2)$ as $E_N/N \to \text{minimum of } \mathcal{E}^{\text{GP}}$. In regimes where the interaction strength $g \in \mathbb{R}$ (related to the scattering length of the interaction potential) dominates over the kinetic term one may neglect that term and keep only the last two terms, again leaving a functional of $\varrho = |\psi|^2$ of the Thomas–Fermi type with $g \leftrightarrow 2\pi N$ (this is only a formal analogy at the level of functionals, since the two mechanisms of the approximations are very different).

In view of the above, could one then suggest the following interpolating functional for $0 < \alpha \le 1$, describing an "average-field" approximation for anyons

$$\mathcal{E}^{\mathrm{af}}[\varrho] \approx \int_{\mathbb{R}^2} \left[2\pi \alpha \varrho(\mathbf{x})^2 + V(\mathbf{x})\varrho(\mathbf{x}) \right] d\mathbf{x} \quad ? \tag{29}$$

Indeed, in a mean-field⁹ (**Hartree**) ansatz for the ground state in the bosonic representation, consider:

$$\Psi(\mathbf{x}) = \psi(\mathbf{x}_1)\psi(\mathbf{x}_2)\dots\psi(\mathbf{x}_N), \qquad (30)$$

and we may think of a single particle $\psi \in L^2(\mathbb{R}^2)$ in the magnetic field of the others (identically distributed):

$$B(\mathbf{x}) \approx 2\pi\alpha (N-1)|\psi(\mathbf{x})|^2 \approx 2\pi\alpha \varrho_{\Psi}(\mathbf{x}).$$
(31)

If the density $\rho_{\Psi} = N |\psi|^2$ is approximately constant on an area $L \times L$ where N bosons condense into the lowest Landau level of the field B (cf. (64) below), then one indeed obtains an energy per unit area

$$E_N/L^2 \approx BN/L^2 \approx 2\pi\alpha\bar{\varrho}^2, \qquad \bar{\varrho} = N/L^2, \qquad (32)$$

thus suggesting (29). For the ideal anyon gas this argument is not rigorous, because one cannot apply such an ansatz in the singular magnetic field (it is not in the appropriate domain of \hat{T}_{α}). This difficulty can be overcome using regularization however, and we return to it in Section III B in conjunction with the extended anyon gas. Thus, after accepting some refinements to the "average-field" approach in that context, it has been successfully applied in the limits $\alpha \to 0$ and $\alpha \to 1$ as $N \to \infty$.

To be precise, if we wish to refer to the approximation as above with an almost constant field (on the scale of a trap) then it is perhaps more appropriate to call it a "constant-field" approximation. We also note that if (29) would be a valid description for proper anyons then it suggests that for the harmonic oscillator problem

$$E_N \stackrel{?}{\approx} \inf_{\substack{\varrho \ge 0\\ \int \varrho = N}} \int_{\mathbb{R}^2} \left[\frac{\hbar^2 \pi \alpha}{m} \varrho(\mathbf{x})^2 + \frac{m\omega^2}{2} |\mathbf{x}|^2 \varrho(\mathbf{x}) \right] d\mathbf{x}$$
$$= \frac{\sqrt{8}}{3} \sqrt{\alpha} \, \hbar \omega N^{3/2}, \qquad 0 < \alpha \le 1.$$
(33)

⁹ We differentiate between "average-field" and "mean-field", partly to distinguish the anyonic context, but also because of more technical reasons that will be clarified below.

Early applications of variants of **density functional theory** (DFT) to the anyon context were made by Chitra and Sen, who proposed the refinement as in Fig. 5 due to the bosons' hard core [Chitra and Sen 1992], and by Li, Bhaduri and Murthy who considered an extension of TF theory to excited states in the spectrum [Li et al. 1992]. Again, only some aspects have been rigorously justified, in the limits close to bosons and fermions.

G. Local exclusion principle and degeneracy pressure for the ideal anyon gas

An approach that has been successful for an increased qualitative understanding of the genuine ideal anyon gas at arbitrary α is to take a strictly *local* route to exclusion via the wider concept of **statistical repulsion**. Namely, one may note that statistical repulsion manifests itself in the anyon gas in (at least) three ways:

1. Effective scalar pairwise repulsion. The exchange conditions (7) for each pair of particles transpire to yield a pair repulsion between particles. Its simplest form can be stated as a lower bound (known in mathematics as a many-particle Hardy inequality [Hoffmann-Ostenhof et al. 2008, Lundholm and Solovej 2013a]) for the kinetic energy: for any N,

$$\hat{T}_{\alpha} \geq \frac{4\alpha_N^2}{N} \sum_{j < k} \frac{1}{|\mathbf{x}_j - \mathbf{x}_k|^2}, \qquad (34)$$

interpreted in the sense of forms or expectation values, and where the so-called *N*-fractionality of α , defined

$$\alpha_N := \min_{p=0,1,...,N-2} \min_{q \in \mathbb{Z}} |(2p+1)\alpha - 2q|$$

is the arcwise distance (in units of π) on the unit circle from the set $\{e^{i(2p+1)\alpha\pi}\}_{p=0,1,\ldots,N-2}$ of all possible pairexchange phases to the point +1 of bosons. Taking the limit as $N \to \infty$ one finds (see Fig. 6)

$$\alpha_{N \to \infty} = \begin{cases} \frac{1}{\nu}, \text{ if } \alpha = \frac{\mu}{\nu} \text{ odd-numerator reduced rational,} \\ 0, \text{ otherwise,} \end{cases}$$

an odd variant of Thomae's "popcorn function".

The origin of the scalar lower bound (34) lies in the effective angular momentum barrier of each pair of particles at distance r > 0:

$$V_{\text{stat}}(r) = |(2p+1)\alpha - 2q|^2 \frac{1}{r^2} \ge \frac{\alpha_N^2}{r^2}.$$
 (36)

(35)

Namely, in an exchange of two particles, as in (7) where p other particles happen to be ensnared, an anyonic phase shift $e^{i(2p+1)\alpha\pi} \neq 1$ leaves a residual angular momentum according to (36). Heuristically, we can anticipate this effect in the two-particle energy form (12), where the quantization condition (10) in the angular variable φ is



FIG. 6. ∞ -fractionality of α given by the odd popcorn function (35) [Lundholm 2017, Lundholm and Solovej 2013a].

appropriately modified to account for the presence of any extra particles, assuming all other position variables are fixed and the relative angular variable undergoes a full loop $\varphi \rightarrow \varphi + \pi$ in configuration space. The anyonic phase then shifts the relative angular momentum of the pair, which however is quantized over the even integers due to the half-circle symmetry for identical particles.¹⁰

We expect the repulsive barrier (36) to be stronger than the r.h.s. on average as the number p fluctuates. Indeed, the coupling constant of the pair-interaction potential on the r.h.s. of (34) can be strengthened to a nearest-neighbor version, which only depends on the 2fractionality α_2 (periodized α), if it is at the same time weakened by yet another factor of N [Lundholm and Qvarfordt 2020, Rougerie and Yang 2023]. Anyway, the repulsion offered by the potential is strong enough to imply that $\Psi \to 0$ at Δ_N if $\alpha_2 \neq 0$, and can therefore be viewed as a **generalized Pauli principle** for anyons [Lundholm 2017, Lundholm and Qvarfordt 2020, Lundholm and Solovej 2013b] (other variants were considered in [Fröhlich and Marchetti 1988, 1989, Goldin and Majid 2004, Goldin and Sharp 1996]).

2. Local exclusion principle. The bound (34) (with its various refinements [Larson and Lundholm 2018, Lundholm and Qvarfordt 2020, Lundholm and Solovej 2013a]) can also be applied locally on subsets of the plane and is strong enough to yield a linear bound $E_N \gtrsim N-1$ for the local (Neumann) energy [Lundholm and Seiringer 2018, Lundholm and Solovej 2013b]. Again, this holds modulo a (positive) constant that either depends on α_N ,

¹⁰ The operator $\hat{\ell} = -i\partial_{\varphi}$ on $[0,\pi]$ with twisted boundary conditions $\psi(\pi) = e^{i\alpha\pi}\psi(0)$ has eigenvalues $\{\alpha - 2q\}_{q\in\mathbb{Z}}$. Therefore the minimum of $\hat{\ell}^2$ is $\min_{q\in\mathbb{Z}} |\alpha - 2q|^2 = \alpha_2^2$. Compare also (20).



FIG. 7. Lower bounds to the homogeneous ideal anyon gas energy [Larson and Lundholm 2018, Lundholm and Seiringer 2018]. The blue points are numerical lower bounds for $E_2(\alpha) = 4\pi\alpha + O(\alpha^{4/3})$ at $\alpha = \alpha_{N\to\infty}$, while the orange curve describes $F_2(\alpha_2) = \frac{1}{4} \min\{E_2(\alpha_2), 0.147\}$, correspondingly.

or a typically weaker one (still positive) that only depends on α_2 (cf. Fig. 7).

For comparison, take e.g. the unit square $Q = [0, 1]^2$ and consider fermions which there satisfy the bound

$$E_N(\alpha = 1, Q) \ge \pi^2 (N - 1) \quad \text{for } N \ge 1,$$
 (37)

since each added particle is orthogonal to the ground state (the constant function) and thus adds at least the energy $E_2 = \pi^2$ of a first-excited state $\psi(x, y) \propto \cos(\pi x)$ or $\cos(\pi y)$.

For anyons, we denote the N-particle **Neumann energy on** Q by

$$E_N(\alpha, Q) := \inf_{0 \neq \Psi \in L^2_{\text{sym}}(Q^N)} \langle \hat{T}_{\text{sym} \to \alpha} \rangle_{\Psi}.$$
 (38)

It is known that [Larson and Lundholm 2018, Lundholm and Seiringer 2018]

$$E_2(\alpha, Q) = 4\pi\alpha_2 + O(\alpha_2^{4/3}) \quad \text{for any } \alpha, \qquad (39)$$

and furthermore, all higher-particle energies can be bounded uniformly in terms of this energy, or a suitable approximation¹¹ \tilde{E}_2 to it: for $N \ge 2$,

$$E_N \ge (N-1)E_2(\alpha_N),\tag{40}$$

as well as

$$E_N/N \ge \frac{1}{4}\min\{E_2, E_3, E_4\} \ge \frac{1}{4}\min\{E_2(\alpha_2), 0.147\}.$$
(41)

¹¹ $\tilde{E}_2(\alpha) \approx 2\pi (j'_{\alpha_2})^2 \geq 4\pi\alpha_2$, where j'_{α} denotes the first zero of the derivative of the Bessel function J_{α} [Larson and Lundholm 2018].

This r.h.s. we denote by $F_2(\alpha_2)$; cf. Fig. 7. An interpretation of the above is that the nearest-neighbor energy E_2 is a relatively good approximation at every scale (at least as a lower bound) due to a balancing of uncertainty and exclusion in the gas.

3. **Degeneracy pressure.** Lastly, a combination of the above local exclusion principle and the uncertainty principle (also in a local formulation) yields a convenient measure of the degeneracy pressure in the anyon gas.

Compare the TF functional (27) which shows that the effect of the exclusion principle for fermions is a selfenergy for the density that penalizes localization and thus forces a spread of the profile into the more expensive regions of the trapping potential. Indeed, the ideal Fermi gas satisfies the following kinetic energy inequality (known in mathematics as a **Lieb–Thirring inequality** [Lieb and Thirring 1975]) which captures this type of degeneracy pressure: for any number N of particles and any N-fermion state $\Psi \in L^2_{asym}(\mathbb{R}^{2N})$,

$$\langle \hat{T}_{asym} \rangle_{\Psi} \gtrsim \int_{\mathbb{R}^2} \varrho_{\Psi}(\mathbf{x})^2 \, d\mathbf{x}$$
 (42)

(modulo a universal constant¹²), where ρ_{Ψ} is the onebody density associated to Ψ , obtained by marginalizing N-1 particles. Inequalities of this type have been useful to prove qualitative properties of large fermionic systems such as their thermodynamic stability w.r.t. Coulomb interaction with a mixture of charges [Dyson and Lenard 1967, Lieb and Seiringer 2010, Lieb and Thirring 1975]. In the case that ν species of fermions or spin states are considered, the constant in (42) is weakened by a factor $1/\nu$.

The degeneracy pressure for anyons is measured by a similar inequality [Lundholm and Solovej 2013a,b] (see [Lundholm and Qvarfordt 2020] for review): for any number N of particles and any N-anyon state $\Psi \in L^2_{\alpha}$,

$$\langle \hat{T}_{\alpha} \rangle_{\Psi} \gtrsim \alpha_2 \int_{\mathbb{R}^2} \varrho_{\Psi}(\mathbf{x})^2 d\mathbf{x}$$
 (43)

(again modulo a universal constant). Its strength thus depends to leading order on the 2-fractionality, i.e. the nearest-neighbor exchange statistics α . It then follows, like for fermions, that charged anyonic systems (except bosons) with Coulomb interaction are also stable [Lundholm and Seiringer 2018, Lundholm and Solovej 2014].

Applied to the harmonic trap (19), the degeneracy pressure (43) yields the bound

$$E_N(\alpha, \text{harm.osc.}) \gtrsim \sqrt{\alpha_2} \,\hbar\omega N^{3/2},$$
 (44)

¹² Deciding on the best constant in this inequality is a long-standing open problem, but it is conjectured [Lieb and Thirring 1976] to be given by that of its restriction to N = 1 (a Sobolev constant) and is strictly smaller than the TF constant 2π . See [Seiringer and Solovej 2023] for recent progress.



FIG. 8. Universal energy phenomena in 2D, from [Lankhorst et al. 2018]: lowest energy branch of a 100×100 square Josephson junction array (left/blue) resp. the Hofstadter butterfly (right/orange), which also is proportional to the critical temperature of a superconducting square network. (For comparison, $\alpha \sim 2f$.)

which indeed matches the constant-field approximation (33), and Fig. 5, up to the value of the universal constant.

On a disk geometry D with unit area, the analysis of Chitra and Sen again suggests a large number of level crossings as $N \to \infty$, and thus possibly nonanalyticities in the limiting g.s.e. per particle and unit density of the **homogeneous ideal anyon gas**¹³

$$e(\alpha) := \liminf_{N \to \infty} E_N(\alpha, D) / N^2.$$
(45)

Most details about this function are still open, but from a further analysis of the homogeneous ideal anyon gas on the unit square Q (dividing it into smaller squares), one can at least derive the rigorous uniform bounds (cf. Fig. 7) [Lundholm and Seiringer 2018]

$$\frac{1}{4}\max\{\tilde{E}_2(\alpha_N), F_2(\alpha_2)\} \le e(\alpha) \le 2\pi^2, \qquad (46)$$

and

$$\alpha_2 \lesssim e(\alpha) \lesssim \alpha_2,\tag{47}$$

modulo lower and upper universal constants (certainly not optimal). Subject to the assumption that $e(\alpha)$ indeed has a fractal structure which captures universal aspects of planar geometry, a conjecture on its possible form can be given as in Fig. 8; cf. [Lankhorst et al. 2018].



FIG. 9. From [Lundholm 2017]. A coloring of N = 12 particles with $\nu = 3$ colors into K = 4 3-clusters \mathcal{V}_q^* . Each colored edge $(j,k) \in \mathcal{E}_q$ corresponds to one unit $-\mu$ of pairwise angular momentum. Also shown is the contribution to the magnetic potential $\alpha \mathbf{A}_j$ and the current \mathbf{J}_j of particle jdue solely to the 3-cluster \mathcal{V}_1^* .

H. Some states of particular interest

We saw above that there is a nontrivial and delicate balance between exclusion and uncertainty in the ground state energy. Certain types of states have been proposed [Lundholm 2017, Lundholm and Solovej 2013b] to minimize the kinetic energy $\hat{T}_{\text{sym}\to\alpha}$ at particular values of the statistics parameter: in the case of $\alpha = \mu/\nu \in [0, 1]$ being an *even*-numerator reduced fraction, and for $N = \nu K$, $K = 1, 2, \ldots$, a suitable sequence of particle numbers, let

$$\Psi_{\alpha} := \prod_{j < k} |z_{jk}|^{-\alpha} \mathcal{S}\left[\prod_{q=1}^{\nu} \prod_{(j,k) \in \mathcal{E}_q} (\bar{z}_{jk})^{\mu}\right] \prod_{l=1}^{N} \psi_0(\mathbf{x}_l),$$
(48)

while for *odd* numerators μ , let

$$\Psi_{\alpha} := \prod_{j < k} |z_{jk}|^{-\alpha} \mathcal{S} \left[\prod_{q=1}^{\nu} \prod_{(j,k) \in \mathcal{E}_q} (\bar{z}_{jk})^{\mu} \bigwedge_{k=0}^{K-1} \psi_k \left(\mathbf{x}_{l \in \mathcal{V}_q} \right) \right].$$
(49)

Here $z_{jk} := z_j - z_k$ are the pairwise relative complex coordinates and we have grouped, or 'colored', the particles into ν different colors where $G_q = (\mathcal{V}_q, \mathcal{E}_q)$ denotes the complete graph over each such group of $|\mathcal{V}_q| = K$ vertices=particles (see Fig. 9). The symmetrization \mathcal{S} over all the particles then amounts to symmetrization over all such colorings, and can be viewed as passing from a set of distinguishable particles (by color) to indistinguishable (cf. [Regnault et al. 2008]). The ψ_k , $k = 0, 1, 2, \ldots$, are the eigenstates (ordered by increasing energy) of the corresponding one-body Hamiltonian \hat{H}_1 . The states Ψ_{α} are moderately singular and so we should actually consider $\Psi = \Phi \Psi_{\alpha} \in L^2_{\text{sym}}(\mathbb{R}^{2N})$ with Φ a suitably chosen regularization factor to allow inclusion into the domain of $\hat{T}_{\text{sym}\to\alpha}$.

Some of the reasons why these states appear to be of interest in the many-anyon problem include:

¹³ Although it has not been rigorously shown, we may expect the thermodynamic limit to be independent of the shape of the domain (and otherwise take the infimum over reasonable shapes).

• They have the right average angular momentum (24) and are (for harmonic trap and certain K) homogeneous with a degree suitable to optimize the bound (23).

• They enjoy certain clustering properties that are beneficial to minimize the pair repulsion $V_{\text{stat}}(r)$ on large scales r. States of the form (48) for $\mu = 2$, $\nu =$ $3, 5, \ldots$ are known in the FQHE context as Read–Rezayi states and are known to form clusters of ν particles bound to μ vortices [Cappelli et al. 2001, Read and Rezayi 1999], while (49) for $\mu = 1$, $\nu = 2$ correspond to Dyson–Moore–Read/BCS/Pfaffian states subject to pairing [Dyson 1967, Moore and Read 1991]. Further, excitations on top of these states appear to have nonabelian character [Bonderson et al. 2011, Nayak et al. 2008, Stern 2008].

• They manifest certain structural stabilities with the smallness of $1 \le \mu \le \nu$, similar to the Laughlin states of the FQHE and indicative of a scale of robustness.

• Some of the states (48) are exact but moderately singular/generalized low-energy eigenfunctions of the Hamiltonian with singular boundary conditions (cf. point interactions below, Sec. IIIE) [Chou 1991, Murthy et al. 1992].

III. THE NONIDEAL ANYON GAS

The "average(/constant)-field" approach, suggesting (29), leads to difficulties when applied to strictly ideal and pointlike particles, and it is both realistic and helpful to instead consider extended particles as well as other nonideal models of anyon gases which can more easily incorporate the average effects of exchange phases or flux. Heuristic approaches to ensure averaged influence of the statistics include a "self-consistency argument" wherein a typical cyclotron orbit of the generated field ought to contain a large number of particles, suggesting a better approximation if [Chen et al. 1989, Trugenberger 1992b]

$$\alpha_2 \ll 1$$
 or $(1 - \alpha_2)^2 \ll 1$ (50)

(i.e. slightly better around fermions due to Pauli repulsion), as well as a similar orbital argument but based instead on the scattering angles of combined Aharonov–Bohm and hard-disk interaction [Caenepeel and MacKenzie 1994], again suggesting the conditions (50).

A. The extended anyon gas

In the **extended anyon gas**, we work in the magnetic perspective and replace the point magnetic fluxes of ideal anyons by extended fluxes. For simplicity (and not completely ad hoc from the perspective of emergent models) we may consider the flux to be constantly spread over a disk-shaped magnetic field attached to each particle, with radius R > 0. We then replace the anyon magnetic potentials (15) by

$$\mathbf{A}_{j}^{R}(\mathbf{x}) := \sum_{k \neq j} \frac{(\mathbf{x}_{j} - \mathbf{x}_{k})^{\perp}}{|\mathbf{x}_{j} - \mathbf{x}_{k}|_{R}^{2}}, \qquad |\mathbf{x}|_{R} := \max\{|\mathbf{x}|, R\},$$
(51)

so that, if $\mathbb{1}_{D(\mathbf{y},R)}$ denotes the indicator function on a disk of radius R centered at \mathbf{y} , the field is

$$\operatorname{curl}_{\mathbf{x}_j} \alpha \mathbf{A}_j^R = 2\pi\alpha \sum_{k \neq j} \frac{\mathbbm{1}_{D(\mathbf{x}_k, R)}(\mathbf{x}_j)}{\pi R^2} \xrightarrow{R \to 0} 2\pi\alpha \sum_{k \neq j} \delta_{\mathbf{x}_k}(\mathbf{x}_j)$$

The kinetic energy of the extended anyon gas with bosons as reference is then

$$\hat{T}_{\mathrm{sym}\to\alpha}^{R} := \frac{\hbar^2}{2m} \sum_{j=1}^{N} \left(-i\nabla_{\mathbf{x}_j} + \alpha \mathbf{A}_j^{R} \right)^2,$$

acting on L^2_{sym} , while if we use fermions as reference then

$$\hat{T}_{\text{asym}\to\alpha}^{R} := \frac{\hbar^2}{2m} \sum_{j=1}^{N} \left(-i\nabla_{\mathbf{x}_j} + (\alpha - 1)\mathbf{A}_j^R \right)^2,$$

acting on L^2_{asym} .

There is now an additional natural dimensionless parameter in the problem that can be defined as the ratio of the size of the magnetic flux disk to the average interparticle distance,

$$\bar{\gamma} := R\bar{\varrho}^{1/2}.$$
(52)

This density parameter has been called the **magnetic** filling ratio in [Larson and Lundholm 2018, Trugenberger 1992a,b]. The limit $\bar{\gamma} = 0$ corresponds to ideal anyons, while $\bar{\gamma} \sim 1$ is the "smearing" limit when particles typically begin to overlap, and $\bar{\gamma} \to \infty$ would describe the constant-field limit.

At any fixed choice of the two parameters $\alpha \in \mathbb{R}$, $\bar{\gamma} \geq 0$, the energy per particle and unit of density of the **homo-geneous extended anyon gas** may thusly be defined via the thermodynamic limit

$$e_{\underset{\text{asym}}{\text{sym}}}(\alpha,\bar{\gamma}) := \liminf_{\substack{N,L\to\infty\\N/L^2=\bar{\varrho}}} \frac{E_N(\alpha,R,LQ)}{\bar{\varrho}N}, \qquad (53)$$

where $E_N(\alpha, R, LQ)$ denotes the g.s. energy of the respective operator $\hat{T}^R_{\text{sym/asym}\to\alpha}$ on $L^2_{\text{sym/asym}}(LQ^N)$. A corresponding TF-type functional for a **local-density approximation** within a trapping potential V is then

$$\mathcal{E}_{\substack{\text{sym}/\\\text{asym}}}^{\alpha,R}[\varrho] := \int_{\mathbb{R}^2} \left[e_{\substack{\text{sym}/\\\text{asym}}} (\alpha, R\varrho(\mathbf{x})^{1/2}) \varrho(\mathbf{x})^2 + V(\mathbf{x})\varrho(\mathbf{x}) \right] d\mathbf{x}$$
(54)

with $\int_{\mathbb{R}^2} \varrho = N$.

The qualitative analysis of local exclusion effects of the ideal gas discussed in Section IIG has been extended to

the extended gas, both in the bosonic and the fermionic references. Up to constants, one finds the qualitative bounds [Larson and Lundholm 2018]

$$e_{\rm sym}(\alpha,\bar{\gamma}) \gtrsim \begin{cases} |\ln \bar{\gamma}|^{-1} + \alpha_{\infty}, & \bar{\gamma} \ll 1 \ (\alpha \text{ fixed}), \\ |\alpha|, & \bar{\gamma} \gtrsim 1, \end{cases}$$

where the first regime is a combination of a dilute 2D Bose gas [Lieb and Yngvason 2001, Schick 1971] and the strongest of the two lower bounds (46) for the ideal gas at odd numerators, $\tilde{E}_2(\alpha_{\infty}) \gtrsim \alpha_{\infty}$, while the second regime indeed captures the constant-field energy (29) for arbitrarily large α . The full regime $\bar{\gamma} \lesssim 1$ is more difficult to describe, since it covers both soft-core and hard-core behavior (see Sec. III F below), and we should expect to replace α_{∞} with α_2 in a suitable limit. Trugenberger [Trugenberger 1992b] suggested that this gas with fixed extension parameter R > 0 will at low temperatures prefer to adjust its density to $\bar{\gamma} \gtrsim 1$, where the average-field approximation is good and the gas becomes a superfluid, while for high temperatures the density drops, $\bar{\gamma} \ll 1$, the average-field loses validity and superfluidity is lost.

In the fermionic reference [Girardot and Rougerie 2022]

$$e_{\rm asym}(\alpha,\bar{\gamma})\gtrsim\begin{cases} \alpha_2, & \bar{\gamma}\lesssim 1,\\ 1, & \bar{\gamma}\gtrsim 1. \end{cases}$$

In this case the inherent Pauli principle improves the dependence to nearest neighbor due to its scale independence. Note that both energies $e_{\text{sym/asym}}$ must be periodic in α at $\bar{\gamma} = 0$, but not at any finite $\bar{\gamma} > 0$, which therefore requires an interesting interpolation between the regimes.

B. Almost-bosonic anyons

The average-field approach (29) can be made fully rigorous in the limit $N \to \infty$ close to bosons:

$$\alpha = \beta / N \to 0$$
, at fixed $\beta \in \mathbb{R}$,

in which the interaction may indeed be treated in a meanfield sense if anyons are also extended/smeared out to a finite size R > 0, as above. It is even possible to simultaneously take $R \sim N^{-\eta} \rightarrow 0$ at some sufficiently slow rate η (an "almost-ideal" limit). The idea is then as a final step to take $\beta = \alpha N$ (total flux) large (then "less bosonic") and study the appropriate ground-state energy functionals in this limit. Because of this particular order of limits, it is not completely clear whether this indeed describes the ideal anyon gas as defined above, or perhaps some nonideal model (see Sec. III F). We shall simply refer to it as "almost-bosonic" anyons.

In any case, the first steps above yield the seemingly *correct* **average-field functional** w.r.t. bosons for any fixed $\beta \in \mathbb{R}$ [Lundholm and Rougerie 2015]:

$$\mathcal{E}_{\beta}^{\mathrm{af}}[\psi] = \int_{\mathbb{R}^2} \left[\left| (-i\nabla + \beta \mathbf{A}[|\psi|^2])\psi \right|^2 + V|\psi|^2 \right].$$
(55)

The magnetic potential $\mathbf{A}[\varrho = |\psi|^2]$ is chosen such that it generates exactly the field $2\pi \varrho$:

$$\operatorname{curl} \beta \mathbf{A}[\varrho](\mathbf{x}) = 2\pi\beta \varrho(\mathbf{x}), \qquad \int_{\mathbb{R}^2} \varrho = 1.$$
 (56)

Thus, the *scalar* self-interaction of the GP model (28) is here replaced with a *magnetic* self-interaction. This produces a peculiar new balancing problem, namely in order to lower its energy in the self-generated magnetic field, as the coupling β (total number of flux units) grows there needs to be an increasing phase and angular momentum in $\psi = |\psi|e^{i\phi}$ to cancel this magnetic field:¹⁴

$$\left| (-i\nabla + \beta \mathbf{A})\psi \right|^2 = \left| \nabla |\psi| \right|^2 + \left| \nabla \phi + \beta \mathbf{A} \right|^2 |\psi|^2.$$
 (57)

Further, in order to preserve regularity, any nontrivial such phase circulation must appear in the form of quantized vortices. On a typical disk $D(\mathbf{x}, R)$, the phase circulation

$$\Phi(\mathbf{x}, R) := \oint_{\partial D(\mathbf{x}, R)} \beta \mathbf{A} \cdot d\mathbf{r} \approx -\int_{D(\mathbf{x}, R)} \operatorname{curl} \nabla \phi \, d\mathbf{y}$$
(58)

then needs to minimize the residual angular momentum,

$$\int_{0}^{2\pi} \left| \partial_{\varphi} \phi + \beta A_{\varphi} \right|^{2} d\varphi \ge \min_{q \in \mathbb{Z}} \left| q - \Phi(\mathbf{x}, R) \right|^{2}.$$
(59)

However, the amount of flux to be cancelled by each vortex also depends on the typical value of the density ρ there, which therefore sets a varying length scale for vortex formation [Correggi et al. 2017]. It is thus expected (see also [Chen et al. 1989] for an early conjecture) and indeed confirmed in numerical studies [Correggi et al. 2019, Girardot 2021] that there is the formation of an Abrikosov-like vortex lattice. However, unlike in the Abrikosov setting with homogeneous rotation/field, the density of vortices here follows the average density profile in the trap V, which for $\beta \gg 1$ turns out to be given again by the minimizer of an effective **Thomas–Fermi-type** functional [Correggi et al. 2017]:

$$\mathcal{E}_{\beta}^{\mathrm{aTF}}[\varrho] := \int_{\mathbb{R}^2} \left[C_{\mathrm{aTF}} |\beta| \varrho(\mathbf{x})^2 + V(\mathbf{x}) \varrho(\mathbf{x}) \right] d\mathbf{x}.$$
(60)

Indeed, numerical simulations [Correggi et al. 2019, Girardot 2021] (see Fig. 10) confirm the theoretical prediction of [Correggi et al. 2017, 2018] that the g.s. ψ_{β}^{af} of (55) exhibits an approximately triangular vortex lattice distribution with scales set by the TF profile (minimizer) $\varrho_{\beta}^{\text{aTF}}$ of (60), and it is further estimated numerically that [Correggi et al. 2019]

$$C_{\rm aTF} \approx 4\pi^{3/2}/3 \approx 1.18 \times 2\pi.$$
 (61)

¹⁴ Due to the reasons described below, the cancellation is not complete, but both terms in (57) contribute an energy of order β .



FIG. 10. Numerical simulation of the ground-state density $|\psi_{\beta}^{\rm af}|^2$ in the average-field functional (55) for $\beta = 90$ in a harmonic trap, averaged over rotations and then compared with the exact minimizer $\varrho_{\beta}^{\rm aTF}$ of (60). From [Correggi et al. 2019].

The fact that this constant is strictly bigger than the fermionic TF constant and constant-field prediction¹⁵ 2π can be understood from the microscopic lattice inhomogeneity of $|\psi_{\beta}^{\text{af}}|^2$ as compared to its average density $\rho_{\beta}^{\text{aTF}}$, which effectively raises the energy by a factor which is at least as big as the Abrikosov constant (≈ 1.1596 ; cf. [Aftalion et al. 2006a,b]).

C. Almost-fermionic anyons

A similar regularization approach can be taken close to fermions:

$$\alpha = 1 - \beta / \sqrt{N} \to 1, \qquad \beta, \hbar \text{ fixed},$$

for which the leading terms of the kinetic energy can be compared with the Fermi gas $E_N(\alpha = 1) \sim 2\pi N^2$, $N \to \infty.$ By rescaling the parameters, it is equivalent to study a semiclassical mean-field limit

$$\alpha = 1 - \beta/N \to 1, \qquad \hbar \sim N^{-1/2} \to 0.$$

Also here we can allow "almost-ideal" or "virtually" extended anyons, $0 < R \sim N^{-\eta} \rightarrow 0$, if the rate η is small enough¹⁶.

This **almost-fermionic** approach takes us back to the actual TF functional for fermions (27), which remains correct for any finite β in this limit [Girardot 2021, Girardot and Rougerie 2021]. More precisely, one obtains a semiclassical **Vlasov functional**:

$$\mathcal{E}_{\beta}^{\mathrm{Vla}}[\mu] := \frac{1}{(2\pi)^2} \int_{\mathbb{R}^4} \left| \mathbf{p} + \beta \mathbf{A}[\varrho] \right|^2 \mu(\mathbf{x}, \mathbf{p}) \, d\mathbf{x} d\mathbf{p} + \int_{\mathbb{R}^2} V \varrho \, d\mathbf{x}$$

for $0 \leq \mu(\mathbf{x}, \mathbf{p}) \leq 1$ a measure on phase space \mathbb{R}^4 which is subject to the Pauli principle and normalized $\int_{\mathbb{R}^4} \mu = (2\pi)^2$. The minimizing measure μ of this functional is the corresponding filled Fermi sea:

$$\mu(\mathbf{x}, \mathbf{p}) = \mathbb{1}\left\{ |\mathbf{p} + \beta \mathbf{A}[\varrho](\mathbf{x})|^2 \le 4\pi \varrho(\mathbf{x}) \right\}, \qquad (62)$$

with spatial density $\rho = \rho^{\text{TF}}$ independent of β :

$$\varrho(\mathbf{x}) = \frac{1}{(2\pi)^2} \int_{\mathbb{R}^2} \mu(\mathbf{x}, \mathbf{p}) \, d\mathbf{p} = (4\pi)^{-1} (\lambda^{\text{TF}} - V(\mathbf{x}))_+.$$

The momentum density $t = t_{\beta}^{\text{TF}}$,

$$t(\mathbf{p}) = \frac{1}{(2\pi)^2} \int_{\mathbb{R}^2} \mu(\mathbf{x}, \mathbf{p}) \, d\mathbf{x},$$

does depend on β however, in a nonlocal way. The energy $\mathcal{E}_{\beta}^{\text{Vla}}[\mu] = \mathcal{E}^{\text{TF}}[\varrho^{\text{TF}}]$ is locally a constant at $\alpha \approx 1$ in this approximation, thus confirming this aspect of Fig. 5.

The almost-fermionic regime was also studied numerically by Girardot et al. in [Girardot 2021] using a **Hartree–Fock** approach, i.e. on Slater determinants, and with a self-generated field $B = 2\pi\beta\varrho/N$ (total flux $2\pi\beta$), showing good agreements to the above. In fact, it can be interpreted as a multicomponent version of the bosonic functional (55) on orthogonal states ψ_k , $k = 1, 2, \ldots, N$. These are then able to complement each other's densities so as to fill in any holes left by vortices, and thus eventually smoothen out the overall density profile:

$$\varrho(\mathbf{x}) = \sum_{k=1}^{N} |\psi_k(\mathbf{x})|^2 \approx \varrho^{\mathrm{TF}}(\mathbf{x}), \qquad (63)$$

thereby validating the "constant-field" (local density) approximation in this regime when β is not too large.

¹⁵ Also in comparison to Fig. 5, Chitra and Sen, who estimated a substantially larger factor [Chitra and Sen 1992], assumed rotational symmetry for the density to simplify their functional, plus added a hard-core delta regularization that itself amounts to the constant-field energy $2\pi |\beta| \varrho^2$.

 $^{^{16}}$ The rate is better than for bosons because of the Pauli principle; cf. (50).

D. Magnetic TF theory

For higher ratios of the magnetic field to the density one may expect [Girardot 2021] a **magnetic Thomas-Fermi theory** [Lieb et al. 1995, McEuen et al. 1992] to become valid. Namely, for a field $B(\mathbf{x})$ which is approximately constant on the scale of the trapping potential V, we can think of distributing the density

$$\varrho(\mathbf{x}) = \sum_{n=0}^{\infty} \varrho_n(\mathbf{x})$$

locally into the Landau levels (LLs) \mathcal{H}_n of the corresponding one-body Landau Hamiltonian¹⁷

$$\hat{H}_{1}^{\text{Lan}} = (-i\nabla_{\mathbf{x}} + B\mathbf{x}^{\perp}/2)^{2} = \sum_{n=0}^{\infty} |B|(2n+1)\mathbb{1}_{\mathcal{H}_{n}}.$$
 (64)

The degeneracy (per unit area) of each level is the number of flux units (per unit area), $d_B := |B|/(2\pi)$. Thus, defining the corresponding expected local magnetic energy at a given density ρ ,

$$j_B(\varrho) := \sum_{n=0}^{\infty} |B|(2n+1)\varrho_n, \qquad 0 \le \varrho_n \le d_B,$$

the **magnetic TF functional** for N fermions in a fixed *external* field B > 0 is then

$$\mathcal{E}^{\mathrm{mTF}}[\varrho] := \int_{\mathbb{R}^2} \left[j_B(\varrho(\mathbf{x})) + V(\mathbf{x})\varrho(\mathbf{x}) \right] d\mathbf{x}, \quad \int_{\mathbb{R}^2} \varrho = N.$$
(65)

At fixed B and ρ , in order to minimize $j_B(\rho)$ we fill all $N_B := \lfloor \rho/d_B \rfloor$ lower Landau levels maximally:

$$\varrho_n = \begin{cases} d_B, & 0 \le n \le N_B - 1, \\ \varrho - N_B d_B, & n = N_B, \\ 0, & n \ge N_B + 1, \end{cases}$$

so that in the limit $B \to 0$, implying $d_B \to 0$,

$$j_B(\varrho) \approx \sum_{n=0}^{N_B - 1} |B|(2n+1)d_B = |B|N_B^2 d_B$$
$$\approx |B|\varrho^2/d_B = 2\pi \varrho^2,$$

returning us to the usual TF for the free Fermi gas. On the other hand, if B is large enough that $d_B \ge \rho$, then all particles will fit into the lowest Landau level (LLL):

$$j_B(\varrho) = |B|\varrho. \tag{66}$$

Now, considering a sufficiently extended anyon gas and formally taking the anyonic (average) magnetic field $B(\mathbf{x}) = 2\pi\beta \varrho(\mathbf{x})$, where $\beta = 1 - \alpha \in [0, 1]$ is the number of flux units per particle attached to fermions, we then obtain a **magnetic TF self-interaction functional** for anyons:

$$\mathcal{E}_{\beta}^{\mathrm{amTF}}[\varrho] := \int_{\mathbb{R}^2} \left[j_{B=2\pi\beta\varrho(\mathbf{x})}(\varrho(\mathbf{x})) + V(\mathbf{x})\varrho(\mathbf{x}) \right] d\mathbf{x}.$$
(67)

At any point $\mathbf{x} \in \mathbb{R}^2$ where $\rho(\mathbf{x}) \neq 0$ let us consider there the fractions of density $f_n := \rho_n(\mathbf{x})/\rho(\mathbf{x})$ in the LLs:

$$\frac{j_B(\varrho)}{2\pi\varrho^2} = \sum_{n=0}^\infty \beta(2n+1)f_n, \qquad 0 \le f_n \le \beta \le 1,$$

which again is minimized by filling the lowest $N_B = \lfloor \beta^{-1} \rfloor$ levels:

$$f_n = \begin{cases} \beta, & 0 \le n \le \lfloor \beta^{-1} \rfloor - 1, \\ 1 - \beta \lfloor \beta^{-1} \rfloor, & n = \lfloor \beta^{-1} \rfloor, \\ 0, & n \ge \lfloor \beta^{-1} \rfloor + 1. \end{cases}$$

Hence, our minimum is

$$\frac{j_B(\varrho)}{2\pi\varrho^2} = \sum_{n=0}^{\lfloor\beta^{-1}\rfloor-1} (2n+1)\beta^2 + \beta(2\lfloor\beta^{-1}\rfloor+1)(1-\beta\lfloor\beta^{-1}\rfloor),$$

which simplifies to

$$\frac{j_B(\varrho)}{2\pi\varrho^2} - 1 = \beta^2 (1 - \{\beta^{-1}\})\{\beta^{-1}\} =: M(\beta) \in [0, \beta^2/4),$$

where $\{\beta^{-1}\} = \beta^{-1} - \lfloor \beta^{-1} \rfloor \in [0, 1)$ is the fractional part of β^{-1} . Therefore our proposed functional for the g.s. (a pointwise lower bound to (67)) takes the form

$$\tilde{\mathcal{E}}_{\beta}^{\mathrm{amTF}}[\varrho] := \int_{\mathbb{R}^2} \left[2\pi \varrho(\mathbf{x})^2 \left(1 + M(\beta) \right) + V(\mathbf{x})\varrho(\mathbf{x}) \right] d\mathbf{x},$$
(68)

thus refining both the "constant-field" approximation (29) around fermions as well as the above almost-fermionic approximation.

The corresponding energy landscape (see Fig. 11) is now indicative of **stability** at the Fermi point $\beta = 0$. A possible interpretation for the gas is that the discretization of the Fermi sea into Landau levels imposes a more rigid structure in phase space, i.e., as the attached flux β per particle increases, particles come from plane-wave states which homogenize easily in space and then need to start organizing into the respective cyclotron orbits with decreasing radii as the field increases. Thus the Fermi sea becomes less fluid and more crystalline, with the most synergetic situation when a whole number of levels are approximately filled ($\{\beta^{-1}\} \approx 0 \text{ or } 1$) so adjustments to the density can be accommodated at all levels in parallel while keeping the fractions f_n constant, and least synergetic when half a level is filled $(\{\beta^{-1}\} \approx 1/2)$ and a larger fraction $\beta\{\beta^{-1}\}$ of particles need to organize their motion into that specific LL. However, we cannot expect this picture to be valid all the way to bosons at $\beta = 1$ (compare

¹⁷ In our units the cyclotron frequency is $\omega_c = 2|B|$.



FIG. 11. The factor $M(\beta) = \beta^2 (1 - \{\beta^{-1}\}) \{\beta^{-1}\} \le \beta^2/4$ (r.h.s. dashed) in the magnetic TF functional (68) as a function of the relative statistics parameter $\beta = 1 - \alpha$ around fermions $\beta = 0$.

also (50)) because there will also be correlations across Landau levels, leading to effective attractions, clustering and condensation, and for example, if the LLL is filled and there are only few particles in the first-excited LL then one might consider the statistics transmutation situation discussed below where the excited particles would effectively rather be treated as bosons than fermions.

An analysis at these seemingly beneficial fractions $\alpha = 1 - 1/n$ was conducted by Chen et al. [Chen et al. 1989] in the homogeneous setting (building on [Fetter et al. 1989]), where they indeed observed characteristics of **superfluidity**. For example, the energy there increases linearly when a small external magnetic field is applied in either direction, which is interpreted as a Meissner effect. Recently another approach to a DFT for anyons, via Kohn–Sham theory with neglected exchange correlation (then similar to self-consistent Hartree–Fock), was taken by Hu et al. to study these effects [Hu et al. 2021]. Their numerics agree well with the semiclassical TF approximation close to fermions and show significant deviations only when the magnetic field is strong enough that particles in the bulk occupy only the lowest few LLs. For trapped systems they also observe a boundary effect which can compensate for the otherwise necessary fractional filling of LLs, but which vanishes in the thermodynamic limit. Indeed, due to the many parallels to the FQHE, Chen et al. suggested that "the physics of the anyon gas at general values of θ is likely to be quite rich and to depend quite strongly on 'number-theoretic' properties of θ ."

E. Point-interacting anyons

Introducing **point or contact interactions** corresponds to specifying the behavior of wave functions at the diagonal (contact) set Δ_N , i.e. imposing boundary conditions there so as to render the Hamiltonian selfadjoint, while allowing a different behavior than that of the canonical/free choice (Friedrichs; cf. Sec. IIC). This is equivalent to considering *all* of the possible boundary conditions, or self-adjoint extensions, of the operator T_{α} starting from its initial definition on wave functions vanishing at Δ_N and having finite expected energy in the standard sense (11). For general N this is difficult even for distinguishable particles, but one may reduce this problem significantly by adding assumptions such as translation and scale covariance. Our present understanding of general point-interacting (and pointlike, i.e. non-extended) anyons is currently rather limited (cf. [Dell'Antonio et al. 1997]), however for N = 2 anyons the situation is now well understood [Bourdeau and Sorkin 1992, Correggi and Oddis 2018, Coutinho et al. 1992, Grundberg et al. 1991, Manuel and Tarrach 1991, Murthy et al. 1992, Oddis 2020], and extends the corresponding analysis of the standard non-magnetic two-body pointinteraction problem in various dimensions (see, e.g., [Albeverio et al. 2005]).

For $\alpha_2 \neq 1$ (non-fermions) the problem admits a circle of different possibilities for point interaction, all of which are attractive and correspond to allowing wave functions with singularities at the diagonals. One simple way to understand this is via the form (12), which in the s-wave (radially symmetric) with $\Psi = r^{\pm \alpha_2} \phi(r)$ turns out to be equivalent to [Manuel and Tarrach 1991, Oddis 2020]

$$\int_0^\infty |\phi'(r)|^2 r^{d_\alpha - 1} dr,\tag{69}$$

for a parameter giving an effective "intermediate dimension"

$$2 \le d_{\alpha} := 2 + 2\alpha_2 \le 4. \tag{70}$$

Thus, for bosons the problem is indeed two-dimensional and has a circle of extensions, where a unique one $T_{\rm sym}$ (Friedrichs, $\Psi \sim 1$ as $r \to 0$) is nonnegative. For fermions it is more akin to four dimensions where the corresponding bosonic problem is essentially self-adjoint, i.e. has a unique extension \hat{T}_{asym} (Friedrichs, $\Psi \sim r$). For proper anyons the situation is intermediate and actually closest akin to three dimensions. In this case there is again a circle of extensions, but the circle now splits into two halves (cf. Fig. 2): in one half one has negative extensions and a scale of increasingly negative bound states, and in the other a scale of decreasingly nonnegative (i.e. increasingly attractive) extensions. The largest (least attractive / "free") such extension is the usual free kinetic energy $\hat{T}_{\alpha} = \hat{T}_{\alpha}^{\mathrm{F}}$ (Friedrichs, $\Psi \sim r^{\alpha_2}$), while the smallest $\hat{T}_{\alpha}^{\mathrm{K}}$ (most attractive, $\Psi \sim r^{-\alpha_2}$, and known as the **Krein** extension [Alonso and Simon 1980]) is also singled out as scale covariant while all the other intermediate extensions have scale.

If these observations extend to $N \ge 3$ particles then it is reasonable to expect such "**kreinyons**" to be of interest in physics, and in fact may even correspond to some of the moderately singular "noninterpolating" exact eigenstates [Murthy et al. 1992] found in the harmonic oscillator problem (cf. Section II H). Other relevant connections of anyons to geometry and quantum mechanics on cone-like manifolds were reviewed in [Jackiw 1990].

F. Other interactions and fields

One can obviously also include other (scalar) interactions into the magnetic models for anyons, as well as an external magnetic field (coupled by a charge -q), e.g.

$$\hat{H}_N := \frac{\hbar^2}{2m} \sum_{j=1}^N \left(-i \nabla_{\mathbf{x}_j} + q \mathbf{A}_{\text{ext}} + \alpha \mathbf{A}_j^R \right)^2 + \sum_{j=1}^N V(\mathbf{x}_j) + \sum_{j < k} W(\mathbf{x}_j - \mathbf{x}_k). \quad (71)$$

Reasonable external fields have been incorporated into some parts of the average-field theory [Girardot 2020, Girardot and Rougerie 2021], as well as point interactions [Correggi and Fermi 2021, Oddis 2020], and interacting anyons are certainly relevant in emergent models, as we will see below. Coulomb interactions and the thermodynamic stability of "anyonic matter" was addressed in [Lundholm 2019, Lundholm and Seiringer 2018, Lundholm and Solovej 2014]. We note that since scalar interactions can show features akin to statistics, such as effective TF-type functionals and degeneracy pressure [Larson et al. 2021, Lieb et al. 2005, Lundholm 2019, Lundholm et al. 2015, 2016], it is important to isolate the features that are due to the actual quantum statistics (cf. [Lambert et al. 2023]).

Another way to view the effect of the magnetic flux disks in the average-field approximation is as an effective scalar potential

$$W(\mathbf{x}_j - \mathbf{x}_k) = \frac{2\pi |\alpha|}{\pi R^2} \mathbb{1}_{D(\mathbf{x}_k, R)}(\mathbf{x}_j)$$
(72)

added to anyons (in the bosonic reference this is indeed correct as a lower bound [Larson and Lundholm 2018]). One may then consider the combination of parameters

$$\frac{|\alpha|}{\bar{\rho}R^2} = |\alpha|\bar{\gamma}^{-2},\tag{73}$$

i.e. the height of the potential compared to the average density, as a dimensionless measure of interaction strength, and for fixed α and $\bar{\gamma} \ll 1$ this then corresponds to a hard-core interaction, while if $|\alpha|\bar{\gamma}^{-2} \ll 1$, such as if $\sqrt{|\alpha|} \ll \bar{\gamma}$ where either side may be taken to grow or decline, then W instead corresponds to a weak soft-core interaction. With the convergence rates η currently allowed in the derivations of almost-bosonic and almost-fermionic models ($0 < \eta < 1/4$ [Girardot 2020] resp. $0 < \eta < 1/3$ [Girardot and Rougerie 2022]), we find that on a fixed homogeneous trap, $\bar{\gamma} \sim N^{1/2-\eta} \to \infty$ and $|\alpha|\bar{\gamma}^{-2} \sim N^{2\eta-2} \to 0$ resp. $|1 - \alpha|\bar{\gamma}^{-2} \sim N^{2\eta-3/2} \to 0$

as $N \to \infty$, that is a combined high-density and weak soft-core limit, indeed beneficial to the average-field approximation.

G. Lowest Landau level (LLL) anyons

If the anyons are ideal but in a strong external constant magnetic field, then all anyons fall into the LLL (cf. (66)), and the model actually becomes exactly solvable. The reason is that their wave function becomes (anti)analytic (modulo a Gaussian factor) and corresponds to changing the Jastrow factors in (48)-(49) from $|z_{jk}|^{-\alpha}$ to $(\bar{z}_{jk})^{\alpha}$, $\alpha \in [0, 1]$. It results in a description in terms of **fractional exclusion statistics** (cf. [Haldane 1991]) with one-body occupation numbers (angular momenta) shifted by α , and a relation to Calogero– Sutherland models. We refer to [Ouvry 2007] for an overview of this approach.

IV. THE NONABELIAN ANYON GAS

Let us now generalize our context further and consider an ideal *N*-anyon wave function as locally a map¹⁸

$$\Psi \colon \mathcal{C}^N \to \mathcal{F}, \quad \text{subject to} \quad \Psi(\sigma X) = \rho(\sigma)\Psi(X), \quad (74)$$

where the **fiber** \mathcal{F} is a Hilbert space of 'internal' (nonspatial) degrees of freedom on which B_N acts unitarily¹⁹:

$$\rho \colon B_N \to \mathrm{U}(\mathcal{F}).$$

Now $\rho(\sigma)$ is an **exchange operator** on \mathcal{F} corresponding to the braid $\sigma \in B_N$. There may be other observables of the considered system acting in \mathcal{F} , but for simplicity we here assume that it is finite dimensional, i.e. $\mathcal{F} = \mathbb{C}^D$ for some $D \in \mathbb{N}$, and $U(\mathbb{C}^D) = U(D)$ is the group of $D \times D$ unitary matrices. We then identify three possibilities for these representations ('reps'):

1. Irreducible abelian: $\mathcal{F} = \mathbb{C}$, which is the case we treated above, with elementary exchange phase

$$o(\sigma_j) = e^{i\alpha\pi} \quad \text{for all } j,$$

for a fixed statistics parameter α of abelian anyons.

2. Reducible abelian: $\mathcal{F} = \mathbb{C}^D$, D > 1, where all elementary exchange operators $\rho(\sigma_i)$ commute and thus are

¹⁸ Again, one way to concretize this is to regard the multivalued function Ψ as a ρ -equivariant function on the covering space of \mathcal{C}^N , or as a section of a vector bundle $E \to \mathcal{C}^N$ with fiber \mathcal{F} [Lundholm and Qvarfordt 2020].

¹⁹ We can even let go of the unitarity condition on reps if one is willing to consider multivalued probability distributions (cf. [Abramsky and Brandenburger 2011]).

simultaneously diagonalizable:²⁰

$$\rho(\sigma_j) \cong \operatorname{diag}(e^{i\alpha(1)\pi}, \dots, e^{i\alpha(D)\pi}) \quad \text{for all } j,$$

for statistics parameters $\alpha(k)$ corresponding to different sectors of particles that do not interact quantum statistically.

3. Nonabelian: $\mathcal{F} = \mathbb{C}^D$, D > 1, where not all elementary exchange operators commute:

$$\rho(\sigma_j)\rho(\sigma_k) \neq \rho(\sigma_k)\rho(\sigma_j)$$
 for some $j \neq k$.

Actually, it turns out that

$$D \ge N - 2 \qquad \text{if } N \ge 7, \tag{75}$$

because otherwise the rep is necessarily abelian (this is a theorem in the rep theory of the braid group [Formanek 1996]; cf. [Weinberger 2015], [Lundholm and Qvarfordt 2020, Theorem 3.13]). Furthermore, if $N \ge 7$ and D = N - 2 or D = N - 1 then the rep is either abelian or, up to an abelian factor, the nonabelian (reduced) Burau representation. For any other reps we need $D \ge N$.

Because of this last remark, we cannot simply fix the fiber Hilbert space \mathcal{F} or the rank of the representation $D = \dim \mathcal{F}$ and then study growing numbers N of anyons, because they will then eventually abelianize. Further, in the nonabelian case it is, as far as this author is aware, not certain whether statistics transmutation is available, i.e. whether one may always pass from the geometric to the magnetic perspective (this is a question in topology; see [Lundholm and Qvarfordt 2020, Maciazek and Sawicki 2019] for further discussion). Some models have magnetic formulations however [Kohno 1987, Lee and Oh 1994, Wen 1991], but for simplicity in this overview we will remain in the geometric perspective and furthermore pull in a small sample of results from the **al**gebraic perspective as well, where certain reps can be systematically and consistently constructed from elementary fusion and braiding rules, and exchange operators can be computed explicitly. The topic is quite complex and extends far outside the scope of this overview; instead we refer to Beer et al. 2018, Lahtinen and Pachos 2017, Lundholm and Qvarfordt 2020, Masaki et al. 2023, Nayak et al. 2008, Stern 2008] for more detail on the algebraic and diagrammatic aspects of anyon models (and note that the representation theory of the braid group is a still incomplete and very active field of mathematics). Anyons with nonabelian braid group representations are generally called "nonabelions" or "plektons".

A many-anyon model is a sequence of fiber Hilbert spaces \mathcal{F}_N , $D_N = \dim \mathcal{F}_N$, and braid group representations

$$\rho_N \colon B_N \to \mathrm{U}(\mathcal{F}_N), \qquad N = 1, 2, 3, \dots, \quad (76)$$

together with a sequence of kinetic energy operators

$$\hat{T}_{\rho_N} = \frac{\hbar^2}{2m} \sum_{j=1}^N (-i\nabla_{\mathbf{x}_j}^{(\rho_N)})^2$$
(77)

which implement these reps (by means of parallel transport) as the corresponding exchange operators in (74).²¹

The free kinetic energy operator \hat{T}_{ρ_N} is again locally the same as the free Laplacian \hat{T}_{dist} , supplemented with holonomies $\rho_N(\sigma) \in U(D_N)$ w.r.t. any nontrivial exchange loops $\sigma \in B_N$. Instead of the abelian pairexchange conditions (7), one now has

$$\Psi(\mathbf{x}_1, \dots, \mathbf{x}_k, \dots, \mathbf{x}_j, \dots, \mathbf{x}_N) = U_{N,p} \Psi(\mathbf{x}_1, \dots, \mathbf{x}_j, \dots, \mathbf{x}_k, \dots, \mathbf{x}_N), \quad (78)$$

with pair-exchange operators²²

$$U_{N,p} \cong \rho_N(\sigma_1 \sigma_2 \dots \sigma_p \sigma_{p+1} \sigma_p \dots \sigma_2 \sigma_1)$$
(79)

for any counterclockwise two-particle exchange loop in which $p \leq N-2$ other particles are enclosed, corresponding to the second figure in Fig. 3. We denote their eigenvalues by

spec
$$U_{N,p} = \{e^{i\pi\gamma_k}\}_{k=1,2,...,D_N}$$

where we normalize the full range of statistics parameters $\gamma_k = \gamma_k(N, p)$ in the interval (-1, 1].

Nonabelions are still harder to treat as perturbations close to bosons or fermions, since typically the eigenvalues of $U_{N,p}$ are separated from (or coincide with) ± 1 . Also, because of a lack of general magnetic formulations, it is even more unclear what we should mean with an extended gas of nonabelions, except as derived from emergent models and Berry connections (see below). However, concerning the qualitative features and local exclusion properties for the ideal gas, somewhat analogous results as for the abelian gas may be obtained [Lundholm and Qvarfordt 2020], as follows.

The *N*-fractionality is now replaced by

$$\alpha_{N,n} := \min_{p=0,1,\dots,n-2} \beta_{N,p}, \qquad n \le N,$$
(80)

where the **pair-exchange parameter** with $p \leq N - 2$ particles enclosed,

$$\beta_{N,p} := \min_{k=1,2,\dots,D_N} |\gamma_k(N,p)| = \operatorname{dist}(\operatorname{spec}(U_{N,p}), +1),$$

is again the arcwise distance (in units of π) from the spectrum of $U_{N,p}$ to +1. Then, the statistical repulsion as

 $^{^{20}}$ We write $A\cong B$ if two matrices or operators are similar, i.e. unitarily equivalent.

²¹ One might also require a compatibility between reps ρ_N w.r.t. the embedding sequence $B_{N-1} \hookrightarrow B_N$ of braid groups, which may be nontrivial (see [Delaney et al. 2016, Rowell and Wang 2012]).

²² Defined up to similarity; note e.g. that $\rho(\sigma_j) \cong \rho(\sigma_k)$ with similarities depending on j and k [Lundholm and Qvarfordt 2020].

given both by the Hardy inequality (34) (and its variants, depending on $\alpha_{N,n}$), the local exclusion principle (40)-(41), as well as the degeneracy pressure (43), generalize to the ideal nonabelion gas:

$$\langle \hat{T}_{\rho_N} \rangle_{\Psi} \gtrsim \alpha_{N,2} \int_{\mathbb{R}^2} \varrho_{\Psi}(\mathbf{x})^2 d\mathbf{x}.$$
 (81)

Thus, it is again the nearest-neighbor pair exchange (in the rep ρ_N) that sets the main behavior of the gas w.r.t. fractional exclusion statistics.

Three classes of representations of special interest were considered in some detail in [Lundholm and Qvarfordt 2020]: Burau, Ising and Fibonacci. Note that variations of these may arise by composing with abelian models. Some important features have been obtained also for other reps, such as scattering and dilute thermodynamics based on the nearest-neighbor pair exchange [Mancarella et al. 2013a,b, Verlinde 1991, Wilczek and Wu 1990].

A. Burau

The Burau rep [Burau 1935] is a simple deformation of the defining rep of the permutation group, $S_N \to U(N)$, that also must be reduced $(D_N = N - 1 \text{ or } N - 2)$ and unitarized to fit in the above framework. We are only aware of the simplest case N = 3, $\mathcal{F} = \mathbb{C}^2$ having been worked out explicitly so far [Weinberger 2015], and then

spec
$$U_{3,0} = \{1, -w^2\},$$
 spec $U_{3,1} = \{w^3, -w^3\}$

for a parameter $w = e^{i\alpha\pi}$, $|\alpha| < 1/3$, of the representation. Here we find $\alpha_{3,2}(w) = 0$ for all such w and indeed there are states that do not vanish on the diagonal because of a lack of exclusion there [Lundholm and Qvarfordt 2020].

B. Ising

Here $D_N = \dim \mathcal{F}_N$ grows roughly as $(\sqrt{2})^N$, which is related to reps of Clifford algebras [Nayak and Wilczek 1996] (and there are distinctions between even/odd numbers of such anyons and of fermions which also appear in the models). The spectra of all pair-exchange operators have been computed [Lundholm and Qvarfordt 2020]:

spec
$$U_{N,p} =$$

$$\begin{cases} \{e^{-i\pi/8}, e^{i\pi3/8}\}, & \text{if } p = 0, \\ \{e^{-i\pi/8}, e^{i\pi7/8}\}, & \text{if } p \ge 1 \text{ is odd}, \\ \{e^{-i\pi5/8}, e^{-i\pi/8}, e^{i\pi3/8}, e^{i\pi7/8}\}, & \text{if } p \ge 2 \text{ is even}, \end{cases}$$

(in one rep, and their complex conjugates in another rep, and with multiplicities that grow with N). Thus,

$$\alpha_{N,n} = \beta_{N,0} = 1/8 \qquad \text{for all } N, n \ge 2,$$

so the Ising gas has an exclusion at the diagonals and a degeneracy pressure. Since these n-fractionalities are

independent of n (and N), we do not see any benefits of clustering to minimize statistical repulsion, at this level of inquiry.

C. Fibonacci

Here $D_N = \dim \mathcal{F}_N$ grows with the Fibonacci sequence, so that $D_{N+1}/D_N \xrightarrow{N \to \infty} \phi$, the golden ratio. Furthermore, the spectra of all pair-exchange operators have been computed [Qvarfordt 2017]:

spec
$$U_{N,p} = \begin{cases} \{e^{-i\pi 3/5}, e^{i\pi 4/5}\}, & \text{if } p = 0, \\ \{e^{-i\pi/5}, e^{i\pi/5}, e^{i\pi 4/5}\}, & \text{if } p = 1, \\ \{e^{-i\pi 3/5}, e^{-i\pi/5}, e^{i\pi/5}, e^{i\pi 4/5}\}, & \text{if } p \ge 2, \end{cases}$$

(in one rep, and their complex conjugates in another rep, and with multiplicities that grow with N, also along the Fibonacci sequence). Thus,

$$\alpha_{N,n} = \begin{cases} \beta_{N,0} = 3/5, & \text{if } n = 2, \\ \beta_{N,1} = 1/5, & \text{if } n \ge 3, \end{cases}$$

and therefore also the Fibonacci gas is subject to exclusion at the diagonals and a degeneracy pressure. Since $\alpha_{N,n} < \alpha_{N,2}$ for $n \ge 3$, this may be indicative of clustering in this gas in order to minimize statistical repulsion over long ranges.

V. EMERGENCE OF THE ANYON GAS

In the study of emergent quantum statistics, we typically consider two distinct species of particle, one of which is numerous and numbered by e.g. $N \gg 1$, which we shall refer to as the **bath**, and another species which is much less numerous, $1 \leq n \ll N$, which we refer to as **tracers** or **impurities**. These two species could be physically the same type of particle but effectively distinguished by some quantum number, such as their Landau level index for example. The bath forms a background and a collective coherence for the tracers, so that its macroscopically well-defined state provides a limiting reference frame with respect to which tracers may transmute their properties.

A. The quantum Hall setting

In the quantum Hall (QH) setting we assume that all particles are confined to a plane and subject to a large and approximately constant magnetic field $\mathbf{B} = -b\mathbf{e}_z$, b > 0, i.e. the relevant one-body operator is again the Landau Hamiltonian (64)

$$\hat{H}_1^{\text{Lan}} = \frac{1}{2m} \left(-i\hbar \nabla_{\mathbf{x}} + qb\mathbf{x}^{\perp}/2 \right)^2 = \hbar\omega_c (\hat{a}^{\dagger}\hat{a} + 1/2),$$

 $\omega_c = |qb|/m$, and particles then distribute into the Landau levels (LLs) of this field. The lowest Landau level (LLL) is the kernel of \hat{a} (let us now fix $\hbar = 1$ and q = -1):

$$LLL := \mathcal{H}_0$$

= $\left\{ \psi \in L^2(\mathbb{R}^2) : \psi(\mathbf{x}) = f(z)e^{-\frac{b}{4}|z|^2}, f \text{ analytic} \right\}$
(82)

and spanned by the orthonormal basis of Fock–Darwin states

$$\psi_l(z) = c_l z^l e^{-\frac{b}{4}|z|^2}, \qquad l = 0, 1, 2, \dots,$$

for normalization constants $c_l > 0$. We assume electron spin is out of the picture due to the strong band separation, and also that there is a radial trapping potential centered at z = 0, penalizing higher angular momenta l.

The first concrete prediction of emergent anyonic statistics was made in the context of the *fractional* quantum Hall effect (FQHE) for electrons [Tsui et al. 1982], i.e. fractional filling of the LLL, for which Laughlin proposed the following trial states in the electrons' coordinates $z_j \in \mathbb{C}$, $z = (z_j)_{j=1,...,N}$, to explain the effect [Laughlin 1983, 1999]:²³

$$\Psi^{\rm Lau}_{\mu}({\bf z}) \propto \prod_{j < k} z^{\mu}_{jk} e^{-\frac{b}{4}|{\bf z}|^2}$$

For electrons, μ is an odd integer, so that $\Psi^{\text{Lau}}_{\mu} \in L^2_{\text{asym}}$. In the case that $\mu = 1$, suitable for an *integer* QH effect, the state corresponds to a Fermi sea filling of LLL with the least angular momentum: $\Psi^{\text{Lau}}_{\mu=1}(z)$ is proportional to

$$\bigwedge_{l=0}^{N-1} \psi_l(\mathbf{z}) \propto \det \begin{bmatrix} 1 & 1 & \dots & 1 \\ z_1 & z_2 & \dots & z_N \\ \vdots & \vdots & & \vdots \\ z_1^{N-1} & z_2^{N-1} & \dots & z_N^{N-1} \end{bmatrix} e^{-\frac{b}{4}|\mathbf{z}|^2},$$

by Vandermonde's formula for the determinant. For fractional filling $1/\mu$, and to reduce Coulomb interaction energy among electrons, Laughlin thus proposed taking the determinant above to a power $\mu > 1$. On top of such states, Laughlin also considered states of quasiholes pinned to positions $w_k \in \mathbb{C}$, $w = (w_k)_{k=1,\dots,n}$:

$$\Psi_{\mu}^{\rm qh}(\mathbf{w};\mathbf{z}) \propto \prod_{k=1}^{n} \prod_{j=1}^{N} (w_k - z_j) \Psi_{\mu}^{\rm Lau}(\mathbf{z})$$

(all states with appropriate normalization, and $\mathbf{y}_k \leftrightarrow w_k$, $\mathbf{x}_j \leftrightarrow z_j$ respective real and complex coordinates).

1. Berry phases

Halperin [Halperin 1984] suggested that if the quasiholes are treated as quantum particles then they ought to have fractional anyonic statistics, and Arovas, Schrieffer and Wilczek [Arovas et al. 1984] subsequently proposed a **Berry-phase approach** to determine their statistics. Namely, if we view Ψ_{μ}^{gh} as a map

$$\mathbb{C}^n \to L^2_{\mathrm{asym}}(\mathbb{R}^{2N}), \qquad \mathbf{w} \mapsto \Psi^{\mathrm{qh}}_{\mu}(\mathbf{w}; \cdot)$$

from the parameters w into the (fiber) Hilbert space of electrons, then Berry associates to such a geometry a canonical connection [Berry 1984, Simon 1983]:

$$\mathbf{A}_{j}(\mathbf{y}) := \langle -i\nabla_{\mathbf{y}_{j}} \rangle_{\Psi^{\mathrm{qh}}(\mathbf{y};\cdot)} \\ = -i \int_{\mathbb{C}^{N}} \overline{\Psi^{\mathrm{qh}}_{\mu}(\mathbf{y};\mathbf{z})} \nabla_{\mathbf{y}_{j}} \Psi^{\mathrm{qh}}_{\mu}(\mathbf{y};\mathbf{z}) \, d\mathbf{z}.$$
(83)

The holonomies of the connection w.r.t. loops on the parameters, $\sigma : [0, 1] \to \mathbb{C}^n$,

$$\rho(\sigma) = e^{i\Phi_{\sigma}}, \qquad \Phi_{\sigma} = \oint_{\sigma} \sum_{j=1}^{n} \tilde{\mathbf{A}}_{j} \cdot d\mathbf{y}_{j}, \qquad (84)$$

are then interpreted to be the exchange phases (1) (now including an "external" field of constant curvature) effectively associated to the motion of the quasiholes in the bath of electrons. Thus, under the adiabatic assumption that the electrons move much faster than the holes, one can take statistical averages over $\mathbf{z} \in \mathbb{C}^N$ and use an analogy with a classical charged plasma to compute $|\Psi^{qh}(\mathbf{y};\cdot)|^2$ and Φ_{σ} , and then indeed conclude that there is, on top of geometric holonomies corresponding to a constant field $B/\mu = -b/\mu$ (i.e. effectively coupled to a fractional charge $+1/\mu$ that *reduces* the field), also topological exchange phases corresponding to the localized (but somewhat extended) attachment of fractional flux $-2\pi/\mu$ to the quasiholes. For a relatively recent review of the approach, including possible extensions to nonabelian scenarios (for which the quasihole map Ψ^{qh} is multidimensional), see [Bonderson et al. 2011].

2. Transmuted Hamiltonian

Forte [Forte 1991] argued that the above approach is not necessarily sufficient to determine the anyonic statistics because, as we have noted above with the freedom of statistics transmutation, the knowledge of the specific Hamiltonian for the quasiholes is essential, since that is what ultimately sets the scale of validity of any adiabatic treatment. Thus, a different approach was suggested in [Lambert et al. 2023, Lundholm and Rougerie 2016] to remedy this situation and supply the missing dynamical information, by considering quasiholes pinned to already present quantum particles.

In brief, one may consider the following idealized situation (see Fig. 12):

²³ Note that these also coincide with the anyonic trial states (48)-(49) if extended to $\alpha = \mu$ with trivial Jastrow factors [Lundholm 2017].



FIG. 12. Statistics transmutation in the QH setting. Bath particles (red) are repelled from tracers (black) by short-range interactions (green), and all particles move in an external homogeneous magnetic field (blue arrows). Subtracting the almost-homogeneous effective field generated by the bath as $b \sim N \gg 1$ then leaves a reduced/zero field plus remnant field (blue disks) at the holes punched in the bath by the tracers.

- 1. large 2D bath of noninteracting fermions, $N \gg 1$,
- 2. *n* tracers/impurities (initially can be either bosons or fermions), $1 \le n \ll N$,
- 3. strong external transverse magnetic field, $b \to \infty$,
- 4. strong short-range repulsive bath-tracer interaction, at coupling $g \to \infty$.

Taking the bath particles \mathbf{x}_j to have unit mass and unit negative charge, the tracers \mathbf{y}_k to have mass m and charge -q, and allowing for interaction and trapping of tracers in a joint potential W, the Hamiltonian of the joint system is:

$$H_{n\oplus N} := \frac{1}{2} \sum_{j=1}^{N} \left(-i\nabla_{\mathbf{x}_j} - b\mathbf{x}_j^{\perp}/2 \right)^2$$

+
$$\frac{1}{2m} \sum_{k=1}^{n} \left(-i\nabla_{\mathbf{y}_k} - qb\mathbf{y}_k^{\perp}/2 \right)^2$$

+
$$g \sum_{j=1}^{N} \sum_{k=1}^{n} \delta(\mathbf{x}_j - \mathbf{y}_k) + W(\mathbf{y}_1, \dots, \mathbf{y}_n),$$

acting on states

$$\Psi(\mathbf{y}_1,\ldots,\mathbf{y}_n;\mathbf{x}_1,\ldots,\mathbf{x}_N)\in L^2(\mathbb{R}^{2n})\otimes L^2_{\mathrm{asym}}(\mathbb{R}^{2N}).$$

If one takes $b \sim N$ and $N \gg n$, and assumes the bath then sits entirely in the LLL, thus forming a disk-shaped QH droplet of size ~ 1 , the Hilbert space is reduced to

$$\mathcal{H}^{n\oplus N}_{\mathrm{sym/asym}} := L^2_{\mathrm{sym/asym}}(\mathbb{R}^{2n}) \otimes \bigwedge^N LLL$$

(allowing for either bosonic or fermionic tracers), with g.s.e.

$$E_{n\oplus N} := \inf_{0 \neq \Psi \in \mathcal{H}^{n\oplus N}_{\text{sym/asym}}} \langle H_{n\oplus N} \rangle_{\Psi}.$$

For $g \to \infty$, we then take as our ansatz for the ground state of the system a state in the kernel of the interaction:

$$\Psi_{\Phi}(\mathbf{y};\mathbf{x}) := \Phi(\mathbf{w})c^{\mathbf{qh}}(\mathbf{w})\Psi^{\mathbf{qh}}(\mathbf{w};\mathbf{z}), \tag{85}$$

with $\Phi \in L^2_{\text{sym/asym}}(\mathbb{C}^n)$, $c^{\text{qh}} > 0$ and

$$\int_{\mathbb{C}^n} |\Phi(\mathbf{w})|^2 \, d\mathbf{w} = 1, \ c^{qh}(\mathbf{w})^{-2} := \int_{\mathbb{C}^N} |\Psi^{qh}(\mathbf{w}; \mathbf{z})|^2 d\mathbf{z},$$

ensuring the correct normalization $\int_{\mathbb{C}^{n+N}} |\Psi_{\Phi}|^2 dw dz = 1.$

The bath in LLL forces an *analytic* vanishing as $z_j \rightarrow w_k$, i.e. $\mathbf{x}_j \rightarrow \mathbf{y}_k$. It suggests to take $\Psi^{\text{qh}} = \Psi^{\text{qh}}_{\mu}$ as above, corresponding to fractional filling $1/\mu$ of the LLL, or

$$\Psi_{\mu,p}^{\rm qh}(\mathbf{w};\mathbf{z}) := \prod_{k=1}^{n} \prod_{j=1}^{N} (w_k - z_j)^p \, \Psi_{\mu}^{\rm Lau}(\mathbf{z})$$

thus also allowing for non-simple quasiholes if p > 1.

It was conjectured in [Lambert et al. 2023, Lundholm and Rougerie 2016] that in this ansatz

$$\langle H_{n\oplus N} \rangle_{\Psi_{\Phi}} = \frac{bN}{2} + \langle H_n^{\text{eff}} \rangle_{\Phi} + \operatorname{error}_{\Phi}(n/N), \quad (86)$$

for an effective **statistics-transmuted Hamiltonian** for the tracers:

$$\begin{split} H_n^{\text{eff}} &:= \frac{bn}{2m} \frac{p}{\mu} + \frac{1}{2m} \sum_{j=1}^n \Bigl(-i \nabla_{\mathbf{y}_j} - \bigl(q - \frac{p}{\mu}\bigr) \frac{b}{2} \mathbf{y}_j^\perp - \frac{p^2}{\mu} \mathbf{A}_j \Bigr)^2 \\ &+ W(\mathbf{y}), \end{split}$$

thus implying an effective renormalized charge $-q + p/\mu$ and shifted statistics parameter by $-p^2/\mu$.

In [Lambert et al. 2023] it was proved mathematically that (86) is indeed a correct conclusion, with an error term of order o(N) at fixed n and reasonable Φ , if $\mu = 1$ and p = 1, corresponding to transmutation of bosons into fermions and vice versa (cf. Sec. II D). Many crucial simplifications occur for these specific parameters, having to do with the Vandermonde structure of the state, and in the proof it is also required that tracers have sufficient extra repulsion to not cluster too strongly at the diagonal set Δ_n , which is ensured either by them being initially bosons and then eventually (free, if q = 1) fermions, or that W includes an additional and sufficiently strong intra-species repulsion.

B. Impurities in the plane and polarons

Another route to statistics transmutation is to place tracers/impurities in a bath of bosons, such as the



FIG. 13. Statistics transmutation in the polaron setting. An initial ladder of even-integer fluxes/vortices (composite bosons or fermions) is in the adiabatic limit transmuted into a ladder of fractional fluxes/vortices (anyons).

phonon vibrational modes of a crystal lattice, or photonic optical modes. Such systems are well known to enable transmutation of the impurities into **polarons**, i.e. bound states of impurity particles and coherent states of phonons/photons [Fröhlich 1954, Landau 1933, Pekar 1946]. In [Yakaboylu and Lemeshko 2018, Yakaboylu et al. 2020] it was observed that, subject to suitable external fields and rotation of such systems, the polarons can also transmute their statistics.

As an illustrative toy model, we consider a single bosonic collective degree of freedom:

$$[\hat{a}, \hat{a}^{\dagger}] = 1, \quad \hat{N} := \hat{a}^{\dagger} \hat{a}, \text{ with eigenstates } |N\rangle, N \ge 0,$$

and an initial n-particle Hamiltonian

$$\hat{H}_0 = \sum_{j=1}^n (-i\nabla_{\mathbf{x}_j})^2 + W(\mathbf{x})$$

for bosons or fermions on $\mathcal{H}_0 = L^2_{\text{sym/asym}}(\mathbb{R}^{2n})$. Let us then define the interaction Hamiltonian

$$\hat{H}_{\omega,\gamma} := \hat{H}_0 + \omega \hat{a}^{\dagger} \hat{a} + \gamma \omega (F \hat{a}^{\dagger} + F^{-1} \hat{a}) + \gamma^2 \omega, \quad (87)$$

with two parameters $\omega > 0$ and $\gamma \in \mathbb{R}$. The interpretation of the terms added to \hat{H}_0 is that ω is the energy associated to an attachment or detachment of two units of flux or vorticity to each of the particles, which is done by the respective hopping terms with coupling $\gamma \omega$ and such that γ defines the ratio between these two effects. The last term is only a constant added for convenience.

We now denote $Z = \prod_{j < k} z_{jk}$, and consider two possible choices for F (or their complex conjugates, depending on which way the orientation symmetry is to be broken):

1. $F = (Z/|Z|)^2 = U^2$ corresponding to flux attachment,

2. $F = Z^2$ corresponding to vortex attachment.

By acting with Fa^{\dagger} on the reference system $\mathcal{H}^{0} = \mathcal{H}_{0}|0\rangle$ one obtains a ladder of Hilbert spaces of **composite** bosons/fermions (see Fig. 13):

$$\mathcal{H}^N := F^N \mathcal{H}_0 |N\rangle, \qquad N = 0, 1, 2, \dots$$

Upon taking the "adiabatic limit" $\omega \to \infty$ at fixed γ , we claim that in the bottom of the spectrum of $\hat{H}_{\omega,\gamma}$ we then obtain a ladder of transmuted anyons with statistics parameter $\alpha = 2\gamma^2 + 2N$ [Yakaboylu et al. 2020]. These anyons are either *interacting* by means of a scalar potential $\sum_j \mathbf{A}_j^2$ (for the more realistic, unitary choice $F = U^2$), or free (for the more algebraic, non-unitary choice $F = Z^2$, yielding a non-hermitian Hamiltonian with $i\mathbf{A}_j = \nabla_{\mathbf{x}_j} \log Z$). Namely, the transformations

$$\hat{S} = F^{\hat{N}} = e^{\hat{N}\log F}, \qquad \hat{U} = e^{-\gamma(\hat{a}^{\dagger} - \hat{a})}, \qquad (88)$$

which diagonalize $\hat{H}_{\omega,\gamma}$,

$$\hat{U}^{-1}\hat{S}^{-1}\hat{H}_{\omega,\gamma}\hat{S}\hat{U}$$

= $\omega\hat{a}^{\dagger}\hat{a} + \sum_{j=1}^{N} \left[-i\nabla_{\mathbf{x}_{j}} + 2\mathbf{A}_{j}(\hat{a}^{\dagger} - \gamma)(\hat{a} - \gamma)\right]^{2} + W(\mathbf{x}),$

create coherent-state mixtures

$$\hat{S}\hat{U}|0\rangle = e^{-\gamma^2/2} \sum_{N=0}^{\infty} \frac{(-\gamma)^N}{\sqrt{N!}} F^N|N\rangle$$
(89)

over the initial ladder of integer composites and enable a shift of the expected amount of flux/vorticity by $\alpha_0 = 2\gamma^2 \ge 0$, possibly to non-integer values. This transmutation procedure has indeed been used (with the choice $F = Z^2$) to compute the numerical spectra seen in Fig. 4.

A possible plane polaron model with the above form is obtained upon applying a constant magnetic field **B** and shifting to a counter-rotating frame at half the cyclotron frequency $\Omega = B/(2m)$ (cf. Fig. 14, upper):

$$\begin{split} \hat{H}_N &= \frac{1}{2m} \sum_{j=1}^n \hat{\mathbf{p}}_j^2 + W^{\Omega}(\mathbf{x}) + \sum_{\mathbf{k}} \omega_{\mathbf{k}}^{\Omega} \hat{b}_{\mathbf{k}}^{\dagger} \hat{b}_{\mathbf{k}} \\ &+ \sum_{\mathbf{k}} \lambda_{\mathbf{k}}(\mathbf{x}) \left(e^{-i\beta_{\mathbf{k}}(\mathbf{x})} b_{\mathbf{k}}^{\dagger} + e^{i\beta_{\mathbf{k}}(\mathbf{x})} b_{\mathbf{k}} \right), \end{split}$$

for phonon vibration modes $b_{\mathbf{k}}, \mathbf{k} \in \mathbb{R}^2$ with dispersion $\omega_{\mathbf{k}}^{\Omega}$ (which is their bare dispersion shifted by their angular momentum coupled to Ω), and phonon-impurity



FIG. 14. Two possible experimental probes for emergent statistics transmutation: rotating plane in which Fröhlich polarons turn into anyons (upper, from [Yakaboylu et al. 2020]) and the exchange of linear molecules (lower, from [Brooks et al. 2021a]).

interaction parameters $\lambda_{\mathbf{k}}$, $\beta_{\mathbf{k}}$. For n = 2 and a suitable interaction, realizable within the Fröhlich plane polaron model with quasi-2D Coulomb forces [Yakaboylu et al. 2020], the emergent gauge field

$$\alpha \mathbf{A}_{j}(\mathbf{x}) \approx \langle -i\nabla_{\mathbf{x}_{j}} \rangle_{\hat{S}\hat{U}|0\rangle} \approx -\sum_{\mathbf{k}} (\lambda_{\mathbf{k}}(\mathbf{x})/\omega_{\mathbf{k}}^{\Omega})^{2} \nabla_{\mathbf{x}_{j}} \beta_{\mathbf{k}}(\mathbf{x})$$
⁽⁹⁰⁾

is then approximately of the correct Aharonov-Bohm type and such as to transmute the statistics by $\alpha \approx \alpha(\Omega)$, ascertained by the system's correlations to the collective reference rotation/field.

C. Impurities on the sphere and angulons

A similar approach [Brooks et al. 2021a,b] can be taken for particles moving effectively on a sphere S^2 , namely one may e.g. replace the planar kinetic energy (4) by the rotation energy

$$T_{\rm rot} = \sum_{j=1}^{n} \mathbf{L}_j^2 \tag{91}$$

(with units normalized suitably) of n identical linear molecules (cf. Fig. 14, lower). The resulting model for the Hamiltonian coupled to a bath of rotational angular momenta is called an **angulon** model [Schmidt and Lemeshko 2015]. This transmutation method has also been used to compute the energy spectrum of two anyons on a sphere [Brooks et al. 2021a]; cf. [Ouvry and Polychronakos 2019, Polychronakos and Ouvry 2020], and see [Einarsson 1992] concerning anyons in various topologies.

D. Other emergent models for anyons

Other approaches to investigate statistics transmutation in the QH setting include effects of anyonic shifted angular momentum on the energy spectrum [Cooper and Simon 2015], and on pair correlation and other aspects of the density [Dubček et al. 2018, Graß et al. 2020, Morampudi et al. 2017, Umucahlar et al. 2018, Zhang et al. 2014], including nonabelian scenarios [Baldelli et al. 2021, Zhang et al. 2015]. Emergent anyons can also be expected on spin networks and lattices [Bachmann 2017, Bachmann et al. 2020, 2023]. Certain classes of nonabelions have been proposed to emerge in the physics of neutrons and neutron stars [Masaki et al. 2023]. Even in the quantum gravity context might we expect nonabelions, as they can capture the relevant degrees of freedom on the black-hole event horizon [Pithis and Ruiz Euler 2015].

VI. CONCLUSIONS AND OUTLOOK

In light of recent experiments concerning anyons [Bartolomei et al. 2020, Fan et al. 2022, Google Quantum AI and Collaborators 2022, Nakamura et al. 2020], it is increasingly important to gain a firm theoretical understanding of the basic properties of the anyon gas. We have here focused on its precise definition in typical ideal and nonideal contexts, its most basic physics as regards the connection between exchange and exclusion statistics, as well as its inevitable emergence in a few conceptually simple scenarios involving relatively well-understood bosonic and fermionic systems with manifest breaking of orientation symmetry, such as the rotated polaron and the quantum Hall setup.

In preparation for future experiments and theoretical research, one could make a few observations and highlight some potential obstacles. First, it is hoped that accurate density functional theories for large systems of anyons will provide more robust, detectable signatures, such as spatial and momentum density profiles, as compared to the typically extremely fragile phases probed by interferometry (cf. [Camino et al. 2005, Nakamura et al. 2020]). Second, the development of impurity models and the pinning of quasiholes, fluxes and vortices to already well-understood quantum particles promises to eliminate some of the ambiguities inherent in conventional Berry phase and adiabatic treatments (cf. [Forte 1991, Jain 2007, Lambert et al. 2023, Lundholm and Rougerie 2016, Myrheim 1999). Third, with the steadily growing literature on anyons and an increasing interest in their fascinating physics, even from the general public, it is important to be aware of potential confusion on how the term

"anyon" is used. Hopefully, this overview has helped to clarify both the nontrivial relationship between exchange and exclusion statistics, as well as the crucial differences between: a) the **choice or construction** of a concrete braid group representation $\rho_N \colon B_N \to U(\mathcal{F}_N)$ (algebra, kinematics, or simply an action of a permutation on a system); b) its **practical realization** such as by means of motion of classical parameters (magnetics, or computation using a classical or quantum computer; cf. [Noh et al. 2020]); and c) **actual anyons** (or bosons or fermions) endowed with quantum kinetic energy \hat{T}_{ρ_N} (geometrodynamics), uncertainty and exclusion.

Acknowledgments. The study of the magnetic TF functional for anyons was done in collaboration with Dinh Thi Nguyen, following helpful discussions with Nicolas Rougerie and Théotime Girardot. Further, I thank the editor Tapash Chakraborty for the timely invitation to write this overview, as well as Vincent Cavalier, Ask Ellingsen, Gerald Goldin and Wolfgang Staubach for useful comments on the manuscript. I am thankful to numerous other colleagues for insightful discussions on this topic over the last 13 years, in particular Jan Philip Solovej, who raised my interest in it, as well as Eddy Ardonne, Morris Brooks, Michele Correggi, Romain Dubosco, Luca Fresta, Jürg Fröhlich, Thors Hans Hansson, Gaultier Lambert, Simon Larson, Jon Magne Leinaas, Tomasz Maciazek, Per Moosavi, Luca Oddis, Stéphane Ouvry, Viktor Qvarfordt, Robert Seiringer, Andrea Trombettoni, Susanne Viefers, and Enderalp Yakaboylu. The research project was funded by the Swedish Research Council (grant no. 2021-05328, "Mathematics of anyons and intermediate quantum statistics").

S. Abramsky and A. Brandenburger, *The sheaf-theoretic structure of non-locality and contextuality*, New Journal of Physics **13** (2011), no. 11, 113036, doi.

A. Aftalion, X. Blanc, and F. Nier, *Vortex distribution in the lowest Landau level*, Phys. Rev. A **73** (2006), 011601(R), doi.

A. Aftalion, X. Blanc, and F. Nier, Lowest Landau level functional and Bargmann spaces for Bose-Einstein condensates,
J. Funct. Anal. 241 (2006), no. 2, 661–702, doi.

Y. Aharonov and D. Bohm, Significance of electromagnetic potentials in the quantum theory, Phys. Rev. **115** (1959), 485–491, doi.

S. Albeverio, F. Gesztesy, R. Høegh-Krohn, and H. Holden, *Solvable models in quantum mechanics*, second ed., AMS Chelsea Publishing, Providence, RI, 2005, With an appendix by Pavel Exner.

A. Alonso and B. Simon, *The Birman-Krein-Vishik theory of* self-adjoint extensions of semibounded operators, Journal of Operator Theory (1980), 251–270.

D. Arovas, J. R. Schrieffer, and F. Wilczek, *Fractional statistics and the quantum Hall effect*, Phys. Rev. Lett. **53** (1984), 722–723, doi.

D. P. Arovas, R. Schrieffer, F. Wilczek, and A. Zee, *Statistical mechanics of anyons*, Nuclear Physics B **251** (1985), 117 – 126, doi.

E. Artin, *Theory of braids*, Ann. of Math. (2) **48** (1947), 101–126, doi.

E. Artin, *Theorie der Zöpfe*, Abh. Math. Sem. Univ. Hamburg 4 (1925), no. 1, 47–72, doi.

S. Bachmann, Local disorder, topological ground state degeneracy and entanglement entropy, and discrete anyons, Rev. Math. Phys. **29** (2017), no. 06, 1750018, doi.

S. Bachmann, A. Bols, W. De Roeck, and M. Fraas, *Manybody Fredholm index for ground-state spaces and abelian anyons*, Phys. Rev. B **101** (2020), 085138, doi.

S. Bachmann, B. Nachtergaele, and S. Vadnerkar, *Dynamical abelian anyons with bound states and scattering states*, arXiv e-prints, 2023, arXiv:2303.07379.

N. Baldelli, B. Juliá-Díaz, U. Bhattacharya, M. Lewenstein, and T. Graß, *Tracing non-abelian anyons via impurity parti-* cles, Phys. Rev. B 104 (2021), 035133, doi.

H. Bartolomei, M. Kumar, R. Bisognin, A. Marguerite, J.-M. Berroir, E. Bocquillon, B. Plaçais, A. Cavanna, Q. Dong, U. Gennser, Y. Jin, and G. Fève, *Fractional statistics in anyon collisions*, Science **368** (2020), no. 6487, 173–177, doi.
K. Beer, D. Bondarenko, A. Hahn, M. Kalabakov, N. Knust,

L. Niermann, T. J. Osborne, C. Schridde, S. Seckmeyer, D. E. Stiegemann, et al., *From categories to anyons: a travelogue*, arXiv e-prints, 2018, arXiv:1811.06670.

M. V. Berry, *Quantal phase factors accompanying adiabatic changes*, Proc. Roy. Soc. London, Ser. A **392** (1984), 45–57, doi.

L. Biedenharn, E. Lieb, B. Simon, and F. Wilczek, *The ancestry of the 'Anyon'*, Physics Today **43** (1990), no. 8, 90, doi.

J. S. Birman, *Braids, links, and mapping class groups*, Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1974, Annals of Mathematics Studies, No. 82.

P. Bonderson, V. Gurarie, and C. Nayak, *Plasma analogy* and non-abelian statistics for Ising-type quantum Hall states, Phys. Rev. B 83 (2011), 075303, doi.

M. Bourdeau and R. D. Sorkin, When can identical particles collide?, Phys. Rev. D 45 (1992), 687–696, doi.

M. Brooks, M. Lemeshko, D. Lundholm, and E. Yakaboylu, Molecular impurities as a realization of anyons on the twosphere, Phys. Rev. Lett. **126** (2021), 015301, doi.

M. Brooks, M. Lemeshko, D. Lundholm, and E. Yakaboylu, Emergence of anyons on the two-sphere in molecular impurities, Atoms 9 (2021), no. 4, 106, doi.

W. Burau, Über Zopfgruppen und gleichsinnig verdrillte Verkettungen, Abh. Math. Sem. Univ. Hamburg **11** (1935), no. 1, 179–186, doi.

D. Caenepeel and R. MacKenzie, *Parity violation, anyon scattering, and the mean field approximation*, Phys. Rev. D **50** (1994), 5418–5424, doi.

F. Calogero, Ground state of a one-dimensional N-body system, J. Math. Phys. **10** (1969), no. 12, 2197–2200, doi.

F. Calogero, Solution of a three-body problem in one dimension, J. Math. Phys. **10** (1969), no. 12, 2191–2196, doi.

F. E. Camino, W. Zhou, and V. J. Goldman, Realization of a Laughlin quasiparticle interferometer: Observation of fractional statistics, Phys. Rev. B **72** (2005), 075342, doi.

G. S. Canright and M. D. Johnson, *Fractional statistics: alpha to beta*, J. Phys. A: Math. Gen. **27** (1994), no. 11, 3579, doi.

A. Cappelli, L. S. Georgiev, and I. T. Todorov, *Parafermion Hall states from coset projections of abelian conformal theories*, Nucl. Phys. B **599** (2001), no. 3, 499–530, doi.

Y. H. Chen, F. Wilczek, E. Witten, and B. I. Halperin, *On anyon superconductivity*, Int. J. Mod. Phys. B **03** (1989), 1001–1067, doi.

R. Chitra and D. Sen, Ground state of many anyons in a harmonic potential, Phys. Rev. B 46 (1992), 10923-10930, doi.

C. Chou, Multianyon spectra and wave functions, Phys. Rev. D 44 (1991), 2533-2547, doi.

N. R. Cooper and S. H. Simon, Signatures of fractional exclusion statistics in the spectroscopy of quantum Hall droplets, Phys. Rev. Lett. **114** (2015), 106802, doi.

M. Correggi and D. Fermi, *Magnetic perturbations of anyonic* and Aharonov-Bohm Schrödinger operators, J. Math. Phys. **62** (2021), 032101, doi.

M. Correggi and L. Oddis, *Hamiltonians for twoanyon systems*, Rend. Mat. Appl. **39** (2018), 277-292, http://www1.mat.uniroma1.it/ricerca/rendiconti/ 39_2_(2018)_277-292.html.

M. Correggi, D. Lundholm, and N. Rougerie, *Local density* approximation for the almost-bosonic anyon gas, Analysis & PDE **10** (2017), 1169–1200, doi.

M. Correggi, D. Lundholm, and N. Rougerie, *Local density approximation for almost-bosonic anyons*, Proceedings of QMath13, Atlanta, October 8–11, 2016, Mathematical problems in quantum physics (F. Bonetto, D. Borthwick, E. Harrell, and M. Loss, eds.), Contemp. Math., vol. 717, 2018, pp. 77–92, doi.

M. Correggi, R. Duboscq, D. Lundholm, and N. Rougerie, *Vortex patterns in the almost-bosonic anyon gas*, EPL (Europhysics Letters) **126** (2019), 20005, doi.

F. A. B. Coutinho, Y. Nogami, and J. Fernando Perez, *Self-adjoint extensions of the Hamiltonian for a charged particle in the presence of a thread of magnetic flux*, Phys. Rev. A **46** (1992), 6052–6055, doi.

G. Date and M. V. N. Murthy, *Classical dynamics of anyons and the quantum spectrum*, Phys. Rev. A **48** (1993), 105–110, doi.

G. Date, M. V. N. Murthy, and R. Vathsan, *Classical and quantum mechanics of anyons*, arXiv e-prints, 2003, arXiv: cond-mat/0302019.

C. Delaney, E. C. Rowell, and Z. Wang, *Local unitary repre*sentations of the braid group and their applications to quantum computing, Rev. Colombiana Mat. **50** (2016), no. 2, 207– 272, doi.

G. Dell'Antonio, R. Figari, and A. Teta, *Statistics in space dimension two*, Lett. Math. Phys. **40** (1997), no. 3, 235–256, doi.

P. A. M. Dirac, *Lectures on quantum mechanics*, Belfer Graduate School of Science Monographs Series, vol. 2, Belfer Graduate School of Science, New York; produced and distributed by Academic Press, Inc., New York, 1967, Second printing of the 1964 original. J. S. Dowker, Quantum mechanics and field theory on multiply connected and on homogeneous spaces, J. Phys. A 5 (1972), no. 7, 936, doi.

J. S. Dowker, *Remarks on nonstandard statistics*, J. Phys. A **18** (1985), no. 18, 3521–3530, doi.

T. Dubček, B. Klajn, R. Pezer, H. Buljan, and D. Jukić, *Quasimomentum distribution and expansion of an anyonic gas*, Phys. Rev. A **97** (2018), 011601, doi.

F. J. Dyson, Ground-state energy of a finite system of charged particles, J. Math. Phys. 8 (1967), no. 8, 1538–1545, doi.

F. J. Dyson and A. Lenard, *Stability of matter. I*, J. Math. Phys. 8 (1967), no. 3, 423–434, doi.

W. Ehrenberg and R. E. Siday, *The refractive index in electron optics and the principles of dynamics*, Proceedings of the Physical Society. Section B **62** (1949), no. 1, 8.

T. Einarsson, Anyons and antiferromagnets: Two twodimensional topics, Ph.D. thesis, Institute of Theoretical Physics, Chalmers University of Technology, Gothenburg, 1992, https://research.chalmers.se/en/publication/ 176319.

Y.-a. Fan, Y. Li, Y. Hu, Y. Li, X. Long, H. Liu, X. Yang, X. Nie, J. Li, T. Xin, et al., *Experimental realization of a topologically protected Hadamard gate via braiding Fibonacci anyons*, arXiv e-prints, 2022, arXiv:2210.12145.

E. Fermi, Un metodo statistico per la determinazione di alcune priorieta dell'atome, Rend. Accad. Naz. Lincei **6** (1927), 602–607.

A. L. Fetter, C. B. Hanna, and R. B. Laughlin, *Random-phase approximation in the fractional-statistics gas*, Phys. Rev. B **39** (1989), 9679–9681, doi.

E. Formanek, Braid group representations of low degree, Proc. London Math. Soc. (3) **73** (1996), no. 2, 279–322, doi.

S. Forte, Berry's phase, fractional statistics and the Laughlin wave function, Mod. Phys. Lett. A **6** (1991), no. 34, 3152–3162, doi.

S. Forte, Quantum mechanics and field theory with fractional spin and statistics, Rev. Mod. Phys. **64** (1992), 193–236, doi.

K. Fredenhagen, K.-H. Rehren, and B. Schroer, Superselection sectors with braid group statistics and exchange algebras. I. General theory, Comm. Math. Phys. **125** (1989), no. 2, 201–226, http://projecteuclid.org/euclid.cmp/1104179464.

M. H. Freedman, A. Kitaev, M. J. Larsen, and Z. Wang, *Topological quantum computation*, Bull. Amer. Math. Soc. (N.S.) **40** (2003), no. 1, 31–38, Mathematical challenges of the 21st century (Los Angeles, CA, 2000), doi.

H. Fröhlich, *Electrons in lattice fields*, Advances in Physics **3** (1954), 325–361, doi.

J. Fröhlich and F. Gabbiani, *Braid statistics in local quantum theory*, Rev. Math. Phys. **2** (1990), no. 3, 251–353, doi.

J. Fröhlich and P. A. Marchetti, *Quantum field theory of anyons*, Lett. Math. Phys. **16** (1988), no. 4, 347–358, doi.

J. Fröhlich and P. A. Marchetti, *Quantum field theories of vortices and anyons*, Commun. Math. Phys. **121** (1989), no. 2, 177–223, doi.

J. Fröhlich, New super-selection sectors ("soliton-states") in two dimensional Bose quantum field models, Commun. Math. Phys. 47 (1976), no. 3, 269–310.

J. Fröhlich, Statistics of fields, the Yang-Baxter equation, and the theory of knots and links, Nonperturbative quantum field theory (Cargèse, 1987), NATO Adv. Sci. Inst. Ser. B Phys., vol. 185, Plenum, New York, 1988, pp. 71-100.

J. Fröhlich, *Quantum statistics and locality*, Proceedings of the Gibbs Symposium (New Haven, CT, 1989), Amer. Math. Soc., Providence, RI, 1990, pp. 89–142.

J. Fröhlich, Spin, or actually: Spin and quantum statistics, The Spin: Poincaré Seminar 2007 (B. Duplantier, J.-M. Raimond, and V. Rivasseau, eds.), Birkhäuser Basel, Basel, 2009, pp. 1–60, doi.

J. Fröhlich and T. Kerler, *Quantum groups, quantum cate*gories and quantum field theory, Berlin-Heidelberg-New York: Springer-Verlag, 1993, doi.

G. Gentile, Osservazioni sopra le statistiche intermedie, Il Nuovo Cimento **17** (1940), no. 10, 493–497, doi.

G. Gentile, Le statistiche intermedie e le proprietà dell'elio liquido, Il Nuovo Cimento **19** (1942), no. 4, 109–125, doi.

M. Girardeau, Relationship between systems of impenetrable bosons and fermions in one dimension, J. Math. Phys. 1 (1960), 516–523, doi.

M. D. Girardeau, Permutation symmetry of many-particle wave functions, Phys. Rev. **139** (1965), B500–B508, doi.

T. Girardot, Average field approximation for almost bosonic anyons in a magnetic field, J. Math. Phys. **61** (2020), no. 7, 071901, 23, doi.

T. Girardot, Mean-field approximation for the anyon gas, Ph.D. thesis, Université Grenoble Alpes & CNRS, l'École Doctorale Mathématiques, Sciences et technologies de l'information, Informatique, 2021, https://www.theses.fr/ 2021GRALM031.

T. Girardot and N. Rougerie, *Semiclassical limit for almost fermionic anyons*, Commun. Math. Phys. **387** (2021), no. 1, 427–480, doi.

T. Girardot and N. Rougerie, A Lieb-Thirring inequality for extended anyons, arXiv e-prints, 2022, arXiv:2209.02543.

G. A. Goldin, R. Menikoff, and D. H. Sharp, *Representations of a local current algebra in nonsimply connected space and the Aharonov-Bohm effect*, J. Math. Phys. **22** (1981), no. 8, 1664–1668, doi.

G. Goldin, R. Menikoff, and D. Sharp, *Particle statistics from induced representations of a local current group*, J. Math. Phys. **21** (1980), no. 4, 650–664, doi.

G. A. Goldin, The prediction of anyons: Its history and wider implications, arXiv e-prints, 2022, arXiv:2212.12632.

G. A. Goldin and S. Majid, On the Fock space for nonrelativistic anyon fields and braided tensor products, J. Math. Phys. **45** (2004), no. 10, 3770–3787, doi.

G. A. Goldin and D. H. Sharp, *Diffeomorphism groups, anyon fields, and q commutators*, Phys. Rev. Lett. **76** (1996), 1183–1187, doi.

G. A. Goldin, R. Menikoff, and D. H. Sharp, *Diffeomorphism* groups, gauge groups, and quantum theory, Phys. Rev. Lett. **51** (1983), 2246–2249, doi.

G. A. Goldin, R. Menikoff, and D. H. Sharp, *Comments* on "General Theory for Quantum Statistics in Two Dimensions", Phys. Rev. Lett. **54** (1985), 603–603, doi.

Google Quantum AI and Collaborators, Observation of nonabelian exchange statistics on a superconducting processor, arXiv e-prints, 2022, arXiv:2210.10255.

T. Graß, B. Juliá-Díaz, N. Baldelli, U. Bhattacharya, and M. Lewenstein, *Fractional angular momentum and anyon statistics of impurities in Laughlin liquids*, Phys. Rev. Lett. 125 (2020), 136801, doi.

H. S. Green, A generalized method of field quantization, Phys. Rev. **90** (1953), 270–273, doi.

E. Gross, Structure of a quantized vortex in boson systems, Nuovo Cimento **20** (1961), no. 3, 454–477.

J. Grundberg, T. Hansson, A. Karlhede, and J. Leinaas, *On singular anyon wavefunctions*, Mod. Phys. Lett. B **05** (1991), no. 07, 539–546, doi.

F. D. M. Haldane, "Fractional statistics" in arbitrary dimensions: A generalization of the Pauli principle, Phys. Rev. Lett. **67** (1991), 937–940, doi.

B. I. Halperin, Statistics of quasiparticles and the hierarchy of fractional quantized Hall states, Phys. Rev. Lett. **52** (1984), 1583–1586, doi.

M. Hoffmann-Ostenhof, T. Hoffmann-Ostenhof, A. Laptev, and J. Tidblom, *Many-particle Hardy Inequalities*, J. London Math. Soc. **77** (2008), 99–114, doi.

Y. Hu, G. Murthy, S. Rao, and J. K. Jain, *Kohn-Sham density functional theory of abelian anyons*, Phys. Rev. B **103** (2021), 035124, doi.

R. Iengo and K. Lechner, Anyon quantum mechanics and Chern-Simons theory, Phys. Rep. **213** (1992), 179–269, doi.

R. Jackiw, *Topics in planar physics*, Nuclear Phys. B Proc. Suppl. **18A** (1990), 107–170, Integrability and quantization (Jaca, 1989), doi.

J. K. Jain, *Composite fermions*, Cambridge Univ. Press, 2007. A. Khare, *Fractional Statistics and Quantum Theory*, 2nd ed., World Scientific, Singapore, 2005.

A. Y. Kitaev, Fault-tolerant quantum computation by anyons, Ann. Physics **303** (2003), no. 1, 2–30, doi.

A. Kitaev, Anyons in an exactly solved model and beyond, Ann. Physics **321** (2006), no. 1, 2–111, doi.

B. Klaiber, *The Thirring model*, Lectures in Theoretical Physics, Vol. X-A: Quantum Theory and Statistical Physics (A. O. Barut and W. E. Brittin, eds.), Gordon and Breach, New York, 1968, Presented at the Theoretical Physics Institute, University of Colorado, Summer 1967, pp. 141–176.

T. Kohno, Monodromy representations of braid groups and Yang-Baxter equations, Ann. Inst. Fourier (Grenoble) **37** (1987), no. 4, 139–160, http://www.numdam.org/item?id=AIF_1987__37_4_139_0.

C. Korff, G. Lang, and R. Schrader, *Two-particle scattering theory for anyons*, J. Math. Phys. **40** (1999), no. 4, 1831–1869, doi.

A. Kundu, Exact solution of double δ function bose gas through an interacting anyon gas, Phys. Rev. Lett. **83** (1999), 1275–1278, doi.

V. Lahtinen and J. Pachos, A short introduction to topological quantum computation, SciPost Physics 3 (2017), no. 3, 021, doi.

M. G. G. Laidlaw and C. M. DeWitt, Feynman functional integrals for systems of indistinguishable particles, Phys. Rev. D 3 (1971), 1375–1378, doi.

G. Lambert, D. Lundholm, and N. Rougerie, *Quantum statistics transmutation via magnetic flux attachment*, to appear in Prob. Math. Phys., 2023, arXiv:2201.03518.

L. Landau, Über die Bewegung der Elektronen in Kristallgitter, Physik Z. Sowjetunion **3** (1933), 644–645.

M. Lankhorst, A. Brinkman, H. Hilgenkamp, N. Poccia,

and A. Golubov, Annealed Low Energy States in Frustrated Large Square Josephson Junction Arrays, Condens. Matter **3** (2018), 19, doi.

S. Larson and D. Lundholm, *Exclusion bounds for extended anyons*, Arch. Ration. Mech. Anal. **227** (2018), 309–365, doi.

S. Larson, D. Lundholm, and P. T. Nam, *Lieb-Thirring in*equalities for wave functions vanishing on the diagonal set, Ann. Henri Lebesgue 4 (2021), 251–282, doi.

R. B. Laughlin, Anomalous quantum Hall effect: An incompressible quantum fluid with fractionally charged excitations, Phys. Rev. Lett. **50** (1983), no. 18, 1395–1398, doi.

R. B. Laughlin, Nobel lecture: Fractional quantization, Rev. Mod. Phys. 71 (1999), 863–874, doi.

T. Lee and P. Oh, Non-abelian Chern-Simons quantum mechanics and non-abelian Aharonov-Bohm effect, Ann. Physics 235 (1994), no. 2, 413–434, doi.

J. M. Leinaas and J. Myrheim, On the theory of identical particles, Nuovo Cimento B **37** (1977), 1–23, doi.

J. M. Leinaas and J. Myrheim, *Heisenberg quantization for* systems of identical particles, International Journal of Modern Physics A 8 (1993), no. 21, 3649–3695, doi.

J. M. Leinaas and J. Myrheim, *Fractional statistics in low-dimensional systems*, For the Encyclopedia of Condensed Matter Physics, 2e., 2022.

A. Lerda, Anyons, Springer-Verlag, Berlin-Heidelberg, 1992.
S. Li, R. K. Bhaduri, and M. V. N. Murthy, Thomas-fermi approximation for confined anyons, Phys. Rev. B 46 (1992), 1228-1231, doi.

E. H. Lieb and W. Liniger, Exact analysis of an interacting Bose gas. I. The general solution and the ground state, Phys. Rev. (2) **130** (1963), 1605–1616, doi.

E. H. Lieb and R. Seiringer, *The Stability of Matter in Quantum Mechanics*, Cambridge Univ. Press, 2010.

E. H. Lieb and W. E. Thirring, Bound for the kinetic energy of fermions which proves the stability of matter, Phys. Rev. Lett. **35** (1975), 687–689, doi.

E. H. Lieb and W. E. Thirring, *Inequalities for the moments of the eigenvalues of the Schrödinger hamiltonian and their relation to Sobolev inequalities*, Studies in Mathematical Physics, pp. 269–303, Princeton Univ. Press, 1976.

E. H. Lieb and J. Yngvason, *The ground state energy of a dilute two-dimensional Bose gas*, J. Statist. Phys. **103** (2001), no. 3-4, 509–526, Special issue dedicated to the memory of Joaquin M. Luttinger, **doi**.

E. H. Lieb, J. P. Solovej, and J. Yngvason, *Ground states of large quantum dots in magnetic fields*, Phys. Rev. B **51** (1995), 10646–10665, doi.

E. H. Lieb, R. Seiringer, J. P. Solovej, and J. Yngvason, *The* mathematics of the Bose gas and its condensation, Oberwolfach Seminars, Birkhäuser, 2005, arXiv:cond-mat/0610117.

D. Lundholm, Many-anyon trial states, Phys. Rev. A 96 (2017), 012116, doi.

D. Lundholm, Methods of modern mathematical physics: Uncertainty and exclusion principles in quantum mechanics, KTH and LMU graduate course textbook (latest version at http://www.mathematik.uni-muenchen.de/ ~lundholm/methmmp.pdf), 2019, arXiv:1805.03063.

D. Lundholm and V. Qvarfordt, Exchange and exclusion in the non-abelian anyon gas, arXiv e-prints, 2020, arXiv:2009. 12709.

D. Lundholm and N. Rougerie, *The average field approximation for almost bosonic extended anyons*, J. Stat. Phys. **161** (2015), no. 5, 1236–1267, doi.

D. Lundholm and N. Rougerie, *Emergence of fractional statistics for tracer particles in a Laughlin liquid*, Phys. Rev. Lett. **116** (2016), 170401, doi.

D. Lundholm and R. Seiringer, *Fermionic behavior of ideal anyons*, Lett. Math. Phys. **108** (2018), 2523–2541, doi.

D. Lundholm and J. P. Solovej, *Hardy and Lieb-Thirring in-equalities for anyons*, Comm. Math. Phys. **322** (2013), 883–908, doi.

D. Lundholm and J. P. Solovej, Local exclusion principle for identical particles obeying intermediate and fractional statistics, Phys. Rev. A 88 (2013), 062106, doi.

D. Lundholm and J. P. Solovej, *Local exclusion and Lieb-Thirring inequalities for intermediate and fractional statistics*, Ann. Henri Poincaré **15** (2014), 1061–1107, doi.

D. Lundholm, F. Portmann, and J. P. Solovej, *Lieb-Thirring bounds for interacting Bose gases*, Comm. Math. Phys. **335** (2015), no. 2, 1019–1056, doi.

D. Lundholm, P. T. Nam, and F. Portmann, *Fractional Hardy-Lieb-Thirring and related inequalities for interacting systems*, Arch. Ration. Mech. Anal. **219** (2016), no. 3, 1343–1382, doi.

T. Maciazek and A. Sawicki, Non-abelian quantum statistics on graphs, Commun. Math. Phys. **371** (2019), 921–973, doi.

F. Mancarella, A. Trombettoni, and G. Mussardo, *Statistical mechanics of an ideal gas of non-abelian anyons*, Nucl. Phys. B **867** (2013), no. 3, 950–976, doi.

F. Mancarella, A. Trombettoni, and G. Mussardo, *Statistical interparticle potential of an ideal gas of non-abelian anyons*, J. Phys. A: Math. Theor. **46** (2013), no. 27, 275001, doi.

C. Manuel and R. Tarrach, *Contact interactions of anyons*, Phys. Lett. B **268** (1991), no. 2, 222–226, doi.

Y. Masaki, T. Mizushima, and M. Nitta, Non-abelian anyons and non-abelian vortices in topological superconductors, For the Encyclopedia of Condensed Matter Physics, 2e., 2023, arXiv:2301.11614.

P. L. McEuen, E. B. Foxman, J. Kinaret, U. Meirav, M. A. Kastner, N. S. Wingreen, and S. J. Wind, *Self-consistent addition spectrum of a Coulomb island in the quantum Hall regime*, Phys. Rev. B **45** (1992), 11419–11422, doi.

A. M. L. Messiah and O. W. Greenberg, Symmetrization postulate and its experimental foundation, Phys. Rev. **136** (1964), B248–B267, doi.

G. Moore and N. Read, Nonabelions in the fractional quantum Hall effect, Nucl. Phys. B **360** (1991), no. 2, 362 – 396, doi.

S. C. Morampudi, A. M. Turner, F. Pollmann, and F. Wilczek, *Statistics of fractionalized excitations through threshold spectroscopy*, Phys. Rev. Lett. **118** (2017), 227201, doi.

A. Mouchet, Path integrals in a multiply-connected configuration space (50 years after), Foundations of Physics **51** (2021), no. 6, 107, doi.

J. Mund and R. Schrader, *Hilbert spaces for nonrelativistic and relativistic "free" plektons (particles with braid group statistics)*, Advances in dynamical systems and quantum physics (Capri, 1993), World Sci. Publ., River Edge, NJ, 1995, pp. 235–259, arXiv:hep-th/9310054.

J. Mund, The spin-statistics theorem for anyons and plektons in d=2+1, Commun. Math. Phys. **286** (2009), no. 3, 1159–

30

1180, **doi**.

M. V. N. Murthy, J. Law, M. Brack, and R. K. Bhaduri, Quantum spectrum of three anyons in an oscillator potential, Phys. Rev. Lett. **67** (1991), 1817–1820, doi.

M. V. N. Murthy, J. Law, R. K. Bhaduri, and G. Date, On a class of noninterpolating solutions of the many-anyon problem, J. Phys. A: Math. Gen. **25** (1992), no. 23, 6163, doi.

J. Myrheim, *Anyons*, Topological aspects of low dimensional systems (A. Comtet, T. Jolicœur, S. Ouvry, and F. David, eds.), Les Houches - Ecole d'Ete de Physique Theorique, vol. 69, (Springer-Verlag, Berlin, Germany), 1999, pp. 265–413, doi.

J. Nakamura, S. Liang, G. C. Gardner, and M. J. Manfra, Direct observation of anyonic braiding statistics, Nature Phys. 16 (2020), 931–936, doi.

C. Nayak and F. Wilczek, 2n-quasihole states realize 2^{n-1} dimensional spinor braiding statistics in paired quantum Hall states, Nuclear Phys. B **479** (1996), no. 3, 529–553, doi.

C. Nayak, S. H. Simon, A. Stern, M. Freedman, and S. Das Sarma, *Non-abelian anyons and topological quantum computation*, Rev. Mod. Phys. **80** (2008), 1083–1159, doi.

J. Noh, T. Schuster, T. Iadecola, S. Huang, M. Wang, K. P. Chen, C. Chamon, and M. C. Rechtsman, *Braiding photonic topological zero modes*, Nature Physics **16** (2020), no. 9, 989–993, doi.

L. Oddis, *Two-anyon schrödinger operators*, Ph.D. thesis, Università di Roma, Sapienza, Facoltà di Scienze Matematiche, Fisiche e Naturali, 2020, https://phd.uniroma1.it/ web/LUCA-ODDIS_nT1669369_EN.aspx.

S. Ouvry, Anyons and lowest Landau level anyons, Séminaire Poincaré 11 (2007), 77–107, doi.

S. Ouvry and A. P. Polychronakos, Anyons on the sphere: Analytic states and spectrum, Nuclear Physics B **949** (2019), 114797, doi.

S. Pancharatnam, Generalized theory of interference, and its applications: Part i. coherent pencils, Proceedings of the Indian Academy of Sciences-Section A, vol. 44, Springer India New Delhi, 1956, pp. 247–262.

S. Pekar, Local quantum states of electrons in an ideal ion crystal, Zhurnal Eksperimentalnoi I Teoreticheskoi Fiziki 16 (1946), no. 4, 341–348.

L. P. Pitaevskii, Vortex lines in an imperfect bose gas, Zh. Eksper. Teor. fiz. **40** (1961), no. 40, 646–651.

A. G. A. Pithis and H.-C. Ruiz Euler, Anyonic statistics and large horizon diffeomorphisms for loop quantum gravity black holes, Phys. Rev. D **91** (2015), 064053, doi.

A. P. Polychronakos, *Non-relativistic bosonization and fractional statistics*, Nuclear Physics B **324** (1989), no. 3, 597– 622, doi.

A. P. Polychronakos and S. Ouvry, *Two anyons on the sphere: Nonlinear states and spectrum*, Nuclear Physics B **951** (2020), 114906, doi.

V. Qvarfordt, Non-abelian anyons: Statistical repulsion and topological quantum computation, MSc thesis, KTH, 2017, http://urn.kb.se/resolve?urn=urn%3Anbn% 3Ase%3Akth%3Adiva-207177.

N. Read and E. Rezayi, Beyond paired quantum Hall states: Parafermions and incompressible states in the first excited Landau level, Phys. Rev. B **59** (1999), 8084–8092, doi.

N. Regnault, M. O. Goerbig, and T. Jolicoeur, Bridge between

abelian and non-abelian fractional quantum Hall states, Phys. Rev. Lett. **101** (2008), 066803, doi.

N. Rougerie and Q. Yang, *Dimensional reduction for a system* of 2D anyons, arXiv e-prints, 2023, arXiv:2305.06670.

E. C. Rowell and Z. Wang, *Localization of unitary braid group* representations, Comm. Math. Phys. **311** (2012), no. 3, 595–615, doi.

M. Schick, Two-dimensional system of hard-core bosons, Phys. Rev. A **3** (1971), 1067–1073, doi.

R. Schmidt and M. Lemeshko, *Rotation of quantum impurities* in the presence of a many-body environment, Phys. Rev. Lett. **114** (2015), 203001, doi.

L. Schulman, A path integral for spin, Phys. Rev. **176** (1968), 1558–1569, doi.

R. Seiringer and J. P. Solovej, A simple approach to Lieb-Thirring type inequalities, arXiv e-prints, 2023, arXiv:2303. 04504.

B. Simon, Holonomy, the quantum adiabatic theorem, and Berry's phase, Phys. Rev. Lett. **51** (1983), 2167–2170, doi.

J.-M. Souriau, *Structure des systèmes dynamiques*, Maîtrises de mathématiques, Dunod, Paris, 1970, English translation by R. H. Cushman and G. M. Tuynman, Progress in Mathematics, 149, Birkhäuser Boston Inc., Boston, MA, 1997, http://www.jmsouriau.com/structure_des_systemes_ dynamiques.htm.

M. Sporre, J. J. M. Verbaarschot, and I. Zahed, Numerical solution of the three-anyon problem, Phys. Rev. Lett. **67** (1991), 1813–1816, doi.

M. Sporre, J. J. M. Verbaarschot, and I. Zahed, *Four anyons in a harmonic well*, Phys. Rev. B **46** (1992), 5738–5741, doi.

A. Stern, Anyons and the quantum Hall effect – A pedagogical review, Ann. Phys. **323** (2008), no. 1, 204–249, January Special Issue 2008, doi.

H. L. Stormer, Nobel lecture: The fractional quantum Hall effect, Rev. Mod. Phys. **71** (1999), 875–889, doi.

R. F. Streater and I. F. Wilde, *Fermion states of a boson field*, Nuclear Physics B **24** (1970), no. 3, 561–575, doi.

B. Sutherland, Quantum Many-Body Problem in One Dimension: Ground State, J. Math. Phys. 12 (1971), 246-250, doi.
L. H. Thomas, The calculation of atomic fields, Proc. Camb.

Philos. Soc. 23 (1927), no. 05, 542–548, doi.

L. Tonks, The complete equation of state of one, two and three-dimensional gases of hard elastic spheres, Phys. Rev. **50** (1936), 955–963, doi.

C. Trugenberger, The anyon fluid in the Bogoliubov approximation, Phys. Rev. D 45 (1992), 3807–3817, doi.

C. Trugenberger, Ground state and collective excitations of extended anyons, Phys. Lett. B **288** (1992), 121–128, doi.

A. Tsuchiya and Y. Kanie, Vertex operators in the conformal field theory on p1 and monodromy representations of the braid group, Lett. Math. Phys. **13** (1987), 303–312, doi.

A. Tsuchiya and Y. Kanie, Vertex operators in conformal field theory on p1 and monodromy representations of braid group, Conformal Field Theory and Solvable Lattice Models, Advanced Studies in Pure Math **16** (1988), 297–372, doi.

D. C. Tsui, H. L. Stormer, and A. C. Gossard, Twodimensional magnetotransport in the extreme quantum limit, Phys. Rev. Lett. 48 (1982), 1559–1562, doi.

D. C. Tsui, Nobel lecture: Interplay of disorder and inter-

action in two-dimensional electron gas in intense magnetic fields, Rev. Mod. Phys. **71** (1999), 891–895, doi.

R. O. Umucahlar, E. Macaluso, T. Comparin, and I. Carusotto, *Time-of-flight measurements as a possible method to observe anyonic statistics*, Phys. Rev. Lett. **120** (2018), 230403, doi.

E. P. Verlinde, A note on braid statistics and the non-Abelian Aharonov-Bohm effect, Modern Quantum Field Theory: proceedings. (S. Das, A. Dhar, S. Mukhi, A. Raina, and A. Sen., eds.), World Scientific, 1991, Proc. of Conference: International Colloquium on Modern Quantum Field Theory, Bombay, 1990, pp. 450–461, https://lib-extopc.kek.jp/ preprints/PDF/1991/9106/9106160.pdf.

O. Weinberger, *The braid group, representations and non-abelian anyons*, BSc thesis, KTH, 2015, http://urn.kb.se/resolve?urn=urn%3Anbn%3Ase%3Akth%3Adiva-167993.

X. G. Wen, Non-abelian statistics in the fractional quantum Hall states, Phys. Rev. Lett. **66** (1991), 802–805, doi.

F. Wilczek, Magnetic flux, angular momentum, and statistics, Phys. Rev. Lett. 48 (1982), 1144–1146, doi.

F. Wilczek, Quantum mechanics of fractional-spin particles,

Phys. Rev. Lett. 49 (1982), 957-959, doi.

F. Wilczek, Fractional Statistics and Anyon Superconductivity, World Scientific, Singapore, 1990.

F. Wilczek and Y.-S. Wu, Space-time approach to holonomy scattering, Phys. Rev. Lett. **65** (1990), 13–16, doi.

Y.-S. Wu, General theory for quantum statistics in two dimensions, Phys. Rev. Lett. **52** (1984), 2103–2106, doi.

Y.-S. Wu, Multiparticle quantum mechanics obeying fractional statistics, Phys. Rev. Lett. 53 (1984), 111–114, doi.

E. Yakaboylu and M. Lemeshko, Anyonic statistics of quantum impurities in two dimensions, Phys. Rev. B **98** (2018), 045402, doi.

E. Yakaboylu, A. Ghazaryan, D. Lundholm, N. Rougerie, M. Lemeshko, and R. Seiringer, *Quantum impurity model for anyons*, Phys. Rev. B **102** (2020), 144109, doi.

Y. Zhang, G. J. Sreejith, N. D. Gemelke, and J. K. Jain, *Fractional angular momentum in cold atom systems*, Phys. Rev. Lett. **113** (2014), 160404, doi.

Y. Zhang, G. J. Sreejith, and J. K. Jain, *Creating and manipulating non-Abelian anyons in cold atom systems using auxiliary bosons*, Phys. Rev. B. **92** (2015), 075116, doi.