Decomposition of $\mathcal{N}=1$ superconformal minimal models and their fractional quantum Hall wavefunctions

Yichen Hu,^a Sirui Ning,^b Yehao Zhou^c

yehao.zhou@ipmu.jp

ABSTRACT: $\mathcal{N} = 1$ superconformal minimal models are the first series of unitary conformal field theories (CFTs) extending beyond Virasoro algebra. Using coset constructions, we characterize CFTs in $\mathcal{N} = 1$ superconformal minimal models using combinations of a parafermion theory, an Ising theory and a free boson theory. Supercurrent operators in the original theory also becomes sums of operators from each constituent theory. If we take our $\mathcal{N} = 1$ superconformal theories as the neutral part of the edge theory of a fractional quantum Hall state, we present a systematic way of calculating its ground state wavefunction using free field methods. Each ground state wavefunction is known previously as a sum of polynomials with distinct clustering behaviours. Based on our decomposition, we find explicit expressions for each summand polynomial. A brief generalization to S_3 minimal models using coset construction is also included.

^aDepartment of Physics, Princeton University,

Princeton, New Jersey 08544, USA

^bThe Rudolf Peierls Centre for Theoretical Physics, University of Oxford,

Oxford OX1 3PU, United Kingdom

^cKavli Institute for the Physics and Mathematics of the Universe (WPI), the University of Tokyo, Kashiwa, Chiba 277-8583, Japan

E-mail: yh1553@princeton.edu, sirui.ning@physics.ox.ac.uk,

Contents

I	Introduction	1
2	$\mathcal{N}=1$ superconformal minimal models and its coset constructions	2
	2.1 Decomposition	3
3	Fractional Quantum Hall wavefunction	5
	3.1 Correlators of supercurrent operator G	6
	3.2 Clustering behaviours of the wavefunction	13
4	Discussion	19
A	S_3 minimal model	20
B	Wavefunction comparison	21

1 Introduction

Unitary conformal field theories [1] have played important roles in many different fields of physics ranging from string theory [2, 3] to critical statistical mechanical systems [3, 4]. Recent studies have also shown their intimate connections to topological phases of matter from symmetry protected topological phases [5–8] to fractional quantum Hall states [9–16]. Specifically, for quantum Hall states, following many pioneering works, wavefunctions of fractional quantum Hall states are shown to be closely related to correlators of primariy fields of the unitary conformal field theory describing its low-energy edge physics. This is a manifestisation of the general "bulk-boundary" correspondence of a topologically ordered state where the bulk hosts a topological field theory and the boundary hosts a conformal field theory. Wavefunctions from unitary conformal field theories associated with fractional quantum Hall states provide deep insights into understandings of physical properties of these exotic phases of matter, such as anyonic excitaions, fractional statistics and even non-Abelian fusion structures among anyons.

 $\mathcal{N} = 1$ superconformal field theories (SCFTs) are the first series of unitary field theories nontrivially extending Virasoro algebra [17–19]. Each member SCFT has a $\Delta = \frac{3}{2}$ supercurrent operator G (also called $\mathbb{Z}_2^{(r)}$ parafermion in [20] because of its fusion structure: $G \times G = \mathbb{I}$) that together with the usual $\Delta = 2$ stress-energy tensor T forming the $\mathcal{N} = 1$ superconformal algebra. This algebra coincides with $WB_1(\beta)$ algebra [21], where β is related to the central charge by $c = 15/2 - 3\beta - 3\beta^{-1}$. Note that $WB_1(\beta)$ minimal models correspond to $\beta = p/q$, where p, q are coprime integers. Other generalizations beyond Virasoro algebra involving operators with fractional spin $\Delta = \frac{N+1}{N}$ can be found in [22–24] (N = 3 corresponds to S_3 minimal models [25]). Our motivations to study this particular series of CFTs are two fold. Recent works following [26–29] have revealed an intricate structure in the theory space of rational CFTs. Different rational CFTs with highly similar topological modular data, fusion structures and anyon contents, are related by number theoretical maps at the level of characters. One can trace these different classes of topological modular data back to realizations in either some parafermion or minimal model M(p,q) theories (usually non-unitary $q \neq p + 1$). Along this line of thinking, we want to understand how our $\mathcal{N} = 1$ minimal models can be decomposed into more "primitive" CFTs. Furthermore, we go beyond characters and try to find explicit decompositions of primary fields with Abelian fusion structure. On the other hand, certain anyons are essential for topological quantum computation [30]. Member CFTs in $\mathcal{N} = 1$ series are known to host these anyons that are capable of performing universal quantum computation by braiding alone. For example, the tri-critical Ising model [3], also known as the edge theory of the "anti-Fibonacci" phase [31], hosts Fibonacci anyons with scaling dimension $\Delta = \frac{3}{5}$ and fusion rule $\tau \times \tau = \mathbb{I} + \tau$. A comprehensive understanding of the ground state and quasiparticle wavefunctions of these topological phases are crucial for their quantum information applications.

This paper is organized as follows: in section 2, we present explicit characterization of the $\mathcal{N} = 1$ superconformal minimal models using coset construction. This also gives us an unique decomposition of the supercurrent operators. In section 3, we take our $\mathcal{N} = 1$ superconformal field theories as the neutral part of a fractional quantum Hall state. We work out its ground state wavefunction based on correlators of the supercurrent operators. The main tool is to use the fact that correlators of supercurrent operators depends on the central charge in a polynomial way, so correlators of supercurrent operators in various free field theories with different central charges uniquely determine the correlators themselves. By fixing the limit theory of our minimal models at $c = \frac{3}{2}$, we can further organize our wavefunction polynomial into sums of symmetric polynomials with distinct clustering behaviours following [20]. Our approach gives explicit expression for each summand symmetric polynomial in a concise format.

2 $\mathcal{N} = 1$ superconformal minimal models and its coset constructions

 $\mathcal{N}=1$ superconformal minimal models are a series of conformal field theories with $\mathcal{N}=1$ super-Virasoro algebra generated by

$$T(z)T(z') = \frac{c/2}{(z-z')^4} + \frac{2T(z')}{(z-z')^2} + \frac{\partial T(z')}{z-z'} + \mathcal{O}(1)$$

$$T(z)G(z') = \frac{(3/2)G(z')}{(z-z')^2} + \frac{\partial G(z')}{z-z'} + \mathcal{O}(1)$$

$$G(z)G(z') = \frac{(2c/3)}{(z-z')^3} + \frac{2T(z')}{z-z'} + \mathcal{O}(1)$$

(2.1)

where $z = x + i\tau$ denoting the (1 + 1)d Euclidean space-time coordinates compactified on a infinite cylinder and G is the scaling dimension $\Delta = \frac{3}{2}$ supercurrent operator that generates the $\mathcal{N} = 1$

supersymmetry. They have central charge $c = \frac{3}{2}(1 - \frac{8}{m(m+2)})$ with $m = 3, 4, \cdots$ where m = 3 is the tri-critical Ising CFT. Each member SCFT has a known coset construction [3, 17]

$$\frac{SU(2)_2 \times SU(2)_{m-2}}{SU(2)_m}$$
(2.2)

2.1 Decomposition

In order to decompose our $\mathcal{N} = 1$ theories, let us first large our SCFTs by pairing them with a \mathbb{Z}_m parafermion CFT $\frac{SU(2)_m}{U(1)_{2m}}$. Reshuffling factors in cosets of this product theory, we get the following relations:

$$\frac{SU(2)_m}{U(1)_{2m}} \times \frac{SU(2)_2 \times SU(2)_{m-2}}{SU(2)_m} = \frac{SU(2)_{m-2}}{U(1)_{2(m-2)}} \times \frac{SU(2)_2}{U(1)_4} \times U(1)_{4m(m-2)}$$
(2.3)

This equality holds at the level of stress-energy tensor. The U(1) factors in Eq. (2.3) needs a little explanation. Let us reshuffle U(1) theories as following:

Left
$$(U(1)_{2m})^{-1} \times U(1)_{2(m-2)} \times U(1)_4$$
, Right $U(1)_{4m(m-2)}$. (2.4)

To establish an equivalence relation between the left and right hand side, we first write out their Lagrangian densities:

$$\mathcal{L}_{l} = \frac{1}{4\pi} \sum_{I,J=1}^{3} K_{l}^{IJ} \partial_{x} \phi_{I} (\partial_{x} + \partial_{t}) \phi_{J}, \quad \mathcal{L}_{r} = \frac{1}{4\pi} K_{r} \partial_{x} \phi_{1} (\partial_{x} + \partial_{t}) \phi_{1}$$
(2.5)

where $K_l = \begin{pmatrix} -2m & 0 & 0 \\ 0 & 2(m-2) & 0 \\ 0 & 0 & 4 \end{pmatrix}$ and $K_r = 4m(m-2)$. It is easy to see that there exists a matrix

M with integer entries

$$M = \begin{pmatrix} 2 - m - m & 0 \\ 1 & 1 & 1 \\ -1 & -1 & 1 \end{pmatrix}, \quad \det M = 4 > 0$$
(2.6)

such that

$$M^{T}\begin{pmatrix} -2m & 0 & 0\\ 0 & 2(m-2) & 0\\ 0 & 0 & 4 \end{pmatrix})M = \left(4m(m-2)\right) \bigoplus 8\sigma_{z}.$$
 (2.7)

The right hand side of Eq. (A.7) is equivalent to \mathcal{L}_r as we can add a backscattering term to the two counter-propagating modes

$$2\cos(\sqrt{2}(\phi_2 - \phi_3))$$
 (2.8)

to gap out $8\sigma_z$ degrees of freedom (ϕ_2 and ϕ_3)[32, 33]. We have established an equivalence relation in Eq. (2.4).

Eq. (2.3) means we can combine a \mathbb{Z}_m parafermion theory with the *m*th member of the $\mathcal{N} = 1$ SCFT theory and turn them into a linear combination of a \mathbb{Z}_{m-2} parafermion theory, a U(1) free boson theory and an Ising (\mathbb{Z}_2 parafermion) theory. Focusing on stress-energy tensor, we have

$$T_{min} = \gamma T_{\mathbb{Z}_{m-2}} + \alpha T_{\mathbb{Z}_2} + \beta T_{U(1)} + \delta t$$

$$T_{\mathbb{Z}_m} = (1 - \gamma) T_{\mathbb{Z}_{m-2}} + (1 - \alpha) T_{\mathbb{Z}_2} + (1 - \beta) T_{U(1)} - \delta t$$
(2.9)

where γ , α , β and δ are coefficients representing weights for each constituent stress-energy tensor. t is an extra $\Delta = 2$ auxiliary field made out of Abelian primary fields from each constituent CFT (γ from Ising theory, φ_1 ($\overline{\varphi_1}$) from parafermion theory and $e^{i\sqrt{\frac{m}{m-2}}\phi}$ from U(1) theory), that is key to make this decomposition work. It satisfies

$$t(z)t(w) \sim \frac{1}{(z-w)^4} + \frac{1}{(z-w)^2} \left(2T_{\mathbb{Z}_2} + \frac{m}{m-2}T_{U(1)} + \frac{m}{m-2}T_{\mathbb{Z}_{m-2}} \right)$$
(2.10)

since consistency of operator product expansion between stress-energy tensor T's and t requires coefficient in front of $T_{\mathbb{Z}_2}$ is $\frac{2\Delta(\gamma)}{c_{\mathbb{Z}_2}} = 2$, $T_{U(1)}$ is $\frac{2\Delta(e^i\sqrt{\frac{m}{m-2}}\phi)}{c_{U(1)}} = \frac{m}{m-2}$ and $T_{\mathbb{Z}_{m-2}}$ is $\frac{2\Delta(\varphi_1(\bar{\varphi_1}))}{c_{\mathbb{Z}_{m-2}}} = \frac{m}{m-2}$. Using this fact together with the usual stress-energy tensor OPE:

$$T_{\min/\mathbb{Z}_m}(z)T_{\min/\mathbb{Z}_m}(w) \sim \frac{c/2}{(z-w)^4} + \frac{2T_{\min/\mathbb{Z}_m}(w)}{(z-w)^2},$$
 (2.11)

we arrive at the following set of equations for these coefficients:

$$\frac{1}{2}\alpha + \frac{m}{2(m-2)}\beta + \frac{m-3}{m-2}\gamma = 1 \text{ (for } t \text{ coefficient)}$$

$$\alpha^{2} + \delta^{2} = \alpha \text{ (for } \mathbb{Z}_{2} \text{ coefficient)}$$

$$\beta^{2} + \frac{m}{2(m-2)}\delta^{2} = \beta \text{ (for } U(1) \text{ coefficient)}$$

$$\gamma^{2} + \frac{m}{2(m-2)}\delta^{2} = \gamma \text{ (for } \mathbb{Z}_{m-2} \text{ coefficient)}$$
(2.12)

They have two solutions which corresponds to the two stress-energy tensors T_{min} and $T_{\mathbb{Z}_m}$:

$$T_{min} = \frac{m-2}{m+2} T_{\mathbb{Z}_2} + \frac{m}{m+2} T_{U(1)} + \frac{2}{m+2} T_{\mathbb{Z}_{m-2}} \pm \frac{2\sqrt{m-2}}{m+2} t$$

$$T_{\mathbb{Z}_m} = \frac{4}{m+2} T_{\mathbb{Z}_2} + \frac{2}{m+2} T_{U(1)} + \frac{m}{m+2} T_{\mathbb{Z}_{m-2}} + \mp \frac{2\sqrt{m-2}}{m+2} t$$
(2.13)

Now, we further decompose the supercurrent operator G in terms of operators from different sectors. To this end, we have

$$T_{min}(z)G(w) \sim \frac{\frac{3}{2}G(w)}{(z-w)^2} \quad T_{\mathbb{Z}_m}(z)G(w) \sim 0$$

$$G(z)G(w) \sim \frac{\frac{2c}{3}}{(z-w)^3}$$
(2.14)

These relations give us a unique decomposition of G with the exception of m = 4 (it has $\mathcal{N} = 2$ superconformal algebra and therefore the decomposition of G is not unique due to the emergent U(1) R-symmetry). For m odd,

$$G = \pm i \sqrt{\frac{2(m-2)}{m(m+2)}} \left(\varphi_1 e^{i\sqrt{\frac{m}{m-2}}\phi} - \bar{\varphi}_1 e^{-i\sqrt{\frac{m}{m-2}}\phi} \right) \mp i \sqrt{\frac{m-2}{m+2}} \gamma \partial \phi$$

$$t = \frac{i}{\sqrt{2}} \left(\varphi_1 \gamma e^{i\sqrt{\frac{m}{m-2}}\phi} + \bar{\varphi}_1 \gamma e^{-i\sqrt{\frac{m}{m-2}}\phi} \right)$$
(2.15)

and for m even,

$$G = \pm \sqrt{\frac{2(m-2)(m+4)}{m(m+2)(m-4)}} \left(\varphi_1 e^{i\sqrt{\frac{m}{m-2}}\phi} - \bar{\varphi}_1 e^{-i\sqrt{\frac{m}{m-2}}\phi}\right) \pm i\sqrt{\frac{(m-2)(m+4)}{(m+2)(m-4)}}\gamma\partial\phi$$

$$t = \frac{1}{\sqrt{2}} \left(\varphi_1\gamma e^{i\sqrt{\frac{m}{m-2}}\phi} + \bar{\varphi}_1\gamma e^{-i\sqrt{\frac{m}{m-2}}\phi}\right)$$
(2.16)

where $\{\varphi_0 = \mathbb{I}, \varphi_1, \varphi_2, \cdots, \varphi_{m-3} = \bar{\varphi}_1\}$ are the \mathbb{Z}_{m-2} parafermion primary fields, γ is a Majorana fermion from Ising CFT and ϕ is a U(1) boson field. Parafermions (including $\mathbb{Z}_{m-2=2}$, Majorana fermion) φ_i has scaling dimension $\Delta = \frac{i(m-2-i)}{m-2}$ and fusion rule $\varphi_i \times \varphi_j = \varphi_{(i+j) \mod (m-2)}$. Boson field has compactification radius $\phi = \phi + 2\pi$. Its vertex operators $e^{ia\phi}$ has scaling dimension $\Delta = \frac{a^2}{2}$. From these relations, we can conclude that $\Delta(\varphi_1 e^{i\sqrt{\frac{m}{m-2}}\phi}) = \Delta(\bar{\varphi}_1 e^{-i\sqrt{\frac{m}{m-2}}\phi}) = \Delta(\gamma\partial\phi) = \frac{3}{2}$.

Curiously, $\varphi_1 e^{i\sqrt{\frac{m}{m-2}}\phi} (\bar{\varphi}_1 e^{-i\sqrt{\frac{m}{m-2}}\phi})$ appeared in Eq. (2.15) is the same as the electron operator of the (m-2)th member in Read-Rezayi quantum Hall states with filling factor $\nu = \frac{m-2}{m}$ even though $\varphi_1 e^{i\sqrt{\frac{m}{m-2}}\phi} (\bar{\varphi}_1 e^{-i\sqrt{\frac{m}{m-2}}\phi})$ does not carry any physical charge. In low energy, it is known that Read-Rezayi states at filling factor $\nu = \frac{m-2}{m}$ possess emergent $\mathcal{N} = 2$ supersymmetry with the physical U(1) charge being its *R*-symmetry [34]. It is therefore not surprising to see the appearance of $\varphi_1 e^{i\sqrt{\frac{m}{m-2}}\phi} - \bar{\varphi}_1 e^{-i\sqrt{\frac{m}{m-2}}\phi}$ in decomposition of $\mathcal{N} = 1$ supercurrent operator *G*. The other generator for the other half of supersymmetry in this rotated basis is $\varphi_1 e^{i\sqrt{\frac{m}{m-2}}\phi} + \bar{\varphi}_1 e^{-i\sqrt{\frac{m}{m-2}}\phi}$. We observe that the following "particle-hole" (since there is no real physical charge) symmetry

$$\phi \Longleftrightarrow -\phi, \ \varphi_1 \Longleftrightarrow -\bar{\varphi_1}, \ \gamma \Longleftrightarrow -\gamma \tag{2.17}$$

takes $G \to G$ and forbids $\varphi_1 e^{i\sqrt{\frac{m}{m-2}}\phi} + \bar{\varphi}_1 e^{-i\sqrt{\frac{m}{m-2}}\phi}$.

Lastly, from our decomposition, we see that as $m \to \infty$, central charge of SCFTs goes as $c \to \frac{3}{2}$. Operator-wise, $T_{min}(z) \to T_{\mathbb{Z}_2} + T_{U(1)}$ and $G \to \gamma \partial \phi$ which makes it clear that the conformal field theory in the infinity limit is the three Majorana fermion theory which is on the moduli space of $c = \frac{3}{2}$ circle line [35]. This is analogous to the unitary minimal models where they become the U(1) free boson theory on the moduli space of c = 1 circle line in the $m \to \infty$ limit [36].

3 Fractional Quantum Hall wavefunction

To study fractional quantum Hall states based on $\mathcal{N} = 1$ minimal models, we need to combine our $\mathcal{N} = 1$ minimal models (the charge neutral sector) with a U(1) physical charge sector. First, let us define an electron operator that has trivial fusion structure and also braid trivially with all other quasi-particles. Formally, we can write this electron operator as $\psi_e = Ge^{i\sqrt{p/q}\Phi}$ where G is the supercurrent operator of our $\mathcal{N} = 1$ theories and Φ is the boson field representing the U(1) charge. p, q coprime $\in \mathbb{Z}^+$ gives a filling factor of $\frac{q}{p}$ for our FQH state (assuming our ground state only occupy the lowest Landau level). By the bulk-boundary correspondence of quantum Hall physics, its ground state wavefunction is the same as the chiral correlation function from the edge $\mathcal{N} = 1$ SCFT and the U(1) theory:

$$\langle \psi_e(z_1)\cdots\psi_e(z_N)\rangle = \langle G(z_1)\cdots G(z_N)\rangle \langle e^{i\sqrt{p/q}\Phi(z_1)}\cdots e^{i\sqrt{p/q}\Phi(z_N)}\rangle = C_N \prod_{i< j}^N (z_i - z_j)^{p/q}$$
(3.1)

where p/q is chosen such that $\langle \psi_e(z_1) \cdots \psi_e(z_N) \rangle$ is a polynomial with no pole and branch-cut. From this, we see that the task of computing this wavefunction is really about computing $C_N \equiv \langle G(z_1) \cdots G(z_N) \rangle$.

3.1 Correlators of supercurrent operator G

In this section, we compute the n point correlators of supercurrent operators G using free field methods [37–39]. Physically, free field methods utilize free bosonic, fermionic or ghost theories to represent a two dimensional conformal field theory. The simplest example is the well-known "Coulomb-Gas" formalism. Free field methods allow us to compute correlation functions of conformal fields via Wick's theorem.

Before calculations for the supercurrent G, as a demonstration of principle, we apply free field methods on stress-energy tensor T to calculate its n correlation functions. We start from Virasoro algebra, which is generated by:

$$T(z)T(z') = \frac{c/2}{(z-z')^4} + \frac{2T(z')}{(z-z')^2} + \frac{\partial T(z')}{z-z'} + \mathcal{O}(1)$$
(3.2)

For the n point correlator of stress energy tensors, we have the following theorem:

Theorem 1.

$$\langle T(z_1)T(z_2)\cdots T(z_n)\rangle = \sum_{\substack{\sigma\in S_n\\\sigma=(l_1)\cdots(l_s)}} \prod_{i=1}^s f_{l_i},$$
(3.3)

where the perumutation $\sigma = (l_1)...(l_s) \in S_n$ is equivalent to product of cyclic permutations of length at least 2, and for each cyclic permutation $(l_i) = (i_1 i_2 ... i_{m_l})$, define

$$f_{l_i}(z_{i_1}, z_{i_2}, \cdots, z_{i_{m_l}}) = \frac{c/2}{(z_{i_1} - z_{i_2})^2 (z_{i_2} - z_{i_3})^2 \cdots (z_{i_{m_l}} - z_{i_1})^2}.$$
(3.4)

We start the proof with two lemmas:

Lemma 3.1.1. $\langle T(z_1)T(z_2)\cdots T(z_n)\rangle$ is a rational function in variables z_1, z_2, \cdots, z_n and central charge *c*.

Proof. This lemma can be proven by mathematical induction. For n = 2, the two point correlators of stress-energy tensor is $\langle T(z_1)T(z_2)\rangle = \frac{c/2}{(z_1-z_2)^4}$, which by definition is a rational function in variables z_1, z_2 and c.

Suppose the conclusion holds for any n-1 point correlators $T_{n-1}(z_2, z_3, ..., z_n) = \langle T(z_2)T(z_3)...T(z_n) \rangle$. By Eq (3.2), the *n* point correlators satisfy the following recursion relation:

$$T_n(z_1,...z_n) = \sum_{i=2}^n \left(\frac{c/2}{(z_1 - z_i)^4} T_{n-2}(\hat{z_1}, \hat{z_i}) + \left(\frac{2}{(z_1 - z_i)^2} + \frac{\partial_i}{z_1 - z_i} \right) T_{n-1}(z_2, z_3, ...z_n) \right).$$
(3.5)

Where $T_n(z_1,...,z_n) = \langle T(z_1)T(z_2)...T(z_n) \rangle$ and $T_{n-2}(\hat{z}_1, \hat{z}_i) = \langle T(z_2)...T(z_{i-1}), T(z_{i+1})...T(z_n) \rangle$. By mathematical induction, $T_{n-2}(\hat{z}_1, \hat{z}_i)$ and $T_{n-1}(z_2, z_3, \cdots, z_n)$ are both rational function, $T_n(z_1, \cdots, z_n)$ is also a rational function. Therefore the conclusion holds for n point correlators. By mathematical induction, this conclusion holds for all integers $n \ge 2$.

Lemma 3.1.2. Let $f(x_1, x_2, \dots, x_M)$ and $g(x_1, x_2, \dots, x_M)$ be two rational functions over $\{x_i\}_{i=1}^M \in \mathbb{C}$. If

$$f(y, x_2, \cdots, x_M) = g(y, x_2, \cdots, x_M)$$

holds for infinitely many $y \in \mathbb{C}$, then f = g identically.

Proof. Fixing x_2, \dots, x_M , the function $f(x_1, x_2, \dots, x_M) - g(x_1, x_2, \dots, x_M)$ is a rational function in x_1 , and it has infinitely many zeroes, but the number of zeroes of a rational function is bounded by the degree of its numerator, thus $f(x_1, x_2, \dots, x_M) - g(x_1, x_2, \dots, x_M) = 0$ identically. \Box

Let

$$F(c, z_1, z_2, \cdots, z_n) = \sum_{\substack{\sigma \in S_n \\ \sigma = (l_1) \cdots (l_s)}} \prod_{i=1}^n f_{l_i}.$$
(3.6)

If we can show that

$$\langle T(z_1)T(z_2)\cdots T(z_n)\rangle = F(c, z_1, z_2, \cdots, z_n)$$

for infinitely many central charges c, then the theorem is proven.

Proof. We construct the free field realization of Virasoro algebra via free bosons ϕ_i , $i = 1, \dots, N$ with OPE:

$$\partial \phi_i(z) \partial \phi_j(w) = \frac{\delta_{ij}}{(z-w)^2} + O(1).$$
(3.7)

The action for our free field realization is:

$$\int d^2 z \sum_{i=1}^{N} \frac{1}{2} \partial \phi_i \,\overline{\partial} \,\phi_i \tag{3.8}$$

with stress energy tensor $T(z) = \sum_{i=1}^{N} : \frac{1}{2} \partial \phi_i(z) \partial \phi_i(z) :$. One can check that the OPE of this stress energy tensor satisfy Eq (3.2) with central charge c = N. Then the correlator $\langle T(z_1)T(z_2)\cdots T(z_n) \rangle$



Figure 1. Every line represents a contraction between different pair of fields, l_i labels a loop. By Wick's theorem, summing over all the possible products of loops gives the n point amplitude.

can be computed using Wick's theorem, namely we sum over all possible contractions between pair of free bosonic fields:

$$\frac{1}{2^n} \langle : \partial \phi_{i_1}(z_1) \partial \overline{\phi_{i_1}(z_1)} :: \partial \phi_{i_2}(z_2) \partial \overline{\phi_{i_2}(z_2)} : \dots : \partial \phi_{i_k}(z_k) \partial \phi_{i_k}(z_k) : \rangle.$$
(3.9)

Note that graphically, for all possible contractions, each vertex z_i (i = 1, 2, ..., n) must connect with other two vertices z_j and z_k $(j, k \neq i)$. Therefore, for each possible contraction, we first decompose the *n* vertices into different clusters, then we connect all the vertices in each cluster to form a loop. Physically, the amplitude associated with each contraction is the product of the amplitude of each loop, which is:

$$\frac{N/2}{(z_{i_1} - z_{i_2})^2 \cdots (z_{i_{m_l}} - z_{i_1})^2} \times \frac{N/2}{(z_{j_1} - z_{j_2})^2 \cdots (z_{j_{m_j}} - z_{j_1})^2} \times \cdots \times \frac{N/2}{(z_{s_1} - z_{s_2})^2 \cdots (z_{s_{m_s}} - z_{s_1})^2}.$$
(3.10)

The overall n-point amplitude is the sum of amplitudes of all possible contractions. Figure 1 gives a graph representation of each possible contraction.

We give an interpretation to the factor N/2 associated with amplitude for each loop. It is easy to understand the factor of N because there are N free bosonic fields in the representation and the OPE between different fields vanishes. For a loop with m vertices, each energy momentum tensor carries a factor of 1/2, which contributes a factor of $\frac{1}{2^m}$, and each double contraction gives a factor of 2, however, because the diagram is a loop, so all the double contractions contribute a factor of 2^{m-1} . Multiply them together leads to a factor of 1/2. Therefore the total factor is N/2. Therefore $\langle T(z_1) \cdots T(z_n) \rangle_{c=N} = F(c = N, z_1, \cdots, z_n)$ for infinitely many $N \in \mathbb{Z}_{>0}$, thus the theorem holds by the Lemma 3.1.2.

Following Eq (3.3), an equivalent way of writing the correlator is:

$$\langle T(z_1)\cdots T(z_n)\rangle = \sum_{\substack{I_1\cup I_2\cdots \cup I_s = (12\cdots n)\\ \text{cluster decomposition}}} \left(\frac{c}{2}\right)^s \prod_{i=1}^n \hat{f}_{I_i},\tag{3.11}$$

where

$$\hat{f}_{I_i}(z_{i_1},\cdots,z_{i_{M_i}}) = \frac{1}{M_i} \sum_{\sigma \in S_{M_i}} \frac{1}{(z_{\sigma(i_1)} - z_{\sigma(i_2)})^2 (z_{\sigma(i_2)} - z_{\sigma(i_3)})^2 \cdots (z_{\sigma(i_{M_i})} - z_{\sigma(i_1)})^2}.$$
(3.12)

 $(z_{i_1}, z_{i_2}, ..., z_{i_{M_i}}) \in I_i$. We divide the *n* vertices into *s* clusters labeled as $I_i (i = 1, 2, ...s)$, where each I_i contains $|I_i| = M_i$ elements. Each cluster I_i contains at least two vertices. Below we give two examples:

$$n = 3, s = 1, I_1 = (123)$$

$$n = 4, s = 2, I_1 = (12), I_2 = (34), \quad I_1 = (13), I_2 = (24), \quad I_1 = (14), I_2 = (23)$$
(3.13)

The factor $\frac{1}{M_i}$ is because that the Feynman diagram corresponding to each cluster I_i is invariant under M_i elements in permutation group S_{M_i} , where each element $\sigma_{j=1,2,...,M_i}$ is:

$$\sigma_j(z_k) = z_{M_i - j + k}, \quad \text{for} \quad k = 1, 2, \dots j$$

$$\sigma_j(z_k) = z_{k-j}, \quad \text{for} \quad k = j + 1, \dots, M_i.$$
(3.14)

Now we use free field techniques to calculate the n point correlation functions of supercurrent G. We start from $\mathcal{N} = 1$ super Virasoro algebra, which is generated by:

$$T(z)T(w) = \frac{c/2}{(z-w)^4} + \frac{2T(w)}{(z-w)^2} + \frac{\partial T(w)}{z-w} + \mathcal{O}(1)$$

$$T(z)G(w) = \frac{(3/2)G(w)}{(z-w)^2} + \frac{\partial G(w)}{z-w} + \mathcal{O}(1)$$

$$G(z)G(w) = \frac{(2c/3)}{(z-w)^3} + \frac{2T(w)}{z-w} + \mathcal{O}(1)$$

(3.15)

Our goal is to compute

$$\langle G(z_1)G(z_2)\cdots G(z_n)\rangle.$$
 (3.16)

Similar to the stress-energy correlators, we have the following:

Lemma 3.1.3. $\langle G(z_1)G(z_2)\cdots G(z_n)\rangle$ is a rational function in variables z_1, z_2, \cdots, z_n and central charge c.

Proof. This lemma can be proven by mathematical induction. For n = 1, the one point correlator of supercurrent vanishes, i.e $\langle G(z_1) = 0 \rangle$. For n = 2, the two point correlators of supercurrent is $\langle G(z_1)G(z_2) \rangle = \frac{2c/3}{(z_1-z_2)^3}$, which is a rational function in variables z_1, z_2 and c.

Suppose the conclusion holds for the n-2 point correlators $G_{n-2}(z_3, z_4, ..., z_n) = \langle G(z_3)G(z_4)...G(z_n) \rangle$, by Eq (3.15), the n point correlators satisfy the following recursion relation:

$$G_n(z_1, \dots z_n) = \sum_{i=2}^n \left((-1)^i \frac{2c/3}{(z_1 - z_i)^3} + (-1)^i \frac{1}{z_1 - z_i} \sum_{j \neq i, 1}^n \left(\frac{3}{(z_i - z_j)^2} + \frac{\partial_j}{z_i - z_j} \right) \right) G_{n-2}(\hat{z}_1, \hat{z}_i),$$
(3.17)

where $G_n(z_1, ..., z_n) = \langle G(z_1)G_(z_2)...G(z_n) \rangle$ and $G_{n-2}(\hat{z_1}, \hat{z_i}) = \langle G(z_2)...G(z_{i-1})G(z_{i+1})...G(z_n) \rangle$. By mathematical induction, $G_{n-2}(\hat{z_1}, \hat{z_i})$ is a rational function, $G_n(z_1, ..., z_n)$ is also a rational function. Therefore the conclusion holds for n point correlators. By mathematical induction, this conclusion holds for all integers $n \geq 3$. Therefore the lemma is proved.

Note that as a by-product, since $\langle G_1(z_1) \rangle$ vanishes, by this recursion relation, $G_n(z_1, ..., z_n) = 0$ if n is odd.

To begin with, we construct free field representation of $\mathcal{N} = 1$ super Virasoro algebra via free ghost fields [40]: $\beta \gamma - bc$ systems which combines two anticommuting fields bc with two commuting fields $\beta \gamma$. Their weights are:

$$h_{b_i} = \lambda, \quad h_{c^i} = 1 - \lambda, \quad i = 1, 2, \cdots, N$$

 $h_{\beta_j} = \lambda - \frac{1}{2}, \quad h_{\gamma^j} = \frac{3}{2} - \lambda, \quad j = 1, 2, \cdots, N$
(3.18)

The action for our free field realization is:

$$S = \frac{1}{2\pi} \int d^2 z \sum_{i=1}^{N} (b_i \overline{\partial} c^i + \beta_i \overline{\partial} \gamma^i), \qquad (3.19)$$

Their OPEs are:

$$b_i(z)c^j(w) = \frac{\delta_i^j}{z - w} + \mathcal{O}(1)$$

$$\gamma^i(z)\beta_j(w) = \frac{\delta_i^j}{z - w} + \mathcal{O}(1)$$
(3.20)

The stress-energy tensor and supercurrent read:

$$T = (\partial b_i(z))c^i(z) - \lambda \partial (b_i(z)c^i(z)) + (\partial \beta_i(z))\gamma^i(z) - \frac{1}{2}(2\lambda - 1)\partial (\beta_i(z)\gamma^i(z))$$

$$G = -\frac{1}{2}(\partial \beta_i)(z)c^i(z) + \frac{2\lambda - 1}{2}\partial (\beta_i(z)c^i(z)) - 2b_i(z)\gamma^i(z)$$
(3.21)

One can check that their OPE satisfy Eq (3.15). Without loss of generality we choose $\lambda = \frac{1}{2}$, then the supercurrent reads:

$$G(z) = -\frac{1}{2} : \partial \beta_i(z) c^i(z) : -2 : b_i(z) \gamma^i(z) :$$
(3.22)

with central charge c = 3N.

Theorem 2.

$$\langle G(z_1)G(z_2)...G(z_{2n})\rangle = \sum_{\substack{\sigma \in S_{2n} \\ \sigma = (l_1)\cdots(l_s)}} \prod_{i=1}^s \operatorname{sign} \begin{pmatrix} 1 & 2 & \cdots & 2n \\ l_1 & l_2 & \cdots & l_s \end{pmatrix} g_{l_i},$$
(3.23)

where the perumutation $\sigma = (l_1)...(l_s) \in S_{2n}$ is equivalent to product of cyclic permutations of length at least 2, and for each cyclic permutation $(l_i) = (i_1 i_2 ... i_{2m_l})$, define

$$g_{l_i}(z_{i_1}\cdots z_{i_{2m}}) = \frac{c/3}{(z_{i_1}-z_{i_2})(z_{i_2}-z_{i_3})^2\cdots(z_{i_{2m-1}}-z_{i_{2m}})(z_{i_{2m}}-z_{i_1})^2} - \frac{c/3}{(z_{i_1}-z_{i_2})^2(z_{i_2}-z_{i_3})\cdots(z_{i_{2m-1}}-z_{i_{2m}})^2(z_{i_{2m}}-z_{i_1})}.$$
(3.24)

Let

$$H(c, z_1, z_2, \dots z_{2n}) = \sum_{\substack{\sigma \in S_{2n} \\ \sigma = (l_1) \cdots (l_s)}} \prod_{i=1}^s \operatorname{sign} \begin{pmatrix} 1 & 2 & \cdots & 2n \\ l_1 & l_2 & \cdots & l_s \end{pmatrix} g_{l_i}.$$
 (3.25)

If we can show that:

$$\langle G(z_1)G(z_2)...G(z_{2n})\rangle = H(c, z_1, z_2, ..., z_{2n})$$
(3.26)

for infinitely many central charges c, then the theorem is proven.

Proof. We know by free field realization $\langle G(z_1) \cdots G(z_n) \rangle$ can be computed by Wick's theorem. Note that there are two differences now: first, the *n* point correlator of *G* vanishes for odd *n*, so we only need to consider the case when n = 2M. Each loop must contain even number of vertices, otherwise the amplitude associated with the loop will vanish in the OPE. Second, for each contraction between pair of supercurrents, as the supercurrent Eq (3.22) has two terms, there are two propagators between each pair of vertices. We write them down explicity:

$$\langle : \partial \beta_{i_1}(z_1)c_{i_1}(z_1) :: b_{i_2}(z_2)\gamma_{i_2}(z_2) : \dots : \partial \beta_{i_{2M-1}}(z_{2M-1})c_{i_{2M-1}}(z_{2M-1})b_{i_{2M}}(z_{2M})\gamma_{i_{2M}}(z_{2M}) : \rangle,$$

$$\langle : b_{i_1}(z_1)\gamma_{i_1}(z_1) :: \partial \beta_{i_2}(z_2)c_{i_2}(z_2) : \dots : b_{i_{2M-1}}(z_{2M-1})\gamma_{i_{2M-1}}(z_{2M-1})\partial \beta_{i_{2M}}(z_{2M})c_{i_{2M}}(z_{2M}) : \rangle.$$
(3.27)

Figure 2 gives the corresponding Feynman diagram.

The amplitude associated with each Feynman diagram is

$$\operatorname{sign} \times g_{l_1}(z_{1_1}z_{1_2}\cdots z_{1_{2m_1}}) \times g_{l_2} \times \cdots \times g_{l_s}.$$
(3.28)

where sign is the signature of $\begin{pmatrix} 1 & 2 & \cdots & 2n \\ l_1 & l_2 & \cdots & l_s \end{pmatrix}$ which is a reflection of anti-communitivity of *bc* fields. And

$$g_{l_i}(z_{i_1}\cdots z_{i_{2m_i}}) = \frac{N}{(z_{i_1}-z_{i_2})(z_{i_2}-z_{i_3})^2\cdots(z_{i_{2m_i-1}}-z_{i_{2m_i}})(z_{i_{2m_i}}-z_{i_1})^2} - \frac{N}{(z_{i_1}-z_{i_2})^2(z_{i_2}-z_{i_3})\cdots(z_{i_{2m_i-1}}-z_{i_{2m_i}})^2(z_{i_{2m_i}}-z_{i_1})}.$$
(3.29)



Figure 2. Orange line represents contraction between fields b, c, blue line represents contraction between fields $\partial\beta, \gamma$. l_i labels a loop. For each loop, we add two graphs because they have the same underlying graph but different propagators. By Wick's theorem, summing over all the possible products of loops gives the 2n point amplitude of supercurrent operators.

where the factor of N is a result of summing over all ghost fields. Since c = 3N in this free field realization, $\langle G(z_1)G(z_2)...G(z_{2n})\rangle_{c=3N} = H(c = 3N, z_1, z_2, ..., z_{2n})$ holds for infitely many $N \in \mathbb{Z}_{>0}$, thus the theorem holds by the **Lemma 3.1.2**.

Below we give an equivalent way of writing the 2n points correlators of supercurrent:

Theorem 3.

$$\langle G(z_1)\cdots G(z_{2n})\rangle = \sum_{\substack{I_1\cup I_2\cdots \cup I_s=(12\cdots 2n)\\cluster\ decomposition}}} \left(\frac{2c}{3}\right)^s \times \operatorname{sign}\begin{pmatrix} 1 & 2 & \cdots & 2n\\I_1 & I_2 & \cdots & I_s \end{pmatrix} \prod_{i=1}^s \hat{g}_{I_i}$$
(3.30)

where each cluster has $|I_i| = 2M_i$ elements, $n = M_1 + M_2 + \cdots + M_s$. Moreover, we have

$$\hat{g}_{I_i}(z_{i_1}, z_{i_2}, \cdots, z_{i_{2M_i}}) = \frac{1}{2M_i} \sum_{\sigma \in S_{2M_i}} \frac{\operatorname{sign}(\sigma)}{(z_{\sigma(i_1)} - z_{\sigma(i_2)})(z_{\sigma(i_2)} - z_{\sigma(i_3)})^2 \cdots (z_{\sigma(i_{2M_i-1})} - z_{\sigma(i_{2M_i})})(z_{\sigma(i_{2M_i})} - z_{\sigma(i_1)})^2},$$
(3.31)

where $(z_{i_1}, z_{i_2}, ..., z_{i_{2M_i}}) \in I_i$.

Proof. This is a direct result from Eq (3.29). Note that the first and second term of Eq (3.29) are related via a permutation σ from $(i_1, i_2, \dots, i_{2M_i})$ to $(i_2, i_3, \dots, i_{2M_i}, i_1)$.

Every I_i is ordered compared with l_i , for example: if $l_i = (3, 1, 2, 4)$, then $I_i = (1, 2, 3, 4)$. Their sign are related:

$$\operatorname{sign}\begin{pmatrix} 1 & 2 & \cdots & 2n \\ l_1 & l_2 & \cdots & l_s \end{pmatrix} = \operatorname{sign}\begin{pmatrix} 1 & 2 & \cdots & 2n \\ I_1 & I_2 & \cdots & I_s \end{pmatrix} \times \prod_{\substack{\sigma(i) \in S_{2M_i} \\ 1 \le i \le s}} \operatorname{sign}(\sigma(i))$$
(3.32)

The factor of $\frac{1}{2M_i}$ is because that the Feynman diagram corresponding to each cluster I_i is invariant under M_i elements in permutation group S_{2M_i} , where each element $\sigma_{j=1,2,...,M_i}$ is:

$$\sigma_j(z_k) = z_{2M_i - 2j + k}, \quad \text{for} \quad k = 1, 2, \dots 2j$$

$$\sigma_j(z_k) = z_{k-2j}, \quad \text{for} \quad k = 2j + 1, \dots, 2M_i.$$
(3.33)

Note here that for supercurrent, the number of vertices in each cluster $|I_i|(i = 1, 2, ..., s)$ must be even. However, for stress-energy tensor, $|I_i|$ could be even or odd.

3.2 Clustering behaviours of the wavefunction

From the results in previous section, we now explore how the structure of supercurrent correlation functions affect clustering behaviours of their fractional quantum Hall wavefunctions. Without loss of generality, let us set p = 3, q = 1 for U(1) charge sector such that wavefunctions are symmetric polynomials.

Corollary 1. Let $1 \le k \le n$, then

$$\lim_{z_{2i-1}\to z_{2i}} \prod_{i=1}^{k} (z_{2i-1} - z_{2i})^3 \langle G(z_1) \cdots G(z_{2n}) \rangle = \left(\frac{2c}{3}\right)^k \langle G(z_{2k+1}) \cdots G(z_{2n}) \rangle$$
(3.34)

Proof. If z_{2i-1} and z_{2i} are not in the same cluster, then Eq (3.34) will vanish when we take the limit $z_{2i-1} \rightarrow z_{2i}$. If z_{2i-1} and z_{2i} are in the same cluster I_i , and the length of I_i is more than 2 (i.e, at least 4 vertices in this group), the the terms related to $(z_{2i-1} - z_{2i})$ are either proportional to $\frac{1}{z_{2i-1}-z_{2i}}$ or $\frac{1}{(z_{2i-1}-z_{2i})^2}$, which also vanishes in the limit when $z_{2i-1} \rightarrow z_{2i}$. The only non zero contribution is if there exist a $I_i = (z_{2i-1}, z_{2i})$, in this case:

$$(z_{2i-1} - z_{2i})^3 g_{I_i}(z_{2i-1}, z_{2i}) = (z_{2i-1} - z_{2i})^3 \frac{\frac{2c}{3}}{(z_{2i-1} - z_{2i})^3} = \frac{2c}{3},$$
(3.35)

Therefore, the only non zero term in this limit is the following cluster decomposition:

$$I_1 \cup I_2 \cup \dots \cup I_k \cup I_{k+1} \cup \dots \cup I_s, \tag{3.36}$$

where for $1 \le i \le k$, $I_i = (z_{\sigma(2i-1)}, z_{\sigma(2i)}), \sigma \in S_{2k}$. Then the conclusion follows from the definition of \hat{g}_I .

Corollary 2. Let

$$\phi(z_1, \cdots, z_{2n}) = \prod_{1 \le i < j \le 2n} (z_i - z_j)^3 \langle G(z_1) \cdots G(z_{2n}) \rangle,$$
(3.37)

then

$$\phi = \sum_{s=1}^{n} \left(\frac{2c}{3}\right)^{s} Q_{2n}^{s}(z_1, \cdots, z_{2n}), \tag{3.38}$$

where $Q_{2n}^s(z_1, \cdots, z_{2n})$ are symmetric polynomials determined by

$$Q_{2n}^{1} = \frac{1}{2n} \sum_{\sigma \in S_{2n}} \frac{\prod_{1 \le i < j \le 2n} (z_{\sigma(i)} - z_{\sigma(j)})^{3}}{(z_{\sigma(1)} - z_{\sigma(2)})(z_{\sigma(2)} - z_{\sigma(3)})^{2} \cdots (z_{\sigma(2n-1)} - z_{\sigma(2n)})(z_{\sigma(2n)} - z_{\sigma(1)})^{2}}, \quad (3.39)$$

and

$$Q_{2n}^{s} = \sum_{\substack{I_1 \cup I_2 \dots \cup I_s = (12\dots 2n)\\ cluster \, decomposition, |I_i| = 2m_i}} \prod_{i=1}^{s} Q_{2m_i}^1 \prod_{\substack{j \in I_r, k \in I_t\\r < t}} (z_j - z_k)^3.$$
(3.40)

Proof. Following Eq (3.30), Eq (3.37) and Eq (3.38), we have:

$$\sum_{s=1}^{n} \left(\frac{2c}{3}\right)^{s} Q_{2n}^{s}(z_{1}, \cdots, z_{2n}) = \prod_{1 \le i < j \le 2n} (z_{i} - z_{j})^{3} \sum_{\substack{I_{1} \cup I_{2} \cdots \cup I_{s} = (12 \cdots 2n) \\ \text{cluster decomposition}}} \left(\frac{2c}{3}\right)^{s} \times \operatorname{sign} \left(\frac{1}{I_{1}} \frac{2}{I_{2}} \cdots \frac{2n}{I_{s}}\right) \prod_{i=1}^{s} \hat{g}_{I_{i}}.$$
(3.41)

Comparing the polynomial expansion of ϕ and G gives Eq (3.39) and Eq (3.40). For s=1, it gives:

$$Q_{2n}^{1} = \prod_{1 \le i < j \le 2n} (z_{i} - z_{j})^{3} \hat{g}_{I_{1}}$$

$$= \frac{1}{2n} \sum_{\sigma \in S_{2n}} \frac{\operatorname{sign}(\sigma) \prod_{1 \le i < j \le 2n} (z_{i} - z_{j})^{3}}{(z_{\sigma(i_{1})} - z_{\sigma(i_{2})})(z_{\sigma(i_{2})} - z_{\sigma(i_{3})})^{2} \cdots (z_{\sigma(i_{2n-1})} - z_{\sigma(i_{2n})})(z_{\sigma(i_{2n})} - z_{\sigma(i_{1})})^{2}}$$

$$= \frac{1}{2n} \sum_{\sigma \in S_{2n}} \frac{\prod_{1 \le i < j \le 2n} (z_{\sigma(i_{1})} - z_{\sigma(i_{2})})(z_{\sigma(i_{2})} - z_{\sigma(i_{3})})^{2} \cdots (z_{\sigma(i_{2n-1})} - z_{\sigma(i_{2n})})(z_{\sigma(i_{2n})} - z_{\sigma(i_{1})})^{2}}{(z_{\sigma(i_{1})} - z_{\sigma(i_{2})})(z_{\sigma(i_{2})} - z_{\sigma(i_{3})})^{2} \cdots (z_{\sigma(i_{2n-1})} - z_{\sigma(i_{2n})})(z_{\sigma(i_{2n})} - z_{\sigma(i_{1})})^{2}},$$
(3.42)

where we use the identity:

$$\prod_{1 \le i < j \le 2n} (z_{\sigma(i)} - z_{\sigma(j)})^3 = \operatorname{sign}(\sigma) \prod_{1 \le i < j \le 2n} (z_i - z_j)^3.$$
(3.43)

For $2 \le s \le n$, we have:

$$Q_{2n}^{s} = \prod_{1 \le i < j \le 2n} (z_{i} - z_{j})^{3} \sum_{\substack{I_{1} \cup I_{2} \cdots \cup I_{s} = (12 \cdots 2n) \\ \text{cluster decomposition}, |I_{i}| = 2m_{i}}} \operatorname{sign} \begin{pmatrix} 1 & 2 & \cdots & 2n \\ I_{1} & I_{2} & \cdots & I_{s} \end{pmatrix} \prod_{i=1}^{s} \hat{g}_{I_{i}}$$

$$= \sum_{\substack{I_{1} \cup I_{2} \cdots \cup I_{s} = (12 \cdots 2n) \\ \text{cluster decomposition}, |I_{i}| = 2m_{i}}} \prod_{i=1}^{s} Q_{2m_{i}}^{1} \prod_{\substack{j \in I_{r}, k \in I_{t} \\ 1 \le r < t \le s}} (z_{j} - z_{k})^{3},$$
(3.44)

where the following identity is used:

$$\prod_{1 \le i < j \le 2n} (z_i - z_j)^3 = \prod_{\substack{a_i < b_i z_{a_i}, z_{b_i} \in I_i \\ 1 \le i \le s}} (z_{a_i} - z_{b_i})^3 \prod_{\substack{j \in I_r, k \in I_t \\ j < k}} (z_j - z_k)^3$$

$$= \prod_{\substack{a_i < b_i z_{a_i}, z_{b_i} \in I_i \\ 1 \le i \le s}} (z_{a_i} - z_{b_i})^3 \operatorname{sign} \begin{pmatrix} 1 & 2 & \cdots & 2n \\ I_1 & I_2 & \cdots & I_s \end{pmatrix} \prod_{\substack{j \in I_r, k \in I_t \\ 1 \le r < t \le s}} (z_j - z_k)^3$$
(3.45)

and

$$Q_{2m_i}^1 = \prod_{a_i < b_i, a_i, b_i \in I_i} (z_{a_i} - z_{b_i})^3 \hat{g}_{I_i}$$
(3.46)

It is easy to deduce from Corollary 1 and Eq (3.37) that

$$\phi(z_1 = z_2 = Z_1, \cdots, z_{2l-1} = z_{2l} = Z_l, \cdots)$$

$$= \left(\frac{2c}{3}\right)^l \prod_{1 \le j < k \le l} (Z_j - Z_k)^{12} \prod_{j=1}^l \prod_{k=2l+1}^{2n} (Z_j - z_k)^6 \phi(z_{2l+1}, \cdots, z_{2n}),$$
(3.47)

then do similar comparasion of polynomial coefficient of Eq (3.38) shows that Q_{2n}^s satisfy the following clustering properties:

$$Q_{2n}^{s}(z_{1} = z_{2} = Z_{1}, \cdots, z_{2l-1} = z_{2l} = Z_{l}, \cdots)$$

$$= \prod_{1 \le j < k \le l} (Z_{j} - Z_{k})^{12} \prod_{j=1}^{l} \prod_{k=2l+1}^{2n} (Z_{j} - z_{k})^{6} Q_{2n-2l}^{s-l}(z_{2l+1}, \cdots, z_{2n}).$$
(3.48)

Following [10, 20], the clustering properties of wave function Q_{2n}^s is a reflection of \mathbb{Z}^s symmetry. Note that Eq (3.48) is invariant under permutations, namely:

$$Q_{2n}^{s}(z_{1} = z_{2} = Z_{1}, \cdots, z_{2l-1} = z_{2l} = Z_{l}, z_{2l+1}, \dots, z_{2n})$$

= $Q_{2n}^{s}(z_{\sigma(1)} = z_{\sigma(2)} = Z_{1}, \cdots, z_{\sigma(2l-1)} = z_{\sigma(2l)} = Z_{l}, z_{\sigma(2l+1)}, \dots, z_{\sigma(2n)}).$ (3.49)

which corresponds to the clustering decomposition:

$$I_1 \cup I_2 \cup \dots \cup I_l \cup I_{l+1} \cup \dots \cup I_s, \tag{3.50}$$

where for $1 \le j \le l$, $I_j = (z_{\sigma(2j-1)} = Z_j, z_{\sigma(2j)} = Z_j)$, $\sigma \in S_l$. For $1 \le s \le n-1$, one can prove that $l \le s-1$ must hold by using proof by contradiction. If l = s, then the clustering decomposition that make Q_{2n}^s non zero is:

$$I_1 \cup I_2 \cup \dots \cup I_s = (1, 2, \dots 2n),$$
 (3.51)

the length of each I_i is 2, however, since $s \le n - 1$, the total length of this clustering decomposition is $2s \le 2n - 2 < 2n$, therefore for $2 \le s \le n - 1$, $1 \le l \le s - 1$. For s = n, the clustering decomposition that make Q_{2n}^n non zero is:

$$I_1 \cup I_2 \cup \dots \cup I_n = (1, 2, \dots, 2n), \tag{3.52}$$

the length of each I_i is 2, so in this case, $1 \le l \le n$. Mathematically, each $Q_{2n}^s(z_1, \dots, z_{2n})$ is a symmetric polynomial with 2n coordinates such that whenever k + 1 = 3 particles coincide, the polynomial vanishes as power r = 6, which is a direct result from Eq (3.48). Individually, for $1 \le s \le n - 1$, $Q_{2n}^s(z_1, \dots, z_{2n})$ vanishes whenever s clusters of two particles coincide but not vanishing if only s - 1 clusters of two particles coincide as shown in Eq.(3.48). For s = n, Q_{2n}^n will be non vanishing even for n clusters of two particles coincide.

Below we give some simple examples of these symmetric polynomials:

Example 3.2.1. Let $z_{ij} = z_i - z_j$, then

$$Q_{2}^{1} = 1,$$

$$Q_{4}^{1} = 3z_{12}^{2}z_{13}^{2}z_{14}^{2}z_{23}^{2}z_{24}^{2}z_{34}^{2},$$

$$Q_{6}^{2} = 3\left(\prod_{a < b}^{6} z_{ab}^{2}\right)\left(\sum_{i < j}^{6} \frac{1}{z_{ij}^{2}} \prod_{k \neq i,j}^{6} z_{ik}z_{jk}\right).$$
(3.53)

Corollary 3. $Q_4^1 = 3z_{12}^2 z_{13}^2 z_{14}^2 z_{23}^2 z_{24}^2 z_{34}^2$.

Proof. Following Eq(3.39), we have:

$$Q_4^1 = \frac{1}{4} \prod_{i < j}^4 z_{ij}^2 \sum_{\sigma} \frac{z_{\sigma(1)\sigma(3)} z_{\sigma(2)\sigma(4)}}{z_{\sigma(1)\sigma(4)} z_{\sigma(2)\sigma(3)}}$$
(3.54)

For the cross ratio, take $x = \frac{z_{13}z_{24}}{z_{14}z_{23}}$ as an example, one can prove that the cross ratio over the 4! = 24 permutations have 6 distinct value:

$$x, \quad 1-x, \quad \frac{1}{x}, \quad 1-\frac{1}{x}, \quad \frac{1}{1-x}, \quad \frac{x}{x-1}$$
 (3.55)

The stabilizer for the cross ratio $\frac{z_{\sigma(1)\sigma(3)}z_{\sigma(2)\sigma(4)}}{z_{\sigma(1)\sigma(4)}z_{\sigma(2)\sigma(3)}}$ is:

$$\begin{aligned} & (\sigma(1)\sigma(2)\sigma(3)\sigma(4)), & (\sigma(2)\sigma(1)\sigma(4)\sigma(3)) \\ & (\sigma(4)\sigma(3)\sigma(2)\sigma(1)), & (\sigma(3)\sigma(4)\sigma(1)\sigma(2)) \end{aligned}$$
(3.56)

Therefore:

$$\sum_{\sigma} \frac{z_{\sigma(1)\sigma(3)} z_{\sigma(2)\sigma(4)}}{z_{\sigma(1)\sigma(4)} z_{\sigma(2)\sigma(3)}}$$

= 4(x + 1 - x + $\frac{1}{x}$ + 1 - $\frac{1}{x}$ + $\frac{1}{1 - x}$ + $\frac{x}{x - 1}$)
= 12. (3.57)

which implies that:

$$Q_4^1 = 3z_{12}^2 z_{13}^2 z_{14}^2 z_{23}^2 z_{24}^2 z_{34}^2. aga{3.58}$$

Corollary 4. $Q_{2n}^n = \operatorname{Pf}\left(z_{ij}^{-3}\right) \prod_{i < j}^{2n} z_{ij}^3.$

Proof. For a skew-symmetric matrix A, Pfaffian of A satisfies $Pf(A)^2 = det(A)$. Let $A = (a_{ij})$ be a $2n \times 2n$ skew-symmetric matrix, the explicit expression of its Pfaffian is:

$$Pf(A) = \frac{1}{2^n n!} \sum_{\sigma \in S_{2n}} sign(\sigma) \prod_{i=1}^n a_{\sigma(2n-1), \sigma(2n)}.$$
(3.59)

which leads to the expression of $Pf\left(z_{ij}^{-3}\right)$:

$$Pf\left(z_{ij}^{-3}\right) = \frac{1}{2^n n!} \sum_{\sigma \in S_{2n}} \operatorname{sign}(\sigma) \prod_{i=1}^n z_{\sigma(2i-1)\sigma(2i)}^{-3}.$$
(3.60)

Following Eq (3.40), for Q_{2n}^n , the cluster decomposition is given by:

$$I_1 \cup I_2 \cup \dots \cup I_n = (1, 2, \dots, 2n)$$
(3.61)

Since the length of each I_i is at least 2 and the total length is 2n. We have for $1 \le i \le n$, $|I_i| = 2$. Note that $Q_2^1 = 1$, by Eq (3.40) we have:

$$Q_{2n}^{n} = \sum_{\substack{I_{1} \cup I_{2} \cdots \cup I_{n} = (12 \cdots 2n) \\ \text{cluster decomposition,} |I_{i}| = 2}} \prod_{i=1}^{n} Q_{I_{i}}^{1} \prod_{\substack{j \in I_{r}, k \in I_{t} \\ r < t}} (z_{j} - z_{k})^{3}$$

$$= \sum_{\substack{I_{1} \cup I_{2} \cdots \cup I_{n} = (12 \cdots 2n) \\ \text{cluster decomposition,} |I_{i}| = 2}} \prod_{\substack{j \in I_{r}, k \in I_{t} \\ r < t}} (z_{j} - z_{k})^{3}.$$

$$= \sum_{\substack{I_{1} \cup I_{2} \cdots \cup I_{n} = (12 \cdots 2n) \\ \text{cluster decomposition,} |I_{i}| = 2}} \prod_{\substack{i < j \\ r < t}} \sum_{\substack{I_{i} \in I_{r} \\ r = 1, 2, \dots, n}} (z_{lk})^{-3}$$

$$= \left(\sum_{\substack{I_{1} \cup I_{2} \cdots \cup I_{n} = (12 \cdots 2n) \\ \text{cluster decomposition,} |I_{i}| = 2}} \prod_{\substack{I, k \in I_{r} \\ r = 1, 2, \dots, n}} (z_{lk})^{-3}\right) \prod_{i < j}^{2n} z_{ij}^{3}$$

$$= \left(\sum_{\substack{I_{1} \cup I_{2} \cdots \cup I_{n} = (12 \cdots 2n) \\ \text{cluster decomposition,} |I_{i}| = 2}} \prod_{\substack{I, k \in I_{r} \\ r = 1, 2, \dots, n}} (z_{lk})^{-3}\right) \prod_{i < j}^{2n} z_{ij}^{3}$$

Following Eq (3.60), there is a subgroup of S_{2n} which leaves the expression sign(σ) $\prod_{i=1}^{n} z_{\sigma(2i-1)\sigma(2i)}^{-3}$ invariant. For example, if we take the cluster decomposition to be:

$$I_i = (z_{2i-1}, z_{2i}) \quad i = 1, 2, ..., n$$
(3.63)

The following equation holds for a subgroup of S_{2n} :

$$\operatorname{sign}(\sigma) \prod_{i=1}^{n} z_{\sigma(2i-1)\sigma(2i)}^{-3} = \prod_{i=1}^{n} z_{(2i-1)(2i)}^{-3},$$
(3.64)

where the generators of this subgroup are given by:

$$a_{i} = \sigma(z_{2i-1}, z_{2i}) = (z_{2i}, z_{2i-1}) \quad i = 1, 2, ..., n$$

$$b_{j} = \sigma(z_{1}, z_{2}, z_{2j-1}, z_{2j}) = (z_{2j-1}, z_{2j}, ..., z_{1}, z_{2}) \quad j = 1, 2, ..., n - 1,$$
(3.65)

One can check that the order of this subgroup is $2^n n!$. Therefore:

$$Pf\left(z_{ij}^{-3}\right) = \frac{1}{2^{n}n!} \sum_{\sigma \in S_{2n}} sign(\sigma) \prod_{i=1}^{n} z_{\sigma(2i-1)\sigma(2i)}^{-3}$$

$$= \left(\sum_{\substack{I_1 \cup I_2 \dots \cup I_n = (12\dots 2n) \\ cluster \, decomposition, |I_i| = 2}} \prod_{\substack{l,k \in I_r \\ r=1,2,\dots n}} (z_{lk})^{-3}\right).$$
(3.66)

thus the Corollary holds by Eq (3.62).

There is an equivalent representation of the ground state wave function if we expand it with another set of symmetric polynomials with same clustering properties. A nice feature for choosing this new set of symmetric polynomials is that it makes direct connections with our limiting case when $m \to \infty$ and $c \to \frac{3}{2}$. One can check that, for the following symmetric polynomials:

$$P_{2n}^{s}(z_1, \cdots, z_{2n}) = \frac{1}{3} \sum_{j=1}^{s} Q_{2n}^{j}(z_1, \cdots, z_{2n}) \quad s = 1, 2, \dots n - 1,$$
(3.67)

$$P_{2n}^{n}(z_1,\cdots,z_{2n}) = \frac{1}{3} \operatorname{Pf}^3\left(z_{ij}^{-1}\right) \prod_{i< j}^{2n} z_{ij}^3.$$
(3.68)

We have:

Corollary 5.

$$\phi(z_1,\cdots,z_{2n}) = \left(\frac{2c}{3}\right)^n 3P_{2n}^n(z_1,\cdots,z_{2n}) + \sum_{s=1}^{n-1} \left(\frac{2c}{3}\right)^s (3-2c)P_{2n}^s(z_1,\cdots,z_{2n}), \quad (3.69)$$

where P_{2n}^s share the same clustering properties with Q_{2n}^s , namely,

$$P_{2n}^{s}(z_{1} = z_{2} = Z_{1}, \cdots, z_{2l-1} = z_{2l} = Z_{l}, \cdots)$$

$$= \prod_{1 \le j < k \le l} (Z_{j} - Z_{k})^{12} \prod_{j=1}^{l} \prod_{k=2l+1}^{2n} (Z_{j} - z_{k})^{6} P_{2n-2l}^{s-l}(z_{2l+1}, \cdots, z_{2n}).$$
(3.70)

Proof. First of all, we show that $\phi(z_1, \dots, z_{2n}) = Pf^3(z_{ij}^{-1}) \prod_{i < j}^{2n} z_{ij}^3$ when c = 3/2. Consider a CFT of three Majorana fermions ψ_1, ψ_2, ψ_3 with OPEs

$$\psi_a(z)\psi_b(w) = \frac{\delta_{ab}}{z-w} + \mathcal{O}(1).$$
(3.71)

This system has $\mathcal{N} = 1$ superconformal symmetry with c = 3/2, namely

$$T(z) = \frac{1}{2} \sum_{a=1}^{3} : \partial \psi_a(z) \psi_a(z) : \quad G(z) = i \psi_1(z) \psi_2(z) \psi_3(z)$$
(3.72)

satisfy the $\mathcal{N} = 1$ super Virasoro algebra. Since ψ_1, ψ_2, ψ_3 are independent, the correlator $\langle G(z_1) \cdots G(z_{2n}) \rangle$ is simply the product of three correlators $\langle \psi_1(z_1) \cdots \psi_1(z_{2n}) \rangle \langle \psi_2(z_1) \cdots \psi_2(z_{2n}) \rangle \langle \psi_3(z_1) \cdots \psi_3(z_{2n}) \rangle$, which can be easily computed:

$$\langle \psi_a(z_1) \cdots \psi_a(z_{2n}) \rangle = \Pr\left(z_{ij}^{-1}\right), \ a = 1, 2, 3.$$
 (3.73)

Thus $\phi(z_1, \dots, z_{2n}) = Pf^3\left(z_{ij}^{-1}\right) \prod_{i < j}^{2n} z_{ij}^3$ when c = 3/2. Finally, the clustering properties follow from the equation (3.47) and the definition of P_{2n}^s .

The dominant clustering behaviours of our wavefunctions for any 2n particles, vanishing whenever three particles coincide, is fixed by $Pf^3\left(z_{ij}^{-1}\right)\prod_{i< j}^{2n} z_{ij}^3$ which is associated to the three Majorana fermion unitary CFT. This is in contrast to previous cases studied where Jack polynomials associated with non-unitary CFTs plus "healing" polynomials are used for generating wavefunctions of unitary CFTs.

4 Discussion

In this paper, we achieved two things in our studies of $\mathcal{N} = 1$ SCFTs. We found explicit characterizations of our $\mathcal{N} = 1$ theories in terms of a parafermion theory, an Ising theory and a free boson theory by first pairing our $\mathcal{N} = 1$ theory with a parafermion theory. Supercurrent operator G can also be written as linear combinations of operators from its constituent CFTs. By utilizing free field methods, we worked out explicit ground state wavefunctions of fractional quantum Hall states based on our $\mathcal{N} = 1$ theories. We have also shown clustering properties of these ground state wavefunctions. Several questions remain. In our case, even though we have worked out an explicit decomposition of the supercurrent operator G into parafermion operators, Majorana fermion operators and U(1) vertex operators, we do not immediately see advantage of this decomposition in calculating correlation functions of G. Our preliminary calculations do not show much simplification from operator decomposition approach. On the other hand, we believe decomposition worked out here should be applicable to boundary conforma field theories with $\mathcal{N} = 1$ superconformal algebra [41–44]. Speaking of wavefunctions, previous studies of Jack polynomials and Read-Rezayi series of fractional quantum Hall states [11–13, 15, 16, 20, 45] have demonstarted that distinct clustering behaviours indicate distinct topological orders. In our case, at least for ground state, wavefunctions are the same in terms of their clustering behaviours. The only changing parameter is central charge c. There are two immediate questions: 1. How to understand different topological orders sharing the same clustering behaviours in their wavefunctions? Original studies relate different pseudo-potential [46] or Kivelson-Trugman type potential Hamiltonians [47] with Jack polynomial ground state wavefunctions with different clustering behaviours. 2. In thermodynamic limit in which $2n \to \infty$, each ground state wavefunction should be orthogonal to each other since they represent different topological orders. How to show or give evidences to such tendency? Finally, for all other fractional spin ($\Delta = \frac{N+1}{N}$) generalization of the minimal models, can we always use free field methods to work out their correlation functions?

Acknowledgments. We thank for the helpful discussions with Paul Fendley, Steven Simon, Taro Kimura, Davide Gaiotto, Nicolas Regnault, Benoit Estienne, Biao Lian, Prashant Kumar, Jie Wang, Ching Hung Lam and Chongying Dong. SN wants to specially thank Joseph Conlon for his help and encouragement during the work. This work is supported by the Alfred P. Sloan Foundation, and NSF through the Princeton University's Materials Research Science and Engineering Center DMR-2011750. Additional support was provided by the Gordon and Betty Moore Foundation through Grant GBMF8685 towards the Princeton theory program. SN wants to acknowledge funding support from the China Scholarship Council-FaZheng Group- University of Oxford. Kavli Institute for the Physics and Mathematics of the Universe is supported by World Premier International Research Center Initiative (WPI), MEXT, Japan. YZ would like to thank Perimeter Institute for Theoretical Physics, where part of YZ's work was done as a graduate student there.

A S_3 minimal model

 S_3 minimal models have central charge $c = 2\left(1 - \frac{12}{(m-2)(m+2)}\right)$ with $m = 5, 6, \cdots$. It has extended algebra beyond Virosora algebra which is generated by

$$T(z)T(w) = \frac{1}{(z-w)^4} \left\{ \frac{c}{2} + 2(z-w)^2 T(w) + (z-w)^3 \partial T(w) + \dots \right\}$$

$$T(z)G^{\pm}(w) = \frac{1}{(z-w)^2} \left\{ \frac{4}{3} G^{\pm}(w) + (z-w) \partial G^{\pm}(w) + \dots \right\},$$

$$G^{+}(z)G^{+}(w) = \frac{\lambda^{+}}{(z-w)^{4/3}} \left\{ G^{-}(w) + \frac{1}{2}(z-w) \partial G^{-}(w) + \dots \right\},$$

$$G^{-}(z)G^{-}(w) = \frac{\lambda^{-}}{(z-w)^{4/3}} \left\{ G^{+}(w) + \frac{1}{2}(z-w) \partial G^{+}(w) + \dots \right\},$$

$$G^{+}(z)G^{-}(w) = \frac{1}{(z-w)^{8/3}} \left\{ \frac{3c}{8} + (z-w)^2 T(w) + \dots \right\}.$$
(A.1)

where G^{\pm} are operators with scaling dimension $\Delta = \frac{4}{3}$. This series of conformal field theories include examples such as the \mathbb{Z}_6 parafermion CFT. Each member has a known coset construction

$$\frac{SU(2)_4 \times SU(2)_{m-4}}{SU(2)_m} \tag{A.2}$$

This series of cosets has the following relation

$$\frac{SU(2)_m}{U(1)_{2m}} \times \frac{SU(2)_4 \times SU(2)_{m-4}}{SU(2)_m} = \frac{SU(2)_{m-4}}{U(1)_{2(m-4)}} \times \frac{SU(2)_4}{U(1)_8} \times U(1)_{8m(m-4)}$$
(A.3)

at the level of stress-energy tensor and $\frac{SU(2)_m}{U(1)_{2m}}$ is the coset for \mathbb{Z}_m parafermion. So this means we can combine a \mathbb{Z}_m parafermion theory with the *m*th member of the S_3 minimial model theory and turn them into a combination of a \mathbb{Z}_{m-4} parafermion theory together with a U(1) theory and a \mathbb{Z}_4 parafermion theory.

Levels for U(1) theories need a little bit more explanations. Let us reshuffle U(1) theories as following:

Left
$$(U(1)_{2m})^{-1} \times U(1)_{2(m-4)} \times U(1)_8$$
, Right $U(1)_{8m(m-4)}$. (A.4)

To establish an equivalence relation between the left and right hand side, we first write out their Lagrangian densities:

$$\mathcal{L}_{l} = \frac{1}{4\pi} \sum_{I,J=1}^{3} K_{l}^{IJ} \partial_{x} \phi_{I} (\partial_{x} + \partial_{t}) \phi_{J}, \quad \mathcal{L}_{r} = \frac{1}{4\pi} K_{r} \partial_{x} \phi_{1} (\partial_{x} + \partial_{t}) \phi_{1}$$
(A.5)

where $K_l = \begin{pmatrix} -2m & 0 & 0 \\ 0 & 2(m-4) & 0 \\ 0 & 0 & 8 \end{pmatrix}$ and $K_r = 8m(m-4)$. It is easy to see that there exists a matrix

M with integer entries

$$M = \begin{pmatrix} 4 - m - m \ 0 \\ 1 & 1 & 1 \\ -1 & -1 & 1 \end{pmatrix}, \quad \det M = 8 > 0$$
(A.6)

such that

$$M^{T}\begin{pmatrix} -2m & 0 & 0\\ 0 & 2(m-4) & 0\\ 0 & 0 & 8 \end{pmatrix})M = \left(8m(m-4)\right) \bigoplus 16\sigma_{z}.$$
 (A.7)

The right hand side of Eq. (A.7) is equivalent to \mathcal{L}_r as we can add a backscattering term to the two counter-propagating modes

$$2\cos(2(\phi_2 - \phi_3))$$
 (A.8)

to gap out $16\sigma_z$ degrees of freedom (ϕ_2 and ϕ_3)[32, 33]. We have established an equivalence relation in Eq. (A.4).

Wavefunction comparison B

In [13], the author gives a formula of the 2n point correlators of $\mathcal{N} = 1$ superconformal currents. In this section, we compare with his results and give a proof on the equivalence for n = 2 and n = 3.

$$\phi_{2n} = \frac{\left(\frac{c}{3}\right)^{\frac{n}{2}(3-n)}}{n!} \sum_{\sigma \in S_{2n}} \prod_{1 \le r < s \le n} \chi(z_{\sigma(2r-1)}, z_{\sigma(2r)}; z_{\sigma(2s-1)}, z_{\sigma(2s)}), \tag{B.1}$$

where the function χ is:

$$\chi(z_1, z_2; z_3, z_4) = z_{13}^3 z_{14}^3 z_{23}^3 z_{24}^3 \left(\frac{c}{3} + \frac{z_{12} z_{34}}{z_{14} z_{23}}\right).$$
(B.2)

For n = 2, we have:

$$\phi_{4} = \frac{c}{6} \sum_{P \in S_{4}} \chi(z_{P(1)}, z_{P(2)}; z_{P(3)}, z_{P(4)})$$

$$= \frac{c}{6} \sum_{\sigma \in S_{4}} z_{\sigma(1)\sigma(3)}^{3} z_{\sigma(1)\sigma(4)}^{3} z_{\sigma(2)\sigma(3)}^{3} z_{\sigma(2)\sigma(4)}^{3} \left(\frac{c}{3} + \frac{z_{\sigma(1)\sigma(2)} z_{\sigma(3)\sigma(4)}}{z_{\sigma(1)\sigma(4)} z_{\sigma(2)\sigma(3)}}\right).$$
(B.3)

Note that:

$$z_{\sigma(1)\sigma(3)}^{3} z_{\sigma(1)\sigma(4)}^{3} z_{\sigma(2)\sigma(3)}^{3} z_{\sigma(2)\sigma(4)}^{3} \frac{z_{\sigma(1)\sigma(2)} z_{\sigma(3)\sigma(4)}}{z_{\sigma(1)\sigma(4)} z_{\sigma(2)\sigma(3)}} = \prod_{1 \le i < j \le 4} z_{ij}^{2} \frac{z_{\sigma(1)\sigma(3)} z_{\sigma(2)\sigma(4)}}{z_{\sigma(1)\sigma(2)} z_{\sigma(3)\sigma(4)}}$$
(B.4)

The sum is over all elements in S_4 . $\prod_{1 \le i < j \le 4} z_{ij}^2$ is invariant under permutation. If we permute $\sigma(2)$ and $\sigma(4)$ with each other, the expression reads:

$$\frac{c}{6} \prod_{1 \le i < j \le 4} z_{ij}^2 \frac{z_{\sigma(1)\sigma(3)} z_{\sigma(2)\sigma(4)}}{z_{\sigma(1)\sigma(4)} z_{\sigma(2)\sigma(3)}} = \frac{2c}{3} Q_4^1$$
(B.5)

where we use Eq (3.54). One can also check that:

$$\frac{c^{2}}{18} \sum_{\sigma \in S_{4}} z^{3}_{\sigma(1)\sigma(3)} z^{3}_{\sigma(1)\sigma(4)} z^{3}_{\sigma(2)\sigma(3)} z^{3}_{\sigma(2)\sigma(4)}
= \frac{c^{2}}{18} \times 8 \times (z^{3}_{13} z^{3}_{14} z^{3}_{23} z^{3}_{24} - z^{3}_{12} z^{3}_{14} z^{3}_{23} z^{3}_{34} + z^{3}_{12} z^{3}_{13} z^{3}_{24} z^{3}_{34})
= (\frac{2c}{3})^{2} Q^{2}_{4},$$
(B.6)

where we use Corollary 4. So $\phi_4(z_1, z_2, z_3, z_4) = (\frac{2c}{3})^2 Q_4^2 + \frac{2c}{3} Q_4^1$.

For n = 3, we have:

$$\begin{split} \phi_{6} &= \frac{1}{6} \sum_{\sigma \in S_{6}} \prod_{1 \leq r < s \leq 6} \chi(z_{\sigma(2r-1)}, z_{\sigma(2r)}; z_{\sigma(2s-1)}, z_{\sigma(2s)}) \\ &= \frac{1}{6} \sum_{\sigma \in S_{6}} z_{\sigma(1)\sigma(3)}^{3} z_{\sigma(1)\sigma(4)}^{3} z_{\sigma(2)\sigma(3)}^{3} z_{\sigma(2)\sigma(4)}^{3} (\frac{c}{3} + \frac{z_{\sigma(1)\sigma(2)} z_{\sigma(3)\sigma(4)}}{z_{\sigma(1)\sigma(4)} z_{\sigma(2)\sigma(3)}}) z_{\sigma(1)\sigma(5)}^{3} z_{\sigma(1)\sigma(6)}^{3} z_{\sigma(2)\sigma(5)}^{3} z_{\sigma(2)\sigma(6)}^{3} \\ &(\frac{c}{3} + \frac{z_{\sigma(1)\sigma(2)} z_{\sigma(5)\sigma(6)}}{z_{\sigma(1)\sigma(6)} z_{\sigma(2)\sigma(5)}}) z_{\sigma(3)\sigma(5)}^{3} z_{\sigma(3)\sigma(6)}^{3} z_{\sigma(4)\sigma(5)}^{3} z_{\sigma(4)\sigma(6)}^{3} (\frac{c}{3} + \frac{z_{\sigma(3)\sigma(4)} z_{\sigma(5)\sigma(6)}}{z_{\sigma(3)\sigma(6)} z_{\sigma(4)\sigma(5)}}) \end{split}$$
(B.7)

For term relating c^3 , it is easy to prove that it is equivalent with Q_6^3 . So we start from term that proportional to c^2 :

$$\begin{aligned} &\frac{1}{6} \frac{c^2}{9} \sum_{\sigma \in S_6} \prod_{1 \le i < j \le 6} z_{ij}^2 \left(\frac{z_{\sigma(1)\sigma(3)} z_{\sigma(2)\sigma(4)}}{z_{\sigma(1)\sigma(2)} z_{\sigma(3)\sigma(4)}} \frac{\prod_{k \ne 5,6} z_{\sigma(k)\sigma(5)} z_{\sigma(k)\sigma(6)}}{z_{\sigma(5)\sigma(6)}^2} \right. \\ &+ \frac{z_{\sigma(1)\sigma(5)} z_{\sigma(2)\sigma(6)}}{z_{\sigma(1)\sigma(2)} z_{\sigma(5)\sigma(6)}} \frac{\prod_{k \ne 3,4} z_{\sigma(k)\sigma(3)} z_{\sigma(k)\sigma(4)}}{z_{\sigma(3)\sigma(4)}^2} + \frac{z_{\sigma(3)\sigma(5)} z_{\sigma(4)\sigma(6)}}{z_{\sigma(3)\sigma(4)} z_{\sigma(5)\sigma(6)}} \frac{\prod_{k \ne 1,2} z_{\sigma(k)\sigma(1)} z_{\sigma(k)\sigma(2)}}{z_{\sigma(1)\sigma(2)}^2} \right) \\ &= \frac{4c^2}{3} \prod_{1 \le a < b \le 6} z_{ab}^2 \left(\sum_{i < j} \frac{1}{z_{ij}^2} \prod_{k \ne i,j}^6 z_{ik} z_{jk} \right) \\ &= \left(\frac{2c}{3} \right)^2 Q_6^2 \end{aligned} \tag{B.8}$$

This is because, consider the summation over S_6 for the following term:

$$\frac{c^2}{54} \sum_{\sigma \in S_6} \prod_{1 \le i < j \le 6} z_{ij}^2 \frac{z_{\sigma(1)\sigma(3)} z_{\sigma(2)\sigma(4)}}{z_{\sigma(1)\sigma(2)} z_{\sigma(3)\sigma(4)}} \frac{\prod_{k \ne 5,6} z_{\sigma(k)\sigma(5)} z_{\sigma(k)\sigma(6)}}{z_{\sigma(5)\sigma(6)}^2}$$
(B.9)

For a given σ , this term is invariant under the permutation between $\sigma(5)$ and $\sigma(6)$, so summation over this subgroup gives a factor of 2. Also if we restrict to a subgroup S_4 of S_6 , which is the permutation between $\sigma(1), \sigma(2), \sigma(3), \sigma(4)$, then the summation over this subgroup gives a factor of 12. Since:

$$6! = 720 = 2 \times 24 \times 15 = 2 \times 24 \times C_6^2 \tag{B.10}$$

So this summation equals:

$$2 \times 12 \times \frac{c^2}{54} \prod_{1 \le a < b \le 6} z_{ab}^2 \left(\sum_{i < j}^6 \frac{1}{z_{ij}^2} \prod_{k \ne i,j}^6 z_{ik} z_{jk} \right)$$

$$= \frac{4c^2}{9} \prod_{1 \le a < b \le 6} z_{ab}^2 \left(\sum_{i < j}^6 \frac{1}{z_{ij}^2} \prod_{k \ne i,j}^6 z_{ik} z_{jk} \right)$$
(B.11)

The summation result is the same for other two terms, so the final result is:

$$\frac{4c^2}{3} \prod_{1 \le a < b \le 6} z_{ab}^2 \left(\sum_{i < j}^6 \frac{1}{z_{ij}^2} \prod_{k \ne i,j}^6 z_{ik} z_{jk} \right)$$
(B.12)

For term that proportional to *c*, the result is:

$$\frac{1}{6} \frac{c}{3} \sum_{\sigma \in S_{6}} \prod_{1 \leq i < j \leq 6} z_{ij}^{2} \left(\frac{z_{\sigma(1)\sigma(3)} z_{\sigma(2)\sigma(4)}}{z_{\sigma(1)\sigma(2)} z_{\sigma(3)\sigma(4)}} \frac{z_{\sigma(1)\sigma(2)} z_{\sigma(5)\sigma(6)}}{z_{\sigma(1)\sigma(6)} z_{\sigma(2)\sigma(5)}} \frac{\prod_{k \neq 5,6} z_{\sigma(k)\sigma(5)} z_{\sigma(k)\sigma(6)}}{z_{\sigma(5)\sigma(6)}^{2}} + \frac{z_{\sigma(1)\sigma(5)} z_{\sigma(2)\sigma(6)}}{z_{\sigma(3)\sigma(6)} z_{\sigma(3)\sigma(4)} z_{\sigma(5)\sigma(6)}} \frac{\prod_{k \neq 3,4} z_{\sigma(k)\sigma(3)} z_{\sigma(k)\sigma(4)}}{z_{\sigma(3)\sigma(4)}^{2}} + \frac{z_{\sigma(3)\sigma(5)} z_{\sigma(4)\sigma(6)}}{z_{\sigma(3)\sigma(4)} z_{\sigma(5)\sigma(6)}} \frac{z_{\sigma(1)\sigma(2)} z_{\sigma(3)\sigma(4)}}{z_{\sigma(1)\sigma(4)} z_{\sigma(2)\sigma(3)}} \frac{\prod_{k \neq 1,2} z_{\sigma(k)\sigma(1)} z_{\sigma(k)\sigma(2)}}{z_{\sigma(1)\sigma(2)}^{2}} \right)$$
(B.13)

Consider the term individually:

$$\frac{1}{6} \frac{c}{3} \sum_{\sigma \in S_6} \prod_{1 \le i < j \le 6} z_{ij}^2 \left(\frac{z_{\sigma(1)\sigma(3)} z_{\sigma(2)\sigma(4)}}{z_{\sigma(1)\sigma(2)} z_{\sigma(3)\sigma(4)}} \frac{z_{\sigma(1)\sigma(2)} z_{\sigma(5)\sigma(6)}}{z_{\sigma(1)\sigma(6)} z_{\sigma(2)\sigma(5)}} \frac{\prod_{k \ne 5,6} z_{\sigma(k)\sigma(5)} z_{\sigma(k)\sigma(6)}}{z_{\sigma(5)\sigma(6)}^2} \right)$$
(B.14)

For the permutation between $\sigma(5)$ and $\sigma(6)$, we have:

$$\frac{z_{\sigma(1)\sigma(2)} z_{\sigma(5)\sigma(6)}}{z_{\sigma(1)\sigma(6)} z_{\sigma(2)\sigma(5)}} + \frac{z_{\sigma(1)\sigma(2)} z_{\sigma(6)\sigma(5)}}{z_{\sigma(1)\sigma(5)} z_{\sigma(2)\sigma(6)}} \\
= \frac{z_{\sigma(1)\sigma(2)}^2 z_{\sigma(5)\sigma(6)}^2}{z_{\sigma(1)\sigma(5)} z_{\sigma(1)\sigma(6)} z_{\sigma(2)\sigma(5)} z_{\sigma(2)\sigma(6)}}$$
(B.15)

For the permutation between $\sigma(3)$ and $\sigma(4)$, we have:

$$\frac{z_{\sigma(1)\sigma(3)}z_{\sigma(2)\sigma(4)}}{z_{\sigma(1)\sigma(2)}z_{\sigma(3)\sigma(4)}} - \frac{z_{\sigma(1)\sigma(4)}z_{\sigma(2)\sigma(3)}}{z_{\sigma(1)\sigma(2)}z_{\sigma(3)\sigma(4)}}$$

$$= 1$$
(B.16)

For the permutation between $\sigma(1)$ and $\sigma(2)$, we have an extra factor of 2.

$$720 = 2 \times 2 \times 2 \times 90 = 2 \times 2 \times 2 \times C_6^2 \times C_4^2$$
 (B.17)

$$\frac{1}{6} \frac{c}{3} \sum_{\sigma \in S_6} \prod_{1 \le i < j \le 6} z_{ij}^2 \left(\frac{z_{\sigma(1)\sigma(3)} z_{\sigma(2)\sigma(4)}}{z_{\sigma(1)\sigma(2)} z_{\sigma(3)\sigma(4)}} \frac{z_{\sigma(1)\sigma(2)} z_{\sigma(5)\sigma(6)}}{z_{\sigma(1)\sigma(6)} z_{\sigma(2)\sigma(5)}} \frac{\prod_{k \ne 5,6} z_{\sigma(k)\sigma(5)} z_{\sigma(k)\sigma(6)}}{z_{\sigma(5)\sigma(6)}^2} \right) \\
= \frac{c}{9} \prod_{1 \le i < j \le 6} z_{ij}^2 \left(\sum_{a < b, m < n \ne a, b}^6 z_{ab}^2 \prod_{k \ne a, b, m, n}^6 z_{km} z_{kn} \right)$$
(B.18)

The result is the same for other two term, so the final result is:

$$\frac{1}{6} \frac{c}{3} \sum_{\sigma \in S_{6}} \prod_{1 \leq i < j \leq 6} z_{ij}^{2} \left(\frac{z_{\sigma(1)\sigma(3)} z_{\sigma(2)\sigma(4)}}{z_{\sigma(1)\sigma(2)} z_{\sigma(3)\sigma(4)}} \frac{z_{\sigma(1)\sigma(2)} z_{\sigma(5)\sigma(6)}}{z_{\sigma(1)\sigma(6)} z_{\sigma(2)\sigma(5)}} \frac{\prod_{k \neq 5,6} z_{\sigma(k)\sigma(5)} z_{\sigma(k)\sigma(6)}}{z_{\sigma(5)\sigma(6)}} \right) \\
+ \frac{z_{\sigma(1)\sigma(5)} z_{\sigma(2)\sigma(6)}}{z_{\sigma(1)\sigma(2)} z_{\sigma(5)\sigma(6)}} \frac{z_{\sigma(3)\sigma(4)} z_{\sigma(5)\sigma(6)}}{z_{\sigma(3)\sigma(6)} z_{\sigma(4)\sigma(5)}} \frac{\prod_{k \neq 3,4} z_{\sigma(k)\sigma(3)} z_{\sigma(k)\sigma(4)}}{z_{\sigma(3)\sigma(4)}^{2}} \\
+ \frac{z_{\sigma(3)\sigma(5)} z_{\sigma(4)\sigma(6)}}{z_{\sigma(3)\sigma(4)} z_{\sigma(5)\sigma(6)}} \frac{z_{\sigma(1)\sigma(2)} z_{\sigma(3)\sigma(4)}}{z_{\sigma(1)\sigma(4)} z_{\sigma(2)\sigma(3)}} \frac{\prod_{k \neq 1,2} z_{\sigma(k)\sigma(1)} z_{\sigma(k)\sigma(2)}}{z_{\sigma(1)\sigma(2)}^{2}} \\
= \frac{c}{3} \prod_{1 \leq i < j \leq 6} z_{ij}^{2} \left(\sum_{a < b, m < n \neq a, b}^{6} z_{ab}^{2} \prod_{k \neq a, b, m, n}^{6} z_{km} z_{kn} \right)$$
(B.19)

which corresponds to the numerical results produced by Eq (3.39). For term that does not depend on c, one can check that it is actually antisymmetic and it will vanish over permutation sum.

So far we proved the equivalence for n = 2 and n = 3. For general n, we leave it as an exercise for enthusiastic readers to find an elementary proof of the equivalence between Simon's[13] and our formulae.

References

- A.A. Belavin, A.M. Polyakov and A.B. Zamolodchikov, *Infinite Conformal Symmetry in Two-Dimensional Quantum Field Theory*, *Nucl. Phys. B* 241 (1984) 333.
- [2] D. Friedan, E. Martinec and S. Shenker, Conformal invariance, supersymmetry and string theory, Nuclear Physics B 271 (1986) 93.
- [3] P. Di Francesco, P. Mathieu and D. Senechal, *Conformal Field Theory*, Graduate Texts in Contemporary Physics, Springer-Verlag, New York (1997), 10.1007/978-1-4612-2256-9.
- [4] J. Cardy, Conformal field theory and statistical mechanics, 2008. 10.48550/ARXIV.0807.3472.
- [5] B. Han, A. Tiwari, C.-T. Hsieh and S. Ryu, *Boundary conformal field theory and symmetry-protected topological phases in* 2 + 1 *dimensions*, *Phys. Rev. B* **96** (2017) 125105.
- [6] G.Y. Cho, K. Shiozaki, S. Ryu and A.W.W. Ludwig, *Relationship between symmetry protected topological phases and boundary conformal field theories via the entanglement spectrum, Journal of Physics A: Mathematical and Theoretical* 50 (2017) 304002.
- [7] X. Chen, A. Roy, J.C.Y. Teo and S. Ryu, *From orbifolding conformal field theories to gauging topological phases*, *Phys. Rev. B* **96** (2017) 115447.
- [8] Y.-Z. You, Z. Bi, A. Rasmussen, K. Slagle and C. Xu, Wave function and strange correlator of short-range entangled states, Phys. Rev. Lett. 112 (2014) 247202.
- [9] B.I. Halperin and J.K. Jain, *Fractional Quantum Hall Effects*, WORLD SCIENTIFIC (Jan., 2020), 10.1142/11751.

- [10] G.W. Moore and N. Read, Nonabelions in the fractional quantum Hall effect, Nucl. Phys. B 360 (1991) 362.
- [11] N. Read, Non-abelian adiabatic statistics and hall viscosity in quantum hall states and $p_x + ip_y$ paired superfluids, Phys. Rev. B 79 (2009) 045308.
- [12] S.H. Simon, E.H. Rezayi, N.R. Cooper and I. Berdnikov, *Construction of a paired wave function for* spinless electrons at filling fraction $\nu = 2/5$, *Phys. Rev. B* **75** (2007) 075317.
- [13] S.H. Simon, Correlators of $\mathcal{N} = 1$ superconformal currents, Journal of Physics A: Mathematical and Theoretical 42 (2009) 055402.
- [14] S.H. Simon, E.H. Rezayi and N. Regnault, *Quantum hall wave functions based on S₃ conformal field theories*, *Phys. Rev. B* 81 (2010) 121301.
- [15] N. Read and E. Rezayi, Beyond paired quantum Hall states: Parafermions and incompressible states in the first excited Landau level, Phys. Rev. B 59 (1999) 8084 [cond-mat/9809384].
- [16] B.A. Bernevig and F.D.M. Haldane, *Model fractional quantum hall states and jack polynomials*, *Physical Review Letters* **100** (2008).
- [17] P. Goddard, A. Kent and D. Olive, Unitary representations of the virasoro and super-virasoro algebras, Communications In Mathematical Physics 103 (1986) 105.
- [18] Z. Qiu, Supersymmetry, two-dimensional critical phenomena and the tricritical ising model, Nuclear Physics B 270 (1986) 205.
- [19] D. Friedan, Z. Qiu and S. Shenker, Superconformal invariance in two dimensions and the tricritical ising model, Physics Letters B 151 (1985) 37.
- [20] B. Estienne, B.A. Bernevig and R. Santachiara, *Electron-quasihole duality and second-order differential equation for read-rezayi and jack wave functions*, *Physical Review B* 82 (2010).
- [21] S.L. Lukyanov and V.A. Fateev, Additional symmetries and exactly soluble models in two-dimensional conformal field theory (1990).
- [22] S.-w. Chung, E. Lyman and S.H.H. Tye, Fractional supersymmetry and minimal coset models in conformal field theory, Int. J. Mod. Phys. A 7 (1992) 3339.
- [23] P.C. Argyres, J.M. Grochocinski and S.-H. Tye, *Construction of the k = 8 fractional superconformal algebras*, *Nuclear Physics B* **391** (1993) 409.
- [24] P.C. Argyres and S.H.H. Tye, Tree scattering amplitudes of the spin-⁴/₃ fractional superstring. i. the untwisted sectors, Phys. Rev. D 49 (1994) 5326.
- [25] V. Fateev and A. ZamolodchikovTheot. Math. Phys. 71 45 (1987).
- [26] J.A. Harvey and Y. Wu, *Hecke relations in rational conformal field theory*, *Journal of High Energy Physics* **2018** (2018).
- [27] J.A. Harvey, Y. Hu and Y. Wu, Galois symmetry induced by hecke relations in rational conformal field theory and associated modular tensor categories, Journal of Physics A: Mathematical and Theoretical 53 (2020) 334003.
- [28] J.-B. Bae, J.A. Harvey, K. Lee, S. Lee and B.C. Rayhaun, *Conformal field theories with sporadic group symmetry*, *Communications in Mathematical Physics* **388** (2021) 1.

- [29] Z. Duan, K. Lee and K. Sun, *Hecke relations, cosets and the classification of 2d RCFTs, Journal of High Energy Physics* **2022** (2022).
- [30] M.H. Freedman, M. Larsen and Z. Wang, A modular functor which is universal for quantum computation, Communications in Mathematical Physics 227 (2002) 605.
- [31] Y. Hu and C.L. Kane, Fibonacci topological superconductor, Phys. Rev. Lett. 120 (2018) 066801.
- [32] M.P.A. Fisher and L.I. Glazman, *Transport in a one-dimensional luttinger liquid*, 1996. 10.48550/ARXIV.COND-MAT/9610037.
- [33] C.L. Kane and M.P.A. Fisher, *Transport in a one-channel luttinger liquid*, *Phys. Rev. Lett.* **68** (1992) 1220.
- [34] E. Sagi and R.A. Santos, *Supersymmetry in the fractional quantum hall regime*, *Phys. Rev. B* **95** (2017) 205144.
- [35] L. Dixon, P. Ginsparg and J. Harvey, superconformal field theory, Nuclear Physics B 306 (1988) 470.
- [36] P. Ginsparg, Applied conformal field theory, .
- [37] I.B. Frenkel and Y. Zhu, Vertex operator algebras associated to representations of affine and Virasoro algebras, Duke Mathematical Journal **66** (1992) 123.
- [38] M.R. Gaberdiel and P. Goddard, Axiomatic conformal field theory, Commun. Math. Phys. 209 (2000) 549 [hep-th/9810019].
- [39] M.R. Gaberdiel, An Introduction to conformal field theory, Rept. Prog. Phys. 63 (2000) 607 [hep-th/9910156].
- [40] J. Polchinski, String theory. Vol. 2: Superstring theory and beyond, Cambridge Monographs on Mathematical Physics, Cambridge University Press (12, 2007), 10.1017/CBO9780511618123.
- [41] C. Chen and J. Maciejko, *Revisiting the ramond sector of the* $\mathcal{N} = 1$ *superconformal minimal models*, *Phys. Rev. D* **102** (2020) 121701.
- [42] I. Makabe and G.M. Watts, Defects in the tri-critical ising model, Journal of High Energy Physics 2017 (2017).
- [43] C. Richard and P.A. Pearce, Integrable lattice realizations of n=1 superconformal boundary conditions, Nuclear Physics B 631 (2002) 447.
- [44] R.I. Nepomechie, Consistent superconformal boundary states, Journal of Physics A: Mathematical and General 34 (2001) 6509.
- [45] B.A. Bernevig, V. Gurarie and S.H. Simon, Central charge and quasihole scaling dimensions from model wavefunctions: toward relating jack wavefunctions to -algebras, Journal of Physics A: Mathematical and Theoretical 42 (2009) 245206.
- [46] F.D.M. Haldane, Fractional quantization of the hall effect: A hierarchy of incompressible quantum fluid states, *Phys. Rev. Lett.* **51** (1983) 605.
- [47] S.A. Trugman and S. Kivelson, *Exact results for the fractional quantum hall effect with general interactions*, *Phys. Rev. B* **31** (1985) 5280.