

On Inflationary Models in $f(R, T)$ Gravity with a Kinetic Coupling Term

A. Belhaj^{*}, M. Benali[†], Y. Hassouni[‡], M. Lamaaoune^{§¶}

Département de Physique, Equipe des Sciences de la matière et du rayonnement, ESMaR
Faculté des Sciences, Université Mohammed V de Rabat, Rabat, Morocco

March 23, 2023

Abstract

We investigate inflationary models in $f(R, T)$ modified gravity with a kinetic coupling term $\omega^2 G^{\mu\nu} \partial_\mu \phi \partial_\nu \phi$ having a positive factor needed to remove the ghosts. Taking $f(R, T) = R + 2\beta T$, we calculate and analyse the relevant observable quantities including the spectral index n_s and the tensor-to-scalar ratio r using the slow-roll approximations. Concretely, we consider two scenarios described by the decoupling and the coupling behaviors between the scalar potential and the $f(R, T)$ gravity via the moduli space by dealing with two potentials being the quartic one $V(\phi) = \lambda\phi^4$ and the small field inflation $V(\phi) = V_0(1 - (\frac{\phi}{\mu})^\alpha)$. For the quartic inflation model, we consider a decoupling behavior. For the small field inflation, however, we present the parameter decoupling and coupling scenarios. For both scenarios, we compute and inspect n_s and r showing interesting results. For three different values of the number of e-folds $N = 60, 65$ and 70 , we find that the coupling between $f(R, T)$ and the scalar potential via the moduli space provides an excellent agreement with the observational findings. In the last part of this work, we provide a possible discussion on the amplitude of scalar power spectrum needed to provide a viability of the proposed theory. Considering the second potential form in the parameter coupling scenario, we find acceptable values in certain points of the moduli space.

Keywords: Inflation, Kinetic coupling terms, Cosmological observables, Slow-roll computations, Observational data.

^{*}a-belhaj@um5r.ac.ma

[†]mohamed_benali4@um5.ac.ma

[‡]y.hassouni@um5r.ac.ma

[§]mustapha_lamaaoune@um5.ac.ma

[¶]Authors in alphabetical order.

1 Introduction

Inflationary models have been extensively investigated by considering various gravity theories including the modified ones [1–5]. Certain elaborated ones have brought roads to understand the universe evolution features such as horizon, flatness and large structure issues [6–8]. Several theories have been proposed by considering the single and the multiple scalar fields via the potentials in order to discuss standard cosmological problems within general relativity (GR). The simplest ones involve only a single field being interpreted as a dominated element to drive inflation. Such an object has been approached by dealing with various theories even the ones appearing in higher dimensions [9–12].

Developments in string theory has been also exploited to engineer inflationary models relying on the D3-brane physics using the Randall-Sundrum II (RS-2) mechanism. In this way, interesting scalar potentials involving certain parameters such as the brane tension have been studied where the stringy corrections of the involved quantities have been derived [13–17]. In particular, the compactifications of superstring models and M-theory have been explored producing many scalar fields. These objects are associated with geometric deformations of the metric and other non trivial tensor fields coupled to the brane objects [18–20]. The corresponding cosmological quantities have been approached and examined. These scalar fields have been exploited to confront the theoretical predictions with observational data provided by cosmic microwave backgrounds (CMB) and the Planck experimental results [21–23].

Alternatively, GR modified models have been explored showing interesting inflationary results. The most dealt with ones are $f(R)$ modified gravity theories where R denotes the Ricci scalar being largely studied by taking several scalar potentials [24–26]. Based on the slow-roll analysis, the associated spectral index n_s and the tensor-to-scalar ratio r have been calculated with specific values of the number of e-folds needed to bridge the resulting models with the observational findings. These models have been extended by adding others quantities including the trace T of the stress-energy tensor. The resulting is called $f(R, T)$ gravity theory being extensively studied providing certain inflationary models [27, 28]. A close examination reveals that the modified gravity theories are motivated by the study of dark energy (DE). It has been remarked that the latter could be exploited to bring certain explanations of the accelerating expansion aspect of the universe. Concretely, the modified gravity theories could generate certain dynamical behaviors which can be interpreted as DE contributions [29–31]. These behaviors go beyond the cosmological constant in the GR context.

Alternatively, a kinetic coupling term has been proposed to generate inflationary models motivated by Higgs field physics. In particular, the relevant observable quantities have been computed and analyzed [32–38]. In the absence parameter coupling scenarios, however, certain models do not provide observational features.

It has been observed that the form of the scalar potential can play a primordial role in the building of inflationary models derived from various theories including superstring models, and M-theory. The choice of the potential form usually depends on motivations relating

on known models such as the standard model (SM) of the particle physics [36, 37]. It has been remarked that famous illustrations are the chaotic inflation potential and the minimal supersymmetric standard model (MSSM) inflation potential. In addition to these studied models, other forms of the scalar potential have been dealt with in connections with black hole physics and quintessential DE scenarios [39–41].

The main objective of this work is to contribute to these activities by studying inflationary models within the $f(R, T)$ modified gravity by means of a kinetic coupling mechanism via the term $\omega^2 G^{\mu\nu} \partial_\mu \phi \partial_\nu \phi$, with a positive factor needed to remove the ghosts. Taking $f(R, T) = R + 2\beta T$, we reconsider the study of the quartic potential and the small field one given by $V(\phi) = \lambda\phi^4$ and $V(\phi) = V_0(1 - (\frac{\phi}{\mu})^\alpha)$, respectively. Within such a mechanism, we compute and examine the relevant observables such as the spectral index n_s and the tensor-to-scalar ratio r by the help of the slow-roll approximations. Precisely, we provide two situations described by the decoupling and the coupling between the scalar potential and the modified gravity via the involved moduli space. In the first model associated with the quartic potential, we investigate a decoupling behavior. In the second one corresponding to the small field inflation, however, we study the parameter decoupling and coupling scenarios. In both scenarios, we investigate the spectral index n_s and the tensor-to-scalar ratio r . Precisely, we illustrate the associated cosmological behaviors using graphic presentations. For three different values of the e-folding number $N = 60, 65$ and 70 , we find that the coupling between the $f(R, T)$ modified gravity and the scalar potential via the moduli space provides an excellent agreement with observational findings. We end this work by elaborating a possible discussion on the amplitude of scalar power spectrum which could be needed to provide a viability of the proposed theory.

The organization of this paper is as follows. In section 2, we study $f(R, T) = R + 2\beta T$ gravity action within a kinetic coupling scenario and establish the field equations of motion. In section 3, we investigate the relevant inflation parameters in such a modified gravity model with a generic scalar potential. In section 4, we make contact with observational data by considering two models being the small field and the quartic potential inflations using the parameter decoupling and coupling scenarios. In section 5, we give concluding remarks. In this work, we take $G = c = 1$ corresponding to the gravitational constant and the light speed, respectively.

2 Field equations in $f(R, T)$ gravity with a kinetic coupling term

In this section, we give a concise discussion on $f(R, T)$ gravity with a kinetic coupling term. Then, we present the associated field equations needed to investigate inflation models in the resulting theory. To start, we consider the following action

$$S = \int dx^4 \sqrt{-g} \left(\frac{f(R, T)}{16\pi} + \mathcal{L}_m \right) \quad (2.1)$$

where $f(R, T)$ is an arbitrary function of the Ricci scalar R and T denoting the trace of the stress-energy tensor $T_{\mu\nu}$ of the matter sector and where g is the determinant of the metric $g_{\mu\nu}$ [42–47]. In the absence of the kinetic coupling terms, \mathcal{L}_m describes a matter Lagrangian density given by

$$\mathcal{L}_m = -\frac{1}{2}g^{\mu\nu}\nabla_\mu\phi\nabla_\nu\phi - V(\phi) \quad (2.2)$$

where ϕ indicates the dynamical scalar field controlled by a potential $V(\phi)$. In this situation, the stress-energy tensor $T_{\mu\nu}$ of the matter reads as

$$T_{\mu\nu} = -\frac{2}{\sqrt{-g}}\frac{\delta(\sqrt{-g}\mathcal{L}_m)}{\delta g^{\mu\nu}}. \quad (2.3)$$

In this work, we combine two scenarios to provide inflationary models. The first scenario that we follow here is based on the introduction of a kinetic coupling term in the previous action. This term has been extensively studied in connections with Higgs theory coupled to the gravity which could appear in heterotic superstring models and related topics [48]. In this way, we consider the matter sector described by the following Lagrangian density

$$\mathcal{L}_m^k = -\frac{1}{2}(g^{\mu\nu} - \omega^2 G^{\mu\nu})\nabla_\mu\phi\nabla_\nu\phi - V(\phi) \quad (2.4)$$

where $G_{\mu\nu}$ is the Einstein tensor and ω is a parameter having the dimension of the mass scale. In present analysis, we take a positive sign in the action corresponding to the factor $+\omega^2$. The positive sign of the kinetic coupling term ensures the absence of the ghost in the underlying models [36, 37]. A close inspection shows that the kinetic terms have been introduced in order to reduce the tensor-to-scalar ratio r providing certain arguments with CMB observations. Varying the modified action with respect to the metric $g_{\mu\nu}$, we can obtain the following field equation

$$\begin{aligned} f_R(R, T)R_{\mu\nu} &= \frac{1}{2}g_{\mu\nu}f(R, T) - (g_{\mu\nu}\square - \nabla_\mu\nabla_\nu)f_R(R, T) \\ &= -f_T(R, T)T_{\mu\nu} - f_T(R, T)\Theta_{\mu\nu} + 8\pi(T_{\mu\nu}^{(\phi)} + \omega^2 A_{\mu\nu}) \end{aligned} \quad (2.5)$$

where we have used the notion $f_T(R, T) = \frac{\partial f(R, T)}{\partial T}$ and $f_R(R, T) = \frac{\partial f(R, T)}{\partial R}$. The involved quantities are found to be

$$\Theta_{\mu\nu} = g^{\alpha\beta}\frac{\delta T_{\alpha\beta}}{\delta g^{\mu\nu}} \quad (2.6)$$

$$T_{\mu\nu}^{(\phi)} = \nabla_\mu\phi\nabla_\nu\phi - \frac{1}{2}g_{\mu\nu}\nabla_\rho\phi\nabla^\rho\phi + g_{\mu\nu}V(\phi) \quad (2.7)$$

$$\begin{aligned} A_{\mu\nu} &= -\frac{1}{2}\nabla_\mu\phi\nabla_\nu\phi R + 2\nabla_\alpha\phi\nabla_{(\mu}\phi R_{\nu)}^\alpha + \nabla^\alpha\phi\nabla^\beta\phi R_{\mu\alpha\nu\beta} \\ &+ \nabla_\mu\nabla^\alpha\phi\nabla_\nu\nabla_\alpha\phi - \nabla_\mu\nabla_\nu\phi\square\phi - \frac{1}{2}(\nabla\phi)^2 G_{\mu\nu} \\ &+ g_{\mu\nu}\left[-\frac{1}{2}\nabla^\alpha\phi\nabla^\beta\phi\nabla_\alpha\phi\nabla_\beta\phi + \frac{1}{2}(\square\phi)^2 - \nabla_\alpha\phi\nabla_\beta\phi R^{\alpha\beta}\right]. \end{aligned} \quad (2.8)$$

Contracting Eq.(2.5), the Ricci scalar R and the trace T of the stress-energy tensor are linked via the relation

$$f_R(R, T)R - \frac{1}{2}f(R, T) = -f_T(R, T)T - f_T(R, T)\Theta + 8\pi(T^{(\phi)} + \omega^2 A) \quad (2.9)$$

where one has used $A = A^\mu_\mu$ and $\Theta = \Theta^\mu_\mu$. In this framework, it has been remarked that the covariant derivative of the energy-momentum tensor is not null $\nabla_\mu T_{\mu\nu} \neq 0$. In order to obtain the modified continuity relation, the covariant derivative of (2.5) should be performed. Indeed, one writes

$$\begin{aligned} \nabla^\mu \left[f_R(R, T) R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} f(R, T) - (g_{\mu\nu} \square - \nabla_\mu \nabla_\nu) f_R(R, T) \right. \\ \left. = -f_T(R, T) T_{\mu\nu} - f_T(R, T) \Theta_{\mu\nu} + 8\pi (T_{\mu\nu}^{(\phi)} + \omega^2 A_{\mu\nu}) \right]. \end{aligned} \quad (2.10)$$

Using the identities $\nabla R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu} = 0$, and $(\nabla_\nu \square - \square \nabla_\nu) f_R(R, T) = R_{\mu\nu} \nabla^\mu f_R$ and considering the following relation

$$\nabla^\mu \left[f_R(R, T) R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} f(R, T) - (g_{\mu\nu} \square - \nabla_\mu \nabla_\nu) f_R(R, T) \right] = 0 \quad (2.11)$$

the divergence of the stress-energy tensor $T_{\mu\nu}$ takes the form

$$\nabla^\mu T_{\mu\nu} = -\frac{1}{f_T(R, T)} \left[(T_{\mu\nu} + \Theta_{\mu\nu}) \nabla^\mu f_T(R, T) - 8\pi (\nabla^\mu T_{\mu\nu}^{(\phi)} + \omega^2 \nabla^\mu A_{\mu\nu}) \right]. \quad (2.12)$$

For $\omega = 0$, we recover the usual expressions [42, 43]. Exploiting the pressure p , we consider the following situation

$$\Theta_{\mu\nu} = -2T_{\mu\nu} - p g_{\mu\nu} \quad (2.13)$$

associated with a homogeneous and an isotropic universe which can be modeled by a perfect fluid [42–44].

A close examination reveals that a generic form of $f(R, T)$ generate complex computations leading to a very hard task. However, certain models have been proposed by dealing with special functions recovering the GR limit where $f(R, T)$ reduces to R . Ignoring the RT mixing terms, such models are based on the following forms

$$f(R, T) = R + 2g(T) \quad (2.14)$$

where $g(T)$ is an arbitrary function of T . Motivated by DE activities, the second scenario relay on a possible extension of GR by restricting the present investigation to a linear form of $f(R, T)$ gravity given by

$$f(R, T) = R + 2\beta T \quad (2.15)$$

where β is a new positive parameter being independent of T and R . It can be considered as a relevant modified gravity parameter. As advantages, this form can be considered as an economic extension of GR without RT mixing terms which need more complicated calculations. Moreover, it involves only one extra parameter reducing the generic moduli space. As envisaged, this parameter will be exploited to suggest a new coupling between the gravity and the dynamical scalar field through the potential $V(\phi)$. This coupling parameter mechanism being elaborated by the implementation of β in $V(\phi)$ could open possible roads to establish

bridges with DE modeling from inflation investigations. Using $g_{\mu\nu}$ and ϕ variations via the Friedman-Lemaitre-Robertson-Walker (FLRW) metric, we get the equations of motion

$$3H^2 = 4\pi\dot{\phi}^2(1 + 9\omega^2 H^2) + (8\pi + 6\beta)V(\phi) - 4\beta\dot{\phi}^2 \quad (2.16)$$

$$2\dot{H} + 3H^2 = -4\pi\dot{\phi}^2(1 - \omega^2(2\dot{H} + 3H^2 + 4H\ddot{\phi}\dot{\phi}^{-1})) + (8\pi + 6\beta)V(\phi) - 4\beta\dot{\phi}^2 \quad (2.17)$$

$$(\ddot{\phi} + 3H\dot{\phi}) + 3\omega^2(H^2\ddot{\phi} + 3H^3\dot{\phi} + 2H\dot{H}\dot{\phi}) = -V'(\phi) \quad (2.18)$$

where the dot is the derivative with respect to the cosmic time t , and one has used $V' = \frac{dV}{d\phi}$. H is the Hubble parameter defined by $H = \frac{\dot{a}}{a}$ where $a(t)$ is a scalar factor. For $\beta = 0$, we recover the equations of motion in the RG limit described by $f(R, T) = R$ [34, 35]. A close examination shows that the cosmological observables can be obtained using certain approximations. To make contact with the observational data [21–23], the appropriate approximations will be exploited.

3 Slow-roll computations

To get the observable quantities in an acceptable way, we could use certain approximations simplifying the previous equations of motion by removing the most slowly changing contributions [49]. Under the slow-roll approximations which could provide exact solutions of the equations of motion, the source of the energy is the scalar potential being the dominated pieces in the inflation scenario. Moreover, the acceleration scalar field terms could be neglected in such equations of motion even in generic situations where the scalar potential is not specified. Concretely, we calculate the relevant inflation parameters associated with the above action involving the proposed $f(R, T)$ gravity function in the kinetic term mechanism. In the inflation phase, the slow-roll parameters are given by

$$\epsilon = -\frac{\dot{H}}{H^2}, \quad \eta = \frac{\dot{\epsilon}}{H\epsilon}, \quad \kappa_0 = 12\pi\kappa\dot{\phi}^2, \quad \kappa_1 = \frac{\dot{\kappa}_0}{H\kappa_0}. \quad (3.1)$$

According to the slow-roll inflation mechanism, these parameters are constrained as follows

$$\epsilon, \eta, \kappa_0, \kappa_1 \ll 1. \quad (3.2)$$

In such slow-roll approximations, the field equations of motion reduce to

$$3H\dot{\phi} = -9\omega^2 H^3 \dot{\phi} - V'(\phi) \quad (3.3)$$

$$3H^2 = (8\pi + 6\beta)V(\phi) \quad (3.4)$$

$$\dot{H} = -4\pi\dot{\phi}^2(1 + 3\omega^2 H^2). \quad (3.5)$$

Calculations reveal that these field equations can be solved as follows

$$H^2 = (2\beta + \frac{8\pi}{3})V(\phi) \quad (3.6)$$

$$\dot{H} = -\frac{2\pi V'(\phi)^2}{3(6\beta + 8\pi)V(\phi)(1 + (6\beta + 8\pi)\omega^2 V(\phi))} \quad (3.7)$$

$$\dot{\phi} = -\frac{V'(\phi)}{\sqrt{6(9\beta + 24\pi)}\sqrt{V(\phi)}(1 + (6\beta + 8\pi)\omega^2 V(\phi))}. \quad (3.8)$$

Exploiting the slow-roll parameters, Eq.(3.8), Eq.(3.7), and Eq.(3.6), we can get the relevant inflation parameters as functions of $V(\phi)$, β and ω . Indeed, we obtain

$$\epsilon = \frac{\pi V'(\phi)^2}{(3\beta + 4\pi)^2 V(\phi)^2 (1 + (6\beta + 8\pi)\omega^2 V(\phi))} \quad (3.9)$$

$$\eta = \frac{V'(\phi)^2 (1 + (9\beta + 12\pi)\omega^2 V(\phi)) - V(\phi)V''(\phi) (1 + (6\beta + 8\pi)\omega^2 V(\phi))}{(3\beta + 4\pi)V(\phi)^2 (1 + (6\beta + 8\pi)\omega^2 V(\phi))^2}. \quad (3.10)$$

In the inflationary building models, the slow-roll parameters provide the scalar field value at the end of the expansion ϕ_E by imposing the constraint $\epsilon(\phi_E) = 1$. Moreover, we can find the beginning of inflation from the total logarithmic phase. It is convenient to express the above slow-roll parameters in terms of the e-folding number corresponding to the duration of inflation. This number can be computed via the following relation

$$N = \int_{t_I}^{t_E} H dt = \int_{\phi_I}^{\phi_E} \frac{H}{\dot{\phi}} d\phi \quad (3.11)$$

where the subscript I and E denote the associated values at the onset and the offset time of inflation, respectively. The calculation gives

$$N = (6\beta + 8\pi) \int_{\phi_I}^{\phi_E} \frac{V(\phi) (1 + (6\beta + 8\pi)\omega^2 V(\phi))}{V'(\phi)} d\phi. \quad (3.12)$$

In order to solve the big-bang issues, it is usually required to consider $N \geq 60$.

Roughly, the main goal of the gravitational theories is to provide consistent predictions matching with the observational data. It has been remarked that the implementation of $f(R, T)$ in the action does not effect the primordial cosmological quantities n_s and r [27, 28]. However, the non-minimal kinetic term coupling can affect such quantities. According to [32, 33], the scalar spectral index n_s and the tensor-to-scalar ratio r are expressed as follows

$$n_s - 1 = -2\epsilon - \eta \quad (3.13)$$

$$r = 16\epsilon. \quad (3.14)$$

In the present modified gravity model, these quantities are found to be

$$n_s = 1 - \frac{2\pi V'(\phi)^2}{(3\beta + 4\pi)^2 V(\phi)^2 (1 + (6\beta + 8\pi)\omega^2 V(\phi))} \quad (3.15)$$

$$+ \frac{V(\phi)V''(\phi) (1 + (6\beta + 8\pi)\omega^2 V(\phi)) - V'(\phi)^2 (1 + 3(3\beta + 4\pi)\omega^2 V(\phi))}{(3\beta + 4\pi)V(\phi)^2 (1 + (6\beta + 8\pi)\omega^2 V(\phi))^2}$$

$$r = \frac{16\pi V'(\phi)^2}{(3\beta + 4\pi)^2 V(\phi)^2 (1 + (6\beta + 8\pi)\omega^2 V(\phi))}. \quad (3.16)$$

Having calculated the primordial observables, we move to study certain models specified by the potential form. To test such models, we provide bridges to observational findings and results.

4 Inflation modeling in $f(R, T)$ gravity with a kinetic coupling scenario

In this section, we would like to investigate concrete inflation models with a kinetic coupling term in the $f(R, T)$ modified gravity. It has been remarked that the previous computations depend on β and the potential function of the single scalar field. However, interesting ones should provide smallness numerical values of the involved cosmological quantities. In the present investigations, the proposed modified gravity will be controlled by varying β in the scalar potential $V(\phi)$. This idea could provide inflationary models going beyond the previous ones generating interesting results compared with models where the gravity parameters and the scalar potential ones are not related. In the parameter coupling mechanism of the modified gravity with the matter sector via the potential, the cosmological observables are modified in terms of β . Instead of being general, we deal with two different potentials and make contact with observational data. The choice such potentials could be motivated by certain parallel investigations including either the coupling of the standard models with the gravity or DE activities. It has been observed that the possible choices could not only rely on the known scalar field potentials but also on the corresponding possible contact with observational constraints obtained in different inflation scenarios. Motivated by the potentials explored in the study of Higgs inflation with a kinetic term [11, 50], we investigate two models known by the quartic potential and the potential associated with the small field inflation. Then, we provide a comparative discussion.

4.1 Quartic potential in kinetic coupling scenarios

The first model that we would like to deal with is the inflation potential investigated in [50]. This concerns the quartic potential given by

$$V(\phi) = \lambda \phi^4 \quad (4.1)$$

where λ is a dimensionless coupling parameter. This parameter can be interpreted as the scalar self-coupling. In connection with SM, this potential has been exploited to make contact with the Higgs field considered as the only scalar field in the underlying physics. In this way, the dynamical scalar ϕ could be identified with the Higgs field. It is worth noting that other scenarios could be possible via intermediate interactions. To get the scalar spectral index n_s and the tensor-to-scalar ratio r as functions of V_0 and ϕ , we should compute the e-folding number. For this model, it takes the following form

$$N = \frac{1}{24} B (\phi_I^2 - \phi_E^2) (3 + Bx (\phi_E^4 + \phi_E^2 \phi_I^2 + \phi_I^4)) \quad (4.2)$$

where one has used $B = 8\pi + 6\beta$ and $x = \lambda\omega^2$.

The scalar spectral index n_s and the tensor-to-scalar ratio r are found to be

$$r = \frac{1024\pi}{B^2\phi^2(1+Bx\phi^4)} \quad (4.3)$$

$$n_s = 1 - \frac{128\pi}{B^2\phi^2(1+Bx\phi^4)} - \frac{8(1+3Bx\phi^4)}{B\phi^2(1+Bx\phi^4)^2}. \quad (4.4)$$

To test the obtained results, we would like to make contact with the observational data including the Planck 2018 and the recently released BICEP/Keck data [21–23]. In Fig(1), we illustrate the $n_s - r$ curve behaviors for different values of N , β , and x .




Figure 1: The behavior of one dimensional curves $n_s - r$ by varying the Parameter $\beta = 1, \dots, 50$ and the parameter $x = 0.001, \dots, 1$. The plots from left to right correspond to the number e-folding $N = 60$, $N = 65$ and $N = 70$, respectively. The green and the orange contour constraints represent 68% and 95% confidential levels of Planck results ($TT, TE, EE+lowE+lensing+BK15+BAO$), respectively. The yellow contour is associated with the Planck results ($TT, TE, EE+lowE+lensing$).

Varying the parameters β and x , these curves are plotted by implementing the Planck contour constraints. It follows from this figure that r decreases by increasing the parameter β . Taking large values of β , the range of n_s decreases. Moreover, the spectral index n_s increases by increasing the parameter x . Fixing β and x , n_s increases by increasing the e-folding number N . It has been remarked that the quartic potential produces interesting numerical values of the spectral index n_s and the tensor-to-scalar ratio r for specific values of β and x . In particular, this potential gives a good compatibility with the Planck results ($TT, TE, EE+lowE+lensing$) [21–23]. For generic values of the involved relevant parameters β , x and N , the range of r is $[0.013, 0.123]$ while n_s varies in the range $[0.963, 0.983]$. The quartic potential can generate an inflation model associated with the expansion of the universe. However, this potential does not provide an excellent compatibility with the Planck and the recent released BICEP/Keck data. In the next subsection, we study the small field inflation potential coupled to such a modified gravity in order to inspect the coupling parameter scenario impact on the compatibility with the observational data including the recent Planck results.

4.2 Small field inflation with kinetic coupling behaviors

Here, we consider the small field inflation described by the following potential

$$V(\phi) = V_0 \left(1 - \left(\frac{\phi}{\mu} \right)^\alpha \right) \quad (4.5)$$

depending on two parameters α and μ . The later is a mass scale parameter. However, α is a non zero dimensionless index [11]. V_0 is a free parameter which can be considered as M^4 , where M is a mass scale. This scalar potential has been used to provide plausible particle physics inflationary models. A rapid examination shows that α and μ enlarge now the inflation moduli space coordinated by β, ω, α and μ . Many regions and sub-spaces can be dealt with either by fixing certain parameters or by identifying other ones. Taking $\beta = \alpha$ in the scalar potential can be considered as a coupling between the scalar field and the modified gravity through the function $f(R, T)$. This kind of couplings via the moduli space could bring interesting results which can be compared with the Planck data. In order to compute the scalar spectral index n_s and the tensor-to-scalar ratio r as functions of the involved parameters describing the inflation moduli space, it is useful to exploit the e-folding number. In the present work, we have taken a fixed value of μ being $\mu = 1$. In this way, we get

$$N = \frac{B}{2\alpha} \left(\phi_E^2 \left(-\frac{2\phi_E^{-\alpha}(By+1)}{\alpha-2} + \frac{2By\phi_E^\alpha}{\alpha+2} - By - 1 \right) \right) - \frac{B}{2\alpha} \left(\phi_I^2 \left(-\frac{2\phi_I^{-\alpha}(By+1)}{\alpha-2} + \frac{2By\phi_I^\alpha}{\alpha+2} - By - 1 \right) \right) \quad (4.6)$$

where we have used $y = V_0\omega^2$. This e-folding number imposes a constraint on the parameter α . It has been remarked that this parameter should be different to 2 and -2 . The scalar spectral index n_s and the tensor-to-scalar ratio r are found to be

$$n_s = 1 - \frac{8\pi\alpha^2\phi^{2\alpha-2}}{B^2(1-\phi^\alpha)^2(By(1-\phi^\alpha)+1)} - \frac{\alpha\phi^{\alpha-2}(2(\phi^\alpha-1)(By(1-\phi^\alpha)+1)+\alpha(2-By(\phi^{2\alpha}+\phi^\alpha-2)))}{B(1-\phi^\alpha)^2(By(1-\phi^\alpha)-1)^2} \quad (4.7)$$

$$r = \frac{64\pi\alpha^2\phi^{2\alpha-2}}{B^2(1-\phi^\alpha)^2(1+By(1-\phi^\alpha))}. \quad (4.8)$$

The scalar potential form and these observable quantities impose certain conditions of the parameter α . These constraints will be considered in the following discussions. To inspect the obtained results, we should make contact with observational data including the Planck 2018 and the recently released BICEP/Keck data [21–23].

In Fig(2), we examine the $n_s - r$ curves by varying N, β, y and fixing $\alpha = 1$. These behaviors are plotted by implementing the Planck contour conditions by varying the parameters β and x with $\alpha = 1$. The $n_s - r$ curves have been examined by taking β in the interval [3, 10]. It is clear from this figure that r decreases by increasing the parameter β . Taking large values




Figure 2: The behavior of one dimensional $n_s - r$ curves by varying the Parameter $\beta = 3, \dots, 10$, $\alpha = 1$ and the parameter $y = 0.001, \dots, 1$. The plots from left to right correspond to the number e-folding $N = 60$, $N = 65$ and $N = 70$, respectively. The green and the orange contour constraints represent 68% and 95% confidential levels of Planck results ($TT, TE, EE+lowE+lensing+BK15+BAO$), respectively. The yellow contour is associated with the Planck results ($TT, TE, EE+lowE+lensing$).

of β , the range of n_s decreases. Moreover, the spectral index n_s increases by increasing the parameter y . Fixing β and y , n_s increases by increasing N . For certain specific models where the scalar field potential and the modified gravity $f(R, T)$ with the kinetic couplings are not coupled, the numerical values for the spectral index n_s and the tensor-to-scalar ratio r do not belong to the range of the Planck data and the recent released BICEP/Keck data for all regions of the involved moduli space [21–23]. For generic values of the involved parameters including the degree polynomial of the potential and N , the range of r is $[0.001, 0.043]$ while n_s varies in the range $[0.978, 0.983]$.

In what follows, we couple the scalar field potential with the $f(R, T)$ gravity via the moduli space by taking $\alpha = \beta$. This corresponds to a line in such a moduli space. In Fig(3), we illustrate the $n_s - r$ curve behaviors for different values of N , β , and y .




Figure 3: The behavior of one dimensional $n_s - r$ curves by varying the parameter $\beta = 2.1, \dots, 10$ and the parameter $y = 0.001, \dots, 1$. The plots from left to right correspond to the number e-folding $N = 60$, $N = 65$ and $N = 70$, respectively. The green and the orange contour constraints represent 68% and 95% confidential levels of Planck results ($TT, TE, EE+lowE+lensing+BK15+BAO$), respectively.

Varying the parameters β and y , these behaviors are plotted by implementing the Planck contour conditions. The $n_s - r$ curves are examined by taking β in the interval $[2.1, 10]$. It follows from this figure that r decreases by increasing the gravity parameter β . Taking large

values of β , the range of n_s decreases. Moreover, the spectral index n_s increases by increasing the parameter y . Fixing β and y , n_s increases by increasing N . A close inspection reveals that the coupling between a specific scalar field potential and the modified gravity $f(R, T)$ with the kinetic couplings provides interesting numerical values for the spectral index n_s and the tensor-to-scalar ratio r which are in a good range of the Planck data and the recent released BICEP/Keck data for all regions of the involved moduli space [21–23]. For generic values of the relevant parameters including the polynomial degree of the potential and N , the range of r is [0.001, 0.014] while n_s varies in the range [0.947, 0.98].

To end this work, we would like to discuss the amplitude of scalar power spectrum needed to provide a viability of the proposed theory. Following [35], the slow-roll analysis can be exploited to bring an expression for such an amplitude. It is given approximatively by

$$P_\zeta \approx \frac{H^2}{8\pi^2\epsilon}. \quad (4.9)$$

To approach such a quantity, we consider the second potential given by Eq.(4.5) in the parameter coupling scenario by taking $\alpha = \beta$. The calculation gives

$$P_\zeta \approx \frac{(3\beta + 8\pi)^3 V_0 \left(\frac{\phi}{\mu}\right)^{2-2\beta} \left(1 - \left(\frac{\phi}{\mu}\right)^\beta\right)^3 \left(1 + 3y(3\beta + 8\pi) \left(1 - \left(\frac{\phi}{\mu}\right)^\beta\right)\right)}{96\pi^2\beta^2}. \quad (4.10)$$

It is observed that this cosmological observable depends on the matter and the gravity parameters. In some points of the inflation moduli space, we could find certain arguments with the observational data. Considering $\beta = 10$, $\mu = 1$, $V_0 = 0.001$, $y = 0.001$, and $\phi = 0.99$, for instance, we find $P_\zeta \approx 1.96 \times 10^{-6}$, being an acceptable value.

5 Conclusion

In this paper, we have investigated the $f(R, T)$ modified gravity theory in the kinetic coupling inflation scenarios. In particular, we have dealt with the form $f(R, T) = R + 2\beta T$ function by considering a positive factor for the kinetic coupling term in order to avoid the ghost in the resulting models. Using the field variation methods, we have obtained the corresponding equations of motion for a generic potential form. We have exploited the slow-roll approximations to compute and inspect the cosmological quantities. To make contact with observational data, we have considered two different scalar potentials given by the quartic form $V(\phi) = \lambda\phi^4$ and the small field inflation potential $V(\phi) = V_0(1 - (\frac{\phi}{\mu})^\alpha)$, respectively. Concretely, we have dealt with two situations described by the decoupling and the coupling between the scalar potential and the gravity via the cosmological moduli space. In the first model associated with the quartic potential, we have presented the decoupling behavior. In the second one corresponding to the small field potential, however, we have considered the decoupling behavior. Then, we have discussed the coupling between the scalar potential and the modified gravity by taking the line $\alpha = \beta$ in the moduli space. In such models, we have computed and examined the spectral index n_s and the tensor-to-scalar ratio r . Precisely, we

have plotted the associated cosmological behaviors. For three different values of the e-folding number $N = 60, 65$ and 70 , we have found that the coupling between $f(R, T)$ and the scalar potential via the moduli space provides a good agreement with observational findings. This has been illustrated for the small field potential giving an excellent match with the Planck data. We anticipate that the coupling between the scalar potential and the modified gravity with a kinetic term via the function $f(R, T)$ could provide new inflation models. In the end of this work, we have presented a possible discussion on the amplitude of scalar power spectrum. This could be needed to provide a viability of the proposed theory. Precisely, we have considered the second potential form in the parameter coupling scenario producing acceptable values of such a quantity in certain points of the inflation moduli space.

This work comes up with many open questions. A natural question is to go beyond such a $f(R, T)$ function by introducing other scalar potentials in the D-brane physics context.

Acknowledgments

The authors would like to thank H. El Moumni, M. Oualaid, and M.B. Sedra for collaborations on related subjects. They would like also to thank the editor and the anonymous referees for remarks, suggestions and scientific helps. This work is partially supported by the ICTP through AF.

References

- [1] S. D. Odintsov and V. K. Oikonomou, *Inflationary α -attractors from $F(R)$ gravity*, Phys. Rev. D **94** (2016)124026, [arXiv:1612.01126](#).
- [2] I. Sawicki and W. Hu, *Stability of Cosmological Solution in $f(R)$ Models of Gravity*, Phys. Rev. D **75** (2007) 127502, [arXiv:astro-ph/0702278](#).
- [3] T. P. Sotiriou, *$f(R)$ gravity and scalar-tensor theory*, Class. Quant. Grav. **23** (2006) 5117, [arXiv:gr-qc/0604028](#).
- [4] V. K. Oikonomou, *Singular Bouncing Cosmology from Gauss-Bonnet Modified Gravity*, Phys. Rev. D **92** (2015) 124027, [arXiv:1509.05827](#).
- [5] V. K. Oikonomou, *A refined Einstein–Gauss–Bonnet inflationary theoretical framework*, Class. Quant. Grav. **38** (2021)195025, [arXiv:2108.10460](#).
- [6] A. H. Guth and P. J. Steinhardt, *The inflationary universe*, *Scientific American*, **250** (1984) 129.
- [7] A. H. Guth, *The Inflationary Universe: A Possible Solution to the Horizon and Flatness Problems*, Phys. Rev. D **23** (1981) 356.

- [8] A. D. Linde, *A New Inflationary Universe Scenario: A Possible Solution of the Horizon, Flatness, Homogeneity, Isotropy and Primordial Monopole Problems*, Phys. Lett. B **108** (1982) 393.
- [9] J. Lin, Q. Gao and Y. Gong, *The reconstruction of inflationary potentials*, Mon. Not. Roy. Astron. Soc. **64** (2016) 4037, [arXiv:1508.07145](#).
- [10] M. Gamonal, *Slow-roll inflation in $f(R, T)$ gravity and a modified Starobinsky-like inflationary model*, Phys. Dark Univ. **31** (2021) 100768, [arXiv:2010.03861](#).
- [11] J. Martin, C. Ringeval and V. Vennin, *Encyclopædia Inflationaris*, Phys. Dark Univ. **6** (2014) 235, [arXiv:1303.3787](#).
- [12] A. G. Cadavid and A. E. Romano, *Effects of discontinuities of the derivatives of the inflaton potential*, Eur. Phys. J. C **75** (2015) 589, [arXiv:1404.2985](#).
- [13] G. R. Dvali and T.S.H. Henry, *Brane inflation*, Phys. Lett. B **450** (1999) 82, [arXiv:hep-ph/9812483](#).
- [14] H. V. Peiris, D. Baumann, B. Friedman and A. Cooray, *Phenomenology of D-Brane Inflation with General Speed of Sound*, Phys. Rev. D **76** (2007) 103517, [arXiv:0706.1240](#).
- [15] A. Borzou, H.R. Sepangi, S. Shahidi and R. Yousefi, *Brane $f(R)$ gravity*, EPL **88** (2009) 29001, [arXiv:0910.1933](#).
- [16] A. Belhaj, M. Benali, Y. Hassouni, M. Oualaid and M. B. Sedra, *On brane cosmological behaviors of Starobinsky inflationary model*, Int. J. Mod. Phys. A **37** (2022) 2250043.
- [17] A. Belhaj, Y. Hassouni, M. Oualaid and M. B. Sedra, *On stringy inflation potentials*, Mod. Phys. Lett. A **36** (2021) 2150225.
- [18] S. Nojiri, S. D. Odintsov and M. Sami, *Dark energy cosmology from higher-order, string-inspired gravity and its reconstruction*, Phys. Rev. D **74** (2006) 046004, [arXiv:hep-th/0605039](#).
- [19] E. D. Stewart, *Inflation, supergravity and superstrings*, Phys. Rev. D **51** (1995) 6847, [arXiv:hep-ph/9405389](#).
- [20] T. J. Li, J.L. Lopez and D.V. Nanopoulos, *Compactifications of M theory and their phenomenological consequences*, Phys. Rev. D **56** (1997) 2606, [arXiv:hep-ph/9704247](#).
- [21] Y. Akrami *et al.*, *Planck 2018 results. X. Constraints on inflation*, Astron. Astrophys. **641** (2020) 10, [arXiv:1807.06211](#).
- [22] N. Aghanim *et al.*, *Planck 2018 results. VI. Cosmological parameters*, Astron. Astrophys. **641** (2020) 6, [arXiv:1807.06209](#).

- [23] P. A. R. Ade *et al.*, *Improved Constraints on Primordial Gravitational Waves using Planck, WMAP, and BICEP/Keck Observations through the 2018 Observing Season*, Phys. Rev. Lett. **127** (2021) 151301, [arXiv:2110.00483](#).
- [24] V. K. Oikonomou, *Unifying inflation with early and late dark energy epochs in axion $F(R)$ gravity*, Phys. Rev. D **103** (2021) 044036, [arXiv:2012.00586](#).
- [25] B. Li and J.D. Barrow, *The Cosmology of $f(R)$ gravity in metric variational approach*, Phys. Rev. D **75** (2007) 084010, [arXiv:gr-qc/0701111](#).
- [26] K. Bamba, S. Nojiri, S. D. Odintsov and D. Sáez-Gómez, *Inflationary universe from perfect fluid and $F(R)$ gravity and its comparison with observational data*, Phys. Rev. D **90** (2014) 124061, [arXiv:1410.3993](#).
- [27] B. Deb and A. Deshamukhya, *Inflation in $f(R, T)$ gravity with double-well potential*, Int. J. Mod. Phys. A **37** (2022) 2250127, [arXiv:2201.04378](#).
- [28] Ashmita, P. Sarkar and D. K. Prasanta, *Inflationary Cosmology in the Modified $f(R, T)$ Gravity*, Int. J. Mod. Phys. D **31** (2022) 2250120, [arXiv:2208.11042](#).
- [29] G. N. Gadbail, S. Arora, and P. K. Sahoo, *Reconstruction of $f(Q, T)$ Lagrangian for various cosmological scenario*, Phys. Lett. B **838** (2023) 137710, [arXiv:2301.08876](#).
- [30] S. Nojiri, S. D. Odintsov, and M. Sasaki, *Gauss-Bonnet dark energy*, Phys. Rev. D **71** (2005) 123509, [arXiv:hep-th/0504052](#).
- [31] S. Nojiri, S. D. Odintsov, and M. Sami, Phys. Rev. D **74** (2006) 046004, [arXiv:hep-th/0605039](#).
- [32] J. Matsumoto and S. V. Sushkov, *Cosmology with nonminimal kinetic coupling and a Higgs-like potential*, JCAP **11**(2015)047, [arXiv:1510.03264](#).
- [33] S. Tsujikawa *Observational tests of inflation with a field derivative coupling to gravity*, Phys. Rev. D **85**(2012) 083518.
- [34] N. Avdeev and A. Toporensky, *On viability of inflation in non-minimal kinetic coupling theory*, Gravit. Cosmol. **27** (2021)269, [arXiv:2103.00556](#).
- [35] N. A. Avdeev and A.V. Toporensky, *Ruling out an inflation driven by a power law potential: kinetic coupling does not help*, Gravit. Cosmol. **28** (2022)416, [arXiv:2203.14599](#).
- [36] C. Germani, and A. Kehagias, *New Model of Inflation with Non-minimal Derivative Coupling of Standard Model Higgs Boson to Gravity*, Phys. Rev. Lett. **105** (2010) 011302, [arXiv:1003.2635](#).
- [37] C. Germani, A. Kehagias, *Cosmological Perturbations in the New Higgs Inflation*, JCAP **1005** (2010) 019, [arXiv:1003.4285](#).

- [38] S. Tsujikawa, *Observational tests of inflation with a field derivative coupling to gravity*, Phys. Rev. D **85** (2012) 083518, [arXiv:1201.5926](#).
- [39] G. Bao-Min, S. Fu-Wen, K. Yang and Z. Yu-Peng, *Primordial black holes from valley*, [arXiv:2207.09968](#).
- [40] D. Bettoni and J. Rubio, *Galaxies* **10** (2022) 22, [arXiv:2112.11948](#).
- [41] J. de Haro and L. A. Salo, *Galaxies* **9** (2021)73, [arXiv:2108.11144](#).
- [42] H. Tiberiu, S. N. L. Francisco, N. Shinichi and S.D. Odintsov, *$f(R, T)$ gravity* Phys. Rev. D **84** (2011) 024020, [arXiv:1104.2669](#).
- [43] P. H. R. S. Moraes, J. D. V. Arbañil, and M. Malheiro, *Stellar equilibrium configurations of compact stars in $f(R, T)$ gravity*, JCAP **06** (2016) 005, [arXiv:1511.06282](#).
- [44] F. G. Alvarenga, A. dela Cruz-Dombriz, M. J. S. Houndjo, M. E. Rodrigues, and D. Sáez-Gómez, *Dynamics of scalar perturbations in $f(R, T)$ gravity*, Phys. Rev. D **10** **87** (2013) 103526, [arXiv:1302.1866](#).
- [45] H. Velten and T. R. P. Caramês, *Cosmological inviability of $f(R, T)$ gravity*, Phys. Rev. D **95** **12** (2017) 123536, [arXiv:1702.07710](#).
- [46] S. Taghavi, K. Saaidi and Z. Ossouliau, *Holographic inflation in $f(R, T)$ gravity*, [arXiv:2301.02631](#).
- [47] S. Bhattacharjee, J. R. L. Santos, P. H. R. S. Moraes and P. K. Sahoo, *Inflation in $f(R, T)$ gravity*, Eur. Phys. J. Plus **135** **7** (2020) 576, [arXiv:2006.04336](#).
- [48] D. J. Gross and J. H. Sloan, *The quartic effective action for the heterotic string*, Nucl. Phys. B **291** (1987)41.
- [49] A.R. Liddle, P. Parsons and J. D. Barrow, *Formalizing the slow-roll approximation in inflation*, Phys. Rev. D **50** (1994) 7222.
- [50] M. Jérôme, C. Ringeval, R. Trotta and V. Vennin, *The Best Inflationary Models After Planck*, JCAP **03** (2014)39, [arXiv:1312.3529](#).