

Kullback-Leibler divergence for the Fréchet extreme-value distribution

Jean-Christophe PAIN^{1,2,*}

¹CEA, DAM, DIF, F-91297 Arpajon, France

²Université Paris-Saclay, CEA, Laboratoire Matière en Conditions Extrêmes,
91680 Bruyères-le-Châtel, France

March 28, 2023

Abstract

We derive a closed-form solution for the Kullback-Leibler divergence between two Fréchet extreme-value distributions. The resulting expression is rather simple and involves the Euler-Mascheroni constant.

1 Introduction

In probability theory and statistics, the Fréchet law is a particular case of a generalized extremum law (which is useful to represent phenomena with extreme values) on the same footing as the Gumbel [1] and Weibull [2] laws. The law was named after Maurice Fréchet [3].

In hydrology for instance, the Fréchet distribution is applied to extreme events such as annually maximum one-day rainfalls and river discharges [4].

The one-parameter Fréchet distribution (probability distribution function) reads

$$f(x; \alpha) = \alpha x^{-\alpha-1} e^{-x^{-\alpha}}, \quad (1)$$

with $\alpha > 0$. Its moments are respectively (see for instance Ref. [5]):

$$\mu_k = \int_0^\infty x^k f(x; \alpha) dx = \int_0^\infty t^{-\frac{k}{\alpha}} e^{-t} dt = \Gamma\left(1 - \frac{k}{\alpha}\right), \quad (2)$$

where Γ is the usual Gamma function and $k \geq 1$. The moments μ_k are defined for $k < \alpha$. The centered moments are defined as

$$\mu_{k,c} = \int_0^\infty (x - \mu_1)^k f(x; \alpha) dx \quad (3)$$

*jean-christophe.pain@cea.fr

and the reduced centered moments

$$\zeta_{k,c} = \frac{\mu_{k,c}}{(\mu_{2,c})^{n/2}}. \quad (4)$$

The skewness of the Fréchet distribution is

$$\zeta_{3,c} = \frac{\Gamma(1 - \frac{3}{\alpha}) - 3\Gamma(1 - \frac{2}{\alpha})\Gamma(1 - \frac{1}{\alpha}) + 2\Gamma^3(1 - \frac{1}{\alpha})}{\sqrt{(\Gamma(1 - \frac{2}{\alpha}) - \Gamma^2(1 - \frac{1}{\alpha}))^3}} \quad (5)$$

for $\alpha > 3$ and $+\infty$ otherwise. The excess kurtosis (kurtosis minus three) reads

$$\zeta_{4,c} - 3 = -6 + \frac{\Gamma(1 - \frac{4}{\alpha}) - 4\Gamma(1 - \frac{3}{\alpha})\Gamma(1 - \frac{1}{\alpha}) + 3\Gamma^2(1 - \frac{2}{\alpha})}{[\Gamma(1 - \frac{2}{\alpha}) - \Gamma^2(1 - \frac{1}{\alpha})]^2} \quad (6)$$

for $\alpha > 4$ and ∞ otherwise.

The Kullback-Leibler (KL) divergence [6,7] provides a non-symmetric measure of the similarity of two probability distributions p and q . In the case where both distributions are continuous, it is defined as

$$\mathcal{D}_{\text{KL}}(p|q) = \int_{-\infty}^{\infty} p(x) \ln \left(\frac{p(x)}{q(x)} \right) dx, \quad (7)$$

where $p(x)$ and $q(x)$ denote the corresponding probability densities. The KL divergence is a measure of relative entropy. It can be understood as the information loss if p is modeled by means of q . Accordingly, the smaller the KL divergence is, the more similar are p and q .

The KL divergence was already investigated in the framework of Weibull [8,9] and Gumbel [10,11] distributions.

In the next section, we present the KL divergence for the one-parameter Fréchet distribution.

2 Calculation of the Kullback-Leibler divergence

Let us consider two Fréchet distributions $f(x; \alpha_1)$ and $f(x; \alpha_2)$. The KL divergence reads

$$\mathcal{D}_{\text{KL}} = \int_0^{\infty} f(x; \alpha_1) \ln \left(\frac{f(x; \alpha_1)}{f(x; \alpha_2)} \right) dx. \quad (8)$$

We have to calculate

$$\mathcal{D}_{\text{KL}} = \int_0^{\infty} f(x; \alpha_1) \ln \left(\frac{f(x; \alpha_1)}{f(x; \alpha_2)} \right) dx, \quad (9)$$

i.e.

$$\mathcal{D}_{\text{KL}} = \int_0^{\infty} \alpha_1 x^{-\alpha_1-1} e^{-x^{-\alpha_1}} \ln \left(\frac{\alpha_1 x^{-\alpha_1-1} e^{-x^{-\alpha_1}}}{\alpha_2 x^{-\alpha_2-1} e^{-x^{-\alpha_2}}} \right) dx, \quad (10)$$

which is equal to

$$\mathcal{D}_{\text{KL}} = \alpha_1 \int_0^{\infty} x^{-\alpha_1-1} e^{-x^{-\alpha_1}} \ln \left(\frac{\alpha_1}{\alpha_2} x^{-\alpha_1+\alpha_2} e^{-x^{-\alpha_1}+x^{-\alpha_2}} \right) dx, \quad (11)$$

yielding

$$\begin{aligned}\mathcal{D}_{\text{KL}} &= \alpha_1 \ln \left(\frac{\alpha_1}{\alpha_2} \right) \int_0^\infty x^{-\alpha_1-1} e^{-x^{-\alpha_1}} dx \\ &\quad + \alpha_1 (\alpha_2 - \alpha_1) \int_0^\infty x^{-\alpha_1-1} e^{-x^{-\alpha_1}} \ln(x) dx \\ &\quad - \alpha_1 \int_0^\infty x^{-\alpha_1-1} e^{-x^{-\alpha_1}} (x^{-\alpha_1} + x^{-\alpha_2}) dx\end{aligned}\quad (12)$$

or

$$\begin{aligned}\mathcal{D}_{\text{KL}} &= \alpha_1 \ln \left(\frac{\alpha_1}{\alpha_2} \right) \int_0^\infty x^{-\alpha_1-1} e^{-x^{-\alpha_1}} dx \\ &\quad + \alpha_1 (\alpha_2 - \alpha_1) \int_0^\infty x^{-\alpha_1-1} e^{-x^{-\alpha_1}} \ln(x) dx \\ &\quad - \alpha_1 \int_0^\infty x^{-2\alpha_1-1} e^{-x^{-\alpha_1}} dx \\ &\quad + \alpha_1 \int_0^\infty x^{-\alpha_2-\alpha_1-1} e^{-x^{-\alpha_1}} dx.\end{aligned}\quad (13)$$

Let us make the change of variable $t = e^{-x^{-\alpha_1}}$. The first integral in Eq. (13) becomes

$$\int_0^\infty x^{-\alpha_1-1} e^{-x^{-\alpha_1}} dx = \frac{1}{\alpha_1} \int_0^\infty e^{-t} dt = \frac{1}{\alpha_1}. \quad (14)$$

The second integral in Eq. (13) is

$$\int_0^\infty x^{-\alpha_1-1} e^{-x^{-\alpha_1}} \ln(x) dx = -\frac{1}{\alpha_1^2} \int_0^\infty \ln t e^{-t} dt = \frac{\gamma}{\alpha_1^2}, \quad (15)$$

where

$$\gamma = - \int_0^\infty \ln t e^{-t} dt = -\Gamma'(1) \quad (16)$$

is the usual Euler-Mascheroni constant.

The fourth integral in Eq. (13) is

$$\int_0^\infty x^{-\alpha_1-\alpha_2-1} e^{-x^{-\alpha_1}} dx = \frac{1}{\alpha_1} \int_0^\infty t^{\alpha_2/\alpha_1} e^{-t} dt = \frac{1}{\alpha_1} \Gamma \left(\frac{\alpha_1 + \alpha_2}{\alpha_1} \right). \quad (17)$$

The third integral in Eq. (13) is given by expression (17) setting $\alpha_2 = \alpha_1$. It reads therefore

$$\int_0^\infty x^{-2\alpha_1-1} e^{-x^{-\alpha_1}} dx = \frac{1}{\alpha_1} \int_0^\infty t e^{-t} dt = \frac{1}{\alpha_1} \Gamma(2) = \frac{1}{\alpha_1}. \quad (18)$$

Inserting expressions (14), (15), (17) and (18) in Eq. (13) gives the final result

$$\boxed{\mathcal{D}_{\text{KL}} = \ln \left(\frac{\alpha_1}{\alpha_2} \right) + \frac{(\alpha_2 - \alpha_1)}{\alpha_1} \gamma - 1 + \Gamma \left(\frac{\alpha_1 + \alpha_2}{\alpha_1} \right)}. \quad (19)$$

3 Conclusion

In this note, we derived a closed-form solution for the Kullback-Leibler divergence between two Fréchet extreme-value distributions. The present results can be easily extended to the generalized Fréchet law, obtained by introducing a parameter m (position of the maximum) and a scale parameter $s > 0$. The cumulative distribution function reads then $e^{-\left(\frac{x-m}{s}\right)^{-\alpha}}$ (instead of e^{-x^α} in the one-parameter case) if $x > m$ and 0 otherwise. It corresponds to the probability distribution function

$$g(x; \alpha, s, m) = \frac{\alpha}{s} \left(\frac{x-m}{s} \right)^{-1-\alpha} e^{-\left(\frac{x-m}{s}\right)^{-\alpha}}. \quad (20)$$

References

- [1] E. J. Gumbel, *Statistics of extremes* (New York: Columbia University Press, 1958).
- [2] A. Papoulis and S. U. Pillai, *Probability, random variables, and stochastic processes* (Boston: McGraw-Hill, 2002).
- [3] M. Fréchet, *Sur la loi de probabilité de l'écart maximum*, Ann. Soc. Polon. Math. **6**, 3 (1927) [in french].
- [4] S. Coles, *An introduction to statistical modeling of extreme values* (Springer-Verlag, 2001).
- [5] G. Muraleedharan, C. Guedes Soares and C. Lucas, *Characteristic and moment generating functions of generalized extreme value distribution (GEV)*, chapter 13, in: *Sea level rise, coastal engineering, shorelines and tides*, edited by Linda L. Wright (Nova Science Publishers, Inc, 2009).
- [6] S. Kullback et R. Leibler, *On information and sufficiency*, Ann. Math. Stat. **22**, 79 (1951).
- [7] S. Kullback, *Information theory and statistics* (New York, John Wiley and Sons, 1959).
- [8] G. Yari, A. Mirhabibi and A. Saghaei, *Estimation of the Weibull parameters by Kullback-Leibler divergence of survival functions*, Appl. Math. Inf. Sci. **7**, 187 (2013).
- [9] Ch. Bauckhage, *Computing the Kullback-Leibler divergence between two Weibull distributions*, arXiv 1310.3713 (2013).
<https://arxiv.org/pdf/1310.3713.pdf>
- [10] M. Gil, *On Rényi divergence measures for continuous alphabet sources*, Masters of Science Thesis, Department of Mathematics and Statistics, Queen's University, Kingston, Ontario, Canada (2011).
- [11] S. A. Al-Subh, *Goodness of fit test for Gumbel distribution based on Kullback-Leibler information using several different estimators*, Appl. Math. Sci. **8**, 4703 (2014).