BOOTSTRAP-ASSISTED INFERENCE FOR GENERALIZED GRENANDER-TYPE ESTIMATORS

By Matias D. Cattaneo^{1,a}, Michael Jansson^{2,b} and Kenichi Nagasawa^{3,c}

¹Department of Operations Research and Financial Engineering, Princeton University, ^acattaneo@princeton.edu

²Department of Economics, University of California at Berkeley, ^bmjansson@econ.berkeley.edu

Westling and Carone (2020) proposed a framework for studying the large sample distributional properties of generalized Grenander-type estimators, a versatile class of nonparametric estimators of monotone functions. The limiting distribution of those estimators is representable as the left derivative of the greatest convex minorant of a Gaussian process whose monomial mean can be of unknown order (when the degree of flatness of the function of interest is unknown). The standard nonparametric bootstrap is unable to consistently approximate the large sample distribution of the generalized Grenander-type estimators even if the monomial order of the mean is known, making statistical inference a challenging endeavour in applications. To address this inferential problem, we present a bootstrap-assisted inference procedure for generalized Grenander-type estimators. The procedure relies on a carefully crafted, yet automatic, transformation of the estimator. Moreover, our proposed method can be made "flatness robust" in the sense that it can be made adaptive to the (possibly unknown) degree of flatness of the function of interest. The method requires only the consistent estimation of a single scalar quantity, for which we propose an automatic procedure based on numerical derivative estimation and the generalized jackknife. Under random sampling, our inference method can be implemented using a computationally attractive exchangeable bootstrap procedure. We illustrate our methods with examples and we also provide a small simulation study. The development of formal results is made possible by some technical results that may be of independent interest.

1. Introduction. Monotone function estimators have received renewed attention in statistics, biostatistics, econometrics, machine learning, and other data science disciplines. See Groeneboom and Jongbloed (2014, 2018) for a textbook introduction and a review article, respectively, the latter being published in a special issue devoted to nonparametric inference under shape constraints. More recently, Westling and Carone (2020) expanded the scope and applicability of monotone function estimators by embedding many such estimators in a unified framework of generalized Grenander-type estimators. Estimation problems covered by Westling and Carone's (2020) general theory include many practically relevant examples such as monotone density, regression and hazard estimation, possibly with censoring and/or covariate adjustment.

The large sample theory developed by Westling and Carone (2020) offers a general distributional approximation involving the left derivative of the Greatest Convex Minorant (GCM) of a Gaussian process whose mean and covariance kernel depend on unknown functions. Furthermore, both the convergence rate of the estimator and the shape of the mean appearing in the representation of the limiting distribution depend on whether the unknown function of interest exhibits certain degeneracies. For these reasons, the large sample distributional approximation for generalized Grenander-type estimators can be difficult to employ in practice

³Department of Economics, University of Warwick, ^ckenichi.nagasawa@warwick.ac.uk

for inference purposes. In their concluding remarks, Westling and Carone (2020) recognize these limitations and pose the question of whether it would be possible to employ bootstrap-assisted methods to conduct automatic/robust statistical inference within their framework.

As is well documented, the standard nonparametric bootstrap does not provide a valid distributional approximation for the generalized Grenander-type estimators (Kosorok, 2008; Sen, Banerjee and Woodroofe, 2010). This fact has led scholars to rely on other bootstrap schemes such as subsampling (Politis and Romano, 1994), the smoothed bootstrap (Kosorok, 2008; Sen, Banerjee and Woodroofe, 2010), the m-out-of-n bootstrap (Sen, Banerjee and Woodroofe, 2010; Lee and Yang, 2020), or the numerical bootstrap (Hong and Li, 2020). See also Cavaliere and Georgiev (2020), and references therein, for some related recent results. Those approaches could in principle be used to construct bootstrap-based inference methods for (some members of the class of) generalized Grenander-type estimators, but they all would require employing specific regularized multidimensional bootstrap distributions or related quantities involving multiple smoothing and tuning parameters, rendering those approaches potentially difficult to implement in practice. Furthermore, those methods would not be robust to unknown degeneracies determining the convergence rate and shape of the limiting distribution without additional modifications. For example, subsampling methods require knowledge of the precise convergence rate of the statistic, or estimation thereof, as a preliminary step (Politis, Romano and Wolf, 1999). Another resampling approach was recently proposed by Kuchibhotla, Balakrishnan and Wasserman (2024) and Mallick, Sarkar and Kuchibhotla (2024), which offers valid confidence intervals under an approximate symmetry condition of the limiting distribution of the estimator.

We complement existing methods by introducing a novel bootstrap-assisted inference approach that restores validity of bootstrap methods by reshaping of the ingredients of the generalized Grenander-type estimator. Our approach is motivated by a constructive interpretation of the source of the bootstrap failure. As a by-product, our interpretation explicitly isolates the role of unknown degeneracies determining the precise form of the limiting distribution, a feature of the interpretation which allow us to develop an automatic inference method that is robust to such degeneracies, ultimately resulting in a more robust bootstrap-assisted inference approach. In the case of random sampling, we show that our method can be implemented using a computationally attractive exchangeable bootstrap procedure. For completeness, we also discuss implementation issues, offering a fully automatic (i.e., data-driven) valid inference method for generalized Grenander-type estimators. Some of the ideas underlying our approach are similar to ideas used in Cattaneo, Jansson and Nagasawa (2020), where we introduced a bootstrap-based distributional approximation for M-estimators with possibly non-standard Chernoff (1964)-type asymptotic distributions (Kim and Pollard, 1990; Seo and Otsu, 2018). Generalized Grenander-type estimators are not M-estimators, however, and as further explained in the next paragraph the analysis of generalized Grenander-type estimators turns out to necessitate the development of technical tools that play no role in the analysis of M-estimators.

Although valid distributional approximations for monotone estimators can be obtained in a variety of ways (Groeneboom and Jongbloed, 2014), by far the most common approach is to employ the *switch relation* (Groeneboom, 1985) to re-express the cumulative distribution function (cdf) of the suitably normalized monotone estimator in terms of a probability statement about the maximizer of a certain stochastic process whose large sample properties can in turn be analyzed by employing standard empirical process methods (van der Vaart and Wellner, 1996). From a technical perspective, this approach requires (at least) two ingredients in order to be successful, namely results establishing (i) validity of the switch relation and (ii) continuity of the limiting cdf of the maximizer of a stochastic process. In the process of developing our main results, we shed new light on both (i) and (ii).

First, we show by example that Westling and Carone's (2020, Supplement) generalization of the switch relation is incomplete as stated and then propose a modification. Our modification of Westling and Carone's (2020, Supplement) Lemma 1 shows that the conclusion of that lemma is valid once an additional assumption is made. The additional assumption seems very mild, being satisfied in all examples considered by Westling and Carone (2020) and all other examples of which we are aware, including some new examples we consider in this paper. Second, we present a lemma establishing continuity of the cdf of the maximizer of a Gaussian process, a result which can in turn be used to establish continuity of the cdf of the suitably normalized generalized Grenander-type estimator. Interestingly, although these continuity properties are important when deriving limiting distributions with the help of the switch relation and justifying bootstrap-type inference procedures, respectively, it would appear that explicit statements of them are unavailable in the existing literature. (A prominent exception is the one where the limiting distribution is a scaled Chernoff distribution, which is known to be absolutely continuous.) For further details on (i) and (ii), see Appendix A.

In the remainder of this introductory section we outline key notation and definitions used throughout the paper. Section 2 then recalls the setup of Westling and Carone (2020) and presents a version of their main distributional approximation result for generalized Grenander-type estimators. Section 3 contains our main results about bootstrap-assisted distributional approximations, while Section 4 discusses implementation issues, including tuning parameter selection and a computationally attractive weighted bootstrap procedure. Section 5 illustrates our general theory by means of prominent examples, while Section 6 reports numerical results from a small-scale simulation experiment. Appendix A reports the two technical results alluded to in the previous paragraph. All proofs and other technical details are given in the supplemental appendix (Cattaneo, Jansson and Nagasawa, 2024).

1.1. Notation and Definitions. For an interval $I \subseteq \mathbb{R}$ and a function $f: I \to \mathbb{R}$, $\mathsf{GCM}_I(f)$ denotes its greatest convex minorant (on I) and if f is non-decreasing and right-continuous, then f^- denotes its generalized inverse; that is, $f^-(x) = \inf\{u \in I: f(u) \geq x\}$, where the dependence of f^- on I has been suppressed (and where the infimum of the empty set is $\sup I$). Also, if f is non-decreasing and right-continuous, we write $f(x-) = \lim_{\eta \downarrow 0} f(x-\eta)$ and $f^-(y+) = \lim_{\eta \downarrow 0} f^-(y+\eta)$. Assuming the relevant derivatives exist, ∂^q denotes the qth partial derivative (operator) and ∂_- denotes the left derivative (operator). In addition, $f \circ g$ denotes the composition of f and g; that is, $(f \circ g)(x) = f(g(x))$. Finally, for $a, b \in \mathbb{R}$, we write $a \wedge b = \min(a, b), a \vee b = \max(a, b)$, and $a^+ = \max(a, 0)$, and $\lfloor a \rfloor$ denotes the integer part of a.

Limits are taken as $n \to \infty$, unless otherwise stated. For two (possibly) random sequences $\{a_n\}$ and $\{b_n\}$, $a_n = O_{\mathbb{P}}(b_n)$ is shorthand for $\limsup_{\epsilon \to \infty} \limsup_n \mathbb{P}[|a_n/b_n| \ge \epsilon] = 0$, $a_n = o_{\mathbb{P}}(b_n)$ is shorthand for $\limsup_{\epsilon \to 0} \limsup_n \mathbb{P}[|a_n/b_n| \ge \epsilon] = 0$, and the subscript "P" on "O" and "o" is often omitted when $\{a_n\}$ and $\{b_n\}$ are non-random. We use $\to_{\mathbb{P}}$ to denote convergence in probability and \leadsto to denote weak convergence, where, for a stochastic process indexed by \mathbb{R} , convergence is in the topology of uniform convergence on compacta. When analyzing the bootstrap, \mathbb{P}_n^* denotes the probability measure under the bootstrap distribution conditional on the original data and $\leadsto_{\mathbb{P}}$ denotes weak convergence in probability conditionally on the original data. For more details, see van der Vaart and Wellner (1996).

2. Setup. Our setup is that of Westling and Carone (2020). The goal is to conduct inference on $\theta_0(x)$, where, for some interval $I \subseteq \mathbb{R}$, θ_0 is non-decreasing on I and x is an interior point of I. Assuming it is well defined, the function Θ_0 given by

$$\Theta_0(x) = \int_{\inf I}^x \theta_0(v) dv$$

is convex on I and therefore enjoys the property that if θ_0 is left-continuous at x, then

$$\theta_0(x) = \partial_- \mathsf{GCM}_I(\Theta_0)(x).$$

An estimator of $\theta_0(x)$ obtained by replacing Θ_0 and I in the preceding display with estimators is said to be of the Grenander-type, a canonical example of this class of estimators being the celebrated Grenander (1956) estimator of a non-decreasing density.

EXAMPLE 1 (Monotone Density Estimation). Suppose $X_1, ..., X_n$ are i.i.d. copies of a continuously distributed non-negative random variable X whose density f_0 is non-decreasing on its support $[0, u_0]$, where u_0 is (possibly) unknown. For $x \in (0, u_0)$, the Grenander (1956) estimator of $f_0(x)$ is

$$\widehat{f}_n(\mathbf{x}) = \partial_{-}\mathsf{GCM}_{[0,\widehat{u}_n]}(\widehat{F}_n)(\mathbf{x}),$$

where $\widehat{u}_n = \max_{1 \leq i \leq n} X_i \vee x$ and where $\widehat{F}_n(x) = n^{-1} \sum_{i=1}^n \mathbb{1}(X_i \leq x)$ is the empirical cdf. Section 5.1 and the supplemental appendix discuss more general monotone density estimators allowing for censoring and/or covariate adjustment (e.g., van der Laan and Robins, 2003).

To define the class of generalized Grenander-type estimators, let $\psi_0 = \theta_0 \circ \Phi_0^-$, where Φ_0 is non-negative, non-decreasing, and continuous on I. Defining

$$\Gamma_0 = \Psi_0 \circ \Phi_0, \qquad \Psi_0(x) = \int_0^x \psi_0(v) dv,$$

and assuming that $\Phi_0(x) < \Phi_0(x) < u_0$ for every x < x, we have

(1)
$$\theta_0(\mathbf{x}) = \partial_{-}\mathsf{GCM}_{[0,u_0]}(\Gamma_0 \circ \Phi_0^-) \circ \Phi_0(\mathbf{x})$$

whenever θ_0 is left-continuous at x. In the terminology of Westling and Carone (2020), an estimator of $\theta_0(x)$ is of the Generalized Grenander-type if it is obtained by replacing Γ_0 , Φ_0 , and u_0 in the preceding display with estimators $\widehat{\Gamma}_n$, $\widehat{\Phi}_n$, and \widehat{u}_n (say); to be specific, an estimator of the Generalized Grenander-type is of the form

$$\widehat{\theta}_n(\mathbf{x}) = \partial_-\mathsf{GCM}_{[0,\widehat{u}_n]}(\widehat{\Gamma}_n \circ \widehat{\Phi}_n^-) \circ \widehat{\Phi}_n(\mathbf{x}),$$

where $\widehat{\Phi}_n$ is non-negative, non-decreasing, and right-continuous. Of course, Grenander-type estimators are of the generalized Grenander-type (with $\widehat{\Phi}_n$ equal to the identity mapping) whenever the associated estimator of I is of the form $[0,\widehat{u}_n]$, but the class of generalized Grenander-type estimators contains many important estimators that are not of Grenander-type, a canonical example being the celebrated isotonic regression estimator of Brunk (1958).

EXAMPLE 2 (Monotone Regression Estimation). Suppose $(Y_1, X_1), \ldots, (Y_n, X_n)$ are i.i.d. copies of (Y, X), where X is a continuously distributed random variable and where the regression function $\mu_0(x) = \mathbb{E}(Y|X=x)$ is non-decreasing. For x in the interior of the support of X, the Brunk (1958) estimator of $\mu_0(x)$ is

$$\widehat{\mu}_n(\mathsf{x}) = \partial_-\mathsf{GCM}_{[0,1]}(\widehat{\Gamma}_n \circ \widehat{F}_n^-) \circ \widehat{F}_n(\mathsf{x}),$$

where $\widehat{\Gamma}_n(x) = n^{-1} \sum_{i=1}^n Y_i \mathbb{1}(X_i \leq x)$ and where $\widehat{F}_n(x) = n^{-1} \sum_{i=1}^n \mathbb{1}(X_i \leq x)$ is an estimator of F_0 , the cdf of X. Section 5.2 discusses more general monotone regression estimators allowing for covariate adjustment (e.g., Westling, Gilbert and Carone, 2020).

Under regularity conditions, if $\widehat{\theta}_n(x)$ is of the generalized Grenander-type, then its rate of convergence is governed by the flatness of θ_0 around x, as measured by the characteristic exponent

$$\mathfrak{q} = \min\{j \in \mathbb{N} : \partial^j \theta_0(\mathsf{x}) \neq 0\},\$$

where \mathbb{N} is the set of positive integers. (When θ_0 is non-decreasing and suitably smooth, q is necessarily an odd integer.) To be specific, Westling and Carone (2020, Theorem 3) gave conditions under which

$$(2) r_n(\widehat{\theta}_n(\mathsf{x}) - \theta_0(\mathsf{x})) \leadsto \frac{1}{\partial \Phi_0(\mathsf{x})} \partial_-\mathsf{GCM}_{\mathbb{R}}(\mathcal{G}_\mathsf{x} + \mathcal{M}_\mathsf{x}^{\mathfrak{q}})(0), r_n = n^{\mathfrak{q}/(1+2\mathfrak{q})},$$

where \mathcal{G}_x is a scalar multiple of two-sided Brownian motion and where $\mathcal{M}_x^\mathfrak{q}$ is a monomial given by

(3)
$$\mathcal{M}_{\mathsf{x}}^{\mathsf{q}}(v) = \frac{\partial^{\mathsf{q}} \theta_0(\mathsf{x}) \partial \Phi_0(\mathsf{x})}{(\mathsf{q}+1)!} v^{\mathsf{q}+1}.$$

In addition to governing the rate of convergence, the characteristic exponent q also governs the shape of $\mathcal{M}_{x}^{\mathfrak{q}}$. On the other hand, and as the notation suggests, the covariance kernel of \mathcal{G}_x does not depend on \mathfrak{q} . If $\mathfrak{q}=1$, then the distribution of $\partial_-\mathsf{GCM}_\mathbb{R}(\mathcal{G}_x+\mathcal{M}_x^{\mathfrak{q}})(0)$ is a scaled Chernoff distribution. More generally, the distribution of $\partial_{-}GCM_{\mathbb{R}}(\mathcal{G}_{x}+\mathcal{M}_{x}^{\mathfrak{q}})(0)$ is a scaled Chernoff-type distribution (in the terminology of Han and Kato, 2022).

Among other things, the following assumption guarantees validity of the representation (1) and ensures existence of the right hand side of (3).

ASSUMPTION A. For some $\delta > 0$ and some $\mathfrak{s} \geq \mathfrak{q}$, the following are satisfied:

- (A1) $I \subseteq \mathbb{R}$ is an interval and $I_{\mathsf{x}}^{\delta} = \{x \in \mathbb{R} : |x \mathsf{x}| \leq \delta\} \subseteq I$. (A2) θ_0 is non-decreasing and bounded on I, and $\lfloor \mathfrak{s} \rfloor$ times continuously differentiable on I_{x}^{δ} with

$$\sup_{x,x'\in I^{\delta}}\frac{|\partial^{\lfloor \mathfrak{s}\rfloor}\theta_{0}(x)-\partial^{\lfloor \mathfrak{s}\rfloor}\theta_{0}(x')|}{|x-x'|^{\mathfrak{s}-\lfloor \mathfrak{s}\rfloor}}<\infty.$$

(A3) Φ_0 is non-negative, non-decreasing, continuous, and bounded on I, and $\lfloor \mathfrak{s} \rfloor - \mathfrak{q} + 1$ times continuously differentiable on I_x^{δ} with $\partial \Phi_0(x) \neq 0$ and

$$\sup_{x,x'\in I_{\delta}^{\delta}}\frac{\left|\partial^{\left\lfloor \mathfrak{s}\right\rfloor -\mathfrak{q}+1}\Phi_{0}(x)-\partial^{\left\lfloor \mathfrak{s}\right\rfloor -\mathfrak{q}+1}\Phi_{0}(x')\right|}{\left|x-x'\right|^{\mathfrak{s}-\left\lfloor \mathfrak{s}\right\rfloor}}<\infty.$$

3. Bootstrap-Assisted Distributional Approximation. The distributional approximation (2) motivates a plug-in procedure for statistical inference: with consistent estimators of $\mathcal{M}_{\mathsf{x}}^{\mathfrak{q}}$ and the covariance kernel of \mathcal{G}_{x} , the limiting law in (2) can be simulated. This approach requires characterizing the covariance kernel explicitly, and then forming a consistent estimator based on preliminary non-parametric smoothing methods (in addition to the other preliminary nonparametric estimators needed). Also, simulating from the limiting law requires specifying or estimating the unknown characteristic exponent q. As a consequence, although conceptually feasible, the plug-in approach is often computationally cumbersome to employ in practice.

Since resampling methods have the potential to provide automatic and robust distributional approximations, it seems natural to consider such methods as an alternative to plug-in methods. Letting $(\widehat{\Gamma}_n^*, \widehat{\Phi}_n^*, \widehat{u}_n^*)$ denote a generic (not necessarily nonparametric) bootstrap analog of $(\widehat{\Gamma}_n, \widehat{\Phi}_n, \widehat{u}_n)$ and assuming that $\widehat{\Phi}_n^*$ is non-decreasing and right-continuous, the associated bootstrap analog of $\widehat{\theta}_n(\mathsf{x})$ is

$$\widehat{\theta}_n^*(\mathbf{x}) = \partial_-\mathsf{GCM}_{[0,\widehat{u}_n^*]}(\widehat{\boldsymbol{\Gamma}}_n^* \circ \widehat{\boldsymbol{\Phi}}_n^{*-}) \circ \widehat{\boldsymbol{\Phi}}_n^*(\mathbf{x}).$$

As is well documented for the classical Grenander estimator (e.g. Kosorok, 2008; Sen, Banerjee and Woodroofe, 2010), the bootstrap analog of (2) does not (necessarily) hold when $(\widehat{\Gamma}_n^*, \widehat{\Phi}_n^*, \widehat{u}_n^*)$ is obtained by means of the nonparametric bootstrap. In other words,

$$r_n(\widehat{\theta}_n^*(\mathbf{x}) - \widehat{\theta}_n(\mathbf{x})) \not \rightsquigarrow_{\mathbb{P}} \frac{1}{\partial \Phi_0(\mathbf{x})} \partial_{-}\mathsf{GCM}_{\mathbb{R}}(\mathcal{G}_{\mathbf{x}} + \mathcal{M}_{\mathbf{x}}^{\mathfrak{q}})(0)$$

in general; that is, the bootstrap is inconsistent in general. (A precise statement is provided in Theorem A.1 in Appendix A.3.) It turns out, however, that under plausible conditions on $(\widehat{\Gamma}_n^*, \widehat{\Phi}_n^*, \widehat{u}_n^*)$, a valid bootstrap-based distributional approximation can be obtained by employing

$$\widetilde{\theta}_n^*(\mathbf{x}) = \partial_-\mathsf{GCM}_{[0,\widehat{u}_n^*]}(\widetilde{\Gamma}_n^* \circ \widehat{\Phi}_n^{*-}) \circ \widehat{\Phi}_n^*(\mathbf{x}),$$

where, for some judiciously chosen $\widetilde{M}_{x,n}$,

$$\widetilde{\Gamma}_n^*(x) = \widehat{\Gamma}_n^*(x) - \widehat{\Gamma}_n(x) + \widehat{\theta}_n(\mathbf{x}) \widehat{\Phi}_n(x) + \widetilde{M}_{\mathbf{x},n}(x-\mathbf{x}).$$

(As the notation suggests, and as further discussed below, a defining property of $\widetilde{M}_{x,n}$ is that a suitably re-scaled version of it can be interpreted as an estimator of \mathcal{M}_x^q .)

3.1. Heuristics. To explain the source of the bootstrap failure and motivate the functional form of $\widetilde{\Gamma}_n^*$, it is useful to begin by sketching the derivation of (2). For any $t \in \mathbb{R}$ and under mild conditions on $\widehat{\Gamma}_n$ and $\widehat{\Phi}_n$, it follows from the so-called (generalized) switch relation (e.g., Appendix A.1) that

$$\mathbb{P}\left[r_n(\widehat{\theta}_n(\mathsf{x}) - \theta_0(\mathsf{x})) \leq t\right] = \mathbb{P}\left[\underset{v \in \widehat{V}_{\mathsf{x},n}^{\mathfrak{q}}}{\operatorname{argmin}} \{\widehat{G}_{\mathsf{x},n}^{\mathfrak{q}}(v) + M_{\mathsf{x},n}^{\mathfrak{q}}(v) - t\widehat{L}_{\mathsf{x},n}^{\mathfrak{q}}(v)\} \geq \widehat{Z}_{\mathsf{x},n}^{\mathfrak{q}}\right],$$

where $\widehat{V}_{\mathbf{x},n}^{\mathfrak{q}}$ is a "large" subset of \mathbb{R} , $\widehat{G}_{\mathbf{x},n}^{\mathfrak{q}}$ is a stochastic process given by

$$\begin{split} \widehat{G}_{\mathbf{x},n}^{\mathfrak{q}}(v) &= \sqrt{na_n} [\widehat{\Gamma}_n(\mathbf{x} + va_n^{-1}) - \widehat{\Gamma}_n(\mathbf{x}) - \Gamma_0(\mathbf{x} + va_n^{-1}) + \Gamma_0(\mathbf{x})] \\ &- \theta_0(\mathbf{x}) \sqrt{na_n} [\widehat{\Phi}_n(\mathbf{x} + va_n^{-1}) - \widehat{\Phi}_n(\mathbf{x}) - \Phi_0(\mathbf{x} + va_n^{-1}) + \Phi_0(\mathbf{x})], \quad a_n = n^{1/(1+2\mathfrak{q})}, \end{split}$$

 $M_{\mathsf{x},n}^{\mathfrak{q}}$ is a smooth non-random function, $\widehat{L}_{\mathsf{x},n}^{\mathfrak{q}}$ can be interpreted as an estimator of the linear function \mathcal{L}_{x} given by $\mathcal{L}_{\mathsf{x}}(v) = v \partial \Phi_0(\mathsf{x})$, and where $\widehat{Z}_{\mathsf{x},n}^{\mathfrak{q}}$ is zero when $\widehat{\Phi}_n$ is the identity mapping and can be interpreted as an "estimator" of zero more generally. (The exact definitions of $\widehat{V}_{\mathsf{x},n}^{\mathfrak{q}}, M_{\mathsf{x},n}^{\mathfrak{q}}, \widehat{L}_{\mathsf{x},n}^{\mathfrak{q}}$, and $\widehat{Z}_{\mathsf{x},n}^{\mathfrak{q}}$ are given in Appendix A.4.) In the above and the sequel, the superscript \mathfrak{q} indicates that the superscripted object is defined on a "localized" domain.

Under mild conditions, $\widehat{G}_{\mathsf{x},n}^{\mathfrak{q}}$ converges weakly to a scalar multiple of two-sided Brownian motion \mathcal{G}_{x} (say), while $M_{\mathsf{x},n}^{\mathfrak{q}}, \widehat{L}_{\mathsf{x},n}^{\mathfrak{q}}$, and $\widehat{Z}_{\mathsf{x},n}^{\mathfrak{q}}$ converge in probability to $\mathcal{M}_{\mathsf{x}}^{\mathfrak{q}}, \mathcal{L}_{\mathsf{x}}$, and zero, respectively. In other words, we would expect that

(4)
$$(\widehat{G}_{x,n}^{\mathfrak{q}}, \widehat{L}_{x,n}^{\mathfrak{q}}, \widehat{Z}_{x,n}^{\mathfrak{q}}) \leadsto (\mathcal{G}_{x}, \mathcal{L}_{x}, 0),$$

and that

$$(5) M_{\mathsf{x},n}^{\mathfrak{q}} \rightsquigarrow \mathcal{M}_{\mathsf{x}}^{\mathfrak{q}}.$$

It therefore stands to reason that under mild additional conditions (including suitable convergence of $\widehat{V}_{x,n}^{\mathfrak{q}}$ to \mathbb{R}), we have

$$\begin{split} \lim_{n \to \infty} \mathbb{P}\left[r_n(\widehat{\theta}_n(\mathsf{x}) - \theta_0(\mathsf{x})) \leq t\right] &= \mathbb{P}\left[\underset{v \in \mathbb{R}}{\operatorname{argmin}} \{\mathcal{G}_\mathsf{x}(v) + \mathcal{M}_\mathsf{x}^\mathsf{q}(v) - t\mathcal{L}_\mathsf{x}(v)\} \geq 0\right] \\ &= \mathbb{P}\left[\frac{1}{\partial \Phi_0(\mathsf{x})} \partial_- \mathsf{GCM}_\mathbb{R} (\mathcal{G}_\mathsf{x} + \mathcal{M}_\mathsf{x}^\mathsf{q})(0) \leq t\right], \end{split}$$

where the first equality follows from an argmax continuous mapping theorem, and the second equality is obtained by another application of the switch relation.

Next, consider the bootstrap analog of $\widehat{\theta}_n(x)$. For any $t \in \mathbb{R}$, the switch relation gives

$$\mathbb{P}_n^* \left[r_n(\widehat{\theta}_n^*(\mathsf{x}) - \widehat{\theta}_n(\mathsf{x})) \le t \right] = \mathbb{P}_n^* \left[\underset{v \in \widehat{V}_{\mathsf{x},n}^{\mathfrak{q},*}}{\operatorname{argmin}} \{ \widehat{G}_{\mathsf{x},n}^{\mathfrak{q},*}(v) + \widehat{M}_{\mathsf{x},n}^{\mathfrak{q}}(v) - t \widehat{L}_{\mathsf{x},n}^{\mathfrak{q},*}(v) \} \ge \widehat{Z}_{\mathsf{x},n}^{\mathfrak{q},*} \right],$$

where

$$\begin{split} \widehat{G}_{\mathbf{x},n}^{\mathfrak{q},*}(v) &= \sqrt{na_n} [\widehat{\Gamma}_n^*(\mathbf{x} + va_n^{-1}) - \widehat{\Gamma}_n^*(\mathbf{x}) - \widehat{\Gamma}_n(\mathbf{x} + va_n^{-1}) + \widehat{\Gamma}_n(\mathbf{x})] \\ &- \widehat{\theta}_n(\mathbf{x}) \sqrt{na_n} [\widehat{\Phi}_n^*(\mathbf{x} + va_n^{-1}) - \widehat{\Phi}_n^*(\mathbf{x}) - \widehat{\Phi}_n(\mathbf{x} + va_n^{-1}) + \widehat{\Phi}_n(\mathbf{x})], \end{split}$$

and where $\widehat{V}_{\mathsf{x},n}^{\mathfrak{q},*}, \widehat{M}_{\mathsf{x},n}^{\mathfrak{q}}, \widehat{L}_{\mathsf{x},n}^{\mathfrak{q},*}$, and $\widehat{Z}_{\mathsf{x},n}^{\mathfrak{q},*}$ are bootstrap analogs of $\widehat{V}_{\mathsf{x},n}^{\mathfrak{q}}, M_{\mathsf{x},n}^{\mathfrak{q}}, \widehat{L}_{\mathsf{x},n}^{\mathfrak{q}}$, and $\widehat{Z}_{\mathsf{x},n}^{\mathfrak{q}}$, respectively. (The exact definitions of $\widehat{V}_{\mathsf{x},n}^{\mathfrak{q},*}, \widehat{M}_{\mathsf{x},n}^{\mathfrak{q}}, \widehat{L}_{\mathsf{x},n}^{\mathfrak{q},*}$, and $\widehat{Z}_{\mathsf{x},n}^{\mathfrak{q},*}$ are given in Appendix A.4.) If (4) holds, then the following bootstrap counterpart thereof can also be expected to hold:

(6)
$$(\widehat{G}_{\mathsf{x},n}^{\mathfrak{q},*}, \widehat{L}_{\mathsf{x},n}^{\mathfrak{q},*}, \widehat{Z}_{\mathsf{x},n}^{\mathfrak{q},*}) \leadsto_{\mathbb{P}} (\mathcal{G}_{\mathsf{x}}, \mathcal{L}_{\mathsf{x}}, 0).$$

On the other hand, unlike $M_{\mathbf{x},n}^{\mathfrak{q}}$ the process $\widehat{M}_{\mathbf{x},n}^{\mathfrak{q}}$ is (random and) non-smooth, so the bootstrap counterpart of (5) typically fails, implying in turn that the bootstrap is inconsistent; for details, see Theorem A.1 (stated in Appendix A.3) and its proof.

In other words, the sole source of the bootstrap inconsistency is the failure of $\widehat{M}_{\mathbf{x},n}^{\mathfrak{q}}$ to replicate the salient properties of $M_{\mathbf{x},n}^{\mathfrak{q}}$. Recognizing this, our proposed estimator $\widehat{\theta}_n^*(\mathbf{x})$ has been carefully constructed so that it differs from $\widehat{\theta}_n^*(\mathbf{x})$ only in terms of the implied estimator of $\mathcal{M}_{\mathbf{x}}^{\mathfrak{q}}$. To be specific, $\widetilde{\theta}_n^*(\mathbf{x})$ is similar to $\widehat{\theta}_n^*(\mathbf{x})$ insofar as it satisfies the following switch relation: For any $t \in \mathbb{R}$,

$$\mathbb{P}_n^* \left[r_n(\widetilde{\theta}_n^*(\mathsf{x}) - \widehat{\theta}_n(\mathsf{x})) \le t \right] = \mathbb{P}_n^* \left[\underset{v \in \widehat{V}_{\mathsf{x},n}^{\mathfrak{q},*}}{\operatorname{argmin}} \{ \widehat{G}_{\mathsf{x},n}^{\mathfrak{q},*}(v) + \widetilde{M}_{\mathsf{x},n}^{\mathfrak{q}}(v) - t \widehat{L}_{\mathsf{x},n}^{\mathfrak{q},*}(v) \} \ge \widehat{Z}_{\mathsf{x},n}^{\mathfrak{q},*} \right],$$

where $\widetilde{M}_{x,n}^{\mathfrak{q}}$ is the following transformation of $\widetilde{M}_{x,n}$:

$$\widetilde{M}_{\mathsf{x},n}^{\mathfrak{q}}(v) = \sqrt{na_n}\widetilde{M}_{\mathsf{x},n}(va_n^{-1}).$$

As a consequence, if (6) holds and if $\widetilde{M}_{x,n}^{\mathfrak{q}} \rightsquigarrow_{\mathbb{P}} \mathcal{M}_{x}^{\mathfrak{q}}$, then it stands to reason that under mild additional conditions $\widetilde{\theta}_{n}^{*}(x)$ satisfies the following bootstrap counterpart of (2):

(7)
$$r_n(\widetilde{\theta}_n^*(\mathsf{x}) - \widehat{\theta}_n(\mathsf{x})) \leadsto_{\mathbb{P}} \frac{1}{\partial \Phi_0(\mathsf{x})} \partial_{-}\mathsf{GCM}_{\mathbb{R}}(\mathcal{G}_\mathsf{x} + \mathcal{M}_\mathsf{x}^{\mathfrak{q}})(0).$$

In words, our proposed bootstrap-assisted inference procedure can be explained as follows: we employ the bootstrap to approximate the "fluctuations" of the process \mathcal{G}_x , while we use the estimator $\widetilde{M}_{x,n}^{\mathfrak{q}}$ to approximate $\mathcal{M}_x^{\mathfrak{q}}$. It is an interesting question whether our approach offers provably better distributional approximations than the plug-in procedure mentioned previously, but it is beyond the scope of this paper to answer that question.

3.2. Main result. Our heuristic derivation of (2) can be made rigorous by providing conditions under which four properties hold. First, the switch relation(s) must be valid. Second, the convergence properties (4) and (5) must hold. Third, to use (4) and (5) to obtain the result

$$\underset{v \in \widehat{V}_{x,n}^{\mathfrak{q}}}{\operatorname{argmin}} \{ \widehat{G}_{\mathsf{x},n}^{\mathfrak{q}}(v) + M_{\mathsf{x},n}^{\mathfrak{q}}(v) - t \widehat{L}_{\mathsf{x},n}^{\mathfrak{q}}(v) \} \leadsto \underset{v \in \mathbb{R}}{\operatorname{argmin}} \{ \mathcal{G}_{\mathsf{x}}(v) + \mathcal{M}_{\mathsf{x}}^{\mathfrak{q}}(v) - t \mathcal{L}_{\mathsf{x}}(v) \}$$

with the help of a suitable argmax continuous mapping theorem, tightness of the left hand side in the previous display must hold. Finally, to furthermore obtain the conclusion

$$\lim_{n \to \infty} \mathbb{P} \left[\underset{v \in \widehat{V}_{\mathbf{x},n}^{\mathfrak{q}}}{\operatorname{argmin}} \{ \widehat{G}_{\mathbf{x},n}^{\mathfrak{q}}(v) + M_{\mathbf{x},n}^{\mathfrak{q}}(v) - t \widehat{L}_{\mathbf{x},n}^{\mathfrak{q}}(v) \} \ge \widehat{Z}_{\mathbf{x},n}^{\mathfrak{q}} \right]$$

$$= \mathbb{P} \left[\underset{v \in \mathbb{R}}{\operatorname{argmin}} \{ \mathcal{G}_{\mathbf{x}}(v) + \mathcal{M}_{\mathbf{x}}^{\mathfrak{q}}(v) - t \mathcal{L}_{\mathbf{x}}(v) \} \ge 0 \right],$$

the cdf of $\operatorname{argmin}_{v \in \mathbb{R}} \{ \mathcal{G}_{\mathsf{x}}(v) + \mathcal{M}_{\mathsf{x}}^{\mathfrak{q}}(v) - t\mathcal{L}_{\mathsf{x}}(v) \}$ must be continuous at zero.

Conditions under which the second and third properties hold can be formulated with the help of well known empirical process results. We base our formulations on van der Vaart and Wellner (1996) and employ the generalized argmax continuous mapping theorem of Cox (2022), which applies also to settings where the feasibility set varies with n. The first and fourth properties, on the other hand, seem more difficult to verify. Regarding the first property, it turns out that the generalization of the switch relation employed by Westling and Carone (2020) requires additional conditions in order to be valid. To address this concern, we present a modified generalized switch relation whose assumptions include a condition not present in Lemma 1 of Westling and Carone (2020, Supplement). Thankfully, the condition in question seems very mild and having imposed it we are able to preserve the main implication of Lemma 1 of Westling and Carone (2020, Supplement); for details, see Lemma A.1 in Appendix A.1. In the special case where q = 1, the fourth property follows from well known properties of the Chernoff distribution. More generally, however, we are unaware of existing results guaranteeing the requisite continuity property when $q \neq 1$, but fortunately it turns out that the continuity property of interest can be established; for details, see Lemma A.2 in Appendix A.2.

The following assumption collects the conditions under which our verification of the four above-mentioned properties will proceed.

ASSUMPTION B. For some $\delta > 0$, the following are satisfied:

(B1) $\widehat{G}_{x,n}^{\mathfrak{q}} \leadsto \mathcal{G}_x$ and $\widehat{G}_{x,n}^{\mathfrak{q},*} \leadsto_{\mathbb{P}} \mathcal{G}_x$, where \mathcal{G}_x is a centered Gaussian process whose covariance function is $C_{\mathsf{x}}(s,t) = C(|s| \land |t|) \mathbb{1}(\operatorname{sign}(s) = \operatorname{sign}(t))$ for some C > 0.

(B2) There exist $\beta < \mathfrak{q} + 1$ and events A_n with $\lim_{n \to \infty} \mathbb{P}[A_n] = 1$,

$$\sup_{V \in [1, a_n \delta]} \mathbb{E} \left[V^{-\beta} \sup_{|v| \in [V, 2V]} |\widehat{G}_{\mathsf{x}, n}^{\mathfrak{q}}(v)| \mathbb{1}_{\mathcal{A}_n} \right] = O(1)$$

and

$$\sup_{V \in [1, a_n \delta]} \mathbb{E} \left[V^{-\beta} \sup_{|v| \in [V, 2V]} |\widehat{G}_{\mathsf{x}, n}^{\mathfrak{q}, *}(v)| \mathbb{1}_{\mathcal{A}_n} \right] = O(1).$$

(B3)
$$\sup_{x \in I} |\widehat{\Gamma}_n(x) - \Gamma_0(x)| = o_{\mathbb{P}}(1) \text{ and } \sup_{x \in I} |\widehat{\Gamma}_n^*(x) - \widehat{\Gamma}_n(x)| = o_{\mathbb{P}}(1).$$

$$\begin{array}{ll} \text{(B3)} & \sup_{x\in I}|\widehat{\Gamma}_n(x)-\Gamma_0(x)|=o_{\mathbb{P}}(1) \text{ and } \sup_{x\in I}|\widehat{\Gamma}_n^*(x)-\widehat{\Gamma}_n(x)|=o_{\mathbb{P}}(1). \\ \text{(B4)} & \sup_{x\in I}|\widehat{\Phi}_n(x)-\Phi_0(x)|=o_{\mathbb{P}}(1) \text{ and } \sup_{x\in I}|\widehat{\Phi}_n^*(x)-\widehat{\Phi}_n(x)|=o_{\mathbb{P}}(1). \end{array}$$

$$a_n \sup_{x \in I_{\mathbb{X}}^{\delta}} |\widehat{\Phi}_n(x) - \Phi_0(x)| = o_{\mathbb{P}}(1) \quad \text{and} \quad a_n \sup_{x \in I_{\mathbb{X}}^{\delta}} |\widehat{\Phi}_n^*(x) - \widehat{\Phi}_n(x)| = o_{\mathbb{P}}(1).$$

- (B5) For some $u_0 > \Phi_0(x)$, $\widehat{u}_n \ge u_0 + o_{\mathbb{P}}(1)$ and $\widehat{u}_n^* \ge \widehat{u}_n + o_{\mathbb{P}}(1)$.
- (B6) $\widehat{\Phi}_n$ and $\widehat{\Phi}_n^*$ are non-negative, non-decreasing, and right-continuous on I. In addition, $\{0, \widehat{u}_n\} \subseteq \widehat{\Phi}_n(I)$ and $\{0, \widehat{u}_n^*\} \subseteq \widehat{\Phi}_n^*(I)$. Also, $\widehat{\Phi}_n(I) \cap [0, \widehat{u}_n]$ and $\widehat{\Phi}_n^*(I) \cap [0, \widehat{u}_n^*]$ are closed.

(B7)

$$\sqrt{na_n} \sup_{x \in I_s^\delta} |\widehat{\Phi}_n(x) - \widehat{\Phi}_n(x-)| = o_{\mathbb{P}}(1) \quad \text{and} \quad \sqrt{na_n} \sup_{x \in I_s^\delta} |\widehat{\Phi}_n^*(x) - \widehat{\Phi}_n^*(x-)| = o_{\mathbb{P}}(1).$$

Verification of the bootstrap parts of Assumption B will be discussed in Section 4.2 below. When combined with Assumption A, Assumption B suffices in order to establish (2) and (6). In addition, (7) can be shown to hold if $\widetilde{M}_{x,n}$ satisfies the following

ASSUMPTION C. $\widetilde{M}_{x,n}^{\mathfrak{q}} \leadsto_{\mathbb{P}} \mathcal{M}_{x}^{\mathfrak{q}}$ and, for some c > 0 and every K > 0,

$$\liminf_{n \to \infty} \mathbb{P}\left[\inf_{|v| > K^{-1}} \widetilde{M}_{\mathsf{x},n}(v) \ge cK^{-(\mathfrak{q}+1)}\right] = 1.$$

Moreover, continuity of the cdf of $\partial_-\mathsf{GCM}_\mathbb{R}(\mathcal{G}_\mathsf{x}+\mathcal{M}_\mathsf{x}^\mathfrak{q})(0)$ can be shown with the help of the arguments used to show that the cdf of $\mathrm{argmin}_{v\in\mathbb{R}}\{\mathcal{G}_\mathsf{x}(v)+\mathcal{M}_\mathsf{x}^\mathfrak{q}(v)-t\mathcal{L}_\mathsf{x}(v)\}$ is continuous at zero. As a consequence, we obtain the following result.

THEOREM 1. Suppose Assumptions A, B, and C are satisfied. Then (2) and (7) hold, and

(8)
$$\sup_{t \in \mathbb{R}} \left| \mathbb{P}_n^* \left[\widetilde{\theta}_n^*(\mathsf{x}) - \widehat{\theta}_n(\mathsf{x}) \le t \right] - \mathbb{P} \left[\widehat{\theta}_n(\mathsf{x}) - \theta_0(\mathsf{x}) \le t \right] \right| = o_{\mathbb{P}}(1).$$

In an attempt to emphasize the rate-adaptive nature of the consistency property enjoyed by the bootstrap-based distributional approximation based on $\widetilde{\theta}_n^*(x)$, the formulation (8) deliberately omits the rate term r_n present in (2) and (7). Theorem 1 has immediate implications for inference. For instance, it follows from (2) and (8) that for any $\alpha \in (0,1)$, we have

$$\lim_{n\to\infty} \mathbb{P}[\theta_0(\mathbf{x}) \in \mathsf{CI}_{1-\alpha,n}(\mathbf{x})] = 1-\alpha,$$

 $\text{ where, defining } Q_{a,n}^*(\mathbf{x}) = \inf\{Q \in \mathbb{R}: \mathbb{P}_n^*[\widetilde{\theta}_n^*(\mathbf{x}) - \widehat{\theta}_n(\mathbf{x}) \leq Q] \geq a\},$

$$\mathsf{CI}_{1-\alpha,n}(\mathsf{x}) = \left[\widehat{\theta}_n(\mathsf{x}) - Q_{1-\alpha/2,n}^*(\mathsf{x}) \;,\; \widehat{\theta}_n(\mathsf{x}) - Q_{\alpha/2,n}^*(\mathsf{x})\right]$$

is the (nominal) level $1 - \alpha$ bootstrap confidence interval for θ_0 based on the "percentile method" (in the terminology of van der Vaart, 1998).

4. Implementation. Suppose Assumption A is satisfied and suppose $(\widehat{\Gamma}_n, \widehat{\Phi}_n, \widehat{u}_n)$ satisfies the non-bootstrap parts of Assumption B. Then, in order to compute the estimator $\widetilde{\theta}_n^*(\mathbf{x})$ upon which our proposed bootstrap-assisted distributional approximation is based, two implementation issues must be addressed, namely the choice/construction of $\widetilde{M}_{\mathbf{x},n}$ and $(\widehat{\Gamma}_n^*, \widehat{\Phi}_n^*, \widehat{u}_n^*)$, respectively. Section 4.1 demonstrates the plausibility of Assumption C by exhibiting estimators $\widetilde{M}_{\mathbf{x},n}$ satisfying it under Assumptions A and D, the latter being a high-level condition that typically holds whenever the non-bootstrap parts of Assumption B hold. Then, Section 4.2 exhibits easy-to-compute $(\widehat{\Gamma}_n^*, \widehat{\Phi}_n^*, \widehat{u}_n^*)$ satisfying the bootstrap parts of Assumption B under a random sampling assumption and other mild conditions.

4.1. Mean Function Estimation. The ease with which an $\widetilde{M}_{x,n}$ satisfying Assumption C can be constructed depends in particular on whether q is known. To facilitate the subsequent discussion, let $\mathcal{D}_0(x) = 0$ and for $j = 1, ..., |\mathfrak{s}|$, define (recursively)

$$\mathcal{D}_{j}(\mathbf{x}) = \lim_{x \to \mathbf{x}} \frac{\Upsilon_{0}(x) - \Upsilon_{0}(\mathbf{x}) - \sum_{k=0}^{j-1} [(k+1)!]^{-1} \mathcal{D}_{k}(\mathbf{x}) (x-\mathbf{x})^{k+1}}{(x-\mathbf{x})^{j+1}}, \quad \Upsilon_{0} = \Gamma_{0} - \theta_{0}(\mathbf{x}) \Phi_{0}.$$

Noting that $\partial \Gamma_0(x) = \theta_0(x) \partial \Phi_0(x)$ for x near x, we have $\mathcal{D}_0(x) = \cdots = \mathcal{D}_{\mathfrak{q}-1}(x) = 0$ and

$$\mathcal{D}_{j}(\mathsf{x}) = \frac{1}{(j+1)!} \sum_{k=\mathfrak{q}}^{j} \binom{j}{k} \partial^{k} \theta_{0}(\mathsf{x}) \partial^{j+1-k} \Phi_{0}(\mathsf{x}), \qquad j = \mathfrak{q}, \dots, \lfloor \mathfrak{s} \rfloor.$$

In particular, $\mathcal{D}_{\mathfrak{q}}(\mathsf{x}) = \partial^{\mathfrak{q}}\theta_0(\mathsf{x})\partial\Phi_0(\mathsf{x})/(\mathfrak{q}+1)!$ and therefore $\mathcal{M}_\mathsf{x}^{\mathfrak{q}}(v) = \mathcal{D}_{\mathfrak{q}}(\mathsf{x})v^{\mathfrak{q}+1}$.

4.1.1. *Mean Function Estimation with Known* \mathfrak{q} . First, consider the (simpler) case where \mathfrak{q} is known. In this case, if

(9)
$$\widetilde{\mathcal{D}}_{\mathfrak{q},n}(\mathsf{x}) \to_{\mathbb{P}} \mathcal{D}_{\mathfrak{q}}(\mathsf{x}),$$

then Assumption C holds when

(10)
$$\widetilde{M}_{\mathsf{x},n}(x) = \widetilde{\mathcal{D}}_{\mathsf{q},n}(\mathsf{x})x^{\mathsf{q}+1}.$$

Example-specific estimators $\widetilde{\mathcal{D}}_{\mathfrak{q},n}(\mathsf{x})$ satisfying the consistency requirement (9) may be readily available. For instance, in the case of the Grenander (1956) estimator described in Example 1, we have $\mathcal{D}_{\mathfrak{q}}(\mathsf{x}) = \partial^{\mathfrak{q}} f_0(\mathsf{x})/(\mathfrak{q}+1)!$, a consistent estimator of which can be based on any consistent estimator of $\partial^{\mathfrak{q}} f_0(\mathsf{x})$, such as a standard kernel estimator or, if the evaluation point x is near the boundary of the support of X, boundary adaptive versions thereof.

More generic estimators are also available. For specificity, the remainder of this section focuses on estimators of $\mathcal{D}_j(x)$ obtained by applying numerical derivative-type operators to the following (possibly) non-smooth estimator of Υ_0 :

$$\widehat{\Upsilon}_n = \widehat{\Gamma}_n - \widehat{\theta}_n(\mathsf{x})\widehat{\Phi}_n.$$

The fact that $\mathcal{D}_j(\mathsf{x}) = 0$ for $j < \mathfrak{q}$ implies that $\mathcal{D}_{\mathfrak{q}}(\mathsf{x})$ admits the "monomial approximation" representation

$$\mathcal{D}_{\mathfrak{q}}(\mathsf{x}) = \lim_{\epsilon \to 0} \left\{ \epsilon^{-(\mathfrak{q}+1)} [\Upsilon_0(\mathsf{x}+\epsilon) - \Upsilon_0(\mathsf{x})] \right\},$$

motivating the estimator

$$\widetilde{\mathcal{D}}_{\mathfrak{q},n}^{\mathtt{MA}}(\mathbf{x}) = \epsilon_n^{-(\mathfrak{q}+1)} [\widehat{\Upsilon}_n(\mathbf{x}+\epsilon_n) - \widehat{\Upsilon}_n(\mathbf{x})],$$

where $\epsilon_n > 0$ is a tuning parameter. Similarly, the generic "forward difference" representation

$$\mathcal{D}_{\mathfrak{q}}(\mathbf{x}) = \lim_{\epsilon \to 0} \left\{ \epsilon^{-(\mathfrak{q}+1)} \sum_{k=1}^{\mathfrak{q}+1} (-1)^{k+\mathfrak{q}+1} \binom{\mathfrak{q}+1}{k} [\Upsilon_0(\mathbf{x}+k\epsilon) - \Upsilon_0(\mathbf{x})] \right\}$$

motivates the estimator

$$\widetilde{\mathcal{D}}_{\mathfrak{q},n}^{\mathrm{FD}}(\mathbf{x}) = \epsilon_n^{-(\mathfrak{q}+1)} \sum_{k=1}^{\mathfrak{q}+1} (-1)^{k+\mathfrak{q}+1} \binom{\mathfrak{q}+1}{k} [\widehat{\Upsilon}_n(\mathbf{x}+k\epsilon_n) - \widehat{\Upsilon}_n(\mathbf{x})].$$

For $\eta > 0$, let

$$\begin{split} \widehat{G}_{\mathbf{x},n}(v;\eta) &= \sqrt{n\eta^{-1}} [\widehat{\Gamma}_n(\mathbf{x} + v\eta) - \widehat{\Gamma}_n(\mathbf{x}) - \Gamma_0(\mathbf{x} + v\eta) + \Gamma_0(\mathbf{x})] \\ &- \theta_0(\mathbf{x}) \sqrt{n\eta^{-1}} [\widehat{\Phi}_n(\mathbf{x} + v\eta) - \widehat{\Phi}_n(\mathbf{x}) - \Phi_0(\mathbf{x} + v\eta) + \Phi_0(\mathbf{x})], \end{split}$$

and

$$\widehat{R}_{\mathbf{x},n}(v;\eta) = \eta^{-1} [\widehat{\Phi}_n(\mathbf{x} + v\eta) - \widehat{\Phi}_n(\mathbf{x}) - \Phi_0(\mathbf{x} + v\eta) + \Phi_0(\mathbf{x})].$$

Using this notation, the first part of (B1) can be restated as

$$\widehat{G}_{\mathsf{x},n}(\cdot;a_n^{-1}) \leadsto \mathcal{G}_{\mathsf{x}}$$

and the second displayed condition of (B4) implies that

$$\widehat{R}_{\mathsf{x},n}(\cdot;a_n^{-1}) = o_{\mathbb{P}}(1).$$

In the preceding displays, one can typically replace $a_n^{-1}=n^{-1/(1+2\mathfrak{q})}$ by any $\eta_n>0$ with $\eta_n=o(1)$ and $a_n^{-1}\eta_n^{-1}=O(1)$. As a consequence, validity of the following assumption usually follows as a by-product of the arguments used to justify (B1) and (B4).

ASSUMPTION D. For every $\eta_n > 0$ with $\eta_n = o(1)$ and $a_n^{-1} \eta_n^{-1} = O(1)$,

$$\widehat{G}_{\mathsf{x},n}(1;\eta_n) = O_{\mathbb{P}}(1)$$
 and $\widehat{R}_{\mathsf{x},n}(1;\eta_n) = O_{\mathbb{P}}(1)$.

In turn, Assumption D is useful for the purposes of analyzing $\widetilde{\mathcal{D}}_{j,n}^{\mathtt{MA}}$, and $\widetilde{\mathcal{D}}_{j,n}^{\mathtt{FD}}$.

LEMMA 1. Suppose Assumptions A and D are satisfied, that $r_n(\widehat{\theta}_n(\mathsf{x}) - \theta_0(\mathsf{x})) = O_{\mathbb{P}}(1)$, and that $\epsilon_n \to 0$ and $n\epsilon_n^{1+2\mathfrak{q}} \to \infty$. If $\widetilde{\mathcal{D}}_{\mathfrak{q},n}^{\mathsf{MA}} \in \{\widetilde{\mathcal{D}}_{\mathfrak{q},n}^{\mathsf{MA}}, \widetilde{\mathcal{D}}_{\mathfrak{q},n}^{\mathsf{FD}}\}$, then (9) is satisfied and Assumption C holds with $\widetilde{M}_{\mathsf{x},n}$ in (10).

4.1.2. Mean Function Estimation: Adaptation to Unknown Degeneracy. Next, consider the somewhat more complicated case where \mathfrak{q} is unknown. In this case, we are able to obtain a positive result under the (additional) assumption that there are known integers $\bar{\mathfrak{q}}$ and $\underline{\mathfrak{s}}$ for which $\mathfrak{q} \leq \bar{\mathfrak{q}} \leq \underline{\mathfrak{s}} < \mathfrak{s}$. As the notation suggests, $\bar{\mathfrak{q}}$ and $\underline{\mathfrak{s}}$ are upper and lower bounds on \mathfrak{q} and \mathfrak{s} , respectively. The stated condition furthermore implies that $\mathfrak{s} > \mathfrak{q}$, a stronger smoothness condition than the condition $\mathfrak{s} \geq \mathfrak{q}$ of Assumption A.

For any $\bar{\mathfrak{q}} \geq \mathfrak{q}$, noting that $\mathcal{D}_j(\mathsf{x}) = 0$ for $j < \mathfrak{q}$ and that $\mathcal{D}_{\mathfrak{q}}(\mathsf{x}) > 0$ with \mathfrak{q} an odd integer, $\mathcal{M}_{\mathsf{x}}^{\mathfrak{q}}$ can be written as

$$\mathcal{M}_{\mathsf{x}}^{\mathfrak{q}}(v) = \sum_{\ell=1}^{\lfloor (\bar{\mathfrak{q}}+1)/2 \rfloor} \mathbb{1}(2\ell-1 \leq \mathfrak{q}) \mathcal{D}_{2\ell-1}(\mathsf{x})^+ v^{2\ell}.$$

Dropping the indicator function term from each summand, we obtain

$$\bar{\mathcal{M}}_{\mathsf{x}}(v) = \sum_{\ell=1}^{\lfloor (\bar{\mathfrak{q}}+1)/2\rfloor} \mathcal{D}_{2\ell-1}(\mathsf{x})^+ v^{2\ell}.$$

The majorant $\bar{\mathcal{M}}_x$ is an "adaptive" approximation to $\mathcal{M}_x^{\mathfrak{q}}$ in the sense that it does not depend on \mathfrak{q} , yet satisfies the local approximation property

$$\bar{\mathcal{M}}_{\mathsf{x}}^{\mathfrak{q}} \leadsto \mathcal{M}_{\mathsf{x}}^{\mathfrak{q}}, \qquad \bar{\mathcal{M}}_{\mathsf{x}}^{\mathfrak{q}}(v) = \sqrt{na_n} \bar{\mathcal{M}}_{\mathsf{x}}(va_n^{-1}).$$

Moreover, the following ("global") positivity property automatically holds: for some c > 0,

$$\inf_{|v|>K^{-1}} \bar{\mathcal{M}}_{\mathsf{x}}(v) > cK^{-(\mathfrak{q}+1)} \qquad \text{for every } K > 0.$$

As a consequence, it seems plausible that a "plug-in" estimator of $\bar{\mathcal{M}}_{x}$ would satisfy Assumption C under reasonable conditions. Indeed, if

(11)
$$a_n^{\mathfrak{q}-(2\ell-1)}(\widetilde{\mathcal{D}}_{2\ell-1,n}(\mathsf{x}) - \mathcal{D}_{2\ell-1}(\mathsf{x})) = o_{\mathbb{P}}(1), \qquad \ell = 1, \dots, |(\bar{\mathfrak{q}}+1)/2|,$$

then Assumption C is satisfied by

(12)
$$\widetilde{M}_{\mathbf{x},n}(v) = \sum_{\ell=1}^{\lfloor (\bar{\mathbf{q}}+1)/2 \rfloor} \widetilde{\mathcal{D}}_{2\ell-1,n}(\mathbf{x})^+ v^{2\ell}.$$

For "small" ℓ (namely, for $2\ell-1 < \mathfrak{q}$), the precision requirement (11) is stronger than consistency, but fortunately it turns out that the requirement can be met as long as \mathfrak{s} admits an integer $\underline{\mathfrak{s}}$ satisfying the condition $\overline{\mathfrak{q}} \leq \underline{\mathfrak{s}} < \mathfrak{s}$. To describe estimators satisfying (11) under this condition, let the constants $c_1, \ldots, c_{\underline{\mathfrak{s}}+1}$ be such that invertibility holds on the part of the $(\underline{\mathfrak{s}}+1)\times (\underline{\mathfrak{s}}+1)$ matrix with element (k,p) given by c_k^p and let the defining property of $\{\lambda_i^{\mathrm{BR}}(k): 1 \leq k \leq \underline{\mathfrak{s}}+1\}$ be

$$\sum_{k=1}^{\underline{\mathfrak{s}}+1} \lambda_j^{\mathrm{BR}}(k) c_k^p = \mathbb{1}(p=j+1), \qquad p=1,\ldots,\underline{\mathfrak{s}}+1.$$

Then, for any $j = 1, \dots, \underline{\mathfrak{s}}$, the "bias-reduced" estimator

$$\widetilde{\mathcal{D}}_{j,n}^{\mathrm{BR}}(\mathbf{x}) = \epsilon_n^{-(j+1)} \sum_{k=1}^{\underline{\mathfrak{s}}+1} \lambda_j^{\mathrm{BR}}(k) [\widehat{\Upsilon}_n(\mathbf{x} + c_k \epsilon_n) - \widehat{\Upsilon}_n(\mathbf{x})]$$

is motivated by the fact that as $\epsilon \to 0$,

$$\mathcal{D}_{j}(\mathbf{x}) = \epsilon^{-(j+1)} \sum_{k=1}^{\underline{\mathfrak{s}}+1} \lambda_{j}^{\mathtt{BR}}(k) [\Upsilon_{0}(\mathbf{x} + c_{k}\epsilon) - \Upsilon_{0}(\mathbf{x})] + O(\epsilon^{\min(\underline{\mathfrak{s}}+1,\mathfrak{s})-j})$$

when $\mathfrak{s}>j$. Relative to $\widetilde{\mathcal{D}}_{j,n}^{\mathrm{MA}}(\mathsf{x})$ and $\widetilde{\mathcal{D}}_{j,n}^{\mathrm{FD}}(\mathsf{x})$, this is a distinguishing feature of $\widetilde{\mathcal{D}}_{j,n}^{\mathrm{BR}}(\mathsf{x})$ and as it turns out this feature will enable us to formulate sufficient conditions for (11).

LEMMA 2. Suppose Assumptions A and D are satisfied, that $r_n(\widehat{\theta}_n(\mathsf{x}) - \theta_0(\mathsf{x})) = O_{\mathbb{P}}(1)$, and that $\epsilon_n \to 0$. If $\mathfrak{q} \leq \overline{\mathfrak{q}} \leq \underline{\mathfrak{s}} < \mathfrak{s}$, then

$$a_n^{\mathfrak{q}-j}(\widetilde{\mathcal{D}}_{j,n}^{\mathrm{BR}}(\mathbf{x})-\mathcal{D}_j(\mathbf{x}))=O(a_n^{\mathfrak{q}-j}\epsilon_n^{\min(\underline{\mathfrak{s}}+1,\mathfrak{s})-j})+O_{\mathbb{P}}[(a_n\epsilon_n)^{-(j+1/2)}+(a_n\epsilon_n)^{-j}]$$

for $j = 1, ..., \underline{\mathfrak{s}}$, implying in particular that if also

$$n\epsilon_n^{(1+2\bar{\mathfrak{q}})\min(\underline{\mathfrak{s}},\mathfrak{s}-1)/(\bar{\mathfrak{q}}-1)} \to 0 \quad and \quad n\epsilon_n^{1+2\bar{\mathfrak{q}}} \to \infty,$$

then (11) is satisfied and Assumption C holds with $\widetilde{M}_{x,n}$ in (12) if $\widetilde{\mathcal{D}}_{2\ell-1,n}=\widetilde{\mathcal{D}}^{BR}_{2\ell-1,n}$.

As alluded to previously, the ability of $\widetilde{\mathcal{D}}^{BR}$ to satisfy (11) is attributable in large part to its bias properties. In an attempt to highlight this, the first display of the lemma gives a stochastic expansion wherein the $O(a_n^{\mathfrak{q}-j}\epsilon_n^{\min(\underline{\mathfrak{s}}+1,\mathfrak{s})-j})$ term is a (possibly) negligible bias term. For $\widetilde{\mathcal{D}} \in \{\widetilde{\mathcal{D}}^{MA}, \widetilde{\mathcal{D}}^{FD}\}$, the analogous stochastic expansions are of the form

$$a_n^{\mathfrak{q}-j}(\widetilde{\mathcal{D}}_{j,n}(\mathsf{x}) - \mathcal{D}_j(\mathsf{x})) = O(a_n^{\mathfrak{q}-j}\epsilon_n^{\mathfrak{q}-j}) + O_{\mathbb{P}}[(a_n\epsilon_n)^{-(j+1/2)} + (a_n\epsilon_n)^{-j}],$$

the $O(a_n^{\mathfrak{q}-j}\epsilon_n^{\mathfrak{q}-j})$ term also being a bias term. When $a_n\epsilon_n\to\infty$ (as is required for the "noise" term in the stochastic expansion to be $o_{\mathbb{P}}(1)$), this bias term is non-negligible and the estimators $\widetilde{\mathcal{D}}^{\mathtt{MA}}$ and $\widetilde{\mathcal{D}}^{\mathtt{FD}}$ therefore do not satisfy (11).

Under additional assumptions (including $\mathfrak{s} \geq \underline{\mathfrak{s}} + 1$ and additional smoothness on Φ_0), $\widetilde{\mathcal{D}}^{BR}$ admits a Nagar-type mean squared error (MSE) expansion that can be used to select ϵ_n . The resulting approximate MSE formula is of the form

$$\epsilon_n^{2(\underline{\mathfrak{s}}+1-j)}\mathsf{B}_j^{\mathtt{BR}}(\mathsf{x})^2 + \frac{1}{n\epsilon_n^{1+2j}}\mathsf{V}_j^{\mathtt{BR}}(\mathsf{x}),$$

where the bias constant is

$$\mathsf{B}_{j}^{\mathrm{BR}}(\mathsf{x}) = \mathcal{D}_{\underline{\mathfrak{s}}+1}(\mathsf{x}) \sum_{k=1}^{\underline{\mathfrak{s}}+1} \lambda_{j}^{\mathrm{BR}}(k) c_{k}^{\underline{\mathfrak{s}}+2}$$

and the variance constant is

$$\mathsf{V}^{\mathtt{BR}}_{j}(\mathsf{x}) = \sum_{k=1}^{\underline{\mathfrak{s}}+1} \sum_{l=1}^{\underline{\mathfrak{s}}+1} \lambda^{\mathtt{BR}}_{j}(k) \lambda^{\mathtt{BR}}_{j}(l) \mathcal{C}_{\mathsf{x}}(c_{k}, c_{l});$$

for details, see Section SA.1.7 in the Supplemental Appendix (Cattaneo, Jansson and Nagasawa, 2024). Assuming $B_i^{BR}(x) \neq 0$, the approximate MSE is minimized by

$$\epsilon_{j,n}^{\mathrm{BR}}(\mathbf{x}) = \left(\frac{1+2j}{2(\underline{\mathfrak{s}}+1-j)}\frac{\mathsf{V}_{j}^{\mathrm{BR}}(\mathbf{x})}{\mathsf{B}_{j}^{\mathrm{BR}}(\mathbf{x})^{2}}\right)^{1/(3+2\underline{\mathfrak{s}})}n^{-1/(3+2\underline{\mathfrak{s}})},$$

a feasible version of which can be constructed by replacing $\mathcal{D}_{\underline{s}+1}(x)$ and \mathcal{C}_x with estimators in the expressions for $\mathsf{B}^{\mathtt{BR}}_j(x)$ and $\mathsf{V}^{\mathtt{BR}}_j(x)$, respectively.

4.2. Bootstrapping. Suppose inference is to be based on a random sample $\mathbf{Z}_1,\ldots,\mathbf{Z}_n$ from the distribution of some \mathbf{Z} . In all examples of which we are aware, the bootstrap parts of Assumption \mathbf{B} are satisfied when $(\widehat{\Gamma}_n^*,\widehat{\Phi}_n^*,\widehat{u}_n^*)$ is given by the nonparametric bootstrap analog of $(\widehat{\Gamma}_n,\widehat{\Phi}_n,\widehat{u}_n)$. Nevertheless, computationally simpler alternatives are often available and in what follows we will present one such alternative. To motivate our proposal, it is instructive to begin by revisiting Example 1.

EXAMPLE 1 (Monotone Density Estimation, continued). In this example, $\mathbf{Z} = X$. Moreover, defining $\gamma_0(x; \mathbf{z}) = \mathbb{1}(\mathbf{z} \le x)$ and $\phi_0(x; \mathbf{z}) = x$, we have the representations

$$\Gamma_0(x) = \mathbb{E}[\gamma_0(x; \mathbf{Z})]$$
 and $\Phi_0(x) = \mathbb{E}[\phi_0(x; \mathbf{Z})],$

and the estimators $\widehat{\Gamma}_n$ and $\widehat{\Phi}_n$ are linear in the sense that they are of the form

$$\widehat{\Gamma}_n(x) = \frac{1}{n} \sum_{i=1}^n \gamma_0(x; \mathbf{Z}_i) \qquad \text{and} \qquad \widehat{\Phi}_n(x) = \frac{1}{n} \sum_{i=1}^n \phi_0(x; \mathbf{Z}_i),$$

respectively. Finally, $\widehat{u}_n = \max_{1 \leq i \leq n} \mathbf{Z}_i \vee \mathsf{x}$.

Letting $\mathbf{Z}_{1,n}^*, \dots, \mathbf{Z}_{n,n}^*$ be a random sample from the empirical distribution of $\mathbf{Z}_1, \dots, \mathbf{Z}_n$, $\widehat{u}_n^* = \max_{1 \leq i \leq n} \mathbf{Z}_{i,n}^* \vee \mathbf{x}$ is the nonparametric bootstrap analog of \widehat{u}_n , while

$$\widehat{\Gamma}_{n}^{*}(x) = \frac{1}{n} \sum_{i=1}^{n} \gamma_{0}(x; \mathbf{Z}_{i,n}^{*})$$
 and $\widehat{\Phi}_{n}^{*}(x) = \frac{1}{n} \sum_{i=1}^{n} \phi_{0}(x; \mathbf{Z}_{i,n}^{*})$

are the nonparametric bootstrap analogs of $\widehat{\Gamma}_n$ and $\widehat{\Phi}_n$, respectively.

In the case of \widehat{u}_n , the alternative bootstrap analog $\widehat{u}_n^* = \widehat{u}_n$ is computationally trivial and automatically satisfies the bootstrap parts of (B5) and (B6). As for $\widehat{\Gamma}_n$ and $\widehat{\Phi}_n$, their nonparametric bootstrap analogs admit the weighted bootstrap representations

$$\widehat{\Gamma}_n^*(x) = \frac{1}{n} \sum_{i=1}^n W_{i,n} \gamma_0(x; \mathbf{Z}_i) \qquad \text{and} \qquad \widehat{\Phi}_n^*(x) = \frac{1}{n} \sum_{i=1}^n W_{i,n} \phi_0(x; \mathbf{Z}_i),$$

where, conditionally on $\mathbf{Z}_1, \dots, \mathbf{Z}_n$, $(W_{1,n}, \dots, W_{n,n})$ is multinomial distributed with probabilities (n^{-1}, \dots, n^{-1}) and number of trials n. (In turn, the weighted bootstrap interpretation

of the nonparametric bootstrap analog of $(\widehat{\Gamma}_n, \widehat{\Phi}_n)$ in this example is interesting partly because it can be used to embed the nonparametric bootstrap in a class of bootstraps also containing the Bayesian bootstrap and the wild bootstrap.)

Looking beyond Example 1, finding a computationally trivial \widehat{u}_n^* satisfying the bootstrap part of (B5) and (B6) is usually straightforward. On the other hand, although a weighted bootstrap interpretation of the nonparametric bootstrap version of the estimators $\widehat{\Gamma}_n$ and $\widehat{\Phi}_n$ is available whenever they are linear (e.g., in Example 2), there is no shortage of examples for which linearity does not hold. Nevertheless, the weighted bootstrap representation of the nonparametric bootstrap in Example 1 turns out be useful for our purposes, as it is suggestive of computationally attractive alternatives to the nonparametric bootstrap in more complicated examples.

When the non-bootstrap part of (B1) holds, the estimators $\widehat{\Gamma}_n$ and $\widehat{\Phi}_n$ are typically asymptotically linear in the sense that they admit (possibly) unknown functions γ_0 and ϕ_0 (satisfying $\Gamma_0(x) = \mathbb{E}[\gamma_0(x; \mathbf{Z})]$ and $\Phi_0(x) = \mathbb{E}[\phi_0(x; \mathbf{Z})]$, respectively) for which the approximations

$$\widehat{\Gamma}_n(x) \approx \overline{\Gamma}_n(x) = \frac{1}{n} \sum_{i=1}^n \gamma_0(x; \mathbf{Z}_i)$$
 and $\widehat{\Phi}_n(x) \approx \overline{\Phi}_n(x) = \frac{1}{n} \sum_{i=1}^n \phi_0(x; \mathbf{Z}_i)$

are suitably accurate. Assuming also that γ_0 and ϕ_0 admit sufficiently well behaved estimators $\widehat{\gamma}_n$ and $\widehat{\phi}_n$ (say), it then stands to reason that the salient properties of $\widehat{\Gamma}_n$ and $\widehat{\Phi}_n$ are well approximated by those of the easy-to-compute exchangeable bootstrap-type pair

$$\widehat{\Gamma}_n^*(x) = \frac{1}{n} \sum_{i=1}^n W_{i,n} \widehat{\gamma}_n(x; \mathbf{Z}_i) \quad \text{and} \quad \widehat{\Phi}_n^*(x) = \frac{1}{n} \sum_{i=1}^n W_{i,n} \widehat{\phi}_n(x; \mathbf{Z}_i),$$

where $W_{1,n}, \ldots, W_{n,n}$ denote exchangeable random variable (independent of $\mathbf{Z}_1, \ldots, \mathbf{Z}_n$). To give a precise statement, let

$$\psi_x(v; \mathbf{z}) = \gamma_0(x + v; \mathbf{z}) - \gamma_0(x; \mathbf{z}) - \theta_0(x) [\phi_0(x + v; \mathbf{z}) - \phi_0(x; \mathbf{z})]$$

and define

$$\check{\Gamma}_n(x) = \frac{1}{n} \sum_{i=1}^n \widehat{\gamma}_n(x; \mathbf{Z}_i)$$
 and $\check{\Phi}_n(x) = \frac{1}{n} \sum_{i=1}^n \widehat{\phi}_n(x; \mathbf{Z}_i)$.

In addition, for any function class \mathfrak{F} , let $N_U(\varepsilon,\mathfrak{F})$ denote the associated uniform covering numbers relative to L_2 ; that is, for any $\varepsilon > 0$, let

$$N_U(\varepsilon, \mathfrak{F}) = \sup_{Q} N(\varepsilon \|\bar{F}\|_{Q,2}, \mathfrak{F}, L_2(Q)),$$

where \bar{F} is the minimal envelope function of \mathfrak{F} , $\|\cdot\|_{Q,2}$ is the $L_2(Q)$ norm, $N(\cdot)$ is the covering number, and the supremum is over all discrete probability measure Q with $\|\bar{F}\|_{Q,2} > 0$.

ASSUMPTION E. For some $\delta > 0$, the following are satisfied:

- (E1) $\mathbf{Z}_1, \dots, \mathbf{Z}_n$, are independent and identically distributed.
- (E2) $W_{1,n}, \ldots, W_{n,n}$ are exchangeable random variables independent of $\mathbf{Z}_1, \ldots, \mathbf{Z}_n$, $\widehat{\gamma}_n$, and $\widehat{\phi}_n$. In addition, for some $\mathfrak{r} > (4\mathfrak{q} + 2)/(2\mathfrak{q} 1)$,

$$\frac{1}{n} \sum_{i=1}^{n} W_{i,n} = 1, \quad \frac{1}{n} \sum_{i=1}^{n} (W_{i,n} - 1)^2 \to_{\mathbb{P}} 1, \quad and \quad \mathbb{E}[|W_{1,n}|^{\mathfrak{r}}] = O(1).$$

(E3)
$$\sup_{x\in I} |\widehat{\Gamma}_n(x) - \bar{\Gamma}_n(x)| = o_{\mathbb{P}}(1)$$
 and

$$\frac{1}{n} \sum_{i=1}^{n} \sup_{x \in I} |\widehat{\gamma}_n(x; \mathbf{Z}_i) - \gamma_0(x; \mathbf{Z}_i)|^2 = o_{\mathbb{P}}(1).$$

Also,

$$\frac{a_n}{n} \sum_{i=1}^n \sup_{x \in I_x^i} |\widehat{\gamma}_n(x; \mathbf{Z}_i) - \gamma_0(x; \mathbf{Z}_i)|^2 = o_{\mathbb{P}}(1),$$

and there exist $\beta_{\gamma} \in [1/2, \mathfrak{q}+1)$ and random variables $A_{\gamma,n} = o_{\mathbb{P}}(1)$ and $B_{\gamma,n} = o_{\mathbb{P}}(a_n^{\beta_{\gamma}})$ such that, for $V \in (0, 2\delta]$,

$$\sqrt{na_n} \sup_{|v| < V} \left| \widehat{\Gamma}_n(\mathsf{x} + v) - \widehat{\Gamma}_n(\mathsf{x}) - \bar{\Gamma}_n(\mathsf{x} + v) + \bar{\Gamma}_n(\mathsf{x}) \right| \le A_{\gamma,n} + V^{\beta_\gamma} B_{\gamma,n},$$

$$\sqrt{na_n} \sup_{|v| < V} \left| \check{\Gamma}_n(\mathsf{x} + v) - \check{\Gamma}_n(\mathsf{x}) - \bar{\Gamma}_n(\mathsf{x} + v) + \bar{\Gamma}_n(\mathsf{x}) \right| \le A_{\gamma,n} + V^{\beta_{\gamma}} B_{\gamma,n}.$$

In addition, for some $\rho_{\gamma} \in (0,2)$,

$$\limsup_{\varepsilon \downarrow 0} \frac{\log N_U(\varepsilon, \mathfrak{F}_{\gamma})}{\varepsilon^{-\rho_{\gamma}}} < \infty, \quad \mathbb{E}[\bar{F}_{\gamma}(\mathbf{Z})^2] < \infty, \quad \limsup_{\varepsilon \downarrow 0} \frac{\log N_U(\varepsilon, \widehat{\mathfrak{F}}_{\gamma, n})}{\varepsilon^{-\rho_{\gamma}}} = O_{\mathbb{P}}(1),$$

where $\mathfrak{F}_{\gamma} = \{ \gamma_0(x; \cdot) : x \in I \}$, \bar{F}_{γ} is its minimal envelope, and $\widehat{\mathfrak{F}}_{\gamma,n} = \{ \widehat{\gamma}_n(x; \cdot) : x \in I \}$. Also,

$$\limsup_{\eta \downarrow 0} \frac{\mathbb{E}[\bar{D}_{\gamma}^{\eta}(\mathbf{Z})^{2} + \bar{D}_{\gamma}^{\eta}(\mathbf{Z})^{4}]}{\eta} < \infty,$$

where \bar{D}_{γ}^{η} is the minimal envelope of $\{\gamma_0(x;\cdot) - \gamma_0(x;\cdot) : x \in I_{\mathsf{x}}^{\eta}\}$.

(E4) $\sup_{x \in I} |\widehat{\Phi}_n(x) - \bar{\Phi}_n(x)| = o_{\mathbb{P}}(1)$ and

$$\frac{1}{n}\sum_{i=1}^{n}\sup_{x\in I}|\widehat{\phi}_n(x;\mathbf{Z}_i)-\phi_0(x;\mathbf{Z}_i)|^2=o_{\mathbb{P}}(1).$$

Also, $a_n|\widehat{\Phi}_n(\mathsf{x}) - \bar{\Phi}_n(\mathsf{x})| = o_{\mathbb{P}}(1), a_n|\widecheck{\Phi}_n(\mathsf{x}) - \bar{\Phi}_n(\mathsf{x})| = o_{\mathbb{P}}(1),$

$$\frac{a_n}{n} \sum_{i=1}^n \sup_{x \in I^{\delta}} \left| \widehat{\phi}_n(x; \mathbf{Z}_i) - \phi_0(x; \mathbf{Z}_i) \right|^2 = o_{\mathbb{P}}(1),$$

and there exist $\beta_{\phi} \in [1/2, \mathfrak{q}]$ and random variables $A_{\phi,n} = o_{\mathbb{P}}(1)$ and $B_{\phi,n} = o_{\mathbb{P}}(a_n^{\beta_{\phi}})$ such that, for $V \in (0, 2\delta]$,

$$\sqrt{na_n} \sup_{|v| \le V} \left| \widehat{\Phi}_n(\mathsf{x} + v) - \widehat{\Phi}_n(\mathsf{x}) - \bar{\Phi}_n(\mathsf{x} + v) + \bar{\Phi}_n(\mathsf{x}) \right| \le A_{\phi,n} + V^{\beta_\phi} B_{\phi,n},$$

$$\sqrt{na_n} \sup_{|v| < V} \left| \check{\Phi}_n(\mathsf{x} + v) - \check{\Phi}_n(\mathsf{x}) - \bar{\Phi}_n(\mathsf{x} + v) + \bar{\Phi}_n(\mathsf{x}) \right| \le A_{\phi,n} + V^{\beta_\phi} B_{\phi,n}.$$

In addition, for some $\rho_{\phi} \in (0,2)$,

$$\limsup_{\varepsilon \downarrow 0} \frac{\log N_U(\varepsilon, \mathfrak{F}_{\phi})}{\varepsilon^{-\rho_{\phi}}} < \infty, \quad \mathbb{E}[\bar{F}_{\phi}(\mathbf{Z})^2] < \infty, \quad \limsup_{\varepsilon \downarrow 0} \frac{\log N_U(\eta, \widehat{\mathfrak{F}}_{\phi,n})}{\varepsilon^{-\rho_{\phi}}} = O_{\mathbb{P}}(1),$$

where $\mathfrak{F}_{\phi} = \{\phi_0(x;\cdot) : x \in I\}$, \bar{F}_{ϕ} is its minimal envelope, and $\widehat{\mathfrak{F}}_{\phi,n} = \{\widehat{\phi}_n(x;\cdot) : x \in I\}$. Also,

$$\limsup_{\eta\downarrow 0} \frac{\mathbb{E}[\bar{D}_\phi^\eta(\mathbf{Z})^2 + \bar{D}_\phi^\eta(\mathbf{Z})^4]}{\eta} < \infty,$$

where \bar{D}^{η}_{ϕ} is the minimal envelope of $\{\phi_0(x;\cdot) - \phi_0(x;\cdot) : x \in I^{\eta}_{\mathsf{x}}\}$. (E5) For every $\eta_n > 0$ with $a_n \eta_n = O(1)$,

$$\sup_{v,v' \in [-\eta_n,\eta_n]} \frac{\mathbb{E}[|\psi_{\mathsf{x}}(v;\mathbf{Z}) - \psi_{\mathsf{x}}(v';\mathbf{Z})|]}{|v - v'|} = O(1)$$

and, for all $v, u \in \mathbb{R}$, and for some C_{\times} (as defined in (B1)),

$$\frac{\mathbb{E}[\psi_{\mathsf{x}}(\eta_n v; \mathbf{Z})\psi_{\mathsf{x}}(\eta_n u; \mathbf{Z})]}{\eta_n} \to \mathcal{C}_{\mathsf{x}}(v, u).$$

LEMMA 3. Suppose Assumptions A and E are satisfied. Then (B1)-(B4) are satisfied.

5. Examples. We apply our main results to two distinct sets of examples, both previously analyzed in Westling and Carone (2020) and Westling, Gilbert and Carone (2020), and references therein. In the supplemental appendix, we also consider two other set of examples, namely monotone hazard estimation (Huang and Wellner, 1995) and monotone distribution estimation (van der Vaart and van der Laan, 2006). All the proofs are found in the supplemental appendix.

For future reference, we collect the conditions on the bootstrap weights in an assumption.

ASSUMPTION BW. The bootstrap weights $W_{1,n}, \ldots, W_{n,n}$ are exchangeable, independent of the random sample $\mathbb{Z}_1, \ldots, \mathbb{Z}_n$, and satisfy, for some $\mathfrak{r} > 6$,

$$\frac{1}{n} \sum_{i=1}^{n} W_{i,n} = 1, \qquad \frac{1}{n} \sum_{i=1}^{n} (W_{i,n} - 1)^2 \to_{\mathbb{P}} 1, \qquad and \qquad \mathbb{E}[|W_{1,n}|^{\mathfrak{r}}] = O(1).$$

5.1. Monotone Density Estimation. Consider the problem of estimating the density of a non-negative, continuously distributed random variable, possibly with censoring. Let $\mathbf{Z}_1, \ldots, \mathbf{Z}_n$ be i.i.d. copies of $\mathbf{Z} = (\check{X}, \Delta)'$, with $\check{X} = X \wedge C$ and $\Delta = \mathbb{1}(X \leq C)$. Assuming that f_0 , the density of X, (exists and) is non-decreasing on $I = [0, u_0]$ where u_0 is some point in the support of X, the parameter of interest is $\theta_0(\mathsf{x})$ for some $\mathsf{x} \in (0, u_0)$, where $\theta_0 = f_0$. This setup generalizes Example 1. We take $\Phi_0(x) = x$ and therefore have $\Gamma_0 = F_0$, where F_0 is the cdf of X.

The canonical case of no censoring (i.e., $\mathbb{P}[C \geq X] = 1$) has been considered in Example 1. Assumptions under which this example is covered by our general theory is given in the following result.

COROLLARY 1. Let u_0 be the supremum of the support of X. Suppose \times is in the interior of I, θ_0 satisfies (A2), and Assumption BW holds. Then, Assumptions A, B, and D hold with

$$\widehat{\Gamma}_n(x) = \frac{1}{n} \sum_{i=1}^n \mathbb{1}(X_i \le x), \quad \widehat{\Gamma}_n^*(x) = \frac{1}{n} \sum_{i=1}^n W_{i,n} \mathbb{1}(X_i \le x),$$

$$\widehat{\Phi}_n(x) = \widehat{\Phi}_n^*(x) = x, \quad \widehat{u}_n = \widehat{u}_n^* = \max_{1 \le i \le n} X_i \lor \mathsf{x},$$

$$\mathcal{C}_\mathsf{x}(s,t) = f_0(\mathsf{x})(|s| \land |t|) \mathbb{1}(\mathrm{sign}(s) = \mathrm{sign}(t)), \quad \mathcal{D}_{\mathsf{q}}(\mathsf{x}) = \frac{\partial^{\mathsf{q}} f_0(\mathsf{x})}{(\mathsf{q}+1)!}.$$

Under the assumptions of Corollary 1, it follows from Theorem 1 that the bootstrap consistency result (8) holds for any $\widetilde{M}_{\mathbf{x},n}$ satisfying Assumption C. In particular, any pointwise consistent estimator of $\partial^{\mathbf{q}} f_0(\mathbf{x})$ could be used to estimate $\mathcal{D}_{\mathbf{q}}(\mathbf{x})$. Alternatively, since Assumption D also holds, $\widetilde{M}_{\mathbf{x},n}$ in (10) with $\widetilde{\mathcal{D}}_{\mathbf{q},n} \in \{\widetilde{\mathcal{D}}_{\mathbf{q},n}^{\mathrm{MA}}, \widetilde{\mathcal{D}}_{\mathbf{q},n}^{\mathrm{FD}}, \widetilde{\mathcal{D}}_{\mathbf{q},n}^{\mathrm{BR}}\}$ can also be used, provided that $\epsilon_n \to 0$ and $n\epsilon_n^{1+2\mathbf{q}} \to \infty$.

Suppose now that censoring occurs completely at random; that is, suppose $X \perp C$ (e.g., Huang and Wellner, 1995, and references therein). Let \widehat{S}_n and \widehat{G}_n be the Kaplan-Meier estimators of the survival functions $S_0(\cdot) = \mathbb{P}[X > \cdot]$ and $G_0(\cdot) = \mathbb{P}[C > \cdot]$, and define $\widehat{F}_n = 1 - \widehat{S}_n$ and $\widehat{\Lambda}_n(x) = \int_0^x \widehat{S}_n(u)^{-1} d\widehat{F}_n(u)$. In anticipation of the following result, we note that $\Gamma_0(x) = F_0(x) = \mathbb{E}[\gamma_0(x; \mathbf{Z})]$, where

$$\gamma_0(x; \mathbf{Z}) = F_0(x) + S_0(x) \left[\frac{\Delta \mathbb{1}(\check{X} \le x)}{S_0(\check{X}) G_0(\check{X})} - \int_0^{\check{X} \wedge x} \frac{\Lambda_0(du)}{S_0(u) G_0(u)} \right],$$

with $\Lambda_0(x) = \int_0^x S_0(u)^{-1} dF_0(u)$.

COROLLARY 2. Let u_0 be an interior point in the support of X. Suppose \times is in the interior of I, θ_0 satisfies (A2), $C \perp X$, C is absolutely continuous on I with bounded density, and $S_0(u_0)G_0(u_0) > 0$. Then, Assumptions A, B, and D hold with

$$\widehat{\Gamma}_n(x) = \widehat{F}_n(x), \quad \widehat{\Gamma}_n^*(x) = \frac{1}{n} \sum_{i=1}^n W_{i,n} \widehat{\gamma}_n(x; \mathbf{Z}_i)$$

$$\widehat{\gamma}_n(x; \mathbf{Z}) = \widehat{F}_n(x) + \widehat{S}_n(x) \left[\frac{\Delta \mathbb{1}(\check{X} \leq x)}{\widehat{S}_n(\check{X})\widehat{G}_n(\check{X})} - \int_0^{\check{X} \wedge x} \frac{d\widehat{\Lambda}_n(u)}{\widehat{S}_n(u)\widehat{G}_n(u)} \right],$$

$$\widehat{\Phi}_n(x) = \widehat{\Phi}_n^*(x) = x, \quad \widehat{u}_n = \widehat{u}_n^* = u_0,$$

$$C_{\mathsf{x}}(s,t) = \frac{f_0(\mathsf{x})}{G_0(\mathsf{x})} (|s| \wedge |t|) \mathbb{1}(\mathrm{sign}(s) = \mathrm{sign}(t)), \quad \mathcal{D}_{\mathsf{q}}(\mathsf{x}) = \frac{\partial^{\mathsf{q}} f_0(\mathsf{x})}{(\mathsf{q}+1)!}.$$

Theorem 1 implies that inference based on our proposed bootstrap-assisted method is asymptotically valid for this example as well. Additionally, Assumption D also holds under the above conditions, and we can use one of the numerical derivative estimators discussed in Section 4.1.

In the supplemental appendix, we also consider estimation of monotone density under conditionally independent right-censoring, where covariates are present. We do not discuss this example here because it involves additional notation and technicalities.

5.2. Monotone Regression Estimation. Consider now the problem of regression estimation, possibly with covariate adjustment. We assume that $\mathbf{Z}_1,\ldots,\mathbf{Z}_n$ are i.i.d. copies of $\mathbf{Z}=(Y,X,\mathbf{A}')'$, where X is continuous on its support I with density f_0 and $\mathrm{cdf}\, F_0$. Defining $\mu_0(x,\mathbf{a})=\mathbb{E}[Y|X=x,\mathbf{A}=\mathbf{a}]$, the parameter of interest is $\theta_0(x)$ for some x in the interior of I, where $\theta_0(\cdot)=\mathbb{E}[\mu_0(\cdot,\mathbf{A})]$ is assumed to be non-decreasing on I. (If there are no covariates \mathbf{A} , then $\theta_0(\cdot)=\mu_0(\cdot)=\mathbb{E}[Y|X=\cdot]$.) We take $\Phi_0=F_0$ and set $u_0=1$.

The classical monotone regression estimator has been considered in Example 2. Assumptions under which this example is covered by our general theory is given in the following result. To state it, let $\varepsilon = Y - \mathbb{E}[Y|X]$ and define $\sigma_0^2(x) = \mathbb{E}[\varepsilon^2|X = x]$.

COROLLARY 3. Suppose that x is in the interior of I, θ_0 satisfies (A2), Φ_0 satisfies (A3), and Assumption BW holds. If $\sigma_0^2(x)$ is continuous and positive at x, $\mathbb{E}[\varepsilon^2] < \infty$, and if $\sup_{x\in I^{\delta}} \mathbb{E}[\varepsilon^4|X=x] < \infty$ for some $\delta > 0$, then Assumptions A, B, and D hold with

$$\widehat{\Gamma}_n(x) = \frac{1}{n} \sum_{i=1}^n Y_i \mathbb{1}(X_i \le x), \quad \widehat{\Gamma}_n^*(x) = \frac{1}{n} \sum_{i=1}^n W_{i,n} Y_i \mathbb{1}(X_i \le x),$$

$$\widehat{\Phi}_n(x) = \frac{1}{n} \sum_{i=1}^n \mathbb{1}(X_i \le x), \quad \widehat{\Phi}_n^*(x) = \frac{1}{n} \sum_{i=1}^n W_{i,n} \mathbb{1}(X_i \le x), \quad \widehat{u}_n = \widehat{u}_n^* = 1,$$

$$\mathcal{C}_{\mathsf{x}}(s,t) = f_0(\mathsf{x})\sigma_0^2(\mathsf{x})(|s| \wedge |t|)\mathbb{1}(\mathrm{sign}(s) = \mathrm{sign}(t)), \qquad \mathcal{D}_{\mathfrak{q}}(\mathsf{x}) = \frac{f_0(\mathsf{x})\partial^{\mathfrak{q}}\mu_0(\mathsf{x})}{(\mathfrak{q}+1)!}.$$

Under the assumptions of Corollary 3, it follows from Theorem 1 that the bootstrap consistency result (8) holds for any $M_{x,n}$ satisfying Assumption C. In this example, any pointwise consistent estimators of $f_0(x)$ and $\partial^q \mu_0(x)$ could be used to estimate $\mathcal{D}_q(x)$. Alternatively, instead of using two distinct estimators, we can use the numerical derivative-type estimators since Assumption D also holds.

Next, consider the case of monotone regression estimation with covariate-adjustment. Introducing covariates A, let $f_{X|A}(\cdot|\mathbf{a})$ denote the conditional Lebesgue density of X given $\mathbf{A} = \mathbf{a}$ and define $g_0(x, \mathbf{a}) = f_{X|\mathbf{A}}(x|\mathbf{a})/f_0(x)$. Noting that $\Gamma_0(x) = \mathbb{E}[\gamma_0(x; \mathbf{Z})]$, where

$$\gamma_0(x; \mathbf{Z}) = \mathbb{1}(X \le x) \left[\frac{Y - \mu_0(X, \mathbf{A})}{g_0(X, \mathbf{A})} - \theta_0(X) \right],$$

we assume that $\widehat{\Gamma}_n$ is of the form $\widehat{\Gamma}_n(x) = n^{-1} \sum_{i=1}^n \widehat{\gamma}_n(x; \mathbf{Z}_i)$, where

$$\widehat{\gamma}_n(x; \mathbf{Z}) = \mathbb{1}(X \le x) \left[\frac{Y - \widehat{\mu}_n(X, \mathbf{A})}{\widehat{g}_n(X, \mathbf{A})} + \frac{1}{n} \sum_{j=1}^n \widehat{\mu}_n(X, \mathbf{A}_j) \right],$$

with $\widehat{\mu}_n$ and \widehat{g}_n being preliminary estimators of μ_0 and g_0 based on $\mathbf{Z}_1, \dots, \mathbf{Z}_n$. As discussed by Westling, Gilbert and Carone (2020), there are a number of candidates for $\widehat{\mu}_n$ and \widehat{g}_n . To accommodate a relatively wide class of estimators, the following assumption employs highlevel conditions on $\widehat{\mu}_n$ and \widehat{g}_n . To state the assumption, let $\varepsilon = Y - \mathbb{E}[Y|X, \mathbf{A}]$ and define $\sigma_0^2(x, \mathbf{a}) = \mathbb{E}[\varepsilon^2 | X = x, \mathbf{A} = \mathbf{a}].$

ASSUMPTION MRC. For some $\delta > 0$, the following are satisfied:

- (i) $\mathbb{E}[g_0(\mathbf{x},\mathbf{A})^{-1}\sigma_0^2(\mathbf{x},\mathbf{A})]>0$, $\mathbb{E}[\varepsilon^2]<\infty$, and $\sup_{x\in I_\mathbf{x}^\delta}\mathbb{E}[\varepsilon^4|X=x]<\infty$. (ii) $f_{X|\mathbf{A}}$ is bounded and g_0 is bounded away from zero.
- (iii) There exist random variables $A_n = o_{\mathbb{P}}(1)$ and $B_n = O_{\mathbb{P}}(a_n^{1/2})$ such that

$$\sqrt{na_n} \sup_{|v| \le V} |\widehat{\Gamma}_n(\mathbf{x} + v) - \widehat{\Gamma}_n(\mathbf{x}) - \bar{\Gamma}_n(\mathbf{x} + v) + \bar{\Gamma}_n(\mathbf{x})| \le A_n + VB_n, \qquad V \in (0, 2\delta]$$

In addition,

$$\frac{a_n}{n} \sum_{i=1}^n |\widehat{\mu}_n(X_i, \mathbf{A}_i) - \mu_0(X_i, \mathbf{A}_i)|^2 = o_{\mathbb{P}}(1),$$

$$\frac{a_n}{n^2} \sum_{i=1}^n \sum_{j=1}^n |\widehat{\mu}_n(X_i, \mathbf{A}_j) - \mu_0(X_i, \mathbf{A}_j)|^2 = o_{\mathbb{P}}(1),$$

and

$$\frac{a_n}{n} \sum_{i=1}^n \varepsilon_i^2 |\widehat{g}_n(X_i, \mathbf{A}_i) - g_0(X_i, \mathbf{A}_i)|^2 = o_{\mathbb{P}}(1).$$

(iv) $\mathbb{E}[\bar{\mu}(\mathbf{A})^2] < \infty$, where

$$\bar{\mu}(\mathbf{A}) = \sup_{|x-x'| \le \delta} \frac{|\mu_0(x, \mathbf{A}) - \mu_0(x', \mathbf{A})|}{|x - x'|}.$$

(v) $\mathbb{E}[\bar{\sigma}^2(\mathbf{A})] < \infty$, where, for some function ω with $\lim_{\eta \downarrow 0} \omega(\eta) = 0$,

$$\left| \frac{\sigma_0^2(x, \mathbf{A}) f_0(x)}{g_0(x, \mathbf{A})} - \frac{\sigma_0^2(\mathsf{x}, \mathbf{A}) f_0(\mathsf{x})}{g_0(\mathsf{x}, \mathbf{A})} \right| \le \omega(|x - \mathsf{x}|) \bar{\sigma}^2(\mathbf{A}), \qquad x \in I_\mathsf{x}^\delta.$$

Part (iii) of the assumption contains high-level conditions. In the supplemental appendix, we provide primitive sufficient conditions using nonparametric estimators based on sample splitting.

COROLLARY 4. Suppose that x is in the interior of I, θ_0 satisfies (A2), Φ_0 satisfies (A3), and Assumption BW holds. If Assumption MRC holds, then Assumptions A and B hold with

$$\widehat{\Gamma}_n^*(x) = \frac{1}{n} \sum_{i=1}^n W_{i,n} \widehat{\gamma}_n(x; \mathbf{Z}_i),$$

$$\widehat{\Phi}_n(x) = \frac{1}{n} \sum_{i=1}^n \mathbb{1}(X_i \le x), \quad \widehat{\Phi}_n^*(x) = \frac{1}{n} \sum_{i=1}^n W_{i,n} \mathbb{1}(X_i \le x), \quad \widehat{u}_n = \widehat{u}_n^* = 1,$$

$$\mathcal{C}_{\mathsf{x}}(s,t) = f_0(\mathsf{x}) \mathbb{E}\left[\frac{\sigma_0^2(\mathsf{x},\mathbf{A})}{g_0(\mathsf{x},\mathbf{A})}\right] (|s| \wedge |t|) \mathbb{1}(\mathrm{sign}(s) = \mathrm{sign}(t)), \quad \mathcal{D}_{\mathfrak{q}}(\mathsf{x}) = \frac{f_0(\mathsf{x}) \partial^{\mathfrak{q}} \theta_0(\mathsf{x})}{(\mathfrak{q}+1)!}.$$

As a consequence, if $\widetilde{M}_{\mathbf{x},n}$ satisfies Assumption C, then the "reshaped" bootstrap estimator $\widetilde{\theta}_n^*(\mathbf{x}) = \partial_- \mathsf{GCM}_{[0,1]}(\widetilde{\Gamma}_n^* \circ \widehat{\Phi}_n^{*-}) \circ \widehat{\Phi}_n^*(\mathbf{x})$ gives a bootstrap-assisted distributional approximation satisfying (8). In this example, the covariance kernel contains $\mathbb{E}[g_0(\mathbf{x}, \mathbf{A})^{-1}\sigma_0^2(\mathbf{x}, \mathbf{A})]$ and the plug-in procedure discussed in Section 3 would involve estimating this nuisance parameter, which in turn requires estimation of $\sigma_0^2(\mathbf{x},\cdot)$. When \mathbf{A} is of moderate dimension, this would introduce a challenging estimation problem on top of estimating μ_0 and g_0 . Our bootstrap-assisted procedure circumvents estimation of this additional nuisance parameter.

6. Simulations. We consider the canonical case of monotone regression (i.e., Example 2). We estimate the non-decreasing regression function $\theta_0(\cdot) = \mathbb{E}[Y|X=\cdot]$ at an interior point x using a random sample of observations. Three distinct data generating processes (DGPs) are considered. For each DGP, we let X be a uniform (0,1) random variable and we let $Y = \theta_0(X) + \sigma_0(X)\widetilde{\varepsilon}$, where $\widetilde{\varepsilon}$ is a standard normal random variable independent of X. DGP 1 sets $\theta_0(x) = 2\exp(x-0.5)$ and $\sigma_0 = 1$, DGP 2 sets $\theta_0(x) = 2\exp(x-0.5)$ and $\sigma_0 = \exp(x)$, and DGP 3 sets $\theta_0(x) = 24\exp(X-0.5) - 24(X-0.5) - 12(X-0.5)^2$ and $\sigma_0 = 0.1$. In all three cases, we set x = 0.5. The second DGP exhibits heteroskedastic regression errors, and the third DGP features a regression function whose first derivative equals zero at the evaluation point x; that is, DGP 3 exhibits a degree of degeneracy and if inference is conducted without knowledge of the fact that $\mathfrak{q}=3$, then this feature makes the inference problem more challenging.

The Monte Carlo experiment employs a sample size n = 1,000 with 2,000 bootstrap replications and 4,000 simulations, and compares four types of bootstrap-based inference procedures: the standard non-parametric bootstrap, m-out-of-n bootstrap, the "plug-in method" which simulates the limit law with estimated nuisance parameters, and our proposed bootstrap-assisted inference method implemented using the bias-reduced numerical derivative estimator discussed in Section 4.1. For the plug-in method, we estimate the covariance kernel and the nuisance parameter $\partial \Phi_0(x)$ using local linear kernel estimators, and we estimate the mean function by the same numerical derivative estimator as for our bootstrapassisted procedure. For m-out-of-n bootstrap and the plug-in method, researchers need to specify the characteristic exponent q, and we use the true (infeasible) value. For the numerical derivative estimator, we developed a rule of thumb for the step size ϵ_n to operationalize the procedure. For further details, see Section SA.6 of Cattaneo, Jansson and Nagasawa (2024). For our proposed method, we report results for three implementations: (i) (infeasible) procedure using the true value of $\mathcal{D}_0(x)$ (referred to as "oracle"), (ii) implementation using the numerical derivative estimator with a correct specification of q (referred to as "known q"), and (iii) "robust" implementation only assuming $q \in \{1, 3\}$ (referred to as "robust").

Table 1 presents the numerical results. We report empirical coverage for nominal 95% confidence intervals and their average interval length. For all of the DGPs considered, our proposed bootstrap-assisted inference method leads to confidence intervals with excellent empirical coverage and average interval length. The infeasible "oracle" procedure attains empirical coverage very close to the nominal 95%, which aligns with our theoretical results, and feasible procedures using the numerical derivative estimators perform almost identical to the infeasible oracle version.

In DGPs 1 and 2, our procedures outperform both the standard non-parametric bootstrap (which is inconsistent) and the m-out-of-n bootstrap (which is consistent) in empirical coverage and average length. In DGP 3, the m-out-of-n with the subsample size $m = \lceil n^{1/2} \rceil$ performs comparable to our procedure, but two caveats should be noted. First, the m-out-of-n bootstrap performance is sensitive to the choice of the subsample size. Therefore, to operationalize the m-out-of-n bootstrap procedure, one needs to develop a reliable procedure to choose the subsample size. Another caveat, arguably more important in this context, is that the m-out-of-n bootstrap procedure requires the knowledge of the convergence rate of the estimator. In our simulation, we use the true convergence rate for m-out-of-n bootstrap, but in practice, one needs to assume or estimate from data the convergence rate. Since the convergence rate and the limit distribution of the generalized Grenander-type estimator crucially hinge on unknown \mathfrak{q} , this feature of the m-out-of-n bootstrap may be unappealing. In contrast, our proposed bootstrap-based procedure only requires specifying an upper bound on \mathfrak{q} and it automatically adapts to the unknown convergence rate.

The plug-in method performs on par with our method, which is not unexpected because (i) the covariance kernel in this setting takes a relatively simple form and estimating it is not challenging with sample size n=1,000 and (ii) we use the true $\mathfrak q$ to simulate the limit law. In other examples where covariates are present, estimating covariance kernels typically involves nonparametric estimation of additional, possibly high-dimensional, nuisance functions. Our proposed method reduces the number of nuisance parameters that need to be estimated. Moreover, the good performance of the plug-in method for DGP 3 crucially depends on correctly specifying $\mathfrak q$, whereas our "robust" procedure only requires specifying an upper bound on $\mathfrak q$, and this is another desirable feature of our proposed method.

APPENDIX A: TECHNICAL RESULTS AND OMITTED DETAILS

This appendix collects several technical results and details omitted from the main paper to improve exposition. First, in Section A.1 we present a corrected version of the generalization

	DGP 1					DGP 2				DGP 3			
	$\tilde{\mathcal{D}}_{1,n}$	$\tilde{\mathcal{D}}_{3,n}$	Coverage	Length	$\tilde{\mathcal{D}}_{1,n}$	$\tilde{\mathcal{D}}_{3,n}$	Coverage	Length	$\tilde{\mathcal{D}}_{1,n}$	$\tilde{\mathcal{D}}_{3,n}$	Coverage	Length	
Standard													
			0.828	0.373			0.838	0.519			0.912	0.029	
m-out-of-n													
$m = \lceil n^{1/2} \rceil$			0.896	0.413			0.915	0.583			0.939	0.031	
$m = \lceil n^{2/3} \rceil$			0.868	0.400			0.884	0.557			0.928	0.029	
$m = \lceil n^{4/5} \rceil$			0.851	0.393			0.864	0.545			0.918	0.029	
Plug-in													
	1.053	0.000	0.950	0.399	1.052	0.000	0.946	0.553	0.000	1.003	0.936	0.028	
Reshaped													
Oracle	1.000	0.000	0.941	0.395	1.000	0.000	0.951	0.550	0.000	1.000	0.946	0.029	
ND known q	1.053	0.000	0.949	0.398	1.052	0.000	0.945	0.547	0.000	1.003	0.936	0.028	
ND robust	1.053	0.599	0.950	0.401	1.052	0.938	0.954	0.559	0.014	1.003	0.960	0.030	

Table 1
Simulations, Monotone Regression Estimator, 95% Confidence Intervals.

Notes: (i) Panel **Standard** refers to standard non-parametric bootstrap, Panel **m-out-of-n** refers to *m*-out-of-*n* non-parametric bootstrap with subsample *m*, Panel **Plug-in** refers to the plug-in method, Panel **Reshaped** refers to our proposed bootstrap-assisted procedure.

- (ii) Columns " $\tilde{\mathcal{D}}_{1,n}$ " and " $\tilde{\mathcal{D}}_{3,n}$ " report the averages of the estimated $\mathcal{D}_1,\mathcal{D}_3$ across simulations, and Columns "Coverage" and "Length" report empirical coverage and average length of bootstrap-based 95% percentile confidence intervals, respectively.
- (iii) "Oracle" corresponds to the infeasible version of our proposed procedure using the true value of $\mathcal{D}_{\mathfrak{q}}$, "ND known \mathfrak{q} " corresponds to our proposed procedure using the bias-reduced numerical derivative estimator with a correct specification of \mathfrak{q} , and "ND robust" corresponds to our proposed procedure only assuming $\mathfrak{q} \in \{1,3\}$. The step size choice for the numerical derivative estimator is described in the supplemental appendix.
- (iv) The sample size is 1,000, the number of bootstrap iterations is 2,000, and the number of Monte Carlo simulations is 4,000.

of the switch relation stated by Westling and Carone (2020, Supplement). Second, in Section A.2 we present a lemma that can be used to establish continuity of the cdf of the maximizer of a Gaussian process whose covariance kernel is that of two-sided Brownian motion. Both lemmas are used in the proof of Theorem 1 and may be of independent interest as well. Third, in Section A.3 we establish inconsistency of the plug-in nonparametric bootstrap for a large class of Generalized Grenander-type Estimators, following early work by Kosorok (2008) and Sen, Banerjee and Woodroofe (2010). Finally, in Section A.4 we report omitted formulas from Section 3.1.

A.1. Generalized Switch Relation. For a real-valued function f defined on a set $X \subset \mathbb{R}$, $\operatorname{argmax}_{x \in X} \{f(x)\}$ denotes the (possibly empty) set of maximizers of f over X. If $\operatorname{argmax}_{x \in X} \{f(x)\}$ is non-empty and contains a largest element, we denote that element by $\operatorname{max} \operatorname{argmax}_{x \in X} \{f(x)\}$.

In their analysis of generalized Grenander-type estimators, Westling and Carone (2020) relied on a generalization of the *switch relation* (Groeneboom, 1985). Their generalized switch relation is given in Lemma 1 of Westling and Carone (2020, Supplement). For the purposes of comparing it with Lemma A.1 below, it is convenient to restate that lemma as follows:

Statement GSR (Westling and Carone, 2020). Let Φ and Γ be real-valued functions defined on an interval $I \subseteq \mathbb{R}$ and suppose that Φ is non-decreasing and right-continuous. Fix l, u in $\Phi(I)$ with l < u and let $\theta = \partial_- \mathsf{GCM}_{[l,u]}(\Gamma \circ \Phi^-) \circ \Phi$. If I is closed, $\Phi(I) \subseteq [l,u]$, and if Γ and $\Gamma \circ \Phi^-$ are lower semi-continuous, then

(13)
$$\theta(\mathsf{x}) > t \quad \Longleftrightarrow \quad \sup_{x \in \Phi^{-}([l,u])} \{t\Phi(x) - \Gamma(x)\} < \Phi^{-}(\Phi(\mathsf{x}))$$

for any $t \in \mathbb{R}$ and any $x \in I$ with $\Phi(x) \in (l, u)$.

The main purpose of the following example is to show that without further restrictions, the argmax in (13) can be empty and the relation (13) can be violated (if we interpret $\sup \operatorname{argmax}_{x \in X} \{f(x)\}$ as $\inf X$ when $\operatorname{argmax}_{x \in X} \{f(x)\}$ is empty). In other words, Statement GSR can fail to hold.

EXAMPLE A.1. Let $I = [l, u] = [0, 1], \Gamma(x) = \gamma x$ (for some $\gamma \in \mathbb{R}$), and let

$$\Phi(x) = \begin{cases} x & \text{if } 0 \le x < 1/2\\ 1 & \text{if } 1/2 \le x \le 1. \end{cases}$$

Then

$$t\Phi(x) - \Gamma(x) = \begin{cases} (t - \gamma)x & \text{if } 0 \le x < 1/2 \\ t - \gamma/2 & \text{if } 1/2 \le x \le 1, \end{cases}$$

so $\operatorname{argmax}_{x \in \Phi^{-}([l,u])} \{t\Phi(x) - \Gamma(x)\}$ is empty when $0 > t > \gamma$. In particular, if $0 > t > \gamma$ and if $x \in (0,1/2)$, then

$$\sup_{x\in\Phi^-([l,u])} \{t\Phi(x) - \Gamma(x)\} = 0 < \mathsf{x} = \Phi^-(\Phi(\mathsf{x})),$$

whereas $\theta(\mathsf{x}) = \partial_-\mathsf{GCM}_{[l,u]}(\Gamma \circ \Phi^-) \circ \Phi(\mathsf{x}) = \gamma < t$, so (13) is violated. For future reference, we note that $\Phi(I) = \Phi(I) \cap [l,u] = [0,1/2) \cup \{1\}$ in this example.

The problems highlighted by the example are attributable to the fact that $\Phi(I) \cap [l,u]$ is not closed. Fortunately, it turns out that one can obtain a result in the spirit of (13) as long as $\Phi(I) \cap [l,u]$ is closed. The following lemma gives a precise statement.

LEMMA A.1 (Generalized Switch Relation). Let Φ and Γ be real-valued functions defined on an interval $I \subseteq \mathbb{R}$ and suppose that Φ is non-decreasing and right-continuous. Fix $l, u \in \Phi(I)$ with l < u and let $\theta = \partial_- \mathsf{GCM}_{[l,u]}(\Gamma \circ \Phi^-) \circ \Phi$. If $\Phi(I) \cap [l,u]$ is closed, then

(14)
$$\theta(\mathsf{x}) > t \iff \max_{x \in \Phi^{-}([l,u])} \{t\Phi(x) - \mathsf{LSC}_{\Phi}(\Gamma)(x)\} < \Phi^{-}(\Phi(\mathsf{x}))$$

for any $t \in \mathbb{R}$ and any $x \in I$ with $\Phi(x) \in (l, u)$, where

$$\mathsf{LSC}_{\Phi}(\Gamma)(x) = \mathsf{LSC}(\Gamma \circ \Phi^{-}) \circ \Phi(x),$$

and where LSC(\cdot) denotes the greatest lower semi-continuous minorant.

The assumptions of the lemma seem mild. In particular, the assumption that $\Phi(I) \cap [l,u]$ is closed is satisfied not only in the examples of Section 5, but in all examples of which we are aware.

The lemma does not assume lower semi-continuity of Γ or $\Gamma \circ \Phi^-$. Instead, the conclusion involves the function $\mathsf{LSC}_\Phi(\Gamma)$. As the notation suggests, $\mathsf{LSC}_\Phi(\cdot) = \mathsf{LSC}(\cdot)$ when Φ is the identity. More generally, $\mathsf{LSC}_\Phi(\cdot) = \mathsf{LSC}(\cdot)$ when Φ is continuous and strictly increasing and whether or not these properties hold it follows from Lemma 4.3 of van der Vaart and van der Laan (2006) that $\mathsf{LSC}_\Phi(\Gamma)$ admits the following characterization:

$$\mathsf{LSC}_{\Phi}(\Gamma)(x) = \liminf_{y \to \Phi(x)} (\Gamma \circ \Phi^{-})(y).$$

A.2. Continuity of argmax. Let $\{\mathbb{G}(v):v\in\mathbb{R}\}$ be a Gaussian process with $\mathbb{E}[\mathbb{G}(v)]=\mu(v)$ and $\mathbb{C}\mathrm{ov}[\mathbb{G}(v),\mathbb{G}(u)]=C(|v|\wedge|u|)\mathbb{1}(\mathrm{sign}(v)=\mathrm{sign}(u))$ for some C>0, i.e. \mathbb{G} is a scalar multiple of two-sided Brownian motion with mean shift. Under conditions on μ stated below, there exists a unique maximizer of $\mathbb{G}(v)$ over $v\in\mathbb{R}$ with probability one (Kim and Pollard, 1990). We complement this known fact with a result showing that the cdf of $\mathrm{argmax}_{v\in\mathbb{R}}\{\mathbb{G}(v)\}$ is continuous.

LEMMA A.2. Suppose that for each $v \in \mathbb{R}$, $\lim_{\delta \downarrow 0} [\mu(v+\delta) - \mu(v)]/\sqrt{\delta} = 0$ and that $\limsup_{|v| \to \infty} \mu(v)/|v|^c = -\infty$ for some c > 1/2. Then $x \mapsto \mathbb{P}[\operatorname{argmax}_{v \in \mathbb{R}} \{\mathbb{G}(v)\} \leq x]$ is continuous.

Under Assumptions A and B of the main text, $-\mathcal{M}_{\times}^{\mathfrak{q}} + t\mathcal{L}_{\times}$ satisfies the hypothesis of Lemma A.2 for any $t \in \mathbb{R}$. It therefore follows from the lemma that the function

$$x \mapsto \mathbb{P}\left[\underset{v \in \mathbb{R}}{\operatorname{argmin}} \{\mathcal{G}_{\mathsf{x}}(v) + \mathcal{M}_{\mathsf{x}}^{\mathsf{q}}(v) - t\mathcal{L}_{\mathsf{x}}(v)\} \ge x\right]$$

is continuous at x=0 for any $t \in \mathbb{R}$. We utilize that fact in our proof of (2) and note in passing that most of the existing literature on monotone function estimators seems to implicitly utilize a similar continuity result.

A.3. Bootstrap Inconsistency. Employing the exchangeable bootstrap setup introduced in Section 4.2, we show the inconsistency of the plug-in bootstrap-based distributional approximation. More precisely, we consider the "naïve" (plug-in) bootstrap estimator

$$\widehat{\theta}_n^*(\mathbf{x}) = \partial_-\mathsf{GCM}_{[0,\widehat{u}_n^*]}(\widehat{\Gamma}_n^* \circ \widehat{\Phi}_n^{*-}) \circ \widehat{\Phi}_n^*(\mathbf{x})$$

with

$$\widehat{\Gamma}_n^*(x) = \frac{1}{n} \sum_{i=1}^n W_{i,n} \widehat{\gamma}_n(x; \mathbf{Z}_i), \qquad \widehat{\Phi}_n^*(x) = \frac{1}{n} \sum_{i=1}^n W_{i,n} \widehat{\phi}_n(x; \mathbf{Z}_i),$$

and establish the following negative result.

THEOREM A.1. Suppose Assumptions A, (B5)-(B7), and E hold. Then

$$\sup_{t\in\mathbb{R}} \left| \mathbb{P}_n^* \left[\widehat{\theta}_n^*(\mathbf{x}) - \widehat{\theta}_n(\mathbf{x}) \leq t \right] - \mathbb{P}\left[\widehat{\theta}_n(\mathbf{x}) - \theta_0(\mathbf{x}) \leq t \right] \right| \neq o_{\mathbb{P}}(1).$$

Theorem A.1 accommodates exchangeable bootstrap schemes and a wide class of generalized Grenander-type estimators and contains as a special case the well-known fact that the nonparametric bootstrap approximation to the distribution of the classical Grenander estimator is inconsistent (e.g., Kosorok, 2008; Sen, Banerjee and Woodroofe, 2010).

A.4. Omitted Formulas. The objects $\widehat{V}_{x,n}^{\mathfrak{q}}, M_{x,n}^{\mathfrak{q}}, \widehat{L}_{x,n}^{\mathfrak{q}}$, and $\widehat{Z}_{x,n}^{\mathfrak{q}}$ appearing in Section 3.1 are defined as follows:

$$\widehat{V}_{\mathbf{x},n}^{\mathfrak{q}} = \{a_n(x - \mathbf{x}) : x \in \widehat{\Phi}_n^-([0, \widehat{u}_n^*])\},$$

$$M_{\mathbf{x},n}^{\mathfrak{q}}(v) = \sqrt{na_n}[\Gamma_0(\mathbf{x} + va_n^{-1}) - \Gamma_0(\mathbf{x})] - \theta_0(\mathbf{x})\sqrt{na_n}[\Phi_0(\mathbf{x} + va_n^{-1}) - \Phi_0(\mathbf{x})],$$

and

$$\widehat{L}_{\mathbf{x},n}^{\mathfrak{q}}(v) = a_n[\widehat{\Phi}_n(\mathbf{x} + va_n^{-1}) - \widehat{\Phi}_n(\mathbf{x})], \qquad \widehat{Z}_{\mathbf{x},n}^{\mathfrak{q}} = a_n[(\widehat{\Phi}_n^- \circ \widehat{\Phi}_n)(\mathbf{x}) - \mathbf{x}].$$

Similarly, the bootstrap analogs of $\widehat{V}_{x,n}^{\mathfrak{q}}, M_{x,n}^{\mathfrak{q}}, \widehat{L}_{x,n}^{\mathfrak{q}}$, and $\widehat{Z}_{x,n}^{\mathfrak{q}}$ are given by

$$\widehat{V}_{\mathbf{x},n}^{\mathfrak{q},*} = \{ a_n(x - \mathbf{x}) : x \in \widehat{\Phi}_n^{*-}([0, \widehat{u}_n^*]) \},$$

$$\widehat{M}_{\mathbf{x},n}^{\mathbf{q}}(v) = \sqrt{na_n}[\widehat{\Gamma}_n(\mathbf{x} + va_n^{-1}) - \widehat{\Gamma}_n(\mathbf{x})] - \widehat{\theta}_n(\mathbf{x})\sqrt{na_n}[\widehat{\Phi}_n(\mathbf{x} + va_n^{-1}) - \widehat{\Phi}_n(\mathbf{x})],$$

and

$$\widehat{L}_{\mathbf{x},n}^{\mathfrak{q},*}(v) = a_n[\widehat{\Phi}_n^*(\mathbf{x} + va_n^{-1}) - \widehat{\Phi}_n^*(\mathbf{x})], \qquad \widehat{Z}_{\mathbf{x},n}^{\mathfrak{q},*} = a_n[(\widehat{\Phi}_n^{*-} \circ \widehat{\Phi}_n^*)(\mathbf{x}) - \mathbf{x}],$$

respectively.

Acknowledgments. We are grateful to Boris Hanin, Jason Klusowski, Whitney Newey, and seminar participants at various institutions for their comments. We specially thank Marco Carone and Ted Westling for insightful discussions. The co-Editor, an Associate Editor, and two reviewers offered detailed suggestions that improved our paper.

Funding. Cattaneo gratefully acknowledges financial support from the National Science Foundation through grants SES-1947805 and DMS-2210561, and Jansson gratefully acknowledges financial support from the National Science Foundation through grant SES-1947662.

SUPPLEMENTARY MATERIAL

Supplement to "Bootstrap-Assisted Infernce for Generalized Grenander-Type Estimators."

The supplementary material contains proofs of Theorems 1 and A.1, Corollaries 1-4, and Lemmas 1-3 and A.1-A.2. It discusses additional examples that our main results cover.

REFERENCES

Brunk, H. D. (1958). On the Estimation of Parameters Restricted by Inequalities. *Annals of Mathematical Statistics* **29** 437–454.

CATTANEO, M. D., JANSSON, M. and NAGASAWA, K. (2020). Bootstrap-Based Inference for Cube Root Asymptotics. *Econometrica* 88 2203–2219.

CATTANEO, M. D., JANSSON, M. and NAGASAWA, K. (2024). Supplement to "Bootstrap-Assisted Inference for Generalized Grenander-Type Estimators".

CAVALIERE, G. and GEORGIEV, I. (2020). Inference under Random Limit Bootstrap Measures. *Econometrica* 88 2547–2574.

CHERNOFF, H. (1964). Estimation of the Mode. Annals of the Institute of Statistical Mathematics 16 31-41.

Cox, G. (2022). A Generalized Argmax Theorem with Applications. arXiv:2209.08793.

GRENANDER, U. (1956). On the Theory of Mortality Measurement: Part II. Scandinavian Actuarial Journal 39 125–153.

GROENEBOOM, P. (1985). Estimating a monotone density. In *Proceedings of the Berkeley Conference in Honor of Jerzy Neyman and Jack Kiefer, Vol. II* 539–555. Wadsworth Statist./Probab.

GROENEBOOM, P. and JONGBLOED, G. (2014). *Nonparametric Estimation under Shape Constraints*. Cambridge University Press.

GROENEBOOM, P. and JONGBLOED, G. (2018). Some Developments in the Theory of Shape Constrained Inference. *Statistical Science* **33** 473–492.

HAN, Q. and KATO, K. (2022). Berry-Esseen Bounds for Chernoff-type Non-standard Asymptotics in Isotonic Regression. Annals of Applied Probability 32 1459–1498.

HONG, H. and LI, J. (2020). The Numerical Bootstrap. Annals of Statistics 48 397-412.

HUANG, J. and WELLNER, J. A. (1995). Estimation of a Monotone Density or Monotone hazard under Random Censoring. *Scandinavian Journal of Statistics* **22** 3–33.

KIM, J. and POLLARD, D. (1990). Cube Root Asymptotics. Annals of Statistics 18 191-219.

KOSOROK, M. R. (2008). Bootstrapping the Grenander Estimator. In *Beyond Parametrics in Interdisciplinary Research: Festschrift in Honor of Professor Pranab K. Sen* 282–292. Institute of Mathematical Statistics.

- KUCHIBHOTLA, A. K., BALAKRISHNAN, S. and WASSERMAN, L. (2024). The HulC: Confidence Regions from Convex Hulls. *Journal of the Royal Statistical Society (Series B)*.
- LEE, S. M. S. and YANG, P. (2020). Bootstrap Confidence Regions Based on M-Estimators under Nonstandard Conditions. Annals of Statistics 48 274–299.
- MALLICK, S., SARKAR, S. and KUCHIBHOTLA, A. K. (2024). New Asymptotic Limit Theory and Inference for Monotone Regression. *arXiv:2310.20058*.
- POLITIS, D. N. and ROMANO, J. P. (1994). Large Sample Confidence Regions Based on Subsamples under Minimal Assumptions. *Annals of Statistics* **22** 2031–2050.
- POLITIS, D. N., ROMANO, J. P. and WOLF, M. (1999). Subsampling. Springer Science & Business Media.
- SEN, B., BANERJEE, M. and WOODROOFE, M. (2010). Inconsistency of Bootstrap: The Grenander Estimator. *Annals of Statistics* **38** 1953–1977.
- SEO, M. H. and OTSU, T. (2018). Local M-Estimation with Discontinuous Criterion for Dependent and Limited Observations. *Annals of Statistics* **46** 344–369.
- VAN DER LAAN, M. J. and ROBINS, J. M. (2003). *Unified Methods for Censored longitudinal Data and Causality*. Springer Science & Business Media.
- VAN DER VAART, A. W. (1998). Asymptotic Statistics. Cambridge University Press.
- VAN DER VAART, A. W. and VAN DER LAAN, M. J. (2006). Estimating a Survival Distribution with Current Status Data and High-Dimensional Covariates. *International Journal of Biostatistics* 2.
- VAN DER VAART, A. W. and WELLNER, J. A. (1996). Weak Convergence and Empirical Processes. Springer.
- WESTLING, T. and CARONE, M. (2020). A Unified Study of Nonparametric Inference for Monotone Functions. Annals of Statistics 48 1001–1024.
- WESTLING, T., GILBERT, P. and CARONE, M. (2020). Causal isotonic regression. *Journal of the Royal Statistical Society, Series B* 82 719–747.

SUPPLEMENT TO "BOOTSTRAP-ASSISTED INFERENCE FOR GENERALIZED GRENANDER-TYPE ESTIMATORS"

By Matias D. Cattaneo^{1,a}, Michael Jansson^{2,b} and Kenichi Nagasawa^{3,c}

 $^{1} Department\ of\ Operations\ Research\ and\ Financial\ Engineering,\ Princeton\ University,\ ^{a}cattaneo@princeton.edu$

Contents

SA.1 Pr	oofs	2
SA.1.1	Proof of Lemma A.1	2
SA.1.2	Proof of Lemma A.2	2
SA.1.3	Technical Lemmas	4
SA.1.4	Proof of Theorem 1	7
SA.1.5	Proof of Lemma 1	15
SA.1.6	Proof of Lemma 2	16
SA.1.7	Higher-order expansion of the bias-reduced estimator	16
SA.1.8	Proof of Lemma 3	17
SA.1.9	Proof of Theorem A.1	24
SA.1.10	Remarks on verifying conditions in applications	25
SA.1.11	Proof of Corollary 1	25
SA.1.12	Proof of Corollary 2	26
SA.1.13	Proof of Corollary 3	28
SA.1.14	Proof of Corollary 4	29
SA.2 Pr	imitive sufficient conditions for Assumption MRC (iii)	31
	Iditional example: monotone density function with conditionally inde-	2
	ndent right-censoring	34
SA.3.1	Proof of Corollary SA-1	35
SA.4 Ad	lditional example: monotone hazard function	36
SA.4.1	Proof of Corollary SA-2	37
~		
	Iditional example: distribution function estimation with current sta-	20
	s data	38
SA.5.1	Independent right-censoring	39
SA.5.2		39
SA.5.3	Proof of Corollaries SA-3 and SA-4	40
SA.6 Ru	ale-of-thumb step size selection	40
References		41

²Department of Economics, University of California at Berkeley, ^bmjansson@econ.berkeley.edu

³Department of Economics, University of Warwick, ^ckenichi.nagasawa@warwick.ac.uk

SA.1. Proofs.

SA.1.1. *Proof of Lemma A.1*. Lemma 4.2 of van der Vaart and van der Laan (2006) implies that, on (l, u),

$$\mathsf{GCM}_{[l,u]}(\Gamma \circ \Phi^-) = \mathsf{GCM}_{[l,u]}(\mathsf{LSC}(\Gamma \circ \Phi^-)).$$

Since $\Phi(x) \in (l, u)$, we therefore have

$$\theta(\mathsf{x}) = \partial_{-}\mathsf{GCM}_{[l,u]}(\Gamma \circ \Phi^{-}) \circ \Phi(\mathsf{x}) = \partial_{-}\mathsf{GCM}_{[l,u]}(\mathsf{LSC}(\Gamma \circ \Phi^{-})) \circ \Phi(\mathsf{x}),$$

so Lemma 4.1 of van der Vaart and van der Laan (2006) implies that

$$\theta(x) > t \iff y^* < \Phi(x),$$

where

$$y^{\star} = \max \underset{y \in [l,u]}{\operatorname{argmax}} \left\{ ty - \mathsf{LSC}(\Gamma \circ \Phi^{-})(y) \right\}.$$

Suppose $y^{\star} \in \Phi(I)$. Then $y^{\star} = \Phi(x^{\star})$, where

$$x^* = \Phi^-(y^*) \in \underset{x \in \Phi^-([l,u])}{\operatorname{argmax}} \{t\Phi(x) - \mathsf{LSC}_{\Phi}(\Gamma)(x)\}.$$

In particular,

$$\theta(\mathbf{x}) > t \iff y^* = \Phi(x^*) < \Phi(\mathbf{x}) \iff x^* < \Phi^-(\Phi(\mathbf{x})),$$

where, in fact,

$$x^* = \max_{x \in \Phi^-([l,u])} \left\{ t\Phi(x) - \mathsf{LSC}_{\Phi}(\Gamma)(x) \right\},$$

because if

$$x^* < x' \in \underset{x \in \Phi^-([l,u])}{\operatorname{argmax}} \{ t\Phi(x) - \mathsf{LSC}_{\Phi}(\Gamma)(x) \},$$

then, contradicting the definition of y^* , we have

$$y' = \Phi(x') \in \operatorname*{argmax}_{y \in [l,u]} \left\{ ty - \mathsf{LSC}(\Gamma \circ \Phi^-)(y) \right\}$$
 and $y' > y^\star$.

The proof can therefore be completed by showing that $y^* \in \Phi(I)$.

If $\Phi(I) \supseteq [l,u]$, then there is nothing to show, so suppose $\Phi(I) \not\supseteq [l,u]$. If $y \in \Phi(I)^c \cap [l,u]$, then, since $\Phi(I) \cap [l,u]$ is closed and $l,u \in \Phi(I)$, we have $[y-\eta,y+\eta] \cap \Phi(I) = \emptyset$ for some $\eta > 0$ with $[y-\eta,y+\eta] \subset [l,u]$. Therefore, the function $\mathsf{LSC}(\Gamma \circ \Phi^-)$ is constant on the interval $[y-\eta/2,y+\eta/2]$, implying in particular that

$$y^{\star} = \max \mathop{\mathrm{argmax}}_{y' \in [l,u]} \left\{ ty' - \mathsf{LSC}(\Gamma \circ \Phi^{-})(y') \right\} \neq y. \quad \Box$$

SA.1.2. Proof of Lemma A.2. $\mathbb{G}(v)$ is a Gaussian process with $\mathbb{E}[\mathbb{G}(v)] = \mu(v)$ for all $v \in \mathbb{R}$ and $\mathbb{C}\text{ov}[\mathbb{G}(v), \mathbb{G}(v')] = C\min\{|v|, |v'|\}\mathbb{1}(\text{sign}(v) = \text{sign}(v')) =: \mathcal{K}(v, v')$ for some positive constant C and for all $v, v' \in \mathbb{R}$. The form of the covariance kernel \mathcal{K} implies that for every $\tau > 0$ and every $v, v' \in \mathbb{R}$,

$$\mathcal{K}(v\tau, v'\tau) = \mathcal{K}(v, v')\tau$$

and

$$\mathcal{K}(v + v', v + v') - 2\mathcal{K}(v + v', v') + \mathcal{K}(v', v') = \mathcal{K}(v, v).$$

In addition, $\mathcal{K}(1,1) > 0$ and $\lim_{v \downarrow 0} \mathcal{K}(1,v) / \sqrt{v} = 0.$

We begin by adapting the arguments of Kim and Pollard (1990, pp. 196-198) to show that a maximizer of $\mathbb{G}(v)$ over $v \in \mathbb{R}$ exists and is unique with probability one. Let $\mathbb{G}^{\mu}(v) = \mathbb{G}(v) - \mu(v)$ be the centered process and suppose that, for the same c > 1/2 as in the hypothesis,

(SA.1)
$$\mathbb{P}\left[\limsup_{|v|\to\infty}\frac{\mathbb{G}^{\mu}(v)}{|v|^c}>\eta\right]=0 \quad \text{for any } \eta>0.$$

Then, with probability one, $\mathbb{G}(v) \to -\infty$ as $|v| \to \infty$, implying in turn that a maximizer of $\mathbb{G}(v)$ exists (because sample paths are continuous). Also, since

$$\mathbb{V}[\mathbb{G}(v) - \mathbb{G}(v')] = \mathcal{K}(v,v) + \mathcal{K}(v',v') - 2\mathcal{K}(v,v-v') = \mathcal{K}(v-v',v-v') > 0$$

for $v \neq v'$, Lemma 2.6 of Kim and Pollard (1990) implies that this maximizer is unique with probability one. In turn, (SA.1) follows from the Borel-Cantelli lemma because

$$\begin{split} \sum_{k=2}^{\infty} \mathbb{P} \left[\sup_{k-1 \le |v| \le k} \frac{\mathbb{G}^{\mu}(v)}{|v|^{c}} > \eta \right] &\leq \sum_{k=2}^{\infty} \mathbb{P} \left[\sup_{|v| \le k} \mathbb{G}^{\mu}(v) > (k/2)^{c} \eta \right] \\ &= \sum_{k=2}^{\infty} \mathbb{P} \left[\sup_{|v| \le 1} \mathbb{G}^{\mu}(v) > k^{c-1/2} 2^{-c} \eta \right] \\ &\leq \frac{2^{4c/(2c-1)} \mathbb{E} \left[\sup_{|v| \le 1} |\mathbb{G}^{\mu}(v)|^{4c/(2c-1)} \right]}{\eta^{4c/(2c-1)}} \sum_{k=2}^{\infty} k^{-2} < \infty, \end{split}$$

where the equality uses the rescaling property $\mathcal{K}(v\tau, v'\tau) = \tau \mathcal{K}(v, v')$ and where the last inequality uses Jain and Marcus (1978, Corollary 4.7).

To show continuity of the function $x \mapsto \mathbb{P}[\operatorname{argmax}_{v \in \mathbb{R}} \{\mathbb{G}(v)\} \leq x]$, it suffices to show that $\mathbb{P}[\operatorname{argmax}_{v \in \mathbb{R}} \{\mathbb{G}(v)\} = x] = 0$ for every $x \in \mathbb{R}$. Fix $x \in \mathbb{R}$ and define $\widetilde{\mathbb{G}}_x(0) = 0$ and

$$\widetilde{\mathbb{G}}_x(v) = \frac{\mathbb{G}(x+v) - \mathbb{G}(x)}{\sqrt{\mathcal{K}(v,v)}}, \quad v \neq 0.$$

Then $\max_{v \in \mathbb{R}} \widetilde{\mathbb{G}}_x(v) \geq 0$ and, for any set $\mathcal{V} \subset \mathbb{R}$,

$$(\mathrm{SA.2}) \qquad \mathbb{P}\left[\underset{v \in \mathbb{R}}{\operatorname{argmax}}\{\mathbb{G}(v)\} = x\right] = \mathbb{P}\left[\underset{v \in \mathbb{R}}{\max}\,\widetilde{\mathbb{G}}_x(v) \leq 0\right] \leq \mathbb{P}\left[\underset{v \in \mathcal{V}}{\max}\,\widetilde{\mathbb{G}}_x(v) \leq 0\right].$$

In the sequel, we show that the majorant in (SA.2) can be made arbitrarily small by choice of \mathcal{V} . In particular, for $\varepsilon \in (0,1)$, we construct $v_{\varepsilon} > 0$ such that

(SA.3)
$$\mathbb{E}\left[\widetilde{\mathbb{G}}_x(v_{\varepsilon}^i)\right] \geq -\varepsilon \quad \text{for every } i \geq 1,$$

and

(SA.4)
$$\left| \mathbb{C}\text{ov}\left(\widetilde{\mathbb{G}}_x(v_{\varepsilon}^i), \widetilde{\mathbb{G}}_x(v_{\varepsilon}^j) \right) \right| \leq \varepsilon \quad \text{ for every } j > i \geq 1.$$

For any $N\in\mathbb{N}$, defining $\mathcal{V}_{\varepsilon,N}=\{v_{\varepsilon},\dots,v_{\varepsilon}^{N}\}$, we therefore have

$$\mathbb{P}\left[\max_{v\in\mathbb{R}}\widetilde{\mathbb{G}}_x(v)\leq 0\right]\leq \liminf_{\varepsilon\downarrow 0}\mathbb{P}\left[\max_{v\in\mathcal{V}_{\varepsilon,N}}\widetilde{\mathbb{G}}_x(v)\leq 0\right]\leq \left|\int_{-\infty}^0\frac{\exp(-x^2/2)}{\sqrt{2\pi}}dx\right|^N=2^{-N},$$

¹As a side note, if the covariance kernel satisfies the two displays, then the Gaussian process is two-sided Brownian motion, up to mean shift and scaling by a positive constant.

where the second inequality uses the fact that convergence of means and covariances of normal random vectors implies convergence in distribution. Since N is arbitrary, the left-hand side in the preceding display is zero. Letting $\varepsilon \in (0,1)$ be given, the proof can therefore be completed by exhibiting v_{ε} satisfying (SA.3) and (SA.4).

Because $\mathcal{K}(\tau,\tau) = \tau \mathcal{K}(1,1)$ and $\lim_{\tau \downarrow 0} [\mu(x+\tau) - \mu(x)]/\sqrt{\tau} = 0$, there exists $\bar{\tau}_{\varepsilon}' \in (0,1)$ such that

$$\mathbb{E}\left[\widetilde{\mathbb{G}}_x(\tau)\right] = \frac{\mu(x+\tau) - \mu(x)}{\sqrt{\mathcal{K}(\tau,\tau)}} = \frac{\left[\mu(x+\tau) - \mu(x)\right]/\sqrt{\tau}}{\sqrt{\mathcal{K}(1,1)}} \ge -\varepsilon$$

for every $\tau \in (0, \bar{\tau}_{\varepsilon}']$. Also, because $\mathcal{K}(\tau_i, \tau_j) = \tau_i \mathcal{K}(1, \tau_j/\tau_i)$ and $\lim_{\tau \downarrow 0} \mathcal{K}(1, \tau)/\sqrt{\tau} = 0$, there exists $\bar{\tau}_{\varepsilon}'' \in (0, 1)$ such that

$$\mathbb{C}\mathrm{ov}\left(\widetilde{\mathbb{G}}_{x}(\tau_{i}),\widetilde{\mathbb{G}}_{x}(\tau_{j})\right) = \frac{\mathcal{K}(\tau_{i},\tau_{j})}{\sqrt{\mathcal{K}(\tau_{i},\tau_{i})\mathcal{K}(\tau_{j},\tau_{j})}} = \frac{\mathcal{K}(1,\tau_{j}/\tau_{i})/\sqrt{\tau_{j}/\tau_{i}}}{\mathcal{K}(1,1)} \in [-\varepsilon,\varepsilon]$$

for all $\tau_i, \tau_j > 0$ with $\tau_j / \tau_i \leq \bar{\tau}_{\varepsilon}^{''}$. If $v_{\varepsilon} \in (0, \bar{\tau}_{\varepsilon}^{'} \wedge \bar{\tau}_{\varepsilon}^{''})$, then (SA.3) and (SA.4) are satisfied. \square

SA.1.3. *Technical Lemmas*. In preparation for the proof of Theorem 1, this section presents six technical lemmas. The first lemma is a switching lemma, which will be used when characterizing the limiting distributions obtained in Theorem 1.

LEMMA SA-1. Let $\Gamma : \mathbb{R} \to \mathbb{R}$ be a lower semi-continuous function that is bounded from below and satisfies $\lim_{|v| \to \infty} \Gamma(v)/|v| = \infty$. Then, for any $x, t \in \mathbb{R}$,

$$\partial_{-}\mathsf{GCM}_{\mathbb{R}}(\Gamma)(\mathsf{x}) > t \qquad \Longleftrightarrow \qquad \max \operatorname*{argmax}_{v \in \mathbb{R}} \left\{ vt - \Gamma(v) \right\} < \mathsf{x}.$$

PROOF. Because $\lim_{|v|\to\infty} \Gamma(v)/|v| = \infty$, there exists a $K > |\mathbf{x}|$ such that if $|v| \ge K$, then $vt - \Gamma(v) < -\Gamma(0)$, implying in particular that

$$\operatorname*{argmax}_{v \in \mathbb{R}} \left\{ vt - \Gamma(v) \right\} = \operatorname*{argmax}_{v \in [-c,c]} \left\{ vt - \Gamma(v) \right\}$$

for every $c \geq K$. Also, by Lemma A.1. of Sen, Banerjee and Woodroofe (2010) there exists a c > K such that $\mathsf{GCM}_{\mathbb{R}}(\Gamma) = \mathsf{GCM}_{[-c,c]}(\Gamma)$ on [-K,K], implying in particular that

$$\partial_{-}\mathsf{GCM}_{\mathbb{R}}(\Gamma)(\mathsf{x}) = \partial_{-}\mathsf{GCM}_{[-c,c]}(\Gamma)(\mathsf{x}).$$

For any such c, the conclusion of the lemma is equivalent to the statement

$$\partial_{-}\mathsf{GCM}_{[-c,c]}(\Gamma)(\mathsf{x}) > t \qquad \iff \qquad \max \underset{v \in [-c,c]}{\operatorname{argmax}} \left\{ vt - \Gamma(v) \right\} < \mathsf{x},$$

whose validity follows from Lemma 4.1 of van der Vaart and van der Laan (2006).

The proof of Theorem 1 furthermore employs various approximations to functionals of the form $LSC_{\Phi}(f)$. The approximations in question are obtained using Lemmas SA-2, SA-3, SA-5, and SA-6. In all cases, the approximations are based on the representation

(SA.5)
$$\mathsf{LSC}_{\Phi}(f)(x) = \liminf_{\epsilon \downarrow 0} \inf_{x' \in \mathcal{X}_{\Phi}^{\epsilon}(x)} f(x'),$$

where
$$\mathcal{X}^{\epsilon}_{\Phi}(x) = (\Phi^{-}(\Phi(x) - \epsilon), \Phi^{-}(\Phi(x))] \cup (\Phi^{-}(\Phi(x) + \epsilon), \Phi^{-}(\Phi(x) + \epsilon)).$$

The following lemma uses (SA.5) and the special structure of Γ_0 to obtain a simple "global" bound on the error of the approximation LSC $_{\Phi}(\Gamma_0) \approx \Gamma_0$.

LEMMA SA-2. Suppose Assumption A holds and suppose Φ is non-decreasing and right-continuous on I. Then, for every $x \in I$,

$$|\mathsf{LSC}_{\Phi}(\Gamma_0)(x) - \Gamma_0(x)| \le 2 \left(\sup_{x' \in I} |\theta_0(x')| \right) \sup_{x' \in I} |\Phi(x') - \Phi_0(x')|.$$

PROOF. By (SA.5) and continuity of Γ_0 ,

$$\mathsf{LSC}_{\Phi}(\Gamma_0)(x) = \min \left[\Gamma_0(\Phi^-(\Phi(x))), \Gamma_0(\Phi^-(\Phi(x)+)) \right],$$

while, by Assumption A,

$$|\Gamma_0(x') - \Gamma_0(x)| \le \left(\sup_{x'' \in I} |\theta_0(x'')|\right) |\Phi_0(x') - \Phi_0(x)|.$$

Now, using $\Phi \circ \Phi^- \circ \Phi = \Phi$,

$$|\Phi_0(\Phi^-(\Phi(x))) - \Phi_0(x)| = |\Phi_0(\Phi^-(\Phi(x))) - \Phi(\Phi^-(\Phi(x))) + \Phi(x) - \Phi_0(x)|$$

$$\leq 2 \sup_{x' \in I} |\Phi(x') - \Phi_0(x')|.$$

Also,

$$|\Phi_0(\Phi^-(\Phi(x)+)) - \Phi_0(x)| \le 2 \sup_{x' \in I} |\Phi(x') - \Phi_0(x')|$$

because, for every $\eta > 0$,

$$\begin{split} 0 & \leq \Phi_0(\Phi^-(\Phi(x)+)) - \Phi_0(x) \leq \Phi_0(\Phi^-(\Phi(x)+\eta)) - \Phi_0(x) \\ & \leq \Phi_0\left(\Phi^-\left(\Phi_0(x) + \sup_{x' \in I} |\Phi(x') - \Phi_0(x')| + \eta\right)\right) - \Phi_0(x) \\ & \leq \Phi_0\left(\Phi_0^-\left(\Phi_0(x) + 2\sup_{x' \in I} |\Phi(x') - \Phi_0(x')| + \eta\right)\right) - \Phi_0(x) \\ & \leq 2\sup_{x' \in I} |\Phi(x') - \Phi_0(x')| + \eta, \end{split}$$

where the last inequality uses continuity of Φ_0 .

A simple "global" bound on the error of the approximation $LSC_{\Phi}(f) \approx f$ is available also in the important special case where f is proportional to Φ .

LEMMA SA-3. Suppose Assumption A holds and suppose Φ is non-decreasing and right-continuous on I. Then, for every $x \in I$ and every $\theta \in \mathbb{R}$,

$$|\mathsf{LSC}_{\Phi}(\theta\Phi)(x) - \theta\Phi(x)| \le |\theta| \sup_{x' \in I} |\Phi(x') - \Phi(x'-)|.$$

PROOF. First, if $\theta > 0$, then the result follows from the fact that, by (SA.5),

$$\mathsf{LSC}_{\Phi}(\theta\Phi)(x) = \theta\Phi(\Phi^-(\Phi(x)) -) \leq \theta\Phi(\Phi^-(\Phi(x))) = \theta\Phi(x),$$

where

$$\Phi(x) = \Phi(\Phi^{-}(\Phi(x))) \le \Phi(\Phi^{-}(\Phi(x)) - 1) + \sup_{x' \in I} |\Phi(x') - \Phi(x')|.$$

Next, if $\theta < 0$, then the result follows from the fact that, by (SA.5),

$$\mathsf{LSC}_{\Phi}(\theta\Phi)(x) = \theta\Phi(\Phi^{-}(\Phi(x)+)) < \theta\Phi(\Phi^{-}(\Phi(x))) = \theta\Phi(x),$$

where

$$\begin{split} \Phi(\Phi^{-}(\Phi(x)+)) &\leq \liminf_{\eta \downarrow 0} \Phi(\Phi^{-}(\Phi(x)+\eta)-) + \sup_{x' \in I} |\Phi(x') - \Phi(x'-)| \\ &\leq \Phi(x) + \sup_{x' \in I} |\Phi(x') - \Phi(x'-)|. \end{split}$$

Next, we give a "local" approximation to $\Phi^- \circ \Phi$. That approximation will later be used in combination with (SA.5) to obtain "local" approximations to LSC $_{\Phi}(f)$, but the approximation is also useful in its own right and we therefore state it as a separate lemma.

Lemma SA-4. Suppose Assumption A holds and suppose Φ is non-decreasing and right-continuous on I. Also, suppose

$$\frac{2\sup_{x'\in I_x^{\delta+\epsilon}} |\Phi(x') - \Phi_0(x')|}{\inf_{x'\in I_x^{\delta+\epsilon}} \partial \Phi_0(x')} < \epsilon$$

for some $\delta, \epsilon > 0$ with $I_{\mathsf{x}}^{\delta+\epsilon} \subseteq I$. Then, for every $x \in I_{\mathsf{x}}^{\delta}$,

$$|\Phi^{-}(\Phi(x)) - x| < \epsilon$$
 and $|\Phi^{-}(\Phi(x)+) - x| < \epsilon$.

PROOF. First, suppose $|\Phi^-(\Phi(x)) - x| \ge \epsilon$. Then $\Phi^-(\Phi(x)) \le x - \epsilon$, implying in particular that $\Phi(x - \epsilon) = \Phi(x)$ and therefore also

$$[\Phi(x - \epsilon) - \Phi_0(x - \epsilon)] - [\Phi(x) - \Phi_0(x)] = \Phi_0(x) - \Phi_0(x - \epsilon).$$

Now, if $x \in I_{\mathsf{x}}^{\delta}$, then $[x - \epsilon, x] \subseteq I_{\mathsf{x}}^{\delta + \epsilon}$, so

$$\Phi_0(x) - \Phi_0(x - \epsilon) \ge \epsilon \inf_{x' \in I_x^{\delta + \epsilon}} \partial \Phi_0(x') > 2 \sup_{x' \in I_x^{\delta + \epsilon}} |\Phi(x') - \Phi_0(x')|,$$

whereas

$$|[\Phi(x-\epsilon) - \Phi_0(x-\epsilon)] - [\Phi(x) - \Phi_0(x)]| \le 2 \sup_{x' \in L^{\delta+\epsilon}} |\Phi(x') - \Phi_0(x')|.$$

In other words, $x \notin I_{\mathsf{x}}^{\delta}$.

Next, suppose $|\Phi^-(\Phi(x)+)-x| \ge \epsilon$. Then, for every $\eta, \eta' > 0$, $\Phi^-(\Phi(x)+\eta) \ge x + \epsilon$ and therefore

$$\Phi(x+\epsilon-\eta') - \sup_{x'\in I_x^{\delta+\epsilon}} |\Phi(x') - \Phi_0(x')| - \eta < \Phi(x) - \sup_{x'\in I_x^{\delta+\epsilon}} |\Phi(x') - \Phi_0(x')|,$$

where, for $x \in I_{\times}^{\delta}$,

$$\Phi(x) - \sup_{x' \in I_x^{\delta + \epsilon}} |\Phi(x') - \Phi_0(x')| \le \Phi_0(x),$$

whereas

$$\lim_{\eta,\eta'\downarrow 0} \left[\Phi(x+\epsilon-\eta') - \sup_{x'\in I_x^{\delta+\epsilon}} |\Phi(x') - \Phi_0(x')| - \eta \right] \\
\geq \lim_{\eta'\downarrow 0} \inf \left[\Phi_0(x+\epsilon-\eta') - 2 \sup_{x'\in I_x^{\delta+\epsilon}} |\Phi(x') - \Phi_0(x')| \right] \\
\geq \Phi_0(x) + \left(\inf_{x'\in I_x^{\delta+\epsilon}} \partial\Phi_0(x') \right) \lim_{\eta'\downarrow 0} \inf \left[\epsilon - \frac{2 \sup_{x'\in I_x^{\delta+\epsilon}} |\Phi(x') - \Phi_0(x')|}{\inf_{x'\in I_x^{\delta+\epsilon}} \partial\Phi_0x'} - \eta' \right] > \Phi_0(x).$$

In other words, $x \notin I_{\times}^{\delta}$.

Next, we obtain two "local" approximations to LSC $_{\Phi}(f)$. The first of these is a generic approximation obtained by simply combining (SA.5) and Lemma SA-4, but for later reference we state the result as a separate lemma.

LEMMA SA-5. Suppose the assumptions of Lemma SA-4 hold. Then, for every $x \in I_{\times}^{\delta}$ and every $f: I \to \mathbb{R}$,

$$\inf_{|x'-x| \le \epsilon} f(x') \le \mathsf{LSC}_{\Phi}(f)(x) \le \sup_{|x'-x| \le \epsilon} f(x')$$

and

$$|\mathsf{LSC}_{\Phi}(f)(x) - f(x)| \le \sup_{|x' - x| \le \epsilon} |f(x') - f(x)|.$$

The final lemma is concerned with the special case where f is proportional to Φ . In that case, the following "local" analog of Lemma SA-3 shows that the bound(s) obtained in Lemma SA-5 can be improved under mild conditions on Φ .

LEMMA SA-6. Suppose the assumptions of Lemma SA-4 hold. Then, for every $x \in I_{\times}^{\delta}$ and every $\theta \in \mathbb{R}$,

$$|\mathsf{LSC}_{\Phi}(\theta\Phi)(x) - \theta\Phi(x)| \le |\theta| \sup_{x' \in I_x^{\delta + \epsilon}} |\Phi(x') - \Phi(x'-)|.$$

PROOF. First, if $\theta > 0$, then the result follows from the fact that, by (SA.5),

$$\mathsf{LSC}_{\Phi}(\theta\Phi)(x) = \theta\Phi(\Phi^{-}(\Phi(x)) -) \le \theta\Phi(\Phi^{-}(\Phi(x))) = \theta\Phi(x),$$

where, if $x \in I_{\mathsf{x}}^{\delta}$, then $\Phi^{-}(\Phi(x)) \in I_{\mathsf{x}}^{\delta+\epsilon}$ by Lemma SA-4, and therefore

$$\Phi(x) = \Phi(\Phi^{-}(\Phi(x))) \le \Phi(\Phi^{-}(\Phi(x)) - 1) + \sup_{x' \in I_{\delta}^{\delta + \epsilon}} |\Phi(x') - \Phi(x')|.$$

Next, if $\theta < 0$, then the result follows from the fact that, by (SA.5),

$$\mathsf{LSC}_{\Phi}(\theta\Phi)(x) = \theta\Phi(\Phi^{-}(\Phi(x)+)) \le \theta\Phi(\Phi^{-}(\Phi(x))) = \theta\Phi(x),$$

where, if $x \in I_{\mathsf{x}}^{\delta}$, then $\Phi^{-}(\Phi(x)+) \in I_{\mathsf{x}}^{\delta+\epsilon}$ by Lemma SA-4, and therefore

$$\begin{split} \Phi(\Phi^-(\Phi(x)+)) &\leq \liminf_{\eta \downarrow 0} \Phi(\Phi^-(\Phi(x)+\eta)-) + \sup_{x' \in I_{\mathbf{x}}^{\delta+\epsilon}} |\Phi(x') - \Phi(x'-)| \\ &\leq \Phi(x) + \sup_{x' \in I_{\mathbf{x}}^{\delta+\epsilon}} |\Phi(x') - \Phi(x'-)|. \end{split}$$

SA.1.4. Proof of Theorem 1.

Proof of (2). Let $t \in \mathbb{R}$ be given. By Lemma A.1 and change of variables,

$$\begin{split} & \mathbb{P}\left[r_n(\widehat{\theta}_n(\mathsf{x}) - \theta_0(\mathsf{x})) > t\right] \\ & = \mathbb{P}\left[\max_{x \in \widehat{\Phi}_n^-([0,\widehat{u}_n])} \left\{ [\theta_0(\mathsf{x}) + tr_n^{-1}] \widehat{\Phi}_n(x) - \mathsf{LSC}_{\widehat{\Phi}_n}(\widehat{\Gamma}_n)(x) \right\} < \widehat{\Phi}_n^- \circ \widehat{\Phi}_n(\mathsf{x}) \right] \\ & = \mathbb{P}\left[\max_{v \in \widehat{V}_{\mathsf{x},n}^{\mathfrak{q}}} \widehat{H}_{\mathsf{x},n}^{\mathfrak{q}}(v;t) < \widehat{Z}_{\mathsf{x},n}^{\mathfrak{q}} \right], \end{split}$$

where

$$\widehat{V}_{\mathbf{x},n}^{\mathfrak{q}} = \left\{ a_n(x - \mathbf{x}) : x \in \widehat{\Phi}_n^-([0, \widehat{u}_n]) \right\},$$

$$\widehat{H}_{\mathbf{x},n}^{\mathfrak{q}}(v;t) = \mathsf{LSC}_{\widehat{L}_{\mathbf{x},n}^{\mathfrak{q}}} \left(\widehat{G}_{\mathbf{x},n}^{\mathfrak{q}} + M_{\mathbf{x},n}^{\mathfrak{q}} + r_n \theta_0(\mathbf{x}) \widehat{L}_{\mathbf{x},n}^{\mathfrak{q}} \right)(v) - [r_n \theta_0(\mathbf{x}) + t] \widehat{L}_{\mathbf{x},n}^{\mathfrak{q}}(v),$$

and

$$\widehat{Z}_{\mathbf{x},n}^{\mathfrak{q}} = a_n \left[\widehat{\Phi}_n^- \circ \widehat{\Phi}_n(\mathbf{x}) - \mathbf{x} \right],$$

with

$$\begin{split} \widehat{G}_{\mathbf{x},n}^{\mathbf{q}}(v) &= \sqrt{na_n} \left[\widehat{\Gamma}_n(\mathbf{x} + va_n^{-1}) - \widehat{\Gamma}_n(\mathbf{x}) - \Gamma_0(\mathbf{x} + va_n^{-1}) + \Gamma_0(\mathbf{x}) \right] \\ &\quad - \theta_0(\mathbf{x}) \sqrt{na_n} \left[\widehat{\Phi}_n(\mathbf{x} + va_n^{-1}) - \widehat{\Phi}_n(\mathbf{x}) - \Phi_0(\mathbf{x} + va_n^{-1}) + \Phi_0(\mathbf{x}) \right], \\ M_{\mathbf{x},n}^{\mathbf{q}}(v) &= \sqrt{na_n} \left[\Gamma_0(\mathbf{x} + va_n^{-1}) - \Gamma_0(\mathbf{x}) \right] - \theta_0(\mathbf{x}) \sqrt{na_n} \left[\Phi_0(\mathbf{x} + va_n^{-1}) - \Phi_0(\mathbf{x}) \right], \\ \widehat{L}_{\mathbf{x},n}^{\mathbf{q}}(v) &= a_n \left[\widehat{\Phi}_n(\mathbf{x} + va_n^{-1}) - \widehat{\Phi}_n(\mathbf{x}) \right]. \end{split}$$

By (B4) and Lemma SA-4, $\widehat{Z}_{\mathbf{x},n}^{\mathfrak{q}} = o_{\mathbb{P}}(1)$. Suppose also that

(SA.6)
$$\max \underset{v \in \widehat{V}_{\mathbf{x},n}^{\mathfrak{q}}}{\operatorname{argmin}} \, \widehat{H}_{\mathbf{x},n}^{\mathfrak{q}}(v;t) \leadsto \underset{v \in \mathbb{R}}{\operatorname{argmin}} \, \mathcal{H}_{\mathbf{x}}^{\mathfrak{q}}(v;t),$$

where $\mathcal{H}_{\mathsf{x}}^{\mathfrak{q}}(v;t) = \mathcal{G}_{\mathsf{x}}(v) + \mathcal{M}_{\mathsf{x}}^{\mathfrak{q}}(v) - t\partial\Phi_{0}(\mathsf{x})v$. Then

$$\mathbb{P}\left[r_{n}(\widehat{\theta}_{n}(\mathsf{x}) - \theta_{0}(\mathsf{x})) > t\right] = \mathbb{P}\left[\max \underset{v \in \widehat{V}_{\mathsf{x},n}}{\operatorname{argmin}} \widehat{H}_{\mathsf{x},n}^{\mathsf{q}}(v;t) < \widehat{Z}_{\mathsf{x},n}^{\mathsf{q}}\right]$$

$$\to \mathbb{P}\left[\underset{v \in \mathbb{R}}{\operatorname{argmin}} \mathcal{H}_{\mathsf{x}}^{\mathsf{q}}(v;t) < 0\right]$$

$$= \mathbb{P}\left[\frac{1}{\partial \Phi_{0}(\mathsf{x})} \partial_{-}\mathsf{GCM}_{\mathbb{R}}(\mathcal{G}_{\mathsf{x}} + \mathcal{M}_{\mathsf{x}}^{\mathsf{q}}(0) > t\right],$$

where the second line uses Lemma A.2 and where the last equality uses Lemma SA-1. The proof of (2) can therefore be completed by showing (SA.6).

We shall do so by means of the argmax continuous mapping theorem of Cox (2022). To be specific, using that theorem it can be shown that (SA.6) holds if

(SA.7)
$$\widehat{v}_n(t) = \max \underset{v \in \widehat{V}_{\mathbf{x},n}}{\operatorname{argmin}} \widehat{H}_{\mathbf{x},n}^{\mathfrak{q}}(v;t) = O_{\mathbb{P}}(1)$$

and if

(SA.8)
$$\widehat{H}_{\mathbf{x},n}^{\mathfrak{q}}(\cdot;t) \rightsquigarrow \mathcal{H}_{\mathbf{x}}^{\mathfrak{q}}(\cdot;t).$$

We begin by showing (SA.8). First, by (B1), $\widehat{G}_{x,n}^{\mathfrak{q}} \rightsquigarrow \mathcal{G}_{x}$. Also, by (A2) and (A3), as $u \to 0$,

$$\begin{split} \frac{\Gamma_0(\mathbf{x}+u) - \theta_0(\mathbf{x})\Phi_0(\mathbf{x}+u) - \Gamma_0(\mathbf{x}) + \theta_0(\mathbf{x})\Phi_0(\mathbf{x})}{u^{\mathfrak{q}+1}} \to \lim_{u \to 0} \frac{\left[\theta_0(\mathbf{x}+u) - \theta_0(\mathbf{x})\right]\partial\Phi_0(\mathbf{x}+u)}{(\mathfrak{q}+1)u^{\mathfrak{q}}} \\ &= \frac{\partial^{\mathfrak{q}}\theta_0(\mathbf{x})\partial\Phi_0(\mathbf{x})}{(\mathfrak{q}+1)!}, \end{split}$$

where the first equality uses L'Hôpital's rule and

$$\partial \left[\Gamma_0(\mathsf{x} + u) - \theta_0(\mathsf{x}) \Phi_0(\mathsf{x} + u) - \Gamma_0(\mathsf{x}) + \theta_0(\mathsf{x}) \Phi_0(\mathsf{x}) \right] = \left[\theta_0(\mathsf{x} + u) - \theta_0(\mathsf{x}) \right] \partial \Phi_0(\mathsf{x} + u).$$

As a consequence, $M_{\mathsf{x},n}^{\mathfrak{q}} \leadsto \mathcal{M}_{\mathsf{x}}^{\mathfrak{q}}$. Moreover, $\widehat{L}_{\mathsf{x},n}^{\mathfrak{q}} - L_{\mathsf{x},n}^{\mathfrak{q}} \leadsto 0$ by (B4) and $L_{\mathsf{x},n}^{\mathfrak{q}} \leadsto \mathcal{L}_{\mathsf{x}}$ by (A3), where $L_{\mathsf{x},n}^{\mathfrak{q}}(v) = a_n \left[\Phi_0(\mathsf{x} + v a_n^{-1}) - \Phi_0(\mathsf{x}) \right]$. In particular, $\widehat{L}_{\mathsf{x},n}^{\mathfrak{q}} \leadsto \mathcal{L}_{\mathsf{x}}$ and therefore

$$(\widehat{G}_{\mathsf{x},n}^{\mathfrak{q}}, M_{\mathsf{x},n}^{\mathfrak{q}}, \widehat{L}_{\mathsf{x},n}^{\mathfrak{q}}) \leadsto (\mathcal{G}_{\mathsf{x}}, \mathcal{M}_{\mathsf{x}}^{\mathfrak{q}}, \mathcal{L}_{\mathsf{x}}).$$

Because $\widehat{G}_{x,n}^{\mathfrak{q}} + M_{x,n}^{\mathfrak{q}}$ is asymptotically equicontinuous,

$$\mathsf{LSC}_{\widehat{L}_{\mathsf{x},n}^{\mathfrak{q}}}\left(\widehat{G}_{\mathsf{x},n}^{\mathfrak{q}} + M_{\mathsf{x},n}^{\mathfrak{q}}\right) - \left(\widehat{G}_{\mathsf{x},n}^{\mathfrak{q}} + M_{\mathsf{x},n}^{\mathfrak{q}}\right) \leadsto 0$$

by (B4) and Lemma SA-5. Also, by (B7) and Lemma SA-6,

$$\mathsf{LSC}_{\widehat{L}_{\mathsf{x},n}^{\mathfrak{q}}}\left(r_n\theta_0(\mathsf{x})\widehat{L}_{\mathsf{x},n}^{\mathfrak{q}}\right) - \left(r_n\theta_0(\mathsf{x})\widehat{L}_{\mathsf{x},n}^{\mathfrak{q}}\right) \leadsto 0.$$

The result (SA.8) follows from the three preceding displays and the fact that, on $\hat{V}_{x,n}^{\mathfrak{q}}$

$$\begin{split} 0 &\geq \mathsf{LSC}_{\widehat{L}_{\mathsf{x},n}^{\mathfrak{q}}} \left(\widehat{G}_{\mathsf{x},n}^{\mathfrak{q}} + M_{\mathsf{x},n}^{\mathfrak{q}} + r_n \theta_0(\mathsf{x}) \widehat{L}_{\mathsf{x},n}^{\mathfrak{q}} \right) - \left(\widehat{G}_{\mathsf{x},n}^{\mathfrak{q}} + M_{\mathsf{x},n}^{\mathfrak{q}} + r_n \theta_0(\mathsf{x}) \widehat{L}_{\mathsf{x},n}^{\mathfrak{q}} \right) \\ &\geq \mathsf{LSC}_{\widehat{L}_{\mathsf{x},n}^{\mathfrak{q}}} \left(\widehat{G}_{\mathsf{x},n}^{\mathfrak{q}} + M_{\mathsf{x},n}^{\mathfrak{q}} \right) - \left(\widehat{G}_{\mathsf{x},n}^{\mathfrak{q}} + M_{\mathsf{x},n}^{\mathfrak{q}} \right) + \mathsf{LSC}_{\widehat{L}_{\mathsf{x},n}^{\mathfrak{q}}} \left(r_n \theta_0(\mathsf{x}) \widehat{L}_{\mathsf{x},n}^{\mathfrak{q}} \right) - \left(r_n \theta_0(\mathsf{x}) \widehat{L}_{\mathsf{x},n}^{\mathfrak{q}} \right). \end{split}$$

Next, to show (SA.7), we first define $\theta_n(x;t) = \theta_0(x) + tr_n^{-1}$ and note that

$$\widehat{v}_n(t) = a_n \left[\max \underset{\mathbf{x} \in \widehat{\Phi}_n^-([0,\widehat{u}_n])}{\operatorname{argmax}} \left\{ \theta_n(\mathbf{x};t) \widehat{\Phi}_n(x) - \mathsf{LSC}_{\widehat{\Phi}_n} \left(\widehat{\Gamma}_n \right)(x) \right\} - \mathsf{x} \right].$$

Now, if $|\widehat{v}_n(t)| > a_n \delta > 0$, then

$$\sup_{x\not\in I_{\mathbf{x}}^{\delta}}\left\{\theta_{n}(\mathbf{x};t)\widehat{\Phi}_{n}(x)-\mathsf{LSC}_{\widehat{\Phi}_{n}}\left(\widehat{\boldsymbol{\Gamma}}_{n}\right)(x)\right\}\geq\theta_{n}(\mathbf{x};t)\widehat{\Phi}_{n}(\mathbf{x})-\mathsf{LSC}_{\widehat{\Phi}_{n}}\left(\widehat{\boldsymbol{\Gamma}}_{n}\right)(\mathbf{x}),$$

where $|\theta_n(x;t) - \theta_0(x)| = O(r_n^{-1}) = o(1)$, and, by (B4),

$$\sup_{x \in I} \left| \widehat{\Phi}_n(x) - \Phi_0(x) \right| = o_{\mathbb{P}}(1).$$

Also, using (B3), (B4), and Lemma SA-2,

$$\begin{split} &\sup_{x\in I} \left| \mathsf{LSC}_{\widehat{\Phi}_n} \left(\widehat{\Gamma}_n \right)(x) - \Gamma_0(x) \right| \\ &\leq \sup_{x\in I} \left| \mathsf{LSC}_{\widehat{\Phi}_n} \left(\widehat{\Gamma}_n \right)(x) - \mathsf{LSC}_{\widehat{\Phi}_n} \left(\Gamma_0 \right)(x) \right| + \sup_{x\in I} \left| \mathsf{LSC}_{\widehat{\Phi}_n} \left(\Gamma_0 \right)(x) - \Gamma_0(x) \right| \\ &\leq \sup_{x\in I} \left| \widehat{\Gamma}_n(x) - \Gamma_0(x) \right| + 2 \left(\sup_{x\in I} |\theta_0(x)| \right) \sup_{x\in I} |\widehat{\Phi}_n(x) - \Phi_0(x)| = o_{\mathbb{P}}(1). \end{split}$$

As a consequence, $\widehat{v}_n(t) = o_{\mathbb{P}}(a_n)$: For any $\delta > 0$,

$$\mathbb{P}\left[\left|\widehat{v}_n(t)\right| > a_n \delta\right]$$

$$\leq \mathbb{P}\left[\sup_{x \notin I_{\mathsf{x}}^{\delta}} \left\{\theta_{0}(\mathsf{x})\Phi_{0}(x) - \Gamma_{0}(x)\right\} \geq \theta_{0}(\mathsf{x})\Phi_{0}(\mathsf{x}) - \Gamma_{0}(\mathsf{x}) + o_{\mathbb{P}}(1)\right] = o(1),$$

where the equality uses the fact, noted by Westling and Carone (2020, Supplement, proof of Lemma 3), that the function $v \mapsto \theta_0(x)\Phi_0(v) - \Gamma_0(v)$ is unimodal and maximized at v = x.

Next, defining $\widehat{V}_{\mathsf{x},n}^{\mathfrak{q}}(j) = \{v \in \widehat{V}_{\mathsf{x},n}^{\mathfrak{q}} : 2^j < |v| \le 2^{j+1}\}$ and using $\widehat{v}_n(t) = o_{\mathbb{P}}(a_n)$, we have, for any K, any positive δ' , and any sequence of events $\{\mathcal{A}'_n\}$ with $\lim_{n \to \infty} \mathbb{P}[\mathcal{A}'_n] = 1$,

$$\limsup_{n \to \infty} \mathbb{P}\left[|\widehat{v}_n(t)| > 2^K\right] \leq \limsup_{n \to \infty} \sum_{j \geq K: 2^j \leq a_n \delta'} \mathbb{P}\left[2^j < |\widehat{v}_n(t)| \leq 2^{j+1} \cap \mathcal{A}'_n\right]$$

$$\leq \limsup_{n \to \infty} \sum_{j \geq K: 2^j \leq a_n \delta'} \mathbb{P} \left[\inf_{v \in \widehat{V}_{\mathsf{x},n}^{\mathfrak{q}}(j)} \widehat{H}_{\mathsf{x},n}^{\mathfrak{q}}(v;t) \leq \widehat{H}_{\mathsf{x},n}^{\mathfrak{q}}(0;t) \cap \mathcal{A}'_n \right].$$

The proof of (SA.7) can therefore be completed by showing that the majorant side in the display can be made arbitrarily small by choice of K, δ' , and $\{A'_n\}$.

To do so, we begin by analyzing each term in the basic bound

$$\begin{split} \widehat{H}_{\mathbf{x},n}^{\mathfrak{q}}(v;t) - \widehat{H}_{\mathbf{x},n}^{\mathfrak{q}}(0;t) &\geq \mathsf{LSC}_{\widehat{L}_{\mathbf{x},n}^{\mathfrak{q}}} \left(\widehat{G}_{\mathbf{x},n}^{\mathfrak{q}} \right)(v) + \mathsf{LSC}_{\widehat{L}_{\mathbf{x},n}^{\mathfrak{q}}} \left(M_{\mathbf{x},n}^{\mathfrak{q}} \right)(v) - t \widehat{L}_{\mathbf{x},n}^{\mathfrak{q}}(v) \\ &+ \mathsf{LSC}_{\widehat{L}_{\mathbf{x},n}^{\mathfrak{q}}} \left(r_n \theta_0(\mathbf{x}) \widehat{L}_{\mathbf{x},n}^{\mathfrak{q}} \right)(v) - r_n \theta_0(\mathbf{x}) \widehat{L}_{\mathbf{x},n}^{\mathfrak{q}}(v) - \widehat{H}_{\mathbf{x},n}^{\mathfrak{q}}(0;t). \end{split}$$

Because $\widehat{H}_{\mathsf{x},n}^{\mathfrak{q}}(0;t) \rightsquigarrow \mathcal{H}_{\mathsf{x}}^{\mathfrak{q}}(0;t) = 0$ and because, by (B7) and Lemma SA-6, there is a positive δ' such that

$$\sup_{|v| \leq a_n \delta'} \left| \mathsf{LSC}_{\widehat{L}^{\mathfrak{q}}_{\mathsf{x},n}} \left(r_n \theta_0(\mathsf{x}) \widehat{L}^{\mathfrak{q}}_{\mathsf{x},n} \right)(v) - r_n \theta_0(\mathsf{x}) \widehat{L}^{\mathfrak{q}}_{\mathsf{x},n}(v) \right| = o_{\mathbb{P}}(1),$$

we may assume that, on $\{A'_n\}$ and for some C_0 ,

$$\sup_{|v| < a_n \delta'} \left| \mathsf{LSC}_{\widehat{L}^{\mathfrak{q}}_{\mathsf{x},n}} \left(r_n \theta_0(\mathsf{x}) \widehat{L}^{\mathfrak{q}}_{\mathsf{x},n} \right)(v) - r_n \theta_0(\mathsf{x}) \widehat{L}^{\mathfrak{q}}_{\mathsf{x},n}(v) - \widehat{H}^{\mathfrak{q}}_{\mathsf{x},n}(0;t) \right| \leq C_0.$$

Also, because, by (B4) and (A3), there is a positive δ' such that

$$\sup_{|v| \leq a_n \delta'} \left| \widehat{L}_{\mathsf{x},n}^{\mathfrak{q}}(v) - L_{\mathsf{x},n}^{\mathfrak{q}}(v) \right| = o_{\mathbb{P}}(1) \qquad \text{and} \qquad \sup_{1 \leq |v| \leq a_n \delta'} \left| \frac{L_{\mathsf{x},n}^{\mathfrak{q}}(v)}{\mathcal{L}_{\mathsf{x}}(v)} \right| < \infty,$$

we may assume that, on $\{A'_n\}$ and for some C_L ,

$$\sup_{1 \le |v| \le a_n \delta'} \left| \frac{\widehat{L}_{\mathsf{x},n}^{\mathfrak{q}}(v)}{v} \right| \le C_L.$$

Next, by (B4) and Lemma SA-5, with probability approaching one,

$$\mathsf{LSC}_{\widehat{L}_{\mathsf{x},n}^{\mathfrak{q}}}\left(M_{\mathsf{x},n}^{\mathfrak{q}}\right)(v) \geq \inf_{|v|/2 < |v'| < 2|v|} M_{\mathsf{x},n}^{\mathfrak{q}}(v') \qquad \text{for every } |v| \geq 2,$$

while, by (A2) and (A3), there is a positive δ' such that

$$\inf_{1 \le |v| \le \eta a_n \delta'} \frac{M_{\mathsf{x},n}^{\mathsf{q}}(v)}{\mathcal{M}_{\mathsf{q}}^{\mathsf{q}}(v)} > 0.$$

We may therefore assume that, on $\{A'_n\}$ and for some positive C_M ,

$$\inf_{2 \leq |v| \leq a_n \delta'} \frac{\mathsf{LSC}_{\widehat{L}^{\mathfrak{q}}_{\mathsf{x},n}}\left(M^{\mathfrak{q}}_{\mathsf{x},n}\right)(v)}{v^{\mathfrak{q}+1}} \geq C_M.$$

Finally, by (B4) and Lemma SA-5, with probability approaching one,

$$\left|\mathsf{LSC}_{\widehat{L}_{\mathsf{x},n}^{\mathfrak{q}}}\left(\widehat{G}_{\mathsf{x},n}^{\mathfrak{q}}\right)(v)\right| \leq \sup_{|v|/2 \leq |v'| \leq 2|v|} \left|\widehat{G}_{\mathsf{x},n}^{\mathfrak{q}}(v')\right| \qquad \text{for every } |v| \geq 2,$$

and we may therefore assume that, on $\{A'_n\}$,

$$\sup_{v \in \widehat{V}_{\mathbf{x},n}^{\mathfrak{q}}(j)} \left| \mathsf{LSC}_{\widehat{L}_{\mathbf{x},n}^{\mathfrak{q}}} \left(\widehat{G}_{\mathbf{x},n}^{\mathfrak{q}} \right)(v) \right| \leq \sup_{v \in V_2(j)} \left| \widehat{G}_{\mathbf{x},n}^{\mathfrak{q}}(v) \right| \qquad \text{for every } j \geq 2 \text{ with } 2^j \leq a_n \delta',$$

where $V_{\eta}(j) = \{v \in \mathbb{R} : \eta^{-1}2^{j} \le |v| \le \eta 2^{j+1}\}.$

As a consequence, by the Markov inequality,

$$\begin{split} & \mathbb{P}\left[\inf_{v \in \widehat{V}_{\mathbf{x},n}^{\mathfrak{q}}(j)} \widehat{H}_{\mathbf{x},n}^{\mathfrak{q}}(v;t) \leq \widehat{H}_{\mathbf{x},n}^{\mathfrak{q}}(0;t) \cap \mathcal{A}_{n}'\right] \\ & \leq \mathbb{P}\left[\sup_{v \in V_{2}(j)} \left|\widehat{G}_{\mathbf{x},n}^{\mathfrak{q}}(v)\right| \geq \inf_{v \in V_{1}(j)} \left[C_{M}v^{\mathfrak{q}+1} - C_{L}|v| - C_{0}\right] \cap \mathcal{A}_{n}'\right] \\ & \leq \frac{\mathbb{E}\left[\sup_{v \in V_{2}(j)} \left|\widehat{G}_{\mathbf{x},n}^{\mathfrak{q}}(v)\right| \mathbbm{1}_{\mathcal{A}_{n}'}\right]}{\inf_{v \in V_{1}(j)} \left[C_{M}v^{\mathfrak{q}+1} - C_{L}|v| - C_{0}\right]} \quad \text{ for every } j \geq 2 \text{ with } 2^{j} \leq a_{n}\delta', \end{split}$$

where, by (B2), we may assume that, for some C_G ,

$$\mathbb{E}\left[\sup_{v \in V_2(j)} \left| \widehat{G}_{\mathsf{x},n}^{\mathfrak{q}}(v) \right| \mathbb{1}_{\mathcal{A}_n'} \right] \leq \sum_{j-1 < j' < j+1} \mathbb{E}\left[\sup_{v \in V_1(j')} \left| \widehat{G}_{\mathsf{x},n}^{\mathfrak{q}}(v) \right| \mathbb{1}_{\mathcal{A}_n'} \right] \leq C_G 2^{j\beta},$$

and where, for all sufficiently large j,

$$\inf_{v \in V_1(j)} \left[C_M v^{\mathfrak{q}+1} - C_L |v| - C_0 \right] \ge \frac{1}{2} C_M 2^{j(\mathfrak{q}+1)}.$$

In other words, for large K,

$$\limsup_{n \to \infty} \sum_{j > K: 2^j < a_n \delta'} \mathbb{P}\left[\inf_{v \in \widehat{V}_{\mathbf{x},n}^{\mathfrak{q}}(j)} \widehat{H}_{\mathbf{x},n}^{\mathfrak{q}}(v;t) \leq \widehat{H}_{\mathbf{x},n}^{\mathfrak{q}}(0;t) \cap \mathcal{A}_n'\right] \leq \frac{2C_G}{C_M} \sum_{j > K} 2^{j[\beta - (\mathfrak{q} + 1)]},$$

which can be made arbitrarily small by choice of K.

Proof of (7). We proceed as in the proof of (2). Let $t \in \mathbb{R}$ be given. By Lemma A.1 and change of variables,

$$\begin{split} &\mathbb{P}_n^* \left[r_n(\widetilde{\theta}_n^*(\mathbf{x}) - \widehat{\theta}_n(\mathbf{x})) > t \right] \\ &= \mathbb{P}_n^* \left[\max \underset{x \in \widehat{\Phi}_n^{*-}([0,\widehat{u}_n^*])}{\operatorname{argmax}} \left\{ [\widehat{\theta}_n(\mathbf{x}) + tr_n^{-1}] \widehat{\Phi}_n^*(x) - \mathsf{LSC}_{\widehat{\Phi}_n^*}(\widetilde{\Gamma}_n^*)(x) \right\} < \widehat{\Phi}_n^{*-} \circ \widehat{\Phi}_n^*(\mathbf{x}) \right] \\ &= \mathbb{P}_n^* \left[\max \underset{v \in \widehat{V}_{\mathbf{x},n}^{\mathfrak{q},*}}{\operatorname{argmin}} \widehat{H}_{\mathbf{x},n}^{\mathfrak{q},*}(v;t) < \widehat{Z}_{\mathbf{x},n}^{\mathfrak{q},*} \right], \end{split}$$

where

$$\begin{split} \widehat{V}_{\mathbf{x},n}^{\mathfrak{q},*} &= \left\{ a_n(x-\mathbf{x}) : x \in \widehat{\Phi}_n^{*-}([0,\widehat{u}_n^*]) \right\}, \\ \widehat{H}_{\mathbf{x},n}^{\mathfrak{q},*}(v;t) &= \mathsf{LSC}_{\widehat{L}_{\mathbf{x},n}^{\mathfrak{q},*}} \left(\widehat{G}_{\mathbf{x},n}^{\mathfrak{q},*} + \widetilde{M}_{\mathbf{x},n}^{\mathfrak{q}} + r_n \widehat{\theta}_n(\mathbf{x}) \widehat{L}_{\mathbf{x},n}^{\mathfrak{q},*} \right)(v) - [r_n \widehat{\theta}_n(\mathbf{x}) + t] \widehat{L}_{\mathbf{x},n}^{\mathfrak{q},*}(v) \end{split}$$

and

$$\widehat{Z}_{\mathbf{x},n}^{\mathfrak{q},*} = a_n \left[\widehat{\Phi}_n^{*-} \circ \widehat{\Phi}_n^*(\mathbf{x}) - \mathbf{x} \right],$$

with

$$\begin{split} \widehat{G}_{\mathbf{x},n}^{\mathfrak{q},*}(v) &= \sqrt{na_n} \left[\widehat{\Gamma}_n^*(\mathbf{x} + va_n^{-1}) - \widehat{\Gamma}_n^*(\mathbf{x}) - \widehat{\Gamma}_n(\mathbf{x} + va_n^{-1}) + \widehat{\Gamma}_n(\mathbf{x}) \right] \\ &\qquad \qquad - \widehat{\theta}_n(\mathbf{x}) \sqrt{na_n} \left[\widehat{\Phi}_n^*(\mathbf{x} + va_n^{-1}) - \widehat{\Phi}_n^*(\mathbf{x}) - \widehat{\Phi}_n(\mathbf{x} + va_n^{-1}) + \widehat{\Phi}_n(\mathbf{x}) \right], \\ \widetilde{M}_{\mathbf{x},n}^{\mathfrak{q}}(v) &= \sqrt{na_n} \widetilde{M}_{\mathbf{x},n}(va_n^{-1}), \qquad \widehat{L}_{\mathbf{x},n}^{\mathfrak{q},*}(v) = a_n \left[\widehat{\Phi}_n^*(\mathbf{x} + va_n^{-1}) - \widehat{\Phi}_n^*(\mathbf{x}) \right]. \end{split}$$

By (B4) and Lemma SA-4, $\widehat{Z}_{\mathbf{x},n}^{\mathfrak{q},*} = o_{\mathbb{P}}(1)$. Suppose also that

(SA.9)
$$\max \underset{v \in \widehat{\mathbb{V}}_{\mathbf{x},n}^{\mathfrak{q},*}}{\operatorname{argmin}} \widehat{H}_{\mathsf{x},n}^{\mathfrak{q},*}(v;t) \leadsto_{\mathbb{P}} \underset{v \in \mathbb{R}}{\operatorname{argmin}} \mathcal{H}_{\mathsf{x}}^{\mathfrak{q}}(v;t).$$

Then, as in the proof of (2),

$$\mathbb{P}_n^* \left[r_n(\widetilde{\theta}_n^*(\mathbf{x}) - \widehat{\theta}_n(\mathbf{x})) > t \right] \to_{\mathbb{P}} \mathbb{P} \left[\frac{1}{\partial \Phi_0(\mathbf{x})} \partial_- \mathsf{GCM}_{\mathbb{R}}(\mathcal{G}_{\mathbf{x}} + \mathcal{M}_{\mathbf{x}}^{\mathfrak{q}}(0) > t \right].$$

The proof of (7) can therefore be completed by showing (SA.9).

We shall do so by showing that

$$\widehat{v}_n^*(t) = \max \underset{v \in \widehat{V}_{\mathbf{x},n}^{\mathfrak{q},*}}{\operatorname{argmin}} \widehat{H}_{\mathbf{x},n}^{\mathfrak{q},*}(v;t) = O_{\mathbb{P}}(1)$$

and

(SA.11)
$$\widehat{H}_{\mathsf{x},n}^{\mathfrak{q},*}(\cdot;t) \leadsto_{\mathbb{P}} \mathcal{H}_{\mathsf{x}}^{\mathfrak{q}}(\cdot;t).$$

We begin by showing (SA.11). First, by (B1), $\widehat{G}_{x,n}^{q,*} \leadsto_{\mathbb{P}} \mathcal{G}_{x}$. Also, by Assumption C, $\widetilde{M}_{x,n}^{q,*} \leadsto_{\mathbb{P}} \mathcal{M}_{x}^{q}$. Moreover, $\widehat{L}_{x,n}^{q,*} - \widehat{L}_{x,n}^{q} \leadsto_{\mathbb{P}} 0$ by (B4), where, as shown in the proof of (SA.6), $\widehat{L}_{x,n}^{q} \leadsto_{\mathbb{P}} \mathcal{L}_{x}$. In particular, $\widehat{L}_{x,n}^{q,*} \leadsto_{\mathbb{P}} \mathcal{L}_{x}$ and therefore

$$(\widehat{G}_{\mathsf{x},n}^{\mathfrak{q},*}, \widetilde{M}_{\mathsf{x},n}^{\mathfrak{q}}, \widehat{L}_{\mathsf{x},n}^{\mathfrak{q},*}) \leadsto_{\mathbb{P}} (\mathcal{G}_{\mathsf{x}}^{\mathfrak{q}}, \mathcal{M}_{\mathsf{x}}^{\mathfrak{q}}, \mathcal{L}_{\mathsf{x}}).$$

Because $\widehat{G}_{x,n}^{q,*} + \widetilde{M}_{x,n}^{q}$ is asymptotically equicontinuous,

$$\mathsf{LSC}_{\widehat{C}_{\mathsf{X},n}^{\mathfrak{q},*}}\left(\widehat{G}_{\mathsf{X},n}^{\mathfrak{q},*}+\widetilde{M}_{\mathsf{X},n}^{\mathfrak{q}}\right)-\left(\widehat{G}_{\mathsf{X},n}^{\mathfrak{q},*}+\widetilde{M}_{\mathsf{X},n}^{\mathfrak{q}}\right)\leadsto_{\mathbb{P}}0$$

by (B4) and Lemma SA-5. Also, by (B7) and Lemma SA-6,

$$\mathsf{LSC}_{\widehat{L}_{\mathsf{x},n}^{\mathfrak{q},*}}\left(r_n\widehat{\theta}_n(\mathsf{x})\widehat{L}_{\mathsf{x},n}^{\mathfrak{q},*}\right) - \left(r_n\widehat{\theta}_n(\mathsf{x})\widehat{L}_{\mathsf{x},n}^{\mathfrak{q},*}\right) \leadsto_{\mathbb{P}} 0.$$

The result (SA.11) follows from the three preceding displays and the fact that, on $\hat{V}_{x,n}^{q,*}$,

$$\begin{split} 0 &\geq \mathsf{LSC}_{\widehat{L}_{\mathsf{x},n}^{\mathfrak{q},*}} \left(\widehat{G}_{\mathsf{x},n}^{\mathfrak{q},*} + \widetilde{M}_{\mathsf{x},n}^{\mathfrak{q}} + r_n \widehat{\theta}_n(\mathsf{x}) \widehat{L}_{\mathsf{x},n}^{\mathfrak{q},*} \right) - \left(\widehat{G}_{\mathsf{x},n}^{\mathfrak{q},*} + \widetilde{M}_{\mathsf{x},n}^{\mathfrak{q}} + r_n \widehat{\theta}_n(\mathsf{x}), \widehat{L}_{\mathsf{x},n}^{\mathfrak{q},*} \right) \\ &\geq \mathsf{LSC}_{\widehat{L}_{\mathsf{x},n}^{\mathfrak{q},*}} \left(\widehat{G}_{\mathsf{x},n}^{\mathfrak{q},*} + \widetilde{M}_{\mathsf{x},n}^{\mathfrak{q}} \right) - \left(\widehat{G}_{\mathsf{x},n}^{\mathfrak{q},*} + \widetilde{M}_{\mathsf{x},n}^{\mathfrak{q}} \right) \\ &+ \mathsf{LSC}_{\widehat{L}_{\mathsf{x},n}^{\mathfrak{q},*}} \left(r_n \widehat{\theta}_n(\mathsf{x}) \widehat{L}_{\mathsf{x},n}^{\mathfrak{q},*} \right) - \left(r_n \widehat{\theta}_n(\mathsf{x}) \widehat{L}_{\mathsf{x},n}^{\mathfrak{q},*} \right). \end{split}$$

Next, to show (SA.10), we first define $\widehat{\theta}_n(\mathbf{x};t) = \widehat{\theta}_n(\mathbf{x}) + tr_n^{-1}$ and note that

$$\widehat{v}_n^*(t) = a_n \left[\max \underset{x \in \widehat{\Phi}_n^{*-}([0,\widehat{u}_n^*])}{\operatorname{argmax}} \left\{ \widehat{\theta}_n(\mathsf{x};t) \widehat{\Phi}_n^*(x) - \mathsf{LSC}_{\widehat{\Phi}_n^*} \left(\widetilde{\Gamma}_n^* \right)(x) \right\} - \mathsf{x} \right].$$

Now, if $|\widehat{v}_n^*(t)| > a_n \delta > 0$, then

$$\sup_{x\not\in I_{\diamond}^{\ast}}\left\{\widehat{\theta}_{n}(\mathbf{x};t)\widehat{\Phi}_{n}^{\ast}(x)-\mathsf{LSC}_{\widehat{\Phi}_{n}^{\ast}}\left(\widetilde{\Gamma}_{n}^{\ast}\right)(x)\right\}\geq\widehat{\theta}_{n}(\mathbf{x};t)\widehat{\Phi}_{n}^{\ast}(\mathbf{x})-\mathsf{LSC}_{\widehat{\Phi}_{n}^{\ast}}\left(\widetilde{\Gamma}_{n}^{\ast}\right)(\mathbf{x}),$$

where $|\widehat{\theta}_n(\mathbf{x};t) - \theta_0(\mathbf{x})| = O_{\mathbb{P}}(r_n^{-1}) = o_{\mathbb{P}}(1)$, and, by (B4),

$$\sup_{x \in I} \left| \widehat{\Phi}_n^*(x) - \Phi_0(x) \right| = o_{\mathbb{P}}(1).$$

Therefore, defining $\widehat{\mathbf{x}}_n^* = \widehat{\Phi}_n^{*-}(\widehat{\Phi}_n^*(\mathbf{x})) = \mathbf{x} + o_{\mathbb{P}}(1)$ and using (SA.5),

$$\begin{split} \widehat{\theta}_{n}(\mathbf{x};t)\widehat{\Phi}_{n}^{*}(\mathbf{x}) - \mathsf{LSC}_{\widehat{\Phi}_{n}^{*}}\left(\widetilde{\Gamma}_{n}^{*}\right)(\mathbf{x}) \\ &\geq \widehat{\theta}_{n}(\mathbf{x};t)\widehat{\Phi}_{n}^{*}(\mathbf{x}) - \widetilde{\Gamma}_{n}^{*}(\widehat{\mathbf{x}}_{n}^{*}) \\ &= \widehat{\theta}_{n}(\mathbf{x};t)\widehat{\Phi}_{n}^{*}(\mathbf{x}) - \widehat{\theta}_{n}(\mathbf{x})\widehat{\Phi}_{n}(\widehat{\mathbf{x}}_{n}^{*}) - \widehat{\Gamma}_{n}^{*}(\widehat{\mathbf{x}}_{n}^{*}) + \widehat{\Gamma}_{n}(\widehat{\mathbf{x}}_{n}^{*}) - \widetilde{M}_{\mathbf{x},n}^{\mathfrak{q}}(\widehat{Z}_{\mathbf{x},n}^{\mathfrak{q},*})/\sqrt{na_{n}} = o_{\mathbb{P}}(1), \end{split}$$

where the last equality uses (B3), $\widehat{Z}_{\mathsf{x},n}^{\mathfrak{q},*} = o_{\mathbb{P}}(1)$, and Assumption C. Also, using (B3), (B4), and Assumption C, we have, uniformly in $x \notin I_{\mathsf{x}}^{\delta}$ and for some c > 0,

$$\widetilde{\Gamma}_n^*(x) - \widehat{\theta}_n(x)\widehat{\Phi}_n^*(x) = \widehat{\Gamma}_n^*(x) - \widehat{\Gamma}_n(x) - \widehat{\theta}_n(x) \left(\widehat{\Phi}_n^*(x) - \widehat{\Phi}_n(x)\right) + \widetilde{M}_n(x - x)$$

$$\geq c\delta^{q+1} + o_{\mathbb{P}}(1),$$

and therefore, by Lemma SA-3,

$$\sup_{x\not\in I_{s}^{\delta}}\left\{\widehat{\theta}_{n}(\mathbf{x};t)\widehat{\Phi}_{n}^{*}(x)-\mathsf{LSC}_{\widehat{\Phi}_{n}^{*}}\left(\widetilde{\Gamma}_{n}^{*}\right)(x)\right\}\leq -c\delta^{\mathsf{q}+1}+o_{\mathbb{P}}(1).$$

As a consequence, $\widehat{v}_n^*(t) = o_{\mathbb{P}}(a_n)$: For any $\delta > 0$,

$$\mathbb{P}\left[|\widehat{v}_n^*(t)| > a_n \delta\right] \le \mathbb{P}\left[-c \delta^{\mathfrak{q}+1} \ge o_{\mathbb{P}}(1)\right] = o(1).$$

Next, defining $\widehat{V}_{\mathsf{x},n}^{\mathsf{q},*}(j) = \{v \in \widehat{V}_{\mathsf{x},n}^{\mathsf{q},*} : 2^j < |v| \le 2^{j+1}\}$ and using $\widehat{v}_n^*(t) = o_{\mathbb{P}}(a_n)$, we have, for any K, any positive δ' , and any sequence of events $\{\mathcal{A}_n'\}$ with $\lim_{n \to \infty} \mathbb{P}[\mathcal{A}_n'] = 1$,

$$\limsup_{n \to \infty} \mathbb{P}\left[|\widehat{v}_n^*(t)| > 2^K\right] \leq \limsup_{n \to \infty} \sum_{j > K: 2^j < a_n \delta'} \mathbb{P}\left[2^j < |\widehat{v}_n^*(t)| \leq 2^{j+1} \cap \mathcal{A}_n'\right]$$

$$\leq \limsup_{n \to \infty} \sum_{j \geq K: 2^j \leq a_n \delta'} \mathbb{P} \left[\inf_{v \in \widehat{V}_{\mathbf{x},n}^{\mathfrak{q},*}(j)} \widehat{H}_{\mathbf{x},n}^{\mathfrak{q},*}(v;t) \leq \widehat{H}_{\mathbf{x},n}^{\mathfrak{q},*}(0;t) \cap \mathcal{A}'_n \right].$$

The proof of (SA.10) can therefore be completed by showing that the majorant side in the display can be made arbitrarily small by choice of K, δ' , and $\{A'_n\}$.

To do so, we begin by analyzing each term in the basic bound

$$\begin{split} \widehat{H}_{\mathbf{x},n}^{\mathfrak{q},*}(v;t) - \widehat{H}_{\mathbf{x},n}^{\mathfrak{q},*}(0;t) &\geq \mathsf{LSC}_{\widehat{L}_{\mathbf{x},n}^{\mathfrak{q},*}} \left(\widehat{G}_{\mathbf{x},n}^{\mathfrak{q},*} \right)(v) + \mathsf{LSC}_{\widehat{L}_{\mathbf{x},n}^{\mathfrak{q},*}} \left(\widetilde{M}_{\mathbf{x},n}^{\mathfrak{q}} \right)(v) - t \widehat{L}_{\mathbf{x},n}^{\mathfrak{q},*}(v) \\ &+ \mathsf{LSC}_{\widehat{L}_{\mathbf{x},n}^{\mathfrak{q},*}} \left(r_n \widehat{\theta}_n(\mathbf{x}) \widehat{L}_{\mathbf{x},n}^{\mathfrak{q},*} \right)(v) - r_n \widehat{\theta}_n(\mathbf{x}) \widehat{L}_{\mathbf{x},n}^{\mathfrak{q},*}(v) - \widehat{H}_{\mathbf{x},n}^{\mathfrak{q},*}(0;t). \end{split}$$

Because $\widehat{H}_{\mathsf{x},n}^{\mathfrak{q},*}(0;t) \leadsto_{\mathbb{P}} \mathcal{H}_{\mathsf{x}}^{\mathfrak{q}}(0;t) = 0$ and because, by (B7) and Lemma SA-6, there is a positive δ' such that

$$\sup_{|v| \leq a_n \delta'} \left| \mathsf{LSC}_{\widehat{L}_{\mathsf{x},n}^{\mathfrak{q},*}} \left(r_n \widehat{\theta}_n(\mathsf{x}) \widehat{L}_{\mathsf{x},n}^{\mathfrak{q},*} \right)(v) - r_n \widehat{\theta}_n(\mathsf{x}) \widehat{L}_{\mathsf{x},n}^{\mathfrak{q},*}(v) \right| = o_{\mathbb{P}}(1),$$

we may assume that, on $\{A'_n\}$ and for some C_0 ,

$$\sup_{|v| < a_n \delta'} \left| \mathsf{LSC}_{\widehat{L}^{\mathfrak{q},*}_{\mathsf{x},n}} \left(r_n \widehat{\theta}_n(\mathsf{x}) \widehat{L}^{\mathfrak{q},*}_{\mathsf{x},n} \right)(v) - r_n \widehat{\theta}_n(\mathsf{x}) \widehat{L}^{\mathfrak{q},*}_{\mathsf{x},n}(v) - \widehat{H}^{\mathfrak{q},*}_{\mathsf{x},n}(0;t) \right| \leq C_0.$$

Also, because, by (B4) and (A3), there is a positive δ' such that

$$\sup_{|v| \leq a_n \delta'} \left| \widehat{L}_{\mathsf{x},n}^{\mathfrak{q},*}(v) - L_{\mathsf{x},n}^{\mathfrak{q}}(v) \right| = o_{\mathbb{P}}(1) \qquad \text{and} \qquad \sup_{1 \leq |v| \leq a_n \delta'} \left| \frac{L_{\mathsf{x},n}^{\mathfrak{q}}(v)}{\mathcal{L}_{\mathsf{x}}(v)} \right| < \infty,$$

we may assume that, on $\{A'_n\}$ and for some C_L

$$\sup_{1 \le |v| \le a_n \delta'} \left| \frac{\widehat{L}_{\mathsf{x},n}^{\mathsf{q},*}(v)}{v} \right| \le C_L.$$

Next, by (B4) and Lemma SA-5, with probability approaching one,

$$\mathsf{LSC}_{\widehat{L}_{\mathsf{x},n}^{\mathfrak{q},*}}\left(\widetilde{M}_{\mathsf{x},n}^{\mathfrak{q}}\right)(v) \geq \inf_{|v|/2 < |v'| < 2|v|} \widetilde{M}_{\mathsf{x},n}^{\mathfrak{q}}(v') \qquad \text{for every } |v| \geq 2,$$

while, by Assumption C, there is a positive c such that, with probability approaching one,

$$\inf_{|v|>1} \frac{\widetilde{M}_{\mathsf{x},n}^{\mathfrak{q}}(v)}{v^{\mathfrak{q}+1}} > c.$$

We may therefore assume that, on $\{A'_n\}$ and for some positive C_M ,

$$\inf_{2\leq |v|\leq a_n\delta'}\frac{\mathsf{LSC}_{\widehat{L}_{\mathsf{x},n}^{\mathfrak{q},*}}\left(\widetilde{M}_{\mathsf{x},n}^{\mathfrak{q}}\right)(v)}{v^{\mathfrak{q}+1}}\geq C_M.$$

Finally, by (B4) and Lemma SA-5, with probability approaching one,

$$\left|\mathsf{LSC}_{\widehat{L}^{\mathfrak{q},*}_{\mathsf{x},n}}\left(\widehat{G}^{\mathfrak{q},*}_{\mathsf{x},n}\right)(v)\right| \leq \sup_{|v|/2 < |v'| < 2|v|} \left|\widehat{G}^{\mathfrak{q},*}_{\mathsf{x},n}(v')\right| \qquad \text{for every } |v| \geq 2,$$

and we may therefore assume that, on $\{A'_n\}$,

$$\sup_{v \in \widehat{V}_{\mathbf{x},n}^{\mathfrak{q},*}(j)} \left| \mathsf{LSC}_{\widehat{L}_{\mathbf{x},n}^{\mathfrak{q},*}} \left(\widehat{G}_{\mathbf{x},n}^{\mathfrak{q},*} \right)(v) \right| \leq \sup_{v \in V_2(j)} \left| \widehat{G}_{\mathbf{x},n}^{\mathfrak{q},*}(v) \right| \qquad \text{for every } j \geq 2 \text{ with } 2^j \leq a_n \delta'.$$

As a consequence, by the Markov inequality,

$$\begin{split} & \mathbb{P}\left[\inf_{v \in \widehat{V}_{\mathbf{x},n}^{\mathfrak{q},*}(j)} \widehat{H}_{\mathbf{x},n}^{\mathfrak{q},*}(v;t) \leq \widehat{H}_{\mathbf{x},n}^{\mathfrak{q},*}(0;t) \cap \mathcal{A}_{n}'\right] \\ & \leq \mathbb{P}\left[\sup_{v \in V_{2}(j)} \left|\widehat{G}_{\mathbf{x},n}^{\mathfrak{q},*}(v)\right| \geq \inf_{v \in V_{1}(j)} \left[C_{M}v^{\mathfrak{q}+1} - C_{L}|v| - C_{0}\right] \cap \mathcal{A}_{n}'\right] \\ & \leq \frac{\mathbb{E}\left[\sup_{v \in V_{2}(j)} \left|\widehat{G}_{\mathbf{x},n}^{\mathfrak{q},*}(v)\right| \mathbbm{1}_{\mathcal{A}_{n}'}\right]}{\inf_{v \in V_{1}(j)} \left[C_{M}v^{\mathfrak{q}+1} - C_{L}|v| - C_{0}\right]} \quad \text{for every } j \geq 2 \text{ with } 2^{j} \leq a_{n}\delta', \end{split}$$

where, by (B4), we may assume that, for some C_G ,

$$\mathbb{E}\left[\sup_{v\in V_2(j)}\left|\widehat{G}_{\mathsf{x},n}^{\mathfrak{q},*}(v)\right|\mathbb{1}_{\mathcal{A}_n'}\right] \leq \sum_{j-1\leq j'\leq j+1}\mathbb{E}\left[\sup_{v\in V_1(j')}\left|\widehat{G}_{\mathsf{x},n}^{\mathfrak{q},*}(v)\right|\mathbb{1}_{\mathcal{A}_n'}\right] \leq C_G 2^{j\beta},$$

and where, for all sufficiently large j,

$$\inf_{v \in V_1(j)} \left[C_M v^{q+1} - C_L |v| - C_0 \right] \ge \frac{1}{2} C_M 2^{j(q+1)}.$$

In other words, for large K,

$$\limsup_{n\to\infty} \sum_{j\geq K: 2^j \leq a_n \delta'} \mathbb{P}\left[\inf_{v\in \widehat{V}_{\mathbf{x},n}^{\mathfrak{q},*}(j)} \widehat{H}_{\mathbf{x},n}^{\mathfrak{q},*}(v;t) \leq \widehat{H}_{\mathbf{x},n}^{\mathfrak{q},*}(0;t) \cap \mathcal{A}_n'\right] \leq \frac{2C_G}{C_M} \sum_{j\geq K} 2^{j[\beta-(\mathfrak{q}+1)]},$$

which can be made arbitrarily small by choice of K.

Proof of (8). The bootstrap consistency result (8) follows from (2), (7), Polya's theorem, and the fact that, by Lemma A.2, the limiting distribution in (2) and (7) has a continuous cdf. \Box

SA.1.5. *Proof of Lemma 1*. For the monomial approximation estimator, we have

$$\begin{split} \widetilde{\mathcal{D}}_{\mathfrak{q},n}^{\text{MA}}(\mathbf{x}) &= \epsilon_n^{-(\mathfrak{q}+1)} [\Gamma_0(\mathbf{x}+\epsilon_n) - \Gamma_0(\mathbf{x}) - \theta_0(\mathbf{x}) \{\Phi_0(\mathbf{x}+\epsilon_n) - \Phi_0(\mathbf{x})\}] \\ &+ \epsilon_n^{-(\mathfrak{q}+1/2)} n^{-1/2} \widehat{G}_{\mathbf{x},n}(1;\epsilon_n) \\ &- \epsilon_n^{-\mathfrak{q}} [\widehat{\theta}_n(\mathbf{x}) - \theta_0(\mathbf{x})] \widehat{R}_{\mathbf{x},n}(1;\epsilon_n) \\ &- \epsilon_n^{-(\mathfrak{q}+1)} [\widehat{\theta}_n(\mathbf{x}) - \theta_0(\mathbf{x})] [\Phi_0(\mathbf{x}+\epsilon_n) - \Phi_0(\mathbf{x})] \\ &= \frac{\partial^{\mathfrak{q}} \theta_0(\mathbf{x}) \partial \Phi_0(\mathbf{x})}{(\mathfrak{q}+1)!} + o(1) + O_{\mathbb{P}} [(n\epsilon_n^{1+2\mathfrak{q}})^{-1/2} + (n\epsilon_n^{1+2\mathfrak{q}})^{-\mathfrak{q}/(1+2\mathfrak{q})}] \\ &= \frac{\partial^{\mathfrak{q}} \theta_0(\mathbf{x}) \partial \Phi_0(\mathbf{x})}{(\mathfrak{q}+1)!} + o_{\mathbb{P}}(1), \end{split}$$

where the second equality uses $\epsilon_n \to 0$ and the last equality uses $n\epsilon_n^{1+2\mathfrak{q}} \to \infty$. Similarly, for the forward difference estimator, we have

$$\begin{split} \widetilde{\mathcal{D}}_{\mathfrak{q},n}^{\text{FD}}(\mathbf{x}) &= \epsilon_{n}^{-(\mathfrak{q}+1)} \sum_{k=1}^{\mathfrak{q}+1} (-1)^{k+\mathfrak{q}+1} \binom{\mathfrak{q}+1}{k} [\Upsilon_{0}(\mathbf{x}+k\epsilon_{n}) - \Upsilon_{0}(\mathbf{x})] \\ &+ \epsilon_{n}^{-(\mathfrak{q}+1/2)} n^{-1/2} \sum_{k=1}^{\mathfrak{q}+1} (-1)^{k+\mathfrak{q}+1} \binom{\mathfrak{q}+1}{k} \widehat{G}_{\mathbf{x},n}(k;\epsilon_{n}) \\ &- \epsilon_{n}^{-\mathfrak{q}} [\widehat{\theta}_{n}(\mathbf{x}) - \theta_{0}(\mathbf{x})] \sum_{k=1}^{\mathfrak{q}+1} (-1)^{k+\mathfrak{q}+1} \binom{\mathfrak{q}+1}{k} \widehat{R}_{\mathbf{x},n}(k;\epsilon_{n}) \\ &- \epsilon_{n}^{-(\mathfrak{q}+1)} [\widehat{\theta}_{n}(\mathbf{x}) - \theta_{0}(\mathbf{x})] \sum_{k=1}^{\mathfrak{q}+1} (-1)^{k+\mathfrak{q}+1} \binom{\mathfrak{q}+1}{k} [\Phi_{0}(\mathbf{x}+k\epsilon_{n}) - \Phi_{0}(\mathbf{x})] \\ &= \frac{\partial^{\mathfrak{q}} \theta_{0}(\mathbf{x}) \partial \Phi_{0}(\mathbf{x})}{(\mathfrak{q}+1)!} + o(1) + O_{\mathbb{P}} [(n\epsilon_{n}^{1+2\mathfrak{q}})^{-1/2} + (n\epsilon_{n}^{1+2\mathfrak{q}})^{-\mathfrak{q}/(1+2\mathfrak{q})}] \\ &= \frac{\partial^{\mathfrak{q}} \theta_{0}(\mathbf{x}) \partial \Phi_{0}(\mathbf{x})}{(\mathfrak{q}+1)!} + o_{\mathbb{P}}(1). \quad \Box \end{split}$$

SA.1.6. *Proof of Lemma 2*. Proceeding as in the proof of Lemma 1, we have

$$\begin{split} \widetilde{\mathcal{D}}_{j,n}^{\mathrm{BR}}(\mathbf{x}) &= \epsilon_n^{-(j+1)} \sum_{k=1}^{\underline{\mathfrak{s}}+1} \lambda_j^{\mathrm{BR}}(k) [\Upsilon_0(\mathbf{x} + c_k \epsilon_n) - \Upsilon_0(\mathbf{x})] \\ &+ \epsilon_n^{-(j+1/2)} n^{-1/2} \sum_{k=1}^{\underline{\mathfrak{s}}+1} \lambda_j^{\mathrm{BR}}(k) \widehat{G}_{\mathbf{x},n}(c_k; \epsilon_n) \\ &- \epsilon_n^{-j} [\widehat{\theta}_n(\mathbf{x}) - \theta_0(\mathbf{x})] \sum_{k=1}^{\underline{\mathfrak{s}}+1} \lambda_j^{\mathrm{BR}}(k) \widehat{R}_{\mathbf{x},n}(c_k; \epsilon_n) \\ &- \epsilon_n^{-(j+1)} [\widehat{\theta}_n(\mathbf{x}) - \theta_0(\mathbf{x})] \sum_{k=1}^{\underline{\mathfrak{s}}+1} \lambda_j^{\mathrm{BR}}(k) [\Phi_0(\mathbf{x} + c_k \epsilon_n) - \Phi_0(\mathbf{x})] \\ &= \frac{\partial^{j+1} \Upsilon_0(\mathbf{x})}{(j+1)!} + O(\epsilon_n^{\min\{\underline{\mathfrak{s}},\underline{\mathfrak{s}}+1\}-j}) + O_{\mathbb{P}}(\epsilon_n^{-(j+1/2)} n^{-1/2} + \epsilon_n^{-j} a_n^{-\mathfrak{q}}) \\ &= \mathcal{D}_j(\mathbf{x}) + O(\epsilon_n^{\min\{\underline{\mathfrak{s}},\underline{\mathfrak{s}}+1\}-j}) + a_n^{j-\mathfrak{q}} O_{\mathbb{P}}[(a_n \epsilon_n)^{-(j+1/2)} + (a_n \epsilon_n)^{-j}], \end{split}$$

where the second equality uses $\epsilon_n \to 0$ and the defining property of $\{\lambda_j^{\rm BR}(k) : k=1,\ldots,\underline{\mathfrak{s}}\}$. The second part of the lemma follows from the fact that if

$$n\epsilon_n^{(1+2\bar{\mathfrak{q}})\min(\underline{\mathfrak{s}},\mathfrak{s}-1)/(\bar{\mathfrak{q}}-1)}\to 0\quad\text{and}\quad n\epsilon_n^{1+2\bar{\mathfrak{q}}}\to\infty,$$

then

$$a_n^{\mathfrak{q}-j}\epsilon_n^{\min\{\mathfrak{s},\underline{\mathfrak{s}}+1\}-j} \to 0$$
 and $a_n\epsilon_n \to \infty$.

SA.1.7. Higher-order expansion of the bias-reduced estimator . In addition to the assumptions of Lemma 2, suppose that $\widehat{R}_{\mathsf{x},n}(1;\eta_n) = O_{\mathbb{P}}(a_n^{-1/2})$ for $a_n^{-1}\eta_n^{-1} = O(1)$ and that, for some $\delta > 0$, θ_0 is $(\underline{\mathfrak{s}}+1)$ -times continuously differentiable and Φ_0 is $(\underline{\mathfrak{s}}+2)$ -times continuously differentiable on I_{x}^{δ} . Then, the first term in the stochastic expansion of $\widetilde{\mathcal{D}}_{j,n}^{\mathsf{BR}}(\mathsf{x})$ satisfies

$$\begin{split} \epsilon_n^{-(j+1)} \sum_{k=1}^{\underline{\mathfrak{s}}+1} \lambda_j^{\mathrm{BR}}(k) [\Upsilon_0(\mathbf{x} + c_k \epsilon_n) - \Upsilon_0(\mathbf{x})] &- \frac{\partial^{j+1} \Upsilon_0(\mathbf{x})}{(j+1)!} \\ &= \epsilon_n^{\underline{\mathfrak{s}}+1-j} \frac{\partial^{\underline{\mathfrak{s}}+2} \Upsilon_0(\mathbf{x})}{(\underline{\mathfrak{s}}+2)!} \sum_{k=1}^{\underline{\mathfrak{s}}+1} \lambda_j^{\mathrm{BR}}(k) c_k^{\underline{\mathfrak{s}}+2} + o(\epsilon_n^{\underline{\mathfrak{s}}+1-j}) = \epsilon_n^{\underline{\mathfrak{s}}+1-j} \mathsf{B}_j^{\mathrm{BR}}(\mathbf{x}) + o(\epsilon_n^{\underline{\mathfrak{s}}+1-j}). \end{split}$$

Also, the approximate variance of

$$\epsilon_n^{-(j+1/2)} n^{-1/2} \sum_{k=1}^{\underline{\mathfrak{s}}+1} \lambda_j^{\mathtt{BR}}(k) \widehat{G}_{\mathsf{x},n}(c_k;\epsilon_n)$$

is

$$\frac{1}{n\epsilon_n^{1+2j}} \sum_{k=1}^{\underline{\mathfrak{s}}+1} \sum_{l=1}^{\underline{\mathfrak{s}}+1} \lambda_j^{\mathrm{BR}}(k) \lambda_j^{\mathrm{BR}}(l) \mathcal{C}_{\mathsf{x}}(c_k,c_l) = \frac{1}{n\epsilon_n^{1+2j}} \mathsf{V}_j^{\mathrm{BR}}(\mathsf{x}).$$

Finally, the third term in the stochastic expansion of $\widetilde{\mathcal{D}}_{i,n}^{BR}(\mathsf{x})$ is asymptotically negligible under the condition that $\widehat{R}_{x,n}(1;\eta_n) = O_{\mathbb{P}}(a_n^{-1/2})$ for $a_n^{-1}\eta_n^{-1} = O(1)$, while the fourth term exhibits only a higher-order dependence on ϵ_n (relative to the dependence exhibited by the first two terms).

SA.1.8. Proof of Lemma 3. We verify that Assumptions A and E imply Assumptions (B1)-(B4). Define

$$\bar{\Gamma}_n^*(x) = \frac{1}{n} \sum_{i=1}^n W_{i,n} \gamma_0(x; \mathbf{Z}_i) \qquad \text{and} \qquad \bar{\Phi}_n^*(x) = \frac{1}{n} \sum_{i=1}^n W_{i,n} \phi_0(x; \mathbf{Z}_i).$$

Verifying Assumption (B1).

Non-bootstrap weak convergence. We first prove $\widehat{G}_{x,n}^{\mathfrak{q}} \leadsto \mathcal{G}_{x}$. By Assumption (E3)-(E4),

$$\sqrt{na_n} \sup_{|v| < K} \left| \widehat{\Gamma}_n(\mathbf{x} + va_n^{-1}) - \widehat{\Gamma}_n(\mathbf{x}) - \bar{\Gamma}_n(\mathbf{x} + va_n^{-1}) + \bar{\Gamma}_n(\mathbf{x}) \right| = o_{\mathbb{P}}(1),$$

$$\sqrt{na_n} \sup_{|v| \le K} \left| \widehat{\Phi}_n(\mathbf{x} + va_n^{-1}) - \widehat{\Phi}_n(\mathbf{x}) - \bar{\Phi}_n(\mathbf{x} + va_n^{-1}) + \bar{\Phi}_n(\mathbf{x}) \right| = o_{\mathbb{P}}(1),$$

for each K > 0, and thus, defining $\bar{\psi}_{x,n}(v; \mathbf{Z}_i) = \sqrt{a_n} \psi_x(va_n^{-1}; \mathbf{Z}_i)$,

$$\sup_{|v| \le K} \left| \widehat{G}_{\mathsf{x},n}^{\mathfrak{q}}(v) - \frac{1}{\sqrt{n}} \sum_{i=1}^{n} \left\{ \bar{\psi}_{\mathsf{x},n}(v; \mathbf{Z}_i) - \mathbb{E}[\bar{\psi}_{\mathsf{x},n}(v; \mathbf{Z})] \right\} \right| = o_{\mathbb{P}}(1).$$

Next, we prove weak convergence of the empirical process indexed by $\{\bar{\psi}_{x,n}(v;\cdot):|v|\leq K\}$ by verifying finite-dimensional weak convergence and stochastic equicontinuity.

Letting $\eta_n = Ka_n^{-1}$,

etting
$$\eta_n = K a_n^{-1}$$
,
 $n^{-1} \sup_{|v| \le K} \mathbb{E}[|\bar{\psi}_{\mathbf{x},n}(v; \mathbf{Z})|^4] \le (1 + |\theta_0(\mathbf{x})|)^4 n^{-1} a_n^2 \mathbb{E}[\bar{D}_{\gamma}^{\eta_n}(\mathbf{Z})^4 + \bar{D}_{\phi}^{\eta_n}(\mathbf{Z})^4] = o(1).$

Also, convergence of the covariance kernel is imposed in Assumption (E5). Thus, the Lyapunov central limit theorem implies finite-dimensional weak convergence.

For stochastic equicontinuity, proceeding as in Kim and Pollard (1990, Lemma 4.6) and using $a_n \mathbb{E}[\bar{D}_{\gamma}^{\eta_n}(\mathbf{Z})^2 + \bar{D}_{\phi}^{\eta_n}(\mathbf{Z})^2] = O(1)$, it suffices to show that

$$\sup_{\substack{|v-v'|\leq \epsilon_n\\|v|\vee|v'|\leqslant K}} \frac{1}{n} \sum_{i=1}^n |\bar{\psi}_{\mathsf{x},n}(v;\mathbf{Z}_i) - \bar{\psi}_{\mathsf{x},n}(v';\mathbf{Z}_i)|^2 = o_{\mathbb{P}}(1)$$

 $\text{for any } \epsilon_n = o(1). \text{ For any } M > 0, \text{ using } |\bar{\psi}_{\mathsf{x},n}(v;\mathbf{z})| \leq \sqrt{a_n} (1+\theta_0(\mathsf{x})) (\bar{D}_{\gamma}^{\eta_n}(\mathbf{z}) + \bar{D}_{\phi}^{\eta_n}(\mathbf{z}))$

$$\sup_{\substack{|v-v'|\leq\epsilon_n\\|v|\vee|v'|\leq K}}\frac{1}{n}\sum_{i=1}^n|\bar{\psi}_{\mathsf{x},n}(v;\mathbf{Z}_i)-\bar{\psi}_{\mathsf{x},n}(v';\mathbf{Z}_i)|^2$$

$$\leq 4(1+|\theta_0(\mathsf{x})|)^2 \frac{a_n}{n} \sum_{i=1}^n [\bar{D}_{\gamma}^{\eta_n}(\mathbf{Z}_i) + \bar{D}_{\phi}^{\eta_n}(\mathbf{Z}_i)]^2 \mathbb{1}(\bar{D}_{\gamma}^{\eta_n}(\mathbf{Z}_i) + \bar{D}_{\phi}^{\eta_n}(\mathbf{Z}_i) > M)$$

$$+ \bar{M} \sqrt{a_n} \sup_{\substack{|v-v'| \leq \epsilon_n \\ |v| \lor |v'| \leq K}} \mathbb{E} \left[|\bar{\psi}_{\mathsf{x},n}(v; \mathbf{Z}_i) - \bar{\psi}_{\mathsf{x},n}(v'; \mathbf{Z}_i)| \right]$$

$$+ \bar{M} \sqrt{a_n} \sup_{\substack{|v-v'| \leq \epsilon_n \\ |v| \lor |v'| < K}} \frac{1}{n} \sum_{i=1}^n \left\{ |\bar{\psi}_{\mathsf{x},n}(v; \mathbf{Z}_i) - \bar{\psi}_{\mathsf{x},n}(v'; \mathbf{Z}_i)| - \mathbb{E}\left[|\bar{\psi}_{\mathsf{x},n}(v; \mathbf{Z}_i) - \bar{\psi}_{\mathsf{x},n}(v'; \mathbf{Z}_i)| \right] \right\}$$

where $\bar{M} = 2M(1 + |\theta_0(x)|)$, the expectation of the first term after the inequality can be made arbitrarily small by making M large using $a_n \mathbb{E}[\bar{D}_{\gamma}^{\eta_n}(\mathbf{Z})^4 + \bar{D}_{\phi}^{\eta_n}(\mathbf{Z})^4] = O(1)$, the second term is o(1) by Assumption (E5), and the third term is $O_{\mathbb{P}}(\sqrt{a_n/n})$ by Theorem 4.2 of Pollard (1989).

Bootstrap weak convergence. We next prove $\widehat{G}_{x,n}^{\mathfrak{q}} \leadsto_{\mathbb{P}} \mathcal{G}_{x}$. As shown below, we have

$$\sqrt{na_n} \sup_{|v| \leq K} |\widehat{\Gamma}_n^*(\mathbf{x} + va_n^{-1}) - \widehat{\Gamma}_n^*(\mathbf{x}) - \bar{\Gamma}_n^*(\mathbf{x} + va_n^{-1}) + \bar{\Gamma}_n^*(\mathbf{x})| = o_{\mathbb{P}}(1),$$

(SA.12)
$$\sqrt{na_n} \sup_{|v| < K} |\widehat{\Phi}_n^*(\mathbf{x} + va_n^{-1}) - \widehat{\Phi}_n^*(\mathbf{x}) - \bar{\Phi}_n^*(\mathbf{x} + va_n^{-1}) + \bar{\Phi}_n^*(\mathbf{x})| = o_{\mathbb{P}}(1),$$

for each K > 0. Therefore, using $\widehat{\theta}_n(x) \to_{\mathbb{P}} \theta_0(x)$ and the fact that, uniformly over $|v| \le K$,

$$\sqrt{\frac{a_n}{n}} \sum_{i=1}^n W_{i,n} \{ \phi_0(\mathbf{x} + va_n^{-1}; \mathbf{Z}_i) - \phi_0(\mathbf{x}; \mathbf{Z}_i) - \frac{1}{n} \sum_{j=1}^n [\phi_0(\mathbf{x} + va_n^{-1}; \mathbf{Z}_j) - \phi_0(\mathbf{x}; \mathbf{Z}_j)] \} = O_{\mathbb{P}}(1),$$

we obtain

$$\sup_{|v| \le K} \left| \widehat{G}_{\mathsf{x},n}^{\mathsf{q},*}(v) - \frac{1}{\sqrt{n}} \sum_{i=1}^{n} W_{i,n} \widehat{\psi}_{\mathsf{x},n}(v; \mathbf{Z}_i) \right| = o_{\mathbb{P}}(1),$$

where $\widehat{\psi}_{\mathbf{x},n}(v;\mathbf{Z}) = \overline{\psi}_{\mathbf{x},n}(v;\mathbf{Z}) - n^{-1}\sum_{j=1}^n \overline{\psi}_{\mathbf{x},n}(v;\mathbf{Z}_j)$. To prove finite-dimensional weak convergence, we apply Lemma 3.6.15 of van der Vaart and Wellner (1996). Assumption (E2) implies that $n^{-1}\sum_{i=1}^n (W_{i,n}-1)^2 \to_{\mathbb{P}} 1$ and $n^{-1}\max_{1\leq i\leq n}W_{i,n}^2 = 0$ $o_{\mathbb{P}}(1)$. Since

$$\frac{1}{n} \sum_{i=1}^{n} \widehat{\psi}_{\mathbf{x},n}(v; \mathbf{Z}_{i}) \widehat{\psi}_{\mathbf{x},n}(u; \mathbf{Z}_{i}) = \frac{1}{n} \sum_{i=1}^{n} \overline{\psi}_{\mathbf{x},n}(v; \mathbf{Z}_{i}) \overline{\psi}_{\mathbf{x},n}(u; \mathbf{Z}_{i}) \\
- \frac{1}{n} \sum_{i=1}^{n} \overline{\psi}_{\mathbf{x},n}(v; \mathbf{Z}_{i}) \frac{1}{n} \sum_{i=1}^{n} \overline{\psi}_{\mathbf{x},n}(u; \mathbf{Z}_{i})$$

and $\sup_{|v|<\eta} |\psi_{\mathsf{x}}(v;\mathbf{Z})| \leq \bar{D}^{\eta}_{\gamma}(\mathbf{Z}) + |\theta_{0}(\mathsf{x})| \bar{D}^{\eta}_{\phi}(\mathbf{Z})$, for any $v, u \in \mathbb{R}$,

$$\frac{1}{n} \sum_{i=1}^{n} \widehat{\psi}_{\mathsf{x},n}(v; \mathbf{Z}_i) \widehat{\psi}_{\mathsf{x},n}(u; \mathbf{Z}_i) - \mathbb{E}[\bar{\psi}_{\mathsf{x},n}(v; \mathbf{Z}) \bar{\psi}_{\mathsf{x},n}(u; \mathbf{Z})] = o_{\mathbb{P}}(1).$$

Also, $n^{-1}\sum_{i=1}^n \widehat{\psi}_{\mathbf{x},n}^4(v; \mathbf{Z}_i) = O_{\mathbb{P}}(1)$, verifying the hypothesis of the lemma.

For stochastic equicontinuity, let $\epsilon_n = o(1)$ and $\eta_n = Ka_n^{-1}$. Lemma 3.6.7 of van der Vaart and Wellner (1996) implies that for any $n_0 \in \{1, \dots, n\}$, there is a fixed constant C > 0 such that

$$\mathbb{E}\left[\sup_{\substack{|v-v'|\leq\epsilon_{n}\\|v|\vee|v'|\leq K}}\left|\frac{1}{\sqrt{n}}\sum_{i=1}^{n}W_{i,n}\left[\widehat{\psi}_{\mathbf{x},n}(v;\mathbf{Z}_{i})-\widehat{\psi}_{\mathbf{x},n}(v';\mathbf{Z}_{i})\right]\right|\left\{\mathbf{Z}_{i}\right\}_{i=1}^{n}\right]$$

$$\leq C\frac{\sqrt{a_{n}}}{n}\sum_{i=1}^{n}\left[\bar{D}_{\gamma}^{\eta_{n}}(\mathbf{Z}_{i})+\bar{D}_{\phi}^{\eta_{n}}(\mathbf{Z}_{i})\right](n_{0}-1)\frac{1}{\sqrt{n}}\mathbb{E}\max_{1\leq i\leq n}|W_{i,n}|$$

$$+C\max_{n_{0}\leq k\leq n}\mathbb{E}\left[\sup_{\substack{|v-v'|\leq\epsilon_{n}\\|v|\vee|v'|\leq K}}\left|\frac{1}{\sqrt{k}}\sum_{i=n_{0}}^{k}\left[\widehat{\psi}_{\mathbf{x},n}(v;\mathbf{Z}_{R_{i}})-\widehat{\psi}_{\mathbf{x},n}(v';\mathbf{Z}_{R_{i}})\right]\right|\left\{\mathbf{Z}_{i}\right\}_{i=1}^{n}\right]$$

where (R_1,\ldots,R_n) is uniformly distributed on the set of all permutations of $\{1,\ldots,n\}$, independent of $\{\mathbf{Z}_i\}_{i=1}^n$. Choose n_0 such that $n^{1/2-1/\mathfrak{r}}/n_0\to\infty$ and $n_0/a_n\to\infty$ (which is possible by $\mathfrak{r}>(4\mathfrak{q}+2)/(2\mathfrak{q}-1)$), and the first term after the inequality in the above display is $o_{\mathbb{P}}(1)$. For the second term, following the argument of van der Vaart and Wellner (1996, Theorem 3.6.13), it suffices to bound

$$\max_{n_0 \le k \le n} \mathbb{E}_n^* \left[\sup_{\substack{|v-v'| \le \epsilon_n \\ |v| \lor |v'| \le K}} \left| \frac{1}{\sqrt{k}} \sum_{i=1}^k \left[\widehat{\psi}_{\mathsf{x},n}(v; \mathbf{Z}_i^*) - \widehat{\psi}_{\mathsf{x},n}(v'; \mathbf{Z}_i^*) \right] \right| \right],$$

where $\{\mathbf{Z}_i^*\}_{i=1}^k$ denotes a random sample from the empirical cdf and \mathbb{E}_n^* is the expectation under this empirical bootstrap law. Following the argument of Kim and Pollard (1990, Lemma 4.6), it suffices to show that

$$\max_{n_0 \le k \le n} \mathbb{E}_n^* \left[\sup_{\substack{|v-v'| \le \epsilon_n \\ |v| \lor |v'| \le K}} \frac{1}{k} \sum_{i=1}^k \left| \widehat{\psi}_{\mathsf{x},n}(v; \mathbf{Z}_i^*) - \widehat{\psi}_{\mathsf{x},n}(v'; \mathbf{Z}_i^*) \right|^2 \right] = o_{\mathbb{P}}(1).$$

For $k \in \{n_0, ..., n\}$ and M > 0,

$$\mathbb{E}_{n}^{*} \left[\sup_{\substack{|v-v'| \leq \epsilon_{n} \\ |v| \lor |v'| \leq K}} \frac{1}{k} \sum_{i=1}^{k} \left| \widehat{\psi}_{\mathsf{x},n}(v; \mathbf{Z}_{i}^{*}) - \widehat{\psi}_{\mathsf{x},n}(v'; \mathbf{Z}_{i}^{*}) \right|^{2} \right]$$

$$\leq 2\mathbb{E}_{n}^{*} \left[\sup_{\substack{|v-v'| \leq \epsilon_{n} \\ |v| \lor |v'| \leq K}} \frac{1}{k} \sum_{i=1}^{k} \left| \overline{\psi}_{\mathsf{x},n}(v; \mathbf{Z}_{i}^{*}) - \overline{\psi}_{\mathsf{x},n}(v'; \mathbf{Z}_{i}^{*}) \right|^{2} \right]$$

$$+ 2 \sup_{\substack{|v-v'| \leq \epsilon_{n} \\ |v| \lor |v'| \leq K}} \left| \frac{1}{n} \sum_{i=1}^{n} \overline{\psi}_{\mathsf{x},n}(v; \mathbf{Z}_{i}) - \overline{\psi}_{\mathsf{x},n}(v'; \mathbf{Z}_{i}) \right|^{2}$$

where the second term after the inequality is shown to be $o_{\mathbb{P}}(1)$ in the non-bootstrap case above. For the first term after the inequality,

$$\mathbb{E}_{n}^{*} \left[\sup_{\substack{|v-v'| \leq \epsilon_{n} \\ |v| \vee |v'| \leq K}} \frac{1}{k} \sum_{i=1}^{k} \left| \bar{\psi}_{\mathbf{x},n}(v; \mathbf{Z}_{i}^{*}) - \bar{\psi}_{\mathbf{x},n}(v'; \mathbf{Z}_{i}^{*}) \right|^{2} \right] \\
\leq 4(1 + |\theta_{0}(\mathbf{x})|)^{2} \frac{a_{n}}{n} \sum_{i=1}^{n} \left[\bar{D}_{\gamma}^{\eta_{n}}(\mathbf{Z}_{i}) + \bar{D}_{\phi}^{\eta_{n}}(\mathbf{Z}_{i}) \right]^{2} \mathbb{I}(\bar{D}_{\gamma}^{\eta_{n}}(\mathbf{Z}_{i}) + \bar{D}_{\phi}^{\eta_{n}}(\mathbf{Z}_{i}) > M) \\
+ \bar{M} \sqrt{a_{n}} \sup_{\substack{|v-v'| \leq \epsilon_{n} \\ |v| \vee |v'| \leq K}} \frac{1}{n} \sum_{i=1}^{n} \left| \bar{\psi}_{\mathbf{x},n}(v; \mathbf{Z}_{i}) - \bar{\psi}_{\mathbf{x},n}(v'; \mathbf{Z}_{i}) \right| \\
+ \bar{M} \sqrt{a_{n}} \mathbb{E}_{n}^{*} \left[\sup_{\substack{|v-v'| \leq \epsilon_{n} \\ |v| \vee |v'| \leq K}} \frac{1}{k} \sum_{i=1}^{k} \left| \bar{\psi}_{\mathbf{x},n}(v; \mathbf{Z}_{i}^{*}) - \bar{\psi}_{\mathbf{x},n}(v'; \mathbf{Z}_{i}^{*}) \right| - \mathbb{E}_{n}^{*} \left[\left| \bar{\psi}_{\mathbf{x},n}(v; \mathbf{Z}^{*}) - \bar{\psi}_{\mathbf{x},n}(v'; \mathbf{Z}^{*}) \right| \right] \right]$$

where $M = 2M(1 + |\theta_0(x)|)$. The first term after the inequality does not depend on k and its expectation can be made arbitrarily small by taking M sufficiently large. The second term

is independent of k and we can handle this term by adding and subtracting the expectation inside the summation. For the third term, applying Theorem 4.2 of Pollard (1989) again, it is bounded by a constant multiple of

$$\sqrt{\frac{a_n}{k}} \sqrt{\frac{a_n}{n} \sum_{i=1}^n \left[\bar{D}_{\gamma}^{\eta_n}(\mathbf{Z}_i)^2 + \bar{D}_{\phi}^{\eta_n}(\mathbf{Z}_i)^2 \right]} = O_{\mathbb{P}} \left(\sqrt{a_n/k} \right),$$

which is $o_{\mathbb{P}}(1)$ by the choice of n_0 .

Verifying (SA.12). We focus on the first display. By adding and subtracting the bootstrap means,

$$\begin{split} \widehat{\Gamma}_n^*(\mathbf{x} + va_n^{-1}) - \widehat{\Gamma}_n^*(\mathbf{x}) - \bar{\Gamma}_n^*(\mathbf{x} + va_n^{-1}) + \bar{\Gamma}_n^*(\mathbf{x}) \\ &= \frac{1}{n} \sum_{i=1}^n W_{i,n} \check{\gamma}_n(v; \mathbf{Z}) + \left[\check{\Gamma}_n(\mathbf{x} + va_n^{-1}) - \check{\Gamma}_n(\mathbf{x}) - \bar{\Gamma}_n(\mathbf{x} + va_n^{-1}) + \bar{\Gamma}_n(\mathbf{x}) \right] \end{split}$$

where

$$\check{\gamma}_n(v; \mathbf{Z}) = \widehat{\gamma}_n(\mathsf{x} + va_n^{-1}; \mathbf{Z}_i) - \widehat{\gamma}_n(\mathsf{x}; \mathbf{Z}_i) - \gamma_0(\mathsf{x} + va_n^{-1}; \mathbf{Z}_i) + \gamma_0(\mathsf{x}; \mathbf{Z}_i)
- \check{\Gamma}_n(\mathsf{x} + va_n^{-1}) + \check{\Gamma}_n(\mathsf{x}) - \bar{\Gamma}_n(\mathsf{x} + va_n^{-1}) + \bar{\Gamma}_n(\mathsf{x}).$$

By Assumption (E3),

$$\sqrt{na_n} \sup_{|v| \le K} |\check{\Gamma}_n(\mathsf{x} + va_n^{-1}) - \check{\Gamma}_n(\mathsf{x}) - \bar{\Gamma}_n(\mathsf{x} + va_n^{-1}) + \bar{\Gamma}_n(\mathsf{x})| = o_{\mathbb{P}}(1).$$

Identical to above, Lemma 3.6.7 and the argument in Theorem 3.6.13 of van der Vaart and Wellner (1996) imply that for some fixed C > 0,

$$\sqrt{na_n} \mathbb{E} \left[\sup_{|v| \le K} \left| \frac{1}{n} \sum_{i=1}^n W_{i,n} \breve{\gamma}_n(v; \mathbf{Z}_i) \right| \right] \left\{ \mathbf{Z}_i \right\}_{i=1}^n \right] \\
\le C \frac{\sqrt{a_n}}{n} \sum_{i=1}^n \sup_{|v| \le K} \left| \breve{\gamma}_n(v; \mathbf{Z}_i) \right| \frac{n_0 n^{\mathfrak{r}}}{\sqrt{n}} + C \sqrt{a_n} \max_{n_0 \le k \le n} \mathbb{E}_n^* \left[\sup_{|v| \le K} \left| \frac{1}{\sqrt{k}} \sum_{i=1}^k \breve{\gamma}_n(v; \mathbf{Z}_i^*) \right| \right].$$

By Assumption (E3),

$$\frac{\sqrt{a_n}}{n} \sum_{i=1}^n \sup_{|v| \le K} |\check{\gamma}_n(v; \mathbf{Z}_i)|$$

$$\le \frac{\sqrt{a_n}}{n} \sum_{i=1}^n \sup_{|v| \le K} |\widehat{\gamma}_n(\mathsf{x} + va_n^{-1}; \mathbf{Z}_i) - \widehat{\gamma}_n(\mathsf{x}; \mathbf{Z}_i) - \gamma_0(\mathsf{x} + va_n^{-1}; \mathbf{Z}_i) + \gamma_0(\mathsf{x}; \mathbf{Z}_i)|$$

$$+ \sqrt{a_n} \sup_{|v| \le K} |\check{\Gamma}_n(\mathsf{x} + va_n^{-1}) - \check{\Gamma}_n(\mathsf{x}) - \bar{\Gamma}_n(\mathsf{x} + va_n^{-1}) + \bar{\Gamma}_n(\mathsf{x})| = o_{\mathbb{P}}(1).$$

Also, Corollary 4.3 of Pollard (1989) implies that for some fixed C > 0,

$$\mathbb{E}_{n}^{*} \left[\sup_{|v| \leq K} \left| \frac{1}{\sqrt{k}} \sum_{i=1}^{k} \check{\gamma}_{n}(v; \mathbf{Z}_{i}^{*}) \right| \right]^{2}$$

$$\leq C \frac{1}{n} \sum_{i=1}^{n} \sup_{|v| \leq K} \left| \widehat{\gamma}_{n}(\mathsf{x} + va_{n}^{-1}; \mathbf{Z}_{i}) - \widehat{\gamma}_{n}(\mathsf{x}; \mathbf{Z}_{i}) - \gamma_{0}(\mathsf{x} + va_{n}^{-1}; \mathbf{Z}_{i}) + \gamma_{0}(\mathsf{x}; \mathbf{Z}_{i}) \right|^{2},$$

which is $o_{\mathbb{P}}(a_n^{-1})$ by Assumption (E3).

Verifying Assumption (B2). Defining

$$\begin{split} \bar{G}_{\mathbf{x},n}^{\mathbf{q}}(v) &= \sqrt{na_n} \left[\bar{\Gamma}_n(\mathbf{x} + va_n^{-1}) - \bar{\Gamma}_n(\mathbf{x}) - \Gamma_0(\mathbf{x} + va_n^{-1}) + \Gamma_0(\mathbf{x}) \right] \\ &- \theta_0(\mathbf{x}) \sqrt{na_n} \left[\bar{\Phi}_n(\mathbf{x} + va_n^{-1}) - \bar{\Phi}_n(\mathbf{x}) - \Phi_0(\mathbf{x} + va_n^{-1}) + \Phi_0(\mathbf{x}) \right] \end{split}$$

we have

$$\begin{split} \widehat{G}_{\mathbf{x},n}^{\mathfrak{q}}(v) - \bar{G}_{\mathbf{x},n}^{\mathfrak{q}}(v) &= \sqrt{na_n} \left[\widehat{\Gamma}_n(\mathbf{x} + va_n^{-1}) - \widehat{\Gamma}_n(\mathbf{x}) - \bar{\Gamma}_n(\mathbf{x} + va_n^{-1}) + \bar{\Gamma}_n(\mathbf{x}) \right] \\ &- \theta_0(\mathbf{x}) \sqrt{na_n} \left[\widehat{\Phi}_n(\mathbf{x} + va_n^{-1}) - \widehat{\Phi}_n(\mathbf{x}) - \bar{\Phi}_n(\mathbf{x} + va_n^{-1}) + \bar{\Phi}_n(\mathbf{x}) \right]. \end{split}$$

Assumptions (E3) and (E4) imply that for $V \in [1, a_n \delta]$,

$$\sqrt{na_n} \sup_{|v| \in [V,2V]} |\widehat{\Gamma}_n(\mathbf{x} + va_n^{-1}) - \widehat{\Gamma}_n(\mathbf{x}) - \bar{\Gamma}_n(\mathbf{x} + va_n^{-1}) + \bar{\Gamma}_n(\mathbf{x})| \le (2V)^\beta a_n^{-\beta} B_n + A_n$$

and

$$\sqrt{na_n} \sup_{|v| \in [V, 2V]} |\widehat{\Phi}_n(\mathbf{x} + va_n^{-1}) - \widehat{\Phi}_n(\mathbf{x}) - \bar{\Phi}_n(\mathbf{x} + va_n^{-1}) + \bar{\Phi}_n(\mathbf{x})| \le (2V)^{\beta} a_n^{-\beta} B_n + A_n,$$

where $\beta = \beta_{\gamma} \vee \beta_{\phi}$, $A_n = A_{\gamma,n} \vee A_{\phi,n} = o_{\mathbb{P}}(1)$, and $B_n = B_{\gamma,n} \vee B_{\phi,n} = o_{\mathbb{P}}(a_n^{\beta})$. As a consequence, there exists $\eta'_n = o(1)$ such that

$$\lim_{n\to\infty} \mathbb{P}\left[\bigcap_{V\in[1,a_n\delta]}\left\{\sup_{|v|\in[V,2V]}|\widehat{G}_{\mathsf{x},n}^{\mathfrak{q}}(v)-\bar{G}_{\mathsf{x},n}^{\mathfrak{q}}(v)|\leq [V^\beta+1]\eta_n'\right\}\right]=1;$$

we take the event in the display to be A_n . Also,

$$\limsup_{\eta\downarrow 0} \mathbb{E}[\bar{D}_{\gamma}^{\eta}(\mathbf{Z})^2/\eta] + \mathbb{E}[\bar{D}_{\phi}^{\eta}(\mathbf{Z})^2/\eta] < \infty$$

and, using Corollary 4.3 of Pollard (1989),

$$\mathbb{E}\left[\sup_{|v| \le V} |\bar{G}_{\mathbf{x},n}^{\mathfrak{q}}(v)|\right] \le C\sqrt{V} \quad \text{for } V \in (0, a_n \delta].$$

Therefore,

$$\mathbb{E}\left[\sup_{|v|\in[V,2V]}|\widehat{G}_{\mathsf{x},n}^{\mathsf{q}}(v)|\mathbb{1}_{\mathcal{A}_n}\right] \leq C\sqrt{V} + [V^{\beta} + 1]\eta_n',$$

implying in particular that

$$\sup_{V\in[1,a_n\delta]}\mathbb{E}\left[V^{-\beta}\sup_{|v|\in[V,2V]}|\widehat{G}_{\mathbf{x},n}^{\mathfrak{q}}(v)|\mathbbm{1}_{\mathcal{A}_n}\right]=O(1).$$

For the bootstrap counterpart, defining

$$\begin{split} \bar{G}_{\mathbf{x},n}^{\mathfrak{q},*}(v) &= \sqrt{na_n} \left[\bar{\Gamma}_n^*(\mathbf{x} + va_n^{-1}) - \bar{\Gamma}_n^*(\mathbf{x}) - \bar{\Gamma}_n(\mathbf{x} + va_n^{-1}) + \bar{\Gamma}_n(\mathbf{x}) \right] \\ &- \theta_0(\mathbf{x}) \sqrt{na_n} \left[\bar{\Phi}_n^*(\mathbf{x} + va_n^{-1}) - \bar{\Phi}_n^*(\mathbf{x}) - \bar{\Phi}_n(\mathbf{x} + va_n^{-1}) + \bar{\Phi}_n(\mathbf{x}) \right], \end{split}$$

we have

$$\begin{split} &(\text{SA.13}) \quad \widehat{G}_{\mathbf{x},n}^{\mathfrak{q},*}(v) - \bar{G}_{\mathbf{x},n}^{\mathfrak{q},*}(v) \\ &= \sqrt{na_n} \left[\widehat{\Gamma}_n^*(\mathbf{x} + va_n^{-1}) - \widehat{\Gamma}_n^*(\mathbf{x}) - \bar{\Gamma}_n^*(\mathbf{x} + va_n^{-1}) + \bar{\Gamma}_n^*(\mathbf{x}) \right] \\ &- \widehat{\theta}_n(\mathbf{x}) \sqrt{na_n} \left[\widehat{\Phi}_n^*(\mathbf{x} + va_n^{-1}) - \widehat{\Phi}_n^*(\mathbf{x}) - \bar{\Phi}_n^*(\mathbf{x} + va_n^{-1}) + \bar{\Phi}_n^*(\mathbf{x}) \right] \\ &+ \sqrt{na_n} \left[\bar{\Gamma}_n(\mathbf{x} + va_n^{-1}) - \bar{\Gamma}_n(\mathbf{x}) - \widehat{\Gamma}_n(\mathbf{x} + va_n^{-1}) + \widehat{\Gamma}_n(\mathbf{x}) \right] \\ &- \widehat{\theta}_n(\mathbf{x}) \sqrt{na_n} \left[\bar{\Phi}_n(\mathbf{x} + va_n^{-1}) - \bar{\Phi}_n(\mathbf{x}) - \widehat{\Phi}_n(\mathbf{x} + va_n^{-1}) + \widehat{\Phi}_n(\mathbf{x}) \right] \\ &+ \sqrt{a_n} [\widehat{\theta}_n(\mathbf{x}) - \theta_0(\mathbf{x})] \sqrt{n} \left[\bar{\Phi}_n^*(\mathbf{x} + va_n^{-1}) - \bar{\Phi}_n^*(\mathbf{x}) - \bar{\Phi}_n(\mathbf{x} + va_n^{-1}) + \bar{\Phi}_n(\mathbf{x}) \right], \end{split}$$

where bounds for the third and fourth terms were derived in the non-bootstrap case and where the last term is $o_{\mathbb{P}}(1)$ uniformly over $|v| \leq a_n \delta$ as $\sqrt{a_n} [\widehat{\theta}_n(\mathsf{x}) - \theta_0(\mathsf{x})] = o_{\mathbb{P}}(1)$ and $\sqrt{n} \sup_{|v| \leq \delta} |\bar{\Phi}_n^*(\mathsf{x}+v) - \bar{\Phi}_n(\mathsf{x}+v)| = O_{\mathbb{P}}(1)$. For the first term after the equality in (SA.13),

$$\begin{split} &\sqrt{na_n} \left[\widehat{\Gamma}_n^* (\mathbf{x} + va_n^{-1}) - \widehat{\Gamma}_n^* (\mathbf{x}) - \bar{\Gamma}_n^* (\mathbf{x} + va_n^{-1}) + \bar{\Gamma}_n^* (\mathbf{x}) \right] \\ &= \frac{1}{\sqrt{n}} \sum_{i=1}^n W_{i,n} \widecheck{\gamma}_n (va_n^{-1}; \mathbf{Z}_i) + \sqrt{na_n} \left[\widecheck{\Gamma}_n (\mathbf{x} + va_n^{-1}) - \widecheck{\Gamma}_n (\mathbf{x}) - \bar{\Gamma}_n (\mathbf{x} + va_n^{-1}) + \bar{\Gamma}_n (\mathbf{x}) \right] \end{split}$$

where, by Assumption (E3),

$$\sqrt{na_n} \sup_{|v| \le V} |\check{\Gamma}_n(\mathsf{x} + va_n^{-1}) - \check{\Gamma}_n(\mathsf{x}) - \bar{\Gamma}_n(\mathsf{x} + va_n^{-1}) + \bar{\Gamma}_n(\mathsf{x})| = [1 + V^{\beta}]o_{\mathbb{P}}(1)$$

uniformly over $V \in [1, a_n \delta]$ and where, applying Lemma 3.6.7 of van der Vaart and Wellner (1996), we have, for any $n_0 \in \{1, ..., n\}$,

$$\mathbb{E}\left[\sup_{|v| \le \delta} \left| \frac{1}{\sqrt{n}} \sum_{i=1}^{n} W_{i,n} \breve{\gamma}_n(v; \mathbf{Z}_i) \right| \middle| \{\mathbf{Z}_i\}_{i=1}^n \right]$$

$$\leq C \frac{1}{n} \sum_{i=1}^{n} \sup_{|v| \leq \delta} |\check{\gamma}_n(v; \mathbf{Z}_i)| \frac{n_0}{\sqrt{n}} \mathbb{E} \left[\max_{1 \leq i \leq n} |W_{i,n}| \right] + C \max_{n_0 \leq k \leq n} \mathbb{E}_n^* \left[\sup_{|v| \leq \delta} \left| \frac{1}{\sqrt{k}} \sum_{i=1}^k \check{\gamma}_n(v; \mathbf{Z}_i^*) \right| \right],$$

where C > 0 is a fixed constant. Letting n_0 be a diverging sequence (dependent on n) such that $n_0 n^{\mathfrak{r}} \sqrt{a_n/n} = o(1)$, the first term on the majorant side is bounded by

$$C\frac{1}{n}\sum_{i=1}^{n}\sup_{|x-\mathbf{x}|\leq\delta}|\widehat{\gamma}_n(x;\mathbf{Z}_i)-\gamma_0(x;\mathbf{Z}_i)|+C\sup_{|v|\leq\delta}|\check{\Gamma}_n(\mathbf{x}+v)-\bar{\Gamma}_n(\mathbf{x}+v)|=o_{\mathbb{P}}(1),$$

and, by Corollary 4.3 of Pollard (1989), the second term is bounded (up to a constant) by

$$\sqrt{a_n} \left| \frac{1}{n} \sum_{i=1}^n \sup_{x \in I_s^{\delta}} |\widehat{\gamma}_n(x; \mathbf{Z}_i) - \gamma_0(x; \mathbf{Z}_i)|^2 \right|^{1/2}$$

which is $o_{\mathbb{P}}(1)$ by Assumption (E3). As a consequence, there exists a sequence of random variables $A'_n = o_{\mathbb{P}}(1)$ such that for $V \in [1, a_n \delta]$,

$$\sqrt{na_n} \sup_{|v| \le [V,2V]} \left| \widehat{\Gamma}_n^*(\mathbf{x} + va_n^{-1}) - \widehat{\Gamma}_n^*(\mathbf{x}) - \bar{\Gamma}_n^*(\mathbf{x} + va_n^{-1}) + \bar{\Gamma}_n^*(\mathbf{x}) \right| \le [V^{\beta} + 1]A_n'.$$

By identical arguments, an analogous bound holds for the second term after the inequality in (SA.13). Therefore, there exists $\eta'_n = o(1)$ and events \mathcal{A}_n such that $\lim_{n \to \infty} \mathbb{P}[\mathcal{A}_n] = 1$ and

$$\mathbb{E}\left[\sup_{|v|\in[V,2V]}\left|\widehat{G}_{\mathsf{x},n}^{\mathsf{q},*}(v) - \bar{G}_{\mathsf{x},n}^{\mathsf{q},*}(v)\right|\mathbb{1}_{\mathcal{A}_n}\right] \leq [V^{\beta} + 1]\eta_n'$$

for any $V \in [1, a_n \delta]$. Finally, proceeding as in the proof of stochastic equicontinuity, we obtain the bound $\mathbb{E}[\sup_{|v| \in [V, 2V]} |\bar{G}_{\mathbf{x}, n}^{\mathbf{q}, *}(v)|] \leq C\sqrt{V}$.

Verifying Assumptions (B3)-(B4). We have

$$\sup_{x \in I} \left| \widehat{\Gamma}_n(x) - \Gamma_0(x) \right| \le \sup_{x \in I} \left| \widehat{\Gamma}_n(x) - \overline{\Gamma}_n(x) \right| + \sup_{x \in I} \left| \overline{\Gamma}_n(x) - \Gamma_0(x) \right|,$$

where the first term after the inequality is assumed to be $o_{\mathbb{P}}(1)$ and the second term is $o_{\mathbb{P}}(1)$ by standard arguments. Similarly, $\sup_{x\in I}|\widehat{\Phi}_n(x)-\Phi_0(x)|=o_{\mathbb{P}}(1)$. Also,

$$\begin{split} \sup_{x \in I_{\mathsf{x}}^{\delta}} \left| \widehat{\Phi}_{n}(x) - \Phi_{0}(x) \right| &\leq \sup_{x \in I_{\mathsf{x}}^{\delta}} \left| \widehat{\Phi}_{n}(x) - \widehat{\Phi}_{n}(\mathsf{x}) - \bar{\Phi}_{n}(x) + \bar{\Phi}_{n}(\mathsf{x}) \right| \\ &+ \sup_{x \in I_{\mathsf{x}}^{\delta}} \left| \bar{\Phi}_{n}(x) - \Phi_{0}(x) \right| + \left| \widehat{\Phi}_{n}(\mathsf{x}) - \bar{\Phi}_{n}(\mathsf{x}) \right| \end{split}$$

where the last two terms are $o_{\mathbb{P}}(a_n^{-1})$ and where Assumption (E4) implies

$$\sup_{\mathbf{x}\in I_{\diamond}^{\delta}}\left|\widehat{\Phi}_{n}(\mathbf{x})-\widehat{\Phi}_{n}(\mathbf{x})-\bar{\Phi}_{n}(\mathbf{x})+\bar{\Phi}_{n}(\mathbf{x})\right|\leq (na_{n})^{-1/2}[A_{\phi,n}+\delta^{\beta_{\phi}}B_{\phi,n}]=o_{\mathbb{P}}(a_{n}^{-1}),$$

the last equality using $B_{\phi,n} = o_{\mathbb{P}}(a_n^{\beta_{\phi}})$ and $\beta_{\phi} \leq \mathfrak{q}$.

Now we look at the bootstrap objects. For $\widehat{\Gamma}_n^*$,

$$\sup_{x \in I} \left| \widehat{\Gamma}_n^*(x) - \widehat{\Gamma}_n(x) \right| \leq \sup_{x \in I} \left| \widehat{\Gamma}_n^*(x) - \bar{\Gamma}_n^*(x) \right| + \sup_{x \in I} \left| \bar{\Gamma}_n^*(x) - \bar{\Gamma}_n(x) \right| + \sup_{x \in I} \left| \bar{\Gamma}_n(x) - \widehat{\Gamma}_n(x) \right|$$

where the last term is assumed to be $o_{\mathbb{P}}(1)$ and where

$$\sup_{x \in I} \left| \widehat{\Gamma}_{n}^{*}(x) - \overline{\Gamma}_{n}^{*}(x) \right| \leq \frac{1}{n} \sum_{i=1}^{n} |W_{i,n}| \sup_{x \in I} |\widehat{\gamma}_{n}(x; \mathbf{Z}_{i}) - \gamma_{0}(x; \mathbf{Z}_{i})|$$

$$\leq \sqrt{\frac{1}{n} \sum_{i=1}^{n} |W_{i,n}|^{2} \frac{1}{n} \sum_{i=1}^{n} \sup_{x \in I} |\widehat{\gamma}_{n}(x; \mathbf{Z}_{i}) - \gamma_{0}(x; \mathbf{Z}_{i})|^{2}} = o_{\mathbb{P}}(1).$$

For $\bar{\Gamma}_n^* - \bar{\Gamma}_n$, using Lemma 3.6.7 of van der Vaart and Wellner (1996) and the same argument as for verifying Assumption (B1), it suffices to show that

$$n^{-1/2} \max_{\lfloor \sqrt{n} \rfloor \le k \le n} \mathbb{E}_n^* \left[\sup_{x \in I} \left| \frac{1}{\sqrt{k}} \sum_{i=1}^k \bar{\gamma}_n(x; \mathbf{Z}_i^*) \right| \right] = o_{\mathbb{P}}(1),$$

as can be done using Corollary 4.3 of Pollard (1989).

For $\widehat{\Phi}_n^*$, $\sup_{x\in I} |\widehat{\Phi}_n^*(x) - \widehat{\Phi}_n(x)| = o_{\mathbb{P}}(1)$ follows from the same argument as for $\widehat{\Gamma}_n^*$. For $a_n \sup_{x\in I_\delta^*} |\widehat{\Phi}_n^*(x) - \widehat{\Phi}_n(x)| = o_{\mathbb{P}}(1)$,

$$\begin{split} \sup_{x \in I_{\mathbf{x}}^{\delta}} \left| \widehat{\Phi}_{n}^{*}(x) - \widehat{\Phi}_{n}(x) \right| &\leq \sup_{x \in I_{\mathbf{x}}^{\delta}} \left| \widehat{\Phi}_{n}^{*}(x) - \bar{\Phi}_{n}^{*}(x) \right| + \sup_{x \in I_{\mathbf{x}}^{\delta}} \left| \bar{\Phi}_{n}^{*}(x) - \bar{\Phi}_{n}(x) \right| \\ &+ \sup_{x \in I_{\mathbf{x}}^{\delta}} \left| \bar{\Phi}_{n}(x) - \Phi_{0}(x) \right| + \sup_{x \in I_{\mathbf{x}}^{\delta}} \left| \widehat{\Phi}_{n}(x) - \Phi_{0}(x) \right| \end{split}$$

where the last term is $o_{\mathbb{P}}(a_n^{-1})$ as shown above and the second and third terms after the inequality are $O_{\mathbb{P}}(n^{-1/2})$ by standard arguments. For the remaining term,

$$\sup_{x \in I_{\mathsf{x}}^{\delta}} \left| \widehat{\Phi}_{n}^{*}(x) - \bar{\Phi}_{n}^{*}(x) \right| \leq \sup_{x \in I_{\mathsf{x}}^{\delta}} \left| \frac{1}{n} \sum_{i=1}^{n} W_{i,n} \check{\phi}_{n}(x; \mathbf{Z}_{i}) \right| + \left| \check{\Phi}_{n}(\mathsf{x}) - \bar{\Phi}_{n}(\mathsf{x}) \right|$$
$$+ \sup_{x \in I_{\mathsf{x}}^{\delta}} \left| \check{\Phi}_{n}(x) - \check{\Phi}_{n}(\mathsf{x}) - \bar{\Phi}_{n}(\mathsf{x}) + \bar{\Phi}_{n}(\mathsf{x}) \right|$$

where $\check{\phi}_n(x;\mathbf{Z}) = \widehat{\phi}_n(x;\mathbf{Z}) - \phi_0(x;\mathbf{Z}) - [\check{\Phi}_n(x) - \bar{\Phi}_n(x)]$. The last two terms are $o_{\mathbb{P}}(a_n^{-1})$ by Assumption (E4). Using Lemma 3.6.7 of van der Vaart and Wellner (1996) and the argument similar to above, the remaining term is $o_{\mathbb{P}}(a_n^{-1})$.

SA.1.9. *Proof of Theorem A.1*. The proof is by contradiction and follows Kosorok (2008). We omit some details in cases where the arguments are almost identical to those for Theorem 1 and Lemma 3.

Suppose that the bootstrap approximation is consistent; that is, suppose

$$r_n\left(\widehat{\theta}_n^*(\mathbf{x}) - \widehat{\theta}_n(\mathbf{x})\right) \leadsto_{\mathbb{P}} \frac{1}{\partial \Phi_0(\mathbf{x})} \partial_-\mathsf{GCM}_{\mathbb{R}}\{\mathcal{G}_{\mathbf{x}} + \mathcal{M}_{\mathbf{x}}^{\mathfrak{q}}\}(0) = Y.$$

Then, by Theorem 2.2 of Kosorok (2008), we have

(SA.14)
$$r_n\left(\widehat{\theta}_n^*(\mathsf{x}) - \theta_0(\mathsf{x})\right) \rightsquigarrow Y_1 + Y_2 =_d \sqrt{2}Y$$

where $=_d$ denotes the distributional equality, Y_1 and Y_2 are independent copies of Y, and where the convergence in distribution is unconditional.

Using the switching lemma, $\mathbb{P}\left[r_n\left(\widehat{\theta}_n^*(\mathbf{x}) - \theta_0(\mathbf{x})\right) > t\right]$ equals

$$\mathbb{P}\left[\max\underset{x\in\widehat{\Phi}_n^{*^-}([0,\widehat{u}_n^*])}{\operatorname{argmax}}\left\{[\theta_0(\mathbf{x})+r_n^{-1}t]\widehat{\Phi}_n^*(x)-\mathsf{LSC}_{\widehat{\Phi}_n^*}\left(\widehat{\Gamma}_n^*\right)(x)\right\}<\widehat{\Phi}_n^{*^-}\circ\widehat{\Phi}_n^*(\mathbf{x})\right].$$

By the arguments used in the proof of Theorem 1, to characterize the limiting distribution of $r_n(\widehat{\theta}_n^*(x) - \theta_0(x))$, it suffices to look at

$$-\check{G}_{\mathbf{x},n}^{\mathfrak{q},*}(v) - \check{M}_{\mathbf{x},n}^{\mathfrak{q}}(v) + t\widehat{L}_{\mathbf{x},n}^{\mathfrak{q},*}(v)$$

where

$$\check{G}_{\mathbf{x},n}^{\mathfrak{q},*}(v) = \sqrt{na_n} \left[\widehat{\Gamma}_n^*(\mathbf{x} + va_n^{-1}) - \widehat{\Gamma}_n^*(\mathbf{x}) - \check{\Gamma}_n(\mathbf{x} + va_n^{-1}) + \check{\Gamma}_n(\mathbf{x}) \right] \\
- \theta_0(\mathbf{x}) \sqrt{na_n} \left[\widehat{\Phi}_n^*(\mathbf{x} + va_n^{-1}) - \widehat{\Phi}_n^*(\mathbf{x}) - \check{\Phi}_n(\mathbf{x} + va_n^{-1}) + \check{\Phi}_n(\mathbf{x}) \right]$$

and

$$\check{M}_{\mathbf{x},n}^{\mathbf{q}}(v) = \sqrt{na_n} \left[\check{\Gamma}_n(\mathbf{x} + va_n^{-1}) - \check{\Gamma}_n(\mathbf{x}) - \theta_0(\mathbf{x}) \{ \check{\Phi}_n(\mathbf{x} + va_n^{-1}) - \check{\Phi}_n(\mathbf{x}) \} \right].$$

It can be shown that $\check{G}_{\mathsf{x},n}^{\mathfrak{q},*} \leadsto_{\mathbb{P}} \mathcal{G}_{\mathsf{x}}, \widehat{L}_{\mathsf{x},n}^{\mathfrak{q},*} \leadsto \mathcal{L}_{\mathsf{x}}$, and that $\check{M}_{\mathsf{x},n}^{\mathfrak{q}} \leadsto \mathcal{G}_{\mathsf{x}} + \mathcal{M}_{\mathsf{x}}^{\mathfrak{q}}$. Thus,

$$\mathbb{P}\left[r_n\left(\widehat{\theta}_n^*(\mathbf{x}) - \theta_0(\mathbf{x})\right) > t\right] \to \mathbb{P}\left[\underset{v \in \mathbb{R}}{\operatorname{argmin}}\left\{\mathcal{G}_{\mathbf{x},1}(v) + \mathcal{G}_{\mathbf{x},2}(v) + \mathcal{M}_{\mathbf{x}}^{\mathfrak{q}}(v) - t\partial\Phi_0(\mathbf{x})v\right\} < 0\right],$$

where $\mathcal{G}_{x,1}$ and $\mathcal{G}_{x,2}$ are independent copies of \mathcal{G}_x . Noting that $\mathcal{G}_x(av) =_d \sqrt{|a|} \mathcal{G}_x(v)$ and using the change of variable $v = u2^{\frac{1}{2q+1}}$, the limit distribution equals

$$\begin{split} & \mathbb{P}\left[2^{\frac{1}{2\mathfrak{q}+1}} \operatorname*{argmin}_{u \in \mathbb{R}} \left\{ \mathcal{G}_{\mathsf{X}}(u) + \mathcal{M}_{\mathsf{X}}^{\mathfrak{q}}(u) - 2^{-\frac{\mathfrak{q}}{2\mathfrak{q}+1}} t \partial \Phi_{0}(\mathsf{x}) u \right\} < 0 \right] \\ & = \mathbb{P}\left[2^{\frac{\mathfrak{q}}{2\mathfrak{q}+1}} (\partial \Phi_{0}(\mathsf{x}))^{-1} \partial_{-} \mathsf{GCM}_{\mathbb{R}} \{ \mathcal{G}_{\mathsf{X}} + \mathcal{M}_{\mathsf{X}}^{\mathfrak{q}} \}(0) > t \right]. \end{split}$$

As a consequence,

$$r_n\left(\widehat{\theta}_n^*(\mathbf{x}) - \theta_0(\mathbf{x})\right) \leadsto 2^{\frac{\mathfrak{q}}{2\mathfrak{q}+1}}(\partial \Phi_0(\mathbf{x}))^{-1}\partial_-\mathsf{GCM}_{\mathbb{R}}\{\mathcal{G}_{\mathbf{x}} + \mathcal{M}_{\mathbf{x}}^{\mathfrak{q}}\}(0),$$

contradicting (SA.14) because $2^{\frac{q}{2q+1}} \neq \sqrt{2}$.

In other words, the bootstrap estimator $\widehat{\theta}_n^*(x)$ fails to approximate the limit distribution. \square

SA.1.10. *Remarks on verifying conditions in applications*. Below we verify the hypothesis of Theorem 1 for various examples. For this purpose, it suffices to verify Assumptions A, (B5)-(B7), and E since Assumption E implies (B1)-(B4) by Lemma 3.

When γ_0 is known, it is natural to take $\widehat{\Gamma}_n = \check{\Gamma}_n = \bar{\Gamma}_n$, in which case (E3) reduces to the requirement that, for some $\rho_{\gamma} \in (0,2)$, (SA.15)

$$\limsup_{\varepsilon \downarrow 0} \frac{\log N_U(\varepsilon, \mathfrak{F}_{\gamma})}{\varepsilon^{-\rho_{\gamma}}} < \infty, \quad \mathbb{E}[\bar{F}_{\gamma}(\mathbf{Z})^2] < \infty, \quad \limsup_{\eta \downarrow 0} \frac{\mathbb{E}[\bar{D}_{\gamma}^{\eta}(\mathbf{Z})^2 + \bar{D}_{\gamma}^{\eta}(\mathbf{Z})^4]}{\eta} < \infty.$$

An identical remark applies to ϕ_0 and (E4).

SA.1.11. *Proof of Corollary 1*. Assumption A and (E1)-(E2) follow from the hypothesis. (E3): In this example, $\gamma_0(x; \mathbf{Z}) = \mathbb{1}(X \leq x)$ is known, so it suffices to verify (SA.15). The uniform covering number of $\{\mathbb{1}(\cdot \leq x) : x \in \mathbb{R}\}$ grows linearly, and an envelope function can be taken to be 1. For an envelope function of $\{\mathbb{1}(\cdot \leq x) : |x - x| \leq \eta\}$, we can take $\mathbb{1}(-\eta + x \leq \cdot \leq x + \eta)$ and the moment bound is satisfied as $\mathbb{E}[\mathbb{1}(-\eta + x \leq X \leq x + \eta)] \leq C\eta$.

(E4) trivially holds as $\widehat{\Phi}_n(x) = \widehat{\Phi}_n^*(x) = x$.

(E5): Here $\psi_{\mathsf{x}}(v; \mathbf{Z}) = \mathbb{1}(X \le \mathsf{x} + v) - \mathbb{1}(X \le \mathsf{x}) - \Phi_0(\mathsf{x})v$. Then,

$$\frac{\mathbb{E}[|\psi_{\mathsf{x}}(v;\mathbf{Z}) - \psi_{\mathsf{x}}(v';\mathbf{Z})|]}{|v - v'|} \le \frac{\mathbb{E}[\mathbb{1}(\mathsf{x} + v \wedge v' < X \le \mathsf{x} + v \vee v')]}{|v - v'|} + f_0(\mathsf{x}) \le C.$$

Also,
$$\psi_{\mathsf{x}_n}(s\eta_n; \mathbf{Z}) = \mathbb{1}(\mathsf{x}_n \wedge (\mathsf{x}_n + s\eta_n) < X \le \mathsf{x}_n \vee (\mathsf{x}_n + s\eta_n)) - f_0(\mathsf{x})s\eta_n$$
 and

$$\psi_{\mathsf{x}_{n}}(s\eta_{n}; \mathbf{Z})\psi_{\mathsf{x}_{n}}(t\eta_{n}; \mathbf{Z}) = \mathbb{1}(\mathsf{x}_{n} < X \le \mathsf{x}_{n} + \eta_{n}(s \wedge t))\mathbb{1}(s > 0, t > 0)$$

$$+ \mathbb{1}(\mathsf{x}_{n} + (s \vee t) < X \le \mathsf{x}_{n})\mathbb{1}(s < 0, t < 0)$$

$$- \mathbb{1}(\mathsf{x}_{n} \wedge (\mathsf{x}_{n} + s\eta_{n}) < X \le \mathsf{x}_{n} \vee (\mathsf{x}_{n} + s\eta_{n}))f_{0}(\mathsf{x}_{n})t\eta_{n}$$

$$- \mathbb{1}(\mathsf{x}_{n} \wedge (\mathsf{x}_{n} + t\eta_{n}) < X \le \mathsf{x}_{n} \vee (\mathsf{x}_{n} + t\eta_{n}))f_{0}(\mathsf{x}_{n})s\eta_{n}$$

$$+ f_{0}(\mathsf{x}_{n})^{2}st\eta_{n}^{2}.$$

Then, for any $s, t \in \mathbb{R}$ and $x_n \to x$, using continuity of f_0 at x,

$$\eta_n^{-1} \mathbb{E}[\psi_{\mathsf{x}_n}(s\eta_n; \mathbf{Z})\psi_{\mathsf{x}_n}(t\eta_n; \mathbf{Z})]
= \eta_n^{-1} \int_{\mathsf{x}_n}^{\mathsf{x}_n + \eta_n(s \wedge t)} f_0(u) du \mathbb{1}(s > 0, t > 0)
+ \eta_n^{-1} \int_{\mathsf{x}_n + \eta_n(s \vee t)}^{\mathsf{x}_n} f_0(u) du \mathbb{1}(s < 0, t < 0) + o(1)
= f_0(\mathsf{x})[(s \wedge t) \mathbb{1}(s > 0, t > 0) - (s \vee t) \mathbb{1}(s < 0, t < 0)] + o(1)
= f_0(\mathsf{x})(|s| \wedge |t|) \mathbb{1}(\mathrm{sign}(s) = \mathrm{sign}(t)) + o(1).$$

(B5) holds since $\hat{u}_n = \hat{u}_n^*$ converges in probability to u_0 , the supremum of the support of X by i.i.d. assumption.

(B6) and (B7) hold trivially since $\widehat{\Phi}_n$ and $\widehat{\Phi}_n^*$ are the identity map.

Assumption D follows from (E3) and empirical process theory arguments.

SA.1.12. Proof of Corollary 2. Assumption A and (E1)-(E2) follow from the hypothesis. (E3): We have $\widehat{\Gamma}_n = 1 - \widehat{S}_n$ with \widehat{S}_n the Kaplan-Meier estimator. By Theorem 1 of Lo and Singh (1986),

$$\sup_{x \in I} \left| \widehat{\Gamma}_n(x) - \frac{1}{n} \sum_{i=1}^n \gamma_0(x; \mathbf{Z}) \right| = O_{\mathbb{P}} \left(\left| \frac{\log n}{n} \right|^{3/4} \right).$$

Since $\sqrt{na_n} \le n^{2/3}$ for $\mathfrak{q} \ge 1$, $\sup_{x \in I} |\widehat{\Gamma}_n(x) - \Gamma_0(x)| = o_{\mathbb{P}}(1)$ and $\sqrt{na} \sup_{|v| \le \delta} |\widehat{\Gamma}_n(\mathsf{x} + v) - \widehat{\Gamma}_n(\mathsf{x})| = o_{\mathbb{P}}(1)$ hold.

$$\widehat{\gamma}_n(x; \mathbf{Z}) - \gamma_0(x; \mathbf{Z})$$

$$=\widehat{F}_{n}(x) - F_{0}(x) + \left[\widehat{S}_{n}(x) - S_{0}(x)\right] \left[\frac{\mathbb{1}(\check{X} \leq x)\Delta}{\widehat{S}_{n}(\check{X})\widehat{G}_{n}(\check{X})} - \int_{0}^{\check{X} \wedge x} \frac{\widehat{\Lambda}_{n}(du)}{\widehat{S}_{n}(u)\widehat{G}_{n}(u)}\right]$$

$$+ S_{0}(x)\mathbb{1}(\check{X} \leq x)\Delta \frac{S_{0}(\check{X})G_{0}(\check{X}) - \widehat{S}_{n}(\check{X})\widehat{G}_{n}(\check{X})}{S_{0}(\check{X})G_{0}(\check{X})\widehat{S}_{n}(\check{X})\widehat{G}_{n}(\check{X})}$$

$$- S_{0}(x) \int_{0}^{\check{X} \wedge x} \frac{S_{0}(u)G_{0}(u) - \widehat{S}_{n}(u)\widehat{G}_{n}(u)}{S_{0}(u)\widehat{G}_{n}(u)}\widehat{\Lambda}_{n}(du) - S_{0}(x) \int_{0}^{\check{X} \wedge x} \frac{[\widehat{\Lambda}_{n} - \Lambda_{0}](du)}{S_{0}(u)G_{0}(u)}.$$

By $S_0(u_0)G_0(u_0) > 0$, we have $\sqrt{n}\sup_{x \in I} |\widehat{S}_n(x) - S_0(x)| = O_{\mathbb{P}}(1)$, $\sqrt{n}\sup_{x \in I} |\widehat{G}_n(x) - G_0(x)| = O_{\mathbb{P}}(1)$, and $\sqrt{n}\sup_{x \in I} |\widehat{\Lambda}_n(x) - \Lambda_0(x)| = O_{\mathbb{P}}(1)$, which in turn implies

$$\frac{1}{n}\sum_{i=1}^n\sup_{x\in I}\left|\widehat{\gamma}_n(x;\mathbf{Z}_i)-\gamma_0(x;\mathbf{Z}_i)\right|^2=o_{\mathbb{P}}(1),\ a_n\frac{1}{n}\sum_{i=1}^n\sup_{x\in I_s^\delta}\left|\widehat{\gamma}_n(x;\mathbf{Z}_i)-\gamma_0(x;\mathbf{Z}_i)\right|^2=o_{\mathbb{P}}(1).$$

Let $\delta = \min\{\mathbf{x}, (u_0 - \mathbf{x})\}/4$, $R_{1n}(v) = |\widehat{F}_n(\mathbf{x} + v) - \widehat{F}_n(\mathbf{x}) - F_0(\mathbf{x} + v) + F_0(\mathbf{x})|$, $R_{2n} = |\widehat{F}_n(\mathbf{x}) - F_0(\mathbf{x})|$, $R_{3n} = \sup_{x \in I} |[\widehat{S}_n(x)\widehat{G}_n(x)]^{-1} - [S_0(x)G_0(x)]^{-1}|$, and $R_{4n}(x_1, x_2) = |\int_{x_1}^{x_2} \frac{\widehat{\Lambda}_n(du)}{\widehat{S}_n(u)\widehat{G}_n(u)} - \int_{x_1}^{x_2} \frac{\Lambda_0(du)}{S_0(u)G_0(u)}|$. For $|v| \leq \delta$,

$$\left| \check{\Gamma}_n(\mathsf{x} + v) - \check{\Gamma}_n(\mathsf{x}) - \bar{\Gamma}_n(\mathsf{x} + v) + \bar{\Gamma}_n(\mathsf{x}) \right|$$

$$\leq R_{1n}(v) \left[1 + \frac{1 + \widehat{\Lambda}_n(x)}{\widehat{S}_n(x)\widehat{G}_n(x)} \right] + R_{2n} \left[\widehat{S}_n(\mathsf{x} - \delta)\widehat{G}_n(\mathsf{x} - \delta) \right]^{-1} \frac{1}{n} \sum_{i=1}^n \mathbb{1}(|X_i - \mathsf{x}| \leq v)$$

$$+ R_{2n} \int_{\mathsf{x}-v}^{\mathsf{x}+v} \frac{\widehat{\Lambda}_n(du)}{\widehat{S}_n(u) \widehat{G}_n(u)} + \sup_{x \in I_s^{\delta}} f_0(x) |v| \left[R_{3n} + \frac{1}{n} \sum_{i=1}^n R_{4n}(0, \check{X}_i \wedge (\mathsf{x}+v)) \right]$$

$$+ R_{3n} \frac{1}{n} \sum_{i=1}^{n} \mathbb{1}(|X_i - \mathsf{x}| \le v) + \frac{1}{n} \sum_{i=1}^{n} R_{4n}(\check{X}_i \wedge \mathsf{x}, \check{X}_i \wedge (\mathsf{x} + v)).$$

As noted above, $\sup_{|v| \le \delta} R_{1n}(v) = o_{\mathbb{P}}((na_n)^{-1/2}), R_{2n} = O_{\mathbb{P}}(n^{-1/2}), \text{ and } R_{3n} = O_{\mathbb{P}}(n^{-1/2})$ by $S_0(u_0)G_0(u_0) > 0$. Also, uniformly over $V \in (0, 2\delta]$, $\sup_{|v| \le V} |\frac{1}{n} \sum_{i=1}^n (\mathbb{1}(X_i \le \mathsf{x} + v) - \mathbb{1}(X_i \le \mathsf{x}))| \le CV + O_{\mathbb{P}}(n^{-1/2})$. If

(SA.16)
$$\sup_{|v| \le \delta} \frac{1}{n} \sum_{i=1}^{n} R_{4n}(0, \check{X}_i \wedge (\mathsf{x} + v)) = O_{\mathbb{P}}(n^{-1/2})$$

and

(SA.17)
$$\frac{1}{n} \sum_{i=1}^{n} R_{4n}(\check{X}_i \wedge \mathsf{x}, \check{X}_i \wedge (\mathsf{x} + v)) \le |v| O_{\mathbb{P}}(n^{-1/2}) + o_{\mathbb{P}}((na_n)^{-1/2})$$

uniformly over $|v| \le 2\delta$, then there exist random variables $A_n = o_{\mathbb{P}}(1)$ and $B_n = O_{\mathbb{P}}(\sqrt{a_n})$ independent of v such that for $V \in (0, 2\delta]$,

$$\sqrt{na_n} \sup_{|v| \le V} \left| \check{\Gamma}_n(\mathsf{x} + v) - \check{\Gamma}_n(\mathsf{x}) - \bar{\Gamma}_n(\mathsf{x} + v) + \bar{\Gamma}_n(\mathsf{x}) \right| \le A_n + VB_n;$$

that is, $\beta_{\gamma} = 1$ in the notation of (E3). To show (SA.16) and (SA.17), for $x_1, x_2 \in I$,

$$R_{4n}(x_1, x_2) \le R_{3n} |\widehat{\Lambda}_n(x_2) - \widehat{\Lambda}_n(x_1)| + \left| \int_{x_1}^{x_2} \frac{[\widehat{\Lambda}_n - \Lambda_0](du)}{S_0(u)G_0(u)} \right|.$$

Let $J_0(u) = 1/[S_0(u)G_0(u)]$ and integration by parts implies

$$\begin{split} \int_{x_1}^{x_2} \frac{[\widehat{\Lambda}_n - \Lambda_0](du)}{S_0(u)G_0(u)} &= \frac{\widehat{\Lambda}_n(x_2) - \Lambda_0(x_2)}{S_0(x_2)G_0(x_2)} - \frac{\widehat{\Lambda}_n(x_1) - \Lambda_0(x_1)}{S_0(x_1)G_0(x_1)} - \int_{x_1}^{x_2} [\widehat{\Lambda}_n(u) - \Lambda_0(u)] J_0(du) \\ &= [\widehat{\Lambda}_n(x_2) - \Lambda_0(x_2)] \left[\frac{1}{S_0(x_2)G_0(x_2)} - \frac{1}{S_0(x_1)G_0(x_1)} \right] \\ &+ \frac{\widehat{\Lambda}_n(x_2) - \Lambda_0(x_2) - \widehat{\Lambda}_n(x_1) + \Lambda_0(x_1)}{S_0(x_1)G_0(x_1)} - \int_{x_1}^{x_2} [\widehat{\Lambda}_n(u) - \Lambda_0(u)] J_0(du). \end{split}$$

The first term after the second equality is bounded by $O_{\mathbb{P}}(n^{-1/2})|x_2-x_1|$. For the second term, Theorem 1 of Burke, Csörgő and Horváth (1988) implies that on a suitable probability space there exists a sequence of standard Brownian motion W_n such that $\sqrt{a_n}\sup_{|x_1-x_2|\leq v}|\sqrt{n}[\widehat{\Lambda}_n(x_1)-\Lambda_0(x_2)-\widehat{\Lambda}_n(x_1)+\Lambda_0(x_2)]-W_n(d(x_1))+W_n(d(x_2))|=o_{\mathbb{P}}(1),$ where $d(x)=\int_0^x\frac{F_0(du)}{S_0(u)^2G_0(u)}.$ By Theorem 3.2 of Pollard (1989), there is some fixed constant C>0 such that $\mathbb{E}[\sup_{|x_1-x_2|\leq v}|W_n(d(x_1))-W_n(d(x_2))|]\leq Cv.$ Finally, $|\int_{x_1}^{x_2}[\widehat{\Lambda}_n(u)-\Lambda_0(u)]J_0(du)|\leq \sup_{x'\in[x_1,x_2]}|\widehat{\Lambda}_n(x')-\Lambda_0(x')|[J_0(x_2)-J_0(x_1)]\leq O_{\mathbb{P}}(n^{-1/2})|x_2-x_1|.$ Thus, (SA.16) and (SA.17) hold.

For the function class \mathfrak{F}_{γ} , we can take $\bar{F}_{\gamma}(\mathbf{Z}) = 1 + [S_0(u_0)G_0(u_0)]^{-1}[1 + \Lambda_0(u_0)]$ as a constant envelope. For the function class $\{S_0(x): x \in I\}$, given $m \in \mathbb{N}$, there exists $\{x_1,\ldots,x_{m+1}\} \subset I$ such that $\sup_{x \in I} \min_{l=1,\ldots,m+1} |S_0(x_l) - S_0(x)| \leq 1/m$, which implies the uniform covering number is bounded by a linear function. The covering numbers of $\{\mathbb{1}(\cdot \leq s): s \in I\}$ and $\{\int_0^{\cdot \wedge s} [S_0(u)G_0(u)]^{-1}\Lambda_0(du): s \in I\}$ are also bounded by a linear function. By Lemma 5.1 of van der Vaart and van der Laan (2006), there exists $\rho \in (0,2)$ such that $\limsup_{n \downarrow 0} \log N_U(\eta, \mathfrak{F}_{\gamma})\eta^{\rho} < \infty$ holds.

Now consider the uniform covering number of $\widehat{\mathfrak{F}}_{\gamma}$. Given a realization of $(\widehat{S}_n,\widehat{G}_n)$, the mapping $x\mapsto \int_0^{x\wedge s} [\widehat{S}_n(u)\widehat{G}_n(u)]^{-1}\widehat{\Lambda}_n(du)$ is a composition of $x\mapsto x\wedge s$ and $x\mapsto \int_0^x [\widehat{S}_n(u)\widehat{G}_n(u)]^{-1}\widehat{\Lambda}_n(du)$. The latter mapping is monotone, and the first mapping is a VC-subgraph class, and Lemma 2.6.18 of van der Vaart and Wellner (1996) implies $\{\int_0^{\cdot \wedge s} [\widehat{S}_n(u)\widehat{G}_n(u)]^{-1}\widehat{\Lambda}_n(du): s\in I\}$ is a VC-subgraph class. Note that since S_0, G_0 are bounded away from zero, $\widehat{S}_n, \widehat{G}_n$ are bounded away from zero with probability approaching one. Thus, for some $\rho\in(0,2)$, $\limsup_{\eta\downarrow 0}\log N_U(\eta,\widehat{\mathfrak{F}}_{\gamma})\eta^{\rho}=O_{\mathbb{P}}(1)$ holds. For $s\leq t\in I$,

$$|\gamma_0(s; \mathbf{Z}) - \gamma_0(t; \mathbf{Z})| \le C|F_0(s) - F_0(t)| + C|\mathbb{1}(\check{X} \le s) - \mathbb{1}(\check{X} \le t)|\Delta + \int_{\check{X} \land s}^{\check{X} \land t} \frac{\Lambda_0(du)}{S_0(du)G_0(du)}$$

and we can take $D_{\gamma}^{\eta}(\mathbf{Z})$ to be a constant multiple of $\sup_{|s| \leq \eta} |F_0(\mathbf{x}+s) - F_0(\mathbf{x})| + \Delta \mathbb{1}(|\check{X} - \mathbf{x}| \leq \eta) + \int_{x-\eta}^{x+\eta} \Lambda_0(du)/S_0(u)G_0(u)$. For $\eta > 0$ small enough, there is some fixed C > 0 with

$$\mathbb{E}[D_{\gamma}^{\eta}(\mathbf{Z})^2 + D_{\gamma}^{\eta}(\mathbf{Z})^4] \le C f_0(\mathsf{x} + \eta)\eta.$$

(E4) trivially holds as $\widehat{\Phi}_n(x) = \widehat{\Phi}_n^*(x) = x$.

(E5): We have

$$\psi_{\mathsf{x}}(v; \mathbf{Z}) = S_0(\mathsf{x}) \frac{(\mathbb{1}(\check{X} \le \mathsf{x} + v) - \mathbb{1}(\check{X} \le \mathsf{x}))\Delta}{S_0(\check{X})G_0(\check{X})} + O(|v|)$$

where O(|v|) is uniformly over small enough |v|. Since

$$\mathbb{E}[|\mathbb{1}(\check{X} \le \mathsf{x} + v) - \mathbb{1}(\check{X} \le \mathsf{x} + v')|\Delta] = \int_{\mathsf{x} + v \wedge v'}^{\mathsf{x} + v \vee v'} G_0(u) f_0(u) du \le C|v - v'|,$$

the first display in (E5) is satisfied. For the covariance kernel,

$$\begin{split} \mathbb{E}[\psi_{\mathsf{x}_n}(s\eta_n;\mathbf{Z})\psi_{\mathsf{x}_n}(t\eta_n;\mathbf{Z})] \\ &= S_0(\mathsf{x}_n)^2 \bigg(\int_{\mathsf{x}_n}^{\mathsf{x}_n + \eta_n(s \wedge t)} \frac{f_0(u)}{S_0(u)^2 G_0(u)} du \mathbb{1}(s > 0, t > 0) \\ &+ \int_{\mathsf{x}_n + \eta_n(s \vee t)}^{\mathsf{x}_n} \frac{f_0(u)}{S_0(u)^2 G_0(u)} du \mathbb{1}(s < 0, t < 0) \bigg) + O(\eta_n^2) \\ &= \frac{S_0(\mathsf{x}_n)^2 f_0(\mathsf{x})}{S_0(\mathsf{x})^2 G_0(\mathsf{x})} (s \wedge t) \eta_n \mathbb{1}(s > 0, t > 0) \\ &- \frac{S_0(\mathsf{x}_n)^2 f_0(\mathsf{x})}{S_0(\mathsf{x})^2 G_0(\mathsf{x})} (s \vee t) \eta_n \mathbb{1}(s < 0, t < 0) + o(\eta_n) \end{split}$$

where the last equality uses continuity of (S_0, G_0, f_0) at \times i.e., $\int_{\mathsf{x}_n}^{\mathsf{x}_n + \eta_n} \left[\frac{f_0(u)}{S_0(u)^2 G_0(u)} - \frac{f_0(\mathsf{x})}{S_0(\mathsf{x})^2 G_0(\mathsf{x})} \right] du = o(1) \eta_n$.

(B5), (B6), and (B7) hold since in this example, $\widehat{u}_n = \widehat{u}_n^* = u_0$ and $\widehat{\Phi}_n$, $\widehat{\Phi}_n^*$ are the identity map.

Assumption D. As noted when verifying (E3), $\widehat{G}_n(1;\eta_n) = o_{\mathbb{P}}(1)$ for any $\eta_n = o(1)$ with $a_n^{-1}\eta_n^{-1} = O(1)$. $\widehat{\Phi}_n = \Phi_0$ is the identity map and the desired result holds.

SA.1.13. Proof of Corollary 3. Assumption A and (E1)-(E2) follow from the hypothesis. (E3): In this example, $\gamma_0(x; \mathbf{Z}) = Y\mathbbm{1}(X \le x)$ is known, so it suffices to verify (SA.15). The uniform covering number bound is straightforward as $\{\mathbbm{1}(\cdot \le x) : x \in \mathbbm{R}\}$ is a VC-subgraph class. An envelope function is |Y|, whose second moment is finite. For $x \in I_{\times}^{\eta}$, $|\gamma_0(x; \mathbf{Z}) - \gamma_0(x; \mathbf{Z})| \le |Y|\mathbbm{1}(x - \eta \le X \le x + \eta)$, which we can take as $\bar{D}_{\gamma}^{\eta}(\mathbf{Z})$. Then, for j = 2, 4,

$$\mathbb{E}[\bar{D}_{\gamma}^{\eta}(\mathbf{Z})^{j}] \le 2^{j-1} \int_{\mathbf{x}-\eta}^{\mathbf{x}+\eta} (|\mu_{0}(x)|^{j} + \mathbb{E}[\varepsilon^{j}|X=x]) f_{0}(x) dx \le C\eta$$

and the desired bound holds.

(E4): $\phi_0(x; \mathbf{Z}) = \mathbb{1}(X \le x)$ is known, so it suffices to verify the analogue of (SA.15). The argument is the same as for checking (E3) in monotone density estimation with no censoring. (E5): We have

$$\psi_{\mathsf{x}}(v; \mathbf{Z}) = \varepsilon (\mathbb{1}(X \le \mathsf{x} + v) - \mathbb{1}(X \le \mathsf{x})) + (\mu_0(X) - \mu_0(\mathsf{x}))(\mathbb{1}(X \le \mathsf{x} + v) - \mathbb{1}(X \le \mathsf{x})).$$

Then,

$$\mathbb{E}[|\psi_{\mathsf{x}}(v;\mathbf{Z}) - \psi_{\mathsf{x}}(v';\mathbf{Z})|] \le \int_{\mathsf{x}+v \wedge v'}^{\mathsf{x}+v \vee v'} [\sigma_0(x) + |\mu_0(x) - \mu_0(\mathsf{x})|] f_0(x) dx \le C|v - v'|$$

and the first display holds. For the covariance kernel, note $|(\mu_0(X) - \mu_0(\mathsf{x}_n))(\mathbbm{1}(X \le \mathsf{x}_n + v) - \mathbbm{1}(X \le \mathsf{x}_n)| \le |v| \sup_{|x-\mathsf{x}| \le 2\eta} |\partial \mu_0(x)|$ for $|x_n - \mathsf{x}| \lor |v| \le \eta$ for $\eta > 0$ small enough. Then,

$$\mathbb{E}[\psi_{\mathsf{x}_{n}}(s\eta_{n};\mathbf{Z})\psi_{\mathsf{x}_{n}}(t\eta_{n};\mathbf{Z})] \\
= \mathbb{E}[\varepsilon^{2}(\mathbb{1}(X \leq \mathsf{x}_{n} + s\eta_{n}) - \mathbb{1}(X \leq \mathsf{x}_{n}))(\mathbb{1}(X \leq \mathsf{x}_{n} + t\eta_{n}) - \mathbb{1}(X \leq \mathsf{x}_{n}))] + O(\eta_{n}^{2}) \\
= \int_{\mathsf{x}_{n}}^{\mathsf{x}_{n} + \eta_{n}(s \wedge t)} \sigma_{0}^{2}(x)f_{0}(x)dx\mathbb{1}(s > 0, t > 0) \\
+ \int_{\mathsf{x}_{n} + \eta_{n}(s \vee t)}^{\mathsf{x}_{n}} \sigma_{0}^{2}(x)f_{0}(x)dx\mathbb{1}(s < 0, t < 0) + O(\eta_{n}^{2})$$

and

$$\eta_n^{-1} \mathbb{E}[\psi_{\mathsf{x}_n}(s\eta_n; \mathbf{Z})\psi_{\mathsf{x}_n}(t\eta_n; \mathbf{Z})]$$

$$\to \sigma_0^2(\mathsf{x}) f_0(\mathsf{x}) \big[(s \wedge t) \mathbb{1}(s > 0, t > 0) - (s \vee t) \mathbb{1}(s < 0, t < 0) \big]$$

$$= \sigma_0^2(\mathsf{x}) f_0(\mathsf{x}) (|s| \wedge |t|) \mathbb{1}(\mathrm{sign}(s) = \mathrm{sign}(t)),$$

as desired.

(B5) trivially holds since $\widehat{u}_n = \widehat{u}_n^* = 1$ in this example.

(B6): $\widehat{\Phi}_n, \widehat{\Phi}_n^*$ are (empirical) cdfs, so they are non-negative, non-decreasing, and right-continuous. $\{0, \widehat{u}_n\} \subset \widehat{\Phi}_n(I)$ and $\{0, \widehat{u}_n^*\} \subset \widehat{\Phi}_n^*(I)$ hold as $\widehat{u}_n = \widehat{u}_n^* = 1$, $\widehat{\Phi}_n(\min_i X_i -) = 0 = \widehat{\Phi}_n^*(\min_i X_i -)$, and $\widehat{\Phi}_n(\max_i X_i) = 1 = \widehat{\Phi}_n^*(\max_i X_i)$. The sets $\widehat{\Phi}_n(I), \widehat{\Phi}_n^*(I)$ are finite and thus closed.

(B7): With probability one, all X_i 's are distinct. If x is one of X_i 's, $\widehat{\Phi}_n^*(x) - \widehat{\Phi}_n^*(x-) = n^{-1}W_{j,n}$ for some j, and $\widehat{\Phi}_n^*(x) - \widehat{\Phi}_n^*(x-) = 0$ otherwise.

Given $\mathbb{E}|W_{1,n}|^{\mathfrak{r}}<\infty$, $n^{-1}\max_{1\leq i\leq n}|W_{i,n}|=o_{\mathbb{P}}(n^{-5/6})$, which implies

$$\sqrt{na_n} \sup_{x \in I} |\widehat{\Phi}_n^*(x) - \widehat{\Phi}_n^*(x-)| = o_{\mathbb{P}}(1).$$

The argument for $\widehat{\Phi}_n$ is similar.

Assumption D follows from (E3)-(E4) and empirical process theory arguments.

SA.1.14. *Proof of Corollary 4*. Assumption A and (E1)-(E2) follow from the hypothesis. (E3): In this example, $\tilde{\Gamma}_n = \hat{\Gamma}_n$.

$$|\widehat{\gamma}_n(x; \mathbf{Z}) - \gamma_0(x; \mathbf{Z})| \leq \mathbb{I}(X \leq x) \left[|\varepsilon| \left| \widehat{g}_n(X, \mathbf{A})^{-1} - g_0(X, \mathbf{A})^{-1} \right| + \frac{|\widehat{\mu}_n(X, \mathbf{A}) - \mu_0(X, \mathbf{A})|}{\widehat{g}_n(X, \mathbf{A})} + \frac{1}{n} \sum_{j=1}^n |\widehat{\mu}_n(X, \mathbf{A}_j) - \mu_0(X, \mathbf{A}_j)| + \left| \frac{1}{n} \sum_{j=1}^n \mu_0(X, \mathbf{A}_j) - \theta_0(X) \right| \right].$$

The last sum is bounded by $\sup_{x \in I} |\frac{1}{n} \sum_{j=1}^n \mu_0(x, \mathbf{A}_j) - \theta_0(x)|$, and this object is $O_{\mathbb{P}}(n^{-1/2})$: to see this claim, first note that Assumption MRC (iv) and Theorem 2.7.11 of van der Vaart and Wellner

(1996) imply $\limsup_{\epsilon\downarrow 0}\log N_U(\epsilon,\{\mu(x,\cdot):x\in I\})\epsilon^V<\infty$ for some $V\in(0,2)$ and Theorem 4.2 of Pollard (1989) implies $\sup_{x\in I}|\frac{1}{n}\sum_{j=1}^n\mu_0(x,\mathbf{A}_j)-\theta_0(x)|=O_{\mathbb{P}}(n^{-1/2})$. Together with Assumption MRC (iii), $a_n\frac{1}{n}\sum_{i=1}^n\sup_{x\in I}|\widehat{\gamma}_n(x;\mathbf{Z})-\gamma_0(x;\mathbf{Z})|^2=o_{\mathbb{P}}(1)$ holds. By Assumption MRC (iii), uniformly over $V\in(0,2\delta]$

$$\sqrt{na_n} \sup_{|v| \le V} \left| \widehat{\Gamma}_n(\mathbf{x} + v) - \widehat{\Gamma}_n(\mathbf{x}) - \bar{\Gamma}_n(\mathbf{x} + v) + \bar{\Gamma}_n(\mathbf{x}) \right| \le o_{\mathbb{P}}(1) + O_{\mathbb{P}}(\sqrt{a_n})V$$

and the desired inequality holds.

The uniform covering numbers of \mathfrak{F}_{γ} , $\hat{\mathfrak{F}}_{\gamma,n}$ are the same order as for $\{\mathbb{1}(\cdot \leq x) : x \in I\}$. For $x \in I_{\mathsf{x}}^{\eta}$, $|\gamma_0(x; \mathbf{Z}) - \gamma_0(\mathsf{x}; \mathbf{Z})| \leq \mathbb{1}(\mathsf{x} - \eta \leq X \leq \mathsf{x} + \eta)(|\varepsilon|c^{-1} + \theta_0(\mathsf{x} + \eta))$. Then, $\limsup_{\eta \downarrow 0} \mathbb{E}[\bar{D}_{\gamma}^{\eta}(\mathbf{Z})^j]\eta^{-1} < \infty$ holds for j = 2, 4.

(E4): $\phi_0(x; \mathbf{Z}) = \mathbb{1}(X \le x)$ is known and the same as in the classical case, so the same argument applies.

(E5): We have

$$\psi_{\mathsf{x}}(v;\mathbf{Z}) = (\mathbb{1}(X \le \mathsf{x} + v) - \mathbb{1}(X \le \mathsf{x})) \left[\frac{\varepsilon}{g_0(X,\mathbf{A})} + \theta_0(X) - \theta_0(\mathsf{x}) \right].$$

Then, for $v, v' \in [-\eta, \eta]$ with sufficiently small $\eta > 0$,

$$|\psi_{\mathsf{x}}(v;\mathbf{Z}) - \psi_{\mathsf{x}}(v';\mathbf{Z})| \leq |\mathbb{1}(X \leq \mathsf{x} + v) - \mathbb{1}(X \leq \mathsf{x} + v')| \left(c^{-1}|\varepsilon| + |X - \mathsf{x}| \sup_{x \in L^p} |\partial \theta_0(x)|\right)$$

and $\sup_{v\neq v'\in[-\eta_n,\eta_n]}\mathbb{E}[|\psi_{\mathsf{X}}(v;\mathbf{Z})-\psi_{\mathsf{X}}(v';\mathbf{Z})|]/|v-v'|=O(1)$ holds.

For $s\eta_n$ small enough, $\psi_{\mathsf{x}}(s\eta_n;\mathbf{Z})=(\mathbbm{1}(X\leq\mathsf{x}+s\eta_n)-\mathbbm{1}(X\leq\mathsf{x}))\varepsilon g_0(X,\mathbf{A})^{-1}+O(\eta_n)$ and

$$\mathbb{E}[\psi_{\mathsf{x}}(s\eta_n;\mathbf{Z})\psi_{\mathsf{x}}(t\eta_n;\mathbf{Z})]$$

$$= \mathbb{E}\left[\frac{\sigma_0^2(X,\mathbf{A})}{g_0(X,\mathbf{A})^2}(\mathbb{1}(X\leq \mathsf{x}+s\eta_n)-\mathbb{1}(X\leq \mathsf{x}))(\mathbb{1}(X\leq \mathsf{x}+t\eta_n)-\mathbb{1}(X\leq \mathsf{x}))\right]+O(\eta_n^2)$$

and

$$\mathbb{E}\left[\frac{\sigma_0^2(X,\mathbf{A})}{g_0(X,\mathbf{A})^2}(\mathbb{1}(X\leq \mathsf{x}+s\eta_n)-\mathbb{1}(X\leq \mathsf{x}))(\mathbb{1}(X\leq \mathsf{x}+t\eta_n)-\mathbb{1}(X\leq \mathsf{x}))\right] \\
=\mathbb{E}\left[\int_{\mathsf{x}}^{\mathsf{x}+\eta_n(s\wedge t)} \frac{\sigma_0^2(x,\mathbf{A})}{g_0(x,\mathbf{A})^2} f_{X|\mathbf{A}}(x|\mathbf{A}) dx\right] \mathbb{1}(s>0,t>0) \\
+\mathbb{E}\left[\int_{\mathsf{x}+\eta_n(s\vee t)}^{\mathsf{x}} \frac{\sigma_0^2(x,\mathbf{A})}{g_0(x,\mathbf{A})^2} f_{X|\mathbf{A}}(x|\mathbf{A}) dx\right] \mathbb{1}(s<0,t<0) \\
=\mathbb{E}\left[\frac{\sigma_0^2(\mathsf{x},\mathbf{A})}{g_0(\mathsf{x},\mathbf{A})^2} f_{X|\mathbf{A}}(\mathsf{x}|\mathbf{A})\right] \left(\eta_n(s\wedge t)\mathbb{1}(s>0,t>0)-\eta_n(s\vee t)\mathbb{1}(s<0,t<0)\right) + o(\eta_n).$$

Since
$$\frac{f_{X|\mathbf{A}}(x|\mathbf{A})}{g_0(x,\mathbf{A})^2} = \frac{f_0(x)}{g_0(x,\mathbf{A})}$$
, we have

$$\eta_n^{-1} \mathbb{E}[\psi_{\mathsf{x}}(s\eta_n; \mathbf{Z})\psi_{\mathsf{x}}(t\eta_n; \mathbf{Z})] \to f_0(\mathsf{x}) \mathbb{E}\left[\frac{\sigma_0^2(\mathsf{x}, \mathbf{A})}{g_0(\mathsf{x}, \mathbf{A})}\right] \Big((s \wedge t)\mathbb{1}(s, t > 0) - (s \vee t)\mathbb{1}(s, t < 0)\Big),$$

as desired.

(B5) (B6) (B7): Verifying these conditions is the same as in the classical monotone regression case.

SA.2. Primitive sufficient conditions for Assumption MRC (iii). Here we provide primitive sufficient conditions for Assumption MRC (iii) by focusing on specific estimators $\widehat{\mu}_n$ and \widehat{g}_n . As discussed by Westling, Gilbert and Carone (2020), cross-fitting avoids restrictions on uniform entropy, allowing for a large class of flexible preliminary estimators. Here we use sample splitting to simplify exposition, but the proposed procedure can be straightforwardly modified for cross-fitting.

Suppose there is a separate random sample $\tilde{\mathbf{Z}}_1,\ldots,\tilde{\mathbf{Z}}_n$ drawn from the distribution of \mathbf{Z} , which is independent of $\mathbf{Z}_1,\ldots,\mathbf{Z}_n$. Preliminary estimators $\widehat{\mu}_n$ and \widehat{g}_n are constructed from $\tilde{\mathbf{Z}}_1,\ldots,\tilde{\mathbf{Z}}_n$. For concreteness, we consider a partitioning-based least squares estimator $\widehat{\mu}_n$ (Cattaneo, Farrell and Feng, 2020) and local polynomial kernel-based estimators $\widehat{f}_{X|\mathbf{A},n}(x|\mathbf{a})$ and $\widehat{f}_n(x)$ of $f_{X|\mathbf{A}}(x|\mathbf{a})$ and $f_0(x)$ (Cattaneo, Jansson and Ma, 2020; Cattaneo et al., 2024), from which we construct $\widehat{g}_n(x,\mathbf{a}) = \widehat{f}_{X|\mathbf{A},n}(x|\mathbf{a})/\widehat{f}_n(x)$.

Let $d = \dim(\mathbf{A})$. For simplicity, suppose the support of $(X, \mathbf{A}')'$ equals $[0, 1]^{1+d}$. Let $\mathbf{p}(x, \mathbf{a})$ be a k_n -dimensional vector of bounded basis functions of order m on S which are locally supported e.g., splines (see Cattaneo, Farrell and Feng, 2020, for details and examples of basis functions). We consider the estimator

$$\widehat{\mu}_n(x,\mathbf{a}) = \mathbf{p}(x,\mathbf{a})' \left(\sum_{i=1}^n \mathbf{p}(\widetilde{X}_i, \widetilde{\mathbf{A}}_i) \mathbf{p}(\widetilde{X}_i, \widetilde{\mathbf{A}}_i)' \right)^{-1} \sum_{i=1}^n \mathbf{p}(\widetilde{X}_i, \widetilde{\mathbf{A}}_i) \widetilde{Y}_i.$$

For the estimator of $f_{X|\mathbf{A}}(x|\mathbf{a})$, letting $\widehat{F}_{X|\mathbf{A},n}(\cdot|\mathbf{a})$ be an estimator of $\mathbb{P}[X \leq \cdot | \mathbf{A} = \mathbf{a}]$ specified below, $\widehat{f}_{X|\mathbf{A},n}(x|\mathbf{a})$ is obtained by local polynomial regression:

$$\widehat{f}_{X|\mathbf{A},n}(x|\mathbf{a}) = \mathbf{e}_2' \underset{\mathbf{u} \in \mathbb{R}^{\mathfrak{p}_1+1}}{\operatorname{argmin}} \sum_{i=1}^n \left(\widehat{F}_{X|\mathbf{A},n}(\widetilde{X}_i|\mathbf{a}) - \mathbf{q}_1(\widetilde{X}_i - x)'\mathbf{u} \right)^2 K_h(\widetilde{X}_i - x)$$

where $\mathfrak{p}_1 \geq 1$ is the order of the polynomial basis $\mathbf{q}_1(x) = (1, x/1!, x^2/2!, \dots, x^{\mathfrak{p}_1}/\mathfrak{p}_1!)'$, \mathbf{e}_l is the conformable unit vector whose lth element is unity, and $K_h(x) = K(x/h)/h$ for some kernel function K and some positive bandwidth h. The estimator $\widehat{F}_{X|\mathbf{A},n}(x|\mathbf{a})$ is constructed via local polynomial regression of order $\mathfrak{p}_2 = \mathfrak{p}_1 - 1$:

$$\widehat{F}_{X|\mathbf{A},n}(x|\mathbf{a}) = \mathbf{e}_1' \underset{\mathbf{v} \in \mathbb{R}^{k_{\mathfrak{p}_2}}}{\operatorname{argmin}} \sum_{i=1}^n \left(\mathbb{1}(\widetilde{X}_i \leq x) - \mathbf{q}_2(\widetilde{\mathbf{A}}_i - \mathbf{a})' \mathbf{v} \right)^2 L_h(\widetilde{\mathbf{A}}_i - \mathbf{a})$$

where, using standard multi-index notation, $\mathbf{q}_2(\mathbf{a})$ denotes the $k_{\mathfrak{p}_2}$ -dimensional vector collecting the polynomials $\mathbf{a}^{\mathbf{m}}/\mathbf{m}!$ for $0 \leq |\mathbf{m}| \leq \mathfrak{p}_2$ with $\mathbf{a}^{\mathbf{m}} = a_1^{m_1} a_2^{m_2} \dots a_d^{m_d}$, $|\mathbf{m}| = \sum_{j=1}^d m_j$, and $k_{\mathfrak{p}_2} = \frac{(d+\mathfrak{p}_2)!}{d!\mathfrak{p}_2!} + 1$, and $L_h(\mathbf{a}) = L(\mathbf{a}/h)/h^d$ for $L(\mathbf{a}) = \prod_{j=1}^d K(a_j)$ i.e., product kernel. The estimator $\widehat{f}_n(x)$ is constructed in a similar manner. First, the empirical cdf \widehat{F}_n of $\{\widetilde{X}_i\}$ is constructed and then $\widehat{f}_n(x)$ is formed via local polynomial regression:

$$\widehat{f}_n(x) = \mathbf{e}'_2 \underset{\mathbf{w} \in \mathbb{R}^{\mathfrak{p}_1} + 1}{\operatorname{argmin}} \sum_{i=1}^n \left(\widehat{F}_n(\tilde{X}_i) - \mathbf{q}_1(\tilde{X}_i - x)' \mathbf{w} \right)^2 K_b(\tilde{X}_i - x)$$

where b > 0 is some bandwidth.

Now we state sufficient conditions for Assumption MRC (ii) based on the partitioning-based series estimator $\hat{\mu}_n$ and the kernel-based estimator \hat{g}_n .

Primitive Conditions MRC

(i) $\tilde{\mathbf{Z}}_1, \dots, \tilde{\mathbf{Z}}_n$ are independent of $\mathbf{Z}_1, \dots, \mathbf{Z}_n$.

- (ii) The support of $(X, \mathbf{A}')'$ is $[0, 1]^{1+d}$, and the distribution of $(X, \mathbf{A}')'$ is absolutely continuous. The Lebesgue density of $(X, \mathbf{A}')'$ and the conditional variance of Y given $(X, \mathbf{A}')'$ are bounded away from zero and continuous on $[0, 1]^{1+d}$. μ_0 is (m+1)-times continuously differentiable on $[0, 1]^{1+d}$.
- (iii) The vector of basis functions p satisfies Assumptions 2, 3, and 4 of Cattaneo, Farrell and Feng (2020).
- (iv) $f_{X|\mathbf{A}}(x|\mathbf{a})$ and $f_0(x)$ are \mathfrak{p}_1 -times continuously differentiable in x, and $f_{X|\mathbf{A}}(x|\mathbf{a})$ is \mathfrak{p}_1 -times continuously differentiable in \mathbf{a} .
- (v) K is a symmetric, Lipschitz continuous probability density function supported on [-1,1].

As verified by Cattaneo, Farrell and Feng (2020), (iii) holds for widely used local basis functions such as splines and wavelets.

LEMMA SA-7. Suppose $\tilde{\mathbf{Z}}_1,\ldots,\tilde{\mathbf{Z}}_n$ is a random sample drawn from the distribution of \mathbf{Z} and Primitive Conditions MRC hold. In addition, with $\tau_n=n^{-1}\log n$, $k_n=O(\tau_n^{-\frac{d+1}{2m+d+1}})$, $h=O(\tau_n^{\frac{d+1}{2p_1+d+1}})$, $b=O(\tau_n^{\frac{1}{2p_1+d+1}})$, $\frac{m}{2m+d+1}+\frac{\mathfrak{p}_1}{2\mathfrak{p}_1+d+1}\geq \frac{1}{2}$, and $\min\{\frac{2m}{2m+d+1},\frac{2\mathfrak{p}_1}{2\mathfrak{p}_1+d+1}\}>\frac{1}{2\mathfrak{q}+1}$. Then, $\widehat{\mu}_n$ and \widehat{g}_n described above and $\widehat{\Gamma}_n$ based on the $\widehat{\mu}_n,\widehat{g}_n$ satisfy Assumption MRC (iii) with $\delta=\min\{\mathsf{x},1-\mathsf{x}\}/4$. In particular,

(SA.18)
$$\sqrt{n} \sup_{|v| \le V} \left| \widehat{\Gamma}_n(\mathsf{x} + v) - \widehat{\Gamma}_n(\mathsf{x}) - \overline{\Gamma}_n(\mathsf{x} + v) + \overline{\Gamma}_n(\mathsf{x}) \right| \le VO_{\mathbb{P}}(1) + o_{\mathbb{P}}(a_n^{-1})$$

uniformly over $V \in (0, 2\delta]$.

PROOF. By Theorem 4.3 of Cattaneo, Farrell and Feng (2020),

$$\sup_{(x,\mathbf{a})\in[0,1]^{1+d}} |\widehat{\mu}_n(x,\mathbf{a}) - \mu_0(x,\mathbf{a})| = O_{\mathbb{P}}\left(\sqrt{\frac{k_n \log n}{n}} + k_n^{-\frac{m}{d+1}}\right)$$

and by Theorem 1 of Cattaneo et al. (2024),

$$\sup_{(x,\mathbf{a})\in[0,1]^{1+d}}|\widehat{f}_{X|\mathbf{A},n}(x|\mathbf{a}) - f_{X|\mathbf{A}}(x|\mathbf{a})| = O_{\mathbb{P}}\left(\sqrt{\frac{\log n}{nh^{1+d}}} + h^{\mathfrak{p}_1}\right).$$

Also, one can show

$$\sup_{x \in [0,1]} |\widehat{f}_n(x) - f_0(x)| = O_{\mathbb{P}}\left(\sqrt{\frac{\log n}{nb}} + b^{\mathfrak{p}_1}\right).$$

Then, with the specified rate of k_n , h, b,

$$\sup_{(x,\mathbf{a})\in[0,1]^{1+d}}|\widehat{\mu}_n(x,\mathbf{a})-\mu_0(x,\mathbf{a})|=O_{\mathbb{P}}\left(\tau_n^{\frac{m}{2m+d+1}}\right),$$

$$\sup_{(x,\mathbf{a})\in[0,1]^{1+d}}|\widehat{g}_n(x,\mathbf{a})-g_0(x,\mathbf{a})|=O_{\mathbb{P}}\Big(\tau_n^{\frac{\mathfrak{p}_1}{2\mathfrak{p}_1+d+1}}\Big).$$

Since $\min\{\frac{2m}{2m+d+1}, \frac{2\mathfrak{p}_1}{2\mathfrak{p}_1+d+1}\} > \frac{1}{2\mathfrak{q}+1}$, it follows $a_n \frac{1}{n} \sum_{i=1}^n |\widehat{\mu}_n(X_i, \mathbf{A}_i) - \mu_0(X_i, \mathbf{A}_i)|^2 = o_{\mathbb{P}}(1), a_n \frac{1}{n^2} \sum_{i=1}^n \sum_{j=1}^n |\widehat{\mu}_n(X_i, \mathbf{A}_j) - \mu_0(X_i, \mathbf{A}_j)|^2 = o_{\mathbb{P}}(1), \text{ and } a_n \frac{1}{n} \sum_{i=1}^n \varepsilon_i^2 |\widehat{g}_n(X_i, \mathbf{A}_i) - g_0(X_i, \mathbf{A}_i)|^2 = o_{\mathbb{P}}(1).$ Also, by $\frac{m}{2m+d+1} + \frac{\mathfrak{p}_1}{2\mathfrak{p}_1+d+1} \ge \frac{1}{2}$, (SA.19)

$$\sup_{(x,\mathbf{a})\in[0,1]^{1+d}} |\widehat{\mu}_n(x,\mathbf{a}) - \mu_0(x,\mathbf{a})| \sup_{(x,\mathbf{a})\in[0,1]^{1+d}} |\widehat{g}_n(x,\mathbf{a}) - g_0(x,\mathbf{a})| = O_{\mathbb{P}}\Big(n^{-1/2}\Big).$$

Decompose $\widehat{\gamma}_n$ into $\widehat{\gamma}_{1,n}$ and $\widehat{\gamma}_{2,n}$ where

$$\widehat{\gamma}_{1,n}(x; \mathbf{Z}) = \mathbb{1}(X \le x) \frac{Y - \widehat{\mu}_n(X, \mathbf{A})}{\widehat{g}_n(X, \mathbf{A})}, \quad \widehat{\gamma}_{2,n}(x; \mathbf{Z}) = \mathbb{1}(X \le x) \frac{1}{n} \sum_{j=1}^n \widehat{\mu}_n(X, \mathbf{A}_j)$$

and let $\widehat{\Gamma}_{k,n}(x) = \frac{1}{n} \sum_{i=1}^n \widehat{\gamma}_{1,n}(x; \mathbf{Z}_i)$ for k=1,2. Define $\gamma_{k,0}, \overline{\Gamma}_{k,n}$ k=1,2 in the same manner.

Letting $\tilde{\mathfrak{Z}}_n$ be the σ -field generated by $\tilde{\mathbf{Z}}_1, \ldots, \tilde{\mathbf{Z}}_n$,

$$\mathbb{V}\left[\widehat{\Gamma}_{1,n}(\mathsf{x}+v) - \widehat{\Gamma}_{1,n}(\mathsf{x}) - \overline{\Gamma}_{1,n}(\mathsf{x}+v) + \overline{\Gamma}_{1,n}(\mathsf{x}) \middle| \widetilde{\mathfrak{Z}}_{n} \right] \\
\leq n^{-1}\mathbb{E}\left[\left(\mathbb{1}(X \leq \mathsf{x}+v) - \mathbb{1}(X \leq \mathsf{x}) \right)^{2} \left(\frac{Y - \widehat{\mu}_{n}(X,\mathbf{A})}{\widehat{g}_{n}(X,\mathbf{A})} - \frac{Y - \mu_{0}(X,\mathbf{A})}{g_{0}(X,\mathbf{A})} \right)^{2} \middle| \widetilde{\mathfrak{Z}}_{n} \right] \\
\leq n^{-1}C\mathbb{E}\left[\left| \mathbb{1}(X \leq \mathsf{x}+v) - \mathbb{1}(X \leq \mathsf{x}) \middle| \varepsilon^{2} \right] \sup_{|x-\mathsf{x}| \leq |v|} \sup_{\mathbf{a} \in [0,1]^{d}} |\widehat{g}_{n}(x,\mathbf{a}) - g_{0}(x,\mathbf{a})|^{2} \\
+ n^{-1}C\mathbb{E}\left[\left| \mathbb{1}(X \leq \mathsf{x}+v) - \mathbb{1}(X \leq \mathsf{x}) \middle| \right] \sup_{|x-\mathsf{x}| \leq |v|} \sup_{\mathbf{a} \in [0,1]^{d}} |\widehat{\mu}_{n}(x,\mathbf{a}) - \mu_{0}(x,\mathbf{a})|^{2} \\
= o_{\mathbb{P}}\left((na_{n})^{-1} \right)$$

where the inequalities hold with probability one.

Note
$$\widehat{\Gamma}_{2,n}(x) = \frac{1}{n^2} \sum_{1 \leq i \neq j \leq n} \mathbb{1}(X_i \leq x) \widehat{\mu}_n(X_i, \mathbf{A}_j) + O_{\mathbb{P}}(n^{-1})$$
, and

$$\mathbb{V}\left[\frac{1}{n(n-1)} \sum_{1 \le i \ne j \le n} \left(\mathbb{1}(X_i \le x + v) - \mathbb{1}(X_i \le x)\right) \left(\widehat{\mu}_n(X_i, \mathbf{A}_j) - \mu_0(X_i, \mathbf{A}_j)\right) \middle| \widetilde{\mathfrak{Z}}_n \right] \\
\le n^{-1} C|v| \sup_{|x-x| \le |v|} \sup_{\mathbf{a} \in [0,1]^d} |\widehat{\mu}_n(x, \mathbf{a}) - \mu_0(x, \mathbf{a})|^2 \\
+ Cn^{-2}|v| \sup_{|x-x| \le |v|} \sup_{\mathbf{a} \in [0,1]^d} |\widehat{\mu}_n(x, \mathbf{a}) - \mu_0(x, \mathbf{a})|^2 = o_{\mathbb{P}}\left((na_n)^{-1}\right)$$

where we use Hoeffding decomposition and the inequality holds with probability one.

To complete the proof, it suffices to show that there is a sequence of random variables $A'_n = O_{\mathbb{P}}(1)$ such that for $V \in (0, a_n \delta]$,

$$\sqrt{n} \sup_{|v| < V} \left| \mathbb{E} \left[\widehat{\Gamma}_n(\mathsf{x} + v) - \widehat{\Gamma}_n(\mathsf{x}) - \overline{\Gamma}_n(\mathsf{x} + v) + \overline{\Gamma}_n(\mathsf{x}) \middle| \widetilde{\mathfrak{Z}}_n \right] \right| \le A_n' V.$$

With $f_{\mathbf{A}}(\mathbf{a})$ denoting the Lebesgue density of \mathbf{A} ,

$$\mathbb{E}\left[\widehat{\Gamma}_{n}(\mathbf{x}+v)-\widehat{\Gamma}_{n}(\mathbf{x})\middle|\widetilde{\mathfrak{Z}}_{n}\right] \\
= \int \int \left(\mathbb{1}(u \leq \mathbf{x}+v)-\mathbb{1}(u \leq \mathbf{x})\right) \frac{g_{0}(u,\mathbf{a})}{\widehat{g}_{n}(u,\mathbf{a})} \left[\mu_{0}(u,\mathbf{a})-\widehat{\mu}_{n}(u,\mathbf{a})\right] f_{0}(u) du f_{\mathbf{A}}(\mathbf{a}) d\mathbf{a} \\
+ \int \left(\mathbb{1}(u \leq \mathbf{x}+v)-\mathbb{1}(u \leq \mathbf{x})\right) \int \widehat{\mu}_{n}(u,\mathbf{a}) f_{\mathbf{A}}(\mathbf{a}) d\mathbf{a} du \\
\mathbb{E}\left[\overline{\Gamma}_{n}(\mathbf{x}+v)-\overline{\Gamma}_{n}(\mathbf{x})\middle|\widetilde{\mathfrak{Z}}_{n}\right] \\
= \int \left(\mathbb{1}(u \leq \mathbf{x}+v)-\mathbb{1}(u \leq \mathbf{x})\right) \int \mu_{0}(u,\mathbf{a}) f_{\mathbf{A}}(\mathbf{a}) d\mathbf{a} f_{0}(u) du.$$

Then, for $v \ge 0$.

$$\begin{split} &\mathbb{E}\left[\widehat{\Gamma}_n(\mathbf{x}+v) - \widehat{\Gamma}_n(\mathbf{x}) - \bar{\Gamma}_n(\mathbf{x}+v) + \bar{\Gamma}_n(\mathbf{x}) \middle| \widetilde{\mathfrak{Z}}_n \right] \\ &= \int_{\mathbf{x} < u < \mathbf{x}+v} \left[\widehat{\mu}_n(u, \mathbf{a}) - \mu_0(u, \mathbf{a}) \right] \left[1 - \frac{g_0(u, \mathbf{a})}{\widehat{g}_n(u, \mathbf{a})} \right] f_0(u) f_{\mathbf{A}}(\mathbf{a}) d(u, \mathbf{a}). \end{split}$$

A similar expression holds for v < 0. Then, for some fixed C > 0,

$$\left| \mathbb{E} \left[\widehat{\Gamma}_n(\mathbf{x} + v) - \widehat{\Gamma}_n(\mathbf{x}) - \overline{\Gamma}_n(\mathbf{x} + v) + \overline{\Gamma}_n(\mathbf{x}) \middle| \widetilde{\mathbf{3}}_n \right] \right|$$

$$\leq C|v| \sup_{|x-\mathbf{x}| \leq |v|} \sup_{\mathbf{a} \in [0,1]^d} |\widehat{\mu}_n(x, \mathbf{a}) - \mu_0(x, \mathbf{a})| \sup_{|x-\mathbf{x}| \leq |v|} \sup_{\mathbf{a} \in [0,1]^d} |\widehat{g}_n(x, \mathbf{a}) - g_0(x, \mathbf{a})|$$

and the desired result follows from (SA.19).

Remark. Using (SA.18), one can verify the first part of Assumption D i.e., $\widehat{G}_{x,n}(1;\eta_n) = O_{\mathbb{P}}(1)$. The second part is easy to verify using standard empirical process theory arguments.

SA.3. Additional example: monotone density function with conditionally independent right-censoring. We consider the problem of estimating the density of a non-negative, continuously distributed random variable with censoring. We use the same notation as in Section 5.1 of the main paper. Relative to Section 5.1, we consider the additional complication of censoring being informative about the "survival time" X i.e., $X \not\perp C$. With covariates **A**, we consider the setting of censoring at random: $X \perp C \mid A$. See van der Laan and Robins (2003); Zeng (2004) and references therein for existing analysis of this problem. We have

$$\gamma_0(x; \mathbf{Z}) = F_0(x|\mathbf{A}) + S_0(x|\mathbf{A}) \left[\frac{\Delta \mathbb{1}(\check{X} \le x)}{S_0(\check{X}|\mathbf{A})G_0(\check{X}|\mathbf{A})} - \int_0^{\check{X} \wedge x} \frac{\Lambda_0(du|\mathbf{A})}{S_0(u|\mathbf{A})G_0(u|\mathbf{A})} \right]$$

where $F_0(x|A) = 1 - S_0(x|A)$, $S_0(x|\mathbf{A}) = \mathbb{P}[X > x|\mathbf{A}]$, $G_0(c|\mathbf{A}) = \mathbb{P}[C > c|\mathbf{A}]$, and $\Lambda_0(x|\mathbf{A}) = \int_0^x \frac{f_0(u|\mathbf{A})}{S_0(u|\mathbf{A})} du$ with f_0 being the Lebesgue density of X. Denote by $\widehat{S}_n(\cdot|\cdot)$, $\widehat{G}_n(\cdot|\cdot)$, $\widehat{\Lambda}_n(\cdot|\cdot)$ preliminary estimates of S_0, G_0, Λ_0 , respectively.

Assumption SA.3. Let \mathfrak{S}_n , \mathfrak{G}_n , \mathfrak{L}_n be sequences of function classes that contain $S_0(\cdot|\cdot)$, $G_0(\cdot|\cdot)$, respectively.

- (i) x is in the interior of $I = [0, u_0]$, $X \perp C | A$, and $\theta_0 = f_0$ satisfies Assumption (A2).
- (ii) There exist $c, c_1, c_2 > 0$, $\rho_{\gamma} \in (0, 2)$ such that for $n \geq 1$, for any $S \in \mathfrak{S}_n$, $G \in \mathfrak{G}_n$, and $\Lambda \in \mathfrak{L}_n$, the following hold: $\log N_U(\varepsilon, \{S(x|\cdot) : x \in I\}) \leq c\varepsilon^{-\rho_{\gamma}}$ for $\varepsilon \in (0, 1)$, where N_U is as defined in Section 4.2 of the main paper, and $c_1 \leq S(x|\mathbf{A}) \leq c_2$, $c_1 \leq G(x|\mathbf{A}) \leq c_2$ for $x \in I$, and $\Lambda(u_0|\mathbf{A}) \leq c_2$ with probability one.
- (iii) There exist $\delta > 0$, $\beta_{\gamma} \in [1/2,2)$ such that for $V \in (0,2\delta]$, $\sqrt{na_n} \sup_{|v| \leq V} |\widehat{\Gamma}_n(\mathsf{x} + v) \widehat{\Gamma}_n(\mathsf{x})| \leq o_{\mathbb{P}}(1) + V^{\beta_{\gamma}} o_{\mathbb{P}}(a_n^{\beta_{\gamma}})$ where $o_{\mathbb{P}}$ terms do not depend on V.
- (iv) With probability approaching one, $\widehat{S}_n \in \mathfrak{S}_n$, $\widehat{G}_n \in \mathfrak{G}_n$, $\widehat{\Lambda}_n \in \mathfrak{L}_n$. For $(\widehat{h}_n, h_0) \in \{(\widehat{S}_n, S_0), (\widehat{G}_n, G_0), (\widehat{\Lambda}_n, \Lambda_0)\}$,

$$a_n \frac{1}{n} \sum_{i=1}^n \sup_{x \in I} |\widehat{h}_n(x|\mathbf{A}_i) - h_0(x|\mathbf{A}_i)|^2 = o_{\mathbb{P}}(1).$$

(v) The conditional distribution of X given \mathbf{A} has bounded Lebesgue density $f_{X|\mathbf{A}}$, $\mathbb{E}[\frac{f_{X|\mathbf{A}}(\mathbf{x}|\mathbf{A})}{G_0(\mathbf{x}|\mathbf{A})}] > 0$, and there are real-valued functions B, ω such that $\mathbb{E}[B(\mathbf{A})] < \infty$, $\lim_{\eta \downarrow 0} \omega(\eta) = 0$, and for $|x - \mathbf{x}|$ sufficiently small, $|\frac{f_{X|\mathbf{A}}(\mathbf{x}|\mathbf{A})}{S_0(\mathbf{x}|\mathbf{A})G_0(\mathbf{x}|\mathbf{A})} - \frac{f_{X|\mathbf{A}}(\mathbf{x}|\mathbf{A})}{S_0(\mathbf{x}|\mathbf{A})G_0(\mathbf{x}|\mathbf{A})}| \leq \omega(|x - \mathbf{x}|)B(\mathbf{A})$.

The condition (iii) is high-level, and there are a few different approaches to verify them. See Westling and Carone (2020) for details.

COROLLARY SA-1. Under Assumption SA.3, Assumptions A and B hold with

$$\widehat{\Gamma}_n(x) = \frac{1}{n} \sum_{i=1}^n \widehat{\gamma}_n(x; \mathbf{Z}_i), \quad \widehat{\Gamma}_n^*(x) = \frac{1}{n} \sum_{i=1}^n W_{i,n} \widehat{\gamma}_n(x; \mathbf{Z}_i)$$

$$\widehat{\gamma}_n(x; \mathbf{Z}) = \widehat{F}_n(x|\mathbf{A}) + \widehat{S}_n(x|\mathbf{A}) \left[\frac{\Delta \mathbb{1}(\check{X} \leq x)}{\widehat{S}_n(\check{X}|\mathbf{A})\widehat{G}_n(\check{X}|\mathbf{A})} - \int_0^{\check{X} \wedge x} \frac{\widehat{\Lambda}_n(du|\mathbf{A})}{\widehat{S}_n(u|\mathbf{A})\widehat{G}_n(u|\mathbf{A})} \right],$$

where $\widehat{F}_n = 1 - \widehat{S}_n$,

$$\widehat{\Phi}_n(x) = \widehat{\Phi}_n^*(x) = x, \quad \widehat{u}_n = \widehat{u}_n^* = u_0,$$

$$C_{\mathsf{x}}(s,t) = \mathbb{E}\left[\frac{f_{X|\mathbf{A}}(\mathsf{x}|\mathbf{A})}{G_0(\mathsf{x}|\mathbf{A})}\right](|s| \wedge |t|)\mathbb{1}(\mathrm{sign}(s) = \mathrm{sign}(t)), \quad \mathcal{D}_{\mathfrak{q}}(\mathsf{x}) = \frac{\partial^{\mathfrak{q}} f_0(\mathsf{x})}{(\mathfrak{q}+1)!}.$$

SA.3.1. *Proof of Corollary SA-1*. In this example, $\widehat{\Phi}_n(x) = \widehat{\Phi}_n^*(x) = x = \Phi_0(x)$. Assumptions A and (E1)-(E2) follow from the hypothesis. (E3): In this example, $\widetilde{\Gamma}_n = \widehat{\Gamma}_n$. For $x \in I$,

$$\widehat{\gamma}_{n}(x; \mathbf{Z}) - \gamma_{0}(x; \mathbf{Z}) = \widehat{F}_{n}(x|\mathbf{A}) - F_{0}(x|\mathbf{A})$$

$$+ \left[\widehat{S}_{n}(x|\mathbf{A}) - S_{0}(x|\mathbf{A})\right] \left[\frac{\Delta \mathbb{1}(\check{X} \leq x)}{\widehat{S}_{n}(\check{X}|\mathbf{A})\widehat{G}_{n}(\check{X}|\mathbf{A})} - \int_{0}^{\check{X} \wedge x} \frac{\widehat{\Lambda}_{n}(du|\mathbf{A})}{\widehat{S}_{n}(u|\mathbf{A})\widehat{G}_{n}(u|\mathbf{A})}\right]$$

$$+ S_{0}(x|\mathbf{A})\Delta \mathbb{1}(\check{X} \leq x) \left[\left(\widehat{S}_{n}(\check{X}|\mathbf{A})\widehat{G}_{n}(\check{X}|\mathbf{A})\right)^{-1} - \left(S_{0}(\check{X}|\mathbf{A})G_{0}(\check{X}|\mathbf{A})\right)^{-1}\right]$$

$$- S_{0}(x|\mathbf{A}) \left[\int_{0}^{\check{X} \wedge x} \frac{\widehat{\Lambda}_{n}(du|\mathbf{A})}{\widehat{S}_{n}(u|\mathbf{A})\widehat{G}_{n}(u|\mathbf{A})} - \int_{0}^{\check{X} \wedge x} \frac{\Lambda_{0}(du|\mathbf{A})}{S_{0}(u|\mathbf{A})G_{0}(u|\mathbf{A})}\right]$$

and

$$\left| \int_{0}^{\check{X} \wedge x} \frac{\widehat{\Lambda}_{n}(du|\mathbf{A})}{\widehat{S}_{n}(u|\mathbf{A})\widehat{G}_{n}(u|\mathbf{A})} - \int_{0}^{\check{X} \wedge x} \frac{\Lambda_{0}(du|\mathbf{A})}{S_{0}(u|\mathbf{A})G_{0}(u|\mathbf{A})} \right|$$

$$\leq \sup_{x' \in I} \left| \left(\widehat{S}_{n}(x'|\mathbf{A})\widehat{G}_{n}(x'|\mathbf{A}) \right)^{-1} - \left(S_{0}(x'|\mathbf{A})G_{0}(x'|\mathbf{A}) \right)^{-1} \right| \widehat{\Lambda}_{n}(u_{0}|\mathbf{A})$$

$$+ \left| \frac{\widehat{\Lambda}_{n}(\check{X} \wedge x|\mathbf{A}) - \Lambda_{0}(\check{X} \wedge x|\mathbf{A})}{S_{0}(\check{X} \wedge x|\mathbf{A})G_{0}(\check{X} \wedge x|\mathbf{A})} \right| + \left| \frac{\widehat{\Lambda}_{n}(0|\mathbf{A}) - \Lambda_{0}(0|\mathbf{A})}{S_{0}(0|\mathbf{A})G_{0}(0|\mathbf{A})} \right|$$

$$+ \left| \int_{0}^{\check{X} \wedge x} [\widehat{\Lambda}_{n}(u|\mathbf{A}) - \Lambda_{0}(u|\mathbf{A})] J_{0}(du|\mathbf{A}) \right|$$

using integration by parts, where $J_0(u|\mathbf{a}) = [S_0(u|\mathbf{a})G_0(u|\mathbf{a})]^{-1}$. Thus, there is a fixed C > 0 such that

$$|\widehat{\gamma}_n(x; \mathbf{Z}) - \gamma_0(x; \mathbf{Z})| \le C \left[\sup_{x \in I} |\widehat{S}_n(x|\mathbf{A}) - S_0(x|\mathbf{A})| + \sup_{x \in I} |\widehat{G}_n(x|\mathbf{A}) - G_0(x|\mathbf{A})| + \sup_{x \in I} |\widehat{\Lambda}_n(x|\mathbf{A}) - \Lambda_0(x|\mathbf{A})| \right],$$

From the hypothesis,

$$\frac{1}{n} \sum_{i=1}^{n} \sup_{x \in I} |\widehat{\gamma}_n(x; \mathbf{Z}_i) - \gamma_0(x; \mathbf{Z}_i)|^2 = o_{\mathbb{P}}(1), \ a_n \frac{1}{n} \sum_{i=1}^{n} \sup_{x \in I_x^{\delta}} |\widehat{\gamma}_n(x; \mathbf{Z}_i) - \gamma_0(x; \mathbf{Z}_i)|^2 = o_{\mathbb{P}}(1)$$

follow.

For uniform covering numbers, it suffices to show that each of $\{S(x|\cdot):x\in I\}$, $\{\mathbb{1}(\cdot\leq x):x\in I\}$, and $\{\int_0^{\cdot\wedge x}\frac{\Lambda(du|\cdot)}{S(u|\cdot)G(u|\cdot)}:x\in I\}$ has an appropriate bound on the uniform covering number by Lemma 5.1 of van der Vaart and van der Laan (2006) (see examples after the lemma). For $\{\int_0^{\cdot\wedge x}\frac{\Lambda(du|\cdot)}{S(u|\cdot)G(u|\cdot)}:x\in I\}$ with $(S,G,\Lambda)\in\mathfrak{S}_n\times\mathfrak{G}_n\times\mathfrak{L}_n$, the mapping $x\mapsto\int_0^x\frac{\Lambda(du|\cdot)}{S(u|\cdot)G(u|\cdot)}$ is monotone (by the non-decreasing property of Λ and $S,G\geq c_1>0$) and Lemma 2.6.18 of van der Vaart and Wellner (1996) implies the desired result.

There is a fixed C > 0 such that for $x \in I_{\times}^{\eta}$,

$$|\gamma_0(x; \mathbf{Z}) - \gamma_0(\mathsf{x}; \mathbf{Z})| \le C \left[|S_0(x|\mathbf{A}) - S_0(\mathsf{x}|\mathbf{A})| + C\Delta |\mathbb{1}(\check{X} \le x) - \mathbb{1}(\check{X} \le \mathsf{x})| \right]$$

and using $1 - S_0(x|\cdot) = \int_0^x f_{X|\mathbf{A}}(u|\cdot)du$ with $f_{X|\mathbf{A}}$ being bounded, we can take

$$\bar{D}_{\gamma}^{\eta}(\mathbf{Z}) = C\Delta \mathbb{1}(\mathsf{x} - \eta \le \check{X} \le \mathsf{x} + \eta) + C\eta,$$

which satisfies the desired bound condition.

(E4) trivially holds since $\widehat{\Phi}_n(x) = \widehat{\Phi}_n^*(x) = x$.

(E5): We have

$$\psi_{\mathsf{x}}(v; \mathbf{Z}) = S_0(\mathsf{x}|\mathbf{A}) \frac{(\mathbb{1}(\check{X} \leq \mathsf{x} + v) - \mathbb{1}(\check{X} \leq \mathsf{x}))\Delta}{S_0(\check{X}|\mathbf{A})G_0(\check{X}|\mathbf{A})} + O(|v|)$$

and the first display follows as in the independent censoring case. For the covariance kernel,

$$\mathbb{E}[\psi_{\mathsf{x}_n}(s\eta_n; \mathbf{Z})\psi_{\mathsf{x}_n}(t\eta_n; \mathbf{Z})] = \mathbb{E}\left[S_0(\mathsf{x}_n|\mathbf{A})^2 \int_{\mathsf{x}_n}^{\mathsf{x}_n + \eta_n(s \wedge t)} \frac{f_{X|\mathbf{A}}(u|\mathbf{A})}{S_0(u|\mathbf{A})^2 G_0(u|\mathbf{A})} du \mathbb{1}(s, t > 0)\right] + \mathbb{E}\left[S_0(\mathsf{x}_n|\mathbf{A})^2 \int_{\mathsf{x}_n + \eta_n(s \vee t)}^{\mathsf{x}_n} \frac{f_{X|\mathbf{A}}(u|\mathbf{A})}{S_0(u|\mathbf{A})^2 G_0(u|\mathbf{A})} du \mathbb{1}(s, t < 0)\right] + O(\eta_n^2)$$

and $\eta_n^{-1}\mathbb{E}[\psi_{\mathsf{x}_n}(s\eta_n;\mathbf{Z})\psi_{\mathsf{x}_n}(t\eta_n;\mathbf{Z})]$ converges to $\mathbb{E}[\frac{f_{X|\mathbf{A}}(\mathsf{x}|\mathbf{A})}{G_0(\mathsf{x}|\mathbf{A})}](|s| \wedge |t|)\mathbb{1}(\mathrm{sign}(s) = \mathrm{sign}(t))$. (B5), (B6), and (B7) hold since in this example, $\widehat{u}_n = \widehat{u}_n^* = u_0$ and $\widehat{\Phi}_n$, $\widehat{\Phi}_n^*$ are the identity map.

SA.4. Additional example: monotone hazard function. Let X be a non-negative random variable, f_0 be its Lebesgue density, and $S_0(x) = \mathbb{P}[X > x]$ be its survival function. We consider the parameter of estimating the hazard function of X, $\theta_0(x) = f_0(x)/S_0(x)$, with possible right-censoring as in the monotone density function example. Observations $\mathbf{Z}_1, \ldots, \mathbf{Z}_n$ come from a random sample of $\mathbf{Z} = (\check{X}, \Delta)'$ where $\check{X} = \min\{X, C\}$ and $\Delta = \mathbb{I}(X \leq C)$, C being a random censoring time. As pointed out by Westling and Carone (2020), with strictly increasing Φ_0 with $\Phi_0(0) = 0$, the function Γ_0 takes the form $\Gamma_0(x) = \int_0^x \frac{f_0(u)}{S_0(u)} \Phi_0(du)$, and by taking $\Phi_0(x) = \int_0^x S_0(u) du$, $\Gamma_0(x) = F_0(x) = \mathbb{P}[X \leq x]$. Since Γ_0 is identical to the monotone density case with the choice $\Phi_0 = \int_0^x S_0(u) du$, we can leverage the analysis for the monotone density. The interval I equals $[0, u_0^{\text{MD}}]$ where u_0^{MD} is u_0 in the monotone density example. The u_0 for the monotone hazard function estimation is $u_0 = \Phi_0(u_0^{\text{MD}})$.

Consider the case of completely random censoring i.e., $X \perp C$. As in the setup for Corollary 2, let $\widehat{S}_n(x)$ be the Kaplan-Meier estimator for $S_0(x) = 1 - F_0(x) = \mathbb{P}[X > x]$, $\widehat{F}_n = 1 - \widehat{S}_n$, and \widehat{G}_n be the Kaplan-Meier estimator for $G_0(x) = \mathbb{P}[C > x]$. Also,

$$\gamma_0(x; \mathbf{Z}) = F_0(x) + S_0(x) \left[\frac{\Delta \mathbb{1}(\check{X} \le x)}{S_0(\check{X})G_0(\check{X})} - \int_0^{\check{X} \land x} \frac{\Lambda_0(du)}{S_0(u)G_0(u)} \right]$$

and $\phi_0(x; \mathbf{Z}) = x - \int_0^x \gamma_0(u; \mathbf{Z}) du$.

COROLLARY SA-2. Suppose that the hypothesis of Corollary 2 and Assumption BW hold. Then, Assumptions A and B hold with

$$\begin{split} \widehat{\Gamma}_n(x) &= 1 - \widehat{S}_n(x), \quad \widehat{\Gamma}_n^*(x) = \frac{1}{n} \sum_{i=1}^n W_{i,n} \widehat{\gamma}_n(x; \mathbf{Z}_i), \\ \widehat{\gamma}_n(x; \mathbf{Z}) &= \widehat{F}_n(x) + \widehat{S}_n(x) \left[\frac{\Delta \mathbb{1}(\check{X} \leq x)}{\widehat{S}_n(\check{X}) \widehat{G}_n(\check{X})} - \int_0^{\check{X} \wedge x} \frac{\widehat{\Lambda}_n(du)}{\widehat{S}_n(u) \widehat{G}_n(u)} \right], \\ \widehat{\Phi}_n(x) &= \int_0^x \widehat{F}_n(u) du, \quad \widehat{\Phi}_n^*(x) = \int_0^x [1 - \widehat{\Gamma}_n^*(u)] du = \frac{1}{n} \sum_{i=1}^n W_{i,n} \widehat{\phi}_n(x; \mathbf{Z}_i), \\ \widehat{\phi}_n(x; \mathbf{Z}) &= x - \int_0^x \widehat{\gamma}_n(u; \mathbf{Z}) du, \quad \widehat{u}_n = \widehat{u}_n^* = \widehat{\Phi}_n(u_0^{\text{MD}}), \\ \mathcal{C}_{\mathsf{X}}(s,t) &= \frac{f_0(\mathsf{X})}{G_0(\mathsf{X})} (|s| \wedge |t|) \mathbb{1}(\mathrm{sign}(s) = \mathrm{sign}(t)), \quad \mathcal{D}_{\mathfrak{q}}(\mathsf{X}) = \frac{S_0(\mathsf{X}) \partial^{\mathfrak{q}} f_0(\mathsf{X})}{(\mathfrak{q}+1)!}. \end{split}$$

SA.4.1. *Proof of Corollary SA-2*. We use the same $\widehat{\gamma}_n$ function and assumptions as in the monotone density setting. Also, the covariance kernels are the same as in the monotone density case. We focus on (E4)-(E5) and (B5)-(B7). (E4): Since

$$\widehat{\phi}_n(x; \mathbf{Z}) - \phi_0(x; \mathbf{Z}) = -\int_0^x [\widehat{\gamma}_n(u; \mathbf{Z}) - \gamma_0(u; \mathbf{Z})] du,$$

$$\frac{1}{n} \sum_{i=1}^{n} \sup_{x \in I} |\widehat{\phi}_n(x; \mathbf{Z}_i) - \phi_0(x; \mathbf{Z}_i)|^2 = o_{\mathbb{P}}(1), \ a_n \frac{1}{n} \sum_{i=1}^{n} \sup_{x \in I_{\delta}^{\delta}} |\widehat{\phi}_n(x; \mathbf{Z}_i) - \phi_0(x; \mathbf{Z}_i)|^2 = o_{\mathbb{P}}(1)$$

follow from $a_n \frac{1}{n} \sum_{i=1}^n \sup_{x \in I} |\widehat{\gamma}_n(x; \mathbf{Z}_i) - \gamma_0(x; \mathbf{Z}_i)|^2$, which was verified in Section SA.1.12. To check $\sup_{x \in I} |\widehat{\Phi}_n(x) - \Phi_0(x)| = o_{\mathbb{P}}(1)$, $\sup_{x \in I} |\frac{1}{n} \sum_{i=1}^n \phi_0(x; \mathbf{Z}_i) - \Phi_0(x)| = o_{\mathbb{P}}(1)$ follows from Glivenko-Cantelli, and

$$\sup_{x \in I} \left| \frac{1}{n} \sum_{i=1}^{n} \left[\widehat{\phi}_n(x; \mathbf{Z}_i) - \phi_0(x; \mathbf{Z}_i) \right] \right| \le \frac{1}{n} \sum_{i=1}^{n} \sup_{x \in I} \left| \widehat{\gamma}_n(x; \mathbf{Z}_i) - \gamma_0(x; \mathbf{Z}_i) \right| = o_{\mathbb{P}}(1),$$

where the last equality follows from $\frac{1}{n}\sum_{i=1}^n\sup_{x\in I}|\widehat{\gamma}_n(x;\mathbf{Z}_i)-\gamma_0(x;\mathbf{Z}_i)|^2=o_{\mathbb{P}}(1)$. Now $\sup_{x\in I}|\widehat{\Phi}_n(x)-\Phi_0(x)|=o_{\mathbb{P}}(1)$ follows by the triangle inequality.

For $|v| \leq V$

$$\begin{split} \left|\widehat{\Phi}_n(\mathbf{x}+v) - \widehat{\Phi}_n(\mathbf{x}) - \bar{\Phi}_n(\mathbf{x}+v) + \bar{\Phi}_n(\mathbf{x})\right| &= \left|\int_{\mathbf{x}-v}^{\mathbf{x}+v} [\widehat{\Gamma}_n(u) - \bar{\Gamma}_n(u)] du\right| \\ &\leq 2|V| \Big(\sup_{|v| \leq |V|} \left|\widehat{\Gamma}_n(\mathbf{x}+v) - \widehat{\Gamma}_n(\mathbf{x}) - \bar{\Gamma}_n(\mathbf{x}+v) + \bar{\Gamma}_n(\mathbf{x})\right| + |\widehat{\Gamma}_n(\mathbf{x}) - \bar{\Gamma}_n(\mathbf{x})| \Big) \end{split}$$

Using the argument in Section SA.1.12, we can bound $\sup_{|v|<|V|} |\widehat{\Gamma}_n(x+v) - \widehat{\Gamma}_n(x) - \overline{\Gamma}_n(x+v)|$ $|v| + \bar{\Gamma}_n(x)|$. Then, $\sqrt{n}|\hat{\Gamma}_n(x) - \bar{\Gamma}_n(x)| = O_{\mathbb{P}}(1)$ implies

$$\sqrt{na_n} \sup_{|v| \le V} \left| \widehat{\Phi}_n(\mathbf{x} + v) - \widehat{\Phi}_n(\mathbf{x}) - \bar{\Phi}_n(\mathbf{x} + v) + \bar{\Phi}_n(\mathbf{x}) \right| \le o_{\mathbb{P}}(1) + VO_{\mathbb{P}}\left(\sqrt{a_n}\right)$$

uniformly over $V \in (0, 2\delta]$. Theorem 1 of Lo and Singh (1986) implies $\sqrt{na_n} \sup_{x \in I} |\check{\Phi}_n(x) - \hat{\Phi}_n(x)|$ $\check{\Phi}_n(\mathsf{x}) - \bar{\Phi}_n(x) + \bar{\Phi}_n(\mathsf{x})| = o_{\mathbb{P}}(1).$

The conditions on the uniform covering number hold because γ_0 and $\hat{\gamma}_n$ are bounded (for $\widehat{\gamma}_n$, with probability approaching one) and thus $|\phi_0(x_1; \mathbf{Z}) - \phi_0(x_2; \mathbf{Z})| \le C|x_1 - x_2|$ and $|\widehat{\phi}_n(x_1; \mathbf{Z}) - \widehat{\phi}_n(x_2; \mathbf{Z})| \leq C|x_1 - x_2|$ with probability approaching one. By this Lipschitz property, the condition on $\bar{D}^{\eta}_{\phi}(\mathbf{Z})$ also holds.

(E5): Let $\psi_{\mathbf{x}}^{\mathtt{MD}}(v; \mathbf{Z}) = \gamma_0(\mathbf{x} + v; \mathbf{Z}) - \gamma_0(\mathbf{x}; \mathbf{Z}) - \theta_0(\mathbf{x})v$ be the $\psi_{\mathbf{x}}$ function for the monotone density. Then, for x sufficiently close to \times and |v| small enough,

$$\psi_x(v; \mathbf{z}) = \gamma_0(x + v; \mathbf{z}) - \gamma_0(x; \mathbf{z}) - \theta_0(x) [\phi_0(x + v; \mathbf{z}) - \phi_0(x; \mathbf{z})]$$
$$= \psi_x^{\text{MD}}(v; \mathbf{Z}) + \theta_0(x) \int_x^{x+v} \gamma_0(u; \mathbf{Z}) du = \psi_x^{\text{MD}}(v; \mathbf{Z}) + O(|v|).$$

Then, the same argument as in the monotone density case implies the desired result.

(B5) follows from consistency of $\widehat{\Phi}_n$ and $\widehat{\Phi}_n^*$.
(B6): $\widehat{\Phi}_n(x) = \int_0^x \widehat{F}_n(u) du$, $\widehat{\Phi}_n^*(x) = \int_0^x 1 - \widehat{\Gamma}_n^*(u) du$ are non-negative since $\widehat{F}_n \geq 0$ and $1-\widehat{\Gamma}_n^* \ge 0$ with probability approaching one. This property also implies the non-decreasing property as $\widehat{\Phi}_n$, $\widehat{\Phi}_n^*$ are integrals. The continuity property also follows from the integral representation. By definition, $\widehat{\Phi}_n(0) = 0 = \widehat{\Phi}_n^*(0)$ and $\widehat{\Phi}_n(u_0^{\text{MD}}) = \widehat{u}_n = \widehat{u}_n^* = \widehat{\Phi}_n^*(u_0^{\text{MD}})$ with $I = [0, u_0^{\text{MD}}]$. The closedness of the range follows from continuity and I being a compact interval.

(B7) follows from continuity of $\widehat{\Phi}_n$ and $\widehat{\Phi}_n^*$.

SA.5. Additional example: distribution function estimation with current status data.

We consider the problem of estimating the cdf of X at x, $\theta_0(x) = F_0(x)$. Observations $\mathbf{Z}_1, \dots, \mathbf{Z}_n$ come from a random sample of $\mathbf{Z} = (\Delta, C, \mathbf{A}')'$ where $\Delta = \mathbb{1}(X \leq C), C$ is a random censoring time, and A is a vector of covariates. In this example, we do not observe $X = X \wedge C$. Instead, we observe the censoring time and whether the observation was censored. This setup is often referred to as current status data. Let $H_0(x) = \mathbb{P}[C \le x]$ be the cdf of C. We can use $\Gamma_0(x) = \int_0^x F_0(u) H_0(du)$ and $\Phi_0(x) = H_0(x)$. The interval I is the support of X and $u_0 = 1$. We also assume H_0 admits a Lebesgue density h_0 . The structure of the estimation problem turns out to be identical to the one for the monotone regression example, and we can leverage the common structure.

SA.5.1. Independent right-censoring. First we consider the case of completely at random censoring $X \perp C$. See Groeneboom and Wellner (1992) for existing analysis. In this exaple, we do not use covariates A. We set $\gamma_0(x; \mathbf{Z}) = \Delta \mathbb{1}(C \leq x)$ and $\phi_0(x; \mathbf{Z}) = \mathbb{1}(C \leq x)$. Note that if the notation is mapped by $(\Delta, C) \leftrightarrow (Y, X)$, then these functions are identical to those of the classical monotone regression problem (Corollary 3). Thus, the following result is identical to Corollary 3, up to notation and some changes due to boundedness of Δ .

COROLLARY SA-3. Let $\varepsilon = \Delta - \mathbb{E}[\Delta|C]$ and \times be an interior point of I. Suppose that Assumption BW holds, $\theta_0 = F_0$ satisfies Assumption (A2), the cdf $\Phi_0 = H_0$ satisfies Assumption (A3), and $\sigma_0^2(x) = \mathbb{E}[\varepsilon^2|C=x]$ is continuous and positive at \times . Then Assumptions A and B hold with

$$\widehat{\Gamma}_n(x) = \frac{1}{n} \sum_{i=1}^n \Delta \mathbb{1}(C \le x), \quad \widehat{\Gamma}_n^*(x) = \frac{1}{n} \sum_{i=1}^n W_{i,n} \Delta \mathbb{1}(C \le x),$$

$$\widehat{\Phi}_n(x) = \frac{1}{n} \sum_{i=1}^n \mathbb{1}(C \le x), \quad \widehat{\Phi}_n^*(x) = \frac{1}{n} \sum_{i=1}^n W_{i,n} \mathbb{1}(C \le x), \quad \widehat{u}_n = \widehat{u}_n^* = 1,$$

$$\mathcal{C}_{\mathsf{x}}(s,t) = h_0(\mathsf{x})\sigma_0^2(\mathsf{x})(|s| \wedge |t|)\mathbb{1}(\mathrm{sign}(s) = \mathrm{sign}(t)), \qquad \mathcal{D}_{\mathfrak{q}}(\mathsf{x}) = \frac{h_0(\mathsf{x})\partial^{\mathfrak{q}}F_0(\mathsf{x})}{(\mathfrak{q}+1)!}.$$

SA.5.2. Conditionally independent right-censoring. We consider the case where right-censoring is conditionally independent i.e., $X \perp C | \mathbf{A}$. van der Vaart and van der Laan (2006) analyzed this example as well as settings with time-varying covariates. We are focusing on time-invariant covariates. Define $F_0(C,\mathbf{A}) = \mathbb{E}[\Delta|C,\mathbf{A}]$ and $g_0(C,\mathbf{A}) = \frac{h_{C|\mathbf{A}}(C|\mathbf{A})}{h_0(C)}$ where $h_{C|\mathbf{A}}$ is a conditional Lebesgue density of C given \mathbf{A} and h_0 is a Lebesgue density of C. Let $\widehat{F}_n(c,\mathbf{a})$ and $\widehat{g}_n(c,\mathbf{a})$ be preliminary estimators for $F_0(c,\mathbf{a})$ and $g_0(c,\mathbf{a})$, respectively. Identical to the censoring completely at random case, with appropriate changes in the notation (i.e., $(\Delta,C) \leftrightarrow (Y,X)$), the setup is equivalent to that of the monotone regression with covariates.

Assumption SA.5.2. Let $\varepsilon = \Delta - \mathbb{E}[\Delta|C, \mathbf{A}]$, $\sigma_0^2(C, \mathbf{A}) = \mathbb{E}[\varepsilon^2|C, \mathbf{A}]$, and $\delta > 0$ be some fixed number.

- (i) $\mathbb{E}\left[\frac{\sigma_0^2(\mathsf{x},\mathbf{A})}{q_0(\mathsf{x},\mathbf{A})}\right] > 0.$
- (ii) $h_{C|A}$ is bounded and g_0 is bounded away from zero.
- (iii) There exist random variables $A_n = o_{\mathbb{P}}(1)$ and $B_n = O_{\mathbb{P}}(a_n^{1/2})$ such that

$$\sqrt{na_n} \sup_{|v| \le V} |\widehat{\Gamma}_n(\mathsf{x} + v) - \widehat{\Gamma}_n(\mathsf{x}) - \Gamma_0(\mathsf{x} + v) + \Gamma_0(\mathsf{x})| \le A_n + B_n V, \qquad V \in (0, 2\delta].$$

In addition,

$$\frac{a_n}{n} \sum_{i=1}^n |\widehat{F}_n(C_i, \mathbf{A}_i) - F_0(C_i, \mathbf{A}_i)|^2 = o_{\mathbb{P}}(1),$$

$$\frac{a_n}{n^2} \sum_{i=1}^n \sum_{j=1}^n |\widehat{F}_n(C_i, \mathbf{A}_j) - F_0(C_i, \mathbf{A}_j)|^2 = o_{\mathbb{P}}(1),$$

and

$$\frac{a_n}{n} \sum_{i=1}^n |\widehat{g}_n(C_i, \mathbf{A}_i) - g_0(C_i, \mathbf{A}_i)|^2 = o_{\mathbb{P}}(1).$$

(iv) $\mathbb{E}[\bar{F}(\mathbf{A})^2] < \infty$, where

$$\bar{F}(\mathbf{A}) = \sup_{|c-c'|<\delta} \frac{|F_0(c, \mathbf{A}) - F_0(c', \mathbf{A})|}{|c-c'|}.$$

(v) $\mathbb{E}[\bar{\sigma}(\mathbf{A})^2] < \infty$, where, for some function ω with $\lim_{\eta \downarrow 0} \omega(\eta) = 0$,

$$\left| \frac{\sigma_0^2(x, \mathbf{A}) h_0(x)}{g_0(x, \mathbf{A})} - \frac{\sigma_0^2(\mathbf{x}, \mathbf{A}) h_0(x)}{g_0(\mathbf{x}, \mathbf{A})} \right| \le \omega(|x - \mathbf{x}|) \bar{\sigma}(\mathbf{A}), \qquad x \in I_{\mathbf{x}}^{\delta}.$$

Note that $|\varepsilon| \leq 1$.

COROLLARY SA-4. Suppose that \times is in the interior of I, θ_0 satisfies (A2), Φ_0 satisfies (A3), and Assumption BW holds. If Assumption Assumption SA.5.2 holds, Assumptions A and B hold with

$$\widehat{\Gamma}_n(x) = \frac{1}{n} \sum_{i=1}^n \widehat{\gamma}_n(x; \mathbf{Z}_i), \quad \widehat{\Gamma}_n^*(x) = \frac{1}{n} \sum_{i=1}^n W_{i,n} \widehat{\gamma}_n(x; \mathbf{Z}_i),$$

$$\widehat{\gamma}_n(x; \mathbf{Z}) = \mathbb{1}(C \le x) \left[\frac{\Delta - \widehat{F}_n(C, \mathbf{A})}{\widehat{g}_n(C, \mathbf{A})} + \frac{1}{n} \sum_{i=1}^n \widehat{F}_n(C, \mathbf{A}_i) \right],$$

$$\widehat{\Phi}_n(x) = \frac{1}{n} \sum_{i=1}^n \mathbb{1}(C_i \le x), \quad \widehat{\Phi}_n^*(x) = \frac{1}{n} \sum_{i=1}^n W_{i,n} \mathbb{1}(C_i \le x), \quad \widehat{u}_n = \widehat{u}_n^* = 1,$$

$$\mathcal{C}_{\mathsf{x}}(s,t) = h_0(\mathsf{x}) \mathbb{E}\left[\frac{\sigma_0^2(\mathsf{x},\mathbf{A})}{g_0(\mathsf{x},\mathbf{A})}\right] (|s| \wedge |t|) \mathbb{1}(\mathrm{sign}(s) = \mathrm{sign}(t)), \qquad \mathcal{D}_{\mathfrak{q}}(\mathsf{x}) = \frac{h_0(\mathsf{x}) \partial^{\mathfrak{q}} F_0(\mathsf{x})}{(\mathfrak{q}+1)!}.$$

SA.5.3. *Proof of Corollaries SA-3 and SA-4*. As noted above, by mapping the notation $(\Delta, C) \leftrightarrow (Y, X)$, the arguments in Sections SA.1.13 and SA.1.14 directly apply to the current status estimators.

SA.6. Rule-of-thumb step size selection. Here we develop a rule-of-thumb procedure to choose a step size for the bias-reduced numerical derivative estimator in the context of isotonic regression without covariates. Specifically, we consider the numerical derivative estimator

$$\widetilde{\mathcal{D}}_{j,n}^{\mathrm{BR}}(\mathbf{x}) = \epsilon_n^{-(j+1)} \sum_{k=1}^{\underline{\mathfrak{s}}+1} \lambda_j^{\mathrm{BR}}(k) [\widehat{\Upsilon}_n(\mathbf{x} + c_k \epsilon_n) - \widehat{\Upsilon}_n(\mathbf{x})]$$

with $\underline{\mathfrak{s}} = 3$, $c_1 = 1$, $c_2 = -1$, $c_3 = 2$, $c_4 = -2$. Then,

$$\lambda_1^{\rm BR}(1) = \frac{2}{3} = \lambda_1^{\rm BR}(2), \quad \lambda_1^{\rm BR}(3) = -\frac{1}{24} = \lambda_1^{\rm BR}(4),$$

$$\lambda_3^{\rm BR}(1) = -\frac{1}{6} = \lambda_3^{\rm BR}(2), \quad \lambda_3^{\rm BR}(3) = \frac{1}{24} = \lambda_3^{\rm BR}(4).$$

We use the (asymptotic) MSE-optimal step size discussed in the main paper. See also SA.1.7. Yet, with the choice of c_k 's, part of the bias constant $\sum_{k=1}^{\underline{s}+1} \lambda_j^{\mathrm{BR}}(k) c_k^{\underline{s}+2}$ equals zero, and we need to turn to the next leading term of the bias, which is

$$\epsilon_n^{\underline{\mathfrak{s}}+2-j} \frac{\partial^{\underline{\mathfrak{s}}+3} \Upsilon_0(\mathsf{x})}{(\underline{\mathfrak{s}}+3)!} \sum_{k=1}^{\underline{\mathfrak{s}}+1} \lambda_j^{\mathrm{BR}}(k) c_k^{\underline{\mathfrak{s}}+3}.$$

Then, letting $\mathsf{B}_j^{\mathrm{BR}}(\mathsf{x}) = rac{\partial^6 \Upsilon_0(\mathsf{x})}{6!} \sum_{k=1}^4 \lambda_j^{\mathrm{BR}}(k) c_k^6$, the MSE-optimal step size is

$$\epsilon_{j,n}^{\rm BR} = \left(\frac{(2j+1)\mathsf{V}_j^{\rm BR}(\mathsf{x})}{2(5-j)\mathsf{B}_j^{\rm BR}(\mathsf{x})^2}\right)^{1/11} n^{-1/11}.$$

The bias and variance constants depend on unknown features of the data generating process. Specifically, $\mathsf{B}_j^{\mathsf{BR}}(\mathsf{x})$ depends on the regression function θ_0 , the Lebesgue density of X, and their derivatives at $X=\mathsf{x}$ while $\mathsf{V}_j^{\mathsf{BR}}(\mathsf{x})$ is determined by the density of X and the conditional variance of the regression error $\varepsilon=Y-\theta_0(X)$ at $X=\mathsf{x}$. To implement the construction of the step size, we posit a simple parametric model:

$$\mathbb{E}[Y|X] = \gamma_0 + \sum_{k=1}^{5} \gamma_k (X - \mathsf{x}_0)^k, \quad X \sim \text{Normal}(\mu, \sigma^2)$$

where $\{\gamma_0, \gamma_1, \gamma_2, \gamma_4, \gamma_4, \gamma_5, \mu, \sigma\}$ are parameters to be estimated. Once we estimate the parameters of this reference model, we can construct a rule-of-thumb step size $\epsilon_{j,n}^{\text{ROT}}$ by replacing $\mathsf{B}_j^{\text{BR}}(\mathsf{x})$ and $\mathsf{V}_j^{\text{BR}}(\mathsf{x})$ with their estimates. Note that although the bias and variance constant estimators may not be consistent for the true $\mathsf{B}_j^{\text{BR}}(\mathsf{x})$ and $\mathsf{V}_j^{\text{BR}}(\mathsf{x})$, the rate of $\epsilon_{j,n}^{\text{ROT}}$ is MSE-optimal, and the numerical derivative estimator converges to $\mathcal{D}_j(\mathsf{x})$ sufficiently fast to satisfy Equation (11) in the main paper.

REFERENCES

BURKE, M. D., CSÖRGŐ, S. and HORVÁTH, L. (1988). A Correction to and Improvement of 'Strong Approximations of Some Biometric Estimates under Random Censorship'. *Probability Theory and Related Fields* **79** 51–57

CATTANEO, M. D., FARRELL, M. and FENG, Y. (2020). Large Sample Properties of Partitioning-Based Series Estimators. *Annals of Statistics* **48** 1718-1741.

CATTANEO, M. D., JANSSON, M. and MA, X. (2020). Simple Local Polynomial Density Estimators. *Journal of the American Statistical Association* 115 1449-1455.

CATTANEO, M. D., CHANDAK, R., JANSSON, M. and MA, X. (2024). Boundary Adaptive Local Polynomial Conditional Density Estimators. *Bernoulli*.

Cox, G. (2022). A Generalized Argmax Theorem with Applications. arXiv:2209.08793.

GROENEBOOM, P. and WELLNER, J. A. (1992). Information Bounds and Nonparametric Maximum Likelihood Estimation. Birkhäuser Basel.

JAIN, N. C. and MARCUS, M. B. (1978). Continuity of Subgaussian Processes. In Probability in Banach Spaces. Advances in Probability and Related Topics, Volume 4 81–196. Dekker.

KIM, J. and POLLARD, D. (1990). Cube Root Asymptotics. Annals of Statistics 18 191-219.

KOSOROK, M. R. (2008). Bootstrapping the Grenander Estimator. In *Beyond Parametrics in Interdisciplinary Research: Festschrift in Honor of Professor Pranab K. Sen* 282–292. Institute of Mathematical Statistics.

Lo, S.-H. and SINGH, K. (1986). The Product-Limit Estimator and the Bootstrap: Some Asymptotic Representations. *Probability Theory and Related Fields* **71** 455–465.

POLLARD, D. (1989). Asymptotics via Empirical Processes. Statistical Science 4 341–354.

SEN, B., BANERJEE, M. and WOODROOFE, M. (2010). Inconsistency of Bootstrap: The Grenander Estimator. *Annals of Statistics* **38** 1953–1977.

VAN DER LAAN, M. J. and ROBINS, J. M. (2003). *Unified Methods for Censored longitudinal Data and Causality*. Springer Science & Business Media.

VAN DER VAART, A. W. and VAN DER LAAN, M. J. (2006). Estimating a Survival Distribution with Current Status Data and High-Dimensional Covariates. *International Journal of Biostatistics* 2.

VAN DER VAART, A. W. and WELLNER, J. A. (1996). Weak Convergence and Empirical Processes. Springer.

WESTLING, T. and CARONE, M. (2020). A Unified Study of Nonparametric Inference for Monotone Functions. Annals of Statistics 48 1001–1024.

WESTLING, T., GILBERT, P. and CARONE, M. (2020). Causal isotonic regression. *Journal of the Royal Statistical Society, Series B* 82 719–747.

ZENG, D. (2004). Estimating Marginal Survival Function by Adjusting for Dependent Censoring using Many Covariates. *Annals of Statistics* **32** 1533–1555.