Optimal transport and Wasserstein DISTANCES FOR CAUSAL MODELS*

Patrick Cheridito Department of Mathematics ETH Zurich, Switzerland Stephan Eckstein Department of Mathematics University of Tübingen, Germany

ABSTRACT

In this paper, we introduce a variant of optimal transport adapted to the causal structure given by an underlying directed graph G. Different graph structures lead to different specifications of the optimal transport problem. For instance, a fully connected graph yields standard optimal transport, a linear graph structure corresponds to causal optimal transport between the distributions of two discrete-time stochastic processes, and an empty graph leads to a notion of optimal transport related to CO-OT, Gromov–Wasserstein distances and factored OT. We derive different characterizations of G-causal transport plans and introduce Wasserstein distances between causal models that respect the underlying graph structure. We show that average treatment effects are continuous with respect to G-causal Wasserstein distances and small perturbations of structural causal models lead to small deviations in G-causal Wasserstein distance. We also introduce an interpolation between causal models based on G-causal Wasserstein distance and compare it to standard Wasserstein interpolation.

Keywords: Average treatment effect, causality, directed graphs, optimal transport, Wasserstein distance

1 Introduction

Originally, optimal transport problems were introduced by Monge [43] and, in a more general form, by Kantorovich [33] to study the most efficient way to transport and allocate resources. But in addition to this immediate application, optimal transport theory has also lead to the the notion of Wasserstein distance [34, 65, 66], which defines a metric on probability distributions. Over the years, optimal transport has found applications in different areas of economics [16, 26, 53], probability theory [55, 56], statistics [25, 27, 44], differential geometry [22, 24, 62], robust optimization [10, 42, 70], machine learning and data science [4, 19, 50, 61], just to name a few. At the same time, various variants and extensions of optimal transport have emerged, like multi-marginal versions [2, 23, 47], optimal transport with additional constraints [9, 15, 14, 20, 36, 45], optimal transport between measures of different masses [17, 60], relaxations [12, 39] and regularizations [18, 40]. On the other hand, causal research examines causal relationships between different events; see e.g. [30, 48, 49, 59]. In probabilistic the-

^{*}We are grateful to Beatrice Acciaio, Julio Backhoff-Veraguas, Daniel Bartl, Mathias Beiglböck, Songyan Hou, Nicolai Meinshausen, Alexander Neitz, Gudmund Pammer and Drago Plečko for interesting discussions and helpful comments.

ories of causality, the underlying causal structure is typically described by a directed graph, leading to graphical causal models [11].

In this paper we introduce a version of optimal transport that is adapted to the causal structure given by an underlying directed graph and investigate corresponding Wasserstein distances between different causal models. Our main goal is to obtain a version of the optimal transport problem which yields topological and geometric properties consistent with the structure of the underlying directed graph. We build on existing literature studying this question for distributions of stochastic processes such as [7, 6, 8, 37]. This string of literature demonstrates different advantages of incorporating the temporal structure into the definition of optimal transport distances between distributions of stochastic processes compared to using standard Wasserstein distances, which are agnostic to this structure. For instance, it improves the stability of dynamic optimization problems that are unstable under standard Wasserstein distances (see [7]), and it leads to geometric structures compatible with certain properties of stochastic processes (e.g. it has been shown in [8] that sets of martingales are geodesically convex). In this paper, we show that it has similar benefits if a suitably adapted concept of optimal transport is used for general causal structures. For such a version of optimal transport, the structure of the underlying graph G determines the exact specification of the optimal transport problem. A fully connected graph corresponds to standard optimal transport, which consists in finding an optimal transport plan transforming a given distribution into another one. Missing edges translate into additional constraints, which increase the G-causal Wasserstein distance and therefore, lead to a finer topology on the corresponding set of causal models. For instance, a linear graph requires transport plans to respect the temporal structure. This results in causal transport problems between distributions of stochastic processes, which have been studied by e.g. [7, 6, 8, 37]. An empty graph restricts transport plans the most and is related to CO-OT problems [63], the Gromov-Wasserstein distance [41] and factored OT [64]. For a general directed graph G, we call the corresponding transport plans G-causal and demonstrate their suitability for downstream tasks which depend on the information structure given by G. For instance, we show that while one of the most fundamental causal statistics, the average treatment effect, is not continuous in the standard Wasserstein metric, it becomes Lipschitz continuous if distances between distributions are measured with a G-causal Wasserstein distance resulting from G-causal optimal transport. Furthermore, we demonstrate that under suitable assumptions, G-causal Wasserstein interpolation between two G-compatible distributions preserves G-compatibility, which in general, is not true for standard Wasserstein interpolation.

The remainder of this paper is organized as follows: In Section 2 we introduce the setup and provide our definition of G-causal transport plans together with the corresponding optimal transport problems in Definitions 2.1 and 2.2. In Section 2.4 we showcase the relation to existing concepts of optimal transport from the literature. In Section 3 we focus on G-causal transport maps for directed acyclic graphs. We derive alternative characterizations of G-causal transport plans in Theorem 3.4 and Corollary 3.5 and study the structure of sets of G-causal transport plans in Proposition 3.6. In Section 4 we introduce Wasserstein distances between causal models that respect the structure of the underlying graph G. We show that they satisfy all properties of a metric except for the triangle inequality in Proposition 4.3. In Section 4.1, we prove that, under suitable assumptions, average treatment effects, which estimate the causal effect of a treatment or intervention, are Lipschitz continuous with respect to G-causal Wasserstein distance under a change of the underlying probability model while they are not continuous for standard Wasserstein distances. As a consequence, bounds on the error resulting from estimating average treatment effects from data can be obtained by controlling the G-causal Wasserstein estimation error. On the other hand, we show in Section 4.2 that small perturbations of a structural causal model correspond to small deviations in G-causal Wasserstein distance. In particular,

convergence of structural models implies convergence of the corresponding probability models with respect to G-causal Wasserstein distance. Finally, in Section 4.3 we study a Wasserstein interpolation between causal models that respects the causal structure and compare it to standard Wasserstein interpolation.

2 Notation and definitions

2.1 Causal structures described by directed graphs

We consider a finite set $V = \{1, ..., n\}$ for some $n \in \mathbb{N}$, endowed with a causal structure given by a set of *directed edges* $E \subseteq \{(i, j) \in V^2 : i \neq j\}$. This turns G = (V, E) into a *directed graph*. For $j \in V$, we denote by pa_j the parents of j, given by

$$pa_i := \{i \in V : (i, j) \in E\},\$$

and for a subset $A \subseteq V$, we define

$$\operatorname{pa}_A := \bigcup_{j \in A} \operatorname{pa}_j.$$

We order both sets pa_i and pa_A according to the natural ranking of the elements of $V = \{1, \ldots, n\}$.

2.2 Spaces and measures

We consider two product spaces of the form $\mathcal{X} = \mathcal{X}_1 \times \cdots \times \mathcal{X}_n$ and $\mathcal{Y} = \mathcal{Y}_1 \times \cdots \times \mathcal{Y}_n$ for nonempty Polish spaces $\mathcal{X}_i, \mathcal{Y}_i, i = 1, ..., n$. We endow \mathcal{X}, \mathcal{Y} and $\mathcal{X} \times \mathcal{Y}$ with the product topologies, which makes them again Polish; see e.g. [13, Chapter IX, §6.1, Proposition I]. By $\mathcal{P}(\mathcal{X}), \mathcal{P}(\mathcal{Y})$ and $\mathcal{P}(\mathcal{X} \times \mathcal{Y})$ we denote the set of all Borel probability measures on \mathcal{X}, \mathcal{Y} and $\mathcal{X} \times \mathcal{Y}$, respectively. If we write $X \sim \mu$ for a measure $\mu \in \mathcal{P}(\mathcal{X})$, we mean that X is an \mathcal{X} -valued random variable with distribution μ defined on some underlying probability space. Moreover, we use the notation $X_{1:i}$ to denote the tuple (X_1, \ldots, X_i) for $i \in V = \{1, \ldots, n\}$, and similarly, X_A for any ordered subset A of $V = \{1, \ldots, n\}$. $X_{1:0}$ as well as X_{pa_i} in cases where pa_i is empty, will be understood as empty tuple.

2.3 Transport plans and *G*-causal couplings

A transport map between two probability measures $\mu \in \mathcal{P}(\mathcal{X})$ and $\nu \in \mathcal{P}(\mathcal{Y})$ in the sense of Monge [43] is a measurable mapping $T: \mathcal{X} \to \mathcal{Y}$ such that $T_{\#}\mu = \nu$, where $T_{\#}\mu$ denotes the push-forward $\mu \circ T^{-1}$ of μ along T. A Monge transport map does not always exist, e.g. if μ is a Dirac measure and ν is not. Therefore, Kantorovich [33] extended the set of transport plans by considering couplings between μ and ν , which are probability measures $\pi \in \mathcal{P}(\mathcal{X} \times \mathcal{Y})$ with marginals μ and ν . The set $\Pi(\mu, \nu)$ of all couplings between μ and ν always contains the product measure $\mu \otimes \nu$, and therefore, is non-empty. Moreover, for every Monge transport map $T: \mathcal{X} \to \mathcal{Y}$ between μ and ν , the distribution of the mapping $(\mathrm{id}, T): \mathcal{X} \to \mathcal{X} \times \mathcal{Y}$ under μ is a coupling in $\Pi(\mu, \nu)$. On the other hand, every coupling $\pi \in \Pi(\mu, \nu)$ can be realized as a randomized transport map. Indeed, since \mathcal{Y} is Polish, π admits a disintegration of the form $\pi(dx, dy) = \mu(dx)\pi(dy \mid x)$, where $\pi(dy \mid x)$ is a regular version of the conditional distribution of $y \in \mathcal{Y}$ given $x \in \mathcal{X}$ with respect to π ; see e.g. [32, Theorem 3.4]. Therefore, one obtains from the kernel representation [32, Lemma 4.22] that π can be realized as the distribution of a pair of random variables (X, Y) such that $X \sim \mu$ and Y is of the form Y = g(X, U) for a measurable map $g: \mathcal{X} \times \mathbb{R} \to \mathcal{Y}$ and an \mathbb{R} -valued random variable U defined on the same probability space as and independent of X.

To motivate our definition of G-causal couplings, we first consider the case of Monge transport maps $T: \mathcal{X} = \mathcal{X}_1 \times \cdots \times \mathcal{X}_n \to \mathcal{Y} = \mathcal{Y}_1 \times \ldots \mathcal{Y}_n$. Intuitively, one might want to define G-causality of T by saying that T is G-causal if it is of the form $T = (T_1, \ldots, T_n)$, where

$$T_i(x) = g_i(x_i, x_{pa_i}) \quad \text{for all } x \in \mathcal{X} \text{ and } i = 1, \dots, n,$$
(2.1)

for measurable mappings $g_i: \mathcal{X}_i \times \mathcal{X}_{pa_i} \to \mathcal{Y}_i, i = 1, ..., n$. More generally, one could call $T = (T_1, \ldots, T_n)$ G-causal if

$$T_i(x) = g_i(x_i, x_{\operatorname{pa}_i}, T_{\operatorname{pa}_i}(x)) \quad \text{for all } x \in \mathcal{X} \text{ and } i = 1, \dots, n,$$
(2.2)

for measurable mappings $g_i: \mathcal{X}_i \times \mathcal{X}_{pa_i} \times \mathcal{Y}_{pa_i} \to \mathcal{Y}_i$, i = 1, ..., n. It is easy to see that in the following three cases, conditions (2.1) and (2.2) are equivalent:

- if the graph G = (V, E) is fully connected, that is, $E = \{(i, j) \in V^2 : i \neq j\}$,
- if the graph G = (V, E) is empty, that is, $E = \emptyset$, or
- if the graph G = (V, E) has the linear structure $E = \{(i, j) \in V^2 : i < j\}$.

On the other hand,

• if the graph G = (V, E) has the Markovian structure $E = \{(i - 1, i) : 2 \le i \le n\},\$

(2.2) is slightly more flexible than (2.1). In general, (2.1) is too restrictive for our purposes. For instance, a generalization of (2.1) to randomized transport maps would still not guarantee the existence of G-causal couplings between any two probability measures μ and ν that are compatible with respect to a directed acyclic graph G in the sense of Definition 3.1 below. Therefore, we use (2.2) to define G-causal Monge transport maps, which leads to the following general definition.

Definition 2.1 (*G*-causal couplings). Let G = (V, E) be a directed graph. We call a coupling $\pi \in \Pi(\mu, \nu)$ between two measures $\mu \in \mathcal{P}(\mathcal{X})$ and $\nu \in \mathcal{P}(\mathcal{Y})$ *G*-causal, if there exists a pair of random variables $(X, Y) \sim \pi$ together with measurable mappings

$$g_i: \mathcal{X}_i \times \mathcal{X}_{\mathrm{pa}_i} \times \mathcal{Y}_{\mathrm{pa}_i} \times \mathbb{R} \to \mathcal{Y}_i, \quad i = 1, \dots, n,$$

and \mathbb{R} -valued random variables U_1, \ldots, U_n defined on the same probability space as (X, Y) such that X, U_1, \ldots, U_n are independent and

$$Y_i = g_i(X_i, X_{pa_i}, Y_{pa_i}, U_i)$$
 for all $i = 1, ..., n.$ (2.3)

We call π *G*-bicausal, if π and $\tilde{\pi} \sim (Y, X)$ are *G*-causal. We denote the set of *G*-causal and *G*-bicausal couplings between μ and ν by $\Pi_G(\mu, \nu)$ and $\Pi_G^{bc}(\mu, \nu)$, respectively.

 $\Pi_G^{\rm bc}(\mu,\nu)$ is a symmetrized version of $\Pi_G(\mu,\nu)$ consisting of transport plans that are *G*-causal in both directions.

In Theorem 3.4 and Corollary 3.5 below we give alternative characterizations of G-causal and G-bicausal transport plans in the case where G is a directed acyclic graph. Proposition 3.6 shows that $\Pi_G(\mu,\nu)$ and $\Pi_G^{bc}(\mu,\nu)$ are non-empty for all probability measures μ and ν respecting the structure of G. The following transport problems can be defined for arbitrary directed graphs G.

Definition 2.2 (*G*-causal optimal transport). Let G = (V, E) be a directed graph. For $\mu \in \mathcal{P}(\mathcal{X})$, $\nu \in \mathcal{P}(\mathcal{Y})$ and a measurable cost function $c : \mathcal{X} \times \mathcal{Y} \to [0, \infty)$, we define the *G*-causal and *G*-bicausal optimal transport problems as

$$\inf_{\pi \in \Pi_G(\mu,\nu)} \int_{\mathcal{X} \times \mathcal{Y}} c \, d\pi \quad \text{and} \quad \inf_{\pi \in \Pi_G^{\mathrm{bc}}(\mu,\nu)} \int_{\mathcal{X} \times \mathcal{Y}} c \, d\pi, \tag{2.4}$$

respectively.

2.4 Special cases and related concepts

2.4.1 Fully connected graphs and standard optimal transport

If *E* equals the set $\{(i, j) \in V^2 : i \neq j\}$ of all possible edges, a coupling $\pi \in \Pi(\mu, \nu)$ between two measures $\mu \in \mathcal{P}(\mathcal{X})$ and $\nu \in \mathcal{P}(\mathcal{Y})$ satisfies Definition 2.1 if and only if there exists a pair of random variables $(X, Y) \sim \pi$ together with measurable mappings

$$g_i: \mathcal{X} \times \mathcal{Y}_{V \setminus \{i\}} \times \mathbb{R} \to \mathcal{Y}_i, \quad i = 1, \dots, n,$$

and \mathbb{R} -valued random variables U_1, \ldots, U_n defined on the same probability space as (X, Y) such that X, U_1, \ldots, U_n are independent and

$$Y_i = g_i(X, Y_{V \setminus \{i\}}, U_i)$$
 for all $i = 1, ..., n.$ (2.5)

Since \mathcal{X} and \mathcal{Y}_i , i = 1, ..., n, are Polish spaces, any coupling $\pi \in \Pi(\mu, \nu)$ admits an iterated disintegration of the form

$$\pi(dx, dy) = \mu(dx)\pi(dy_1 \mid x)\pi(dy_2 \mid x, y_1)\cdots\pi(dy_n \mid x, y_{1:n-1})$$

for regular conditional distributions $\pi(dy_1 \mid x)$ and $\pi(dy_i \mid x, y_{1:i-1})$, i = 2, ..., n; see e.g. [32, Theorem 3.4]. So, it follows from the kernel representation [32, Lemma 4.22] that π has a realization of the form (2.5), which shows that

$$\Pi_G(\mu,\nu) = \Pi_G^{\rm bc}(\mu,\nu) = \Pi(\mu,\nu).$$

That is, the two optimal transport problems in (2.4) reduce to standard optimal transport.

2.4.2 Empty graphs, CO-OT and Gromov–Wasserstein distances

If the graph has no edges, that is, $E = \emptyset$, then a coupling $\pi \in \Pi(\mu, \nu)$ satisfies Definition 2.1 if and only if there exists a pair of random variables $(X, Y) \sim \pi$ together with measurable mappings

$$g_i: \mathcal{X}_i \times \mathbb{R} \to \mathcal{Y}_i, \quad i = 1, \dots, n,$$

and \mathbb{R} -valued random variables U_1, \ldots, U_n defined on the same probability space as (X, Y) such that X, U_1, \ldots, U_n are independent and

$$Y_i = g_i(X_i, U_i)$$
 for all $i = 1, ..., n.$ (2.6)

Since $\mathcal{X}_i, \mathcal{Y}_i, i = 1, ..., n$, are Polish spaces, every measure $\pi_i \in \Pi(\mu_i, \nu_i)$ can be disintegrated as

$$\pi_i(dx_i, dy_i) = \mu_i(dx_i)\pi_i(dy_i \mid x_i)$$

for a regular conditional distribution $\pi_i(dy_i \mid x_i)$; see e.g. [32, Theorem 3.4]. Therefore, we obtain from the kernel representation [32, Lemma 4.22] that π_i has a representation of the form (2.6). In particular, if $\mu = \mu_1 \otimes \cdots \otimes \mu_n$ and $\nu = \nu_1 \otimes \cdots \otimes \nu_n$ are product measures, one has

$$\Pi_G(\mu,\nu) = \Pi_G^{\mathrm{bc}}(\mu,\nu) = \left\{ \pi_1 \otimes \cdots \otimes \pi_n : \pi_i \in \Pi(\mu_i,\nu_i) \text{ for all } i = 1, \dots, n \right\}.$$

This set of couplings is related to CO-OT problems [63], the Gromov–Wasserstein distance [41] and factored OT [64]. CO-OT problems also optimize over couplings of marginals of product form. They aim to simultaneously optimize two transport plans between features and labels of two datasets. Therefore, n = 2 is used. The Gromov–Wasserstein distance measures the discrepancy between two spaces. It considers n = 2 and marginal measures of the special form $\mu = \mu_1 \otimes \mu_1$, $\nu = \nu_1 \otimes \nu_1$. Factored OT is a generalization of CO-OT to a multi-marginal setting based on couplings of the form

 $\{\pi_1 \otimes \cdots \otimes \pi_n : \pi_i \in \Pi(\mu_{i,1}, \ldots, \mu_{i,n_i}) \text{ for } i = 1, \ldots, n\}.$

2.4.3 Linear graph structure and transport between distributions of stochastic processes

If the graph has the linear structure $E = \{(i, j) \in V^2 : i < j\}$, then a coupling $\pi \in \Pi(\mu, \nu)$ satisfies Definition 2.1 if and only if there exists a pair of random variables $(X, Y) \sim \pi$ together with measurable mappings

$$g_1: \mathcal{X}_1 \times \mathbb{R} \to \mathcal{Y}_1, \quad g_i: \mathcal{X}_{1:i} \times \mathcal{Y}_{1:i-1} \times \mathbb{R} \to \mathcal{Y}_i, \quad i = 2, \dots, n,$$

and \mathbb{R} -valued random variables U_1, \ldots, U_n defined on the same probability space as (X, Y) such that X, U_1, \ldots, U_n are independent and

$$Y_1 = g_1(X_1, U_1), \quad Y_i = g_i(X_{1:i}, Y_{1:i-1}, U_i), \quad i = 2, \dots, n.$$

If μ and ν are the distributions of two discrete-time stochastic processes with indices i = 1, ..., n, the coupling π can be viewed as a transport plan that respects the temporal structure. Such couplings, sometimes treated in more generality through the use of filtrations, have been studied, for instance, by [8, 37, 51] and have applications in robust finance [1, 6, 7, 52] as well as machine learning [68, 69]. For relations to different topologies on spaces of distributions of stochastic processes, see e.g. [3, 5, 29, 46]. The resulting optimal transport problems are called causal, bicausal or adapted.

In relation to this literature, we mention that interesting extensions of the G-causal optimal transport problem may be possible by using a more flexible definition of G-causal couplings through more precise encodings of the information structures, similarly to how [8] extends the space of stochastic processes using filtrations.

2.4.4 Markovian graph structure

If the graph G = (V, E) has the Markovian structure $E = \{(i - 1, i) : 2 \le i \le n\}$, a coupling $\pi \in \Pi(\mu, \nu)$ satisfies Definition 2.1 if and only if if there exists a pair of random variables $(X, Y) \sim \pi$ together with measurable mappings

$$g_1: \mathcal{X}_1 \times \mathbb{R} \to \mathcal{Y}_1, \quad g_i: \mathcal{X}_{i-1:i} \times \mathcal{Y}_{i-1} \times \mathbb{R} \to \mathcal{Y}_i, \quad i = 2, \dots, n,$$

and \mathbb{R} -valued random variables U_1, \ldots, U_n defined on the same probability space as (X, Y) such that X, U_1, \ldots, U_n are independent and

$$Y_1 = g_1(X_1, U_1), \quad Y_i = g_i(X_{i-1:i}, Y_{i-1}, U_i), \quad i = 2, \dots, n.$$

2.4.5 Related concepts

Different variants of optimal transport have been proposed in the literature that do not exactly fit into the framework of this paper but are related. [54] proposes a coupling between two populations, such as male and female job applicants, along a given graph with the goal of achieving a fair embedding of one group into the other. [35] proposes a GAN framework based on structural assumptions similar to (2.3). [21] studies collections of couplings between conditional distributions.

3 *G*-causal couplings for DAGs

In causal modelling it is often assumed that the causal structure is given by a *directed acyclic graph* (DAG), which is a directed graph with no *cycles*. By reordering the vertices, a DAG can always assumed to be *sorted*, which means that it only contains edges (i, j) such that i < j; see e.g. [11, 38, 48]. If G is a DAG, it is possible to provide alternative characterizations of G-causal and G-bicausal couplings, from which simple properties of the sets of all such couplings can be derived.

We start by recalling the notions of conditional independence and compatibility of a probability measure with a given DAG. For more details, we refer to [11, 38, 48]. For random variables S, W, Zdefined on the same probability space $(\Omega, \mathcal{F}, \mathbb{P})$ with values in measurable spaces S, W, Z, we say Sis independent of W given Z and write $S \perp\!\!\!\perp_Z W$ if

$$\mathbb{P}[S \in A, W \in B \mid Z] = \mathbb{P}[S \in A \mid Z] \mathbb{P}[W \in B \mid Z] \quad \mathbb{P}\text{-a.s.}$$
(3.1)

for all measurable subsets $A \subseteq S$ and $B \subseteq W$, which is equivalent to

$$\mathbb{P}[S \in A \mid W, Z] = \mathbb{P}[S \in A \mid Z] \quad \mathbb{P}\text{-a.s.}$$
(3.2)

for all measurable subsets $A \subseteq S$. Since the conditional probabilities in (3.1)–(3.2) can all be written as g(Z) or h(W, Z) for measurable functions $g: Z \to [0, 1]$ or $h: W \times Z \to [0, 1]$ (see e.g. [32, Lemma 1.14]), (3.1)–(3.2) can equivalently be formulated as

$$\rho[A \times B \times \mathcal{Z} \mid \mathcal{F}_{\mathcal{Z}}] = \rho[A \times \mathcal{W} \times \mathcal{Z} \mid \mathcal{F}_{\mathcal{Z}}] \rho[\mathcal{V} \times B \times \mathcal{Z} \mid \mathcal{F}_{\mathcal{Z}}] \quad \rho\text{-a.s}$$

and

$$\rho[A \times \mathcal{W} \times \mathcal{Z} \mid \mathcal{F}_{\mathcal{W}, \mathcal{Z}}] = \rho[A \times \mathcal{W} \times \mathcal{Z} \mid \mathcal{F}_{\mathcal{Z}}] \quad \rho\text{-a.s.},$$

where ρ is the distribution of (S, W, Z) and $\mathcal{F}_{\mathcal{Z}}$, $\mathcal{F}_{W,Z}$ are the σ -algebras generated by the projections from $S \times W \times Z$ to Z and $W \times Z$, respectively. This shows that conditions (3.1)–(3.2) only depend on the distribution of (S, W, Z).

If S and W are Polish spaces equipped with their Borel σ -algebras, (S, W) has a regular conditional distribution $\rho(ds, dw \mid z)$ given Z = z with respect to ρ (see e.g. [32, Theorem 3.4]). In this case, conditions (3.1)–(3.2) can equivalently be written as

$$\rho(ds, dw \mid z) = \rho(ds \mid z)\rho(dw \mid z) \quad \text{for } Z_{\#}\mathbb{P}\text{-almost all } z \in \mathcal{Z}.$$

Definition 3.1 (G-compatible probability measures). A measure $\mu \in \mathcal{P}(\mathcal{X})$ is said to be compatible with a sorted DAG G = (V, E) if any of the following three equivalent conditions hold. We denote the set of G-compatible measures in $\mathcal{P}(\mathcal{X})$ by $\mathcal{P}_G(\mathcal{X})$. (i) There exists a random variable $X \sim \mu$ together with measurable functions $f_i : \mathcal{X}_{pa_i} \times \mathbb{R} \to \mathcal{X}_i$, i = 1, ..., n, and independent \mathbb{R} -valued random variables $U_1, ..., U_n$ such that

$$X_i = f_i(X_{\text{pa}_i}, U_i) \quad \text{for all } i = 1, \dots, n.$$
(3.3)

(ii) For every random variable $X \sim \mu$, one has

$$X_i \perp \!\!\!\perp_{X_{\text{pa}_i}} X_{1:i-1}$$
 for all $i = 2, \dots, n.$ (3.4)

(iii) The measure μ can be disintegrated as

$$\mu(dx_1,\ldots,dx_n) = \prod_{i=1}^n \mu\left(dx_i \mid x_{\mathsf{pa}_i}\right) \tag{3.5}$$

for $\mu(dx_1 \mid x_{pa_1}) = \mu(dx_1)$ and regular conditional distributions $\mu(dx_i \mid x_{pa_i})$, i = 2, ..., n, called *causal mechanisms* in causal modelling.

Remark 3.2. The equivalence of conditions (i)–(iii) in Definition 3.1 follows from the definitions (3.1)–(3.2) of conditional independence together with some well-known properties of probability measures on Polish spaces. For the sake of completeness, we give a short proof:

(i) \Leftrightarrow (ii): If (i) holds, there exists an \mathcal{X} -valued random variable $X = (X_1, \ldots, X_n) \sim \mu$ such that the X_i are of the form $X_i = f_i(X_{pa_i}, U_i)$ for measurable mappings $f_i: \mathcal{X}_{pa_i} \times \mathbb{R} \to \mathcal{X}_i$ and independent \mathbb{R} -valued random variables $U_i, i = 1, \ldots, n$, defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. It follows that $\mathbb{P}[X_i \in A \mid X_{1:i-1}] = \mathbb{P}[X_i \in A \mid X_{pa_i}] \mathbb{P}$ -a.s. for all measurable subsets $A \subseteq \mathcal{X}_i$, which, by (3.2), shows that (ii) is satisfied.

On the other hand, if (ii) holds, one obtains from [32, Proposition 8.20] that, possibly after extending the underlying probability space, the X_i have representations of the form $X_i = f_i(X_{pa_i}, U_i)$ \mathbb{P} -a.s. for measurable functions $f_i: \mathcal{X}_{pa_i} \times \mathbb{R} \to \mathcal{X}_i$ and U(0, 1)-distributed random variables U_i satisfying $U_i \perp X_{1:i-1}$ for all i = 2, ..., n. So X is of the form (3.3), except that the representation only holds \mathbb{P} -a.s. and the random variables $U_1, ..., U_n$ are not necessarily independent. But this can easily be remedied by considering independent U(0, 1)-distributed random variables $\tilde{U}_1, ..., \tilde{U}_n$ on a new probability space and iteratively defining $\tilde{X}_i = f_i(\tilde{X}_{pa_i}, \tilde{U}_i), i = 1, ..., n$. Then \tilde{X}_1 has the same distribution as X_1 , and for all $i \ge 2$, the conditional distribution of \tilde{X}_i given $\tilde{X}_{1:i-1}$ is equal to the conditional distribution of X_i given $X_{1:i-1}$. So, it follows by induction over i that $\tilde{X} = (\tilde{X}_1, ..., \tilde{X}_n)$ has the same distribution as $X = (X_1, ..., X_n)$ and satisfies condition (i).

(ii) \Leftrightarrow (iii): Since $\mathcal{X}_1, \ldots, \mathcal{X}_n$ are Polish spaces, one obtains from [32, Theorem 3.4] that μ admits an iterated disintegration of the form

$$\mu(dx_1, \dots, dx_n) = \mu(dx_1)\mu(dx_2 \mid x_1) \cdots \mu(dx_n \mid x_{1:n-1})$$

for regular conditional distributions $\mu(dx_i \mid x_{1:i-1})$, i = 2, ..., n, from which it can be seen that (3.4) is equivalent to (3.5).

The following result provides some elementary properties of the set $\mathcal{P}_G(\mathcal{X})$.

Proposition 3.3. Let G = (V, E) be a sorted DAG. Then

- (i) $\mathcal{P}_G(\mathcal{X})$ is non-empty,
- (ii) $\mathcal{P}_G(\mathcal{X})$ is closed in total variation,

- (iii) if \mathcal{X} is a discrete² Polish space, $\mathcal{P}_G(\mathcal{X})$ is weakly closed,
- (iv) but there exist Polish spaces $\mathcal{X} = \mathcal{X}_1 \times \cdots \times \mathcal{X}_n$ and graphs G such that $\mathcal{P}_G(\mathcal{X})$ is not weakly closed.

Proof. According to Definition 3.1.(iii), the product measure $\mu_1 \otimes \cdots \otimes \mu_n$ belongs to $\mathcal{P}_G(\mathcal{X})$ for arbitrary marginal measures $\mu_i \in \mathcal{P}(\mathcal{X}_i)$, i = 1, ..., n. This shows (i).

Moreover, it follows from Lemma A.1 in Appendix A below that Definition 3.1.(ii) is stable under convergence in total variation, which proves (ii).

(iii) follows from (ii) since in case \mathcal{X} is a discrete Polish space, weak convergence is equivalent to convergence in total variation.

To show (iv), we consider the Markovian graph G = (V, E) given by $V = \{1, 2, 3\}$ and $G = \{(1, 2), (2, 3)\}$. Let $\mathcal{X}_1 = \mathcal{X}_2 = \mathcal{X}_3 = [-1, 1]$ and consider the product space $\mathcal{X} = [-1, 1]^3$ with the Borel σ -algebra. The measures $\mu^k = \frac{1}{2}\delta_{(1,1/k,1)} + \frac{1}{2}\delta_{(-1,-1/k,-1)}$, $k \in \mathbb{N}$, belong to $\mathcal{P}_G(\mathcal{X})$ for all $k \in \mathbb{N}$ and weakly converge to the measure $\frac{1}{2}\delta_{(1,0,1)} + \frac{1}{2}\delta_{(-1,0,-1)}$, which does not belong to $\mathcal{P}_G(\mathcal{X})$. This proves (iv).

The following is the main structural result of this paper. It gives different characterizations of G-causal couplings between two measures $\mu \in \mathcal{P}_G(\mathcal{X})$ and $\nu \in \mathcal{P}(\mathcal{Y})$. Note that for a sorted DAG G, one has $\Pi_G(\mu, \nu) \subseteq \Pi_{G_l}(\mu, \nu)$, where G_l is the linear graph from Section 2.4.3, which leads to causal optimal transport between distributions of stochastic processes, for which related results have been shown by e.g. [6, Proposition 2.4].

Theorem 3.4. Let G = (V, E) be a sorted DAG, and consider measures $\mu \in \mathcal{P}_G(\mathcal{X})$, $\nu \in \mathcal{P}(\mathcal{Y})$ and $\pi \in \Pi(\mu, \nu)$. Then, the following are equivalent:

- (i) $\pi \in \Pi_G(\mu, \nu)$,
- (ii) for $(X, Y) \sim \pi$, one has

 $Y_i \perp X_{i,X_{\text{pa}_i},Y_{\text{pa}_i}}(X,Y_{1:i-1})$ for all $i = 1, \ldots, n$, where $Y_{1:0}$ is the empty tuple,

(iii) for $(X, Y) \sim \pi$, one has

$$X_i \perp X_{pa_i}(X_{1:i-1}, Y_{1:i-1})$$
 and $Y_i \perp X_{i, X_{pa_i}, Y_{pa_i}}(X_{1:i}, Y_{1:i-1})$ for all $i = 2, \dots, n$,

(iv) π is jointly *G*-compatible, that is,

$$\pi(dx_1, dy_1, \ldots, dx_n, dy_n) = \prod_{i=1}^n \pi(dx_i, dy_i \mid x_{\operatorname{pa}_i}, y_{\operatorname{pa}_i}),$$

for regular conditional distributions $\pi(dx_i, dy_i \mid x_{pa_i}, y_{pa_i})$, and

 $\pi(dx_i \mid x_{pa_i}, y_{pa_i}) = \mu(dx_i \mid x_{pa_i})$ for all i = 1, ..., n and π -almost all $(x, y) \in \mathcal{X} \times \mathcal{Y}$, where $\mu(dx_i \mid x_{pa_i})$ are regular conditional versions of μ .

²i.e. \mathcal{X} is a countable discrete topological space

Before turning to the proof of Theorem 3.4, we note that it implies the following alternative characterizations of G-bicausal couplings between two measures $\mu \in \mathcal{P}_G(\mathcal{X})$ and $\nu \in \mathcal{P}_G(\mathcal{Y})$. Particularly the characterization in part (iii) of Corollary 3.5 is worth emphasizing, as this characterization is very natural and hence serves as validation of Definition 2.1. The characterization states that, just like the marginals, a G-bicausal coupling also has to be G-compatible. And on top of that, the causal mechanisms of the marginals are coupled by the causal mechanisms of the coupling. In this way, G-bicausal couplings can intuitively be understood as fine-grained couplings, which put the essential building blocks of causal models, the causal mechanisms, into relation.

Corollary 3.5. Let G = (V, E) be a sorted DAG, $\mu \in \mathcal{P}_G(\mathcal{X})$, $\nu \in \mathcal{P}_G(\mathcal{Y})$ and $\pi \in \Pi(\mu, \nu)$. Then, the following are equivalent:

- (i) $\pi \in \Pi^{\mathrm{bc}}_G(\mu, \nu)$,
- (ii) for $(X, Y) \sim \pi$, one has

$$(X_i, Y_i) \perp X_{pa_i}, Y_{pa_i}, (X_{1:i-1}, Y_{1:i-1}), \quad X_i \perp X_{pa_i}, Y_{pa_i} \quad and \quad Y_i \perp Y_{pa_i}, X_{pa_i} \quad for \ all \ i = 2, \dots, n$$

(iii) π is jointly *G*-compatible, that is,

$$\pi(dx_1, dy_1, \dots, dx_n, dy_n) = \prod_{i=1}^n \pi(dx_i, dy_i \mid x_{\operatorname{pa}_i}, y_{\operatorname{pa}_i})$$

for regular conditional distributions $\pi(dx_i, dy_i \mid x_{pa_i}, y_{pa_i})$, and

$$\pi(dx_i, dy_i \mid x_{\operatorname{pa}_i}, y_{\operatorname{pa}_i}) \in \Pi(\mu(dx_i \mid x_{\operatorname{pa}_i}), \nu(dy_i \mid y_{\operatorname{pa}_i}))$$

for all i = 1, ..., n and π -almost all $(x, y) \in \mathcal{X} \times \mathcal{Y}$, where $\mu(dx_i \mid x_{pa_i})$ and $\nu(dy_i \mid y_{pa_i})$ are regular conditional versions of μ and ν , respectively.

Proof of Theorem 3.4. (i) \Rightarrow (ii): If (i) holds, there exists $(X, Y) \sim \pi$ such that the Y_i are of the form $Y_i = g_i(X_i, X_{\text{pa}_i}, Y_{\text{pa}_i}, U_i)$ for measurable functions $g_i \colon \mathcal{X}_i \times \mathcal{X}_{\text{pa}_i} \times \mathcal{Y}_{\text{pa}_i} \times \mathbb{R} \to \mathcal{Y}_i$ and \mathbb{R} -valued random variables U_i , i = 1, ..., n defined on the same probability space as (X, Y) such that X, U_1, \ldots, U_n are independent. It follows that $U_i \perp (X, Y_{1:i-1})$, and therefore, $\mathbb{P}[Y_i \in A \mid X, Y_{1:i-1}] = \mathbb{P}[Y_i \in A \mid X_i, X_{\text{pa}_i}, Y_{\text{pa}_i}] \mathbb{P}$ -a.s. for every *i* and all measurable subsets $A \subseteq \mathcal{Y}_i$, which by (3.2), shows that (ii) is satisfied.

(ii) \Rightarrow (i): If (X, Y) fulfils (ii), one obtains from [32, Proposition 8.20] that, possibly after extending the underlying probability space, the Y_i can be represented as $Y_i = g_i(X_i, X_{\text{pa}_i}, Y_{\text{pa}_i}, U_i)$ \mathbb{P} -a.s. for measurable functions $g_i: \mathcal{X}_i \times \mathcal{X}_{\text{pa}_i} \times \mathcal{Y}_{\text{pa}_i} \times \mathbb{R} \rightarrow \mathcal{Y}_i$ and U(0, 1)-distributed random variables U_i satisfying $U_i \perp (X, Y_{1:i-1})$ for all $i = 1, \ldots, n$. So (X, Y) is of the form (2.3), except that the representation only holds \mathbb{P} -a.s. and the random variables U_1, \ldots, U_n are not necessarily independent. However, by extending the probability space further if necessary, one can assume that there exist U(0, 1)-distributed random variables $\tilde{U}_1, \ldots, \tilde{U}_n$ such that $X, \tilde{U}_1, \ldots, \tilde{U}_n$ are independent. If one iteratively defines $\tilde{Y}_i = g_i(X_i, X_{\text{pa}_i}, \tilde{Y}_{\text{pa}_i}, \tilde{U}_i), i = 1, \ldots, n$, then the conditional distribution of \tilde{Y}_i given $(X_i, X_{\text{pa}_i}, \tilde{Y}_{\text{pa}_i})$ is for all i equal to the conditional distribution of Y_i given $(X_i, X_{\text{pa}_i}, Y_{\text{pa}_i})$. It follows by induction over i that (X, \tilde{Y}) has the same distribution as (X, Y), and it satisfies all requirements of Definition 2.1. So, (i) holds.

(ii) \Rightarrow (iii): The second condition of (iii) is immediate from (ii). In addition, one obtains from (ii) that $Y_j \perp X_j, X_{pa_j}, Y_{pa_j} X_i$ for all $1 \le i, j \le n$. In particular, $X_i \perp X_{1:i-1}, Y_{1:j-1} Y_j$ for all $1 \le j < j$ $i \leq n$, which, by the chain rule [32, Theorem 8.12] yields $X_i \perp I_{X_{1:i-1}} Y_{1:i-1}$. This together with $X_i \perp I_{X_{pa_i}} X_{1:i-1}$, which holds since $\mu \in \mathcal{P}_G(\mu)$, gives $X_i \perp I_{X_{pa_i}} (X_{1:i-1}, Y_{1:i-1})$ for all $i = 2, \ldots, n$, showing that also the first condition of (iii) holds.

(iii) \Rightarrow (ii): The first condition of (iii) implies $X_j \perp \!\!\!\perp_{X_{1:j-1}} Y_{1:i}$, and therefore, $Y_i \perp \!\!\!\perp_{X_{1:j-1},Y_{1:i-1}} X_j$ for all $1 \le i < j \le n$. This, together with the second condition of (iii) and the chain rule [32, Theorem 8.12], yields (ii).

(iii) \Leftrightarrow (iv): Since \mathcal{X}_i , \mathcal{Y}_i , i = 1, ..., n, are Polish spaces, it follows from [32, Theorem 3.4] that π can be disintegrated as

$$\pi(dx_1, dy_1, \dots, dx_n, dy_n) = \prod_{i=1}^n \pi(dx_i \mid x_{1:i-1}, y_{1:i-1}) \pi(dy_i \mid x_{1:i}, y_{1:i-1}),$$

where $x_{1:0}$ and $y_{1:0}$ are empty tuples. Since $\pi \in \Pi(\mu, \nu)$, one has

$$X_i \perp \!\!\!\perp_{X_{\mathsf{pa}_i}} (X_{1:i-1}, Y_{1:i-1}) \quad \Leftrightarrow \quad \pi(dx_i \mid x_{1:i-1}, y_{1:i-1}) = \mu(dx_i \mid x_{\mathsf{pa}_i}) \quad \pi\text{-almost surely}$$

and

 $Y_{i} \perp \!\!\!\perp_{X_{i}, X_{\mathrm{pa}_{i}}, Y_{\mathrm{pa}_{i}}}(X_{1:i}, Y_{1:i-1}) \quad \Leftrightarrow \quad \pi \left(dy_{i} \mid x_{1:i}, y_{1:i-1} \right) = \pi \left(dy_{i} \mid x_{i}, x_{\mathrm{pa}_{i}}, y_{\mathrm{pa}_{i}} \right) \quad \pi\text{-almost surely,}$

which shows that (iii) is equivalent to

$$\pi(dx_1, dy_1, \ldots, dx_n, dy_n) = \prod_{i=1}^n \mu\left(dx_i \mid x_{\operatorname{pa}_i}\right) \pi\left(dy_i \mid x_i, x_{\operatorname{pa}_i}, y_{\operatorname{pa}_i}\right),$$

which, in turn, is equivalent to (iv).

Proof of Corollary 3.5. (i) \Leftrightarrow (iii) follows directly from the equivalence of (i) and (iv) of Theorem 3.4.

(ii) \Leftrightarrow (iii): As in the proof of Theorem 3.4, we obtain from [32, Theorem 3.4] that π can be disintegrated as

$$\pi(dx_1, dy_1, \dots, dx_n, dy_n) = \prod_{i=1}^n \pi(dx_i, dy_i \mid x_{1:i-1}, y_{1:i-1}),$$

where $x_{1:0}$ and $y_{1:0}$ are empty tuples, from which it can be seen that (ii) is equivalent to (iii).

In the following result we derive some elementary properties of the sets $\Pi_G(\mu, \nu)$ and $\Pi_G^{bc}(\mu, \nu)$. **Proposition 3.6.** Let G = (V, E) be a sorted DAG and $\mu \in \mathcal{P}_G(\mathcal{X})$, $\nu \in \mathcal{P}_G(\mathcal{Y})$. Then

- (i) $\Pi_G(\mu,\nu)$ and $\Pi_G^{bc}(\mu,\nu)$ are both non-empty,
- (ii) $\Pi_G(\mu, \nu)$ and $\Pi_G^{bc}(\mu, \nu)$ are closed in total variation,
- (iii) if μ and ν are finitely supported, $\Pi_G(\mu, \nu)$ and $\Pi_G^{bc}(\mu, \nu)$ are weakly closed,
- (iv) but in general, there exist \mathcal{X} , \mathcal{Y} and G such that $\Pi_G(\mu, \nu)$ and $\Pi_G^{bc}(\mu, \nu)$ are not weakly closed.

Proof. Since μ and ν are both compatible with G, they admit a decomposition of the form (3.5). So, it can be seen from Corollary 3.5 that the product measure $\mu \otimes \nu$ belongs to $\Pi_G^{bc}(\mu, \nu)$, which is contained in $\Pi_G(\mu, \nu)$. This shows (i).

By Lemma A.1 in Appendix A, conditions (ii) and (iii) of Theorem 3.4 are stable under convergence in total variation, which implies (ii).

Regarding (iii), if μ and ν have finite supports $S_{\mu} \subseteq \mathcal{X}$ and $S_{\nu} \subseteq \mathcal{Y}$, respectively, every measure $\pi \in \Pi(\mu, \nu)$ is supported by the finite subset $S_{\mu} \times S_{\nu} \subseteq \mathcal{X} \times \mathcal{Y}$. Since weak convergence among such measures is equivalent to convergence in total variation, (iii) follows from (ii).

To show (iv), we consider a Markovian graph G = (V, E) with $V = \{1, 2, 3\}$ and $E = \{(1, 2), (2, 3)\}$. Consider the measures $\mu = \nu = \theta \otimes \eta \otimes \theta$ on $[0, 1]^3$, where $\theta = \frac{1}{2}(\delta_0 + \delta_1)$ and η is the uniform distribution on [0, 1]. We show that $\Pi_G^{\mathrm{bc}}(\mu, \nu)$ is not weakly closed. Then $\Pi_G(\mu, \nu)$ cannot be weakly closed either. Otherwise, $\Pi_G(\nu, \mu)$ and, as a consequence, $\Pi_G^{\mathrm{bc}}(\mu, \nu)$ would also have to be weakly closed. To prove that $\Pi_G^{\mathrm{bc}}(\mu, \nu)$ is not weakly closed, we denote by $\gamma \in \Pi(\eta, \eta)$ the uniform distribution on the diagonal $D = \{(x, x) \in [0, 1]^2 : x \in [0, 1]\}$ and by $\gamma^{1,k}, \gamma^{2,k} \in \Pi(\eta, \eta), k \geq 3$, the uniform distributions on the shifted diagonals $D^{1,k}, D^{2,k}$ given by

$$\begin{split} D^{1,k} &= \left\{ (x,x+1/k) : x \in [0,1-1/k] \right\} \cup \left\{ (x,x+1/k-1) : x \in (1-1/k,1] \right\}, \\ D^{2,k} &= \left\{ (x,x+1-1/k) : x \in [0,1/k] \right\} \cup \left\{ [x,x-1/k) : x \in [1/k,1] \right\}. \end{split}$$

Clearly, $\gamma^{1,k}$ and $\gamma^{2,k}$ both weakly converge to γ for $k \to \infty$. Now, let us define the measures π^k , $k \ge 3$, on $([0,1]^2)^3$ through

$$\pi^{k}(dz_{1}, dz_{2}, dz_{3}) = \pi^{k}_{1}(dz_{1})\pi^{k}_{1,2}(z_{1}, dz_{2})\pi^{k}_{2,3}(z_{2}, dz_{3})$$

for

$$\pi_1^k = \frac{1}{2}(\delta_{(0,0)} + \delta_{(1,1)}), \quad \pi_{1,2}^k(0,0) = \gamma^{1,k}, \quad \pi_{1,2}^k(1,1) = \gamma^{2,k}$$

and

$$\pi_{2,3}^k(z_2) = \begin{cases} \frac{1}{2}(\delta_{(0,0)} + \delta_{(1,1)}) \text{ if } z_2 \in D^{1,k} \\ \frac{1}{2}(\delta_{(0,1)} + \delta_{(1,0)}) \text{ if } z_2 \in D^{2,k}. \end{cases}$$

Then, $\pi^k \in \Pi^{\mathrm{bc}}_G(\mu, \nu)$ for all $k \geq 3$, and for $k \to \infty$, π^k weakly converges to π given by

$$\pi(dz_1, dz_2, dz_3) = \pi_1(dz_1)\pi_{1,2}(dz_2)\pi_{2,3}(z_1, dz_3)$$

for

$$\pi_1 = \frac{1}{2}(\delta_{(0,0)} + \delta_{(1,1)}), \quad \pi_{1,2} = \gamma$$

and

$$\pi_{2,3}(z_1) = \begin{cases} \frac{1}{2}(\delta_{(0,0)} + \delta_{(1,1)}) \text{ if } z_1 = (0,0) \\ \frac{1}{2}(\delta_{(0,1)} + \delta_{(1,0)}) \text{ if } z_1 = (1,1). \end{cases}$$

So π is no longer Markovian, and therefore, $\pi \notin \Pi_G^{bc}(\mu, \nu)$, which proves (iv).

4 *G*-causal Wasserstein distances

In this section, we study Wasserstein distances arising from G-bicausal couplings. We show that they define semimetrics on appropriate spaces of G-compatible probability measures but in general do not

satisfy the triangle inequality. In Section 4.1 we prove that under suitable assumptions, average treatment effects are continuous with respect to G-causal Wasserstein distance, and in Section 4.2 we show that small perturbations of structural causal models lead to small deviations in G-causal Wasserstein distance. In Subsection 4.3 we investigate the interpolation of two G-compatible measures such that the causal structure given by G is respected. In the whole section, we let $\mathcal{X} = \mathcal{Y}$ and denote by $d_{\mathcal{X}}$ a metric $d_{\mathcal{X}}: \mathcal{X} \times \mathcal{X} \to \mathbb{R}_+$ generating³ the given Polish topology on \mathcal{X} .

Remark 4.1 (Standard Wasserstein distances). For a $p \in [1, \infty)$, we denote by $\mathcal{P}_p(\mathcal{X})$ the set of all measures $\mu \in \mathcal{P}(\mathcal{X})$ satisfying

$$\int_{\mathcal{X}} d_{\mathcal{X}}(x, x_0)^p \mu(dx) < \infty \quad \text{for some } x_0 \in \mathcal{X}.$$
(4.1)

Note that (4.1) implies

$$\int_{\mathcal{X}} d_{\mathcal{X}}(x, x'_0)^p \mu(dx) \le 2^{p-1} \left(\int_{\mathcal{X}} d_{\mathcal{X}}(x, x_0)^p \mu(dx) + d_{\mathcal{X}}(x_0, x'_0)^p \right) < \infty \quad \text{for every } x'_0 \in \mathcal{X}.$$

So, if (4.1) holds for one, it holds for all $x_0 \in \mathcal{X}$. The standard *p*-Wasserstein distance $W_p(\mu, \nu)$ between two measures μ and ν in $\mathcal{P}_p(\mathcal{X})$ is given by

$$W_p^p(\mu,\nu) := \inf_{\pi \in \Pi(\mu,\nu)} \int_{\mathcal{X} \times \mathcal{X}} d_{\mathcal{X}}(x,y)^p \, \pi(dx,dy).$$

It is well known (see e.g. [67]) that the infimum is attained and W_p defines a metric on $\mathcal{P}_p(\mathcal{X})$. Moreover, $\Pi(\mu, \nu)$ is a weakly compact subset of $\mathcal{P}(\mathcal{X} \times \mathcal{X})$.

In the following definition we introduce Wasserstein distances based on *G*-bicausal couplings. For causal Wasserstein distances between distributions of stochastic processes, see [5] and the references therein.

Definition 4.2 (*G*-causal Wasserstein distances). Let G = (V, E) be a sorted DAG and $p \in [1, \infty)$. We denote $\mathcal{P}_{G,p}(\mathcal{X}) = \mathcal{P}_G(\mathcal{X}) \cap \mathcal{P}_p(\mathcal{X})$ and define the *p*-th order *G*-causal Wasserstein distance $W_{G,p}(\mu, \nu)$ between two measures μ and ν in $\mathcal{P}_{G,p}(\mathcal{X})$ by

$$W^p_{G,p}(\mu,\nu) := \inf_{\pi \in \Pi^{\mathrm{bc}}_G(\mu,\nu)} \int_{\mathcal{X} \times \mathcal{X}} d_{\mathcal{X}}(x,y)^p \, \pi(dx,dy).$$
(4.2)

The following result provides some elementary properties of G-causal Wasserstein distances. **Proposition 4.3.** Let G = (V, E) be a sorted DAG and $p \in [1, \infty)$. Then

(i) for another directed graph G' = (V, E') with $E' \subseteq E$, one has $W_{G,p} \leq W_{G',p}$, and therefore,

$$\{\nu \in \mathcal{P}(\mathcal{X}) : W_{G',p}(\mu,\nu) \le r\} \subseteq \{\nu \in \mathcal{P}(\mathcal{X}) : W_{G,p}(\mu,\nu) \le r\}$$

for all $\mu \in \mathcal{P}(\mathcal{X})$ and r > 0,

- (ii) $W_{G,p}$ is a semimetric⁴ on $\mathcal{P}_{G,p}(\mathcal{X})$ and
- (iii) if $\mu, \nu \in \mathcal{P}_{G,p}(\mathcal{X})$ are finitely supported, the infimum in (4.2) is attained.

³we assume $d_{\mathcal{X}}$ generates the topology, but \mathcal{X} is not necessarily complete with respect to $d_{\mathcal{X}}$

⁴that is, it has all properties of a metric except the triangle inequality

Proof. (i): If $E' \subseteq E$, one has $\operatorname{pa}_i' \subseteq \operatorname{pa}_i$, $i = 1, \ldots, n$, for the corresponding parent sets. Therefore, it can be seen from Definition 2.1 that $\Pi_{G'}^{\operatorname{bc}}(\mu, \nu) \subseteq \Pi_{G}^{\operatorname{bc}}(\mu, \nu)$ for all $\mu, \nu \in \mathcal{P}(\mathcal{X})$, which implies (i).

(ii) It is clear that $W_{G,p}$ is symmetric and non-negative. Moreover, it follows from Proposition 3.6.(i) that for all $\mu, \nu \in \mathcal{P}_{G,p}$, there exists a coupling $\pi \in \Pi^{\mathrm{bc}}_{G}(\mu, \nu)$. Therefore, one has for any $x_0 \in \mathcal{X}$,

$$\begin{split} W^p_G(\mu,\nu) &\leq \int_{\mathcal{X}\times\mathcal{X}} (d_{\mathcal{X}}(x,y))^p \pi(dx,dy) \leq \int_{\mathcal{X}\times\mathcal{X}} (d_{\mathcal{X}}(x,x_0) + d_{\mathcal{X}}(y,x_0))^p \pi(dx,dy) \\ &\leq 2^{p-1} \left(\int_{\mathcal{X}} (d_{\mathcal{X}}(x,x_0))^p \mu(dx) + \int_{\mathcal{X}} (d_{\mathcal{X}}(x,x_0))^p \nu(dx) \right) < \infty. \end{split}$$

Next, note that for $X \sim \mu \in \mathcal{P}_{G,p}(\mathcal{X})$, the distribution of (X, X) belongs to $\Pi_G^{\mathrm{bc}}(\mu, \mu)$, from which one obtains $W_{G,p}(\mu, \mu) = 0$. Finally, since $\Pi_G^{\mathrm{bc}}(\mu, \nu) \subseteq \Pi(\mu, \nu)$, one has

$$W_{G,p}(\mu,\nu) \ge W_p(\mu,\nu) > 0$$
 if $\mu \ne \nu$,

where the last inequality holds since the standard p-Wasserstein distance W_p is a metric.

(iii) If $\mu, \nu \in \mathcal{P}_{G,p}(\mathcal{X})$ are finitely supported, choose a sequence $(\pi^k)_{k\geq 1}$ in $\Pi_G^{\mathrm{bc}}(\mu, \nu)$ such that

$$\int_{\mathcal{X}} d_{\mathcal{X}}(x,y)^p d\pi^k(dx,dy) \to W^p_{G,p}(\mu,\nu) \quad \text{for } k \to \infty.$$
(4.3)

Since $\Pi(\mu, \nu)$ is a weakly compact subset of $\mathcal{P}(\mathcal{X} \times \mathcal{X})$, there exists a subsequence, again denoted $(\pi^k)_{k\geq 1}$, which weakly converges to a measure $\pi \in \Pi(\mu, \nu)$. By Proposition 3.6.(iii), π belongs to $\Pi_G^{\mathrm{bc}}(\mu, \nu)$, and, by (4.3), it minimizes (4.2).

We show in Appendix B below that in general, $W_{G,p}$ does not satisfy the triangle inequality.

4.1 Continuity of average treatment effects with respect to G-causal Wasserstein distance

The average treatment effect is a measure of the causal effect of a treatment or intervention (cf. [57]). In the following we show that under suitable assumptions, it is Lipschitz continuous with respect to $W_{G,1}$ under a change of the underlying probability model while it is not continuous with respect to the standard Wasserstein distance. I.e., we are interested in the mapping from probability measures to average treatment effect. An important implication of this continuity result is that one can obtain bounds on the error resulting from estimating average treatment effects from data by controlling the *G*-causal Wasserstein estimation error. For recent studies on how average treatment effects can be estimated from data, see e.g. [58, 31].

Let us assume there exist two indices $j, k \in V = \{1, ..., n\}$ such that $j < k, \mathcal{X}_j = \{0, 1\}$ and \mathcal{X}_k is a compact subset of \mathbb{R} . X_j indicates whether a treatment is applied or not, and X_k describes a resulting outcome. Suppose the metric $d_{\mathcal{X}}$ on $\mathcal{X} = \mathcal{X}_1 \times \cdots \times \mathcal{X}_n$ is given by $d_{\mathcal{X}}(x, y) := \sum_{i=1}^n d_{\mathcal{X}_i}(x_i, y_i)$, where $d_{\mathcal{X}_i}(x_i, y_i) = |x_i - y_i|$ for i = j, k. Using Pearl's do-notation (cf. the back-door adjustment [48, Theorem 3.3.2] applied with pa_j), the average treatment effect under a model $\mu \in \mathcal{P}_G(\mathcal{X})$ is given by

$$\psi^{\mu} := \int_{\mathcal{X}_{k}} x_{k} \, \mu(dx_{k} \mid \operatorname{do}(x_{j} = 1)) - \int_{\mathcal{X}_{k}} x_{k} \, \mu(dx_{k} \mid \operatorname{do}(x_{j} = 0)) \\ = \int_{\mathcal{X}_{\operatorname{pa}_{i}}} \int_{\mathcal{X}_{k}} x_{k} \, \mu(dx_{k} \mid x_{j} = 1, x_{\operatorname{pa}_{j}}) \, \mu(dx_{\operatorname{pa}_{j}}) - \int_{\mathcal{X}_{\operatorname{pa}_{i}}} \int_{\mathcal{X}_{k}} x_{k} \, \mu(dx_{k} \mid x_{j} = 0, x_{\operatorname{pa}_{j}}) \, \mu(dx_{\operatorname{pa}_{j}}),$$

$$(4.4)$$

where for the purposes of this paper, the second line serves as definition of the do-notation in the first line. In the following, we fix a constant $\delta > 0$ and consider the set $\mathcal{P}_G^{\delta,j}(\mathcal{X})$ of models $\mu \in \mathcal{P}_G(\mathcal{X})$ such that the propensity score $\mu(x_j = 1 \mid x_{\text{pa}_j})$ satisfies

$$\delta \le \mu(x_j = 1 \mid x_{\operatorname{pa}_j}) \le 1 - \delta$$
 for μ -almost all x_{pa_j}

Then the following holds.

Proposition 4.4. Let G = (V, E) be a sorted DAG, $\delta > 0$ and j < k, \mathcal{X}_j , \mathcal{X}_k , $d_{\mathcal{X}}$ as above. Then there exists a constant $C \ge 0$ such that

$$|\psi^{\mu} - \psi^{\nu}| \le C W_{G,1}(\mu, \nu)$$

for all $\mu, \nu \in \mathcal{P}_{G}^{\delta, j}(\mathcal{X})$.

Proof. Note that

$$\begin{split} &\int_{\mathcal{X}_{k}} x_{k} \, \mu(dx_{k} \mid \operatorname{do}(x_{j} = 1)) \\ &= \int_{\mathcal{X}_{pa_{j}}} \int_{\mathcal{X}_{k}} x_{k} \, \mu(dx_{k} \mid x_{j} = 1, x_{pa_{j}}) \frac{\mu(x_{j} = 1 \mid x_{pa_{j}})}{\mu(x_{j} = 1 \mid x_{pa_{j}})} \, \mu(dx_{pa_{j}}) \\ &= \int_{\mathcal{X}_{pa_{j}}} \int_{\mathcal{X}_{j}} \int_{\mathcal{X}_{k}} \frac{x_{k} \mathbf{1}_{\{x_{j} = 1\}}}{\mu(x_{j} = 1 \mid x_{pa_{j}})} \, \mu(dx_{k} \mid x_{j}, x_{pa_{j}}) \mu(dx_{j} \mid x_{pa_{j}}) \, \mu(dx_{pa_{j}}) \\ &= \int_{\mathcal{X}_{pa_{j}}} \int_{\mathcal{X}_{j}} \int_{\mathcal{X}_{k}} \frac{x_{k} \mathbf{1}_{\{x_{j} = 1\}}}{\mu(x_{j} = 1 \mid x_{pa_{j}})} \, \mu(dx_{k}, dx_{j}, dx_{pa_{j}}) \\ &= \int_{\mathcal{X}} \frac{x_{k} \mathbf{1}_{\{x_{j} = 1\}}}{\mu(x_{j} = 1 \mid x_{pa_{j}})} \, \mu(dx) \\ &= \int_{\mathcal{X}} fgh \, \mu(dx), \end{split}$$

where $f(x) = 1_{\{x_j=1\}} = x_j$, $g(x) = x_k$ and $h(x) = \mu(x_j = 1 | x_{pa_j})^{-1}$. Moreover, one has

$$\begin{aligned} &f(x)g(x)h(x) - f(y)g(y)h(y) \\ &= & [f(x) - f(y)] \, g(x)h(x) + f(y)[g(x) - g(y)]h(x)) + f(y)g(y)[h(x) - h(y)]. \end{aligned}$$

Since by assumption, \mathcal{X}_k is compact and $\mu(x_j = 1 \mid x_{\text{pa}_j}) \ge \delta$, there exists a constant $K \ge 0$ such that

$$\begin{split} |g(x)h(x)| \leq K, \quad |f(y)h(x)| \leq K \quad \text{and} \quad |f(y)g(y)| \leq K \quad \text{for all} \quad x,y \in \mathcal{X}. \\ \text{Let } \varepsilon > 0 \text{ and choose a coupling } \pi \in \Pi_G^{\mathrm{bc}}(\mu,\nu) \text{ such that} \end{split}$$

$$\int_{\mathcal{X}^2} d_{\mathcal{X}}(x, y) \pi(dx, dy) \le W_{G,1}(\mu, \nu) + \varepsilon.$$

Then,

$$\begin{aligned} \left| \int_{\mathcal{X}_{k}} x_{k} \, \mu(dx_{k} \mid \operatorname{do}(x_{j} = 1)) - \int_{\mathcal{X}_{k}} x_{k} \, \nu(dx_{k} \mid \operatorname{do}(x_{j} = 1)) \right| &= \left| \int_{\mathcal{X}} fgh \, d\mu - \int_{\mathcal{X}} fgh \, d\nu \right| \\ &= \left| \int [f(x)g(x)h(x) - f(y)g(y)h(y)] \, \pi(dx, dy) \right| \\ &\leq K \int_{\mathcal{X}} \left(|f(x) - f(y)| + |g(x) - g(y)| + |h(x) - h(y)| \right) \, \pi(dx, dy) \end{aligned}$$

It is clear that $\int_{\mathcal{X}} \left(|f(x) - f(y)| + |g(x) - g(y)| \right) \pi(dx, dy) \leq W_{G,1}(\mu, \nu) + \varepsilon$. Moreover, by the lower bound on the propensity score and since

$$\left|\frac{1}{a} - \frac{1}{b}\right| \le \frac{1}{\delta^2} |a - b| \text{ for all } a, b \ge \delta,$$

we obtain from Corollary 3.5 that

$$\begin{split} \delta^{2} \int_{\mathcal{X}^{2}} |h(x) - h(y)| \, \pi(dx, dy) &\leq \int_{\mathcal{X}^{2}} \left| \mu(x_{j} = 1 \mid x_{\mathrm{pa}_{j}}) - \nu(y_{j} = 1 \mid y_{\mathrm{pa}_{j}}) \right| \, \pi(dx, dy) \\ &= \int_{\mathcal{X}^{2}} \left| \pi(x_{j} = 1 \mid x_{\mathrm{pa}_{j}}, y_{\mathrm{pa}_{j}}) - \pi(y_{j} = 1 \mid x_{\mathrm{pa}_{j}}, y_{\mathrm{pa}_{j}}) \right| \, \pi(dx, dy) \\ &= \int_{\mathcal{X}^{2}_{\mathrm{pa}_{j}}} \left| \int_{\mathcal{X}^{2}_{j}} (x_{j} - y_{j}) \pi(dx_{j}, dy_{j} \mid y_{\mathrm{pa}_{j}}, y_{\mathrm{pa}_{j}}) \right| \, \pi(dx_{\mathrm{pa}_{j}}, dy_{\mathrm{pa}_{j}}) \\ &\leq \int_{\mathcal{X}^{2}} \left| x_{j} - y_{j} \right| \, \pi(dx, dy) \leq W_{G,1}(\mu, \nu) + \varepsilon. \end{split}$$

Since $\varepsilon > 0$ was arbitrary, this shows that

$$\int_{\mathcal{X}_k} x_k \, \mu(dx_k \mid \operatorname{do}(x_j = 1)) - \int_{\mathcal{X}_k} x_k \, \nu(dx_k \mid \operatorname{do}(x_j = 1)) \bigg| \le K \left(1 + \frac{1}{\delta^2}\right) W_{G,1}(\mu, \nu). \quad (4.5)$$

Analogously, one deduces from the upper bound on the propensity score that

$$\left| \int_{\mathcal{X}_k} x_k \,\mu(dx_k \mid \operatorname{do}(x_j = 0)) - \int_{\mathcal{X}_k} x_k \,\nu(dx_k \mid \operatorname{do}(x_j = 0)) \right| \le K \left(1 + \frac{1}{\delta^2} \right) W_{G,1}(\mu, \nu). \quad (4.6)$$
we the proposition follows from a combination of (4.5) and (4.6).

Now, the proposition follows from a combination of (4.5) and (4.6).

Remark 4.5. It is easy to see that Proposition 4.4 does not hold with the standard 1-Wasserstein distance W_1 instead of $W_{G,1}$ since the topology generated by W_1 is not fine enough. Indeed, measures that are close with respect to W_1 may have completely different transition probabilities $\mu(dx_k \mid x_i)$ $(1, x_{pa_i})$, which play a crucial role in the definition of the average treatment effect ψ^{μ} ; see (4.4).

To construct a simple counterexample, we consider the temporal graph G with three nodes (i.e., $E = \{(1,2), (1,3), (2,3)\}$, where the second variable is the treatment variable and the third the output variable. For $p, H \in (0, 1), q = (1 - p), L = (1 - H)$ and $\varepsilon > 0$, let μ^{ε} be given by

$$\mu^{\varepsilon} = pH^{2}\delta_{(\varepsilon,1,1)} + pHL\delta_{(\varepsilon,1,0)} + pLH\delta_{(\varepsilon,0,1)} + pL^{2}\delta_{(\varepsilon,0,0)} + qH^{2}\delta_{(0,0,0)} + qHL\delta_{(0,0,1)} + qLH\delta_{(0,1,0)} + qL^{2}\delta_{(0,1,1)}$$

Under μ^{ε} , treatment and outcome variable are independent given the first variable. In other words, the treatment has no direct effect on the outcome and one readily finds $\psi^{\mu^{\varepsilon}} = 0$. On the other hand, consider the weak limit of μ^{ε} for $\varepsilon \to 0$, which is given by

$$\mu^{0} = (pH^{2} + qL^{2})\delta_{(0,1,1)} + (pL^{2} + qH^{2})\delta_{(0,0,0)} + LH\delta_{(0,1,0)} + LH\delta_{(0,0,1)}.$$

For instance, setting $p = q = \frac{1}{2}$ and H = 0.9, L = 0.1, we see that now there is a strong positive dependence between treatment and outcome variable, even conditional on the first variable (since conditioning on the first variable is now irrelevant). Indeed, with these values, one finds that $\psi^{\mu^0} =$ 0.64.

The above example may intuitively be framed as modeling the question whether "drinking wine causes health benefits", where the first variable is the common cause "being wealthy or not", but the common cause is no longer included in the information structure for $\varepsilon = 0$.

4.2 Perturbation of structural causal models

We know from Proposition 4.3.(i) that the topology generated by $W_{G,p}$ is the finer the fewer edges the graph G has. In Section 4.1 we saw that average treatment effects are continuous with respect to $W_{G,1}$ but not with respect to the standard 1-Wasserstein distance W_1 , which corresponds to a fully connected graph. In this section we show that the topology generated by $W_{G,1}$ is not too fine for practical purposes by proving that small perturbations of structural causal modes lead to distributions that are close with respect to $W_{G,1}$.

Proposition 4.6. Let G = (V, E) be a sorted DAG and consider for each i = 1, ..., n, a metric $d_{\mathcal{X}_i}$ on \mathcal{X}_i generating the topology on \mathcal{X}_i . Define the metric $d_{\mathcal{X}}$ on $\mathcal{X} = \mathcal{X}_1 \times \cdots \times \mathcal{X}_n$ by $d_{\mathcal{X}}(x, y) = \sum_{i=1}^n d_{\mathcal{X}_i}(x_i, y_i)$ and consider \mathcal{X} -valued random variables $X \sim \mu \in \mathcal{P}_G(\mathcal{X})$ and $Y \sim \nu \in \mathcal{P}_G(\mathcal{X})$ such that for all i = 1, ..., n,

$$X_i = f_i(X_{\operatorname{pa}_i}, U_i)$$
 and $Y_i = g_i(Y_{\operatorname{pa}_i}, V_i)$

for measurable functions $f_i, g_i \colon \mathcal{X}_{pa_i} \times \mathbb{R} \to \mathcal{X}_i$ and \mathbb{R} -valued random variables U_i, V_i satisfying

- (i) f_i is L_i -Lipschitz for a constant $L_i \ge 0$,
- (ii) U_1, \ldots, U_n are independent and
- (iii) V_1, \ldots, V_n are independent.

Then there exists a constant $C \ge 0$ depending on G and L_1, \ldots, L_n such that

$$W_{G,1}(\mu,\nu) \le C \sum_{i=1}^{n} \left\{ \|f_i - g_i\|_{\infty} + W_1(\mathcal{L}(U_i), \mathcal{L}(V_i)) \right\},\tag{4.7}$$

where $\mathcal{L}(U_i)$ and $\mathcal{L}(V_i)$ are the distributions of U_i and V_i , respectively.

Proof. It is well-known that there exist independent pairs of random variables $(\tilde{U}_i, \tilde{V}_i)$ such that $\mathcal{L}(\tilde{U}_i) = \mathcal{L}(U_i), \mathcal{L}(\tilde{V}_i) = \mathcal{L}(V_i)$ and

$$\mathbb{E} |\tilde{U}_i - \tilde{V}_i| = W_1(\mathcal{L}(U_i), \mathcal{L}(V_i)) \quad \text{for all } i = 1, \dots, n;$$

see e.g. [67]. Let us define iteratively

$$(\tilde{X}_i, \tilde{Y}_i) = (f_i(\tilde{X}_{pa_i}, \tilde{U}_i), g_i(\tilde{Y}_{pa_i}, \tilde{V}_i)), \quad i = 1, \dots, n.$$
 (4.8)

Then, $\tilde{X} \sim X$ and $\tilde{Y} \sim Y$. Moreover, it can be seen from (4.8) that

$$(\tilde{X}_i, \tilde{Y}_i) \perp\!\!\!\perp_{\tilde{X}_{\mathsf{pa}_i}, \tilde{Y}_{\mathsf{pa}_i}} (\tilde{X}_{1:i-1}, \tilde{Y}_{1:i-1}), \quad \tilde{X}_i \perp\!\!\!\perp_{\tilde{X}_{\mathsf{pa}_i}} \tilde{Y}_{\mathsf{pa}_i} \quad \text{and} \quad \tilde{Y}_i \perp\!\!\!\perp_{\tilde{Y}_{\mathsf{pa}_i}} \tilde{X}_{\mathsf{pa}_i} \quad \text{for all } i = 2, \dots, n.$$

So it follows from Corollary 3.5 that the distribution π of (\tilde{X}, \tilde{Y}) belongs to $\Pi_G^{\rm bc}(\mu, \nu)$.

If we can show that

$$\mathbb{E} d_{\mathcal{X}_i}(\tilde{X}_i, \tilde{Y}_i) \le C_i \sum_{j=1}^i \left\{ \|f_j - g_j\|_\infty + W_1(\mathcal{L}(U_j), \mathcal{L}(V_j)) \right\}$$
(4.9)

for constants $C_i \ge 0$ depending on G and L_1, \ldots, L_i , we obtain (4.7) with $C = \sum_{i=1}^n C_i$. We prove (4.9) by induction over $i = 1, \ldots, n$. First, note that

$$\mathbb{E} d_{\mathcal{X}_1}(\tilde{X}_1, \tilde{Y}_1) \le \mathbb{E} \Big[d_{\mathcal{X}_1}(f_1(\tilde{U}_1), f_1(\tilde{V}_1)) + d_{\mathcal{X}_1}(f_1(\tilde{V}_1), g_1(\tilde{V}_1)) \Big]$$

$$\le L_1 \mathbb{E} |\tilde{U}_1 - \tilde{V}_1| + ||f_1 - g_1||_{\infty} = L_1 W_1(\mathcal{L}(U_1), \mathcal{L}(V_1)) + ||f_1 - g_1||_{\infty},$$

showing that for i = 1, (4.9) holds with $C_1 = \max \{L_1, 1\}$. For $i \ge 2$, assuming that (4.9) holds for all j < i, one obtains

$$\begin{split} & \mathbb{E} \, d_{\mathcal{X}_{i}}(\tilde{X}_{i}, \tilde{Y}_{i}) \leq \mathbb{E} \Big[d_{\mathcal{X}_{i}}(f_{i}(\tilde{X}_{\mathrm{pa}_{i}}, \tilde{U}_{i}), f_{i}(\tilde{Y}_{\mathrm{pa}_{i}}, \tilde{V}_{i})) + d_{\mathcal{X}_{i}}(f_{i}(\tilde{Y}_{\mathrm{pa}_{i}}, \tilde{V}_{i}), g_{i}(\tilde{Y}_{\mathrm{pa}_{i}}, \tilde{V}_{i}) \Big] \\ \leq \quad L_{i} \, \mathbb{E} \Bigg[\sum_{j \in \mathrm{pa}_{i}} d_{\mathcal{X}_{j}}(\tilde{X}_{j}, \tilde{Y}_{j}) + |\tilde{U}_{i} - \tilde{V}_{i}| \Bigg] + \|f_{i} - g_{i}\|_{\infty} \\ \leq \quad L_{i} \, \sum_{j \in \mathrm{pa}_{i}} C_{j} \, \sum_{k=1}^{j} \big\{ \|f_{k} - g_{k}\|_{\infty} + W_{1}(\mathcal{L}(U_{k}), \mathcal{L}(V_{k})) \big\} + L_{i} W_{1}(\mathcal{L}(U_{i}), \mathcal{L}(V_{i})) + \|f_{i} - g_{i}\|_{\infty} \end{split}$$

So (4.9) holds with

$$C_i = \max\left\{L_i \sum_{j \in \mathsf{pa}_i} C_j, L_i, 1\right\}, \quad i = 1, \dots, n$$

and the proposition follows.

4.3 *G*-causal Wasserstein interpolation

In this section, we assume that in addition to a metric $d_{\mathcal{X}}$ compatible with the Polish topology, \mathcal{X} is endowed with a vector space structure so that the vector space operations are continuous.

Note that for a sorted DAG G = (V, E), $p \in [1, \infty)$ and given measures $\mu, \nu \in \mathcal{P}_{G,p}(\mathcal{X})$, there exists a sequence $(\pi^k)_{k\geq 1}$ of couplings in $\Pi_G^{\mathrm{bc}}(\mu, \nu)$ such that

$$\int_{\mathcal{X}\times\mathcal{X}} d_{\mathcal{X}}(x,y)^p d\pi^k(dx,dy) \to W^p_{G,p}(\mu,\nu) \quad \text{for } k \to \infty.$$

Since $\Pi_G^{bc}(\mu, \nu) \subseteq \Pi(\mu, \nu)$, and the latter is a weakly compact subset of $\mathcal{P}(\mathcal{X} \times \mathcal{X})$, there exists a subsequence, again denoted $(\pi^k)_{k\geq 1}$, which weakly converges to a measure $\pi \in \Pi(\mu, \nu)$. This leads us to the following notion of *G*-causal interpolation, building on the concept of standard Wasserstein geodesics (cf. [67]) and more recent work studying causal Wasserstein interpolation between distributions of stochastic processes; see, e.g. [8, Section 5.4]). We aim for a *G*-causal interpolation concept which, under suitable assumptions, ensures that the interpolation between two *G*-compatible distributions preserves *G*-compatibility.

Definition 4.7. Let G = (V, E) be a sorted DAG and $p \in [1, \infty)$. For given $\mu, \nu \in \mathcal{P}_{G,p}(\mathcal{X})$, let $(\pi^k)_{k\geq 1}$ be a sequence of couplings in $\Pi_G^{\mathrm{bc}}(\mu, \nu)$ such that

$$\int_{\mathcal{X}\times\mathcal{X}} d_{\mathcal{X}}(x,y)^p d\pi^k(dx,dy) \to W^p_{G,p}(\mu,\nu) \quad \text{for } k \to \infty,$$

and π^k weakly converges to some $\pi \in \Pi(\mu, \nu)$. For $\lambda \in [0, 1]$, denote by κ_{λ} the distribution of $(1 - \lambda)X + \lambda Y$, where $(X, Y) \sim \pi$. Then we call $(\kappa_{\lambda})_{\lambda \in [0, 1]}$ a $W_{G, p}$ -interpolation between μ and ν .

The measure π in Definition 4.7 is a limit of measures $\pi^k \in \Pi_G^{bc}(\mu, \nu)$. But in general, it might not belong to $\Pi_G^{bc}(\mu, \nu)$ (see Proposition 3.6.(iv)), and even if it does, the measures κ_{λ} are not necessarily *G*-compatible (see Example 4.11 below). Nevertheless, there are situations where the interpolations κ_{λ} are *G*-compatible, which we study below.

The following result gives conditions under which the distribution of a convex combination of the form $(1 - \lambda)X + \lambda Y$ is G-compatible.

Proposition 4.8. Let G = (V, E) be a sorted DAG and $\mu, \nu \in \mathcal{P}_G(\mathcal{X})$. Assume $(X, Y) \sim \pi$ for a distribution $\pi \in \Pi_G^{\mathrm{bc}}(\mu, \nu)$. Then, for any $\lambda \in [0, 1]$, the distribution of $(X, Y, (1 - \lambda)X + \lambda Y)$ is G-compatible. Moreover, if there exists a measurable set $A \subseteq \mathcal{X} \times \mathcal{X}$ with $\pi(A) = 1$ such that for each i = 2, ..., n, the map

$$h_{\lambda,i}: \mathcal{X}_{\mathrm{pa}_i} \times \mathcal{X}_{\mathrm{pa}_i} \to \mathcal{X}_{\mathrm{pa}_i}, \ (x_{\mathrm{pa}_i}, y_{\mathrm{pa}_i}) \mapsto (1-\lambda)x_{\mathrm{pa}_i} + \lambda y_{\mathrm{pa}_i}$$

is injective on the projection of A to $\mathcal{X}_{pa_i} \times \mathcal{X}_{pa_i}$, then the distribution of $\lambda X + (1 - \lambda)Y$ is G-compatible.

Proof. Since π is in $\Pi_G^{\rm bc}(\mu, \nu)$, we obtain from Corollary 3.5 that

$$(X_i, Y_i) \perp _{X_{pa_i}, Y_{pa_i}} (X_{1:i-1}, Y_{1:i-1})$$

which implies

$$(X_i, Y_i, Z_{\lambda,i}) \perp \!\!\!\perp_{X_{\text{pa}_i}, Y_{\text{pa}_i}, Z_{\lambda, \text{pa}_i}} (X_{1:i-1}, Y_{1:i-1}, Z_{\lambda, 1:i-1})$$
(4.10)

for all $\lambda \in [0,1]$ and $Z_{\lambda} = (1 - \lambda)X + \lambda Y$. By Definition 3.1, this shows that the distribution of (X, Y, Z_{λ}) is *G*-compatible.

Now, let us assume there exists a measurable set $A \subseteq \mathcal{X} \times \mathcal{X}$ with $\pi(A) = 1$ such that for all $i = 2, \ldots, n, h_{\lambda,i}$ is injective on the projection of A to $\mathcal{X}_{pa_i} \times \mathcal{X}_{pa_i}$. If $(\Omega, \mathcal{F}, \mathbb{P})$ is the probability space on which (X, Y) is defined, we can, without loss of generality, assume that $\Omega = (X, Y)^{-1}(A)$ since restricting (X, Y) to $(X, Y)^{-1}(A)$ does not change its distribution. But then, $(X_{pa_i}(\omega), Y_{pa_i}(\omega)) \mapsto (1-\lambda)X_{pa_i}(\omega) + \lambda Y_{pa_i}(\omega)$ is injective for all $\omega \in \Omega$ and $i = 2, \ldots, n$. It follows that $\sigma(X_{pa_i}, Y_{pa_i}) = \sigma(Z_{\lambda,pa_i})$, which together with (4.10) gives $Z_{\lambda,i} \perp \mathbb{Z}_{\lambda,pa_i} Z_{\lambda,1:i-1}$ for all $i = 2, \ldots, n$. This shows that Z_{λ} satisfies condition (ii) of Definition 3.1 and therefore, is G-compatible.

Corollary 4.9. Let G = (V, E) be a sorted DAG and $\mu, \nu \in \mathcal{P}_G(\mathcal{X})$. Assume $(X, Y) \sim \pi$ for a distribution $\pi \in \Pi_G^{\mathrm{bc}}(\mu, \nu)$.

- (i) If µ and ν are finitely supported, the distribution of (1 − λ)X + λY is G-compatible for all but finitely many λ ∈ [0, 1].
- (ii) If μ and ν are countably supported, the distribution of $(1 \lambda)X + \lambda Y$ is *G*-compatible for all but countably many $\lambda \in [0, 1]$.

Proof. We show (i). The proof of (ii) is analogous. Assume μ and ν are of the form $\mu = \sum_{j=1}^{J} p_j \delta_{x_i}$ and $\nu = \sum_{j=1}^{J} q_i \delta_{y_j}$ for $p_j, q_j \ge 0$ and $x_j, y_j \in \mathcal{X}, j = 1, \dots, J$. Then, for all $i \ge 2, x, x' \in \{x_1, \dots, x_J\}$ and $y, y' \in \{y_1, \dots, y_J\}$ such that $(x_{pa_i}, y_{pa_i}) \ne (x'_{pa_i}, y'_{pa_i})$, there exists at most one $\lambda \in [0, 1]$ satisfying

$$(1 - \lambda)x_{\mathrm{pa}_i} + \lambda y_{\mathrm{pa}_i} = (1 - \lambda)x'_{\mathrm{pa}_i} + \lambda y'_{\mathrm{pa}_i}$$
$$\Leftrightarrow \quad (1 - \lambda)(x_{\mathrm{pa}_i} - x'_{\mathrm{pa}_i}) = \lambda(y'_{\mathrm{pa}_i} - y_{\mathrm{pa}_i}).$$

It follows that for every i = 2, ..., n and all but finitely many $\lambda \in [0, 1]$, the map

 $h_{\lambda,i}: \mathcal{X}_{\mathrm{pa}_i} \times \mathcal{X}_{\mathrm{pa}_i} \to \mathcal{X}_{\mathrm{pa}_i}, \ (x_{\mathrm{pa}_i}, y_{\mathrm{pa}_i}) \mapsto (1-\lambda)x_{\mathrm{pa}_i} + \lambda y_{\mathrm{pa}_i}$

is injective on the projection of $A = \{x_1, \ldots, x_J\} \times \{y_1, \ldots, y_J\}$ to $\mathcal{X}_{pa_i} \times \mathcal{X}_{pa_i}$. Since $\pi \in \Pi_G^{\mathrm{bc}}(\mu, \nu)$, we have $\pi(A) = 1$ and therefore, obtain from Proposition 4.8 that the distribution of $\lambda X + (1 - \lambda)Y$ is *G*-compatible.

Corollary 4.10. Let G = (V, E) be a sorted DAG, $p \in [1, \infty)$ and μ, ν two finitely supported measures in $\mathcal{P}_G(\mathcal{X})$ (and therefore also in $\mathcal{P}_{G,p}(\mathcal{X})$). Then there exists a measure $\pi \in \Pi_G^{\mathrm{bc}}(\mu, \nu)$ such that

$$\int_{\mathcal{X}\times\mathcal{X}} d_{\mathcal{X}}(x,y)^p d\pi(dx,dy) = W^p_{G,p}(\mu,\nu).$$
(4.11)

Moreover, for $(X, Y) \sim \pi$ *, the distribution of* $(1 - \lambda)X + \lambda Y$ *is G-compatible for all but finitely many* $\lambda \in [0, 1]$.

Proof. That there is a measure $\pi \in \Pi_G^{bc}(\mu, \nu)$ satisfying (4.11) follows directly from Proposition 4.3.(iii). Moreover, for $(X, Y) \sim \pi$, we obtain from Corollary 4.9.(i) that the distribution of $(1 - \lambda)X + \lambda Y$ is *G*-compatible for all but finitely many $\lambda \in [0, 1]$.

Example 4.11. Consider the Markovian graph G = (V, E) with $V = \{1, 2, 3\}$ and $E = \{(1, 2), (2, 3)\}$, and take $\mathcal{X} = \mathbb{R} \times \mathbb{R} \times \mathbb{R}$ endowed with the Euclidean distance. The distributions $\mu = \frac{1}{2}(\delta_{(0,0,0)} + \delta_{(1,1,1)})$ and $\nu = \frac{1}{2}(\delta_{(0,1,0)} + \delta_{(1,0,1)})$ are both Markovian, and clearly, the optimal W_2 -transport plan transports (0, 0, 0) to (0, 1, 0) and (1, 1, 1) to (1, 0, 1). It can directly be seen from Definition 2.1 that this transport plan belongs to $\Pi_G^{\mathrm{bc}}(\mu, \nu)$. Therefore,

$$\kappa_{\lambda} = \frac{1}{2} \left(\delta_{(0,\lambda,0)} + \delta_{(1,1-\lambda,1)} \right), \quad \lambda \in [0,1],$$

is the unique $W_{G,2}$ -interpolation between μ and ν . Note that κ_{λ} is Markovian for all $\lambda \in [0,1]$ except for $\lambda = 1/2$. But for $\lambda = 1/2$, it is still a weak limit of Markovian distributions.

Next, we study an example where standard Wasserstein interpolation between two G-compatible measures leads to distributions that are far from G-compatible.

Example 4.12. Consider the same setup as in Example 4.11. Denote by μ the distribution of the random walk (X_1, X_2, X_3) given by $X_i = \sum_{j=1}^i \xi_j$ for i.i.d. innovations ξ_j with distribution $\mathbb{P}[\xi_j = \pm 1] = 1/2$ and ν the distribution of (Y_1, Y_2, Y_3) given by $Y_i = \sum_{j=1}^i \eta_j$ for i.i.d. innovations η_j with distribution $\mathbb{P}[\eta_j = -1, 0, 1] = 1/3$. The first two pictures in Figure 1 show all possible trajectories of $X = (X_1, X_2, X_3)$ and $Y = (Y_1, Y_2, Y_3)$, respectively.

For the standard Wasserstein distance $W_2(\mu, \nu)$, a numerically obtained optimal coupling π can be seen to satisfy

$$\pi((X_3, Y_3) = (1, 0) \mid (X_1, Y_1) = (-1, 0), (X_2, Y_2) = (0, 0))$$

< $\pi((X_3, Y_3) = (1, 0) \mid (X_1, Y_1) = (1, 0), (X_2, Y_2) = (0, 0)),$

showing that $(X_i, Y_i)_{i=1,2,3}$ is not Markovian under π . It follows that, for all but finitely many $\lambda \in [0,1]$, the resulting interpolations κ_{λ} under this coupling π are also not Markovian.⁵

⁵This follows from the same arguments as in the proofs of Proposition 4.8 and Corollary 4.9, where the main argument is that for $Z_k^{\lambda} = (1 - \lambda)X_k + \lambda Y_k$, one has $\sigma(Z_k^{\lambda}) = \sigma(X_k, Y_k)$ for all but finitely many $\lambda \in [0, 1]$.



Figure 1: The binomial model μ on the left, the trinomial model ν in the middle, and the interpolating measure $\kappa_{1/3}$ under $W_{G,2}$ on the right as given by Corollary 4.10 using the Euclidean distance on \mathbb{R}^3 as the underlying metric.

On the other hand, using a $W_{G,p}$ -interpolation as given by Corollary 4.10, the resulting interpolating measures κ_{λ} are Markovian for all but finitely many $\lambda \in [0, 1]$. The last picture in Figure 1 shows an illustration of the numerically computed G-causal (i.e., Markov) Wasserstein interpolation $\kappa_{1/3}$ using $W_{G,1}$.

A Stability of conditional independence

Lemma A.1. Let S, W, Z be measurable spaces and S, W, Z the projections from $S \times W \times Z$ to S, W, Z, respectively. Denote by $\mathcal{P}(S \times W \times Z)$ the collection of all probability measures on $S \times W \times Z$ equipped with the product σ -algebra. Then the set

$$\{\pi \in \mathcal{P}(\mathcal{S} \times \mathcal{W} \times \mathcal{Z}) : S \amalg_Z W \text{ with respect to } \pi\}$$
(A.1)

is closed under convergence in total variation.

Proof. Let $(\pi^k)_{k\in\mathbb{N}}$ be a sequence in (A.1) converging to a measure $\pi \in \mathcal{P}(\mathcal{S} \times \mathcal{W} \times \mathcal{Z})$ in total variation. By (3.2), one has

$$\pi^{k}[A \times \mathcal{W} \times \mathcal{Z} \mid W, Z] = \pi^{k}[A \times \mathcal{W} \times \mathcal{Z} \mid Z] \quad \pi^{k}\text{-a.s.}$$
(A.2)

for every measurable subset $A \subseteq S$ and all $k \in \mathbb{N}$. Let us denote

$$H_k = \pi^k [A \times \mathcal{W} \times \mathcal{Z} \mid Z]$$
 and $H = \pi [A \times \mathcal{W} \times \mathcal{Z} \mid Z]$

If \mathbb{E}^k denotes expectation with respect to π^k and \mathbb{E} expectation with respect to π , one has

$$\left|\mathbb{E}^{k}U - \mathbb{E}U\right| = \left|\int_{0}^{1} \left(\pi^{k}[U > u] - \pi[U > u]\right) du\right| \le \int_{0}^{1} \left|\pi^{k}[U > u] - \pi[U > u]\right| du$$

for all random variables $U: S \times W \times Z \rightarrow [0, 1]$, which shows that for every $\varepsilon > 0$, there exists a $k_0 \in \mathbb{N}$ such that

$$\left|\mathbb{E}^{k}U - \mathbb{E}U\right| \leq \varepsilon$$
 for every $k \geq k_{0}$ and all such U .

In particular,

$$\left|\mathbb{E}^{k}(1_{A\times\mathcal{W}\times\mathcal{Z}}-H)^{2}-\mathbb{E}\left(1_{A\times\mathcal{W}\times\mathcal{Z}}-H\right)^{2}\right|\leq\varepsilon$$

and

$$\mathbb{E}^{k}(1_{A\times\mathcal{W}\times\mathcal{Z}}-H_{k})^{2}-\mathbb{E}\left(1_{A\times\mathcal{W}\times\mathcal{Z}}-H_{k}\right)^{2}\Big|\leq\varepsilon$$

for all $k \ge k_0$. Moreover, since the H_k can be viewed as L^2 -projections of $1_{A \times W \times Z}$ on the space of square-integrable Z-measurable random variables, one has

$$\mathbb{E}^k (1_{A \times \mathcal{W} \times \mathcal{Z}} - H_k)^2 \le \mathbb{E}^k (1_{A \times \mathcal{W} \times \mathcal{Z}} - H)^2.$$

Together with Pythagoras's theorem this gives

$$\begin{split} \mathbb{E}(H_k - H)^2 &= \mathbb{E}(1_{A \times \mathcal{W} \times \mathcal{Z}} - H_k)^2 - \mathbb{E}(1_{A \times \mathcal{W} \times \mathcal{Z}} - H)^2 \\ &\leq \mathbb{E}^k (1_{A \times \mathcal{W} \times \mathcal{Z}} - H_k)^2 + \varepsilon - \mathbb{E}(1_{A \times \mathcal{W} \times \mathcal{Z}} - H)^2 \\ &\leq \mathbb{E}^k (1_{A \times \mathcal{W} \times \mathcal{Z}} - H)^2 + \varepsilon - \mathbb{E}(1_{A \times \mathcal{W} \times \mathcal{Z}} - H)^2 \\ &\leq \mathbb{E}(1_{A \times \mathcal{W} \times \mathcal{Z}} - H)^2 + 2\varepsilon - \mathbb{E}(1_{A \times \mathcal{W} \times \mathcal{Z}} - H)^2 = 2\varepsilon \quad \text{for all } k \ge k_0, \end{split}$$

showing that $H_k \to H$ in $L^2(\pi)$. Analogously, one obtains that $\pi^k[A \times \mathcal{W} \times \mathcal{Z} \mid W, Z]$ converges to $\pi[A \times \mathcal{W} \times \mathcal{Z} \mid W, Z]$ in $L^2(\pi)$. So, equation (A.2) is stable under convergence in total variation, which proves the lemma.

B Counterexample to the triangle inequality

In this section, we provide an example showing that in general, $W_{G,1}$ does not satisfy the triangle inequality. We consider the Markovian graph G = (V, E) with vertices $V = \{1, 2, 3\}$ and edges $\{(1, 2), (2, 3)\}$ and choose $\mathcal{X} = \mathcal{X}_1 \times \mathcal{X}_2 \times \mathcal{X}_3$, where $\mathcal{X}_1, \mathcal{X}_2, \mathcal{X}_3$ are all equal to the same abstract discrete Polish space $\{x_1, x_2, x_3, x_4\}$. Define μ, ν, η as follows, which we note are Markovian:

$$\mu = \frac{1}{4} \left(\delta_{(x_1, x_1, x_1)} + \delta_{(x_1, x_1, x_2)} + \delta_{(x_2, x_1, x_1)} + \delta_{(x_2, x_1, x_2)} \right) \nu = \frac{1}{4} \left(\delta_{(x_1, x_2, x_1)} + \delta_{(x_1, x_2, x_2)} + \delta_{(x_2, x_3, x_1)} + \delta_{(x_2, x_3, x_2)} \right) \eta = \frac{1}{4} \left(\delta_{(x_3, x_4, x_3)} + \delta_{(x_3, x_4, x_4)} + \delta_{(x_4, x_4, x_3)} + \delta_{(x_4, x_4, x_4)} \right).$$

To gain some intuition why $W_{G,1}$ violates the triangle inquality in this example, consider stochastic processes $(X_1, X_2, X_3) \sim \mu$, $(Y_1, Y_2, Y_3) \sim \nu$, $(Z_1, Z_2, Z_3) \sim \eta$ and note that X_2 and Z_2 are constant while $\sigma(Y_2) = \sigma(Y_1, Y_2)$. So it can be seen from Corollary 3.5 that transport plans in $\Pi_G^{\text{bc}}(\mu, \eta)$ are more restricted than those in $\Pi_G^{\text{bc}}(\mu, \nu)$ and $\Pi_G^{\text{bc}}(\nu, \eta)$, which, for a suitable metric on \mathcal{X} , implies that $W_{G,1}(\mu, \eta)$ is large compared to $W_{G,1}(\mu, \nu)$ and $W_{G,1}(\nu, \eta)$. We specify the distances between the 12 sequences $(x_1, x_1, x_1), (x_1, x_1, x_2), \ldots$ used to define the measures μ, ν, η with the following 12×12 matrix M:

0	0.53	1.08	1.33	1.29	0.78	0.64	0.44	1.15	1.3	1.92	1.38
0.53	0	0.7	1.05	1.18	0.91	0.11	0.97	1.1	1.83	1.9	1.91
1.08	0.7	0	0.98	1.11	0.49	0.59	0.64	0.86	1.88	1.82	1.58
1.33	1.05	0.98	0	0.13	1.15	0.94	0.98	1.52	1.25	1.87	1.92
1.29	1.18	1.11	0.13	0	1.02	1.07	0.85	1.39	1.38	2	1.79
0.78	0.91	0.49	1.15	1.02	0	1.02	1.13	0.37	1.74	1.77	1.22
0.64	0.11	0.59	0.94	1.07	1.02	0	1.08	1.21	1.94	1.79	2.02
0.44	0.97	0.64	0.98	0.85	1.13	1.08	0	1.5	1.24	1.86	0.94
1.15	1.1	0.86	1.52	1.39	0.37	1.21	1.5	0	1.37	1.4	0.85
1.3	1.83	1.88	1.25	1.38	1.74	1.94	1.24	1.37	0	0.62	1.2
1.92	1.9	1.82	1.87	2	1.77	1.79	1.86	1.4	0.62	0	1.69
1.38	1.91	1.58	1.92	1.79	1.22	2.02	0.94	0.85	1.2	1.69	0

That is, $M_{1,2}$ is the distance between (x_1, x_1, x_1) and (x_1, x_1, x_2) , $M_{1,3}$ is the distance between (x_1, x_1, x_1) and (x_2, x_1, x_1) and so on. We obtained M by simulating symmetric matrices with zeros on the diagonal and positive entries off the diagonal and iteratively updating the entries as long as the triangle inequality was violated. More precisely, we started with a symmetric random matrix M^0 with zeros on the diagonal and positive off-diagonal entries. Then we iteratively set

$$M_{i,j}^{k} = \min\left\{M_{i,j}^{k-1}, M_{i,1}^{k-1} + M_{1,j}^{k-1}, M_{i,2}^{k-1} + M_{2,j}^{k-1}, \dots, M_{i,12}^{k-1} + M_{12,j}^{k-1}\right\}$$

until $M^k = M^{k-1}$. This guarantees that M defines a metric on $(x_1, x_1, x_1), (x_1, x_1, x_2) \dots$ From there it can be extended to a metric on \mathcal{X} by Fréchet embedding. Finally, we used Gurobi [28] to compute $W_{G,1}(\mu, \nu) = 0.585, W_{G,1}(\nu, \eta) = 2.24$ and $W_{G,1}(\mu, \eta) = 2.925$, showing that

$$W_{G,1}(\mu,\eta) > W_{G,1}(\mu,\nu) + W_{G,1}(\nu,\eta).$$

References

- Beatrice Acciaio, Julio Backhoff-Veraguas, and Junchao Jia. Cournot–Nash equilibrium and optimal transport in a dynamic setting. *SIAM Journal on Control and Optimization*, 59(3):2273– 2300, 2021.
- [2] Martial Agueh and Guillaume Carlier. Barycenters in the Wasserstein space. *SIAM Journal on Mathematical Analysis*, 43(2):904–924, 2011.
- [3] David Aldous. Weak convergence and the general theory of processes. *Incomplete Draft*, 1981. URL https://www.stat.berkeley.edu/~aldous/Papers/weak-gtp.pdf.
- [4] Martin Arjovsky, Soumith Chintala, and Léon Bottou. Wasserstein generative adversarial networks. In *International Conference on Machine Learning*, pages 214–223. PMLR, 2017.
- [5] Julio Backhoff-Veraguas, Daniel Bartl, Mathias Beiglböck, and Manu Eder. All adapted topologies are equal. *Probability Theory and Related Fields*, 178(3):1125–1172, 2020.

- [6] Julio Backhoff-Veraguas, Mathias Beiglböck, Yiqing Lin, and Anastasiia Zalashko. Causal transport in discrete time and applications. *SIAM Journal on Optimization*, 27(4):2528–2562, 2017.
- [7] Julio Backhoff-Veraguas and Gudmund Pammer. Stability of martingale optimal transport and weak optimal transport. *The Annals of Applied Probability*, 32(1):721–752, 2022.
- [8] Daniel Bartl, Mathias Beiglböck, and Gudmund Pammer. The Wasserstein space of stochastic processes. *arXiv Preprint 2104.14245*, 2021.
- [9] Mathias Beiglböck and Nicolas Juillet. On a problem of optimal transport under marginal martingale constraints. *The Annals of Probability*, 44(1):42–106, 2016.
- [10] Jose Blanchet and Karthyek Murthy. Quantifying distributional model risk via optimal transport. *Mathematics of Operations Research*, 44(2):565–600, 2019.
- [11] Stephan Bongers, Patrick Forré, Jonas Peters, and Joris M Mooij. Foundations of structural causal models with cycles and latent variables. *The Annals of Statistics*, 49(5):2885–2915, 2021.
- [12] Nicolas Bonneel and David Coeurjolly. Spot: sliced partial optimal transport. *ACM Transactions on Graphics (TOG)*, 38(4):1–13, 2019.
- [13] Nicolas Bourbaki. *Elements of Mathematics: General Topology, Part 2*. Actualités Scientifiques et Industrielles. Hermann, 1966.
- [14] Patrick Cheridito, Matti Kiskii, David Prömel, and Mete Soner. Martingale optimal transport duality. *Mathematische Annalen*, 379:1685–1712, 2021.
- [15] Patrick Cheridito, Michael Kupper, and Ludovic Tangpi. Duality formulas for robust pricing and hedging in discrete time. *SIAM Journal on Financial Mathematics*, 8(1):738–765, 2017.
- [16] Pierre-André Chiappori, Robert J McCann, and Lars P Nesheim. Hedonic price equilibria, stable matching, and optimal transport: equivalence, topology, and uniqueness. *Economic Theory*, pages 317–354, 2010.
- [17] Lenaic Chizat, Gabriel Peyré, Bernhard Schmitzer, and François-Xavier Vialard. Unbalanced optimal transport: Dynamic and Kantorovich formulations. *Journal of Functional Analysis*, 274 (11):3090–3123, 2018.
- [18] Marco Cuturi. Sinkhorn distances: lightspeed computation of optimal transport. Advances in Neural Information Processing Systems, 26, 2013.
- [19] Marco Cuturi and Arnaud Doucet. Fast computation of Wasserstein barycenters. In *International Conference on Machine Learning*, pages 685–693. PMLR, 2014.
- [20] Luca De Gennaro Aquino and Stephan Eckstein. Minmax methods for optimal transport and beyond: regularization, approximation and numerics. *Advances in Neural Information Processing Systems*, 33:13818–13830, 2020.
- [21] Lucas De Lara, Alberto González-Sanz, Nicholas Asher, and Jean-Michel Loubes. Transportbased counterfactual models. arXiv Preprint 2108.13025, 2021.

- [22] Guido De Philippis and Alessio Figalli. The Monge–Ampère equation and its link to optimal transportation. *Bulletin of the American Mathematical Society*, 51(4):527–580, 2014.
- [23] Paul Embrechts, Giovanni Puccetti, and Ludger Rüschendorf. Model uncertainty and VaR aggregation. *Journal of Banking & Finance*, 37(8):2750–2764, 2013.
- [24] Alessio Figalli, Francesco Maggi, and Aldo Pratelli. A mass transportation approach to quantitative isoperimetric inequalities. *Inventiones Mathematicae*, 182(1):167–211, 2010.
- [25] Nicolas Fournier and Arnaud Guillin. On the rate of convergence in Wasserstein distance of the empirical measure. *Probability Theory and Related Fields*, 162(3):707–738, 2015.
- [26] Alfred Galichon. Optimal Transport Methods in Economics. Princeton University Press, 2018.
- [27] Promit Ghosal and Bodhisattva Sen. Multivariate ranks and quantiles using optimal transport: consistency, rates and nonparametric testing. *The Annals of Statistics*, 50(2):1012–1037, 2022.
- [28] Gurobi Optimization, LLC. Gurobi Optimizer Reference Manual, 2022. URL https://www.gurobi.com.
- [29] Martin Hellwig. Sequential decisions under uncertainty and the maximum theorem. *Journal of Mathematical Economics*, 25(4):443–464, 1996.
- [30] Paul W Holland. Statistics and causal inference. *Journal of the American Statistical Association*, 81(396):945–960, 1986.
- [31] Yiyan Huang, Cheuk Hang Leung, Qi Wu, Xing Yan, Shumin Ma, Zhiri Yuan, Dongdong Wang, and Zhixiang Huang. Robust causal learning for the estimation of average treatment effects. In 2022 International Joint Conference on Neural Networks (IJCNN 2022). IEEE, 2022.
- [32] Olav Kallenberg. Foundations of Modern Probability, 3rd Edition. Springer.
- [33] Leonid V Kantorovich. On the translocation of masses. In *Dokl. Akad. Nauk. USSR (NS)*, volume 37, pages 199–201, 1942.
- [34] Leonid V Kantorovich. Mathematical methods of organizing and planning production. *Management Science*, 6(4):366–422, 1960.
- [35] Murat Kocaoglu, Christopher Snyder, Alexandros G Dimakis, and Sriram Vishwanath. Causalgan: learning causal implicit generative models with adversarial rraining. In *International Conference on Learning Representations*, 2018.
- [36] Jonathan Korman and Robert McCann. Optimal transportation with capacity constraints. *Transactions of the American Mathematical Society*, 367(3):1501–1521, 2015.
- [37] Rémi Lassalle. Causal transport plans and their Monge–Kantorovich problems. Stochastic Analysis and Applications, 36(3):452–484, 2018.
- [38] Steffen L Lauritzen. Graphical Models, volume 17. Clarendon Press, 1996.
- [39] Jia Li and Lin Lin. Optimal transport with relaxed marginal constraints. *IEEE Access*, 9:58142– 58160, 2021.

- [40] Dirk A Lorenz, Paul Manns, and Christian Meyer. Quadratically regularized optimal transport. *Applied Mathematics & Optimization*, 83(3):1919–1949, 2021.
- [41] Facundo Mémoli. Gromov–Wasserstein distances and the metric approach to object matching. *Foundations of Computational Mathematics*, 11(4):417–487, 2011.
- [42] Peyman Mohajerin Esfahani and Daniel Kuhn. Data-driven distributionally robust optimization using the Wasserstein metric: Performance guarantees and tractable reformulations. *Mathematical Programming*, 171(1-2):115–166, 2018.
- [43] Gaspard Monge. Mémoire sur la théorie des déblais et des remblais. Mem. Math. Phys. Acad. Royale Sci., pages 666–704, 1781.
- [44] Jonathan Niles-Weed and Quentin Berthet. Minimax estimation of smooth densities in Wasserstein distance. *The Annals of Statistics*, 50(3):1519–1540, 2022.
- [45] Marcel Nutz and Ruodu Wang. The directional optimal transport. The Annals of Applied Probability, 32(2):1400–1420, 2022.
- [46] Gudmund Pammer. A note on the adapted weak topology in discrete time. *arXiv Preprint* 2205.00989, 2022.
- [47] Brendan Pass. Multi-marginal optimal transport: theory and applications. *ESAIM: Mathematical Modelling and Numerical Analysis*, 49(6):1771–1790, 2015.
- [48] Judea Pearl. Causality. Cambridge university press, 2009.
- [49] Jonas Peters, Peter Bühlmann, and Nicolai Meinshausen. Causal inference by using invariant prediction: identification and confidence intervals. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 78(5):947–1012, 2016.
- [50] Gabriel Peyré and Marco Cuturi. *Computational Optimal Transport: With Applications to Data Science*. Now Publishers and Trends, 2019.
- [51] Georg Ch Pflug and Alois Pichler. A distance for multistage stochastic optimization models. *SIAM Journal on Optimization*, 22(1):1–23, 2012.
- [52] Georg Ch Pflug and Alois Pichler. Dynamic generation of scenario trees. *Computational Optimization and Applications*, 62(3):641–668, 2015.
- [53] Georg Ch Pflug, Alois Pichler, and David Wozabal. The 1/n investment strategy is optimal under high model ambiguity. *Journal of Banking & Finance*, 36(2):410–417, 2012.
- [54] Drago Plečko and Nicolai Meinshausen. Fair data adaptation with quantile preservation. *Journal* of Machine Learning Research, 21(242):1–44, 2020.
- [55] Svetlozar T Rachev. The Monge–Kantorovich mass transference problem and its stochastic applications. *Theory of Probability & Its Applications*, 29(4):647–676, 1985.
- [56] Svetlozar T Rachev and Ludger Rüschendorf. *Mass Transportation Problems. Vol. I Theory, Vol. II Applications*. Springer Verlag, Berlin, 1998.

- [57] Paul R Rosenbaum and Donald B Rubin. The central role of the propensity score in observational studies for causal effects. *Biometrika*, 70:41–55, 1983.
- [58] Fredrik Sävje, Peter Aronow, and Michael Hudgens. Average treatment effects in the presence of unknown interference. *Annals of Statistics*, 49(2):673, 2021.
- [59] Bernhard Schölkopf. Causality for machine learning. In *Probabilistic and Causal Inference: The Works of Judea Pearl*, pages 765–804. 2022.
- [60] Thibault Séjourné, Gabriel Peyré, and François-Xavier Vialard. Unbalanced optimal transport, from theory to numerics. *arxiv Preprint 2211.08775*, 2022.
- [61] Bharath K Sriperumbudur, Kenji Fukumizu, Arthur Gretton, and Schölkopf. On the empirical estimation of integral probability metrics.
- [62] V. N. Sudakov. Geometric problems of the theory of infinite-dimensional probability distributions. *Trudy Mat. Inst. Steklov*, 141:3–191, 1976.
- [63] Vayer Titouan, Ievgen Redko, Rémi Flamary, and Nicolas Courty. Co-optimal transport. Advances in Neural Information Processing Systems, 33:17559–17570, 2020.
- [64] Quang Huy Tran, Hicham Janati, Ievgen Redko, Flamary Rémi, and Nicolas Courty. Factored couplings in multi-marginal optimal transport via difference of convex programming. *NeurIPS* 2021 Optimal Transport and Machine Learning Workshop (OTML), 2021.
- [65] Sergey S Vallender. Calculation of the Wasserstein distance between probability distributions on the line. *Theory of Probability & Its Applications*, 18(4):784–786, 1974.
- [66] Leonid N Vaserstein. Markov processes over denumerable products of spaces, describing large systems of automata. *Problemy Peredachi Informatsii*, 5(3):64–72, 1969.
- [67] Cédric Villani. Optimal Transport: Old and New, volume 338. Springer, 2009.
- [68] Tianlin Xu and Beatrice Acciaio. Quantized conditional cot-gan for video prediction. *arXiv Preprint 2106.05658*, 2021.
- [69] Tianlin Xu, Li Kevin Wenliang, Michael Munn, and Beatrice Acciaio. Cot-gan: generating sequential data via causal optimal transport. *Advances in Neural Information Processing Systems*, 33:8798–8809, 2020.
- [70] Insoon Yang. A convex optimization approach to distributionally robust Markov decision processes with Wasserstein distance. *IEEE Control Systems Letters*.