Difference-based covariance matrix estimate in time series nonparametric regression with applications to specification tests

Lujia Bai Center for Statistical Science, Department of Industrial Engineering, Tsinghua University Weichi Wu Center for Statistical Science, Department of Industrial Engineering, Tsinghua University

Abstract

Long-run covariance matrix estimation is the building block of time series inference. The corresponding difference-based estimator, which avoids detrending, has attracted considerable interest due to its robustness to both smooth and abrupt structural breaks and its competitive finite sample performance. However, existing methods mainly focus on estimators for the univariate process while their direct and multivariate extensions for most linear models are asymptotically biased. We propose a novel difference-based and debiased long-run covariance matrix estimator for functional linear models with time-varying regression coefficients, allowing time series non-stationarity, long-range dependence, stateheteroscedasticity and their mixtures. We apply the new estimator to (i) the structural stability test, overcoming the notorious non-monotonic power phenomena caused by piecewise smooth alternatives for regression coefficients, and (ii) the nonparametric residual-based tests for long memory, improving the performance via the residual-free formula of the proposed estimator. The effectiveness of the proposed method is justified theoretically and demonstrated by superior performance in simulation studies, while its usefulness is elaborated via real data analysis. Our method is implemented in the R package mlrv.

Keywords: Debias; Difference statistic; Local stationarity; Long-run variance; Monotonic power; Time-varying linear model;

1 Introduction

The long-run variance plays a central role in the statistical inference of time series linear models. Consider the following functional linear model ((Zhou and Wu, 2010)) for the time series observations $(y_{i,n}, x_{i,n})$ (i = 1, ..., n),

$$y_{i,n} = x_{i,n}^{\top} \beta_{i,n} + e_{i,n}, \quad \beta_{i,n} = \beta(i/n) \quad (i = 1, \dots, n),$$
(1.1)

¹E-mail addresses: blj20@mails.tsinghua.edu.cn(L.Bai), wuweichi@mail.tsinghua.edu.cn(W.Wu)

where $x_{i,n}$ is a p-dimensional covariate process whose first element is 1 and $e_{i,n}$ is the error process, both of which can be non-stationary, $(\beta_{i,n})$ are the time-varying regression coefficients and $\beta(\cdot): [0,1] \to \mathbb{R}^p$ is the regression function. If $\hat{\beta}_{i,n}$ is assumed to be constant or smoothly changing, it can be estimated by ordinary least squares or nonparametric methods ((Chan and Zhang, 2010), (Li et al., 2011), (Su et al., 2019)) and the variation of such estimates is determined by the *p*-dimensional possibly time-varying long-run covariance matrix of $(x_{i,n}e_{i,n})$, see (2.1) for exact definition. Classic estimation of the long-run covariance matrix in linear models, see for instances (Newey and West, 1987) and (Andrews, 1991), requires consistent estimation of regression coefficients, which is difficult to achieve under structural changes. A prevalent approach to overcome the obstacle is the difference-based estimation, which is built on $y_{i+m,n} - y_{i,n}$ where m is a diverging tuning parameter such that m = o(n) so that most $y_{i+m,n} - y_{i,n}$ are approximately zeromean under both smooth and abrupt structural changes. Most of the existing results assume stationarity, see for example, (Tecuapetla-Gómez and Munk, 2017) and (Chan, 2022). Exceptions include (Dette and Wu, 2019), which allows non-stationary errors. To the best of the author's knowledge, the existing difference-based long-run variance estimators for time series data are designed for p = 1. When $(e_{i,n})$ are independent and identically distributed, the difference-based estimators for variance has been studied by for example (Müller and Stadtmuller, 1987), (Hall et al., 1990) and (Brown and Levine, 2007).

For p > 1, the long-run variance is referred as the long-run covariance matrix, whose estimation is much more involved, see (Jansson, 2002), (Hirukawa, 2021) and reviews therein. Most existing methods are plug-in methods that utilize estimated residuals which have been widely applied to the goodness of fit tests, tests of structural breaks ((Aue and Horváth, 2013), (Wu and Zhou, 2018), (Kao et al., 2018)), detecting gradual changes ((Vogt and Dette, 2015)), simultaneous confidence bands for coefficient functions ((Zhou and Wu, 2010)), tests for long memory ((Beran et al., 2013), (Bai and Wu, 2024)), etc. Since they depend critically on the accurate pre-estimation of regression coefficients, they are inconsistent under abrupt structural breaks, causing the notorious non-monotonic power ((Kejriwal, 2009)) when applied to structural stability tests. Moreover, most nonparametric specification tests involve both $\hat{\beta}_{i,n}$ and the long-run covariance matrix, while $\hat{\beta}_{i,n}$ is also used in the formula of the plug-in estimator for the long-run covariance matrix. Therefore, the plug-in estimate tends to sensitize those tests to the tuning parameters chosen for $\hat{\beta}_{i,n}$.

2 Preliminaries

In this paper, we consider a general form of non-stationary called local stationarity for the covariate and the error processes of (1.1), which has received substantial attention in the literature. We employ the definition of locally stationary processes based on Bernoulli shift processes, see (Wu, 2005) and (Zhou and Wu, 2010), while there are also many other formulations, see (Dahlhaus, 1997), (Nason et al., 2000), and (Dahlhaus et al., 2019) for a comprehensive review.

We start by introducing necessary notation that will be used in the rest of the paper. Let $1(\cdot)$ denote the indicator function. Define $\lfloor a \rfloor$ as the largest integer smaller than a, and $\lambda_{\min}(A)$ as the smallest eigenvalue of any symmetric squared matrix A. Let $|\cdot|$ denote the absolute value for scalars and the Frobenius norm

for matrices. Let Z denote the set of integers. Let $\mathcal{F}_i = (\dots, \eta_{i-1}, \eta_i)$, where $(\eta_i)_{i \in \mathbb{Z}}$ are independent and identically distributed random elements, and the couple process $\mathcal{F}_{i,\{0\}} = (\dots, \eta_{-1}, \eta'_0, \eta_1, \dots, \eta_i)$, where $(\eta'_i)_{i \in \mathbb{Z}}$ is the independent and identically distributed copy of $(\eta_i)_{i \in \mathbb{Z}}$. We write $f(\cdot) \in C^q[0, 1]$ if $f(\cdot)$ has *q*th order continuous derivative. Consider the data generating mechanism $L(t, \mathcal{F}_i) \in \mathbb{R}^p$, where L is a filter function such that $L(t, \mathcal{F}_i)$ is well-defined. We say $L(t, \mathcal{F}_0) \in \operatorname{Lip}_s(I)$ if there exists a constant c > 0such that for any $t_1, t_2 \in I, t_1 < t_2$,

$$||L(t_1, \mathcal{F}_0) - L(t_2, \mathcal{F}_0)||_s \le c|t_1 - t_2|,$$

where $||X||_s = \{E(|X|^s)\}^{1/s}$. To measure the dependence of time series, we adopt the physical dependence ((Wu, 2005)). For $L(t, \mathcal{F}_i)$ on interval I, the physical dependence in \mathcal{L}^r norm is defined by

$$\delta_r(L,k,I) = \sup_{t \in I} \|L(t,\mathcal{F}_k) - L(t,\mathcal{F}_{k,\{0\}})\|_r.$$

In the following, we give the definition of local stationarity and short-range dependence.

Definition 2.1. The process $G(t, \mathcal{F}_i)$ $(i \in Z)$ is of r-order short-range dependence on interval I if $\sup_{t \in I} \|G(t, \mathcal{F}_0)\|_r < \infty$, $\delta_r(H, k, I) = O(\chi^k)$, for some $\chi \in (0, 1)$, $r \ge 1$, and s-order locally stationary on interval I, $s \ge 2$, if $G(t, \mathcal{F}_0) \in \operatorname{Lip}_s(I)$.

In the functional linear model (1.1), we consider $x_{i,n} = W(t_i, \mathcal{F}_i)$, $e_{i,n} = H(t_i, \mathcal{F}_i)$, where $t_i = i/n$, , W and H are measurable nonlinear filters mapping from $[0,1] \times \mathbb{R}^Z$ to \mathbb{R}^p and $(-\infty,1] \times \mathbb{R}^Z$ to \mathbb{R} , respectively. Permitting i to approach $-\infty$ enables us to develop our proposed method under long memory, see Section 5.2 for detailed discussion. We further assume $e_{i,n}$ has the following state-heteroscedastic form, i.e., $e_{i,n} = \tilde{H}(t_i, \mathcal{H}_i)\tilde{G}(t_i, \mathcal{G}_i)$ for $-\infty < i \leq n$, where the nonlinear filters $\tilde{H}(t, \cdot)$ and $\tilde{G}(t, \cdot)$ are \mathcal{H}_i and \mathcal{G}_i measurable functions for $t \in (-\infty, 1]$, and filtrations \mathcal{G}_i and \mathcal{H}_i are sub- σ -fields of \mathcal{F}_i , independent of each other. Furthermore, $\mathcal{H}_i \subset \sigma(x_{1,n}, ..., x_{i,n})$ if $1 \leq i \leq n$, and for any $t \in (-\infty, 1]$, $E(\tilde{G}(t, \mathcal{G}_i)) = 0$. The above formulation admits the heteroscedastic errors considered in (He and Zhu, 2003) and (Kulik and Wichelhaus, 2012), where they assume $\tilde{H}(t_i, \mathcal{H}_i)$ to have the form of $s(x_{i,n})$ for some unknown smooth function $s(\cdot)$. Moreover, $\tilde{G}(t_i, \mathcal{G}_i)$ allows the conditional heteroscedasticity as considered by assumption 1(b) of (Cavaliere et al., 2017). Further, let $U(t_i, \mathcal{F}_i) = W(t_i, \mathcal{F}_i)H(t_i, \mathcal{F}_i)$ such that $x_{i,n}e_{i,n} = U(t_i, \mathcal{F}_i)$ (i = 1, ..., n). The time-varying long-run covariance function for the functional linear model (1.1) is defined by

$$\Sigma(t) = \sum_{j=-\infty}^{\infty} \operatorname{cov} \left\{ U\left(t, \mathcal{F}_{0}\right), U\left(t, \mathcal{F}_{j}\right) \right\} \quad (t \in [0, 1]).$$

$$(2.1)$$

For stationary $x_{i,n}$ and $e_{i,n}$, $\Sigma(t)$ will be time invariant. In this case for linear models with no structural changes, (Zhou and Shao, 2013) proposes a self-normalization method for statistical inference so that the estimation of the long-run covariance matrix can be avoided. However, their method relies crucially on strict stationarity. Therefore, for the statistical inference of non-stationary time series linear models, the accurate and robust estimation of $\Sigma(t)$ is essential.

3 The debiased difference-based estimator

When p = 1, (Dette and Wu, 2019) proposes the following difference-based estimator $\Sigma(t)$ based on the difference of $x_{i,n}y_{i,n}$ for the long run variance of $e_{i,n}$. Let $Q_{k,m} = \sum_{i=k}^{k+m-1} x_{i,n}y_{i,n}$, and for $t \in [m/n, 1-m/n]$,

$$\dot{\Sigma}(t) = \sum_{j=m}^{n-m} \frac{m\Delta_j \Delta_j^{\top}}{2} \omega(t,j), \quad \Delta_j = \frac{Q_{j-m+1,m} - Q_{j+1,m}}{m},$$

where for some bandwidth τ_n and the kernel function $K(\cdot)$ with support (-1, 1),

$$\omega(t,i) = K_{\tau_n}(t_i - t) / \sum_{i=1}^n K_{\tau_n}(t_i - t), \quad K_{\tau_n}(\cdot) = K(\cdot/\tau_n).$$

For $t \in [0, m/n)$, $\dot{\Sigma}(t) = \dot{\Sigma}(m/n)$ and for $t \in (1 - m/n, 1]$, $\dot{\Sigma}(t) = \dot{\Sigma}(1 - m/n)$.

Unfortunately, in Appendix F.2, we find that the difference-based estimator $\hat{\Sigma}(t)$ is asymptotically biased for $p \geq 2$. The bias is negligible when the covariates are fixed and continuous as assumed in Assumption 1 of (Zhou et al., 2015), since the bias is caused by the stochastic variation of covariates. We explain this in detail in Section A of the supplemental material. In Appendix F.2, we also show that $E(\hat{\Sigma}(t)) - \Sigma(t)$ can be uniformly approximated by the expectation of the following $\Sigma^A(t)$:

$$\Sigma^{A}(t) = \sum_{j=m}^{n-m} \frac{mA_{j,m}A_{j,m}^{\top}}{2} \omega(t,j) \quad (t \in [m/n, 1 - m/n]),$$

where for $j = m, \ldots, n$,

$$A_{j,m} = \frac{1}{m} \sum_{i=j-m+1}^{j} \{ x_{i,n} x_{i,n}^{\top} \beta(t_i) - x_{i+m,n} x_{i+m,n}^{\top} \beta(t_{i+m}) \},\$$

and $\Sigma^{A}(t) = \Sigma^{A}(m/n)$ for $t \in [0, m/n)$, $\Sigma^{A}(t) = \Sigma^{A}(1 - m/n)$ for $t \in (1 - m/n, 1]$. To utilize $\Sigma^{A}(t)$, we shall substitute $\beta(\cdot)$ in $A_{j,m}$ via an estimator without introducing additional tuning parameters. For this purpose, define $\tilde{Y}_{i,m} = x_{i,n}y_{i,n} - x_{i+m,n}y_{i+m,n}$, $\tilde{X}_{i,m} = x_{i,n}x_{i,n}^{\top} - x_{i+m,n}x_{i+m,n}^{\top}$, and $\tilde{E}_{j,m} = \frac{1}{m}\sum_{i=j-m+1}^{j}(x_{i,n}e_{i,n} - x_{i+m,n}e_{i+m,n})$. By the continuity of $\beta(t)$,

$$\frac{1}{m} \sum_{i=j-m+1}^{j} \tilde{Y}_{i,m} = A_{j,m} + \tilde{E}_{j,m} \approx \frac{1}{m} \sum_{i=j-m+1}^{j} \tilde{X}_{i,m} \beta(t_i) + \tilde{E}_{j,m}$$
$$\approx \frac{1}{m} \beta(t_j) \sum_{i=j-m+1}^{j} \tilde{X}_{i,m} + \tilde{E}_{j,m}.$$
(3.1)

Since $E(\tilde{E}_{j,m}) = 0$, the random vector $\sqrt{m\tilde{E}_{j,m}}$ is $O_p(1)$. Treating $\sum_{i=j-m+1}^{j} \tilde{Y}_{i,m}/\sqrt{m}$ as the response variable, $\sum_{i=j-m+1}^{j} \tilde{X}_{i,m}/\sqrt{m}$ as covariates and $\sqrt{m\tilde{E}_{j,m}}$ as errors, (3.1) motives us to approximate $\beta(t)$

in $A_{j,m}$ by

$$\check{\beta}(t) = \Omega^{-1}(t)\varpi(t), \tag{3.2}$$

where $\Omega(t)$ and $\varpi(t)$ are the smoothed versions of

$$\dot{\Delta}_j/2 = \frac{1}{2m} \sum_{i=j-m+1}^j \tilde{X}_{i,m} \tilde{X}_{i,m}^{\top} \text{ and } \breve{\Delta}_j/2 = \frac{1}{2m} \sum_{i=j-m+1}^j \tilde{X}_{i,m}^{\top} \tilde{Y}_{i,m},$$

i.e., for $t \in [m/n, 1 - m/n]$,

$$\Omega(t) = \sum_{j=m}^{n-m} \Delta_j \omega(t,j)/2, \quad \varpi(t) = \sum_{j=m}^{n-m} \breve{\Delta}_j \tilde{\omega}(t,j)/2$$

where $\tilde{\omega}(t,i) = K\{(t_i - t)/\tau_n^{3/2}\}/\sum_{i=1}^n K\{(t_i - t)/\tau_n^{3/2}\}$, while for $t \in [0, m/n)$, $\Omega(t) = \Omega(m/n)$, $\varpi(t) = \varpi(m/n)$ and for $t \in (1 - m/n, 1]$, $\Omega(t) = \Omega(1 - m/n)$, $\varpi(t) = \varpi(1 - m/n)$. The estimator $\check{\beta}(t)$, which is also based on difference series, is accurate except in the vicinity of abrupt changes. Fortunately, the effect of abrupt changes can be mitigated by the local averaging in the formula of $A_{j,m}$. Replacing $\beta(t)$ by $\check{\beta}(t)$ in $A_{j,m}$ we obtain

$$\hat{A}_{j,m} = \frac{1}{m} \sum_{i=j-m+1}^{j} \{ x_{i,n} x_{i,n}^{\top} \breve{\beta}(t_i) - x_{i+m,n} x_{i+m,n}^{\top} \breve{\beta}(t_{i+m}) \},\$$

and the corresponding debiased difference-based estimator $\hat{\Sigma}(t)$ for $t \in [0, 1]$:

$$\hat{\Sigma}(t) = \hat{\Sigma}(t) - \breve{\Sigma}(t), \quad \breve{\Sigma}(t) = \sum_{j=m}^{n-m} \frac{m\hat{A}_{j,m}\hat{A}_{j,m}^{\top}}{2}\omega(t,j), \quad (3.3)$$

which is robust to structural breaks in regression coefficient functions due to differencing, and except for m, τ_n used for $\Sigma(t)$, the correction $\Sigma(t)$ does not involve additional tuning parameters. Thanks to the residual-free formula, the estimator (3.3) can preserve its consistency even when it is challenging to estimate $\beta(t)$ accurately.

4 Consistency under smooth structural changes

In this section, we discuss the uniform convergence of the debiased difference-based long-run covariance matrix estimator (3.3) for the functional linear model (1.1) under smooth structural changes and shortrange dependence with locally stationary predictors and errors, which accommodates many null hypotheses of nonparametric specification tests. The convergence of $\hat{\Sigma}(t)$ when both smooth and abrupt changes occur is deferred to Section 5.1, and the performance of $\hat{\Sigma}(t)$ under long memory is postponed to Section 5.2.

Assumption 4.1. Each coordinate of $\beta(t)$ lies in $C^{3}[0,1]$.

Assumption 4.1 imposes smooth structural change, i.e., the coefficient function $\beta(\cdot)$ is smooth. In Section 5.1, we shall relax Assumption 4.1 to allow abrupt structural changes.

Assumption 4.2. The value $\lambda_{\min}(\Sigma(t))$ is bounded away from 0 uniformly on [0, 1], and each element of $\Sigma(t) \in C^2[0, 1]$.

Assumption 4.2 guarantees the non-degeneracy of the long-run covariance matrix of the process of $U(t, \mathcal{F}_i)$ and each component of $\Sigma(t)$ is smooth, which are common in the analysis of functional linear models of locally stationary time series, see for example (Zhou and Wu, 2010). Let $J(t, \mathcal{F}_0) = W(t, \mathcal{F}_0)W^{\top}(t, \mathcal{F}_0)$. Define $\mu_W(t) = E\{W(t, \mathcal{F}_0)\}, M(t) = E\{J(t, \mathcal{F}_0)\}, t \in [0, 1].$

Assumption 4.3. Each element of the functions $M(t) \in C^1[0,1]$, $\mu_W(t) \in C^1[0,1]$, and $\inf_{t \in [0,1]} \lambda_{\min}(M(t)) > 0$. The covariate process $W(t, \mathcal{F}_i)$ (i = 1, ..., n) is of 4-order local stationarity and 16 κ -order short-range dependence on [0,1] for some $\kappa \geq 1$.

Assumption 4.3 requires that M(t) is non-degenerate, implies that $\sup_{t \in [0,1]} ||J(t, \mathcal{F}_0)||_{8\kappa} < \infty$, $\delta_{8\kappa}(J, k) = O(\chi_1^k)$ for some $\chi_1 \in (0,1)$, and ensures that each element of $(x_{i,n} x_{i,n}^{\top})$ is 2-order locally stationary.

The following assumption ensures the invertibility of $\Omega(t)$ in (3.2).

Assumption 4.4. Each element of the covariance function $\operatorname{cov} \{J(t, \mathcal{F}_0), J(t, \mathcal{F}_0)\} \in C^2[0, 1]$ and its smallest eigenvalue is strictly positive on [0, 1].

Assumption 4.5. The process $H(t, \mathcal{F}_i)$ $(i \in \mathbb{Z})$ satisfies 16κ -order of short-range dependence and 4-order of local stationarity on [0, 1].

Assumptions 4.3 and 4.5 ensure that $U(t, \mathcal{F}_i)$ $(i \in Z)$ is of 2-order local stationarity and 8κ -order of short-range dependence on [0, 1]. The existence of 16 κ th moments of covariates and errors are assumed for technical convenience, and it is satisfied by sub-exponential random variables. We conjecture that it can be relaxed by substantially more involved mathematical arguments, see the simulation study of (Bai and Wu, 2024). Specifically, the condition of 16 κ -order dependence can be replaced by sub-exponential moment conditions and dependence measure in \mathcal{L}^1 norm. Further technical discussion on Assumption 5 can be found in Appendix C of the supplement. Assumption 4.4 is mild, and it excludes the scenario in which all of the time series covariates reduce to deterministic smooth trends; in this case, we recommend using the direct multivariate extension of the difference-based long-run covariance estimator of (Dette and Wu, 2019), namely $\hat{\Sigma}(t)$. The following assumption gives the properties of the kernel function. The use of different kernels is discussed in Appendix B.

Assumption 4.6. The kernel function $K(\cdot)$ is a continuously differentiable, symmetric density function and supported on (-1, 1).

Let $I = [\gamma_n, 1 - \gamma_n] \subset (0, 1)$, where $\gamma_n = \tau_n + (m+1)/n$.

Theorem 4.1. Under Assumptions 4.1, 4.2, 4.3, 4.4, 4.6 and 4.5 with constant $\kappa \geq 1$, suppose $m = O(n^{1/3}), m \to \infty, \tau_n \to 0, m/(n\tau_n^{3/2+2/\kappa}) \to 0, \tau_n^{3-1/\kappa}\sqrt{m} \to 0, m\tau_n \to \infty$, we have

$$\sup_{t \in I} \left| \hat{\Sigma}(t) - \Sigma(t) \right| = O_p \left(\frac{m^{1/2}}{n^{1/2} \tau_n^{3/4 + 1/\kappa}} + \frac{1}{m} + m^{1/2} \tau_n^{3 - 1/\kappa} \right).$$

The above equation and the bandwidth conditions for Theorem 4.1 ensure the uniform consistency of $\hat{\Sigma}(t)$. As shown by Lemma F.1 in the supplement, the estimator $\check{\beta}(t)$ is consistent when there are no jump points. As pointed out by a referee, it is viable to use any pilot estimator of $\beta(t)$ that satisfies the conditions in Appendix C of the supplement in $\check{\Sigma}(t)$ for debiasing.

We compare our results with (Zhou and Wu, 2010) which proposes a plug-in estimator of long-run covariance matrix using nonparametric residuals. According to Theorem 5 in (Zhou and Wu, 2010), their best approximation rate can be close to but not faster than $n^{-1/4}$. In contrast, our uniform rate in Theorem 4.1 is $n^{-4/15+2/(15\kappa)}$ by taking $m \simeq n^{4/15}$, $\tau_n \simeq n^{-2/15}$, which is better when κ is sufficiently large. In unreported studies, we find that the performance of plug-in estimators using $\breve{\beta}(\cdot)$ is superior to (Zhou and Wu, 2010) but inferior to our proposed estimator. This estimator is also available in R package mlrv.

5 Applications

5.1 Structural change detection with monotonic power

The first application is the detection of structural changes in the stochastic linear regression

$$y_{i,n} = x_{i,n}^{\top} \beta_{i,n} + e_{i,n} \quad (i = 1, \dots, n),$$

where $(x_{i,n})_{i=1}^n$ is the *p*-dimensional covariate process and $(e_{i,n})_{i=1}^n$ is the error process. The test for structural changes of $(\beta_{i,n})$ considers the hypothesis $H_0: \beta_{1,n} = \beta_{2,n} = \cdots = \beta_{n,n}$. Allowing general non-stationarity in the covariates and the errors, (Wu and Zhou, 2018) proposes to use the test statistic

$$T_n = \max_{1 \le j \le n} \left| \sum_{i=1}^{j} \hat{e}_{i,n} x_{i,n} / \sqrt{n} \right|, \quad \hat{e}_{i,n} = y_{i,n} - x_{i,n}^{\top} \hat{\beta}_n \quad (i = 1, \dots, n),$$

where $\hat{\beta}_n = \operatorname{argmin}_{\beta} \sum_{i=1}^n (y_{i,n} - x_{i,n}^{\top}\beta)^2$. For the alternative hypothesis, we consider

$$y_{i,n} = x_{i,n}^{\top} \beta(i/n) + e_{i,n}, \quad \beta(t) = \beta + L_n g(t) \quad (i = 1, \dots, n; \ t \in [0, 1]),$$
(5.1)

where $\beta, g(t) \in \mathbb{R}^p$, g(0) = 0, L_n is a positive real sequence, $\beta(t)$ has potential abrupt changes, i.e., $\beta(t) = \sum_{j=0}^{q_n} b_j(t) 1(a_j \leq t < a_{j+1}), \ 0 = a_0 < a_1 < \cdots < a_{q_n} < a_{q_n+1} = 1, \ b_j(t) \in \mathcal{C}^3(a_j, a_{j+1}),$ $\sup_{0 \leq j \leq q_n} |b_j(a_j) - b_j(a_j^-)| < \infty$, and q_n is the number of abrupt changes. (Wu and Zhou, 2018) proposes a general bootstrap statistics according to (5.2), which relies on the residual $\hat{e}_{i,n} = y_{i,n} - x_{i,n}^{\top} \hat{\beta}_n$. Their test can be applied to piecewise locally stationary covariates and errors, and is unified for testing structural changes in general M estimation. They show their power approaches 1 under local alternatives $L_n = n^{-1/2}$ or $L_n = o(1), \sqrt{nL_n} \to \infty$. However, if L_n does not vanish, $\hat{\beta}_n$ is not consistent and the power of the test in (Wu and Zhou, 2018) in this case is not theoretically guaranteed. If we focus on the locally stationary process which is quite general and on least squares regression which is arguably the most widely applied M-estimator in practice, we could improve the power of the test for structural changes via our proposed long-run covariance matrix estimator $\hat{\Sigma}(t)$ and an alternative bootstrap procedure to (Wu and Zhou, 2018). For this purpose, we define

$$F_r = \max_{m \le i \le n-m+1} |\Psi_{i,m}^{(r)} - \hat{\Lambda}(i/n)\hat{\Lambda}^{-1}(1)\Psi_{n-m+1,m}^{(r)}|,$$

where $\Psi_{i,m}^{(r)} = n^{-1/2} \sum_{j=1}^{i} \hat{\Sigma}^{1/2}(t_j) R_j^{(r)}$, $\hat{\Lambda}(i/n) = \sum_{j=1}^{i} x_{j,n} x_{j,n}^{\top}/n$, $(R_j^{(r)})_{j=1}^n$ are the independent and identically distributed standard normal random variables independent of data and are independently generated in the r_{th} bootstrap iteration, and $\hat{\Sigma}(t)$ is the proposed difference-based estimator of $\Sigma(t)$. Let B denote the number of bootstrap iterations. Let $F_{(1)} \leq F_{(2)} \leq \cdots \leq F_{(B)}$ be the order statistics of (F_r) . We reject the structural stability test at the significance level of α if T_n is greater than the $F_{\lfloor (1-\alpha)B \rfloor}$. We proceed to relax Assumption 4.1 in Theorem 4.1 allowing for the possible presence of abrupt changes and discuss the property of the bootstrap procedure.

Theorem 5.1. Under the Assumptions 4.2, 4.3, 4.4, 4.6 and 4.5 with constant $\kappa \geq 1$, and the bandwidth conditions $m = O(n^{1/3}), \tau_n \to 0, n\tau_n^{7/2}/m \to \infty, m\tau_n^3 \to \infty$, assuming $q_n = o\{\min(\tau_n^{-1/2}, n\tau_n/m^2)\}$, under the alternative hypothesis (5.1), we have

$$\sup_{t \in \mathcal{I}} |\hat{\Sigma}(t) - \Sigma(t)| = o_p(1),$$

and as a result there exists a p-dimensional zero-mean Gaussian process Z(t) with covariance function $\gamma(t,s) = \int_0^{\min(t,s)} \Sigma(r) dr$ such that

$$F_r \Rightarrow \sup_{t \in (0,1]} |G(t)| = \sup_{t \in (0,1]} |Z(t) - \Lambda(t)\Lambda^{-1}(1)Z(1)|, \quad \Lambda(t) = \int_0^t M(s)ds.$$

Taking $m \simeq n^{4/15}$, $\tau_n \simeq n^{-2/15}$, which can achieve $q_n = o(n^{1/15})$ for sufficiently large κ , which allows the number of abrupt jumps diverges as $n \to \infty$, though the estimator $\breve{\beta}(t)$ is inconsistent due to the inconsistency of smoothing in the neighborhood of discontinuous points. Define

$$\Lambda(t) = \int_0^t M(s)ds, \quad \Lambda(s, g(\cdot)) = \int_0^s M(r)g(r)dr$$

and $F(t, g(\cdot)) = \Lambda(t, g(\cdot)) - \Lambda(t)\Lambda^{-1}(1)\Lambda(1, g(\cdot))$. The following proposition gives the limiting distribution of the test statistic T_n under the null hypothesis and ensures the monotonic power of the bootstrap procedure under $H_A: \beta(t) = \beta + L_n g(t)$.

Proposition 5.1. (i) Under the conditions of Theorem 4.1 and the null hypothesis of no structural changes, we have

$$T_n \Rightarrow \sup_{t \in (0,1]} |G(t)|, \tag{5.2}$$

where G(t) is as defined in Theorem 5.1.

(ii) Under the conditions of Theorem 5.1 and the alternative hypothesis (5.1) with $L_n = O(1)$, $n^{1/2}L_n \to \infty$, we have $T_n \to \infty$ in probability at the rate $\sqrt{nL_n}$, and

$$\left| \operatorname{pr}(T_n \ge \hat{q}_{1-\alpha}) - \operatorname{pr}\left(\sup_{t \in [0,1]} |G(t) + n^{1/2} L_n F(t, g(\cdot))| \ge \hat{q}_{1-\alpha} \right) \right| = o(1),$$

where $\hat{q}_{1-\alpha}$ is the bootstrap critical value of F_r at the significance level α .

Proposition 5.1 shows that T_n is of order $\max(1, \sqrt{nL_n})$ under null hypothesis. Theorem 5.1 and Proposition 5.1 guarantee the asymptotic correctness of the bootstrap procedure with the differencebased estimator and that its asymptotic power approaches 1 under the fixed alternative. Meanwhile, the bootstrap procedure with the difference-based estimator can detect the local alternatives at the parametric rate \sqrt{n} in the sense that if $L_n = n^{-1/2}$, $T_n \Rightarrow \sup_{t \in [0,1]} |G(t) + F(t,g(\cdot))|$. In contrast, Proposition B.1 of (Wu and Zhou, 2018) shows that with the ordinary least squares residuals, the magnitude of F_r using their bootstrap procedure is $\sqrt{m} \max(L_n, \log^2 n/\sqrt{n}) \log n$ when $\sqrt{mL_n} \to \infty$, $L_n \to 0$, implying power loss due to the divergence of F_r . Therefore, the bootstrap procedure equipped with the difference-based estimator will be more powerful than that with ordinary least squares residuals and overcome non-monotonic power caused by the inflation of the bootstrap statistics in (Wu and Zhou, 2018). In earlier work on the remedy of the non-monotonic power for the test of smooth structural changes, (Juhl and Xiao, 2009) proposes to estimate the long-run variance via plugging in the nonparametric residuals for p = 1. However, the improvement in the power of their approach does not carry over in the presence of abrupt structural changes.

5.2 Testing for long memory

Another application is testing for long memory in the functional linear model

$$y_{i,n} = x_{i,n}^{\top} \beta(t_i) + (1 - \mathcal{B})^{-d} e_{i,n} \quad (i = 1, \dots, n),$$

where \mathcal{B} is the lag operator, $d \in [0, 1/2)$ is the long-memory parameter. When d = 0, the error process of the model is locally stationary and short-range dependent. We are interested in the following hypothesis testing problem

$$H_0: d = 0$$
 versus $H_A: 0 < d < 1/2.$ (5.3)

The rejection of H_0 implies that the short-memory linear model is inadequate for the data and long-range dependence should be considered. (Bai and Wu, 2024) proposes to test H_0 using the jackknife corrected nonparametric residuals. They obtain the local linear estimate of $\beta(\cdot)$, i.e.,

$$(\hat{\beta}_{b_n}(t), \hat{\beta}'_{b_n}(t)) = \underset{\eta_0, \eta_1 \in \mathbb{R}^p}{\arg\min} \sum_{i=1}^n \{y_{i,n} - x_{i,n}^\top \eta_0 - x_{i,n}^\top \eta_1(t_i - t)\}^2 K_{b_n}(t_i - t),$$
(5.4)

where K(t) is a kernel function with finite support (-1, 1), b_n is a bandwidth. Then, they consider the jackknife estimator $\tilde{\beta}_{b_n}(t) = 2\hat{\beta}_{b_n/\sqrt{2}}(t) - \hat{\beta}_{b_n}(t)$ of which the asymptotic bias terms involving $\beta''(\cdot)$ in the formula of $\hat{\beta}_{b_n}$ and $\hat{\beta}_{b_n/\sqrt{2}}$ are canceled. Let $K^*(\cdot)$ denote the jackknife equivalent kernel $2\sqrt{2K}(\sqrt{2x}) - K(x)$. For the sake of simplicity, we write n' as $\lfloor nb_n \rfloor$ for short. Define the nonparametric residuals and their partial sum as $\tilde{e}_{i,n} = y_{i,n} - x_{i,n}^{\top} \tilde{\beta}_{b_n}(t_i)$ and $\tilde{S}_{r,n} = \sum_{i=n'+1}^r \tilde{e}_{i,n}, r = n'+1, \ldots, n-n'$, respectively. The KPSS, R/S, V/S and K/S-type test statistics of (Bai and Wu, 2024) are

- 1. KPSS-type statistic $K_n = \frac{1}{n(n-2n')} \sum_{r=n'+1}^{n-n'} \left(\tilde{S}_{r,n} \right)^2$.
- 2. R/S-type statistic $Q_n = \max_{n'+1 \le k \le n-n'} \tilde{S}_{k,n} \min_{n'+1 \le k \le n-n'} \tilde{S}_{k,n}$.
- 3. V/S-type statistic $M_n = \frac{1}{n(n-2n')} \left\{ \sum_{k=n'+1}^{n-n'} \tilde{S}_{k,n}^2 \frac{1}{n-2n'} \left(\sum_{k=n'+1}^{n-n'} \tilde{S}_{k,n} \right)^2 \right\}.$
- 4. K/S-type statistic $G_n = \max_{n'+1 \le k \le n-n'} \left| \tilde{S}_{k,n} \right|$.

(Bai and Wu, 2024) proposes to implement the above tests via the following bootstrap-assisted procedure. Define $\hat{M}(t) = \sum_{i=1}^{n} x_{i,n} x_{i,n}^{\top} K_{\eta_n}(t_i - t^*)/(n\eta_n)$, where $t^* = \max\{\eta_n, \min(t, 1 - \eta_n)\}$ for some bandwidth $\eta_n \to 0$, $n\eta_n^2 \to \infty$. Let $\hat{\Sigma}^*(\cdot)$ be any consistent long-run covariance matrix estimator satisfying the regularity condition 5.1 in their paper, and $\hat{\sigma}_H^{*2}(t) = (\hat{\Sigma}^*(t))_{1,1}$. Generate *B* independent and identically distributed copies of $N(0, I_p)$ vectors $V_i^{(r)} = (V_{i,1}^{(r)}, ..., V_{i,p}^{(r)})^{\top}$ (r = 1, ..., B), and for each *r* calculate

$$\tilde{G}_{k}^{(r)} = -\sum_{j=1}^{n} \left\{ \frac{1}{nb_{n}} \sum_{i=n'+1}^{k} x_{i,n}^{\top} \hat{M}^{-1}(t_{i}) K_{b_{n}}^{*}(t_{i}-t_{j}) \right\} \hat{\Sigma}^{*,1/2}(t_{j}) V_{j}^{(r)} + \sum_{i=n'+1}^{k} \hat{\sigma}_{H}^{*}(t_{i}) V_{i,1}^{(r)}$$

as well as the bootstrap statistics: $\widetilde{K}_n^{(r)}$, $\widetilde{RS}_n^{(r)}$, $\widetilde{VS}_n^{(r)}$ and $\widetilde{KS}_n^{(r)}$ which can be obtained by substituting $\widetilde{S}_{k,n}$ in the corresponding statistics by $\widetilde{G}_k^{(r)}$. Let $\widetilde{K}_{n,(1)} \leq \widetilde{K}_{n,(2)} \leq \cdots \leq \widetilde{K}_{n,(B)}$ be the ordered statistics of $\widetilde{K}_n^{(r)}$ $(r = 1, \ldots, B)$, and $B^* = \max\{r : \widetilde{K}_{n,(r)} \leq K_n\}$. Then the *p*-value of the KPSS-type test is $1 - B^*/B$, and the *p*-values of R/S, V/S, and K/S-type tests can be obtained similarly. Given a nominal level α , if the *p*-value is smaller than α , we reject the null hypothesis of short memory.

In this paper we propose to set $\hat{\Sigma}^*(t) = \hat{\Sigma}(t)$. With the new difference-based long-run covariance matrix estimator, our testing procedure will be more robust than that using the plug-in estimator for $\hat{\Sigma}^*(t)$, since for the latter procedure, the parameter b_n in (5.4) for the estimation of the regression coefficients will additionally affect the estimate of the long-run covariance matrix through the nonparametric residuals $\tilde{e}_{i,n}$ as well as the selection of m and τ_n , as indicated by the discussion of Theorem 5 in (Zhou and Wu, 2010) that m and τ_n should be chosen from an interval determined implicitly by b_n . Moreover, adopting $\hat{\Sigma}(t)$ leads to more accurate type-I error control due to the faster convergence rate, see the discussion below Theorem 4.1.

In the following, we show the validity of $\hat{\Sigma}(t)$ via studying the asymptotic behavior of $\hat{\Sigma}(t)$ under the fixed and local alternatives for the testing problem (5.3), which is essential for the consistency of the aforementioned bootstrap tests.

Assumption 5.1. Assumption 4.5 holds over $(-\infty, 1]$, $H(t, \mathcal{F}_i)$ $(i \in Z)$ is of 2κ -order local stationarity

on $(-\infty, 1]$, and its long-run variance function

$$\sigma_{H}^{2}(t) = \sum_{k=-\infty}^{\infty} \operatorname{cov} \left\{ H\left(t, \mathcal{F}_{0}\right), H\left(t, \mathcal{F}_{k}\right) \right\} \quad (t \in (-\infty, 1]),$$

satisfies that $\inf_{t \in (-\infty,1]} \sigma_H^2(t) > 0$, $\sup_{t \in (-\infty,1]} \sigma_H^2(t) < \infty$, and $\sigma_H^2(\cdot)$ is twice continuous differentiable on [0,1].

Theorem 5.2. Under Assumptions 4.1, 4.3, 4.4, 4.6 and 5.1, assuming $m\tau_n^{3/2}/\log n \to \infty$, $\tau_n \to 0$, $n\tau_n^3 \to \infty$, $m/(n\tau_n^3) \to 0$, $\tau_n^{3-1/\kappa}\sqrt{m} \to 0$, $m = O(n^{1/3})$, $\kappa \ge \max\{4/(1/2-d), 2/(3d), 4\}$, it follows that under H_A

$$\sup_{t\in I} \left| m^{-2d} \hat{\Sigma}(t) - \kappa_2(d) \sigma_H^2(t) \mu_W(t) \mu_W^\top(t) \right| = o_p(1),$$

where $\kappa_2(d) = \Gamma^{-2}(d+1) \int_0^\infty \{t^d - (t-1)^d_+\} \{2t^d - (t-1)^d_+ - (t+1)^d\} dt.$

Theorem 5.2 shows that the proposed difference-based estimator $\hat{\Sigma}(t)$ in (3.3) inflates at the rate of m^{2d} under long-range dependence with parameter d > 0, while its limit normalized by m^{2d} depends on the $\mu_W(t)$ along with the long-run variance of $e_{i,n}$ and the long-memory parameter d. The long-memory parameter d also affects the theoretical properties of the long-run variance-covariance estimate through the moment condition of κ . The exact convergence rate is displayed in Step 6 of the proof of Theorem 5.2 in supplement due to page limit. In the following Theorem 5.3, we investigate the performance of the estimator $\hat{\Sigma}(t)$ under the local alternatives $d_n = c/\log n$ for some constant c > 0. For this purpose, we define the long-run cross covariance vector between the locally stationary processes $U(t, \mathcal{F}_i)$ $(i \in Z)$ and $H(t, \mathcal{F}_j)$ $(j \in Z)$.

Definition 5.1. Define the long-run cross-covariance vector $s_{UH}(t) \in \mathbb{R}^p$ by

$$s_{UH}(t) = \sum_{j=-\infty}^{\infty} \operatorname{Cov}\{U(t, \mathcal{F}_0), H(t, \mathcal{F}_j)\} \quad (t \in [0, 1]).$$

For given constants c > 0 and $\alpha_1 \in (0, 1)$, define for $0 \le t \le 1$, the symmetric matrix

$$\check{\Sigma}(t) = \Sigma(t) + (e^{c\alpha_1} - 1)^2 \sigma_H^2(t) \mu_W(t) \mu_W^\top(t) + (e^{c\alpha_1} - 1) \{s_{UH}(t) \mu_W^\top(t) + \mu_W(t) s_{UH}^\top(t)\}.$$

The following Assumption 5.2 guarantees that $\check{\Sigma}(t)$ is smooth and non-degenerate.

Assumption 5.2. $\check{\Sigma}(\cdot) \in C^2[0,1]$, and $\lambda_{\min}\{\check{\Sigma}(t)\}$ is bounded above 0 on [0,1].

Since $\Sigma(t)$ and $(e^{c\alpha} - 1)^2 \sigma_H^2(t) \mu_W(t) \mu_W^{\top}(t)$ are positive definite, by Weyl's inequality Assumption 5.2 is satisfied for sufficiently small positive c.

Theorem 5.3. Let Assumptions 4.1, 4.3, 4.4, 4.6, 5.1 and 5.2 be satisfied. If $m\tau_n^{3/2} \to \infty$, $\tau_n \to 0$, $m/(n\tau_n^3) \to 0$, $\tau_n^{3-1/\kappa} \sqrt{m} \to 0$, $m = \lfloor n^{\alpha_1} \rfloor$, $\alpha_1 \in (0, 1/3)$, we have

$$\sup_{t \in I} \left| \hat{\Sigma}(t) - \check{\Sigma}(t) \right| = o_p(1).$$

Theorem 5.2 and Theorem 5.3 lead to the desired limiting distribution of the bootstrap statistics under the fixed and local alternatives achieving satisfactory power performance in finite samples, see (Bai and Wu, 2024) for theoretical justification and numerical evidence. In finite samples, we demonstrate that long memory tests with difference-based long-run covariance matrix estimates can achieve sizes closer to the nominal level and are more powerful than their counterparts using plug-in estimates, see Section 6.

6 Simulation

6.1 Setting

We elaborate the procedure of tuning parameter selection, which is available in the R package mlrv, and display the values of the parameters selected in Appendix B. Let $(\varepsilon_l)_{l\in \mathbb{Z}}, (\zeta_l)_{l\in \mathbb{Z}}, (\eta_l)_{l\in \mathbb{Z}}$ be N(0,1), $\vartheta_i = (\eta_i + \varepsilon_i)/2$, and consider the filtrations

$$\mathcal{F}_j = (\dots, \zeta_{j-1}, \zeta_j), \quad \mathcal{G}_j = (\dots, \varepsilon_{j-1}, \varepsilon_j), \quad \mathcal{H}_j = (\dots, \varepsilon_j, \eta_j) \quad (j = -\infty, \dots, n).$$

6.2 Testing for structural changes

We generate the locally stationary process $x_{i,n,1}$ from $G_1(t, \mathcal{H}_i) = \sum_{j=0}^{\infty} (0.5 - 0.5t)^j \vartheta_{i-j}$, the locally stationary process $x_{i,n,2}$ from $G_2(t, \mathcal{H}_i) = \sum_{j=0}^{\infty} \{0.25 + 0.5(t - 0.5)^2\}^j \epsilon_{i-j}$, and the locally stationary process $u_{i,n}$ from $G(t, \mathcal{F}_i) = 0.65 \cos(2\pi t) G(t, \mathcal{F}_{i-1}) + \zeta_i$. We consider the following heteroscedastic linear regression model:

$$y_{i,n} = 1 + m_{i,n} + x_{i,n,1} + x_{i,n,2} + e_{i,n}, \quad e_{i,n} = (1 + 0.1x_{i,n,1})u_{i,n}, \quad (i = 1, \dots, n),$$

where the function $m_{i,n}$ (i = 1, ..., n) includes the following scenarios

CP1: $m_{i,n} = 2\delta \sin(2\pi t_i) x_{i,n,1} 1(0.5 \le t_i \le 1).$

CP2: $m_{i,n} = \delta \sin(2\pi t_i) 1 (0 \le t_i \le 0.4) + \delta x_{i,n,1} 1 (0.7 \le t_i \le 1)/2.$

CP4: $m_{i,n} = 1.5\delta \sin(2\pi t_i) 1 (0 \le t_i \le 0.2 \text{ or } 0.4 \le t_i \le 0.6 \text{ or } 0.8 \le t_i \le 1).$

We conduct our simulation with a sample size 300. As shown in Figure 6.1, when there are 4 change points, the block bootstrap test based on ordinary least squares residuals ((Wu and Zhou, 2018)) suffers from low and non-monotonic power that can not approach 1. By contrast, the newly proposed differencebased long-run covariance matrix estimator enhances the simulated power significantly and addresses the non-monotonic power issue by taking the difference. To further illustrate the impact of long-run covariance matrix estimators in bootstrap tests of structural breaks, we investigate the estimation accuracy of both long-run covariance matrix estimators and find that our proposed estimator halves the empirical mean



Figure 6.1: The empirical rejection rates of gradient-based structural change point tests as δ increases from 0 to 1 with sample size n = 300 and simulation times 2000, under three scenarios CP1(blue), CP2(orange), CP4(red), using blocks of ordinary least squares residuals (small-dashes), and difference-based long-run covariance matrix estimator (solid). Left panel: nominal size 0.05; Right panel: nominal size 0.1.

square error in the presence of change points, see Appendix E of the supplement for extra simulation results and sensitivity analysis.

6.3 Testing for long-range dependence

Consider the following heteroscedastic functional linear model,

$$y_{i,n} = \beta_1(t_i) + \beta_2(t_i)x_{i,n} + (1 - \mathcal{B})^{-d}e_{i,n}, \quad (i = 1, \dots, n),$$

where \mathcal{B} is the lag operator, $\beta_1(t) = 4\sin(\pi t), \beta_2(t) = 4\exp\{-2(t-0.5)^2\}, x_{i,n} = W(t_i, \mathcal{F}_i) \ (i = 1, ..., n),$ and $e_{j,n} = H(t_j, \mathcal{F}_j, \mathcal{G}_j) \ (j = 1, ..., n),$ where

$$H(t, \mathcal{F}_i, \mathcal{G}_i) = B(t, \mathcal{G}_i) \{ 1 + W^2(t, \mathcal{F}_i) \}^{1/2} \quad (i \in \mathbb{Z}; \ t \in [0, 1]),$$

 $W(t, \mathcal{F}_i) = \{0.1+0.1\cos(2\pi t)\}W(t, \mathcal{F}_{i-1})+0.2\zeta_i+0.7(t-0.5)^2$, and $B(t, \mathcal{G}_i) = \{0.3-0.4(t-0.5)^2\}B(t, \mathcal{G}_{i-1})+0.8\varepsilon_i$. As demonstrated by Figure 6.2, the difference-based long-run covariance matrix estimator yields uniform improvement for the power of KPSS, V/S, R/S, and K/S tests against $0 < d \leq 1/2$ in finite samples. Notably, equipped with the difference-based estimator the simulated power of K/S, R/S, and KPSS tests can reach 1 with the sample size 1500 as d increases to 0.5, while using the plug-in long-run covariance matrix estimator the power is much lower and stays far below 1, except for the V/S test. The corresponding sensitive analysis is in Appendix E.

7 Data analysis

We apply our newly proposed long-run covariance matrix estimator (3.3) to the analysis of Hong Kong hospital data, including structural change detection and tests for long memory. The data set consists of daily hospital admissions in Hong Kong as well as daily measurements of pollutants between January 1,



Figure 6.2: Empirical rejection rates of KPSS(orange), K/S(blue), R/S(red), and V/S(green) tests under different d's with sample size 1500, using the plug-in method (small-dashes) and the difference-based method (solid). Left panel: nominal size 0.05; Right panel: nominal size 0.1.

1994, and December 31, 1995. The sample size is 730. Consider the functional linear model for this data set, i.e.,

$$y_{i,n} = \beta_1(t_i) + \sum_{p=2}^{4} \beta_p(t_i) x_{i,p,n} + \varepsilon_{i,n}, \quad (i = 1, \dots, n),$$

where $(y_{i,n})$ is the series of daily total number of hospital admissions of circulation and respiration and $(x_{i,p,n})$, p = 2, 3, 4, are the series of daily levels of SO₂, NO₂ and dust, respectively, in micrograms per cubic meter.

As illustrated in Section 5 of (Wu and Zhou, 2018), it is of practical concern to test whether $\beta(\cdot) = (\beta_p(\cdot), 1 \leq p \leq 4)^{\top}$ is a constant vector. The test for structural changes equipped with the differencebased estimator yields *p*-value 0.006, which rejects the null hypothesis of no structural change, while the test procedure proposed in (Wu and Zhou, 2018) based on ordinary least squares residuals yields *p*-value greater than 0.1. The different testing results can be attributed to the power loss of (Wu and Zhou, 2018) under structural change, as shown in Figure 6.1.

We then consider the test for the long memory of $\varepsilon_{i,n}$. (Bai and Wu, 2024) performs long-memory tests on each covariate process and concludes that they are short-range dependent. Therefore, we could apply the tests introduced in Section 5.2 to this data, and compare the *p*-values of the tests equipped with the difference-based estimator (3.3) and with the plug-in estimator of (Zhou and Wu, 2010), respectively. The bandwidth b_n in Section 5.2 are selected by the GCV method advocated by (Zhou and Wu, 2010) and (Bai and Wu, 2024). As in Table 7.1, the *p*-values of four types of tests for long memory based on

method	KPSS	R/S	V/S	K/S	method	KPSS	R/S	V/S	K/S
plug	0.300	0.171	0.079	0.356	diff	0.810	0.888	0.835	0.907

Table 7.1: *p*-values of tests for long memory

plug-in estimates are much smaller than those based on $\hat{\Sigma}(t)$. The smaller *p*-values might result from the inaccurate size performance associated with the plug-in estimator, which is evidenced by extra simulation results in Table E.1 of the supplemental material showing that the methods with plug-in estimates tend

to over-reject and result in smaller *p*-values. A further sensitivity check shows that when using $1.2 \times \text{GCV}$ bandwidths, the R/S test with the plug-in estimator yields *p*-value 0.08 rejecting the null hypothesis at the significance level of 10%, while the *p*-values of tests equipped with the difference-based estimator remain large leading to the same decision of accepting the null for all four tests.

8 Conclusion

Additional potential applications can be found in Appendix D of the supplement. The optimal long-run variance for time series with stationary errors has been thoroughly discussed recently by (Chan, 2022). However, the approach therein is not applicable when non-stationarity is present. We leave the optimal estimation of the long-run covariance matrix under time series non-stationarity as rewarding future work. In addition, the generalization of our method beyond linear models will also be of great importance.

Acknowledgement

Weichi Wu is the corresponding author and is supported by National Natural Science Foundation of China 12271287. The authors thank the editors, associate editors and referees for constructive comments.

Supplementary material

In the supplement, we present implementation details, extra simulation studies, the proofs of the findings in this paper as well as auxiliary technical results.

Supplement to "Difference-based covariance matrix estimate in time series nonparametric regression with applications to specification tests"

We organize the supplementary material as follows: Appendix A gives the intuition of the bias in the difference-based estimator. The implementation details including the procedure of selection of tuning parameters are in Appendix B. Appendix C provides some discussion on the assumptions. Appendix D offers other applications of the proposed difference-based estimator. We investigate the sensitivity of finitesample performance of the tests with respect to the smoothing parameters, extra simulation with smaller sample size, and the performance of different estimates in the presence of change points in Appendix E. Appendix F presents proofs of the results in the main paper. Appendix G provides auxiliary results which are used in the proofs.

A Bias in the difference-based estimator

To see this, consider the simple case where x_i and e_i are independent and *i.i.d.* random variables. Then, for the differenced series we have

$$E\{(e_i + \beta_0(i/n) + \beta(i/n)x_i - e_{i-1} - \beta_0((i-1)/n) - \beta((i-1)/n)x_{i-1})^2\}$$

= E\{(e_i - e_{i-1})^2\} + E\{(\beta_0(i/n) - \beta_0((i-1)/n))^2\} + E\{(\beta(i/n)x_i - \beta((i-1)/n)x_{i-1})^2\}

where the third term is close to $2\beta(i/n)\operatorname{var}(x_i)$. Notice that this term is $O(n^{-2})$ if x_i is deterministic and smooth instead. In addition, we examine the bias via empirical studies. We consider the following dependent and independent settings and compare the differenced data simulated from models with the stochastic trend and with only the deterministic smooth trend for a simple illustration:

A. Independent scenario.

A.1 Stochastic trend

$$y_i = 4(i/n - 0.5)^2 + 0.5x_{1,i} + 0.4x_{2,i} + e_i$$
 $(i = 1, ..., n)_i$

where $x_{1,i}$ are independent and identically distributed N(2,1) random variables, $x_{2,i}$ and e_i are independent and identically distributed standard Gaussian variables. Note that this setting is allowed by our Assumptions 1-5.

A.2 Deterministic smooth trend

$$y_i = 4(i/n - 0.5)^2 + 1 + e_i$$
 $(i = 1, ..., n),$

where e_i are independent and identically distributed standard Gaussian variables.

B. Dependent scenario



Figure A.1: The sample path of $(x_{i+1}y_{i+1} - x_iy_i)^2 - (x_{i+1}e_{i+1} - x_ie_i)^2$ with stochastic trend (solid lines), and with only deterministic smooth trend (dotted lines). The left panel shows the gaps between A.1 and A.2 and the right panel shows that between B.1 and B.2, respectively.

B.1 Stochastic trend

$$y_i = 4(i/n - 0.5)^2 + \sin(\pi i/n)x_{1,i} + x_{2,i} + e_i \quad (i = 1, \dots, n),$$
(A.1)

where $x_{1,i}$ is an Autoregressive Moving Average process with autoregressive coefficients 0.8897 and -0.4858, moving average coefficients -0.2279 and 0.2488, i.e.,

$$x_{1,i} - 0.8897x_{1,i-1} + 0.4858x_{1,i-2} = z_i - 0.2279z_{i-1} + 0.2488z_{i-2},$$

where (z_i) are independent and identically distributed standard Gaussian variables, $(x_{2,i})$ are independent and identically distributed standard Gaussian variables, and (e_i) is an autoregressive process with coefficient 0.5, i.e.,

$$e_i = 0.5e_{i-1} + \eta_i,$$

with η_i being independent and identically distributed standard Gaussian variables. Note that this setting is allowed by our Assumption 1-5.

B.2. Deterministic smooth trend

$$y_i = 4(i/n - 0.5)^2 + e_i$$
 $(i = 1, ..., n)$

where e_i is as defined in (A.1).

Figure A.1 displays the sample path of the gap between the differences between $(x_{i+1}y_{i+1} - x_iy_i)^2$ and $(x_{i+1}e_{i+1} - x_ie_i)^2$ under the two scenarios described above. As shown by Figure A.1, the paths of the stochastic trend model are much more jagged than those of the deterministic smooth trend model even in the independent setting, since the deterministic smooth trend is almost eliminated by differencing. Similar gaps can also be observed between the differences of $(x_{i+k}y_{i+k} - x_iy_i)^2$ and $(x_{i+k}e_{i+k} - x_ie_i)^2$ when k > 1. These illustrate the influence of stochastic covariates on the difference-based statistics in approximating

the difference of true errors weighted by covariates, which is in fact the source of the non-negligible bias.

B Implementation details

B.1 Selection of tuning parameters

For refinement, we recommend the following extended minimum volatility method as proposed in Chapter 9 of (Politis et al., 1999) which works quite well in our empirical studies. The extended minimum volatility method has the advantage of robustness under complex dependence structures and does not depend on any parametric assumptions of the time series. To be concrete, we first propose a grid of possible block sizes and bandwidths $\{m_1, m_2, \dots, m_{M_1}\}$, $\{\tau_1, \tau_2, \dots, \tau_{M_2}\}$ from $[\lfloor c_1 n^{4/15} \rfloor, \lfloor c_2 n^{4/15} \rfloor]$ and $[c_3 n^{-2/15}, c_4 n^{-2/15}]$, respectively, where c_1, \dots, c_4 are constants set as default in the package. Define s_{m_i,τ_j}^2 as the sample variance of the bootstrap statistics, say $\tilde{T}_{n,(1)}, \dots, \tilde{T}_{n,(100)}$ calculated from 100 bootstrap runs with parameters m_i and τ_j . The formula of the bootstrap statistics is determined by the tests. For example, in the structural stability test, we use

$$\tilde{T}_{n,(r)} = \max_{m \le i \le n-m+1} |\Psi_{i,m}^{(r)} - \hat{\Lambda}(i/n)\hat{\Lambda}^{-1}(1)\Psi_{n-m+1,m}^{(r)}|,$$

where $\Psi_{i,m}^{(r)} = n^{-1/2} \sum_{j=1}^{i} \hat{\Sigma}^{1/2}(t_j) R_j^{(r)}$, $\hat{\Lambda}(i/n) = \sum_{j=1}^{i} x_{j,n} x_{j,n}^{\top}/n$, $(R_j^{(r)})_{j=1}^n$ are the independent and identically distributed standard normal random variables independent of data and are independently generated in the *r*th bootstrap iteration, and $\hat{\Sigma}(t)$ is an estimator of $\Sigma(t)$. For testing long memory, we can use $\tilde{K}_n^{(r)}$, $\tilde{RS}_n^{(r)}$, $\tilde{VS}_n^{(r)}$ and $\tilde{KS}_n^{(r)}$ in Section 5.2 for $\tilde{T}_{n,(r)}$ to choose smoothing parameters for different tests. Then we calculate

$$MV(i,j) := SE\left\{ \bigcup_{r_1=-1}^{1} \{s_{m_i,\tau_{j+r_1}}^2\} \cup \bigcup_{r_2=-1}^{1} \{s_{m_{i+r_2},\tau_j}^2\} \right\},$$
(B.1)

where SE stands for standard error. Finally, we select the pair (m_{i^*}, τ_{j^*}) where (i^*, j^*) minimizes MV(i, j). The extended minimum volatility selection criterion (B.1) is similar in spirit to the classical one except that (B.1) is built on the bootstrap test statistics instead of using solely long-run covariance estimators. Therefore, the extended minimum volatility selection criterion (B.1) is adaptive to various types of hypothesis testing problems. In our simulation studies, we recommend $c_1 = 3/7$, $c_2 = 11/7$, M_1 is the number of the points and τ_n from $(2/3)n^{-2/15}$ to $n^{-2/15}$ with grid 0.05, i.e., $c_3 = 2/3$, $c_4 = 1$, and $M_2 = \lfloor n^{-2/15}/0.15 \rfloor$ is the number of grid points of τ_n . The choices of the constants c_1, \ldots, c_4 and M_1 and M_2 are constants replying on the dependence and smoothness of the time series. In practical, one can also choose the constants according to the prior knowledge. The selection procedures of the tuning parameters considered in our paper are implemented in our package, while the package also supports user-specific choices of m and τ_n . In practice, we recommend choosing m from

$$\max(\lfloor (3/7)n^{4/15} \rfloor - 1, 1), \max(\lfloor (3/7)n^{4/15} \rfloor - 1, 1) + 1, \dots, \\\max(\lfloor (11/7)n^{4/15} \rfloor + 1, \max(\lfloor (3/7)n^{4/15} \rfloor - 1, 1) + 2)),$$

to make sure there are enough neighborhood points for extended minimum volatility selection at the rate between $\lfloor (3/7)n^{4/15} \rfloor$ and $\lfloor (11/7)n^{4/15} \rfloor$. The terms max, +1, -1, and +2 make the grid appropriate when the sample size is small.

The full algorithm including data-driven choices of m and τ_n is as follows

- 1. First propose a grid of possible block sizes and bandwidths $\{m_1, m_2, \cdots, m_{M_1}\}, \{\tau_1, \tau_2, \cdots, \tau_{M_2}\}$ for m and τ , say the grid for m is $\max(\lfloor (3/7)n^{4/15} \rfloor 1, 1), \max(\lfloor (3/7)n^{4/15} \rfloor 1, 1) + 1, \ldots, \max(\lfloor (11/7)n^{4/15} \rfloor + 1, \max(\lfloor (3/7)n^{4/15} \rfloor 1, 1) + 2))$ and the grid for τ_n is $(2/3)n^{-2/15}, (2/3)n^{-2/15} + 0.05, \ldots, n^{-2/15}$.
- 2. Compute s_{m_i,τ_j}^2 , the sample variance of the bootstrap statistics, say $\tilde{T}_{n,(1)}, \ldots, \tilde{T}_{n,(100)}$ calculated from 100 bootstrap runs with parameters m_i and τ_j . For example, in the structural stability test, we use

$$\tilde{T}_{n,(r)} = \max_{m \le i \le n-m+1} |\Psi_{i,m}^{(r)} - \hat{\Lambda}(i/n)\hat{\Lambda}^{-1}(1)\Psi_{n-m+1,m}^{(r)}|$$

where $\Psi_{i,m}^{(r)} = n^{-1/2} \sum_{j=1}^{i} \hat{\Sigma}^{1/2}(t_j) R_j^{(r)}$, $\hat{\Lambda}(i/n) = \sum_{j=1}^{i} x_{j,n} x_{j,n}^{\top} / n$, $(R_j^{(r)})_{j=1}^n$ are the independent and identically distributed standard normal random variables independent of data and are independently generated in the *r*th bootstrap iteration, and $\hat{\Sigma}(t)$ is an estimator of $\Sigma(t)$.

3. Calculate

$$MV(i,j) := SE\left\{\bigcup_{r_1=-1}^{1} \{s_{m_i,\tau_{j+r_1}}^2\} \cup \bigcup_{r_2=-1}^{1} \{s_{m_{i+r_2},\tau_j}^2\}\right\},\$$

where SE stands for standard error.

- 4. Select the pair (m_{i^*}, τ_{j^*}) where (i^*, j^*) minimizes MV(i, j)
- 5. For $t \in [m/n, 1 m/n]$, compute the estimator using m_{i^*} for m, and τ_{j^*} for τ_n ,

$$\hat{\Sigma}(t) = \hat{\Sigma}(t) - \breve{\Sigma}(t), \quad \breve{\Sigma}(t) = \sum_{j=m}^{n-m} \frac{m\hat{A}_{j,m}\hat{A}_{j,m}^{\top}}{2}\omega(t,j),$$

where

 $\tilde{\omega}(t,i)$

$$\hat{A}_{j,m} = \frac{1}{m} \sum_{i=j-m+1}^{j} \{ x_{i,n} x_{i,n}^{\top} \breve{\beta}(t_i) - x_{i+m,n} x_{i+m,n}^{\top} \breve{\beta}(t_{i+m}) \}, \ \breve{\beta}(t) = \Omega^{-1}(t) \varpi(t),$$
$$\Omega(t) = \sum_{j=m}^{n-m} \Delta_j \omega(t,j)/2, \ \varpi(t) = \sum_{j=m}^{n-m} \Delta_j \widetilde{\omega}(t,j)/2,$$
$$(t_i - t)/\tau_n^{3/2}) / [\sum_{i=1}^n K\{(t_i - t)/\tau_n^{3/2}\}], \ \omega(t,i) = K\{(t_i - t)/\tau_n\}/[\sum_{i=1}^n K\{(t_i - t)/\tau_n\}].$$

6. For $t \in [0, m/n)$, set $\hat{\Sigma}(t) = \hat{\Sigma}(m/n)$. For $t \in (1 - m/n, 1]$, set $\hat{\Sigma}(t) = \hat{\Sigma}(1 - m/n)$.



Figure B.1: A boxplot of the selected smoothing parameter m of V/S, R/S, KPSS, K/S tests and the test of structural stability.

Using the default choices of the tuning parameters, the summary of selected m's and τ_n 's in the time series regression setting with d = 0 for the four types of long memory tests and the regression model with no change points for the structural ability tests are displayed in Table B.1 and Figure B.1, which partly demonstrates that our proposed estimator is not sensitive to the choices of the smoothing parameters. From the results, we find that our tuning parameter selection approach recommends different but similar tuning parameters for those tests, especially for the long memory tests V/S, R/S, KPSS, and K/S tests which share the same statistical model and the same null hypothesis, implying the stability of our selection procedure. Together with the simulation studies on the simulated rejection rates, the results indicate that our selection procedure works reasonably well.

	m		$ au_n$		
	Median	Max	Mean	Max	
VS	8	10	0.328	0.352	
\mathbf{RS}	8	10	0.329	0.352	
KPSS	7	10	0.327	0.352	
\mathbf{KS}	7	10	0.328	0.352	
CP	7	9	0.399	0.424	

Table B.1: Selected values of m and τ_n in the long-memory tests (V/S, R/S, KPSS, K/S) and the test for structural stability (CP).

B.2 The use of kernels

We investigate a group of common kernels that satisfy Assumption 6, including quartic $(15/16(1-u^2)^2)$, triweight $(35/32(1-u^2)^3)$ with bounded support $|u| \leq 1$, and other kernels which are differentiable but are not continuously differentiable at some points in (-1, 1) are continuously differentiable almost everywhere in (-1, 1) except a few points. See Figure B.2 for our simulation results which check the performance of change points detection using our proposed long-run covariance matrix estimator with different kernels. The simulation result shows that the performance is reasonably well using different kernels satisfying Assumption 6 and some kernels partially fullfill Assumption 6.



Figure B.2: Empirical rejection rates of the structural stability test with triangular(solid), Epanechnikov (dotted), quartic(small dashes), triweight (dashes) and tricube (long dashes) kernels, respectively.

In the R package mlrv, we also offer the options of employing different kernels in the estimation of long-run covariance matrix, including triangular kernel, Epanechnikov kernel, quartic kernel, triweight kernel and tricube kernel.

B.3 Positive definite estimators

In practice when the estimated covariance matrix, which proves to be consistent, is not positive definite, there are two possibilities: collinearity or the small sample size. The former implies that the estimand, namely the covariance matrix, is not positive definite, which is excluded by Assumption 2. In the latter case, the covariance matrix is positive definite, but the estimated covariance matrix can be non-positive definite due to the stochastic variation at a small sample size.

Therefore, the remedy of the non-positive definite estimator when the sample size is small is of primary and practical concern. As discussed in the literature, one can use threshold method ((Politis, 2011)) or penalization ((Rothman, 2012)) for the remedy. We only discuss the threshold approach employed by (Politis, 2011), (Dette and Wu, 2020) among others due to the page limit, and leave further exploration of the modification as a rewarding future work.

We follow the threshold procedure as discussed in (Politis, 2011). In particular, let s_n denote the stochastic upper bound under possible smooth and abrupt changes or long memory in Theorem 1, Theorem 2 and Theorem 3, respectively. We choose the threshold level for the eigenvalues to be 1/n, so that $\rho(\hat{\Sigma}^{\rm pd}(t) - \hat{\Sigma}(t)) = o(s_n)$, where $\hat{\Sigma}^{\rm pd}(t)$ denotes the estimator after using threshold for eigenvalues, i.e., $\hat{\Sigma}^{\rm pd}(t) = U(t)\Lambda^{\rm pd}(t)U(t)^{\top}$, $\Lambda^{\rm pd}(t)$ is the diagonal matrix with diagonal elements $\lambda_i^+(t) = \max(1/n, \lambda_i(t))$ ($i = 1, \ldots, p$), where $\lambda_i(t)$ is the ordered eigenvalue of $\hat{\Sigma}(t)$, and U(t) is the matrix consisting of the corresponding eigenvectors of $\hat{\Sigma}(t)$.

C Discussion on assumptions

C.1 high-level assumptions

It is possible to formulate high-level assumptions for the pilot estimator so that the debias effect can be achieved at least theoretically. In the *absence* of jump points, the high-level assumption of the pilot estimator that will lead to the consistency of the difference-based long-run covariance matrix estimator is that for the event $G_n = \{\breve{\beta}_0(t) \text{ is well-defined for } t \in I\}$ and $P(G_n) \to 1$,

$$\sup_{t \in \mathcal{I}} \|\{\beta(t) - \breve{\beta}_0(t)\} \mathbb{1}(G_n)\|_{4\kappa} = O\{\tau_n^3 + (n\tau_n^{3/2})^{-1/2}\}.$$

Let $C_m = \{j : j - m \leq i \leq j + m, \beta'''(t_i) \text{ exists and continuous}\}, C_n = \{j : j - n\tau_n^{3/2} \leq i \leq j + n\tau_n^{3/2}, \beta'''(t_i) \text{ exists and continuous}\}$ and the number of q_n jump points should satisfy the condition of Theorem 2. In the presence of jump points, the high-level assumption will be

$$\sup_{j\in\mathcal{C}_m} \|\{\check{\beta}_0(t_{j+m}) - \check{\beta}_0(t_j)\}1(G_n)\|_{4\kappa} = O\left\{\frac{mq_n}{n\tau_n^{3/2}} + (n\tau_n^{3/2})^{-1/2}\right\},\$$

and

$$\sup_{j \in \mathcal{C}_n} \|\{\check{\beta}_0(t_j) - \beta_0(t_j)\} \mathbb{1}(G_n)\|_{4\kappa} = O\left\{\tau_n^3 + (n\tau_n^{3/2})^{-1/2}\right\}.$$

For the case of long memory, from Lemma C.1 and Lemma C.2, we can obtain the high-level assumption is

$$\sup_{t \in \mathcal{I}} \|\{\beta(t) - \breve{\beta}_0(t)\} \mathbb{1}(G_n)\|_{4\kappa} = O\{\tau_n^3 + (n\tau_n^{3/2})^{d-1/2}\}.$$

Therefore, it is possible to find other estimators that satisfy the high-level assumptions to achieve a similar debias effect. However, in non-parametric estimation and inference, it is more convenient in practice to use tuning parameters as few as possible. We recommend the statistic $\check{\beta}(\cdot)$ in the paper, mainly because it can satisfy all the high-level assumptions without introducing extra smoothing parameters.

C.2 Discussion on Assumption 5

Recall our definition of short-range dependence in the paper:

Definition C.1. The process $G(t, \mathcal{F}_i)$ is of r-order short-range dependence on interval I if $\sup_{t \in I} ||G(t, \mathcal{F}_0)||_r < \infty$, $\delta_r(H, k, I) = O(\chi^k)$, for some $\chi \in (0, 1)$, $r \ge 1$, and s-order locally stationary on interval I, $s \ge 2$, if $G(t, \mathcal{F}_0) \in \operatorname{Lip}_s(I)$.

Therefore, there are two restrictions of Assumption 5, which are the moment constraint and the

constraints in order r of the physical dependence measure

$$\delta_{r}(L,k,I) = \sup_{t \in I} \left\| L\left(t,\mathcal{F}_{k}\right) - L\left(t,\mathcal{F}_{k,\{0\}}\right) \right\|_{r}.$$

Under the conditions of short-range dependence and no jump points, we allow $\kappa = 1$, i.e., 16-order moment is required mainly because of the corresponding non-parametric smoothing of time series as well as the use of physical dependence. In general, the requirement of 16 is hard to be reduced because we estimate $\check{\beta}(\cdot)$ using second-order series and formulation of $\check{\Sigma}(\cdot)$ involving the square of $x_i x_i^{\top} \check{\beta}(t_i)$. Recall that

$$\check{\Sigma}(t) = \sum_{j=m}^{n-m} \frac{m\hat{A}_{j,m}\hat{A}_{j,m}^{\mathrm{T}}}{2} \omega(t,j), \quad \hat{A}_{j,m} = \frac{1}{m} \sum_{i=j-m+1}^{j} \left\{ x_{i,n} x_{i,n}^{\mathrm{T}} \breve{\beta}\left(t_{i}\right) - x_{i+m,n} x_{i+m,n}^{\mathrm{T}} \breve{\beta}\left(t_{i+m}\right) \right\},$$

and

$$\breve{\beta}(t) = \Omega^{-1}(t) \varpi(t) \quad (t \in [0, 1]),$$

where $\Omega(t)$ and $\varpi(t)$ are the smoothed versions of

$$\dot{\Delta}_j/2 = \frac{1}{2m} \sum_{i=j-m+1}^{j} \tilde{X}_{i,m} \tilde{X}_{i,m}^{\top} \text{ and } \breve{\Delta}_j/2 = \frac{1}{2m} \sum_{i=j-m+1}^{j} \tilde{X}_{i,m}^{\top} \tilde{Y}_{i,m},$$

where $\tilde{Y}_{i,m} = x_{i,n}y_{i,n} - x_{i+m,n}y_{i+m,n}$, $\tilde{X}_{i,m} = x_{i,n}x_{i,n}^{\top} - x_{i+m,n}x_{i+m,n}^{\top}$. Therefore, the moment condition of order 16 is difficult to relax since we require $\check{\Sigma}(\cdot)$ to have a finite second moment.

Although Assumption 5 is seemingly strong for technical convenience, empirical studies show that our proposed estimator could still be consistent allowing $H(\cdot, \mathcal{F}_i)$ to have a heavier tail than Assumption 5 under considered scenarios, though with possibly slower convergence rate. In Figure C.1, we display the empirical rejection rates of different types of innovations, i.e., normal, t(5) and t(6) for ζ_i in

$$u_{i,n} = G(t_i, \mathcal{F}_i) = 0.65 \cos(2\pi t_i) G(t_i, \mathcal{F}_{i-1}) + \zeta_i,$$

in the model in Section 5.2 of the paper for the tests for structural stability:

$$y_{i,n} = 1 + m_{i,n} + x_{i,n,1} + x_{i,n,2} + e_{i,n}, \quad e_{i,n} = (1 + 0.1x_{i,n,1})u_{i,n}$$

The results show that the tests for structural stability and long memory can still work reasonably well under some scenarios for example the CP1 case when we relax the moment conditions using t(5) or t(6).

In Figure C.2, the empirical rejection rates of different types of innovations, i.e, normal, t(4) and t(6) for ε_i in

$$B(t, \mathcal{G}_i) = \{0.3 - 0.4(t - 0.5)^2\}B(t, \mathcal{G}_{i-1}) + 0.8\varepsilon_i\}$$



Figure C.1: Empirical rejection rates of tests for structural stability with respect to different innovations: normal(solid), t(5)(small dashes) and t(6)(dashes) with sample size n = 300.



Figure C.2: Empirical rejection rates of tests for long memory with respect to difference innovations: normal (solid), t(4) (dashes) and t(6) (dotted) with sample size n = 1500.

in the model in Section 5.3 of the paper for the tests for long memory, i.e.,

$$y_{i,n} = \beta_1(t_i) + \beta_2(t_i)x_{i,n} + (1 - \mathcal{B})^{-d}e_{i,n}, \quad e_{i,n} = H(t_i, \mathcal{F}_i, \mathcal{G}_i) = B(t_i, \mathcal{G}_i) \{1 + W^2(t_i, \mathcal{F}_i)\}^{1/2}$$

where $W(t, \mathcal{F}_i) = \{0.1 + 0.1 \cos(2\pi t)\}W(t, \mathcal{F}_{i-1}) + 0.2\zeta_i + 0.7(t-0.5)^2$ for $i = 1 \cdots, n$. The bandwidth selection procedure is identical to the one used in the paper. We can find the influences of moment conditions in both the structural stability test and the long memory detection, i.e., the empirical power of the former is reduced when the tails are heavier, while the empirical sizes of the latter increase with the heavier tails.

Our current setting allows polynomial tailed distribution. High-order physical dependence measure condition can be omitted when sub-exponential moment condition is assumed, where we only need order-1 physical dependence measure, see the following lemma.

Lemma C.1. Suppose $\exists t_0 > 0$, $C = \sup_{t \in [0,1]} E\{\exp(t_0|H(t,\mathcal{F}_0)|)\} < \infty$, $\delta_1(H,l,[0,1]) = O(\chi^l)$ for some

 $\chi \in (0,1)$. Then, we have the following holds for q > 2: (i) $\sup_{t \in [0,1]} E(|H(t,\mathcal{F}_0)|^q) \leq Ct_0^{-q}q^q$. (ii) There exists a positive constant C' such that $\delta_q(H,l,[0,1]) = \sup_{t \in [0,1]} ||H(t,\mathcal{F}_l) - H(t,\mathcal{F}_{l,\{0\}})||_q \leq C'q\chi^{l/q}$.

Proof. For $p \ge P$, P > 0, elementary calculation gives that $x > p \log(x/p)$, x > 0. For a sufficiently large constant C, we have

$$\sup_{t \in [0,1]} E(|H(t,\mathcal{F}_0)|^q) < \left(\frac{q}{t_0}\right)^q \sup_{t \in [0,1]} E\{\exp(t_0|H(t,\mathcal{F}_0)|)\} = t_0^{-q} q^q C$$

By Hölder inequality,

$$\sup_{t \in [0,1]} \|H(t,F_l) - H(t,F_{l,\{0\}})\|_q^2 = [E\{|H(t,F_l) - H(t,F_{l,\{0\}})|^{q-1/2}|H(t,F_l) - H(t,F_{l,\{0\}})|^{1/2}\}]^2$$

$$\leq E\{|H(t,F_l) - H(t,F_{l,\{0\}})|^{2q-1}\}E\{|H(t,F_l) - H(t,F_{l,\{0\}})|\}$$

$$\leq \|H(t,F_l) - H(t,F_{l,\{0\}})\|_{2q-1}^{2q-1}\delta_1(H,l,[0,1])$$

$$\leq C_1(2q-1)^{2q-1}t_0^{-2q+1}\chi^l,$$

where C_1 is a sufficiently large constant. Therefore, we have $\delta_q(H, l, [0, 1]) \leq C' \chi^{l/q} q$.

D Other potential applications

Under local stationarity, our proposed estimator can be used in many practical scenarios, such as constructing simultaneous confidence bands for time-varying regression coefficients, deriving preliminary estimation and visualization of the long memory parameter d for locally stationary long memory process, as well as many other inference problems that involve the estimation of the long-run covariance matrix, such as testing for white noises, generalized likelihood ratio test and squared integrated tests for time-varying regression coefficient functions, see (Zhou, 2014b) for instance. In the following we list several detailed examples.

Visualization of long memory. The long-run covariance estimator can serve as a simple and heuristic tool for visualizing and assessing the presence of long memory, see Section 1.2 of (Beran et al., 2013). By Theorem 3, under the fixed alternative, we have

$$\sup_{t\in I} \left| m^{-2d} \hat{\Sigma}(t) - \kappa_2(d) \sigma_H^2(t) \mu_W(t) \mu_W^\top(t) \right| = o_p(1).$$

Suppose we have a grid of m's, i.e., m_1, \ldots, m_M , and the corresponding long-run covariance estimator $\hat{\Sigma}_1(\cdot), \ldots, \hat{\Sigma}_M(\cdot)$ calculated using m_1, \ldots, m_M , respectively. Taking $x_i = \log m_i, y_i = \sum_{j=1}^n \log |\hat{\Sigma}_i(t_j)|/n$, $(i = 1, \ldots, M)$, where $|\cdot|$ denotes the Frobenious norm, and we have

$$y_i \approx (2d)x_i + \log \kappa_2(d) + \sum_{j=1}^n \log \sigma_H^2(t_j)/n + \sum_{j=1}^n \log |\mu_W(t_j)\mu_W^\top(t_j)|/n$$

= $(2d)x_i + f(d),$ (D.1)

where the quantity f(d) is independent of *i*. Therefore, one can visualize *d* by drawing a regression line $y_i \sim x_i$. For illustration, we generate a data set from the functional linear model in Section 6.3 of the main paper with d = 0.2, i.e.,

$$y_{i,n} = \beta_1(t_i) + \beta_2(t_i)x_{i,n} + (1 - \mathcal{B})^{-d}e_{i,n},$$

where \mathcal{B} is the lag operator, $\beta_1(t) = 8\sin(\pi t)$, $\beta_2(t) = 4\exp\{-2(t-0.5)^2\}$, $x_{i,n} = W(t_i, \mathcal{F}_i)$ (i = 1, ..., n), and $e_{j,n} = H(t_j, \mathcal{F}_j, \mathcal{G}_j)$ (j = 1, ..., n) with

$$H(t, \mathcal{F}_i, \mathcal{G}_i) = B(t, \mathcal{G}_i) \{ 1 + W^2(t, \mathcal{F}_i) \}^{1/2} \quad (i \in Z; \ t \in [0, 1]),$$

where $W(t, \mathcal{F}_i) = 0.1 \cos(2\pi t) W(t, \mathcal{F}_{i-1}) + 0.2\zeta_i + 0.4(t-0.5)^2$, and $B(t, \mathcal{G}_i) = \{0.3 - 0.4(t-0.5)^2\} B(t, \mathcal{G}_{i-1}) + 0.6\varepsilon_i$.

Figure D.1 displays an instance for the visualization. The y-axis is the average logarithm of Frobenius norm of the long-run covariance estimator, while the x-axis is the logarithm of the parameter m. The displayed fitted regression line of (D.1) is y = -0.82 + 0.41x, and the estimated d is close to the half of the slope, i.e. 0.205. Furthermore, 100 times of simulations yield the average estimated d being 0.204(0.006).



Figure D.1: Regression using data from the long-run covariance matrix estimator. The estimated d is half of the slope.

Simultaneous confidence tubes. We consider a similar functional linear model as in Section 6.3 of the main article with d = 0, namely

$$y_{i,n} = \beta_1(t_i) + \beta_2(t_i)x_{i,n} + e_{i,n} \quad (i = 1, \dots, n),$$

where $\beta_1(t) = 4\sin(\pi t), \beta_2(t) = 4\exp\{-2(t-0.5)^2\}, x_{i,n} = W(t_i, \mathcal{F}_i)$ with $W(t, \mathcal{F}_i) = \{0.25 + 0.25\cos(2\pi t)\}W(t, \mathcal{F}_{i-1}, 0.2\zeta_i)$, and $e_{i,n} = H(t_i, \mathcal{F}_i, \mathcal{G}_i)$, where $t_i = i/n$ and

$$H(t, \mathcal{F}_i, \mathcal{G}_i) = B(t, \mathcal{G}_i) \left\{ 1 + W^2(t, \mathcal{F}_i) \right\}^{1/2},$$

where $B(t, \mathcal{G}_i) = \{0.2 - 0.4(t - 0.5)^2\}B(t, \mathcal{G}_{i-1}) + 0.8\varepsilon_i$. (Zhou and Wu, 2010) considers the simultaneous confidence tubes for regression coefficient functions using plug-in estimators, see estimator (17) of their paper, which is also available in our R package. We compare the performance of simultaneous confidence tubes jointly for $\beta_1(t)$ and $\beta_2(t)$, using our estimator and the plug-in estimator advocated by (Zhou and Wu, 2010) in Table D.1, which shows the advantage of our estimator. Notice that $H(t, \mathcal{F}_i, \mathcal{G}_i)$ depends on the covariates in a nonlinear way, which has not been investigated empirically by (Zhou and Wu, 2010).

	Oı	ırs	Plug-in		
b/nominal	95%	90%	95%	90%	
0.2500	89.5	83.8	87.0	78.5	
0.2750	89.9	84.8	86.8	77.2	
0.2875	90.9	85.1	85.7	78.5	
0.3000	92.1	86.3	87.1	77.8	
0.3125	92.8	87.3	86.5	78.2	
0.3250	92.2	86.5	87.4	79.8	
0.3375	93.3	87.6	88.0	78.8	
0.3500	91.2	85.3	86.6	78.1	

Table D.1: Empirical coverage rates (in %) via 1000 times of simulations for the simultaneous confidence tubes of $(\beta_1(t), \beta_2(t))$ with sample size 500 using our estimator and the plug-in estimator advocated by (Zhou and Wu, 2010).

E Extra simulation

E.1 Sensitivity analysis

For the sensitivity check, we examine the performance of our method with different choices of m's and τ_n 's under both null and various alternative hypotheses and compare it to that of baseline methods using nonparametric or ordinary least square residuals for estimating the long-run covariance matrix.

- Figure E.1 demonstrates the power performance of the tests equipped with the debiased differencebased estimator compared with the baseline method based on ordinary least square residuals $\hat{e}_{i,n} = y_{i,n} - x_{i,n}^{\top} \hat{\beta}_n$ in the CP1 model in the main paper, where we compare fixed *m*'s (10 and 20 with $\tau_n = n^{-2/15}$) and fixed τ_n 's (0.2, 0.3, 0.4 with m = 10) as well as *m* selected via extended minimum volatility over different ranges. In the paper, we choose *m* from 6 ($\lfloor 10/7n^{4/15} \rfloor$) to 9 ($\lfloor 15/7n^{4/15} \rfloor$). In Figure E.1, we show the results when choosing *m* from 6 ($\lfloor 10/7n^{4/15} \rfloor$) to 30 ($\lfloor 50/7n^{4/15} \rfloor$) and from 10 ($\lfloor 18/7n^{4/15} \rfloor$) to 50 ($\lfloor 75/7n^{4/15} \rfloor$). Using our difference-based debiased long-run covariance matrix estimator, the test outperforms that based on ordinary least square residuals by a large margin under different choices of tuning parameters.
- Similar improvement can also be found in Figure E.2 in the CP4 model.
- In Figure E.3, the roles of tuning parameters in the detection of long memory are investigated. The bootstrap tests equipped with the debiased difference-based estimator under different smoothing



Figure E.1: Comparing structural stability tests of using plug-in estimators of ordinary least square residuals (ols, dotted), the proposed difference-based estimator using extended minimum volatility selection procedure (MV, dashes), choosing m from 6 ($\lfloor 10/7n^{4/15} \rfloor$) to 9 ($\lfloor 15/7n^{4/15} \rfloor$) versus choosing m from 6 ($\lfloor 10/7n^{4/15} \rfloor$) to 30 ($\lfloor 50/7n^{4/15} \rfloor$), and 10 ($\lfloor 18/7n^{4/15} \rfloor$) to 50 ($\lfloor 75/7n^{4/15} \rfloor$), as well as the proposed difference-based estimator with several fixed m's with $\tau_n = n^{-2/15}$ and τ_n 's with m = 10 (fixed, solid) when there is one change point (CP1).

parameters are compared with the baseline method based on plugging in nonparametric residuals in (Zhou and Wu, 2010) as in the main paper. Long-memory tests with the proposed difference-based estimator achieve much better trade-offs in type-I and type-II errors under various choices of tuning parameters.

E.2 Simulated rejected rates with sample size 750

The following Table E.1 reports the simulated sizes of KPSS, R/S, V/S and K/S-type tests when n = 750.

	KI	\mathbf{PSS}	\mathbf{R}_{i}	$/\mathrm{S}$	V	$/\mathrm{S}$	K	$/\mathrm{S}$
	5%	10%	5%	10%	5%	10%	5%	10%
plug-in	8.40	15.00	21.00	32.80	52.00	67.10	13.30	22.30
diff	4.00	7.50	6.10	11.50	10.00	15.50	3.30	7.30

Table E.1: Simulated rejection rates of KPSS, R/S, V/S and K/S-type tests for long memory when d = 0, n = 750.

E.3 Long-run covariance estimates with change points

To further illustrate the role of long-run variance estimators in bootstrap tests, we conduct 1000 times simulation, where we find the MSE of long-run variance estimators using ordinary least square residuals is 974.12, more than twice the magnitude of the MSE of that of difference-based method 275.94 when there are four change points in the time series structure (scenario CP4 with $\delta = 1$). Moreover, two-sample t test also shows that under the alternative with $\delta = 1$, the difference-based method yields long-run variance estimate with much smaller MSE than the ordinary least square method with *p*-value smaller than 0.01%.



Figure E.2: Comparing structural stability tests of using plug-in estimators of ordinary least square residuals (ols, dotted), the proposed difference-based estimator using extended minimum volatility selection procedure (MV, dashes), choosing m from 6 ($\lfloor 10/7n^{4/15} \rfloor$) to 9 ($\lfloor 15/7n^{4/15} \rfloor$) versus choosing m from 6 ($\lfloor 10/7n^{4/15} \rfloor$) to 30 ($\lfloor 50/7n^{4/15} \rfloor$), and 10 ($\lfloor 18/7n^{4/15} \rfloor$) to 50 ($\lfloor 75/7n^{4/15} \rfloor$) as well as the proposed difference-based estimator with several fixed m's with $\tau_n = n^{-2/15}$ and τ_n 's with m = 10 (fixed, solid) when there are 4 change points (CP4).



Figure E.3: Comparing V/S-type tests of using plug-in estimators of nonparametric residuals (plug-in, dashes), the proposed difference-based estimator choosing m from 3 to 11 versus from 6 to 30, and from 7 to 16 (MV, dotted) as well as using several fixed m's with $\tau_n = n^{-2/15}$ and τ_n 's with m = 10 (fixed, solid).

F Proof

F.1 Notation

We first introduce some notation that will be frequently used in the mathematical argument of this section. In the following proofs, we will omit the index n in $e_{i,n}, x_{i,n}, y_{i,n}$ for simplicity. Define filtration $\mathcal{F}_i = (\varepsilon_{-\infty}, ..., \varepsilon_i)$ for independent and identically distributed random variables $(\varepsilon_i)_{i \in \mathbb{Z}}$. For a random vector $(v_i)_{i=1}^n \in \mathcal{F}_s$, let $v_{i,\{s\}}$ denote the series replacing the ε_s with its independent and identically distributed copy. For a random matrix $(A_i)_{i=1}^n \in \mathcal{F}_s$, define $A_{j,\{s\}}$ as the random matrix replacing ε_s in A_j with its independent and identically distributed copy. Recall that $e_{i,n}^{(d)} = \sum_{j=0}^{\infty} \psi_j(d)e_{i-j,n}$, $e_{i,n}^{(d_n)} = \sum_{j=0}^{\infty} \psi_j(d_n)e_{i-j,n}$. For the sake of simplicity, we use ψ_j to represent $\psi_j(d)$ when we discuss the fixed alternatives and $\psi_j(d_n)$ for the theory of the local alternatives. Recall $t_i = i/n$, and $K^*(x)$ denotes the jackknife equivalent kernel $2\sqrt{2K}(\sqrt{2x}) - K(x)$. Let $0 \times \infty = 0$, $a_n \sim b_n$ denote $\lim_{n\to\infty} a_n/b_n = 1$ for real sequences a_n and b_n . Let $I = [\gamma_n, 1 - \gamma_n] \subset (0, 1)$, $\gamma_n = \tau_n + (m+1)/n$. Recall

$$A_{j,m} = \frac{1}{m} \sum_{i=j-m+1}^{j} \{ x_i x_i^{\top} \beta(t_i) - x_{i+m} x_{i+m}^{\top} \beta(t_{i+m}) \}, \quad \Sigma^A(t) = \sum_{j=m}^{n-m} \frac{m A_{j,m} A_{j,m}^{\top}}{2} \omega(t,j),$$

$$\hat{A}_{j,m} = \frac{1}{m} \sum_{i=j-m+1}^{j} \{ x_i x_i^\top \breve{\beta}(t_i) - x_{i+m} x_{i+m}^\top \breve{\beta}(t_{i+m}) \}, \quad \breve{\Sigma}(t) = \sum_{j=m}^{n-m} \frac{m \hat{A}_{j,m} \hat{A}_{j,m}^\top}{2} \omega(t,j),$$

where $\omega(t, i) = K_{\tau_n}(t_i - t) / \sum_{i=1}^n K_{\tau_n}(t_i - t)$. Let

$$\dot{A}_{j,m} = \frac{1}{m} \sum_{i=j-m+1}^{j} (x_i x_i^{\top} - x_{i+m} x_{i+m}^{\top}) \{ x_i x_i^{\top} \beta(t_i) - x_{i+m} x_{i+m}^{\top} \beta(t_{i+m}) \},$$
(F.1)

and

$$\dot{\Delta}_j = \frac{1}{m} \sum_{i=j-m+1}^j (x_i x_i^\top - x_{i+m} x_{i+m}^\top) (x_i e_i - x_{i+m} e_{i+m}).$$
(F.2)

We define below the counterparts of $Q_{k,m}$, Δ_j and $\hat{\Sigma}(\cdot)$ in Section 3. Define for $m \ge 2$, $t \in [m/n, 1 - m/n]$,

$$\tilde{Q}_{k,m} = \sum_{i=k}^{k+m-1} x_i e_i, \quad \tilde{\Delta}_j = \frac{\tilde{Q}_{j-m+1,m} - \tilde{Q}_{j+1,m}}{m}, \quad \tilde{\Sigma}(t) = \sum_{j=m}^{n-m} \frac{m\tilde{\Delta}_j\tilde{\Delta}_j^\top}{2}\omega(t,j).$$
(F.3)

F.2 Proof of Theorem 4.1

The following lemma presents the consistency of bias correction for the difference-based estimator with time series covariates, which is crucial to establish the consistency result of Theorem 4.1.

Lemma F.1. Under the condition of Theorem 4.1, we have

$$\sup_{t \in \mathcal{I}} \left| \breve{\Sigma}(t) - \Sigma^{A}(t) \right| = O_{\mathbb{P}} \left(\sqrt{m\tau_{n}^{3-1/\kappa}} + \sqrt{\frac{m}{n\tau_{n}^{3/2+2\kappa}}} \right)$$

Proof of Lemma F.1. Let l_n be a sequence of real numbers so that $l_n \to \infty$ arbitrarily slow. Define $A_n = \{\sup_{t \in \mathcal{I}} |\Omega(t) - M^+(t)| \le l_n \{(m\tau_n^{3/2})^{-1/2} + m/(n\tau_n^{3/4}) + \tau_n^{3/4}\}\}$. By (F.9), $\lim_{n\to\infty} \mathbb{P}(A_n) = 1$. Since $0 \le \sum_{j=m}^{n-m} \omega(t,j) \le 1$, we have

$$\begin{split} \sup_{t\in\mathcal{I}} \left\| (\check{\Sigma}(t) - \Sigma^{A}(t)) \mathbf{1}(A_{n}) \right\|_{\kappa} \\ &\leq \sup_{t\in\mathcal{I}} \sum_{j=m}^{n-m} \frac{m\omega(t,j)}{2} \left\| (\hat{A}_{j,m} - A_{j,m}) \mathbf{1}(A_{n}) \right\|_{2\kappa} \left(\left\| (\hat{A}_{j,m} - A_{j,m}) \mathbf{1}(A_{n}) \right\|_{2\kappa} + 2 \|A_{j,m}\|_{2\kappa} \right) \\ &\leq m \max_{m\leq j\leq n-m} \left\| (\hat{A}_{j,m} - A_{j,m}) \mathbf{1}(A_{n}) \right\|_{2\kappa} \\ &\times \left(\max_{m\leq j\leq n-m} \left\| (\hat{A}_{j,m} - A_{j,m}) \mathbf{1}(A_{n}) \right\|_{2\kappa} + 2 \max_{m\leq j\leq n-m} \|A_{j,m}\|_{2\kappa} \right). \end{split}$$
(F.4)

First, we shall show that

$$\max_{m \le j \le n-m} \|A_{j,m}\|_{2\kappa} = O(m^{-1/2} + m/n) = O(m^{-1/2}).$$
(F.5)

Define $B_{j,m} = \frac{1}{m} \sum_{i=j-m+1}^{j} x_i x_i^{\top}$. Notice that

$$\|A_{j,m}\|_{2\kappa} \le \sup_{m \le i \le n-m} \|\beta(t_i) - \beta(t_{i+m})\| \|B_{j,m}\|_{2\kappa} + \sup_{t \in [0,1]} |\beta(t)| \|B_{j,m} - B_{j+m,m}\|_{2\kappa} = A_1 + A_2.$$

Since $\beta(t)$ is Lipschitz continuous, and under Assumption 4.3, $\max_{m \leq j \leq n} \|B_{j,m}\|_{2\kappa}$ is bounded, we have

$$A_1 = O(m/n). \tag{F.6}$$

For the calculation of A_2 , notice that

$$||B_{j,m} - B_{j+m,m}||_{2\kappa} \le ||B_{j,m} - \mathbb{E}(B_{j,m})||_{2\kappa} + ||B_{j+m,m} - \mathbb{E}(B_{j+m,m})||_{2\kappa} + ||\mathbb{E}(B_{j,m}) - M(t_j)||_{2\kappa} + ||\mathbb{E}(B_{j+m,m}) - M(t_j)||_{2\kappa}.$$

Similar to Lemma 6 in (Zhou and Wu, 2010), using rectangular kernel with bandwidth m/n, under Assumption 4.3, we have

$$\sup_{m \le j \le n} \|B_{j,m} - \mathbb{E}(B_{j,m})\|_{2\kappa} = O(m^{-1/2}).$$

Since $\mathbb{E}(B_{j,m}) = \frac{1}{m} \sum_{i=j-m+1}^{j} M(t_j)$ and M(t) is Lipschitz continuous, $\|\mathbb{E}(B_{j,m}) - M(t_j)\|_{2\kappa} = O(m/n)$.

Finally, by the boundedness of $\sup_{t \in [0,1]} |\beta(t)|$, we have

$$A_2 = O(m^{-1/2} + m/n).$$
 (F.7)

Therefore, by (F.6) and (F.7), we have shown (F.5). Second, by triangle inequality, we have

$$\max_{m \le j \le n-m} \left\| (\hat{A}_{j,m} - A_{j,m}) \mathbf{1}(A_n) \right\|_{2\kappa} \le 2 \max_{1 \le i \le n} \left\| x_i x_i^\top \right\|_{4\kappa} \| \{ \beta(t_i) - \breve{\beta}(t_i) \} \mathbf{1}(A_n) \|_{4\kappa}.$$

Since under Assumption 4.3, $\max_{1 \le i \le n} \left\| x_i x_i^\top \right\|_{4\kappa} = O(1)$, we shall show that

$$\sup_{t \in \mathcal{I}} \|\{\beta(t) - \breve{\beta}(t)\} \mathbb{1}(A_n)\|_{4\kappa} = O\left\{\tau_n^3 + (n\tau_n^{3/2})^{-1/2}\right\}.$$
 (F.8)

Let $M^+(t) = \mathbb{E}\{\bar{J}(t, \mathcal{F}_0)\bar{J}^\top(t, \mathcal{F}_0)\}$. Following similar arguments in Lemma 6 of (Zhou and Wu, 2010) and Theorem 5.2 of (Dette and Wu, 2019), under Assumptions 4.3 and 4.4, we have

$$\sup_{m \le j \le n} \| \hat{\Delta}_j / 2 - M^+(t_j) \| = O(m^{-1/2} + m/n).$$

Then, it follows that

$$\sup_{t \in \mathcal{I}} \|\Omega(t) - M^+(t)\| = O(m^{-1/2} + m/n + \tau_n^{3/2}).$$

By the chaining argument in Proposition B.1 in Section B.2 in (Dette et al., 2019), we have

$$\sup_{t \in \mathcal{I}} |\Omega(t) - M^+(t)| = O_{\mathbb{P}}\{(m\tau_n^{3/2})^{-1/2} + m/(n\tau_n^{3/4}) + \tau_n^{3/4}\}.$$
 (F.9)

Note that $\Omega(t)$ is invertible on A_n . Then, for a sufficiently large constant C, we have

$$\begin{aligned} \|\{\breve{\beta}(t) - \beta(t)\} \mathbf{1}(A_n)\|_{4\kappa} &= \|\Omega^{-1}(t)\{\varpi(t) - \Omega(t)\beta(t)\} \mathbf{1}(A_n)\|_{4\kappa} \\ &\leq \|\rho(\Omega^{-1}(t))\|\varpi(t) - \Omega(t)\beta(t)\|\mathbf{1}(A_n)\|_{4\kappa} \leq C\|\varpi(t) - \Omega(t)\beta(t)\|_{4\kappa}. \end{aligned}$$

Then, it's sufficient to show that

$$\sup_{t \in \mathcal{I}} \|\varpi(t) - \Omega(t)\beta(t)\|_{4\kappa} = O\left\{\tau_n^3 + (n\tau_n^{3/2})^{-1/2}\right\}$$

Recall the definition of $\dot{\Delta}_j$ and $\dot{A}_{j,m}$ in (F.2) and (F.1) respectively. Observe that

$$\varpi(t) = \sum_{j=m}^{n-m} \frac{\tilde{\omega}(t,j)}{2} \dot{A}_{j,m} + \sum_{j=m}^{n-m} \frac{\tilde{\omega}(t,j)}{2} \dot{\Delta}_j := W_1(t) + W_2(t),$$
(F.10)

where $W_1(t), W_2(t)$ are defined in the obvious way. Recall that $\hat{\Delta}_j = \frac{1}{m} \sum_{i=j-m+1}^j (x_{i,n} x_{i,n}^\top - x_{i+m,n} x_{i+m,n}^\top)^2$.

By triangle inequality, we have

$$\begin{split} \sup_{t\in\mathcal{I}} \|W_1(t) - \Omega(t)\beta(t)\|_{4\kappa} &\leq \sup_{t\in\mathcal{I}} \left\|W_1(t) - \sum_{j=m}^{n-m} \frac{\dot{\Delta}_j \tilde{\omega}(t,j)}{2} \beta(t_j)\right\|_{4\kappa} + \sup_{t\in\mathcal{I}} \left\|\sum_{j=m}^{n-m} \frac{\dot{\Delta}_j \tilde{\omega}(t,j)}{2} \{\beta(t_j) - \beta(t)\}\right\|_{4\kappa} \\ &= W_{11} + W_{12}. \end{split}$$

Again, by triangle inequality, under Assumptions 4.1 and 4.3, we obtain

$$W_{11} \leq \max_{m \leq j \leq n-m} \|\dot{A}_{j,m} - \dot{\Delta}_{j}\beta(t_{j})\|_{4\kappa},$$

$$\leq \frac{1}{m} \max_{m \leq j \leq n-m} \left(\sum_{i=j-m+1}^{j+m} \left\| x_{i}x_{i}^{\top} - x_{i+m}x_{i+m}^{\top} \right\|_{8\kappa} \|x_{i}x_{i}^{\top}\|_{8\kappa} |\beta(t_{i}) - \beta(t_{j})| \right)$$

$$= O(m/n).$$
(F.11)

Under Assumption 4.3, we have $\max_{m \leq j \leq n-m} \| \hat{\Delta}_j \|_{4\kappa} = O(1)$. Then, by similar arguments in Lemma 3 of (Zhou and Wu, 2010) and the continuity of $M^+(t)$, $m\tau_n^{3/2} \to \infty$, we obtain

$$W_{12} = \sup_{t \in \mathcal{I}} \left\| \sum_{j=m}^{n-m} \frac{\dot{\Delta}_{j} \tilde{\omega}(t,j)}{2} \{\beta'(t)(t_{j}-t) + O(\tau_{n}^{3})\} \right\|_{4\kappa}$$

$$\leq \sup_{t \in \mathcal{I}} \left\| \sum_{j=m}^{n-m} \{\dot{\Delta}_{j}/2 - M^{+}(t_{j})\} \tilde{\omega}(t,j)\beta'(t)(t_{j}-t) \right\|_{4\kappa}$$

$$+ \sup_{t \in \mathcal{I}} \left\| \sum_{j=m}^{n-m} M^{+}(t_{j}) \tilde{\omega}(t,j)\beta'(t)(t_{j}-t) \right\|_{4\kappa} + O(\tau_{n}^{3})$$

$$= \sup_{t \in \mathcal{I}} \left\| \sum_{j=m}^{n-m} \tilde{\omega}(t,j)\beta'(t)M^{+}(t)(t_{j}-t) \right\|_{4\kappa} + O\{\tau_{n}^{3/2}(m^{-1/2}+m/n) + \tau_{n}^{3}\}$$

$$= O(\tau_{n}^{3}).$$
(F.12)

Therefore, combining (F.11) and (F.12), since $n\tau_n^3 \to \infty$, $m/(n\tau_n^3) \to 0$, we have

$$\sup_{t \in \mathcal{I}} \|W_1(t) - \Omega(t)\beta(t)\|_{4\kappa} = O\left(\tau_n^3\right).$$
(F.13)

To proceed, we shall show that

$$\sup_{t \in \mathcal{I}} \|W_2(t)\|_{4\kappa} = O\{(n\tau_n^{3/2})^{-1/2} + \chi^m\}.$$
 (F.14)

Under Assumption 4.3, we have

$$\left|\mathbb{E}(\dot{\Delta}_{j})\right| = \frac{1}{m} \left|\sum_{i=j-m+1}^{j} \mathbb{E}\left(x_{i}x_{i}^{\top}x_{i+m}e_{i+m}\right) + \mathbb{E}\left(x_{i+m}x_{i+m}^{\top}x_{i}e_{i}\right)\right| = O(\chi^{m}).$$
(F.15)

Then, by Burkholder's inequality, for a sufficiently large C, we have

.

$$\left\|\sum_{j=m}^{n-m} \frac{\tilde{\omega}(t,j)}{2} \left\{ \hat{\Delta}_{j} - \mathbb{E}(\hat{\Delta}_{j}) \right\} \right\|_{4\kappa} = \left\|\sum_{s=-m}^{\infty} \sum_{j=m}^{n-m} \frac{\tilde{\omega}(t,j)}{2} \mathcal{P}_{j-s} \hat{\Delta}_{j} \right\|_{4\kappa} \le C \sum_{s=-m}^{\infty} \left\{ \sum_{j=m}^{n-m} \frac{\tilde{\omega}^{2}(t,j)}{4} \left\| \mathcal{P}_{j-s} \hat{\Delta}_{j} \right\|_{4\kappa}^{2} \right\}^{1/2}.$$
(F.16)

Under Assumptions 4.3 and 4.5, using similar techniques in Lemma 3 of (Zhou and Wu, 2010), we have

$$\left\| \mathcal{P}_{j-s} \dot{\Delta}_j \right\|_{4\kappa} \le \left\| \dot{\Delta}_j - \dot{\Delta}_{j,\{j-s\}} \right\|_{4\kappa} \le \frac{1}{m} \sum_{i=j-m+1}^{j+m} \{ \delta_{8\kappa}(J, i-j+s) + \delta_{8\kappa}(U, i-j+s) \}.$$
(F.17)

Then, (F.14) follows from (F.15), (F.16) and (F.17). Combining (F.13) and (F.14), since $m = O(n^{1/3})$, we have (F.8). Hence, by (F.4), under conditions $m = O(n^{1/3})$, we have

$$\sup_{t \in \mathcal{I}} \left\| \{ \breve{\Sigma}(t) - \Sigma^A(t) \} \mathbb{1}(A_n) \right\|_{\kappa} = O\left(\sqrt{m\tau_n^3} + \sqrt{\frac{m}{n\tau_n^{3/2}}} \right).$$

Finally, the result follows from the chaining argument in Proposition B.1 in Section B.2 in (Dette et al., 2019) and Proposition A.1 in (Wu and Zhou, 2018).

Theorem 4.1. Recall the definition of $\tilde{\Sigma}(t)$ in (F.3). Observe that

$$\hat{\Sigma}(t) - \tilde{\Sigma}(t) = \sum_{j=m}^{n-m} \frac{m\left(\Delta_j \Delta_j^\top - \tilde{\Delta}_j \tilde{\Delta}_j^\top\right)}{2} \omega(t, j) - \breve{\Sigma}(t)$$
$$= \Sigma^A(t) - \breve{\Sigma}(t) + \frac{m}{2} \sum_{j=m}^{n-m} \left(A_{j,m} \tilde{\Delta}_j^\top + \tilde{\Delta}_j A_{j,m}^\top\right) \omega(t, j).$$

Then, we have

$$\sup_{t \in \mathcal{I}} \left| \hat{\Sigma}(t) - \tilde{\Sigma}(t) \right| \le \sup_{t \in \mathcal{I}} \left| \Sigma^{A}(t) - \breve{\Sigma}(t) \right| + m \sup_{t \in \mathcal{I}} \left| \sum_{j=m}^{n-m} \omega(t,j) \tilde{\Delta}_{j} A_{j,m}^{\top} \right|.$$
(F.18)

Take note that $\Sigma^{A}(t)$ is the leading term of bias, so we introduce the correction. By Lemma F.1, we have

$$\sup_{t\in\mathcal{I}} \left| \Sigma^A(t) - \breve{\Sigma}(t) \right| = O_{\mathbb{P}} \left(\sqrt{m\tau_n^{3-1/\kappa}} + \sqrt{\frac{m}{n\tau_n^{3/2+2/\kappa}}} \right).$$
(F.19)

To proceed, define

$$h_s(t) = \sum_{j=m}^{n-m} \omega(t,j) \mathcal{P}_{j-s}(\tilde{\Delta}_j A_{j,m}^{\top}) = \sum_{j=m}^{n-m} h_{s,j}(t).$$

Under Assumption 4.5, it's straightforward that $\mathbb{E}(\tilde{\Delta}_j A_{j,m}^{\top}) = 0$, for $m \leq j \leq n - m$. Then, we can write $\sum_{j=m}^{n-m} \omega(t,j) \tilde{\Delta}_j A_{j,m}^{\top}$ as a summation of martingale differences, i.e.

$$\sum_{j=m}^{n-m} \omega(t,j) \tilde{\Delta}_j A_{j,m}^{\top} = \sum_{s=-m}^{\infty} h_s(t), \qquad (F.20)$$

Next, we shall show that $||h_{s,j}(t)||_{\kappa} = O\{(1/n + m^{-3/2})\min(\chi^{s-m}, 1)\}$. Under Assumptions 4.4 and 4.5, similar to (36) in Lemma 3 in (Zhou and Wu, 2010), we have

$$\delta_{2\kappa}(\tilde{\Delta}(m),k) := \sup_{1 \le j \le n} \|\tilde{\Delta}_j - \tilde{\Delta}_{j,\{j-k\}}\|_{2\kappa} = O\left\{\frac{1}{m} \sum_{i=-m+1}^m \delta_{2\kappa}(U,k+i)\right\}$$
$$= O\{\min(\chi^{k-m},1)/m\}.$$
(F.21)

Using similar arguments in (F.21), by the boundedness of $\beta(\cdot)$, we have

$$\sup_{1 \le j \le n} \left\| A_{j,m} - A_{j,m,\{j-k\}} \right\|_{2\kappa} \le \frac{M}{m} \sum_{i=-m+1}^{m} \delta_{4\kappa}(W,k+i) = O\{\min(\chi^{k-m},1)/m\},$$
(F.22)

where M is a sufficiently large positive constant. Following similar arguments in Theorem 1 of (Wu, 2007), $\sup_{j} \|\tilde{\Delta}_{j}\|_{2\kappa} = O(m^{-1/2})$. Similar to (F.5), we have uniformly for $m \leq j \leq n - m$,

$$\|A_{j,m}\|_{2\kappa} = O(m/n + m^{-1/2}).$$
(F.23)

Under Assumption 4.2, from (F.21), (F.22) and (F.23), we obtain uniformly for $m \le j \le n-m$, $s \ge -m$,

$$\begin{split} \|h_{s,j}\|_{\kappa}/\omega(t,j) &= \|\mathcal{P}_{j-s}\{\tilde{\Delta}_{j}A_{j,m}^{\top}\}\|_{\kappa} \\ &\leq \|A_{j,m,\{j-s\}} - A_{j,m}\|_{2\kappa} \left\|\tilde{\Delta}_{j}^{\top}\right\|_{2\kappa} + \|A_{j,m,\{j-s\}}\|_{2\kappa} \left\|\tilde{\Delta}_{j}^{\top} - \tilde{\Delta}_{j,\{j-s\}}^{\top}\right\|_{2\kappa} \\ &= O\{(1/n + m^{-3/2})\min(\chi^{s-m}, 1)\}. \end{split}$$

Since $h_{s,j}(t)$ are martingale differences with respect to j, we have for $t \in [m/n, 1 - m/n]$,

$$\|h_s(t)\|_{\kappa}^2 = \sum_{j=m}^{n-m} \|h_{s,j}(t)\|_{\kappa}^2 = O\left\{ (n\tau_n)^{-1} (1/n^2 + m^{-3}) \min(\chi^{2s-2m}, 1) \right\}.$$
 (F.24)

By (F.20) and (F.24), we obtain

$$\left\| m \sum_{j=m}^{n-m} \omega(t,j) \tilde{\Delta}_j A_{j,m}^{\mathsf{T}} \right\|_{\kappa} \le m \sum_{s=-m}^{m} \|h_s(t)\|_{\kappa} + m \sum_{s=m+1}^{\infty} \|h_s(t)\|_{\kappa} = O\left(\sqrt{\frac{m}{n\tau_n}}\right).$$
(F.25)

By the chaining argument in Proposition B.1 in Section B.2 in (Dette et al., 2019), we have

$$\sup_{t \in I} \left| m \sum_{j=m}^{n-m} \omega(t,j) \tilde{\Delta}_j A_{j,m}^{\top} \right| = O_{\mathbb{P}} \left(\sqrt{\frac{m}{n\tau_n^{1+2/\kappa}}} \right).$$
(F.26)

Combining (F.18), (F.19) and (F.26), we obtain

$$\sup_{t \in I} \left| \hat{\Sigma}(t) - \tilde{\Sigma}(t) \right| = O_{\mathbb{P}} \left(\sqrt{m\tau_n^{3-1/\kappa}} + \sqrt{\frac{m}{n\tau_n^{3/2+2/\kappa}}} \right).$$
(F.27)

By Lemma 3 in (Zhou and Wu, 2010), Proposition B.1 in Section B.2 in (Dette et al., 2019), we have

$$\sup_{t \in I} \left| \tilde{\Sigma}(t) - E \tilde{\Sigma}(t) \right| = O_{\mathbb{P}} \left(\sqrt{\frac{m}{n\tau_n^{1+2/\kappa}}} \right).$$
(F.28)

Using similar techniques in Lemma 4 and Lemma 5 of (Zhou and Wu, 2010), which hold uniformly for $t \in (0, 1)$, (F.28) leads to

$$\sup_{t\in I} \left| \tilde{\Sigma}(t) - \Sigma(t) \right| = O_{\mathbb{P}} \left(\sqrt{\frac{m}{n\tau_n^{3/2+2/\kappa}}} + \frac{1}{m} + \sqrt{m\tau_n^{3-1/\kappa}} \right),\tag{F.29}$$

With (F.27) and (F.29), the supreme bound is thus proved.

F.3 Proof of Theorem 5.1

Let $C_n = \{j : j - n\tau_n^{3/2} \le i \le j + n\tau_n^{3/2}, \beta'''(t_i) \text{ exists and continuous}\}, C_m = \{j : j - m \le i \le j + m, \beta'''(t_i) \text{ exists and continuous}\}.$ Recall in Lemma F.1 $A_n = \{\sup_{t \in \mathcal{I}} |\Omega(t) - M^+(t)| \le l_n \{(m\tau_n^{3/2})^{-1/2} + m/(n\tau_n^{3/4}) + \tau_n^{3/4}\}\}$, and $\lim_{n \to \infty} \mathbb{P}(A_n) = 1$.

Lemma F.2. Under the conditions of Theorem 5.1, we have

$$\sup_{j \in \mathcal{C}_m} \sup_{m_1 = 1, \dots, m} \|\{\check{\beta}(t_{j+m}) - \check{\beta}(t_j)\} \mathbb{1}(A_n)\|_{4\kappa} = O\left\{\frac{mq_n}{n\tau_n^{3/2}} + (n\tau_n^{3/2})^{-1/2}\right\}.$$

Proof of Lemma F.2. Following the arguments in Lemma F.1, since $\Omega(t)$ is invertible and continuous on A_n , and $\sup_{j \in \mathcal{C}_m} \|\varpi(t_j)\|_{8\kappa} = O(1)$, we have

$$\sup_{j \in \mathcal{C}_{m}} \sup_{m_{1}=1,...,m} \|\{\check{\beta}(t_{j+m_{1}}) - \check{\beta}(t_{j})\}1(A_{n})\|_{4\kappa} \\ = \sup_{j \in \mathcal{C}_{m}} \sup_{m_{1}=1,...,m} \|\{\Omega^{-1}(t_{j+m_{1}})\varpi(t_{j+m_{1}}) - \Omega^{-1}(t_{j})\varpi(t_{j})\}1(A_{n})\|_{4\kappa} \\ \leq \sup_{j \in \mathcal{C}_{m}} \sup_{m_{1}=1,...,m} \sqrt{p} \|\rho\{\Omega^{-1}(t_{j+m_{1}})\}|\varpi(t_{j+m_{1}}) - \varpi(t_{j})|1(A_{n})\|_{4\kappa} \\ + \sup_{j \in \mathcal{C}_{m}} \sup_{m_{1}=1,...,m} \sqrt{p} \|\rho\{\Omega^{-1}(t_{j+m_{1}})\}|\Omega(t_{j+m_{1}}) - \Omega(t_{j})|\rho\{\Omega^{-1}(t_{j})\}\varpi(t_{j})1(A_{n})\|_{4\kappa} \\ \leq C_{1} \sup_{j \in \mathcal{C}_{m}} \sup_{m_{1}=1,...,m} \||\varpi(t_{j+m_{1}}) - \varpi(t_{j})|1(A_{n})\|_{4\kappa} + C_{2}m/n,$$
(F.30)

where C_1 and C_2 are sufficiently large constants.

By (F.30), it is sufficient to show

$$\sup_{r \in \mathcal{C}_m} \sup_{m_1 = 1, \dots, m} \|\{\varpi(t_{r+m_1}) - \varpi(t_r)\} \mathbb{1}(A_n)\|_{4\kappa} = O\{(mq_n)/(n\tau_n^{3/2}) + (n\tau_n^{3/2})^{-1/2}\}.$$
 (F.31)

By triangle inequality, we have

$$\sup_{r \in \mathcal{C}_{m}} \sup_{m_{1}=1,...,m} \|\{\varpi(t_{r+m_{1}}) - \varpi(t_{r})\}1(A_{n})\|_{4\kappa} \\ \leq \sup_{r \in \mathcal{C}_{m}} \sup_{m_{1}=1,...,m} \left\|\left\{\varpi(t_{r+m_{1}}) - \sum_{j=m}^{n-m} \frac{\Delta_{j}\tilde{\omega}(t_{r+m_{1}},j)}{2}\beta(t_{j})\right\}1(A_{n})\right\|_{4\kappa} \\ + \sup_{r \in \mathcal{C}_{m}} \sup_{m_{1}=1,...,m} \left\|\left\{\varpi(t_{r}) - \sum_{j=m}^{n-m} \frac{\Delta_{j}\tilde{\omega}(t_{r},j)}{2}\beta(t_{j})\right\}1(A_{n})\right\|_{4\kappa} \\ + \sup_{r \in \mathcal{C}_{m}} \sup_{m_{1}=1,...,m} \left\|\left[\sum_{j=m}^{n-m} \Delta_{j}\{\tilde{\omega}(t_{r},j) - \tilde{\omega}(t_{r+m_{1}},j)\}\beta(t_{j})/2\right]1(A_{n})\right\|_{4\kappa} \\ \leq 2\sup_{t \in I} \left\|\left\{\varpi(t) - \sum_{j=m}^{n-m} \frac{\Delta_{j}\tilde{\omega}(t,j)}{2}\beta(t_{j})\right\}1(A_{n})\right\|_{4\kappa} \\ + \sup_{r \in \mathcal{C}_{m}} \sup_{m_{1}=1,...,m} \left\|\left[\sum_{j=m}^{n-m} \Delta_{j}\{\tilde{\omega}(t_{r},j) - \tilde{\omega}(t_{r+m_{1}},j)\}\beta(t_{j})/2\right]1(A_{n})\right\|_{4\kappa} \right\|_{4\kappa} .$$
(F.32)

Based on (F.32), we break the proof for (F.31) into two parts. First, we investigate the first term in (F.32). Recall the definition of $\dot{\Delta}_j$ and $\dot{A}_{j,m}$ in (F.2) and (F.1) and we have from (F.10) that

$$\varpi(t) = \sum_{j=m}^{n-m} \frac{\tilde{\omega}(t,j)}{2} \dot{A}_{j,m} + \sum_{j=m}^{n-m} \frac{\tilde{\omega}(t,j)}{2} \dot{\Delta}_j = W_1(t) + W_2(t),$$

where we directly have from (F.14) that

$$\sup_{t \in \mathcal{I}} \|W_2(t)\|_{4\kappa} = O\{(n\tau_n^{3/2})^{-1/2}\}.$$

Therefore, we have

$$\sup_{t \in I} \left\| \left\{ \varpi(t) - \sum_{j=m}^{n-m} \frac{\Delta_{j} \tilde{\omega}(t,j)}{2} \beta(t_{j}) \right\} 1(A_{n}) \right\|_{4\kappa} \\
\leq \sup_{t \in I} \sum_{j=m}^{n-m} \frac{\tilde{\omega}(t,j)}{2m} \left(\sum_{i=j-m+1}^{j} \left\| x_{i} x_{i}^{\top} - x_{i+m} x_{i+m}^{\top} \right\|_{8\kappa} \\
\times \left\| x_{i} x_{i}^{\top} (\beta(t_{i}) - \beta(t_{j})) - x_{i+m} x_{i+m}^{\top} (\beta(t_{i+m}) - \beta(t_{j})) \right\|_{8\kappa} \right) + \sup_{t \in \mathcal{I}} \left\| W_{2}(t) \right\|_{4\kappa} \\
\leq \sup_{t \in I} \left(\sum_{j \in \mathcal{C}_{m}, m \leq j \leq n-m} + \sum_{j \notin \mathcal{C}_{m}, m \leq j \leq n-m} \right) \frac{\tilde{\omega}(t,j)}{2m} \left(\sum_{i=j-m+1}^{j} \left\| x_{i} x_{i}^{\top} - x_{i+m} x_{i+m}^{\top} \right\|_{8\kappa} \\
\times \left\| x_{i} x_{i}^{\top} (\beta(t_{i}) - \beta(t_{j})) - x_{i+m} x_{i+m}^{\top} (\beta(t_{i+m}) - \beta(t_{j})) \right\|_{8\kappa} \right) + \sup_{t \in \mathcal{I}} \left\| W_{2}(t) \right\|_{4\kappa} \\
= O\left\{ \frac{m}{n} + \frac{mq_{n}}{n\tau_{n}^{3/2}} + (n\tau_{n}^{3/2})^{-1/2} + \chi^{m} \right\}.$$
(F.33)

Secondly, by the continuity of the kernel function $K(\cdot)$ and since $\sup_{j=m,\dots,n-m} \| \acute{\Delta}_j \| = O(1)$, we have

$$\sup_{r \in \mathcal{C}_m} \sup_{m_1 = 1, \dots, m} \left\| \left[\sum_{j=m}^{n-m} \Delta_j \{ \tilde{\omega}(t_r, j) - \tilde{\omega}(t_{r+m_1}, j) \} \beta(t_j) / 2 \right] 1(A_n) \right\|_{4\kappa} = O\{m/(n\tau_n^{3/2})\}.$$
(F.34)

Then, (F.31) follows from (F.33) and (F.34). The lemma follows from (F.30) and (F.31). \Box

Proof of Theorem 2. For sufficiently large constants C_1 and C_2 , we have

$$\sup_{t\in\mathcal{I}} \left| \{ \check{\Sigma}(t) - \Sigma^{A}(t) \} \mathbb{1}(A_{n}) \right| \\
\leq \sup_{t\in\mathcal{I}} \left| \sum_{j\in\mathcal{C}_{n}, m\leq j\leq n-m} \frac{m\omega(t,j)}{2} (\hat{A}_{j,m} \hat{A}_{j,m}^{\top} - A_{j,m} A_{j,m}^{\top}) \mathbb{1}(A_{n}) \right| \\
+ C_{1} \sum_{j\notin\mathcal{C}_{n}, j\in\mathcal{C}_{m} m\leq j\leq n-m} \frac{m}{n\tau_{n}} |(\hat{A}_{j,m} \hat{A}_{j,m}^{\top} - A_{j,m} A_{j,m}^{\top}) \mathbb{1}(A_{n})| \\
+ C_{2} \sum_{j\notin\mathcal{C}_{m}, m\leq j\leq n-m} \frac{m}{n\tau_{n}} |(\hat{A}_{j,m} \hat{A}_{j,m}^{\top} - A_{j,m} A_{j,m}^{\top}) \mathbb{1}(A_{n})|.$$
(F.35)

For $j \in \mathcal{C}_m$, by (F.23), we have $||A_{j,m}||_{2\kappa} = O(m^{-1/2} + m/n)$. We shall show that

$$\sup_{j \in \mathcal{C}_m} \left\| (\hat{A}_{j,m} - A_{j,m}) \mathbf{1}(A_n) \right\|_{2\kappa} = O(m^{-1/2}), \tag{F.36}$$

and

$$\sup_{j \in \mathcal{C}_n} \| (\hat{A}_{j,m} - A_{j,m}) \mathbf{1}(A_n) \|_{2\kappa} = O\{ (mq_n)/(n\tau_n^{3/2}) + (n\tau_n^{3/2})^{-1/2} + m^{-1/2}\tau_n^3 \}.$$

By Lemma F.2, we have

$$\begin{split} \sup_{j \in \mathcal{C}_{m}} \| (\hat{A}_{j,m} - A_{j,m}) \mathbf{1}(A_{n}) \|_{2\kappa} \\ &\leq \sup_{j \in \mathcal{C}_{m}} \left\| m^{-1} \sum_{i=j-m+1}^{j} x_{i} x_{i}^{\top} \left[\{\check{\beta}(t_{i}) - \check{\beta}(t_{i+m})\} - \{\beta(t_{i}) - \beta(t_{i+m})\} \right] \mathbf{1}(A_{n}) \right\|_{2\kappa} \\ &+ \sup_{j \in \mathcal{C}_{m}} \left\| m^{-1} \sum_{i=j-m+1}^{j} (x_{i} x_{i}^{\top} - x_{i+m} x_{i+m}^{\top}) \{\check{\beta}(t_{i+m}) - \beta(t_{i+m})\} \mathbf{1}(A_{n}) \right\|_{2\kappa} \\ &\leq C_{1} \left\{ \frac{mq_{n}}{n\tau_{n}^{3/2}} + (n\tau_{n}^{3/2})^{-1/2} \right\} + C_{2}m/n + \sup_{j \in \mathcal{C}_{m}} \left\| m^{-1} \sum_{i=j-m+1}^{j} (x_{i} x_{i}^{\top} - x_{i+m} x_{i+m}^{\top}) \right\|_{4\kappa} \| \{\check{\beta}(t_{j}) - \beta(t_{j})\} \mathbf{1}(A_{n}) \|_{2\kappa} \\ &= O(m^{-1/2}), \end{split}$$
(F.37)

where the last line follows from similar arguments in proving (F.7). Therefore, we have shown (F.36). Following the arguments in (F.37), for the indices $j \in C_n \subset C_m$, by (F.8), we have

$$\sup_{j \in \mathcal{C}_n} \|(\hat{A}_{j,m} - A_{j,m}) \mathbf{1}(A_n)\|_{2\kappa} = O\{(mq_n)/(n\tau_n^{3/2}) + (n\tau_n^{3/2})^{-1/2} + m^{-1/2}\tau_n^3\}.$$

For the first term of (F.35), note that

$$\left\| \sum_{j \in \mathcal{C}_{n}, m \leq j \leq n-m} \frac{m\omega(t,j)}{2} (\hat{A}_{j,m} \hat{A}_{j,m}^{\top} - A_{j,m} A_{j,m}^{\top}) \mathbf{1}(A_{n}) \right\|_{\kappa}$$

$$\leq \sum_{j \in \mathcal{C}_{n}, m \leq j \leq n-m} \frac{m\omega(t,j)}{2} \left\| (\hat{A}_{j,m} - A_{j,m}) \mathbf{1}(A_{n}) \right\|_{2\kappa}$$

$$\times \left(\left\| (\hat{A}_{j,m} - A_{j,m}) \mathbf{1}(A_{n}) \right\|_{2\kappa} + 2 \|A_{j,m}\|_{2\kappa} \right)$$

$$= O\left(\frac{m^{3/2}q_{n}}{n\tau_{n}^{3/2}} + \tau_{n}^{3} + \sqrt{\frac{m}{n\tau_{n}^{3/2}}} \right).$$
(F.38)

By the chaining argument of Proposition B.1 of (Dette et al., 2019), by we have

$$\sup_{t \in \mathcal{I}} \left| \sum_{j \in \mathcal{C}_n, m \le j \le n-m} \frac{m\omega(t,j)}{2} (\hat{A}_{j,m} \hat{A}_{j,m}^\top - A_{j,m} A_{j,m}^\top) 1(A_n) \right| = O_{\mathbb{P}} \left(\sqrt{\frac{m}{n\tau_n^{3/2+2/\kappa}}} + \tau_n^{3-1/\kappa} + \frac{m^{3/2}q_n}{n\tau_n^{3/2+1/\kappa}} \right)$$

For the second term of (F.35), by (F.36), we have

$$\sum_{\substack{j \notin \mathcal{C}_{n}, j \in \mathcal{C}_{m}m \leq j \leq n-m \\ j \notin \mathcal{C}_{n}, j \in \mathcal{C}_{m}m \leq j \leq n-m }} \frac{m}{n\tau_{n}} \mathbb{E} |(\hat{A}_{j,m} \hat{A}_{j,m}^{\top} - A_{j,m} A_{j,m}^{\top}) \mathbf{1}(A_{n})|$$

$$\leq \sum_{\substack{j \notin \mathcal{C}_{n}, j \in \mathcal{C}_{m}m \leq j \leq n-m \\ n\tau_{n}}} \frac{m}{n\tau_{n}} \left\| (\hat{A}_{j,m} - A_{j,m}) \mathbf{1}(A_{n}) \right\| \left(\left\| (\hat{A}_{j,m} - A_{j,m}) \mathbf{1}(A_{n}) \right\| + 2 \|A_{j,m}\| \right)$$

$$= O(q_{n}\tau_{n}^{1/2}).$$
(F.39)

Since for $j \notin C_m$, $||A_{j,m}||_{2\kappa} = O(1)$ and $\left\| (\hat{A}_{j,m} - A_{j,m}) \mathbb{1}(A_n) \right\|_{2\kappa} = O(1)$, for the third term of (F.35), we have

$$\sum_{j \notin \mathcal{C}_m, m \le j \le n-m} \frac{m}{n\tau_n} \mathbb{E}|(\hat{A}_{j,m} \hat{A}_{j,m}^\top - A_{j,m} A_{j,m}^\top) \mathbf{1}(A_n)| = O\left(\frac{m^2 q_n}{n\tau}\right).$$
(F.40)

Combining (F.35), (F.38), (F.39) and (F.40), since $\kappa \ge 1$, $m\tau_n^3 \to \infty$, we have

$$\sup_{t \in \mathcal{I}} \left| \breve{\Sigma}(t) - \Sigma^{A}(t) \right| = O_{\mathbb{P}} \left(q_{n} \tau_{n}^{1/2} + \sqrt{\frac{m}{n\tau_{n}^{3/2+2/\kappa}}} + \frac{m^{2}q_{n}}{n\tau_{n}} + \tau_{n}^{3-1/\kappa} + \frac{m^{3/2}q_{n}}{n\tau_{n}^{3/2+1/\kappa}} \right) \\ = O_{\mathbb{P}} (q_{n} \tau_{n}^{1/2} + \sqrt{\frac{m}{n\tau_{n}^{7/2}}} + \frac{m^{2}q_{n}}{n\tau_{n}} + \tau_{n}^{2})$$
(F.41)

We proceed to investigate Theorem 4.1 in the presence of change points in $\beta(t)$. Recall that from (F.18),

$$\sup_{t\in\mathcal{I}} \left| \hat{\Sigma}(t) - \tilde{\Sigma}(t) \right| \le \sup_{t\in\mathcal{I}} \left| \Sigma^{A}(t) - \breve{\Sigma}(t) \right| + m \sup_{t\in\mathcal{I}} \left| \sum_{j=m}^{n-m} \omega(t,j) \tilde{\Delta}_{j} A_{j,m}^{\top} \right|.$$
(F.42)

Inspecting (F.24), we have

$$\begin{aligned} \|h_s(t)\|_{\kappa}^2 &= \sum_{j \in \mathcal{C}_m, m \le j \le n-m} \|h_{s,j}(t)\|_{\kappa}^2 + \sum_{j \notin \mathcal{C}_m, m \le j \le n-m} \|h_{s,j}(t)\|_{\kappa}^2 \\ &= O\left\{ (n\tau_n)^{-1} (1/n^2 + m^{-3}) \min(\chi^{2s-2m}, 1) + \frac{q_n}{m(n\tau_n)^2} \min(\chi^{2s-2m}, 1) \right\} \end{aligned}$$

Similar to (F.25), we have

$$\left\| m \sum_{j=m}^{n-m} \omega(t,j) \tilde{\Delta}_j A_{j,m}^{\mathsf{T}} \right\|_{\kappa} \le m \sum_{s=-m}^{m} \|h_s(t)\|_{\kappa} + m \sum_{s=m+1}^{\infty} \|h_s(t)\|_{\kappa}$$
$$= O\left\{ \sqrt{\frac{m}{n\tau_n}} + \sqrt{\frac{q_n m^3}{(n\tau_n)^2}} \right\} = O\left(\sqrt{\frac{m}{n\tau_n}}\right)$$

By the chaining argument in Proposition B.1 in Section B.2 in (Dette et al., 2019), similar to (F.26), we have

$$\sup_{t \in I} \left| m \sum_{j=m}^{n-m} \omega(t,j) \tilde{\Delta}_j A_{j,m}^{\top} \right| = O_{\mathbb{P}} \left(\sqrt{\frac{m}{n\tau_n^{1+2/\kappa}}} \right).$$
(F.43)

Finally, combining (F.41), (F.42), and (F.43), following similar argument of Theorem 4.1, we have shown the first part of the results. The second part follows from Lemma G.1 and Theorem 3.4 of (Wu and Zhou, 2018). \Box

F.4 Proof of Proposition 5.1

Proof of Proposition 5.1. Proposition 5.1 follows from similar but easier arguments of Theorem 3.1 and Theorem 3.2 of (Wu and Zhou, 2018). \Box

F.5 Proof of Theorem 5.2

For the clarity of presentation, write $\hat{\Sigma}_d(t)$ for $\hat{\Sigma}(t)$ defined in (3.3) under the fixed alternative. Define

$$\Sigma_d(t) = \kappa_2(d)\sigma_H^2(t)\mu_W(t)\mu_W^{\top}(t), \quad t \in [0, 1].$$

For the quantities under the fixed alternatives, let $\tilde{Q}_{k,m}^{(d)} = \sum_{i=k}^{k+m-1} x_i e_i^{(d)}$,

$$\tilde{\Delta}_j^{(d)} = \frac{\tilde{Q}_{j-m+1,m}^{(d)} - \tilde{Q}_{j+1,m}^{(d)}}{m}, \quad \tilde{\Sigma}_d(t) = \sum_{j=m}^{n-m} \frac{m \tilde{\Delta}_j^{(d)} \tilde{\Delta}_j^{(d),\top}}{2} \omega(t,j),$$

and

$$\dot{\Delta}_{j}^{(d)} = \frac{1}{m} \sum_{i=j-m+1}^{j} (x_{i}x_{i}^{\top} - x_{i+m}x_{i+m}^{\top})(x_{i}e_{i}^{(d)} - x_{i+m}e_{i+m}^{(d)}).$$

Let $\hat{A}^{(d)}, \varpi^{(d)}(\cdot), \breve{\beta}^{(d)}(\cdot), \breve{\Sigma}_{d}(\cdot)$ denote the counterparts of $\hat{A}, \varpi(\cdot), \breve{\beta}(\cdot)$ and $\breve{\Sigma}(\cdot)$ in (3.3) under the fixed alternatives. Define $H^{(d)}(t_i, \mathcal{F}_i) = \sum_{k=0}^{\infty} \psi_k H(t_{i-k}, \mathcal{F}_{i-k}), U^{(d)}(t, \mathcal{F}_i) = W(t, \mathcal{F}_i) H^{(d)}(t, \mathcal{F}_i)$. The following proposition is from Theorem 4.2 of (Bai and Wu, 2024).

Proposition F.1. Under Assumptions 4.3 and 5.1, we have

.

$$\max_{\lfloor nb_n \rfloor + 1 \le r \le n - \lfloor nb_n \rfloor} \left| \sum_{i=\lfloor nb_n \rfloor + 1}^r x_{i,n} e_{i,n}^{(d)} - \sum_{i=\lfloor nb_n \rfloor + 1}^r \mu_W(t_i) e_{i,n}^{(d)} \right| = O_{\mathbb{P}}(\sqrt{n}(\log n)^d).$$

Corollary F.1. Under the conditions of Proposition F.1, assume $l/\log n \to \infty$, $l/n \to 0$, we have

$$\max_{1 \le k \le n-l+1} \left\| \sum_{i=k}^{k+l-1} (x_i - \mu_W(t_i)) e_i^{(d)} \right\| = O(\sqrt{l(\log n)^d}).$$

Proof of Corollary F.1. The corollary follows from a careful check of the proof of Theorem 4.2 of (Bai and Wu, 2024). \Box

Lemma F.3. Under the conditions of Theorem 5.2, we have

$$\sup_{t\in\mathcal{I}} \left| \breve{\Sigma}_d(t) - \Sigma^A(t) \right| = O_{\mathbb{P}} \left[\sqrt{m} \{ (n\tau_n^{3/2})^{d-1/2} + \tau_n^3 \} \tau_n^{-1/\kappa} + m \{ (n\tau_n^{3/2})^{2d-1} + \tau_n^6 \} \tau_n^{-1/\kappa} \right] = o_{\mathbb{P}}(m^{2d}).$$

Proof of Lemma F.3. After a careful check of the proof of Lemma F.1, the behavior of $W_1(t)$ is unchanged under the fixed alternatives and it's sufficient to show that $W_2^{(d)}(t)$, $W_2(t)$ under the fixed alternatives, s.t.

$$\sup_{t \in \mathcal{I}} \left\| W_2^{(d)}(t) \right\|_{4\kappa} = O\{ (n\tau_n^{3/2})^{d-1/2} \}.$$
 (F.44)

Then the lemma will follow from the similar steps in Lemma F.1. Similar to (F.17), under Assumptions 4.3 and 4.5, following similar arguments in Lemma G.3, we have

$$\left\| \mathcal{P}_{j-s} \dot{\Delta}_{j}^{(d)} \right\|_{4\kappa} \leq \frac{1}{m} \sum_{i=j-m+1}^{j+m} \left\{ \delta_{8\kappa}(J, i-j+s) + \delta_{8\kappa}(U^{(d)}, i-j+s) \right\} = O\left\{ \frac{1}{m} \sum_{i=-m+1}^{m} (i+s)^{d-1} \right\}.$$

Let $N = N_n = \lfloor n \tau_n^{3/2} \rfloor$. Note that under Assumption 4.5, we can write

$$W_2^{(d)}(t) = \sum_{s=0}^N \sum_{j=m}^{n-m} \frac{\tilde{\omega}(t,j)}{2} \mathcal{P}_{j-s} \dot{\Delta}_j^{(d)} + \sum_{j=m}^{n-m} \frac{\tilde{\omega}(t,j)}{2} \sum_{s=N+1}^{\infty} \mathcal{P}_{j-s} \dot{\Delta}_j^{(d)} = W_{21}^{(d)}(t) + W_{22}^{(d)}(t).$$

Since $\mathcal{P}_{j-s}\dot{\Delta}_{j}^{(d)}$ are martingale differences with respect to j, for $0 \leq s \leq N$, we have

$$\sup_{t \in \mathcal{I}} \left\| \sum_{j=m}^{n-m} \frac{\tilde{\omega}(t,j)}{2} \mathcal{P}_{j-s} \dot{\Delta}_{j}^{(d)} \right\|_{4\kappa}^{2} = \sum_{j=m}^{n-m} \frac{\tilde{\omega}^{2}(t,j)}{4} \left\| \mathcal{P}_{j-s} \dot{\Delta}_{j}^{(d)} \right\|_{4\kappa}^{2} \\ = O\left[\frac{1}{n\tau_{n}^{3/2}m^{2}} \left\{ \sum_{i=-m+1}^{m} (i+s)^{d-1} \right\}^{2} \right].$$
(F.45)

Therefore, by (F.45), we have

$$\sup_{t \in \mathcal{I}} \left\| W_{21}^{(d)}(t) \right\|_{4\kappa} = O(N^d / \sqrt{n\tau_n^{3/2}}) = O\{(n\tau_n^{3/2})^{d-1/2}\}.$$
 (F.46)

Since $\mathcal{P}_{j-s} \dot{\Delta}_{j}^{(d)}$ are martingale differences with respect to s, elementary calculations shows

$$\left\|\sum_{s=N+1}^{\infty} \mathcal{P}_{j-s} \dot{\Delta}_{j}^{(d)}\right\|_{4\kappa}^{2} = O\left[m^{-2} \sum_{s=N+1}^{\infty} \left\{\sum_{i=-m+1}^{m} (i+s)^{d-1}\right\}^{2}\right] = O(N^{2d-1}).$$
(F.47)

Therefore, by (F.47) and triangle inequality we have

$$\sup_{t \in \mathcal{I}} \left\| W_{22}^{(d)}(t) \right\|_{4\kappa} = O(N^{d-1/2}) = O\{ (n\tau_n^{3/2})^{d-1/2} \}.$$
(F.48)

Finally, (F.44) follows from (F.46) and (F.48).

Proof of Theorem 5.2. Recall $\tilde{Q}_{k,m}^{(d)} = \sum_{i=k}^{k+m-1} x_i e_i^{(d)}$, and

$$\tilde{\Delta}_{j}^{(d)} = \frac{\tilde{Q}_{j-m+1,m}^{(d)} - \tilde{Q}_{j+1,m}^{(d)}}{m}, \quad \tilde{\Sigma}_{d}(t) = \sum_{j=m}^{n-m} \frac{m\tilde{\Delta}_{j}^{(d)} \{\tilde{\Delta}_{j}^{(d)}\}^{\top}}{2} \omega(t,j).$$

We break the proof into 6 steps.

Step 1: We shall prove that under the bandwidth conditions $\kappa \geq 4/(1/2 - d)$, $m/(n\tau_n^3) \rightarrow 0$, $\sqrt{m\tau_n^{3-1/\kappa}} \rightarrow 0$, $m = O(n^{1/3})$, $m \rightarrow \infty$, $n\tau_n^3 \rightarrow \infty$,

$$\sup_{t \in [0,1]} \left| \hat{\Sigma}_d(t) - \tilde{\Sigma}_d(t) \right| = O_{\mathbb{P}} \left[\sqrt{m} \{ (n\tau_n^{3/2})^{d-1/2} + \tau_n^3 \} \tau_n^{-1/\kappa} + m \{ (n\tau_n^{3/2})^{2d-1} + \tau_n^6 \} \tau_n^{-1/\kappa} \right]$$
$$= o_{\mathbb{P}}(m^{2d}).$$
(F.49)

Recall that $\Sigma^{A}(t) = \sum_{j=m}^{n-m} \frac{m\omega(t,j)}{2} A_{j,m} A_{j,m}^{\top}$, and similar to (F.18), we have

$$\sup_{t\in\mathcal{I}} \left| \hat{\Sigma}_d(t) - \tilde{\Sigma}_d(t) \right| \le \sup_{t\in\mathcal{I}} \left| \breve{\Sigma}_d(t) - \Sigma^A(t) \right| + \sup_{t\in\mathcal{I}} \left| \sum_{j=m}^{n-m} m\omega(t,j) \tilde{\Delta}_j^{(d)} A_{j,m}^\top \right|,$$

where the first term has been investigated in Lemma F.3.

Define $h_{s,j}^{(d)}(t) = \mathcal{P}_{j-s}(\tilde{\Delta}_j^{(d)}A_{j,m}^{\top})$. Let $N = N_n = \lfloor n\tau_n \rfloor$. Observe that under Assumption 4.5,

$$\sum_{j=m}^{n-m} \omega(t,j) \tilde{\Delta}_j^{(d)} A_{j,m}^{\top} = \sum_{s=0}^N \sum_{j=m}^{n-m} \omega(t,j) h_{s,j}^{(d)} + \sum_{j=m}^{n-m} \omega(t,j) \sum_{s=N+1}^{\infty} h_{s,j}^{(d)} = S_1 + S_2,$$

where S_1 and S_2 are defined in the obvious way. To proceed, we first calculate $\|h_{s,j}^{(d)}(t)\|_{\kappa}$.

By Lemma F.5, $\sup_j \|\tilde{\Delta}_j^d\|_{2\kappa} = O(m^{d-1/2})$. Then, we have

$$\|h_{s,j}^{(d)}(t)\|_{\kappa} = O\left\{m^{d-3/2}\min(chi^{s-m}, 1) + (1/n + m^{-3/2})\sum_{i=-m+1}^{m}\psi_{s+i}\right\}.$$
 (F.50)

Since $h_{s,h}^{(d)}(t)$ are martingale differences with respect to j, since $m/(n\tau_n) \to 0$, we have for $t \in \mathcal{I}$,

$$\|S_1\|_{\kappa} \le \sum_{s=0}^{N} \left\| \sum_{j=m}^{n-m} \omega(t,j) h_{s,j}^{(d)} \right\|_{\kappa} = O\left[\sum_{s=0}^{N} \left\{ \sum_{j=m}^{n-m} \omega^2(t,j) \|h_{s,j}^{(d)}\|_{\kappa}^2 \right\}^{1/2} \right] = O\{m^{-1/2} (n\tau_n)^{d-1/2}\}.$$
 (F.51)

By (F.50) and triangle inequality, elementary calculation shows that

$$\|S_2\|_{\kappa} \le \sum_{j=m}^{n-m} \omega(t,j) \left\| \sum_{s=N+1}^{\infty} h_{s,j}^{(d)} \right\|_{\kappa} = O\left\{ \left(\sum_{s=N+1}^{\infty} \max_{m \le j \le n-m} \|h_{s,j}^{(d)}\|_{\kappa}^2 \right)^{1/2} \right\} = O\{m^{-1/2} (n\tau_n)^{d-1/2}\}.$$
(F.52)

where the first big O follows from the fact that $h_{s,j}^{(d)}$ are martingale differences with respect to s. Combining (F.51) and (F.52), by chaining argument in Proposition B.1 in (Dette et al., 2019), we have

$$\sup_{t\in\mathcal{I}} \left| \sum_{j=m}^{n-m} m\omega(t,j)\tilde{\Delta}_j^{(d)} A_{j,m}^{\top} \right| = O_{\mathbb{P}} \left\{ \sqrt{\frac{m}{(n\tau_n)^{1-2d}}} \tau_n^{-1/\kappa} \right\}.$$
(F.53)

Combining Lemma F.4 and (F.53), we have shown (F.49).

Step 2: Define $\bar{Q}_{k,m}^{(d)} = \sum_{i=k}^{k+m-1} \mu_W(t_i) e_i^{(d)} (k = 1, \dots, n-m+1),$

$$\bar{\Delta}_{j}^{(d)} = \frac{\bar{Q}_{j-m+1,m}^{(d)} - \bar{Q}_{j+1,m}^{(d)}}{m}, \quad \bar{\Sigma}_{d}(t) = \sum_{j=m}^{n-m} \frac{m\bar{\Delta}_{j}^{(d)}(\bar{\Delta}_{j}^{(d)})^{\top}}{2} \omega(t,j).$$

We shall show that under the bandwidth condition $m\tau_n^{3/2}/\log n \to \infty, \ \kappa \ge 2/(3d),$

$$\sup_{t\in\mathcal{I}} \left| \tilde{\Sigma}_d(t) - \bar{\Sigma}_d(t) \right| = O_{\mathbb{P}}\{ m^d (\log n)^d \tau_n^{-1/\kappa} \} = o_{\mathbb{P}}(m^{2d}).$$
(F.54)

Following similar arguments in Corollary F.1, we have

$$\max_{1 \le k \le n-m+1} \left\| \bar{Q}_{k,m}^{(d)} - \tilde{Q}_{k,m}^{(d)} \right\|_{2\kappa} = O(\sqrt{m(\log n)^d}).$$
(F.55)

Using (F.55), and the fact $\sup_{j} \|\tilde{\Delta}_{j}^{(d)}\|_{2\kappa} = O(m^{d-1/2}), \sup_{j} \|\bar{\Delta}_{j}^{(d)}\|_{2\kappa} = O(m^{d-1/2})$, we have $\left\|\tilde{\Delta}_{j}^{(d)}\tilde{\Delta}_{j}^{(d),\top} - \bar{\Delta}_{j}^{(d)}\bar{\Delta}_{j}^{(d),\top}\right\|_{\kappa} \leq \left\|\tilde{\Delta}_{j}^{(d)} - \bar{\Delta}_{j}^{(d)}\right\|_{2\kappa} \left\|\bar{\Delta}_{j}^{(d),\top}\right\|_{2\kappa}$

$$+ \left\| \tilde{\Delta}_{j}^{(d)} \right\|_{2\kappa} \left\| \left(\tilde{\Delta}_{j}^{(d)} - \bar{\Delta}_{j}^{(d)} \right)^{\top} \right\|_{2\kappa} \right\|_{2\kappa}$$
$$= O\{m^{d-1}(\log n)^{d}\}.$$

Since $m\tau_n^{3/2}/\log n \to \infty$, (F.54) follows from triangle inequality and Proposition B.1 in (Dette et al., 2019).

Step 3: Let $\zeta_j = \sum_{i=j}^{\infty} \mathcal{P}_j e_i, \zeta_j^\circ = \zeta_j(t_j) = \sum_{i=j}^{\infty} \mathcal{P}_j H(t_j, \mathcal{F}_i)$. Define $\overline{Z}_{k,m} = \sum_{j=0}^{L} \psi_j \sum_{i=k}^{k+m-1} \mu_W(t_i) \zeta_{i-j}^\circ$,

$$\Delta_j^{(d),\circ} = \frac{\overline{Z}_{j-m+1,m} - \overline{Z}_{j+1,m}}{m}, \quad \Sigma_d^{\circ}(t) = \sum_{j=m}^{n-m} \frac{m\Delta_j^{(d),\circ}(\Delta_j^{(d),\circ})^{\top}}{2} \omega(t,j).$$

Let $L = Mm^{1+\frac{1}{2d+1}}\tau_n^{1/2}$, where M is a sufficiently large constant. We will show that

$$\sup_{t \in \mathcal{I}} |\bar{\Sigma}_d(t) - \Sigma_d^{\circ}(t)| = O_{\mathbb{P}} \left\{ m^{2d} \left(m^{-\frac{1/2 - d}{2d + 1}} \tau_n^{d/2 - 1/4 - 1/\kappa} \right) \right\} = o_{\mathbb{P}}(m^{2d}).$$
(F.56)

Since $m\tau_n \to \infty$, $m^2 \tau_n^{1/2} / n = O(n^{-1/3} \tau_n^{1/2}) = o(1)$, then $L/m \to \infty$, $L/m^2 \to 0$, $m^{1+1/(2d)}/L \to \infty$, $L(\log n)^2/n \to 0$. Observe that

$$\begin{split} \left\| \bar{\Sigma}_{d}(t) - \Sigma_{d}^{\circ}(t) \right\|_{\kappa} \\ &\leq \sum_{j=m}^{n-m} \frac{m\omega(t,j)}{2} \left\| \bar{\Delta}_{j}^{(d)} \left(\bar{\Delta}_{j}^{(d)} \right)^{\top} - \Delta_{j}^{(d),\circ} \left(\Delta_{j}^{(d),\circ} \right)^{\top} \right\|_{\kappa} \\ &\leq m \max_{m \leq j \leq n-m} \left(\left\| \bar{\Delta}_{j}^{(d)} \right\|_{2\kappa} \left\| \left(\bar{\Delta}_{j}^{(d)} \right)^{\top} - \left(\Delta_{j}^{(d),\circ} \right)^{\top} \right\|_{2\kappa} + \left\| \bar{\Delta}_{j}^{(d)} - \Delta_{j}^{(d),\circ} \right\|_{2\kappa} \left\| \left(\Delta_{j}^{(d),\circ} \right)^{\top} \right\|_{2\kappa} \right), \quad (F.57) \end{split}$$

where

$$\left\|\bar{\Delta}_{j}^{(d)} - \Delta_{j}^{(d),\circ}\right\|_{2\kappa} \leq \frac{1}{m} \left(\left\|\bar{Q}_{j-m+1,m}^{(d)} - \overline{Z}_{j-m+1,m}\right\|_{2\kappa} + \left\|\bar{Q}_{j+1,m}^{(d)} - \overline{Z}_{j+1,m}\right\|_{2\kappa} \right).$$

Define

$$\overline{W}_{k,m} = \sum_{j=0}^{L} \psi_j \sum_{i=k}^{k+m-1} \mu_W(t_i) e_{i-j}, \quad 1 \le k \le n-m+1.$$

Then, we have for $1 \le k \le n - m + 1$ that

$$\begin{split} \left\| \bar{Q}_{k-m+1,m}^{(d)} - \overline{Z}_{k-m+1,m} \right\|_{2\kappa} &\leq \left\| \bar{Q}_{k-m+1,m}^{(d)} - \overline{W}_{k-m+1,m} \right\|_{2\kappa} + \left\| \overline{W}_{k-m+1,m} - \overline{Z}_{k-m+1,m} \right\|_{2\kappa} \\ &= C_{2\kappa,1} + C_{2\kappa,2}, \end{split}$$
(F.58)

where $C_{2\kappa,1}$ and $C_{2\kappa,2}$ are defined in the obvious way.

Under conditions 4.5 and 4.3, by Burkholder's inequality, we have for $k \ge m$,

$$C_{2\kappa,1} = \left\| \sum_{j=L+1}^{\infty} \left\{ \sum_{i=0}^{\min\{m,j-L\}-1} \psi_{j-i} \mu_W\left(\frac{k-i}{n}\right) \right\} e_{k-j} \right\|_{2\kappa}$$

$$\leq \sum_{t=0}^{\infty} \left[\sum_{j=L+1}^{\infty} \left\{ \sum_{i=0}^{\min\{m,j-L\}-1} \psi_{j-i} \mu_W\left(\frac{k-i}{n}\right) \right\}^2 \|\mathcal{P}_{k-j-t}e_{k-j}\|_{2\kappa}^2 \right]^{1/2}$$

$$\leq \left[\sum_{j=L+1}^{\infty} \left\{ \sum_{i=0}^{m} \psi_{j-i} \mu_W\left(\frac{k-i}{n}\right) \right\}^2 \right]^{1/2} \sum_{t=0}^{\infty} \delta_{2\kappa}(H,t,(-\infty,1])$$

$$= O(L^{d-1/2}m). \quad (F.59)$$

Then, we consider the upper bound of $C_{2\kappa,2}$. Let $p_{j,k,m} = \sum_{i=(j-L)_+}^{(m-1)\wedge j} \psi_{j-i}\mu_W\left(\frac{k-i}{n}\right), m \leq k \leq n-m$. Then, for $m \leq k \leq n-m$, we can write

$$\overline{W}_{k-m+1,m} - \overline{Z}_{k-m+1,m} = \sum_{j=0}^{L+m-1} p_{j,k,m} (e_{k-j} - \zeta_{k-j}^{\circ}).$$

After a careful check on Lemma 2 in (Wu and Zhou, 2011), we have

$$\max_{0 \le l \le L+m-1} \left\| \sum_{j=0}^{l} (e_{k-j} - \zeta_{k-j}) \right\|_{2\kappa}^{2} \le M \sum_{i=1}^{L+m} \left\{ \sum_{j=i}^{\infty} \delta_{2\kappa}(H, j, (-\infty, 1]) \right\}^{2} = O(1),$$
(F.60)

where M is a sufficiently large constant. Following the proof of Corollary 2 in (Wu and Zhou, 2011), under Assumption 5.1, we obtain

$$\|\zeta_i - \zeta_i^{\circ}\|_{2\kappa} = O\{(\log n)^2/n\}.$$
 (F.61)

Observe that

$$\sum_{l=0}^{L+m-1} |p_{l,k,m} - p_{l-1,k,m}| = O(L^d).$$
(F.62)

Then, by the summation-by-parts formula, combining (F.60), (F.61) and (F.62), since $L(\log n)^2/n \to 0$, we have

$$\|\overline{W}_{k-m+1,m} - \overline{Z}_{k-m+1,m}\|_{2\kappa} = \left\|\sum_{j=0}^{L+m-1} p_{j,k,m}(e_{k-j} - \zeta_{k-j}^{\circ})\right\|_{2\kappa} = O(L^d).$$
(F.63)

Since $\sup_{j} \|\bar{\Delta}_{j}^{(d)}\|_{2\kappa} = O(m^{d-1/2}), \ \sup_{j} \|\Delta_{j}^{(d),\circ}\|_{2\kappa} = O(m^{d-1/2}), \ \text{by (F.57), (F.58), (F.59) and (F.63),}$

since $L/m^2 \to 0$, we obtain

$$\|\bar{\Sigma}_d(t) - \Sigma_d^{\circ}(t)\|_{\kappa} = O(L^{d-1/2}m^{d+1/2})$$

Under the conditions $m\tau_n^{3/2} \to \infty$ and $\kappa \ge 4/(1/2 - d)$, (F.56) then follows from Proposition B.1 in (Dette et al., 2019).

Step 4: We shall show that under condition $m\tau_n^{3/2} \to \infty$,

$$\sup_{t \in \mathcal{I}} |\Sigma_d^{\circ}(t) - \mathbb{E}\Sigma_d^{\circ}(t)| = O_{\mathbb{P}} \left\{ m^{2d} (m\tau_n^{3/2})^{-1/2} \right\} = o_{\mathbb{P}}(m^{2d}).$$
(F.64)

Following similar arguments in the proof of Theorem 3.1 (Wu and Shao, 2006), for $k = 0, \ldots, \lfloor \frac{n-m}{2L} \rfloor$, let $D_{k,i} = \Delta_{2kL+i}^{(d),\circ} (\Delta_{2kL+i}^{(d),\circ})^{\top} - \mathbb{E} \{ \Delta_{2kL+i}^{(d),\circ} (\Delta_{2kL+i}^{(d),\circ})^{\top} | \mathcal{F}_{2kL+i-2L}) \}, (i = 0, \ldots, 2L - 1), \text{ and}$

$$E_{h} = \mathbb{E}\{\Delta_{h}^{(d),\circ}(\Delta_{h}^{(d),\circ})^{\top} | \mathcal{F}_{h-2L}\} - \mathbb{E}\{\Delta_{h}^{(d),\circ}(\Delta_{h}^{(d),\circ})^{\top}\}, \quad (h = m, \dots, n-m).$$

Let $D_{k,i} = 0$, if 2kL + i < m or 2kL + i > n - m. Then, we have

$$\Sigma_{d}^{\circ}(t) - \mathbb{E}\Sigma_{d}^{\circ}(t) = \sum_{h=m}^{n-m} \frac{m\omega(t,h)}{2} E_{h} + \sum_{i=0}^{2L-1} \sum_{k=0}^{\lfloor n/(2L) \rfloor} \frac{m\omega(t,2kL+i)}{2} D_{k,i}.$$
 (F.65)

Recall that $\Delta_h^{(d),\circ} = \frac{\overline{Z}_{h-m+1,m} - \overline{Z}_{h+1,m}}{m}$, in which

$$\overline{Z}_{h,m} = \sum_{j=0}^{L} \psi_j \sum_{i=h}^{h+m-1} \mu_W(t_i) \zeta_{i-j}^{\circ} = \sum_{j=0}^{L+m-1} p_{j,h+m-1,m} \zeta_{h+m-1-j}^{\circ},$$

where $p_{j,h+m-1,m} = \sum_{i=(j-L)_+}^{(m-1)\wedge j} \psi_{j-i}\mu_W\left(\frac{h+m-1-i}{n}\right), \{\zeta_j^o\}$ are martingale differences.

Under the geometric measure contraction condition, for j = 0, ..., L, we have

$$\|\mathbb{E}\{(\zeta_{r-j}^{\circ})^{2}|\mathcal{F}_{r-2L}\} - \mathbb{E}\{(\zeta_{r-j}^{\circ})^{2})\}\| = O(\chi^{L}).$$
(F.66)

By Lemma G.2 and elementary calculation, we have

$$\sum_{j=0}^{L+m-1} |p_{j,s,m} p_{j,s,m}^{\top}| = O(m^{2d+1}), \quad \sum_{j=0}^{L-1} |p_{j,s,m} p_{j+m,s,m}^{\top}| = O(m^{2d+1}).$$
(F.67)

Therefore, combining (F.66) and (F.67), we derive

$$||E_h|| = O(m^{2d-1}\chi^L).$$
 (F.68)

By Burkholder's inequality, uniformly for all i = 0, ..., 2L - 1,

$$\left\| \sum_{k=1}^{\lfloor n/(2L) \rfloor} \omega(t, 2kL+i) D_{k,i} \right\|^{2} \leq C \sum_{k=1}^{\lfloor n/(2L) \rfloor} \omega^{2}(t, 2kL+i) \|D_{k,i}\|^{2}$$
$$\leq 2C \sum_{k \in \{r: |2rL+i-nt| \le n\tau_{n}\}} (\|\Delta_{2kL+i}^{(d),\circ}\|_{4}\| (\Delta_{2kL+i}^{(d),\circ})^{\top}\|_{4})^{2} / (n\tau_{n})^{2}$$
$$= O\{(n\tau_{n})^{-1}L^{-1}m^{4d-2}\},$$
(F.69)

where C is a sufficiently large constant. Therefore, since $L/(n\tau_n) = m^{1+\frac{1}{2d+1}}/(n\tau_n^{1/2}) = O\{1/(m\tau_n^{1/2})\},$ by (F.65), (F.68) and (F.69), and Proposition B.1 in (Dette et al., 2019), we have shown (F.64).

Step 5: Recall that $L = Mm^{1+\frac{1}{2d+1}}\tau_n^{1/2}, m \to \infty, m = O(n^{1/3})$. It follows that $m^{1+\frac{1}{d+1}}/L \to \infty, L^2/(mn) = O(m^3\tau_n/n) = o(1)$. We shall show that uniformly for $s \in \mathcal{I}$,

$$m^{-2d} \mathbb{E}\Sigma_d^{\circ}(s) = \kappa_2(d)\mu_W(s)\mu_W^{\top}(s)\sigma_H^2(s) + O(f_n),$$
(F.70)

where $\kappa_2(d) = \Gamma^{-2}(d+1) \int_0^\infty \{t^d - (t-1)_+^d\} \{2t^d - (t-1)_+^d - (t+1)^d\} dt$, and $f_n = m^{-d} + \tau_n^2 + L^{d+1}/m^{d+2} + L^2/(mn) + (L/m^2)^{-\frac{1}{d-2}} = o(1)$. Note that $\overline{Z}_{k-m+1,m} = \sum_{j=0}^{L+m-1} p_{j,k,m} \zeta_{k-j}^\circ$. Recall

$$p_{j,k,m} = \sum_{i=(j-L)_{+}}^{(m-1)\wedge j} \psi_{j-i}\mu_{W}\left(\frac{k-i}{n}\right) = \begin{cases} \sum_{i=0}^{j-1} \psi_{j-i}\mu_{W}\left(\frac{k-i}{n}\right) + \mu_{W}\left(\frac{k-j}{n}\right), & j = 0, \dots, m-1\\ \sum_{i=0}^{m-1} \psi_{j-i}\mu_{W}\left(\frac{k-i}{n}\right), & j = m, \dots, L\\ \sum_{i=j-L}^{m-1} \psi_{j-i}\mu_{W}\left(\frac{k-i}{n}\right) = O(mL^{d-1}), & j = L+1, \dots, L+m. \end{cases}$$

Then, approximate $p_{j,k,m}$ by integrals. When $j = 0, \ldots, m-1$, by the continuity of μ_W and Lemma G.2,

$$m^{-d}\Gamma(d)p_{j,k,m} = d^{-1}\mu_W (k/n) (j/m)^d + O(m^{-d} + m/n).$$

When $j = m, \ldots, L$,

$$m^{-d}\Gamma(d)p_{j,k,m} = d^{-1}\mu_W(k/n)\left\{(j/m)^d - ((j+1)/m - 1)^d\right\} + O\{m/n + m^{-1}(j/m - 1)^{d-1}\}.$$

Since (ζ_i°) are martingale differences, and $\sigma_H(t_j) = \|\sum_{i=j}^{\infty} \mathcal{P}_j H(t_j, \mathcal{F}_i)\| = \|\zeta_j^\circ\|$, (F.70) then follows from elementary calculation.

 $\begin{array}{l} \mathbf{Step } \stackrel{\circ}{\mathbf{6}:} \ \mathrm{Let} \ g_{\kappa,n} = m^{-d} (\log n)^{d} \tau_{n}^{-1/\kappa} + \sqrt{m} \{ (n\tau_{n}^{3/2})^{d-1/2} + \tau_{n}^{3} \} \tau_{n}^{-1/\kappa} + m \{ (n\tau_{n}^{3/2})^{2d-1} + \tau_{n}^{6} \} \tau_{n}^{-1/\kappa} + m^{-\frac{1/2-d}{2d+1}} \tau_{n}^{d/2-1/4-1/\kappa} + (m\tau_{n}^{3/2})^{-1/2} + f_{n}. \end{array}$ Summarizing Step 1-5, we have

$$\sup_{t\in\mathcal{I}} \left| m^{-2d} \hat{\Sigma}_d(t) - \kappa_2(d) \sigma_H^2(t) \mu_W(t) \mu_W^\top(t) \right| = O_{\mathbb{P}}(g_{\kappa,n}) = o_{\mathbb{P}}(1).$$

	-	-	
L			
L			
L			

F.6 Proof of Theorem 5.3

We define the notation under the local alternatives by replacing d with d_n . In the following, Lemma F.4 studies the asymptotic behavior of the bias correction term under the local alternatives. Lemma F.5 investigates the physical dependence of $U^{(d_n)}(t, \mathcal{F}_i)$ as well as the order of its partial sum process under the local alternatives.

Lemma F.4. Under the conditions of Theorem 5.3, we have

$$\sup_{t\in\mathcal{I}} \left| \breve{\Sigma}_{d_n}(t) - \Sigma^A(t) \right| = O_{\mathbb{P}} \left(\sqrt{m\tau_n^{3-1/\kappa}} + \sqrt{\frac{m}{n\tau_n^{3/2+2\kappa}}} \right) = o_{\mathbb{P}}(1).$$

of Lemma F.4. Letting $d_n = c/\log n$, the proof follows from similar steps in Lemma F.3.

Lemma F.5. Under Assumptions 4.3 and 5.1, $m \to \infty$, m = O(n), we have

$$\sup_{1 \le k \le n-m+1} \left\| \sum_{i=k}^{k+m-1} x_i e_i^{(d_n)} \right\|_4 = O(\sqrt{m}).$$

Proof of Lemma F.5. Define $\tilde{Q}_{k,m}^{(d_n)} = \sum_{i=k}^{k+m-1} x_i e_i^{(d_n)}$. By similar arguments in Lemma G.3, we obtain

$$\delta_4(U^{(d_n)}, k) = O\{\psi_k(d_n)\}, \quad k \ge 0,$$
(F.71)

and $\delta_4(U^{(d_n)}, k) = 0$, for k < 0. Then, uniformly for $1 \le k \le n - m + 1$, by Burkholder's inequality we have

$$\left\|\tilde{Q}_{k,m}^{(d_n)}\right\|_{4}^{2} \le B_{4}^{2} \left\|\sum_{l=-\infty}^{\infty} \left|\mathcal{P}_{l}\tilde{Q}_{k,m}^{(d_n)}\right|^{2}\right\| \le B_{4}^{2} \sum_{l=-\infty}^{k+m-1} \left\|\mathcal{P}_{l}\tilde{Q}_{k,m}^{(d_n)}\right\|_{4}^{2} \le B_{4}^{2} \sum_{l=-\infty}^{k+m-1} \left\{\sum_{i=k-l}^{k+m-1-l} \delta_{4}(U^{(d_n)},i)\right\}^{2}, \quad (F.72)$$

where B_4 is a constant. Therefore, combining (F.71) and (F.72), it follows from Lemma G.4 that

$$\max_{1 \le k \le n-m+1} \left\| \tilde{Q}_{k,m}^{(d_n)} \right\|_4^2 = O\left[\sum_{l=-\infty}^k \left\{ (k+m-l)^{d_n} - (k-l+1)^{d_n} \right\}^2 + \sum_{l=k+1}^{k+m-1} (k+m-l)^{2d_n} \right] = O(m).$$

Proof of Theorem 5.3. Recall that

$$\tilde{\Sigma}_{d_n}(t) = \sum_{j=m}^{n-m} \frac{m\tilde{\Delta}_j^{(d_n)}\tilde{\Delta}_j^{(d_n),\top}}{2} \omega(t,j), \quad \tilde{\Delta}_j^{(d_n)} = \frac{\tilde{Q}_{j-m+1,m}^{(d_n)} - \tilde{Q}_{j+1,m}^{(d_n)}}{m}$$

where $\tilde{Q}_{k,m}^{(d_n)} = \sum_{i=k}^{k+m-1} x_i e_i^{(d_n)}$.

We break the proof in the following 8 steps.

Step 1: Following the proof of Theorem 5.2 by replacing d by d_n , since $\kappa \ge 4$, we have

$$\sup_{t\in\mathcal{I}} \left| \hat{\Sigma}_{d_n}(t) - \tilde{\Sigma}_{d_n}(t) \right| = O_{\mathbb{P}}\left(\sqrt{m\tau_n^{3-1/\kappa}} + \sqrt{\frac{m}{n\tau_n^{3/2+2\kappa}}} \right) = o_{\mathbb{P}}(1).$$

Step 2: Let $L = m^2 \tau_n^{1/2}$, $\check{e}_{i,L}^{(d_n)} = \sum_{j=0}^L \psi_j e_{i-j}$. Define

$$\check{Q}_{k,m}^{(d_n)} =: \sum_{i=k}^{k+m-1} x_i \check{e}_{i,L}^{(d_n)}, \quad \check{\Delta}_j^{(d_n)} = \frac{\check{Q}_{j-m+1,m}^{(d_n)} - \check{Q}_{j+1,m}^{(d_n)}}{m}, \quad \check{\Sigma}_{d_n}(t) = \sum_{j=m}^{n-m} \frac{m\check{\Delta}_j^{(d_n)}\check{\Delta}_j^{(d_n),\top}}{2} \omega(t,j).$$

In this step, we shall show that under bandwidth condition $m\tau_n^{3/2} \to \infty$,

$$\sup_{t\in\mathcal{I}} \left| \tilde{\Sigma}_{d_n}(t) - \check{\Sigma}_{d_n}(t) \right| = O_{\mathbb{P}}(m^{-1/2}\tau_n^{-3/4}) = o_{\mathbb{P}}(1).$$
(F.73)

Observe that

$$e_i^{(d_n)} = \check{e}_{i,L}^{(d_n)} + \tilde{e}_{i,L}^{(d_n)}$$
, where $\check{e}_{i,L}^{(d_n)} = \sum_{j=0}^L \psi_j e_{i-j}$, $\tilde{e}_{i,L}^{(d_n)} = \sum_{j=L+1}^\infty \psi_j e_{i-j}$

By Proposition G.1, we have $\|\tilde{e}_{i,L}^{(d_n)}\|_4^2 = O\{\sum_{s=L+1}^{\infty} (s+1)^{2d_n-2}\} = O(L^{-1})$. Then, under Assumption 4.3, we have uniformly for $1 \le k \le n-m+1$,

$$\left\| \check{Q}_{k,m}^{(d_n)} - \check{Q}_{k,m}^{(d_n)} \right\| \le m \max_{1 \le i \le n} \left\| x_i \tilde{e}_{i,L}^{(d_n)} \right\| \le m \max_{1 \le i \le n} \| x_i \|_4 \left\| \check{e}_{i,L}^{(d_n)} \right\|_4 = O(m/\sqrt{L}).$$
(F.74)

By Lemma F.5 and (F.74), we have

$$\left\|\check{\Sigma}_{d_n}(t) - \check{\Sigma}_{d_n}(t)\right\| \le m \max_{m \le j \le n-m} \left\|\check{\Delta}_j^{(d_n)} - \check{\Delta}_j^{(d_n)}\right\|_4 \left(\left\|\check{\Delta}_j^{(d_n)} - \check{\Delta}_j^{(d_n)}\right\|_4 + 2\left\|\check{\Delta}_j^{(d_n)}\right\|_4\right) = O\left(\sqrt{m/L}\right).$$

By Proposition B.1 in (Dette et al., 2019), since $m/(L\tau_n) = m^{-1}\tau_n^{-3/2} \to 0$, (F.73) is proved. **Step 3** : Define $\bar{Q}_{k,m}^{(d_n)} =: \sum_{i=k}^{k+m-1} \{x_i e_i + \mu_W(t_i) (\check{e}_{i,L}^{(d_n)} - e_i)\},$

$$\bar{\Delta}_{j}^{(d_{n})} = \frac{\bar{Q}_{j-m+1,m}^{(d_{n})} - \bar{Q}_{j+1,m}^{(d_{n})}}{m}, \quad \bar{\Sigma}_{d_{n}}(t) = \sum_{j=m}^{n-m} \frac{m\bar{\Delta}_{j}^{(d_{n})}\bar{\Delta}_{j}^{(d_{n}),\top}}{2}\omega(t,j).$$

We shall show that

$$\sup_{t\in\mathcal{I}} \left| \check{\Sigma}_{d_n}(t) - \bar{\Sigma}_{d_n}(t) \right| = O_{\mathbb{P}}\left(\sqrt{\frac{m}{n\tau_n^{1+2\kappa}}} + d_n \right) = o_{\mathbb{P}}(1).$$

Observe that

$$\check{Q}_{k,m}^{(d_n)} - \bar{Q}_{k,m}^{(d_n)} = \sum_{j=1}^{L} \sum_{i=k}^{k+m-1} (x_i - \mu_W(t_i))\psi_j e_{i-j} = \sum_{j=1}^{L} \sum_{i=k}^{k+m-1} \psi_j \bar{x}_i e_{i-j},$$

where $\bar{x}_i = x_i - \mu_W(t_i)$. Let

$$\vartheta_{k,m} = \frac{1}{m} \sum_{j=1}^{L} \sum_{i=k-m+1}^{k} \psi_j (\bar{x}_i e_{i-j} - \bar{x}_{i+m} e_{i+m-j}).$$

Then, it follows that

$$\check{\Sigma}_{d_n}(t) - \bar{\Sigma}_{d_n}(t) = \sum_{j=m}^{n-m} \frac{m\omega(t,j)}{2} (\check{\Delta}_j^{(d_n)}\check{\Delta}_j^{(d_n),\top} - \bar{\Delta}_j^{(d_n)}\bar{\Delta}_j^{(d_n),\top}),$$
(F.75)

and

$$\check{\Delta}_{j}^{(d_{n})}\check{\Delta}_{j}^{(d_{n}),\top} - \bar{\Delta}_{j}^{(d_{n})}\bar{\Delta}_{j}^{(d_{n}),\top} = \vartheta_{k,m}\vartheta_{k,m}^{\top} + \vartheta_{k,m}\bar{\Delta}_{j}^{(d_{n}),\top} + \bar{\Delta}_{j}^{(d_{n})}\vartheta_{k,m}^{\top}.$$
(F.76)

Step 3.1 We first show that

$$\sup_{t \in \mathcal{I}} \left| \mathbb{E}\{ \check{\Sigma}_{d_n}(t) - \bar{\Sigma}_{d_n}(t) \} \right| = O(d_n) = o(1).$$
(F.77)

Observe that $(\bar{x}_i)_{i=1}^n$, $(e_i)_{i=-\infty}^n$ are two centered sequences. Under Assumptions 4.3 and 5.1, by Lemma 7 in (Zhou, 2014a) we have for l, j > 0,

$$|\mathbb{E}(\bar{x}_i e_{i-j} \bar{x}_{i+k}^\top e_{i+k-l})| = O(\chi^{\rho^*}),$$

where following the lines in the proof of Theorem 2 in (Zhou, 2014a), we have

$$\rho^* \ge \frac{1}{2} \min\{\max(|k|, |k-l+j|), \max(|k-l|, |k+j|)\},\$$

where we define the right hand side as $\rho_{k,l,j}$. Then, we are able to bound the expectation of (F.76), for $1 \le p \le L, 1 \le q \le L$,

$$\begin{split} & |\mathbb{E}(\vartheta_{k,m}\vartheta_{k,m}^{\top})| \\ & \leq \frac{1}{m^2} \sum_{p,q=1}^{L} \psi_p \psi_q \left| \mathbb{E}\left[\left\{ \sum_{i=k-m+1}^{k} (\bar{x}_i e_{i-p} - \bar{x}_{i+m} e_{i+m-p}) \right\} \left\{ \sum_{j=k-m+1}^{k} (\bar{x}_j e_{j-q} - \bar{x}_{j+m} e_{j+m-q}) \right\}^{\top} \right] \right| \\ & = O\left(\frac{1}{m^2} \sum_{p,q=1}^{L} \psi_p \psi_q \sum_{i,j=k-m+1}^{k+m} \chi^{\rho_{j-i,q,p}} \right). \end{split}$$

Consider $q \leq p$, since when $q \geq 1$, $\psi_q = O(d_n(1+q)^{d_n-1})$, we have

$$\frac{1}{m^2} \sum_{q=1}^{L} \sum_{p=q}^{L} \psi_p \psi_q \sum_{i,j=k-m+1}^{k+m} \chi^{\rho_{j-i,q,p}} \\
= \frac{1}{m^2} \sum_{q=1}^{L} \sum_{p=q}^{L} \psi_p \psi_q \sum_{i=k-m+1}^{k+m} \left(\sum_{j>i+(q-p)/2} \chi^{(j-i-q+p)/2} + \sum_{j\le i+(q-p)/2} \chi^{(i-j)/2} \right) \\
= O\left(\sum_{q=1}^{L} \psi_q^2/m \right) = O(d_n/m).$$

Similarly,

$$\frac{1}{m^2} \sum_{q=1}^{L} \sum_{p=1}^{q-1} \psi_p \psi_q \sum_{i,j=k-m+1}^{k+m} \chi^{\rho_{j-i,q,p}} = O(d_n/m).$$

Then, we have

$$|\mathbb{E}(\vartheta_{k,m}\vartheta_{k,m}^{\top})| = O(d_n/m).$$
(F.78)

Following similar arguments in Lemma F.5, $\|\bar{\Delta}_{j}^{(d_{n})}\| = O(m^{-1/2})$. Then, it follows that

$$|\mathbb{E}(\bar{\Delta}_{j}^{(d_{n})}\vartheta_{k,m}^{\top})| \le \|\bar{\Delta}_{j}^{(d_{n})}\| \|\vartheta_{k,m}^{\top}\| = O(d_{n}^{1/2}/m).$$
(F.79)

Therefore, by (F.76), (F.75), (F.78), and (F.79), we have (F.77).

Step 3.2 We proceed to show that

$$\sup_{t\in\mathcal{I}} \left|\check{\Sigma}_{d_n}(t) - \bar{\Sigma}_{d_n}(t) - \mathbb{E}(\check{\Sigma}_{d_n}(t) - \bar{\Sigma}_{d_n}(t))\right| = O_{\mathbb{P}}\left(\sqrt{\frac{m}{n\tau_n^{1+2\kappa}}}\right) = o_{\mathbb{P}}(1).$$
(F.80)

Notice that $\check{e}_{i,L}^{(d_n)} - e_i$ has summable physical dependence. Specifically,

$$\begin{aligned} \|\vartheta_{k,m} - \vartheta_{k,m,\{k-s\}}\|_{2\kappa} &= O\left[\frac{1}{m}\sum_{j=1}^{L}\psi_{j}\sum_{i=k-m+1}^{k+m} \{\delta_{4\kappa}(W,i-k+s) + \delta_{4\kappa}(H,i-j-k+s)\}\right] \\ &= O\left[\frac{d_{n}}{m}\sum_{i=-m+1}^{m} \min\left\{\chi^{i-L+s}L^{d_{n}-1},(i+s)^{d_{n}-1}\right\}\mathbf{1}(i+s>0)\right],\end{aligned}$$

where in the last equality, we use the fact $\delta_{4\kappa}(H,k) = 0$, if $k \leq 0$ and $\sum_{j=1}^{L} \psi_j \chi^{L-j} = O(\psi_L)$. From (F.78), we have $\sup_j \|\vartheta_{j,m}\|_{2\kappa} = O(m^{-1/2})$.

For simplicity, write $r_{i,s,n} = \frac{d_n}{m} \min\left\{\chi^{i-L+s}L^{d_n-1}, (i+s)^{d_n-1}\right\} 1(i+s > 0)$. Then, we obtain for

 $m\leq j\leq n,\,s\geq 0,$

$$\begin{aligned} \left\| \mathcal{P}_{j-s} \vartheta_{j,m} \vartheta_{j,m}^{\top} \right\|_{\kappa} &\leq \left\| \vartheta_{j,m} \right\|_{2\kappa} \left\| \vartheta_{j,m}^{\top} - \vartheta_{j,m,\{j-s\}}^{\top} \right\|_{2\kappa} + \left\| \vartheta_{j,m} - \vartheta_{j,m,\{j-s\}} \right\|_{2\kappa} \left\| \vartheta_{j,m,\{j-s\}}^{\top} \right\|_{2\kappa} \\ &= O\left(m^{-1/2} r_{i,s,n} \right). \end{aligned}$$
(F.81)

Under Assumption 4.4, similar to (36) in Lemma 3 in (Zhou and Wu, 2010), we have

$$\delta_{2\kappa}(\tilde{\Delta}(m),k) := \sup_{1 \le j \le n} \|\tilde{\Delta}_j - \tilde{\Delta}_{j,\{j-k\}}\|_{2\kappa} = O\left\{\frac{1}{m} \sum_{i=-m+1}^m \delta_{2\kappa}(U,k+i)\right\} = O\{\min(\chi^{k-m},1)/m\}.$$

Similar to (F.81), and by (F.21), we have

$$\left\|\bar{\Delta}_{k}^{(d_{n})} - \bar{\Delta}_{k,\{k-s\}}^{(d_{n})}\right\|_{2\kappa} \leq \left\|\tilde{\Delta}_{k} - \tilde{\Delta}_{k,\{k-s\}}\right\|_{2\kappa} + \frac{1}{m} \sum_{i=k-m+1}^{k+m} \sum_{j=1}^{L} \psi_{j} \left\|\mu_{W}(t_{i})(e_{i-j} - e_{i-j,\{k-s\}})\right\|_{2\kappa} \\ = O\left\{\min(\chi^{s-m}, 1)/m + \sum_{i=-m+1}^{m} r_{i,s,n}\right\}.$$
(F.82)

Since $\sup_j \|\bar{\Delta}_j^{(d_n)}\|_{\kappa} = O(m^{-1/2})$, by (F.82), similar to (F.81), we obtain

$$\left\| \mathcal{P}_{j-s} \bar{\Delta}_{j}^{(d_{n})} \vartheta_{j,m}^{\top} \right\|_{\kappa} = O\left\{ m^{-1/2} r_{i,s,n} + m^{-3/2} \min(\chi^{s-m}, 1) \right\}.$$
 (F.83)

By Burkholder's inequality, by (F.75) and (F.76), combining (F.81) and (F.83), we have for $t \in \mathcal{I}$,

$$\begin{split} \left\| \check{\Sigma}_{d_n}(t) - \bar{\Sigma}_{d_n}(t) - \mathbb{E}(\check{\Sigma}_{d_n}(t) - \bar{\Sigma}_{d_n}(t)) \right\|_{\kappa} \\ &= O\left\{ \sum_{s=0}^{\infty} \left(\sum_{j=m}^{n-m} \omega^2(t,j) m^2 \left\| \mathcal{P}_{j-s} \vartheta_{j,m} \vartheta_{j,m}^\top + \mathcal{P}_{j-s} \bar{\Delta}_j^{(d_n)} \vartheta_{j,m}^\top + \mathcal{P}_{j-s} \bar{\Delta}_j^{(d_n)} \vartheta_{j,m}^\top \right\|_{\kappa}^2 \right)^{1/2} \right\} \\ &= O\left(\sqrt{\frac{m}{n\tau_n}}\right), \end{split}$$

where in the last equality, we consider 0 < i + s < L, and $i + s \ge L$ separately and use the fact $\sum_{i=1}^{L} i^{-1} = O(\log L)$. Then, (F.80) follows from the chaining argument in Proposition B.1 in (Dette et al., 2019).

Step 4: Decomposition Recall that $\tilde{Q}_{k,m} = \sum_{i=k}^{k+m-1} x_i e_i$,

$$\tilde{\Delta}_j = \frac{\tilde{Q}_{j-m+1,m} - \tilde{Q}_{j+1,m}}{m}, \quad \tilde{\Sigma}(t) = \sum_{j=m}^{n-m} \frac{m\tilde{\Delta}_j \tilde{\Delta}_j^\top}{2} \omega(t,j).$$

Define
$$\check{\Delta}_{j}^{(d_{n})} = \bar{\Delta}_{j}^{(d_{n})} - \tilde{\Delta}_{j} = \frac{1}{m} \sum_{i=j-m+1}^{j} \mu_{W}(t_{i}) (\check{e}_{i,L}^{(d_{n})} - e_{i}) - \mu_{W}(t_{i+m}) (\check{e}_{i+m,L}^{(d_{n})} - e_{i+m}).$$
 Let
 $\tilde{s}_{1}(t) = \sum_{j=m}^{n-m} \frac{m\omega(t,j)}{2} \check{\Delta}_{j}^{(d_{n})} \check{\Delta}_{j}^{(d_{n}),\top}, \quad \tilde{s}_{2}(t) = \sum_{j=m}^{n-m} \frac{m\omega(t,j)}{2} \check{\Delta}_{j} \check{\Delta}_{j}^{(d_{n}),\top}.$

Observe that

$$\bar{\Sigma}_{d_n}(t) = \tilde{\Sigma}(t) + \tilde{s}_1(t) + \tilde{s}_2(t) + \tilde{s}_2^{\top}(t).$$

Step 5: Martingale approximation

Let

$$z_j = \sum_{i=j}^{\infty} \mathcal{P}_j(x_i e_i), \quad z_j^\circ = z_j(t_j) = \sum_{i=j}^{\infty} \mathcal{P}_j U(t_j, \mathcal{F}_i).$$

Recall that in Theorem 5.2, $\zeta_j = \sum_{i=j}^{\infty} \mathcal{P}_j e_i$, $\zeta_j^\circ = \zeta_j(t_j) = \sum_{i=j}^{\infty} \mathcal{P}_j H(t_j, \mathcal{F}_i)$. Let $z_{j,1}$ denote the first element in z_i . Then, it follows that $z_{j,1} = \zeta_j$, $z_{j,1}^\circ = \zeta_j^\circ$. Define $\overline{Z}_{k,m}^{(d_n)} = \sum_{i=k}^{k+m-1} \left\{ z_i^\circ + \sum_{j=1}^L \psi_j \mu_W(t_i) \zeta_{i-j}^\circ \right\}$,

$$\Delta_{j}^{(d_{n}),\circ} = \frac{\overline{Z}_{j-m+1,m}^{(d_{n})} - \overline{Z}_{j+1,m}^{(d_{n})}}{m}, \quad \Sigma_{d_{n}}^{\circ}(t) = \sum_{j=m}^{n-m} \frac{m \Delta_{j}^{(d_{n}),\circ} (\Delta_{j}^{(d_{n}),\circ})^{\top}}{2} \omega(t,j).$$

Similarly to $p_{j,k,m}$ defined in Step 3 of Theorem 5.2, we define $\underline{p}_{j,k,m} = \sum_{i=(j-L)_+}^{(m \wedge j)-1} \psi_{j-i} \mu_W\left(\frac{k-i}{n}\right)$. By (F.63) and similar arguments in Theorem 1(ii) of (Wu, 2007), we have uniformly for $1 \le k \le n - m + 1$,

$$\begin{split} \|\bar{Q}_{k,m}^{(d_n)} - \overline{Z}_{k,m}^{(d_n)}\|_4 &\leq \left\| \sum_{j=1}^{L+m-1} \underline{p}_{j,k+m-1,m} (e_{k+m-1-j} - \zeta_{k+m-1-j}^{\circ}) \right\|_4 \\ &+ \left\| \sum_{i=k}^{k+m-1} x_i e_i - \sum_{i=k}^{k+m-1} z_i^{\circ} \right\|_4 = O(1). \end{split}$$

Since $\sup_j \|\bar{\Delta}_j^{(d_n)}\| = O(m^{-1/2})$, by triangle inequality and Cauchy–Schwarz inequality, we have for $t \in \mathcal{I}$,

$$\|\bar{\Sigma}_{d_n}(t) - \Sigma_{d_n}^{\circ}(t)\| \le \sum_{j=m}^{n-m} \frac{m\omega(t,j)}{2} \left\|\bar{\Delta}_j^{(d_n)}(\bar{\Delta}_j^{(d_n)})^{\top} - \Delta_j^{(d_n),\circ}(\Delta_j^{(d_n),\circ})^{\top}\right\| = O(m^{-1/2}).$$

By chaining argument in Proposition B.1 in (Dette et al., 2019), we have

$$\sup_{t \in \mathcal{I}} |\bar{\Sigma}_{d_n}(t) - \Sigma_{d_n}^{\circ}(t)| = O_{\mathbb{P}}\{(m\tau_n)^{-1/2}\} = o_{\mathbb{P}}(1).$$

Step 6 Observe that

$$\overline{Z}_{k,m}^{(d_n)} = \sum_{i=k}^{k+m-1} \left\{ z_i^{\circ} + \sum_{j=1}^{L} \psi_j \mu_W(t_i) \zeta_{i-j}^{\circ} \right\} = \sum_{i=k}^{k+m-1} z_i^{\circ} + \sum_{j=1}^{L+m-1} \underline{p}_{j,k+m-1,m} \zeta_{k+m-1-j}^{\circ}$$

After a careful inspection of Step 4 of Theorem 5.2, we have

$$\sup_{t \in \mathcal{I}} |\Sigma_{d_n}^{\circ}(t) - \mathbb{E}\Sigma_{d_n}^{\circ}(t)| = O_{\mathbb{P}}\left(\sqrt{\frac{1}{m\tau_n^{3/2}}}\right) = o_{\mathbb{P}}(1)$$

Step 7 Recall that $\check{\Sigma}(t) = \Sigma(t) + (e^{c\alpha_1} - 1)^2 \sigma_H^2(t) \mu_W(t) \mu_W^{\top}(t) + (e^{c\alpha_1} - 1) s_{UH}(t) \mu_W^{\top}(t) + (e^{c\alpha_1} - 1) \mu_W(t) s_{UH}^{\top}(t)$. We shall show that uniformly for $t \in \mathcal{I}$,

$$\mathbb{E}\Sigma_{d_n}^{\circ}(t) = \check{\Sigma}(t) + O\{(\log n)^{-1}\}.$$

Let $\check{\Delta}_{k}^{(d_{n}),\circ} = \frac{1}{m} \sum_{i=k-m+1}^{k} \sum_{j=1}^{L} \psi_{j} \left\{ \mu_{W}(t_{i})\zeta_{i-j}^{\circ} - \mu_{W}(t_{i+m})\zeta_{i+m-j}^{\circ} \right\}, \ \Delta_{k}^{\circ} = \frac{1}{m} \sum_{i=k-m+1}^{k} (z_{i}^{\circ} - z_{i+m}^{\circ}).$ Define for $t \in [0,1], \ \tilde{\Sigma}^{\circ}(t) = \sum_{j=m}^{n-m} \frac{m\omega(t,j)}{2} \Delta_{j}^{\circ} \Delta_{j}^{\circ,\top},$

$$\tilde{s}_1^{\circ}(t) = \sum_{j=m}^{n-m} \frac{m\omega(t,j)}{2} \breve{\Delta}_j^{(d_n),\circ} (\breve{\Delta}_j^{(d_n),\circ})^{\top}, \text{ and } \tilde{s}_2^{\circ}(t) = \sum_{j=m}^{n-m} \frac{m\omega(t,j)}{2} \Delta_j^{\circ} (\breve{\Delta}_j^{(d_n),\circ})^{\top}.$$

Then it follows that

$$\mathbb{E}\{\Sigma_{d_n}^{\circ}(t)\} = \mathbb{E}\{\tilde{\Sigma}^{\circ}(t)\} + \mathbb{E}\{\tilde{s}_1^{\circ}(t)\} + \mathbb{E}\{\tilde{s}_2^{\circ}(t)\} + \mathbb{E}\{\tilde{s}_2^{\circ,\top}(t)\}.$$

Following similar arguments in Step 5 of Theorem 5.2, by the continuity of μ_W , we have when $j \leq m$,

$$\underline{p}_{j,k,m} = (e^{c\alpha_1} - 1)\mu_W(k/n) + O(m/n + d_n).$$
(F.84)

when $j \ge m+1$, since $L/n \to 0$,

$$\underline{p}_{j,k,m} = \mu_W \left(k/n \right) \left\{ j^{d_n} - (j - m + 1)^{d_n} \right\} + O(m/n + d_n + m/L).$$
(F.85)

Since z_i° are martingale differences, and $\sigma_H(t_j) = \|\sum_{i=j}^{\infty} \mathcal{P}_j H(t_j, \mathcal{F}_i)\| = \|\zeta_j^{\circ}\|$,

$$\mathbb{E}(\overline{Z}_{k-m+1,m}^{(d_n),\top}\overline{Z}_{k-m+1,m}^{(d_n),\top}) = \sum_{j=1}^{L+m-1} \underline{p}_{j,k,m} \underline{p}_{j,k,m}^{\top} \mathbb{E}\{(\zeta_{k-j}^{\circ})^2\} + \sum_{i=k-m+1}^{k} \mathbb{E}(z_i^{\circ} z_i^{\circ,\top}) + \sum_{j=1}^{m-1} \underline{p}_{j,k,m} \mathbb{E}(\zeta_{k-j}^{\circ} z_{k-j}^{\circ,\top}) + \sum_{j=1}^{m-1} \mathbb{E}(z_{k-j}^{\circ} \zeta_{k-j}^{\circ}) \underline{p}_{j,k,m}^{\top} = Z_1 + Z_2 + Z_3 + Z_4.$$
(F.86)

By Lemma G.4, we have

$$Z_1/m = \frac{1}{m} \sum_{j=1}^{m-1} \underline{p}_{j,k,m} \underline{p}_{j,k,m}^\top \mathbb{E}\{(\zeta_{k-j}^\circ)^2\} + \frac{1}{m} \sum_{j=m}^{L+m-1} \underline{p}_{j,k,m} \underline{p}_{j,k,m}^\top \mathbb{E}\{(\zeta_{k-j}^\circ)^2\}$$
$$= (e^{c\alpha_1} - 1)^2 \mu_W(k/n) \mu_W^\top(k/n) \sigma_H^2(k/n) + O\{m/n + d_n + (\log m)^{-1}\}.$$

Under Assumption 4.2, we have

$$Z_2/m = \Sigma(t) + O(m/n).$$

Observe that

$$\mathbb{E}(z_j^{\circ}\zeta_j^{\circ}) = \sum_{k=-\infty}^{\infty} \operatorname{Cov}\{U(t_j, \mathcal{F}_0), H(t_j, \mathcal{F}_k)\} = s_{UH}(t_j).$$

Under Assumption 5.2, similar arguments in the calculation of Z_1 and Z_2 imply,

$$Z_3/m = (e^{c\alpha_1} - 1)\mu_W(k/n)s_{UH}^{\top}(k/n) + O(m/n + d_n),$$

and

$$Z_4/m = (e^{c\alpha_1} - 1)s_{UH}(k/n)\mu_W^\top(k/n) + O(m/n + d_n).$$

By Lemma G.4 (a), (F.84) and (F.85), similar techniques of (F.86) show that

$$\mathbb{E}(\overline{Z}_{k+1,m}^{(d_n),\top}\overline{Z}_{k-m+1,m}^{(d_n),\top}) = \sum_{j=1}^{L-1} \underline{p}_{j+m,k+m,m} \underline{p}_{j,k,m}^{\top} \mathbb{E}\{(\zeta_{k-j}^{\circ})^2\} + \sum_{j=1}^{m} \underline{p}_{j+m-1,k+m,m} \mathbb{E}(\zeta_{k+1-j}^{\circ} z_{k+1-j}^{\circ,\top}) \\ = O\{m(\log m)^{-1}\}.$$
(F.87)

Therefore, by (F.86) and (F.87), we have

$$\mathbb{E}\Sigma_{d_n}^{\circ}(t) = \sum_{j=m}^{n-m} \check{\Sigma}(t_j)\omega(t,j) + O\{m/n + d_n + (\log m)^{-1}\} = \check{\Sigma}(t) + O\{(\log n)^{-1}\}.$$

Step 8 Summarizing Step 1 - Step 7, we have

$$\sup_{t \in \mathcal{I}} |\hat{\Sigma}_{d_n}(t) - \check{\Sigma}(t)| = O_{\mathbb{P}} \left\{ \sqrt{\frac{m}{n\tau_n^{3/2 + 2\kappa}}} + \sqrt{m\tau_n^{3-1/\kappa}} + \sqrt{\frac{1}{m\tau_n^{3/2}}} + (\log n)^{-1} \right\} = o_{\mathbb{P}}(1).$$

G Auxiliary results

Proposition G.1. Suppose $Q_i = L(t_i, \mathcal{F}_i), t_i \in \mathcal{I}$, for $q \ge 1$, we have

$$\|\mathcal{P}_{i-l}Q_i\|_q \le \delta_q(L,l,\mathcal{I}).$$

Proof of Proposition G.1. The proposition follows after a careful investigation of Theorem 1 in (Wu, 2005). \Box

Proposition G.2. Under Assumption 5.1, we have uniformly for $l \ge 0$, 0 < d < 1/2,

$$\delta_p(H^{(d)}, l, (-\infty, 1]) = O\{(1+l)^{d-1}\}.$$

Proof of Proposition G.2. Under Assumption 5.1, by Lemma 3.2 of (Kokoszka and Taqqu, 1995) and Proposition G.1, we have

$$\delta_p(H^{(d)}, l, (-\infty, 1]) \le \sum_{k=0}^l \psi_k(d) \delta_p(H, l-k, (-\infty, 1]) = O\{(1+l)^{d-1}\}.$$

Lemma G.1. Suppose $\|\sup_{t \in [0,1]} |\hat{\Sigma}(t) - \Sigma(t)|\| = O(s_n)$, where $\Sigma(t)$ is a covariance matrix with its eigenvalues bounded from zero, dim $(\Sigma(t)) = p < \infty$. Then, we have

$$\sup_{t \in [0,1]} \left| \hat{\Sigma}^{1/2}(t) - \Sigma^{1/2}(t) \right| = O_{\mathbb{P}}(s_n^{1/2}).$$

Proof of Lemma G.1. Without loss of generality, suppose $\Sigma(t)$ has eigenvalues $\lambda_1(t) \geq \cdots \geq \lambda_p(t)$, and eigenvector matrix $V(t) = (v_1(t), \dots, v_p(t))$, $\Sigma(t)v_j(t) = \lambda_j(t)v_j(t)$, $\Lambda(t) = \text{diag}\{\lambda_1(t) \geq \cdots \geq \lambda_p(t)\}$. Suppose $\hat{\Sigma}(t)$ has eigenvalues $\hat{\lambda}_1(t) \geq \cdots \geq \hat{\lambda}_p(t)$, and eigenvector matrix $\hat{V}(t) = (\hat{v}_1(t), \dots, \hat{v}_p(t))$, $\hat{\Sigma}(t)\hat{v}_j(t) = \hat{\lambda}_j(t)\hat{v}_j(t)$, $\hat{\Lambda}(t) = \text{diag}\{\hat{\lambda}_1(t) \geq \cdots \geq \hat{\lambda}_p(t)\}$. Suppose $\Sigma(t)$ has q distinct eigenvalues, $\hat{\lambda}_1(t) > \cdots > \hat{\lambda}_q(t)$. Let $Q(t) = \{k : \exists j \neq i, \lambda_j(t) = \lambda_i(t) = \hat{\lambda}_k(t)\}$. Let

$$\Sigma^{\circ}(t) = \hat{V}(t)\Lambda(t)\hat{V}(t)^{\top}.$$

Then, we have

$$\mathbb{E} \sup_{t \in [0,1]} \left| \hat{\Sigma}^{1/2}(t) - \Sigma^{1/2}(t) \right| \leq \mathbb{E} \sup_{t \in [0,1]} \left| \hat{\Sigma}^{1/2}(t) - (\Sigma^{\circ})^{1/2}(t) \right| + \mathbb{E} \sup_{t \in [0,1]} \left| (\Sigma^{\circ})^{1/2}(t) - \Sigma^{1/2}(t) \right| \\
= S_1 + S_2,$$
(G.1)

where S_1 and S_2 are defined in the obvious way. $\hat{V}(t)$ is orthogonal, and $|\cdot|$ is the Frobenius norm, then

$$S_{1} = \mathbb{E} \sup_{t \in [0,1]} |\hat{V}(t) \{ \Lambda^{1/2}(t) - \hat{\Lambda}^{1/2}(t) \} \hat{V}(t)^{\top} \|$$

$$\leq \left\| \sup_{t \in [0,1]} \left| \Lambda^{1/2}(t) - \hat{\Lambda}^{1/2}(t) \right| \right\| \left\| \sup_{t \in [0,1]} \left| \hat{V}(t) \right| \right\| = O(s_{n}^{1/2}).$$
(G.2)

By Corollary 1 in (Yu et al., 2015), if $k \notin Q(t)$, suppose $\lambda_i(t) = \tilde{\lambda}_k(t)$. Then, we have

$$|\hat{v}_{i}(t) - v_{i}(t)| \leq \frac{2^{3/2} \rho \left\{ \hat{\Sigma}(t) - \Sigma(t) \right\}}{\min\{\lambda_{i-1}(t) - \lambda_{i}(t), \lambda_{i}(t) - \lambda_{i+1}(t)\}}.$$
(G.3)

If $j \in Q(t)$, suppose $\lambda_{r-1}(t) > \lambda_r(t) = \cdots = \tilde{\lambda}_j(t) = \cdots = \lambda_s(t) > \lambda_{s+1}(t)$, and let $V_j(t) = (v_r(t), \dots, v_s(t))$. Let $\hat{V}_j(t) = (\hat{v}_r(t), \dots, \hat{v}_s(t))$. By Theorem 2 in (Yu et al., 2015), $\exists \hat{O}_j(t) \in \mathbb{R}^{(s-r+1)\times(s-r+1)}$ which is orthogonal, s.t.

$$\left| \hat{V}_{j}(t)\hat{O}_{j}(t) - V_{j}(t) \right| \leq \frac{2^{3/2} \min\left\{ \left(s - r + 1 \right)^{1/2} \rho \left| \hat{\Sigma}(t) - \Sigma(t) \right|, \left| \hat{\Sigma}(t) - \Sigma(t) \right| \right\}}{\min\left(\lambda_{r-1}(t) - \lambda_{r}(t), \lambda_{s}(t) - \lambda_{s+1}(t) \right)}.$$
 (G.4)

Without loss of generality, suppose $\lambda_1(t) > \cdots > \lambda_s(t) > \lambda_{s+1}(t) = \cdots + \lambda_{s+n_{s+1}}(t) > \lambda_{s+n_{s+1}+1}(t) = \cdots = \lambda_{s+n_{s+1}+n_{s+2}}(t) > \cdots > \lambda_{s+\sum_{i=s+1}^{q-1} n_i+1}(t) = \cdots = \lambda_p(t)$, where n_i is algebraic multiplicity of $\tilde{\lambda}_i$, and $\sum_{i=s+1}^{q} n_i = p - s$. Let

$$\hat{O}(t) = \begin{pmatrix} I_s & & & \\ & \hat{O}_{s+1}(t) & & \\ & & & \ddots & \\ & & & & \hat{O}_q(t) \end{pmatrix},$$

where $\hat{O}_q(t) \in \mathbb{R}^{n_q \times n_q}$. From (G.3) and (G.4), we have

$$\left| \hat{V}(t)\hat{O}(t) - V(t) \right| \le \frac{2^{3/2} p^{3/2} \left| \hat{\Sigma}(t) - \Sigma(t) \right|}{\min_{1 \le s \le q+1} \left\{ \tilde{\lambda}_{s-1}(t) - \tilde{\lambda}_s(t) \right\}},\tag{G.5}$$

where $\lambda_0(t) = \infty$, $\lambda_{q+1}(t) = -\infty$. On the other hand,

$$\hat{V}(t)\hat{O}(t)\Lambda^{1/2}(t)\hat{O}^{\top}(t)\hat{V}^{\top}(t) = \hat{V}(t)\Lambda^{1/2}(t)\hat{V}^{\top}(t).$$
(G.6)

Therefore, by (G.5) and (G.6), we have

$$S_{2} \leq \mathbb{E} \left(\sup_{[0,1]} \left| \hat{V}(t) \Lambda^{1/2}(t) \hat{V}^{\top}(t) - V(t) \Lambda^{1/2}(t) V^{\top}(t) \right| \right)$$

$$\leq \mathbb{E} \left(\sup_{[0,1]} \left| \{ \hat{V}(t) \hat{O}(t) - V(t) \} \Lambda^{1/2}(t) \hat{O}^{\top}(t) \hat{V}^{\top}(t) \right| \right) + \mathbb{E} \left(\sup_{[0,1]} \left| V(t) \Lambda^{1/2}(t) \{ \hat{O}^{\top}(t) \hat{V}^{\top}(t) - V^{\top}(t) \} \right| \right)$$

$$\leq C \left\| \sup_{[0,1]} \left| \hat{\Sigma}(t) - \Sigma(t) \right| \right\| = O(s_{n}),$$
(G.7)

where C is a sufficiently large positive constant. Combining (G.1), (G.2) and (G.7), we have

$$\sup_{t \in [0,1]} \left| \hat{\Sigma}^{1/2}(t) - \Sigma^{1/2}(t) \right| = O_{\mathbb{P}}(s_n^{1/2}).$$

Lemma G.2. The following argument shows the properties of long memory coefficient $\psi_j = \psi_j(d)$. $\psi_0 = 1$, and for $j \ge 1$,

$$\psi_j = j^{d-1} l_d(j),$$

where $l_d(j) = 1/\Gamma(d)\{1 + O(1/j)\}.$

Proof of Lemma G.2. By Stirling's formula,

$$\frac{\Gamma(j+d)}{\Gamma(j+1)} = \frac{\sqrt{\frac{2\pi}{j+d}}(\frac{j+d}{e})^{j+d}\{1+O(\frac{1}{j+d})\}}{\sqrt{\frac{2\pi}{j+1}}(\frac{j+1}{e})^{j+1}\{1+O(\frac{1}{j+1})\}} = \frac{j^{d-1}(1+d/j)^j(1+d/j)^{d-1/2}\{1+O(1/j)\}}{e^{d-1}(1+1/j)^j(1+1/j)^{1/2}\{1+O(1/j)\}} = j^{d-1} + O(j^{d-2}).$$

Since $\ln \Gamma(z) \sim z \ln z - z + \frac{1}{2} \ln \frac{2\pi}{z} + \sum_{n=1}^{N-1} \frac{B_{2n}}{2n(2n-1)z^{2n-1}}$, the constant in the big O of $\Gamma(z) = \sqrt{\frac{2\pi}{z}} \left(\frac{z}{e}\right)^z \left\{1 + O\left(\frac{1}{z}\right)\right\}$ is always $B_2/2 = 1/12$.

Lemma G.3. Assuming that $\sup_{t \in (-\infty,1]} \|H(t,\mathcal{F}_0)\|_{2p} < \infty$, $\delta_{2p}(H,k,(-\infty,1]) = O(\chi^k)$, $\delta_{2p}(W,k) = O(\chi^k)$, $\chi \in (0,1)$, $\sup_{t \in [0,1]} \|W(t,\mathcal{F}_0)\|_{2p} < \infty$, we have

$$\delta_p(U^{(d)}, k) = O(k^{d-1}).$$

Proof of Lemma G.3. Note that for $j \leq i$,

$$\delta_p(U^{(d)}, i-j) \le \|W(t_i, \mathcal{F}_i)\|_{2p} \delta_{2p}(H^{(d)}, i-j) + \|H^{(d)}(t_i, \mathcal{F}^*_{i-j})\|_{2p} \delta_{2p}(W, i-j).$$
(G.8)

By Burkholder's inequality and Proposition G.2, we have

$$\|H^{(d)}(t_i, \mathcal{F}_i)\|_{2p}^2 \le M \left\| \sum_{j \in \mathbb{Z}} \left\{ \mathcal{P}_j H^{(d)}(t_i, \mathcal{F}_i) \right\}^2 \right\|_p \le M \sum_{j \in \mathbb{Z}} \left\| \mathcal{P}_j H^{(d)}(t_i, \mathcal{F}_i) \right\|_{2p}^2 = O(1),$$

where M is a sufficiently large constant.

Then by Proposition G.2 and Equation (G.8), we have proved the desired result.

Lemma G.4. Let ψ_j denote $\psi_j(d_n)$. (a) For $h = o(n), h \to \infty$ we have

$$\sum_{l=0}^{\infty} \left(\sum_{j=l}^{h+l} \psi_j \right)^2 \sim \sum_{l=0}^{\infty} \{ (h+l)^{d_n} - l^{d_n} \}^2 = O\{h/(\log h)\}.$$

(b) If we further assume $h = \lfloor n^{\alpha_1} \rfloor$, $\alpha_1 \in (0, 1)$, we have

$$h^{-1} \sum_{l=0}^{h-1} \left(\sum_{j=0}^{l} \psi_j \right)^2 \to e^{2c\alpha_1}.$$

Remark G.1. These two results correspond to the conclusions in Lemma 2 of (Shao and Wu, 2007).

Proof of Lemma G.4. Proof of (a). We first show

$$\sum_{l=0}^{\infty} \{ (h+l)^{d_n} - l^{d_n} \}^2 = O\{ h/(\log h) \}.$$

Let $N_1 = \lfloor h^{1-\alpha_h} \rfloor$, $N_2 = \lfloor h^{1+\alpha_h} \rfloor$, $\alpha_h = (\log h)^{-1} \log \log h$. Then, $h/N_1 = O(h^{\alpha_h}) = O(\log h)$. $N_1^{d_n} = O(1)$, $N_2^{d_n} = O(1)$. By Lemma G.2 and Taylor's expansion, we have

$$\sum_{l=0}^{\infty} \{(h+l)^{d_n} - l^{d_n}\}^2 = \sum_{l=0}^{N_1} \{(h+l)^{d_n} - l^{d_n}\}^2 + \sum_{l=N_1+1}^{N_2} \{(h+l)^{d_n} - l^{d_n}\}^2 + \sum_{l=N_2+1}^{\infty} \{(h+l)^{d_n} - l^{d_n}\}^2 = O\{h(\log h)^{-1}\}.$$

Then by Lemma G.2, we have

$$\sum_{l=0}^{\infty} \left(\sum_{j=l}^{h+l} \psi_j \right)^2 = \sum_{l=0}^{\infty} \{ (h+l)^{d_n} - l^{d_n} \}^2 + O\{ d_n h (\log h)^{-1} \}$$

Proof of (b). By Lemma G.2, we have

$$\sum_{l=0}^{h-1} \left(\sum_{j=0}^{l} \psi_j \right)^2 = 1 + \sum_{l=1}^{h-1} l^{2d_n} + O\left(d_n \sum_{l=1}^{h-1} l^{2d_n} \right), \tag{G.9}$$

where for h = O(n),

$$\sum_{l=1}^{h} l^{2d_n} = e^{2c\alpha_1} \sum_{l=1}^{h} (l/h)^{2d_n} = e^{2c\alpha_1} h \int_0^1 t^{2d_n} dt + O(1)$$
$$= e^{2c\alpha_1} h/(2d_n + 1) + O(1) = e^{2c\alpha_1} h + O(1).$$
(G.10)

Combining (G.9) and (G.10), we have shown the desired result.

References

- Andrews, D. (1991). Heteroskedasticity and autocorrelation consistent covariant matrix estimation. *Econo*metrica, 59(3):817–858.
- Aue, A. and Horváth, L. (2013). Structural breaks in time series. J. Time Ser. Anal., 34(1):1–16.
- Bai, L. and Wu, W. (2024). Detecting long-range dependence for time-varying linear models. *Bernoulli*, page to appear.
- Beran, J., Feng, Y., Ghosh, S., and Kulik, R. (2013). Long-Memory Processes. Springer.
- Brown, L. D. and Levine, M. (2007). Variance estimation in nonparametric regression via the difference sequence method. *Ann. Statist.*, 35(5):2219 2232.
- Cavaliere, G., Nielsen, M. Ø., and Taylor, A. R. (2017). Quasi-maximum likelihood estimation and bootstrap inference in fractional time series models with heteroskedasticity of unknown form. J. Econometrics, 198(1):165–188.
- Chan, K. W. (2022). Optimal difference-based variance estimators in time series: A general framework. Ann. Statist., 50(3):1376–1400.
- Chan, S.-C. and Zhang, Z. (2010). Local polynomial modeling and variable bandwidth selection for time-varying linear systems. *IEEE Trans. Instrum. Meas*, 60(3):1102–1117.
- Dahlhaus, R. (1997). Fitting time series models to nonstationary processes. Ann. Statist., 25(1):1–37.
- Dahlhaus, R., Richter, S., and Wu, W. B. (2019). Towards a general theory for nonlinear locally stationary processes. *Bernoulli*, 25(2):1013 1044.
- Dette, H. and Wu, W. (2019). Detecting relevant changes in the mean of nonstationary processes—a mass excess approach. Ann. Statist., 47(6):3578–3608.
- Dette, H. and Wu, W. (2020). Prediction in locally stationary time series. J. Bus. Econom. Statist., pages 1–12.

- Dette, H., Wu, W., and Zhou, Z. (2019). Change point analysis of correlation in non-stationary time series. *Statist. Sinica*, 29(2):611–643.
- Hall, P., Kay, J. W., and Titterington, D. M. (1990). Asymptotically optimal difference-based estimation of variance in nonparametric regression. *Biometrika*, 77(3):521–528.
- He, X. and Zhu, L.-X. (2003). A lack-of-fit test for quantile regression. J. Amer. Statist. Assoc., 98(464):1013–1022.
- Hirukawa, M. (2021). Robust covariance matrix estimation in time series: A review. Econ. Stat.
- Jansson, M. (2002). Consistent covariance matrix estimation for linear processes. *Econom. Theory*, 18(6):1449–1459.
- Juhl, T. and Xiao, Z. (2009). Tests for changing mean with monotonic power. J. Econometrics, 148(1):14–24.
- Kao, C., Trapani, L., and Urga, G. (2018). Testing for instability in covariance structures. *Bernoulli*, 24(1):740 771.
- Kejriwal, M. (2009). Tests for a mean shift with good size and monotonic power. *Economics Letters*, 102(2):78–82.
- Kokoszka, P. S. and Taqqu, M. S. (1995). Fractional arima with stable innovations. *Stoch. Process. their* Appl., 60(1):19 47.
- Kulik, R. and Wichelhaus, C. (2012). Conditional variance estimation in regression models with long memory. J. Time Ser. Anal., 33(3):468–483.
- Li, D., Chen, J., and Gao, J. (2011). Non-parametric time-varying coefficient panel data models with fixed effects. *Econom. J.*, 14(3):387–408.
- Müller, H.-G. and Stadtmuller, U. (1987). Estimation of heteroscedasticity in regression analysis. Ann. Statist., 15(2):610–625.
- Nason, G. P., Von Sachs, R., and Kroisandt, G. (2000). Wavelet processes and adaptive estimation of the evolutionary wavelet spectrum. J. R. Stat. Soc. Ser. B. Stat. Methodol., 62(2):271–292.
- Newey, W. K. and West, K. D. (1987). A simple, positive semi-definite, heteroskedasticity and autocorrelation consistent covariance matrix. *Econometrica*, 55(3):703–708.
- Politis, D. N. (2011). Higher-order accurate, positive semidefinite estimation of large-sample covariance and spectral density matrices. *Econom. Theory*, 27(4):703–744.
- Politis, D. N., Romano, J. P., and Wolf, M. (1999). Subsampling. Springer Science & Business Media.

- Rothman, A. J. (2012). Positive definite estimators of large covariance matrices. *Biometrika*, 99(3):733–740.
- Shao, X. and Wu, W. B. (2007). Local asymptotic powers of nonparametric and semiparametric tests for fractional integration. *Stoch. Process. their Appl.*, 117(2):251–261.
- Su, L., Wang, X., and Jin, S. (2019). Sieve estimation of time-varying panel data models with latent structures. J. Bus. Econom. Statist., 37(2):334–349.
- Tecuapetla-Gómez, I. and Munk, A. (2017). Autocovariance estimation in regression with a discontinuous signal and m-dependent errors: A difference-based approach. *Scand. J. Stat.*, 44(2):346–368.
- Vogt, M. and Dette, H. (2015). Detecting gradual changes in locally stationary processes. Ann. Statist., 43(2):713–740.
- Wu, W. and Zhou, Z. (2018). Gradient-based structural change detection for nonstationary time series M-estimation. Ann. Statist., 46(3):1197 – 1224.
- Wu, W. B. (2005). Nonlinear system theory: Another look at dependence. Proc. Natl. Acad. Sci. U.S.A, 102(40):14150–14154.
- Wu, W. B. (2007). Strong invariance principles for dependent random variables. Ann. Probab., 35(6):2294–2320.
- Wu, W. B. and Shao, X. (2006). Invariance principles for fractionally integrated nonlinear processes. In Recent developments in nonparametric inference and probability, pages 20–30. Institute of Mathematical Statistics.
- Wu, W. B. and Zhou, Z. (2011). Gaussian approximations for non-stationary multiple time series. Statist. Sinica, 21(3):1397–1413.
- Yu, Y., Wang, T., and Samworth, R. J. (2015). A useful variant of the davis–kahan theorem for statisticians. *Biometrika*, 102(2):315–323.
- Zhou, Y., Cheng, Y., Wang, L., and Tong, T. (2015). Optimal difference-based variance estimation in heteroscedastic nonparametric regression. *Statist. Sinica*, pages 1377–1397.
- Zhou, Z. (2014a). Inference of weighted V-statistics for nonstationary time series and its applications. Ann. Statist., 42(1):87 – 114.
- Zhou, Z. (2014b). Nonparametric specification for non-stationary time series regression. *Bernoulli*, 20(1):78 108.
- Zhou, Z. and Shao, X. (2013). Inference for linear models with dependent errors. J. R. Stat. Soc. Ser. B. Stat. Methodol., 75(2):323–343.
- Zhou, Z. and Wu, W. B. (2010). Simultaneous inference of linear models with time varying coefficients. J. R. Stat. Soc. Ser. B. Stat. Methodol., 72(4):513–531.