

Principal Geodesic Analysis applied to climate time series

Nozomi Sugiura

Reserch Insitute for Global Change, JAMSTEC, Japan

29 March 2023

Abstract

Principal Geodesic Analysis (PGA) is applied to a climate time series. First, we transform each multidimensional sequence into the path signature. Since the signature lives in a curved space, usual principal component analysis (PCA) is not applicable. Instead, we treat the signature space as a geodesic manifold. By replacing the notion of straight lines with that of geodesics, the domain of PCA can be extended to the curved space. Then, the first principal component is derived as the geodesic that minimizes the unexplained variance in the data. As an application, we computed the leading modes for one-year NINO SST time series, which are divided into segments that represent annual variations of monthly averages. It is interesting that some months in a year reveal characteristic undulations that could indicate the early signs of upcoming El Niño events.

1 Introduction

The usefulness of the principal component analysis (PCA; Pearson (1901)) in data analysis is indisputable. In particular, it has been used in the earth sciences to extract characteristic structures within observational datasets. In this field, it is also called the EOF analysis (e.g., Stephenson and Benestad, 2000).

On the other hand, multidimensional sequential data, such as time series data and profile observation data, frequently appear in data analysis. It is known that such sequential data can be efficiently represented as paths by converting them into tensors called signatures (e.g., Lyons et al., 2007). Signature spaces are known to have rich algebraic and geometric structures, especially as a Lie group (e.g., Friz and Victoir, 2010).

Principal component analysis on the data constituting Lie groups has been vigorously investigated by Pennec and his group. It is essential that geodesics be made concrete rather than introducing specific connections to Lie groups. This makes the principal geodesic analysis (PGA) by Fletcher et al. (2004) feasible for structured data.

In this study, we describe how to apply PGA to sequential data through signature transformation. We also illustrate its usefulness by applying PGA to real climate time series (NOAA, 2021).

2 Theoretical background

2.1 Signatures constitute a Lie group

- A path is a function with bounded variation from unit interval $[0, 1] \subset \mathbb{R}$ to \mathbb{R}^d , defined as $X : t \mapsto (X_t^{(1)}, \dots, X_t^{(d)})$. The 0-th and 1-st iterated integrals (e.g., Lyons et al., 2007; Friz and Victoir, 2010) of path X are defined as

$$\mathcal{S}^{(0)}(X)_{0,t} = 1, \quad (1)$$

$$\mathcal{S}^{(i)}(X)_{0,t} = \int_0^t dX_s^{(i)} = X_t^{(i)} - X_0^{(i)}, \quad i = 1, \dots, d. \quad (2)$$

For $k = 2, 3, \dots$, the k -th iterated integral is defined recursively as

$$\mathcal{S}^{(i_1 \dots i_k)}(X)_{0,t} = \int_0^t \mathcal{S}^{(i_1 \dots i_{k-1})}(X)_{0,s} dX_s^{(i_k)}, \quad (3)$$

where $i_1, \dots, i_k = 1, \dots, d$.

- Then, step- n signature x (e.g., Lyons et al., 2007; Friz and Victoir, 2010) is defined as an element of tensor algebra:

$$\mathcal{T}_n(\mathbb{R}^d) = \bigoplus_{k=0}^n (\mathbb{R}^d)^{\otimes k} \ni x = 1 + \sum_{i_1=1, \dots, d} x^{(i_1)} e_{i_1} + \dots + \sum_{i_1, \dots, i_n=1, \dots, d} x^{(i_1 \dots i_n)} e_{i_1} \dots e_{i_n}, \quad (4)$$

where $x^{(i_1 \dots i_k)} = \mathcal{S}^{(i_1 \dots i_k)}(X)_{0,1}$ is the iterated integral of path X , and e_i is the basis of \mathbb{R}^d .

1. The I -th coordinate for the signature is $x^{(I)}$, where I is a multiindex. In this sense, we can regard $x \in \mathbb{R}^N$, $N = 1 + d + \dots + d^n = (d^{n+1} - 1)/(d - 1)$.
2. Since the iterated integrals should satisfy the shuffle relation (e.g., Lyons et al., 2007), e.g. $x^{(1)}x^{(2)} = x^{(12)} + x^{(21)}$, the signatures are located in a limited area in \mathbb{R}^N .

This implies that the set of order- n signature has a manifold structure (not flat).

- Signature is also an element of a **group** G . Because, for $x = \mathcal{S}_n(X) \in G$, $y = \mathcal{S}_n(Y) \in G$, we have
 1. Product: If $x, y \in G$, then $xy = \mathcal{S}_n(X * Y) \in G$.
 2. Unit: $e = \mathcal{S}_n(\bullet) = 1 \in G$ that satisfies $ex = xe = x$.
 3. Inverse: $x^{-1} = \mathcal{S}_n(\overleftarrow{X}) \in G$ that satisfies $x^{-1}x = xx^{-1} = e$.
- Since the signature is in both Group and Manifold, it is in a Lie group.
- The group has exponential map: $\exp : T_e G \rightarrow G$ and its inverse: $\log : G \rightarrow T_e G$. In particular, Log signature $v = \log x$ is an element of tangent space at the origin: $v \in T_e G$.
- A simple example with $n = 2$, $d = 2$ is as follows.

- **Signature space** G is a 3-dimensional manifold embedded in \mathbb{R}^6 :

$$G = \left\{ 1 + x^{(1)}e_1 + x^{(2)}e_2 + x^{(11)}e_{11} + x^{(12)}e_{12} + x^{(21)}e_{21} + x^{(22)}e_{22} \mid x^{(i)}x^{(j)} = x^{(ij)} + x^{(ji)} \right\} \\ \cong \left\{ \left(a, b, \frac{a^2}{2}, \frac{ab}{2} + c, \frac{ab}{2} - c, \frac{b^2}{2} \right) \mid a, b, c \in \mathbb{R} \right\}.$$

Moreover, if $x, y \in G$, then $xy \in G$ and $x^{-1} \in G$.

- **Log-signature space** \mathfrak{g} is a 3-dimensional vector space:

$$\mathfrak{g} = \left\{ x^{(1)}e_1 + x^{(2)}e_2 + \frac{x^{(12)} - x^{(21)}}{2}(e_{12} - e_{21}) \right\} \cong \{(a, b, c) \mid a, b, c \in \mathbb{R}\}.$$

Moreover, if $u, v \in \mathfrak{g}$, then $[u, v] \in \mathfrak{g}$.

- Therefore, G is regarded as a 3-dimensional manifold (not necessarily endowed with a Riemannian metric), and furthermore a Lie group.

2.2 Motivation toward Pennec's theory

- It is not possible to define a bi-invariant metric for most Lie groups, including the signature space.
- Since the **tangent PCA** (Fletcher et al., 2004) uses a metric in $T_e G$, it is not a legitimate method.
- However, without using a metric, we can extend the notion of principal component analysis to a curved space.
- In that geometry, the only thing we should employ is **geodesics**.
- By replacing straight lines to geodesics, we can perform the **principal geodesic analysis (PGA)** (Fletcher et al., 2004; Sommer et al., 2014) instead of PCA.

2.3 Essence of Pennec's theory

- A geodesic $\gamma(t)$ satisfies $\nabla_{\dot{\gamma}(t)}\dot{\gamma}(t) = 0$ (autoparallel). In particular, we consider geodesics starting from origin $\gamma(0) = e$.
- A 1-parameter subgroup $\beta(t)$ starting from origin e is determined by $\dot{\beta}(0) = X \in T_e G$, and written as $\beta(t) = \exp(tX)$, using the exponential map of the Lie group G . It is an integral curve of the left-invariant vector field \tilde{X} determined by $X \in T_e G$.
- We assume that the manifold is endowed with a Cartan-Schouten connection (CSC): $\nabla_{\tilde{X}}\tilde{Y} = \frac{1}{2}[\tilde{X}, \tilde{Y}]$, which satisfies $\nabla_{\tilde{X}}\tilde{X} = 0$.
- This connection is bi-invariant; namely, the translation of \tilde{X} and \tilde{Y} is equivalent to that of $\nabla_{\tilde{X}}\tilde{Y}$:

$$\nabla_{dL_m\tilde{X}}dL_m\tilde{Y} = dL_m(\nabla_{\tilde{X}}\tilde{Y}), \quad \nabla_{dR_m\tilde{X}}dR_m\tilde{Y} = dR_m(\nabla_{\tilde{X}}\tilde{Y}).$$

- Under the CSC, the geodesic from the origin with initial velocity $\dot{\gamma}(0) = X$ coincides with the 1-parameter subgroup $\beta(t)$.
- Therefore, we can write the geodesic as $\gamma(t) = \beta(t) = \exp(tX)$.
- Also, a geodesic starting from $m \in G$ with initial velocity $U \in T_m G$ is written as $\gamma(t) = L_m \exp(tdL_m^{-1}U)$.

$$\nabla_{\pi(x_\ell)}d^2(\pi(x_\ell), x_\ell) = -2\pi(x_\ell) \log(\pi(x_\ell)^{-1}x_\ell), \quad (5)$$

for signature x_ℓ and its projection $\pi(x_\ell)$, and logarithm map, \log , in tensor algebra.




Figure 1: Concept of Principal Geodesic Analysis. Black and blue curves are geodesics. Vector v is chosen to minimize the unexplained variance $\sum_\ell d^2(\pi(x_\ell), x_\ell)$, where $\pi(x_\ell) = m e^{t_\ell v}$.

2.4 Group mean

- Suppose we have data points $\{x_\ell | \ell = 1, \dots, L\}$.
- Group mean m satisfies $\nabla_m \sum_{\ell=1}^L d^2(m, x_\ell) = (-2)m \sum_{\ell=1}^L \log(m^{-1}x_\ell) = 0$.




Figure 2: In the right fig., the blue curves are geodesics $\gamma(t) = \exp(t \log(m^{-1}x_\ell))$. Group mean m is chosen so that the tangent vectors $v_\ell = \log(m^{-1}x_\ell)$ at the origin are in balance. Locally, it is equivalent to minimizing $\sum_\ell d^2(m, x_\ell)$.

Fréchet (1948) defined the mean m of data $\{x_\ell\}_{\ell=1}^L$ as

$$m = \arg \min_{m \in G} \sum_{\ell=1}^L d^2(m, x_\ell). \quad (6)$$

Notice that the gradient of the squared distance has an explicit form:

$$\nabla_m d^2(m, x) = -2 \text{Log}_m x, \quad (7)$$

where $\text{Log}_m x$ represents the initial velocity $\dot{\gamma}(0)$ of the geodesic from $\gamma(0) = m$ to $\gamma(1) = x$. See Appendix for derivation. Using this, Karcher (1977) characterized the mean as a stationary point:

$$\nabla_m \sum_{\ell=1}^L d^2(m, x_\ell) = \sum_{\ell=1}^L (-2) \text{Log}_m x_\ell = 0, \quad (8)$$

In the case where x and m are in the signature space G equipped with the CSC, we have $\text{Log}_m x = m \log(m^{-1}x)$. Therefore, Eq. (8) reads

$$\sum_{\ell=1}^L \log(m^{-1}x_\ell) = 0. \quad (9)$$

Equation (9) characterizes the group mean m , and it can be solved by fixed-point iteration:

$$m^{\text{new}} = m \exp\left(\frac{1}{L} \sum_{\ell=1}^L \log(m^{-1}x_\ell)\right). \quad (10)$$

2.5 Principal Geodesic Analysis

- Suppose we have data $\{x_\ell\}_{\ell=1, \dots, L}$ with their group mean m .
- We draw a geodesic, equivalently 1-parameter subgroup, passing through m . In the case of signature, it reads

$$\gamma_v(t) = m \exp(tv). \quad (11)$$

- We project each data x_ℓ onto the geodesic. The projected point is written as $\pi_v(x_\ell) = \gamma_v(t_\ell)$ with an appropriate t_ℓ .

- Then, we find the vector $v \in T_e G$ of the geodesic that minimizes the distance from the data point to the projected point (Guigui et al., 2023).

$$v = \arg \min_v \sum_{\ell=1}^L d^2(\pi_v(x_\ell), x_\ell). \quad (12)$$

- Since the summation in (12) is the variance that the geodesic cannot explain, we call it the unexplained variance.
- In this regard, the geodesic determined by (12) can be regarded as the mode that best describe the data.

2.6 Stationary value problem

We have data $\{x_\ell\}_{\ell=1, \dots, L}$ with their group mean m . We want to find the modes that minimize the unexplained variance in a geodesic space.

1. We project data x_ℓ onto the geodesic passing through the mean m ; the projected point is $\pi(x_\ell) = m \exp(t_\ell v)$. Then, find the initial velocity v of the geodesic that minimizes the distance from the data point to the projected point.

$$\forall \ell) \quad t_\ell = \arg \min_{t_\ell \in \mathbb{R}} d^2(m \exp(t_\ell v), x_\ell), \quad (13)$$

$$v = \arg \min_v \sum_{\ell=1}^L d^2(m \exp(t_\ell v), x_\ell). \quad (14)$$

2. When the signature space is regarded as a geodesic manifold, we don't have explicitly the distance, but its gradient, $\nabla_y d^2(y, x) = -2 \text{Log}_y x$. Therefore, without using a distance, we can solve it as a stationary value problem:

$$\forall \ell) \quad \nabla_{t_\ell} d^2(m \exp(t_\ell v), x_\ell) = 0, \quad (15)$$

$$\nabla_v \sum_{\ell=1}^L d^2(m \exp(t_\ell v), x_\ell) = 0. \quad (16)$$

2.7 PCA as a special case of PGA

We want to minimize the unexplained variance in Euclidean space.

- The squared distance is $d^2(m, x_\ell) = |m - x_\ell|^2$. The gradient of squared distance is $\nabla_m d^2(m, x_\ell) = (-2)(x_\ell - m)$.
- Therefore, the mean should satisfy $\nabla_m \sum_{\ell=1}^L d^2(m, x_\ell) = (-2) \sum_{\ell=1}^L (x_\ell - m) = 0$, which is solved by $m = L^{-1} \sum_{\ell=1}^L x_\ell$.
- According to Eqs. (19) and (20), the first mode v and coefficients $\{t_\ell\}$ are derived through iterations:

$$\forall \ell) \quad t_\ell^{\text{new}} = \frac{\langle v, x_\ell - m \rangle}{|v|^2}, \quad (17)$$

$$v^{\text{new}} = \frac{\sum_{\ell=1}^L t_\ell (x_\ell - m)}{\sum_{\ell=1}^L t_\ell^2}. \quad (18)$$

- Solution v coincides with the first eigenvector of covariance matrix ¹: $C = L^{-1} \sum_{\ell=1}^L (x_\ell - m)(x_\ell - m)^T$.

¹It is not the case in curved spaces.




Figure 3: Concept of Principal Component Analysis. Black and blue curves are straight lines. Vector v is chosen to minimize the unexplained variance $\sum_{\ell} d^2(\pi(x_{\ell}), x_{\ell})$, where $\pi(x_{\ell}) = m + t_{\ell}v$.

3 Methods

3.1 Climate Data

The theory will be demonstrated by using climate data from NOAA (NOAA, 2021). The spatio-temporal data to be processed are the following.

- We use time series $\{x_{\ell}\}_{\ell=1}^L$, each of which is 4-dimensional path, consisting of 13 points of monthly mean value (January to the next January).
- Number of time series is 121 for year 1900 to 2020.
- $x_{\ell}^{(0)} = \text{time}$, $x_{\ell}^{(1)} = \text{NINO1+2}$, $x_{\ell}^{(2)} = \text{NINO3}$, $x_{\ell}^{(3)} = \text{NINO4}$.
- Each time series is converted into step-4 signature. Thus, the dimension for signature is $N = (4^{4+1} - 1)/(4 - 1) = 341$.

3.2 Problem setting

1. The first mode

The first mode v is derived by solving the following stationary value problems.

$$\forall \ell) \quad \nabla_{t_{\ell}} d^2(m \exp(t_{\ell}v), x_{\ell}) = 0, \quad (19)$$

$$\nabla_v \sum_{\ell=1}^L d^2(m \exp(t_{\ell}v), x_{\ell}) = 0. \quad (20)$$

2. The second mode

Similarly, the second mode v' is derived by solving the following stationary value problems.

$$\forall \ell) \quad \nabla_{t'_\ell} d^2(m \exp(t_\ell v + t'_\ell v'), x_\ell) = 0, \quad (21)$$

$$\nabla_{v'} \sum_{\ell=1}^L d^2(m \exp(t_\ell v + t'_\ell v'), x_\ell) = 0, \quad (22)$$

where we assume v and $\{t_\ell\}$ are already derived by solving Eqs. (19) and (20). This amounts to finding the best projections onto the best 2-dimensional span $\{v, v'\}$ in $T_e G$.

3.3 Gradient Descent

1. The first mode is solved by the following iterations. Initially, v is set to the first tangent PC.

$$\forall \ell) \quad t_\ell^{\text{new}} = t_\ell - \epsilon v^T (\partial \exp|_{t_\ell v})^T (\partial_2 \text{prod}|_{m, \exp(t_\ell v)})^T (-2) p_\ell \log(p_\ell^{-1} x_\ell), \quad (23)$$

$$v^{\text{new}} = v - \epsilon' C \sum_{\ell=1}^L t_\ell (\partial \exp|_{t_\ell v})^T (\partial_2 \text{prod}|_{m, \exp(t_\ell v)})^T (-2) p_\ell \log(p_\ell^{-1} x_\ell), \quad (24)$$

where $p_\ell := m \exp(t_\ell v)$. The reasoning for this setting is detailed below.

2. The transposed derivatives of two maps $u \mapsto \exp(u)$ and $u, w \mapsto u \cdot w$, where \cdot is the product in the tensor algebra, can be derived by using the automatic differentiation (adjoint operator) of $(\partial_u \exp|_u)^T$ and $(\partial_2 \text{prod}|_{u, w})^T$.
3. With respect to $v \in T_e G$ (contra-variant), the transposed derivative is $(\partial_v d^2(m e^{tv}, x))^T \in T_e^* G$ (covariant). We thus need a map $C \in T_e G \otimes T_e G$ to get an update in $T_e G$.
4. For this purpose, we set

$$\nabla_v d^2(m e^{tv}, x) = C (\partial_v d^2(m e^{tv}, x))^T \in T_e G, \quad (25)$$

$$C = \frac{1}{L} \sum_{\ell=1}^L u_\ell u_\ell^T, \quad (26)$$

$$u_\ell = \log(m^{-1} x_\ell) \in T_e G, \quad (27)$$

where C is the covariance matrix for the data.

5. Note that C serves as a projection operator onto $\text{span}\{u_1, \dots, u_L\}$. Since u_ℓ 's are legitimate log-signatures that belong to the Lie algebra, so the update is.
6. Furthermore, our climate data have enough dimension: $\text{rank}(C) = 89$, which is comparable to that for the algebra of log-signatures: $\dim(\mathfrak{g}_4(\mathbb{R}^4)) = 90$ (cf. Reizenstein, 2017, Table 2). The degeneracy by one is due to the constant first iterated integral corresponding to the time interval common to the samples.
7. After all, the solution of Eq. (20) is found by minimizing scalar

$$[\partial_v d^2(m e^{tv}, x)] C [\partial_v d^2(m e^{tv}, x)]^T.$$

8. Similar solution procedure is employed for the 2nd mode.

4 Results

4.1 Principal modes

We draw typical time series that correspond to the eigen modes of the signatures. Before the analysis, each path $X_\ell : [0, 1] \rightarrow \mathbb{R}^4$ is converted into the step-4 signature $x_\ell = \mathcal{S}_4(X_\ell) \in \mathcal{T}_4(\mathbb{R}^4)$. In the following, v_j is the vector for the j -th mode, and σ_j is the standard deviation of intensities in the j -th mode.

- **PCA** in \mathbb{R}^{341} (control case)

- Mean (black): $m = L^{-1} \sum_{\ell=1}^L x_{\ell}$.
- First modes (red): $m \pm \sigma_1 v_1$.
- First and second modes (blue or purple): $m \pm \sigma_1 v_1 \pm \sigma_2 v_2$.

- **PGA** in Lie group

- Group mean (black): m , such that $\sum_{\ell=1}^L \log(m^{-1} x_{\ell}) = 0$.
- First modes (red): $m \exp(\pm \sigma_1 v_1)$.
- First and second modes (blue or purple): $m \exp(\pm \sigma_1 v_1 \pm \sigma_2 v_2)$.

After the analysis, each signature is converted back into a path. The first two principal modes in PCA are shown as multi-dimensional time series in Figs. 4,5,6,7, and 8. On the other hand, The first two principal geodesic modes are shown as multi-dimensional time series in Figs. 9,10,11,12, and 13.




Figure 4: Principal modes in PCA viewed as NINO1+2 time series




Figure 5: Principal modes in PCA viewed as NINO3 time series




Figure 6: Principal modes in PCA viewed as NINO4 time series




Figure 7: Principal modes in PCA viewed as NINO12vs4 time series




Figure 8: Principal modes in PCA viewed as NINO3vs4 time series




Figure 9: Principal Geodesic modes viewed as NINO1+2 time series




Figure 10: Principal Geodesic modes viewed as NINO3 time series




Figure 11: Principal Geodesic modes viewed as NINO4 time series




Figure 12: Principal Geodesic modes viewed as NINO12vs4 time series




Figure 13: Principal Geodesic modes viewed as NINO3vs4 time series

4.2 Comparison of reconstruction errors

- **PCA** in \mathbb{R}^{341} (control case #1)

$$\pi_1(x_\ell) = m + \tau_{1,\ell}u_1 + \tau_{2,\ell}u_2, \quad (28)$$

$$\tau_{i,\ell} = (u_i, x_\ell - m) \quad (29)$$

where u_i 's are eigenvectors of $\frac{1}{L} \sum_{\ell=1}^L (x_\ell - m) \otimes (x_\ell - m)$.

- **tPCA** (control case #2)

$$\pi_2(x_\ell) = m \exp(\mu_{1,\ell}w_1 + \mu_{2,\ell}w_2), \quad (30)$$

$$\mu_{i,\ell} = (w_i, \log(m^{-1}x_\ell)), \quad (31)$$

where w_i 's are eigenvectors of $\frac{1}{L} \sum_{\ell=1}^L \log(m^{-1}x_\ell) \otimes \log(m^{-1}x_\ell)$.

- **PGA** in Lie group

$$\pi(x_\ell) = m \exp(t_{1,\ell}v_1 + t_{2,\ell}v_2), \quad (32)$$

where $t_{i,\ell}$ and v_i are the solutions of Eqs. (19) and (20).

Reconstructed time series with the first two leading modes are shown in Figs. 15, 16, 17, 18, and 19.

5 Discussion

5.1 Reconstruction errors

To compare the efficacy of the mode detections, we plot the reconstruction errors by measuring a distance between the projection onto the first two modes and the original data. The points in Fig. 14 are mainly distributed in the upper left region, which shows on the whole the reconstruction by PGA is better than that by PCA or tPCA. Note that the reconstruction from PCA does not necessarily satisfies the Chen's relation.





Figure 14: Comparison of the reconstruction errors for PGA vs PCA or tPCA. Each point x represents the error: $\epsilon^2 = \|x - \pi(x)\|^2 := \sum_{k=1}^4 \left[\sum_{i_1, \dots, i_k=1}^d (x^{(i_1 \dots i_k)} - \pi(x)^{(i_1 \dots i_k)})^2 \right]^{1/k}$, where x is a signature of 1year-path from the climate data.


5.2 El Niño modes

- Obviously, PG1 and PG1±PG2 are considered as El Niño modes (Fig. 10 and 11).
- El Niño modes have characteristic undulations in May (Fig. 11) that could indicate the early signs of upcoming El Niño events.
- Regarding El Niño modes in PCA and PGA graphs (Figs. 8 and 13), we see time goes **counterclockwise** and then **clockwise**. Whereas, in PG1+PG2 time goes just **counterclockwise**. The latter feature is clearer in PGA (Fig. 13) than in PCA (Fig. 8).



samples from real data


projection onto 2PCs



projection onto 2PGs


projection onto 2tPCs

Figure 15: Several samples with their projections (NINO1+2)



samples from real data


projection onto 2PCs




projection onto 2PGs

projection onto 2tPCs


Figure 16: Several samples with their projections (NINO3)




samples from real data



projection onto 2PCs




projection onto 2PGs




projection onto 2tPCs


Figure 17: Several samples with their projections (NINO4)




samples from real data



projection onto 2PCs



projection onto 2PGs



projection onto 2tPCs

Figure 18: Several samples with their projections (NINO1+2 vs 4)




Figure 19: Several samples with their projections (NINO3 vs 4)

6 Conclusion

- We applied Penneç's theory to the Principal Geodesic Analysis (PGA) for the path signatures.
- By this method, the group mean and the leading eigenmodes can be identified by iterations for given time series data.
- We demonstrated the PGA with climate time series, and successfully identified the climatological mean and the first two leading variations.
- Our result offers the first example of rigorous analysis for the principal modes of sequential data.

7 Acknowledgments

This study was funded by JST-PROJECT-20218919.

A Why the gradient of distance squared is logarithm?

A.1 Canonical Cartan-Schouten Connection

We assume that the Lie group G is endowed with the canonical Cartan-Schouten connection (hereafter, CCS) (e.g., Cogliati and Mastrolià, 2018). Namely, the covariant derivative under CCS, $\nabla^{(0)}$, is determined by $\nabla_a^{(0)} b = \frac{1}{2}[a, b]$ for $a, b \in T_e G$.

A.2 Geodesic starting from the origin $e \in G$

Then, the geodesic starting from the origin $e \in G$ with initial velocity $v \in T_e G$ is $\gamma_v(t) = \exp(tv)$, where \exp is exponential map in Lie group. This is nothing but a one parameter subgroup. Indeed, since $\nabla^{(0)}$

is a bi-invariant connection, we have the following auto-parallelism.

$$\begin{aligned}\nabla_{\dot{\gamma}_v(t)}^{(0)} \dot{\gamma}_v(t) &= \nabla_{dL_{\gamma_v(t)} \dot{\gamma}(0)}^{(0)} dL_{\gamma_v(t)} \dot{\gamma}_v(0) = dL_{\gamma_v(t)} \nabla_{\dot{\gamma}_v(0)}^{(0)} \dot{\gamma}_v(0) \\ &= dL_{\gamma_v(t)} \frac{1}{2} [\dot{\gamma}_v(0), \dot{\gamma}_v(0)] = dL_{\gamma_v(t)} \frac{1}{2} [v, v] = 0,\end{aligned}\tag{A.33}$$

where $dL_{\gamma_v(t)}$ is the derivative of left translation by $\gamma_v(t)$.

A.3 Geodesic starting from $a \in G$

Furthermore, for $a \in G$ and $v \in T_e G$, $\gamma(t) = a \exp^{(tv)}$ is a geodesic starting from a . Indeed, because of $\dot{\gamma}(t) = dL_a \dot{\gamma}_v(t)$ and Eq. (A.33), we have the following auto-parallelism.

$$\nabla_{\dot{\gamma}(t)}^{(0)} \dot{\gamma}(t) = dL_a \nabla_{\dot{\gamma}_v(t)}^{(0)} \dot{\gamma}_v(t) = 0.\tag{A.34}$$

Note that we can also write the geodesic as

$$\gamma(t) = a \exp(tdL_{a^{-1}} \dot{\gamma}(0))\tag{A.35}$$

by using $\dot{\gamma}(0) \in T_a G$ (Theorem 5.9 in Penneec and Lorenzi, 2020).

A.4 Definition of $\text{Log}_a(b)$ in Geodesic manifold

Since a geodesic is described by a second-order ODE, initial point $\gamma(0)$ and terminal point $\gamma(1)$ uniquely determine a geodesic. For such geodesic, we define the initial velocity as

$$\text{Log}_{\gamma(0)}(\gamma(1)) := \dot{\gamma}(0).\tag{A.36}$$

Inversely, the terminal point is defined as

$$\text{Exp}_{\gamma(0)}(\dot{\gamma}(0)) := \gamma(1).\tag{A.37}$$

A.5 Log in Lie group

From Eqs. (A.35) and (A.36), we have

$$\text{Log}_a(b) = dL_a \log(a^{-1}b),\tag{A.38}$$

for $a, b \in G$ and logarithm map, \log , in Lie group.

A.6 Log in signature space

Suppose G is a signature space. By using the left translation by a of curve $\exp(tv)$, which has initial velocity v , the derivative of the left translation reads

$$dL_a v = \frac{d}{dt} L_a \exp(tv)|_{t=0} = \frac{d}{dt} a(1+tv)|_{t=0} = av,\tag{A.39}$$

for signature $a \in G$, log-signature $v \in T_e G$, and exponential map, \exp , in tensor algebra. From Eqs. (A.38) and (A.39), we get

$$\text{Log}_a(b) = a \log(a^{-1}b),\tag{A.40}$$

for signatures a and b , and logarithm map, \log , in tensor algebra.

A.7 Gradient of Squared Distance

As proven in Appendix, we have the following property for general Riemannian manifold:

$$\nabla_a d^2(a, b) = -2\text{Log}_a(b).\tag{A.41}$$

Combining Eq. (A.41) with Eq. (A.40), we get

$$\nabla_a d^2(a, b) = -2a \log(a^{-1}b),\tag{A.42}$$

for signatures a and b , and logarithm map, \log , in tensor algebra.

B Gradient of Squared Distance

B.1 Variation of Energy

Let $\gamma : [a, b] \rightarrow M$ be a smooth curve, and $f : [a, b] \times (-\epsilon, \epsilon) \rightarrow M$ be a smooth map ($\epsilon > 0$) such that $f(t, 0) = \gamma(t)$ for all $t \in [a, b]$. Energy functional is defined as $E(\gamma) = \frac{1}{2} \int_a^b |\dot{\gamma}(t)|^2 dt$.

Then, we have the first variation:

$$\begin{aligned}
 \frac{dE(\gamma_s)}{ds} &= \frac{1}{2} \frac{d}{ds} \int_a^b \langle f_t, f_t \rangle dt = \int_a^b \langle \nabla_{f_s} f_t, f_t \rangle dt \\
 &= \int_a^b \langle \nabla_{f_t} f_s, f_t \rangle dt \\
 &= \int_a^b \frac{d}{dt} \langle f_s, f_t \rangle dt - \int_a^b \langle f_s, \nabla_{f_t} f_t \rangle dt \quad \because \text{integration by parts} \\
 &= \langle f_s, f_t \rangle(b, s) - \langle f_s, f_t \rangle(a, s) - \int_a^b \langle f_s, \nabla_{f_t} f_t \rangle dt,
 \end{aligned} \tag{B.43}$$

where ∇ is covariant derivative, and $\gamma_s := f(\cdot, s)$, and f_s and $f_t : [a, b] \times (-\epsilon, \epsilon) \rightarrow T_\bullet M$ are defined as $f_s = df(\frac{\partial}{\partial s})$ and $f_t = df(\frac{\partial}{\partial t})$, respectively.

In particular, if γ_s are geodesics, and the endpoint at $t = b$ is fixed, we have

$$\left. \frac{dE(\gamma_s)}{ds} \right|_{s=0} = -\langle f_s(a, 0), \dot{\gamma}(a) \rangle. \tag{B.44}$$

B.2 Length vs Energy

For the length $L(\gamma) = \int_a^b |\dot{\gamma}(t)| dt$ of curve $\gamma : [a, b] \rightarrow M$, we have

$$L(\gamma)^2 \leq 2(b-a)E(\gamma). \tag{B.45}$$

In particular, the equality is achieved when γ is a geodesic.

B.3 Distance and geodesic

If γ is a geodesic, we have $d(\gamma(a), \gamma(b)) = L(\gamma)$.

B.4 Variation of squared distance

Hence, applying Eqs. (B.44) and (B.45) to geodesics γ_s with $a = 0$, $b = 1$ and $\gamma_s(0) = m(s)$ (variable), $\gamma_s(1) = x$ (fixed), we have

$$\begin{aligned}
 \left. \frac{d}{ds} d^2(m(s), x) \right|_{s=0} &= 2 \left. \frac{d}{ds} E(\gamma_s) \right|_{s=0} = -2 \left\langle \frac{dm}{ds}(0), \text{Log}_m x \right\rangle \\
 &= \left\langle \frac{dm}{ds}(0), -2\text{Log}_m x \right\rangle.
 \end{aligned} \tag{B.46}$$

B.5 Gradient of squared distance

In Eq. (B.46), the second term in the bracket is nothing but the gradient with respect to the first variable in the distance, which implies

$$\nabla_m d^2(m, x) = -2\text{Log}_m x. \tag{B.47}$$

Intuitively, this means that the distance increases the most in the tangential direction of the geodesic (Fig. 20).




Figure 20: The distance increases the most in the tangential direction of the geodesic.

References

- Cogliati, A. and Mastrolia, P. (2018). Cartan, Schouten and the search for connection. *Historia Mathematica*, 45(1):39–74.
- Fletcher, P. T., Lu, C., Pizer, S. M., and Joshi, S. (2004). Principal geodesic analysis for the study of nonlinear statistics of shape. *IEEE transactions on medical imaging*, 23(8):995–1005.
- Friz, P. K. and Victoir, N. B. (2010). *Multidimensional stochastic processes as rough paths: theory and applications*, volume 120. Cambridge University Press.
- Guigui, N., Miolane, N., Pennec, X., et al. (2023). Introduction to Riemannian Geometry and Geometric Statistics: from basic theory to implementation with Geomstats. *Foundations and Trends® in Machine Learning*, 16(3):329–493.
- Lyons, T. J., Caruana, M., and Lévy, T. (2007). *Differential Equations Driven by Rough Paths*, volume 1908 of *Lecture Notes in Mathematics*. Springer.
- NOAA (2021). Climate Timeseries at PSL.
- Pearson, K. (1901). LIII. On lines and planes of closest fit to systems of points in space. *The London, Edinburgh, and Dublin philosophical magazine and journal of science*, 2(11):559–572.
- Pennec, X. and Lorenzi, M. (2020). Beyond Riemannian geometry: The affine connection setting for transformation groups. In *Riemannian Geometric Statistics in Medical Image Analysis*, pages 169–229. Elsevier.
- Reizenstein, J. (2017). Calculation of iterated-integral signatures and log signatures. *arXiv preprint arXiv:1712.02757*.
- Sommer, S., Lauze, F., and Nielsen, M. (2014). Optimization over geodesics for exact principal geodesic analysis. *Advances in Computational Mathematics*, 40:283–313.
- Stephenson, D. and Benestad, R. E. (2000). Environmental statistics for climate researchers. *Tutorial Version*, 14:2006.