### INFERENCE ON EIGENVECTORS OF NON-SYMMETRIC MATRICES

JEROME R. SIMONS

ABSTRACT. Please click here for the latest version. This paper argues that the symmetrisability condition in Tyler (1981) is not necessary to establish asymptotic inference procedures for eigenvectors. We establish distribution theory for a Wald and *t*-test for full-vector and individual coefficient hypotheses, respectively. Our test statistics originate from eigenprojections of non-symmetric matrices. Representing projections as a mapping from the underlying matrix to its spectral data, we find derivatives through analytic perturbation theory. These results demonstrate how the analytic perturbation theory of Sun (1991) is a useful tool in multivariate statistics and are of independent interest. As an application, we construct confidence sets for Bonacich centralities estimated from adjacency matrices induced by directed graphs.

#### 1. INTRODUCTION

Finding eigenvectors of matrices is a fundamental task in many disciplines. It is common for eigenvectors to appear as stochastic objects. We consider hypothesis testing for eigenvectors and the construction of confidence regions of a deterministic matrix of which we have a random estimate. Thereby the popular matrix denoising model applies in our setting, with the exception that we focus on fixed-size matrices.

We assume that we observe a sample of n matrices  $M_1, \ldots, M_n$ , such that the columns of  $a_n (M_n - M)$  converge weakly to a multivariate normal distribution centered at the columns of  $M \in \mathbb{R}^{p \times p}$  for an increasing sequence  $a_n$ . We are interested in testing the associated null hypothesis of v belonging to an eigenspace of M indexed by a collection of roots  $\mathcal{L}$  according to

$$H_0: v \in \operatorname{eig}_{\mathcal{L}} M$$

against the one-sided alternative  $H_1 : v \notin \operatorname{eig}_{\mathcal{L}} M$ . Associating eigenvectors with a set of roots, we assume that eigenvalues split according to

$$\mathcal{L} := \mathcal{L}_I \cup \mathcal{L}_J.$$

A case distinction arises from the rank of v:

Case 1.  $\operatorname{rk} v = r \leq |\mathcal{L}_I|$ , i.e. the number of linearly independent columns in v is less than or equal to the dimension of the invariant subspace spanned by the vectors associated with roots in  $\mathcal{L}_I$ . To accommodate this case, define the null hypothesis

(1.1) 
$$H_0: v \in \operatorname{eig}_{\mathcal{L}_I} M$$

against the alternative  $H_1$ :  $v \notin \operatorname{eig}_{\mathcal{L}_I} M$ , where  $\operatorname{eig}_{\mathcal{L}_I} M$  refers to the eigenspace associated with roots in  $\mathcal{L}_I$ .

Case 2.  $\operatorname{rk} v = r > |\mathcal{L}_I|$ , i.e. the number of linearly independent columns in v exceeds the number of roots in  $\mathcal{L}_I$ . We define the null hypothesis

$$H_0^*$$
 : eig<sub>L</sub>  $M \in \operatorname{sp} v$ ,

Date: 27 March '23.

which can be tested by constructing the  $p \times q$  matrix  $v_{\perp}$  such that  $v_{\perp}^{\mathsf{T}} v = 0$ . Consequently, we write

(1.2) 
$$H_0^* : v_\perp \in \operatorname{eig}_{\mathcal{L}_{J \neq I}} M.$$

Case 1 is intuitive because the columns of v span a lower-dimensional subspace than  $\operatorname{eig}_{\mathcal{L}} M$  while Case 2 is effectively an overdetermination of the eigenspace of interest so that the estimated eigenspace  $\operatorname{eig}_{\mathcal{L}_I} M$ , under the null hypothesis, is wholly contained in the candidate space sp v. Tyler (1981) implements Case 2 by inverting the hypothesis (1.1) so that Case 2 becomes (1.2) and thereby reduces to Case 1. The orthocomplement,  $v_{\perp}$ , is always easy to compute from vusing, e.g. an LU factorization or the Gram-Schmidt algorithm. Moreover, for the benefit of the reader wanting to appreciate the details of the perturbation theory, the notation of  $H_0^*$  implies simpler notation. We shall therefore write our arguments in the form of Case 2, where inference on  $v_{\perp}$  belonging to  $\operatorname{eig}_{\mathcal{L}_I}$  implies inference on v without loss of generality.

The main limitation of the existing approaches is that we have to assume that M is such that  $\Gamma M$  is symmetric for a positive definite symmetric matrix  $\Gamma$ . We illustrate this issue with

**Example 1.** The matrix  $M_1$  is symmetric in the metric of  $\Gamma$  for

$$\Gamma = \begin{bmatrix} 2 & 1 \\ 1 & 6 \end{bmatrix} \qquad \qquad M_1 = \begin{bmatrix} 1 & 3 \\ 1 & 1 \end{bmatrix}$$

whereas  $M_2$  is not symmetric in the metric of  $\Gamma$ 

$$M_2 = \begin{bmatrix} 1 & -3 \\ -1 & 1 \end{bmatrix}.$$

Any  $2 \times 2$  matrix  $\Gamma$  would have to be negative definite for  $M_1\Gamma$  to be symmetric. Therefore, inferential procedures for non-symmetric matrices apply to  $M_1$  but not to  $M_2$ . In the following, we shall establish that no such requirement is necessary for inference on eigenspaces of nonsymmetric matrices.

1.1. Related literature on eigenspace inference. Statistical analysis of eigenvectors and eigenvalues finds use in diverse applications such as quasi-cointegration analysis, the study of centrality in social networks (Bonacich 1987), or in radio signal processing (Penna, Garello, and Spirito 2009).

The study of inference and estimation of eigenvectors of more general, dense square random matrices  $M \in \mathbb{R}^{p \times p}$  with p fixed began with the seminal study of Davis (1977) in the non-normal case. James (1977) posed the problem of eigenvector inference for general eigenvectors without associating them with a particular set of eigenvalues. Tyler (1981) considers null hypotheses of eigenvectors associated with particular eigenvalues. His study served as the inspiration for our paper and requires quasi-symmetry of M, which we relax.

The first study to examine spectra of symmetric random matrices is the seminal paper by Anderson (1963), which provided joint density functions for eigenvectors and eigenvalues of covariance matrices and a testing procedure for equality of eigenvalues. Subsequently, James (1964) offered a general method to obtain the distribution of statistics derived from random matrices. Since then, the field of eigenspace statistics based on random matrices has progressed along two strands.

The first branch of the literature considers the spectra of Wishart matrices. To see how they arise, consider a matrix  $A \in \mathbb{R}^{p \times n}$  whose columns  $A_i$  are independent and normal according to

 $A_i \sim N_p(0, \Omega)$  where  $\Omega \in \mathbb{R}^{p \times p}$  is a positive-definite covariance matrix. The variate  $M \coloneqq AA^{\mathsf{T}}$  will then have a central, *p*-variate Wishart distribution with *n* degrees of freedom, usually denoted by  $M \sim W_p(\Omega, n)$ . A representative study with a further overview of the development of density functions of such matrices is given in Zanella, Chiani, and Win (2009).<sup>1</sup> Further refinements include expanding or restricting the spectral structure such as the study by Takemura and Sheena (2005) who derive the asymptotic distribution of eigenvalues and eigenvectors of Wishart matrices in the case of infinitely dispersed population eigenvalues.

A second strand of inference on eigenspaces are the spectra of symmetric square matrices  $M \in \mathbb{R}^{p \times p}$  with  $p \to \infty$ . The often sparse nature of this type of matrix implies that one can truncate spectral data or singular value decomposition to estimate low-rank representations of such matrices. This literature began with the semi-circle law of Wigner (1958) for the joint eigenvalue distribution and Tracy and Widom (1994) for the distribution of the largest eigenvalue. For applied researchers, Chen, Cheng, and Fan (2020) develop a method to conduct inference on low-rank representations of random matrices that obtain from deterministic symmetric matrices that are (potentially asymmetrically) perturbed. As a further refinement of this case, Cheng, Wei, and Chen (2021) consider matrices with finely spaced eigenvalues while Iain M Johnstone, Onatski, et al. (2020) develop a testing framework for high-dimensional matrices with a single, dominant eigenvalue, or *spike*. That latter phenomenon arises for example in auto-covariance matrices of embedded time series if the process possesses a root of unity. In the class of low-rank matrices with  $p \to \infty$ , Cheng, Wei, and Chen (2021) offer an estimation and inference procedure for eigenvectors of such random matrices. Bura and Pfeiffer (2008) add theory for inference on singular vectors.

A subspecialty of the aforementioned literature studies estimation and inference procedures for eigenvectors and principal component vectors of covariance matrices, as they arise in in factor models. If the column dimension of  $A_n$  diverges as  $n \to \infty$ , central limit theorems imply that a suitably normalized  $M_n := A_n A_n^{\mathsf{T}}$  converges to a central Wishart matrix as above. This result nests the limiting distributions of a large class of covariance matrix estimators. Tyler (1983) provides a procedure to compute asymptotic confidence sets for principal components which obtain from eigenspaces of covariance matrices. Iain M. Johnstone (2001) and Lee and Schnelli (2016) contribute results on the largest eigenvalue statistics of principal component estimators and covariance matrices. Koltchinskii, Löffler, Nickl, et al. (2020) consider a representative functional extension and focus on efficiently estimating linear functionals of principal components, which have symmetric underlying covariance matrices.

A related literature studies the singular value decomposition of rectangular random matrices  $M \in \mathbb{R}^{p \times d}$  where  $p, d < \infty$  and p and d are fixed. Cai et al. (2019) offer a method to estimate the column space via a singular value decomposition and Girko (1999) provides conditions for and derive a convergence law for associated singular values.

# 2. Assumptions

We work with the matrix  $M \in \mathbb{R}^{p \times p}$  with associated estimators  $\hat{M}_n$ . To simplify the subsequent exposition and to more easily define test statistics, we vectorise the matrix M for most of the arguments. The following defines our framework:

<sup>&</sup>lt;sup>1</sup>Owing to their use as representations of physical systems, studies in eigenstatistics of Wishart matrices tend to prioritize those with complex entries. For the social sciences, this literature is therefore less useful because results derived using complex matrices do not, in general, apply to real matrices.

**Assumption 2.** Let  $\hat{M}_n$  be an estimator of M and  $\hat{\Omega}$  be an estimator of  $\Omega$ . Then,

- (1)  $\sqrt{n} \operatorname{vec} \left( \hat{M}_n M \right) \rightsquigarrow N(0, \Omega)$  for a full rank covariance matrix  $\Omega \in \mathbb{R}^{p^2 \times p^2}$ . The columns of  $\sqrt{n}\left(\hat{M}_n - M\right) \rightsquigarrow N\left(0, \Omega_M\right)$  for a full rank  $\Omega \in \mathbb{R}^{p \times p}$  so that  $\Omega = I_p \otimes \Omega_M$ .
- (2)  $\hat{M}_n \xrightarrow{p} M$ .
- (3) For a covariance matrix estimator,  $\hat{\Omega} \xrightarrow{p} \Omega$ .
- (4) The set of roots of interest,  $\mathcal{L}_I$ , is closed under conjugation, i.e. for any  $z \in \mathcal{L}_I$ , the complex conjugate  $\bar{z} \in \mathcal{L}_I$ .
- (5) Estimators of right and left eigenvectors  $\hat{R}_n$  and  $\hat{L}_n$  converge in probability to their population analogues.
- (6) The eigenvalues of M correspond to unit blocks in the Jordan normal form of M.
- (7)  $\mathcal{L}_I \cap \mathcal{L}_J = \emptyset.$

Assumption 2.5 may be relaxed with very little additional work. We do not pursue this extension to focus on the perturbation theory. Assumption 2.7 is fairly innocuous and simply means that we can discriminate between eigenvalues of interest and remaining ones. Assumption 2.4 simplifies our arguments but is also not strictly necessary.

## 3. Eigenprojections as functionals of underlying matrix

Before introducing our test statistics, let us study eigenprojections as our key ingredient for a test statistic. Define the eigenprojection over the subspace spanned by eigenvectors associated with roots in  $\mathcal{L}_I$ , for right and left eigenvectors associated with root  $\lambda$ ,  $r_{\lambda}$  and  $l_{\lambda}$ , via

$$P_I := \sum_{\lambda \in \mathcal{L}_I} \lambda r_\lambda l_\lambda^{\mathsf{T}}.$$

To see eigenprojections in practice, we consider

### Example 3. Let

(3.1) 
$$M = \begin{bmatrix} 0.8 & 0.5 \\ 0 & 0.4 \end{bmatrix}$$

Then, right eigenvectors are  $r_1^{\mathsf{T}} = \begin{bmatrix} 1 & 0 \end{bmatrix}$  and  $r_2^{\mathsf{T}} = \begin{bmatrix} -0.781 & 0.625 \end{bmatrix}$  and eigenvalues are  $\lambda_1 = 0.8$  and  $\lambda_2 = 0.4$ . Its left eigenvectors are  $l_1^{\mathsf{T}} = [1 \ 1.25]$  and  $l_2^{\mathsf{T}} = [0 \ 1.6]$ . The associated eigenprojections  $i_i l_i^{\mathsf{T}}$ 

$$(3.2) P_{\lambda_i} = r$$

for  $i \in \{1, 2\}$ 

$$P_{\lambda_1} = \begin{bmatrix} 1 & 1.25 \\ 0 & 00 \end{bmatrix} \qquad \qquad P_{\lambda_2} = \begin{bmatrix} 0 & -1.25 \\ 0 & 1 \end{bmatrix}$$

We see that  $P_{\lambda_i}$  are rank-1 projection matrices for which  $P_{\lambda_i}P_{\lambda_j} = \delta_{ij}$  and if v belongs to an eigenspace associated with root  $\lambda_i$ , we have  $\nu P_{\lambda_i} = P_{\lambda_i} \nu$  and  $P_{\lambda_1} + P_{\lambda_2} = I$ . Finally, under the maintained null hypothesis,

$$P_{\lambda_i}v = v,$$

from which we can build a test statistic.

For the normalisation of test statistics, we shall need a generalized inverse for a matrix A. which is a matrix  $A^+$  such that  $AA^+A = A$ . For a matrix of rank s, let the eigenprojection be as in Example 3. Then, a generalized inverse

(3.3) 
$$A^+ \coloneqq \sum_{i=1}^s \lambda_i^{-1} P_{\lambda_i}$$

Respecting the division of eigenvalues

$$M = R_I \Lambda_I L_I^\mathsf{T} + R_J \Lambda_J L_J^\mathsf{T}$$

(3.4) 
$$R = \begin{bmatrix} R_I R_I \end{bmatrix} L = \begin{bmatrix} L_I L_J \end{bmatrix} \Lambda = \begin{bmatrix} \Lambda_I & 0 \\ 0 & \Lambda_J \end{bmatrix},$$

we introduce the skew projection

$$(3.5) P_I \coloneqq R_I L_I^{\dagger}.$$

If the roots in the set  $\mathcal{L}_I$  are closed under conjugation, under Assumption 2.4, then we may take  $R_I$  and  $L_I$  to be real without loss of generality. We proceed with testing  $H_0: v \in \text{eig}_{\mathcal{L}_I}$  via requiring  $v_{\perp}^{\mathsf{T}} P_I = 0$ .

Invariant subspace functionals. Our study proceeds by considering the map  $\psi$  from M to its invariant subspaces, which we specialise to eigenspaces. Building on the perturbation theory in Stewart and Sun (1991), we compute first and second order derivatives of  $\psi$ . This approach has the advantage that its conclusions apply to any invariant subspace map  $\psi$ . For a member of an invariant subspace  $X_1 \in \mathbb{R}^{p \times r}$ , the defining equality holds

for some matrix  $A_1 \in \mathbb{R}^{r \times r}$ . Thereby, we nest the Schur decomposition for  $A_1$  being a Schur form of M. Let the ring of matrices satisfying Assumption 2 be  $\mathscr{Q}$ . Then, define the map  $\psi : \mathscr{Q} \mapsto \mathbb{R}^{r \times q}$  by

(3.7) 
$$\psi(M) \coloneqq v_{\perp}^{\mathsf{T}} P_{I}(M) .$$

Introduce the following notation for convenience

(3.8) 
$$\hat{\Delta}_{n,\mathrm{IJ}} = L_J^{\mathsf{T}} \left( \hat{M}_n - M \right) R_I$$

and the operator  $\boldsymbol{S}$  :  $\mathbb{R}^{r \times q} \to \mathbb{R}^{r \times q}$ 

(3.9) 
$$\boldsymbol{S}(Q) = Q\Lambda_I - \Lambda_J Q,$$

where  $Q \in \mathbb{R}^{r \times q}$ . Then, the first-order Fréchet derivative of  $\psi$  is

(3.10) 
$$\dot{\psi}(M) = R_I \boldsymbol{S}^{-1} \left( \hat{\Delta}_{\mathrm{JI},n} \right) L_J^{\mathsf{T}} \boldsymbol{\upsilon}.$$

Before concluding this section with our main result, we define the Jacobian matrix of a vectorized, matrix-valued function  $F(X) : \mathbb{R}^{p \times p} \to \mathbb{R}^{r \times q}$  via first differentials as J satisfying

$$\mathrm{d}F = J\,\mathrm{dvec}\,X.$$

We conclude with

**Lemma 4.** Let the map  $\psi(M)$  be as in (3.7). Then, the Jacobian matrix with respect to the eigenspace spanned by the columns of  $L_J$  is B such that dvec  $\psi = B \operatorname{dvec} R_I$  and

(3.11) 
$$B = \left(L_I \otimes v_{\perp}^{\mathsf{T}} R_J\right) \left\{ \left(\Lambda_I^{\mathsf{T}} \otimes I_J\right) - \left(I_I \otimes \Lambda_J\right) \right\}^{-1} \left(I_q \otimes L_J^{\mathsf{T}}\right).$$

### 4. Asymptotic distribution of estimators

We define the estimator of the eigenprojection  $v_{\perp}^{\mathsf{T}} \hat{P}_{I} \left( \hat{M} \right) := v_{\perp}^{\mathsf{T}} \hat{R}_{I} \hat{L}_{I}$ . Then, we obtain

Proposition 5. Suppose Assumption 2 holds. Then,

$$\sqrt{n} \operatorname{vec} v_{\perp}^{\mathsf{T}} \hat{P}_{I}\left(\hat{M}\right) \rightsquigarrow N\left(0, B\Omega B\right).$$

Proposition 5 forms the basis for the Wald statistic and is our prototype statistic. In some cases, it is convenient to have a unique orthocomplement estimator with a normalization of the test matrices v and  $v_{\perp}$  that effects interpretable coefficients.

Eigenvectors are only unique up to a normalisation. We therefore introduce the normalisation for a coefficient matrix  $D \in \mathbb{R}^{q \times r}$ . The normalization of v is such that  $v_{\perp}^{\mathsf{T}} N v_{\perp} = I_p$  for  $N = \begin{bmatrix} I_r & 0 \\ 0 & 0 \end{bmatrix} \in \mathbb{R}^{p \times p}$ . The test matrices become

(4.1) 
$$v \coloneqq \begin{bmatrix} D\\ I_r \end{bmatrix}$$

and

(4.2) 
$$v_{\perp} \coloneqq \begin{bmatrix} I_r \\ -D \end{bmatrix}$$

Unfortunately, uniqueness comes at the steep cost of making the second order moment potentially unbounded as Anderson (2010) discusses. Letting  $\Omega_M$  be as in Assumption 2.1, we have the alternative normalization  $v_{k,\perp}^{\mathsf{T}} \Omega_M v_{k,\perp} = I_p$  for every column k in v.<sup>2</sup>

We define the set estimator

(4.3) 
$$\widehat{v}_{\perp,n} \coloneqq \left\{ v_{\perp} \in \mathbb{R}^p : v_{\perp}^{\mathsf{T}} \widehat{P}_{i,n} = 0 \right\},$$

where  $\hat{P}_{I,n} \coloneqq \hat{R}_{I,n} \hat{L}_{I,n}^{\mathsf{T}}$  uses sample analogues of  $R_I$  and  $L_I$ . For a closed-form expression of  $\hat{D}_n$ under the normalization (4.2), partition  $R_I$  conformably for  $R_{I,1} \in \mathbb{R}^{r \times q}$  and  $R_{I,2} \in \mathbb{R}^{q \times q}$ .

(4.4) 
$$\begin{bmatrix} R_{I,1} \\ R_{I,2} \end{bmatrix} \coloneqq R_I$$

Consequently, define

(4.5) 
$$\hat{D}_n = \left(\hat{R}_{I,2,n}\right)^{-1} \hat{R}_{I,1,n}^{\mathsf{T}}$$

and observe that (4.5) defines  $\hat{D}_n$  uniquely. Then, we obtain

**Theorem 6.** Let  $D_n$  be as in (4.5) and suppose Assumption 2 holds. Then,

$$n^{1/2}vec \left\{ \hat{D}_n^{\mathsf{T}} - D^{\mathsf{T}} \right\} \rightsquigarrow N\left( 0, B_A \Omega_M B_A^{\mathsf{T}} \right)$$

for

$$B_A \coloneqq \left( R_{I,2}^{-1} \otimes v_{\perp}^{\mathsf{T}} R_J \right) \left\{ \left( \Lambda_I^{\mathsf{T}} \otimes I_J \right) - (I_I \otimes \Lambda_J) \right\}^{-1} (R_I^{\mathsf{T}} \otimes L_J^{\mathsf{T}}).$$

Finally, for covariance matrix estimators we record

**Lemma 7.** Suppose Assumption 2 holds, then, whenever  $\hat{B} \xrightarrow{p} B$ ,  $\hat{B}\hat{\Omega}\hat{B}^{\mathsf{T}} \xrightarrow{p} B\Omega_M B$ .

 $<sup>^{2}</sup>$ For an approach that avoids normalizations see Silin and Fan (2020) in the context of symmetric matrices.

#### 5. Tests

We define a Wald test statistic for  $H_0^* : v_{\perp}^{\mathsf{T}} P_I = 0$ . Based on Proposition 5 we define a Wald test statistic for  $\hat{\Omega}_W^+$  denoting a generalised inverse of  $\hat{\Omega}_W$ ,

(5.1) 
$$\hat{W}_n(v_{\perp}) \coloneqq \operatorname{vec} \left(v_{\perp}^{\mathsf{T}} \hat{P}_I\right)^{\mathsf{T}} \hat{\Omega}_W^+ \operatorname{vec} \left(v_{\perp}^{\mathsf{T}} \hat{P}_I\right)$$

and an associated covariance matrix estimator  $\hat{\Omega}_W = \hat{B}_W \hat{\Omega} \hat{B}_W$  for

(5.2) 
$$\hat{B}_W \coloneqq \left(\hat{L}_I \otimes \upsilon_{\perp}^{\mathsf{T}} \hat{R}_J\right) \left\{ \left(\hat{\Lambda}_I^{\mathsf{T}} \otimes I_J\right) - \left(I_I \otimes \hat{\Lambda}_J\right) \right\}^{-1} \left(\hat{R}_I^{\mathsf{T}} \otimes \hat{L}_J^{\mathsf{T}}\right).$$

Theorem 8. Suppose Assumption 2 holds and let. Then,

(5.3) 
$$\hat{W}_n(v_\perp) \rightsquigarrow \chi^2_{qm}.$$

The test proposed in Theorem 8 is for a full-vector hypothesis on the eigenspace. To be able to conduct subset inference, we wish to consider a null hypothesis about individual coefficients.

Let  $\hat{R}_{I,2}^{-1}$  be the empirical analogue of  $R_{I,2}^{-1}$  defined in (4.4) and define analogously to (5.2),

$$\hat{B}_D \coloneqq \left(\hat{R}_{I,2}^{-1} \otimes \hat{v}_{\perp}^{\mathsf{T}} \hat{R}_J\right) \left\{ \left(\Lambda_I^{\mathsf{T}} \otimes I_J\right) - \left(I_I \otimes \Lambda_J\right) \right\}^{-1} \left(\hat{R}_I^{\mathsf{T}} \otimes \hat{L}_J^{\mathsf{T}}\right)$$

and define a t-test statistic for inference on the ijth coefficient of D via

(5.4) 
$$t_n(a_0) \coloneqq \frac{d_{ij} - d_0}{\hat{\sigma}_{ij}},$$

where  $\hat{\sigma}_{ij}^2 \coloneqq \left(\hat{\Omega}_D^{-1}\right)_{ij}$  and  $\hat{\Omega}_D \coloneqq \hat{B}_D \Omega \hat{B}_D^{\mathsf{T}}$ .

Theorem 9. Suppose Assumption 2 holds, then

 $t_{i,n}\left(d_{0}\right) \rightsquigarrow N\left(0,1\right).$ 

## 6. Application

Interpret A as representing a graph with vertices indexed by i and j and entries representing trade flows  $q_{ij} > 0$  if firm i is a net seller to firm j and  $q_{ij} < 0$  otherwise. Then, the matrix A captures the directed graph of the trade network. The importance score  $s_i$ ,

(6.1) 
$$s_i \coloneqq \frac{1}{\lambda} \sum_{j \in N_i} s_j$$

measures the relative importance of firm i in the network. The sum runs over all j that are direct neighbours of firm i, denoted by  $N_i$  and  $\lambda$  is a constant. The measure  $s_i$  is known as the Katz or eigenvector centrality, and was first introduced by Bonacich (1987). For the full vector of scores s, we obtain the relationship

$$(6.2) As = \lambda s,$$

which defines an eigenvector problem. If the matrix A is estimated with a degree of uncertainty or measurement error, we will be able to test hypotheses of the form

 $H_0$ : s is a vector of importance scores.

which we construct via positing that  $s \in \mathbb{R}^{p \times 1}$  is an eigenvector of the adjacency matrix A belonging to the largest eigenvalue  $\lambda$  against the alternative that some entries do not contain

importance scores. To derive an asymptotically valid confidence set of importance scores, we use Theorem 8 for an inversion of the Wald test. Find the orthocomplement  $s_{\perp} \in \mathbb{R}^{p \times (p-1)}$  such that  $s_{\perp}^{\mathsf{T}} s = 0$  and define a level  $\alpha$  confidence set as

$$\mathcal{C} := \left\{ s \text{ such that } \hat{W}_n(s_{\perp}) \leq \chi_{1-\alpha}^2 \right\}.$$

If  $\lambda \in \mathbb{C}$  and  $|\lambda|$  is maximal across  $\mathcal{L}_A$  then we take the two eigenvectors corresponding to  $\lambda$  and its complex conjugate, and set  $w_i := |s_i|$  with  $w_i$  replacing  $s_i$  everywhere.

An extension left for a future development of this section is the simulation of this method to study the performance numerically.

## 7. CONCLUSION

This paper has extended the theory of Tyler (1981) to cover non-symmetric matrices. In addition to testing full vector hypotheses, we also provide a t-test on individual coefficients of the vector and derive the asymptotic distribution of the estimator for normalised eigenvectors. We caution the reader that normalised estimates may suffer from a potentially unbounded second moment. However, we believe that the benefits of being able to normalise coefficients outweigh its drawbacks.

We have also shown how to apply the analytic perturbation theory of Sun (1991) to the problem of inference on eigenvectors and hope that our exposition can aid researchers in similar settings. Naturally, the perturbation theory of Kato (1995) delivers the same results. For a pedagogic introduction to the subject, we refer the reader to Greenbaum, Li, and Overton (2019).

The leading application of our results is to graph Laplacians or adjacency matrices that directed graphs induce. A future version of this paper will offer further statistical development of this application.

#### 8. Proofs

8.1. Expansions of eigenprojections. We remark that the perturbation theory employed in this paper does not stipulate a normalisation for eigenvectors, which is innocuous; derivatives of normalised maps deliver the same results (Duffy and Simons 2020).

In addition to the map  $\psi$ , define the following notation:

(8.1) 
$$\begin{bmatrix} \hat{\Delta}_{n,I} \\ \hat{\Delta}_{n,SL} & \hat{\Delta}_{n,J} \end{bmatrix} \coloneqq \begin{bmatrix} L_I^\mathsf{T} \left( \hat{M}_n - M \right) R_I \\ L_J^\mathsf{T} \left( \hat{M}_n - M \right) R_I & L_J^\mathsf{T} \left( \hat{M}_n - M \right) R_J \end{bmatrix}.$$

We collect first and second-order derivatives in

Lemma 10. Suppose Assumption 2 holds. Then the first and second order Fréchet derivatives of  $\psi(M)$  at M are

(8.2) 
$$\dot{\psi}\left(\hat{M}_n - M\right) = v_{\perp}^{\mathsf{T}} R_J S^{-1}\left(\hat{\Delta}_{n,SL}\right) L_I^{\mathsf{T}}$$

(8.3) 
$$\ddot{\psi}\left(\hat{M}_n - M\right) = 2v_{\perp}^{\mathsf{T}} R_J \boldsymbol{S}^{-1} \left(\left(\hat{\Delta}_{n,J} \boldsymbol{S}^{-1}\left(\hat{\Delta}_{n,IJ}\right) - \boldsymbol{S}^{-1}\left(\hat{\Delta}_{n,IJ}\right)\hat{\Delta}_{n,I}\right)\right) L_I^{\mathsf{T}}.$$

Record the following result for the expansion of  $\psi$  and denote by r(.) the remainder term of the asymptotic expansion.

(1) We have the expansion Lemma 11.

$$\sqrt{n}v_{\perp}^{\mathsf{T}}\left(\hat{P}_{n,I}-P_{I}\right) = \sqrt{n}v_{\perp}^{\mathsf{T}}\hat{R}_{n,J}\boldsymbol{S}^{-1}\left(\Delta_{n,\mathrm{IJ}}\right)\hat{L}_{n,I}^{\mathsf{T}} + \sqrt{n}r\left(\hat{M}_{n}-M\right)$$

(2) The remainder term obeys  $r\left(\hat{M}_n - M\right) = O\left(n^{-1}\right)$ .

8.2. Asymptotic distribution of eigenprojection and Wald test statistic. Further, define  $B_{IJ}^{\mathsf{T}} \coloneqq R_I \otimes L_J$ . Then, we have

**Lemma 12.** Let  $\Delta_{n,IJ}$  be as in Lemma 10. Then, for  $v_{\perp} \in \mathbb{R}^{p \times (p-m)}$ ,

- (1)  $\sqrt{n} \operatorname{vec} \Delta_{n,IJ} \rightsquigarrow N\left(0, B_{IJ}\Omega B_{IJ}^{\mathsf{T}}\right).$ (2)  $\sqrt{n} \operatorname{vec} \left\{ v_{\perp}^{\mathsf{T}} \hat{R}_{n,J} \boldsymbol{S}^{-1} \left(\Delta_{n,IJ}\right) \hat{L}_{n,I}^{\mathsf{T}} \right\} \rightsquigarrow N\left(0, \Omega_{W}\right)$
- (3)  $\operatorname{rk}\Omega_W = qm$ .

Proposition 5 follows from Lemma 10, Lemma 11, and Lemma 12. Lemma 7 follows from the continuous mapping theorem.

To prove Theorem 8, recall that

$$\boldsymbol{\upsilon}_{\perp}^{\mathsf{T}}\hat{P}_{n,I} = \boldsymbol{\upsilon}_{\perp}^{\mathsf{T}}\left(\hat{P}_{n,I} - P_{I}\right)$$

because  $v_{\perp}^{\mathsf{T}} P_I = 0$  under the maintained hypothesis. By Lemma 111, we obtain the expansion

$$\boldsymbol{v}_{\perp}^{\mathsf{T}}\left(\hat{P}_{n,I}-P_{I}\right)=\sqrt{n}\boldsymbol{v}_{\perp}^{\mathsf{T}}R_{n,J}\boldsymbol{S}^{-1}\left(\hat{\Delta}_{n,\mathrm{IJ}}\right)L_{n,I}^{\mathsf{T}}+\sqrt{n}r\left(\hat{M}_{n}-M\right).$$

The first term in the preceding display converges according to Lemma 12, while the second converges in probability to zero by Lemma 112.

For convergence rates, we record:

**Corollary 13.** Let  $\hat{\Delta}_{n,i}$  with  $i \in \{I, IJ, J\}$  be as in (8.1). Then,

$$\begin{bmatrix} \hat{\Delta}_{n,I} \\ \hat{\Delta}_{n,IJ} & \hat{\Delta}_{n,J} \end{bmatrix} = \begin{bmatrix} O_p \left( n^{-1/2} \right) \\ O_p \left( n^{-1/2} \right) & O_p \left( n^{-1/2} \right) \end{bmatrix}.$$

8.3. Asymptotic distributions of test statistics. To prove Theorem 6, apply to  $\sqrt{n} \left( \hat{D}_n^{\mathsf{T}} - D^{\mathsf{T}} \right)$  the expression given in the following

**Lemma 14.** Let the estimator  $\hat{D}_n = \hat{R}_{n,2,I} \left( \hat{R}_{n,1,I} \right)^{-1}$ . Then, the centered estimator

(8.4) 
$$\left(\hat{D}_n - D\right)^{\mathsf{T}} = v_{\perp}^{\mathsf{T}} \left(\hat{R}_I \hat{R}_{2,I}^{-1} - R_I R_{2,I}^{-1}\right).$$

The following result is a corollary to Lemma 11.

**Corollary 15.** (1) We have the expansion

$$\sqrt{n}\left(\hat{D}_{n}^{\mathsf{T}}-D^{\mathsf{T}}\right)=\sqrt{n}v_{\perp}^{\mathsf{T}}\hat{R}_{n,J}\boldsymbol{S}^{-1}\left(\Delta_{n,\mathrm{IJ}}\right)R_{n,2,I}^{-1}+\sqrt{n}r\left(\hat{M}_{n}-M\right)$$

(2) The remainder term obeys  $r\left(\hat{M}_n - M\right) = O\left(n^{-1}\right)$ .

Then, we obtain the desired result by applying Lemma 12 to the expansion obtained in Corollary 15 by substituting  $R_{n,2,I}^{-1}$  for  $L_{n,I}^{\mathsf{T}}$ .

In the following, we show two results that should be compared to their analogues in Lemma 10 and Lemma 11. Moreover, define in analogy to (3.7) the map  $\psi_D : \mathscr{Q} \mapsto \mathbb{R}^{r \times q}$  with  $\psi_D(M) := v_{\perp}^{\mathsf{T}} R_I(M) R_{2,I}^{-1}$ . Then, we have

**Corollary 16.** The Fréchet derivative of  $\psi_D$ ,

$$\dot{\psi}_D\left(\hat{M}_n - M\right) = \upsilon_{\perp}^{\mathsf{T}} R_{n,J} \mathbf{S}^{-1}\left(\hat{\Delta}_{n,SL}\right) L_{n,I}^{\mathsf{T}} R_{n,2,I}^{-1}.$$

Theorem 6 follows from Lemma 14, Corollary 16, and Corollary 15.

### 8.4. Proofs for perturbation expansions.

Proof of Lemma 10. We wish to construct the first and second derivatives of the map

$$\psi\left(M\left(t\right)\right) = v_{\perp}^{\mathsf{T}} R_{I}\left(t\right) L_{I}^{\mathsf{T}}\left(t\right) = v_{\perp}^{\mathsf{T}} P_{I}\left(t\right) + v_{\perp}^{\mathsf{T$$

where we have made the dependence on t explicit and M(0) = M. Then based on Sun (1991, Remark 4.2), the Fréchet and Gateaux derivatives coincide and we can write

$$\dot{\psi}_M\left(\hat{M}_n - M\right) = \upsilon_{\perp}^{\mathsf{T}} \dot{P}_I\left(t\right)|_{t=0},$$

which by the product rule implies

$$v_{\perp}^{\mathsf{T}}\dot{P}(0) = v_{\perp}^{\mathsf{T}}\dot{R}_{I}(0) L_{I}^{\mathsf{T}} + v_{\perp}^{\mathsf{T}}R_{I}\dot{L}_{I}^{\mathsf{T}}(0) = v_{\perp}^{\mathsf{T}}\dot{R}_{I}(0) L_{I}^{\mathsf{T}}$$
  
because  $v_{\perp}^{\mathsf{T}}R_{I}(0) = v_{\perp}^{\mathsf{T}}R_{I} = 0.$ 

*Proof of Corollary 13.* The result follows from Assumption 2(1) and the definition in (8.1).

Proof of Lemma 12. (1) By Lemma 10,

vec 
$$\hat{\Delta}_{n,\mathrm{IJ}} = \left( R_I^\mathsf{T} \otimes L_J^\mathsf{T} \right) \operatorname{vec} \left( \hat{M}_n - M \right)$$

The result follows by the continuous mapping theorem and Assumption 2.

(2) By the preceding argument, and the definition of  $S^{-1}$  in Lemma 10, we see that

$$v_{\perp}^{\mathsf{T}} R_{n,J} \mathbf{S}^{-1} \left( \hat{\Delta}_{n,\mathrm{IJ}} \right) L_{n,I}^{\mathsf{T}}$$

is a linear and continuous transformation of  $\hat{\Delta}_{n,IJ}$ . Vectorizing  $S^{-1}\left(\hat{\Delta}_{n,IJ}\right)$  and taking the expectation of the outer product provides the result.

(3) We construct an adjacent matrix and shall argue by continuity to establish the rank. In particular, if we can show that  $\Omega$  is of rank qm, we can argue by continuity to establish the result. For  $\Lambda_I \approx I_q$ , the matrix  $\Omega_W$  is close to  $\Sigma$ 

$$\Sigma := B_S \Omega_M B_S^{\mathsf{T}}$$
$$B_S := \left( L_I R_I^{\mathsf{T}} \otimes v_{\perp}^{\mathsf{T}} R_J \left( I_J - \Lambda_J \right)^{-1} L_J^{\mathsf{T}} \right)$$

We begin by showing that  $\operatorname{rk} v_{\perp}^{\mathsf{T}} R_J (I_J - \Lambda_J)^{-1} L_J^{\mathsf{T}} = m$ . By Thm. 2.4 in Gohberg, Lancaster, and Rodman (1982) we know that

$$M^{-1}(\mu) = R_I (I_I - \Lambda_I)^{-1} L_I^{\mathsf{T}} + R_J (I_J - \Lambda_J)^{-1} L_J^{\mathsf{T}}$$

with  $\mu \notin \mathcal{L}$ . Now, we wish to restrict ourselves to the subspace  $\mathcal{R}_I^{\perp} \coloneqq \operatorname{sp} R_I^{\perp}$  and hence consider

$$\boldsymbol{\upsilon}_{\perp}^{\mathsf{T}}\boldsymbol{M}^{-1}\left(\boldsymbol{\mu}\right) = \boldsymbol{\upsilon}_{\perp}^{\mathsf{T}}\boldsymbol{R}_{J}\left(\boldsymbol{I}_{J}\boldsymbol{\mu}-\boldsymbol{\Lambda}_{J}\right)^{-1}\boldsymbol{L}_{J}^{\mathsf{T}}.$$

Postmultiply by  $M(\mu)$  to obtain

$$\boldsymbol{v}_{\perp}^{\mathsf{T}} = \boldsymbol{v}_{\perp}^{\mathsf{T}} R_J \left( I_J \boldsymbol{\mu} - \boldsymbol{\Lambda}_J \right)^{-1} \boldsymbol{L}_J^{\mathsf{T}} \boldsymbol{M}(\boldsymbol{\mu}),$$

for  $\mu \notin \mathcal{L}$ . By continuity of M as  $\mu \to 1$  we can extend the allowable  $\mu$ . Now, since

$$\operatorname{rk}\left(v_{\perp}^{\mathsf{T}}R_{J}\left(I_{J}-\Lambda_{J}\right)^{-1}L_{J}^{\mathsf{T}}M(1)\right)$$

$$\leq \min\left\{\operatorname{rk}v_{\perp}^{\mathsf{T}}R_{J}\left(I_{J}\mu-\Lambda_{J}\right)^{-1}L_{J}^{\mathsf{T}},\operatorname{rk}M(1)\right\}$$

and  $\operatorname{rk} M(1) \geq m$  and as a result of the preceding displayed equation we arrive at

$$\operatorname{rk}\left(v_{\perp}^{\mathsf{T}}R_{J}\left(I_{J}-\Lambda_{J}\right)^{-1}L_{J}^{\mathsf{T}}M(1)\right)=m.$$

We have  $\operatorname{rk} v_{\perp}^{\mathsf{T}} R_J (I_J \mu - \Lambda_J)^{-1} L_J^{\mathsf{T}} \geq m$ . Finally, we observe that

$$\operatorname{rk} v_{\perp}^{\mathsf{T}} R_J \left( I_J \mu - \Lambda_J \right)^{-1} L_J^{\mathsf{T}} \le m$$

because it is  $m \times p - q$  and  $m \le p - q$ . Now, under Assumption 2,  $\Omega_M$  has full rank and the result follows.

# Proof of Lemma 11.

(1) We Taylor-expand

$$\sqrt{n}v_{\perp}^{\mathsf{T}}\hat{P}_{n,I} = \sqrt{n}\psi\left(\hat{M}_{n}\right) \\
=_{(2)}\sqrt{n}\left(\psi\left(\hat{M}_{n}\right) - \psi\left(M\right)\right) \\
= \sqrt{n}\psi\left(\hat{M}_{n} - M\right) + \sqrt{n}r\left(\hat{M}_{n} - M\right) \\
=_{(3)}\sqrt{n}v_{\perp}^{\mathsf{T}}R_{n,J}\boldsymbol{S}^{-1}\left(\hat{\Delta}_{n,\mathrm{IJ}}\right)L_{n,I}^{\mathsf{T}} + \sqrt{n}r\left(\hat{M}_{n} - M\right)$$

where  $=_{(2)}$  follows from application of the null hypothesis and  $=_{(3)}$  follows from Lemma 10.

(2) We obtain a remainder term

$$r\left(\hat{M}-M\right) \coloneqq \psi\left(\hat{M}\right) - \psi\left(M\right) - \dot{\psi}\left(\hat{M}-M\right).$$

To ensure that the first order term dominates the remainder, we need to verify that standardisation by  $\sqrt{n}$  does not cause dominant second order terms. By Assumption 2,  $\|\hat{M} - M\| = O_p(n^{-1/2})$ . By Taylor's theorem,  $r(\hat{M} - M) = O(n^{-1/2})$ . For simplicity, we can treat  $(\hat{M}_n - M)$  as a deterministic sequence and then apply Vaart (2000, Lemma 2.12, Ch. 2) to obtain the relevant statistical result. First, recognize that  $S^{-1}(a_n M) = a_n S^{-1}(M)$  for any scalar sequence  $a_n$  and admissible argument M. Then, rewrite r as the sum of the next higher order term and another unspecified remainder term of known order via application of Lemma 10:

$$r\left(\hat{M}_n - M\right) = 2\upsilon_{\perp}^{\mathsf{T}} R_J \boldsymbol{S}^{-1} \left( \left( \hat{\Delta}_{n,J} V - V \hat{\Delta}_{n,I} \right) \right) L_I^{\mathsf{T}} + r_2 \left( \hat{M}_n - M \right),$$

where  $r_2$  collects higher order terms. By Corollary 13 and Lemma 10, we obtain the tighter bound  $n^{1/2}r\left(\hat{M}_n - M\right) = O\left(n^{-1/2}\right)$ , which implies that normalization by  $\sqrt{n}$  is innocuous and does not affect convergence.

Proof of Lemma 14. Add and subtract  $v_{\perp}^{\mathsf{T}}$  to obtain  $\left(\left[\hat{D}_{n}^{\mathsf{T}} I_{r}\right] \pm \left[D^{\mathsf{T}} I_{r}\right]\right) R_{n,I} = 0$ . Therefore,

$$\left(\hat{D}_{n}^{\mathsf{T}}-D^{\mathsf{T}}\right)\hat{R}_{n,2,I}=\left[\hat{D}_{n}^{\mathsf{T}}\quad I_{r}\right]\left(\hat{R}_{n,I}-R_{I}R_{2,I}^{-1}\hat{R}_{n,2,I}\right),$$

whence the result follows.

*Proof of Corollary 16.* In Lemma 10, replace  $L_I^{\mathsf{T}}$  by  $R_{2,I}^{-1}$  and the result follows.

*Proof of Corollary 15.* Apply Corollary 16 to the expression in Lemma 14 and the result follows.

8.5. Perturbation of invariant subspaces. To prove Lemma 10, we find an expansion of basis vectors of invariant subspaces of a matrix in response to a small perturbation of that matrix. To aid with the general exposition of the idea behind invariant subspaces and associated perturbations, we adopt the notation of Sun (1991). Similar results to those considered here are available in Kato (1995, Ch. 2).

Let  $X_1 \in \mathbb{R}^{p \times q}$  with  $\operatorname{rk} X_1 = q$ ,  $X_1^{\mathsf{T}} X_1 = I_q$  and  $AX_1 = X_1 A_1$  for some matrix  $A_1 \in \mathbb{R}^{q \times q}$ . Then sp  $X_1$  is an invariant subspace of  $A \in \mathbb{R}^{p \times p}$  if and only if there exists a non-singular matrix  $X = [X_1 X_2] \in \mathbb{R}^{p \times p}$  with  $X_2^{\mathsf{T}} X_2 = I_{p-q}$  such that

$$X^{-1}AX = \begin{bmatrix} A_{11} & A_{12} \\ 0 & A_{22} \end{bmatrix}, A_{11} \in \mathbb{R}^{q \times q}.$$

We now introduce the perturbation matrix of A,

$$C = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}.$$

The block  $C_{21}$  measures the departure from B being upper-triangular. Letting  $t \in \mathbb{C}$  we perturb A and obtain A(t) := A + tC. For the purposes of this study, we are interested in the span of the vectors that are orthogonal to the columns of  $X_1$ . Recall the operator (3.9),

$$\boldsymbol{S}\left(Q\right) = QA_{11} - A_{22}Q$$

and  $Q \in \mathbb{R}^{(p-q) \times q}$ . Then, if  $A_{11}$  and  $A_{22}$  do not share any eigenvalues, Theorem 2.1 in Sun (1991) applies. Introducing the parameter  $t \in \mathbb{C}$  to help us keep track of the order of the expansion. We state Sun (1991, Thm. 1):

**Theorem 17.** There exists a unique q-dimensional invariant subspace sp  $X_1(t)$  of A(t)  $(t \in \mathbb{C})$  such that sp  $X_1(0) = \text{sp } X_1$  and the basis vectors (columns of  $X_1$ ) may be defined to be analytic functions of t in some neighbourhood of the origin of  $\mathbb{C}$ . Further, the analytic matrix-valued function  $X_1(t)$  has the second order perturbation expansion

(8.5) 
$$X_{1}(t) = X_{1} + X_{2} \mathbf{S}^{-1}(C_{21}) t + X_{2} \mathbf{S}^{-1}(C_{22} \mathbf{S}^{-1}(C_{21}) - \mathbf{S}^{-1}(C_{21}) C_{11} - \mathbf{S}^{-1}(C_{21}) A_{12} \mathbf{S}^{-1}(C_{21}) t^{2} + O(t^{3}).$$

We now interpret Theorem 17 to make it amenable for our statistical analysis. Expression (8.5) shows how small perturbations in the matrix A affect the column space of  $X_1$ , which correspond to the first and second-order derivatives. The equivalents of A and  $X_1$  are  $\Lambda$  and  $R_I$  respectively. Furthermore,  $A_{11} := \Lambda_I$  and  $A_{22} := \Lambda_J$ , so that by Assumption 2.7,  $A_{11}$  and  $A_{22}$  have no eigenvalues in common.

### References

Anderson, Theodore W. (1963). "Asymptotic theory for principal component analysis". In: The Annals of Mathematical Statistics 34.1, pp. 122–148.

- Anderson, Theodore W. (2010). "The LIML estimator has finite moments!" In: Journal of Econometrics 157.2, pp. 359-361. ISSN: 0304-4076. DOI: 10.1016/j.jeconom.2010.02.004. URL: http://www.sciencedirect.com/science/article/pii/S0304407610000801.
- Bonacich, Phillip (1987). "Power and centrality: A family of measures". In: American Journal of Sociology 92.5, pp. 1170–1182.
- Bura, E and R Pfeiffer (2008). "On the distribution of the left singular vectors of a random matrix and its applications". In: *Statistics & Probability Letters* 78.15, pp. 2275–2280.
- Cai, Changxiao et al. (2019). "Subspace Estimation from Unbalanced and Incomplete Data Matrices: Statistical Guarantees". In: *arXiv preprint arXiv:1910.04267*.
- Chen, Yuxin, Chen Cheng, and Jianqing Fan (2020). Asymmetry Helps: Eigenvalue and Eigenvector Analyses of Asymmetrically Perturbed Low-Rank Matrices. arXiv: 1811.12804 [math.ST].
- Cheng, Chen, Yuting Wei, and Yuxin Chen (2021). Tackling small eigen-gaps: Fine-grained eigenvector estimation and inference under heteroscedastic noise. arXiv: 2001.04620 [math.ST].
- Davis, AW (1977). "Asymptotic Theory for Principal Component Analysis: Non-normal Case". In: Australian Journal of Statistics 19.3, pp. 206–212.
- Duffy, James A. and Jerome R. Simons (2020). The Cointegrated VAR without Unit Roots: Representation Theory and Asymptotics. arXiv: 2002.08092 [econ.EM].
- Girko, V. L. (1999). "Strong Law for the singular values and eigenvectors of random matrices III. Inequalities for the spectral radius of large random matrices". In: *Random Operators and Stochastic Equations* 7.2, pp. 177-200. DOI: https://doi.org/10.1515/rose.1999.7.2.177.
  URL: https://www.degruyter.com/view/journals/rose/7/2/article-p177.xml.
- Gohberg, I, P. Lancaster, and L. Rodman (1982). *Matrix Polynomials*. Society for Industrial and Applied Mathematics. URL: 341000/082996b96e33bfa23b3ed8decb058625.
- Greenbaum, Anne, Ren-cang Li, and Michael L. Overton (Mar. 2, 2019). "First-order Perturbation Theory for Eigenvalues and Eigenvectors". Version 1. In: arXiv: http://arxiv.org/abs/1903.00785v2 [n URL: http://arxiv.org/abs/1903.00785v2.
- James, Alan T. (1964). "Distributions of Matrix Variates and Latent Roots Derived from Normal Samples". In: The Annals of Mathematical Statistics, pp. 475–501.
- (1977). "Tests for a prescribed subspace of principal components". In: Multivariate Analysis IV, Ed. PR Krishnaiah, North-Holland: Amsterdam, pp. 73–77.
- Johnstone, Iain M, Alexei Onatski, et al. (2020). "Testing in high-dimensional spiked models". In: Annals of Statistics 48.3, pp. 1231–1254.
- Johnstone, Iain M. (2001). "On the Distribution of the Largest Eigenvalue in Principal Components Analysis". In: *The Annals of Statistics* 29.2, pp. 295–327. ISSN: 00905364. URL: http://www.jstor.org/s
- Kato, Tosio (1995). *Perturbation Theory for Linear Operators*. 2nd. Springer-Verlag Berlin and Heidelberg.
- Koltchinskii, Vladimir, Matthias Löffler, Richard Nickl, et al. (2020). "Efficient estimation of linear functionals of principal components". In: *The Annals of Statistics* 48.1, pp. 464–490.
- Lee, Ji Oon and Kevin Schnelli (Dec. 2016). "Tracy–Widom distribution for the largest eigenvalue of real sample covariance matrices with general population". In: Ann. Appl. Probab. 26.6, pp. 3786–3839. DOI: 10.1214/16-AAP1193. URL: https://doi.org/10.1214/16-AAP1193.
- Penna, F., R. Garello, and M. A. Spirito (2009). "Cooperative spectrum sensing based on the limiting eigenvalue ratio distribution in wishart matrices". In: *IEEE Communications Letters* 13.7, pp. 507–509.

- Silin, Igor and Jianqing Fan (2020). "Hypothesis testing for eigenspaces of covariance matrix". In: *arXiv preprint arXiv:2002.09810*.
- Stewart, G. W. and Ji-guang Sun (1991). Matrix perturbation theory. Vol. 33. Academic Press, p. 74. ISBN: 9781466507289. DOI: 10.1016/0378-4754(91)90038-5.
- Sun, Ji-guang (1991). "Perturbation expansions for invariant subspaces". In: *Linear Algebra And Its Applications*.
- Takemura, Akimichi and Yo Sheena (2005). "Distribution of eigenvalues and eigenvectors of Wishart matrix when the population eigenvalues are infinitely dispersed and its application to minimax estimation of covariance matrix". In: Journal of Multivariate Analysis 94.2, pp. 271– 299.
- Tracy, Craig A. and Harold Widom (1994). "Level-spacing distributions and the Airy kernel". In: Comm. Math. Phys. 159.1, pp. 151–174. URL: https://projecteuclid.org:443/euclid.cmp/1104254495.
- Tyler, David E. (1981). "Asymptotic Inference for Eigenvectors". In: *The Annals of Statistics* 9, pp. 725–736. ISSN: 0090-5364. DOI: 10.1214/aos/1176345514.
- (1983). "A Class of Asymptotic Tests for Principal Component Vectors". In: *The Annals of Statistics* 11.4, pp. 1243–1250. DOI: 10.1214/aos/1176346337. URL: http://dx.doi.org/10.1214/aos/1176346337.
- Vaart, Aad W. van der (2000). *Asymptotic Statistics*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press.
- Wigner, Eugene P (1958). "On the distribution of the roots of certain symmetric matrices". In: Annals of Mathematics, pp. 325–327.
- Zanella, Alberto, Marco Chiani, and Moe Z Win (2009). "On the marginal distribution of the eigenvalues of Wishart matrices". In: *IEEE Transactions on Communications* 57.4, pp. 1050–1060.