NONLOCAL ENERGY FUNCTIONALS AND DETERMINANTAL POINT PROCESSES ON NON-SMOOTH DOMAINS

ZHENGJIANG LIN

ABSTRACT. Given a nonnegative integrable function J on \mathbb{R}^n , we relate the asymptotic properties of the nonlocal energy functional

$$\int_{\Omega} \int_{\Omega^c} J\left(\frac{x-y}{t}\right) \, dx dy$$

as $t \to 0^+$ with the boundary properties of a given domain $\Omega \subset \mathbb{R}^n$. Then, we use these asymptotic properties to study the fluctuations of many determinantal point processes, and show that their variances measure the Minkowski dimension of $\partial \Omega$.

1. INTRODUCTION

A point process on \mathbb{R}^n is defined as a measurable map from a probability space $(\Omega, \mathcal{B}(\Omega), \mathcal{P})$ to random configurations of points $\{x_i\}_{i \in I}$, where I is a countable index set. More precisely, a point process on \mathbb{R}^n is a probability measure on the set of locally finite counting measures on \mathbb{R}^n . Usually, a point process X takes the form

(1.0.1)
$$X = \sum_{i \in I} \delta_{x_i},$$

where δ_x is the dirac measure at point $x \in \mathbb{R}^n$, and the configurations of x_i 's follow a probability distribution. Locally finiteness means that, almost surely, $X(K) < \infty$ for any compact set $K \subset \mathbb{R}^n$. We will define this precisely in Section 4 and readers can also see [1] for more details. We define $X(\varphi) \equiv \sum_{i \in I} \varphi(x_i)$ for any appropriate function φ , provided that the summation on the right-hand side is well-defined. In particular, if $A \subset \mathbb{R}^n$ is a measurable set, we define $X(A) \equiv X(\mathbb{1}_A)$ to denote the number of x_i 's that are contained in the set A. Here, $\mathbb{1}_A$ is the indicator function of A.

We are interested in the variance $\operatorname{Var}(X(\varphi))$. In most of the models from either physics or probability, there is a nonnegative integrable function $J = J_X$ on \mathbb{R}^n , depending on the point process X, such that

(1.0.2)
$$\operatorname{Var}(X(\varphi)) = \int_{\mathbb{R}^n \times \mathbb{R}^n} |\varphi(x) - \varphi(y)|^2 \cdot J(x-y) \, dx dy.$$

Such functionals have already been extensively studied in analysis and geometry to investigate the asymptotic behaviors of a particular family of $\{J_t\}$'s as t converges to a critical

Date: June 16, 2023.

number in \mathbb{R} or ∞ . For example, in the fractional Sobolev spaces $W^{t,p}(\mathbb{R}^n)$ (or $W^{t,p}(A)$ with A being a smooth bounded domain in \mathbb{R}^n), a measurable function φ is said to be in $W^{t,p}(A)$ if the functional

(1.0.3)
$$||\varphi||_{W^{t,p}}^{p} = \int_{A} \int_{A} \frac{|\varphi(x) - \varphi(y)|^{p}}{|x - y|^{n + tp}} \, dx dy$$

is finite. One would like to see the asymptotic behaviors (orders) of such functionals when $t \to 1^-$, which diverges for smooth nonconstant φ . For further discussions on $W^{t,p}$ -spaces, one may refer to [2], where the authors associated the limit of the functional (1.0.2) for a general family of $\{J_t\}$ to the $W^{1,2}$ -norm of φ . We will also briefly discuss the Sobolev spaces in Section 3.2.

When $\varphi = \mathbb{1}_{\Omega}$ is the indicator function of a measurable set $\Omega \subset \mathbb{R}^n$, the functional (1.0.2) is closely related to the study of the nonlocal perimeter and nonlocal minimal surfaces [4, 17]. In this manuscript, we also focus on situations where $\varphi = \mathbb{1}_{\Omega}$. Our primary purpose is to study the asymptotic behavior (order) of the functional

(1.0.4)
$$\mathcal{J}_t(\Omega) \equiv \frac{1}{2} \int_{\mathbb{R}^n \times \mathbb{R}^n} \left| \mathbb{1}_{\Omega}(x) - \mathbb{1}_{\Omega}(y) \right|^2 \cdot J\left(\frac{x-y}{t}\right) \, dx dy. = \int_{\Omega} \int_{\Omega^c} J\left(\frac{x-y}{t}\right) \, dx dy,$$

as $t \to 0^+$. Intuitively, if J has a fast-decaying tail, then as $t \to 0^+$, only those pairs (x, y) with $\operatorname{dist}(x, y) \sim t$, or equivalently, near the boundary $\partial\Omega$ within a distance $\sim t$, will contribute to the first order term in $\mathcal{J}_t(\Omega)$. We note that although $\mathbb{1}_{\Omega}$ is not weakly differentiable, it can be in the space of functions of bounded total variation (BV space). This functional space includes indicator functions of bounded domains with Lipschitz boundaries. The asymptotic behavior for functionals with BV test functions was also studied in [6], which we will also discuss in Section 3.1.

In addition to the domains with "locally flat" boundaries mentioned above, one of our novel parts is that we will also discuss cases when $\partial\Omega$ does not locally "look like" a hyperplane, including the well-known Koch snowflake (Figure 1) and a family of fractals constructed in a similar way as the standard Koch snowflake. Also, for general $\Omega \subset \mathbb{R}^n$ whose boundary $\partial\Omega$ has Minkowski dimension (or Hausdorff dimension) larger than n-1, we can give a way to quantify the rate of decay with respect to t, i.e., we will show that $\mathcal{J}_t(\Omega) \sim t^{\beta}$, where β depends on the Minkowski dimension of $\partial\Omega$ and n. Moreover, if the boundary $\partial\Omega$ is selfsimilar in some way, it is possible to show the existence of the limit $\lim_{t\to 0^+} \mathcal{J}_t(\Omega)/t^{\beta}$. See our Theorem 1.1.1 and Theorem 1.1.3.

Our technical sections, Section 2 and Section 3, do not rely on any specific probability models. In Section 1.1, we will present our main results without referring to probability. And we will present the main results in probability language in Section 1.2. At the end of Section 1.2, we will briefly discuss those J which can possibly change signs and give an example in Remark 1.2.3. At the end of this manuscript, Section 4, we will give the proofs of the results in Section 1.2 for the complex Ginibre ensemble.

1.1. Results on Nonlocal Energy Functionals.

Recall that for a bounded measurable set $A \subset \mathbb{R}^n$, we define the α -dimensional upper Minkowski content of A as

(1.1.1)
$$\overline{\mathcal{M}}^{\alpha}(A) \equiv \limsup_{t \to 0^+} \frac{|B(A,t)|}{t^{n-\alpha}},$$

and the α -dimensional lower Minkowski content of Ω as

(1.1.2)
$$\underline{\mathcal{M}}^{\alpha}(A) \equiv \liminf_{t \to 0^+} \frac{|B(A,t)|}{t^{n-\alpha}}$$

Here,

(1.1.3)
$$B(A,t) \equiv \{x \in \mathbb{R}^n \mid \operatorname{dist}(x,A) = \inf_{y \in A} \operatorname{dist}(x,y) < t\},\$$

and |B(A,t)| is the Lebesgue measure (volume) of B(A,t). The upper and lower Minkowski dimensions of A are defined by

(1.1.4)
$$\overline{\dim}_{\mathcal{M}}(A) = \sup\{\alpha \ge 0 \mid \overline{\mathcal{M}}^{\alpha}(A) = \infty\} = \inf\{\alpha \ge 0 \mid \overline{\mathcal{M}}^{\alpha}(A) = 0\},$$

and

(1.1.5)
$$\underline{\dim}_{\mathcal{M}}(A) = \sup\{\alpha \ge 0 \mid \underline{\mathcal{M}}^{\alpha}(A) = \infty\} = \inf\{\alpha \ge 0 \mid \underline{\mathcal{M}}^{\alpha}(A) = 0\}.$$

Now, we assume that $\Omega \subset \mathbb{R}^n$ is a bounded measurable set with $|\Omega| > 0$, and with a topological boundary $\partial\Omega$ such that

(1.1.6)
$$0 < M_2 \leq \underline{\mathcal{M}}^{\alpha}(\partial \Omega) \leq \overline{\mathcal{M}}^{\alpha}(\partial \Omega) \leq M_1 < \infty,$$

where M_1, M_2 are two positive constants and $\alpha \in [n-1, n]$. This actually implies that $\partial \Omega$ is of Minkowski dimension α .

For a nonnegative function $J \in L^1(\mathbb{R}^n)$, let use consider the functional

(1.1.7)
$$\mathcal{J}_t(\Omega) \equiv \frac{1}{2} \int_{\mathbb{R}^n \times \mathbb{R}^n} \left| \mathbb{1}_{\Omega}(x) - \mathbb{1}_{\Omega}(y) \right|^2 \cdot J\left(\frac{x-y}{t}\right) \, dx dy. = \int_{\Omega} \int_{\Omega^c} J\left(\frac{x-y}{t}\right) \, dx dy,$$

where $\Omega^c = \mathbb{R}^n \setminus \Omega$. For all theorems in Section 1.1, we will assume that $\int_{\mathbb{R}^n} J(z) \cdot |z|^n dz < \infty$. Notice that this together with the assumption that $J \in L^1(\mathbb{R}^n)$ shows that there is a constant $C_J > 0$, such that for any $\gamma \in [0, n]$, $\int_{\mathbb{R}^n} J(z) \cdot |z|^{\gamma} dz \leq C_J < \infty$.

Theorem 1.1.1. For the kernel J, we also assume that there are two positive constants c_J, a_J such that when $|x| \leq a_J$, $J(z) \geq c_J > 0$. For the domain Ω , we further assume that there is a positive constant $D_{\partial\Omega}$, such that for all $x \in \partial\Omega$ and all $t \in (0, 1)$, $\min\{|B(x, t) \cap \Omega|, |B(x, t) \cap \Omega^c|\} \geq D_{\partial\Omega} \cdot t^n$. Then, there is a constant O_1 depending on n, C_J , and a constant O_2 depending on $n, a_J, c_J, D_{\partial\Omega}$, such that

(1.1.8)
$$0 < O_2 \cdot M_2 \le \liminf_{t \to 0^+} \frac{\mathcal{J}_t(\Omega)}{t^{2n-\alpha}} \le \limsup_{t \to 0^+} \frac{\mathcal{J}_t(\Omega)}{t^{2n-\alpha}} \le O_1 \cdot M_1 < \infty.$$

We remark that the density lower bounds condition for points on $\partial\Omega$, i.e., the existence of $D_{\partial\Omega}$, is satisfied by a class of domains known as NTA (nontangentially accessible) domains, which encompass domains with Lipschitz boundaries, quasiballs, and many self-similar fractals. This concept was first introduced by Jerison and Kenig in [13], and one can also find more related domains in the recent survey paper [24].

For sets of finite perimeter (also known as Caccioppoli sets; see Definition 3.1.2), the limit was explicitly computed in [6]. Let us cite a theorem in [6]. In particular, sets of finite perimeter include bounded domains with Lipschitz boundaries.

Theorem 1.1.2 ([6], Theorem 1). If Ω is a set of finite perimeter and J(z) = J(|z|) is radially symmetric, then there is a positive dimensional constant K(n), such that

(1.1.9)
$$\lim_{t \to 0^+} \frac{\mathcal{J}_t(\Omega)}{t^{n+1}} = 2K(n) \cdot ||(J(z) \cdot |z|)||_{L^1(\mathbb{R}^n)} \cdot \mathcal{H}^{n-1}(\partial^*\Omega).$$

Here, K(n) is explicitly computed in (3.1.5), and \mathcal{H}^{n-1} denotes the (n-1)-dimensional Hausdorff measure (or surface measure for smooth hypersurfaces). $\partial^*\Omega$ is called the reduced boundary of Ω . By Federer's theorem (see, for example, Theorem 16.2 in [16]), up to a set of zero (n-1)-dimensional Hausdorff measure, this reduced boundary $\partial^*\Omega$ equals the essential boundary $\partial^e\Omega$ of Ω , which is

(1.1.10)
$$\partial^e \Omega \equiv \mathbb{R}^n \setminus (\Omega^{(0)} \cup \Omega^{(1)}),$$

where we define the set of points of density β of Ω for $\beta \in [0, 1]$ as

(1.1.11)
$$\Omega^{(\beta)} \equiv \left\{ x \in \mathbb{R}^n \ \left| \ \lim_{r \to 0^+} \frac{|\Omega \cap B(x,r)|}{|B(x,r)|} = \beta \right\} \right\}$$

See more discussions on BV functions and sets of finite perimeter in Section 3.1.

The last theorem is about a family of self-similar fractals constructed in as similar way as the standard Koch snowflake, which we call snowflakes of scales η . See more detailed discussions on Koch snowflakes in Section 3.3, Figure 1, Figure 2, and Figure 3.

Theorem 1.1.3. Let $\Omega = \Omega(\eta) \subset \mathbb{R}^2$ be the Koch snowflake of a scale $\eta > 1$ with η satisfying the condition (3.3.14), then the limit

(1.1.12)
$$\lim_{t \to 0^+} \frac{\mathcal{J}_t(\Omega)}{t^{4-\alpha(\eta)}}$$

exists, where $\alpha(\eta) \in (1,2)$ is the Minkowski dimension of $\partial\Omega$ and satisfies (3.3.1).

Combining Theorem 1.1.3 with Theorem 1.1.1, the limit in Theorem 1.1.3 is also comparable to the Minkowski contents of $\partial\Omega$. We also remark that this Theorem 1.1.3 is an illustration on how to establish the existence of such a limit, which can be extended to other domains with self-similar fractal boundaries. Moreover, the algebraic condition (3.3.14) is satisfied by almost all $\eta > 1$ and hence $\alpha(\eta)$ can potentially take on almost any value in (1,2) in Theorem 1.1.3. And for other constructions of self-similar fractal boundaries, one can obtain similar algebraic conditions like (3.3.14). The union of all those possible $\alpha(\eta)$ will be the whole (1,2). See more discussions in Section 3.3.

Finally, we note that in the functional $\mathcal{J}_t(\Omega)$, we considered a double integral over a domain Ω and its complement Ω^c . In Section 2 and some parts of Section 3, we will present the lemmas and proofs for double integrals over two disjoint domains Ω_1 and Ω_2 , with similar assumptions near $\partial \Omega_1 \cap \partial \Omega_2$. This provides a more general statement of our results.

1.2. Results on Fluctuations of Determinantal Point Processes.

This section contains the applications of results in Section 1.1 on determinantal point processes, which was first introduced by Macchi [15] and originally called the fermion point process. Let X be a determinantal point process on \mathbb{R}^n , and let A_1, A_2, \ldots, A_p be disjoint measurable sets in \mathbb{R}^n , then

(1.2.1)
$$\mathcal{E}\left[X(A_1)X(A_2)\cdots X(A_p)\right] = \int_{A_1\times A_2\cdots\times A_p} \det\left(K(x_i, x_j)\right)_{1\le i,j\le p} d\mu(x_1)\cdots d\mu(x_p).$$

Here, $\mathcal{E}(\cdot)$ is the expectation, μ is a reference probability measure on \mathbb{R}^n , K is a measurable function, and for many physical models, the matrix $(K(x_i, x_j))_{1 \le i,j \le p}$ is Hermitian for any $p \in \mathbb{Z}_+$. Such a determinantal point process X is called a determinantal point process with kernel K with respect to μ .

Typical examples for determinantal point processes on $\mathbb{R}^2 = \mathbb{C}$ can come from random matrix theory, where we usually consider $X_N = \sum_{i=1}^N \delta_{x_i}$ with x_i 's being eigenvalues of $N \times N$ random matrices. For example, the Gaussian unitary ensemble (GUE) and the complex Ginibre ensemble [1].

On the other hand, given a function K and a reference measure μ on \mathbb{R}^n , if K defines a self-adjoint integral operator from $L^2(\mathbb{R}^n, d\mu)$ to $L^2(\mathbb{R}^n, d\mu)$, and is locally trace class with all eigenvalues in [0, 1], then there exists a determinantal point process, uniquely in law, with such a K satisfying (1.2.1), see Corollary 4.2.21 of [1]. In [19], readers can see three models following this logic. For determinantal point processes on higher dimensional \mathbb{R}^n , readers can see [25], where the authors called them Fermi-shells point processes. Soshnikov [23] discussed many models of determinantal point processes from random matrix theory, statistical mechanics, and quantum mechanics.

For most of determinantal point processes, including those aforementioned models, the variance has a more concise formula: by (1.2.1), for any appropriate test function φ , it has the form

(1.2.2)
$$\operatorname{Var}(X(\varphi)) = \frac{1}{2} \int_{\mathbb{R}^n \times \mathbb{R}^n} |\varphi(x) - \varphi(y)|^2 \cdot |K(x,y)|^2 \, d\mu(x) d\mu(y).$$

In particular, for some of the aforementioned models, such as the planar model in [19], and the Fermi-shells point processes in [25], it can be shown that $\frac{d\mu(x)}{dx} = \xi(x) \in L^1(\mathbb{R}^n, dx)$ is nonnegative, and $|K(x,y)|^2 \xi(x)\xi(y)$ is a function of x - y, i.e., has the form J(x - y) for a nonnegative integrable function J. Here, $\frac{d\mu(x)}{dx} = \xi(x)$ is the Radon-Nikodym derivative of μ

with respect to the Lebesgue measure dx on \mathbb{R}^n . The functionals then become

(1.2.3)
$$\int_{\mathbb{R}^n \times \mathbb{R}^n} |\varphi(x) - \varphi(y)|^2 \cdot J(x-y) \, dx dy,$$

for which we discussed the asymptotic orders of $\mathcal{J}_t(\Omega)$ for the case $\varphi = \mathbb{1}_{\Omega}$ in Section 1.1. On the other hand, the asymptotic behaviors of the variances for a family of $(X_t, \varphi_t = \mathbb{1}_{\Omega_t})$ will imply some forms of central limit theorems. Roughly speaking, under some mild assumptions on those determinantal processes X_t , if one can further obtain that $\operatorname{Var}(X_t(\Omega_t)) \to \infty$ as the parameter t tending to some number in \mathbb{R} or ∞ , then $[X_t(\Omega_t) - \mathcal{E}(X_t(\Omega_t))]/\sqrt{\operatorname{Var}(X_t(\Omega_t))}$ converges in law to the normal distribution. See [5], Theorem 8 of [23], or Theorem 4.2.25 of [1]. We then can obtain the following direct corollaries.

Corollary 1.2.1. Let X be a determinantal point process on \mathbb{R}^n with kernel K with respect to μ , such that $|K(x,y)|^2 \xi(x)\xi(y) = J(x-y)$ is a nonnegative integrable function, where $\xi(x)$ is the derivative of μ with respect to dx on \mathbb{R}^n . Then we have the following cases:

(i) Assume that Ω is a set of finite perimeter in \mathbb{R}^n and J(x) = J(|x|) is radially symmetric. Then, there is a computable positive constant $T = T(n, ||(J(x) \cdot |x|))||_{L^1(\mathbb{R}^n, dx)})$, such that

(1.2.4)
$$\lim_{t \to 0^+} t^{n-1} \cdot \operatorname{Var}(X(\Omega/t)) = T \cdot \mathcal{H}^{n-1}(\partial^*\Omega).$$

Here, $\Omega/t \equiv \{x/t \in \mathbb{R}^n \mid x \in \Omega\}.$

(ii) Assume that $\partial\Omega$ is of Minkowski dimension $\alpha \in [n-1,n]$ and satisfies (1.1.6). We also assume that J and Ω satisfy the assumptions we made for Theorem 1.1.1, then

(1.2.5)
$$0 < \liminf_{t \to 0^+} \left[t^{\alpha} \cdot \operatorname{Var}(X(\Omega/t)) \right] \le \limsup_{t \to 0^+} \left[t^{\alpha} \cdot \operatorname{Var}(X(\Omega/t)) \right] < +\infty$$

Moreover, $t^{\alpha} \cdot \operatorname{Var}(X(\Omega/t))$ is comparable to the Minkowski contents of $\partial \Omega$.

(iii) When n = 2, assume that Ω is the Koch snowflake of a scale $\eta > 1$ with η satisfying the condition (3.3.14), then the limit

(1.2.6)
$$\lim_{t \to 0^+} t^{\alpha(\eta)} \cdot \operatorname{Var}(X(\Omega/t))$$

exists, where $\alpha(\eta)$ is the Minkowski dimension of $\partial\Omega$. Moreover, the limit is comparable to the Minkowski contents of $\partial\Omega$.

In all these three cases, we then obtain that the normalization of $X(\Omega/t)$ converges in law to the normal distribution.

Corollary 1.2.1 is a direct result according to those theorems in Section 1.1 and the aforementioned central limit theorems results in [5, 23, 1]. Also, part (iii) of Corollary 1.2.1 also holds true for some other domains Ω with self-similar fractal boundaries $\partial \Omega$ in \mathbb{R}^n . See the discussions in Section 3.3 and the proofs of our Theorem 3.3.1 Theorem 3.3.2.

In Corollary 1.2.1, the condition that $|K(x,y)|^2 \xi(x)\xi(y)$ has a form of J(x-y) is satisfied in the planar model in [19] and the Fermi-shells point processes in [25]. And in both cases, J(x-y) = J(|x-y|) is radially symmetric. On the other hand, if X_N is the determinantal point process associated with the complex Ginibre emsemble on $\mathbb{C} = \mathbb{R}^2$, the corresponding $|K_N(x,y)|^2 \xi_N(x) \xi_N(y)$ is not a function of x-y anymore. Instead, as $N \to +\infty$, $|K_N(x,y)|^2 \xi_N(x) \xi_N(y)$ will tend to the planar model in [19] when |x| < 1, |y| < 1. See more detailed discussions in Section 4.

Corollary 1.2.2. Let X_N be the determinantal point process associated the complex Ginibre ensemble and let $\Omega \subset \mathbb{C} = \mathbb{R}^2$ be a domain whose closure $\overline{\Omega}$ is contained in the unit ball $B(0,1) \equiv \{x \in \mathbb{R}^2 \mid |x| < 1\}$. Then we have the following cases:

(i) Assume that Ω is a set of finite perimeter, then there is a computable positive constant T such that

(1.2.7)
$$\lim_{N \to \infty} \frac{\operatorname{Var}(X_N(\Omega))}{\sqrt{N}} = T \cdot \mathcal{H}^1(\partial^* \Omega).$$

(ii) Assume that Ω satisfies the assumptions we made for Theorem 1.1.1 and $\partial \Omega$ is of Minkowski dimension $\alpha \in [1, 2]$ and satisfies (1.1.6), then

(1.2.8)
$$0 < \liminf_{N \to +\infty} \frac{\operatorname{Var}(X_N(\Omega))}{N^{(\alpha/2)}} = \limsup_{N \to +\infty} \frac{\operatorname{Var}(X_N(\Omega))}{N^{(\alpha/2)}} < +\infty$$

Moreover, $\frac{\operatorname{Var}(X_N(\Omega))}{N^{(\alpha/2)}}$ is comparable to the Minkowski contents of $\partial\Omega$. (iii) If Ω is the Koch snowflake of a scale $\eta > 1$ with η satisfying the condition (3.3.14), then the limit

(1.2.9)
$$\lim_{i \to \infty} \frac{\operatorname{Var}(X_N(\Omega))}{N^{(\alpha(\eta)/2)}}$$

exists, where $\alpha(\eta)$ is the Minkowski dimension of $\partial\Omega$. Moreover, the limit is comparable to the Minkowski contents of $\partial \Omega$.

In all these three cases, we then obtain that the normalization of $X_N(\Omega)$ converges in law to the normal distribution.

The proof of Corollary 1.2.2 will be shown in Section 4. In Section 4, we will also consider the limit of $\operatorname{Var}(X_N(\varphi))$ as $N \to +\infty$ when $\varphi \in W_0^{1,2}(B(0,\lambda))$ with $\lambda \in (0,1)$, which is a partial generalization of the results in [20]. It would also be intersting to build up similar central limit theorems for $\varphi \in W_0^{1,2}(\mathbb{R}^2)$, but we do not discuss further along this direction in this manuscript for compactness.

Remark 1.2.3. There are other point processes where our methods can also apply, but their variances may not have a nonnegative two-point function J in the functional

(1.2.10)
$$\mathcal{J}_t(\Omega) = \int_{\Omega} \int_{\Omega^c} J\left(\frac{x-y}{t}\right) \, dx dy$$

For example, the zeros of the Gaussian entire function [3] become a point process. The function is defined by $f(z) = \sum_{p=0}^{\infty} a_p e_p(z), z \in \mathbb{C}, a_p$'s are i.i.d.standard complex Gaussian

random variables, $\{e_p = z^p/\sqrt{p!}\}$ and it is an orthonormal basis of $L^2(\mathbb{C}, (e^{-|z|^2}/\pi) d \operatorname{Area}(z))$, where $d \operatorname{Area}(z)$ is the Lebesgue measure when we regard \mathbb{C} as \mathbb{R}^2 . In this example,

(1.2.11)
$$J(z) = J(|z|) = \frac{1}{2} \frac{d^2}{dr^2} \Big|_{r=|z|} r^2 \big[\coth(r) - 1 \big].$$

See, for example, [3, 11, 21]. This function J will change its sign so the method proving Theorem 1.1.1 does not apply to this case. However, we can separate the positive and negative part of this J and apply Theorem 1.1.2 to those $\Omega \subset \mathbb{R}^2$ with finite perimeter. The limit $\lim_{t\to 0^+} \mathcal{J}_t(\Omega)/t^3$ will then depend on $\int_{\mathbb{R}^2} J(z) \cdot |z| d \operatorname{Area}(z)$, which is still positive for the J in this example. The method proving Theorem 1.1.3 also applies to this example when Ω is a domain with a fractal boundary, while we do not know whether the limit is nonzero.

Apart from this Gaussian entire function example, it would also be interesting to consider those functions J such that the integrals $||J(z) \cdot |z|||_{L^1(\mathbb{R}^n)}$ vanish. In [22], the authors also considered another similar nonlocal integral functional with a probability background, where they obtained results similar to Theorem 1.1.2 in the dimension n = 2 case. It would also be intersting to see more applications of the nonlocal functional in [22] in other fields like probability. But these are out of the scope of this manuscript and we leave those questions for further works.

2. Upper and Lower Limits in Theorem 1.1.1

Let $\Omega_1, \Omega_2 \subset \mathbb{R}^n$ be two Lebesgue measurable sets and $\Omega_1 \cap \Omega_2 = \emptyset$. We define

(2.0.1)
$$\Gamma_t \equiv \overline{B(\Omega_2, t)} \cap \Omega_1 = \{ x \in \Omega_1 \mid \operatorname{dist}(x, \Omega_2) \le t \}$$

Because $\Omega_1 \cap \Omega_2 = \emptyset$, Γ_t also equals to $\overline{B(\partial\Omega_2, t)} \cap \Omega_1$. We assume that $|\Omega_1|$ is bounded, where for any measurable set $A \subset \mathbb{R}^n$, we let |A| be its Lebesgue measure. We assume assume that there is a $C_1 > 0$ and an $\alpha \in [0, n]$, such that $|\Gamma_t| \leq C_1 \cdot \min\{t^{n-\alpha}, 1\}$ for all t > 0. Intuitively, this assumption is a little bit weaker than assuming that $\partial\Omega_1 \cap \partial\Omega_2$ is of Minkowski dimension α .

We assume that J(z) for $z \in \mathbb{R}^n$ is a nonnegative function, such that

(2.0.2)
$$\int_{\mathbb{R}^n} J(z) \cdot |z|^{n-\alpha} dz \le C_J < \infty,$$

for a positive constant C_J . For simplicity, readers can keep natural examples like $J(z) \sim e^{-|z|}$ or $J(z) \sim e^{-|z|^2}$ in mind in the following.

Lemma 2.0.1. For all t > 0,

(2.0.3)
$$\int_{\Omega_1 \times \Omega_2} J\left(\frac{x-y}{t}\right) \, dx dy \le C_1 \cdot C_J \cdot t^{2n-\alpha}$$

Proof. Notice that

(2.0.4)
$$\int_{\Omega_1 \times \Omega_2} J\left(\frac{x-y}{t}\right) dx dy$$
$$= \int_{\Omega_1} \int_0^\infty \int_{\partial B(x,r) \cap \Omega_2} J\left(\frac{x-y}{t}\right) d\mathcal{H}^{n-1}(y) dr dx$$

As a function of (x, r) on $\Omega_1 \times (0, +\infty)$,

(2.0.5)
$$\int_{\partial B(x,r)\cap\Omega_2} J\left(\frac{x-y}{t}\right) d\mathcal{H}^{n-1}(y) \leq \mathbb{1}_{\Gamma_r}(x) \cdot \int_{\partial B(x,r)} J\left(\frac{x-y}{t}\right) d\mathcal{H}^{n-1}(y).$$

Hence,

$$(2.0.6) \qquad \int_{\Omega_{1}} \int_{0}^{\infty} \int_{\partial B(x,r)\cap\Omega_{2}} J\left(\frac{x-y}{t}\right) d\mathcal{H}^{n-1}(y) dr dx$$
$$\leq \int_{\Omega_{1}} \int_{0}^{\infty} \int_{\partial B(x,r)} \mathbb{1}_{\Gamma_{r}}(x) J\left(\frac{x-y}{t}\right) d\mathcal{H}^{n-1}(y) dr dx$$
$$= \int_{\Omega_{1}} \int_{\mathbb{R}^{n}} \mathbb{1}_{\Gamma_{|z|}}(x) J\left(\frac{-z}{t}\right) dz dx = \int_{\mathbb{R}^{n}} |\Gamma_{|z|}| J\left(\frac{z}{t}\right) dz$$
$$\leq C_{1} \cdot \int_{\mathbb{R}^{n}} \min\left\{|z|^{n-\alpha}, 1\right\} J\left(\frac{z}{t}\right) dz \leq C_{1} \cdot \int_{\mathbb{R}^{n}} |z|^{n-\alpha} J\left(\frac{z}{t}\right) dz$$
$$\leq C_{1} \cdot C_{J} \cdot t^{2n-\alpha}.$$

Lemma 2.0.1 provided a general approach for obtaining upper bounds. In the following, we will discuss potential lower bounds, subject to some additional natural assumptions for technical reasons. We first assume that $J(z) > c_J > 0$ when $|z| \leq a_J$ for some positive constants c_J, a_J . We also assume that there is a part $E \subset \partial \Omega_1 \cap \partial \Omega_2$, and there is a $D_E > 0$, such that for all $z \in E$ and $t \in (0, 1)$, $|B(z, t) \cap \Omega_2| \geq D_E \cdot t^n$. We may emphasize that this is the noncollapsing density at z on the Ω_2 side, and this is a natural assumption for NTA domains [13, 24]. We also define $N_t(E) \equiv B(E, t) \cap \Omega_1 = \{x \in \Omega_1 \mid \text{dist}(x, E) < t\}$.

Lemma 2.0.2. Suppose that there is a $C_2 > 0$ such that

$$(2.0.7) |N_t(E)| \ge C_2 \cdot t^{n-\alpha},$$

for all $t \in (0,1)$. Then there is a constant c = c(n) > 0, such that for all $t \in (0,1)$,

(2.0.8)
$$\int_{\Omega_1} \int_{\Omega_2} J\left(\frac{x-y}{t}\right) \, dx dy \ge c(n) \cdot C_2 c_J D_E \cdot (a_J t)^{2n-\alpha}.$$

Proof. The proof is direct. For any $x \in N_{(a_J/2) \cdot t}(E)$, there is a $z \in E \cap B(x, (a_J/2) \cdot t)$. And we see that $B(z, (a_J/2) \cdot t) \cap \Omega_2 \subset B(x, a_J \cdot t)$, and $|B(z, (a_J/2) \cdot t) \cap \Omega_2| \ge D_E \cdot ((a_J/2) \cdot t)^n$

by our assumptions. For any $y \in B(z, (a_J/2) \cdot t) \cap \Omega_2$, we also know that $J((x-y)/t) > c_J$. Hence,

$$(2.0.9) \qquad \int_{\Omega_1} \int_{\Omega_2} J\left(\frac{x-y}{t}\right) dx dy \ge \int_{N_{(a_J/2) \cdot t}(E)} \int_{B(z,(a_J/2) \cdot t) \cap \Omega_2} J\left(\frac{x-y}{t}\right) dy dx$$
$$(2.0.9) \qquad \ge \int_{N_{(a_J/2) \cdot t}(E)} \int_{B(z,(a_J/2) \cdot t) \cap \Omega_2} c_J dy dx \ge c_J D_E \left(\frac{a_J t}{2}\right)^n \cdot \left|N_{(a_J/2) \cdot t}(E)\right|$$
$$\ge C_2 c_J D_E \left(\frac{a_J t}{2}\right)^{2n-\alpha}.$$

We assume that $\Omega \subset \mathbb{R}^n$ is a bounded measurable set with $|\Omega| > 0$, and with topological boundary $\partial \Omega$ such that

(2.0.10)
$$0 < M_2 \leq \underline{\mathcal{M}}^{\alpha}(\partial \Omega) \leq \overline{\mathcal{M}}^{\alpha}(\partial \Omega) \leq M_1 < \infty,$$

where M_1, M_2 are two positive constants and $\alpha \in [n-1, n]$. The following Corollary then directly implies Theorem 1.1.1.

Corollary 2.0.3. Under the same assumptions on J(z) in Lemma 2.0.2, we also assume that there is a part $E \subset \partial \Omega$, on which for all $z \in E$ and $t \in (0,1)$, $|B(z,t) \cap \Omega| \ge D_E \cdot t^n$ and $|B(z,t) \cap \Omega^c| \ge D_E \cdot t^n$ for a positive constant D_E . Furthermore, we assume that $\underline{\mathcal{M}}^{\alpha}(E) > 0$. Then,

$$0 < \liminf_{t \to 0^+} \frac{1}{t^{2n-\alpha}} \int_{\Omega} \int_{\Omega^c} J\left(\frac{x-y}{t}\right) \, dx dy \leq \limsup_{t \to 0^+} \frac{1}{t^{2n-\alpha}} \int_{\Omega} \int_{\Omega^c} J\left(\frac{x-y}{t}\right) \, dx dy < \infty,$$

where the lower and upper limits are comparable to the Minkowski contents of $\partial\Omega$.

Proof. The upper limit follows from Lemma 2.0.1 directly. For the lower limit, notice that our assumption $\underline{\mathcal{M}}^{\alpha}(E) > 0$ is slightly weaker than the one we made in Lemma 2.0.2. The proof is exactly the same as the proof for Lemma 2.0.2, and one needs to combine both parts $B(E,t) \cap \Omega$ and $B(E,t) \cap \Omega^c$ to get a similar lower bound in Lemma 2.0.2. We can actually get that

(2.0.12)
$$\frac{1}{t^{2n-\alpha}} \int_{\Omega} \int_{\Omega^c} J\left(\frac{x-y}{t}\right) \, dx dy \ge C \cdot \frac{|B(\partial\Omega, t)|}{t^{n-\alpha}},$$

where the positive constant C depends on some parameters on J and $\partial\Omega$ that we mentioned in Lemma 2.0.2.

Remark 2.0.4. Recall that for Lemma 2.0.2 and Corollary 2.0.3, we assume that $J(z) > c_J > 0$ when $|z| \leq a_J$. The condition that $J(z) > c_J$ on $|z| \leq a_J$ can also be replaced by J(z) bounded from below on an annulus, but then one needs to have volume lower bounds better than $|B(x,t) \cap \Omega| \geq D_E \cdot t^n$ and $|B(x,t) \cap \Omega^c| \geq D_E \cdot t^n$, i.e., one also needs to replace balls B(x,t) by annuluses.

When J(z) has the form $|z|^{-\beta}$, the author of [14] introduced a fractal dimension and discussed its relation with the Minkowski dimension. This singular form of J(z) comes from the definition of fractional Sobolev space, as discussed in Section 1 and [2]. We will revisit the results in [2] in Section3.

3. Limits in Theorem 1.1.2 and Theorem 1.1.3

3.1. Sets of finite perimeter and general BV functions.

Let us first recall the results from [2, 6]. We will make some revisions to ensure that the settings in [2, 6] are compatible with this manuscript. Assume that J(z) for $z \in \mathbb{R}^n$ is nonnegative and is radially symmetric, i.e., J(z) = J(|z|). And we assume that $\int_{\mathbb{R}^n} J(z) \cdot$ $|z| dz = \int_0^\infty J(r) \cdot r^n dr = 1$, i.e., $\alpha = n - 1$ in Section 2. Recall that the functional $\mathcal{J}_t(\Omega)$ we considered in Section 2 has the following more symmetric form. For a measurable set $\Omega \subset \mathbb{R}^n$, (3.1.1)

$$\int_{\Omega} \int_{\Omega^c} J\left(\frac{x-y}{t}\right) dxdy + \int_{\Omega^c} \int_{\Omega} J\left(\frac{x-y}{t}\right) dxdy = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \left|\mathbbm{1}_{\Omega}(x) - \mathbbm{1}_{\Omega}(y)\right| J\left(\frac{x-y}{t}\right) dxdy$$

We can consider a general function f(x) instead of only $\mathbb{1}_{\Omega}(x)$, and we know the following theorem from [2].

Theorem 3.1.1 ([2], Theorem 3'). Let $A \subset \mathbb{R}^n$ be a bounded smooth domain, $f \in L^1(A)$. Then $f \in BV(A)$ if and only if

(3.1.2)
$$\liminf_{t \to 0^+} \frac{1}{t^{n+1}} \int_A \int_A |f(x) - f(y)| \cdot J\left(\frac{x-y}{t}\right) \, dx \, dy < \infty.$$

Here, for a bounded smooth domain $A \subset \mathbb{R}^n$, we say that a function $f \in BV(A)$ if and only if $[f]_{BV(A)} < \infty$, where the BV-seminorm $[f]_{BV(A)}$ is defined by

(3.1.3)
$$[f]_{\mathrm{BV}(A)} \equiv \sup \left\{ \int_{A} f \operatorname{div}(\varphi) \mid \varphi \in C_{0}^{\infty}(A, \mathbb{R}^{n}), ||\varphi||_{\infty} \leq 1 \right\}.$$

This BV-seminorm equals to $\int_A |\nabla f|$ if $f \in W^{1,1}(A)$, which means that BV(A) can be seen as an extension of $W^{1,1}(A)$. In particular, BV(A) contains indicator functions for domains $\Omega \subset A$ with well-behaved topological boundaries $\partial\Omega$. For instance, if $\Omega \subset A$ has a Lipschitz boundary $\partial\Omega$, then the divergence theorem implies that $\mathbb{1}_{\Omega} \in BV(A)$ with $[\mathbb{1}_{\Omega}]_{BV(A)} = \mathcal{H}^{n-1}(\partial\Omega)$, even though $\mathbb{1}_{\Omega} \notin W^{1,1}(A)$.

Definition 3.1.2. Let Ω be a Borel set in \mathbb{R}^n . We call Ω a Caccioppoli set (or a set of locally finite perimeter) if and only if for every bounded open set $A \subset \mathbb{R}^n$, $[\mathbb{1}_{\Omega}]_{BV(A)} < \infty$.

Theorem 3.1.1 provides a direct criterion for determining whether a function belongs to the BV class. Additionally, De Giorgi [7, 8] actually used the limit $\lim_{t\to 0^+} \mathcal{J}_t(\Omega)/t^{n+1}$ with $J(z) = e^{-|z|^2}$ as his definition of perimeter for sets of locally finite perimeter up to a constant. As we have seen, when Ω is a domain with a Lipschitz boundary, we have that $[\mathbb{1}_{\Omega}]_{\mathrm{BV}(A)} = \mathcal{H}^{n-1}(\partial\Omega)$. In general, for every $f \in \mathrm{BV}(A)$, the limit in Theorem 3.1.1 actually exists, as shown in the following theorem.

Theorem 3.1.3 ([6], Theorem 1). Let $A \subset \mathbb{R}^n$ be a bounded domain with a Lipschitz boundary ∂A , and let $f \in BV(A)$. Then,

(3.1.4)
$$\lim_{t \to 0^+} \frac{1}{t^{n+1}} \int_A \int_A |f(x) - f(y)| \cdot J\left(\frac{x-y}{t}\right) \, dx \, dy = K(n) \cdot [f]_{\mathrm{BV}(A)},$$

where

(3.1.5)
$$K(n) = \frac{\Gamma(\frac{n}{2})}{\sqrt{\pi}\Gamma(\frac{n+1}{2})}$$

We then see that Theorem 3.1.3 holds true for $f = \mathbb{1}_{\Omega}$ when Ω is a Caccioppoli set.

Corollary 3.1.4. For the same A, K(n) in Theorem 3.1.3, if Ω is a Caccioppoli set, then

(3.1.6)
$$\lim_{t \to 0^+} \frac{1}{t^{n+1}} \int_{\Omega \cap A} \int_{\Omega^c \cap A} J\left(\frac{x-y}{t}\right) \, dx \, dy = 2K(n) \cdot \mathcal{H}^{n-1}(\partial^* \Omega \cap A)$$

Here, $\partial^*\Omega$ is called the reduced boundary of Ω , which is a subset of $\partial\Omega$ on which the density of Ω in balls centered at those points is 1/2. This is a consequence of De Giorgi's structure theorem. See Corollary 15.8 and Theorem 15.9 in [16], for example. For Ω with a Lipschitz boundary, $\partial^*\Omega$ includes those differentiable points on $\partial\Omega$. We can also generalize Corollary 3.1.4 slightly.

Corollary 3.1.5. For the same A, K(n) in Theorem 3.1.3, if Ω_1 and Ω_2 are two disjoint Caccioppoli sets, then

(3.1.7)
$$\lim_{t \to 0^+} \frac{1}{t^{n+1}} \int_{\Omega_1 \cap A} \int_{\Omega_2 \cap A} J\left(\frac{x-y}{t}\right) dxdy$$
$$= 2K(n) \cdot \mathcal{H}^{n-1}(\partial^*\Omega_1 \cap \partial^*\Omega_2 \cap A).$$

Proof. Notice that because J is radially symmetric, and $\Omega_1 \cap \Omega_2 = \emptyset$,

$$(3.1.8) \qquad 2\int_{\Omega_1 \cap A} \int_{\Omega_2 \cap A} J\left(\frac{x-y}{t}\right) dxdy$$
$$= \int_{\Omega_1 \cap A} \int_{\Omega_1^c \cap A} J\left(\frac{x-y}{t}\right) dxdy + \int_{\Omega_2 \cap A} \int_{\Omega_2^c \cap A} J\left(\frac{x-y}{t}\right) dxdy$$
$$- \int_{(\Omega_1 \cup \Omega_2) \cap A} \int_{(\Omega_1 \cup \Omega_2)^c \cap A} J\left(\frac{x-y}{t}\right) dxdy.$$

And by Corollary3.1.4, we know that the limit is

(3.1.9)
$$2K(n) \cdot (\mathcal{H}^{n-1}(\partial^*\Omega_1 \cap A) + \mathcal{H}^{n-1}(\partial^*\Omega_2 \cap A) - \mathcal{H}^{n-1}(\partial^*(\Omega_1 \cup \Omega_2) \cap A)).$$

Because $\Omega_1 \cap \Omega_2 = \emptyset$, Theorem 16.3 in [16] implies that $\partial^*(\Omega_1 \cup \Omega_2)$ is, up to a set of \mathcal{H}^{n-1} measure zero, $(\partial^*\Omega_1 \setminus \partial^*\Omega_2) \cup (\partial^*\Omega_2 \setminus \partial^*\Omega_1)$. Hence, the above equation equals to (3.1.10) $4K(n) \cdot \mathcal{H}^{n-1}(\partial^*\Omega_1 \cap \partial^*\Omega_2 \cap A)$.

3.2. Sobolev Spaces. There is another result in [2] that deals with the case for the functional

(3.2.1)
$$\int_{A} \int_{A} \left| f(x) - f(y) \right|^{p} \cdot J\left(\frac{x-y}{t}\right) \, dx dy,$$

where 1 . To be more precise, we assume that <math>J(z) for $z \in \mathbb{R}^n$ is nonnegative and is radially symmetric, i.e., J(z) = J(|z|). We further assume that $\int_{\mathbb{R}^n} J(z) \cdot |z|^p dz = \int_0^\infty J(r) \cdot r^{n+p-1} dr = 1$. One can compare these assumptions with those we made in Section 2 and also in Section 3.1.

Definition 3.2.1. Let $A \subset \mathbb{R}^n$ be a bounded smooth domain, $f \in L^p(A)$ and $1 \leq p < \infty$. We say that $f \in W^{1,p}(A)$ if and only if $[f]_{W^{1,p}(A)} < \infty$. Here, the seminorm $[f]_{W^{1,p}(A)}$ is defined by

(3.2.2)
$$[f]_{W^{1,p}(A)} = \left(\int_{A} |\nabla f|^{p} dx\right)^{\frac{1}{p}}$$

where ∇f is the weak derivative of f.

Then, we have the following theorem from [2] when 1 .

Theorem 3.2.2 ([2], Theorem 2). Let $A \subset \mathbb{R}^n$ be a bounded smooth domain, $f \in L^p(A)$ and 1 . Then,

(3.2.3)
$$\lim_{t \to 0^+} \frac{1}{t^{n+p}} \int_A \int_A |f(x) - f(y)|^p \cdot J\left(\frac{x-y}{t}\right) dx dy = K(n,p) \cdot \left([f]_{W^{1,p}(A)}\right)^p$$

with the convention that $[f]_{W^{1,p}(A)} = \infty$ if $f \notin W^{1,p}(A)$. Here, K(n,p) is a constant depending on n, p.

3.3. Koch snowflake and sets of self-similar fractal boundary.

In stead of discussing sets with general self-similar fractal boundaries in detail, we will focus on a specific example, Koch snowflake (Figure 1), in this section. But the idea is actually applicable to many other sets with self-similar boundaries constructed in a similar way to the Koch snowflake. And we will provide an alternative example, constructed using a similar method as the Koch snowflake but with a different scale (Figure 2). Figure 3 shows the first several steps of the construction of these shrunk Koch snowflakes (or Koch curves). In order to maintain consistency with Section 2, we will take n = 2 in this subsection, but continue to use the notation n when referring to the dimension.

The standard Koch snowflake, which we say is of scale $\eta = 3$, is constructed iteratively. The process starts with an equilateral triangle of side length 1. Then, on the middle of each




FIGURE 1. Standard Koch Snowflake

side of this triangle, we add a smaller equilateral triangle of side length $1/\eta = 1/3$, which results in a regular hexagram. We repeat this process iteratively, adding an equilateral triangle of side length $(1/3)^2$ to the middle of each side of the hexagram. Taking the limit of this process yields an open set with a boundary of Minkowski dimension (and also Hausdorff dimension) $\alpha(3) \equiv \log_3(4)$. For further details, see, for example, [9].

A general Koch snowflake of scale $\eta > 1$ is constructed in a similar manner, as illustrated in Figure 3. The iterative process is carried out on each side of an equilateral triangle, where a small equilateral triangle of side length $1/\eta$ is added to the middle of the original line segment (one side of the original equilateral triangle). Then we obtain 4 line segments. On each of these line segments, a smaller equilateral triangle with side length $1/\eta \times$ (side length of this line segment) is then added to its middle, generating a total of 4^2 line segments. The process is repeated by adding smaller equilateral triangles to the new line segments. The limit open set is the Koch snowflake of scale η , whose boundary has Minkowski dimension $\alpha(\eta)$ satisfying

(3.3.1)
$$2 \cdot \left(\frac{\eta - 1}{2\eta}\right)^{\alpha(\eta)} + 2 \cdot \left(\frac{1}{\eta}\right)^{\alpha(\eta)} = 1$$

 $\alpha(\eta)$ satisfies this equation because of the self-similarity. Let us see Figure 2. The boundary part between U_1 and V_1 is $(\eta - 1)/2$ times larger than the boundary part between U_2 and V_2 , while the boundary part between $(U_1 \cup U_2 \cup U_3 \cup U_4)$ and $(V_1 \cup V_2 \cup V_3 \cup V_4)$ is η times larger




FIGURE 2. Snowflake with $\eta = 5$

than the boundary part between U_2 and V_2 . Additionally, the boundary part between $(U_1 \cup U_2 \cup U_3 \cup U_4)$ and $(V_1 \cup V_2 \cup V_3 \cup V_4)$ is the sum of twice the boundary part between U_1 and V_1 and twice the boundary part between U_2 and V_2 .

Notice that the Koch snowflake satisfies the property we need to prove Corollary 2.0.3, i.e., it has positive finite lower and upper bounds on Minkowski contents, together with density lower bounds at the boundary of the Koch snowflake. This is because the Koch snowflake is a quasiball and therefore an NTA domain. For example, see Remark 2.1 in the survey paper by [24]. Additionally, in [18], some 3-dimensional analogs of snowflakes were also shown to be quasiballs. In particular, in the 2-dimensional case, Ahlfors's 3-point condition states that a Jordan curve γ in \mathbb{R}^2 is a quasicircle if and only if for any two points a, b on γ , the smaller arc between a, b has a diameter comparable to the distance |a - b|.

In [12], similar limiting behaviors of Minkowski contents were established for general selfsimilar fractals.

Theorem 3.3.1. Assume that J(z) for $z \in \mathbb{R}^n$ is a nonnegative function in $L^1(\mathbb{R}^n)$ such that

(3.3.2)
$$\int_{\mathbb{R}^n} J(z) \cdot |z|^n \, dz \le C_J < \infty,$$




FIGURE 3. Koch Curve with $\eta = 5$. We do this iterative process on each side of a equilateral triangle as shown in Figure 2.

for a positive constant C_J . Let $\Omega = \Omega(3)$ be the standard Koch snowflake. Then, for any t > 0, the limit

(3.3.3)
$$\lim_{i \to \infty} \left(3^i/t\right)^{2n-\alpha(3)} \int_{\Omega \times \Omega^c} J\left(\frac{x-y}{(t/3^i)}\right) \, dx \, dy$$

exists and is positive.

Proof. The positivity of the limit follows from Lemma 2.0.2. For the existence, we let Ω be the bounded domain in Figure 1 of Koch snowflake, and we let Ω^c be the unbounded part. We evenly partition Ω into U_1, \ldots, U_{12} , and evenly partition Ω^c into V_1, \ldots, V_{12} as shown in Figure 1. This division is chosen to facilitate the computation of the functional since the boundary part of $\partial U_1 \cap \partial V_1$ is a rescaling by 1/3 of the boundary part of $\partial (U_{12} \cup U_1 \cup U_2 \cup U_3) \cap \partial (V_{12} \cup V_1 \cup V_2 \cup V_3)$. Therefore, the functional can be expressed as

$$(3.3.4) \quad \int_{\Omega \times \Omega^c} J\left(\frac{x-y}{t}\right) \, dxdy = \sum_{i=1}^{12} \int_{U_i \times V_i} J\left(\frac{x-y}{t}\right) \, dxdy + \sum_{i \neq j} \int_{U_i \times V_j} J\left(\frac{x-y}{t}\right) \, dxdy.$$

Let us first see the second term in (3.3.4). If $i \neq j$, we see that either dist $(U_i, V_j) > 0$, like U_1 and V_3 , or $\partial U_i \cap \partial V_j$ is a single point, like U_1 and V_2 .

For the pair (U_1, V_3) , one can similarly define Γ_t like we did for Lemma 2.0.1 in Section 2. $\Gamma_t = \emptyset$ when t is small, so there is a $D_1(\Omega) > 0$ such that $|\Gamma_t| < D_1 \cdot \min\{t^{2n}, 1\}$. So, Lemma 2.0.1 shows that this pair will give a term less than order t^{2n} . For the pair (U_1, V_2) (or the pair (U_1, V_{12})), notice that U_1 is included in a cone \widetilde{U}_1 and V_2 is included in a cone \widetilde{V}_2 . Also, \widetilde{U}_1 and \widetilde{V}_2 touch at their vertices. Again, one can similarly define Γ_t , and there is a $D_2(\Omega) > 0$ such that $|\Gamma_t| < D_2 \cdot \min\{t^{2n}, 1\}$. So, Lemma 2.0.1 shows that this pair will give a term less than order t^{2n} .

Hence, there is a $D(\Omega) > 0$ such that the second term in (3.3.4) is less than $C_J \cdot D \cdot t^{2n}$.

Next, let us consider the first term in (3.3.4). Since all $\{U_i\}$'s (and all $\{V_i\}$'s) are the same up to rigid transformations, the first term in (3.3.4) is

(3.3.5)
$$12 \cdot \int_{U_1 \times V_1} J\left(\frac{x-y}{t}\right) \, dx dy,$$

which we denote as 12F(t), where F(t) is this functional involving domains U_1, V_1 . Because $\partial U_1 \cap \partial V_1$ is a rescaling by 1/3 of the boundary part of $\partial (U_{12} \cup U_1 \cup U_2 \cup U_3) \cap \partial (V_{12} \cup V_1 \cup V_2 \cup V_3)$, $\partial (3U_1) \cap \partial (3V_1) = \partial (U_{12} \cup U_1 \cup U_2 \cup U_3) \cap \partial (V_{12} \cup V_1 \cup V_2 \cup V_3)$. Let us consider the pair of domains $(3U_1, 3V_1)$, and the functional

(3.3.6)
$$\int_{3U_1 \times 3V_1} J\left(\frac{x-y}{t}\right) \, dx dy.$$

Compare the pair of domains $(3U_1, 3V_1)$ with the pair of domains $(U_{12} \cup U_1 \cup U_2 \cup U_3, V_{12} \cup V_1 \cup V_2 \cup V_3)$. By using a similar method to estimate the second term in (3.3.4), up to a term less than $C_J \cdot D \cdot t^{2n}$, the above functional of the pair $(3U_1, 3V_1)$ equals to

(3.3.7)
$$\int_{(U_{12}\cup U_1\cup U_2\cup U_3)\times (V_{12}\cup V_1\cup V_2\cup V_3)} J\left(\frac{x-y}{t}\right) dxdy,$$

which is also equal to

(3.3.8)
$$4 \cdot \int_{U_1 \times V_1} J\left(\frac{x-y}{t}\right) \, dx \, dy$$

up to another term less than $C_J \cdot D \cdot t^{2n}$. On the other hand, by change of variables,

(3.3.9)
$$\int_{3U_1 \times 3V_1} J\left(\frac{x-y}{t}\right) \, dx \, dy = 3^4 \cdot \int_{U_1 \times V_1} J\left(\frac{x-y}{(t/3)}\right) \, dx \, dy.$$

So, $4F(t) = 3^4F(t/3) + R(t)$ for an R(t) such that $|R(t)| \leq C_J \cdot D \cdot t^{2n}$. Recall that in our case, n = 2 and $\alpha(3) = \log_3(4)$, hence $3^4/4 = 3^{2n-\alpha(3)}$. Because t > 0 is arbitrary, so for any $i \in \mathbb{N}$,

(3.3.10)
$$(3^{2n-\alpha(3)})^{i} F(t/3^{i}) = (3^{2n-\alpha(3)})^{i+1} F(t/3^{i+1}) + (3^{2n-\alpha(3)})^{i} \cdot R(t/3^{i}).$$

We see that

(3.3.11)
$$\sum_{i=0}^{\infty} \left(3^{2n-\alpha(3)}\right)^{i} \cdot |R(t/3^{i})| \leq \sum_{i=0}^{\infty} \left(3^{2n-\alpha(3)}\right)^{i} \cdot C_{J} \cdot D(\Omega) \cdot t^{2n} \cdot \left(3^{2n}\right)^{-i} < \infty.$$

So, the limit

(3.3.12) $\lim_{i \to \infty} (3^{2n-\alpha(3)})^i F(t/3^i)$

exists for any t > 0.

It is unknown whether the limit

(3.3.13)
$$\lim_{t \to 0^+} \frac{1}{t^{2n-\alpha(3)}} \int_{\Omega \times \Omega^c} J\left(\frac{x-y}{t}\right) \, dx \, dy$$

exists for the standard Koch snowflake. On the other hand, it is known that the lower and upper Minkowski contents of the standard Koch snowflake are not the same.

However, for a snowflake of scale $\eta > 1$, if the logarithm ratio of the two factors $1/\eta$ and $(\eta - 1)/(2\eta)$ is not a rational number, i.e.,

(3.3.14)
$$\frac{\log[(2\eta)/(\eta-1)]}{\log(\eta)} = 1 + \frac{\log(2) - \log(\eta-1)}{\log(\eta)} \notin \mathbb{Q},$$

then the limiting behaviors are actually better in some sense. Indeed, we have the following theorem. Also notice that any transcendental number satisfies the condition (3.3.14).

Theorem 3.3.2. Assume that J(z) for $z \in \mathbb{R}^n$ is a nonnegative function in $L^1(\mathbb{R}^n)$ such that

(3.3.15)
$$\int_{\mathbb{R}^n} J(z) \cdot |z|^n \, dz \le C_J < \infty,$$

for a positive constant C_J . Let $\Omega = \Omega(\eta)$ be the Koch snowflake of scale $\eta > 1$. If η satisfies (3.3.14), then the limit

(3.3.16)
$$\lim_{t \to 0^+} \frac{1}{t^{2n-\alpha(\eta)}} \int_{\Omega \times \Omega^c} J\left(\frac{x-y}{t}\right) \, dx dy$$

exists.

Proof. Let us use Figure 2 to illustrate the proofs, which follow the same strategy as the proof for Theorem 3.3.1. We take domains U_1, U_2, U_3, U_4 and V_1, V_2, V_3, V_4 as shown in Figure 2. And we only need to prove the existence for the limit

(3.3.17)
$$\lim_{t \to 0^+} \frac{1}{t^{2n-\alpha(\eta)}} \int_{(U_1 \cup U_2 \cup U_3 \cup U_4) \times (V_1 \cup V_2 \cup V_3 \cup V_4)} J\left(\frac{x-y}{t}\right) \, dx \, dy.$$

If we let

(3.3.18)
$$F(t) \equiv \int_{(U_1 \cup U_2 \cup U_3 \cup U_4) \times (V_1 \cup V_2 \cup V_3 \cup V_4)} J\left(\frac{x-y}{t}\right) dxdy,$$

then, use the self-similarity as we obtained the dimension $\alpha(\eta)$ in (3.3.1) again, we will get that for all t > 0,

$$(3.3.19) \quad 2\left[F\left(\frac{t}{(1-(1/\eta))/2}\right) \cdot \left(\frac{1-(1/\eta)}{2}\right)^{2n} + F\left(\frac{t}{(1/\eta)}\right) \cdot \left(\frac{1}{\eta}\right)^{2n}\right] = F(t) + R(t),$$

where there is a constant $D = D(\Omega(\eta)) > 0$ such that $|R(t)| \leq C_J \cdot D \cdot t^{2n}$.

Define $G(s) \equiv e^{(2n-\alpha(\eta))s} \cdot F(e^{-s})$, and let $r_1 = (1 - (1/\eta))/2$, $r_2 = 1/\eta$, we see that

$$(3.3.20) \quad 2\left\lfloor G\left(s + \log(r_1)\right) \cdot r_1^{\alpha(\eta)} + G\left(s + \log(r_2)\right) \cdot r_2^{\alpha(\eta)} \right\rfloor = G(s) + e^{(2n - \alpha(\eta))s} R(e^{-s}).$$

Let μ be the probability distribution function which assigns weight $2r_i^{\alpha(\eta)}$ at the point $-\log(r_i)$ for i = 1, 2. Then, the function G(s) satisfies the renewal equation

(3.3.21)
$$G(s) = -e^{(2n-\alpha(\eta))s}R(e^{-s}) + \int_0^s G(s-s') \ d\mu(s').$$

Because we assumed that $\log(r_1)/\log(r_2) \notin \mathbb{Q}$, we can apply the renewal theorem, as was done in [12] for the limits of Minkowski contents. For general statements of the renewal theorem, see Chapter XI of [10]. Therefore, we can show that

(3.3.22)
$$\lim_{s \to +\infty} G(s) = \frac{1}{2r_1^{\alpha(\eta)} \log(r_1) + 2r_2^{\alpha(\eta)} \log(r_2)} \cdot \int_0^{+\infty} e^{(2n - \alpha(\eta))s} R(e^{-s}) \, ds.$$

This completes the proof of Theorem 3.3.2.

Remark 3.3.3. As shown in Lemma 2.0.1 and Lemma 2.0.2, both limits in Theorem 3.3.1 and Theorem 3.3.2 are comparable, i.e., up to a constant depending on the dimension, the kernel J, and the noncollapsing parameter $D_{\partial\Omega}$, to the Minkowski contents of the corresponding $\partial\Omega$'s. And in these specific Koch snowflakes, D_{Ω} can also be chosen explicitly depending on the scale η .

4. Applications in Fluctuations of Determinantal Processes

For simplicity, let us consider random point processes on (\mathbb{R}^n, μ) , where μ is a probability measure on \mathbb{R}^n . For more rigorous definitions, see Section 4.2 of [1]. We let **M** be the space of σ -finite Radon measures on \mathbb{R}^n , and let \mathbf{M}_+ be the subset of **M** consisting of positive measures.

Definition 4.0.1. A random point process on \mathbb{R}^n is a random, integer-valued $X \in \mathbf{M}_+$.

By random, we mean that for any Borel set $A \subset \mathbb{R}^n$, X(A) is an integer-valued random variable.

Definition 4.0.2. Assume that there are locally nonnegative integrable symmetric functions ρ_k on $\mathbb{R}^{n \times k}$ for all $k \ge 1$, such that for any measurable subset A of \mathbb{R}^n ,

(4.0.1)
$$k! \cdot \mathcal{E}\left[\binom{X(A)}{k}\right] = \int_{A^{\otimes k}} \rho_k(x_1, \dots, x_k) \ d\mu(x_1) \dots d\mu(x_k).$$

Here, $\mathcal{E}(\cdot)$ denotes expectations. And we say that ρ_k is symmetric if

(4.0.2)
$$\rho_k(x_1, \dots, x_k) = \rho_k(x_{\sigma(1)}, \dots, x_{\sigma(k)}),$$

for all $\sigma \in S_k$, the k-th symmetry group. And we call the function ρ_k k-th joint intensity or k-th correlation function of the point process X with respect to $(\mathbb{R}^n, |\cdot|)$. Here, $|\cdot|$ denotes the Lebesgue measure on \mathbb{R}^n .

Definition 4.0.3. X is called a determinantal point process if there is a kernel function K on $\mathbb{R}^n \times \mathbb{R}^n$ such that for all $k \ge 1$,

(4.0.3)
$$\rho_k(x_1, \dots, x_k) = \det \left(K(x_i, x_j) \right)_{1 \le i, j \le k}$$

Example 4.0.4. Let $N \in \mathbb{N}$. For the *complex Ginibre ensemble* of dimension N, we mean $N \times N$ random matrices over \mathbb{C} , with entries being i.i.d. complex Gaussian random variables with mean 0 and variance 1/N. Let $(\lambda_1^N, \ldots, \lambda_N^N)$ be the eigenvalues of the Ginibre ensemble of dimension N. Then, the law for those eigenvalues is

(4.0.4)
$$p(z_1, \dots, z_N) = \frac{1}{Z_N} \cdot \prod_{1 \le i < j \le N} |z_i - z_j|^2 \cdot e^{-N \sum_{i=1}^N |z_i|^2} d\mathcal{H}^2(z_1) \dots d\mathcal{H}^2(z_N).$$

Here, each z_i is in \mathbb{C} , and \mathcal{H}^2 is the 2-dimensional Hausdorff measure on \mathbb{C} , which is exactly the Lebesgue measure on $\mathbb{C} = \mathbb{R}^2$. Z_N is a constant such that the integral of p is 1. Then, if we let $X_N = \sum_{i=1}^N \delta_{z_i}$, where δ_z is the Dirac measure at z, then X_N is a determinantal process, such that the kernel function is

(4.0.5)
$$K_N(z,w) = \sum_{m=0}^{N-1} \frac{N^m}{m!} (z\overline{w})^m, \quad z,w \in \mathbb{C},$$

with respect to the measure $d\mu_N(z) = \frac{N}{\pi} e^{-N|z|^2} d\mathcal{H}^2(z)$. Notice that $K_N(w, z) = \overline{K_N(z, w)}$. For more properties of the Ginibre ensemble, one may see [20].

4.1. Asymptotic behaviors of eigenvalues of the Ginibre ensemble.

We will let $N \to \infty$ and study the behavior of X_N , which is the sum of Dirac measures on the eigenvalues of the Ginibre ensemble as defined in Example 4.0.4. First, we have the following lemma for the kernel function K_N in Example 4.0.4.

Lemma 4.1.1. For any $\lambda \in (0, 1)$, there is a positive constant $C(\lambda)$ and a constant $\delta(\lambda) \in (0, 1)$, such that for any $z, w \in \mathbb{C}$ with $|z|, |w| < \lambda$,

(4.1.1)
$$\left| e^{N(z\overline{w})} - K_N(z,w) \right| \le C \cdot \frac{e^{(N \cdot |z\overline{w}|)}}{\sqrt{N}} \cdot \delta^N$$

Proof. For any fixed N and $s \in (0, \lambda)$,

(4.1.2)
$$e^{Ns} - \sum_{m=0}^{N-1} \frac{N^m}{m!} s^m = \sum_{m \ge N} \frac{N^m}{m!} s^m = \frac{(Ns)^N}{N!} \cdot \left(1 + \sum_{m \ge 1} \frac{(Ns)^m}{(N+1)\cdots(N+m)}\right) \le C(\lambda) \cdot \frac{(Ns)^N}{N!}$$

where the last inequality is because $N \leq N + i$ for $i \geq 0$ and $\sum_{m\geq 1} s^m \leq C(\lambda)$. Notice that Stirling's formula showed that there is another dimensional constant C > 0, such that $k! \geq C\sqrt{k}(k/e)^k$ for any $k \in \mathbb{Z}_+$. Hence,

(4.1.3)
$$\frac{(Ns)^N}{N!} \le \frac{(se)^N}{C\sqrt{N}}.$$

Notice that the function for s, se^{1-s} , is increasing in (0, 1). So, $se \leq \lambda e^{1+s-\lambda}$. What's more, $\lambda e^{1-\lambda} < 1e^{1-1} = 1$ since $\lambda \in (0, 1)$. We may then denote the constant $\lambda e^{1-\lambda}$ as $\delta(\lambda) < 1$. Then, $se \leq \delta(\lambda)e^s$. So,

(4.1.4)
$$\left| e^{Ns} - \sum_{m=0}^{N} \frac{N^m}{m!} s^m \right| \le \frac{e^{Ns}}{C\sqrt{N}} \cdot \delta^N.$$

We first identify \mathbb{C} with \mathbb{R}^2 . Then, let us consider a function $\varphi \in L^1(\mathbb{R}^2)$ with the closure of its support $\overline{\operatorname{supp}(\varphi)} \subset \{x \in \mathbb{R}^2 \mid |x| < 1\}$. Then,

(4.1.5)
$$X_N(\varphi) \equiv \sum_{i=1}^N \varphi(z_i)$$

is a random variable and we will study its asymptotic behavior as $N \to \infty$. A typical choice for φ is to take $\varphi = \mathbb{1}_{\Omega}$ for some domain Ω with closure contained in the unit ball of \mathbb{R}^2 . In the following, we assume that there exists a $\lambda \in (0, 1)$ such that $\overline{\operatorname{supp}(\varphi)} \subset \{x \in \mathbb{R}^2 \mid |x| < \lambda\}$, and let $\delta(\lambda) \in (0, 1)$ be the constant we obtained in Lemma 4.1.1.

Lemma 4.1.2. There is a dimensional constant C > 0, such that

(4.1.6)
$$\left| \mathcal{E}(X_N(\varphi)) - \frac{N}{\pi} \int \varphi(x) \, dx \right| \le C \cdot (\sqrt{N}\delta^N) \cdot ||\varphi||_{L^1(\mathbb{R}^2)}$$

Proof. Because X_N is a determinantal process, the definition and Lemma 4.1.1 can give us that

(4.1.7)
$$\left| \begin{aligned} \mathcal{E}(X_N(\varphi)) - \frac{N}{\pi} \int \varphi(x) \, dx \right| &= \left| \frac{N}{\pi} \int_{\mathbb{R}^2} \varphi(x) \cdot (K_N(x, x) - e^{N|x|^2}) e^{-N|x|^2} \, dx \right| \\ &\leq C \cdot N \int_{\mathbb{R}^2} |\varphi| \cdot \frac{\delta^N}{\sqrt{N}} \, dx = C \cdot (\sqrt{N}\delta^N) \cdot ||\varphi||_{L^1(\mathbb{R}^2)}. \end{aligned} \right|$$

Next, we need the formula for the variance of $X_N(\varphi)$. We cite the equation (28) in [19]. For $\varphi \in L^2(\mathbb{R}^2)$,

(4.1.8)
$$\operatorname{Var}(X_N(\varphi)) = \frac{1}{2} \int_{\mathbb{R}^2 \times \mathbb{R}^2} |\varphi(x) - \varphi(y)|^2 \cdot |K_N(x,y)|^2 \, d\mu_N(x) d\mu_N(y).$$

In particular, when $\varphi = \mathbb{1}_{\Omega}$,

(4.1.9)
$$\operatorname{Var}(X_N(\mathbb{1}_{\Omega})) = \int_{\Omega \times \Omega^c} |K_N(x,y)|^2 d\mu_N(x) d\mu_N(y).$$

Recall that we assume that there exists $\lambda \in (0, 1)$ such that $\overline{\operatorname{supp}(\varphi)} \subset \{x \in \mathbb{R}^2 \mid |x| < \lambda\}$, and we can get the constant $\delta(\lambda) \in (0, 1)$ in Lemma 4.1.1. Let us choose a $\lambda_+ \in (\lambda, 1)$. We see that since $\varphi = 0$ outside of $B(0, \lambda)$, we have that

(4.1.10)
$$\operatorname{Var}(X_{N}(\varphi)) = \frac{1}{2} \int_{\mathbb{R}^{2} \times \mathbb{R}^{2}} |\varphi(x) - \varphi(y)|^{2} \cdot |K_{N}(x,y)|^{2} d\mu_{N}(x) d\mu_{N}(y) \\ = \frac{1}{2} \int_{B(0,\lambda_{+}) \times B(0,\lambda_{+})} |\varphi(x) - \varphi(y)|^{2} \cdot |K_{N}(x,y)|^{2} d\mu_{N}(x) d\mu_{N}(y) \\ + \int_{B(0,\lambda_{+}) \times (B(0,\lambda_{+}))^{c}} |\varphi(x)|^{2} \cdot |K_{N}(x,y)|^{2} d\mu_{N}(x) d\mu_{N}(y).$$

Let us compute the second term in the above formula. We use the information of Ginibre ensemble and we see that since $\varphi(x) = 0$ outside of $B(0, \lambda)$,

$$(4.1.11)$$

$$\int_{B(0,\lambda_{+})\times(B(0,\lambda_{+}))^{c}} |\varphi(x)|^{2} \cdot |K_{N}(x,y)|^{2} d\mu_{N}(x)d\mu_{N}(y)$$

$$= \int_{B(0,\lambda)\times(B(0,\lambda_{+}))^{c}} |\varphi(x)|^{2} \cdot |K_{N}(x,y)|^{2} d\mu_{N}(x)d\mu_{N}(y)$$

$$\leq \left(\frac{N}{\pi}\right)^{2} \int_{B(0,\lambda)\times(B(0,\lambda_{+}))^{c}} |\varphi(x)|^{2} \cdot e^{2N|x||y|} \cdot (e^{-N|x|^{2}}e^{-N|y|^{2}}) dxdy$$

$$\leq \left(\frac{N}{\pi}\right)^{2} \int_{B(0,\lambda)\times(B(0,\lambda_{+}))^{c}} |\varphi(x)|^{2} \cdot e^{-N(|y|-\lambda)^{2}} dxdy$$

$$= \left(\frac{N}{\pi}\right)^{2} \cdot ||\varphi||^{2}_{L^{2}(\mathbb{R}^{2})} \cdot (2\pi) \cdot \int_{\lambda_{+}-\lambda}^{\infty} e^{-Nr^{2}}r dr$$

$$= \left(\frac{N}{\pi}\right) \cdot ||\varphi||^{2}_{L^{2}(\mathbb{R}^{2})} \cdot e^{-N(\lambda_{+}-\lambda)^{2}}.$$

Hence, the second term is an exponentially small term. For the first term, we need to use Lemma 4.1.1 to replace $|K_N(x,y)|^2$ with $|e^{N(x\overline{y})}|^2$. By Lemma 4.1.1, we see that when $x, y \in B(0, \lambda_+)$, for suitably large N depending on λ_+ ,

(4.1.12)
$$\left| \left| K_N(x,y) \right|^2 - \left| e^{N(x\overline{y})} \right|^2 \right| \le C(\lambda_+) \cdot \frac{e^{(N \cdot |x||y|)}}{\sqrt{N}} \cdot (\delta(\lambda_+))^N \cdot 3e^{(N \cdot |x||y|)}.$$

So, recall that $d\mu_N(x) = \frac{N}{\pi} e^{-N|x|^2} dx$, we know that

(4.1.13)
$$\int_{B(0,\lambda_{+})\times B(0,\lambda_{+})} |\varphi(x) - \varphi(y)|^{2} \cdot \left| |K_{N}(x,y)|^{2} - |e^{N(x\overline{y})}|^{2} \right| d\mu_{N}(x) d\mu_{N}(y)$$

$$\leq \frac{6}{\pi^{2}} C(\lambda_{+}) \cdot (\delta(\lambda_{+}))^{N} \cdot N^{\frac{3}{2}} \cdot ||\varphi||^{2}_{L^{2}(\mathbb{R}^{2})} \cdot |B(0,\lambda_{+})|,$$

which is also an exponentially small term. Hence, up to a term R_N which is exponentially small as $N \to +\infty$,

(4.1.14)
$$\operatorname{Var}(X_N(\varphi)) = \frac{1}{2} \int_{B(0,\lambda_+) \times B(0,\lambda_+)} \left| \varphi(x) - \varphi(y) \right|^2 \cdot \left| e^{N(x\overline{y})} \right|^2 d\mu_N(x) d\mu_N(y) + R_N$$
$$= \frac{N^2}{2\pi^2} \int_{B(0,\lambda_+) \times B(0,\lambda_+)} \left| \varphi(x) - \varphi(y) \right|^2 \cdot e^{-N|x-y|^2} dxdy + R_N.$$

Therefore, we can directly obtain the following corollary by applying Theorem 3.2.2.

Corollary 4.1.3. There is a dimensional constant $T_1 > 0$, such that for any $\lambda \in (0, 1)$, and any $\varphi \in W_0^{1,2}(B(0, \lambda))$, we have that

(4.1.15)
$$\lim_{N \to \infty} \operatorname{Var}(X_N(\varphi)) = T_1 \int_{\mathbb{R}^2} |\nabla \varphi|^2 \, dx.$$

Notice that similar results with $\varphi \in W^{1,2}$ sense for the Gaussian kernel $e^{-N|x-y|^2}$ were already built up in [19]. Additionally, the authors of [19] discussed other kernels K(x,y)that are not of the form K(x-y) but can be bounded by another kernel J(x-y). For the Ginibre ensemble, in [20], it was shown that for $\varphi \in C_0^1(\mathbb{R}^2)$,

(4.1.16)
$$\lim_{N \to \infty} \operatorname{Var}(X_N(\varphi)) = \frac{1}{4\pi} ||\nabla \varphi||^2_{L^2(\mathbb{R}^2)} + \frac{1}{2} ||\varphi||^2_{H^{1/2}(\partial B(0,1))}$$

So, T_1 in Corollary 4.1.3 is $1/4\pi$. And the authors of [20] also built up a central limit theorem for the normalizations of $X_N(\varphi)$, which did not follow from the methods of [5, 23, 1] because $\operatorname{Var}(X_N(\varphi))$ is bounded now. It would also be interesting to extend those results in [20] to all $\varphi \in W_0^{1,2}(\mathbb{R}^2)$, which requires a deeper analysis of the behavior of the kernel $K_N(x, y)$ for the Ginibre ensemble near the boundary $\partial B(0, 1)$.

Now, consider $\varphi = \mathbb{1}_{\Omega}$ with $\overline{\Omega} \subset B(0,1) \subset \mathbb{R}^2$. Notice that in this case, $|\varphi(x) - \varphi(y)|^2 = |\varphi(x) - \varphi(y)|$. Therefore, if Ω is a set of finite perimeter (bounded Caccioppoli set), Corollary 3.1.4 together with (4.1.14) will imply the following corollaries.

Corollary 4.1.4. There is a dimensional constant $T_2 > 0$, such that

(4.1.17)
$$\lim_{N \to \infty} \frac{\operatorname{Var}(X_N(\mathbb{1}_{\Omega}))}{\sqrt{N}} = T_2 \cdot \mathcal{H}^1(\partial^* \Omega)$$

Or by Corollary 3.1.5, we also have the following.

Corollary 4.1.5. If Ω_1 and Ω_2 are two disjoint sets of finite perimeter with closures contained in $B(0,1) \subset \mathbb{R}^2$, then

(4.1.18)
$$\lim_{N \to \infty} \frac{\operatorname{Cov}(X_N(\mathbb{1}_{\Omega_1}), X_N(\mathbb{1}_{\Omega_2}))}{\sqrt{N}} = T_2 \cdot \mathcal{H}^1(\partial^* \Omega_1 \cap \partial^* \Omega_2).$$

Let $\Omega = \Omega(\eta)$ be the Koch snowflake of scale $\eta > 1$ in Section 3.3 with η satisfying the condition (3.3.14). Then, by Theorem 3.3.2 and (4.1.14), we know the following.

Corollary 4.1.6. There is a positive constant $T(\Omega)$ depending on Ω , which is also compatible with the Minkowski content of $\partial\Omega$, such that,

(4.1.19)
$$\lim_{i \to \infty} \frac{\operatorname{Var}(X_N(\mathbb{1}_{\Omega}))}{N^{(\alpha(\eta)/2)}} = T(\Omega),$$

where $\alpha(\eta) \in (1,2)$ satisfies (3.3.1) and is the Minkowski dimension of $\partial\Omega$.

Finally, if $\Omega \in \mathbb{R}^n$ satisfies the assumptions in Theorem 1.1.1 (equivalently, Corollary 2.0.3), then we can also get part (ii) of Corollary 1.2.2 by using (4.1.14) and Corollary 2.0.3.

Acknowledgements The author would like to thank Professor Paul Bourgade for introducing him to the probabilistic background of this problem and for the insightful discussions. The author's advisor, Professor Fang-Hua Lin, brought [4] and Professor Guido De Philippis brought [7, 8] to the author's attention, and the author also wants to thank them for their continuous help and support at Courant. The author also thanks Oren Yakir for his very helpful comments on the early draft of the manuscript.

References

- G. W. ANDERSON, A. GUIONNET, AND O. ZEITOUNI, <u>An introduction to random matrices</u>, vol. 118 of Camb. Stud. Adv. Math., Cambridge: Cambridge University Press, 2010.
- [2] J. BOURGAIN, H. BREZIS, AND P. MIRONESCU, <u>Another look at Sobolev spaces</u>, in Optimal control and partial differential equations. In honour of Professor Alain Bensoussan's 60th birthday. Proceedings of the conference, Paris, France, December 4, 2000, Amsterdam: IOS Press; Tokyo: Ohmsha, 2001, pp. 439–455.
- J. BUCKLEY AND M. SODIN, <u>Fluctuations of the increment of the argument for the Gaussian entire</u> <u>function</u>, J. Stat. Phys., 168 (2017), pp. 300–330.
- [4] L. CAFFARELLI, J.-M. ROQUEJOFFRE, AND O. SAVIN, <u>Nonlocal minimal surfaces</u>, Commun. Pure Appl. Math., 63 (2010), pp. 1111–1144.
- [5] O. COSTIN AND J. L. LEBOWITZ, <u>Gaussian fluctuation in random matrices</u>, Physical Review Letters, 75 (1995), p. 69.
- [6] J. DÁVILA, On an open question about functions of bounded variation., Calc. Var. Partial Differ. Equ., 15 (2002), pp. 519–527.
- [7] E. DE GIORGI, <u>Definizione ed espressione analitica del perimetro di un insieme</u>, Atti Accad. Naz. Lincei, VIII. Ser., Rend., Cl. Sci. Fis. Mat. Nat., 14 (1953), pp. 390–393.
- [8] —, Su una teoria generale della misura (r-1)-dimensionale in uno spazio ad r dimensioni, Ann. Mat. Pura Appl. (4), 36 (1954), pp. 191–213.
- [9] K. FALCONER, <u>Fractal geometry. Mathematical foundations and applications</u>, Chichester: Wiley, 2nd ed. ed., 2003.

- [10] W. FELLER, <u>An introduction to probability theory and its applications. Vol. II. 2nd ed.</u>, Wiley Ser. Probab. Math. Stat., John Wiley & Sons, Hoboken, NJ, 1971.
- [11] P. J. FORRESTER AND G. HONNER, <u>Exact statistical properties of the zeros of complex random</u> polynomials, J. Phys. A, Math. Gen., 32 (1999), pp. 2961–2981.
- [12] D. GATZOURAS, <u>Lacunarity of self-similar and stochastically self-similar sets</u>, Trans. Am. Math. Soc., 352 (2000), pp. 1953–1983.
- [13] D. S. JERISON AND C. E. KENIG, <u>Boundary behavior of harmonic functions in non-tangentially</u> accessible domains, Adv. Math., 46 (1982), pp. 80–147.
- [14] L. LOMBARDINI, <u>Fractional perimeters from a fractal perspective</u>, Adv. Nonlinear Stud., 19 (2019), pp. 165–196.
- [15] O. MACCHI, <u>The coincidence approach to stochastic point processes</u>, Adv. Appl. Probab., 7 (1975), pp. 83–122.
- [16] F. MAGGI, Sets of finite perimeter and geometric variational problems. An introduction to geometric measure theory, vol. 135 of Camb. Stud. Adv. Math., Cambridge: Cambridge University Press, 2012.
- [17] J. M. MAZÓN, J. D. ROSSI, AND J. TOLEDO, <u>Nonlocal perimeter</u>, curvature and minimal surfaces for measurable sets, J. Anal. Math., 138 (2019), pp. 235–279.
- [18] D. MEYER, Snowballs are quasiballs, Trans. Am. Math. Soc., 362 (2010), pp. 1247–1300.
- [19] B. RIDER AND B. VIRAG, <u>Complex determinantal processes and H¹ noise</u>, Electron. J. Probab., 12 (2007), pp. 1238–1257.
- [20] B. RIDER AND B. VIRÁG, <u>The noise in the circular law and the Gaussian free field</u>, Int. Math. Res. Not., 2007 (2007), p. 32. Id/No rnm006.
- [21] M. SODIN, A. WENNMAN, AND O. YAKIR, <u>The random weierstrass zeta function i. existence</u>, uniqueness, fluctuations, arXiv preprint arXiv:2210.09882, (2022).
- [22] _____, The random weierstrass zeta function ii. fluctuations of the electric flux through rectifiable curves, arXiv preprint arXiv:2211.01312, (2022).
- [23] A. SOSHNIKOV, Determinantal random point fields, Russ. Math. Surv., 55 (2000), pp. 923–975.
- [24] T. TORO, <u>Analysis and geometry on non-smooth domains</u>, Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales, 41 (2017), pp. 521–527.
- [25] S. TORQUATO, A. SCARDICCHIO, AND C. E. ZACHARY, <u>Point processes in arbitrary dimension from fermionic gases, random matrix theory, and number theory</u>, J. Stat. Mech. Theory Exp., 2008 (2008), p. 39. Id/No p11019.

COURANT INSTITUTE OF MATHEMATICAL SCIENCES, NEW YORK UNIVERSITY, 251 MERCER STREET, NEW YORK, NY10012, USA

Email address: malin@nyu.edu