# CLASSIFICATION STRENGTH OF POLISH GROUPS I: INVOLVING $S_{\infty}$

SHAUN ALLISON

ABSTRACT. In recent years, much work has been done to measure and compare the complexity of orbit equivalence relations, especially for certain classes of Polish groups. We start by introducing some language to organize this previous work, namely the notion of classification strength of Polish groups. Broadly speaking, a Polish group G has stronger classification strength than H if every orbit equivalence relation induced by a continuous action of H on a Polish space can be "emulated" by such an action of G in the sense of Borel reduction.

Among the non-Archimedean Polish groups, the groups with the highest classification strength are those that involve  $S_{\infty}$ , the Polish group of permutations of a countably-infinite set. We prove that several properties, including a weakening of the disjoint amalgamation in Fraïssé theory, a weakening of the existence of an absolute set of generating indiscernibles, and not having ordinal rank in a particular coanalytic rank function, are all equivalent to a non-Archimedean Polish group involving  $S_{\infty}$ . Furthermore, we show the equivalence relation  $=^+$ , which is a relatively simple benchmark equivalence relation in the theory of Borel reducibility, can only be classified by such groups that involve  $S_{\infty}$ .

### 1. INTRODUCTION

Invariant descriptive set theory is concerned with definable equivalence relations and definable reductions between them. In particular, we usually consider equivalence relations living on Polish spaces, where the equivalence relations are *analytic*, i.e. analytic as a subset of the product space. The definable reductions that we consider are usually the *Borel* ones. To be specific, given analytic equivalence relations E and F living on Polish spaces X and Y respectively, a **Borel reduction** from E to F is a Borel function  $f: X \to Y$  satisfying x E x' iff f(x) F f(x') for any  $x, x' \in X$ . When such a reduction exists, we say that E is *Borel-reducible* to F, i.e.  $E \leq_B F$ . In the case that E and F represent classification problems in math, then from  $E \leq_B F$  we can conclude that the problem E is no harder than F.

One common way to show that E does not Borel reduce to F is by showing that E is generically-ergodic with respect to F. A function  $f: X \to Y$  is a homomorphism from E to F iff for every  $x, y \in X$ , if  $x \in y$  then  $f(x) \in f(y)$ . Then we say that E is generically-ergodic with respect to F iff for every Borel homomorphism  $f: X \to Y$  from E to F, there is a comeager  $C \subseteq X$  such that for any  $x, y \in C$  we have  $f(x) \in f(y)$ . In the case that E does not have a comeager class, this precludes the existence of a Borel reduction.

While comparing equivalence relations has value, it is perhaps more valuable to give an absolute notion of complexity of a given equivalence relation, which is invariant under Borel reducibility. One very successful program along these lines has been, in the case of Borel

Date: April 4, 2023.

<sup>2020</sup> Mathematics Subject Classification. Primary 54H05, 37B02, 54H11; Secondary 03E15, 03C15.

Key words and phrases. Polish group, Borel reduction, Knight's model, countable model, Fraïssé limit.

equivalence relations, to consider their place in the Borel hierarchy via a notion of *potential* Borel complexity. This was essentially initiated in the seminal [HKL98], where a connection to potential Borel complexity was made to the set-theoretic complexity of any definable assignment of invariants. In particular, in the large class of equivalence relations that they consider, an equivalence relation is potentially  $\Pi_2^0$  iff there is a Borel assignment of reals as invariants, potentially  $\Pi_3^0$  iff there is a Borel assignment of sets of reals as invariants, potentially  $\Pi_4^0$  iff there is a Borel assignment of sets of reals as invariants, and so on.

The case of **orbit equivalence relations** gives another opportunity to produce a meaningful hierarchy of complexity, this time by studying the acting ("classifying") groups. Many definable equivalence relations that arise in practice are orbit equivalence relations. Given a Polish group G, a Polish G-space is a Polish space X along with a continuous action  $G \curvearrowright X$ . We use  $E_X^G$  to denote the orbit equivalence relation, which is analytic, and moreover every orbit is Borel.

Formally, given an equivalence relation E on a Polish space X, we say that a Polish group G classifies E iff there is a Polish G-space Y such that  $E \leq_B E_Y^G$ . This notion can be used to separate equivalence relations up to Borel reduction. We identify three examples of this phenomenon of particular importance.

(1) We say that an equivalence relation E is classifiable by countable structures iff it is classifiable by  $S_{\infty}$ , the Polish group of automorphisms of a countably-infinite set with the natural topology.

An important benchmark equivalence relation is  $=^+$ , the Friedman-Stanley jump of equality, which has particular importance in the study of potential Borel complexity in [HKL98]. It is defined on  $\mathbb{R}^{\omega}$  by

$$(x_n) =^+ (y_n) \quad \text{iff} \quad \{x_n \mid n \in \omega\} = \{y_n \mid n \in \omega\}$$

This is easily seen to be classifiable by countable structures as witnessed by the natural Bernoulli-shift action  $S_{\infty} \curvearrowright \mathbb{R}^{\omega}$ .

Hjorth showed that any orbit equivalence relation that is generically-ergodic with respect to  $=^+$  is also generically-ergodic with respect to any action of  $S_{\infty}$ . This implies in any case where there isn't a comeger orbit that the equivalence relation is not classifiable by countable structures. Hjorth also isolated a dynamical property of some orbit equivalence relations called *turbulence*, which precludes them from being classifiable by countable structures (see [Hjo00] and [Hjo02a]).

(2) Among the orbit equivalence relations classifiable by countable structures, further separations can be found. Recall that a Polish group is *complete left-invariant* (or cli) iff it has a complete metric  $d_G$  which compatible in the sense that it generates the topology, and furthermore is left-invariant, i.e.

$$d_G(g,g') = d_G(hg,hg')$$

for every group elements g, g', h. In [Hjo99], Hjorth identifies a metamathematical property of equivalence relations that preclude them from being classifiable by cli Polish groups, and shows that  $=^+$  exhibits this property. This property was further explored and dubbed being "unpinned" in [Kan08].

(3) A stronger invariance notion than a Polish group being cli is being two-sided invariant (or tsi). A Polish group is tsi iff there is a compatible complete metric  $d_G$  satisfying

$$d_G(g,g') = d_G(hg,hg') = d_G(gh,g'h)$$

for every group elements g, g', h.

A natural example of an equivalence relation classifiable by a tsi Polish group is  $E_{\infty}$ , defined as the orbit equivalence relation induced by the Bernoulli-shift action of  $F_2 \curvearrowright \mathbb{R}^{F_2}$ , where  $F_2$  is the free group on two generators with the discrete topology. Since  $F_2$  is a countable discrete group, it is tsi for trivial reasons.

In [All], it was shown that an orbit equivalence relation that is generically ergodic with respect to  $E_{\infty}$  is generically ergodic with respect to any orbit equivalence relation induced by a tsi non-Archimedean Polish group. This strongly parallels Hjorth's result relating generic ergodicity with respect to  $=^+$  and orbit equivalence relations induced by any non-Archimedean Polish group.

In [CC22] Clemens-Coskey introduced an equivalence relation  $E_0^{[\mathbb{Z}]}$ , called the  $\mathbb{Z}$ -jump of  $E_0$ , which is classifiable by a cli Polish group. They asked if it is in fact classifiable by a tsi Polish group. However they showed that it is generically ergodic with respect to  $E_{\infty}$  and furthermore has all meager classes, thus by [All] it is not classifiable by a tsi non-Archimedean Polish group.

This was generalized by the author and Panagiotopoulos in [AP21] to general tsi Polish groups, and a purely dynamical property was identified similar to Hjorth's notion of turbulence which serves as an obstruction to classifiability by tsi Polish groups. In particular,  $E_0^{[\mathbb{Z}]}$  is not classifiable by *any* tsi Polish group. We will not need the definition of  $E_0^{[\mathbb{Z}]}$  in this work, but its definition, as well as an exploration of an interesting hierarchy of similar equivalence relations, can be found in [CC22].

We can observe that by considering properties of the classifying group we can now start to see that it produces a meaningful hierarchy of equivalence relations which are classifiable by countable structures:

# classifiable by non-Archimedean TSI $\subsetneq$ classifiable by non-Archimedean CLI $\subsetneq$ classifiable by $S_{\infty}$ .

In another upcoming paper, we are showing with Panagiotopoulos that there is a finer hierarchy below classifiability by CLI, and in this paper we expose another hierarchy which exists above classifiability by CLI. With this picture in mind, the following definition seems natural:

**Definition 1.1.** Say that G is stronger in classification strength than H iff for every Polish H-space X, the orbit equivalence relation  $E_X^H$  is classifiable by G.

The following weak restatement of a result of Mackey and Hjorth further motivates this definition. Recall that G involves H iff there is a closed subgroup  $G_0$  of G and a continuous surjective homomorphism from  $G_0$  onto H. Note that this is also sometimes said "H divides G".

**Lemma 1.2** (Mackey, Hjorth). If G involves H, then G is stronger in classification strength than H.

The non-Archimedean Polish groups are exactly those that are isomorphic to closed subgroups of  $S_{\infty}$ . Thus the non-Archimedean Polish groups which involve  $S_{\infty}$  has maximal classification strength among the non-Archimedean Polish groups. A result of Hjorth implies the converse [Hjo01] (see also the more recent [LUb] which recovers this result using a different strategy.)

Hjorth's result gives a metamathematical sufficient condition for a non-Archimedean Polish group to involve  $S_{\infty}$ . On the other hand, a paper of Baldwin-Friedman-Koerwien-Laskowski contains an argument that the automorphism group of the limit of any Fraïssé class which satisfies disjoint amalgamation involves  $S_{\infty}$ . However, not much else was known, and there had not yet been any comprehensive study of the division between the Polish groups which do and do not involve  $S_{\infty}$ .

In this paper, we identify several seemingly unconnected properties of non-Archimedean Polish groups, which are equivalent to involving  $S_{\infty}$ . This tells us, we believe, that such groups have an interesting and deep structure and are worthy of further study.

Our main result is the following, where the terms mentioned in equivalences (2), (3), (4), and (5) are yet to be defined.

**Theorem 1.3.** Let  $G = Aut(\mathcal{M})$  be a non-Archimedean Polish group. Then TFAE:

- (1) G does not involve  $S_{\infty}$ ;
- (2) Every disjointifying closure operator on G is trivial;
- (3)  $Krk(\mathcal{M}) < \infty;$
- (4)  $Krk(\mathcal{M}) < \omega_1;$
- (5) There is no indiscernible support function on  $\mathcal{M}$ ;
- (6) G does not classify  $=^+$ ;

In the sequel, we will add several more equivalences onto the list, as well as initiate a finer study of the hierarchy of Polish groups not involving  $S_{\infty}$  which arises from the rank function Krk.

1.1. Acknowledgements. The author would like to thank Omer Ben-Neria, Clinton Conley, Aristotelis Panagiotopoluos, and Spencer Unger for many valuable conversations. This work was partially funded by the Israel Science Foundation (grant no. 1832/19).

# 2. Preliminaries

2.1. Countable model theory. We briefly review a few very basic concepts and notation from the model theory of countable structures. We will always use  $\mathcal{L}$  to refer to a countable relational language. Given an  $\mathcal{L}$ -structure  $\mathcal{M}$ , we will write M to refer to the underlying set of  $\mathcal{M}$ . If  $a, a' \in M$  and  $B \subseteq M$ , we write  $a \cong_B a'$  iff there is an automorphism  $\pi$  of  $\mathcal{M}$ satisfying  $\pi(a) = a'$  and  $\pi(b) = b$  for every  $b \in B$ . We write  $\mathcal{N} \leq_{\mathcal{L}} \mathcal{M}$  to denote that  $\mathcal{N}$  is an  $\mathcal{L}$ -substructure of  $\mathcal{M}$ , and we write  $\mathcal{N} \prec_{\mathcal{L}_{\omega_1,\omega}}$  $\mathcal{M}$  iff  $\mathcal{N}$  is an  $\mathcal{L}_{\omega_1,\omega}$ -elementary substructure of  $\mathcal{M}$ . Equivalently,  $\mathcal{N} \prec_{\mathcal{L}_{\omega_1,\omega}} \mathcal{M}$  iff for every finite  $B \subseteq N$  and  $a \in M$ , there is some  $a' \in N$  such that  $a' \cong_B a$ .

We say  $\mathcal{M}$  is *ultrahomogeneous* iff for any two tuples  $\bar{a}$  and  $\bar{b}$  from M, if they have the same quantifier-free type, then there is an automorphism sending one to the other.

2.2. Non-Archimedean Polish groups. A Polish group is called *non-Archimedean* if it has a countable local basis of the identity of open subgroups. The most important such group is  $S_{\infty}$ , the Polish group of permutations of a countably-infinite set (with the discrete topology), equipped with the pointwise convergence topology. There are several useful equivalent formulations of a Polish group being non-Archimedean:

Fact 2.1. Let G be a Polish group. The following are equivalent:

(1) G is non-Archimedean

(2) G is isomorphic to  $Aut(\mathcal{M})$  for a countable structure  $\mathcal{M}$  in a countable language;

- (3) G is isomorphic to a closed subgroup of  $S_{\infty}$ ;
- (4) G has a compatible complete ultrametric.

See [Gao09, Theorem 2.4.1, Theorem 2.4.4] for proofs.

We will make the most use of equivalence (2), as it allows the use of language and techniques from model theory, even though our use of model theory will not particularly deep. An important point is that we can assume that  $\mathcal{M}$  is ultrahomogeneous by adding new relations without changing the set of automorphisms. This is done as follows: for each tuple  $\bar{b}$  from  $\mathcal{M}$ , add a new relation  $R_{\bar{b}}$  to the language and declare  $R_{\bar{b}}^{\mathcal{M}}(\bar{a})$  iff there is an automorphism of  $\mathcal{M}$  sending  $\bar{a}$  to  $\bar{b}$ .

2.3. Non-Archimedean Polish groups that are cli and not cli. Recall that a Polish group is cli iff it has a complete left-invariant metric which is compatible with the topology. In the case of a non-Archimedean Polish group, we have some nice characterizations due to Gao.

**Fact 2.2** (Gao). Let  $G = Aut(\mathcal{M})$  be a non-Archimedean Polish group. Then TFAE:

- (1) G is cli
- (2)  $\mathcal{M}$  has no nontrivial  $\mathcal{L}_{\omega_1,\omega}$ -elementary substructure;
- (3) there is no uncountable model of the Scott sentence of  $\mathcal{M}$

There is also a nice rank function due to Deissler which characterizes when G is cli, which we will define and discuss later.

A Polish group is tsi iff it has a complete two-sided metric which is compatible with the topology. An example of a Polish group which is cli but not tsi is the automorphism group of the linear order with order-type  $\mathbb{Z} * \mathbb{Z}$ , which is just the lex order on  $\mathbb{Z} \times \mathbb{Z}$ . A natural action of this group induces the equivalence relation  $E_0^{[\mathbb{Z}]}$  mentioned in the introduction, discussed further in [CC22], [All], [AP21].

The most important non-cli Polish group from our perspective is the automorphism group of Knight's model. It is significant in the sense that was the simplest known Polish group which is not cli but does not involve  $S_{\infty}$ . Knight's model  $\mathcal{K}$  is a countable structure in the language  $\mathcal{L} = \{\langle, f_n\}_{n \in \omega}, \text{ where } \langle \rangle$  is a binary relation and each  $f_n$  is a unary function which satisfies:

- (1) < is a linear order on K;
- (2) for every  $a \in K$ ,  $\{b \in K \mid b < a\} = \{f_n(a) \mid n \in \omega\};$
- (3)  $\mathcal{K}$  is 1-transitive (there is an automorphism between any two pairs of elements);
- (4) there is a non-trivial  $\mathcal{L}_{\omega_1,\omega}$ -substructure of  $\mathcal{K}$ .

By Gao's characterization, (4) implies that  $\operatorname{Aut}(\mathcal{K})$  is not cli. Hjorth showed [Hjo99] that it does not classify  $=^+$  and thus does not involve  $S_{\infty}$  (it also follows from our results in this paper).

2.4. Examples of non-Archimedean Polish groups involving  $S_{\infty}$ . The original of Knight's model was to produce a countable structure whose Scott sentence has a model of cardinality  $\aleph_1$ , but no larger. It is said that the Scott sentence of Knight's model "characterizes"  $\aleph_1$ . In [Hjo02b], Hjorth constructed, for each  $\alpha$ , a countable structure characeterizing  $\aleph_{\alpha}$ . Rather than a generalization of Knight's model, Hjorth's construction relies on a very different construction method which resulted in structures whose automorphism groups involve  $S_{\infty}$ .

Hjorth's model characterizing  $\aleph_1$ , which we denote by  $\mathcal{H}$ , is a countable ultrahomogeneous structure in a language with binary relations  $S_n$  for every n, and k + 2-ary relations  $S_k$  for every k, satisfying

- (1) there is some function  $f: [H]^2 \to \omega$  such that for every a and b in H and  $n \in \omega$ , we have  $S_n^{\mathcal{H}}(a, b)$  iff  $f(\{a, b\}) = n$ ;
- (2) for every a and b, the set  $S(\{a, b\})$  of all c such that  $f(\{a, c\}) = f(\{b, c\})$  is finite, and  $R_k^{\mathcal{H}}(a, b, \bar{c})$  iff  $\bar{c}$  enumerates  $S(\{a, b\})$ .

It's straightforward to check that every  $\mathcal{L}_{\omega,\omega}$ -substructure any model of the Scott sentence of  $\mathcal{H}$  must be countable. Thus there are no models of the Scott sentence of  $\mathcal{H}$  of cardinality  $\aleph_2$  or higher. On the other hand, Hjorth showed that the automorphism group of  $\mathcal{H}$  involves  $S_{\infty}$  (in the process of showing that its Scott sentence has an uncountable model). This is done by exploiting the fact that  $\mathcal{H}$  satisfies disjoint amalgamation. The fact that the automorphism groups of limits of Fraïssé structures satisfying disjoint amalgamation is stated and proved more explicitly in [BFKL16]. Another proof of this, in even greater detail, appeared recently in [LUa]. We briefly sketch this argument in Section 3, as we will build on this idea.

A classic example of an automorphism group that involves  $S_{\infty}$  is  $Aut(\mathbb{Q}, <)$ , the automorphism group of the rational linear order. To see that it involves  $S_{\infty}$ , partition  $\mathbb{Q}$  into countably-many dense subsets  $A_n$ . Let  $H \leq Aut(\mathbb{Q})$  be the closed subgroup of automorphisms  $\pi$  satisfying that for every  $n, m \in \omega$  and  $a, b \in A_n$ , then  $\pi(a) \in A_m$  iff  $\pi(b) \in A_m$ . Then define a continuous homomorphism  $f : H \to S_{\infty}$  where  $f(\pi)$  is defined to be the unique  $\sigma \in S_{\infty}$  such that  $a \in A_m$  iff  $\pi(a) \in A_{\sigma(m)}$  for every a and m. To see that f is indeed surjective, apply a back-and-forth argument.

We will construct one more example, which we will actually make use of later. Let  $\Delta$  be some countably-infinite group, and consider a free action  $\Delta \curvearrowright I$  on a countably-infinite set I with infinitely-many  $\Delta$ -orbits. Now consider the closed subgroup  $P \leq S_I$  of permutations  $\pi$  of I satisfying  $\pi(\delta \cdot x) = \delta \cdot \pi(x)$  for every  $x \in I$ .

This group involves  $S_{\infty}$ . To see this, fix a transversal  $T \subseteq I$ , i.e. a set which intersects every  $\Delta$ -orbit exactly once. Since T is a countably-infinite set, the Polish group  $S_T$  is isomorphic to  $S_{\infty}$ . Moreover, the map  $f : P \to S_T$  sending each  $\pi \in P$  to the unique  $\sigma \in S_T$  satisfying  $\pi(x) \in \Delta \cdot \sigma(x)$  for every  $x \in T$ , is easily seen to be a continuous surjective homomorphism.

2.5. Baire-measurable homomorphisms and the orbit continuity lemma. Given a Polish group G, recall that a Polish G-space is a Polish space X along with a continuous action  $G \curvearrowright X$ . We write the induced orbit equivalence relation as  $E_X^G$ . Given two such orbit equivalence relations  $E_X^G$  and  $E_Y^H$ , a homomorphism is a function  $f: X \to Y$  such that  $f(x) E_Y^H f(y)$  whenever  $x E_X^H y$ .

The following so-called "orbit continuity lemma", due to Hjorth and Lupini-Panagiotopoulos, is central. This is the main way to extract information from the existence of Baire-measurable homomorphisms that don't trivialize on a comeager set. Our statement differs slightly from the statement in [LP18], so we give an argument for how to recover this one from theirs.

**Lemma 2.3** (Hjorth, Lupini-Panagiotopoulos). Suppose  $f : X \to Y$  is a Baire-measurable homomorphism from  $E_X^G$  to  $E_Y^H$  and  $G_0$  is a countable dense subgroup of G. Then there is a comeager subset  $C \subseteq X$  such that

- (1) f is continuous on C;
- (2) for every  $x \in C$ , the set of g such that  $g \cdot x \in C$  is comeager;
- (3) for every  $x \in C$  and for every  $g \in G_0, g \cdot x \in C$ ;
- (4) for every  $x_0 \in C$  and every nonempty open neighborhood  $V \subseteq H$  of the identity, there is a neighborhood  $U_0$  of  $x_0$  and a nonempty open neighborhood  $W \subseteq G$  of the identity such that for any  $x \in C \cap U$  and for every  $w \in W$  with  $w \cdot x \in C \cap U_0$ , we have  $f(w \cdot x) \in V \cdot f(x)$ ; and
- (5) for every  $x_0 \in C$  and  $g \in G_0$  and nonempty open  $W \subseteq H$  such that  $f(g \cdot x) \in W \cdot f(x)$ , there is an open neighborhood  $U_0$  of  $x_0$  such that for every  $x \in C \cap U_0$ ,  $g \cdot x \in C$  and  $f(g \cdot x) \in W \cdot f(x)$ ;

*Proof.* Let  $C_0$  be a comeager set satisfying the first three conditions as in [LP18, Lemma 2.5]. For every  $g \in G_0$  and basic open W, let  $C_{g,W}$  be a comeager set on which the Baire-measurable function  $f: X \to 2$  defined by

$$f(x) = 1 \iff f(gx) \in Wf(x)$$

is continuous. Let C' be the intersection of  $C_0$  and each  $C_{g,W}$  and define

 $C = \{ x \in C' \mid \forall^* g, gx \in C' \text{ and } \forall g \in G_0, gx \in C' \}.$ 

We argue that this works.

The fact that (1)-(3) and (5) are satisfied are immediate, so we proceed to (4). Clause (4) is almost the same as the clause (3) of [LP18, Lemma 2.5], with the difference is that we quantify over all  $w \in W$ , not just a set comeager in W, with the additional assumption that  $w \cdot x \in U_0$ . Let  $x_0 \in C$  and  $V \subseteq H$  be an open neighborhood of the identity of H. Let  $\hat{V} \subseteq H$  be a symmetric open neighborhood of the identity of H such that  $\hat{V}^2 \subseteq V$ . From our choice of C, there is an open neighborhood  $U_0 \ni x_0$  and an open neighborhood  $W \subseteq G$ of the identity of G such that for every  $x \in C \cap U$  there is a comeager set of  $w \in W$  such that  $f(w \cdot x) \in \hat{V} \cdot f(x)$ .

Let  $x \in U_0 \cap C$  and  $w \in W$  such that  $w \cdot x \in U_0 \cap C$ . Let  $D_0$  be the set of  $w' \in W$  such that  $f(w' \cdot x) \in \hat{V} \cdot f(x)$  and let  $D_1$  be the set of  $w' \in W$  such that  $f(w'w \cdot x) \in \hat{V} \cdot f(w \cdot x)$ . The set  $D_0$  is comeager in W and  $D_1w$  is nonmeager in W, thus we may fix some  $w' \in D_0 \cap D_1w$ . Then we have that  $f(w' \cdot x) \in \hat{V} \cdot x$  and  $f(w'w^{-1} \cdot (w \cdot x)) \in \hat{V} \cdot (w \cdot x)$ . Thus  $f(w \cdot x) \in \hat{V}^{-1} \cdot f(x) \subseteq V \cdot f(x)$  as desired.  $\Box$ 

## 3. A weakening of disjoint amalgamation

In [Hj002b], and more explicitly in [BFKL16], disjoint amalgamation is identified as a sufficient condition for the automorphism group of a structure to involve  $S_{\infty}$ . In this section, we introduce the appropriate weakening of disjoint amalgamation. We first give a brief review of the Fraïssé theory of classes of finite structures, and review the ideas of [BFKL16]. Then we introduce the weakening of disjoint amalgamation and prove that it is necessary and sufficient.

3.1. Fraïssé theory. We start with a brief presentation of the generalized Fraïssé theory which we will later need. A general survey of generalized Fraïssé theory can be found in [Kru15].

Let  $\mathcal{L}$  be a countable relational language. Let  $\mathcal{F}$  be a class of finite  $\mathcal{L}$ -structures closed under isomorphism. Let  $\leq_{\mathcal{F}}$  be a notion of "strong substructure" on  $\mathcal{F}$ , i.e. a transitive and reflexive binary relation satisfying

- (1) if  $\mathcal{A} \preceq_{\mathcal{F}} \mathcal{B}$  then  $\mathcal{A} \preceq_{\mathcal{L}} \mathcal{B}$ ; and
- (2) if  $\mathcal{A} \leq_{\mathcal{F}} \mathcal{B}$  and  $\pi : \mathcal{B} \cong \mathcal{C}$  is an isomorphism, then  $\pi[\mathcal{A}] \leq_{\mathcal{F}} \mathcal{C}$ .

We write  $\mathcal{B} \cong \mathcal{C}$  iff there is an isomorphism  $\pi : B \to C$  between  $\mathcal{B}$  and  $\mathcal{C}$ . More generally, if  $A \subseteq B \cap C$ , then we write  $\mathcal{B} \cong_A \mathcal{C}$  iff there is an isomorphism  $\pi : B \to C$  between  $\mathcal{B}$  and  $\mathcal{C}$  satisfying  $\pi(a) = a$  for every  $a \in A$ .

The following is standard.

**Proposition 3.1.** Suppose  $(\mathcal{F}, \preceq_{\mathcal{F}})$  satisfies the following:

- (i) the empty structure  $\emptyset$  is in  $\mathcal{F}$  and  $\emptyset \preceq_{\mathcal{F}} \mathcal{A}$  for every  $\mathcal{A} \in F$ ; and
- (ii) (amalgamation property) for any  $\mathcal{A}, \mathcal{B}, \mathcal{C} \in \mathcal{F}$ , if  $\mathcal{A} \preceq_{\mathcal{F}} \mathcal{B}$  and  $\mathcal{A} \preceq_{\mathcal{F}} \mathcal{C}$  then there is some  $\mathcal{C}' \in \mathcal{F}$  satisfying  $\mathcal{C}' \cong_{\mathcal{A}} \mathcal{C}$  and  $\mathcal{D} \in \mathcal{F}$  such that  $\mathcal{B} \preceq_{\mathcal{F}} \mathcal{D}$  and  $\mathcal{C}' \preceq_{\mathcal{F}} \mathcal{D}$ .

Then there is a unique (up to isomorphism) countable  $\mathcal{L}$ -structure  $\mathcal{M}$  satisfying:

- (1) for any finite  $D \subseteq M$ , there is some  $\mathcal{A} \in \mathcal{F}$  with  $D \subseteq A$  and  $\mathcal{A} \preceq_{\mathcal{L}} M$ ;
- (2) if  $\mathcal{A} \preceq_{\mathcal{F}} \mathcal{B}$  and  $\mathcal{A} \preceq_{\mathcal{L}} M$ , then there is some  $\mathcal{B}' \in \mathcal{F}$  with  $\mathcal{A} \preceq_{\mathcal{F}} \mathcal{B}', \mathcal{B}' \preceq_{\mathcal{L}} M$  and  $\mathcal{B}' \cong_{\mathcal{A}} \mathcal{B}$ ; and
- (3) if  $\mathcal{A} \preceq_{\mathcal{F}} \mathcal{B}$  and  $\mathcal{A} \preceq_{\mathcal{F}} \mathcal{B}'$  and  $\mathcal{A}, \mathcal{B}, \mathcal{B}' \preceq_{\mathcal{L}} M$ , if  $\mathcal{B} \cong_{A} \mathcal{B}'$  then there is an automorphism  $\pi$  of  $\mathcal{M}$  such that  $\pi[B] = B'$  and  $\pi(a) = a$  for every  $a \in A$ .

We note that any class  $(\mathcal{F}, \preceq_{\mathcal{F}})$  satisfying the assumptions of the proposition is a generalized Fraïssé class as defined in [Kru15], however the notion of generalized Fraïssé class introduced there is much more general. Classically, the definition of Fraïssé class might include other axioms. The joint embedding property axiom follows from the amalgamation property and the fact that we require the empty structure to be a strong substructure of every structure in  $\mathcal{F}$ . The hereditary property is commonly required, but isn't necessary (or desired) in this context. Often it is required that  $\mathcal{F}$  satisfies the essential countability axiom, which already follows in our case from the fact that  $\mathcal{L}$  is countable and every structure in  $\mathcal{F}$ is finite. For the rest of this section, we will refer to any  $(\mathcal{F}, \preceq_{\mathcal{F}})$  satisfying the assumptions of Proposition 3.1 as a Fraïssé class.

We call  $\mathcal{M}$  in the conclusion of the proposition as the **limit** of  $(\mathcal{F}, \preceq_{\mathcal{F}})$ . The structure  $\mathcal{M}$  is ultrahomogenous, which follows by a back-and-forth argument using property (3) of Proposition 3.1.

Conversely, if  $\mathcal{M}$  is an arbitrary ultrahomogeneous  $\mathcal{L}$ -structure, then  $(\operatorname{Age}(\mathcal{M}), \preceq_{\mathcal{L}})$  is a Fraïssé class and its limit is isomorphic to  $\mathcal{M}$ . Recall that  $\operatorname{Age}(\mathcal{M})$  is the class of all finite substructures of  $\mathcal{M}$ , closed under isomorphism.

3.2. Disjoint amalgamation property. Let  $\mathcal{L}$  be a countable relational language and let  $(\mathcal{F}, \leq_{\mathcal{F}})$  be a Fraïssé class. Say that  $(\mathcal{F}, \leq_{\mathcal{F}})$  satisfies the disjoint amalgamation property (also called the strong amalgamation property) iff for every  $\mathcal{A}, \mathcal{B}, \mathcal{C} \in \mathcal{F}$  with  $\mathcal{A} \leq_{\mathcal{F}} \mathcal{B}$  and  $\mathcal{A} \leq_{\mathcal{F}} \mathcal{C}$ , there is some  $\mathcal{B}', \mathcal{D} \in \mathcal{F}$  such that  $\mathcal{B}' \cong_{\mathcal{A}} \mathcal{B}$  and  $\mathcal{B}' \leq_{\mathcal{F}} \mathcal{D}$  and  $\mathcal{C} \leq_{\mathcal{F}} \mathcal{D}$ , and moreover  $\mathcal{B}' \cap \mathcal{C} = \mathcal{A}$ . Notice that this is property (ii) in Proposition 3.1, with the additional requirement that  $\mathcal{B}'$  and  $\mathcal{C}$  be as disjoint as possible.

**Proposition 3.2** (Baldwin-Friedman-Koerwien-Laskowski, [BFKL16]). If  $(\mathcal{F}, \preceq_{\mathcal{F}})$  is a Fraissé class satisfying the disjoint amalgamation property, and  $\mathcal{M}$  is the limit, then  $Aut(\mathcal{M})$  involves  $S_{\infty}$ .

The proof proceeds by considering the class  $\mathcal{F}^*$  of "colored" versions of structures in  $\mathcal{F}$ . More precisely, we consider pairs  $(\mathcal{A}, \mathbf{c}_{\mathcal{A}})$  where  $\mathcal{A} \in \mathcal{F}$  and  $\mathbf{c}_{\mathcal{A}} : \mathcal{A} \to \omega$ . We could view these formally as  $\mathcal{L}^*$ -structures, where  $\mathcal{L}^* \supseteq \mathcal{L}$  is the language adding countably-many new unary relations. We say  $(\mathcal{A}, \mathbf{c}_{\mathcal{A}})$  is a strong substructure of  $(\mathcal{B}, \mathbf{c}_{\mathcal{B}})$ , which we write as  $(\mathcal{A}, \mathbf{c}_{\mathcal{A}}) \preceq_{\mathcal{F}}^* (\mathcal{B}, \mathbf{c}_{\mathcal{B}})$ , iff  $\mathcal{A} \preceq_{\mathcal{F}} \mathcal{B}$  and  $\mathbf{c}_{\mathcal{B}}(a) = \mathbf{c}_{\mathcal{A}}(a)$  for every  $a \in \mathcal{A}$ . The class  $(\mathcal{F}^*, \preceq_{\mathcal{F}}^*)$  need not satisfy the amalgamation property in general. However, with the additional assumption that  $(\mathcal{F}^*, \preceq_{\mathcal{F}})$  has the disjoint amalgamation property, it is easy to confirm that  $(\mathcal{F}^*, \preceq_{\mathcal{F}}^*)$ , satisfies the amalgamation property. Thus we can compute the limit of  $(\mathcal{F}^*, \preceq_{\mathcal{F}})$ , which would be the pair  $(\mathcal{M}, \mathbf{c})$  for some coloring  $\mathbf{c} : \mathcal{M} \to \omega$ , by the uniqueness of the limit of  $(\mathcal{F}, \preceq_{\mathcal{F}})$ . Next, consider the closed subgroup  $\mathcal{H} \leq \operatorname{Aut}(\mathcal{M})$  of automorphisms of  $\mathcal{M}$  which permute the colors consistently, i.e.  $h \in \mathcal{H}$  iff there is some  $\sigma_h \in S_\infty$  such that  $\mathbf{c}(h \cdot a) =$ 

 $\sigma_h(\mathbf{c}(a))$  for every  $a \in M$ . There is a natural continuous homomorphism  $f : H \to S_\infty$  sending h to  $\sigma_h$ , noting that  $\sigma_h$  is unique as every color appears somewhere in M. The final step is to show that f is surjective, by fixing an arbitrary  $\sigma$  and constructing by a back-and-forth method (utilizing the homogeneity of the coloring  $\mathbf{c}$ ) an automorphism  $h \in H$  such that  $\sigma = \sigma_h$ .

The natural question: is the converse to Proposition 3.2 true? Evidently not, as one could consider the case where  $\mathcal{F}$  is the class of finite equivalence relations where each equivalence class has at most two elements, and  $\leq_{\mathcal{F}}$  is the usual substructure relation. This does not satisfy disjoint amalgamation for trivial reasons, however its automorphism group involves  $S_{\infty}$  as all of the pairs can be permuted arbitrarily.

One could consider the "fix" of only considering the class of substructures of  $\mathcal{M}$  which are definably-closed. This may require us in general to consider classes of infinite structures, and indeed there exists a Fraïssé theory of "finitely-generated" structures which would be useful if we were to consider substructures of  $\mathcal{M}$  which are the definable closures of finite sets. One would need to devise the appropriate notion of the disjoint amalgamation property in this context, but this seems to be a reasonable approach.

This does not, however, produce a necessary condition. One could just as well take the random graph and replace every vertex with some much more complicated structure which is not finitely-generated. The automorphism group of the resulting structure would still involve  $S_{\infty}$ , but would not satisfy any reasonable notion of disjoint amalgamation relative to definable closure. The realization, then, is that we need to work with some notion of closure which is weaker (coarser) than definable closure.

One reasonable candidate is the notion of pseudo-algebraic closure. For a countable atomic  $\mathcal{L}$ -structure  $\mathcal{M}$ , the pseudo-algebraic closure of a set  $A \subseteq M$  is defined to be the set of all  $b \in M$  such that  $b \in N$  for every  $\mathcal{N} \preceq_{\mathcal{L}_{\omega_1,\omega}} \mathcal{M}$  with  $A \subseteq N$ . We will see later that this is not quite the correct closure notion, but it is reasonably close to the correct one, and indeed our theory will significantly parallel the theory of the pseudo-algebraic closure as developed in [Dei77].

In this section we work backwards and reverse-engineer exactly what property we will want our notion of closure to exhibit in order to develop the "right" weakening of disjoint amalgamation. In Section 4, we will instead work from the bottom-up and define the right notion of closure that we want, which will be a very natural coarsening of pseudo-algebraic closure from the perspective of [Dei77]. Here, we use "coarsening" to mean that our closure operator will include possibly more elements in the closure of any set than pseudo-algebraic closure.

In an attempt to keep the theory as simple as possible, we discard the idea of working with classes of finitely-generated infinite structures and restrict ourselves only to classes of finite structures. This will fortunately not be an obstacle to developing the right weakening of disjoint amalgamation. Rather than considering the class of substructures which are finitely-generated with respect to some closure operator and develop a notion of disjoint amalgamation for such structures, we will work with the class of finite substructures and instead use the closure notion to control just how "disjoint" we require the amalgamation to be. 3.3. Closure operators and independence relations. We begin by considering definable closure and pseudo-algebraic closure in an abstract sense, as *closure operators*.

A closure operator on a set I is a function  $cl : \mathcal{P}(I) \to \mathcal{P}(I)$  satisfying for every  $A, B \subseteq I$ :

(1) 
$$A \subseteq \operatorname{cl}(A)$$
;

- (2)  $\operatorname{cl}(A) \subseteq \operatorname{cl}(B)$  whenever  $A \subseteq B$ ; and
- (3)  $\operatorname{cl}(\operatorname{cl}(A)) = \operatorname{cl}(A).$

Note that sometimes the axioms of closure operators demand that  $cl(\emptyset) = \emptyset$ , though we will not (and should not) demand that here. In the case that cl(A) = A, we say that A is **closed**. The closure operators considered in these notes will have **finite character**, meaning that cl(A) is the union of  $cl(A_0)$  where  $A_0$  ranges over finite subsets of A. We say that cl is **non-trivial** iff  $cl(\emptyset) \neq I$ . This will be reflected by the fact that we will only bother to define closure operators in terms of finite subsets. We will adopt the standard notation to sometimes write  $A \cup B$  as simply AB and  $A \cup \{b\}$  as simply Ab, for  $A, B \subseteq I$  and  $b \in I$ .

Given  $B \supseteq C$  finite subsets of I, say that  $b \in B \setminus cl(C)$  is **minimal in** B over C iff for every  $b' \in (B \cap cl(bC)) \setminus cl(C)$ , we have  $b \in cl(b'C)$ .

**Lemma 3.3.** Suppose cl is a closure operator on I and  $B \supseteq C$  are two finite subsets of I such that  $B \not\subseteq cl(C)$ . Then there exists some  $b \in B \setminus cl(C)$  which is minimal in B over C.

*Proof.* Define on  $B \setminus cl(C)$  a binary relation  $\leq$  by  $b \leq b'$  iff  $b \in cl(b'C)$ . This is easily seen to be a preorder. As  $B \setminus cl(C)$  is finite, we may choose a  $\leq$ -minimal element.

Given such a closure operator cl on a set I and finite subsets  $A, B, C \subseteq I$ , write  $A \perp_C B$ iff  $A \cap cl(BC) \subseteq cl(C)$  and  $B \cap cl(AC) \subseteq cl(C)$ . We call  $\downarrow$  the independence relation derived from cl.

**Lemma 3.4.** Suppose igcup is the independence relation derived from a closure operator cl on a set I. For any finite A, B, C with  $A \supseteq C$  and  $B \supseteq C$ , the following hold.

(1) (Symmetry) We have

$$A \underset{C}{\downarrow} B \quad iff \quad B \underset{C}{\downarrow} A$$

(2) (Monotonicity) Whenever  $A' \subseteq A$ ,  $B' \subseteq B$ , and  $C' \supseteq C$ , we have

$$A \underset{C}{\downarrow} B \quad implies \quad A' \underset{C'}{\downarrow} B'$$

(3) (Weak transitivity) If  $b \in B \setminus cl(C)$  is minimal in B over C, then

$$A \underset{C}{\downarrow} B$$
 iff both  $A \underset{C}{\downarrow} b$  and  $A \underset{bC}{\downarrow} B$ .

*Proof.* Let cl be the closure operator from which  $\perp$  is derived. Statements (1) and (2) are immediate. Now we prove (3). The forwards direction follows immediately from statement (2), so we focus on the backwards direction. First, fix an arbitrary  $a \in A \cap cl(B)$ . By

 $A \perp_{bC} B$  we conclude  $a \in cl(bC)$  and then by  $A \perp_C b$  we conclude  $a \in cl(C)$  as desired. Second, fix an arbitrary  $b' \in B \cap cl(A)$  and assume for contradiction  $b' \notin cl(C)$ . Then by  $A \perp_{bC} B$  we have that  $b' \in cl(bC)$ . By minimality of b, we have  $b \in cl(b'C)$ . Thus we have  $b \in cl(A)$  and so by  $A \perp_C B$ , we have  $b \in cl(C)$ . This in turn implies  $b' \in cl(C)$ , which is a contradiction.

3.4. **Invariant closure operators.** We will ultimately be defining closure operators on structures, and we will need them to have invariance properties with respect to automorphisms. In the meantime we will continue our discussion of closure operators in the abstract, but now taking into account the existence of a group action on the underlying set.

When there is some group action  $P \curvearrowright I$ , we say that a closure operator cl on I is P-**invariant** (or just **invariant** when there is no chance of confusion) iff for every  $A \subseteq M$  and  $\pi \in P$ , we have  $\pi[cl(A)] = cl(\pi[A])$ . Given sets  $A, B, C \subseteq I$ , we write  $A \cong_C^P B$  to denote that there is some  $\pi \in P$  such that  $\pi[A] = B$  and  $\pi(c) = c$  for every  $c \in C$ . In practice, we will drop the superscript.

The following property of invariant closure operators comes in handy and is easily verified.

**Lemma 3.5.** Suppose  $P \curvearrowright I$  and cl is an invariant closure operator on I. Then for any  $a, b \in M$  and  $C \subseteq M$  finite, if for every  $a' \cong_C a$  we have  $a' \in cl(bC)$ , then for every  $a' \cong_C a$  and  $b' \cong_C b$  we have  $a' \in cl(b'C)$ .

*Proof.* Let  $a' \cong_C a$  and  $b' \cong_C b$  and let  $\sigma \in P$  such that  $\sigma \upharpoonright C = \operatorname{id}_C$  and  $\sigma(b) = b'$ . By assumption  $\sigma^{-1}(a') \in \operatorname{cl}(bC)$  and by invariance of cl we have  $a' \in \operatorname{cl}(b'C)$ .

We will usually be interested in the case that we have an  $\mathcal{L}$ -structure  $\mathcal{M}$  where  $\mathcal{L}$  is a countable relational language, and cl is a closure operator on  $\mathcal{M}$  which is invariant with respect to the natural action  $\operatorname{Aut}(\mathcal{M}) \curvearrowright \mathcal{M}$ . Definable closure and pseudo-algebraic closure on a structure  $\mathcal{M}$  are invariant closure operators with respect to this action.

3.5. The disjointifying property. We now define a new property of invariant closure operators on sets I with group action  $P \curvearrowright I$ . Let  $\bigcup$  be the independence relation derived from an invariant closure operator cl on I. We say that cl is **disjointifying** iff whenever  $A, B \supseteq C$  are finite subsets of I, there exists a finite subset A' of I such that  $A' \cong_C A$  and  $A' \bigcup_C B$ .

In the following proposition, we prove three additional statements equivalent to an invariant closure operator cl being disjointifying. The equivalence of (2) and (3) below show that in the definition of the disjointifying property, it is enough to check  $A, B \supseteq C$  with  $|A \setminus C| = |B \setminus C| = 1$ . The equivalence of (4) is helpful for verifying the disjointifying property as it allows one to check two easier statements separately.

**Proposition 3.6.** Suppose cl is an invariant closure operator on  $P \curvearrowright I$  with derived independence relation  $\bigcup$ . The following are equivalent:

- (1) cl is disjointifying;
- (2) for any finite  $B \subseteq I$  and  $a \in I$ , there is some  $a' \cong_C a$  such that  $a' \downarrow_C B$ ; and
- (3) for any finite  $C \subseteq I$  and  $a, b \in I$ , there is some  $a' \cong_C a$  such that  $a' \downarrow_C b$ ;

- (4) for any finite  $C \subseteq I$  and  $a, b \in I$  with  $a \notin cl(C)$ , the following both hold:
  - (a) there is some  $a' \cong_C a$  such that  $a' \downarrow_C a$ ; and
  - (b) there is some  $a' \cong_C a$  such that  $a' \notin cl(bC)$ .

*Proof.* Clearly (1) implies (2), (2) implies (3), and (3) implies (4). However showing (4) implies (1) is not easy, and instead we show each step backwards. In order we show (3) implies (2), (2) implies (1), then (4) implies (3)

We prove (3) implies (2). We will induct on  $|B \setminus cl(C)|$ . The claim is immediately true when this is 0, so fix some  $b_0 \in B \setminus cl(C)$  minimal in B over C.

Fix an arbitrary  $a \in I$  and assume  $a \notin cl(C)$  as otherwise there is nothing to prove. Find  $a' \cong_C a$  such that  $a' \downarrow_C b_0$ . By the induction hypothesis, find some  $a'' \cong_{b_0C} a'$  such that  $a'' \downarrow_{b_0C} B$ . Clearly  $a'' \cong_C a$ . By invariance we have  $a'' \downarrow_C b_0$ , and then by Lemma 3.4.(3) we have  $a'' \downarrow_C B$  as desired.

The proof that (2) implies (1) is similar. We induct on  $|A \setminus cl(C)|$ . If this is 0, there is nothing to prove. Otherwise, let  $a_0 \in A \setminus cl(C)$  minimal in A over C. Find  $a'_0 \cong_C a_0$  such that  $a'_0 \, {\color{black} \_}_C B$ , as witnessed by some  $\pi \in P$ . Define  $A' := \pi[A]$  so that  $A' \cong_C A$  and  $a'_0 \in A'$ is minimal in A' over C. By the induction hypothesis, find  $A'' \cong_{a'_0C} A'$  such that  $A'' \, {\color{black} \_}_C B$ . Observe that  $a'_0$  is minimal in A'' over C as well. By Lemma 3.4.(3) we have  $A'' \, {\color{black} \_}_C B$  as desired.

Finally we see that (4) implies (3). We first note that (3) and (4) are equivalent to the following statements, respectively:

- (3\*) Given a, C suppose there exists some  $b \notin cl(C)$  such that for every  $b' \cong_C b$ , either  $a \in cl(b'C)$  or  $b' \in cl(aC)$ . Then  $a \in cl(C)$ .
- $(4^*)$  Given a, C suppose either of the following holds:
  - (a) there exists some b such that for every  $b' \cong_C b$ ,  $a \in cl(b'C)$ ; or
  - (b) for any  $a' \cong_C a$  either  $a \in cl(a'C)$  or  $a' \in cl(aC)$ .

Then  $a \in \operatorname{cl}(C)$ .

Assume (4\*). Suppose we're given a, C and some  $b \notin cl(C)$  such that for every  $b' \cong_C b$ , either  $a \in cl(b'C)$  or  $b' \in cl(aC)$ . Then in fact by automorphism-invariance we have that for every  $a' \cong_C a$  and  $b' \cong_C b$ , either  $a' \in cl(b'C)$  or  $b' \in cl(a'C)$ . Assume for contradiction that  $a \notin cl(C)$ .

Define a relation  $\leq$  on the set of all a' with  $a' \cong_C a$  by

 $a' \leq a''$  iff  $\forall b' \cong_C b$ ,  $(a'' \in \operatorname{cl}(b'C) \to a' \in \operatorname{cl}(b'C))$ .

We will argue that this is in fact a linear preorder. Granting this for now, we show that  $a' \leq a''$  implies that  $a' \in cl(a''C)$ . Since either  $a \leq a'$  or  $a' \leq a$  for every  $a' \cong_C a$ , we would be done because (4\*).(b) would hold and we would have  $a \in cl(C)$ , a contradiction.

Assume  $a' \leq a''$ . We may find some  $b' \cong_C b$  such that  $a'' \in \operatorname{cl}(b'C)$  (otherwise we would have  $b' \in \operatorname{cl}(a''C)$  for every  $b' \cong_C b$ , and thus  $b' \in \operatorname{cl}(a'''C)$  for every  $a''' \cong_C a$  by Lemma 3.5, allowing us to conclude  $b \in \operatorname{cl}(C)$  by (4\*).(a), a contradiction). Then of course we have  $a'' \in \operatorname{cl}(b''C)$  for every  $b'' \cong_{a''C} b'$  and thus  $a' \in \operatorname{cl}(b''C)$  for every  $b'' \cong_{a''C} b'$ . By (4\*).(a), we can conclude  $a' \in \operatorname{cl}(a''C)$  as desired.

All that remains to show that  $\leq$  is a linear preorder. Reflexivity and transitivity of  $\leq$  are clear (and not immediately useful). To see linearity, suppose for contradiction there exist some  $a_0, a_1 \cong_C a$  and some  $b_0, b_1 \cong_C b$  with

$$a_1 \in \operatorname{cl}(b_0 C)$$
 and  $a_0 \notin \operatorname{cl}(b_0 C)$ 

and

$$a_1 \notin \operatorname{cl}(b_1C)$$
 and  $a_0 \in \operatorname{cl}(b_1C)$ .

By the second statement, we have  $b_1 \in cl(a_1C)$  and thus  $a_0 \in cl(a_1C)$ . But that implies  $a_0 \in cl(b_0C)$  by the first clause of the first statement. This contradicts the second clause of the first statement.

If cl is an invariant disjointifying closure operator, then cl is non-trivial iff  $I \neq cl(A)$  for any finite  $A \subseteq I$ . Also, we will implicitly assume from now on that any disjointifying closure operator is invariant.

3.6. Disjointifying closure operators and involving  $S_{\infty}$ . We now finish this section by proving the following theorem:

**Theorem 3.7.** Let I be a countably-infinite set and  $P \leq S_I$  closed with the natural action  $P \curvearrowright I$ . If I has a nontrivial invariant disjointifying closure operator cl, then P involves  $S_{\infty}$ .

Let  $\mathcal{L}$  be the countable relational language with an *n*-ary relation  $R_{\bar{a}}$  for each  $n \in \omega$ and  $\bar{a} \in I^n$ . We define an  $\mathcal{L}$ -structure  $\mathcal{M}$  living on I where for every  $\bar{a}, \bar{b} \in I^{<\omega}$ , we have that  $R_{\bar{a}}(\bar{b})^{\mathcal{M}}$  holds iff there exists some  $g \in P$  with  $g \cdot \bar{a} = \bar{b}$ . It's easy to check that  $\mathcal{M}$ is ultrahomogeneous, thus if we let  $\mathcal{F}$  be the age of  $\mathcal{M}$ , we have that  $(\mathcal{F}, \leq_{\mathcal{L}})$  is a Fraïssé class. For each  $\mathcal{A} \in \mathcal{F}$ , we have the invariant disjointifying closure operator  $cl_{\mathcal{A}}$  on  $\mathcal{A}$  that  $\mathcal{A}$  inherits as a substructure of  $\mathcal{M}$ .

Let  $\mathcal{F}^*$  be the class of pairs  $(\mathcal{A}, \mathbf{c}_{\mathcal{A}})$  where  $\mathcal{A} \in \mathcal{F}$  and  $\mathbf{c}_{\mathcal{A}} : \mathcal{A} \to \omega \cup \{\text{null}\}$  satisfying that  $\mathbf{c}_{\mathcal{A}}(a) = \text{null}$  for every  $a \in \text{cl}_{\mathcal{A}}(\emptyset)$ . We say that  $(\mathcal{A}, \mathbf{c}_{\mathcal{A}})$  is a strong substructure of  $(\mathcal{B}, \mathbf{c}_{\mathcal{B}})$ , denoted  $(\mathcal{A}, \mathbf{c}_{\mathcal{A}}) \preceq^* (\mathcal{B}, \mathbf{c}_{\mathcal{B}})$ , iff

- (a)  $\mathcal{A} \leq_{\mathcal{L}} \mathcal{B};$
- (b)  $\mathbf{c}_{\mathcal{A}}(a) = \mathbf{c}_{\mathcal{B}}(a)$  for every  $a \in A$ ; and
- (c)  $\mathbf{c}_{\mathcal{B}}(b) = \text{null for every } b \in (B \cap \text{cl}_{\mathcal{B}}(A)) \setminus A.$

We remark that  $\preceq^*$  is a notion of strong substructure, and that  $\mathcal{F}^*$  consists of the pairs  $(\mathcal{A}, \mathbf{c}_{\mathcal{A}})$  for which the empty structure  $(\emptyset, \emptyset)$  is a strong substructure.

We wish to show that  $(\mathcal{F}^*, \preceq^*)$  satisfies the assumptions of Proposition 3.1. The only tricky part is showing that it satisfies the amalgamation property. Suppose  $(\mathcal{A}, \mathbf{c}_{\mathcal{A}}), (\mathcal{B}, \mathbf{c}_{\mathcal{B}}), (\mathcal{C}, \mathbf{c}_{\mathcal{C}}) \in$  $\mathcal{F}^*$  satisfy  $(\mathcal{A}, \mathbf{c}_{\mathcal{A}}) \preceq^* (\mathcal{B}, \mathbf{c}_{\mathcal{B}})$  and  $(\mathcal{A}, \mathbf{c}_{\mathcal{A}}) \preceq^* (\mathcal{C}, \mathbf{c}_{\mathcal{C}})$ . Because cl is disjointifying, after replacing  $(\mathcal{C}, \mathbf{c}_{\mathcal{C}})$  with some  $(\mathcal{C}', \mathbf{c}_{\mathcal{C}'}) \cong_{\mathcal{A}} (\mathcal{C}, \mathbf{c}_{\mathcal{C}})$ , we may find  $\mathcal{D} \in \mathcal{F}$  such that  $\mathcal{B} \leq_{\mathcal{L}} \mathcal{D}$  and  $\mathcal{C} \leq_{\mathcal{L}} \mathcal{D}$  and  $\mathrm{cl}_{\mathcal{D}}(B) \cap C \subseteq \mathrm{cl}_{\mathcal{D}}(A)$  and  $\mathrm{cl}_{\mathcal{D}}(C) \cap B \subseteq \mathrm{cl}_{\mathcal{D}}(A)$ . By shrinking  $\mathcal{D}$ , we can assume that  $D = B \cap C$ . Our task now is to find a coloring  $\mathbf{c}_{\mathcal{D}}$  of  $\mathcal{D}$  such that  $(\mathcal{B}, \mathbf{c}_{\mathcal{B}}) \preceq^* (\mathcal{D}, \mathbf{c}_{\mathcal{D}})$ and  $(\mathcal{C}, \mathbf{c}_{\mathcal{C}}) \preceq^* (\mathcal{D}, \mathbf{c}_{\mathcal{D}})$ . This would also imply that  $(\mathcal{D}, \mathbf{c}_{\mathcal{D}}) \in \mathcal{F}^*$ .

The choice of coloring is natural. Define

$$\mathbf{c}_{\mathcal{D}}(d) = \begin{cases} \mathbf{c}_{\mathcal{B}}(b) & b \in B \\ \mathbf{c}_{\mathcal{C}}(c) & c \in C. \end{cases}$$

To see that it is well-defined, observe that if  $d \in B \cap C$ , then  $d \in \operatorname{cl}_{\mathcal{D}}(A)$ . If  $d \in A$ , then of course we have  $\mathbf{c}_{\mathcal{B}}(d) = \mathbf{c}_{\mathcal{C}}(d)$ , otherwise, if  $d \in \operatorname{cl}(A) \setminus A$ , then we also have  $\mathbf{c}_{\mathcal{B}}(d) = \operatorname{null} = \mathbf{c}_{\mathcal{C}}(d)$ . Next, suppose  $d \in \operatorname{cl}_{\mathcal{D}}(B) \setminus B$ , for which we need to check  $\mathbf{c}_{\mathcal{D}}(d) = \operatorname{null}$ . By our choice of  $\mathcal{D}$ , since  $d \notin B$  we must have  $d \in C$ , but since  $d \in \operatorname{cl}_{\mathcal{D}}(B) \cap C$ , we must have  $d \in \operatorname{cl}(A)$ . Clearly  $d \notin A$  (because  $d \notin B$ ), thus  $\mathbf{c}_{\mathcal{D}}(d) = \mathbf{c}_{\mathcal{C}}(d) = \operatorname{null}$ . For any  $d \in \operatorname{cl}_{\mathcal{D}}(C) \setminus C$ , we can confirm that  $\mathbf{c}_{\mathcal{D}}(d) = \operatorname{null}$  by the same argument.

Now we can compute (using Proposition 3.1) the limit of  $(\mathcal{F}^*, \preceq_{\mathcal{F}}^*)$ , which of course will be isomorphic to  $(\mathcal{M}, \mathbf{c})$  for some coloring  $\mathbf{c}$ . Because cl is nontrivial, we have that every color appears in  $(\mathcal{M}, \mathbf{c})$ . Finally, we let  $H \leq P$  be the subgroup of all g such that for some  $\sigma \in S_{\infty}$ , we have that for every  $a \in I$ ,  $\mathbf{c}(g \cdot a) = \sigma(\mathbf{c}(a))$ . To see that H is closed, observe  $h \in H$  iff for every  $a, a' \in I$ , if  $\mathbf{c}(a) = \mathbf{c}(a')$  then  $\mathbf{c}(h \cdot a) = \mathbf{c}(h \cdot a')$ , which is a closed condition. We consider the natural continuous homomorphism  $f : H \to S_{\infty}$  which sends each  $h \in H$  to the unique  $\sigma \in S_{\infty}$  such that  $\mathbf{c}(h \cdot a) = \sigma(\mathbf{c}(a))$  for every  $a \in I$ . Our final task is to show that f is surjective.

Fix some arbitrary permutation  $\sigma \in S_{\infty}$ . Let  $\{c_n \mid n \in \omega\}$  be an enumeration of I. We will define finite substructures  $\mathcal{A}_n \leq \mathcal{M}$  and  $\mathcal{B}_n \leq \mathcal{M}$  and  $g_n \in P$  satisfying:

- (1)  $A_n \subseteq A_{n+1}$  and  $B_n \subseteq B_{n+1}$  for every n;
- (2)  $(\mathcal{A}_n, \mathbf{c} \upharpoonright \mathcal{A}_n) \preceq^* (\mathcal{A}_{n+1}, \mathbf{c} \upharpoonright \mathcal{A}_{n+1})$  and  $(\mathcal{B}_n, \mathbf{c} \upharpoonright \mathcal{B}_n) \preceq^* (\mathcal{B}_{n+1}, \mathbf{c} \upharpoonright \mathcal{B}_{n+1})$  for every n;
- (3)  $c_n \in A_{2n+1}$  and  $c_n \in B_{2n+2}$  for every n;
- (4)  $g_n \cdot A_n = B_n$  for every n;
- (5)  $\mathbf{c}(g_n \cdot a) = \sigma(\mathbf{c}(a))$  for every n and  $a \in A_n$ ;
- (6)  $g_{n+1} \cdot c = g_n \cdot c$  for every n and  $c \in A_n$ .

Having done this, we can easily check that  $g_n \to g_\infty$  for some  $g_\infty \in P$  with  $g_\infty \in H$  and  $f(g_\infty) = \sigma$ . We proceed to the construction. Let  $A_0 = B_0 = \emptyset$ . Having defined  $\mathcal{A}_{2n}$ , by Proposition 3.1.(1) we choose  $\mathcal{A}_{2n+1}$  to be any substructure of  $\mathcal{M}$  satisfying  $c_n \in \mathcal{A}_{2n+1}$  and

$$(\mathcal{A}_{2n}, \mathbf{c} \upharpoonright A_{2n}) \preceq^* (\mathcal{A}_{2n+1}, \mathbf{c} \upharpoonright A_{2n+1}).$$

Now let  $(\hat{\mathcal{B}}, \hat{\mathbf{c}})$  be defined by

$$\mathcal{B} = \mathcal{M} \upharpoonright (g_{2n} \cdot A_{2n+1}) \text{ and } \hat{\mathbf{c}}(b) = (\sigma \circ \mathbf{c})(g_{2n}^{-1} \cdot b).$$

Observe that

$$(\mathcal{B}_{2n},\mathbf{c}\upharpoonright B_{2n})\preceq^* (\mathcal{B},\hat{\mathbf{c}})$$

By Proposition 3.1.(2), there is some  $\mathcal{B}_{2n+1} \leq_{\mathcal{L}} M$  satisfying that

$$(\mathcal{B}_{2n}, \mathbf{c} \upharpoonright B_{2n}) \preceq^* (\mathcal{B}_{2n+1}, \mathbf{c} \upharpoonright B_{2n+1})$$

and

$$(\mathcal{B}_{2n+1},\mathbf{c} \upharpoonright B_{2n+1}) \cong_{B_{2n}} (\hat{\mathcal{B}},\mathbf{c}^*).$$

By Proposition 3.1.(3) there is some  $h \in \operatorname{Stab}_P(B_{2n})$  with  $B_{2n+1} = h \cdot \hat{B}$  and  $\mathbf{c}(b) = \hat{\mathbf{c}}(h^{-1} \cdot b)$ for every  $b \in B_{2n+1}$ . Let  $g_{2n+1} = hg_{2n}$ .

The even case is symmetric, and thus we have showed that P involves  $S_{\infty}$ .

#### 4. A RANK FUNCTION AND THE MINIMAL DISJOINTIFYING CLOSURE OPERATOR

We now identify a rank function which identifies the existence of a disjointifying closure operator. It will also allow us to define a canonical closure operator which will happen to be the minimal disjointifying closure operator if one exists. The rank that we will define will be a natual extension of a rank function which is already in the literature. We will start by discussing this rank function and how it characterizes the pseudo-algebraic closure, and then proceed to defining our new rank function and the analogous way in which it characterizes the minimal disjointifying closure operator.

4.1. **Deissler rank.** This subsection is presentation of the theory developed by Deissler in [Dei77] in somewhat different language.

Let I be a set with an action  $P \curvearrowright I$ . Given a finite set  $B \subseteq I$  and  $a \in I$ , we define an ordinal rank Drk(a, B) as follows. We say  $Drk(a, B) \leq 0$  iff for every  $g \in Stab_P(B)$ ,  $g \cdot a = a$ . In general, for  $\alpha > 0$ , we say  $Drk(a, B) \leq \alpha$  iff there exists some  $c \in I$  such that for every  $c' \cong_B c$ ,  $Drk(a, Bc') < \alpha$ . We then define Drk(a, B) to be the least  $\alpha$  such that  $Drk(a, B) \leq \alpha$ , if such  $\alpha$  exists, otherwise we write  $Drk(a, B) = \infty$ . If  $Drk(a, B) \leq \alpha$  for some ordinal  $\alpha$ , then we write  $Drk(a, B) < \infty$ .

The following is easy to check.

**Lemma 4.1.** Suppose for some finite  $B \subseteq I$  and  $a, c \in I$ , we have  $\text{Drk}(a, c'B) < \infty$  for every  $c' \cong_B c$ . Then  $\text{Drk}(a, B) < \infty$ .

We define an operator Dcl on I as follows: given  $B \subseteq I$  and  $a \in I$ , we declare  $a \in Dcl(B)$ iff  $Drk(a, B_0) < \infty$  for some finite  $B_0 \subseteq B$ .

**Theorem 4.2** (ess. Deissler). Let  $\mathcal{L}$  be a countable relational language and  $\mathcal{M}$  a countable  $\mathcal{L}$ -structure with the natural action  $Aut(\mathcal{M}) \curvearrowright \mathcal{M}$ . Let  $B \subseteq M$  be finite and  $a \in M$ . Then  $a \in Dcl(B)$  iff for every  $\mathcal{M}_0 \prec_{\mathcal{L}_{\omega_1,\omega}} \mathcal{M}$  with  $B \subseteq M_0$ , we have  $a \in M_0$ .

*Proof.* Recall that for any set  $M_0 \subseteq M$ , we have  $\mathcal{M}_0 \prec_{\mathcal{L}_{\omega_1,\omega}} \mathcal{M}$  iff for every finite  $B \subseteq M_0$  and  $a \in M$ , there is some  $a' \in M_0$  with  $a' \cong_B a$ .

For the forward direction, suppose  $a \in \text{Dcl}(B)$  and  $M_0 \prec_{\mathcal{L}_{\omega_1,\omega}} M$  with  $B \subseteq M_0$ . We argue by induction on Drk(a, B). If  $\text{Drk}(a, B) \leq 0$  and  $a' \cong_B a$  satisfies  $a' \in M_0$ , then we must have a' = a and thus  $a \in M_0$ . Otherwise, suppose  $\text{Drk}(a, B) \leq \alpha$  with  $\alpha > 0$  and the claim is true below  $\alpha$ . Fix some  $c \in M$  such that  $\text{Drk}(a, Bc') < \alpha$  for every  $c' \cong_B c$ . Let  $c' \cong_B c$  such that  $c' \in M_0$  in which case by the induction hypothesis we have  $a \in M_0$ . For the backwards direction, suppose  $a \notin \mathrm{Dcl}(B)$ . We will construct a  $\mathcal{L}_{\omega_1,\omega}$ -elementary submodel  $\mathcal{M}_0$  of  $\mathcal{M}$  with  $a \notin M_0$ .

Given  $C \subseteq C' \subseteq M$ , say that C' has closure property (\*) over C iff for every finite  $D \subseteq C$  and  $e \in M$ , there exists some  $e' \cong_D e$  with  $e' \in C'$ . We argue that for any  $C \subseteq M$  with  $a \notin \mathrm{Dcl}(C)$ , there is a set  $C' \subseteq M$  with C' which has closure property (\*) over C and  $a \notin \mathrm{Dcl}(C')$ . Having done this, we can define a sequence

$$\emptyset = C_0 \subseteq C_1 \subseteq C_2 \subseteq \dots$$

such that  $a \notin \operatorname{Dcl}(C_n)$  and  $C_{n+1}$  has property (\*) over  $C_n$ , and define  $M_0 = \bigcup_{n \in \omega} C_n$ .

Let  $C \subseteq M$ . Write C as an increasing union  $\bigcup_{n \in \omega} B_n$  of finite sets. Fix an enumeration  $\{a_n \mid n \in \omega\}$  of I. Let  $h : \omega \to \omega$  be a function such that the preimage of every  $n \in \omega$  is infinite. We recursively define using Lemma 4.1 a sequence  $d_0, d_1, \ldots$  such that for every  $n \in \omega, d_n \cong_{B_n} a_{h(n)}$  and  $a \notin \text{Dcl}(B_n d_0 \ldots d_n)$ . Then let  $C' = C \cup \{d_n \mid n \in \omega\}$ . It is easy to check that this works.

**Proposition 4.3.** The operator Dcl is a locally-finite invariant closure operator, and moreover, for any finite  $B \subseteq I$  and  $a, c \in I$ , if  $a \in Dcl(Bc')$  for every  $c' \cong_B c$ , then  $a \in Dcl(B)$ .

Proof. Local-finiteness and invariance are immediate. The fact that Dcl is a closure operator follows from Theorem 4.2. The additional property (or more accurately, its contrapositive) is also easily proved from Theorem 4.2. Suppose  $a \notin Dcl(B)$ . Then there is some  $\mathcal{M}_0 \prec_{\mathcal{L}_{\omega_1,\omega}} \mathcal{M}$ with  $B \subseteq M_0$  and  $a \notin M_0$ . Then for any c, there must be some  $c' \cong_B c$  with  $c' \in M_0$ . Since  $a \notin M_0$ , we must have  $a \notin Dcl(Bc')$ .

In [Gao98], Gao relates the non-existence of nontrivial  $\mathcal{L}_{\omega_1,\omega}$ -elementary substructures with dynamical properties of the automorphism group. Recall that a Polish group G is cli iff there is a complete metric d on G, compatible with the topology on G, such that d is left-invariant, i.e. d(g, g') = d(hg, hg') for every  $g, g', h \in G$ .

**Theorem 4.4** (Gao, [Gao98]). Let  $\mathcal{L}$  be a countable relational language and  $\mathcal{M}$  a countable  $\mathcal{L}$ -structure. Then the Polish group  $Aut(\mathcal{M})$  is cli iff there is no  $M_0 \subsetneq \mathcal{M}$  such that  $\mathcal{M}_0 \prec_{\mathcal{L}_{\omega_1,\omega}} \mathcal{M}$ .

Putting these equivalences together, we get the following "master" list of equivalences.

**Corollary 4.5** (Deissler, Gao). Let  $\mathcal{L}$  be a countable relational language and  $\mathcal{M}$  a countable  $\mathcal{L}$ -structure. The following are equivalent:

- (1)  $Aut(\mathcal{M})$  is cli;
- (2) there is no  $M_0 \subsetneq M$  such that  $\mathcal{M}_0 \prec_{\mathcal{L}_{\omega_1,\omega}} \mathcal{M}$ ;
- (3) there is no uncountable  $\mathcal{L}$ -structure satisfying the Scott sentence of  $\mathcal{M}$ ;
- (4)  $\operatorname{Dcl}(\emptyset) = M;$
- (5) for every  $a \in M$ ,  $Drk(a, \emptyset) < \omega_1$ .

4.2. **Disjointifying rank.** With the Deissler rank as motivation, we move on to the disjointifying rank.

Let I be a set with a group action  $P \curvearrowright I$ . Given a finite set  $B \subseteq I$  and  $a \in I$ , we define a rank an ordinal rank  $\operatorname{Krk}(a, B)$  as follows.

We say  $\operatorname{Krk}(a, B) \leq 0$  iff iff for every  $\pi \in P$  with  $\pi \upharpoonright B = \operatorname{id}_B$ ,  $\pi(a) = a$ . In general for  $\alpha > 0$ , we say  $\operatorname{Krk}(a, B) \leq \alpha$  iff at least one of the following holds:

(1) there exists some c such that for every  $c' \cong_B c$ ,  $\operatorname{Krk}(a, Bc') < \alpha$ . or

(2) for every  $a' \cong_B a$ , either  $\operatorname{Krk}(a, Ba') < \alpha$  or  $\operatorname{Krk}(a', Ba) < \alpha$ .

We say  $\operatorname{Krk}(a, B) = \infty$  iff  $\operatorname{Krk}(a, B)$  is not less than  $\alpha$  for any ordinal  $\alpha$ . We write  $\operatorname{Krk}(a, B) < \infty$  to mean that  $\operatorname{Krk}(a, B) \leq \alpha$  for some ordinal  $\alpha$ . Ultimately, we define  $\operatorname{Krk}(a, B)$  to be the least ordinal  $\alpha$  such that  $\operatorname{Krk}(a, B) \leq \alpha$ , or  $\infty$  if no such  $\alpha$  exists.

Note that if we were to remove condition (2) in the recursive case of the definition of disjointifying rank, we will just get the Deissler rank.

We list some basic properties of the rank which are easily confirmed.

Lemma 4.6. The following all hold:

- (1) For finite subsets B, C of I satisfying  $C \supseteq B$ , and  $a \in I$ , we have  $Krk(a, C) \leq Krk(a, B)$ ;
- (2) For finite  $B \subseteq I$  and  $a \in I$ , if there exists some  $c \in I$  such that  $Krk(a, c'B) < \infty$  for every  $c' \cong_B c$ , then  $Krk(a, B) < \infty$ ;
- (3) For finite  $B \subseteq I$  and  $a \in I$ , if either  $Krk(a, a'B) < \infty$  or  $Krk(a', aB) < \infty$  for every  $a' \cong_B a$ , then  $Krk(a, B) < \infty$ .

We define a closure operator  $\operatorname{cl}^{\min}$  on M by saying for  $a \in M$  and finite  $B \subseteq M$  that  $a \in \operatorname{cl}^{\min}(B)$  iff  $\operatorname{Krk}(a, B) < \infty$ .

Lemma 4.7. cl<sup>min</sup> is a closure operator.

*Proof.* We show by induction on  $\alpha$  that for any finite  $C \subseteq I$  and  $a, b \in I$ , if  $\operatorname{Krk}(a, C) \leq \alpha$  and  $\operatorname{Krk}(b, aC) < \infty$ , then  $\operatorname{Krk}(b, C) < \infty$ .

For the base case  $\alpha = 0$ , we now induct on  $\beta := \operatorname{Krk}(b, aC)$ . For  $\beta = 0$ , the claim is immediate, so let  $\beta > 0$  and assume the claim is true below  $\beta$ . For the first case, we assume there is some  $d \in I$  such that for every  $d' \cong_{aC} d$ , we have  $\operatorname{Krk}(a, d'aC) < \beta$ . Whenever  $d' \in I$  satisfies  $d' \cong_C d$ , we have that  $d' \cong_{aC} d$  by the fact that  $\operatorname{Krk}(a, C) \leq 0$ . Therefore we have  $\operatorname{Krk}(a, d'C) < \beta$  for every  $d' \cong_C d$ , and thus  $\operatorname{Krk}(a, C) \leq \beta$ . In the second case, we have for every  $b' \cong_{aC} b$ , either  $\operatorname{Krk}(b', baC) < \beta$  or  $\operatorname{Krk}(b, b'aC) < \beta$ . By the same argument as in the first case,  $\operatorname{Krk}(b, C) \leq \beta$  follows.

Now suppose  $\alpha > 0$  and the claim is true below  $\alpha$ . Suppose  $\operatorname{Krk}(a, C) \leq \alpha$  and  $\operatorname{Krk}(b, aC) < \infty$ .

For the first case, we assume there is some  $d \in I$  such that for every  $d' \cong_C d$ , we have  $\operatorname{Krk}(a, d'C) < \alpha$ . We have by the induction hypothesis and Lemma 4.6.(1) that for every  $d' \cong_C d$ ,  $\operatorname{Krk}(b, d'C) < \infty$ . Then by Lemma 4.6.(2), we have  $\operatorname{Krk}(b, C) < \infty$ .

For the second case, we assume that for every  $a' \cong_C a$ , either  $\operatorname{Krk}(a', aC) < \alpha$  or  $\operatorname{Krk}(a, a'C) < \alpha$ . For  $b' \cong_C b$  and  $b'' \cong_C b$ , write  $b' \leq b''$  iff for every  $a' \cong_C a$ , if  $\operatorname{Krk}(b', a'C) < \infty$  then  $\operatorname{Krk}(b'', a'C) < \infty$ . Easily,  $\leq$  is reflexive and transitive on  $\{b' \in I \mid b' \cong_C b\}$ . We argue it is also linear, similar to the proof of Proposition 3.6. Suppose we have such b', b'' such that  $b' \leq b''$  and  $b'' \leq b'$ . Then there are  $a', a'' \cong_C a$  such that

(1) 
$$\operatorname{Krk}(b', a'C) < \infty$$
 and  $\operatorname{Krk}(b'', a'C) = \infty$ 

(2) 
$$\operatorname{Krk}(b', a''C) = \infty \quad \text{and} \quad \operatorname{Krk}(b'', a''C) < \infty.$$

We have either  $\operatorname{Krk}(a', a''C) < \alpha$  or  $\operatorname{Krk}(a'', a'C) < \alpha$ . In the first case, we have  $\operatorname{Krk}(b', a''C) < \infty$  by the induction hypothesis, which contradicts Equation 2. In the second case, we have  $\operatorname{Krk}(b'', a'C) < \infty$  by the induction hypothesis, which contradicts Equation 1. Thus  $\leq$  is a prelinear order.

Now we argue that if  $b' \leq b''$ , then  $\operatorname{Krk}(b'', b'C) < \infty$ . To see this, fix some  $a' \cong_C a$  such that  $\operatorname{Krk}(b', a'C) < \infty$ , which must exist because of  $\operatorname{Krk}(b, aC) < \infty$  and invariance. Then for any  $a'' \cong_{Cb'} a'$ , we have  $\operatorname{Krk}(b', a''C) < \infty$ . Because  $b' \leq b''$ , this means that for any  $a'' \cong_{Cb'} a$ , we have  $\operatorname{Krk}(b'', a''b'C) < \infty$ . By Lemma 4.6.(2), we have  $\operatorname{Krk}(b'', b'C) < \infty$ .

Finally, by Lemma 4.6.(3) we have just proved that  $\operatorname{Krk}(b, C) < \infty$ .

# **Proposition 4.8.** cl<sup>min</sup> is the minimum disjointifying closure operator.

*Proof.* We first check that  $cl^{\min}$  is disjointifying, using equivalence (4) from Proposition 3.6. Let  $C \subseteq I$  be finite and  $a, b \in I$  with  $a \notin cl^{\min}(C)$ . First we check clause (a) from Proposition 3.6.4. Suppose for contradiction that for every  $a' \cong_C a, a' \not\perp_C a$  where  $\bigcup$  is the independence relation derived from  $cl^{\min}$ . This means for every  $a' \cong_C a, a \in cl^{\min}(a'C)$  or  $a' \in cl^{\min}(aC)$ . By definition of  $cl^{\min}$  and Lemma 4.6.3, we have  $a \in cl^{\min}(C)$ , a contradiction. By a similar argument using Lemma 4.6.2, we see that clause (b) holds as well.

Now we see that  $cl^{\min}$  is the minimum. Let cl be some other disjointifying closure operator. We show by induction on  $\alpha$  that for any finite  $C \subseteq I$  and  $a \in I$ , if  $Krk(a, C) \leq \alpha$  then  $a \in cl(B)$ .

If  $\alpha = 0$  and for contradiction we assume  $a \notin cl(B)$ , then we apply clause (4a) from Proposition 3.6 to find some  $a' \cong_C a$  such that  $a' \notin cl(aC)$ . However, as Krk(a, C) = 0, it must be the case that a' = a, a contradiction.

Now let  $\alpha > 0$  and assume the claim is true below  $\alpha$ . Suppose  $\operatorname{Krk}(a, C) \leq \alpha$ . There are two cases in the definition of Krk, so we first consider the first one. Suppose there is some  $b \in I$  such that for every  $b' \cong_C b$ ,  $\operatorname{Krk}(a, b'C) < \alpha$ . In particular, we have  $a \in \operatorname{cl}(b'C)$  for every  $b' \cong_C b$ . This directly contradicts clause (4b) of Proposition 3.6. The second case of the definition is handled in the same way using clause (4a) of Proposition 3.6.  $\Box$ 

#### 5. INDISCERNIBLE SUPPORT FUNCTIONS

We define another technical notion which we will see is equivalent to an automorphism group involving  $S_{\infty}$ , with two goals in mind: to provide a motivating example of where a nontrivial disjointifying closure operator arises, and as a technical tool which we will make us of later.

As usual let I be a set with an action  $P \curvearrowright I$ . We write  $[I]^{<\omega}$  to represent the set of finite subsets of I. A function supp :  $[I]^{<\omega} \to [\omega]^{<\omega}$  is a **support function** iff

(1) for every finite  $A, B \subseteq I$  with  $A \subseteq B$ , we have  $\operatorname{supp}(A) \subseteq \operatorname{supp}(B)$ .

We say supp is **notrivial** iff furthermore

(2)  $\operatorname{supp}(A) \neq \emptyset$  for some finite  $A \subseteq I$ ;

and finally we say supp is **indiscernible** iff

(3) for every finite  $A, B \subseteq I$  with  $A \subseteq B$ , and for every finite  $u, v \subseteq \omega$  with  $\operatorname{supp}(A) = u$ and  $\operatorname{supp}(B) = v$ , and for every  $v' \cong_u v$ , there exists some  $B' \cong_A B$  with  $\operatorname{supp}(B') = v'$ .

We write  $v' \cong_u v$  to indicate  $|v' \setminus u| = |v \setminus u|$ , allowing for the possibility of putting some extra structure on the space of supports.

Note that we do not make any demands that supp is invariant. One could view this as meaning that a support function captures local information. From the existence of such a function supp, we will derive the existence of a nontrivial disjointifying closure operator. An invariant closure operator, on the other hand, is a global object, as in describes relationships between sets which is invariant under automorphisms.

We assume that supp is such a function, and our objective is to show that  $cl^{\min}$  is nontrivial.

**Lemma 5.1.** Suppose  $a \in cl^{min}(B)$ . Then supp(aB) = supp(B).

*Proof.* We prove by induction on  $\alpha$  that if  $\operatorname{Krk}(a, B) \leq \alpha$  then  $\operatorname{supp}(aB) = \operatorname{supp}(B)$ .

Consider the case where  $\alpha = 0$ . By indiscernibility of supp, there is some  $a' \cong_B a$  such that  $\operatorname{supp}(a'B) \cap \operatorname{supp}(aB) = \operatorname{supp}(B)$ . By the definition of  $\operatorname{Krk}(a, B) \leq 0$ , we know that a' = a, which means  $\operatorname{supp}(a'B) = \operatorname{supp}(aB) = \operatorname{supp}(B)$  as desired.

Otherwise let  $\alpha > 0$  and suppose the claim is true below  $\alpha$ .

In the first case of the definition of  $\operatorname{Krk}(a, B) \leq \alpha$ , there is some  $c \in I$  such that  $\operatorname{Krk}(a, c'B) < \alpha$  for every  $c' \cong_B c$ . Fix some  $c' \cong_B c$  such that  $\operatorname{supp}(c'B) \cap \operatorname{supp}(aB) = \operatorname{supp}(B)$ . By the induction hypothesis we have  $\operatorname{supp}(ac'B) = \operatorname{supp}(c'B)$  and thus  $\operatorname{supp}(aB) \subseteq \operatorname{supp}(c'B)$ . Again we have  $\operatorname{supp}(aB) = \operatorname{supp}(B)$ .

The second case of the definition of  $\operatorname{Krk}(a, B) \leq \alpha$  is handled a similar way. Suppose for every  $a' \cong_B a$ , either  $\operatorname{Krk}(a, a'B) < \alpha$  or  $\operatorname{Krk}(a', aB) < \alpha$ . Fix some  $a' \cong_B a$  such that  $\operatorname{supp}(a'B) \cap \operatorname{supp}(aB) = \operatorname{supp}(B)$  and  $|\operatorname{supp}(a'B)| = |\operatorname{supp}(aB)|$ . In the case that  $\operatorname{Krk}(a, a'B) < \alpha$  we have  $\operatorname{supp}(aB) \subseteq \operatorname{supp}(aa'B) = \operatorname{supp}(a'B)$  and thus we can conclude  $\operatorname{supp}(aB) = \operatorname{supp}(B)$ . Finally, if  $\operatorname{Krk}(a', aB) < \alpha$  we have  $\operatorname{supp}(a'B) \subseteq \operatorname{supp}(a'aB) =$  $\operatorname{supp}(aB)$ . Oow we realize we must have  $|\operatorname{supp}(aB)| = |\operatorname{supp}(a'B)| = |\operatorname{supp}(B)|$ , and thus in particular,  $\operatorname{supp}(aB) = \operatorname{supp}(B)$ .  $\Box$ 

Although it is easy to check, we remark that if supp is nontrivial, then there must be some finite B and a such that  $\operatorname{supp}(B) = \emptyset$  and  $\operatorname{supp}(aB) \supseteq \operatorname{supp}(B)$ . This can easily be done by letting C be the smallest set with nonempty support, and choosing a to be some arbitrary element of C and defining  $B = C \setminus \{a\}$ . Thus we can conclude:

**Proposition 5.2.** If  $P \curvearrowright I$  has a nontrivial indiscernible support function, then it has a nontrivial disjointifying closure operator.

And we are done.

5.1. Deriving an indiscernible support function from a Baire-measurable homo**morphism.** Our next goal is to show that if  $P \leq S_I$  classifies  $=^+$ , then there is a nontrivial indiscernible support function on  $P \curvearrowright I$ . We start by finding a presentation of  $=^+$  which is easier for us to work with.

Let  $\Delta \curvearrowright J$  be a free action of a countably-infinite group  $\Delta$  on a countably-infinite set J with infinitely-many orbits. Let  $T \subseteq J$  be a transversal for  $\Delta \curvearrowright J$  (i.e. a set which intersects every  $\Delta$ -orbit exactly once) and fix an enumeration  $T = \{t_n \mid n \in \omega\}$ . Let  $Q \leq S_J$ be the set of permutations  $\pi$  satisfying that  $\delta \cdot \pi(a) = \pi(\delta \cdot a)$  for every  $\delta \in \Delta$  and  $a \in J$ .

Now define Y to be the  $G_{\delta}$  set of all injections  $f: J \to \mathbb{R}$ . This is a Polish space with the pointwise-compactness topology (putting the discrete topology on J), and moreover the natural action  $Q \curvearrowright Y$  defined by

$$(g \cdot p)(a) = p(g^{-1} \cdot a)$$

is continuous with respect to this topology.

**Lemma 5.3.** The equivalence relations  $E_Y^Q$  and  $=^+$  are Borel bi-reducible.

*Proof.* We first see that  $=^{+}\leq_{B} E_{Y}^{Q}$ . Let  $\Delta = \{\delta_{n} \mid n \in \omega\}$  be an enumeration. Let  $g: \mathbb{R} \to \mathbb{R}^{\omega}$  be a Borel function satisfying

$$\{g(x)(n) \mid n \in \omega\} \cap \{g(x')(n) \mid n \in \omega\} = \emptyset$$

for any  $x \neq x' \in \mathbb{R}$ . One may see this as an application of the axiom of choice, but we remark that in our applications we will always take J to be  $\omega$ , in which case both the action of  $\Delta$  and T can be definable. Now define  $f: \mathbb{R}^{\omega} \to Y$  by

$$f(p) = y_p$$
 where  $y_p(\delta_n \cdot t_m) = g(p(m))(n)$ 

for every  $t_m \in T$  and  $\delta_n \in \Delta$ . This is easily seen to be a Borel reduction. Now we see that  $E_Y^Q \leq_B =^+$ . Fix an enumeration  $J = \{x_n \mid n \in \omega\}$  and a Borel bijection  $h: \mathbb{R}^{\Delta} \to \mathbb{R}$ . Define  $f: Y \to \mathbb{R}^{\omega}$  by

$$f(y) = p_y$$
 where  $p_y(n) = h(\delta \mapsto p(\delta \cdot x_n)).$ 

Using the fact that Q consists of only injective functions, this is a reduction, and it is clearly Borel. 

Let  $N \leq Q$  be the closed normal subgroup of  $g \in Q$  satisfying  $g \cdot a \in \Delta \cdot a$  for every  $a \in I$ . Let  $\chi: S_T \to Q$  be the homomorphism such that for every permutation  $\sigma \in S_T$ , the group element  $\chi(\sigma)$  is the unique one such that  $\chi(\sigma) \cdot a = \sigma(a)$  for every  $a \in T$ . Then N is the normal complement of  $\operatorname{Im}(\chi)$ , i.e.  $\operatorname{Im}(\chi) \cap N = \{1\}$  and  $Q = \operatorname{Im}(\chi)N$ . Define  $S_T^{\text{fin}}$  to be the countable subgroup of  $\pi \in S_T$  with finite support, i.e.  $\pi(a) = a$  for cofinitely-many  $a \in T$ . Define  $N_0$  to be the set of  $h \in N$  with "finite-support" in the sense that  $h \cdot a = a$  for cofinitely-many  $x \in T$ . Then define  $Q_0 = \chi[S_T^{\text{fin}}]N_0$ , which is easily seen to be a countable dense subgroup of Q.

Let I be a countably-infinite set and let  $P \leq S_I$  be a closed subgroup with the natural action  $P \curvearrowright I$ . Let X be a Polish P-space. Fix a Baire-measurable homomorphism  $f: Y \rightarrow I$ 

X from  $E_Y^Q$  to  $E_X^P$ . Then the action  $Q \curvearrowright Y$  restricts to an action  $N \curvearrowright Y$  and f is also a homomorphism from  $E_Y^N$  to  $E_X^P$ . From f we will derive an indiscernible support function. Assuming that f is not degenerate in a way which we will soon define, the support function will be non-trivial.

Let  $C \subseteq Y$  be the comeager set satisfying properties (1)-(4) in Lemma 2.3 applied to f as a Baire-measurable homomorphism from  $E_Y^Q$  to  $E_X^P$ , with  $Q_0$  as the countable dense subgroup of Q.

Fix some point  $y_0 \in C$ . For finite  $A \subseteq I$  and  $u \subseteq T$ , say that u **supports** A iff there is a basic open neighborhood  $U \ni y_0$  such that for every  $y \in U \cap C$  and for every  $g \in \text{Stab}_{Q_0}(u)$ , if  $g \cdot y \in U$  then  $f(g \cdot y) \in \text{Stab}_P(A) \cdot f(y)$ . By Lemma 2.3, for every finite A there exists some finite u such that u supports A.

**Claim 1.** For every finite  $A \subseteq I$  and  $u, v \subseteq T$ , if both u and v support A then  $u \cap v$  supports A.

Proof. Let  $U \ni y_0$  witnessing that u and v both support A. We argue that U witnesses that  $u \cap v$  supports A as well. Let  $y \in U \cap C$  and  $g \in \operatorname{Stab}_{Q_0}(u \cap v)$  be arbitrary such that  $g \cdot y \in U$ . Write  $g = \chi(\pi)h$  where  $\pi \in S_T^{\operatorname{fin}} \cap \operatorname{Stab}_{S_T}(u \cap v)$  and  $h \in \operatorname{Stab}_{N_0}(u \cap v)$ . Fix  $h_0 \in \operatorname{Stab}_{N_0}(v)$  and  $h_1 \in \operatorname{Stab}_{N_0}(T \setminus v)$  such that  $h = h_1h_0$ .

Let  $W \subseteq T$  be a large enough set such that  $\Delta \cdot T$  contains the support of  $\pi$  and h, the support of U, as well as u and v.

We first check that there exists  $\pi_0, \pi_2 \in \operatorname{Stab}_{S_T}(v)$  and  $\pi_1 \in \operatorname{Stab}_{S_T}(u)$ , each with finite support, such that

- (1)  $\pi = \pi_2 \circ \pi_1 \circ \pi_0$ ; and
- (2)  $\chi(\pi_0)h_0 \cdot x \in U$ ; and
- (3)  $\chi(\pi_1)h_1\chi(\pi_0)h_0 \cdot x \in U.$

Let  $\sigma \in S_T$  be an involution such that  $\sigma[u \setminus v] \cap W = \emptyset$  with support  $(u \setminus v) \cup \sigma[u \setminus v]$ . By a density argument, given that  $y \in U$  and  $\chi(\pi)h \cdot y \in U$ , we may also ensure  $\chi(\sigma)h_0 \cdot y \in U$ and  $\chi(\sigma)\chi(\pi)h \cdot y \in U$ . Define  $\pi_0 = \pi_2 = \sigma$ .

With the observation that  $h_1$  and  $\chi(\pi_0)$  commute, it is enough to find  $\pi_1$  to satisfy condition (1) and then conditions (2) and (3) would immediately follow.

Define  $\pi_1 \in S_T$  by

$$\pi_1(a) = \begin{cases} \sigma(\pi(\sigma(a))) & a \in \sigma[u \setminus v] \text{ and } \pi(\sigma(a)) \in u \setminus v \\ \pi(\sigma(a)) & a \in \sigma[u \setminus v] \text{ and } \pi(\sigma(a)) \notin u \setminus v \\ \sigma(\pi(a)) & a \notin u \setminus v \text{ and } \pi(a) \in u \setminus v \\ a & a \in u \\ \pi(a) & \text{otherwise.} \end{cases}$$

It is easy to check that  $\pi_1 \in \operatorname{Stab}_{S_T}(u)$  and has finite support. Checking that it is welldefined is straightforward. To see that  $\pi = \pi_2 \circ \pi_1 \circ \pi_0$ , we fix an arbitrary  $a \in J$  and check that  $\pi(a) = (\pi_2 \circ \pi_1 \circ \pi_0)(a)$ . First consider the case that  $a \in u \setminus v$  and  $\pi(a) \notin u \setminus v$ . Then by the definition of  $\pi_1$ , we have

$$\pi_2(\pi_1(\pi_0(a))) = \sigma(\pi_1(\sigma(a))) = \sigma(\pi(a)) = \pi(a).$$

The case that  $a \in u \setminus v$  and  $\pi(a) \in u \setminus v$  is similar. The next case we have is  $a \notin u \setminus v$  and  $\pi(a) \in u \setminus v$  where we have

$$\pi_2(\pi_1(\pi_0(a))) = \sigma(\pi_1(\sigma(a))) = \sigma(\pi_1(a)) = \sigma(\sigma(\pi(a))) = \pi(a).$$

The case  $a \in u \cap v$  is easily checked as a is fixed by  $\pi_2$ ,  $\pi_1$ , and  $\pi_2$ . The final case is that  $a \notin u$  and  $\pi(a) \notin u$ . Then

$$\pi_2(\pi_1(\pi_0(a))) = \sigma(\pi_1(\sigma(a))) = \sigma(\pi_1(a)) = \sigma(\pi(a)) = \pi(a).$$

We can conclude then that

- (a)  $\chi(\pi_1)h_0 \in \operatorname{Stab}_{Q_0}(v)$  and  $\chi(\pi_1)h_0 \cdot y = \chi(\sigma)h_0 \cdot y \in U$  and thus  $f(\chi(\pi_1)h \cdot y) \in \operatorname{Stab}_P(A) \cdot f(y)$ ;
- (b)  $\chi(\pi_2)h_1 \in \operatorname{Stab}_{Q_0}(u)$  and  $\chi(\pi_2)h_1 \cdot (\chi(\pi_1)h_0 \cdot y) = \chi(\sigma)\chi(\pi)h \cdot y \in U$  and thus  $f(\chi(\pi_2)h_1\chi(\pi_1)h_0 \cdot y) = \operatorname{Stab}_P(A) \cdot f(y)$ ; and
- (c)  $\chi(\pi_3) \in \operatorname{Stab}_{Q_0}(v)$  and  $\chi(\pi_3) \cdot (\chi(\pi_2)h_1 \cdot (\chi(\pi_1)h_0 \cdot y)) = \chi(\pi)h \cdot y \in U$ , and thus  $f(\chi(\pi)h \cdot y) \in \operatorname{Stab}_P(A) \cdot f(y)$

as desired.

We conclude in particular that for every A there is a minimal u which supports A. The minimal support of A, denoted supp(A), is **the support of** A. Now we wish to show that supp is indiscernible. We will prove this after the following claim.

**Claim 2.** For any u and A, if u is a support of A witnessed by neighborhood  $U_0 \ni y_0$ , then for any finite-support  $\sigma \in S_T$  and  $h \in N_0$  and  $g \in P$  with  $\chi(\sigma)h \cdot y_0 \in U_0$  and  $f(\chi(\sigma)h \cdot y_0) = g \cdot f(y_0)$ , we have that  $\sigma^{-1} \cdot u$  is a support of  $g^{-1} \cdot A$ .

*Proof.* Let  $v = \sigma^{-1} \cdot u$  and  $B = g^{-1} \cdot A$ . Observe that  $\operatorname{Stab}_P(B) = g^{-1} \operatorname{Stab}_P(A)g$  and  $\operatorname{Stab}_{S_T}(v) = \sigma^{-1} \operatorname{Stab}_{S_T}(u)\sigma$ . We show that v is a support of B.

Find open  $U_1 \ni y_0$  by Lemma 2.3.(4) such that for every  $y \in U_1 \cap C$ ,  $\chi(\sigma)h \cdot y \in U_0$  and  $f(\chi(\sigma)h \cdot y) \in g \operatorname{Stab}_P(B) \cdot f(y)$ . We argue that  $U_1$  witnesses that v supports B.

Let  $y \in U_1 \cap C$  be arbitrary. Let  $g_v \in \operatorname{Stab}_{Q_0}(v)$  arbitrary and suppose  $g_v \cdot y \in U_1$ . We want to show that  $f(g_v y) \in \operatorname{Stab}_P(B) \cdot f(y)$ . Because  $y, g_v \cdot y \in U_1$  we have

(3) 
$$f(\chi(\sigma)h \cdot y) \in g\operatorname{Stab}_P(B) \cdot f(y)$$

and

(4) 
$$f(\chi(\sigma)hg_v \cdot y) \in g\operatorname{Stab}_P(B) \cdot f(g_v \cdot y).$$

Because  $\chi(\sigma)h \cdot y, \chi(\sigma)hg_v \cdot y \in U_0$  and

$$\chi(\sigma)hg_v \cdot y = [(\chi(\sigma)h)g_v(\chi(\sigma)h)^{-1}]\chi(\sigma)hg_v \cdot y,$$

where

$$(\chi(\sigma)h)g_v(\chi(\sigma)h)^{-1} \in \operatorname{Stab}_{Q_0}(u)$$

we have

(5) 
$$f(\chi(\sigma)h \cdot y) \in \operatorname{Stab}_P(A) \cdot f(\chi(\sigma)hg_v \cdot y).$$

Thus putting these together we have

$$f(g_v \cdot y) \in \operatorname{Stab}_P(B)g^{-1} \cdot f(\chi(\sigma)hg_v \cdot y)$$
  

$$\subseteq \operatorname{Stab}_P(B)\pi^{-1}\operatorname{Stab}_P(A) \cdot f(\chi(\sigma)h \cdot y)$$
  

$$\subseteq \operatorname{Stab}_P(B)g^{-1}\operatorname{Stab}_P(A)g\operatorname{Stab}_P(B) \cdot f(y)$$
  

$$= \operatorname{Stab}_P(B) \cdot f(y),$$

where the first inclusion is from Equation 4, the second from Equation 5, and the third from Equation 3 as desired.  $\hfill \Box$ 

Now we see that supp is indiscernible. Let  $A, B \subseteq I$  be finite and  $u, v \subseteq J$  finite such that  $\operatorname{supp}(A) = u$  and  $\operatorname{supp}(B) = v$ , and let  $U_0 \ni y_0$  be the open neighborhood witnessing this. Let  $v' \subseteq J$  such that  $v' \cong_u v$ . Fix a finite-support  $\sigma \in S_T$  such that  $\sigma(a) = a$  for every  $a \in u$ and  $\sigma[v'] = v$ . By a density argument, fix some  $h \in \operatorname{Stab}_{N_0}(u)$  such that  $\chi(\sigma)h \cdot y_0 \in U_0$ . Because  $\chi(\sigma)h \in \operatorname{Stab}_{Q_0}(u)$ , there is some  $g \in \operatorname{Stab}_P(A)$  such that  $f(\chi(\sigma)h \cdot y_0) = g \cdot f(y_0)$ . By the claim, this means that v' is a support of  $g^{-1} \cdot B$ , and because  $g \in \operatorname{Stab}_P(A)$ , we have  $B \cong_A g^{-1} \cdot B$ . A symmetric argument gives us that v' is in fact the minimal support of  $g^{-1} \cdot B$ , as desired.

Our final task is to show that supp is nontrivial, otherwise it carries no meaningful structure.

**Claim 3.** If supp is trivial, then  $f(y) E_X^P f(y')$  for every  $y, y' \in C$ .

*Proof.* Suppose supp is trivial, i.e. the support of every A is empty. In other words, for every finite  $A \subseteq I$ , there is an open neighborhood  $U_A$  of  $y_0$  such that for every  $y \in U_A \cap C$  and every  $h \in Q_0$  with  $g \cdot y \in U_A$ , we have  $f(h \cdot y) \in \operatorname{Stab}_P(A) \cdot y$ .

Write I as an increasing union  $\bigcup_n A_n$  of finite sets. Fix a compatible complete metric d on Y. For each n, let  $U_n$  be an open neighborhood of  $y_0$  with d-diameter less than 1/n and contained in  $U_{A_n}$ .

Let  $y \in Y \cap C$  by arbitrary. We will show that  $f(y) E_X^P f(y_0)$ . For every n, let  $h_n \in Q_0$  such that  $h_n \cdot y \in C \cap U_n$ . We have  $h_n \cdot y \to y_0$ , and so by continuity of f on C, we have  $f(h_n \cdot y) \to f(y_0)$ .

For every *n* we may find  $g_n \in \operatorname{Stab}_P(A_n)$  such that  $f(h_{n+1} \cdot y) = g_n \cdot f(h_n \cdot y)$ . Defining  $g_n^* := g_n \dots g_0$ , we have  $f(h_n \cdot y) = g_n^* \cdot f(y)$  for every *n*. The sequence  $(g_n^*)$  is Cauchy and thus  $g_n^* \to g_\infty$  for some  $g_\infty^* \in P$ . By continuity of the group action, we have  $g_\infty^* \cdot f(y) = f(y_0)$ 

Observe that for any  $x, x' \in I$ , the set of y such that  $y(x) \neq y_0(x')$  is dense and open. Thus there is some  $y \in C$  such that  $y(x) \neq y_0(x')$  for every  $x, x' \in I$ . In particular, this means  $Q \cdot y \neq Q \cdot y_0$ . By essentially the same argument we have that  $E_Y^Q$  is meager. We conclude that if f is not just a homomorphism but a reduction, then supp is nontrivial. The same could be concluded if f witnesses that  $E_Y^Q$  is not generically ergodic with respect to  $E_X^P$ .

Combining the results of this section, we get:

**Theorem 5.4.** Let I be a countable set and  $P \leq S_I$  a closed subgroup with the natural action  $P \curvearrowright I$ . The following are equivalent:

- (1)  $P \curvearrowright I$  has a nontrivial indiscernible support function;
- (2) P classifies =+;
- (3) there is a Polish P-space X such that  $=^+$  is not generically-ergodic with respect to  $E_X^P$ .

We have now proved all of the equivalences of the main theorem.

#### References

- [All] S. Allison. Non-Archimedean TSI Polish groups and their potential Borel complexity spectrum.
- [AP21] S. Allison and A. Panagiotopoulos. Dynamical obstructions to classification by (co)homology and other TSI-group invariants. *Trans. Amer. Math. Soc.*, 374:8793–8811, 2021.
- [BFKL16] J. T. Baldwin, Sy D. Friedman, M. Koerwien, and M. C. Laskowski. Three red herrings around Vaught's conjecture. *Trans. Amer. Math. Soc.*, 368:3673–3694, 2016.
- [CC22] J. D. Clemens and S. Coskey. New jump operators on equivalence relations. J. Math. Log., 22:Paper No. 2250015, 44, 2022.
- [Dei77] R. Deissler. Minimal models. J. Symbolic Logic, 42:254–260, 1977.
- [Gao98] S. Gao. On automorphism groups of countable structures. J. Symbolic Logic, 63:891–896, 1998.
- [Gao09] S. Gao. Invariant descriptive set theory, volume 293 of Pure and Applied Mathematics. CRC Press, 2009.
- [Hjo99] G. Hjorth. Orbit cardinals: on the effective cardinalities arising as quotient spaces of the form X/G where G acts on a Polish space X. Israel J. Math., 111:221–261, 1999.
- [Hjo00] G. Hjorth. Classification and orbit equivalence relations, volume 75 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2000.
- [Hjo01] G. Hjorth. Vaught's conjecture on analytic sets. J. Amer. Math. Soc., 14:125–143, 2001.
- [Hjo02a] G. Hjorth. A dichotomy theorem for turbulence. J. Symbolic Logic, 67:1520–1540, 2002.
- [Hjo02b] G. Hjorth. Knight's model, its automorphism group, and characterizing the uncountable cardinals. J. Math. Log., 2:113–144, 2002.
- [HKL98] G. Hjorth, A. S. Kechris, and A. Louveau. Borel equivalence relations induced by actions of the symmetric group. Ann. Pure Appl. Logic, 92:63–112, 1998.
- [Kan08] V. Kanovei. Borel equivalence relations, volume 44 of University Lecture Series. American Mathematical Society, Providence, RI, 2008.
- [Kru15] A. Kruckman. Notes on generalized Fraissé theory, 2015.
- [LP18] M. Lupini and A. Panagiotopoulos. Games orbits play and obstructions to Borel reducibility. Groups Geom. Dyn., 12:1461–1483, 2018.
- [LUa] M. Laskowski and D. Ulrich. Borel complexity of modules.
- [LUb] M. Laskowski and D. Ulrich. Characterizing the existence of a Borel complete expansion.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO Email address: shaunpallison@gmail.com