HOLOMORPHIC SLOW-FAST SYSTEMS

GABRIEL RONDÓN, PAULO R. DA SILVA AND LUIZ F. S. GOUVEIA

ABSTRACT. In this paper, we are concerned with studying the existence of invariant complex manifolds of two-dimensional holomorphic systems. From the geometric singular perturbation theory we know that if a slow-fast system has associated a normally hyperbolic compact critical manifold, then there exists a smooth locally invariant manifold. However, this smooth manifold does not necessarily have a complex structure. Here, we provide conditions to guarantee the existence of one-dimensional invariant complex manifolds. Consequently, this allows us to establish that the centers, foci, and nodes of the reduced problem are persistent by singular perturbation. The tools used by us are the usual techniques of Fenichel and Briot-Bouquet Theories.

1. Introduction

In the qualitative theory of dynamical systems, there are several studies on singular perturbation problems (see, for instance, [3, 9, 10, 17]). In this paper, we want to study singular perturbation problems in \mathbb{R}^4 , which can be written as two-dimensional holomorphic systems. It is important to highlight that although there are applications that can be modeled by two-dimensional holomorphic systems, see [11, 12], there are few articles on the subject such as [13, 14].

The basic systems we consider are of the form

(SF1)
$$\begin{cases} \varepsilon \dot{z} = f(z, w), \\ \dot{w} = g(z, w), \end{cases}$$

where $f,g:D\to\mathbb{C}$ are holomorphic functions, $D\subseteq\mathbb{C}^2$ is a simply connected domain, $\varepsilon>0$ is a small enough parameter and the dot represents the derivative of the functions $z(\tau)$ and $w(\tau)$ with respect to the real variable τ .

²⁰²⁰ Mathematics Subject Classification. 34C45, 34A09.

Key words and phrases. Holomorphic systems, geometric singular perturbation theory, Fenichel theory, invariant complex manifold.

Emphasize that $C_0 = \{(z, w) \in D : f(z, w) = 0\}$ is the onedimensional critical manifold associated with the system (SF1) provided that $\frac{\partial f}{\partial z}(z, w) \neq 0$ for all $(z, w) \in C_0$. Recall that (SF1) can be written as a real four-dimensional autonomous system in an appropriate domain of \mathbb{R}^4 .

Suppose that $S_0 \subset C_0$ is a normally hyperbolic compact critical manifold. Thus, we can assume that S_0 is given as the graph of a function of z in terms of w. Indeed, since $\frac{\partial f}{\partial z}(z,w) \neq 0$ for all $(z,w) \in S_0$, then from *Implicit Function Theorem* (see [15]), there exist an open set U and a holomorphic function z = h(w) such that f(h(w), w) = 0, for all $w \in U$. Hence, without loss of generality we can assume that $S_0 = \{(z,w) : z = h(w), w \in U\}$.

One of the main goals of this paper is to study the existence of invariant complex manifolds of two-dimensional holomorphic system (SF1). At some point, the reader may think that it is enough to apply Fenichel's theorem (see, for instance, [9]) to obtain a one-dimensional invariant complex manifold. In general, this is not true, due to the fact that there are smooth manifolds that have no complex structure. An obvious impediment is that the dimension of the smooth manifold must be even. Another obstacle is orientability, since every complex manifold is orientable.

Although the geometric singular perturbation theory does not allow us to guarantee the existence of invariant complex manifolds, it is possible to use this theory and the Laurent series to construct examples of smooth manifolds without complex structure. Indeed, if we apply the Fenichel's Theorem to the equivalent C^{∞} four-dimensional real system associated with system (SF1), then there exists a smooth locally invariant manifold $S_{\varepsilon} = \{(z, w) : z = h_{\varepsilon}(w)\}$ of slow-fast system (SF1), which is diffeomorphic to S_0 . In general, this function $h_{\varepsilon}(z)$ does not have to be holomorphic, as we illustrate in the following example. Consider the complex dynamical system

(1)
$$\begin{cases} \varepsilon \dot{z} = z + w, \\ \dot{w} = w^2, \end{cases}$$

notice that $C_0 = \{(z, w) : z = -w\}$ is a normally hyperbolic critical manifold of (1). In addition, if $S_0 \subset C_0$ is a compact set, then the equivalent system to (1) in \mathbb{R}^4 has a smooth locally invariant manifold S_{ε} . Nevertheless, this manifold has no complex structure. Indeed, suppose that there exists a holomorphic function h_{ε} such that $S_{\varepsilon} = \{(z, w) : z = h_{\varepsilon}(w)\}$. Then h_{ε} can be written as a power series. In

Section 5, we will prove that this series is given by

$$h_{\varepsilon}(w) = -\sum_{k=0}^{\infty} \varepsilon^k k! w^{k+1}.$$

Notice that this series diverges for $w \neq 0$, which contradicts the fact that h_{ε} is holomorphic.

At this point, it is natural to ask: under what conditions can we ensure the existence of invariant manifolds with complex structure of system (SF1)? In Section 3, we use the *Briot-Bouquet Theory* and provide conditions to guarantee the existence of one-dimensional invariant complex manifolds. Specifically, in Theorem A we prove that, for $\varepsilon > 0$ sufficiently enough, the system

(SF2)
$$\begin{cases} \varepsilon \dot{z} = \alpha z + \tilde{f}(z, w), \\ \dot{w} = \beta w + \tilde{g}(z, w), \end{cases} \text{ where } \tilde{f}, \tilde{g} = \mathcal{O}_2(z, w) \text{ and } \alpha, \beta \in \mathbb{C},$$

has a unique one-dimensional invariant complex manifold C_{ε} passing through the equilibrium point $q_{\varepsilon} = (0,0)$ provided that the critical manifold is normally hyperbolic.

We employ Theorem A to prove that the centers, foci and nodes of the reduced problem associated with the holomorphic slow-fast system (SF1), whose Jacobian matrix of f and g is diagonalizable at the origin, are preserved by singular perturbation, for more details see Theorem B and Figure 1.




FIGURE 1. Persistence of a focus, node and center of the reduced problem by singular perturbation, respectively.

Also, we are concerned with determining the existence of complex invariant manifolds for certain families of two-dimensional holomorphic systems, whose jacobian matrix of f and g is not necessarily digonalizable at the origin and this point does not need to be an equilibrium point of the system, see Theorems C and D. Specifically, Theorem C

stablishes the existence of one-dimensional invariant complex manifolds associated with slow-fast system (SF1) whose equation of orbits is separable, that is, in the case that there exist holomorphic functions η, κ such that $\frac{g(z,w)}{f(z,w)} = \eta(z)\kappa(w)$ in a suitable domain. Furthermore, in Theorem D we state conditions to ensure when the system

(SF3)
$$\begin{cases} \varepsilon \dot{z} = \alpha z + \beta G(w), \\ \dot{w} = g(w), \end{cases} \text{ where } G' \equiv 1/g \text{ and } \alpha, \beta \in \mathbb{C} \setminus \{0\},$$

has associated a one-dimensional exponentially attractive invariant complex manifold C_{ε} .

Finally, we will present a method to study slow-fast systems of the form

(SF4)
$$\begin{cases} \varepsilon \dot{z} = \alpha z + \beta w, \\ \dot{w} = g(w), \end{cases}, \text{ where } \alpha \in \mathbb{C} \setminus \{0\} \text{ and } \beta \in \mathbb{C},$$

where g is one of the normal forms given in [4], namely: $\dot{w}=1, \dot{w}=(a+ib)w, \ \dot{w}=w^n, \ \dot{w}=\frac{1}{w^n}$ and $\dot{w}=\frac{\gamma w^n}{1+w^{n-1}}$. We will determine which of these families could have associated manifolds with complex structure and in that case we will investigate what happens when a singularity of the reduced problem is perturbed for $\varepsilon>0$. Specifically, in Propositions 10 and 11 we prove that the centers and poles of order n of the reduced problem associated with system (SF4) are persistent by singular perturbation.

The paper is organized in the following form. In Section 2, we present some basic results on the complex manifolds, geometric singular perturbation theory and the Briot-Bouquet Theory. In Section 3, we employ Briot-Bouquet theory to prove Theorem A and B. In Section 4, we state and prove Theorems C and D. Lastly, in Section 5 we use the Laurent series to give complex systems that have no associated complex manifolds.

2. Preliminaries

This section is devoted to establishing some basic results that will be used throughout the paper.

- 2.1. Complex manifolds. A holomorphic function f is a complex-valued function defined in a domain $\mathcal{V} \subseteq \mathbb{C}$ and satisfying that
 - u = Re(f) and v = Im(f) are continuous,
 - there exist the partial derivatives u_x, u_y, v_x, v_y in \mathcal{V} and,

• the partial derivatives satisfy the Cauchy–Riemann equations, see [5],

$$u_x = v_y, \quad u_y = -v_x, \quad \forall z = x + iy \in \mathcal{V}.$$

We recall that Looman-Menchoff's Theorem establishes that the above conditions are sufficient to guarantee the analyticity of f. This result implies that for any $z_0 \in \mathcal{V}$

$$f(z) = A_0 + A_1(z - z_0) + A_2(z - z_0)^2 + \dots, \quad A_k = a_k + ib_k = \frac{f^{(k)}(z_0)}{k!}$$

for $z \in D(z_0, R_{z_0}) \subseteq \mathcal{V}$ where $D(z_0, R_{z_0})$ is the largest possible z_0 -centered disk contained in \mathcal{V} .

If f is holomorphic in a punctured disc $D(0, R) \setminus \{0\}$ and it is not derivable at 0 we say that 0 is a singularity of f. In this case f is equal to its Laurent's series in $D(0, R) \setminus \{0\}$

$$f(z) = \sum_{k=1}^{\infty} \frac{B_k}{z^k} + \sum_{k=0}^{\infty} A_k z^k,$$

where $B_k = \frac{1}{2\pi i} \int_{C_{\varepsilon}} f(z) z^{k-1} dz$, $A_k = \frac{1}{2\pi i} \int_{C_{\varepsilon}} \frac{f(z)}{z^{k+1}} dz$ with C_{ε} parameterized by $z(t) = \varepsilon e^{it}$, $\varepsilon \sim 0$, $t \in [0, 2\pi]$.

If $B_k \neq 0$ for an infinite set of indices k we say that 0 is an essential singularity and if there exists $n \geq 1$ such that $B_n \neq 0$ and $B_k = 0$ for every k > n then we say that 0 is a pole of order n.

The notion of holomorphic function can be extended as follow.

Consider $W \subset \mathbb{C}^2$ an open subset and let $f: W \to \mathbb{C}$ be a continuously differentiable function. Then f is said *holomorphic* if the Cauchy-Riemann equations hold for all coordinates $z_j = x_j + iy_j$, with j = 1, 2, i.e.

$$u_{x_j} = v_{y_j}, \quad u_{y_j} = -v_{x_j}, \quad \forall z_j = x_j + iy_j \in \mathcal{W}.$$

In addition, if f is holomorphic in W, then in each point $\zeta = (\zeta_1, \zeta_2) \in W$ has an open neighborhood $D(\zeta, R_{\zeta})$, such that the function f can be expanded into a power series

$$f(z,w) = \sum_{n=1}^{\infty} \sum_{k_1+k_2=n} c_{k_1,k_2} (z-\zeta_1)^{k_1} (w-\zeta_2)^{k_2},$$

which converges for all $(z, w) \in D(\zeta, R_{\zeta})$.

The definition of a complex manifold is analogous to the smooth manifold, but we require that the charts take on values in \mathbb{C}^2 and that the transition functions be holomorphic (see Figure 2).




FIGURE 2. Complex manifolds require that the transition functions φ_{ij} be holomorphic.

A holomorphic atlas in a topological space $X \subseteq \mathbb{C}^2$ is a collection of pairs (U_i, φ_i) called holomorphic charts where

- Each U_i is an open in X and $X = \bigcup_i U_i$;
- Each $\varphi_i: U_i \to \mathbb{C}^n$ is a homomorphism over an open of \mathbb{C}^n , with $n \leq 2$;
- Whenever $U_i \cap U_j \neq \emptyset$ the transition

$$\varphi_{ij} = \varphi_i \circ \varphi_i^{-1} : \varphi_i(U_i \cap U_j) \to \varphi_i(U_i \cap U_j)$$

is a holomorphic map.

A complex manifold is a topological space X, Hausdorff and with enumerable base, equipped with a maximal holomorphic atlas. The number n is called the complex dimension of X.

The following proposition will give us a way to obtain complex submanifolds. For more details see, for instance, [7, 18].

Proposition 1. Let $\phi: X \subseteq \mathbb{C}^2 \to \mathbb{C}$ be a holomorphic map between complex manifolds. Consider $b \in \phi(X) \subseteq \mathbb{C}$ such that the rank of ϕ is maximal $(rank(\phi) = 1)$, for all $a \in \phi^{-1}(b)$. Then $\phi^{-1}(b)$ is a complex submanifold of X of dimension 1.

2.2. Geometric singular perturbation theory. We consider singularly perturbed systems of differential equations

(2)
$$\begin{cases} \varepsilon \dot{\mathbf{x}} = f(\mathbf{x}, \mathbf{y}, \varepsilon), \\ \dot{\mathbf{y}} = g(\mathbf{x}, \mathbf{y}, \varepsilon), \end{cases}$$

which is called *slow-fast system*, where $\varepsilon \in (0, \varepsilon_0)$, ε_0 small, $(\mathbf{x}, \mathbf{y}) \in \mathcal{M}$, an open set $\mathcal{M} \subset \mathbb{R}^4$ and $f, g : (0, \varepsilon_0) \times \mathcal{M} \to \mathbb{R}^4$ are C^r . The dot represents the derivative of the functions $\mathbf{x}(\tau)$ and $\mathbf{y}(\tau)$ with respect to the variable τ .

If we write $t = \frac{\tau}{\varepsilon}$, then system (2) becomes

(3)
$$\begin{cases} \mathbf{x}' = f(\mathbf{x}, \mathbf{y}, \varepsilon), \\ \mathbf{y}' = \varepsilon g(\mathbf{x}, \mathbf{y}, \varepsilon), \end{cases}$$

in which the apostrophe 'denotes the derivative of the functions $\mathbf{x}(t)$ and $\mathbf{y}(t)$ with respect to the variable t. Notice that the parameter $\varepsilon = \frac{\tau}{t}$ represents the ratio of the time scales.

Consider equation (2) and set $\varepsilon = 0$. We obtain the so called *reduced* problem given by

(4)
$$0 = f(\mathbf{x}, \mathbf{y}, 0), \quad \dot{\mathbf{y}} = g(\mathbf{x}, \mathbf{y}, 0).$$

Observe that (4) is not an ordinary differential equation, but it is an algebraic differential equation.

Solutions of (4) are contained in the set

$$C_0 = \{ (\mathbf{x}, \mathbf{y}) \in \mathcal{M} : f(\mathbf{x}, \mathbf{y}, 0) = 0 \}.$$

The set C_0 is called *critical set*. In the case where C_0 is a manifold, C_0 is called critical manifold.

On the other hand, setting $\varepsilon = 0$ in equation (3) we obtain the so called *layer problem*

(5)
$$\mathbf{x}' = f(\mathbf{x}, \mathbf{y}, 0), \quad \mathbf{y}' = 0.$$

Moreover, the system (5) can be seen as a system of ordinary differential equations, where $\mathbf{y} \in \mathbb{R}^2$ is a parameter and the critical set C_0 is a set of equilibrium points of (5).

The main goal of geometric singular perturbation theory is to study systems (4) and (5) in order to obtain information of the full system (2). Observe that the systems (2) and (3) are equivalent when $\varepsilon > 0$, since they only differ by time scale.

We next introduce the notion of normally hyperbolic points.

Let $\mathbf{x}_0 \in S$, for any set $S \subset \mathcal{M}$. We say that \mathbf{x}_0 is normally hyperbolic if the 2×2 matrix $Df_{\mathbf{x}}(\mathbf{x}_0)$ does not have eigenvalues with zero real part.

In what follows we present a version of *Fenichel's Theorem* whose proof can be found in [16].

Theorem 2. Consider \mathcal{M} a C^{r+1} manifold, $2 \leq r \leq \infty$. Let X_{ε} , $\varepsilon \in (-\varepsilon_0, \varepsilon_0)$ be a C^r family of vector fields on \mathcal{M} , and assume C_0 a C^r submanifold of \mathcal{M} consisting entirely of equilibrium points of X_0 . If $S \subset C_0$ is a j-dimensional compact normally hyperbolic invariant manifold of the reduced vector field X_R with a $j + j^s$ -dimensional local

۶

stable manifold W^s and a $j + j^u$ -dimensional local unstable manifold W^u , then there exists $\varepsilon_1 > 0$ such that

- (i) There exists a C^{r-1} family of manifolds $\{S_{\varepsilon} : \varepsilon \in (-\varepsilon_1, \varepsilon_1)\}$ with $S_0 = S$ and S_{ε} is a normally hyperbolic invariant manifold of X_{ε} .
- (ii) There are C^{r-1} families of $(j+j^s+k^s)$ -dimensional and $(j+j^u+k^u)$ -dimensional manifolds $\{S^{\varepsilon}_{\varepsilon}: \varepsilon \in (-\varepsilon_1,\varepsilon_1)\}$ and $\{S^{u}_{\varepsilon}: \varepsilon \in (-\varepsilon_1,\varepsilon_1)\}$ such that for $\varepsilon > 0$ the manifolds S^{s}_{ε} and S^{u}_{ε} are local stable and unstable manifolds of S_{ε} .

Now, suppose that S is given as in Theorem 2, then $Df_{\mathbf{x}}(\mathbf{x}, \mathbf{y}, 0)$ is invertible for all $(\mathbf{x}, \mathbf{y}) \in S$. From *Implicit Function Theorem* there exists a smooth function h_0 defined on a compact domain $K \subset \mathbb{R}^2$ such that

$$S = \{(\mathbf{x}, \mathbf{y})\} : \mathbf{x} = h_0(\mathbf{y})\}.$$

That is, S is locally the graph of a function of \mathbf{x} in terms of \mathbf{y}

Theorem 3. Assume the same hypotheses of Theorem 2. If $\varepsilon > 0$ is sufficiently small, there exists a function $\mathbf{x} = h_{\varepsilon}(\mathbf{y})$, defined on K, so that the graph

$$S_{\varepsilon} = \{(\mathbf{x}, \mathbf{y})) : \mathbf{x} = h_{\varepsilon}(\mathbf{y})\}$$

is locally invariant under (3). Moreover h_{ε} is C^r , for any $r < \infty$, jointly in \mathbf{y} and ε .

2.3. **Two-dimensional holomorphic slow-fast systems.** Consider the two-dimensional holomorphic slow-fast system (SF1), where $f(z, w) = u_1+iv_1$ and $g(z, w) = u_2+iv_2$ are holomorphic functions of the complex variables $(z, w) = (x_1 + iy_1, x_2 + iy_2) \in D$.

Recall that the fast system associated to system (SF1) is given by

(6)
$$\begin{cases} z' = f(z, w) = u_1 + iv_1, \\ w' = \varepsilon g(z, w) = \varepsilon (u_2 + iv_2), \end{cases}$$

where the apostrophe 'denotes the derivative of the functions z(t) and w(t) with respect to the variable t, with $t = \tau/\varepsilon$. Notice that the critical set associated to system (6) is defined as $C_0 = \{(z, w) \in D : f(z, w) = 0\}$.

In this context the reduced problem is given by

(7)
$$0 = f(z, w), \ \dot{w} = g(z, w),$$

and the layer problem is defined as

(8)
$$z' = f(z, w), \ w' = 0.$$

Observe that the systems (SF1) and (6) are equivalent when $\varepsilon > 0$, since they only differ by time scale. In addition, systems (7) and (8) allow us to obtain information about system (SF1).

Now, we introduce the notion of normal hyperbolicity in the complex context. Let $q_0 = (z_0, w_0) \in C_0$ be an equilibrium point of the system (6) for $\varepsilon = 0$. Since f, g are holomorphic functions, then the Cauchy-Riemann equations hold. Thus, the Jacobian matrix of the system in \mathbb{R}^4 associated to (6) at q_0 is given by

$$(9) J_{\mathbb{R}}(f,0)|_{q_0} = \begin{pmatrix} (u_1)_{x_1} & -(v_1)_{x_1} & (u_1)_{x_2} & -(v_1)_{x_2} \\ (v_1)_{x_1} & (u_1)_{x_1} & (v_1)_{x_2} & (u_1)_{x_2} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

this implies that the eigenvalues of $J_{\mathbb{R}}(f,0)|_{q_0}$ are $\lambda_{\pm} = (u_1)_{x_1} \pm i(v_1)_{x_1}$. Hence, a subset $S \subset C_0$ is called normally hyperbolic provided that $\operatorname{Re}\left(\frac{\partial f}{\partial z}(q)\right) = (u_1)_{x_1}(q) \neq 0$ for all $q \in S$.

2.4. **Briot-Bouquet Systems.** Briot-Bouquet systems have been widely used in the research literature (see, for instance, [6, 13, 14]).

A Briot-Bouquet system is of the following form

(10)
$$zw' = F(z, w), \quad F(0, 0) = 0,$$

where F is a holomorphic function and the apostrophe 'denotes the derivative of the function w(z) with respect to the complex variable z.

The one-dimensional theory is well known since the original work of Briot and Bouquet (see [1]). Here we will use the following result, which was proved in [8, Proposition 1.1.1].

Proposition 4 (Briot-Bouquet criterion). If $F(z, w) = \lambda w + \mathcal{O}(z, w^2)$ and $\lambda \notin \mathbb{N}$, then (10) admits a unique holomorphic solution at z = 0 satisfying w(0) = 0.

3. Persistence of Invariant Manifold

This section aims to prove that centers, foci and nodes of the reduced problem associated with the holomorphic slow-fast system (SF1), whose Jacobian matrix of f and g is diagonalizable at the origin, are preserved by singular perturbation.

For that, first we are going to construct one-dimensional invariant complex manifolds via the *Briot-Bouquet Theory*. Thus, consider the complex differential system given by (SF1). Without loss of generality, assume that $q_{\varepsilon} = (0,0)$ is an equilibrium point of (SF1) in D. Suppose that the associated linearized system is such that the eigenvalues of the

10

Jacobian matrix $J_{\mathbb{C}}(f,g)|_{q_{\varepsilon}}$ of f,g are given by $\alpha,\beta\in\mathbb{C}$ and $J_{\mathbb{C}}(f,g)|_{q_{\varepsilon}}$ is diagonalizable. Hence, we can write (SF1) as (SF2) where

(11)
$$\tilde{f}(z,w) = \sum_{n=2}^{\infty} \sum_{s+l=n} a_{s,l} z^s w^l$$
 and $\tilde{g}(z,w) = \sum_{n=2}^{\infty} \sum_{s+l=n} b_{s,l} z^s w^l$,

which convergent in some neighbourhood of q_{ε} . Here s, l are nonnegative integers and $a_{s,l}, b_{s,l}$ are complex constants.

In this context, the critical set associated with complex system (SF2) is given by $C_0 = \{(z, w) : \alpha z + \tilde{f}(z, w) = 0\}$. Emphasize that if C_0 is a normally hyperbolic critical manifold, then $\text{Re}(\alpha) \neq 0$.

To find an invariant manifold of system (SF2), we consider the following initial value problem

(12)
$$(\alpha z + \tilde{f}(z, w))\frac{dw}{dz} = \varepsilon(\beta w + \tilde{g}(z, w)), \quad w_{\varepsilon}(0) = 0, \quad w'_{\varepsilon}(0) = 0.$$

In what follows we present a fundamental result that will be used to prove the main theorem of this section.

Theorem A. Suppose that C_0 is a normally hyperbolic critical manifold associated with complex system (SF2) and consider $\varepsilon > 0$ a small enough parameter. Then, system (SF2) has a unique one-dimensional invariant complex manifold C_{ε} passing through the equilibrium point q_{ε} , which is given by

$$C_{\varepsilon} = \{(z, w) : w = h_{\varepsilon}(z), \quad h_{\varepsilon}(0) = 0, \quad h'_{\varepsilon}(0) = 0\},$$

where h_{ε} is holomorphic in B = B(0, r).

Proof. First, we prove that the initial value problem (12) has a unique solution $w = h_{\varepsilon}(z)$, which is holomorphic in B. Indeed, consider the holomorphic transformation $w(z) = z\varphi(z)$, where $z \in B$. Then, the initial value problem (12) can be written as

(13)
$$(\alpha z + \tilde{f}(z, z\varphi))[\varphi + z\varphi'] = \varepsilon(\beta z\varphi + \tilde{g}(z, z\varphi)), \quad \varphi_{\varepsilon}(0) = 0.$$

Since C_0 is a normally hyperbolic critical manifold, then $\text{Re}(\alpha) \neq 0$. Thus, dividing equation (13) by αz , we get

(14)
$$\left(1 + \frac{\tilde{f}(z, z\varphi)}{\alpha z}\right) [\varphi + z\varphi'] = \varepsilon \left(\frac{\beta \varphi}{\alpha} + \frac{\tilde{g}(z, z\varphi)}{\alpha z}\right).$$

Now, we define the holomorphic functions $p(z,\varphi):=\frac{\tilde{f}(z,z\varphi)}{\alpha z}$ and $q(z,\varphi):=\frac{\tilde{g}(z,z\varphi)}{\alpha z}$. Then, from (11) and $w=z\varphi$, we obtain

$$\tilde{p}(z,\varphi) = \frac{1}{\alpha} \sum_{n=2}^{\infty} \sum_{s+l=n} a_{s,l} \varphi^l z^{n-1} \quad \text{and} \quad \tilde{q}(z,\varphi) = \frac{1}{\alpha} \sum_{n=2}^{\infty} \sum_{s+l=n} b_{s,l} \varphi^l z^{n-1},$$

which converge in some neighbourhood of q_{ε} . Rewriting equation (14), we get

$$z\varphi' = -\varphi + \varepsilon \frac{\frac{\beta\varphi}{\alpha} + q(z,\varphi)}{1 + p(z,\varphi)} := F_{\varepsilon}(z,\varphi).$$

Thus, putting the linear part of F_{ε} in an explicit way, we obtain that

(15)
$$z\varphi' = -\left(1 - \frac{\varepsilon\beta}{\alpha}\right)\varphi + \frac{\varepsilon b_{20}}{\alpha}z + \varepsilon Q(z,\varphi),$$

where

$$Q(z,\varphi) = \left(q(z,\varphi) - \frac{b_{20}}{\alpha}z\right) - \frac{q(z,\varphi)p(z,\varphi)}{1 + p(z,\varphi)} - \frac{\beta\varphi p(z,\varphi)}{\alpha(1 + p(z,\varphi))}.$$

Recall that F_{ε} is a holomorphic function in a neighborhood of q_{ε} , $F_{\varepsilon}(0,0)=0$ and $Q(z,\varphi)=\mathcal{O}_2(z,\varphi)$. For $\varepsilon>0$ small enough, we have that $\frac{\varepsilon\beta}{\alpha}-1\notin\mathbb{N}$, then from Briot-Bouquet criterion (see Proposition 4) we conclude that (15) has a unique solution $\varphi=\Lambda_{\varepsilon}(z)$, which is holomorphic in B. Using that $w=z\varphi$, then the initial value problem (12) has a unique solution $w=h_{\varepsilon}(z)$, where $h_{\varepsilon}(z)=z\Lambda_{\varepsilon}(z)$ is holomorphic in B.

Now, consider the holomoprhic function $H_{\varepsilon}(z,w) = w - h_{\varepsilon}(z)$, with $H_{\varepsilon}(0,0) = 0$ and $\frac{\partial H_{\varepsilon}}{\partial z}(0,0) = 0$. Define the set $C_{\varepsilon} := \{(z,w) : w = h_{\varepsilon}(z), h_{\varepsilon}(0) = 0, h'_{\varepsilon}(0) = 0\}$. Emphasize that $C_{\varepsilon} = H_{\varepsilon}^{-1}(0)$. By construction, we get that C_{ε} is an invariant set.

Notice that the rank of H is the rank of the matrix 1×2 , which is given by

$$J_{\mathbb{C}}H_{\varepsilon}(z,w) = \left(\frac{\partial H}{\partial z}(z,w), \frac{\partial H}{\partial w}(z,w)\right) = \left(-\frac{\partial h_{\varepsilon}}{\partial z}(z), 1\right).$$

Thus, $J_{\mathbb{C}}H$ has maximal rank (rank 1) in each point of C_{ε} . From Proposition 1, we conclude that C_{ε} is a complex manifold of dimension 1. \square

Remark 5. From the proof of Theorem A, we can deduce that this result still holds when we replace the hypothesis in which C_0 is a normally hyperbolic manifold by $\frac{\partial f}{\partial z}(0,0) = \alpha \neq 0$, which is a weaker hypothesis.

In the sequel, assume that:

(H) the eigenvalues of the $J_{\mathbb{C}}(f,g)|_{q_{\varepsilon}}$ are $\alpha,\beta\in\mathbb{C}\setminus\{0\}$ and that α,β are of the same type, that is, both α and β are pure imaginary complex numbers, real numbers, or complex numbers with nonzero real and imaginary parts.

At this point, we can assume that C_0 is given as the graph of a function of z in terms of w. Indeed, define $\Theta(z,w) = \alpha z + \tilde{f}(z,w)$ and $q_0 = (0,0)$. Since $\Theta(q_0) = 0$ and $\frac{\partial \Theta}{\partial z}(q_0) = \alpha \neq 0$, then from *Implicit*

12

Function Theorem, there exist a neighborhood U_{q_0} of q_0 and a holomorphic function z = L(w) such that L(0) = 0 and $\Theta(L(w), w) = 0$, for all $w \in U_{q_0}$. Therefore, locally the critical manifold is given by $C_0 = \{(z, w) : z = L(w), \quad L(0) = 0, \quad w \in U\}$.

Recall that the dynamics on C_0 is given by

(16)
$$\dot{w} = \beta w + g(L(w), w) =: G(w).$$

From item (b) of [4, Theorem 1.1] we know that G and βw are 0-conformally conjugate.

We are ready to state the main result of this section.

Theorem B. Suppose that the eigenvalues of $J_{\mathbb{C}}(f,g)|_{q_{\varepsilon}}$ satisfy hypothesis **(H)**, where ε is a small enough positive parameter. Then, centers, foci, and nodes of the reduced problem associated to (SF1) are persistent by singular perturbation.

Proof. By Theorem A, we know that, for $\varepsilon > 0$ small enough, system (SF2) has a unique one-dimensional invariant complex manifold C_{ε} at q_{ε} , which is given by

$$C_{\varepsilon} = \{(z, w) : w = h_{\varepsilon}(z), \quad h_{\varepsilon}(0) = 0, \quad h'_{\varepsilon}(0) = 0\},$$

where h_{ε} is holomorphic in B.

Emphasize that the dynamics about C_{ε} is given by

(17)
$$\dot{z} = \frac{\dot{w}}{h'_{\varepsilon}(z)} = \frac{\alpha}{\varepsilon} z + \frac{\tilde{f}(z, h_{\varepsilon}(z))}{\varepsilon} =: \Gamma_{\varepsilon}(z).$$

By item (b) of [4, Theorem 1.1] we can conclude that Γ_{ε} and $\frac{\alpha}{\varepsilon}z$ are 0-conformally conjugate. From (16) and (17) and the fact that α and β are of the same type the result follows.

4. Other families of holomorphic slow-fast systems

In this section, we are concerned with finding one-dimensional invariant complex manifolds of slow-fast systems whose Jacobian matrix is not necessarily diagonalizable at the origin. Furthermore, we do not require that the origin be an equilibrium point of the system.

4.1. **Uncoupled differential systems.** Consider the two-dimensional holomorphic slow-fast system (SF1), where f, g are holomorphic functions defined in a punctured disk $D = D(0, R) \setminus \{0\}$.

We assume that $\frac{\partial f}{\partial w}(z, w) \neq 0$, for all $(z, w) \in C_0$. From Proposition 1, we know that C_0 is a complex manifold. Moreover, we can assume that C_0 is given as the graph of a function of w in terms of z. Indeed, since $\frac{\partial f}{\partial w}(z, w) \neq 0$ for all $(z, w) \in C_0$, then from *Implicit Function Theorem*,

there exist an open set U and a holomorphic function $w = \lambda(z)$ such that $f(z, \lambda(z)) = 0$, for all $z \in U$.

Let us introduce the equations for the orbits of system (SF1):

(18)
$$\frac{dw}{dz} = \varepsilon \frac{g(z, w)}{f(z, w)}.$$

Suppose that there exist holomorphic functions η , κ defined in $\Omega = B \setminus L$ such that $\frac{g(z,w)}{f(z,w)} = \eta(z)\kappa(w)$ for all $z,w \in \Omega$, where L is a ray starting at $0, B = B(0,r) \setminus \{0\}$ and

- either 0 is the single zero of η (resp. κ) in B;
- or 0 is an isolated singularity of η (resp. κ) and η (resp. κ) has no zeros in B.

From [2, Corollary 6.16], we know that the holomorphic functions η, κ : $\Omega \to \mathbb{C}$ have a primitive in Ω . Thus $1/\eta, 1/\kappa$ are also holomorphic functions in Ω and consequently both η, κ and $1/\eta, 1/\kappa$ have primitives in Ω . Integrating the equation $\frac{dw}{dz} = \varepsilon \eta(z)\kappa(w)$, we get that there exist holomorphic functions F, G such that $G(w) = \varepsilon F(z)$, $F'(z) = \eta(z)$ and $G'(w) = 1/\kappa(w)$.

In what follows we state an interesting result about the existence of invariant complex manifolds associated with slow-fast systems whose equation of the orbits is separable.

Theorem C. Consider the slow-fast system (SF1) and assume that there exist holomorphic functions η, κ defined in Ω such that $\frac{g(z,w)}{f(z,w)} = \eta(z)\kappa(w)$, for all $z, w \in \Omega$. Then, there exists a one-dimensional invariant complex manifold C_{ε} associated with the differential system (SF1), such that

- (a) the flow on C_{ε} converges to the slow flow as $\varepsilon \to 0$.
- (b) the dynamics over C_{ε} is given by $\dot{z} = g(z, w)$.

Proof. Consider the holomoprhic function $H_{\varepsilon}(z, w) = G(w) - \varepsilon F(z)$ and the set $C_{\varepsilon} = \{(z, w) : G(w) = \varepsilon F(z)\}$. Recall that $C_{\varepsilon} = H_{\varepsilon}^{-1}(0)$. From (18), we obtain that C_{ε} is an invariant set.

Emphasize that the rank of H_{ε} is the rank of the matrix 1×2 , which is given by

$$J_{\mathbb{C}}H_{\varepsilon}(z,w) = \left(\frac{\partial H_{\varepsilon}}{\partial z}(z,w), \frac{\partial H_{\varepsilon}}{\partial w}(z,w)\right) = \left(-\varepsilon\eta(z), \frac{1}{\kappa(w)}\right).$$

Hence, $J_{\mathbb{C}}H_{\varepsilon}$ has maximal rank (rank 1) in each point of C_{ε} . By Proposition 1, we conclude that C_{ε} is a complex manifold of dimension 1.

Now, we shall prove item (a). Deriving the equation $G(w) = \varepsilon F(z)$, we get that $1 = \varepsilon k(w)\eta(z)$, which implies that $f(z,w) = \varepsilon g(z,w)$. Therefore, $C_{\varepsilon} \subset D_{\varepsilon} = \{(z,w) : f(z,w) = \varepsilon g(z,w)\}$ and $D_0 = C_0$.

Since $\frac{\partial H_{\varepsilon}}{\partial w}(z, w) \neq 0$ for all $(z, w) \in C_{\varepsilon}$, then from *Implicit Function Theorem*, there exist an open set V and a holomorphic function $w = h_{\varepsilon}(z)$ such that $H_{\varepsilon}(z, h_{\varepsilon}(z)) = 0$, for all $z \in V$. Then, expanding $h_{\varepsilon}(z)$ in Taylor series around $\varepsilon = 0$, we get that $h_{\varepsilon}(z) = \lambda(z) + \varepsilon Q(z, \varepsilon)$, for all $\varepsilon \in [0, \varepsilon_0]$ and $z \in V \cap U$, where Q is a holomorphic function and $C_0 = \{(z, w) : w = \lambda(z), z \in U\}$ is locally the critical manifold.

In what follow, we show that the Hausdorff distance between C_0 and C_{ε} is of order ε , that is $d_H(C_0, C_{\varepsilon}) = \mathcal{O}(\varepsilon)$. Recall that the Hausdorff distance d_H between nonempty subsets A and B is given by

$$d_H(A, B) = \max \left\{ \sup_{x \in A} \inf_{y \in B} d(x, y), \sup_{x \in B} \inf_{y \in A} d(x, y) \right\}.$$

Hence, $d_H(C_0, C_{\varepsilon}) = \mathcal{O}(\varepsilon)$ provided that the following statements hold:

- i. for each $p \in C_{\varepsilon}$ there exists $q \in C_0$ such that $d(p,q) = \mathcal{O}(\varepsilon)$;
- ii. for each $q \in C_0$ there exists $p \in C_{\varepsilon}$ such that $d(p,q) = \mathcal{O}(\varepsilon)$.

In the sequel, we shall verify item (i). Item (ii) can be verified analogously.

Consider the point $p = (z, w) \in C_{\varepsilon}$ and take $q = (z, \xi) \in C_0$. Then, $w = h_{\varepsilon}(z) = \lambda(z) + \varepsilon Q(z, \varepsilon)$ and $\xi = \lambda(z)$. Thus,

$$d(p,q) = ||(0,w-\xi)|| = ||(0,\varepsilon Q(z,\varepsilon))|| = |Q(z,\varepsilon)|\,\varepsilon \leq \varepsilon M,$$

where we have used that $|Q(z,\varepsilon)| \leq M$, for all $(z,\varepsilon) \in K_{V \cap U} \times [0,\varepsilon_0]$, with $K_{V \cap U}$ a compact subset of $V \cap U$. This implies that $d(p,q) = \mathcal{O}(\varepsilon)$.

Finally, we prove item (b). Since $H_{\varepsilon}(z,w)=0$ for all $(z,w)\in C_{\varepsilon}$, then $\frac{\partial H_{\varepsilon}}{\partial z}\dot{z}+\frac{\partial H_{\varepsilon}}{\partial w}\dot{w}=0$. Consequently, the dynamics over C_{ε} is determined by

$$\dot{w} = -\frac{\frac{\partial H_{\varepsilon}}{\partial z}\dot{z}}{\frac{\partial H_{\varepsilon}}{\partial w}} = -\frac{-\varepsilon\eta(z)\dot{z}}{\frac{1}{\kappa(w)}} = \varepsilon\underbrace{\eta(z)\kappa(w)}_{}\dot{z} = \frac{g(z,w)}{f(z,w)}f(z,w) = g(z,w).$$

Emphasize that it is possible to adapt the previous result for uncoupled slow-fast systems of the form:

(19)
$$\begin{cases} \varepsilon \dot{z} = f(z), \\ \dot{w} = g(w), \end{cases}$$

where f, g are holomorphic functions defined in Ω . Thus, the following result is a direct consequence of the above.

Corollary 6. Equilibrium points and poles of order n of the reduced problem associated to (19) are persistent by singular perturbation.

Example 7. Let n be natural number with $n \geq 2$. Consider the following system

(20)
$$\begin{cases} \varepsilon \dot{z} = z^n, \\ \dot{w} = g(w), \end{cases}$$

where g is one of the normal forms given in [4, Theorem 1.1]. Recall that $C_0 = \{(z, w) \in \mathbb{C}^2 : z = 0\}$ is the critical manifold associated with system (20), which is not normally hyperbolic. From Theorem C, there exists an invariant complex manifold C_{ε} given by:

- $C_{\varepsilon} = \{(z, w) : z^{n-1} \ln(w) = \varepsilon \frac{\eta}{-n+1} \}$, provided that $g(w) = \eta w$, with $\eta \in \mathbb{C}$.
- $C_{\varepsilon} = \{(z, w) : z^{n-1} = \varepsilon \frac{m-1}{n-1} w^{m-1} \}$, provided that $g(w) = w^m$, with m > 2.
- $C_{\varepsilon} = \{(z, w) : z^{n-1} = \varepsilon \frac{m+1}{-n+1} w^{-(m+1)}\}$, provided that $g(w) = \frac{1}{w^m}$, with $m \ge 2$.
- $C_{\varepsilon} = \left\{ (z, w) : z^{n-1} \left(\frac{w^{-m+1}}{-m+1} + \ln(w) \right) = \varepsilon \frac{\gamma}{-n+1} \right\}, \text{ provided that}$ $g(w) = \frac{\gamma w^m}{1 + w^{m-1}}, \text{ with } m \ge 2 \text{ and } \gamma \in \mathbb{C}.$

Moreover, in either case we have that C_{ε} converges to C_0 when ε tends to 0.

4.2. Coupled differential equations. Consider the system (SF3), where g, G are holomorphic functions defined in Ω such that G'(w) = 1/g(w), $\alpha, \beta \in \mathbb{C} \setminus \{0\}$. In this constext, the critical manifold of (SF3) is $C_0 = \{(z, w) : z = -\frac{\beta G(w)}{\alpha}\}$.

Let us introduce the equations for the orbits of system (SF3):

(21)
$$\varepsilon \frac{dz}{dw} = \frac{\alpha z + \beta G(w)}{g(w)}.$$

Since $\dot{w} = g(w)$, then G(w) = t. From the first equation of (SF3), we get the linear differential equation given by $\varepsilon \dot{z} = \alpha z + \beta t$. Recall that $z = -\frac{\beta}{\alpha^2}(\alpha t + \varepsilon)$ is a solution of previous equation. Thus, we obtain the equation $z = -\frac{\beta}{\alpha^2}(\alpha G(w) + \varepsilon)$, which satisfies the equation of orbits (21).

In the next result we prove that the set of points that satisfies the above equation is in fact a one-dimensional invariant complex manifold. Moreover, if we assume that the critical manifold is not normally hyperbolic, then the geometric singular perturbation theory does not determine the type of stability of the invariant manifold C_{ε} . However,

in the following theorem we are giving conditions to ensure when the invariant manifold C_{ε} is exponentially attractive (see Figure 3).




FIGURE 3. C_{ε} is exponentially attractive in a certain region \mathcal{R} , when C_0 is not normally hyperbolic.

Theorem D. Consider the slow-fast system (SF3). Then, there exists a one-dimensional invariant complex manifold C_{ε} associated with the differential system (SF3), such that

- (a) The flow on C_{ε} converges to the slow flow as $\varepsilon \to 0$.
- (b) The dynamics over C_{ε} is given by $\dot{w} = g(w)$.
- (c) If $\alpha = i\alpha_2$, with $\alpha_2 \in \mathbb{R} \setminus \{0\}$, then the orbit of system (21) with initial condition $z(w_0) = z_0$ stays exponentially close to the invariant manifold C_{ε} in the region $\mathcal{R} = \{(z, w) : \alpha_2 \Im(G(w) G(w_0)) \geq \eta > 0\}$, for some $\eta > 0$.

Proof. Consider the holomoprhic function $H_{\varepsilon}(z,w) = z + \frac{\beta}{\alpha^2}(\alpha G(w) + \varepsilon)$ and the set $C_{\varepsilon} = \{(z,w) : z = -\frac{\beta}{\alpha^2}(\alpha G(w) + \varepsilon)\}$. Recall that $C_{\varepsilon} = H_{\varepsilon}^{-1}(0)$. By construction C_{ε} is an invariant set.

Emphasize that the rank of H_{ε} is the rank of the matrix 1×2 , which is given by

$$J_{\mathbb{C}}H_{\varepsilon}(z,w) = \left(\frac{\partial H_{\varepsilon}}{\partial z}(z,w), \frac{\partial H_{\varepsilon}}{\partial w}(z,w)\right) = \left(1, \frac{\beta}{\alpha g(w)}\right).$$

Hence, $J_{\mathbb{C}}H_{\varepsilon}$ has maximal rank (rank 1) in each point of C_{ε} . By Proposition 1, we conclude that C_{ε} is a complex manifold of dimension 1.

Now, we shall prove item (a). For that, it is enough to show that the Hausdorff distance between C_0 and C_{ε} is of order ε or equivalently that the following statements hold:

- i. for each $p \in C_{\varepsilon}$ there exists $q \in C_0$ such that $d(p,q) = \mathcal{O}(\varepsilon)$;
- ii. for each $q \in C_0$ there exists $p \in C_{\varepsilon}$ such that $d(p,q) = \mathcal{O}(\varepsilon)$.

In the sequel, we shall verify item (i). Item (ii) can be verified analogously.

Consider the point $p = (z, w) \in C_{\varepsilon}$ and take $q = (\xi, w) \in C_0$. Then, $z = -\frac{\beta}{\alpha^2}(\alpha G(w) + \varepsilon)$ and $\xi = -\frac{\beta}{\alpha}G(w)$. Thus,

$$d(p,q) = ||(z - \xi, 0)|| = \left| \left| \left(-\frac{\beta}{\alpha^2} \varepsilon, 0 \right) \right| \right| = \left| \frac{\beta}{\alpha^2} \right| \varepsilon.$$

This implies that $d(p,q) = \mathcal{O}(\varepsilon)$.

Next, we prove item (b). Since $H_{\varepsilon}(z,w)=0$ for all $(z,w)\in C_{\varepsilon}$, then $\frac{\partial H_{\varepsilon}}{\partial z}\dot{z}+\frac{\partial H_{\varepsilon}}{\partial w}\dot{w}=0$. Consequently, the dynamics over C_{ε} is determined by

$$\dot{w} = -\frac{\frac{\partial H_{\varepsilon}}{\partial z}\dot{z}}{\frac{\partial H_{\varepsilon}}{\partial w}}\dot{z} = -\frac{\dot{z}}{\frac{\beta}{\alpha g(w)}} = -\frac{\alpha z + \beta G(w)}{\frac{\varepsilon \beta}{\alpha g(w)}} = -\frac{-\frac{\varepsilon \beta}{\alpha}}{\frac{\varepsilon \beta}{\alpha g(w)}} = g(w).$$

Finally, we prove item (c). Define $h_{\varepsilon}(w) = -\frac{\beta}{\alpha^2}(\alpha G(w) + \varepsilon)$. Hence, the invariant manifold is given by $C_{\varepsilon} = \{(z, w) : z = h_{\varepsilon}(w)\}$.

Now, we perform the change of variables $v = z - h_{\varepsilon}(w)$ in equation (21) obtaining:

(22)
$$\varepsilon \frac{dv}{dw} = \frac{i\alpha_2}{q(w)}v.$$

Recall that the solution of (22) with initial condition $v(w_0) = z(w_0) - h_{\varepsilon}(w_0)$ can be written as $v(w) = v(w_0)e^{\frac{i\alpha_2(G(w) - G(w_0))}{\varepsilon}}$. Thus,

$$|v(w)| = |v(w_0)|e^{-\frac{\alpha_2\Im(G(w) - G(w_0))}{\varepsilon}} \le |v(w_0)|e^{-\frac{\eta}{\varepsilon}}.$$

Since $\eta > 0$, then any solution gets exponentially closer to the invariant manifold C_{ε} .

Example 8. Consider the system defined as

(23)
$$\begin{cases} \varepsilon \dot{z} = iz + w^2, \\ \dot{w} = \frac{1}{2w}, \end{cases}$$

with $(z_0, w_0) = (\varepsilon + i, 1)$. Notice that $G(w) = w^2$ and $\alpha = i$. Since G'(w) = 2w, from Theorem D there exists a one-dimensional invariant complex manifold C_{ε} associated with the differential system (23), which is exponentially attractive on $\mathcal{R} = \{(z, w) \in \mathbb{C}^2 : \operatorname{Re}(w) \operatorname{Im}(w) > 0\}$.

18

5. Fenichel Manifold Approximation

This section is focused on using Laurent series and Fenichel's Theorem to approximate complex manifolds and find smooth manifolds without complex structure of the slow-fast system (SF4). In particular, we are interested in studying the dynamics of the system (SF4).

5.1. Non-existence of smooth manifolds with complex structure. Below we present 2 families of slow-fast systems that, despite having associated smooth locally invariant manifolds, have no complex structure.

5.1.1. Linear- w^n case. Consider the following system

(24)
$$\begin{cases} \varepsilon \dot{z} = \alpha z + \beta w, \\ \dot{w} = w^n, \end{cases}$$

where $\alpha, \beta \in \mathbb{C} \setminus \{0\}$ and $n \geq 2$. Recall that the critical manifold C_0 associated with system (24) is normally hyperbolic provided that $\operatorname{Re}(\alpha) = \alpha_1 \neq 0$. Let S_0 be a compact subset of C_0 .

We now apply the *Fenichel's Theorem* to the equivalent \mathbb{C}^{∞} four-dimensional real system associated with the system (24), then there exists a smooth locally invariant manifold $S_{\varepsilon} = \{(z, w) : z = h_{\varepsilon}(w)\}$ of the slow-fast system (24), which is diffeomorphic to S_0 .

Emphasize that this does not guarantee that the complex function $h_{\varepsilon}(z)$ is a holomorphic function. Indeed, suppose that there exists an invariant manifold $C_{\varepsilon} = \{(z, w) : z = h_{\varepsilon}(w)\}$, where

(25)
$$h_{\varepsilon}(w) = a_0 + \dots + a_n w^n + \dots + a_{2n-1} w^{2n-1} + \mathcal{O}(w^{2n}).$$

Since $\dot{z} = h'_{\varepsilon}(w)\dot{w}$ then we obtain the following equation

(26)
$$\alpha h_{\varepsilon}(w) + \beta w = h'_{\varepsilon}(w)\varepsilon w^{n}.$$

Substituting (25) in (26), we get that $a_i = 0$, for all $i \neq kn - (k-1)$, $a_1 = -\frac{\beta}{\alpha}$, $a_n = -\frac{\beta\varepsilon}{\alpha^2}$, and $a_{kn-(k-1)} = -\frac{\beta^k\varepsilon^k\prod_{j=2}^k[(j-1)n-(j-2)]}{\alpha^{k+1}}$, with $k \geq 2$. Therefore,

$$C_{\varepsilon} = \left\{ (z, w) : z = -\frac{\beta}{\alpha} w - \frac{\beta \varepsilon}{\alpha^2} w^n + \sum_{k=2}^{\infty} a_{kn-(k-1)} w^{kn-(k-1)} \right\}.$$

Notice that the series $\sum_{k=2}^{\infty} a_{kn-(k-1)} w^{kn-(k-1)}$ diverges for $w \neq 0$, which contradicts the existence of the holomorphic function h_{ε} .

5.1.2. $Linear-\frac{\gamma w^n}{1+w^{n-1}}$ case. Consider the following system

(27)
$$\begin{cases} \varepsilon \dot{z} = \alpha z + \beta w, \\ \dot{w} = \frac{\gamma w^n}{1 + w^{n-1}}, \end{cases}$$

where $\alpha, \beta \in \mathbb{C} \setminus \{0\}$ and $n \geq 2$. Recall that the critical manifold C_0 associated with system (27) is normally hyperbolic provided that $\operatorname{Re}(\alpha) = \alpha_1 \neq 0$. Take S_0 a compact subset of C_0 .

We now apply the *Fenichel's Theorem* to the equivalent \mathbb{C}^{∞} fourdimensional real system associated with the system (27), then there exists a smooth locally invariant manifold $S_{\varepsilon} = \{(z, w) : z = h_{\varepsilon}(w)\}$ of the slow-fast system (27), which is diffeomorphic to S_0 .

Emphasize that this does not guarantee that the complex function $h_{\varepsilon}(z)$ is a holomorphic function. Indeed, suppose that there exists an invariant manifold $C_{\varepsilon} = \{(z, w) : z = h_{\varepsilon}(w)\}$, where

(28)
$$h_{\varepsilon}(w) = a_0 + \dots + a_n w^n + \dots + a_{2n-1} w^{2n-1} + \mathcal{O}(w^{2n}).$$

Since $\dot{z} = h'_{\varepsilon}(w)\dot{w}$, then

(29)
$$\alpha h_{\varepsilon}(w) + \beta w = h'_{\varepsilon}(w) \varepsilon \frac{\gamma w^n}{1 + w^{n-1}}.$$

Substituting (28) in (29), we get that $a_i=0$, for all $i\neq kn-(k-1)$, $a_1=-\frac{\beta}{\alpha},\ a_n=-\frac{\beta\gamma\varepsilon}{\alpha^2}$, and $a_{kn-(k-1)}=-\frac{\beta\gamma\varepsilon\prod_{j=2}^k[\alpha-((j-1)n-(j-2))\gamma\varepsilon]}{\alpha^{k+1}}$, with $k\geq 2$. Therefore,

$$C_{\varepsilon} = \left\{ (z, w) : z = -\frac{\beta}{\alpha} w - \frac{\beta \gamma \varepsilon}{\alpha^2} w^n + \sum_{k=2}^{\infty} a_{kn-(k-1)} w^{kn-(k-1)} \right\}.$$

Notice that the series $\sum_{k=2}^{\infty} a_{kn-(k-1)} w^{kn-(k-1)}$ diverges for $w \neq 0$, which contradicts the existence of the holomorphic function h_{ε} .

The following result is a direct consequence of [4, Theorem 1.1] and of cases 5.1.1 and 5.1.2.

Proposition 9. Consider system (SF1) such that g has a zero of order n > 1 and f is a linear holomorphic function, then system (SF1) has no complex manifolds of form $C_{\varepsilon} = \{(z, w) : z = h_{\varepsilon}(w)\}.$

5.2. Approximation of complex invariant manifold. Here we present 2 families of slow-fast systems that, in the case of having associated locally invariant smooth manifolds with complex structure, can be approximated via Laurent series and also the singularity associated with the reduced problem is preserved by singular perturbation.

5.2.1. Linear-linear case. Consider the following system

(30)
$$\begin{cases} \varepsilon \dot{z} = az + bw, \\ \dot{w} = cz + dw, \end{cases}$$

20

where $a \in \mathbb{C} \setminus \{0\}$ and $b, c, d \in \mathbb{C}$. Assume that the critical manifold C_0 associated with system (30) is normally hyperbolic, that is, $\operatorname{Re}(a) \neq 0$.

Notice that the reduced problem associated with system (30) is given by

(31)
$$\begin{cases} 0 = az + bw, \\ \dot{w} = \left(\frac{ad - bc}{a}\right)w =: \widetilde{g}(w). \end{cases}$$

Thus, the equilibrium point is $w_0 = 0$ and $\tilde{g}'(w_0) = \frac{ad-bc}{a} = \alpha + i\beta$, where $\alpha = \text{Re}\left(\frac{ad-bc}{a}\right)$ and $\beta = \text{Im}\left(\frac{ad-bc}{a}\right)$. In addition, the Jacobian matrix at the equilibrium point of system (31) is

$$J_{\mathbb{R}}\widetilde{g}|_{w_0} = \begin{pmatrix} \alpha & -\beta \\ \beta & \alpha \end{pmatrix}.$$

The determinant D and the trace T are

$$D = \alpha^2 + \beta^2$$
 and $T = 2\alpha$.

- $\beta \neq 0$. We have that if $\alpha > 0$ then the origin is a repelling focus of the reduced problem (31), if $\alpha < 0$ then the origin is an attracting focus of (31).
- $\beta = 0$. We have that if $\alpha > 0$ then the origin is a repelling node of the reduced problem (31) and if $\alpha < 0$ then the origin is an attracting node of (31).

Therefore, we get the following table:

α	j^u	j^s
+	2	0
	0	2

where j^u, j^s have been defined in Theorem 2. In this case, the eigenvalues of the Jacobian matrix (9) are $\lambda_{\pm} = \text{Re}(a) \pm i \text{Im}(a)$ Thus, we have the following table:

$\operatorname{Re}\left(a\right)$	k^u	k^s
+	2	0
_	0	2

where k^u , k^s have been defined in Theorem 2. Hence, if α , Re (a) > 0, then $j^u + k^u = 4$ and $j^s + k^s = 0$. if α , Re (a) < 0, then $j^u + k^u = 0$ and

 $j^s + k^s = 4$. If $sign(\alpha) \neq sign(\text{Re}(a))$, then $j^u + k^u = 2$ and $j^s + k^s = 2$. From item (ii) of Theorem 2:

- If α , Re (a) > 0, then the equilibrium point $q_{\varepsilon} = (0,0)$ of system (30) is a global repelling point.
- If α , Re (a) < 0, then the equilibrium point $q_{\varepsilon} = (0,0)$ of system (30) is a global attracting point.
- If $sign(\alpha) \neq sign(\text{Re}(a))$, then the equilibrium point $q_{\varepsilon} = (0,0)$ of system (30) is a saddle point.

Notice that when $\alpha = 0$ we can not use item (ii) of Theorem 2, however, C_0 is normally hyperbolic, thus if we take a compact subset S_0 of C_0 , we can apply the *Fenichel's Theorem* to the equivalent \mathbb{C}^{∞} four-dimensional real system associated with the system (30), then there exists a smooth locally invariant manifold $S_{\varepsilon} = \{(z, w) : z = h_{\varepsilon}(w)\}$ of the slow-fast system (30), which is diffeomorphic to S_0 .

Although this does not guarantee that the complex function $h_{\varepsilon}(z)$ is a holomorphic function, we would like to know the dynamics on the Fenichel manifold if it exists. Hence, suppose that $C_{\varepsilon} = \{(z, w) : z = h_{\varepsilon}(w)\}$, where $h_{\varepsilon}(w) = \lambda_0 + \lambda_1 w + \mathcal{O}(w^2)$. Since $\dot{z} = h'_{\varepsilon}(w)\dot{w}$, then

$$ah_{\varepsilon}(w) + bw = h'_{\varepsilon}(w)\varepsilon((\alpha + i\beta)w).$$

This implies that $\lambda_i = 0$, for all $i \neq 1$ and $\lambda_1 = \frac{-b}{a - \varepsilon(\alpha + i\beta)}$. Therefore,

$$C_{\varepsilon} = \left\{ (z, w) : z = \frac{-b}{a - \varepsilon(\alpha + i\beta)} w \right\}.$$

In particular, if $\alpha = 0$, then

$$C_{\varepsilon}|_{\alpha=0} = \left\{ (z, w) : z = \frac{-b}{a - i\varepsilon\beta} w \right\}.$$

Recall that the dynamics over the manifold $C_{\varepsilon}|_{\alpha=0}$ is given by the following differential equation

$$\dot{w} = i\beta w$$
.

Therefore, if w_0 is a center of reduced problem (31), then w_{ε} is a center of system (30) such that $w_{\varepsilon} \to w_0$ when $\varepsilon \to 0$ (see Figure 4).

The following result is a direct consequence of [4, Theorem 1.1] and of case 5.2.1.

Proposition 10. Consider system (SF1) such that g has a zero w_0 of order one and f is a linear holomorphic function. If system (SF1) admits an invariant manifold with complex structure, then centers of the reduced problem associated to (SF1) are persistent by singular perturbation.




FIGURE 4. Persistence of a center of the reduced problem by singular perturbation.

5.2.2. Linear-pole case. Consider the following system

(32)
$$\begin{cases} \varepsilon \dot{z} = \alpha z + \beta w, \\ \dot{w} = \frac{1}{w^n}, \end{cases}$$

where $\alpha, \beta \in \mathbb{C} \setminus \{0\}$ and $n \geq 1$. Recall that the critical manifold C_0 associated with system (32) is normally hyperbolic provided that $\operatorname{Re}(\alpha) = \alpha_1 \neq 0$. Let S_0 be a compact subset of C_0 .

We now apply the *Fenichel's Theorem* to the equivalent \mathbb{C}^{∞} fourdimensional real system associated with the system (32), then there exists a smooth locally invariant manifold $S_{\varepsilon} = \{(z, w) : z = h_{\varepsilon}(w)\}$ of the slow-fast system (32), which is diffeomorphic to S_0 .

Although this does not guarantee that the complex function $h_{\varepsilon}(z)$ is a holomorphic function, we would like to know the dynamics on the Fenichel manifold if it exists. Hence, suppose that there exists an invariant manifold $C_{\varepsilon} = \{(z, w) : z = h_{\varepsilon}(w)\}$, where

(33)
$$h_{\varepsilon}(w) = a_0 + \dots + a_{n+2}w^{n+2} + \dots + a_{2n+3}w^{2n+3} + \dots + \mathcal{O}(w^{3n+4}).$$

Since $\dot{z} = h'_{\varepsilon}(w)\dot{w}$, then we get the following equation

(34)
$$\alpha h_{\varepsilon}(w) + \beta w = h'_{\varepsilon}(w) \varepsilon w^{-n}.$$

Substituting (33) in (34), we get that $a_i = 0$, for all $i \neq kn + (k+1)$ and $a_{kn+(k+1)} = \frac{\alpha^{k-1}\beta}{\varepsilon^k \prod_{j=1}^k [jn+(j+1)]}$, with $k \geq 1$. Therefore,

$$C_{\varepsilon} = \left\{ (z, w) : z = \sum_{k=1}^{\infty} a_{kn+(k+1)} w^{kn+(k+1)} \right\}.$$

It is easy to see that the series $\sum_{k=1}^{\infty} a_{kn+(k+1)} w^{kn+(k+1)}$ converges for all $w \in \mathbb{C}$. Recall that the dynamics over the manifold C_{ε} is given by the following differential equation

$$\dot{w} = \frac{\beta w + \frac{\alpha\beta}{(n+2)\varepsilon} w^{n+2} + \frac{\alpha^2\beta}{(n+2)(2n+3)\varepsilon^2} w^{2n+3} + \mathcal{O}(1/\varepsilon^3, w^{3n+4})}{\frac{\beta}{\varepsilon} w^{n+1} + \frac{\alpha\beta}{(n+2)\varepsilon^2} w^{2n+2} + \mathcal{O}(1/\varepsilon^3, w^{3n+3})} =: F_{\varepsilon}(w).$$

Thus, expanding F_{ε} around $w_0 = 0$ we get that

$$F_{\varepsilon}(w) = \frac{\varepsilon}{w^n} + \mathcal{O}(1/\varepsilon, w^{n+2}).$$

Therefore, w_0 is a pole of order n of F_{ε} (see Figure 5). From [4, Theorem 1.1], we conclude that F_{ε} and $G \equiv \frac{1}{w^n}$ are 0-conformally conjugated.




FIGURE 5. Persistence of a pole of the reduced problem by singular perturbation.

The following result is a direct consequence of [4, Theorem 1.1] and of case 5.2.2.

Proposition 11. Consider system (SF1) such that g has a pole w_0 of order $n \geq 1$ and f is a linear holomorphic function. If system (SF1) admits an invariant manifold with complex structure, then poles of order n of the reduced problem associated to (SF1) are persistent by singular perturbation.

6. Statements and Declarations

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

7. Acknowledgements

This article was possible thanks to the scholarship granted from the Brazilian Federal Agency for Support and Evaluation of Graduate Education (CAPES), in the scope of the Program CAPES-Print, process number 88887.310463/2018-00, International Cooperation Project number 88881.310741/2018-01. Paulo Ricardo da Silva is also partially supported by São Paulo Research Foundation (FAPESP) grant 2019/10269-3 and CNPq grant 302154/2022-1.

Gabriel Rondón is supported by São Paulo Research Foundation (FAPESP) grants 2020/06708-9 and 2022/12123-9. Luiz Fernando Gouveia is supported by São Paulo Research Foundation (FAPESP) grant 2020/04717-0.

References

- [1] B. J. Briot, C. Recherches sur les propriétés des fonctions définies par des équations différentielles. J. Écol. Imp. Poly., 21:133–197, 1856.
- [2] J. B. Conway. Functions of one complex variable, volume 11 of Graduate Texts in Mathematics. Springer-Verlag, New York, second edition, 1978.
- [3] N. Fenichel. Geometric singular perturbation theory for ordinary differential equations. J. Differential Equations, 31(1):53–98, 1979.
- [4] A. Garijo, A. Gasull, and X. Jarque. Normal forms for singularities of one dimensional holomorphic vector fields. *Electron. J. Differential Equations*, pages No. 122, 7, 2004.
- [5] J. D. Gray and S. A. Morris. When is a function that satisfies the Cauchy-Riemann equations analytic? *Amer. Math. Monthly*, 85(4):246–256, 1978.
- [6] E. Hille. Ordinary differential equations in the complex domain. Pure and Applied Mathematics. Wiley-Interscience [John Wiley & Sons], New York-London-Sydney, 1976.
- [7] D. Huybrechts. *Complex geometry*. Universitext. Springer-Verlag, Berlin, 2005. An introduction.
- [8] K. Iwasaki, H. Kimura, S. Shimomura, and M. Yoshida. From Gauss to Painlevé. Aspects of Mathematics, E16. Friedr. Vieweg & Sohn, Braunschweig, 1991. A modern theory of special functions.
- [9] C. K. R. T. Jones. Geometric singular perturbation theory. In *Dynamical systems (Montecatini Terme, 1994)*, volume 1609 of *Lecture Notes in Math.*, pages 44–118. Springer, Berlin, 1995.
- [10] J. Llibre, P. R. da Silva, and M. A. Teixeira. Regularization of discontinuous vector fields on \mathbb{R}^3 via singular perturbation. *J. Dynam. Differential Equations*, 19(2):309–331, 2007.
- [11] I. M. Moroz. Some complex differential equations arising in telecommunications. In *Singularity theory and its applications*, *Part II (Coventry*, 1988/1989), volume 1463 of *Lecture Notes in Math.*, pages 278–293. Springer, Berlin, 1991.
- [12] I. M. Moroz. Bifurcations in a class of complex differential equations. *European J. Appl. Math.*, 3(3):273–282, 1992.

- [13] D. J. Needham. A centre theorem for two-dimensional complex holomorphic systems and its generalization. *Proc. Roy. Soc. London Ser. A*, 450(1939):225–232, 1995.
- [14] D. J. Needham and S. McAllister. Centre families in two-dimensional complex holomorphic dynamical systems. R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci., 454(1976):2267–2278, 1998.
- [15] V. Scheidemann. Introduction to complex analysis in several variables. Birkhäuser Verlag, Basel, 2005.
- [16] P. Szmolyan. Transversal heteroclinic and homoclinic orbits in singular perturbation problems. *J. Differential Equations*, 92(2):252–281, 1991.
- [17] M. A. Teixeira and P. R. da Silva. Regularization and singular perturbation techniques for non-smooth systems. *Phys. D*, 241(22):1948–1955, 2012.
- [18] R. O. Wells, Jr. Differential and complex geometry: origins, abstractions and embeddings. Springer, Cham, 2017.

SÃO PAULO STATE UNIVERSITY (UNESP), INSTITUTE OF BIOSCIENCES, HUMANITIES AND EXACT SCIENCES. RUA C. COLOMBO, 2265, CEP 15054–000. S. J. RIO PRETO, SÃO PAULO, BRAZIL.

Email address: gabriel.rondon@unesp.br Email address: paulo.r.silva@unesp.br Email address: fernando.gouveia@unesp.br