AFFINIZATIONS, R-MATRICES AND REFLECTION FUNCTORS

MASAKI KASHIWARA, MYUNGHO KIM, SE-JIN OH, AND EUIYONG PARK

ABSTRACT. In this paper we establish affinizations and R-matrices in the language of pro-objects, and as an application, we construct reflection functors over the localizations of quiver Hecke algebras of arbitrary finite types. This reflection functor categorifies the braid group action on the half of a quantum group and the Saito reflection.

Contents

1.	Introduction	3
2.	Preliminaries	12
2.1.	. Module objects	12
2.2.	. Graded categories	13
2.3.	. Pro-objects	14
3.	Affinizations in abelian categories	15

The research of S.-j. Oh was supported by the Ministry of Education of the Republic of Korea and the National Research Foundation of Korea (NRF-2022R1A2C1004045).

The research of E. Park was supported by the National Research Foundation of Korea(NRF) grant funded by the Korea government (MSIT) (RS-2023-00273425 and NRF-2020R1A5A1016126).

Date: February 2, 2024.

²⁰²⁰ Mathematics Subject Classification. 18M05, 16D90, 81R10.

Key words and phrases. Affinization, Localization, Reflection, R-matrices, Quiver Hecke algebra. The research of M. Kashiwara was supported by Grant-in-Aid for Scientific Research (B) 20H01795, Japan Society for the Promotion of Science.

The research of M. Kim was supported by the National Research Foundation of Korea (NRF) Grant funded by the Korea government(MSIP) (NRF-2022R1F1A1076214 and NRF-2020R1A5A1016126).

3.1. Coherent objects in $\operatorname{Pro}(\mathcal{C})$	15
3.2. Flatness	24
4. Duality	27
4.1. Complements	27
4.2. Duality	28
4.3. Affine objects	30
5. Coherent objects in Monoidal categories	34
5.1. Graded monoidal categories	34
5.2. Rigid case	39
6. R-matrices	41
6.1. Affine objects	41
6.2. Rational centers and affinizations	43
6.3. Quasi-rigid Axiom	49
7. Quiver Hecke algebras and Schur-Weyl duality functors	56
7.1. Quiver Hecke algebras	56
7.2. Universal R-matrices	58
7.3. Canonical affinizations in R_{λ} -gmod	60
7.4. Schur-Weyl duality	61
8. Localizations of R -gmod	66
8.1. Categories \mathscr{C}_w and \mathscr{C}_w^*	66
8.2. Localizations	67
9. Resultant algebra and Affinization of Invariants	72
9.1. Resultant algebra	72
9.2. Affinizations of invariants for R -modules	74
9.3. Examples of affinizations	80
10. Saito Reflection functors for R -gmod via localization	80
10.1. $\mathscr{C}_{s_iw_0}$ and $\mathscr{C}^*_{s_iw_0}$	80
10.2. Schur-Weyl datum	82
11. Examples of \mathscr{F}_i	99
11.1. $\mathscr{F}_i \colon R\operatorname{-gmod} \to \widetilde{\mathscr{C}}^*_{s_i w_0}$	99

11.2.
$$\mathscr{F}_i \colon \mathscr{C}_{w_0} \to \mathscr{C}_{w_0}$$

References

1. INTRODUCTION

Affinizations and *R*-matrices are one of the most powerful tools in the representation theory of quiver Hecke algebras and affine quantum groups. One of the most successful applications of them is the proof of the simplicity of the head of the tensor product of simple objects given in [6] (cf. Proposition 6.19).

R-matrices are distinguished homomorphisms between tensor products of modules, which measure the commutativity of tensor products. Affinizations of modules help R-matrices to play their role. The study on affinizations and R-matrices gives rise to the integer invariants Λ , $\tilde{\Lambda}$ and \mathfrak{d} , which have been used crucially in deriving several remarkable results including monoidal categorification of cluster algebras (see [6, 8, 9, 13] and references therein). In the representation theory of quantum affine algebras, R-matrices already occupy an important position. The generalized Schur-Weyl duality functor relates the R-matrices in quiver Hecke algebras and the ones in quantum affine algebras in a natural way, and they enjoy very similar properties in their own categories. See also [1, 4, 5, 7, 10] and references therein.

In the case of symmetric quiver Hecke algebras, there exists a functorial construction of an affinization of a module ([7]). Let R be a symmetric quiver Hecke algebra over a base field \mathbf{k} and M an R-module. As a \mathbf{k} -vector space, the affinization M of Mis isomorphic to $\mathbf{k}[z_{\mathsf{M}}] \otimes M$, and the action of the generators $\mathbf{e}(\nu)$ and τ_l on M is the same as those on M, but the action of x_k is twisted by z_{M} (see [7, Section 1.3]). The indeterminate z_{M} can be understood as a monomorphism in End(M). Once an affinization exists, one can construct the distinguished homomorphisms

$$\mathbf{R}_{\mathsf{M},\mathsf{N}}^{\mathrm{ren}} := (z_{\mathsf{M}} - z_{\mathsf{N}})^{-s} \mathbf{R}_{\mathsf{M},\mathsf{N}}^{\mathrm{uni}} \in \mathrm{Hom}(\mathsf{M} \circ \mathsf{N}, \mathsf{N} \circ \mathsf{M}),$$
$$\mathbf{r}_{M,N} := \mathbf{R}_{\mathsf{M},\mathsf{N}}^{\mathrm{ren}}|_{z_{\mathsf{M}} = z_{\mathsf{N}} = 0} \in \mathrm{Hom}(M \circ N, N \circ M),$$

where $\mathbb{R}_{M,N}^{\text{uni}}$ is the *universal* R-matrix defined by the intertwiners φ_i (see Section 7.2) and s is the largest integer such that $\mathbb{R}_{M,N}^{\text{uni}}(\mathsf{M} \circ \mathsf{N}) \subset (z_{\mathsf{M}} - z_{\mathsf{N}})^s \mathsf{N} \circ \mathsf{M}$. The *renormalized* R-matrix $\mathbb{R}_{M,N}^{\text{ren}}$ commutes with z_{M} and z_{N} , and the R-matrix $\mathbf{r}_{M,N}$ never vanishes.

Unlike the case of symmetric quiver Hecke algebras, the existence of affinizations over a non-symmetric quiver Hecke algebra is not guaranteed, which causes a big difficulty in the study of the representations over non-symmetric quiver Hecke algebras such as monoidal categorification in the non-symmetric case. In [13], the notion of affinizations and R-matrices for arbitrary quiver Hecke algebras was studied. It was shown that, if an affinization exists, the R-matrices can be constructed, and they enjoy very similar properties to those in the symmetric case. Affinizations and renormalized R-matrices allow us to consider a duality datum (see [13, Section 4.1] for the definition) over the quiver Hecke algebra setting, which provides the generalized Schur-Weyl duality between quiver Hecke algebras ([13, Section 4]). This duality can be understood as an analogue of the generalized Schur-Weyl duality from quiver Hecke algebras to quantum affine algebras introduced in [7, Section 3].

In the case of the *localization* of module categories over quiver Hecke algebras, the theory of affinization becomes more complicated. Let R-gmod be the category of finitedimensional graded modules over an *arbitrary* quiver Hecke algebra, and let W be the Weyl group. For any element $w \in W$, let \mathscr{C}_w be the full subcategory of R-gmod generated by the dual PBW vectors corresponding to w (see [8, 9]). Note that the subcategory \mathscr{C}_w categorifies the quantum unipotent coordinate ring $A_q(\mathfrak{n}(w))$ and \mathscr{C}_w gives a monoidal categorification of $A_q(\mathfrak{n}(w))$ as a quantum cluster algebra ([8]). It was shown in [11] that \mathscr{C}_w admits a localization $\widetilde{\mathscr{C}}_w$ via a *real commuting family* of central objects. In the language of cluster algebras, the central objects correspond to the frozen variables and the localization $\widetilde{\mathscr{C}}_w$ categorifies the quantum cluster algebra $A_q(\mathfrak{n}(w))$ at the frozen variables. It turns out that the localization $\widetilde{\mathscr{C}}_w$ is a rigid monoidal category ([11, 12]).

The categorical structure of the localization $\widetilde{\mathscr{C}}_w$ is much more complicated than the original category \mathscr{C}_w . Let $\operatorname{Mod}_{\operatorname{coh}}(R)$ be the category of finitely generated *R*-modules, and let $\mathscr{C}_w^{\operatorname{big}}$ be the full subcategory of $\operatorname{Mod}_{\operatorname{coh}}(R)$ consisting of modules whose simple

subquotients belong to \mathscr{C}_w . Then, we can localize $\mathscr{C}_w^{\text{big}}$ similarly to \mathscr{C}_w and obtain the localization $\widetilde{\mathscr{C}}_w^{\text{big}}$. The problem is that the dual of an affinization of a simple module in \mathscr{C}_w does not belong to $\widetilde{\mathscr{C}}_w^{\text{big}}$ in general. Thus it is not valid to translate directly the classical notion of affinizations and R-matrices to the localization $\widetilde{\mathscr{C}}_w$. In order to overcome this difficulty, we have to introduce the category $\operatorname{Pro}(\widetilde{\mathscr{C}}_w)$ of pro-objects of $\widetilde{\mathscr{C}}_w$. Then the dual of an affinization belongs to $\operatorname{Pro}(\widetilde{\mathscr{C}}_w)$. This categorical approach to affinizations and R-matrices may lead us to new applications in various interesting categories including a new kind of generalized Schur-Weyl duality.

In this paper, we establish the theory of affinizations and R-matrices in the language of pro-objects, and as an application, we construct *reflection functors* over the localizations of quiver Hecke algebras of an *arbitrary finite type*. The main results of the paper can be summarized as follows:

- (i) The first main result is to study the notion of affinization and R-matrices in the language of pro-objects. We define affinizations and R-matrices in the categories with certain conditions, and investigate their properties. We then introduce the invariants D, Λ and δ and prove various properties of them. Thus, in a purely categorical setting, we have recovered key properties appeared in quiver Hecke algebras and quantum affine algebras. We next introduce a *duality datum* at the category level, and construct a generalized Schur-Weyl duality in the category setting by applying the argument given in [7].
- (ii) The second main result is to apply the generalized Schur-Weyl duality to the localized category $\widetilde{\mathscr{C}}^*_{s_iw_0}$ of a quiver Hecke algebra of *arbitrary finite type*. As a result, for any $i \in I$, we obtain an equivalence of monoidal categories

$$\mathscr{S}_i \colon \widetilde{\mathscr{C}}_{s_i w_0} \xrightarrow{\sim} \widetilde{\mathscr{C}}^*_{s_i w_0},$$

which categorifies the braid group action ([20, Chapter 37]) and the *Saito reflec*tion σ_i ([24]).

In the construction, the *affinizations* (or the *lifts*) of the invariants Λ , \mathfrak{d} , $\widetilde{\Lambda}$, etc. take a crucial role (see Section 9).

We conjecture that $\mathscr{S}_i : \widetilde{\mathscr{C}}_{s_i w_0} \xrightarrow{\sim} \widetilde{\mathscr{C}}^*_{s_i w_0}$ induces an equivalence of categories $\mathscr{C}_{s_i w_0} \xrightarrow{\sim} \mathscr{C}^*_{s_i w_0}$ at the level of module categories (see Conjecture 10.12).

In the case of finite ADE type, S. Kato ([15]) constructed the Saito reflection functor using geometry, which categorifies the braid group action on the half of a quantum group (see also [16, 22]). We conjecture that \mathscr{S}_i coincides with the functor induced by the functor constructed by S. Kato. However, in the case of *non-symmetric type*, there is no relevant result as far as the authors know.

Let us explain the main results of the paper in more details.

The first main result is about the notion of affinization and R-matrices in a general category. We shall explain it in three parts, i.e., affinization in categories, affinizations with respect to $\mathbf{k}[z]$ and R-matrices, and generalized Schur-Weyl duality.

Affinization in categories. Let \mathcal{C} be a k-linear category satisfying (3.1) and let $A := \bigoplus_{k \in \mathbb{Z}} A_k$ be a commutative graded k-algebra satisfying (3.2). We first define the subcategory $\operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$ consisting of graded *coherent* A-modules in the category $\operatorname{Pro}_{\operatorname{coh}}(\mathcal{C})$ of all pro-objects of \mathcal{C} (see Definition 3.1 for the precise definition). We investigate the category $\operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$ and obtain several homological properties under the condition when objects are A-flat (see Section 3). We denote by $\operatorname{Aff}_A(\mathcal{C})$ the full subcategory of $\operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$ consisting of A-flat objects (Definition 4.6). For a given anti-equivalence D of categories, we show that D can be lifted to the category $\operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$, denoted by D_A . Then, the functor D_A gives an equivalence of categories between $\operatorname{Aff}_A(\mathcal{C})$ and its opposite category (see Section 4.2). In natural and interesting situations, D can be understood as a duality functor in the category \mathcal{C} .

Affinizations in monoidal categories are also studied in details. We assume further that \mathcal{C} is an abelian **k**-linear graded monoidal category satisfying (5.1). For graded A-modules M and N in $\operatorname{Pro}(\mathcal{C})$, we define a tensor product $\mathsf{M} \otimes_A \mathsf{N} \in \operatorname{Modg}(A, \operatorname{Pro}(\mathcal{C}))$ by the universal property (5.2). Lemma 5.4 says that the category $\operatorname{Aff}_A(\mathcal{C})$ with \otimes_A forms a monoidal category, and Proposition 5.6 tells us that $\operatorname{Aff}_A(\mathcal{C})$ is rigid if \mathcal{C} is a rigid monoidal category. Affinizations with respect to $\mathbf{k}[z]$ and R-matrices. In the case where $A = \mathbf{k}[z]$, we take one more step further. This case exactly appears in the quiver Hecke algebras. We assume that C is a k-linear category satisfying (3.1) and $A = \mathbf{k}[z]$, where z is an indeterminate. We define an *affine object* in C as a pair (M, z) belonging to $\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z], C)$ with the conditions (a)–(d) in Definition 4.7. We then study properties of affine objects. Let $\operatorname{Aff}_{z}(C)$ denote the category of affine objects, and $\operatorname{Raff}_{z}(C)$ the category of $\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z], C)$ modified by making the morphism z invertible (see Definition 4.12 for the precise definition). We show that there is a canonical functor

$$\operatorname{Aff}_{z}(\mathcal{C}) \longrightarrow \operatorname{Raff}_{z}(\mathcal{C}),$$

which is faithful and essentially surjective, and prove several properties (see Section 4.3).

We further assume that C is an abelian **k**-linear graded monoidal category satisfying (5.1). For (M, z_M) and (N, z_N) in Aff_z(C), let $M = M/z_M M$ and $N = N/z_N N$. Under certain assumptions (see Proposition 6.2), we prove that

$$\operatorname{Hom}_{\mathbf{k}[z_{\mathsf{M}},z_{\mathsf{N}}]}(\mathsf{M}\otimes\mathsf{N},\mathsf{M}\otimes\mathsf{N})\simeq\mathbf{k}[z_{\mathsf{M}},z_{\mathsf{N}}]\mathsf{id}_{\mathsf{M}\otimes\mathsf{N}},$$

and there exists $R_{M,N}^{ren} \in Hom_{\mathbf{k}[z_M,z_N]}(\mathsf{M} \otimes \mathsf{N}, \mathsf{N} \otimes \mathsf{M})$ such that

$$\operatorname{Hom}_{\mathbf{k}[z_{\mathsf{M}}, z_{\mathsf{N}}]}(\mathsf{M} \otimes \mathsf{N}, \mathsf{N} \otimes \mathsf{M}) \simeq \mathbf{k}[z_{\mathsf{M}}, z_{\mathsf{N}}] \operatorname{R}^{\operatorname{ren}}_{\mathsf{M}, \mathsf{N}}$$

and $\mathbb{R}_{M,N}^{\text{ren}}|_{z_M=z_N=0} \in \operatorname{Hom}_{\mathcal{C}}(M \otimes N, N \otimes M)$ does not vanish. The morphism $\mathbb{R}_{M,N}^{\text{ren}}$ is called the *renormalized R-matrix*, and the morphism $\mathbf{r}_{M,N} := \mathbb{R}_{M,N}^{\text{ren}}|_{z_M=z_N=0}$ is called the *R-matrix* between M and N (see Definition 6.3). The above isomorphisms are typical properties which affinizations and R-matrices should have. We then introduce a *rational center* in \mathcal{C} using the category $\operatorname{Raff}_z(\mathcal{C})$. Using the notions of affine objects and rational center, we finally define an *affinization* \mathbb{M} of $M \in \mathcal{C}$ (see Definition 6.7). Proposition 7.3 explains that this definition is a generalization of the definition for affinizations (see (7.6)) given in [13, Definition 2.2]. In this categorical setting, the condition being a rational center makes R-matrices satisfy the Yang-Baxter equation (Lemma 6.23). We prove several properties of affinizations and $\mathbb{R}_{M,N}^{\text{ren}}$ and define the invariants in the language of categories:

$$(1.1) \qquad \begin{aligned} \mathscr{D}(\mathsf{M},\mathsf{N}) &\in \mathbf{k}[z_{\mathsf{M}},z_{\mathsf{N}}] \quad \text{such that} \quad \mathbf{R}_{\mathsf{N},\mathsf{M}}^{\mathrm{ren}} \circ \mathbf{R}_{\mathsf{M},\mathsf{N}}^{\mathrm{ren}} = \mathscr{D}(\mathsf{M},\mathsf{N}) \, \mathrm{id}_{\mathsf{M}\otimes\mathsf{N}}, \\ &\Lambda(M,N) := \mathrm{deg}(\mathbf{r}_{M,N}) \in \mathbb{Z}, \\ &\delta(M,N) := \frac{1}{2} \big(\Lambda(M,N) + \Lambda(N,M) \big) = \mathrm{deg}(\mathscr{D}(\mathsf{M},\mathsf{N}))/2 \in \frac{1}{2} \mathbb{Z}_{\geq 0}. \end{aligned}$$

We next introduce the notion of quasi-rigid categories. Note that the module categories of quiver Hecke algebras and quantum affine algebras are examples of quasi-rigid categories (see [6]). The quasi-rigidity plays an important role in proving the simplicity of the head of a tensor product of simples. As a consequence of the quasi-rigidity, the space of morphisms between various tensor products are one-dimensional (see Proposition 6.19 and Corollary 6.21) which guarantees that the renormalized R-matrices exist. It is conjectured that a quasi-rigid monoidal category which satisfies (5.1) is embedded into a rigid monoidal category (Conjecture 6.18). Under the additional assumption that C is quasi-rigid, we prove various properties of R-matrices and the invariants Λ and \mathfrak{d} . Thus, in the purely categorical language, we have recovered the key properties appeared in quiver Hecke algebras and quantum affine algebras.

Generalized Schur-Weyl duality. Let \mathcal{C} be a graded quasi-rigid monoidal category satisfying (6.2), and let R be the quiver Hecke algebra associated with parameter polynomials $Q_{i,j}(u, v)$. We define a *duality datum* $(\{(\widehat{K}_i, z_i)\}_{i \in I}, \{\mathbb{R}_{\widehat{K}_i, \widehat{K}_j}^{\text{ren}}\}_{i, j \in I})$ consisting of a family $\{(\widehat{K}_i, z_i)\}_{i \in I}$ of affinizations in \mathcal{C} satisfying (7.7) and the renormalized \mathbb{R} -matrices $\mathbb{R}_{\widehat{K}_i, \widehat{K}_j}^{\text{ren}} : \widehat{K}_i \otimes \widehat{K}_j \to \widehat{K}_j \otimes \widehat{K}_i$ satisfying (7.8). Applying the argument given in [7, 13] to the duality datum $(\{(\widehat{K}_i, z_i)\}_{i \in I}, \{\mathbb{R}_{\widehat{K}_i, \widehat{K}_j}^{\text{ren}}\}_{i, j \in I})$, we obtain a canonical right exact monoidal functor

$$\mathscr{F}: \operatorname{Modg}_{\operatorname{coh}}(R_{\lambda}) \to \operatorname{Pro}(\mathcal{C})$$

such that

$$\widehat{\mathscr{F}}(\widetilde{L}(i)_{z_i}) \simeq \widehat{\mathbf{K}}_i \quad \text{and} \quad \widehat{\mathscr{F}}(L(i)) \simeq \mathbf{K}_i,
\widehat{\mathscr{F}}(e(i,j)\varphi_1) = \mathbf{R}_{\widehat{\mathbf{K}}_i,\widehat{\mathbf{K}}_j}^{\text{ren}} \in \operatorname{Hom}_{\mathbf{k}[z_i,z_j]}(\widehat{\mathbf{K}}_i \otimes \widehat{\mathbf{K}}_j, \widehat{\mathbf{K}}_j \otimes \widehat{\mathbf{K}}_i),$$

where φ_1 is the intertwiner given in (7.4), and L(i) be the 1-dimensional simple $R(\alpha_i)$ module of degree 0. Then we have the restricted functor

$$\mathscr{F}: R ext{-gmod} \to \mathcal{C}$$

and prove that this functor enjoys the same good properties as the usual generalized Schur-Weyl duality (see Section 7.4).

The second main result is the reflection functor $\mathscr{S}_i : \widetilde{\mathscr{C}}_{s_i w_0} \xrightarrow{\sim} \widetilde{\mathscr{C}}_{s_i w_0}^*$ over the localizations, which categorifies the braid group action on the quantum group. We shall explain it in two parts, i.e., affinizations of the invariants and reflection functors.

Affinizations of the invariants. We lift the integer-valued invariants Λ , $\tilde{\Lambda}$, ϑ , wt, ε_i etc. which are used crucially in quiver Hecke algebras, to functions by using affinizations.

Let $(\mathsf{M}, z_{\mathsf{M}})$ and $(\mathsf{N}, z_{\mathsf{N}})$ be affinizations of simple *R*-modules *M* and *N* respectively. We assume that *M* and *N* are real for convenience. Let us recall $\chi_i(\mathsf{M})$, $\mathscr{E}_i(\mathsf{M})$ and $\mathscr{E}_i^*(\mathsf{M})$ defined in [11, Section 3.1] (see (9.1) for the precise definitions). By properties (see Lemma 4.11 for example) of affinizations, $\chi_i(\mathsf{M})$, $\mathscr{E}_i(\mathsf{M})$ and $\mathscr{E}_i^*(\mathsf{M})$ can be viewed as elements in $\mathbf{k}[t_i, z_{\mathsf{M}}]$. In Definition 9.3, we define $\mathscr{D}(\mathsf{M}, \mathsf{N})$ (see (1.1)) and $\widetilde{\mathscr{L}}(\mathsf{M}, \mathsf{N})$ in $\mathbf{k}[z_{\mathsf{M}}, z_{\mathsf{N}}]$ using the renormalized R-matrix $\mathrm{R}_{\mathsf{M},\mathsf{N}}^{\mathrm{ren}}$. We then define $\mathscr{L}(\mathsf{M}, \mathsf{N})$ and $\mathscr{W}t(\mathsf{M}, \mathsf{N})$ in $\mathbf{k}(x_{\mathsf{M}}, z_{\mathsf{N}})/\mathbf{k}^{\times}$ in terms of $\chi_i(\mathsf{M})$, $Q_{i,j}(t_i, t_j)$, $\widetilde{\mathscr{L}}(\mathsf{M}, \mathsf{N})$. It turns out that these new invariants can be understood as lifts of the well-known integer invariants in the following sense:

$$\begin{split} & \operatorname{deg}\big(\chi_i(\mathsf{M})\big) = n_i(\alpha_i, \alpha_i) \quad \operatorname{where } \operatorname{wt}(M) = -\sum_{i \in I} n_i \alpha_i, \\ & \operatorname{deg}\big(\mathscr{E}_i(\mathsf{M})\big) = \varepsilon_i(M)(\alpha_i, \alpha_i) = 2\widetilde{\Lambda}(L(i), M), \\ & \operatorname{deg}\big(\mathscr{E}_i^*(\mathsf{M})\big) = \varepsilon_i^*(M)(\alpha_i, \alpha_i) = 2\widetilde{\Lambda}(M, L(i)), \\ & \operatorname{deg}\big(\mathscr{D}(\mathsf{M}, \mathsf{N})\big) = 2\mathfrak{d}(M, N), \\ & \operatorname{deg}\big(\widetilde{\mathscr{L}}(\mathsf{M}, \mathsf{N})\big) = 2\widetilde{\Lambda}(M, N), \\ & \operatorname{deg}\big(\mathscr{L}(\mathsf{M}, \mathsf{N})\big) = 2\Lambda(M, N), \\ & \operatorname{deg}\big(\mathscr{W}t(\mathsf{M}, \mathsf{N})\big) = 2\big(\operatorname{wt}(M), \operatorname{wt}(N)\big), \end{split}$$

10

where deg denotes the homogeneous degree. We investigate relations between the lifted invariants and prove several interesting identities which are also interpreted as lifts of the identities between the integer invariants (see Section 9.2). The identities between lifted invariants are described in terms of the *resultant algebra* (see Section 9.1). These identities are used crucially in the construction of the reflection functors.

Reflection functors. We first investigate the canonical functor

$$\Phi_w\colon \mathscr{C}_w \to \widetilde{\mathscr{C}_w} \qquad (w \in \mathsf{W})$$

and prove that the functor Φ_w is fully faithful (Theorem 8.3). We thus show that the full subcategory \mathscr{C}_w of $\widetilde{\mathscr{C}_w}$ is stable by taking subquotients. However \mathscr{C}_w is not stable by taking extensions in $\widetilde{\mathscr{C}_w}$ in general (see Remark 8.6).

Let R be a quiver Hecke algebra of *arbitrary finite* type, and consider the categories $\mathscr{C}_{s_iw_0}$ and $\mathscr{C}^*_{s_iw_0}$ (see (8.1)). Fix an index $i \in I$. For any $j \in I$, we take the real simple objects in $\widetilde{\mathscr{C}}^*_{s_iw_0}$ as follows:


$$\mathbf{K}_{j} := \begin{cases} \mathscr{D} \left(\mathscr{Q}_{s_{i}w_{0}}^{*}(\langle i \rangle) \right) & \text{if } j = i, \\ \langle i^{-\mathbf{c}_{i,j}} \rangle \nabla \langle j \rangle & \text{if } j \neq i, \end{cases}$$

where \mathscr{D} is the right dual functor in $\widetilde{\mathscr{C}}_{s_iw_0}^*$, $\langle i^m \rangle$ is the self-dual simple $R(m\alpha_i)$ -module, and $\mathscr{D}_{s_iw_0}^*$: R-gmod $\to \widetilde{\mathscr{C}}_{s_iw_0}^*$ is the localization functor (see Section 8.2). We then take the canonical affinizations \widehat{K}_j of K_j (see (10.1)), which yield a duality datum

$$\left(\{(\widehat{\mathbf{K}}_j, z_j)\}_{j \in I}, \{\mathbf{R}_{\widehat{\mathbf{K}}_j, \widehat{\mathbf{K}}_k}^{\mathrm{ren}}\}_{j, k \in I}\right).$$

In particular, we take $\widehat{K}_i := \mathscr{D}_{aff}(\mathscr{Q}^*_{s_iw_0}(R(\alpha_i)))$ as the affinization of K_i by using the duality functor \mathscr{D}_{aff} developed in § 4. We then compute the invariants $\Lambda(K_j, K_k)$ and $\mathfrak{d}(K_j, K_k)$ (Proposition 10.1), and the lifted invariants $\mathscr{D}(\widehat{K}_j, \widehat{K}_k)$ (Proposition 10.2). Applying the generalized Schur-Weyl duality to this setting, we obtain the duality functor $\mathscr{F}_i : R$ -gmod $\longrightarrow \widetilde{\mathscr{C}}^*_{s_iw_0}$. Investigating the image of determinantial modules under the functor \mathscr{F}_i , we finally construct a functor $\mathscr{F}_i : \widetilde{\mathscr{C}}_{s_iw_0}$ such that the

following diagram quasi-commutes (Proposition 10.10):



Lemma 10.6 and Proposition 10.8 say that the functor \mathscr{S}_i categorifies the braid group action and the Saito reflection. Theorem 10.11 says that the reflection functor \mathscr{S}_i gives an equivalence of categories. By technical reasons, we separately deal with the case of type A_2 in Remark 10.13.

This paper is organized as follows. In Section 2, we recall necessary mathematical backgrounds for categories. In Section 3 and Section 4, we study affinizations in abelian categories under various conditions. In Section 5, we investigate affinizations in monoidal categories under various conditions, and Section 6 explains R-matrices in categories. In Section 7, we study a generalized Schur-Weyl duality, and in Section 8 we prove that the canonical functor $\Phi_w: \mathscr{C}_w \to \widetilde{\mathscr{C}_w}$ is fully faithful. In Section 9, we develop the lifted invariants using affinizations and Section 10 is devoted to reflection functors over localizations.

Acknowledgments. The authors thank Haruto Murata for valuable questions. The second, third and fourth authors gratefully acknowledge for the hospitality of RIMS (Kyoto University) during their visit in 2023.

Convention. (i) For a statement $P, \delta(P)$ is 1 or 0 whether P is true or not.

- (ii) Unless otherwise stated, a module over a ring is a left module.
- (iii) \mathbf{k} denotes a base field.
- (iv) For a ring A, we denote by Mod(A) the category of A-modules. We denote by $Mod_{coh}(A)$ the category of coherent A-modules.
- (v) For a field \mathbf{k} and a graded \mathbf{k} -algebra A, we denote by A-gmod the category of finite \mathbf{k} -dimensional graded A-modules.

- (vi) For a graded module $X = \bigoplus_{m \in \mathbb{Z}} X_m$, we write $X_{\ge m} = \bigoplus_{k \ge m} X_k$, $X_{\le m} = \bigoplus_{k \le m} X_k$ and $X_{>m} = \bigoplus_{k > m} X_k$.
- (vii) For a commutative **k**-algebra A and $f(z) = \sum_{0 \le k \le m} a_k z^k \in A[z]$, we say that f(z) is a *monic* polynomial of degree m if $a_m = 1$, and that f(z) is a *quasi-monic* polynomial of degree m if $a_m \in \mathbf{k}^{\times}$.
- (viii) For a category \mathcal{C} , \mathcal{C}^{opp} denotes the opposite category of \mathcal{C} .
- (ix) We say that a category I is *filtrant* if I satisfies the conditions
 - (a) $Ob(I) \neq \emptyset$,
 - (b) for any objects $i, j \in I$, there exist $k \in I$ and morphisms $i \to k$ and $j \to k$,
 - (c) for any morphisms $f: i \to j$ and $g: i \to j$, there exists a morphism $h: j \to k$ such that $h \circ f = h \circ g$.

We say that I is *cofiltrant* if I^{opp} is filtrant.

2. Preliminaries

2.1. Module objects. For a ring R and an additive category C, let us denote by Mod(R, C) the category of R-modules in C, i.e., the category of objects $X \in C$ endowed with a ring homomorphism $R \to End_{\mathcal{C}}(X)$. The morphisms between an R-module X and an R-module Y are the morphisms in C which commute with the R-actions.

Assume further that \mathcal{C} is an abelian category. We can define the tensor functor

•
$$\otimes_R \bullet \colon \operatorname{Mod}_{\operatorname{coh}}(R^{\operatorname{opp}}) \times \operatorname{Mod}(R, \mathcal{C}) \to \mathcal{C},$$

which is characterized by the universal property

$$\operatorname{Hom}_{\mathcal{C}}(M \otimes_{R} X, Y) \simeq \operatorname{Hom}_{\operatorname{Mod}(R^{\operatorname{opp}})}(M, \operatorname{Hom}_{\mathcal{C}}(X, Y))$$

functorially in $X \in Mod(R, C)$, $M \in Mod_{coh}(R^{opp})$ and $Y \in C$ (see [14, Remark 8.5.7]). The functor $\bullet \otimes_R \bullet$ is right exact in each variable.

Similarly, we have a functor

$$\mathcal{H}om_R(\bullet, \bullet): \operatorname{Mod}_{\operatorname{coh}}(R)^{\operatorname{opp}} \times \operatorname{Mod}(R, \mathcal{C}) \to \mathcal{C}$$

characterized by the universal property

$$\operatorname{Hom}_{\mathcal{C}}(X, \mathcal{H}om_{R}(M, Y)) \simeq \operatorname{Hom}_{\operatorname{Mod}(R)}(M, \operatorname{Hom}_{\mathcal{C}}(X, Y))$$

for $M \in Mod_{coh}(R)$, $Y \in Mod(R, C)$, $X \in C$, and $Hom_R(\bullet, \bullet)$ is left exact in each variable.

Assume that R^{opp} is noetherian. We say that an R-module X in \mathcal{C} has R-flat dimension $\leq m$ if $\operatorname{Tor}_{i}^{R}(M, X) = 0$ for any i > m and $M \in \operatorname{Mod}_{\operatorname{coh}}(R^{\operatorname{opp}})$, where $\operatorname{Tor}_{i}^{R}(\bullet, X)$ denotes the left derived functor of $\bullet \otimes_{R} X \colon \operatorname{Mod}_{\operatorname{coh}}(R^{\operatorname{opp}}) \to \mathcal{C}$. An R-module X in \mathcal{C} is called R-flat if its R-flat dimension is ≤ 0 , i.e. if $\bullet \otimes_{R} X \colon \operatorname{Mod}_{\operatorname{coh}}(R^{\operatorname{opp}}) \to \mathcal{C}$ is an exact functor.

2.2. Graded categories. Let C be an additive category. It is called a graded category if C is endowed with an auto-equivalence q (called the *grading shift functor*). For an additive graded category C, we set

$$\operatorname{Hom}_{\mathcal{C}}(M,N) := \bigoplus_{k \in \mathbb{Z}} \operatorname{Hom}_{\mathcal{C}}(M,N)_k \quad \text{with } \operatorname{Hom}_{\mathcal{C}}(M,N)_k := \operatorname{Hom}_{\mathcal{C}}(q^k M,N)$$

for $M, N \in \mathcal{C}$. Then \mathcal{C} has a category structure with $\operatorname{Hom}_{\mathcal{C}}$ as a set of morphisms. We say that $f \in \operatorname{Hom}_{\mathcal{C}}(M, N)_k$ is a morphism of degree k.

For a graded ring $R = \bigoplus_{k \in \mathbb{Z}} R_k$, a graded *R*-module in \mathcal{C} is an object *X* endowed with a graded ring homomorphism $R \to \text{End}_{\mathcal{C}}(X) := \text{Hom}_{\mathcal{C}}(X, X)$. The category of graded *R*-modules in \mathcal{C} is an additive graded category and denoted by $\text{Modg}(R, \mathcal{C})$. The morphisms between two graded-*R*-modules in \mathcal{C} are the morphisms in \mathcal{C} which commute with the *R*-actions.

If \mathcal{C} is abelian, we can define

•
$$\otimes_R$$
 • : $\operatorname{Modg}_{\operatorname{coh}}(R^{\operatorname{opp}}) \times \operatorname{Modg}(R, \mathcal{C}) \longrightarrow \mathcal{C}$ and
 $\mathcal{H}om_R(\bullet, \bullet) : \operatorname{Modg}_{\operatorname{coh}}(R)^{\operatorname{opp}} \times \operatorname{Modg}(R, \mathcal{C}) \longrightarrow \mathcal{C}.$

Here $Modg_{coh}(R)$ is the category of coherent graded *R*-modules.

In the sequel, we sometimes neglect grading shifts. For example, we sometimes write that $f: M \to N$ is a morphism when $f \in \text{Hom}(M, N)_k$.

2.3. **Pro-objects.** For a category C, let $\operatorname{Pro}(C)$ be the category of pro-objects of C (see [14, Section 6.1] for the definition and its properties). Then there is a fully faithful functor $C \to \operatorname{Pro}(C)$ and we regard C as a full subcategory of $\operatorname{Pro}(C)$. The category $\operatorname{Pro}(C)$ admits small cofiltrant projective limits, which we denote by "lim"

([14, Theorem 6.1.8]). Every object in $\operatorname{Pro}(\mathcal{C})$ is isomorphic to the projective limit " $\varprojlim_{i \in I}$ " X_i of a small cofiltrant projective system $\{X_i\}_{i \in I}$ in \mathcal{C} . Let \mathbf{k} be a field. Assume

that \mathcal{C} is an abelian (respectively, **k**-linear abelian) category. Then the category $\operatorname{Pro}(\mathcal{C})$ is also an abelian (respectively, **k**-linear abelian) category, and the canonical functor $\mathcal{C} \to \operatorname{Pro}(\mathcal{C})$ is exact. Moreover the functor $\lim_{i \in I}$ is exact ([14, Theorem 6.1.19]).

Namely, for any exact sequence of cofiltrant projective systems $X_i \to Y_i \to Z_i$ in $\operatorname{Pro}(\mathcal{C})$ indexed by a small cofiltrant category I, " $\varprojlim_{i \in I} X_i \to "\varprojlim_{i \in I} Y_i \to "\varprojlim_{i \in I} Z_i$ is

exact in $\operatorname{Pro}(\mathcal{C})$.

The subcategory C of Pro(C) is closed by taking kernels, cokernels and extensions ([14, Proposition 8.6.11]).

If C is an essentially small abelian (respectively, **k**-linear abelian) category, then the category Pro(C) is equivalent to the opposite category of the category of left exact functors from C to the category of abelian groups (respectively, **k**-vector spaces).

If C is an abelian monoidal category with a bi-exact tensor functor \otimes , then so is $\operatorname{Pro}(C)$ with the natural extension of \otimes .

Lemma 2.1. Let C be an abelian category, and let X be an artinian object of C. Then any quotient and subobject of X in Pro(C) are isomorphic to objects in C.

Proof. Let Y be an subobject of X in $\operatorname{Pro}(\mathcal{C})$. Then we can write $Y \simeq \underset{i \in I}{\overset{\text{where}}{\underset{i \in I}{\underset{i : I}{\underset{I}{\underset{i : I}{\underset{i :$

 $\{Y_i\}_{i\in I}$ is a projective system of subobjects of X indexed by a small cofiltrant category I.

Since X is artinian, there exists i_0 in I such that $Y_i \xrightarrow{\sim} Y_{i_0}$ for any $i \to i_0$. Hence $Y \xrightarrow{\sim} Y_{i_0}$ belongs to \mathcal{C} . If Z is a quotient of X in $\operatorname{Pro}(\mathcal{C})$, then $\operatorname{Ker}(X \to Z) \in \mathcal{C}$ and hence $Z \in \mathcal{C}$.

Let $F: \mathcal{C} \to \mathcal{C}'$ be a functor. Then F naturally induces a functor $\operatorname{Pro}(\mathcal{C}) \to \operatorname{Pro}(\mathcal{C}')$. We denote this functor by the same letter F.

3. Affinizations in Abelian Categories

3.1. Coherent objects in $Pro(\mathcal{C})$. Let **k** be a base field. Let \mathcal{C} be a **k**-linear category such that

- \mathcal{C} is abelian,
- (3.1) $\begin{cases} \bullet \ \mathcal{C} \text{ is } \mathbf{k}\text{-linear graded, namely } \mathcal{C} \text{ is endowed with a } \mathbf{k}\text{-linear auto-equivalence } q, \\ \bullet \text{ any object has a finite length,} \\ \bullet \text{ any simple object } S \text{ is absolutely simple, i.e., } \mathbf{k} \xrightarrow{\sim} \text{END}_{\mathcal{C}}(S) := \\ \text{How } (S, S) \end{cases}$

It follows from (3.1) that $\operatorname{Hom}_{\mathcal{C}}(M, N)$ is finite-dimensional over **k** for any $M, N \in \mathcal{C}$ and that $S \not\simeq q^k S$ for any $k \in \mathbb{Z} \setminus \{0\}$ and any non-zero $S \in \mathcal{C}$.

By Lemma 2.1, the full subcategory \mathcal{C} of $\operatorname{Pro}(\mathcal{C})$ is stable by taking subquotients. The grading shift functor is extended to $Pro(\mathcal{C})$.

Let $A = \bigoplus_{k \in \mathbb{Z}} A_k$ be a commutative graded **k**-algebra such that

(3.2)
$$\begin{cases} \bullet \ A_0 \simeq \mathbf{k}, \\ \bullet \ A_k = 0 \text{ for any} \end{cases}$$

 $\begin{cases} \bullet \ A_k = 0 \text{ for any } k < 0, \\ \bullet \ A \text{ is a finitely generated } \mathbf{k}\text{-algebra.} \end{cases}$

In particular, A is a noetherian ring and $\dim_{\mathbf{k}} A_k < \infty$ for all k. For each $m \in \mathbb{Z}$, set

$$A_{\geqslant m} := \bigoplus_{k \ge m} A_k,$$

which is an ideal of A.

Hereafter, we assume that C satisfies (3.1) and A satisfies (3.2).

Recall that $Modg(A, Pro(\mathcal{C}))$ denotes the category of graded A-modules in $Pro(\mathcal{C})$ (see § 2.2).

For $\mathsf{M}, \mathsf{N} \in \mathrm{Modg}(A, \mathrm{Pro}(\mathcal{C}))$, we set $\mathrm{Hom}_A(\mathsf{M}, \mathsf{N}) := \bigoplus_{k \in \mathbb{Z}} \mathrm{Hom}_A(\mathsf{M}, \mathsf{N})_k$ with $\mathrm{Hom}_A(\mathsf{M}, \mathsf{N})_k := \mathrm{Hom}_{\mathrm{Modg}(A, \mathrm{Pro}(\mathcal{C}))}(q^k \mathsf{M}, \mathsf{N})$. Then $\mathrm{Hom}_A(\mathsf{M}, \mathsf{N})$ has a structure of a graded A-module.

Definition 3.1. We denote by $\operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$ the full subcategory of $\operatorname{Modg}(A, \operatorname{Pro}(\mathcal{C}))$ consisting of graded A-modules M in $\operatorname{Pro}(\mathcal{C})$ such that

(a)
$$\mathsf{M}/A_{>0}\mathsf{M} \in \mathcal{C}$$
,

(b) $\mathsf{M} \xrightarrow{\sim} \underset{k}{\overset{\text{"lim"}}{\longrightarrow}} \mathsf{M}/A_{\geq k}\mathsf{M}$, or equivalently $\underset{k}{\overset{\text{"lim"}}{\longrightarrow}} A_{\geq k}\mathsf{M} \simeq 0$.

Here $A_{\geq k} \mathsf{M} := \operatorname{Im}(A_{\geq k} \otimes_A \mathsf{M} \to A \otimes_A \mathsf{M}) \subset \mathsf{M}$. Note that $X \otimes_A \mathsf{M}$ belongs to $\operatorname{Modg}(A, \operatorname{Pro}(\mathcal{C}))$ for any $\mathsf{M} \in \operatorname{Modg}(A, \operatorname{Pro}(\mathcal{C}))$ and any finitely generated graded A-module X.

Lemma 3.2. Let $M \in Modg(A, Pro(\mathcal{C}))$.

(i) For any $m \in \mathbb{Z}_{\geq 0}$, the canonical morphism

$$(3.3) \qquad (A_{\geq m}/A_{>m}) \otimes_{\mathbf{k}} (\mathsf{M}/A_{>0}\mathsf{M}) \simeq (A_{\geq m}/A_{>m}) \otimes_{A} \mathsf{M} \to A_{\geq m}\mathsf{M}/A_{>m}\mathsf{M}$$

is an epimorphism in \mathcal{C} .

- (ii) In particular, if $M/A_{>0}M \in C$, then the object $A_{\geq m}M/A_{>m}M$ belongs to C by Lemma 2.1.
- (iii) If M is A-flat, then the morphism (3.3) is an isomorphism.

Proof. Note that

$$\begin{aligned} (A_{\geq m}/A_{>m}) \otimes_A \mathsf{M} &\simeq \left((A_{\geq m}/A_{>m}) \otimes_{\mathbf{k}} (A/A_{>0}) \right) \otimes_A \mathsf{M} \\ &\simeq (A_{\geq m}/A_{>m}) \otimes_{\mathbf{k}} \left((A/A_{>0}) \otimes_A \mathsf{M} \right) \simeq (A_{\geq m}/A_{>m}) \otimes_{\mathbf{k}} (\mathsf{M}/A_{>0}\mathsf{M}). \end{aligned}$$

We have a commutative diagram in $Modg(A, Pro(\mathcal{C}))$

with exact rows. Since the middle vertical arrow is an epimorphism, so is the right vertical arrow.

If M is A-flat, then the left and the middle vertical arrows in (3.4) are isomorphisms, and hence so is the right vertical arrow.

Proposition 3.3. The full subcategory $\operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$ of $\operatorname{Modg}(A, \operatorname{Pro}(\mathcal{C}))$ has the following properties.

- (i) The category Pro_{coh}(A, C) is stable by taking subquotients and taking extensions in Modg(A, Pro(C)).
- (ii) Any object of $\operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$ is noetherian.
- (iii) For any $\mathsf{M} \in \operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$, we have
 - (a) $A_{\geq m} \mathsf{M} / A_{\geq n} \mathsf{M} \in \mathcal{C}$ for any $m, n \in \mathbb{Z}$ such that $0 \leq m \leq n$,
 - (b) for any $m \in \mathbb{Z}_{\geq 0}$ and any $\mathsf{N} \in \operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$ such that $\mathsf{N} \subset \mathsf{M}$, there exists n > 0 such that $\mathsf{N} \cap A_{\geq n} \mathsf{M} \subset A_{\geq m} \mathsf{N}$,
 - (c) if $\mathsf{M} = A_{>0}\mathsf{M}$, then $\mathsf{M} = 0$.

Proof. (iiia) immediately follows from Lemma 3.2.

Take a set $\{z_i\}_{1 \leq i \leq r}$ of homogeneous generators of the **k**-algebra A with positive degrees. (*l* is used later.) We shall show the statements by induction on r. When r = 0, they are obvious. Assume that r > 0. We set $B := \mathbf{k}[z_2, \ldots, z_r] \subset A$. Then (i)–(iii) with A replaced by B hold by the induction hypothesis. Set $d_1 := \deg(z_1) \in \mathbb{Z}_{>0}$ and $d := \max \{ \deg(z_i) \mid 1 \leq i \leq r \}$.

Then $A_{\geq k} = \sum_{j \in \mathbb{Z}_{\geq 0}} z_1^j B_{\geq k-d_1 j}$. In particular, we have $A_{>0} = z_1 A + B_{>0}$. Since $A_{\geq dk} \subset (A_{>0})^k \subset A_{\geq k}$ for any $k \geq 1$, (iiic) is obvious. It is also obvious that

(3.5) $\operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$ is stable by taking quotients.

Let us first show that

(3.6) Let $\mathsf{M} \in \mathrm{Mod}(A, \mathrm{Pro}(\mathcal{C}))$, and assume that there exists $n \in \mathbb{Z}_{>0}$ such that $z_1^n \mathsf{M} = 0$. Then $\mathsf{M} \in \mathrm{Pro}_{\mathrm{coh}}(A, \mathcal{C})$ if and only if $\mathsf{M} \in \mathrm{Pro}_{\mathrm{coh}}(B, \mathcal{C})$.

Μ

Assume that $\mathsf{M} \in \operatorname{Pro}_{\operatorname{coh}}(B, \mathcal{C})$. Since there is an epimorphism $\mathsf{M}/B_{>0}\mathsf{M} \twoheadrightarrow \mathsf{M}/A_{>0}\mathsf{M}$ in $\operatorname{Pro}(\mathcal{C})$, we have $\mathsf{M}/A_{>0}\mathsf{M} \in \mathcal{C}$. Since

$$A_{\geqslant k} \subset Az_1^n + B_{>k-nd_1}A \quad \text{for } k \in \mathbb{Z}_{\geqslant 0},$$

we have " $\varprojlim_{k} A_{\geq k} \mathsf{M} \subset$ " $\varprojlim_{k} (B_{\geq k-nd_{1}} \mathsf{M}) \simeq 0$. Hence we obtain $\mathsf{M} \in \operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$. Conversely, assume that $\mathsf{M} \in \operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$. Then we have " $\varprojlim_{k} B_{\geq k} \mathsf{M} \subset$ " $\varprojlim_{k} A_{\geq k} \mathsf{M} \simeq$ 0. Let us show $\mathsf{M}/B_{\geq 0} \mathsf{M} \in \mathcal{C}$. Note that $A_{\geq nd_{1}} \subset z_{1}^{n}A + B_{\geq 0}A$. Since $\mathsf{M}/A_{\geq nd_{1}} \mathsf{M} \in \mathcal{C}$ by (iiia), its quotient $\mathsf{M}/(z_{1}^{n}A + B_{\geq 0}A)\mathsf{M} \simeq \mathsf{M}/B_{\geq 0}\mathsf{M}$ belongs to \mathcal{C} . It completes the proof of (3.6).

By (3.5) and (3.6), we have

(3.7) for any
$$\mathsf{M} \in \operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$$
 and $j, k \in \mathbb{Z}_{\geq 0}$ such that $j \leq k$, we have $z_1^j \mathsf{M}/z_1^k \mathsf{M} \in \operatorname{Pro}_{\operatorname{coh}}(B, \mathcal{C}).$

We shall show that

(3.8) for any
$$\mathsf{M} \in \operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$$
 and $\mathsf{N} \in \operatorname{Modg}(A, \operatorname{Pro}(\mathcal{C}))$ such that $\mathsf{N} \subset \mathsf{M}$,
there exists $c \in \mathbb{Z}_{>0}$ such that $\mathsf{N} \cap z_1^{k+c} \mathsf{M} \subset z_1^k \mathsf{N}$ for any $k \in \mathbb{Z}_{\geq 0}$.

Since $\mathsf{M}/z_1\mathsf{M} \in \operatorname{Pro}_{\operatorname{coh}}(B, \mathcal{C})$ is noetherian, the increasing sequence $\{(z_1^k)^{-1}\mathsf{N} + z_1\mathsf{M}\}_{k\in\mathbb{Z}_{\geq 0}}$ of subobjects of M is stationary. Hence there exists $k_0 \in \mathbb{Z}_{\geq 0}$ such that $(z_1^k)^{-1}\mathsf{N}+z_1\mathsf{M} \subset (z_1^{k_0})^{-1}\mathsf{N}+z_1\mathsf{M}$ for any $k \geq k_0$. Hence we have $\mathsf{N} \cap z_1^k\mathsf{M} \subset z_1^{k-k_0}\mathsf{N}+z_1^{k+1}\mathsf{M}$. Let $l \in \mathbb{Z}_{\geq 0}$. Then, we have $\mathsf{N} \cap z_1^k\mathsf{M} \subset z_1^l\mathsf{N}+\mathsf{N} \cap z_1^{k+1}\mathsf{M}$ for any $k \geq k_0+l$. Hence, by induction on k, we have $\mathsf{N} \cap z_1^{k_0+l}\mathsf{M} \subset z_1^l\mathsf{N}+\mathsf{N} \cap z_1^k\mathsf{M}$ for any $k \geq k_0+l$. Since $(\varprojlim_k)^k z_1^k\mathsf{M} \simeq 0$, we

obtain $\mathsf{N} \cap z_1^{k_0+l} \mathsf{M} \subset z_1^l \mathsf{N}$. It shows (3.8).

Let us show

(3.9) $\operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$ is stable by extensions.

Let $0 \to \mathsf{M}' \to \mathsf{M} \to \mathsf{M}'' \to 0$ be an exact sequence in $\operatorname{Mod}(A, \operatorname{Pro}(\mathcal{C}))$ such that $\mathsf{M}', \mathsf{M}'' \in \operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$. Let us show that $\mathsf{M} \in \operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$.

Since we have an exact sequence

$$\mathsf{M}'/A_{>0}\,\mathsf{M}'\to\,\mathsf{M}/A_{>0}\,\mathsf{M}\to\,\mathsf{M}''/A_{>0}\,\mathsf{M}''\to 0$$

in which the left and right terms belong to \mathcal{C} , we have $M/A_{>0}M \in \mathcal{C}$.

Let us show " $\varprojlim_{k} A_{>k} M \simeq 0$. Since the composition $\mathsf{N} :=$ " $\varprojlim_{k} A_{>k} M \to \mathsf{M}''$ factors through " $\varprojlim_{k} A_{>k} \mathsf{M}'' \simeq 0$, it is a zero morphism. Hence $\mathsf{N} \subset \mathsf{M}'$. Hence we

have

$$0 \simeq \underset{k}{\overset{\text{"lim"}}{\longleftarrow}} A_{>k} \mathsf{M}' \supset \underset{k}{\overset{\text{"lim"}}{\longleftarrow}} A_{>k} \mathsf{N} \simeq \underset{k,j}{\overset{\text{"lim"}}{\longleftarrow}} A_{>k} A_{>j} \mathsf{M} \simeq \underset{k}{\overset{\text{"lim"}}{\longleftarrow}} A_{>k} \mathsf{M},$$

which implies that " $\varprojlim_k A_{>k} M \simeq 0$. Thus we obtain (3.9).

Now, we are ready to prove

(3.10)
$$\operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$$
 is stable by taking subobjects.

Let $\mathsf{M} \in \operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$ and $\mathsf{N} \in \operatorname{Modg}(A, \operatorname{Pro}(\mathcal{C}))$ such that $\mathsf{N} \subset \mathsf{M}$. By (3.8), there exists $m \in \mathbb{Z}_{>0}$ such that $\mathsf{N} \cap z_1^m \mathsf{M} \subset z_1 \mathsf{N}$. Since $\mathsf{M}/z_1^m \mathsf{M} \in \operatorname{Pro}_{\operatorname{coh}}(B, \mathcal{C})$ by (3.7), its subobject $\mathsf{N}/(\mathsf{N} \cap z_1^m \mathsf{M})$ belongs to $\operatorname{Pro}_{\operatorname{coh}}(B, \mathcal{C})$ by the induction hypothesis. Hence its quotient $\mathsf{N}' := \mathsf{N}/z_1 \mathsf{N}$ also belongs to $\operatorname{Pro}_{\operatorname{coh}}(B, \mathcal{C})$. Since $\mathsf{N}'/B_{>0}\mathsf{N}' \in \mathcal{C}$ and we have $(\mathsf{N}/z_1\mathsf{N})/B_{>0}(\mathsf{N}/z_1\mathsf{N}) \simeq \mathsf{N}/(z_1\mathsf{N} + B_{>0}\mathsf{N}) \twoheadrightarrow \mathsf{N}/A_{>0}\mathsf{N}$, we conclude that $\mathsf{N}/A_{>0}\mathsf{N} \in \mathcal{C}$. Moreover, " $\varprojlim_k A_{\geq k}\mathsf{N} \subset ``\varprojlim_k A_{\geq k}\mathsf{M}$ vanishes. Hence $\mathsf{N} \in \operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$ and we obtain (2.10)

(3.10).

Thus we have completed the proof of (i).

Let us show (iiib). Since " \varprojlim_{k} " ($\mathbb{N} \cap A_{\geq k} \mathbb{M}$) $\simeq 0$, we have $\mathbb{N} \simeq "\varprojlim_{k}$ " $\mathbb{N}/(\mathbb{N} \cap A_{\geq k} \mathbb{M})$. On the other hand, (iiia) implies that $\mathbb{N}/A_{\geq m}\mathbb{N}$ belongs to \mathcal{C} . Therefore, there exists $n \in \mathbb{Z}_{>0}$ such that the morphism $\mathbb{N} \simeq "\varprojlim_{k}$ " $\mathbb{N}/(\mathbb{N} \cap A_{\geq k} \mathbb{M}) \to \mathbb{N}/A_{\geq m}\mathbb{N}$ factors through $\mathbb{N}/(\mathbb{N} \cap A_{\geq n} \mathbb{M})$. Hence we obtain $\mathbb{N} \cap A_{\geq n} \mathbb{M} \subset A_{\geq m}\mathbb{N}$.

It remains to prove (ii). Let $\{N_j\}_{j \in \mathbb{Z}_{>0}}$ be an increasing sequence of subobjects of $M \in \operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$. We shall show that it is stationary.

The family $\{(z_1^k)^{-1}\mathsf{N}_j + z_1\mathsf{M}\}_{j,k\in\mathbb{Z}_{>0}}$ is stationary since $\mathsf{M}/z_1\mathsf{M} \in \operatorname{Pro}_{\operatorname{coh}}(B,\mathcal{C})$ is noetherian by the induction hypothesis. Hence there exist $k_0, j_0 \in \mathbb{Z}_{>0}$ such that $(z_1^k)^{-1}\mathsf{N}_j + z_1\mathsf{M} = (z_1^{k_0})^{-1}\mathsf{N}_{j_0} + z_1\mathsf{M}$ for $j \ge j_0$ and $k \ge k_0$. Hence we have

$$\mathsf{N}_j \cap z_1^k \mathsf{M} \subset \mathsf{N}_{j_0} + z_1^{k+1} \mathsf{M} \quad \text{for } k \ge k_0 \text{ and } j \ge j_0.$$

It implies that

$$\mathsf{N}_{j_0} + \mathsf{N}_j \cap z_1^k \mathsf{M} \subset \mathsf{N}_{j_0} + \mathsf{N}_j \cap z_1^{k+1} \mathsf{M}$$
 for any $k \ge k_0$.

Hence, by induction, we obtain

$$\mathsf{N}_{j_0} + \mathsf{N}_j \cap z_1^{k_0} \mathsf{M} \subset \mathsf{N}_{j_0} + \mathsf{N}_j \cap z_1^{k+1} \mathsf{M} \quad \text{for any } k \ge k_0.$$

Taking " \varprojlim_k ", we obtain

$$\mathsf{N}_{j_0} + \mathsf{N}_j \cap z_1^{k_0} \mathsf{M} \subset \mathsf{N}_{j_0}.$$

Hence $N_j \cap z_1^{k_0} M \subset N_{j_0}$ for $j \ge j_0$. On the other hand, $\{N_j + z_1^{k_0} M\}_{j \in \mathbb{Z}_{>0}}$ is stationary since $M/z_1^{k_0} M \in \operatorname{Pro}_{\operatorname{coh}}(B, \mathcal{C})$ is noetherian. Hence, there exists $j_1 \ge j_0$ such that $N_j + z_1^{k_0} M = N_{j_1} + z_1^{k_0} M$ for $j \ge j_1$. Hence we have $N_j \subset N_{j_1} + N_j \cap z_1^{k_0} M \subset N_{j_1} + N_{j_0} = N_{j_1}$ for any $j \ge j_1$.

Lemma 3.4. Let $X \in Modg_{coh}(A)$ and $M \in Pro_{coh}(A, C)$. Then we have:

- (i) $X \otimes_A \mathsf{M} \in \operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C}),$
- (ii) if dim_{**k**} $X < \infty$, then $X \otimes_A \mathsf{M} \in \mathcal{C}$.

Proof. (i) Taking an epimorphism $A^{\oplus r} \twoheadrightarrow X$, we conclude (i) since $X \otimes_A \mathsf{M}$ is a quotient of $\mathsf{M}^{\oplus r}$.

(ii) Since dim_{**k**} END_A(X) < ∞ , there exists $m \in \mathbb{Z}$ such that END_A(X)_{$\geq m$} = 0. Then $A_{\geq m}X = 0$, which implies that $A_{\geq m}(X \otimes_A \mathsf{M}) = 0$. Hence $X \otimes_A \mathsf{M} \in \mathcal{C}$ by Proposition 3.3 (iiia).

Lemma 3.5. Let $f: \mathsf{M} \to \mathsf{N}$ be a morphism in $\operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$.

(i) If $\tilde{f} := (A/A_{>0}) \otimes_A f \colon \mathsf{M}/A_{>0}\mathsf{M} \to \mathsf{N}/A_{>0}\mathsf{N}$ is an epimorphism, then f is an epimorphism.

(ii) If N is A-flat and \tilde{f} is an isomorphism, then f is an isomorphism.

Proof. (i) If \tilde{f} is an epimorphism, then $(A/A_{>0}) \otimes_A \operatorname{Coker}(f) \simeq 0$, and hence $\operatorname{Coker}(f) \simeq 0$ by Proposition 3.3.

(ii) Since N is A-flat, we have an exact sequence

$$0 \to (A/A_{>0}) \otimes_A \operatorname{Ker}(f) \to \mathsf{M}/A_{>0}\mathsf{M} \to \mathsf{N}/A_{>0}\mathsf{N} \to 0.$$

Hence $\operatorname{Ker}(f) \simeq 0$.

Proposition 3.6. Let $M, N \in \operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$ and set $M := M/A_{>0}M \in \mathcal{C}$ and $N := N/A_{>0}N \in \mathcal{C}$. Assume that N is A-flat. Let X be a finitely generated graded A-module and $\lambda, \mu \in \mathbb{Z}$. Assume that $\operatorname{Hom}_{\mathcal{C}}(M, N)$ has degree $\geq \lambda$ and X has degree $\geq \mu$. Then we have

(i) $\operatorname{Hom}_A(\mathsf{M}, X \otimes_A \mathsf{N})$ has degree $\geq \mu + \lambda$.

(ii) $\operatorname{Hom}_A(\mathsf{M}, X \otimes_A \mathsf{N})$ is a finitely generated A-module,

(iii) there exists an integer $m \in \mathbb{Z}_{\geq 0}$ such that

$$\operatorname{Hom}_A(\mathsf{M}, X_{\geqslant k+m} \otimes_A \mathsf{N}) \subset A_{\geqslant k} \operatorname{Hom}_A(\mathsf{M}, X \otimes_A \mathsf{N}) \quad for \ any \ k \in \mathbb{Z}_{\geqslant 0},$$

as a subset of $\operatorname{Hom}_A(\mathsf{M}, X \otimes_A \mathsf{N})$.

Proof. Set $L = X \otimes_A N \in Pro_{coh}(A, C)$ and $L_{\geq m} = X_{\geq m} \otimes_A N \subset L$. Since N is A-flat, we have

$$\mathsf{L}_{\geq m}/\mathsf{L}_{>m} \simeq (X_{\geq m}/X_{>m}) \otimes_A \mathsf{N} \simeq X_m \otimes_{\mathbf{k}} N.$$

(i) We have an inclusion

(3.11)

$$\frac{\operatorname{Hom}_{A}(\mathsf{M},\mathsf{L}_{\geq m})}{\operatorname{Hom}_{A}(\mathsf{M},\mathsf{L}_{>m})} \longrightarrow \operatorname{Hom}_{A}(\mathsf{M},\mathsf{L}_{\geq m}/\mathsf{L}_{>m}) \xrightarrow{\sim} X_{m} \otimes_{\mathbf{k}} \operatorname{Hom}_{\mathcal{C}}(M,N).$$

Hence, $\operatorname{Hom}_A(\mathsf{M}, \mathsf{L}_{\geq m}/\mathsf{L}_{>m})$ has degree $\geq m + \lambda$. Therefore, $\operatorname{Hom}_A(\mathsf{M}, \mathsf{L}_{\geq m}/\mathsf{L}_{\geq s})$ has also degree $\geq m + \lambda$ for any $s \geq m$. Hence, we obtain

$$\operatorname{Hom}_{A}(\mathsf{M},\mathsf{L})_{k} \simeq \varprojlim_{s} \operatorname{Hom}_{A}(\mathsf{M},\mathsf{L}_{\geqslant \mu}/\mathsf{L}_{\geqslant s})_{k} \simeq 0 \quad \text{if } k < \mu + \lambda.$$

Thus we obtain (i). Applying (i) to $X_{\geq m}$, we obtain

(3.12) $\operatorname{Hom}_A(\mathsf{M},\mathsf{L}_{\geq m})$ has degree $\geq m + \lambda$.

Let $S = \bigoplus_{m \in \mathbb{Z}} S_m$ be the graded A-module with $S_m = \operatorname{Hom}_A(\mathsf{M}, \mathsf{L}_{\geq m}) / \operatorname{Hom}_A(\mathsf{M}, \mathsf{L}_{>m})$. Then by (3.11) we have a monomorphism

 $S \rightarrow X \otimes_{\mathbf{k}} \operatorname{Hom}_{\mathcal{C}}(M, N),$

which implies that S is a finitely generated graded A-module. Hence there exist $r \in \mathbb{Z}_{\geq \mu}$ and a finite-dimensional graded k-subspace $K_j \subset \text{Hom}_A(\mathsf{M}, \mathsf{L}_{\geq j})$ $(\mu \leq j \leq r)$ such that

$$\operatorname{Hom}_{A}(\mathsf{M},\mathsf{L}_{\geq m}) \subset \sum_{\mu \leq j \leq r} A_{\geq m-j} K_{j} + \operatorname{Hom}_{A}(\mathsf{M},\mathsf{L}_{>m}) \quad \text{for any } m \in \mathbb{Z}.$$

Hence by induction on t, we obtain $\operatorname{Hom}_A(\mathsf{M},\mathsf{L}_{\geq m}) \subset \sum_{j=\mu}^r A_{\geq m-j}K_j + \operatorname{Hom}_A(\mathsf{M},\mathsf{L}_{\geq t})$ for any $m \in \mathbb{Z}$ and $t \in \mathbb{Z}$ such that $t \geq m$. Hence (3.12) implies that $\operatorname{Hom}_A(\mathsf{M},\mathsf{L}_{\geq m}) \subset \sum_{j=0}^r A_{\geq m-j}K_j$ from which (ii) and (iii) follows.

Proposition 3.7. Let $M, N \in Pro_{coh}(A, C)$, and set $M = M/A_{>0}M$ and $N = N/A_{>0}N$. Assume that N is A-flat. Let H_0 be a graded k-subspace of $Hom_A(M, N)$.


- (i) If the composition $H_0 \to \operatorname{Hom}_A(\mathsf{M},\mathsf{N}) \xrightarrow{(A/A_{>0})\otimes_A^{\bullet}} \operatorname{Hom}_{\mathcal{C}}(M,N)$ is injective, then, $A \otimes_{\mathbf{k}} H_0 \to \operatorname{Hom}_A(\mathsf{M},\mathsf{N})$ is injective.
- (ii) If $H_0 \to \operatorname{Hom}_{\mathcal{C}}(M, \mathbb{N})$ is surjective, then, $A \otimes_{\mathbf{k}} H_0 \to \operatorname{Hom}_A(\mathsf{M}, \mathsf{N})$ is surjective.

Proof. Let $s \in \mathbb{Z}$. Then the composition

$$g_m : \left((A_{\geq m}/A_{\geq m+1}) \otimes_{\mathbf{k}} H_0 \right)_s \to \left((A_{\geq m}/A_{\geq m+1}) \otimes_{\mathbf{k}} \operatorname{Hom}_{\mathcal{C}}(M, N) \right)_s \xrightarrow{\sim} \operatorname{Hom}_A(\mathsf{M}, A_{\geq m}\mathsf{N}/A_{\geq m+1}\mathsf{N})_s$$

is injective (resp. surjective) under the assumption in (i) (resp. (ii)). We shall show that $f_m: (A_{\geq m} \otimes_{\mathbf{k}} H_0)_s \to \operatorname{Hom}_A(\mathsf{M}, A_{\geq m}\mathsf{N})_s$ is injective (resp. surjective) by the descending induction on $m \in \mathbb{Z}_{\geq 0}$. If $m \gg 0$, then it is true since $(A_{\geq m} \otimes H_0)_s \simeq$ $\operatorname{Hom}_A(\mathsf{M}, A_{\geq m}\mathsf{N})_s \simeq 0$ by Proposition 3.6.

Now assume that f_{m+1} is injective (resp. surjective). Consider a commutative diagram with exact rows:



Since f_{m+1} and g_m are injective (resp. surjective), so is f_m , and the induction proceeds.

Proposition 3.8. Let $M, N \in Pro_{coh}(A, C)$, and set $M = M/A_{>0}M$ and $N = N/A_{>0}N$. Assume that

- (a) A is an integral domain,
- (b) N is A-flat,
- (c) $\operatorname{Hom}_{\mathcal{C}}(M, N) = \mathbf{k}f$ for a non-zero f of homogeneous degree $\lambda \in \mathbb{Z}$,
- (d) there exists a non-zero $F \in Hom_A(M, N)$.

Then, $\operatorname{Hom}_A(\mathsf{M}, \mathsf{N})$ is a free A-module generated by an element $\widetilde{F} \in \operatorname{Hom}_A(\mathsf{M}, \mathsf{N})_{\lambda}$ such that $(A/A_{>0}) \otimes_A \widetilde{F} = f$.

Proof. First note that $\operatorname{Hom}_A(\mathsf{M}, A_{\geq m}\mathsf{N})$ is of degree $\geq m + \lambda$ by Proposition 3.6.

Let $\psi \in \operatorname{Hom}_A(\mathsf{M},\mathsf{N})$ be a non-zero homomorphism of degree d. Let us take $s \in \mathbb{Z}_{\geq 0}$ such that $\psi \in \operatorname{Hom}_A(\mathsf{M}, A_{\geq s}\mathsf{N})$ and $\psi \notin \operatorname{Hom}_A(\mathsf{M}, A_{>s}\mathsf{N})$. Then, its image $\overline{\psi} \in \operatorname{Hom}_A(\mathsf{M}, A_{\geq s}\mathsf{N}/A_{>s}\mathsf{N})$ does not vanish. Since $\operatorname{Hom}_A(\mathsf{M}, A_{\geq s}\mathsf{N}/A_{>s}\mathsf{N}) \simeq (A_{\geq s}/A_{>s}) \otimes \operatorname{Hom}_{\mathcal{C}}(M, N)$ by (b), there exists $a \in A_s$ such that $\overline{\psi} = af$. In particular, $d = s + \lambda$. Hence, we have $\psi \in \operatorname{Hom}_A(\mathsf{M}, a\mathsf{N} + A_{>s}\mathsf{N})$. We have an exact sequence

$$0 \longrightarrow \operatorname{Hom}_{A}(\mathsf{M}, a\mathsf{N})_{d} \longrightarrow \operatorname{Hom}_{A}(\mathsf{M}, a\mathsf{N} + A_{>s}\mathsf{N})_{d} \longrightarrow \operatorname{Hom}_{A}(\mathsf{M}, (a\mathsf{N} + A_{>s}\mathsf{N})/a\mathsf{N})_{d}.$$

Since $(a\mathsf{N} + A_{>s}\mathsf{N})/a\mathsf{N} \simeq ((aA + A_{>s})/aA) \otimes_A \mathsf{N}$ and $(aA + A_{>s})/aA \simeq A_{>s}/(aA \cap A_{>s})$ has degree $\geq s + 1$, Proposition 3.6 implies that $\operatorname{Hom}_A(\mathsf{M}, (a\mathsf{N} + A_{>s}\mathsf{N})/a\mathsf{N})$ has degree $\geq s + 1 + \lambda = d + 1$. Hence, $\operatorname{Hom}_A(\mathsf{M}, (a\mathsf{N} + A_{>s}\mathsf{N})/a\mathsf{N})_d \simeq 0$, which implies that $\operatorname{Hom}_A(\mathsf{M}, a\mathsf{N})_d \simeq \operatorname{Hom}_A(\mathsf{M}, a\mathsf{N} + A_{>s}\mathsf{N})_d$. Hence we have $\psi \in \operatorname{Hom}_A(\mathsf{M}, a\mathsf{N})_d = a \operatorname{Hom}_A(\mathsf{M}, \mathsf{N})_\lambda$ since A is an integral domain. Hence there exists $\psi' \in \operatorname{Hom}_A(\mathsf{M}, \mathsf{N})_\lambda$ such that $\psi = a\psi'$ and $(A/A_{>0}) \otimes \psi' = f$.

Applying it to $\psi = F$, there exist $a \in A$ and $\widetilde{F} \in \text{Hom}_A(\mathsf{M}, \mathsf{N})_{\lambda}$ such that $F = a\widetilde{F}$. It remains to remark $\text{Hom}_A(\mathsf{M}, \mathsf{N})_{\lambda} \xrightarrow{\sim} \text{Hom}_{\mathcal{C}}(M, N)_{\lambda}$. Hence $\text{Hom}_A(\mathsf{M}, \mathsf{N})$ is gener-

ated by \widetilde{F} as an A-module. Since N is A-flat, $a : \mathbb{N} \to \mathbb{N}$ is injective for any $a \in A$. Thus $a\widetilde{F} = 0$ implies a = 0 so that $\operatorname{Hom}_A(\mathbb{M}, \mathbb{N})$ is free over A.

Remark 3.9. In Proposition 3.8, the condition (a) is necessary. Indeed, if we take $\mathcal{C} = \mathbf{k}$ -gmod, $A = \mathbf{k}[z]/\mathbf{k}[z]z^2$, $\mathsf{M} = Az$ and $\mathsf{N} = A$. Then $\operatorname{Hom}_{\mathcal{C}}(M, N) \simeq \mathbf{k}$, $\operatorname{Hom}_{A}(\mathsf{M}, \mathsf{N}) \simeq A/Az$ and $\operatorname{Hom}_{A}(\mathsf{M}, \mathsf{N}) \to \operatorname{Hom}_{\mathcal{C}}(M, N)$ vanishes.

Here is another example where M and N are A-flat. Let $B = \mathbf{k}[t]/\mathbf{k}[t]t^2$ with deg(t) = 1 and $\mathcal{C} = B$ -gmod and $A = \mathbf{k}[z]/\mathbf{k}[z]z^2$. For $c \in \mathbf{k}^{\times}$, let $\mathsf{X}_c = \mathbf{k}[t]/\mathbf{k}[t]t^2 \in \mathcal{C}$ where $z \in \operatorname{End}(\mathsf{X}_c)$ is given by $\mathsf{X}_c \ni x \mapsto ctx \in \mathsf{X}_c$. Then X_c is A-flat. If $\mathsf{M} = \mathsf{X}_{c_1}$, $\mathsf{N} = \mathsf{X}_{c_2}$ for $c_1 \neq c_2$, then $\operatorname{Hom}_{\mathcal{C}}(M, N) \simeq \mathbf{k}$, $\operatorname{Hom}_A(\mathsf{M}, \mathsf{N}) \simeq Az$ and $\operatorname{Hom}_A(\mathsf{M}, \mathsf{N}) \to \operatorname{Hom}_{\mathcal{C}}(M, N)$ vanishes.

Lemma 3.10. We have equivalences of categories

A-gmod $\xrightarrow{\sim} Mod(A, \mathbf{k}$ -gmod),

 $\operatorname{Modg}_{\operatorname{coh}}(A) \xrightarrow{\sim} \operatorname{Pro}_{\operatorname{coh}}(A, \mathbf{k}\operatorname{-gmod}).$

Proof. The first equivalence is obvious. We define the functor $\Phi: \operatorname{Modg}_{\operatorname{coh}}(A) \to \operatorname{Pro}_{\operatorname{coh}}(A, \mathbf{k}\operatorname{-gmod})$ by $M \to \underset{m}{\overset{\text{im}}{\longleftarrow}} M/M_{\geq m}$, and $\Psi: \operatorname{Pro}_{\operatorname{coh}}(A, \mathbf{k}\operatorname{-gmod}) \to \operatorname{Modg}_{\operatorname{coh}}(A)$ by $\mathsf{M} \to \operatorname{Hom}_A(\Phi(A), \mathsf{M})$. It is easy to check that they are well-defined and quasi-inverse to each other. \Box

3.2. Flatness.

Proposition 3.11. Let A be a commutative graded ring satisfying (3.2), let $z \in A_{>0}$ be a non-zero-divisor, and set B = A/Az. Assume that $M \in Pro_{coh}(A, C)$ satisfies

(a) $z|_{\mathsf{M}}$ is a monomorphism,

(b) M/zM is *B*-flat.

Then M is A-flat.

Proof. It is enough to show that $\operatorname{Tor}_1^A(S, \mathsf{M}) \simeq 0$ for any finitely generated A-module S.

(i) Assume that zS = 0. Then we have $\operatorname{Tor}_1^A(S, \mathsf{M}) \simeq \operatorname{Tor}_1^B(S, \mathsf{M}/z\mathsf{M}) \simeq 0$.

(ii) Assume that $z^m S = 0$ for some $m \in \mathbb{Z}_{>0}$. We shall show $\operatorname{Tor}_1^A(S, \mathsf{M}) \simeq 0$ by induction on m. We have an exact sequence $0 \to \operatorname{Ker}(z|_S) \to S \to zS \to 0$. Since $z^{m-1}|_{zS} = 0$, we have $\operatorname{Tor}_1^A(zS, \mathsf{M}) \simeq 0$ by the induction hypothesis, and $\operatorname{Tor}_1^A(\operatorname{Ker}(z|_S), \mathsf{M}) \simeq 0$ by (i). Hence the exact sequence

$$\operatorname{Tor}_{1}^{A}(\operatorname{Ker}(z|_{S}), \mathsf{M}) \longrightarrow \operatorname{Tor}_{1}^{A}(S, \mathsf{M}) \longrightarrow \operatorname{Tor}_{1}^{A}(zS, \mathsf{M})$$

implies $\operatorname{Tor}_1^A(S, \mathsf{M}) \simeq 0$.

(iii) Assume that $z|_S$ is a monomorphism. Then we have an exact sequence

 $\operatorname{Tor}_1^A(S,\,\mathsf{M}) \xrightarrow{z} \operatorname{Tor}_1^A(S,\,\mathsf{M}) \longrightarrow \operatorname{Tor}_1^A(S/zS,\,\mathsf{M}).$

We have $\operatorname{Tor}_1^A(S/zS, \mathsf{M}) \simeq 0$ by (i). Hence $\operatorname{Tor}_1^A(S, \mathsf{M})/z \operatorname{Tor}_1^A(S, \mathsf{M}) \simeq 0$. Since $\operatorname{Tor}_1^A(S, \mathsf{M}) \in \operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$, Proposition 3.3 (iii) implies $\operatorname{Tor}_1^A(S, \mathsf{M}) \simeq 0$.

(iv) In general we have an exact sequence $0 \to S' \to S \to S'' \to 0$ such that $z^m S' = 0$ for some $m \in \mathbb{Z}_{>0}$, and $z|_{S''}$ is a monomorphism. Hence we have $\operatorname{Tor}_1^A(S, \mathsf{M}) \simeq 0$ by (ii) and (iii).

Corollary 3.12. Let z_k be a homogeneous indeterminate with positive degree (k = 1, ..., n), and let $A := \mathbf{k}[z_1, ..., z_n]$ be the graded polynomial ring. Assume that $\mathsf{M} \in \operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$ satisfies

 (z_1, \ldots, z_n) is M-regular, i.e., for any k such that $1 \leq k \leq n$, $z_k \mid_{\mathsf{M}/\sum_{i=1}^{k-1} z_j \mathsf{M}}$ is a monomorphism.

Then M is A-flat.

Proof. This immediately follows from Proposition 3.11 by induction on n.

Lemma 3.13. Let $m \in \mathbb{Z}_{>0}$ and $A = \mathbf{k}[z]/\mathbf{k}[z]z^m$, and let M be a graded A-module in an abelian \mathbf{k} -linear graded category. Then the following conditions are equivalent: (a) M is A-flat,

(b) $z: z^{k-1}M/z^k M \to z^k M/z^{k+1}M$ is an isomorphism for any k such that $1 \leq k < m$, (c) $z^k: M/zM \to z^k M/z^{k+1}M$ is an isomorphism for any k such that $1 \leq k < m$, (d) $\operatorname{Ker}(z^k|_M) \subset z^{m-k}M$ for any k such that $1 \leq k < m$, (e) $\operatorname{Ker}(z^k|_M) \subset z^{m-k}M$ for some k such that $1 \leq k < m$.

Proof. It follows from the fact that any graded ideal of A is $z^k A$ for some $k \in \mathbb{Z}$ with $0 \leq k \leq m$.

Proposition 3.14. Let A be a commutative graded **k**-algebra satisfying (3.2) and let $M \in Pro_{coh}(A, C)$.

(i) If the sequence

$$0 \to X' \otimes_A \mathsf{M} \to X \otimes_A \mathsf{M} \to X'' \otimes_A \mathsf{M} \to 0$$

is exact for any exact sequence $0 \to X' \to X \to X'' \to 0$ in A-gmod, then M is A-flat.

(ii) M is A-flat if and only if $\operatorname{Tor}_1^A(A/A_{>0}, \mathsf{M}) \simeq 0$.

Proof. (i) Let $0 \to X' \to X \to X'' \to 0$ be an exact sequence in $\operatorname{Modg}_{\operatorname{coh}}(A)$. We shall show that $0 \to X' \otimes_A \mathsf{M} \to X \otimes_A \mathsf{M} \to X'' \otimes_A \mathsf{M} \to 0$ is exact.

Let $0 \to X'_m \to X_m \to X''_m \to 0$ be the exact sequence of A-modules with finite **k**-dimension where $X_m = X/A_{\ge m}X$, $X''_m = X''/A_{\ge m}X''$ and $X'_m = X'/(X' \cap A_{\ge m}X)$. By Proposition 3.3 (iiib), there exists $n \in \mathbb{Z}$ such that $A_{\ge n}X' \subset X' \cap A_{\ge m}X$, which implies that

$$\underset{m}{\overset{\text{``}}{\varprojlim}} X'/A_{\geqslant m}X' \simeq \underset{m}{\overset{\text{``}}{\varprojlim}} X'_{m}$$

Hence

$$\begin{split} 0 &\to \underset{m}{\overset{\underset{m}{\leftarrow}}{\underset{m}{\underset{m}{\longleftarrow}}}} (X'/A_{\geqslant m}X') \otimes_A \mathsf{M} \to \underset{m}{\overset{\underset{m}{\leftarrow}}{\underset{m}{\underset{m}{\longleftarrow}}} (X/A_{\geqslant m}X) \otimes_A \mathsf{M} \to \underset{m}{\overset{\underset{m}{\leftarrow}}{\underset{m}{\underset{m}{\longleftarrow}}} (X'/A_{\geqslant m}X'') \otimes_A \mathsf{M} \to 0 \\ \text{is exact. Since } \underset{m}{\overset{\underset{m}{\leftarrow}}{\underset{m}{\underset{m}{\longleftarrow}}} (X/A_{\geqslant m}X) \otimes_A \mathsf{M} \simeq \underset{m}{\overset{\underset{m}{\leftarrow}}{\underset{m}{\underset{m}{\longleftarrow}}} X \otimes_A (A/A_{\geqslant m}) \otimes_A \mathsf{M} \simeq X \otimes_A \mathsf{M}, \text{ etc.}, \\ \text{we obtain the desired result.} \end{split}$$

(ii) Assume that $\operatorname{Tor}_1^A(A/A_{>0}, \mathsf{M}) \simeq 0$. We shall show that $\operatorname{Tor}_1^A(X, \mathsf{M}) \simeq 0$ for any $X \in A$ -gmod. There exists $m \in \mathbb{Z}_{\geq 0}$ such that $A_{\geq m}X = 0$. We shall argue by induction

on *m*. By the exact sequence $\operatorname{Tor}_1^A(A_{>0}X, \mathsf{M}) \to \operatorname{Tor}_1^A(X, \mathsf{M}) \to \operatorname{Tor}_1^A(X/A_{>0}X, \mathsf{M})$, the left term vanishes by the induction hypothesis, and the right term vanishes since $X/A_{>0}X$ is a direct sum of copies of $A/A_{>0}$.

Now for any exact sequence $0 \to X' \to X \to X'' \to 0$ in A-gmod, we obtain the exact sequence

$$0 \simeq \operatorname{Tor}_{1}^{A}(X'', \mathsf{M}) \longrightarrow X' \otimes_{A} \mathsf{M} \longrightarrow X \otimes_{A} \mathsf{M} \longrightarrow X'' \otimes_{A} \mathsf{M} \longrightarrow 0,$$

as desired.

4. DUALITY

4.1. **Complements.** In this subsection, we give two lemmas. Note that the graded versions of these lemma still hold, although we do not repeat them.

Lemma 4.1. Let B be a commutative noetherian ring and let \mathcal{A} be an abelian category. Let X be a B-flat module in \mathcal{A} , and let M and N are finitely generated B-modules.

$$\mathcal{H}om_B(N, M \otimes_B X) \simeq \operatorname{Hom}_B(N, M) \otimes_B X$$

Proof. We have a morphism

$$\operatorname{Hom}_B(N,M) \otimes_B X \to \mathcal{H}om_B(N,M \otimes_B X).$$

There exists an exact sequence of B-modules

$$L \to L' \to N \to 0,$$

where L and L' are finitely generated free B-modules. Then we have an exact sequence

$$0 \longrightarrow \operatorname{Hom}_B(N, M) \longrightarrow \operatorname{Hom}_B(L', M) \longrightarrow \operatorname{Hom}_B(L, M)$$

In the commutative diagram with exact rows

the middle and right vertical arrows are isomorphism, and hence the left one is also an isomorphism. $\hfill \Box$

Lemma 4.2. Let B be a commutative noetherian ring and let \mathcal{A} and \mathcal{A}' be abelian categories, If $D: \mathcal{A}^{opp} \to \mathcal{A}'$ is an equivalence of categories, then we have

$$D(\mathcal{H}om_B(M, X)) \simeq DX \otimes_B M$$

for any $X \in Mod(B, \mathcal{A})$ and a finitely generated B-module M.

Proof. For any $Y \in \mathcal{A}'$, we have

$$\operatorname{Hom}_{\mathcal{A}'}(\mathrm{D}X \otimes_{B} M, Y) \simeq \operatorname{Hom}_{B}(M, \operatorname{Hom}_{\mathcal{A}'}(\mathrm{D}X, Y))$$
$$\simeq \operatorname{Hom}_{B}(M, \operatorname{Hom}_{\mathcal{A}}(\mathrm{D}^{-1}Y, X))$$
$$\simeq \operatorname{Hom}_{\mathcal{A}}(\mathrm{D}^{-1}Y, \mathcal{H}om_{B}(M, X))$$
$$\simeq \operatorname{Hom}_{\mathcal{A}'}\left(\mathrm{D}\left(\mathcal{H}om_{B}(M, X)\right), Y\right).$$

4.2. **Duality.** Let C and C' be k-linear abelian categories which satisfy (3.1). Let $D: C^{\text{opp}} \to C'$ be an equivalence of categories.

Let A be a commutative graded **k**-algebra which satisfies (3.2). For a graded A-module X, we wet $X^* := \text{Hom}_{\mathbf{k}}(X, \mathbf{k})$ which is a graded A-module.

Lemma 4.3. Let $M \in Modg(A, Pro(\mathcal{C}))$ which is A-flat, and let X and Y be finite \mathbf{k} -dimensional graded A-modules. Then, we have

$$D(X^* \otimes_A \mathsf{M}) \otimes_A Y \simeq D((X \otimes_A Y)^* \otimes_A \mathsf{M}).$$

Proof. We have

$$D(X^* \otimes_A \mathsf{M}) \otimes_A Y \simeq D(\mathcal{H}om_A(Y, X^* \otimes_A \mathsf{M})) \simeq D(\operatorname{Hom}_A(Y, X^*) \otimes_A \mathsf{M}).$$

It remains to remark that $\operatorname{Hom}_A(Y, X^*) \simeq (X \otimes_A Y)^*$.

Proposition 4.5. Assume that $M \in Pro_{coh}(A, C)$ is A-flat. Then we have:

- (i) $D_A(\mathsf{M})$ belongs to $\operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C}')$,
- (ii) $D_A(M)$ is A-flat,
- (iii) for any finite- \mathbf{k} -dimensional graded A-module X, we have

$$D_A(\mathsf{M}) \otimes_A X \simeq D(X^* \otimes_A \mathsf{M}),$$

(iv) $D_A^{-1}D_A(M) \simeq M$.

Proof. For any finite \mathbf{k} -dimensional A-module X, we have

$$D_{A}(\mathsf{M}) \otimes_{A} X \simeq \underset{m}{\overset{\text{``lim''}}{\longleftarrow}} D\left((A/A_{\geq m})^{*} \otimes_{A} \mathsf{M} \right) \otimes_{A} X$$
$$\simeq \underset{m}{\overset{\text{``lim''}}{\longleftarrow}} D\left(\left((A/A_{\geq m}) \otimes_{A} X \right)^{*} \otimes_{A} \mathsf{M} \right) \simeq D(X^{*} \otimes_{A} \mathsf{M})$$

by Lemma 4.3. Hence $D_A(\mathsf{M}) \otimes_A X \in \mathcal{C}'$ and

Hence we have $D_A(\mathsf{M}) \in \operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C}')$. Since the functor $A\operatorname{-gmod} \to \operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C}')$ given by $X \mapsto D_A(\mathsf{M}) \otimes_A X \simeq D(X^* \otimes_A \mathsf{M})$ is exact, the object $D_A(\mathsf{M})$ is A-flat by Proposition 3.14.

Finally we have

$$D_A^{-1}D_A(\mathsf{M}) \simeq \underset{m}{\overset{\text{``lim''}}{\underset{m}{\longrightarrow}}} D^{-1}((A/A_{\geq m})^* \otimes_A D_A(\mathsf{M}))$$
$$\simeq \underset{m}{\overset{\text{``lim''}}{\underset{m}{\longrightarrow}}} D^{-1} \circ D((A/A_{\geq m}) \otimes_A \mathsf{M})) \simeq \mathsf{M}.$$

Definition 4.6. Let us denote by $\operatorname{Aff}_A(\mathcal{C})$ the full subcategory of $\operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$ consisting of A-flat objects.

Then by Proposition 4.5 the functor

$$D_A: Aff_A(\mathcal{C})^{opp} \xrightarrow{\sim} Aff_A(\mathcal{C}')$$

is an equivalence of categories.

4.3. Affine objects.

Definition 4.7. Let z be an indeterminate of homogeneous degree $d \in \mathbb{Z}_{>0}$. An object of $\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z], \mathcal{C})$ is nothing but a pair (M, z) such that

- (a) $\mathsf{M} \in \operatorname{Pro}(\mathcal{C})$ and $z \in \operatorname{End}_{\operatorname{Pro}_{\operatorname{coh}}(\mathcal{C})}(\mathsf{M})_d$,
- (b) $M/zM \in C$,
- (c) $\mathsf{M} \xrightarrow{\sim} ``\varprojlim_n `` \mathsf{M}/z^n \mathsf{M}.$
- If (M, z) satisfies further the following condition
- (d) $z \in \text{End}_{\text{Pro}_{coh}(\mathcal{C})}(\mathsf{M})$ is a monomorphism,
- then we say that (M, z) is an *affine object* (of M/zM).

Note that an affine object (M, z) is nothing but a $\mathbf{k}[z]$ -flat object in $\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z], \mathcal{C})$. We denote by $\operatorname{Aff}_{z}(\mathcal{C})$ the category of affine objects.

Proposition 4.8. The full subcategory $\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z], \mathcal{C})$ of $\operatorname{Mod}(\mathbf{k}[z], \operatorname{Pro}(\mathcal{C}))$ has the following properties.

- (i) $\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z], \mathcal{C})$ is stable by taking subquotients and extensions,
- (ii) any object of $\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z], \mathcal{C})$ is noetherian,
- (iii) for any $(\mathsf{M}, z) \in \operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z], \mathcal{C})$, we have
 - (a) $z^m \mathsf{M}/z^n \mathsf{M} \in \mathcal{C}$ for any $m, n \in \mathbb{Z}$ such that $0 \leq m \leq n$,
 - (b) Ker $z^n \in \mathcal{C}$ for any $n \in \mathbb{Z}_{\geq 0}$, and the sequence $\{\text{Ker}(z^n)\}_{n \in \mathbb{Z}_{\geq 0}}$ is stationary,
 - (c) for any $N \in Mod(\mathbf{k}[z], Pro(\mathcal{C}))$ such that $N \subset M$, there exists n > 0 such that $N \cap z^n M \subset zN$,
 - (d) if $\mathbf{M} = z \mathbf{M}$, then $\mathbf{M} = 0$.

This is nothing but a special case of Proposition 3.3.

For $M \in \operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z], \mathcal{C})$, we set

$$\mathsf{M}_{\mathrm{tor}} := \bigcup_{n \in \mathbb{Z}_{>0}} \mathrm{Ker}(z^n|_{\mathsf{M}}) \subset \mathsf{M} \quad \mathrm{and} \quad \mathsf{M}_{\mathrm{flat}} = \mathsf{M}/\mathsf{M}_{\mathrm{tor}}.$$

Note that M_{tor} is well-defined since M is notherian, and M_{flat} is an affine object.

Definition 4.9. A truncated affine object of $M \in C$ at $m \in \mathbb{Z}_{>0}$ (of degree d) is a pair (M, z) of an object M of C and an endomorphism z of M such that

- (a) z is homogeneous of degree $d \in \mathbb{Z}_{>0}$,
- (b) $z^m = 0$,
- (c) $\mathbf{M}/z\mathbf{M}\simeq M$,

(d) $z: z^{k-1} \mathsf{M}/z^k \mathsf{M} \longrightarrow z^k \mathsf{M}/z^{k+1} \mathsf{M}$ is an isomorphism if $1 \leq k \leq m-1$.

We sometimes say that M is an *m*-truncated affine object.

Note that an *m*-truncated affine object is nothing but a $(\mathbf{k}[z]/\mathbf{k}[z]z^m)$ -flat object of $\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z]/\mathbf{k}[z]z^m, \mathcal{C})$ by Lemma 3.13.

If (M, z) is an affine object, then $M/z^m M$ is an *m*-truncated affine object.

Lemma 4.10. Let M be an object of $Pro_{coh}(\mathbf{k}[z], C)$ such that M := M/zM is a simple object of C. Then (M, z) is either an affine object or a truncated affine object.

Proof. Let us consider the epimorphism

$$f_k: \mathsf{M}/z\mathsf{M} \twoheadrightarrow z^k\mathsf{M}/z^{k+1}\mathsf{M}$$

given by z^k . If f_k is an isomorphism for every $k \in \mathbb{Z}_{\geq 0}$, then (M, z) is an affine object. Otherwise there exists k such that f_k is not an isomorphism. Take the smallest k among such k's. Then $f_k = 0$ and hence $z^k \mathsf{M} = z^{k+1} \mathsf{M}$, which implies that $z^k \mathsf{M} = 0$. Moreover $f_j \colon \mathsf{M}/z \mathsf{M} \to z^j \mathsf{M}/z^{j+1} \mathsf{M}$ is an isomorphism for j < k, since it is non-zero. Hence (M, z) is a k-truncated affine object.

Lemma 4.11. If (M, z) is an affine object of a simple object of \mathscr{C} , then we have

END
$$_{\mathbf{k}[z]}(\mathsf{M}) \simeq \mathbf{k}[z].$$

Proof. It immediately follows from Proposition 3.8.

Definition 4.12. We denote by $\operatorname{Raff}_z(\mathcal{C})$ the category with $\operatorname{Ob}(\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z], \mathcal{C}))$ as the set of objects and with the morphisms defined as follows. For $\mathsf{M}, \mathsf{N} \in \operatorname{Ob}(\operatorname{Raff}_z(\mathcal{C}))$,

$$\operatorname{Hom}_{\operatorname{Raff}_{z}(\mathcal{C})}(\mathsf{M},\mathsf{N}) = \lim_{\substack{k \in \mathbb{Z}_{\geqslant 0}}} \operatorname{Hom}_{\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z],\mathcal{C})} ((z^{k}\mathbf{k}[z]) \otimes_{\mathbf{k}[z]} \mathsf{M},\mathsf{N})$$
$$\simeq \lim_{\substack{k \in \mathbb{Z}_{\geqslant 0}}} \operatorname{Hom}_{\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z],\mathcal{C})}(\mathsf{M},\mathbf{k}[z]z^{-k} \otimes_{\mathbf{k}[z]} \mathsf{N}).$$

Note that we have

 $\operatorname{Hom}_{\operatorname{Raff}_z(\mathcal{C})}(\mathsf{M},\mathsf{N}) \simeq \mathbf{k}[z,z^{-1}] \otimes_{\mathbf{k}[z]} \operatorname{Hom}_{\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z],\mathcal{C})}(\mathsf{M},\mathsf{N}) \quad \text{for } \mathsf{M},\mathsf{N} \in \operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z],\mathcal{C}).$

Hence, any object of $\operatorname{Raff}_z(\mathcal{C})$ is a $\mathbf{k}[z, z^{-1}]$ -module, i.e., $z \in \operatorname{End}(\mathsf{M})$ is invertible for any $\mathsf{M} \in \operatorname{Raff}_z(\mathcal{C})$.

Lemma 4.13. For any $N \in \operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z], \mathcal{C}), N \to N_{\operatorname{flat}}$ is an isomorphism in $\operatorname{Raff}_z(\mathcal{C})$.

Proof. Since $z^m \mathsf{N}_{tor} \simeq 0$ for some $m \in \mathbb{Z}_{>0}$, the canonical morphism

$$\lim_{k \in \mathbb{Z}_{\geq 0}} \operatorname{Hom}_{\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z],\mathcal{C})}(\mathsf{M},\mathbf{k}[z]z^{-k} \otimes_{\mathbf{k}[z]} \mathsf{N}) \to \lim_{k \in \mathbb{Z}_{\geq 0}} \operatorname{Hom}_{\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z],\mathcal{C})}(\mathsf{M},\mathbf{k}[z]z^{-k} \otimes_{\mathbf{k}[z]} \mathsf{N}_{\operatorname{flat}})$$

is an isomorphism for any $\mathsf{M} \in \operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z], \mathcal{C})$.

Hence a canonical functor

$$\operatorname{Aff}_{z}(\mathcal{C}) \to \operatorname{Raff}_{z}(\mathcal{C})$$

is faithful and essentially surjective (i.e., any object of $\operatorname{Raff}_z(\mathcal{C})$ is isomorphic to the image of an object of $\operatorname{Aff}_z(\mathcal{C})$).

Let us denote by $K(\mathcal{C})$ the Grothendieck group of \mathcal{C} . It is a $\mathbb{Z}[q^{\pm 1}]$ -module, where q acts on $K(\mathcal{C})$ by $[M] \mapsto [qM]$. We write $K(\mathcal{C})|_{q=1}$ for $K(\mathcal{C})/(q-1)K(\mathcal{C})$.

Since the following lemma is classical in other contexts, we omit its proof (cf. [2, Section 2.3]).

Lemma 4.14. Let $M \in Aff_z(\mathcal{C})$.

- (i) Then, $[M/zM] \in K(\mathcal{C})|_{q=1}$ depends only on the isomorphism class of M in $\operatorname{Raff}_z(\mathcal{C})$.
- (ii) Let $\mathsf{M}', \mathsf{M}, \mathsf{M}'' \in \operatorname{Aff}_z(\mathcal{C})$. If there exists an exact sequence $0 \to \mathsf{M}' \to \mathsf{M} \to \mathsf{M}'' \to 0$ in $\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z], \mathcal{C})$, then we have

$$[\mathsf{M}/z\mathsf{M}] = [\mathsf{M}'/z\mathsf{M}'] + [\mathsf{M}''/z\mathsf{M}''] \quad in \ K(\mathcal{C})|_{q=1}.$$

Lemma 4.15. The category $\operatorname{Raff}_z(\mathcal{C})$ satisfies the following properties.

- (i) $\operatorname{Raff}_{z}(\mathcal{C})$ is abelian,
- (ii) every object of $\operatorname{Raff}_z(\mathcal{C})$ has finite length.

Proof. Since (i) is elementary, we omit its proof.

Let us show (ii). Since every object of $\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z], \mathcal{C})$ is noetherian, every object M of $\operatorname{Raff}_z(\mathcal{C})$ is noetherian. Let us show that any decreasing sequence $\{\mathsf{N}_j\}_{j\in\mathbb{Z}_{\geq 0}}$ of subobjects of M is stationary. It is represented by a decreasing sequence $\{\mathsf{N}_j\}_{j\in\mathbb{Z}_{\geq 0}}$ of subobjects of M in $\operatorname{Aff}_z(\mathcal{C})$. Hence the decreasing sequence $\ell(\mathsf{N}_j/z\mathsf{N}_j)$ is stationary. Then our assertion follows from

(4.1) Let L', $L \in \operatorname{Aff}_{z}(\mathcal{C})$ satisfies $L' \subset L$ in $\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z], \mathcal{C})$ and $\ell(L'/zL') = \ell(L/zL)$, then $L' \to L$ is an isomorphism in $\operatorname{Raff}_{z}(\mathcal{C})$,

which is a consequence of Lemma 4.14.

Definition 4.16. Let M be an affine object.

- (i) We say that $L \subset M$ is a strict affine subobject of an affine object if M/L is an affine object. In other words, $L \cap zM = zL$.
- (ii) We say that M is a rationally simple affine object if M is simple as an object of Raff_z(C).
- (iii) Let L be an affine object and let $M \to L$ be an epimorphism in $\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z], \mathcal{C})$. If L is a head of M in $\operatorname{Raff}_{z}(\mathcal{C})$, we say that L is a *rational head* of M.

Lemma 4.17. Let (M, z) be an affine object. If M/zM is a simple object, then M is a rationally simple affine object.

Proof. Let N be a non-zero affine object and let $M \to N$ be an epimorphism in $\operatorname{Raff}_{z}(\mathcal{C})$. We may assume that it is an epimorphism in $\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z], \mathcal{C})$, since any epimorphism in

 $\operatorname{Raff}_{z}(\mathcal{C})$ is represented by an epimorphism in $\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z], \mathcal{C})$. Then $\mathsf{M}/z\mathsf{M} \to \mathsf{N}/z\mathsf{N}$ is an epimorphism, and N/zN is non-zero by Proposition 3.3 (iiic), and hence it is an isomorphism. Hence $M \rightarrow N$ is an isomorphism by Lemma 3.5.

Remark 4.18. The converse of Lemma 4.17 is not true (see Proposition 6.29).

5. Coherent objects in Monoidal categories

5.1. Graded monoidal categories. In the sequel, let \mathcal{C} be an abelian k-linear graded monoidal category which satisfies

 $\begin{cases} (a) \ \mathcal{C} \text{ satisfies } (3.1), \\ (b) \otimes \text{ is } \mathbf{k}\text{-bilinear and bi-exact,} \end{cases}$

(5.1)
$$\langle (c) \rangle$$
 the unit object **1** is simple; in particular, END $_{\mathcal{C}}(1) \simeq \mathbf{k}$,

(d) \otimes commutes with the grading shift functor q, i.e., $q(X \otimes Y) \simeq (qX) \otimes Y \simeq X \otimes (qY)$,

- For generalities on monoidal categories, we refer the reader to [14, Chapter 4], [3].
 - By identifying q with the invertible central object $q\mathbf{1} \in \mathcal{C}$, we have $qX \simeq q \otimes X$.

The category $\operatorname{Pro}(\mathcal{C})$ has also a structure of monoidal category in which the tensor product \otimes is bi-exact.

Let A be a commutative graded **k**-algebra satisfying (3.2). The functor **k**-gmod \ni $V \mapsto V \otimes_{\mathbf{k}} \mathbf{1} \in \mathcal{C}$ extends to a fully faithful monoidal functor $\operatorname{Pro}(\mathbf{k}\operatorname{-gmod}) \to \operatorname{Pro}(\mathcal{C})$. In the sequel, we regard $Pro(\mathbf{k}$ -gmod), as well as \mathbf{k} -gmod, as a full subcategory of $\operatorname{Pro}(\mathcal{C})$. For example, A will be identified with " $\operatorname{\underline{\lim}}"(A/A_{\geq m}) \otimes_{\mathbf{k}} \mathbf{1} \in \operatorname{Pro}(\mathcal{C})$ (cf.

Lemma 3.10).

For $\mathsf{M}, \mathsf{N} \in \operatorname{Modg}(A, \operatorname{Pro}(\mathcal{C}))$, define $\mathsf{M} \otimes_A \mathsf{N} \in \operatorname{Modg}(A, \operatorname{Pro}(\mathcal{C}))$ by the universal property:

$$\begin{array}{l} \operatorname{Hom}_{\operatorname{Modg}(A,\operatorname{Pro}(\mathcal{C}))}(\mathsf{M}\otimes_{A}\mathsf{N},\mathsf{L}) \\ (5.2) & \simeq \{f \in \operatorname{Hom}_{\operatorname{Pro}(\mathcal{C})}(\mathsf{M}\otimes\mathsf{N},\mathsf{L}) \mid \\ & (a\operatorname{id}_{\mathsf{L}}) \circ f = f \circ (a\operatorname{id}_{\mathsf{M}} \otimes \operatorname{id}_{\mathsf{N}}) = f \circ (\operatorname{id}_{\mathsf{M}} \otimes a\operatorname{id}_{\mathsf{N}}) \quad \text{for any } a \in A \}. \end{array}$$

It is well-defined since A is a finitely generated **k**-algebra. Note that $Modg(A, Pro(\mathcal{C}))$ is a monoidal category with \otimes_A as its tensor product and $A \in Modg(A, Pro(\mathcal{C}))$ as its unit object. Note also that there is an epimorphism $\mathsf{M} \otimes \mathsf{N} \twoheadrightarrow \mathsf{M} \otimes_A \mathsf{N}$ in $Pro(\mathcal{C})$.

Lemma 5.1. The category $\operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$ has a structure of a monoidal category by \otimes_A . The unit object is $A = \underset{m}{\overset{\text{"lim}}{\longleftarrow}} (A/A_{\geq m}) \otimes_{\mathbf{k}} \mathbf{1} \in \operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C}).$

Proof. Let $\mathsf{M}, \mathsf{N} \in \operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$. For each $m \in \mathbb{Z}_{\geq 0}$ we have

$$\frac{\mathsf{M}\otimes_{A}\mathsf{N}}{A_{\geq m}(\mathsf{M}\otimes_{A}\mathsf{N})}\simeq \frac{\mathsf{M}}{A_{\geq m}\mathsf{M}}\otimes_{A}\frac{\mathsf{N}}{A_{\geq m}\mathsf{N}}\in \mathrm{Modg}(A,\mathcal{C}).$$

In particular, $(\mathsf{M} \otimes_A \mathsf{N})/A_{>0}(\mathsf{M} \otimes_A \mathsf{N}) \in \mathcal{C}$.

Since " \lim_{m} " is exact, we have

$$\underset{m}{``\varprojlim''} \left(\frac{\mathsf{M} \otimes_A \mathsf{N}}{A_{\geq m}(\mathsf{M} \otimes_A \mathsf{N})} \right) \simeq \underset{m}{``\varprojlim''} \left(\frac{\mathsf{M}}{A_{\geq m}\mathsf{M}} \otimes_A \frac{\mathsf{N}}{A_{\geq m}\mathsf{N}} \right) \simeq \mathsf{M} \otimes_A \mathsf{N}$$

Hence $\mathsf{M} \otimes_A \mathsf{N}$ belongs to $\operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$. Now it is immediate that $(\operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C}), \otimes_A)$ is a monoidal category.

Note that \otimes_A is a right exact bi-functor.

Lemma 5.2. Assume that $\dim_{\mathbf{k}} A < \infty$. Let ε denote the projection $A \twoheadrightarrow A/A_{\geq 0} \simeq \mathbf{k}$. (i)

is an exact sequence in Modg(A). Here

$$g_1(a \otimes b \otimes c) = ab \otimes c - a \otimes bc \text{ and } g_0(a \otimes b) = ab.$$

(ii) For $M, N \in Modg(A, C)$, the sequence

$$(5.4) A \otimes A_{>0} \otimes M \otimes N \xrightarrow{f_1(M,N)} A_{>0} \otimes M \otimes N \xrightarrow{f_0(M,N)} M \otimes N$$

is a complex, i.e., the composition vanishes. Here $f_1(M, N)$ is given by

$$A \otimes A_{>0} \ni a \otimes b \mapsto b \otimes a \operatorname{id}_M \otimes \operatorname{id}_N + (a - \varepsilon(a)) \otimes \operatorname{id}_M \otimes b \operatorname{id}_N - ab \otimes \operatorname{id}_M \otimes \operatorname{id}_N \\ \in \operatorname{Hom}_{\mathcal{C}}(M \otimes N, A \otimes M \otimes N),$$

and $f_0(M, N)$ is given by

$$A \ni a \mapsto a \operatorname{id}_M \otimes \operatorname{id}_N - \operatorname{id}_M \otimes a \operatorname{id}_N \in \operatorname{Hom}_{\mathcal{C}}(M \otimes N, M \otimes N).$$

(iii) If M is A-flat, then

is exact. Here $h_1(M)$ is given by $A \otimes A_0 \ni a \otimes b \mapsto b \otimes a \operatorname{id}_M - ab \otimes \operatorname{id}_M \in \operatorname{Hom}_{\mathcal{C}}(M, A_{>0} \otimes M)$, and $h_0(M)$ is the multiplication.

- (iv) $A_{>0} \otimes M \otimes N \xrightarrow{f_0(M,N)} M \otimes N \longrightarrow M \otimes_A N \longrightarrow 0$ is exact.
- (v) If M is A-flat, then (5.4) is exact.
- (vi) If M is A-flat, then $M \otimes_A \bullet : \operatorname{Modg}(A, \mathcal{C}) \to \operatorname{Modg}(A, \mathcal{C})$ is exact.

Proof. (i) follows from $a \otimes b = g_1(1 \otimes a \otimes b) + 1 \otimes g_0(a \otimes b)$.

(ii) is straightforward.

(iii) follows form (i) and $(5.5) \simeq (5.3) \otimes_A M$.

(iv) follows from the fact that $a \operatorname{id}_M \otimes \operatorname{id}_N - \operatorname{id}_M \otimes a \operatorname{id}_N = 0$ if $a \in A_0$.

(v) For $M, N \in Modg(A, C)$, set $S_2(M, N) = A \otimes A_{>0} \otimes M \otimes N$ and $S_1(M, N) = A_{>0} \otimes M \otimes N$.

Assume that M is A-flat. For any $N \in \text{Modg}(A, \mathcal{C})$, there exists $m \in \mathbb{Z}_{>0}$ such that $A_{\geq m}N = 0$. We shall show that (5.4) is exact by induction on m. Assume that m = 1. Then we have $f_k(M, N) = h_k(M) \otimes \text{id}_N$ (k = 0, 1). Hence, $(5.4) \simeq (5.5) \otimes N$ is exact.

Now assume that m > 1.

For an exact sequence $0 \to N' \to N \to N'' \to 0$ in Modg(A, C), we have a commutative diagram with exact rows:

Set $N' = A_{>0}N$ and N'' = N/N'. Then the left and the right column are exact by the induction hypothesis, and hence the middle arrow is exact. Thus the induction proceeds.

(vi) Assume that M is A-flat. It is enough to show that $M \otimes_A N' \to M \otimes_A N$ is a monomorphism for any exact sequence $0 \to N' \to N \to N'' \to 0$ in Modg(A, C). It follows from the fact that the commutative diagram (5.6) has exact rows and columns.

Proposition 5.3. Assume that $\mathsf{M} \in \operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$ is A-flat. Then the functor $\mathsf{M} \otimes_A \bullet$: $\operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C}) \to \operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$ is exact.

Proof. Let us show that

$$(5.7) 0 \to \mathsf{M} \otimes_A \mathsf{N}' \to \mathsf{M} \otimes_A \mathsf{N} \to \mathsf{M} \otimes_A \mathsf{N}'' \to 0$$

is exact if $0 \to \mathsf{N}' \to \mathsf{N} \to \mathsf{N}'' \to 0$ is an exact sequence in $\operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C})$. For $m \in \mathbb{Z}_{>0}$, set $A(m) = A/A_{\ge m}$, $M_m = \mathsf{M}/A_{\ge m}\mathsf{M}$, $N_m = \mathsf{N}/A_{\ge m}\mathsf{N}$, $N_m'' = \mathsf{N}''/A_{\ge m}\mathsf{N}''$ and $N_m' = \mathsf{N}''$

 $\mathsf{N}'/(\mathsf{N}' \cap A_{\geq m}\mathsf{N})$. Then dim $A(m) < \infty$ and $M_m \in \operatorname{Modg}(A(m), \mathcal{C})$ is A(m)-flat. Since $0 \to N'_m \to N_m \to N''_m \to 0$ is an exact sequence in $\operatorname{Modg}(A(m), \mathcal{C})$, the sequence

$$0 \to M_m \otimes_{A(m)} N'_m \to M_m \otimes_{A(m)} N_m \to M_m \otimes_{A(m)} N''_m \to 0$$

is exact by Lemma 5.2. Then taking " \varprojlim_m ", we conclude that (5.7) is exact.

Lemma 5.4. For $M, N \in Aff_A(\mathcal{C})$ (see Definition 4.6), we have $M \otimes_A N \in Aff_A(\mathcal{C})$. Namely $Aff_A(\mathcal{C})$ is a monoidal category.

Proof. Let $X' \to X \to X''$ be an exact sequence in $\operatorname{Mod}_{\operatorname{coh}}(A)$. Then $X' \otimes_A \mathsf{M} \to X \otimes_A \mathsf{M} \to X'' \otimes_A \mathsf{M}$ is exact since M is *A*-flat. Now, Proposition 5.3 implies that $X' \otimes_A \mathsf{M} \otimes_A \mathsf{N} \to X \otimes_A \mathsf{M} \otimes_A \mathsf{N} \to X'' \otimes_A \mathsf{M} \otimes_A \mathsf{N}$ is exact since N is *A*-flat. \Box

Proposition 5.5. Let A' be another commutative graded ring satisfying (3.2), and let $M \in \operatorname{Pro}_{\operatorname{coh}}(A, \mathcal{C}), M' \in \operatorname{Pro}_{\operatorname{coh}}(A', \mathcal{C}).$

- (i) $A \otimes_{\mathbf{k}} A'$ satisfies (3.2).
- (ii) $\mathsf{M} \otimes \mathsf{M}' \in \operatorname{Pro}_{\operatorname{coh}}(A \otimes_{\mathbf{k}} A', \mathcal{C}).$
- (iii) For $X \in Mod_{coh}(A)$, $X' \in Mod_{coh}(A')$ and $m \in \mathbb{Z}_{\geq 0}$, we have

$$\operatorname{Tor}_{m}^{A\otimes_{\mathbf{k}}A'}(X\otimes_{\mathbf{k}}X',\mathsf{M}\otimes\mathsf{M}')\simeq\bigoplus_{a,b\in\mathbb{Z}_{\geq 0},\ a+b=m}\operatorname{Tor}_{a}^{A}(X,\mathsf{M})\otimes\operatorname{Tor}_{b}^{A'}(X',\mathsf{M}').$$

(iv) If M is A-flat and M' is A'-flat, then $M \otimes M'$ is $(A \otimes_{\mathbf{k}} A')$ -flat.

Proof. (i), (ii) and (iii) are immediate. (iv) follows from Proposition 3.14 and

$$\operatorname{Tor}_{1}^{A \otimes_{\mathbf{k}} A'} \left((A \otimes_{\mathbf{k}} A') / (A \otimes_{\mathbf{k}} A')_{>0}, \mathsf{M} \otimes \mathsf{M}' \right) \\ \simeq \operatorname{Tor}_{1}^{A} (A / A_{>0}, \mathsf{M}) \otimes \left((A' / A'_{>0}) \otimes_{A} \mathsf{M}' \right) \\ \oplus \left((A / A_{>0}) \otimes_{A} \mathsf{M} \right) \otimes \operatorname{Tor}_{1}^{A'} (A' / A'_{>0}, \mathsf{M}').$$

9	o	
3	õ	

5.2. **Rigid case.** In this subsection, we assume that C satisfies (5.1) and also that C is a rigid monoidal category, i.e., every object of C has a right dual and a left dual.

We denote by \mathscr{D} the right duality functor. Hence its quasi-inverse \mathscr{D}^{-1} is a left duality functor. Let

$$\operatorname{ev}_M \colon M \otimes \mathscr{D}M \to \mathbf{1}$$
 and $\operatorname{coev}_M \colon \mathbf{1} \to \mathscr{D}M \otimes M$

be the evaluation morphism and the coevaluation morphism, respectively. Recall that for $\mathsf{M} \in \mathrm{Pro}_{\mathrm{coh}}(A, \mathcal{C})$, we define in Definition 4.4

(5.8)
$$\mathscr{D}_{A}(\mathsf{M}) := \underset{m}{\overset{\otimes}{\underset{m}{\vdash}}} \mathscr{D}\left((A/A_{\geq m})^{*} \otimes_{A} \mathsf{M}\right),$$
$$\mathscr{D}_{A}^{-1}(\mathsf{M}) := \underset{m}{\overset{\otimes}{\underset{m}{\vdash}}} \mathscr{D}^{-1}\left((A/A_{\geq m})^{*} \otimes_{A} \mathsf{M}\right).$$

If $\mathsf{M} \in \operatorname{Aff}_{A}(\mathcal{C})$, then $\mathscr{D}_{A}(\mathsf{M})$ and $\mathscr{D}_{A}^{-1}(\mathsf{M})$ also belong to $\operatorname{Aff}_{A}(\mathcal{C})$.

They are a quasi-inverse to each other.

Proposition 5.6. If \mathcal{C} is rigid, then the monoidal category $\operatorname{Aff}_A(\mathcal{C})$ is also rigid, and \mathscr{D}_A and \mathscr{D}_A^{-1} are a right dual and a left dual, respectively.

Proof. We shall show that $\mathscr{D}_A(\mathsf{M})$ is a right dual of $\mathsf{M} \in \operatorname{Aff}_A(\mathcal{C})$.

Let us first construct ev: $\mathsf{M} \otimes_A \mathscr{D}_A(\mathsf{M}) \to A \otimes_k \mathbf{1}$. For $m \in \mathbb{Z}_{>0}$, we have

$$((A/A_{\geq m})^* \otimes_A \mathsf{M}) \otimes \mathscr{D}((A/A_{\geq m})^* \otimes_A \mathsf{M}) \to \mathbf{1}.$$

Set $L_m = A/A_{\geq m}$. Then $\{L_m\}_{m \in \mathbb{Z}_{\geq 0}}$ is a projective system. By using $\mathbf{k} \to L_m \otimes_{\mathbf{k}} L_m^*$, we obtain

$$\begin{split} \mathsf{M} \otimes \mathscr{D} \left(L_m^* \otimes_A \mathsf{M} \right) &\to \left(L_m \otimes_{\mathbf{k}} L_m^* \right) \otimes_{\mathbf{k}} \mathsf{M} \otimes \mathscr{D} \left(L_m^* \otimes_A \mathsf{M} \right) \\ &\to L_m \otimes_{\mathbf{k}} \left(\left(L_m^* \otimes_A \mathsf{M} \right) \otimes \mathscr{D} \left(L_m^* \otimes_A \mathsf{M} \right) \right) \\ &\to L_m \otimes_{\mathbf{k}} \mathbf{1}. \end{split}$$

Since it is A-bilinear, we obtain

$$\mathsf{M} \otimes_A \mathscr{D}_A(\mathsf{M}) \longrightarrow \mathsf{M} \otimes_A \mathscr{D} \left(L_m^* \otimes_A \mathsf{M} \right) \to L_m \otimes_k \mathbf{1}$$

Taking " \varprojlim_{m} , we obtain

ev: $\mathsf{M} \otimes_A \mathscr{D}_A(\mathsf{M}) \to A \otimes_k \mathbf{1}$.

Now let us construct the coevaluation morphism

coev: $A \otimes_{\mathbf{k}} \mathbf{1} \to \mathscr{D}_A(\mathsf{M}) \otimes_A \mathsf{M}.$

For any $m \in \mathbb{Z}_{\geq 0}$, we have

$$f_m: \mathbf{1} \to \mathscr{D}(L_m^* \otimes_A \mathsf{M}) \otimes (L_m^* \otimes_A \mathsf{M}).$$

It satisfies the relation : $(a \otimes 1) \cdot f_m = (1 \otimes a) \cdot f_m$ for any $a \in A$. Hence f_m gives

$$\mathbf{1} \to \mathcal{H}om_{A \otimes_{\mathbf{k}} A} \Big(A, \mathscr{D} \big(L_m^* \otimes_A \mathsf{M} \big) \otimes \big(L_m^* \otimes_A \mathsf{M} \big) \Big).$$

Here, we regard A as the $A \otimes_{\mathbf{k}} A$ -module $(A \otimes_{\mathbf{k}} A) / \sum_{a \in A} (A \otimes_{\mathbf{k}} A)(a \otimes_{\mathbf{k}} 1 - 1 \otimes_{\mathbf{k}} a)$, and $\mathscr{D}(L_m^* \otimes_A \mathsf{M}) \otimes (L_m^* \otimes_A \mathsf{M}) \in \operatorname{Modg}(A \otimes_{\mathbf{k}} A, \mathcal{C})$.

On the other hand, we have

$$\mathscr{D}(L_m^* \otimes_A \mathsf{M}) \otimes (L_m^* \otimes_A \mathsf{M}) \simeq (L_m \otimes_A \mathscr{D}_A(\mathsf{M})) \otimes (L_m^* \otimes_A \mathsf{M})$$
$$\simeq (L_m \otimes_{\mathbf{k}} L_m^*) \otimes_{A \otimes_{\mathbf{k}} A} (\mathscr{D}_A(\mathsf{M}) \otimes \mathsf{M}).$$

Since $\mathscr{D}_A(\mathsf{M}) \otimes \mathsf{M}$ is $(A \otimes_{\mathbf{k}} A)$ -flat by Proposition 5.5, Lemma 4.1 implies that

$$\mathcal{H}om_{A\otimes_{\mathbf{k}}A}\Big(A,\mathscr{D}\big(L_m^*\otimes_A\mathsf{M}\big)\otimes\big(L_m^*\otimes_A\mathsf{M}\big)\Big)$$
$$\simeq \operatorname{Hom}_{A\otimes_{\mathbf{k}}A}(A,L_m\otimes_{\mathbf{k}}L_m^*)\otimes_{A\otimes_{\mathbf{k}}A}\big(\mathscr{D}_A(\mathsf{M})\otimes\mathsf{M}\big).$$

Since $\operatorname{Hom}_{A\otimes_{\mathbf{k}}}(A, L_m\otimes_{\mathbf{k}}L_m^*)\simeq \operatorname{Hom}_A(L_m, L_m)\simeq L_m$, we obtain

$$\operatorname{Hom}_{A\otimes_{\mathbf{k}}^{A}}(A, L_{m}\otimes_{\mathbf{k}}L_{m}^{*})\otimes_{A\otimes_{\mathbf{k}}^{A}}\left(\mathscr{D}_{A}(\mathsf{M})\otimes\mathsf{M}\right)$$
$$\simeq L_{m}\otimes_{A\otimes_{\mathbf{k}}^{A}}\left(\mathscr{D}_{A}(\mathsf{M})\otimes\mathsf{M}\right)\simeq L_{m}\otimes_{A}\left(\mathscr{D}_{A}(\mathsf{M})\otimes_{A}\mathsf{M}\right).$$

Thus we obtain $\mathbf{1} \to L_m \otimes_A (\mathscr{D}_A(\mathsf{M}) \otimes_A \mathsf{M})$. Taking the projective limit " \varprojlim_m ", we obtain $\mathbf{1} \to \mathscr{D}_A(\mathsf{M}) \otimes_A \mathsf{M}$ in $\operatorname{Pro}(\mathcal{C})$, which induces a morphism

$$\operatorname{coev}: A \otimes_{\mathbf{k}} \mathbf{1} \to \mathscr{D}_A(\mathsf{M}) \otimes_A \mathsf{M}$$

in $Modg(A, Pro(\mathcal{C}))$.

Note that $\mathsf{M} \xrightarrow{\operatorname{coev}} \mathsf{M} \otimes_A \mathscr{D}_A(\mathsf{M}) \otimes_A \mathsf{M} \xrightarrow{\operatorname{ev}} \mathsf{M}$ and $\mathscr{D}_A(\mathsf{M}) \xrightarrow{\operatorname{coev}} \mathscr{D}_A(\mathsf{M}) \otimes_A \mathsf{M} \otimes_A \mathscr{D}_A(\mathsf{M}) \xrightarrow{\operatorname{coev}} \mathscr{D}_A(\mathsf{M}) \otimes_A \mathsf{M} \otimes_A$

Lemma 5.7. For $X \in \mathcal{C}$ and $\mathsf{M} \in \mathrm{Aff}_A(\mathcal{C})$, we have

 $\mathscr{D}_{A}^{\pm 1}(\mathsf{M} \otimes X) \simeq \big(\mathscr{D}^{\pm 1}(X) \big) \otimes \big(\mathscr{D}_{A}^{\pm 1}(\mathsf{M}) \big) \quad and \quad \mathscr{D}_{A}^{\pm 1}(X \otimes \mathsf{M}) \simeq \big(\mathscr{D}_{A}^{\pm 1}(\mathsf{M}) \big) \otimes \big(\mathscr{D}^{\pm 1}(X) \big).$

Proof. Let us only show the first isomorphism since the proof of the other is similar. We have

$$\mathscr{D}_{A}^{\pm}(\mathsf{M}\otimes X) \simeq \underset{m}{\overset{\text{lim}}{\longrightarrow}} \mathscr{D}^{\pm}\left((A/A_{\geqslant m})^{*}\otimes_{A}(\mathsf{M}\otimes X)\right)$$
$$\simeq \underset{m}{\overset{\text{lim}}{\longrightarrow}} \mathscr{D}^{\pm}\left(\left((A/A_{\geqslant m})^{*}\otimes_{A}\mathsf{M}\right)\otimes X\right)$$
$$\simeq \underset{m}{\overset{\text{lim}}{\longrightarrow}} (\mathscr{D}^{\pm}X)\otimes \mathscr{D}^{\pm}\left((A/A_{\geqslant m})^{*}\otimes_{A}\mathsf{M}\right)$$
$$\simeq (\mathscr{D}^{\pm}X)\otimes (\mathscr{D}_{A}^{\pm}\mathsf{M}).$$

6. **R-MATRICES**

6.1. Affine objects. Let C be an abelian graded k-linear monoidal category which satisfies (5.1).

Recall that $\operatorname{Aff}_{z}(\mathcal{C})$ is a monoidal category with \otimes_{z} as a tensor product, where we write \otimes_{z} for $\otimes_{\mathbf{k}[z]}$.

Lemma 6.1. Let (M, z_M) be an affine object in C and let $(N, z_N) \in \operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z_N], C)$. Then for any homogeneous polynomial $f(z_M, z_N) \in \mathbf{k}[z_M, z_N]$ which is monic in z_M , the morphism $f(z_M, z_N)|_{M\otimes N}$ is a monomorphism.

Proof. It is enough to show that $f(z_{\mathsf{M}}, z_{\mathsf{N}})|_{\mathsf{M}\otimes(\mathsf{N}/z_{\mathsf{N}}^{m}\mathsf{N})}$ is a monomorphism for any $m \in \mathbb{Z}_{\geq 0}$. We argue by induction on m. If m = 0, then it is obvious. Assume that m > 0. Then, in the commutative diagram with exact rows

the left and the right arrows are monomorphisms and so is the middle.

Proposition 6.2. Let (M, z_M) and (N, z_N) be affine objects. Set $M = M/z_M M$ and $N = N/z_N N$.

(i) Assume that $\operatorname{Hom}_{\mathcal{C}}(M \otimes N, M \otimes N) = \mathbf{k} \operatorname{id}_{M \otimes N}$. Then we have

$$\operatorname{Hom}_{\mathbf{k}[z_{\mathsf{M}}, z_{\mathsf{N}}]}(\mathsf{M} \otimes \mathsf{N}, \mathsf{M} \otimes \mathsf{N}) = \mathbf{k}[z_{\mathsf{M}}, z_{\mathsf{N}}] \operatorname{id}_{\mathsf{M} \otimes \mathsf{N}}.$$

(ii) Assume that $\dim_{\mathbf{k}} \operatorname{Hom}_{\mathcal{C}}(M \otimes N, N \otimes M) = 1$ and $\operatorname{Hom}_{\mathbf{k}[z_{\mathsf{M}}, z_{\mathsf{N}}]}(\mathsf{M} \otimes \mathsf{N}, \mathsf{N} \otimes \mathsf{M}) \neq 0$. Then there exists $\operatorname{R}^{\operatorname{ren}}_{\mathsf{M},\mathsf{N}} \in \operatorname{Hom}_{\mathbf{k}[z_{\mathsf{M}}, z_{\mathsf{N}}]}(\mathsf{M} \otimes \mathsf{N}, \mathsf{N} \otimes \mathsf{M})$ such that

(6.1)
$$\operatorname{Hom}_{\mathbf{k}[z_{\mathsf{M}}, z_{\mathsf{N}}]}(\mathsf{M} \otimes \mathsf{N}, \mathsf{N} \otimes \mathsf{M}) = \mathbf{k}[z_{\mathsf{M}}, z_{\mathsf{N}}] \mathrm{R}_{\mathsf{M}, \mathsf{N}}^{\operatorname{ren}}$$

and $\mathbb{R}_{M,N}^{\text{ren}}|_{z_M=z_N=0} \in \operatorname{Hom}_{\mathcal{C}}(M \otimes N, N \otimes M)$ does not vanish. We call $\mathbb{R}_{M,N}^{\text{ren}}$ the renormalized R-matrix.

Proof. Note that $\mathsf{M} \otimes \mathsf{N} \in \operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z_{\mathsf{M}}, z_{\mathsf{N}}], \mathcal{C})$. Hence it immediately follows from Proposition 3.8.

Definition 6.3. If $M, N \in C$ satisfy dim Hom $(M \otimes N, N \otimes M) = 1$, then a non-zero morphism $\mathbf{r} \in \text{Hom}(M \otimes N, N \otimes M)$ is called the R-matrix between M and N and denoted by \mathbf{r}_{MN} . It is well-defined up to a constant multiple.

If \mathcal{C} is a rigid monoidal category, we write $\mathscr{D}_{aff}^{\pm 1}$ for $\mathscr{D}_{\mathbf{k}[z]}^{\pm 1}$, where \mathscr{D} is the right dual functor of \mathcal{C} (see Definition 4.4). Hence, Proposition 5.6 implies the following proposition.

Proposition 6.4. Assume that \mathcal{C} is rigid. Then $\operatorname{Aff}_z(\mathcal{C})$ is a rigid monoidal category with $\mathscr{D}_{aff}^{\pm 1}$ as their right and left duality functors.

6.2. Rational centers and affinizations. Let \mathcal{C} be a k-linear graded monoidal category which satisfies the following conditions:

(6.2) $\begin{cases} \mathcal{C} \text{ satisfies (5.1) and the following additional condition:} \\ (e) \ \mathcal{C} \text{ has a decomposition } \mathcal{C} = \bigoplus_{\lambda \in \Lambda} \mathcal{C}_{\lambda} \text{ compatible with a monoidal} \\ \text{ structure where } \Lambda \text{ is an abelian monoid, and } \mathbf{1} \in \mathcal{C}_{\lambda} \text{ with } \lambda = 0. \end{cases}$

Let z be a homogeneous indeterminate with degree $d \in \mathbb{Z}_{>0}$.

We have also bi-exact bifunctors:

 $\otimes : \mathcal{C} \times \operatorname{Raff}_{z}(\mathcal{C}) \to \operatorname{Raff}_{z}(\mathcal{C}) \quad \text{and} \quad \otimes : \operatorname{Raff}_{z}(\mathcal{C}) \times \mathcal{C} \to \operatorname{Raff}_{z}(\mathcal{C}).$


Definition 6.5. A rational center in \mathcal{C} is a triple $(\mathsf{M}, \phi, \mathsf{R}_{\mathsf{M}})$ of $\mathsf{M} \in \mathrm{Aff}_{z}(\mathcal{C})$, an additive map $\phi \colon \Lambda \to \mathbb{Z}$ and an isomorphism

$$\mathbf{R}_{\mathsf{M}}(X) \colon q^{\otimes \phi(\lambda)} \otimes \mathsf{M} \otimes X \xrightarrow{\sim} X \otimes \mathsf{M}$$

in $\operatorname{Raff}_z(\mathcal{C})$ functorial in $X \in \mathcal{C}_\lambda$ such that

$$q^{\otimes \phi(\lambda+\mu)} \otimes \mathsf{M} \otimes X \otimes Y \xrightarrow{\mathrm{R}_{\mathsf{M}}(X) \otimes Y} q^{\otimes \phi(\mu)} \otimes X \otimes \mathsf{M} \otimes Y \xrightarrow{}_{\mathrm{R}_{\mathsf{M}}(X \otimes Y)} q^{\otimes \phi(\mu)} \otimes X \otimes \mathsf{M} \otimes Y \xrightarrow{}_{\mathrm{R}_{\mathsf{M}}(X \otimes Y)} q^{\otimes \phi(\mu)} \otimes X \otimes \mathsf{M} \otimes Y$$

and



commute in $\operatorname{Raff}_z(\mathcal{C})$ for any $X \in \mathcal{C}_\lambda$ and $Y \in \mathcal{C}_\mu$ $(\lambda, \mu \in \Lambda)$.

Note that the commutativity of the bottom diagram is a consequence of the one of the top. In the sequel, we neglect grading shifts.

Lemma 6.6. Assume that C is a rigid monoidal category. Let (M, R_M) be a rational center. Then $(\mathscr{D}_{aff}^{\pm 1}(M), R_{\mathscr{D}_{aff}^{\pm 1}(M)})$ is a rational center. Here, $R_{\mathscr{D}_{aff}^{\pm 1}(M)}(X) := \mathscr{D}_{aff}^{\pm 1}(R_M(\mathscr{D}^{\mp 1}(X)))$.

Note that

$$\mathrm{R}_{\mathsf{M}}(\mathscr{D}^{\mp 1}(X)) \colon \mathsf{M} \otimes \mathscr{D}^{\mp 1}(X) \xrightarrow{\sim} \mathscr{D}^{\mp 1}(X) \otimes \mathsf{M}$$

and

where the vertical arrows follow from Lemma 5.7.

Definition 6.7. An *affinization* M of $M \in \mathcal{C}$ is an affine object $(\mathsf{M}, z_{\mathsf{M}})$ with a rational center $(\mathsf{M}, \mathsf{R}_{\mathsf{M}})$ and an isomorphism $\mathsf{M}/z_{\mathsf{M}}\mathsf{M} \simeq M$.

We sometimes simply write M for affinization if no confusion arises. The following lemma is an immediate consequence of Lemma 4.14.

Lemma 6.8. Let M and N be objects of C. Assume that M admits an affinization. Then we have

 $[M] \cdot [N] \equiv [N] \cdot [M] \mod (q-1)K(\mathcal{C}).$

Proposition 6.9. Let (M, R_M) be a rational center in C, and let $L \in C$. Assume that M and L do not vanish. Then there exist $m \in \mathbb{Z}$ and a morphism $R_{M,L}^{ren} \colon M \otimes L \to L \otimes M$ in $\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z], C)$ such that

(a) $\mathrm{R}_{\mathsf{M},L}^{\mathrm{ren}}$ is equal to $z^m \mathrm{R}_{\mathsf{M}}(L)$: $\mathsf{M} \otimes L \to L \otimes \mathsf{M}$ in $\mathrm{Raff}_z(\mathcal{C})$,

(b) $\operatorname{R}_{\mathsf{M},L}^{\operatorname{ren}}|_{z=0}$: $(\mathsf{M}/z\mathsf{M}) \otimes L \to L \otimes (\mathsf{M}/z\mathsf{M})$ does not vanish.

Moreover such an integer m and an $R_{M,L}^{ren}$ are unique.

Similarly, there exist $m \in \mathbb{Z}$ and a morphism $\mathbb{R}_{L,\mathsf{M}}^{\mathrm{ren}} \colon L \otimes \mathsf{M} \to \mathsf{M} \otimes L$ in $\operatorname{Pro}_{\mathrm{coh}}(\mathbf{k}[z], \mathcal{C})$ such that

(c) $\mathrm{R}_{L,\mathsf{M}}^{\mathrm{ren}}$ gives $z^m \mathrm{R}_{\mathsf{M}}(L)^{-1} \colon L \otimes \mathsf{M} \to \mathsf{M} \otimes L$ in $\mathrm{Raff}_z(\mathcal{C})$,

(d) $\operatorname{R}_{L,\mathsf{M}}^{\operatorname{ren}}|_{z=0} \colon L \otimes (\mathsf{M}/z\mathsf{M}) \to (\mathsf{M}/z\mathsf{M}) \otimes L$ does not vanish.

Proof. The assertions (a) and (b) follow by taking m the smallest integer such that $\mathrm{R}_{\mathsf{M}}(L)$ is represented by an $f \in \mathrm{Hom}_{\mathrm{Aff}_{z}(\mathcal{C})}(\mathsf{M} \otimes L, \mathbf{k}[z]z^{-m} \otimes_{\mathbf{k}[z]} (L \otimes \mathsf{M}))$. The proof for (c) and (d) are similar.

Theorem 6.10. Let (M, R_M) be an affinization of $M \in C$, and let (N, z_N) be an affine object of $N \in C$. Assume that dim $\operatorname{Hom}_{\mathcal{C}}(M \otimes N, N \otimes M) = 1$. Then there exist a homogeneous $f(z_M, z_N) \in (\mathbf{k}[z_M, z_M^{-1}])[[z_N]]$ and a morphism $\operatorname{R}_{M,N}^{\operatorname{ren}} \colon M \otimes N \to N \otimes M$ in $\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z_M, z_N], C)$ such that

- (a) $\operatorname{Hom}_{\mathbf{k}[z_{\mathsf{M}},z_{\mathsf{N}}]}(\mathsf{M}\otimes\mathsf{N},\mathsf{N}\otimes\mathsf{M}) = \mathbf{k}[z_{\mathsf{M}},z_{\mathsf{N}}]\operatorname{R}^{\operatorname{ren}}_{\mathsf{M},\mathsf{N}},$
- (b) as an element of $\operatorname{Hom}_{\operatorname{Raff}_{z}(\mathcal{C})}(\mathsf{M}\otimes(\mathsf{N}/z_{\mathsf{N}}{}^{k}\mathsf{N}), (\mathsf{N}/z_{\mathsf{N}}{}^{k}\mathsf{N})\otimes\mathsf{M})$, we have $\operatorname{R}_{\mathsf{M},\mathsf{N}}^{\operatorname{ren}}|_{\mathsf{N}/z_{\mathsf{N}}{}^{k}\mathsf{N}} = f(z_{\mathsf{M}}, z_{\mathsf{N}})\operatorname{R}_{\mathsf{M}}(\mathsf{N}/z_{\mathsf{N}}{}^{k}\mathsf{N})$ for any $k \in \mathbb{Z}_{>0}$,
- (c) $\mathbf{R}_{\mathsf{M},\mathsf{N}}^{\mathrm{ren}}\Big|_{z_{\mathsf{M}}=z_{\mathsf{N}}=0}$ does not vanish,
- (d) $f(z_{\mathsf{M}}, z_{\mathsf{N}})|_{z_{\mathsf{N}}=0}$ is a monomial of z_{M} .

Moreover such $R_{M,N}^{ren}$ and $f(z_M, z_N)$ are unique.

Proof. Take \mathbf{r} such that $\operatorname{Hom}_{\mathcal{C}}(M \otimes N, N \otimes M) = \mathbf{kr}$. Set $\lambda = \operatorname{deg}(\mathbf{r})$. By replacing $\operatorname{R}_{\mathsf{M}}$ with $cz_{\mathsf{M}}^{m}\operatorname{R}_{\mathsf{M}}$ for some $m \in \mathbb{Z}$ and $c \in \mathbf{k}^{\times}$, we may assume from the beginning that $\operatorname{R}_{\mathsf{M}}(N)$ is in $\operatorname{Hom}_{\mathbf{k}[z_{\mathsf{M}}]}(\mathsf{M} \otimes N, N \otimes \mathsf{M})$ and $\operatorname{R}_{\mathsf{M}}(N)|_{z_{\mathsf{M}}=0} = \mathbf{r}$. In particular, $\operatorname{R}_{\mathsf{M}}(N)$ has degree λ .


Set $d = \deg(z_{\mathsf{M}})$ and $d_{\mathsf{N}} = \deg(z_{\mathsf{N}})$. Let C be the ring of homogeneous functions in $\mathbf{k}[z_{\mathsf{M}}^{-1}, z_{\mathsf{N}}]$ with degree 0. For $k \in \mathbb{Z}_{\geq 0}$, let $C(k) = \mathbf{k} \cdot (z_{\mathsf{N}}^{k}/z_{\mathsf{M}}^{kd_{\mathsf{N}}/d})$ or 0 according that kd_{N}/d is an integer or not. Then we have $C = \bigoplus_{k \geq 0} C(k)$. Set $C(\leq k) = \bigoplus_{0 \leq j \leq k} C(j)$.

For $m \in \mathbb{Z}_{\geq 0}$, set $N_m = \mathbb{N}/z_{\mathbb{N}}^m \mathbb{N} \in \text{Mod}(\mathbf{k}[z_{\mathbb{N}}], \mathcal{C})$. Let us show the following statement by induction on m.

(6.3) for any $m \in \mathbb{Z}_{\geq 1}$, there exists $f_m(z_M, z_N) \in C(\leq m-1)$ such that $f_m(z_M, 0) = 1$ and $R_m := f_m(z_M, z_N) \mathbb{R}_M(N_m)$ is a morphism $R_m \colon \mathbb{M} \otimes N_m \to N_m \otimes \mathbb{M}$ in $\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z_M, z_N], \mathcal{C}).$

Since it is trivial for m = 1, assume that m > 1. Then by the induction hypothesis, we have $f_{m-1}(z_{\mathsf{M}}, z_{\mathsf{N}})$ and R_{m-1} as in (6.3). Then $f_{m-1}(z_{\mathsf{M}}, z_{\mathsf{N}}) \mathbb{R}_{\mathsf{M}}(N_m)$ is a morphism in $\operatorname{Raff}_z(\mathcal{C})$. Take the smallest integer $s \ge 0$ which satisfies the following condition: there exists $a \in C(m-1)$ such that $g(z_{\mathsf{M}}, z_{\mathsf{N}}) \mathbb{R}_{\mathsf{M}}(N_m)$: $\mathsf{M} \otimes N_m \to N_m \otimes z_{\mathsf{M}}^{-s} \mathsf{M}$ is a morphism in $\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z_{\mathsf{M}}], \mathcal{C})$, where $g(z_{\mathsf{M}}, z_{\mathsf{N}}) = f_{m-1}(z_{\mathsf{M}}, z_{\mathsf{N}}) + a$.

Then we have a commutative diagram in $\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z_{\mathsf{M}}], \mathcal{C})$



MH

 $(c\mathbf{r} \text{ is erased in the diagram})$

Assume that s > 0. Then the composition

$$\mathsf{M} \otimes N_m \xrightarrow{g \mathsf{R}_{\mathsf{M}}(N_m)} N_m \otimes z_{\mathsf{M}}^{-s} \mathsf{M} \to N_m \otimes (z_{\mathsf{M}}^{-s} \mathsf{M}/z_{\mathsf{M}}^{1-s} \mathsf{M})$$

factors through $z_{\mathsf{N}}{}^{m-1}N_m \otimes (z_{\mathsf{M}}{}^{-s}\mathsf{M}/z_{\mathsf{M}}{}^{1-s}\mathsf{M})$. Hence the resulting morphism $\mathsf{M} \otimes N_m \to z_{\mathsf{N}}{}^{m-1}N_m \otimes (z_{\mathsf{M}}{}^{-s}\mathsf{M}/z_{\mathsf{M}}{}^{1-s}\mathsf{M})$ induces a non-zero morphism $\varphi \colon M \otimes N \to \mathbb{C}$

 $q^{(m-1)d_{\mathsf{N}}-ds}N \otimes M$ of degree λ . Since $\operatorname{Hom}(M \otimes N, N \otimes M)$ is concentrated in degree λ , we have $s = (m-1)d_{\mathsf{N}}/d$. Take $c \in \mathbf{k}$ such that $\varphi = c\mathbf{r}$, Then $R_m := (g - cz_{\mathsf{N}}^{m-1}/z_{\mathsf{M}}^s)\operatorname{R}_{\mathsf{M}}(N_m)$ sends $\mathsf{M} \otimes N_m$ to $N_m \otimes z_{\mathsf{M}}^{1-s}\mathsf{M}$. It contradicts the choice of s.

Hence we have s = 0, and the condition (6.3) is satisfied by setting $f_m = g$. Since f_m in (6.3) is unique by Proposition 3.7 we have $f_m = f_m$, mod z_m^{m-1} as

Since f_m in (6.3) is unique by Proposition 3.7, we have $f_m \equiv f_{m-1} \mod z_N^{m-1}$, and we obtain the desired result.

Remark 6.11. In Theorem 6.10, $f(z_{\mathsf{M}}, z_{\mathsf{N}}) \in (\mathbf{k}[z_{\mathsf{M}}, z_{\mathsf{M}}^{-1}])[[z_{\mathsf{N}}]]$ cannot be weakened by $f(z_{\mathsf{M}}, z_{\mathsf{N}}) \in (\mathbf{k}[z_{\mathsf{M}}, z_{\mathsf{M}}^{-1}])[z_{\mathsf{N}}]$. We shall give two examples.

- (i) Let $\mathfrak{g} = \mathbf{k}t$ be the one-dimensional graded Lie algebra with $\deg(t) = 1$. Let \mathcal{C} be the monoidal category of finite-dimensional graded \mathfrak{g} -modules. Take $\mathsf{M} = \mathbf{k}[z]$ with the trivial action of t, namely $t|_{\mathsf{M}} = 0$. For $X \in \mathcal{C}$, we define $\mathsf{R}_{\mathsf{M}}(X)$: $\mathbf{k}[z] \otimes X \to X \otimes \mathbf{k}[z]$ by $a \otimes b \mapsto e^{t/z}(b \otimes a)$. Then $(\mathsf{M}, \mathsf{R}_{\mathsf{M}})$ is a rational center. Let $\mathsf{N} = \mathbf{k}[t]$ with $z_{\mathsf{N}} = t$. Then, $f(z_{\mathsf{M}}, z_{\mathsf{N}}) = e^{-t/z}$ and $\mathsf{R}_{\mathsf{M},\mathsf{N}}^{\mathrm{ren}}(a \otimes b) = b \otimes a$.
- (ii) Let R be a quiver Hecke algebra (see §7.1 for the definition and notations). Let $\mathcal{C} = R$ -gmod, $i \in I$, and z a homogeneous indeterminate with degree (α_i, α_i) , and $\varphi(w) \in \mathbf{k}[[w]]$ with $\varphi(0) \in \mathbf{k}^{\times}$. Let $\mathsf{M} = \mathbf{k}[z] \in \operatorname{Aff}_z(\mathcal{C})$. For $\beta \in \mathsf{Q}_+$ with $n = \operatorname{ht}(\beta)$ and $X \in R(\beta)$ -gmod, let $\operatorname{R}_{\mathsf{M}}(X) \colon \mathsf{M} \otimes X \to X \otimes \mathsf{M}$ be the morphism $\mathsf{M} \otimes X \simeq \mathbf{k}[z] \otimes_{\mathbf{k}} X \longrightarrow X \otimes \mathsf{M} \simeq \mathbf{k}[z] \otimes_{\mathbf{k}} X$ in $\operatorname{Raff}_z(\mathcal{C})$ given by

$$\sum_{\nu \in I^{\beta}} \prod_{1 \leq k \leq n ; \nu_{k}=i} \varphi(z^{-1}x_{k}e(\nu)).$$

Note that $R_{\mathsf{M}}(X) \in Z(R(\beta))[[z^{-1}]]$. Then $(\mathsf{M}, \mathsf{R}_{\mathsf{M}})$ is a rational center and $R_{\mathsf{M},\mathsf{N}}^{\mathrm{ren}} = \mathsf{id}_{\mathbf{k}[z]\otimes\mathsf{N}}$.

When $\deg(z_{\mathsf{M}}) = \deg(z_{\mathsf{N}})$, since $\mathbb{R}_{\mathsf{M},\mathsf{N}}^{\mathrm{ren}}$ commutes with z_{M} and z_{N} , it induces a morphism in $\operatorname{Aff}_{z}(\mathcal{C})$

(6.4)
$$\mathsf{M} \otimes_{z} \mathsf{N} \xrightarrow{\overline{\mathsf{R}}_{\mathsf{M},\mathsf{N}}^{\mathrm{ren}}} \mathsf{N} \otimes_{z} \mathsf{M}$$

which is denoted by $\overline{\mathbf{R}}_{M,N}^{\text{ren}}$. Here, z acts on M and N by z_{M} and z_{N} , respectively.

Recall that a simple object M in a monoidal abelian category is called *real* if $M \otimes M$ is simple.

Definition 6.12. We say that a simple object $M \in C$ is affreal if M is real and there is an affinization (M, z) of M. If $\deg(z) = d$, we say that M is affreal of degree d.

Proposition 6.13. Let (M, z) be an affinization of a real simple $M \in C$. Then, there exists $c \in \mathbf{k}^{\times}$ and $T \in \operatorname{Hom}_{\operatorname{Pro}(C)}(M \otimes M, M \otimes M)$ such that

$$c \mathbf{R}_{\mathsf{M},\mathsf{M}}^{\mathrm{ren}} - \mathsf{id}_{\mathsf{M}\otimes\mathsf{M}} = (z \otimes \mathsf{id}_{\mathsf{M}} - \mathsf{id}_{\mathsf{M}} \otimes z)T$$

Proof. By Lemma 5.4, $\mathsf{M} \otimes_z \mathsf{M}$ is an affine object such that $(\mathsf{M} \otimes_z \mathsf{M})/z(\mathsf{M} \otimes_z \mathsf{M}) \simeq M \otimes M$. Since $M \otimes M$ is simple, $\operatorname{End}_{\mathbf{k}[z]}(\mathsf{M} \otimes_z \mathsf{M}) = \mathbf{k}[z] \operatorname{id}_{\mathsf{M} \otimes_z \mathsf{M}}$ by Lemma 4.11. Note that $\overline{\mathsf{R}}_{\mathsf{M},\mathsf{M}}^{\operatorname{ren}} \in \operatorname{End}_{\mathbf{k}[z]}(\mathsf{M} \otimes_z \mathsf{M})$ so that $\overline{\mathsf{R}}_{\mathsf{M},\mathsf{M}}^{\operatorname{ren}} = cz^a$ for some $a \in \mathbb{Z}_{\geq 0}$ and $c \in \mathbf{k}^{\times}$. By Theorem 6.10 (c), we have $\overline{\mathsf{R}}_{\mathsf{M},\mathsf{M}}^{\operatorname{ren}}|_{z=0} \neq 0$ and hence a = 0. Thus $\overline{\mathsf{R}}_{\mathsf{M},\mathsf{M}}^{\operatorname{ren}} - c \operatorname{id}_{\mathsf{M} \otimes_z \mathsf{M}} = 0$, from which the assertion follows.

Definition 6.14. Let M and N be simple objects in C.

(i) Assume that dim $\operatorname{Hom}_{\mathcal{C}}(M \otimes N, N \otimes M) = \dim \operatorname{Hom}_{\mathcal{C}}(N \otimes M, M \otimes N) = 1$. Then define (see Definition 6.3)

(6.5)
$$\Lambda(M,N) := \deg(\mathbf{r}_{M,N}) \in \mathbb{Z},$$

(6.6)
$$\mathfrak{d}(M,N) := \frac{1}{2} \big(\Lambda(M,N) + \Lambda(N,M) \big) \in \frac{1}{2} \mathbb{Z}_{\geq 0}$$

(ii) Let $(\mathsf{M}, \mathsf{R}_{\mathsf{M}})$ and $(\mathsf{N}, \mathsf{R}_{\mathsf{N}})$ be affinization of M and N, respectively, and assume that dim END $_{\mathcal{C}}(M \otimes N) = 1$. Then define $\mathscr{D}(\mathsf{M}, \mathsf{N}) \in \mathbf{k}[z_{\mathsf{M}}, z_{\mathsf{N}}]$ by

(6.7)
$$\mathbf{R}_{\mathsf{N},\mathsf{M}}^{\mathrm{ren}} \circ \mathbf{R}_{\mathsf{M},\mathsf{N}}^{\mathrm{ren}} = \mathscr{D}(\mathsf{M},\mathsf{N}) \operatorname{id}_{\mathsf{M} \otimes \mathsf{N}}.$$

Note that $\mathscr{D}(\mathsf{M},\mathsf{N})$ exists by Proposition 6.2, and it is well-defined up to a constant multiple in \mathbf{k}^{\times} .

If moreover dim END $_{\mathcal{C}}(N \otimes M) = 1$, since $\mathbb{R}_{M,N}^{\text{ren}} \circ \mathbb{R}_{N,M}^{\text{ren}} \circ \mathbb{R}_{M,N}^{\text{ren}} = \mathscr{D}(\mathsf{M},\mathsf{N})\mathbb{R}_{M,N}^{\text{ren}} = \mathscr{D}(\mathsf{M},\mathsf{N})\mathbb{R}_{M,N}^{\text{ren}} = \mathscr{D}(\mathsf{N},\mathsf{M})\mathbb{R}_{M,N}^{\text{ren}}$, we have $\mathscr{D}(\mathsf{M},\mathsf{N}) = \mathscr{D}(\mathsf{N},\mathsf{M})$.

Note that $\mathfrak{d}(M, N)$ is an integer in the quiver Hecke algebra case (cf. [13, Proposition 2.5]). As for criterions for the dimension 1 of the hom spaces, see Proposition 6.19 and Corollary 6.21 below.

6.3. Quasi-rigid Axiom.

Definition 6.15. Let \mathcal{A} be a monoidal category. We say that \mathcal{A} is a *quasi-rigid* category if it satisfies:

- (a) \mathcal{A} is abelian and \otimes is bi-exact,
- (b) for any $L, M, N \in \mathcal{A}, X \subset L \otimes M$ and $Y \subset M \otimes N$ such that $X \otimes N \subset L \otimes Y \subset L \otimes M \otimes N$, there exists $K \subset M$ such that $X \subset L \otimes K$ and $K \otimes N \subset Y$,
- (c) for any $L, M, N \in \mathcal{A}, X \subset M \otimes N$ and $Y \subset L \otimes M$ such that $L \otimes X \subset Y \otimes N$, there exists $K \subset M$ such that $X \subset K \otimes N$ and $L \otimes K \subset Y$.

Lemma 6.16. Let C be a quasi-rigid monoidal category such that **1** is simple. Let M and N be objects of C. If $M \otimes N \simeq 0$, then $M \simeq 0$ or $N \simeq 0$.

Proof. Assume that $M \otimes N \simeq 0$. Set $X := M \otimes \mathbf{1} \subset M \otimes \mathbf{1}$ and $Y := 0 \subset \mathbf{1} \otimes N$. Then we have $X \otimes N \subset M \otimes Y \subset M \otimes \mathbf{1} \otimes N$. Hence there exists $K \subset \mathbf{1}$ such that $X \subset M \otimes K$ and $K \otimes N \subset Y$. Since $\mathbf{1}$ is simple, $K \simeq 0$ or $K \simeq \mathbf{1}$. If $K \simeq 0$, then $X \subset M \otimes K$ implies $M \simeq 0$. If $K \simeq \mathbf{1}$, then $K \otimes N \subset Y$ implies $N \simeq 0$.

Lemma 6.17. An abelian rigid monoidal category with bi-exact tensor product is quasirigid.

Proof. Since it is a well-known result and the proof is similar to the one in Lemma 6.24 below, we omit its proof. \Box

Conjecture 6.18. A quasi-rigid monoidal category which satisfies (5.1) is embedded into a rigid monoidal category.

Recall that a simple object M in an abelian monoidal category is called *real* if $M \otimes M$ is simple. In this subsection, we keep the assumption that C is a monoidal abelian category satisfying (6.2).

Recall that, for $M, N \in \mathcal{C}$, we denote by $M \nabla N$ the head of $M \otimes N$ and $M \Delta N$ the socle of $M \otimes N$.

Proposition 6.19. Assume that C is quasi-rigid. Let M be an affreal object of C and N a simple object of C. Then, $M \otimes N$ and $N \otimes M$ have simple heads and simple socles.

Moreover, dim Hom_C $(M \otimes N, N \otimes M)$ = dim Hom_C $(N \otimes M, M \otimes N)$ = 1 and

$$M \nabla N \simeq \operatorname{Im}(\mathbf{r}_{M,N}) \simeq N \Delta M \quad and \quad N \nabla M \simeq \operatorname{Im}(\mathbf{r}_{N,M}) \simeq M \Delta N$$

up to grading shifts.

Recall that $\mathbf{r}_{M,N}$ is an R-matrix between M and N, i.e., a generator of $\operatorname{Hom}_{\mathcal{C}}(M \otimes N, N \otimes M)$.

Proof. Since the same proofs in [6, Theorem 3.2] and [8, Proposition 3.2.9] work, we omit the proof. \Box

Proposition 6.20. Let M be an affreal object of C and N be a simple object of C. Then we have

- (i) $\Lambda(M, M \nabla N) = \Lambda(M, N)$ and $\Lambda(N \nabla M, M) = \Lambda(N, M)$,
- (ii) for any simple subquotient S of the radical $\operatorname{Ker}(M \otimes N \to M \nabla N)$, we have $\Lambda(M,S) < \Lambda(M,N)$,
- (iii) for any simple subquotient S of $(M \otimes N)/(M \Delta N)$, we have $\Lambda(S, M) < \Lambda(N, M)$,
- (iv) for any simple subquotient S of the radical $\operatorname{Ker}(N \otimes M \to N \nabla M)$, we have $\Lambda(S, M) < \Lambda(N, M)$,
- (v) for any simple subquotient S of $(N \otimes M)/(N \Delta M)$, we have $\Lambda(M, S) < \Lambda(M, N)$.

In particular, $M \nabla N$ appears in the composition series of $M \otimes N$ only once (up to a grading).

Proof. The proof is similar to [8, Theorem 4.1].

Corollary 6.21. Assume that $M \in \mathcal{C}$ is affreal and $N \in \mathcal{C}$ is simple. Then we have $\operatorname{End}_{\mathcal{C}}(M \otimes N) = \operatorname{kid}_{M \otimes N}$ and $\operatorname{End}_{\mathcal{C}}(N \otimes M) = \operatorname{kid}_{N \otimes M}$.

Proof. We shall only prove $\operatorname{End}_{\mathcal{C}}(M \otimes N) = \mathbf{k} \operatorname{id}_{M \otimes N}$, since the other assertion can be similarly proved. Let $f \in \operatorname{End}_{\mathcal{C}}(M \otimes N)$. Since $M \otimes N$ has a simple head, the composition $M \otimes N \xrightarrow{f} M \otimes N \to M \nabla N$ factors through $M \nabla N$. Then the resulting endomorphism of $M \nabla N$ should be $\operatorname{cid}_{M \nabla N}$ for some $c \in \mathbf{k}$. Replacing f with $f - \operatorname{cid}_{M \otimes N}$, we may assume that $f(M \otimes N) \subset \operatorname{Ker}(M \otimes N \to M \nabla N)$. If $f(M \otimes N)$ is

non-zero, then a simple quotient S of $f(M \otimes N)$ should be isomorphic to $M \nabla N$. Hence $\Lambda(M, S) = \Lambda(M, M \nabla N) = \Lambda(M, N)$, which contradicts Proposition 6.20 (ii).

Lemma 6.22. Let C be a quasi-rigid monoidal category. Let (M, R_M) be an affinization of a real simple $M \in C$, and let (N, z_N) be an affine object of a simple $N \in C$. Then there exist a homogeneous $f(z_M, z_N), g(z_M, z_N) \in (\mathbf{k}[z_M, z_M^{-1}])[[z_N]]$ and morphisms $R_{M,N}^{ren} \colon M \otimes N \to N \otimes M$ in $\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z_M, z_N], C)$ and $\widetilde{R}_{N,M}^{ren} \colon N \otimes M \to M \otimes N$ in $\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z_M, z_N], C)$ such that


- (a) $f(z_{\mathsf{M}}, z_{\mathsf{N}})|_{z_{\mathsf{N}}=0}$ and $g(z_{\mathsf{M}}, z_{\mathsf{N}})|_{z_{\mathsf{N}}=0}$ are monomials of z_{M} ,
- (b) for any $k \in \mathbb{Z}_{>0}$, we have

$$\mathbf{R}_{\mathsf{M},\mathsf{N}}^{\mathrm{ren}}|_{\mathsf{N}/z_{\mathsf{N}}^{k}\mathsf{N}} = f(z_{\mathsf{M}}, z_{\mathsf{N}})\mathbf{R}_{\mathsf{M}}(\mathsf{N}/z_{\mathsf{N}}^{k}\mathsf{N}) \text{ and } \mathbf{R}_{\mathsf{N},\mathsf{M}}^{\mathrm{ren}}|_{\mathsf{N}/z_{\mathsf{N}}^{k}\mathsf{N}} = g(z_{\mathsf{M}}, z_{\mathsf{N}})\mathbf{R}_{\mathsf{M}}(\mathsf{N}/z_{\mathsf{N}}^{k}\mathsf{N})^{-1},$$

(c) $\operatorname{R}_{\mathsf{M},\mathsf{N}}^{\operatorname{ren}}|_{z_{\mathsf{M}}=z_{\mathsf{N}}=0} \in \operatorname{Hom}_{\mathcal{C}}(M \otimes N, N \otimes M)$ and $\widetilde{\operatorname{R}}_{\mathsf{N},\mathsf{M}}^{\operatorname{ren}}|_{z_{\mathsf{M}}=z_{\mathsf{N}}=0} \in \operatorname{Hom}_{\mathcal{C}}(N \otimes M, M \otimes N)$ do not vanish.

Proof. By Proposition 6.19 we have dim $\operatorname{Hom}_{\mathcal{C}}(M \otimes N, N \otimes M) = 1$, Hence by Theorem 6.10 there exist $\operatorname{R}_{\mathsf{M},\mathsf{N}}^{\operatorname{ren}}$ and $f(z_{\mathsf{M}}, z_{\mathsf{N}})$ satisfying the properties in (a) and (b). The proof of the statement on $\widetilde{\operatorname{R}}_{\mathsf{N},\mathsf{M}}^{\operatorname{ren}}$ is similar.

Lemma 6.23. Assume that C is a quasi-rigid monoidal category. Let (L, R_L) , (M, R_M) and (N, R_N) be affinizations of real simple objects in C. Then the Yang-Baxter equation holds for $(R_{L,M}^{ren}, R_{M,N}^{ren}, R_{L,N}^{ren})$, namely, the following diagram in $Pro_{coh}(\mathbf{k}[z_L, z_M, z_N], C)$ commutes :



Proof. By Proposition 6.19 we have

 $\dim \operatorname{Hom}_{\mathcal{C}}(L \otimes M, M \otimes L) = \dim \operatorname{Hom}_{\mathcal{C}}(L \otimes N, N \otimes L) = \dim \operatorname{Hom}_{\mathcal{C}}(M \otimes N, N \otimes M) = 1.$

Then by Theorem 6.10, we have non-zero morphisms $R_{L,M}^{ren}$, $R_{M,N}^{ren}$, and $R_{L,N}^{ren}$. Since the Yang-Baxter equation holds for the triple $(R_L(M), R_M(N), R_L(N))$ and they are proportional to $R_{L,M}^{ren}$, $R_{M,N}^{ren}$, the assertion follows.

The following two statements are an analogue of results of [6] in the affinization case.

Lemma 6.24. Assume that C is an abelian rigid monoidal category with (5.1). Let (L, z), (M, z), (N, z) be affine objects in C. Let $X \subset L \otimes_z M$ and $Y \subset M \otimes_z N$ be affine objects. If $X \otimes_z N \subset L \otimes_z Y$, then there exists an affine object $Z \subset M$ such that

$$X \subset \mathsf{L} \otimes_{\mathsf{r}} Z$$
 and $Z \otimes_{\mathsf{r}} \mathsf{N} \subset Y$


If we further assume that Y is a strict affine subobject of $M \otimes_z N$, then we may assume that Z is a strict affine subobject of M.

Proof. Let Z be an object in $Pro_{coh}(\mathbf{k}[z], \mathcal{C})$ such that the diagram

is a cartesian square in $\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z], \mathcal{C})$. By Proposition 4.5 and Proposition 5.3, $\bullet \otimes_z \mathscr{D}_{\operatorname{aff}}^{-1}(\mathsf{N})$ is exact and hence the right vertical arrow is a monomorphism, which implies that the left vertical arrow is also a monomorphism. Hence Z is an affine object.

Since $Z \otimes_z \mathbb{N} \to \mathbb{M} \otimes_z \mathbb{N}$ is decomposed into $Z \otimes_z \mathbb{N} \to Y \to \mathbb{M} \otimes_z \mathbb{N}$, we get $Z \otimes_z \mathbb{N} \subset Y$.

By applying $L \otimes_{z} \bullet$, which is exact by Proposition 5.3, to the above square, we obtain the following commutative diagram in which the bottom square is cartesian:



Hence we have $X \to \mathsf{L} \otimes_z Z$ in $\operatorname{Aff}_z(\mathcal{C})$, as desired.

Assume further that Y is a strict affine subobject of $\mathsf{M} \otimes_z \mathsf{N}$. Take $Z' := \operatorname{Ker}(\mathsf{M} \to (\mathsf{M}/Z)_{\text{flat}})$. Then $Z \subset Z' \subset \mathsf{M}$ and $\mathsf{M}/Z' \simeq (\mathsf{M}/Z)_{\text{flat}} \in \operatorname{Aff}_z(\mathcal{C})$. Hence $X \subset \mathsf{L} \otimes_z Z'$ and Z' is a strict affine subobject of M. We have a morphism

$$\left((\mathsf{M}\otimes_{z}\mathsf{N})/(Z\otimes_{z}\mathsf{N})\right)_{\mathrm{flat}} \to \left((\mathsf{M}\otimes_{z}\mathsf{N})/Y\right)_{\mathrm{flat}}.$$

Since the left hand side is $(\mathsf{M} \otimes_z \mathsf{N})/(Z' \otimes_z \mathsf{N})$ and the right hand side is $(\mathsf{M} \otimes_z \mathsf{N})/Y$, we obtain $Z' \otimes_z \mathsf{N} \subset Y$.

Proposition 6.25. Assume that C is an abelian rigid monoidal category with (6.2). Let (M, z) be an affinization and let (N, z) be an affine object. Assume that M/zM is real simple and N/zN is simple. Let $\overline{R}_{M,N}^{ren} \colon M \otimes_z N \to N \otimes_z M$ be the renormalized *R*-matrix in (6.4). Then $\operatorname{Im}(\overline{R}_{M,N}^{ren})$ is a unique rationally simple quotient affine object of $M \otimes_z N$.

Proof. Let $K \subset \mathsf{M} \otimes_z \mathsf{N}$ be a strict affine subobject such that $K \neq \mathsf{M} \otimes_z \mathsf{N}$. It is enough to show that $\overline{\mathsf{R}}_{\mathsf{M},\mathsf{N}}^{\mathrm{ren}}(K) = 0$. Indeed, then $\operatorname{Ker}(\overline{\mathsf{R}}_{\mathsf{M},\mathsf{N}}^{\mathrm{ren}})$ is a unique maximal strict affine subobject of $\mathsf{M} \otimes_z \mathsf{N}$, and it remains to remark that the quotient of an affine object by a maximal strict affine subobject is rationally simple.

Consider the following commutative diagram

$$\begin{array}{c} \mathsf{M} \otimes_{z} K & \longrightarrow & K \otimes_{z} \mathsf{M} \\ \downarrow & & \downarrow \\ \mathsf{M} \otimes_{z} \mathsf{M} \otimes_{z} \mathsf{N} \xrightarrow[\overline{\mathsf{R}_{\mathsf{M},\mathsf{M}}}]{}^{\operatorname{ren}} & \mathsf{M} \otimes_{z} \mathsf{M} \otimes_{z} \mathsf{N} \xrightarrow[\overline{\mathsf{R}_{\mathsf{M},\mathsf{N}}}]{}^{\operatorname{ren}} & \mathsf{M} \otimes_{z} \mathsf{N} \otimes_{z} \mathsf{M}. \end{array}$$

Since $\overline{\mathbb{R}}_{\mathsf{M},\mathsf{M}}^{\operatorname{ren}} = \operatorname{id}_{\mathsf{M}\otimes_{z}\mathsf{M}}$ up to a constant multiple by Proposition 3.8, we have $\mathsf{M} \otimes_{z} \overline{\mathbb{R}}_{\mathsf{M},\mathsf{N}}^{\operatorname{ren}}(K) \subset K \otimes_{z} \mathsf{M}$. Hence, Lemma 6.24 implies that there exists a strict affine subobject Z of N such that $\overline{\mathbb{R}}_{\mathsf{M},\mathsf{N}}^{\operatorname{ren}}(K) \subset Z \otimes_{z} \mathsf{M}$ and $\mathsf{M} \otimes_{z} Z \subset K$. Since $K \neq \mathsf{M} \otimes_{z} \mathsf{N}$, we have $Z \neq \mathsf{N}$. Since N is rationally simple, we have Z = 0. Hence $\overline{\mathbb{R}}_{\mathsf{M},\mathsf{N}}^{\operatorname{ren}}(K) = 0$. \Box

Definition 6.26. Assume that (M, z) and (N, z) be affinizations of real simple objects. If $\operatorname{Im}(\overline{\mathrm{R}}_{M,N}^{\operatorname{ren}})$ is an affinization of $(M/zM) \nabla (N/zN)$, we denote $\operatorname{Im}(\overline{\mathrm{R}}_{M,N}^{\operatorname{ren}})$ by $M \nabla_z N$.

Lemma 6.27. Assume that C is an abelian rigid monoidal category with (6.2). Let (M, z) and (N, z) be affinizations of real simple modules M and N, respectively, and let (L, z) be an affinization of $M \nabla N$. Assume that there is an epimorphism

$$\mathsf{M}\otimes_{\sim}\mathsf{N}\twoheadrightarrow\mathsf{L}$$

in $\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z], \mathcal{C})$. Then, we have $\operatorname{Im}(\overline{\mathbb{R}}_{\mathsf{M},\mathsf{N}}^{\operatorname{ren}}) \simeq \mathsf{L}$.

Proof. It follows from Proposition 6.25.

Proposition 6.28. Assume that C is an abelian rigid monoidal category with (6.2). Let (M, z) be an affinization of a real simple object M. Then $M \otimes_z \mathscr{D}_{aff}(M) \to \mathbf{k}[z]$ is an epimorphism in $\operatorname{Pro}_{coh}(\mathbf{k}[z], C)$.

Proof. It follows from Proposition 3.3 (iiic) and the fact that $M \otimes \mathscr{D}M \to \mathbf{1}$ is an epimorphism.

Proposition 6.29. Assume that C is an abelian rigid monoidal category with (6.2). Let (M, z), (N, z) and (L, z) be an affinization with $\deg(z) = d$ of a real simple M, N and L in C, respectively. Assume that

(a) $\mathfrak{d}(M, N) > 0$,

(b) there exists an epimorphism $\mathsf{M} \otimes_z \mathsf{N} \twoheadrightarrow \mathsf{L}$ in $\mathrm{Aff}_z(\mathcal{C})$.

Then we have

(i) $\operatorname{Im}(\overline{\mathbb{R}}_{M,N}^{\operatorname{ren}}) \simeq \mathsf{L},$ (ii) $\mathscr{D}(\mathsf{M},\mathsf{N}) \in \mathbf{k}[z_{\mathsf{M}}, z_{\mathsf{N}}](z_{\mathsf{M}} - z_{\mathsf{N}}).$ Here $z_{\mathsf{M}} = z|_{\mathsf{M}}$ and $z_{\mathsf{N}} = z|_{\mathsf{N}}.$

Proof. (i) follows from Lemma 6.27.

(ii) Set $z = z_{\mathsf{M}}, w = z_{\mathsf{N}}$ and $f(z, w) = \mathscr{D}(\mathsf{M}, \mathsf{N})$. Assume that f(z, w) is a homogeneous function of degree r in z, w (counting the degrees of z, w as one), i.e., $\mathfrak{d}(M, N) = dr/2$. Then we have $\mathrm{R}_{\mathsf{M},\mathsf{N}}^{\mathrm{ren}} \colon \mathsf{M} \otimes \mathsf{N} \to \mathsf{N} \otimes \mathsf{M}$ and $\mathrm{R}_{\mathsf{N},\mathsf{M}}^{\mathrm{ren}} \colon \mathsf{N} \otimes \mathsf{M} \to \mathsf{M} \otimes \mathsf{N}$ such that

$$\mathbf{R}_{\mathsf{N},\mathsf{M}}^{\mathrm{ren}} \circ \mathbf{R}_{\mathsf{M},\mathsf{N}}^{\mathrm{ren}} = f(z,w) \operatorname{id}_{\mathsf{M}\otimes\mathsf{N}}.$$

Hence, the composition

$$\mathsf{M} \otimes_{z} \mathsf{N} \xrightarrow{\mathrm{R}^{\mathrm{ren}}_{\mathsf{M},\mathsf{N}}|_{z=w}} \mathsf{N} \otimes_{z} \mathsf{M} \xrightarrow{\mathrm{R}^{\mathrm{ren}}_{\mathsf{N},\mathsf{M}}|_{z=w}} \mathsf{M} \otimes_{z} \mathsf{N}$$

is $z^r f(1,1)$. Hence if $f(1,1) \neq 0$, then $\mathsf{M} \otimes_z \mathsf{N} \to \mathsf{N} \otimes_z \mathsf{M}$ is a monomorphism, and hence $\mathsf{M} \otimes_z \mathsf{N} \to \mathsf{L}$ is an isomorphism, which implies that $M \otimes N \simeq L$ is a simple object. Hence $M \otimes N \simeq N \otimes M$ and it contradicts $\mathfrak{d}(M,N) > 0$. Thus we obtain f(1,1) = 0, which implies that z - w is a factor of f(z,w).

Lemma 6.30. Assume that C is an abelian rigid monoidal category with (6.2). Let M, N be simple objects in C such that one of them is affreal. Then

$$\Lambda(M, \mathscr{D}N) = \Lambda(N, M).$$

Proof. This immediately follows from

$$\operatorname{Hom}(N \otimes M, M \otimes N) \simeq \operatorname{Hom}(M \otimes \mathscr{D}N, \mathscr{D}N \otimes M).$$

7. Quiver Hecke Algebras and Schur-Weyl duality functors

7.1. Quiver Hecke algebras. Let **k** be a field and let **C** be a symmetrizable generalized Cartan matrix. We fix a Cartan datum $(C, P, \Pi, \Pi^{\vee}, (\cdot, \cdot))$ consisting of **C** called a generalized Cartan matrix, **P** a free abelian group called the weight lattice, $\Pi = \{\alpha_i \mid i \in I\} \subset P$ called the set of simple roots, $\Pi^{\vee} = \{h_i \mid i \in I\} \subset P^{\vee} := \text{Hom}(P, \mathbb{Z})$ called the set of simple coroots, and (\cdot, \cdot) a Q-valued symmetric bilinear form on P satisfying the following conditions:

(a) C = (⟨h_i, α_j⟩)_{i,j∈I},
(b) (α_i, α_i) ∈ 2Z_{>0} for i ∈ I,
(c) ⟨h_i, λ⟩ = 2(α_i, λ)/(α_i, α_i) for i ∈ I and λ ∈ P,
(d) for each i ∈ I, there exists Λ_i ∈ P such that ⟨h_j, Λ_i⟩ = δ_{ij} for any j ∈ I.

We denote by $\mathfrak{g} := \mathfrak{g}(\mathsf{C})$ the corresponding symmetrizable Kac-Moody algebra and set $\mathsf{P}_+ := \{\lambda \in \mathsf{P} \mid \langle h_i, \lambda \rangle \ge 0 \text{ for any } i \in I\}$ the set of dominant integral weights.

For $i, j \in I$, we choose polynomials $Q_{i,j}(u, v) \in \mathbf{k}[u, v]$ of the form

(7.1)
$$Q_{i,j}(u,v) = \begin{cases} \sum_{\substack{p(\alpha_i,\alpha_i)+q(\alpha_j,\alpha_j)=-2(\alpha_i,\alpha_j)\\0}} t_{i,j;p,q} u^p v^q & \text{if } i \neq j, \end{cases}$$

such that $t_{i,j;-\langle h_i,\alpha_j\rangle,0} \in \mathbf{k}^{\times}$ and

$$Q_{i,j}(u,v) = Q_{j,i}(v,u)$$
 for all $i, j \in I$.

Let $\mathbf{Q} := \bigoplus_{i \in I} \mathbb{Z}\alpha_i$ and $\mathbf{Q}_+ := \bigoplus_{i \in I} \mathbb{Z}_{\geq 0}\alpha_i$ be the root lattice of \mathfrak{g} and the positive root lattice, respectively. For $\beta \in \mathbf{Q}_+$, set

$$I^{\beta} := \Big\{ \nu = (\nu_1, \dots, \nu_n) \in I^n \mid \sum_{k=1}^n \alpha_{\nu_k} = \beta \Big\}.$$

The symmetric group $\mathfrak{S}_n = \langle s_k \mid k = 1, \dots, n-1 \rangle$ acts by place permutations on I^{β} . For $\beta = \sum_{i \in I} b_i \alpha_i \in \mathbb{Q}$, set $\operatorname{ht}(\beta) = \sum_{i \in I} |b_i|$. **Definition 7.1.** For $\beta \in Q_+$ with $ht(\beta) = n$, the quiver Hecke algebra $R(\beta)$ associated with $(\mathsf{C}, \Pi, \mathsf{P}, \Pi^{\vee}, (\cdot, \cdot))$ and $(Q_{i,j}(u, v))_{i,j \in I}$ is the **k**-algebra generated by

$$\{e(\nu) \mid \nu \in I^{\beta}\}, \{x_k \mid 1 \leq k \leq n\}, \{\tau_l \mid 1 \leq l \leq n-1\}$$

satisfying the following defining relations:

$$\begin{aligned} e(\nu)e(\nu') &= \delta_{\nu,\nu'}e(\nu), \ \sum_{\nu \in I^{\beta}} e(\nu) = 1, \ x_{k}e(\nu) = e(\nu)x_{k}, \ x_{k}x_{l} = x_{l}x_{k}, \\ \tau_{l}e(\nu) &= e(s_{l}(\nu))\tau_{l}, \ \tau_{k}\tau_{l} = \tau_{l}\tau_{k} \ if \ |k-l| > 1, \\ \tau_{k}^{2} &= \sum_{\nu \in I^{\beta}} Q_{\nu_{k},\nu_{k+1}}(x_{k}, x_{k+1})e(\nu), \\ \tau_{k}x_{l} - x_{s_{k}(l)}\tau_{k} &= \left(\delta(l = k+1) - \delta(l = k)\right) \sum_{\nu \in I^{\beta}, \ \nu_{k} = \nu_{k+1}} e(\nu), \\ \tau_{k+1}\tau_{k}\tau_{k+1} - \tau_{k}\tau_{k+1}\tau_{k} &= \sum_{\nu \in I^{\beta}, \ \nu_{k} = \nu_{k+2}} \overline{Q}_{\nu_{k},\nu_{k+1}}(x_{k}, x_{k+1}, x_{k+2})e(\nu), \end{aligned}$$

where

$$\overline{Q}_{i,j}(u,v,w) := \frac{Q_{i,j}(u,v) - Q_{i,j}(w,v)}{u-w} \in \mathbf{k}[u,v,w].$$

Now, let $\lambda \colon \mathsf{Q} \times \mathsf{Q} \to \mathbb{Z}$ be a bilinear form such that

(7.2)
$$\lambda(\alpha,\beta) + \lambda(\beta,\alpha) = -2(\alpha,\beta) \text{ for any } \alpha, \beta \in \mathbb{Q}.$$

Then, the algebra $R(\beta)$ is \mathbb{Z} -graded by

(7.3)
$$\deg(e(\nu)) = 0, \quad \deg(x_k e(\nu)) = (\alpha_{\nu_k}, \alpha_{\nu_k}), \quad \deg(\tau_l e(\nu)) = \lambda(\alpha_{\nu_l}, \alpha_{\nu_{l+1}}).$$

We denote by $R_{\lambda}(\beta)$ the graded algebra $R(\beta)$ with the grading (7.3). We also regard $R(\beta)$ as a graded algebra by taking $\lambda = -(\cdot, \cdot)$.

We denote by $\operatorname{Modg}(R_{\lambda}(\beta))$ the graded abelian category of graded $R_{\lambda}(\beta)$ -modules and we set $\operatorname{Modg}(R_{\lambda}) = \bigoplus_{\beta \in \mathbb{Q}_{+}} \operatorname{Modg}(R_{\lambda}(\beta))$.

For $M \in \text{Modg}(R_{\lambda}(\beta))$, we set $\text{wt}(M) := -\beta$.

We denote by $\operatorname{Modg}_{\operatorname{coh}}(R_{\lambda}(\beta))$ the full subcategory of $\operatorname{Modg}(R_{\lambda}(\beta))$ consisting of finitely generated graded $R_{\lambda}(\beta)$ -modules and we set $\operatorname{Modg}_{\operatorname{coh}}(R_{\lambda}) = \bigoplus_{\beta \in \mathbf{Q}_{+}} \operatorname{Modg}_{\operatorname{coh}}(R_{\lambda}(\beta))$.

We denote by $R_{\lambda}(\beta)$ -gmod the full subcategory of $Modg(R_{\lambda}(\beta))$ consisting of Z-graded $R_{\lambda}(\beta)$ -modules with finite **k**-dimension. Set R_{λ} -gmod:= $\bigoplus_{\beta \in \mathbf{Q}_{+}} R_{\lambda}(\beta)$ -gmod.

Note that $R_{\lambda}(\beta)$ -gmod is independent from λ as a graded category after adding $q^{1/2}$ (see [13, Lemma 1.5]). We write Modg(R), R-gmod, etc. for Modg(R_{λ}), R_{λ} -gmod, etc. with $\lambda = -(\cdot, \cdot)$.

For $M \in R$ -gmod, set $M^* := \operatorname{Hom}_{\mathbf{k}}(M, \mathbf{k})$. Then M^* becomes an $R(\beta)$ -module via the grade-preserving antiautomorphism of $R(\beta)$ which fixes the generators $e(\nu)$, x_k , and τ_k 's. We say that M is *self-dual* if $M \simeq M^*$ in R-gmod. For each simple module M in R-gmod, there exists $m \in \mathbb{Z}$ such that $q^m M$ is self-dual.

For each $i \in I$ and $n \ge 1$, $R(n\alpha_i)$ has a unique self-dual simple module which is denoted by $L(i^n)$. Sometimes we will denote it by $\langle i^n \rangle$. We have dim $(\langle i^n \rangle) = n!$.

7.2. Universal R-matrices. The category $\operatorname{Modg}(R_{\lambda})$ is endowed with a monoidal category structure with the convolution product \circ (e.g. see [17, 18, 23] for details). Moreover it is quasi-rigid by [6, Lemma 3.1]. The category R_{λ} -gmod is a graded monoidal category satisfying (5.1). We have a fully faithful monoidal functor

$$\operatorname{Modg}_{\operatorname{coh}}(R_{\lambda}) \to \operatorname{Pro}(R_{\lambda}\operatorname{-gmod})$$

by

$$M \mapsto \lim_{m} M/(R_{\lambda}(R_{\lambda})_{\geq m}M).$$

Hereafter, we identify $\operatorname{Modg}_{\operatorname{coh}}(R_{\lambda})$ as a full subcategory of $\operatorname{Pro}(R_{\lambda}\operatorname{-gmod})$.

Note that an affine object in R_{λ} -gmod is equivalent to a pair (M, z) of a graded R_{λ} -module M and an injective endomorphism z of M of a positive degree such that $\mathsf{M}/z\mathsf{M} \in R_{\lambda}$ -gmod and $\mathsf{M}_k = 0$ for $k \ll 0$.

Let $\beta \in \mathbb{Q}_+$ with $m = \operatorname{ht}(\beta)$. For $k = 1, \ldots, m-1$ and $\nu \in I^{\beta}$, the *intertwiner* $\varphi_k \in R(\beta)$ is defined by

(7.4)
$$\varphi_k e(\nu) = \begin{cases} \left(\tau_k (x_k - x_{k+1}) + 1\right) e(\nu) & \text{if } \nu_k = \nu_{k+1}, \\ \tau_k e(\nu) & \text{otherwise.} \end{cases}$$

Lemma 7.2 ([7, Lemma 1.5]). The intertwiners have the following properties.

- (i) $\varphi_k^2 e(\nu) = \left(Q_{\nu_k,\nu_{k+1}}(x_k, x_{k+1}) + \delta_{\nu_k,\nu_{k+1}} \right) e(\nu).$
- (ii) $\{\varphi_k\}_{k=1,\dots,m-1}$ satisfies the braid relation.
- (iii) For a reduced expression $w = s_{i_1} \cdots s_{i_t} \in \mathfrak{S}_m$, we set $\varphi_w := \varphi_{i_1} \cdots \varphi_{i_t}$. Then φ_w does not depend on the choice of reduced expression of w.
- (iv) For $w \in \mathfrak{S}_m$ and $1 \leq k \leq m$, we have $\varphi_w x_k = x_{w(k)} \varphi_w$.
- (v) For $w \in \mathfrak{S}_m$ and $1 \leq k < m$, if w(k+1) = w(k) + 1, then $\varphi_w \tau_k = \tau_{w(k)} \varphi_w$.

For $m, n \in \mathbb{Z}_{\geq 0}$, we set w[m, n] to be the element of \mathfrak{S}_{m+n} such that

$$w[m,n](k) := \begin{cases} k+n & \text{if } 1 \leq k \leq m, \\ k-m & \text{if } m < k \leq m+n. \end{cases}$$

For an $R(\beta)$ -module M and an $R(\gamma)$ -module N, the $R(\beta) \otimes R(\gamma)$ -linear map $M \otimes N \to N \circ M$ defined by

$$u \otimes v \mapsto \varphi_{w[\operatorname{ht}(\gamma),\operatorname{ht}(\beta)]}(v \otimes u)$$

extends to an $R(\beta + \gamma)$ -module homomorphism (neglecting a grading shift)

$$\mathbf{R}_{M,N}^{\mathrm{uni}} \colon M \circ N \longrightarrow N \circ M$$

We call $\mathbb{R}_{M,N}^{\text{uni}}$ the universal *R*-matrix between *M* and *N*. Since the intertwiners satisfies the braid relations, the universal *R*-matrices $\mathbb{R}_{M,N}^{\text{uni}}$ satisfy the Yang-Baxter equation ([7, (1.9)]).

Let (M, z) be an affine object in R_{λ} -gmod. We set $\mathrm{R}^{\mathrm{uni}}_{\mathsf{M}}(X) = \mathrm{R}^{\mathrm{uni}}_{\mathsf{M},X} \colon \mathsf{M} \circ X \to X \circ \mathsf{M}$ (neglecting the grading shift). If $(\mathsf{M}, \mathrm{R}^{\mathrm{uni}}_{\mathsf{M}})$ is a rational center in R_{λ} -gmod, then we say that $(\mathsf{M}, \mathrm{R}^{\mathrm{uni}}_{\mathsf{M}})$ is a *canonical affinization* of $\mathsf{M}/z\mathsf{M} \in R_{\lambda}$ -gmod. 7.3. Canonical affinizations in R_{λ} -gmod. We recall the notion of affinizations for R-modules introduced in [13]. For $\beta \in \mathbb{Q}_+$ and $i \in I$, let

(7.5)
$$\mathfrak{p}_{i,\beta} := \sum_{\nu \in I^{\beta}} \Big(\prod_{a \in \{1,\dots,\operatorname{ht}(\beta)\}, \nu_a = i} x_a \Big) e(\nu) \in R(\beta).$$

Then $\mathfrak{p}_{i,\beta}$ belongs to the center of $R(\beta)$.

Let M be a simple module in $R_{\lambda}(\beta)$ -gmod. Assume that there exists a graded $R_{\lambda}(\beta)$ -module M with an endomorphism z_{M} of M with degree $d_{\mathsf{M}} \in \mathbb{Z}_{>0}$ such that

(i) $\mathbf{M}/z_{\mathbf{M}}\mathbf{M}\simeq M$,

(7.6) (ii) M is a finitely generated free module over the polynomial ring $\mathbf{k}[z_{\mathsf{M}}]$, (iii) $\mathfrak{p}_{i,\beta}\mathsf{M} \neq 0$ for all $i \in I$.

Note that the conditions (i) and (ii) are equivalent to the statement (M, z_M) is an affine object.

For an indeterminate z and a non-zero homogeneous $w \in \mathbf{k}[z]$, we denote by $\mathsf{M}|_{z_{\mathsf{M}}=w} := \mathbf{k}[z] \otimes_{\mathbf{k}[z_{\mathsf{M}}]} \mathsf{M}$ the object in $\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z], R_{\lambda}\operatorname{-gmod})$. Here, $\mathbf{k}[z]$ is a $\mathbf{k}[z_{\mathsf{M}}]$ algebra by the algebra homomorphism $z_{\mathsf{M}} \mapsto w \in \mathbf{k}[z]$. If $w = cz^{a}$ for some $c \in \mathbf{k}^{\times}, a \in \mathbb{Z}_{>0}$ such that $\deg(z_{\mathsf{M}}) = a \deg(z)$, then $\mathsf{M}|_{z_{\mathsf{M}}=w}$ is an affine object ([13, Example 2.5 (ii)]). Sometimes we will write $\mathsf{M}|_{z_{\mathsf{M}}=w} = \mathsf{M}_{w}$ for short.

Proposition 7.3 ([13, Lemma 2.9]). Let M be a simple module in $R_{\lambda}(\beta)$ -gmod. Assume that an $R_{\lambda}(\beta)$ -module (M, z) satisfies the conditions (7.6). Then (M, R_{M}^{uni}) is a rational center in R_{λ} -gmod.

Proof. It is enough to show that the R-matrix $\mathbb{R}_{M,X}^{\mathrm{uni}} \colon \mathsf{M} \circ X \to X \circ \mathsf{M}$ is an isomorphism in $\operatorname{Raff}_z(R_\lambda\operatorname{-gmod})$ for any $X \in R_\lambda\operatorname{-gmod}$.

The composition $\mathbb{R}_{M,X}^{\mathrm{uni}} \circ \mathbb{R}_{X,M}^{\mathrm{uni}} \colon X \circ \mathsf{M} \to \mathsf{M} \circ X \to X \circ \mathsf{M}$ is given by

$$f(z) := \prod_{i \neq j} \chi_i(\mathsf{M})(t_i) \underset{t_i}{\star} Q_{i,j}(t_i, t_j) \underset{t_j}{\star} \chi_j(X)$$

(see (9.3) below, and for the notation χ_i and \star_t , see § 9.1, § 9.2). Since $\operatorname{End}_{\mathbf{k}[z]}(\mathsf{M}) \simeq \mathbf{k}[z]$ by Lemma 4.11, the homogeneous polynomial $\chi_i(\mathsf{M})(t_i, z) \in \mathbf{k}[t_i, z]$ is monic in t_i and quasi-monic in z by the assumption (iii) (7.6).

Hence f(z) has the form (up to a constant multiple) $f(z) = z^m - a$ for some $m \in \mathbb{Z}_{\geq 0}$ and $a \in \text{END}(X)_{>0}[z]$. Since $a^s = 0$ for some $s \in \mathbb{Z}_{>0}$, if we set $g(z) = (z^m)^{s-1} + az(z^m)^{s-2} + \cdots + a^{s-1}$, then we have

$$f(z)g(z) = g(z)f(z) = z^{ms} - a^s = z^{ms}.$$

Hence $\operatorname{R}_{M,X}^{\operatorname{uni}} \circ \left(\operatorname{R}_{X,M}^{\operatorname{uni}} g(z) \right) = z^{ms} \operatorname{id}_{X \otimes M}$ is invertible in $\operatorname{Raff}_{z}(R_{\lambda}\operatorname{-gmod})$, and hence $\operatorname{R}_{M,X}^{\operatorname{uni}}$ has a right inverse.

Similarly, $\mathbf{R}_{\mathsf{M},X}^{\mathrm{uni}}$ has a left inverse.

Remark 7.4. Let (M, z) be an affine object of a simple $M \in R_{\lambda}$ -gmod. Even if (M, R_{M}^{uni}) is a rational center in R_{λ} -gmod, (M, z) may not satisfy (7.6). For example, $(M \circ C, R_{M}^{uni} \circ R_{C})$ is a rational center if (M, R_{M}^{uni}) is a rational center and (C, R_{C}) is a central object in R_{λ} -gmod.

In the sequel, we use the terminology "affinization" in the sense of Definition 6.7.

7.4. Schur-Weyl duality. Let C be a graded k-linear monoidal category satisfying (6.2). We assume further that C is quasi-rigid. Let R be the quiver Hecke algebra associated with a Cartan matrix C and a set of polynomials $\{Q_{i,j}(u,v)\}$ as in (7.1).

Let $\{(\widehat{K}_i, z_i)\}_{i \in I}$ be a family of affinizations in \mathcal{C} such that

(7.7)
$$\begin{cases} (a) \ \deg(z_i) = 2(\alpha_i, \alpha_i), \\ (b) \ \mathbf{K}_i := \widehat{\mathbf{K}}_i / z_i \widehat{\mathbf{K}}_i \text{ is real simple in } \mathcal{C} \text{ for any } i \in I, \\ (c) \ \mathscr{D}(\widehat{\mathbf{K}}_i, \widehat{\mathbf{K}}_j) = Q_{i,j}(z_i, z_j) \text{ for } i \neq j. \end{cases}$$

We define the bilinear form $\lambda \colon \mathsf{Q} \times \mathsf{Q} \to \mathbb{Z}$ by

$$\lambda(\alpha_i, \alpha_j) = \begin{cases} \Lambda(\mathbf{K}_i, \mathbf{K}_j) & \text{for } i, j \in I \text{ such that } i \neq j. \\ -(\alpha_i, \alpha_j) & \text{if } i = j \in I. \end{cases}$$

Then we have $\lambda(\alpha, \beta) + \lambda(\beta, \alpha) = -2(\alpha, \beta)$ for any $\alpha, \beta \in \mathbb{Q}$.

Let $R_{\widehat{K}_i,\widehat{K}_j}^{\text{ren}}: \widehat{K}_i \otimes \widehat{K}_j \to \widehat{K}_j \otimes \widehat{K}_i$ be a renormalized R-matrix. It is determined up to a constant multiple. We normalize them such that

(7.8)
$$\begin{cases} \text{(a)} \ \mathrm{R}_{\widehat{\mathrm{K}}_{j},\widehat{\mathrm{K}}_{i}}^{\mathrm{ren}} \circ \mathrm{R}_{\widehat{\mathrm{K}}_{i},\widehat{\mathrm{K}}_{j}}^{\mathrm{ren}} = Q_{i,j}(z_{i}, z_{j}) \operatorname{\mathsf{id}}_{\widehat{\mathrm{K}}_{i} \otimes \widehat{\mathrm{K}}_{j}} \text{ for } i \neq j, \\ \text{(b) for any } i \in I, \text{ there exists } T_{i} \in \operatorname{End}_{\operatorname{Pro}(\mathcal{C})}(\widehat{\mathrm{K}}_{i} \otimes \widehat{\mathrm{K}}_{i}) \text{ such that} \\ \mathrm{R}_{\widehat{\mathrm{K}}_{i},\widehat{\mathrm{K}}_{i}}^{\mathrm{ren}} - \operatorname{\mathsf{id}}_{\widehat{\mathrm{K}}_{i} \otimes \widehat{\mathrm{K}}_{i}} = (z_{i} \otimes 1 - 1 \otimes z_{i}) \circ T_{i}. \end{cases}$$

Such normalizations are possible by Proposition 6.13.

We call $(\{(\widehat{K}_i, z_i)\}_{i \in I}, \{R_{\widehat{K}_i, \widehat{K}_j}^{\text{ren}}\}_{i, j \in I})$ a duality datum.

Remark 7.5. Once $\{(\widehat{K}_i, z_i)\}_{i \in I}$ is given, a duality datum $(\{(\widehat{K}_i, z_i)\}_{i \in I}, \{\mathbb{R}_{\widehat{K}_i, \widehat{K}_j}^{\text{ren}}\}_{i, j \in I})$ exists and it is unique up to constant multiples.

More precisely, if $(\{(\widehat{K}_i, z_i)\}_{i \in I}, \{\widetilde{R^{ren}}_{\widehat{K}_i, \widehat{K}_j}\}_{i,j \in I})$ is another duality datum, then there exist $c_{i,j} \in \mathbf{k}^{\times}$ $(i, j \in I)$ such that $\widetilde{R^{ren}}_{\widehat{K}_i, \widehat{K}_j} = c_{i,j} \operatorname{R}^{ren}_{\widehat{K}_i, \widehat{K}_j}$ and $c_{i,j} c_{j,i} = 1, c_{i,i} = 1$.

Proposition 7.6 (cf. [7, § 3.1, § 3.2]). Let $(\{(\widehat{K}_i, z_i)\}_{i \in I}, \{R_{\widehat{K}_i, \widehat{K}_j}^{ren}\}_{i,j \in I})$ be a duality datum.

(i) There exists a right exact monoidal functor

$$\widehat{\mathscr{F}}$$
: Modg_{coh} $(R_{\lambda}) \to \operatorname{Pro}(\mathcal{C})$

such that

(7.9) $\widehat{\mathscr{F}}(\widetilde{L}(i)_{z_i}) \simeq \widehat{K}_i \quad and \quad \widehat{\mathscr{F}}(L(i)) \simeq K_i,$

(7.10)
$$\widehat{\mathscr{F}}(e(i,j)\varphi_1) = \mathrm{R}_{\widehat{\mathrm{K}}_i,\widehat{\mathrm{K}}_j}^{\mathrm{ren}} \in \mathrm{Hom}_{\mathbf{k}[z_i,z_j]}(\widehat{\mathrm{K}}_i \otimes \widehat{\mathrm{K}}_j, \widehat{\mathrm{K}}_j \otimes \widehat{\mathrm{K}}_i).$$

Here $\widetilde{L}(i)_{z_i}$ is the affinization $(R(\alpha_i), z_i) = (\mathbf{k}[x_1], x_1)$ of $L(i) = R(\alpha_i)/R(\alpha_i)x_1$, φ_1 is given in (7.4), and $e(i, j)\varphi_1 \colon R_\lambda(\alpha_i + \alpha_j)e(i, j) \to R_\lambda(\alpha_i + \alpha_j)e(j, i)$ is the morphism by the right multiplication.

Moreover, such a right exact monoidal functor $\widehat{\mathscr{F}}$ is unique up to an isomorphism.

(ii) The functor $\widehat{\mathscr{F}}$ can be restricted to a monoidal functor

$$\mathscr{F}: R_{\lambda}\operatorname{-gmod} \to \mathcal{C}.$$

- (iii) Assume further that C is of finite type. Then the functor $\widehat{\mathscr{F}}$ is exact and we have the following properties.
 - (a) \mathscr{F} sends a simple module in R_{λ} -gmod to a simple objects in \mathcal{C} or 0.
 - (b) If (M, z_M) is an affine object of M in R_{λ} -gmod, then $(\widehat{\mathscr{F}}(M), \widehat{\mathscr{F}}(z_M))$ is an affine object of $\mathscr{F}(M) \in \mathcal{C}$.

Proof. First note that $\{\mathbf{R}_{\hat{\mathbf{K}}_i,\hat{\mathbf{K}}_j}^{\text{ren}}\}_{i,j\in I}$ satisfies the Yang-Baxter equation by Lemma 6.23. For each $\beta \in \mathbf{Q}_+$ and $\nu = (\nu_1, \dots, \nu_m) \in I^{\beta}$, define objects in $\operatorname{Pro}(\mathcal{C})$ by

$$\widehat{\mathrm{K}}(\nu) := \widehat{\mathrm{K}}_{\nu_1} \otimes \cdots \otimes \widehat{\mathrm{K}}_{\nu_m} \quad \text{and} \quad \widehat{\mathrm{K}}(\beta) := \bigoplus_{\nu \in I^\beta} \widehat{\mathrm{K}}(\nu).$$

Set $T_{i,j} = \mathbb{R}_{\widehat{K}_i,\widehat{K}_j}^{\text{ren}}$ if $i \neq j$ and $T_{i,j} = T_i$ if i = j. Here, T_i is given in (7.8).

Then we can endow $\widehat{K}(\beta) \in \operatorname{Pro}(\mathcal{C})$ with a graded $R_{\lambda}(\beta)^{\operatorname{opp}}$ -module structure in $Pro(\mathcal{C})$ as follows (cf. [7, § 3]):

- (1) $e(\nu)$ is the projection $\widehat{K}(\beta) \to \widehat{K}(\nu) \subset \widehat{K}(\beta)$,
- (2) x_k acts on $\widehat{\mathbf{K}}(\nu)$ by

$$\widehat{\mathrm{K}}_{\nu_1}\otimes\cdots\otimes\widehat{\mathrm{K}}_{\nu_{k-1}}\otimes z_{\nu_k}\otimes\widehat{\mathrm{K}}_{\nu_{k+1}}\otimes\cdots\otimes\widehat{\mathrm{K}}_{\nu_m},$$

(3) $\tau_k \colon \widehat{\mathrm{K}}(\nu) \to \widehat{\mathrm{K}}(s_k \nu)$ is given by

$$\widehat{\mathbf{K}}_{\nu_1} \otimes \cdots \otimes \widehat{\mathbf{K}}_{\nu_{k-1}} \otimes T_{\nu_k,\nu_{k+1}} \otimes \widehat{\mathbf{K}}_{\nu_{k+2}} \otimes \cdots \otimes \widehat{\mathbf{K}}_{\nu_m}.$$

Since $R(\beta)$ is left noetherian ([17, Corollary 2.11]), there exists a right exact functor

$$\mathscr{F}_{\beta} \colon \operatorname{Modg}_{\operatorname{coh}}(R_{\lambda}(\beta)) \to \operatorname{Pro}(\mathcal{C})$$

given by

$$M \mapsto \widehat{\mathrm{K}}(\beta) \otimes_{R_{\lambda}(\beta)} M.$$

Note that $\widehat{\mathscr{F}}_{\alpha_i}(\widetilde{L}(i)_{z_i}) \simeq \widehat{\mathscr{F}}_{\alpha_i}(R_{\lambda}(\alpha_i)) \simeq \widehat{\mathrm{K}}_i$ for each $i \in I$. Since $\widehat{\mathscr{F}}_{\beta}$ is right exact and $\widehat{\mathscr{F}}_{\alpha_i}(z_i) = z_i$, we have $\widehat{\mathscr{F}}_{\alpha_i}(L(i)) \simeq \mathrm{K}_i$.

Set

$$\widehat{\mathscr{F}} := \bigoplus_{\beta \in \mathbf{Q}_+} \widehat{\mathscr{F}}_{\beta} : \operatorname{Modg}_{\operatorname{coh}}(R_{\lambda}) \to \operatorname{Pro}(\mathcal{C}).$$

Then, $\widehat{\mathscr{F}}$ is a monoidal functor. Indeed, for any $M \in \operatorname{Mod}_{\operatorname{coh}}(R_{\lambda}(\beta))$ and $N \in \operatorname{Mod}_{\operatorname{coh}}(R_{\lambda}(\gamma))$, we have

$$\widehat{\mathscr{F}}(M \circ N) := \widehat{\mathrm{K}}(\beta + \gamma) \otimes_{R_{\lambda}(\beta + \gamma)} \left(R_{\lambda}(\beta + \gamma)e(\beta + \gamma) \otimes_{R_{\lambda}(\beta)\otimes R_{\lambda}(\gamma)} (M \otimes N) \right)$$
$$\simeq \widehat{\mathrm{K}}(\beta + \gamma)e(\beta + \gamma) \otimes_{R_{\lambda}(\beta)\otimes R_{\lambda}(\gamma)} (M \otimes N)$$
$$\simeq \left(\widehat{\mathrm{K}}(\beta) \otimes \widehat{\mathrm{K}}(\gamma) \right) \otimes_{R_{\lambda}(\beta)\otimes R_{\lambda}(\gamma)} (M \otimes N)$$
$$\simeq \left(\widehat{\mathrm{K}}(\beta) \otimes_{R_{\lambda}(\beta)} M \right) \otimes \left(\widehat{\mathrm{K}}(\gamma) \otimes_{R_{\lambda}(\gamma)} N \right)$$
$$\simeq \widehat{\mathscr{F}}(M) \otimes \widehat{\mathscr{F}}(N).$$

For the uniqueness of $\widehat{\mathscr{F}}$, see Lemma 7.7 below.

Set $\operatorname{Pol}(\beta) = \bigoplus_{\nu \in I^{\beta}} \mathbf{k}[x_1, \dots, x_{\operatorname{ht}(\beta)}] e(\nu)$. Then $\operatorname{Pol}(\beta)$ is a commutative subalgebra of $R(\beta)$. Since $\widehat{K}(\beta) \in \operatorname{Pro}_{\operatorname{coh}}(\operatorname{Pol}(\beta), \mathcal{C})$, Lemma 3.4 implies that

 $\widehat{\mathcal{K}}(\beta) \otimes_{\operatorname{Pol}(\beta)} M \in \mathcal{C}$ for any $M \in R_{\lambda}$ -gmod.

Since $\widehat{\mathcal{K}}(\beta) \otimes_{R_{\lambda}(\beta)} M$ is a quotient of $\widehat{\mathcal{K}}(\beta) \otimes_{\operatorname{Pol}(\beta)} M$, we have $\widehat{\mathscr{F}}(M) \in \mathcal{C}$ by Lemma 2.1. Because $\mathcal{C} \to \operatorname{Pro}(\mathcal{C})$ is fully faithful, we get the restriction

$$\mathscr{F} := \widehat{\mathscr{F}}|_{R_{\lambda}-\mathrm{gmod}} \colon R_{\lambda}-\mathrm{gmod} \to \mathcal{C}.$$

Let us show (iii). Assume that C is of finite type. Then $R(\beta)$ has a finite global dimension by [15, 21]. Hence, [7, Proposition 3.7] implies that $R(\beta)^{\text{opp}}$ -modules is flat over $R(\beta)^{\text{opp}}$ as soon as it is flat over $Pol(\beta)$.

Since \widehat{K}_i 's are affine objects, $\widehat{K}(\beta)$ is flat over $\operatorname{Pol}(\beta)$ by Proposition 5.5 (iv). Hence $\widehat{K}(\beta)$ is $R^{\operatorname{opp}}(\beta)$ -flat, which implies that $\widehat{\mathscr{F}}_{\beta}$ is exact.

Assume that M is simple in $R_{\lambda}(\beta)$ -gmod. Let us show that $\mathscr{F}(M)$ is simple or zero by induction on $ht(\beta)$. We may assume that $\beta \neq 0$. Then there exist $i \in I$

and a simple $R_{\lambda}(\beta - \alpha_i)$ -module N such that $M \simeq N \nabla L(i)$. Then $M \simeq \text{Im}(\mathbf{r})$, where \mathbf{r} is an R-matrix $\mathbf{r} \colon N \circ L(i) \to L(i) \circ N$. Since \mathscr{F} is exact, we have $\mathscr{F}(M) \simeq \mathscr{F}(\text{Im}(\mathbf{r})) \simeq \text{Im} \mathscr{F}(\mathbf{r})$. We may assume that $\mathscr{F}(\mathbf{r})$ is non-zero. Hence $\mathscr{F}(N) \neq 0$. By the induction hypothesis, $\mathscr{F}(N)$ is simple. Then Proposition 6.19 implies that $\text{Im} \mathscr{F}(\mathbf{r})$ is isomorphic to the simple head $\mathscr{F}(N) \nabla \mathscr{F}(L(i))$, since \mathcal{C} is assumed to be quasi-rigid and $\mathscr{F}(L(i)) \simeq K_i$ is affreal. Hence we get (a).

(b) is a direct consequence of the exactness of $\widehat{\mathscr{F}}$.

Lemma 7.7. Let $(\{(\widehat{K}_i, z_i)\}_{i \in I}, \{R_{\widehat{K}_i, \widehat{K}_j}^{\text{ren}}\}_{i, j \in I})$ be a duality datum and let

$$\widehat{\mathscr{F}}$$
: Modg_{coh} $(R_{\lambda}) \to \operatorname{Pro}(\mathcal{C})$

be a right exact monoidal functor satisfying (7.9) and (7.10). Then it is unique up to an isomorphism.

Proof. Since
$$R(\beta)e(\nu) \simeq L(\nu_1)_{z_{\nu_1}} \circ \cdots \circ L(\nu_r)_{z_{\nu_r}}$$
 in $Modg_{coh}(R_{\lambda})$, by (7.9) we have

$$\widehat{\mathscr{F}}(R(\beta)e(\nu)) \simeq \widehat{\mathcal{K}}(\nu) \quad \text{and} \quad \widehat{\mathscr{F}}(R(\beta)) \simeq \widehat{\mathcal{K}}(\beta)$$

for any $\beta \in \mathbb{Q}_+$ and $\nu \in I^{\beta}$. Note that $\widehat{\mathscr{F}}(R(\beta)) \simeq \widehat{K}(\beta)$ has an $R(\beta)^{\text{opp}}$ -module structure, which is uniquely determined by (7.9) and (7.10) since the action of $x_k - x_{k+1}$ on \widehat{K}_{ν} with $\nu_k = \nu_{k+1}$ is a monomorphism.

Hence for each $M \in \operatorname{Modg}_{\operatorname{coh}}(R_{\lambda}(\beta))$ there exists a functorial morphism

(7.11)
$$\widehat{\mathcal{K}}(\beta) \otimes_{R(\beta)} M \longrightarrow \widehat{\mathscr{F}}(M)$$

induced from

$$\operatorname{Hom}_{\operatorname{Pro}(\mathcal{C})}(\widehat{\mathrm{K}}_{\beta} \otimes_{R(\beta)} M, \widehat{\mathscr{F}}(M)) \simeq \operatorname{Hom}_{R(\beta)}(M, \operatorname{Hom}_{\operatorname{Pro}(\mathcal{C})}(\widehat{\mathrm{K}}_{\beta}, \widehat{\mathscr{F}}(M)))$$
$$\simeq \operatorname{Hom}_{R(\beta)}(\operatorname{Hom}_{R(\beta)}(R(\beta), M), \operatorname{Hom}_{\operatorname{Pro}(\mathcal{C})}(\widehat{\mathscr{F}}(R(\beta)), \widehat{\mathscr{F}}(M))).$$

Note that (7.11) is an isomorphism if M is a free $R(\beta)$ -module. Since $\widehat{\mathscr{F}}$ is right exact, one can take a free presentation of M to conclude that (7.11) is an isomorphism for a general M.

8. LOCALIZATIONS OF R-gmod

8.1. Categories \mathscr{C}_w and \mathscr{C}_w^* . Let R be a quiver Hecke algebra as in section 7. For $\alpha, \beta \in \mathbb{Q}_+$, we set

$$e(\alpha,\beta) = \sum_{\substack{\nu \in I^{\alpha+\beta} \\ \sum_{k=1}^{\operatorname{ht}(\alpha)} \alpha_{\nu_k} = \alpha, \ \sum_{k=1}^{\operatorname{ht}(\beta)} \alpha_{\nu_k + \operatorname{ht}(\alpha)} = \beta}} e(\nu) \in R(\alpha+\beta).$$

For $M \in R(\beta)$ -gmod we define

$$W(M) := \{ \gamma \in \mathsf{Q}_+ \cap (\beta - \mathsf{Q}_+) \mid e(\gamma, \beta - \gamma)M \neq 0 \},\$$
$$W^*(M) := \{ \gamma \in \mathsf{Q}_+ \cap (\beta - \mathsf{Q}_+) \mid e(\beta - \gamma, \gamma)M \neq 0 \}.$$

Let W be the Weyl group, that is, the subgroup of Aut(P) generated by $\{s_i\}_{i \in I}$ where $s_i(\lambda) = \lambda - \langle h_i, \lambda \rangle \alpha_i$. Let us denote by > the Bruhat order on W.

For $w \in W$, let us define the full monoidal subcategories of R-gmod by

(8.1)
$$\mathscr{C}_w := \{ M \in R \text{-gmod} \mid \mathsf{W}(M) \subset \mathsf{Q}_+ \cap w\mathsf{Q}_- \}, \\ \mathscr{C}^*_w := \{ M \in R \text{-gmod} \mid \mathsf{W}^*(M) \subset \mathsf{Q}_+ \cap w\mathsf{Q}_- \}.$$

Recall that there is an involutive **k**-algebra automorphism ψ of $R(\beta)$ given by

(8.2)
$$e(\nu_1, \ldots, \nu_n) \mapsto e(\nu_n, \ldots, \nu_1), \quad x_k \mapsto x_{n+1-k}, \quad \tau_k \mapsto -\tau_{n-k},$$

where $n = ht(\beta)$. Then ψ induces an equivalence of monoidal categories:

$$\psi_* \colon R\text{-gmod} \simeq (R\text{-gmod})^{\text{rev}}.$$

Here, for a monoidal category \mathcal{T} , \mathcal{T}^{rev} denotes the monoidal category endowed with the reversed tensor product \otimes_{rev} defined by $M \otimes_{\text{rev}} N := N \otimes M$.

Since we have $\psi(e(\gamma, \beta - \gamma)) = e(\beta - \gamma, \gamma)$ for $\beta, \gamma \in Q_+$ with $\beta - \gamma \in Q_+$, the automorphism ψ induces an equivalence of monoidal categories

$$\mathscr{C}_w^* \simeq (\mathscr{C}_w)^{\operatorname{rev}}.$$

For $\lambda, \mu \in \mathsf{P}$, we define a partial order $\lambda \preccurlyeq \mu$ if there exists a sequence of real positive roots β_k $(1 \leqslant k \leqslant \ell)$ such that $\lambda = s_{\beta_\ell} \cdots s_{\beta_1} \mu$ and $(\beta_k, s_{\beta_{k-1}} \cdots s_{\beta_1} \mu) \ge 0$ for $1 \leqslant k \leqslant \ell$. Here $s_\beta(\lambda) = \lambda - (\beta^{\vee}, \lambda)\beta$ with $\beta^{\vee} = \frac{2}{(\beta,\beta)}\beta$.

Assume that $\lambda, \mu \in W\Lambda$ for some $\Lambda \in \mathsf{P}_+$ and that $\lambda \preccurlyeq \mu$. Then there exists a module $\mathsf{M}(\lambda,\mu)$ in $R(\mu-\lambda)$ -gmod, called the *determinantial module*. We refer [11, Section 3.3] for the definition and properties of determinantial modules. In particular, $\mathsf{M}(\lambda,\mu)$ admits an affinization ([11, Theorem 3.26]).

8.2. Localizations. In this subsection we recall the localization \mathscr{C}_w of the monoidal category \mathscr{C}_w introduced in [11]. Throughout this subsection, $w \in W$ is an element of the Weyl group and we assume that $I_w = I$, where $I_w := \{i_1, \ldots, i_r\}$ for a reduced expression $w = s_{i_1} \cdots s_{i_r}$.

For each $i \in I$, set $\mathsf{C}_i := \mathsf{M}(w\Lambda_i, \Lambda_i)$. Then for each $X \in R(\beta)$ -gmod, there exists a morphism $R_{\mathsf{C}_i}(X) : \mathsf{C}_i \circ X \to q^{\phi_i(\beta)} X \circ \mathsf{C}_i$, where

$$\phi_i(\beta) = -(w\Lambda_i + \Lambda_i, \beta)$$
 for any $\beta \in \mathbf{Q}$.

Moreover, the family $(C_i, R_{C_i}, \phi_i)_{i \in I}$ forms a real commuting family of non-degenerate graded (left) braiders in R-gmod ([11, Proposition 5.1]). Note that $(C_i, R_{C_i}|_{\mathscr{C}_w})$ is a central object of \mathscr{C}_w ([11, Theorem 5.2]), which means that $R_{C_i}(X)$ is an isomorphism for all $X \in \mathscr{C}_w$.

Hence by [11, Theorem 2.12], there exists a localization of \mathscr{C}_w by the family $(\mathsf{C}_i, R_{\mathsf{C}_i}|_{\mathscr{C}_w})$ and a canonical functor

$$\Phi_w \colon \mathscr{C}_w \to \widetilde{\mathscr{C}}_w := \mathscr{C}_w[\mathsf{C}_i^{\circ -1} \mid i \in I].$$

We have the following properties ([11, 12]):

- (i) the category $\widetilde{\mathscr{C}}_w$ is a **k**-linear abelian rigid monoidal category where the tensor product is a natural extension of \circ on \mathscr{C}_w ,
- (ii) the grading shift functor q and the contravariant functor $M \mapsto M^*$ on \mathscr{C}_w are extended to $\widetilde{\mathscr{C}}_w$,
- (iii) for any simple module $M \in \widetilde{\mathscr{C}}_w$, there exists a unique $n \in \mathbb{Z}$ such that $q^n M$ is self-dual,

- (iv) the objects $\Phi_w(\mathsf{C}_i)$ are invertible in $\widetilde{\mathscr{C}}_w$, that is, the functors $\Phi_w(\mathsf{C}_i) \circ \cdot$ and $\cdot \circ \Phi_w(\mathsf{C}_i)$ are equivalences,
- (v) for each $\Lambda \in \mathsf{P}$, there is an invertible object $\mathsf{C}_{\Lambda} \in \widetilde{\mathscr{C}}_w$ such that $\mathsf{C}_{\Lambda} \circ \mathsf{C}_{\Lambda'} \simeq q^{\mathsf{H}(\Lambda,\Lambda')}\mathsf{C}_{\Lambda+\Lambda'}$ for $\Lambda, \Lambda' \in \mathsf{P}$, and $\mathsf{C}_{\Lambda} = \Phi_w(\mathsf{M}(w\Lambda,\Lambda))$ for $\Lambda \in \mathsf{P}_+$, where $\mathsf{H}(\bullet, \bullet)$ is a \mathbb{Z} -bilinear map on Γ determined by $\mathsf{H}(\Lambda_i, \Lambda_j) = (\Lambda_i, w\Lambda_j \Lambda_j)$,
- (vi) for any simple object S of \mathscr{C}_w , the object $\Phi_w(S)$ is simple in \mathscr{C}_w ,
- (vii) every simple object M of \mathscr{C}_w is isomorphic to $\mathsf{C}_{\Lambda} \circ \Phi_w(S)$ for some simple object S of \mathscr{C}_w and $\Lambda \in \mathsf{P}$ and hence $\operatorname{Hom}_{\widetilde{\mathscr{C}}_w}(M, M) = \mathbf{k} \operatorname{id}_M$,
- (viii) every object in $\widetilde{\mathscr{C}}_w$ has finite length,
- (ix) for two simple objects S and S' in \mathscr{C}_w and $\Lambda, \Lambda' \in \mathsf{P}, \mathsf{C}_\Lambda \circ \Phi_w(S) \simeq \mathsf{C}_{\Lambda'} \circ \Phi_w(S')$ in $\widetilde{\mathscr{C}_w}$ if and only if $q^{\mathrm{H}(\Lambda,\mu)}\mathsf{C}_{\Lambda+\mu} \circ S \simeq q^{\mathrm{H}(\Lambda',\mu)}\mathsf{C}_{\Lambda'+\mu} \circ S'$ in \mathscr{C}_w for some $\mu \in \mathsf{P}$ such that $\Lambda + \mu, \Lambda' + \mu \in \mathsf{P}_+$.


If one localizes the category *R*-gmod via the same family $(C_i, R_{C_i}, \phi_i)_{i \in I}$, then the resulting category turns out to be equivalent to the category $\widetilde{\mathscr{C}}_w$ ([11, Theorem 5.9]). Denoting by \mathscr{Q}_w the composition of the equivalence and the canonical functor from *R*-gmod to the localization, we have

$$\mathscr{Q}_w \colon R\text{-gmod} \to \mathscr{C}_w$$

and the following diagram is quasi-commutative:

Similarly, the family $(\psi_*(\mathsf{C}_i), \psi_* \circ R_{\mathsf{C}_i} \circ \psi_*, \phi_i)_{i \in I}$ forms a real commuting family of right braiders in R-gmod so that there exists a localization $\widetilde{\mathscr{C}}_w^*$ of \mathscr{C}_w^* and R-gmod via

the family. Then the localization functors also satisfy



Note that the functor \mathscr{Q}^*_w is the composition

$$\mathscr{Q}_w^* \colon R\operatorname{-gmod} \xrightarrow{\psi_*} (R\operatorname{-gmod})^{\operatorname{rev}} \xrightarrow{\mathscr{Q}_w} (\widetilde{\mathscr{C}_w})^{\operatorname{rev}} \xrightarrow{\psi_*} \widetilde{\mathscr{C}_w^*}.$$


Note that the functor Φ_w is a faithful exact monoidal functor. In particular, a morphism f in \mathscr{C}_w is a monomorphism (resp. epimorphism, isomorphism) if and only if so is $\Phi_w(f)$.

Now let us show that $\mathscr{C}_w \to \widetilde{\mathscr{C}}_w$ is fully faithful. In order to see this, we start by the following proposition.

Proposition 8.1. Let C be a real simple module in R-gmod with an affinization (C, z). Let $X \in R$ -gmod and assume that $\mathbf{r}_{C,X} := \mathbf{R}_{C,X}^{\text{ren}}|_{z=0} : C \circ X \to X \circ C$ is an isomorphism up to a grading shift. Then for any submodule $Y \subset C \circ X$, there exists $X' \subset X$ such that $Y = C \circ X'$.

Proof. Set $\mathbf{r} := \mathbf{r}_{C,X}$.

Let us consider the following commutative diagram



Hence we have $C \circ \mathbf{r}(Y) \subset Y \circ C \subset C \circ X \circ C$. By the quasi-rigid property, there exists $X' \subset X$ such that

$$\mathbf{r}(Y) \subset X' \circ C$$
 and $C \circ X' \subset Y$.

Hence we have

$$\dim(Y) = \dim(\mathbf{r}(Y)) \leqslant \dim(X' \circ C) = \dim(C \circ X') \leqslant \dim(Y).$$

Since $\dim(C \circ X') = \dim(Y)$, we have $Y = C \circ X'$, as desired.

Corollary 8.2. Let $M \in \mathscr{C}_w$. For any $X \subset \Phi_w(M)$, there exists $X' \subset M$ such that $X = \Phi_w(X')$.

Theorem 8.3. $\Phi_w \colon \mathscr{C}_w \to \widetilde{\mathscr{C}}_w$ is fully faithful.

Proof. It is enough to show that

(8.3)
$$\operatorname{Hom}_{R\operatorname{-gmod}}(M, N) \longrightarrow \operatorname{Hom}_{R\operatorname{-gmod}}(\mathsf{C} \circ M, \mathsf{C} \circ N)$$

is bijective for any $M, N \in \mathscr{C}_w$ and $\mathsf{C} = \mathsf{M}(w\Lambda, \Lambda)$. The injectivity is evident. Let us show that any $f \in \operatorname{Hom}_{R\operatorname{-gmod}}(\mathsf{C} \circ M, \mathsf{C} \circ N)$ is in the image of $\operatorname{Hom}_{R\operatorname{-gmod}}(M, N)$.

Let $\varphi \colon \mathsf{C} \circ M \to (\mathsf{C} \circ M) \oplus (\mathsf{C} \circ N)$ be the graph of f, and let $Z \subset (\mathsf{C} \circ M) \oplus (\mathsf{C} \circ N) \simeq$ $\mathsf{C} \circ (M \oplus N)$ be the image of φ . Then, Proposition 8.1 implies that there exists $K \subset M \oplus N$ such that $Z = \mathsf{C} \circ K$. Let $p_1 \colon K \to M$ and $p_2 \colon K \to N$ be the projections. Since the composition $\mathsf{C} \circ M \xrightarrow{\sim} \mathsf{C} \circ K \xrightarrow{p_1} \mathsf{C} \circ M$ is the identity, the faithfulness of Φ_w implies that p_1 is an isomorphism. Since the composition $\mathsf{C} \circ M \xrightarrow{\sim} \mathsf{C} \circ K \xrightarrow{p_2} \mathsf{C} \circ N$ is equal to f, we obtain $f = \mathsf{C} \circ (p_2 p_1^{-1})$.

Corollary 8.4. The full subcategory \mathscr{C}_w of $\widetilde{\mathscr{C}}_w$ is stable by taking subquotients.

Proof. Let $Y \in \widetilde{\mathscr{C}}_w$ be a subobject of $X \in \mathscr{C}_w$. Then there exist $\Lambda \in \mathsf{P}_+$ such that $\mathsf{C}_\Lambda \circ Y \subset \mathsf{C}_\Lambda \circ X$ belongs to \mathscr{C}_w . Then Proposition 8.1 implies that $Y \in \mathscr{C}_w$. \Box

Remark 8.5. Since \mathscr{C}_w is a full subcategory of $\widetilde{\mathscr{C}}_w$, we sometimes regard an object of \mathscr{C}_w as an object of $\widetilde{\mathscr{C}}_w$, namely we identify $\Phi_w(M)$ and M for $M \in \mathscr{C}_w$.

Remark 8.6. The full subcategory \mathscr{C}_w is not stable by taking extensions in \mathscr{C}_w in general. Indeed taking a center $C = \mathsf{M}(w\Lambda, \Lambda)$, we have an extension of **1**:

$$0 \to C \mathrel{\circ} C^{-1} \longrightarrow (\widetilde{C}/z^2\widetilde{C}) \mathrel{\circ} C^{-1} \longrightarrow C \mathrel{\circ} C^{-1} \longrightarrow 0,$$

where (\widetilde{C}, z) is an affinization of C.

Note that $\widetilde{\mathscr{C}}_w$ is a rigid monoidal abelian category satisfying (6.2) with $\Lambda = \mathbb{Q}$. Recall that a real simple object of $\widetilde{\mathscr{C}}_w$ is affreal if it admits an affinization (see Definition 6.7). Then the following proposition immediately follows from Proposition 6.19 and Corollary 6.21.

Proposition 8.7. Let $X, Y \in \widetilde{\mathscr{C}_w}$ be simple objects. Assume that one of them is affreal. Then, we have

$$\operatorname{Hom}_{\widetilde{\mathscr{C}}_{w}}(X \circ Y, X \circ Y) = \mathbf{k} \operatorname{id}_{X \circ Y},$$
$$\operatorname{Hom}_{\widetilde{\mathscr{C}}_{w}}(X \circ Y, Y \circ X) = \mathbf{k} \mathbf{r}_{X Y}$$

for some non-zero $\mathbf{r}_{X,Y}$.

Recall that $\mathbf{r}_{X,Y}$ is the *R*-matrix between X and Y and denote by $\Lambda(X,Y)$ the degree of $\mathbf{r}_{X,Y}$. If $X \simeq X' \circ \mathsf{C}_{\lambda}$ for some $X' \in \mathscr{C}_w$, $\lambda \in \mathsf{P}$, and $Y \in \mathscr{C}_w$, then we have

$$\Lambda(X,Y) = \Lambda(X',Y) + \Lambda(\mathsf{C}_{\lambda},Y).$$

Note that $\Lambda(\mathsf{C}_{\lambda}, Y) = -(w\lambda + \lambda, \operatorname{wt}(Y)).$

Lemma 8.8. Let \mathcal{C} be either R-gmod or \mathscr{C}_w , and let $Q: \mathcal{C} \to \widetilde{\mathscr{C}_w}$ be the localization functor. If $(\mathsf{M}, R_{\mathsf{M}})$ is a rational center in \mathcal{C} , then there is a rational center $(Q(\mathsf{M}), R_{Q(\mathsf{M})})$ in $\widetilde{\mathscr{C}_w}$ such that $R_{Q(\mathsf{M})}(Q(X)) = Q(R_{\mathsf{M}}(X))$ for any $X \in \mathcal{C}$..

Proof. Let \mathcal{A} be the category whose object is a pair (X, R) where $X \in \widetilde{\mathscr{C}}_w$ and $R: Q(\mathsf{M}) \circ X \to X \circ Q(\mathsf{M})$ is an isomorphism in $\operatorname{Raff}_z(\widetilde{\mathscr{C}}_w)$. The morphisms of \mathcal{A} are defined in an obvious way so that we have a faithful functor $\mathcal{A} \to \widetilde{\mathscr{C}}_w$. Then,

 \mathcal{A} is an abelian category. It has a monoidal structure by $(X, R) \otimes (Y \otimes S) = (Z, T)$, where $Z = X \circ Y$ and T is the composition

$$T: Q(\mathsf{M}) \circ Z = Q(\mathsf{M}) \circ X \circ Y \xrightarrow{R} X \circ Q(\mathsf{M}) \circ Y \xrightarrow{S} X \circ Y \circ Q(\mathsf{M}) = Z \circ Q(\mathsf{M}).$$

Now we have a monoidal functor $F: \mathcal{C} \to \mathcal{A}$ defined by by

$$F(X) = (Q(X), Q(R_{\mathsf{M}}(X))) \text{ for } X \in \mathcal{C}.$$

Then it is easy to see that $F(\mathsf{C}_i)$ is an invertible object of \mathcal{A} for any $i \in I$ and $F(R_{\mathsf{C}_i}(X))$ is an isomorphism for any $X \in \mathcal{C}$. Hence, by the universal property of $\mathcal{C} \to \widetilde{\mathscr{C}}_w$, the functor $F: \mathcal{C} \to \mathcal{A}$ extends to $\widetilde{F}: \widetilde{\mathscr{C}}_w \to \mathcal{A}$.

Now, for any $X \in \widetilde{\mathscr{C}}_w$, define $R_{Q(\mathsf{M})}(X)$ by $R_{Q(\mathsf{M})}(X) = R$, where $\widetilde{F}(X) = (X, R) \in \mathcal{A}$. Then $(Q(\mathsf{M}), R_{Q(\mathsf{M})})$ is a rational center in $\widetilde{\mathscr{C}}_w$.

9. Resultant Algebra and Affinization of Invariants

9.1. **Resultant algebra.** Let A be a commutative **k**-algebra. Let z be an indeterminate. Recall that a polynomial $f(z) = \sum_{k=0}^{n} a_k z^{n-k} \in A[z]$ is a quasi-monic (resp. monic) polynomial of degree n if $a_k \in A$ and $a_0 \in \mathbf{k}^{\times}$ (resp. $a_0 = 1$). For a quasi-monic polynomial f(z) of degree m and a quasi-monic polynomial g(z) of degree n, we write their resultant $f \star g$ for

$$\prod_{j=1}^{m} \prod_{k=1}^{n} (x_j - y_k) \equiv \prod_{j=1}^{m} g(x_j) \equiv \prod_{k=1}^{n} f(y_k) \mod \mathbf{k}^{\times},$$

writing formally $f(z) = a \prod_{j=1}^{m} (z - x_j)$ and $g(z) = b \prod_{k=1}^{n} (z - y_k)$ with $a, b \in \mathbf{k}^{\times}$. Here $B \equiv C \mod \mathbf{k}^{\times}$ means that B = cC for some $c \in \mathbf{k}^{\times}$. It is well-defined since any symmetric polynomial of $\{x_j\}_{1 \leq j \leq m}$ is expressed by the coefficients of f(z).

We regard $f \star g$ as an element of A/\mathbf{k}^{\times} .

The resultant product satisfies:

$$f(z) \star g(z) \equiv g(z) \star f(z),$$

$$(f_1(z) \cdot f_2(z)) \star g(z) \equiv (f_1(z) \star g(z)) (f_2(z) \star g(z)).$$

A quasi-monic rational function with coefficients in A is a quotient f(z)/g(z) for quasi-monic homogeneous polynomials f(z) and g(z) with coefficients in A. We can consider them as an element of $A((z^{-1}))$, the ring of Laurent series in z^{-1} with coefficients in A.

We write \star_z when want to emphasize z.

A quasi-monic rational homogeneous function in z and z' with coefficients in A is a quotient f(z, z')/g(z, z') for quasi-monic homogeneous polynomials f(z, z') and g(z, z') in z and z' with coefficients in A.

Let A be a commutative graded **k**-algebra such that $A_{\leq 0} = \mathbf{k}$. For homogeneous indeterminates z_k (k = 1, ..., r) of positive degree, we denote by $\mathcal{M}(z_1, ..., z_r) \subset A[z_1, ..., z_r]/\mathbf{k}^{\times}$ the set of homogeneous polynomials in $z_1, ..., z_r$ with coefficients in A and quasi-monic in each $z_1, ..., z_k$ modulo \mathbf{k}^{\times} . Then we can define

$$\star_{z_k}: \mathcal{M}(z_1, z_2, \dots, z_k) \times \mathcal{M}(z_k, \dots, z_r) \to \mathcal{M}(z_1, \dots, z_{k-1}, z_{k+1}, \dots, z_r).$$

The homogeneous degree of $f \underset{z_k}{\star} g$ for $f \in \mathcal{M}(z_1, z_2, \ldots, z_k)$ and $g \in \mathcal{M}(z_k, \ldots, z_r)$ is given by $\frac{\deg(f) \deg(g)}{\deg(z_k)}$, where $\deg(z_k)$, $\deg(f)$, and $\deg(g)$ denote the homogeneous degrees of z_k , f and g, respectively.

In particular, if z and w be indeterminates with the same degree, $\mathcal{M}(z, w)$ has a structure of semiring (the existence of additive inverse is not assumed): the addition in $\mathcal{M}(z, w)$ is the product in $\mathbf{k}[z, w]$, and the multiplication in $\mathcal{M}(z, w)$ is given as follows: the multiplication h(z, w) of f(z, w) and g(z, w) is given by $h(z_1, z_3) \equiv f(z_1, z_2) \underset{z_2}{\star} g(z_2, z_3)$. We call $\mathcal{M}(z, w)$ the resultant algebra. The multiplication in $\mathcal{M}(z, w)$ is commutative. Let us denote by

$$\deg\colon \mathcal{M}(z,w)\to\mathbb{Z}$$

the map of taking the homogeneous degree. Then $f \mapsto \deg(f)/\deg(z)$ is a semiring homomorphism.

9.2. Affinizations of invariants for R -modules. Let R be a quiver Hecke algebra as in section 7. For $i \in I$ and $n \in \mathbb{Z}_{>0}$, let $P(i^n)$ be the indecomposable projective $R(n\alpha_i)$ -module whose head is isomorphic to $L(i^n)$. Then, for an $R(\beta)$ -module M we define

$$E_i^{(n)}M := \operatorname{Hom}_{R(n\alpha_i)}(P(i^n), e(n\alpha_i, \beta - n\alpha_i)M) \in \operatorname{Mod}(R(\beta - n\alpha_i)),$$

$$F_i^{(n)}M := P(i^n) \circ M \in \operatorname{Mod}(R(\beta + n\alpha_i)).$$

The functors $E_i^{(1)}$ and $F_i^{(1)}$ will be denoted by E_i and F_i .

For $i \in I$ and a non-zero $R(\beta)$ -module M, we define

$$\operatorname{wt}(M) = -\beta, \quad \varepsilon_i(M) = \max\{k \ge 0 \mid E_i^k M \not\simeq 0\}, \quad \varphi_i(M) = \varepsilon_i(M) + \langle h_i, \operatorname{wt}(M) \rangle.$$

We also define E_i^* , F_i^* , ε_i^* , etc. in the same manner as above by replacing $e(n\alpha_i, \beta - n\alpha_i)$, $P(i^n) \circ -$, etc. with $e(\beta - n\alpha_i, n\alpha_i), - \circ P(i^n)$, etc.

Definition 9.1. Let $i \in I$ and let M be an $R(\beta)$ -module. Set $m = \varepsilon_i(M), m' = \varepsilon_i^*(M),$ $n = \operatorname{ht}(\beta)$. We set

(9.1)
$$\begin{cases} \chi_i(M)(t_i) = \left(\sum_{\nu \in I^{\beta}} \prod_{\nu_k = i} (t_i - x_k) e(\nu)\right) \Big|_M \in \operatorname{END}_R(M)[t_i], \\ \mathscr{E}_i(M)(t_i) = (t_i - x_1) \cdots (t_i - x_m) \Big|_{E_i^{(m)}M} \in \operatorname{END}_R(E_i^{(m)}M)[t_i], \\ \mathscr{E}_i^*(M)(t_i) = (t_i - x_{n-m'+1}) \cdots (t_i - x_n) \Big|_{E_i^{*(m')}M} \in \operatorname{END}_R(E_i^{*(m')}M)[t_i]. \end{cases}$$

Note that, with the assignment $\deg(t_i) = (\alpha_i, \alpha_i)$, the monic polynomials $\chi_i(M)$, $\mathscr{E}_i(M)$ and $\mathscr{E}_i^*(M)$ are homogeneous.

For an affine object (M, z_M) of a real simple module, we have

$$\chi_i(\mathsf{M}), \mathscr{E}_i(\mathsf{M}), \mathscr{E}_i^*(\mathsf{M}) \in \mathbf{k}[t_i, z_\mathsf{M}],$$

and they satisfy

$$\chi_{i}(\mathsf{M}) \equiv \mathscr{E}_{i}(\mathsf{M}) \cdot \chi_{i}(E_{i}^{(m)}\mathsf{M}) \equiv \mathscr{E}_{i}^{*}(\mathsf{M}) \cdot \chi_{i}(E_{i}^{*(m')}\mathsf{M})$$

`

where $m = \varepsilon_i(\mathsf{M})$ and $m' = \varepsilon_i^*(\mathsf{M})$ (see [13, Lemma 2.7 (i)] and [11, Lemma 3.3]). Recall that for quasi-monic polynomials f and g, " $f \equiv g$ " means " $f \equiv g \mod \mathbf{k}^{\times n}$.

Lemma 9.2. Let $M \in Mod(R(\beta))$, $N \in Mod(R(\gamma))$ with $\beta, \gamma \in Q_+$, $m = ht(\beta)$, $n = ht(\gamma)$. Then for $u \in M$ and $v \in N$, we have

$$\mathbf{R}_{M,N}^{\mathrm{uni}}(u \boxtimes v) \in \tau_{w[n,m]} \prod_{i \in I} \chi_i(N) \mathop{\star}_{t_i} \chi_i(M) \big(v \boxtimes u \big) + \sum_{w < w[n,m]} \tau_w(N \boxtimes M)$$

Proof. By [7, (1.17)], we have

$$\mathbf{R}_{M,N}^{\mathrm{uni}}(u \boxtimes v) \in \tau_{w[n,m]} \sum_{\nu \in I^{\beta+\gamma}} \prod_{1 \leq a \leq n < b \leq n+m, \atop \nu_a = \nu_b} (x_a - x_b) e(\nu) (v \boxtimes u) + \sum_{w < w[n,m]} \tau_w(N \boxtimes M).$$

Definition 9.3. Let (M, z_M) be an affinization of a real simple module M, and let (N, z_N) be an affinization of a real simple N.

(a) Recall that $\mathscr{D}(\mathsf{M},\mathsf{N}) \in \mathbf{k}[z_{\mathsf{M}},z_{\mathsf{N}}]$ the quasi-monic polynomial given by

$$\mathrm{R}^{\mathrm{ren}}_{\mathsf{N},\mathsf{M}}\circ\mathrm{R}^{\mathrm{ren}}_{\mathsf{M},\mathsf{N}}=\mathscr{D}(\mathsf{M},\mathsf{N})\,\mathsf{id}_{\mathsf{M}\circ\mathsf{N}}$$

(b) Denote by $\widetilde{\mathscr{L}}(\mathsf{M},\mathsf{N}) \in \mathbf{k}[z_{\mathsf{M}}, z_{\mathsf{N}}]$ the quasi-monic polynomial given by

$$\mathbf{R}_{\mathsf{M},\mathsf{N}}^{\mathrm{ren}}(u\boxtimes v)\in\widetilde{\mathscr{S}}(\mathsf{M},\mathsf{N})\tau_{w[n,m]}(v\boxtimes u)+\sum_{w< w[n,m]}\tau_w(\mathsf{N}\boxtimes\mathsf{M})$$

for $u \in \mathsf{M}$ and $v \in \mathsf{N}$. Note that by Proposition 6.2, $\mathbb{R}^{\text{ren}}_{\mathsf{M},\mathsf{N}}$ is proportional to $\mathbb{R}^{\text{uni}}_{\mathsf{M},\mathsf{N}}$ and hence such a quasi-monic polynomial $\widetilde{\mathscr{L}}(\mathsf{M},\mathsf{N})$ exists by Lemma 9.2.

(c) We set

$$\mathscr{W}t(\mathsf{M},\mathsf{N}) := \frac{\prod_{i \in I} (\chi_i(\mathsf{M}) \underset{t_i}{\star} \chi_i(\mathsf{N}))^2}{\prod_{i \neq j} \chi_i(\mathsf{M}) \underset{t_i}{\star} Q_{i,j}(t_i, t_j) \underset{t_j}{\star} \chi_j(\mathsf{N})} \in \mathbf{k}(z_{\mathsf{M}}, z_{\mathsf{N}})/\mathbf{k}^{\times}$$
$$\mathscr{L}(\mathsf{M},\mathsf{N}) := \widetilde{\mathscr{L}}(\mathsf{M},\mathsf{N})^2/\mathscr{W}t(\mathsf{M},\mathsf{N}) \in \mathbf{k}(z_{\mathsf{M}}, z_{\mathsf{N}})/\mathbf{k}^{\times}.$$

Note that $\mathscr{W}t(\mathsf{M},\mathsf{N}) \equiv \mathscr{W}t(\mathsf{N},\mathsf{M})$.

For simples M, N in R-gmod such that one of them is affreal, define

$$\widetilde{\Lambda}(M,N) := \frac{1}{2} \big(\Lambda(M,N) + (\operatorname{wt}(M),\operatorname{wt}(N)) \big).$$

The homogeneous degrees of these invariants are given as follows:

$$\begin{split} & \operatorname{deg} \left(\chi_i(\mathsf{M}) \right) = n_i(\alpha_i, \alpha_i) \quad \operatorname{where } \operatorname{wt}(M) = -\sum_{i \in I} n_i \alpha_i, \\ & \operatorname{deg} \left(\mathscr{E}_i(\mathsf{M}) \right) = \varepsilon_i(M)(\alpha_i, \alpha_i) = 2 \widetilde{\Lambda}(L(i), M), \\ & \operatorname{deg} \left(\mathscr{E}_i^*(\mathsf{M}) \right) = \varepsilon_i^*(M)(\alpha_i, \alpha_i) = 2 \widetilde{\Lambda}(M, L(i)), \\ & \operatorname{deg} \left(\mathscr{D}(\mathsf{M}, \mathsf{N}) \right) = 2 \mathfrak{d}(M, N), \\ & \operatorname{deg} \left(\widetilde{\mathscr{L}}(\mathsf{M}, \mathsf{N}) \right) = 2 \widetilde{\Lambda}(M, N), \\ & \operatorname{deg} \left(\mathscr{U}(\mathsf{M}, \mathsf{N}) \right) = 2 \Lambda(M, N), \\ & \operatorname{deg} \left(\mathscr{W}t(\mathsf{M}, \mathsf{N}) \right) = 2 \left(\operatorname{wt}(M), \operatorname{wt}(N) \right). \end{split}$$

Indeed, the fifth equality is shown in [11, Lemma 3.11]. For the last equality we have $\deg(\mathscr{W}t(\mathsf{M},\mathsf{N}))$

$$= \sum_{i \in I} \left(2 \operatorname{deg}(\chi_i(\mathsf{M})) \operatorname{deg}(\chi_i(\mathsf{N})) / \operatorname{deg}(t_i)) - \sum_{i \neq j} \left(\operatorname{deg}(\chi_i(\mathsf{M})) \operatorname{deg}(\chi_j(\mathsf{N})) \frac{\operatorname{deg}(Q_{i,j}(t_i, t_j))}{\operatorname{deg}(t_i) \operatorname{deg}(t_j)} \right) \\ = 2 \sum_{i,j \in I} \left(\frac{\operatorname{deg}(\chi_i(\mathsf{M}))}{(\alpha_i, \alpha_i)} \frac{\operatorname{deg}(\chi_j(\mathsf{N}))}{(\alpha_j, \alpha_j)} (\alpha_i, \alpha_j) \right) = 2 \left(\operatorname{wt}(\mathsf{M}), \operatorname{wt}(\mathsf{N}) \right).$$

The others follow directly from their definitions. See also [9, Corollary 3.8] for the second and the third equalities.

Therefore, these invariants are affinizations of wt, \mathfrak{d} , $\widetilde{\Lambda}$, etc.

By Lemma 9.2 and Definition 9.3 (b), one has

(9.2)
$$\mathbf{R}_{\mathsf{M},\mathsf{N}}^{\mathrm{uni}} = \frac{\prod_{i \in I} \chi_i(\mathsf{M}) \star_{t_i} \chi_i(\mathsf{N})}{\widetilde{\mathscr{L}}(\mathsf{M},\mathsf{N})} \mathbf{R}_{\mathsf{M},\mathsf{N}}^{\mathrm{ren}}.$$

Lemma 9.4. We have

$$\begin{split} \widetilde{\mathscr{L}}(\mathsf{M},\mathsf{N})\cdot\widetilde{\mathscr{L}}(\mathsf{N},\mathsf{M}) &\equiv \mathscr{D}(\mathsf{M},\mathsf{N})\cdot\mathscr{W}t(\mathsf{M},\mathsf{N}),\\ \mathscr{D}(\mathsf{M},\mathsf{N})^2 &\equiv \mathscr{L}(\mathsf{M},\mathsf{N})\cdot\mathscr{L}(\mathsf{N},\mathsf{M}). \end{split}$$

Proof. By [7, Proposition 1.10 (iv)] for $u \in M$ and $v \in N$, we have

(9.3)
$$R_{\mathsf{N},\mathsf{M}}^{\mathrm{uni}} R_{\mathsf{M},\mathsf{N}}^{\mathrm{uni}}(u \boxtimes v) = \sum_{\substack{1 \leqslant a \leqslant m < b \leqslant m+n, \\ \nu \in I^{\beta+\gamma}, \nu_a \neq \nu_b}} Q_{\nu_a,\nu_b}(x_a, x_b) e(\nu) (u \boxtimes v)$$
$$= \left(\prod_{i \neq j} \chi_i(\mathsf{M}) \underset{t_i}{\star} Q_{i,j}(t_i, t_j) \underset{t_j}{\star} \chi_j(\mathsf{N})\right) (u \boxtimes v)$$

On the other hand, (9.2) implies

$$\mathbf{R}_{\mathsf{N},\mathsf{M}}^{\mathrm{uni}}\mathbf{R}_{\mathsf{M},\mathsf{N}}^{\mathrm{uni}}(u\boxtimes v) = \frac{\prod_{i\in I} \left(\chi_i(\mathsf{M}) \underset{t_i}{\star} \chi_i(\mathsf{N})\right)^2}{\widetilde{\mathscr{L}}(\mathsf{M},\mathsf{N})\widetilde{\mathscr{L}}(\mathsf{N},\mathsf{M})} \mathbf{R}_{\mathsf{N},\mathsf{M}}^{\mathrm{ren}}\mathbf{R}_{\mathsf{M},\mathsf{N}}^{\mathrm{ren}}(u\boxtimes v)$$
$$= \frac{\prod_{i\in I} \left(\chi_i(\mathsf{M}) \underset{t_i}{\star} \chi_i(\mathsf{N})\right)^2}{\widetilde{\mathscr{L}}(\mathsf{M},\mathsf{N})\widetilde{\mathscr{L}}(\mathsf{N},\mathsf{M})} \mathscr{D}(\mathsf{M},\mathsf{N})(u\boxtimes v).$$

Hence, we obtain

$$\prod_{i \neq j} \chi_i(\mathsf{M}) \mathop{\star}_{t_i} Q_{i,j}(t_i, t_j) \mathop{\star}_{t_j} \chi_j(\mathsf{N}) = \frac{\prod_{i \in I} (\chi_i(\mathsf{M}) \mathop{\star}_{t_i} \chi_i(\mathsf{N}))^2}{\widetilde{\mathscr{L}}(\mathsf{M}, \mathsf{N}) \widetilde{\mathscr{L}}(\mathsf{N}, \mathsf{M})} \mathscr{D}(\mathsf{M}, \mathsf{N}),$$

which implies the desired results.

Lemma 9.5. Let (L, w) , (M, z) , (N, z) , (X, z) be affinizations of simples L, M, N, X in R-gmod, respectively, and assume that L is real. If there is an epimorphism $\mathsf{M} \circ \mathsf{N} \to \mathsf{X}$ in $\operatorname{Pro}_{\operatorname{coh}}(\mathbf{k}[z], R$ -gmod), then there exists $a \in \mathbf{k}[z, w] \setminus \{0\}$ such that the diagram below

is commutative.

$$\begin{array}{c|c} L \mathrel{\circ} M \mathrel{\circ} N \xrightarrow[\mathrm{R^{ren}_{L,M}}]{} M \mathrel{\circ} L \mathrel{\circ} N \xrightarrow[\mathrm{R^{ren}_{L,N}}]{} M \mathrel{\circ} N \mathrel{\circ} L \\ & \downarrow \\ \downarrow \\ L \mathrel{\circ} X \xrightarrow[a \mathrel{R^{ren}_{L,X}}]{} X \mathrel{\circ} L. \end{array}$$

In particular, we have $\widetilde{\mathscr{L}}(\mathsf{L},\mathsf{M})\widetilde{\mathscr{L}}(\mathsf{L},\mathsf{N}) = a\widetilde{\mathscr{L}}(\mathsf{L},\mathsf{X})$. If we further assume that $\Lambda(L,X) = \Lambda(L,M) + \Lambda(L,N)$, then $a \equiv 1$ in $\mathbf{k}[z,w]$.

Proof. Since $\operatorname{Hom}_{\mathbf{k}[z,w]}(\mathsf{L} \circ \mathsf{X}, \mathsf{X} \circ \mathsf{L}) = \mathbf{k}[z,w]\operatorname{R}_{\mathsf{L},\mathsf{X}}^{\operatorname{ren}}$ by Proposition 6.2, such an $a \in \mathbf{k}[z,w]$ exists. The other assertions are immediate.

Lemma 9.6. Let $i \in I$, $M \in R(m\alpha_i)$ -gmod, and $N \in R(n\alpha_i)$ -gmod. Assume that $(\mathsf{M}, z_{\mathsf{M}})$ and $(\mathsf{N}, z_{\mathsf{N}})$ are affinizations of M and N respectively. Then $\mathrm{R}_{\mathsf{M},\mathsf{N}}^{\mathrm{ren}} = \mathrm{R}_{\mathsf{M},\mathsf{N}}^{\mathrm{uni}}$ and $\widetilde{\mathscr{L}}(\mathsf{M}, \mathsf{N}) \equiv \chi_i(\mathsf{M}) \underset{t_i}{\star} \chi_i(\mathsf{N})$.

Proof. By (9.3), $\mathbb{R}_{M,N}^{\text{uni}} \circ \mathbb{R}_{N,M}^{\text{uni}} = \text{id}$ and hence we have $\mathbb{R}_{M,N}^{\text{ren}} = \mathbb{R}_{M,N}^{\text{uni}}$. By (9.2), we have $\widetilde{\mathscr{L}}(\mathsf{M},\mathsf{N}) \equiv \chi_i(\mathsf{M}) \underset{t_i}{\star} \chi_i(\mathsf{N})$.

Proposition 9.7. Let M be an affinization of a simple module M in R-gmod. Then for each $i \in I$, we have

(9.4)

$$\mathscr{E}_{i}(\mathsf{M}) \equiv \widetilde{\mathscr{L}}(\widetilde{L}(i)_{t_{i}}, \mathsf{M}),$$

$$\mathscr{E}_{i}^{*}(\mathsf{M}) \equiv \widetilde{\mathscr{L}}(\mathsf{M}, \widetilde{L}(i)_{t_{i}}),$$

$$\mathscr{W}t(\widetilde{L}(i)_{t_{i}}, \mathsf{M}) \equiv \frac{\chi_{i}(\mathsf{M})^{2}}{\prod_{j \neq i} Q_{i,j}(t_{i}, t_{j}) \underset{t_{j}}{\star} \chi_{j}(\mathsf{M})}$$

Here $\widetilde{L}(i)_{t_i}$ is the affinization of L(i) given by $R(\alpha_i)$ with $t_i = x_1$.

Proof. Let $m := \varepsilon_i(M)$, $M_0 := E_i^{(m)}M$, and $\mathsf{M}_0 := E_i^{(m)}\mathsf{M}$ such that M_0 is an affinization of M_0 . There exists an affinization L of $L(i^m)$ such that $\chi_i(\mathsf{L}) = \mathscr{E}_i(\mathsf{M})$ by [11,

Proposition 3.6]. Then we have an epimorphism $L \circ M_0 \rightarrow M$ due to the presentation of L in [11, Remark 3.8]. Hence we have

$$\widetilde{\mathscr{L}}(\langle i \rangle_{t_i}, \mathsf{M}) \equiv \widetilde{\mathscr{L}}(\langle i \rangle_{t_i}, \mathsf{L}) \, \widetilde{\mathscr{L}}(\langle i \rangle_{t_i}, \mathsf{M}_0) \equiv \widetilde{\mathscr{L}}(\langle i \rangle_{t_i}, \mathsf{L}) = \chi_i(\mathsf{L}) = \mathscr{E}_i(\mathsf{M}).$$

Here the first equality follows from [12, Corollary 2.12] and Lemma 9.5 together with the fact that $\langle i \rangle$ and M_0 are unmixed. The second equality follows from Lemma 9.10 and the third equality follows from Lemma 9.6.

The second assertion can be proved in a similar way, and the last assertion is immediate. $\hfill \Box$

Proposition 9.8. Let $i \in I$, and let (M, z_M) be an affinization of a simple module M. Set $m = \varepsilon_i(M)$ and $M_0 = E_i^{(m)} M$. Then we have

$$\mathscr{D}(\mathsf{M},\widetilde{L}(i)_{t_i})\chi_i(\mathsf{M})\chi_i(\mathsf{M}_0) \equiv \mathscr{E}_i^*(\mathsf{M})\prod_{j\neq i}Q_{ij}(t_i,t_j) \underset{t_j}{\star} \chi_j(\mathsf{M}_0).$$

Proof. Set $N = \widetilde{L}(i)_{t_i}$. Then Lemma 9.4 implies

$$\mathscr{L}(\mathsf{M},\mathsf{N})\cdot\mathscr{L}(\mathsf{N},\mathsf{M})\equiv\mathscr{D}(\mathsf{M},\mathsf{N})\cdot\mathscr{W}t(\mathsf{M},\mathsf{N}),$$

which reads as

$$\mathscr{E}_{i}(\mathsf{M})\mathscr{E}_{i}^{*}(\mathsf{M}) \equiv \mathscr{D}(\mathsf{M},\mathsf{N}) \cdot \frac{\chi_{i}(\mathsf{M})^{2}}{\prod_{j \neq i} Q_{i,j}(t_{i},t_{j}) \underset{t_{j}}{\star} \chi_{j}(\mathsf{M})}$$
$$\equiv \mathscr{D}(\mathsf{M},\mathsf{N}) \cdot \frac{\chi_{i}(\mathsf{M})\mathscr{E}_{i}(\mathsf{M})\chi_{i}(\mathsf{M}_{0})}{\prod_{j \neq i} Q_{i,j}(t_{i},t_{j}) \underset{t_{j}}{\star} \chi_{j}(\mathsf{M}_{0})}$$

Hence, we obtain the desired result.

Corollary 9.9. Let $i \in I$, and let $(\mathsf{M}, z_{\mathsf{M}})$ be an affinization of a simple module M. Set $m = \varepsilon_i(M)$ and $\mathsf{M}_0 = E_i^{(m)} \mathsf{M}$. We assume further that $\mathfrak{d}_i(M) = 0$, i.e., M and L(i) commute. (Note that $\mathfrak{d}_i(M) := \varepsilon_i(M) + \varepsilon_i^*(M) + \langle h_i, \operatorname{wt}(M) \rangle$.) Then we have

$$\chi_i(\mathsf{M})\chi_i(\mathsf{M}_0) \equiv \mathscr{E}_i^*(\mathsf{M})\prod_{j\neq i}Q_{ij}(t_i,t_j) \underset{t_j}{\star} \chi_j(\mathsf{M}_0).$$

Proof. It follows from $\mathfrak{d}(M, L(i)) = 0$, which implies $\mathscr{D}(\mathsf{M}, \widetilde{L}(i)_{t_i}) \equiv 1$.

We say that an ordered pair (M, N) of *R*-modules is *unmixed* if

 $\mathsf{W}^*(M) \cap \mathsf{W}(N) \subset \{0\}.$

Lemma 9.10. Let M and N be affinizations in *R*-gmod of real simple modules *M* and *N* respectively. If (M, N) is unmixed, then $\widetilde{\mathscr{L}}(M, N) \equiv 1$.

Proof. By [12, Corollary 2.12], we have $\widetilde{\Lambda}(M, N) = 0$ and hence $\widetilde{\mathscr{L}}(\mathsf{M}, \mathsf{N}) \equiv 1$. \Box

9.3. Examples of affinizations. The following result will be used later.

Lemma 9.11. Let $c \in \mathbb{Z}_{>0}$. Assume that $Q_{i,j}(u, v) = v - u^c$ with $\langle h_i, \alpha_j \rangle = -c$ and $\langle h_j, \alpha_i \rangle = -1$.

For $m \in \mathbb{Z}$ such that $0 \leq m \leq c$, let $L = \langle i^m \rangle$. Let (L, z) be an affinization of L with $\deg(z) = (\alpha_i, \alpha_i)$ such that $\chi_i(\mathsf{L})(t_i) = f(t_i, z)$ where $f(t_i, z)$ is quasi-monic with degree m in t_i and also in z (see [11, Proposition 3.6]).

- (i) $M := L \nabla \langle j \rangle \simeq L \otimes \langle j \rangle$ as an $R(m\alpha_i) \boxtimes R(\alpha_j)$ -module. Moreover, τ_m acts by 0 on $L \nabla \langle j \rangle$.
- (ii) If $f(t_i, z)$ divides $Q_{i,j}(t_i, z^c)$, then $\mathsf{M} := \mathsf{L} \otimes_{\mathbf{k}[z]} \langle j \rangle_{z^c}$ has a structure of $R(m\alpha_i + \alpha_j)$ -module with $\tau_m = 0$. Moreover M is an affinization of M.

Proof. (i) We can check easily the defining relation of $R(m\alpha_i + \alpha_j)$ on M. In particular, the relation $\tau_m^2 = Q_{i,j}(x_m, x_{m+1})$ follows from $x_m^m|_L = 0$.

(ii) We can check easily the defining relation of $R(m\alpha_i + \alpha_j)$ on M. In particular, the relation $\tau_m^2 = Q_{i,j}(x_m, x_{m+1})$ is a consequence of $Q_{i,j}(x_m, x_{m+1})|_{\mathsf{M}} = 0$ which follows from $f(x_m, z)|_{\mathsf{L}} = 0$ and $Q_{i,j}(x_m, x_{m+1}) = Q_{i,j}(x_m, z^c)$ on M.

10. Saito Reflection functors for R-gmod via localization

10.1. $\mathscr{C}_{s_iw_0}$ and $\mathscr{C}^*_{s_iw_0}$. In the sequel, we assume that $C = {c_{i,j}}_{i,j\in I}$ is a Cartan matrix of finite type such that C is not of type A_1 .

Let w_0 be the longest element of the Weyl group W. Let $i \mapsto i^*$ be the involution on I determined by $w_0(\alpha_i) = -\alpha_{i^*}$.

Throughout this section, we fix $i \in I$.

We have

$$\mathscr{C}_{s_iw_0} = \{ M \in R \text{-gmod} \mid E_i M \simeq 0 \},$$
$$\mathscr{C}^*_{s_iw_0} = \{ M \in R \text{-gmod} \mid E_i^* M \simeq 0 \},$$

where E_i and E_i^* denote the functor $R(\beta)$ -gmod $\rightarrow R(\beta - \alpha_i)$ -gmod given by $M \mapsto e(\alpha_i, \beta - \alpha_i)M$ and $M \mapsto e(\beta - \alpha_i, \alpha_i)M$, respectively.

Note that $\mathscr{C}_{s_iw_0} \simeq \mathscr{C}^*_{s_iw_0} \simeq 0$ in case C is of type A_1 .

Then $\{\mathsf{M}(s_i w_0 \Lambda, \Lambda)\}_{\Lambda \in \mathsf{P}_+}$ is a family of central objects in $\mathscr{C}_{s_i w_0}$, and $\{\mathsf{M}(w_0 \Lambda, s_i \Lambda)\}_{\Lambda \in \mathsf{P}_+}$ is a family of central objects in $\mathscr{C}^*_{s_i w_0}$. Note that

$$\mathsf{M}(s_i w_0 \Lambda, \Lambda) = E_i^{(\langle h_i *, \Lambda \rangle)} \mathsf{M}(w_0 \Lambda, \Lambda) \quad \text{and} \quad \mathsf{M}(w_0 \Lambda, s_i \Lambda) = E_i^{*(\langle h_i, \Lambda \rangle)} \mathsf{M}(w_0 \Lambda, \Lambda).$$

Set $C_{\Lambda} := \mathsf{M}(s_i w_0 \Lambda, \Lambda)$ and $\mathsf{C}^*_{\Lambda} := \mathsf{M}(w_0 \Lambda, s_i \Lambda)$ for $\Lambda \in \mathsf{P}_+$. We set $\mathsf{C}_j = \mathsf{C}_{\Lambda_j}$ and $\mathsf{C}^*_j = \mathsf{C}^*_{\Lambda_j}$. Since we have

$$\psi_*(\mathsf{M}(v\Lambda, v'\Lambda)) \simeq \mathsf{M}(-v'\Lambda, -v\Lambda)$$

for any $\Lambda \in \mathsf{P}_+$ and any $v, v' \in \mathsf{W}$ such that $v \ge v'$ ([12, Lemma 2.23]), we have

$$\psi_*(\mathsf{C}_j) \simeq \mathsf{C}^*_{i^*} \quad \text{for any } j \in I.$$

(For ψ , see (8.2).) Then we have (see [11, §5.1] and [9, Corollary 3.8])

$$\Lambda(\mathsf{C}_{\Lambda}, \langle j \rangle) = (s_i w_0 \Lambda + \Lambda, \alpha_j),$$

$$\Lambda(\langle j \rangle, \mathsf{C}^*_{\Lambda}) = -(w_0 \Lambda + s_i \Lambda, \alpha_j),$$
 for any $j \in I.$

We have

$$\begin{aligned} \mathscr{Q}_{s_i w_0}(\langle i \rangle) &\simeq \mathsf{C}_{i^*}^{\circ - 1} \circ \big(\mathsf{C}_{i^*} \nabla \langle i \rangle\big), \\ \mathscr{Q}_{s_i w_0}^*(\langle i \rangle) &\simeq \big(\langle i \rangle \nabla \mathsf{C}_i^*\big) \circ \mathsf{C}_i^{* \circ - 1}. \end{aligned}$$

10.2. Schur-Weyl datum. For any $j \in I$, define an object of $\widetilde{\mathscr{C}}_{s_iw_0}^*$:

$$\mathbf{K}_{j} := \begin{cases} \mathscr{D}\left(\mathscr{Q}_{s_{i}w_{0}}^{*}(\langle i \rangle)\right) \simeq \left(E_{i} \mathbf{C}_{i^{*}}^{*}\right) \circ \mathbf{C}_{i^{*}}^{*} \circ^{-1} & \text{if } j = i, \\ \mathbf{M}(s_{i}s_{j}\Lambda_{j},\Lambda_{j}) \simeq \langle i^{-\mathbf{c}_{i,j}} \rangle \nabla \langle j \rangle & \text{if } j \neq i. \end{cases}$$

Recall that \mathscr{D} is the right dual functor. Note that $K_j \simeq \langle i^{-\mathbf{c}_{i,j}} \rangle \otimes \langle j \rangle$ as a vector space for $j \neq i$ since $E_k^*(K_j) \simeq 0$ for any $k \in I \setminus \{j\}$.

Since $\langle i \rangle = L(i) = R(\alpha_i)/R(\alpha_i)x_1$ and $\langle i^{-c_{i,j}} \rangle \nabla \langle j \rangle \simeq \mathsf{M}(s_i s_j \Lambda_j, \Lambda_j)$ have the canonical affinizations by [11, Theorem 3.26], we have the following affinizations of K_j 's by Lemma 6.6 and Lemma 8.8.

(10.1)
$$\widehat{\mathbf{K}}_{j} := \begin{cases} \mathscr{D}_{\mathrm{aff}} \left(\mathscr{Q}_{s_{i}w_{0}}^{*}(\langle i \rangle_{z_{i}}) \right) & \text{if } j = i, \\ \left(\left\langle i^{-\mathsf{c}_{i,j}} \right\rangle \nabla \left\langle j \right\rangle \right)_{z_{j}} & \text{if } j \neq i. \end{cases}$$

Here the functor $\mathscr{Q}^*_{s_iw_0}$: $\operatorname{Pro}(R\operatorname{-gmod}) \to \operatorname{Pro}(\widetilde{\mathscr{C}^*}_{s_iw_0})$ is induced by $\mathscr{Q}^*_{s_iw_0}$: $R\operatorname{-gmod} \to \widetilde{\mathscr{C}^*}_{s_iw_0}$. Note that deg $z_j = (\alpha_j, \alpha_j)$ for any $j \in I$. For $j \in I \setminus \{i\}$, we have

(10.2)
$$\chi_k(\widehat{\mathbf{K}}_j)(t_k) \equiv \begin{cases} t_j - z_j & \text{if } k = j, \\ Q_{i,j}(t_i, z_j) & \text{if } k = i, \\ 1 & \text{otherwise} \end{cases}$$

Note that $\chi_i(\widehat{\mathbf{K}}_j) \equiv Q_{i,j}(t_i, z_j)$ follows also from Corollary 9.9. We have

$$E_i \mathsf{C}^*_{i^*} \simeq \mathsf{M}(s_i w_0 \Lambda_{i^*}, s_i \Lambda_{i^*})$$

(Note that $s_i \leq s_i w_0$ since $\mathfrak{g} \neq A_1$.)

Note that

$$\Lambda(\langle j \rangle, \langle k \rangle) = -(\alpha_j, \alpha_k) \delta(j \neq k) \quad \text{for any } j, k \in I.$$

Proposition 10.1. For any $j, k \in I$, we have

$$\Lambda(\mathbf{K}_j,\mathbf{K}_k) = \Lambda(\langle j \rangle, \langle k \rangle)$$

except in the case C is of type A_2 and $j \neq k = i$. In the last case, we have $\Lambda(K_j, K_k) = -1$. In particular, we have

$$\mathfrak{d}(\mathbf{K}_j,\mathbf{K}_k) = \mathfrak{d}(\langle j \rangle, \langle k \rangle)$$

unless C is of type A_2 .

Proof. When C is of type A_2 , we can prove directly. Hence we assume that C is not of type A_2 .

(i) Let $j, j' \in I \setminus \{i\}$ satisfy $j \neq j'$. Set $a = -\langle h_i, \alpha_j \rangle$ and $a' = -\langle h_i, \alpha_{j'} \rangle$. Then we have

$$\begin{split} \Lambda(\mathbf{K}_{j},\mathbf{K}_{j'}) &= \Lambda\left(\langle i^{a}\rangle \nabla \langle j\rangle, \langle i^{a'}\rangle \nabla \langle j'\rangle\right) \\ &= \Lambda\left(\langle i^{a}\rangle \nabla \langle j\rangle, \langle i^{a'}\rangle\right) + \Lambda\left(\langle i^{a}\rangle \nabla \langle j\rangle, \langle j'\rangle\right) \\ &= -\Lambda\left(\langle i^{a'}\rangle, \langle i^{a}\rangle \nabla \langle j\rangle\right) + \Lambda\left(\langle i^{a}\rangle \nabla \langle j\rangle, \langle j'\rangle\right) \\ &= -\Lambda\left(\langle i^{a'}\rangle, \langle j\rangle\right) + \Lambda\left(\langle i^{a}\rangle, \langle j'\rangle\right) + \Lambda\left(\langle j\rangle, \langle j'\rangle\right) \\ &= a'(\alpha_{i}, \alpha_{j}) - a(\alpha_{i}, \alpha_{j'}) + \Lambda\left(\langle j\rangle, \langle j'\rangle\right) \\ &= \Lambda\left(\langle j\rangle, \langle j'\rangle\right). \end{split}$$

Here, $=_{(1)}$ and $=_{(2)}$ follow from the fact that $\langle i^a \rangle \nabla \langle j \rangle$ and $\langle i^{a'} \rangle$ commute, The equality $=_{(3)}$ follows from the fact $(\langle i^a \rangle, \langle j' \rangle)$ is unmixed and [12, Corollary 2.3, Corollary 2.12].

(ii) For $j \in I \setminus \{i\}$, let us show that $\Lambda(K_j, K_i) = \Lambda(\langle j \rangle, \langle i \rangle)$. Set $a = -\langle h_i, \alpha_j \rangle$. Since $\langle i \rangle$ commutes with K_j , we have

$$\Lambda(\mathbf{K}_j, \mathbf{C}_{i^*}^*) = \Lambda(\mathbf{K}_j, \langle i \rangle \nabla E_i \mathbf{C}_{i^*}^*) = \Lambda(\mathbf{K}_j, \langle i \rangle) + \Lambda(\mathbf{K}_j, E_i \mathbf{C}_{i^*}^*).$$

Hence

$$\begin{split} \Lambda(\mathbf{K}_{j},\mathbf{K}_{i}) &= \Lambda(\mathbf{K}_{j},E_{i}\mathbf{C}_{i^{*}}^{*}) - \Lambda(\mathbf{K}_{j},\mathbf{C}_{i^{*}}^{*}) \\ &= -\Lambda(\mathbf{K}_{j},\langle i\rangle) = \Lambda(\langle i\rangle,\mathbf{K}_{j}) \\ &= \Lambda(\langle i\rangle,\langle i^{a}\rangle \nabla \langle j\rangle) = \Lambda(\langle i\rangle,\langle j\rangle) = -(\alpha_{i},\alpha_{j}) = \Lambda(\langle j\rangle,\langle i\rangle). \end{split}$$

(iii) For $j \in I \setminus \{i\}$, let us show $\Lambda(\mathbf{K}_i, \mathbf{K}_j) = \Lambda(\langle i \rangle, \langle j \rangle)$. Set $C = \mathsf{C}_{i^*}^*$. We have

$$\left(\operatorname{wt}(E_iC),\operatorname{wt}(\mathbf{K}_j)\right) - \left(\operatorname{wt}(C),\operatorname{wt}(\mathbf{K}_j)\right) = (\alpha_i, -s_i\alpha_j) = (\alpha_i, \alpha_j).$$

Hence we have

$$\begin{split} \Lambda(\mathbf{K}_{i},\mathbf{K}_{j}) &- \Lambda(\langle i \rangle,\langle j \rangle) \\ &= \Lambda(E_{i}C,\mathbf{K}_{j}) - \Lambda(C,\mathbf{K}_{j}) + (\alpha_{i},\alpha_{j}) \\ &= \left(\Lambda(E_{i}C,\mathbf{K}_{j}) + \left(\mathrm{wt}(E_{i}C),\mathrm{wt}(\mathbf{K}_{j})\right)\right) - \left(\Lambda(C,\mathbf{K}_{j}) + \left(\mathrm{wt}(C),\mathrm{wt}(\mathbf{K}_{j})\right)\right) \\ &= 2\widetilde{\Lambda}(E_{i}C,\mathbf{K}_{j}) - 2\widetilde{\Lambda}(C,\mathbf{K}_{j}) \\ &= 2\widetilde{\Lambda}(E_{i}C,\mathbf{K}_{j}) - 2\widetilde{\Lambda}(L(i) \nabla(E_{i}C),\mathbf{K}_{j}) \\ &\leqslant 0. \end{split}$$

Here the last inequality follows from [12, Theorem 2.11]. Thus, we have obtained

 $\Lambda(\mathbf{K}_i, \mathbf{K}_j) \leqslant \Lambda(\langle i \rangle, \langle j \rangle).$

Hence it is enough to show

(10.3) if
$$\Lambda(\mathbf{K}_i, \mathbf{K}_j) < \Lambda(\langle i \rangle, \langle j \rangle)$$
, then **C** is of type A_2 .

Assume that $\Lambda(\mathbf{K}_i, \mathbf{K}_j) < \Lambda(\langle i \rangle, \langle j \rangle)$. Then (ii) implies that

$$\mathfrak{d}(\mathbf{K}_i,\mathbf{K}_j) < \mathfrak{d}(\langle i \rangle,\langle j \rangle) = \delta((\alpha_i,\alpha_j) < 0) \max(\mathsf{d}_i,\mathsf{d}_j).$$

Here $\mathsf{d}_k = (\alpha_k, \alpha_k)/2$. Hence we have $(\alpha_i, \alpha_j) < 0$. Since K_i and C are affreal of degree $2\mathsf{d}_i$, and K_j is affreal of degree $2\mathsf{d}_j$, we have $\mathfrak{d}(K_i, K_j) \in \mathbb{Z} \max(\mathsf{d}_i, \mathsf{d}_j)$. Thus we obtain $\mathfrak{d}(K_i, K_j) = 0$. Hence we have $\mathfrak{d}(E_iC, K_j) = 0$. Hence, $K_j \circ (E_iC)$ is simple. Set $c = -\langle h_i, \alpha_j \rangle > 0$. Then we have $K_j = \langle i^c \rangle \nabla \langle j \rangle$. The composition of

$$(\langle i^c \rangle \nabla \langle j \rangle) \circ (E_i C) \rightarrowtail (\langle i^{c-1} \rangle \nabla \langle j \rangle) \circ \langle i \rangle \circ (E_i C) \twoheadrightarrow (\langle i^{c-1} \rangle \nabla \langle j \rangle) \circ C$$

does not vanish by [8, Lemma 3.1.5]. Since $(\langle i^{c-1} \rangle \nabla \langle j \rangle) \circ C$ is also simple, we obtain

$$(\langle i^c \rangle \nabla \langle j \rangle) \circ (E_i C) \simeq (\langle i^{c-1} \rangle \nabla \langle j \rangle) \circ C.$$

Comparing their dimensions, we obtain

$$c! \cdot \dim(E_iC) \cdot \frac{(c+s)!}{(c+1)! (s-1)!} = (c-1)! \cdot \dim(C) \cdot \frac{(c+s)!}{c! s!}$$

Here, $s := \operatorname{ht}(\operatorname{wt}(C)) = \operatorname{ht}(\Lambda_{i^*} - w_0 \Lambda_{i^*}) - \delta_{i,i^*}$. Note that $\langle i^a \rangle \otimes \langle j \rangle \to \langle i^a \rangle \nabla \langle j \rangle$ is bijective if $0 \leq a \leq -\langle h_i, \alpha_j \rangle$.

Since $E_j C \simeq 0$ for any $j \in I \setminus \{i\}$, we obtain $\dim(E_i C) = \dim C$. Thus we obtain

$$s = \frac{c+1}{c}$$

Hence we obtain c = 1 and s = 2. Note that rank $\mathfrak{g} \leq s + 1 = 3$. Then we can easily check that if $2 \leq \operatorname{rank}(\mathfrak{g}) \leq 3$ and c = 1, then $s \geq 3$ unless $\mathfrak{g} = A_2$.

Proposition 10.2. For any $j, k \in I$ such that $j \neq k$, we have

$$\mathscr{D}(\widehat{\mathbf{K}}_j, \widehat{\mathbf{K}}_k) \equiv Q_{j,k}(z_j, z_k)$$

except in the case C is of type A_2 . In the last case, we have $\mathscr{D}(\widehat{K}_j, \widehat{K}_k) \equiv 1$.

Proof. When C is of type A_2 , we can prove directly. Hence assume that C is not of type A_2 .

(i) Let $j, j' \in I \setminus \{i\}$ satisfy $j \neq j'$. Since $(K_j, K_{j'})$ and $(K_{j'}, K_j)$ are unmixed, by Lemma 9.10 we have

$$\widetilde{\mathscr{L}}(\widehat{\mathbf{K}}_j, \widehat{\mathbf{K}}_{j'}) \equiv \widetilde{\mathscr{L}}(\widehat{\mathbf{K}}_{j'}, \widehat{\mathbf{K}}_j) \equiv 1.$$

By Lemma 9.4 we obtain

$$\mathscr{D}(\widehat{\mathbf{K}}_j, \widehat{\mathbf{K}}_{j'}) \equiv \widetilde{\mathscr{L}}(\widehat{\mathbf{K}}_j, \widehat{\mathbf{K}}_{j'}) \cdot \widetilde{\mathscr{L}}(\widehat{\mathbf{K}}_{j'}, \widehat{\mathbf{K}}_j) \cdot \mathscr{W}t(\widehat{\mathbf{K}}_j, \widehat{\mathbf{K}}_{j'})^{-1} \equiv \mathscr{W}t(\widehat{\mathbf{K}}_j, \widehat{\mathbf{K}}_{j'})^{-1}.$$

By (10.2) we have

$$\mathscr{W}t(\widehat{\mathbf{K}}_{j},\widehat{\mathbf{K}}_{j'}) = \frac{\left(\chi_{i}(\widehat{\mathbf{K}}_{j}) \star \chi_{i}(\widehat{\mathbf{K}}_{j'})\right)^{2}}{A},$$

where

$$A = \left(\chi_i(\widehat{\mathbf{K}}_j) \underset{t_i}{\star} Q_{i,j'}(t_i, t_{j'}) \underset{t_{j'}}{\star} \chi_{j'}(\widehat{\mathbf{K}}_{j'})\right) \cdot \left(\chi_j(\widehat{\mathbf{K}}_j) \underset{t_j}{\star} Q_{i,j}(t_i, t_j) \underset{t_i}{\star} \chi_i(\widehat{\mathbf{K}}_{j'})\right) \\ \cdot \left(\chi_j(\widehat{\mathbf{K}}_j) \underset{t_j}{\star} Q_{j,j'}(t_j, t_{j'}) \underset{t_{j'}}{\star} \chi_{j'}(\widehat{\mathbf{K}}_{j'})\right)$$

On the other hand, we have

$$Q_{i,j}(t_i, z_j) \underset{t_i}{\star} Q_{i,j'}(t_i, z_{j'}) = \chi_i(\widehat{\mathbf{K}}_j) \underset{t_i}{\star} \chi_i(\widehat{\mathbf{K}}_{j'})$$
$$= \chi_i(\widehat{\mathbf{K}}_j) \underset{t_i}{\star} Q_{i,j'}(t_i, t_{j'}) \underset{t_{j'}}{\star} \chi_{j'}(\widehat{\mathbf{K}}_{j'})$$
$$= \chi_j(\widehat{\mathbf{K}}_j) \underset{t_j}{\star} Q_{i,j}(t_i, t_j) \underset{t_i}{\star} \chi_i(\widehat{\mathbf{K}}_{j'})$$

and

$$\chi_j(\widehat{\mathbf{K}}_j) \underset{t_j}{\star} Q_{j,j'}(t_j, t_{j'}) \underset{t_{j'}}{\star} \chi_{j'}(\widehat{\mathbf{K}}_{j'}) = Q_{j,j'}(z_j, z_{j'})$$

Hence we have

$$\mathscr{W}t(\widehat{\mathbf{K}}_j,\widehat{\mathbf{K}}_{j'}) = Q_{j,j'}(z_j,z_{j'})^{-1}$$

and

$$\mathscr{D}(\widehat{\mathrm{K}}_j, \widehat{\mathrm{K}}_{j'}) = Q_{j,j'}(z_j, z_{j'}).$$

(ii) It is obvious $\mathscr{D}(\widehat{\mathbf{K}}_i, \widehat{\mathbf{K}}_j) \equiv 1$ for $j \in I$ such that $(\alpha_i, \alpha_j) = 0$, since $\mathfrak{d}(\mathbf{K}_i, \mathbf{K}_j) = 0$.

(iii) Let us show $\mathscr{D}(\widehat{\mathbf{K}}_i, \widehat{\mathbf{K}}_j) \equiv Q_{i,j}(z_i, z_j)$ for $j \in I$ such that $\langle h_i, \alpha_j \rangle = -1$.

In this case, we have $c := \mathsf{d}_i/\mathsf{d}_j = -\langle h_j, \alpha_i \rangle \in \mathbb{Z}_{\geq 1}$ and we may assume that $Q_{i,j}(z_i, z_j) = z_i - z_j^c$. Set $f(z_i, z_j) := \mathscr{D}(\widehat{\mathsf{K}}_i, \widehat{\mathsf{K}}_j)$. Then we have deg $f(z_i, z_j) = \deg Q_{i,j}(z_i, z_j)$ by Proposition 10.1.

Note that $\widehat{K}_j \simeq \langle i \rangle_{z_j^c} \otimes_{\mathbf{k}[z_j]} \langle j \rangle_{z_j}$ as a $\mathbf{k}[z_j]$ -module on which τ_1 acts by zero. Hence it is a quotient of $\langle i \rangle_{z_j^c} \underset{z_j}{\circ} \langle j \rangle_{z_j}$, so that by Lemma 6.27, we have $\widehat{K}_j \simeq \langle i \rangle_{z_j^c} \nabla_{z_j} \langle j \rangle_{z_j} \rightarrow \langle j \rangle_{z_j} \underset{z_j}{\circ} \langle i \rangle_{z_j^c}$.

We have

e nave

$$\widehat{\mathrm{K}}_{j} \underset{z_{j}}{\circ} \left(\widehat{\mathrm{K}}_{i} |_{z_{i} = z_{j}^{c}} \right) \rightarrowtail \langle j \rangle_{z_{j}} \underset{z_{j}}{\circ} \langle i \rangle_{z_{j}^{c}} \underset{z_{j}}{\circ} \left(\widehat{\mathrm{K}}_{i} |_{z_{i} = z_{j}^{c}} \right) \twoheadrightarrow \langle j \rangle_{z_{j}},$$

which implies an epimorphism

(10.4)
$$\widehat{\mathbf{K}}_{j} \underset{z_{j}}{\circ} \left(\widehat{\mathbf{K}}_{i} |_{z_{i} = z_{j}^{c}} \right) \twoheadrightarrow \langle j \rangle_{z_{j}},$$

since it is so after operating $(\mathbf{k}[z_j]/z_j\mathbf{k}[z_j]) \otimes_{\mathbf{k}[z_i]} \bullet$.

Hence Proposition 6.29 implies that $z_j - w$ divides $f(w^c, z_j)$, i.e., $f(w^c, w) = 0$. Hence $Q_{i,j}(z_i, z_j) = z_i - z_j^c$ divides $f(z_i, z_j)$. Comparing their degrees, we have $f(z_i, z_j) = Q_{i,j}(z_i, z_j)$ up to a constant multiple.

(iv) Let us show $\mathscr{D}(\widehat{K}_i, \widehat{K}_j) \equiv Q_{i,j}(z_i, z_j)$ for $j \in I$ such that $-\langle h_i, \alpha_j \rangle > 1$.

In this case, we have $c := d_j/d_i = -\langle h_i, \alpha_j \rangle \in \mathbb{Z}_{>1}$ and we may assume that $Q_{i,j}(z_i, z_j) = z_j - z_i^c$. Set $f(z_i, z_j) := \mathscr{D}(\widehat{K}_i, \widehat{K}_j)$. Then we have deg $f(z_i, z_j) = \deg Q_{i,j}(z_i, z_j)$ by Proposition 10.1.

Set $\mathsf{M} = \check{\mathsf{K}}_j|_{z_j = w^c}$ with $\deg(w) = \mathsf{d}_i$. By replacing \mathbf{k} with its algebraic closure, we may assume that \mathbf{k} is algebraically closed. Let ω be the *c*-th primitive root of unity. Then we have $\mathsf{M} = \mathsf{L} \nabla_w \langle j \rangle_{w^c}$, where

(10.5)
$$\mathsf{L} = \langle i \rangle_w \mathop{\circ}_w \langle i \rangle_{\omega w} \mathop{\circ}_w \cdots \mathop{\circ}_w \langle i \rangle_{\omega^{c-1} w} \simeq \langle i \rangle_w \mathop{\circ}_w \mathsf{L}' \text{ and } \mathsf{L}' = \langle i \rangle_{\omega w} \mathop{\circ}_w \cdots \mathop{\circ}_w \langle i \rangle_{\omega^{c-1} w}.$$

Then we have $\mathsf{M} = \langle i \rangle_w \nabla_w \left(\mathsf{L}' \nabla_w \langle j \rangle_{w^c} \right) \mapsto \left(\mathsf{L}' \nabla_w \langle j \rangle_{w^c} \right) \underset{w}{\circ} \langle i \rangle_w.$

Hence we have

$$\mathsf{M} \underset{w}{\circ} (\widehat{\mathsf{K}}_{i}|_{z_{i}=w}) \rightarrowtail (\mathsf{L}' \nabla_{w} \langle j \rangle_{w^{c}}) \underset{w}{\circ} \langle i \rangle_{w} \underset{w}{\circ} (\widehat{\mathsf{K}}_{i}|_{z_{i}=w}) \twoheadrightarrow \mathsf{L}' \nabla_{w} \langle j \rangle_{w^{c}}.$$

Then the composition is an epimorphism

(10.6)
$$\left(\widehat{\mathbf{K}}_{j}|_{z_{j}=w^{c}}\right) \underset{w}{\circ} \left(\widehat{\mathbf{K}}_{i}|_{z_{i}=w}\right) \twoheadrightarrow \mathsf{L}' \nabla_{w} \langle j \rangle_{w^{c}},$$

since it is so after operating $(\mathbf{k}[w]/w\mathbf{k}[w]) \otimes_{\mathbf{k}[w]} \bullet$.

By Lemma 9.11, $\mathsf{L}' \nabla_w \langle j \rangle_{w^c}$ is an affinization of $\langle i^{c-1} \rangle \nabla \langle j \rangle$ which is real simple by Lemma 10.4 below. Hence, Proposition 6.29 implies that $Q_{i,j}(z_i, z_j) = z_j - z_i^c$ divides $f(z_i, z_j)$. Comparing their degrees, we have $f(z_i, z_j) = Q_{i,j}(z_i, z_j)$.

Thus we have proved that the datum $\{\widehat{K}_j\}_{j\in I}$ in $\widetilde{\mathscr{C}}^*_{s_iw_0}$ satisfies condition (7.7). Choosing a duality datum $(\{(\widehat{K}_j, z_j)\}_{j\in I}, \{\mathbb{R}^{\text{ren}}_{\widehat{K}_j, \widehat{K}_k}\}_{j,k\in I})$, we can define exact monoidal functors

 $\widehat{\mathscr{F}_i} \colon \operatorname{Mod}_{\operatorname{coh}}(R) \to \operatorname{Pro}(\widetilde{\mathscr{C}}^*_{s_i w_0}) \quad \text{and} \quad \mathscr{F}_i \colon R\operatorname{-gmod} \to \widetilde{\mathscr{C}}^*_{s_i w_0}$

sending $\langle j \rangle_{z_j}$ to $\widehat{\mathbf{K}}_j$, except the A_2 -cases. See Remark 10.13 for the A_2 -cases. Note that

(10.7)
$$\operatorname{wt}(\widehat{\mathscr{F}}_{i}(X)) = s_{i}(\operatorname{wt}(X)) \text{ for any } X \in \operatorname{Mod}_{\operatorname{coh}}(R).$$

Lemma 10.3. Let $M, N \in R$ -gmod be simple modules. Assume that $\mathscr{F}_i(M)$ and $\mathscr{F}_i(N)$ are simples in $\widetilde{\mathscr{C}}^*_{s_iw_0}$. Assume moreover that either M and $\mathscr{F}_i(M)$ are affreal or N and $\mathscr{F}_i(N)$ are affreal.

- (i) We have $\Lambda(\mathscr{F}_i(M), \mathscr{F}_i(N)) \leq \Lambda(M, N)$.
- (ii) The following conditions are equivalent.
 (a) Λ(𝔅_i(M), 𝔅_i(N)) = Λ(M, N),
 (b) 𝔅_i(M ∇ N) ≠ 0.
- (iii) Moreover, if condition (iib) holds, then $\mathscr{F}_i(\mathbf{r}_{M,N}) \neq 0$ and

$$\mathscr{F}_i(M \nabla N) \simeq \mathscr{F}_i(M) \nabla \mathscr{F}_i(N).$$

Proof. Assume that M and $\mathscr{F}_i(M)$ are affreal. Let (M, z) be an affinization of M. Since

$$\mathbf{R}_{N,\mathsf{M}}^{\mathrm{ren}} \circ \mathbf{R}_{\mathsf{M},N}^{\mathrm{ren}} = z^d \; \operatorname{id}_{\mathsf{M} \otimes N},$$

we have a commutative diagram

$$\widehat{\mathscr{F}_{i}(z)^{d}} \xrightarrow{\widehat{\mathscr{F}_{i}(z)^{d}}} \widehat{\mathscr{F}_{i}(\mathsf{M})} \circ \widehat{\mathscr{F}_{i}(\mathsf{M})} \xrightarrow{\widehat{\mathscr{F}_{i}(\mathsf{R}^{\mathrm{ren}}_{N,\mathsf{M}})}} \widehat{\mathscr{F}_{i}(\mathsf{M})} \circ \widehat{\mathscr{F}_{i}(\mathsf{M})} \xrightarrow{\widehat{\mathscr{F}_{i}(\mathsf{M})}} \widehat{\mathscr{F}_{i}(\mathsf{M})} \circ \widehat{\mathscr{F}_{i}(\mathsf{M})} \xrightarrow{\widehat{\mathscr{F}_{i}(\mathsf{M})}} \widehat{\mathscr{F}_{i}(\mathsf{M})} \otimes \widehat{\mathscr{F}_{i}(\mathsf{M})} \otimes \widehat{\mathscr{F}_{i}(\mathsf{M})} \xrightarrow{\widehat{\mathscr{F}_{i}(\mathsf{M})}} \widehat{\mathscr{F}_{i}(\mathsf{M})} \otimes \widehat{\mathscr{F}_{i}(\mathsf{M}$$

Since $\widehat{\mathscr{F}}_{i}(z)$ is a monomorphism, $\widehat{\mathscr{F}}_{i}(\mathbb{R}^{\text{ren}}_{\mathsf{M}_{z},N})$ and $\widehat{\mathscr{F}}_{i}(\mathbb{R}^{\text{ren}}_{N,\mathsf{M}_{z}})$ are non-zero. Note also that dim $\operatorname{Hom}(\mathscr{F}_{i}(M) \circ \mathscr{F}_{i}(N), \mathscr{F}_{i}(N) \circ \mathscr{F}_{i}(M)) = 1$ by Proposition 8.7. It follows

that

$$\widehat{\mathscr{F}}_{i}(\mathbf{R}_{\mathsf{M},N}^{\mathrm{ren}}) = z^{a} \mathbf{R}_{\widehat{\mathscr{F}}_{i}(\mathsf{M}),\widehat{\mathscr{F}}_{i}(N)}^{\mathrm{ren}} \quad \text{and} \quad \widehat{\mathscr{F}}_{i}(\mathbf{R}_{N,\mathsf{M}}^{\mathrm{ren}}) = z^{b} \mathbf{R}_{\widehat{\mathscr{F}}_{i}(N),\widehat{\mathscr{F}}_{i}(\mathsf{M})}^{\mathrm{ren}}$$

for some $a, b \in \mathbb{Z}_{\geq 0}$ up to constant multiples by Proposition 6.2 (ii).

Note also that $\widehat{\mathscr{F}}_i(\mathsf{M})$ is an affine object of $\mathscr{F}_i(M)$. Hence we have

$$\Lambda(M,N) = \deg(\mathbf{R}_{\mathsf{M},N}^{\mathrm{ren}}) = \deg(\widehat{\mathscr{F}_{i}}(\mathbf{R}_{\mathsf{M},N}^{\mathrm{ren}})) \geqslant \deg(\mathbf{R}_{\widehat{\mathscr{F}}_{i}}^{\mathrm{ren}}(\mathbf{M}),\widehat{\mathscr{F}}_{i}(N)) = \Lambda(\widehat{\mathscr{F}_{i}}(M),\widehat{\mathscr{F}_{i}}(N)),$$

and the equality $\deg(\widehat{\mathscr{F}}_{i}(\mathbf{R}_{\mathsf{M},N}^{\mathrm{ren}})) = \deg(\mathbf{R}_{\widehat{\mathscr{F}}_{i}(\mathsf{M}),\widehat{\mathscr{F}}_{i}(N)}^{\mathrm{ren}})$ holds if and only if $\deg(z^{a}) = 0$; i.e., a = 0. In the case, we have

$$\mathscr{F}_{i}(\mathbf{r}_{M,N}) = \widehat{\mathscr{F}_{i}}(\mathbf{R}_{\mathsf{M},N}^{\mathrm{ren}}|_{z=0}) = (\widehat{\mathscr{F}_{i}}(\mathbf{R}_{\mathsf{M},N}^{\mathrm{ren}}))|_{\widehat{\mathscr{F}}_{i}(z)=0} = \mathbf{r}_{\mathscr{F}_{i}(M),\mathscr{F}_{i}(N)} \neq 0$$

and hence

$$M \nabla N \simeq \operatorname{Im} \mathscr{F}_i(\mathbf{r}_{M,N}) = \operatorname{Im} \mathbf{r}_{\mathscr{F}_i(M), \mathscr{F}_i(N)} \simeq \mathscr{F}_i(M) \nabla \mathscr{F}_i(N),$$

which completes the proof.

Lemma 10.4. Let $m, n \in \mathbb{Z}_{\geq 0}$ and $j \in I \setminus \{i\}$, and let $S = \langle j^n \rangle \nabla \langle i^m \rangle$. Set $\lambda = n\Lambda_j - m\Lambda_i$. Then we have (neglecting the degree shifts):

- (i) $s_i \lambda \preccurlyeq \lambda \preccurlyeq s_i \lambda$ and $S \simeq \mathsf{M}(s_i \lambda, s_i \lambda)$,
- (ii) $\varepsilon_i(S) = \max(0, m + n\langle h_i, \alpha_j \rangle),$
- (iii) if $m \ge -n\langle h_i, \alpha_j \rangle$, then $\langle i \rangle$ commutes with S and $\langle j^n \rangle \nabla \langle i^m \rangle \simeq (\langle j^n \rangle \nabla \langle i^{-n\langle h_i, \alpha_j \rangle} \rangle) \circ \langle i^{m+n\langle h_i, \alpha_j \rangle} \rangle$,

(iv) if
$$\langle h_i, s_j \lambda \rangle = -n \langle h_i, \alpha_j \rangle - m \ge 0$$
, then $\mathscr{F}_i(S) \simeq \mathsf{M}(s_i s_j \lambda, \lambda) \simeq \langle i^{-m-n \langle h_i, \alpha_j \rangle} \rangle \nabla \langle j^n \rangle$.

Proof. (i) We have $\langle h_i, s_i \lambda \rangle = m$, $\langle h_j, \lambda \rangle = n$ and $s_j \lambda = (s_j s_i)(s_i \lambda)$. Hence we have $S \simeq \mathsf{M}(s_j \lambda, s_i \lambda)$.

(ii) follows from [12, Proposition 2.16 (iii)].

(iii) follows from the fact that $\psi_*(\mathsf{M}(s_i s_j n \Lambda_j, n \Lambda_j)) \simeq \langle j^n \rangle \nabla \langle i^{-n \langle h_i, \alpha_j \rangle} \rangle$ commutes with $\langle i \rangle$ by [12, Proposition 2.16 (ii)].

(iv) We argue by induction on m. If m = 0, then

$$\begin{aligned} \mathscr{F}_i(S) &= \mathscr{F}_i(\langle j \rangle^{\circ n}) \simeq \mathsf{M}(s_i s_j \Lambda_j, \Lambda_j)^{\circ n} \\ &\simeq \mathsf{M}(s_i s_j n \Lambda_j, n \Lambda_j) \simeq \langle i^{-n \langle h_i, \alpha_j \rangle} \rangle \, \nabla \, \langle j^n \rangle. \end{aligned}$$

Assume that m > 0. Set $c = -\langle h_i, \alpha_j \rangle$. Then we have

$$\begin{aligned} \mathscr{F}_{i}(S) &\simeq \mathscr{F}_{i}\left(\left(\langle j^{n}\rangle \nabla \langle i^{m-1}\rangle\right) \nabla \langle i\rangle\right) \\ &\simeq \mathscr{F}_{i}\left(\langle j^{n}\rangle \nabla \langle i^{m-1}\rangle\right) \nabla \mathscr{F}_{i}\left(\langle i\rangle\right) \\ &\simeq \left(\langle i^{1-m+nc}\rangle \nabla \langle j^{n}\rangle\right) \nabla \mathscr{D}\langle i\rangle \\ &\simeq \left(\langle i\rangle \nabla \left(\langle i^{-m+nc}\rangle \nabla \langle j^{n}\rangle\right)\right) \nabla \mathscr{D}\langle i\rangle \\ &\simeq \langle i^{-m+nc}\rangle \nabla \langle j^{n}\rangle. \end{aligned}$$

Here the second isomorphism $\underset{*}{\simeq}$ follows from Lemma 10.3 and the fact that

$$\Lambda(\langle j^n \rangle \nabla \langle i^{m-1} \rangle, \langle i \rangle) = \Lambda(\langle j^n \rangle, \langle i \rangle) = -n(\alpha_j, \alpha_i)$$

is equal to

$$\begin{split} \Lambda\big(\mathscr{F}_i(\langle j^n \rangle \nabla \langle i^{m-1} \rangle), \mathscr{F}_i(\langle i \rangle)\big) &= \Lambda\big(\langle i^{1-m+nc} \rangle \nabla \langle j^n \rangle, \mathscr{D}\langle i \rangle\big) \underset{(1)}{=} \Lambda\big(\langle i \rangle, \langle i^{1-m+nc} \rangle \nabla \langle j^n \rangle\big) \\ &= \Lambda(\langle i \rangle, \langle j^n \rangle) = -n(\alpha_i, \alpha_j), \end{split}$$

where $=_{(1)}$ follows from Lemma 6.30.

Lemma 10.5. For $w \in W$ such that $s_i w > w$, we have

$$\mathscr{F}_i(\mathsf{M}(w\Lambda,\Lambda)) \simeq \mathsf{M}(s_i w\Lambda, s_i\Lambda) \text{ for any } \Lambda \in \mathsf{P}_+.$$

In particular, we have

$$\mathscr{F}_i(\mathsf{C}_\Lambda) \simeq \mathsf{C}^*_\Lambda \quad \text{for any } \Lambda \in \mathsf{P}_+.$$

Proof. Let us argue by induction on $\ell(w)$. If $\ell(w) \leq 1$, it is obvious. Assume that $\ell(w) > 1$. Take $j \in I \setminus \{i\}$ such that $s_j w < w$.

(i) Assume first that $s_i s_j w > s_j w$. Set $v = s_j w$ and $n = \langle h_j, v\Lambda \rangle \in \mathbb{Z}_{\geq 0}$ and $m = \langle h_i, v\Lambda \rangle \in \mathbb{Z}_{\geq 0}$. Then we have $w\Lambda = v\Lambda - n\alpha_j$ and hence $\langle h_i, w\Lambda \rangle = m - n\langle h_i, \alpha_j \rangle \in \mathbb{Z}_{\geq 0}$. Then we have $\mathsf{M}(w\Lambda, v\Lambda) \simeq \langle j^n \rangle$ and $\mathsf{M}(s_i w\Lambda, v\Lambda) \simeq \langle i^{m-n\langle h_i, \alpha_j \rangle} \rangle \nabla \langle j^n \rangle$. We have

$$\begin{split} \mathsf{M}(s_i w\Lambda, s_i v\Lambda) \nabla \,\mathsf{M}(s_i v\Lambda, v\Lambda) &\simeq \mathsf{M}(s_i w\Lambda, v\Lambda) \\ &\simeq \langle i^{m-n\langle h_i, \alpha_j \rangle} \rangle \,\nabla \,\langle j^n \rangle \simeq \left(\langle i^{-n\langle h_i, \alpha_j \rangle} \rangle \,\nabla \,\langle j^n \rangle \right) \circ \,\langle i^m \rangle \\ &\simeq \left(\langle i^{-\langle h_i, \alpha_j \rangle} \rangle \,\nabla \,\langle j \rangle \right)^{\circ n} \circ \,\mathsf{M}(s_i v\Lambda, v\Lambda), \end{split}$$

where \simeq follows from Lemma 10.4. Hence we have $\mathsf{M}(s_i w \Lambda, s_i v \Lambda) \simeq \left(\langle i^{-\langle h_i, \alpha_j \rangle} \rangle \nabla \langle j \rangle\right)^{\circ n}$, which implies that

$$\mathscr{F}_i(\mathsf{M}(w\Lambda, v\Lambda)) \simeq \mathsf{M}(s_i w\Lambda, s_i v\Lambda).$$

On the other hand, [9, Proposition 4.6] implies that both $\Lambda(\mathsf{M}(w\Lambda, v\Lambda), \mathsf{M}(v\Lambda, \Lambda))$ and $\Lambda(\mathsf{M}(s_iw\Lambda, s_iv\Lambda), \mathsf{M}(s_iv\Lambda, s_i\Lambda))$ are equal to $-(w\Lambda - v\Lambda, v\Lambda - \Lambda)$. Therefore, by Lemma 10.3, we obtain

$$\begin{aligned} \mathscr{F}_i\big(\mathsf{M}(w\Lambda,\Lambda)\big) &\simeq \mathscr{F}_i\big(\mathsf{M}(w\Lambda,v\Lambda) \nabla \mathsf{M}(v\Lambda,\Lambda)\big) \\ &\simeq \mathscr{F}_i\big(\mathsf{M}(w\Lambda,v\Lambda)\big) \nabla \mathscr{F}_i\big(\mathsf{M}(v\Lambda,\Lambda)\big) \\ &\simeq \mathsf{M}(s_iw\Lambda,s_iv\Lambda) \nabla \mathsf{M}(s_iv\Lambda,s_i\Lambda) \\ &\simeq \mathsf{M}(s_iw\Lambda,s_i\Lambda). \end{aligned}$$

(ii) Assume that $s_i s_j w < s_j w$. Set $v = s_i s_j w$. Then we have $s_i v > v$, $\ell(v) < \ell(w)$ and hence by induction hypothesis, $\mathscr{F}_i(\mathsf{M}(v\Lambda,\Lambda)) \simeq \mathsf{M}(s_i v\Lambda, s_i\Lambda)$. Note that $\mathsf{M}(w\Lambda, v\Lambda) \simeq \langle j^n \rangle \nabla \langle i^m \rangle$ where $n = \langle h_j, s_i v\Lambda \rangle$, $m = \langle h_i, v\Lambda \rangle$.

Since $\Lambda(\langle j^n \rangle, \langle i^m \rangle) = -nm(\alpha_j, \alpha_i)$ and

$$\Lambda\big((\langle i^{-\langle h_i,\alpha_j\rangle}\rangle \nabla \langle j\rangle)^{\circ n}, \mathscr{D}(\langle i^m\rangle)\big) = \Lambda\big(\langle i^m\rangle, (\langle i^{-\langle h_i,\alpha_j\rangle}\rangle \nabla \langle j\rangle)^{\circ n}\big) = -nm(\alpha_j,\alpha_i)$$

are equal, we have

$$\begin{split} \mathscr{F}_{i}(\mathsf{M}(w\Lambda,v\Lambda)) &\simeq \left(\langle i^{-\langle h_{i},\alpha_{j}\rangle} \rangle \nabla \langle j \rangle \right)^{\circ n} \nabla \mathscr{D} \langle i^{m} \rangle \\ &\simeq \left(\langle i^{m} \rangle \nabla \left(\langle i^{-m-n\langle h_{i},\alpha_{j}\rangle} \rangle \nabla \langle j^{n} \rangle \right) \right) \nabla \mathscr{D} \langle i^{m} \rangle \\ &\simeq \langle i^{-m-n\langle h_{i},\alpha_{j}\rangle} \rangle \nabla \langle j^{n} \rangle \simeq \mathsf{M}(s_{i}w\lambda,s_{i}v\Lambda). \end{split}$$

Note that $\langle h_i, w\Lambda \rangle = -m - n \langle h_i, \alpha_j \rangle \ge 0.$

Since $\Lambda(\mathsf{M}(w\Lambda, v\Lambda), \mathsf{M}(v\Lambda, \Lambda))$ and $\Lambda(\mathsf{M}(s_iw\Lambda, s_iv\Lambda), \mathsf{M}(s_iv\Lambda, s_i\lambda))$ are equal to $-(w\Lambda - v\Lambda, v\Lambda - \Lambda)$, we have

$$\begin{aligned} \mathscr{F}_i \big(\mathsf{M}(w\Lambda,\Lambda) \big) &\simeq \mathscr{F}_i \big(\mathsf{M}(w\Lambda,v\Lambda) \nabla \,\mathsf{M}(v\Lambda,\Lambda) \big) \\ &\simeq \mathscr{F}_i \big(\mathsf{M}(w\Lambda,v\Lambda) \big) \nabla \,\mathscr{F}_i \big(\mathsf{M}(v\Lambda,\Lambda) \big) \\ &\simeq \,\mathsf{M}(s_i w\Lambda,s_i v\Lambda) \nabla \,\mathsf{M}(s_i v\Lambda,s_i\Lambda) \\ &\simeq \,\mathsf{M}(s_i w\Lambda,s_i\Lambda). \end{aligned}$$

Lemma 10.6. Let $w, v \in W$ such that $v \leq w$ and $s_i v > v$, $s_i w > w$. Then, we have

$$\mathscr{F}_i(\mathsf{M}(w\Lambda, v\Lambda)) \simeq \mathsf{M}(s_i w\Lambda, s_i v\Lambda) \quad \text{for any } \Lambda \in \mathsf{P}_+.$$

Proof. By the preceding lemma, we have

$$\begin{split} \mathsf{M}(s_i w\Lambda, s_i v\Lambda) \nabla \,\mathsf{M}(s_i v\Lambda, s_i\Lambda) &\simeq \mathsf{M}(s_i w\Lambda, s_i\Lambda) \\ &\simeq \mathscr{F}_i \big(\mathsf{M}(w\Lambda, \Lambda)\big) \simeq \mathscr{F}_i \big(\mathsf{M}(w\Lambda, v\Lambda) \nabla \,\mathsf{M}(v\Lambda, \Lambda)\big) \\ &\simeq \mathscr{F}_i \big(\mathsf{M}(w\Lambda, v\Lambda)\big) \nabla \,\mathscr{F}_i \big(\mathsf{M}(v\Lambda, \Lambda) \\ &\simeq \mathscr{F}_i \big(\mathsf{M}(w\Lambda, v\Lambda)\big) \nabla \,\mathsf{M}(s_i v\Lambda, s_i\Lambda), \end{split}$$

where \simeq_{*} holds by Lemma 10.3. Hence we have

$$\mathscr{F}_i(\mathsf{M}(w\Lambda, v\Lambda)) \simeq \mathsf{M}(s_i w\Lambda, s_i v\Lambda).$$

Proposition 10.7. If M is a simple module in $\mathscr{C}_{s_iw_0}$, then $\mathscr{F}_i(M)$ is a simple module in $\mathscr{C}^*_{s_iw_0}$. Moreover the functor \mathscr{F}_i induces a bijection between the set of classes of simple modules in $\mathscr{C}_{s_iw_0}$ and that of $\mathscr{C}^*_{s_iw_0}$.

Proof. Let $\underline{w_0} = s_{i_1} s_{i_2} \cdots s_{i_{l-1}} s_{i_l}$ be a reduced expression of w_0 with $i_1 = i$. Set

(10.8)
$$V_k := \mathsf{M}(s_{i_2} \cdots s_{i_k} \Lambda_{i_k}, s_{i_2} \cdots s_{i_{k-1}} \Lambda_{i_k}) \quad \text{for} \quad 2 \leqslant k \leqslant l \quad \text{and} \\ S_k := \mathsf{M}(s_{i_1} s_{i_2} \cdots s_{i_k} \Lambda_{i_k}, s_{i_1} s_{i_2} \cdots s_{i_{k-1}} \Lambda_{i_k}) \quad \text{for} \quad 1 \leqslant k \leqslant l.$$

Then by Lemma 10.6, we have

$$\mathscr{F}_i(V_k) = S_k \quad \text{for} \quad 2 \leqslant k \leqslant l.$$

Note that $K(\mathscr{C}_{s_iw_0})|_{q=1}$ is the polynomial ring $\mathbb{Z}[[V_2], \ldots, [V_l]]$ and $K(\mathscr{C}_{w_0})|_{q=1}$ is the polynomial ring $\mathbb{Z}[[S_1], [S_2], \ldots, [S_l]]$, where $K(\mathscr{C}_{w_0})$ denotes the Grothendieck ring of the abelian monoidal category \mathscr{C}_{w_0} and $K(\mathscr{C}_{w_0})|_{q=1} := K(\mathscr{C}_{w_0})/(q-1)K(\mathscr{C}_{w_0})$.

Recall that for any simple M in $\mathscr{C}_{w_0} = R$ -gmod, there exists a unique $(a_1, \ldots, a_l) \in \mathbb{Z}_{\geq 0}^l$ such that $S_l^{\circ a_l} \circ S_{l-1}^{\circ a_{l-1}} \circ \cdots \circ S_1^{\circ a_1}$ has a simple head isomorphic to M up to a grading shift. Since $E_i^*(S_k) = 0$ for $2 \leq k \leq l$ and $S_1 = L(i)$, the simple module M belongs to $\mathscr{C}_{s_iw_0}^*$ if and only if $a_1 = 0$. Hence the ring $K(\mathscr{C}_{s_iw_0}^*)|_{q=1}$ is the polynomial ring $\mathbb{Z}[[S_2], \ldots, [S_l]]$.

It follows that the ring homomorphism $[\mathscr{F}_i]: K(\mathscr{C}_{s_iw_o})|_{q=1} \to K(\widetilde{\mathscr{C}}^*_{s_iw_0})|_{q=1}$ is an isomorphism. In particular, for any non-zero module $M, \mathscr{F}_i(M)$ is non-zero, and $[\mathscr{F}_i]$ induces a bijection between the set of simple modules in $\mathscr{C}_{s_iw_0}$ and that of $\mathscr{C}^*_{s_iw_0}$. \Box

For a simple $R(\beta)$ -module M, set

$$\begin{split} \operatorname{wt}(M) &= -\beta, \\ \widetilde{F}_i(M) &= L(i) \,\nabla \, M, \qquad \widetilde{E}_i(M) = \operatorname{hd}(E_i(M)), \\ \varepsilon_i(M) &= \max\{k \ge 0 \mid E_i^{\ k} M \not\simeq 0\}, \quad \varphi_i(M) = \varepsilon_i(M) + \langle h_i, \operatorname{wt}(M) \rangle, \\ \widetilde{F}_i^*(M) &= M \,\nabla \, L(i), \qquad \widetilde{E}_i^*(M) = \operatorname{hd}(E_i^*(M)), \\ \varepsilon_i^*(M) &= \max\{k \ge 0 \mid E_i^{* k} M \not\simeq 0\}, \quad \varphi_i^*(M) = \varepsilon_i^*(M) + \langle h_i, \operatorname{wt}(M) \rangle. \end{split}$$

Here we denote by M its isomorphism class (as ungraded module) for simplicity. Then the set of isomorphism classes of self-dual simple modules in R-gmod together with $(\text{wt}, \tilde{E}_i, \tilde{F}_i, \varepsilon_i, \varphi_i)$ forms a \mathfrak{g} -crystal which is isomorphic to the crystal basis $B(\infty)$ of $U_q^-(\mathfrak{g})$ ([19]).

On the other hand, Yoshihisa Saito ([24]) defined the isomorphism

$$\{b \in B(\infty) \mid \varepsilon_i(b) = 0\} \xrightarrow{\sim} \{b \in B(\infty) \mid \varepsilon_i^*(b) = 0\}$$

given by $b \mapsto \tilde{f}_i^{\varphi_i^*(b)} \tilde{e}_i^{*\varepsilon_i^*(b)} b$. Hence, to a simple $M \in R$ -gmod with $\varepsilon_i(M) = 0$, we can associate a simple module

$$\sigma_i(M) := \widetilde{F}_i^{\varphi_i^*(M)} \widetilde{E}_i^{* \varepsilon_i^*(M)} M$$

which satisfies $\varepsilon_i^*(\sigma_i(M)) = 0$. We call it the Saito reflection of M with respect to i (cf. [24, 15, 16]).

Proposition 10.8. Let M be a simple module in $\mathscr{C}_{s_iw_0}$. Then

$$\mathscr{F}_i(M) \simeq \sigma_i(M)$$

as an ungraded module.

Proof. By [25, Corollary 2.26], it is enough to show that $\mathscr{F}_i(V_k) \simeq S_k \simeq \sigma_i(V_k)$ for $2 \leq k \leq l$, where V_k and S_k are the modules in (10.8).

Let M be a simple module in R-gmod with $\varepsilon_i(M) = 0$. Then, we have

$$\sigma_i(M) = \widetilde{F}_i^{\varphi_i^*(M)} \widetilde{E}_i^{*\varepsilon_i^*(M)} M \simeq \widetilde{E}_i^{*s - \langle h_i, \operatorname{wt}(M) \rangle} \widetilde{F}_i^s M$$

for any $s \ge \varphi_i^*(M)$ by [12, Proposition 2.16, Theorem 2.17]. Let $2 \le k \le l$ and $\lambda = s_{i_2} \cdots s_{i_{k-1}} \Lambda_{i_k}, \mu = s_{i_2} \cdots s_{i_k} \Lambda_{i_k}$ so that $V_k = \mathsf{M}(\lambda, \mu)$. Then $\langle h_i, \lambda \rangle \ge 0, \langle h_i, \mu \rangle \ge 0$ and

$$\varphi_i^*(V_k) = \varepsilon_i^*(V_k) + \langle h_i, \operatorname{wt}(V_k) \rangle \leqslant \langle h_i, \mu \rangle + \langle h_i, \lambda - \mu \rangle = \langle h_i, \lambda \rangle,$$

where the middle inequality follows from [8, Lemma 9.1.5].

Hence by taking $s = \langle h_i, \lambda \rangle$, we have

$$\sigma_{i}(V_{k}) \simeq \widetilde{E}_{i}^{*s-\langle h_{i}, \operatorname{wt}(V_{k}) \rangle} \widetilde{F}_{i}^{s} \mathsf{M}(\lambda, \mu) = \widetilde{E}_{i}^{*\langle h_{i}, \mu \rangle} \widetilde{F}_{i}^{\langle h_{i}, \lambda \rangle} \mathsf{M}(\lambda, \mu)$$
$$\simeq \widetilde{E}_{i}^{*\langle h_{i}, \mu \rangle} \mathsf{M}(s_{i}\lambda, \mu) \simeq \mathsf{M}(s_{i}\lambda, s_{i}\mu) \simeq S_{k},$$


as desired.

Lemma 10.9. For any $j \in I$, we have

$$\Lambda(\mathsf{M}(s_i w_0 \Lambda, \Lambda), \langle j \rangle) = \Lambda(\mathsf{M}(w_0 \Lambda, s_i \Lambda), \mathsf{K}_j).$$

Proof. We have $\Lambda(\mathsf{M}(s_iw_0\Lambda,\Lambda),\langle j\rangle) = (s_iw_0\Lambda+\Lambda,\alpha_j)$ by [11, (5.1)] and $\Lambda(\mathsf{M}(w_0\Lambda,s_i\Lambda),\mathsf{K}_j) = (w_0\Lambda+s_i\Lambda,s_i\alpha_j)$ by [11, Proposition 3.27].

Proposition 10.10. The functor \mathscr{F}_i : R-gmod $\to \widetilde{\mathscr{C}}_{s_iw_0}^*$ factors through $\widetilde{\mathscr{C}}_{s_iw_0}$:



Proof. By Lemma 10.6, we have $\mathscr{F}_i(\mathsf{C}_\Lambda) \simeq \mathsf{C}^*_\Lambda$ for any $\Lambda \in \mathsf{P}_+$. By the universal property of localization ([11, Theorem 2.7]), it remains to show that $\mathscr{F}_i(R_{\mathsf{C}_\Lambda}(X)) : \mathscr{F}_i(\mathsf{C}_\Lambda) \circ \mathscr{F}_i(X) \to \mathscr{F}_i(\mathsf{C}_\Lambda) \circ \mathscr{F}_$

Note that

- (a) if $\mathscr{F}_i(R_{\mathsf{C}_\Lambda}(M))$ and $\mathscr{F}_i(R_{\mathsf{C}_\Lambda}(N))$ are isomorphisms for $M, N \in R$ -gmod, then so is $\mathscr{F}_i(R_{\mathsf{C}_\Lambda}(M \circ N))$.
- (b) for a morphism $f: M \to N$, if $\mathscr{F}_i(R_{\mathsf{C}_\Lambda}(M))$, $\mathscr{F}_i(R_{\mathsf{C}_\Lambda}(N))$ are isomorphisms, then so is $\mathscr{F}_i(R_{\mathsf{C}_\Lambda}(\operatorname{Coker}(f)))$,
- (c) for an exact sequence $0 \to L \to M \to N \to 0$, if $\mathscr{F}_i(R_{\mathsf{C}_\Lambda}(L))$ and $\mathscr{F}_i(R_{\mathsf{C}_\Lambda}(N))$ are isomorphism, then so is $\mathscr{F}_i(R_{\mathsf{C}_\Lambda}(M))$.

Thus it is enough to show that $\mathscr{F}_i(R_{\mathsf{C}_\Lambda}(\langle j \rangle))$ is an isomorphism for any $j \in I$ and it follows from Lemma 10.3 and Lemma 10.9.

Theorem 10.11. The monoidal functor $\mathscr{S}_i : \widetilde{\mathscr{C}}_{s_iw_0} \to \widetilde{\mathscr{C}}^*_{s_iw_0}$ is an equivalence of categories.


Proof. We can define $\mathscr{F}_i^* \colon R$ -gmod $\to \widetilde{\mathscr{C}}_{s_i w_0}$ by the Schur-Weyl datum $\{\widehat{\mathbf{K}}_j^*\}_{j \in I}$ where

$$\mathbf{K}_{j}^{*} = \psi_{*}(\mathbf{K}_{j}) \simeq \begin{cases} \mathscr{D}^{-1} \big(\mathscr{Q}_{s_{i}w_{0}}(\langle i \rangle) \big) \simeq \mathsf{C}_{i}^{-1} \circ (E_{i}^{*}\mathsf{C}_{i}) & \text{if } j = i, \\ \langle j \rangle \nabla \langle i^{-\langle h_{i}, \alpha_{j} \rangle} \rangle & \text{if } j \neq i, \end{cases}$$

and

$$\widehat{\mathbf{K}}_{j}^{*} = \psi_{*}(\widehat{\mathbf{K}}_{j}) \simeq \begin{cases} \mathscr{D}_{\mathrm{aff}}^{-1} \big(\mathscr{Q}_{s_{i}w_{0}}(\langle i \rangle_{z_{i}}) \big) & \text{if } j = i, \\ \big(\langle j \rangle \nabla \langle i^{-\langle h_{i}, \alpha_{j} \rangle} \rangle \big)_{z_{j}} & \text{if } j \neq i. \end{cases}$$

Here the functor $\mathscr{Q}_{s_iw_0}$: $\operatorname{Pro}(R\operatorname{-gmod}) \to \operatorname{Pro}(\widetilde{\mathscr{C}}_{s_iw_0})$ is induced by $\mathscr{Q}_{s_iw_0}$: $R\operatorname{-gmod} \to \widetilde{\mathscr{C}}_{s_iw_0}$. Then, the functor \mathscr{F}_i^* : $R\operatorname{-gmod} \to \widetilde{\mathscr{C}}_{s_iw_0}$ factors through $\widetilde{\mathscr{C}}_{s_iw_0}^*$:



Note that \mathscr{F}_i^* is isomorphic to the composition

$$R\operatorname{-gmod} \xrightarrow{\psi_*} R\operatorname{-gmod} \xrightarrow{\mathscr{F}_i} \widetilde{\mathscr{C}}^*_{s_iw_0} \xrightarrow{\psi_*} \widetilde{\mathscr{C}}_{s_iw_0}.$$

We claim that the composition

$$\operatorname{Modg}_{\operatorname{coh}}(R) \longrightarrow \operatorname{Pro}(R\operatorname{-gmod}) \xrightarrow{\operatorname{Pro}(\mathscr{Q}_{s_iw_0}^*)} \operatorname{Pro}(\widetilde{\mathscr{Q}_{s_iw_0}}) \xrightarrow{\operatorname{Pro}(\mathscr{Q}_i^*)} \operatorname{Pro}(\widetilde{\mathscr{Q}_{s_iw_0}}) \xrightarrow{\operatorname{Pro}(\mathscr{Q}_i)} \operatorname{Pro}(\mathscr{Q}_i) \xrightarrow{Pro}(\mathscr{Q}_i) \xrightarrow{Pro}(\mathscr{Q}_i) \xrightarrow{Pro}(\mathscr{Q}_i) \xrightarrow{Pro}(\mathscr{Q}_i) \xrightarrow{Pro}(\mathscr{Q}_i)} \xrightarrow{Pro}(\mathscr{Q}_i) \xrightarrow{Pro}(\mathscr$$

is isomorphic to the functor associated with the duality datum

$$(\{\mathscr{Q}^*_{s_iw_0}(\langle j \rangle_{z_j})\}_{j \in I}, \{ \mathbb{R}^{\mathrm{ren}}_{\mathscr{Q}^*_{s_iw_0}(\langle j \rangle_{z_j}), \mathscr{Q}^*_{s_iw_0}(\langle k \rangle_{z_k})} \}_{j,k \in I})$$

which is of type C. Then by restriction to R-gmod, we have

$$\mathscr{S}_i \circ \mathscr{S}_i^* \circ \mathscr{Q}_{s_i w_0}^* \simeq \mathscr{Q}_{s_i w_0}^*$$

and by the universal property of the functor $\mathscr{Q}^*_{s_iw_0}$, we get $\mathscr{S}_i \circ \mathscr{S}^*_i \simeq \mathsf{id}_{\widetilde{\mathscr{C}}_{s_iw_0}}$, as desired.

By Lemma 7.7, it is enough to show that

$$(\mathscr{S}_i \circ \mathscr{S}_i^* \circ \mathscr{Q}_{s_i w_0}^*)(\langle j \rangle_{z_j}) \simeq \mathscr{Q}_{s_i w_0}^*(\langle j \rangle_z) \quad \text{for all } j \in I.$$

Since $(\mathscr{S}_i \circ \mathscr{S}_i^* \circ \mathscr{Q}_{s_i w_0}^*)(\langle j \rangle_{z_j}) \simeq (\mathscr{S}_i \circ \mathscr{F}_i^*)(\langle j \rangle_{z_j}) \simeq \mathscr{S}_i(\widehat{\mathbf{K}}_j^*)$, it is enough to show

(10.9) $\mathscr{S}_{i}(\widehat{\mathrm{K}}_{j}^{*}) \simeq \mathscr{Q}_{s_{i}w_{0}}^{*}(\langle j \rangle_{z_{j}})$ for all $j \in I$ up to a grading shift.

If j = i, then we have

$$\mathscr{S}_{i}(\widehat{\mathrm{K}}_{i}^{*}) = \mathscr{S}_{i}(\mathscr{D}_{\mathrm{aff}}^{-1}(\mathscr{Q}_{s_{i}w_{0}}(\langle i \rangle_{z_{i}}))) \simeq \mathscr{D}_{\mathrm{aff}}^{-1}(\mathscr{S}_{i}(\mathscr{Q}_{s_{i}w_{0}}(\langle i \rangle_{z_{i}}))))$$
$$\simeq \mathscr{D}_{\mathrm{aff}}^{-1}(\mathscr{F}_{i}(\langle i \rangle_{z_{i}})) \simeq \mathscr{D}_{\mathrm{aff}}^{-1}(\widehat{\mathrm{K}}_{i}) \simeq \mathscr{Q}_{s_{i}w_{0}}^{*}(\langle i \rangle_{z_{i}}).$$

When $\langle h_i, \alpha_j \rangle = 0$, (10.9) is trivial. If $\langle h_i, \alpha_j \rangle = -1$, then we have

$$\widehat{\mathbf{K}}_{j}^{*} \simeq \langle j \rangle_{z_{j}} \, \nabla_{z_{j}} \, \langle i \rangle_{z_{j}^{c}},$$

where $c = -\langle h_j, \alpha_i \rangle \in \mathbb{Z}_{\geq 1}$. Thus we have an epimorphism

$$\mathscr{S}_{i}(\langle j \rangle_{z_{j}} \underset{z_{j}}{\circ} \langle i \rangle_{z_{j}^{c}}) \twoheadrightarrow \mathscr{S}_{i}(\widehat{\mathrm{K}}_{j}^{*}).$$

On the other hand, we have

$$\mathscr{S}_{i}(\langle j \rangle_{z_{j}} \underset{z_{j}}{\circ} \langle i \rangle_{z_{j}^{c}}) \simeq \mathscr{S}_{i}(\langle j \rangle_{z_{j}}) \underset{z_{j}}{\circ} \mathscr{S}_{i}(\langle i \rangle_{z_{j}^{c}}) \simeq \widehat{\mathrm{K}}_{j} \underset{z_{j}}{\circ} \widehat{\mathrm{K}}_{i}|_{z_{i}=z_{j}^{c}} \twoheadrightarrow \langle j \rangle_{z_{j}},$$

where the last epimorphism is the one in (10.4). Hence we have $\mathscr{S}_i(\widehat{\mathbf{K}}_j^*) \simeq \langle j \rangle_{z_j}$ by Proposition 6.25.

If $\langle h_i, \alpha_j \rangle > 1$, then then we have

$$\widehat{\mathbf{K}}_{j}^{*}|_{z_{j}=w^{c}}\simeq\langle j\rangle_{w^{c}}\,\nabla_{w}\,\mathsf{L}\simeq\langle j\rangle_{w^{c}}\,\nabla_{w}\;(\langle i\rangle_{w}\underset{w}{\circ}\mathsf{L}'),$$

where $c = -\langle h_i, \alpha_j \rangle$ and L, L' are the ones in (10.5). Hence there is an epimorphism

$$\mathscr{S}_{i}(\langle j \rangle_{w^{c}}) \underset{w}{\circ} \mathscr{S}_{i}(\langle i \rangle_{w}) \underset{w}{\circ} \mathscr{S}_{i}(\mathsf{L}') \twoheadrightarrow \mathscr{S}(\widehat{\mathsf{K}}_{j}^{*}|_{z_{j}=w^{c}}).$$

On the other hand, by (10.6) we have

$$\mathscr{S}_{i}(\langle j \rangle_{w^{c}}) \underset{w}{\circ} \mathscr{S}_{i}(\langle i \rangle_{w}) \underset{w}{\circ} \mathscr{S}_{i}(\mathsf{L}') \simeq (\widehat{\mathsf{K}}_{j}|_{z_{j}=w^{c}} \underset{w}{\circ} \widehat{\mathsf{K}}_{i}|_{z_{i}=w}) \underset{w}{\circ} \mathscr{S}_{i}(\mathsf{L}')$$

$$\xrightarrow{(10.6)} (\mathsf{L}' \nabla_{w} \langle j \rangle_{w^{c}}) \underset{w}{\circ} \mathscr{S}_{i}(\mathsf{L}') \twoheadrightarrow (\mathsf{L}' \nabla_{w} \langle j \rangle_{w^{c}}) \nabla_{w} \mathscr{S}_{i}(\mathsf{L}') \simeq \langle j \rangle_{w^{c}},$$

where the last isomorphism follows from that $\mathscr{S}_i(\mathsf{L}') \simeq \mathscr{D}_{\mathrm{aff}}(\mathsf{L}')$. Hence we have $\mathscr{S}_i(\widehat{\mathsf{K}}_i^*) \simeq \langle j \rangle_{z_i}$ by Proposition 6.25, as desired.

Conjecture 10.12. The monoidal functor $\mathscr{S}_i : \widetilde{\mathscr{C}}_{s_iw_0} \to \widetilde{\mathscr{C}}^*_{s_iw_0}$ induces an equivalence of categories $\mathscr{C}_{s_iw_0} \to \mathscr{C}^*_{s_iw_0}$.

We know already that \mathscr{S}_i sends simples in $\mathscr{C}_{s_iw_0}$ to simples in $\mathscr{C}^*_{s_iw_0}$. However, we do not know $\mathscr{S}_i(\mathscr{C}_{s_iw_0}) \subset \mathscr{C}^*_{s_iw_0}$. Remark that $\mathscr{C}^*_{s_iw_0}$ is not stable by extensions in $\widetilde{\mathscr{C}}^*_{s_iw_0}$ (see Remark 8.6).

Remark 10.13. In the A_2 -case, we have $\mathfrak{d}(\mathbf{K}_j, \mathbf{K}_k) = 0 \neq \mathfrak{d}(\langle j \rangle, \langle k \rangle)$. We define \mathscr{F}_i as follows. Take $I = \{1, 2\}, Q_{1,2}(t_1.t_2) = t_1 - t_2$ and i = 1. Then $s_i w_0 = s_2 s_1$.

The renormalized R-matrix $\mathrm{R}_{j,k}^{\mathrm{ren}} \colon \langle j \rangle_{z_j} \circ \langle k \rangle_{z_k} \to \langle k \rangle_{z_k} \circ \langle j \rangle_{z_k}$ is given by $\langle j \rangle_{z_j} \boxtimes \langle k \rangle_{z_k} \mapsto \tau_1(\langle k \rangle_{z_k} \boxtimes \langle j \rangle_{z_k})$ for $j \neq k$. Since $\mathfrak{d}(K_1, K_2) = 0$, we can choose $\mathrm{R}_{j,k} \colon \widehat{\mathrm{K}}_j \circ \widehat{\mathrm{K}}_k \to \widehat{\mathrm{K}}_k \circ \widehat{\mathrm{K}}_j$ $(j \neq k)$ such that $\mathrm{R}_{j,k} \circ \mathrm{R}_{k,j} = \mathrm{id}_{\widehat{\mathrm{K}}_k \circ \widehat{\mathrm{K}}_j}$. Then, we define $\mathscr{F}_1 \colon R\text{-gmod} \to \widetilde{\mathscr{C}}_{s_i w_0}^*$ by:

$$\mathscr{F}_1(\langle j \rangle_{z_j}) = \widehat{\mathcal{K}}_j \ (j = 1, 2), \ \mathscr{F}_1(\mathcal{R}_{1,2}^{\text{ren}}) = Q_{1,2}(z_1, z_2)\mathcal{R}_{1,2} \text{ and } \ \mathscr{F}_1(\mathcal{R}_{2,1}^{\text{ren}}) = \mathcal{R}_{2,1}$$

Then it induces an equivalence $\mathscr{S}_1 \colon \widetilde{\mathscr{C}}_{s_i w_0} \xrightarrow{\sim} \widetilde{\mathscr{C}}^*_{s_i w_0}$.

Set $\langle 12 \rangle := \langle 1 \rangle \nabla \langle 2 \rangle$ and $\langle 21 \rangle := \langle 2 \rangle \nabla \langle 1 \rangle$. Note that $\mathscr{Q}_{s_i w_0}(\langle 1 \rangle) \simeq \langle 2 \rangle \circ \langle 21 \rangle^{-1}$, $\mathscr{Q}_{s_i w_0}(\langle 12 \rangle) \simeq 0$. The functor \mathscr{F}_1 sends:

$$\begin{array}{ll} \langle 1 \rangle \longmapsto \langle 2 \rangle \circ \langle 12 \rangle^{-1}, & \langle 12 \rangle \longmapsto 0, \\ \langle 2 \rangle \longmapsto \langle 12 \rangle, & \langle 21 \rangle \longmapsto \langle 2 \rangle. \end{array}$$

11. Examples of \mathscr{F}_i

Let us give examples of \mathscr{F}_i . We ignore grading shifts.

11.1. $\mathscr{F}_i \colon R\text{-}\mathrm{gmod} \to \widetilde{\mathscr{C}}^*_{s_i w_0}.$

11.1.1. A_2 . i = 1

$$\begin{array}{ll} \langle 1 \rangle \mapsto \langle 2 \rangle \circ \langle 12 \rangle^{-1} & \langle 12 \rangle \mapsto 0 \\ \langle 2 \rangle \mapsto \langle 12 \rangle & \langle 21 \rangle \mapsto \langle 2 \rangle \end{array}$$

11.1.2. A_3 .

 $\begin{array}{ll} (\mathrm{i}) \ i = 1. \\ & \langle 1 \rangle \mapsto \langle 23 \rangle \circ \langle 123 \rangle^{-1} & \langle 12 \rangle \mapsto \langle 2312 \rangle \circ \langle 123 \rangle^{-1} & \langle 132 \rangle \mapsto \langle 3 \rangle \circ \langle 2132 \rangle \circ \langle 123 \rangle^{-1} \\ & \langle 2 \rangle \mapsto \langle 12 \rangle & \langle 21 \rangle \mapsto \langle 2 \rangle & \langle 213 \rangle \mapsto \langle 23 \rangle \\ & \langle 3 \rangle \mapsto \langle 3 \rangle & \langle 23 \rangle \mapsto \langle 123 \rangle & \langle 123 \rangle \mapsto \langle 0 \\ & & \langle 32 \rangle \mapsto \langle 132 \rangle & \langle 321 \rangle \mapsto \langle 32 \rangle \\ & \langle 321 \rangle^{-1} \circ \langle 32 \rangle \mapsto \langle 1 \rangle & \langle 2132 \rangle \mapsto \langle 2132 \rangle \end{array}$

(ii) i = 2.

$$\begin{array}{ll} \langle 1 \rangle \mapsto \langle 21 \rangle & \langle 12 \rangle \mapsto \langle 1 \rangle & \langle 132 \rangle \mapsto \langle 213 \rangle \\ \langle 2 \rangle \mapsto \langle 13 \rangle \circ \langle 213 \rangle^{-1} & \langle 21 \rangle \mapsto \langle 1 \rangle \circ \langle 321 \rangle \circ \langle 213 \rangle^{-1} & \langle 213 \rangle \mapsto \langle 123 \rangle \circ \langle 321 \rangle \circ \langle 213 \rangle^{-1} \\ \langle 3 \rangle \mapsto \langle 23 \rangle & \langle 23 \rangle \mapsto \langle 3 \rangle \circ \langle 123 \rangle \circ \langle 213 \rangle^{-1} & \langle 123 \rangle \mapsto \langle 123 \rangle \\ & \langle 32 \rangle \mapsto \langle 3 \rangle & \langle 321 \rangle \mapsto \langle 321 \rangle \\ & \langle 321 \rangle \mapsto \langle 321 \rangle \\ & \langle 2132 \rangle \mapsto 0 \end{array}$$

(ii) i = 2.

$$\begin{array}{ll} \langle 1 \rangle \mapsto \langle 21 \rangle & \langle 1^2 2 \rangle \mapsto \langle 21^2 \rangle \\ \langle 2 \rangle \mapsto \langle 1^2 \rangle \circ \langle 21^2 \rangle^{-1} & \langle 21^2 \rangle \mapsto \langle 121 \rangle^2 \circ \langle 21^2 \rangle^{-1} \\ \langle 12 \rangle \mapsto \langle 1 \rangle & \langle 121 \rangle \circ \langle 21^2 \rangle^{-1} & \langle 121 \rangle \mapsto \langle 121 \rangle \\ \langle 21 \rangle \mapsto \langle 1 \rangle \circ \langle 121 \rangle \circ \langle 21^2 \rangle^{-1} & \langle 21^2 2 \rangle \mapsto 0 \end{array}$$

11.2. $\mathscr{F}_i \colon \widetilde{\mathscr{C}}_{w_0} \to \widetilde{\mathscr{C}}_{w_0}.$

11.2.1.
$$A_2$$
. $i = 12$, i.e. $\mathscr{F}_1 \mathscr{F}_2$:
 $\langle 1 \rangle \mapsto \mathscr{D} \langle 2 \rangle = \langle 1 \rangle \circ \langle 21 \rangle^{-1} \qquad \langle 12 \rangle \mapsto \langle 21 \rangle^{-1}$
 $\langle 2 \rangle \mapsto \mathscr{D} \langle 1 \rangle = \langle 2 \rangle \circ \langle 12 \rangle^{-1} \qquad \langle 21 \rangle \mapsto \langle 12 \rangle^{-1}$

11.2.2. A_3 .

(i) i = 2.

 $\begin{array}{cccc} \langle 1 \rangle \mapsto \langle 21 \rangle & \langle 12 \rangle \mapsto \langle 1 \rangle & \langle 132 \rangle \mapsto \langle 213 \rangle \\ \langle 2 \rangle \mapsto \langle 132 \rangle \circ \langle 2132 \rangle^{-1} & \langle 21 \rangle \mapsto \langle 12 \rangle \circ \langle 321 \rangle \circ \langle 2132 \rangle^{-1} & \langle 213 \rangle \mapsto \langle 2 \rangle \circ \langle 123 \rangle \circ \langle 321 \rangle \circ \langle 2132 \rangle^{-1} \\ \langle 3 \rangle \mapsto \langle 23 \rangle & \langle 23 \rangle \mapsto \langle 32 \rangle \circ \langle 123 \rangle \circ \langle 2132 \rangle^{-1} & \langle 123 \rangle \mapsto \langle 123 \rangle \\ & \langle 32 \rangle \mapsto \langle 3 \rangle & \langle 321 \rangle \mapsto \langle 321 \rangle \\ & \langle 2132 \rangle \mapsto \langle 123 \rangle \circ \langle 321 \rangle \circ \langle 2132 \rangle^{-1} \end{array}$

$$\begin{array}{ll} \text{(ii)} & i = 13 \\ & \langle 1 \rangle \mapsto \mathscr{D}1 = \langle 23 \rangle \circ \langle 123 \rangle^{-1} & \langle 12 \rangle \mapsto \langle 3 \rangle \circ \langle 2132 \rangle \circ \langle 123 \rangle^{-1} & \langle 132 \rangle \mapsto \langle 213 \rangle \circ \langle 2132 \rangle \circ \langle 123 \rangle^{-1} \circ \langle 321 \rangle^{-1} \\ & \langle 2 \rangle \mapsto \langle 132 \rangle & \langle 21 \rangle \mapsto \langle 32 \rangle & \langle 213 \rangle \mapsto \langle 2\rangle \\ & \langle 3 \rangle \mapsto \mathscr{D}3 = \langle 21 \rangle \circ \langle 321 \rangle^{-1} & \langle 23 \rangle \mapsto \langle 12 \rangle & \langle 123 \rangle & \Rightarrow \langle 2132 \rangle \circ \langle 123 \rangle^{-1} \\ & & \langle 32 \rangle \mapsto \langle 1 \rangle \circ \langle 2132 \rangle \circ \langle 321 \rangle^{-1} & \langle 321 \rangle \mapsto \langle 2132 \rangle \circ \langle 321 \rangle^{-1} \\ & & \langle 2132 \rangle \mapsto \langle 2132 \rangle \circ \langle 321 \rangle^{-1} \\ & & \langle 2132 \rangle \mapsto \langle 2132 \rangle \end{array}$$

$$\begin{aligned} 1 & 2 \\ \text{(i)} \ i &= 1. \end{aligned}$$

$$\begin{aligned} \langle 1 \rangle \mapsto \langle 21 \rangle \circ \langle 121 \rangle^{-1} & \langle 1^2 2 \rangle \mapsto \langle 21^2 \rangle \circ \langle 21^2 2 \rangle \circ \langle 121 \rangle^{-2} \\ \langle 2 \rangle \mapsto \langle 1^2 2 \rangle & \langle 21^2 \rangle \mapsto \langle 2 \rangle \\ \langle 12 \rangle \mapsto \langle 1 \rangle \circ \langle 21^2 2 \rangle \circ \langle 121 \rangle^{-1} & \langle 121 \rangle \mapsto \langle 21^2 2 \rangle \circ \langle 121 \rangle^{-1} \\ \langle 21 \rangle \mapsto \langle 12 \rangle & \langle 21^2 2 \rangle \mapsto \langle 21^2 2 \rangle \end{aligned}$$

(ii) i = 2.

11.2.3. C_2 . $\circ \longleftarrow \circ$

 $\begin{array}{ll} \langle 1 \rangle \mapsto \langle 21 \rangle & \langle 1^2 2 \rangle \mapsto \langle 21^2 \rangle \\ \langle 2 \rangle \mapsto \langle 1^2 2 \rangle \circ \langle 21^2 2 \rangle^{-1} & \langle 21^2 \rangle \mapsto \langle 2 \rangle \circ \langle 121 \rangle^2 \circ \langle 21^2 2 \rangle^{-1} \\ \langle 12 \rangle \mapsto \langle 1 \rangle & \langle 121 \rangle \mapsto \langle 121 \rangle \\ \langle 21 \rangle \mapsto \langle 12 \rangle \circ \langle 121 \rangle \circ \langle 21^2 2 \rangle^{-1} & \langle 21^2 2 \rangle \mapsto \langle 121 \rangle^2 \circ \langle 21^2 2 \rangle^{-1} \\ (\mathscr{S}_1 \circ \mathscr{S}_2)^2 \simeq (\mathscr{S}_2 \circ \mathscr{S}_1)^2. \end{array}$

References

- V. Chari and A. Pressley, A Guide to Quantum Groups, Cambridge University Press, Cambridge, 1994.
- [2] N. Chriss and V. Ginzburg, Representation theory and complex geometry, Birkhäuser Boston, Inc., Boston, MA, 1997.
- [3] P. Etingof, S. Gelaki, D. Nikshych and V. Ostrik, *Tensor Categories*, Mathematical Surveys and Monographs, 205, American Mathematical Society, Providence, RI, 2015. xvi+343 pp.
- [4] P. Etingof and A. A. Moura, Elliptic central characters and blocks of finite dimensional representations of quantum affine algebras, Represent. Theory 7 (2003), 346–373.
- [5] M. Kashiwara, On level zero representations of quantum affine algebras, Duke. Math. J. 112 (2002), 117–175.
- [6] S.-J. Kang, M. Kashiwara, M. Kim and S.-j. Oh, Simplicity of heads and socles of tensor products, Compos. Math. 151 (2015), no. 2, 377–396.
- [7] S.-J. Kang, M. Kashiwara and M. Kim, Symmetric quiver Hecke algebras and R-matrices of quantum affine algebras, Invent. math. 211 (2018), 591–685, arXiv:1304.0323 v3.
- [8] _____, Monoidal categorification of cluster algebras, J. Amer. Math. Soc. 31 (2018), no. 2, 349–426.
- M. Kashiwara, M. Kim, S.-j. Oh, and E. Park, Monoidal categories associated with strata of flag manifolds, Adv. Math. 328 (2018), 959–1009.
- [10] _____, Monoidal categorification and quantum affine algebras, Compos. Math. 156 (2020), no. 5, 1039–1077.
- [11] _____, Localizations for quiver Hecke algebras, Pure Appl. Math. Q. 17 (2021), no. 4, 1465–1548.
- [12] _____, Localizations for quiver Hecke algebras II, Proc. London Math. Soc. (4) 127 (2023) 1134–1184.

- [13] M. Kashiwara and E. Park, Affinizations and R-matrices for quiver Hecke algebras, J. Eur. Math. Soc. 20, (2018), 1161–1193.
- [14] M. Kashiwara and P. Schapira, *Categories and Sheaves*, Grundlehren der Mathematischen Wissenschaften, vol. 332, Springer-Verlag, Berlin (2006).
- [15] S. Kato, Poincaré-Birkhoff-Witt bases and Khovanov-Lauda-Rouquier algebras Duke Math. J. 163 (2014), no. 3, 619–663.
- [16] _____, On the monoidality of Saito reflection functors, Int. Math. Res. Not. 2020, no. 22, 8600– 8623.
- [17] M. Khovanov and A. Lauda, A diagrammatic approach to categorification of quantum groups I, Represent. Theory 13 (2009), 309–347.
- [18] _____, A diagrammatic approach to categorification of quantum groups II, Trans. Amer. Math. Soc. **363** (2011), 2685–2700.
- [19] A. Lauda and M. Vazirani, Crystals from categorified quantum groups, Adv. Math. 228 (2011), no. 2, 803–861.
- [20] G. Lusztig, Introduction to Quantum Groups, Progr. Math. 110 Birkhäuser, 1993.
- [21] P. J. McNamara, Finite dimensional representations of Khovanov-Lauda-Rouquier algebras I: Finite type J. reine angew. Math. 707 (2015), 103–124.
- [22] _____, Monoidality of Kato's reflection functors, arXiv:1712.00173v1.
- [23] R. Rouquier, 2-Kac–Moody algebras, arXiv:0812.5023v1.
- [24] Y. Saito, PBW basis of quantized universal enveloping algebras, Publ. Res. Inst. Math. Sci. 30, (1994), no. 2, 209–232.
- [25] P. Tingley and B. Webster, Mirković-Vilonen polytopes and Khovanov-Lauda-Rouquier algebras, Compos. Math. 152 (2016), no. 8, 1648–1696.

M. KASHIWARA, M. KIM, S.-J. OH, AND E. PARK

(M. Kashiwara) Kyoto University Institute for Advanced Study, Research Institute for Mathematical Sciences, Kyoto University, Kyoto 606-8502, Japan & Korea Institute for Advanced Study, Seoul 02455, Korea

Email address, M. Kashiwara: masaki@kurims.kyoto-u.ac.jp

(M. Kim) DEPARTMENT OF MATHEMATICS, KYUNG HEE UNIVERSITY, SEOUL 02447, KOREA *Email address*, M. Kim: mkim@khu.ac.kr

(S.-j. Oh) Department of Mathematics, Sungkyunkwan University, Suwon 16419 , Korea

Email address, S.-j. Oh: sejin0920gmail.com

(E. Park) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SEOUL, SEOUL 02504, KOREA *Email address*, E. Park: epark@uos.ac.kr