

# The Spectral Einstein functional and Kastler-Kalau-Walze type theorems

Yuchen Yang, Tong Wu\*

*School of Mathematics and Statistics, Northeast Normal University, Changchun, 130024, China*

---

## Abstract

In this paper, on the basis of defining the spectral Einstein functional associated with the Dirac operator for manifolds with boundary, we prove Kastler-Kalau-Walze type theorem for the spectral Einstein functional associated with the Dirac operator on low-dimensional manifolds with boundary.

*Keywords:* The spectral Einstein functional; the Dirac operator; Kastler-Kalau-Walze type theorem;

---

*2010 Mathematics Subject Classification:* 53C40; 53C42.

## 1. Introduction

In recent years, many geometers have made more in-depth study of noncommutative residues. In [6], Guillemin found noncommutative residues are of great importance to the study of noncommutative geometry. In the 1990s, the scholar Connes induced a conformal 4-dimensional Polyakov action analogy by using non-commutative residues[2], he also showed that the non-commutative residues on the compact manifold equals the Dixmier's trace on the pseudo-differential operators with order  $-\dim M$  in [3], so that the non-commutative residues can be used as a non-commutative integral. Connes conjectured that the non-commutative residue of the  $-2$  power of the Dirac operator is proportional to the Einstein-Hilbert action. In 1995, Kastler[8] proved this conjecture. At the same time, Kalau and Walze also proved this conjecture in the normal coordinate system in [7]. In 1996, Ackermann found that the noncommutative residue of the inverse square of the Dirac operator  $\widehat{\text{Wres}}(D^{-2})$  is essentially the second coefficient of the heat kernel expansion of the square of the Dirac operator in [1]. In the same year, Fedosov et al. proposed the idea of noncommutative residue on manifolds with boundary. They defined the non-commutative residue on Boutet de Monvel algebra and proved that it is a unique continuous trace in [5].

On the other hand, Wang provides a kind of method to study the Kastler-Kalau-Walze type theorem for manifolds with boundary. In the early 21st century, Chinese scholar Wang Yong extended the results of Connes to manifolds with boundary in [10, 11], studied the conformal invariant problem and gravity action of manifolds with boundary, and proved the Kastler-Kalau-Walze type theorems of Dirac operator and the Signature operator on low dimensional manifolds with boundary [12]. In [13], Wang defined lower dimensional volumes of spin manifolds with boundary. And computed the lower dimensional volume  $Vol_6^{(2,2)}$  and got a Kastler-Kalau-Walze type theorem in 6-dimensional spin manifolds with boundary and the associated Dirac operator  $D$ . They computed the lower dimensional volume  $Vol_6^{(1,3)}$  and got a Kastler-Kalau-Walze type theorem in 6-dimensional spin manifolds with boundary and the associated Dirac operator  $D$  and  $D^3$  in [9]. In [4], Dabrowski L, Sitarz A and Zalecki P defined bilinear functionals of vector fields and differential forms, the densities of which yield the metric and Einstein tensors on even-dimensional Riemannian manifolds and proved that for the conformally rescaled geometry of the noncommutative two-torus the Einstein functional vanishes. Motivated by [4], we define the spectral Einstein functional associated with the Dirac

---

\*Corresponding author.

*Email addresses:* yangyc580@nenu.edu.cn (Yuchen Yang), wut977@nenu.edu.cn (Tong Wu)

operator for manifolds with boundary, and the motivation of this paper is to compute the noncommutative residue  $\widetilde{\text{Wres}}[\pi^+(\nabla_X^{S(TM)}\nabla_Y^{S(TM)}D^{-1})\circ\pi^+(D^{-3})]$  on 4-dimensional spin manifolds with boundary and  $\widetilde{\text{Wres}}[\pi^+(\nabla_X^{S(TM)}\nabla_Y^{S(TM)}D^{-3})\circ\pi^+(D^{-3})]$  on 6-dimensional spin manifolds with boundary.

The paper is organized in the following way. In Section 2, we define the spectral Einstein functional associated with the Dirac operator and get the noncommutative residue for manifolds without boundary. In Section 3, we prove the Kastler-Kalau-Walze type theorem for the spectral Einstein functional associated with the Dirac operator on 4-dimensional manifolds with boundary and we get a Kastler-Kalau-Walze type theorem associated for 3-dimensional spin manifolds in Section 4. In Section 5, for 6-dimensional manifolds with boundary and the spectral Einstein functional associated with the Dirac operator, we compute the Kastler-Kalau-Walze type theorem.

## 2. The spectral Einstein functional associated with the Dirac operator

Firstly we review the relevant knowledge of the Dirac operator. We have  $M$  is an  $n$ -dimensional ( $n \geq 3$ ) oriented compact Riemannian manifold with a Riemannian metric  $g^M$  and let  $\nabla^L$  be the Levi-Civita connection about  $g^M$ . In the fixed orthonormal frame  $\{e_1, \dots, e_n\}$ , the connection matrix  $(\omega_{s,t})$  is defined by

$$\nabla^L(e_1, \dots, e_n) = (e_1, \dots, e_n)(\omega_{s,t}). \quad (2.1)$$

Let  $\epsilon(e_j^*)$ ,  $\iota(e_j^*)$  be the exterior and interior multiplications respectively, where  $e_j^* = g^{TM}(e_j, \cdot)$ . Write

$$\widehat{c}(e_j) = \epsilon(e_j^*) + \iota(e_j^*); \quad c(e_j) = \epsilon(e_j^*) - \iota(e_j^*), \quad (2.2)$$

which satisfies

$$\begin{aligned} \widehat{c}(e_i)\widehat{c}(e_j) + \widehat{c}(e_j)\widehat{c}(e_i) &= 2g^M(e_i, e_j); \\ c(e_i)c(e_j) + c(e_j)c(e_i) &= -2g^M(e_i, e_j); \\ c(e_i)\widehat{c}(e_j) + \widehat{c}(e_j)c(e_i) &= 0. \end{aligned} \quad (2.3)$$

By [15], we have the Dirac operator

$$D = \sum_{i=1}^n c(e_i)[e_i - \frac{1}{4} \sum_{s,t} \omega_{s,t}(e_i)c(e_s)c(e_t)]. \quad (2.4)$$

We define spin connection by  $\nabla_X^{S(TM)} := X + \frac{1}{4} \sum_{ij} \langle \nabla_X^L e_i, e_j \rangle c(e_i)c(e_j)$ . Let  $A(X) = \frac{1}{4} \sum_{ij} \langle \nabla_X^L e_i, e_j \rangle c(e_i)c(e_j)$ , so we have

$$\begin{aligned} \nabla_X^{S(TM)}\nabla_Y^{S(TM)} &= [X + A(X)][Y + A(Y)] \\ &= XY + X \cdot A(Y) + A(X)Y + A(X)A(Y) \\ &= XY + X[A(Y)] + A(Y)X + A(X)Y + A(X)A(Y), \end{aligned} \quad (2.5)$$

where  $X = \sum_{j=1}^n X_j \partial_{x_j}$ ,  $Y = \sum_{l=1}^n Y_l \partial_{x_l}$  and  $XY = \sum_{j,l=1}^n (X_j Y_l \partial_{x_j} \partial_{x_l} + X_j \frac{\partial Y_l}{\partial x_j} \partial_{x_l})$ .

Let  $g^{ij} = g(dx_i, dx_j)$ ,  $\xi = \sum_k \xi_j dx_j$  and  $\nabla_{\partial_i}^L \partial_j = \sum_k \Gamma_{ij}^k \partial_k$ , we denote that

$$\sigma_i = -\frac{1}{4} \sum_{s,t} \omega_{s,t}(e_i)c(e_s)c(e_t); \quad \xi^j = g^{ij}\xi_i; \quad \Gamma^k = g^{ij}\Gamma_{ij}^k; \quad \sigma^j = g^{ij}\sigma_i; \quad a^j = g^{ij}a_i. \quad (2.6)$$

And by  $\partial_{x_j} = -\sqrt{-1}\xi_j$ , we have the following lemmas.

**Lemma 2.1.** *The following identities hold:*

$$\begin{aligned}
\sigma_0(\nabla_X^{S(TM)} \nabla_Y^{S(TM)}) &= X[A(Y)] + A(X)A(Y); \\
\sigma_1(\nabla_X^{S(TM)} \nabla_Y^{S(TM)}) &= \sqrt{-1} \sum_{j,l=1}^n X_j \frac{\partial Y_l}{\partial x_j} \partial_{x_l} + \sqrt{-1} \sum_j A(Y) X_j \xi_j + \sqrt{-1} \sum_l A(Y) Y_l \xi_l; \\
\sigma_2(\nabla_X^{S(TM)} \nabla_Y^{S(TM)}) &= -\sum_{j,l=1}^n X_j Y_l \xi_j \xi_l.
\end{aligned} \tag{2.7}$$

**Lemma 2.2.** *By[9], the following identities hold:*

$$\begin{aligned}
\sigma_{-1}(D^{-1}) &= \frac{\sqrt{-1}c(\xi)}{|\xi|^2}; \\
\sigma_{-2}(D^{-1}) &= \frac{c(\xi)\sigma_0(D)c(\xi)}{|\xi|^4} + \frac{c(\xi)}{|\xi|^6} \sum_j c(dx_j) (\partial_{x_j}[c(\xi)]|\xi|^2 - c(\xi)\partial_{x_j}(|\xi|^2)); \\
\sigma_{-3}(D^{-3}) &= -\sqrt{-1}|\xi|^{-4}c(\xi); \\
\sigma_{-4}(D^{-3}) &= \frac{c(\xi)\sigma_2(D^3)c(\xi)}{|\xi|^8} + \frac{\sqrt{-1}c(\xi)}{|\xi|^8} \left( |\xi|^4 c(dx_n) \partial_{x_n} c(\xi') - 2h'(0)c(dx_n)c(\xi) \right. \\
&\quad \left. + 2\xi_n c(\xi) \partial_{x_n} c(\xi') + 4\xi_n h'(0) \right);
\end{aligned} \tag{2.8}$$

We write

$$D_x^\alpha = (-i)^{|\alpha|} \partial_x^\alpha; \quad \sigma(D_t^3) = p_3 + p_2 + p_1 + p_0; \quad (\sigma(D_t)^{-3}) = \sum_{j=3}^\infty q_{-j}. \tag{2.9}$$

By the composition formula of pseudodifferential operators, we have

$$\begin{aligned}
1 = \sigma(D^3 \circ D^{-3}) &= \sum_\alpha \frac{1}{\alpha!} \partial_\xi^\alpha [\sigma(D) D_x^\alpha [\sigma(D^{-3})]] \\
&= (p_3 + p_2 + p_1 + p_0)(q_{-3} + q_{-4} + q_{-5} + \dots) \\
&\quad + \sum_j (\partial_{\xi_j} p_3 + \partial_{\xi_j} p_2 + \dots + \partial_{\xi_j} p_1 + \partial_{\xi_j} p_0) (D_{x_j} q_{-3} + D_{x_j} q_{-4} + D_{x_j} q_{-5} + \dots) \\
&= p_3 q_{-3} + (p_3 q_{-4} + p_2 q_{-3} + \sum_j \partial_{\xi_j} p_3 D_{x_j} q_{-3}) + \dots,
\end{aligned} \tag{2.10}$$

so

$$q_{-3} = p_3^{-1}; \quad q_{-4} = -p_3^{-1} [p_2 p_3^{-1} + \sum_j \partial_{\xi_j} p_3 D_{x_j} (p_3^{-1})]. \tag{2.11}$$

Then,

$$\begin{aligned}
\sigma_1(\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-1}) &= -\sqrt{-1} \sum_{j,l=1}^n X_j Y_l \xi_j \xi_l c(\xi) |\xi|^{-2}; \\
\sigma_{-1}(\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-3}) &= \sqrt{-1} \sum_{j,l=1}^n X_j Y_l \xi_j \xi_l c(\xi) |\xi|^{-4}.
\end{aligned} \tag{2.12}$$

According to [4], we have the following theorem

**Theorem 2.3.** *If  $M$  is an  $n$ -dimensional compact oriented manifolds without boundary, and  $n$  is even, then we get the following equality:*

$$\text{Wres}[\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-n}] = \frac{(2\pi)^{\frac{n}{2}}}{3(\frac{n}{2}-1)!} \int_M [\text{Ric}(X, Y) - \frac{1}{2} \text{sg}(X, Y)] d\text{Vol}_M + \frac{(2\pi)^{\frac{n}{2}}}{4(\frac{n}{2}-1)!} \int_M \text{sg}(X, Y) d\text{Vol}_M, \tag{2.13}$$

where  $s$  is the scalar curvature,  $\text{Vol}_M$  is the volume of  $M$  and  $\text{Ric}$  denotes Ricci tensor on  $M$ .

### 3. The noncommutative residue for 4-dimensional manifolds with boundary

In this section, we compute the noncommutative residue  $\widehat{\text{Wres}}[\pi^+(\nabla_X^{S(TM)}\nabla_Y^{S(TM)}D^{-1})\circ\pi^+(D^{-3})]$  on 4-dimensional oriented compact manifolds with boundary. We review some basic facts and formulas about Boutet de Monvel's calculus and the definition of the noncommutative residue for manifolds with boundary which will be used in the following. Details are detailed in the the Section 2 in [12].

Let  $M$  be a 4-dimensional compact oriented manifold with boundary  $\partial M$ . We assume that the metric  $g^M$  on  $M$  has the following form near the boundary,

$$g^M = \frac{1}{h(x_n)}g^{\partial M} + dx_n^2, \quad (3.1)$$

where  $g^{\partial M}$  is the metric on  $\partial M$  and  $h(x_n) \in C^\infty([0, 1]) := \{\widehat{h}|_{[0,1]}\widehat{h} \in C^\infty((-\varepsilon, 1))\}$  for some  $\varepsilon > 0$  and  $h(x_n)$  satisfies  $h(x_n) > 0$ ,  $h(0) = 1$ , where  $x_n$  denotes the normal directional coordinate.

Similar to [12], then we can compute the noncommutative residue

$$\begin{aligned} & \widehat{\text{Wres}}[\pi^+(\nabla_X^{S(TM)}\nabla_Y^{S(TM)}D^{-1})\circ\pi^+(D^{-3})] \\ &= \int_M \int_{|\xi|=1} \text{trace}_{\wedge^*T^*M} \otimes_{\mathbb{C}} [\sigma_{-4}(\nabla_X^{S(TM)}\nabla_Y^{S(TM)}D^{-1}\circ D^{-3})] \sigma(\xi) dx + \int_{\partial M} \Phi, \end{aligned} \quad (3.2)$$

where

$$\begin{aligned} \Phi &= \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \sum_{j,k=0}^{\infty} \sum \frac{(-i)^{|\alpha|+j+k+1}}{\alpha!(j+k+1)!} \times \text{trace}_{\wedge^*T^*M} \otimes_{\mathbb{C}} [\partial_{x_n}^j \partial_{\xi'}^\alpha \partial_{\xi_n}^k \sigma_r^+(\nabla_X^{S(TM)}\nabla_Y^{S(TM)}D^{-1})(x', 0, \xi', \xi_n) \\ &\quad \times \partial_{x'}^\alpha \partial_{\xi_n}^{j+1} \partial_{x_n}^k \sigma_l(D^{-3})(x', 0, \xi', \xi_n)] d\xi_n \sigma(\xi') dx', \end{aligned} \quad (3.3)$$

and the sum is taken over  $r+l-k-j-|\alpha| = -4$ ,  $r \leq 1$ ,  $l \leq -3$ .

By Theorem 2.3, we can compute the interior of  $\widehat{\text{Wres}}[\pi^+(\nabla_X^{S(TM)}\nabla_Y^{S(TM)}D^{-1})\circ\pi^+(D^{-3})]$ , we get

$$\text{Wres}[\nabla_X^{S(TM)}\nabla_Y^{S(TM)}D^{-4}] = \frac{4\pi^2}{3} \int_M [\text{Ric}(X, Y) - \frac{1}{2}sg(X, Y)] d\text{Vol}_M + \pi^2 \int_M sg(X, Y) d\text{Vol}_M. \quad (3.4)$$

Now we need to compute  $\int_{\partial M} \Phi$ . When  $n = 4$ , then  $\text{tr}_{S(TM)}[\text{id}] = \dim(\wedge^*(\mathbb{R}^2)) = 4$ , the sum is taken over  $r+l-k-j-|\alpha| = -3$ ,  $r \leq 0$ ,  $l \leq -2$ , then we have the following five cases:

**case a) I)**  $r = 1$ ,  $l = -2$ ,  $k = j = 0$ ,  $|\alpha| = 1$ .

By (3.3), we get

$$\Phi_1 = - \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \sum_{|\alpha|=1} \text{tr}[\partial_{\xi'}^\alpha \pi_{\xi_n}^+ \sigma_1(\nabla_X^{S(TM)}\nabla_Y^{S(TM)}D^{-1}) \times \partial_{x'}^\alpha \partial_{\xi_n} \sigma_{-3}(D^{-3})](x_0) d\xi_n \sigma(\xi') dx'. \quad (3.5)$$

By Lemma 2.2 in [12], for  $i < n$ , then

$$\partial_{x_i} \sigma_{-3}(D^{-3})(x_0) = \partial_{x_i} (\sqrt{-1}c(\xi)|\xi|^{-4})(x_0) = \sqrt{-1} \frac{\partial_{x_i} c(\xi)}{|\xi|^4} (x_0) + \sqrt{-1} \frac{c(\xi) \partial_{x_i} (|\xi|^4)}{|\xi|^8} (x_0) = 0, \quad (3.6)$$

so  $\Phi_1 = 0$ .

**case a) II)**  $r = 1$ ,  $l = -3$ ,  $k = |\alpha| = 0$ ,  $j = 1$ .

By (3.3), we get

$$\Phi_2 = - \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\partial_{x_n} \pi_{\xi_n}^+ \sigma_1(\nabla_X^{S(TM)}\nabla_Y^{S(TM)}D^{-1}) \times \partial_{\xi_n}^2 \sigma_{-3}(D^{-3})](x_0) d\xi_n \sigma(\xi') dx'. \quad (3.7)$$

By Lemma 2.1 and Lemma 2.2, we have

$$\partial_{\xi_n}^2 \sigma_{-3}((D^{-3}))(x_0) = \partial_{\xi_n}^2 (c(\xi)|\xi|^{-4})(x_0) = \sqrt{-1} \frac{(20\xi_n^2 - 4)c(\xi') + 12(\xi^3 - \xi)c(dx_n)}{(1 + \xi_n^2)^4}, \quad (3.8)$$

and

$$\begin{aligned} \partial_{x_n} \sigma_1(\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-1}) &= \partial_{x_n} (-\sqrt{-1} \sum_{j,l=1}^n X_j Y_l \xi_j \xi_l c(\xi) |\xi|^{-2}) \\ &= \sum_{j,l=1}^n X_j Y_l \xi_j \xi_l \left[ \frac{\partial_{x_n} c(\xi')}{1 + \xi_n^2} + \frac{c(\xi) h'(0) |\xi'|^2}{(1 + \xi_n^2)^2} \right]. \end{aligned} \quad (3.9)$$

Then, we have

$$\begin{aligned} \pi_{\xi_n}^+ \partial_{x_n} \sigma_1(\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-1}) &= \partial_{x_n} \pi_{\xi_n}^+ \sigma_1(\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-1}) \\ &= \sqrt{-1} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l h'(0) |\xi'|^2 \left[ \frac{ic(\xi')}{4(\xi_n - i)} + \frac{c(\xi') + ic(dx_n)}{4(\xi_n - i)^2} \right] \\ &\quad - \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l \frac{\partial_{x_n} c(\xi')}{2(\xi_n - i)} \\ &\quad - \sqrt{-1} X_n Y_n \left\{ \frac{\partial_{x_n} c(\xi')}{2(\xi_n - i)} + h'(0) |\xi'| \left[ -\frac{2ic(\xi') - 3c(dx_n)}{4(\xi_n - i)} \right. \right. \\ &\quad \left. \left. + \frac{[c(\xi') + ic(dx_n)][i(\xi_n - i) + 1]}{4(\xi_n - i)^2} \right] \right\} \\ &\quad - \sum_{j=1}^{n-1} X_j Y_n \xi_j \left[ \sqrt{-1} \frac{\partial_{x_n} c(\xi')}{2(\xi_n - i)} - \frac{\sqrt{-1} h'(0) |\xi'| [c(\xi') + 2ic(dx_n)]}{4(\xi_n - i)} \right. \\ &\quad \left. - \frac{[ic(\xi') - c(dx_n)][i(\xi_n - i) + 1]}{(\xi_n - i)^2} \right] \\ &\quad - \sum_{l=1}^{n-1} X_n Y_l \xi_l \left[ \sqrt{-1} \frac{\partial_{x_n} c(\xi')}{2(\xi_n - i)} - \frac{\sqrt{-1} h'(0) |\xi'| [c(\xi') + 2ic(dx_n)]}{4(\xi_n - i)} \right. \\ &\quad \left. - \frac{[ic(\xi') - c(dx_n)][i(\xi_n - i) + 1]}{(\xi_n - i)^2} \right]. \end{aligned} \quad (3.10)$$

We note that  $i < n$ ,  $\int_{|\xi'|=1} \xi_{i_1} \xi_{i_2} \cdots \xi_{i_{2d+1}} \sigma(\xi') = 0$ , so we omit some items that have no contribution for computing **case a) II**.

Then there is the following formula

$$\begin{aligned} &\text{tr}[\partial_{x_n} \pi_{\xi_n}^+ \sigma_1(\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-1}) \times \partial_{\xi_n}^2 \sigma_{-3}(D^{-3})(x_0)] \\ &= \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l h'(0) \left[ 8i \frac{5\xi_n^2 - 1}{(\xi_n - i)^5 (\xi_n + i)^4} + \frac{4(5\xi_n^2 - 1) + 12i(\xi_n^3 - \xi_n)}{(\xi_n - i)^6 (\xi_n + i)^4} \right] \\ &\quad + X_n Y_n h'(0) \left[ \frac{(4i - 4)(\xi^2 - 1) + 48(\xi_n^3 - \xi_n)}{(\xi_n - i)^5 (\xi_n + i)^4} - \frac{4(5\xi_n^2 - 1) + 12i(\xi_n^3 - \xi_n)}{(\xi_n - i)^6 (\xi_n + i)^4} \right] \\ &\quad + 8 \sum_{j=1}^{n-1} X_j Y_n \xi_j \left[ \frac{(6 - 3ih'(0))(\xi_n^3 - \xi_n) - 2i(5\xi_n^2 - 1)}{(\xi_n - i)^5 (\xi_n + i)^4} + \frac{2(5\xi_n^2 - 1) + 6i(\xi_n^3 - \xi_n)}{(\xi_n - i)^6 (\xi_n + i)^4} \right] \\ &\quad + 8 \sum_{l=1}^{n-1} X_n Y_l \xi_l \left[ \frac{(6 - 3ih'(0))(\xi_n^3 - \xi_n) - 2i(5\xi_n^2 - 1)}{(\xi_n - i)^5 (\xi_n + i)^4} + \frac{2(5\xi_n^2 - 1) + 6i(\xi_n^3 - \xi_n)}{(\xi_n - i)^6 (\xi_n + i)^4} \right]. \end{aligned} \quad (3.11)$$

Therefore, we get

$$\begin{aligned}
\Phi_2 &= \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \left\{ \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l h'(0) \left[ 8i \frac{5\xi_n^2 - 1}{(\xi_n - i)^5 (\xi_n + i)^4} + \frac{4(5\xi_n^2 - 1) + 12i(\xi_n^3 - \xi_n)}{(\xi_n - i)^6 (\xi_n + i)^4} \right] \right. \\
&\quad \left. + X_n Y_n h'(0) \left[ \frac{(4i - 4)(\xi^2 - 1) + 48(\xi_n^3 - \xi_n)}{(\xi_n - i)^5 (\xi_n + i)^4} - \frac{4(5\xi_n^2 - 1) + 12i(\xi_n^3 - \xi_n)}{(\xi_n - i)^6 (\xi_n + i)^4} \right] \right\} d\xi_n \sigma(\xi') dx' \\
&= \sum_{j,l=1}^{n-1} X_j Y_l h'(0) \Omega_3 \int_{\Gamma^+} \left[ 8i \frac{5\xi_n^2 - 1}{(\xi_n - i)^5 (\xi_n + i)^4} + \frac{4(5\xi_n^2 - 1) + 12i(\xi_n^3 - \xi_n)}{(\xi_n - i)^6 (\xi_n + i)^4} \right] \xi_j \xi_l d\xi_n dx' \\
&\quad + X_n Y_n h'(0) \Omega_3 \int_{\Gamma^+} \left[ \frac{(4i - 4)(\xi^2 - 1) + 48(\xi_n^3 - \xi_n)}{(\xi_n - i)^5 (\xi_n + i)^4} - \frac{4(5\xi_n^2 - 1) + 12i(\xi_n^3 - \xi_n)}{(\xi_n - i)^6 (\xi_n + i)^4} \right] d\xi_n dx' \\
&= \sum_{j=1}^{n-1} X_j Y_j \frac{4\pi}{3} h'(0) \Omega_3 \left\{ \frac{2\pi i}{4!} \left[ \frac{8i(5\xi_n^2 - 1)}{(\xi_n + i)^4} \right]^{(4)} \Big|_{\xi_n=i} dx' + \frac{2\pi i}{5!} \left[ \frac{4(5\xi_n^2 - 1)}{(\xi_n + i)^4} \right]^{(5)} \Big|_{\xi_n=i} dx' \right. \\
&\quad \left. + \frac{2\pi i}{5!} \left[ \frac{\xi_n^3 - \xi_n}{(\xi_n + i)^4} \right]^{(5)} \Big|_{\xi_n=i} dx' \right\} \\
&\quad - X_n Y_n h'(0) \Omega_3 \left\{ \frac{2\pi i}{4!} \left[ \frac{(4i - 4)(\xi_n^2 - 1)}{(\xi_n + i)^4} \right]^{(4)} \Big|_{\xi_n=i} dx' + \frac{2\pi i}{4!} \left[ \frac{48(\xi_n^3 - \xi_n)}{(\xi_n + i)^4} \right]^{(4)} \Big|_{\xi_n=i} dx' \right. \\
&\quad \left. - \frac{2\pi i}{5!} \left[ \frac{4(5\xi_n^2 - 1)}{(\xi_n + i)^4} \right]^{(5)} \Big|_{\xi_n=i} dx' - \frac{2\pi i}{5!} \left[ \frac{12i(\xi_n^3 - \xi_n)}{(\xi_n + i)^4} \right]^{(5)} \Big|_{\xi_n=i} dx' \right\} \\
&= - \left[ \frac{592}{3} \pi \sum_{j=1}^{n-1} X_j Y_j + \left( \frac{461}{4} + \frac{23}{4} i \right) X_n Y_n \right] h'(0) \pi \Omega_3 dx', \tag{3.12}
\end{aligned}$$

where  $\Omega_3$  is the canonical volume of  $S^3$ .

**case a) III)**  $r = 1$ ,  $l = -3$ ,  $j = |\alpha| = 0$ ,  $k = 1$ .

By (3.3), we get

$$\begin{aligned}
\Phi_3 &= -\frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_1(\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-1}) \times \partial_{\xi_n} \partial_{x_n} \sigma_{-3}(D^{-3})](x_0) d\xi_n \sigma(\xi') dx' \\
&= \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\partial_{\xi_n}^2 \pi_{\xi_n}^+ \sigma_1(\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-1}) \times \partial_{x_n} \sigma_{-3}(D^{-3})](x_0) d\xi_n \sigma(\xi') dx'. \tag{3.13}
\end{aligned}$$

By Lemma 2.2, we have

$$\partial_{x_n} \sigma_{-3}(D^{-3})(x_0)|_{|\xi'|=1} = \frac{\sqrt{-1} \partial_{x_n} [c(\xi')]}{(1 + \xi_n^2)^4} - \frac{2\sqrt{-1} h'(0) c(\xi) |\xi'|_{g^{\partial M}}^2}{(1 + \xi_n^2)^6}, \tag{3.14}$$

moreover,

$$\begin{aligned}
\pi_{\xi_n}^+ \sigma_1(\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-1}) &= -\frac{c(\xi') + ic(dx_n)}{2(\xi_n - i)} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l - \frac{c(\xi') + ic(dx_n)}{2(\xi_n - i)} X_n Y_n \\
&\quad - \frac{ic(\xi') - c(dx_n)}{2(\xi_n - i)} \sum_{j=1}^{n-1} X_j Y_n \xi_j - \frac{ic(\xi') - c(dx_n)}{2(\xi_n - i)} \sum_{l=1}^{n-1} X_n Y_l \xi_l. \tag{3.15}
\end{aligned}$$

Then, we have

$$\partial_{\xi_n}^2 \pi_{\xi_n}^+ \sigma_1(\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-1}) = -\frac{c(\xi') + ic(dx_n)}{(\xi_n - i)^3} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l - \frac{c(\xi') + ic(dx_n)}{(\xi_n - i)^3} X_n Y_n. \tag{3.16}$$

We note that  $i < n$ ,  $\int_{|\xi'|=1} \xi_{i_1} \xi_{i_2} \cdots \xi_{i_{2d+1}} \sigma(\xi') = 0$ , so we omit some items that have no contribution for computing **case a) III**)

$$\begin{aligned} & \text{tr}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_1(\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-1}) \times \partial_{\xi_n} \partial_{x_n} \sigma_{-3}(D^{-3})](x_0) \\ &= -2 \frac{h'(0)}{(\xi_n - i)^5 (\xi_n + i)^2} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l - 2 \frac{h'(0)}{(\xi_n - i)^5 (\xi_n + i)^2} X_n Y_n. \end{aligned} \quad (3.17)$$

Therefore, we get

$$\begin{aligned} \Phi_3 &= \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \left( -2 \frac{h'(0)}{(\xi_n - i)^5 (\xi_n + i)^2} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l - 2 \frac{h'(0)}{(\xi_n - i)^5 (\xi_n + i)^2} X_n Y_n \right) d\xi_n \sigma(\xi') dx' \\ &= -\sum_{j,l=1}^{n-1} X_j Y_l h'(0) \Omega_3 \int_{\Gamma^+} \frac{1}{(\xi_n - i)^5 (\xi_n + i)^2} \xi_j \xi_l d\xi_n dx' - X_n Y_n h'(0) \Omega_3 \int_{\Gamma^+} \frac{1}{(\xi_n - i)^5 (\xi_n + i)^2} d\xi_n dx' \\ &= -\sum_{j,l=1}^{n-1} X_j Y_l h'(0) \Omega_3 \frac{2\pi i}{4!} \left[ \frac{1}{(\xi_n + i)^2} \right] \Big|_{\xi_n=i}^{dx'} + 2X_n Y_n h'(0) \Omega_3 \frac{2\pi i}{4!} \left[ \frac{1}{(\xi_n + i)^2} \right] \Big|_{\xi_n=i}^{dx'} \\ &= \left( \frac{5\pi i}{6} \sum_{j=1}^{n-1} X_j Y_j + \frac{5i}{8} X_n Y_n \right) h'(0) \pi \Omega_3 dx'. \end{aligned} \quad (3.18)$$

**case b)**  $r = 0$ ,  $l = -3$ ,  $k = j = |\alpha| = 0$ .

By (3.3), we get

$$\begin{aligned} \Phi_4 &= -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\pi_{\xi_n}^+ \sigma_0(\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-1}) \times \partial_{\xi_n} \sigma_{-3}(D^{-3})](x_0) d\xi_n \sigma(\xi') dx' \\ &= i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_0(\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-1}) \times \sigma_{-3}(D^{-3})](x_0) d\xi_n \sigma(\xi') dx'. \end{aligned} \quad (3.19)$$

By Lemma 2.2, we have

$$\partial_{\xi_n} \sigma_{-3}(D^{-3})(x_0)|_{|\xi'|=1} = \frac{ic(dx_n)}{(1 + \xi_n^2)^2} - \frac{4\sqrt{-1}\xi_n c(\xi)}{(1 + \xi_n^2)^3}. \quad (3.20)$$

By Lemma 2.1, we have

$$\begin{aligned} \sigma_0(\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-1}) &= -\sum_{j,l=1}^n X_j Y_l \xi_j \xi_l \left[ \frac{c(\xi) \sigma_0(D) c(\xi)}{|\xi|^4} + \frac{c(\xi)}{|\xi|^6} \sum_j c(dx_j) (\partial_{x_j} [c(\xi)] |\xi|^2 - c(\xi) \partial_{x_j} (|\xi|^2)) \right] \\ &\quad - i \sum_{j,l=1}^n X_j \frac{\partial_{Y_l}}{\partial x_j} \xi_j \frac{c(\xi)}{|\xi|^2} - \sum_j A(Y) X_j \xi_j \frac{c(\xi)}{|\xi|^2} - \sum_l A(Y) Y_l \xi_l \frac{c(\xi)}{|\xi|^2}, \end{aligned} \quad (3.21)$$

moreover,

$$\begin{aligned} \partial_{\xi_n} \pi_{\xi_n}^+ \sigma_0(\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-1}) &= -\sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l [B_1 - B_2] \\ &\quad - X_n Y_n \left\{ \sigma_0 \left[ \frac{[c(\xi') + ic(dx_n)] c(dx_n)}{4(\xi_n - i)} + \frac{[c(\xi') + ic(dx_n)]^2}{4(\xi_n - i)^2} \right] \right. \\ &\quad \left. + c(dx_n) \partial_{x_n} [c(\xi')] \left[ \frac{i[c(\xi') + ic(dx_n)] c(dx_n)}{-4(\xi_n - i)} + \frac{[c(\xi') + ic(dx_n)]}{4(\xi_n - i)^2} \right] \right. \\ &\quad \left. + h'(0) |\xi'|^2 \left[ \frac{i[c(\xi') + ic(dx_n)]^2 - 2c(dx_n) [3c(\xi') + 4ic(dx_n)]}{16(\xi_n - i)} \right] \right\} \end{aligned}$$

$$\begin{aligned}
& + \left. \left\{ \frac{[c(\xi') + ic(dx_n)][c(\xi') + 5ic(dx_n)]}{16(\xi_n - i)^2} - \frac{i[c(\xi') + ic(dx_n)]^2}{8(\xi_n - i)^3} \right\} \right. \\
& - \Sigma_{j=1}^{n-1} X_j Y_n \xi_j \left\{ \sigma_0 \left[ \frac{[-ic(\xi') + ic(dx_n)]c(dx_n)}{2(\xi_n - i)} - \frac{i[c(\xi') + ic(dx_n)]^2}{4(\xi_n - i)^2} \right] \right. \\
& + c(dx_n) \partial_{x_n} [c(\xi')] \left[ \frac{-ic(dx_n)}{4(\xi_n - i)} - \frac{i[c(\xi') + ic(dx_n)]}{4(\xi_n - i)^2} \right] \\
& - c(dx_n) h'(0) |\xi'|^2 \left[ \frac{ic(dx_n)c(\xi')}{8(\xi_n - i)} + \frac{[c(\xi') + ic(dx_n)][c(\xi') + 3ic(dx_n)]}{8(\xi_n - i)^2} \right. \\
& \left. \left. - \frac{[c(\xi') + ic(dx_n)]^2}{8(\xi_n - i)^3} \right] \right\} \\
& - \Sigma_{l=1}^{n-1} X_n Y_l \xi_l \left\{ \sigma_0 \left[ \frac{[-ic(\xi') + ic(dx_n)]c(dx_n)}{2(\xi_n - i)} - \frac{i[c(\xi') + ic(dx_n)]^2}{4(\xi_n - i)^2} \right] \right. \\
& + c(dx_n) \partial_{x_n} [c(\xi')] \left[ \frac{-ic(dx_n)}{4(\xi_n - i)} - \frac{i[c(\xi') + ic(dx_n)]}{4(\xi_n - i)^2} \right] \\
& - c(dx_n) h'(0) |\xi'|^2 \left[ \frac{ic(dx_n)c(\xi')}{8(\xi_n - i)} + \frac{[c(\xi') + ic(dx_n)][c(\xi') + 3ic(dx_n)]}{8(\xi_n - i)^2} \right. \\
& \left. \left. - \frac{[c(\xi') + ic(dx_n)]^2}{8(\xi_n - i)^3} \right] \right\} \\
& - \Sigma_{j,l=1}^n X_j \frac{\partial Y_l}{\partial x_j} \xi_j \frac{c(\xi') + ic(dx_n)}{2(\xi_n - i)^2} - i \Sigma_j A(Y) X_j \xi_j \frac{c(\xi') + ic(dx_n)}{2(\xi_n - i)^2} \\
& - i \Sigma_l A(Y) Y_l \xi_l \frac{c(\xi') + ic(dx_n)}{2(\xi_n - i)^2}. \tag{3.22}
\end{aligned}$$

Similarly to [12], we have

$$B_1 = -\frac{A_1}{4(\xi_n - i)} - \frac{A_2}{4(\xi_n - i)^2} \tag{3.23}$$

where

$$\begin{aligned}
A_1 &= ic(\xi') p_0 c(\xi') + ic(dx_n) p_0 c(dx_n) + ic(\xi') c(dx_n) \partial_{x_n} [c(\xi')]; \\
A_2 &= [c(\xi') + ic(dx_n)] p_0 [c(\xi') + ic(dx_n)] + c(\xi') c(dx_n) \partial_{x_n} [c(\xi')] - i \partial_{x_n} [c(\xi')]; \tag{3.24}
\end{aligned}$$

$$B_2 = \frac{h'(0)}{2} \left[ \frac{c(dx_n)}{4i(\xi_n - i)} + \frac{c(dx_n) - ic(\xi')}{8(\xi_n - i)^2} + \frac{3\xi_n - 7i}{8(\xi_n - i)^3} [ic(\xi') - c(dx_n)] \right]. \tag{3.25}$$

We note that  $i < n$ ,  $\int_{|\xi'|=1} \xi_{i_1} \xi_{i_2} \cdots \xi_{i_{2d+1}} \sigma(\xi') = 0$ , so we omit some items that have no contribution for computing **case b**).

Then, we have

$$\begin{aligned}
& \text{tr}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_0 (\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-1}) \times \sigma_{-3}(D^{-3})](x_0) \\
&= \left[ \frac{-2ic_0}{(1 + \xi_n^2)^2} + h'(0) \frac{\xi_n^2 - i\xi_n - 2}{2(\xi_n - i)(1 + \xi_n^2)^2} - \frac{\sqrt{-1}}{2} h'(0) \frac{-i\xi_n^2 - \xi_n + 4i}{4(\xi_n - i)^3(\xi_n + i)^2} \text{tr}[id] \right] \Sigma_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l \\
&+ \left[ \frac{4 - 4i\xi_n - 3(\xi_n^2 + 1)}{(\xi_n - i)^4(\xi_n + i)^3} + \frac{4i + 4\xi_n - 3i(\xi_n^2 + 1)}{2(\xi_n - i)^5(\xi_n + i)^3} \right] h'(0) X_n Y_n. \tag{3.26}
\end{aligned}$$

Therefore, we get

$$\Phi_4 = i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \left[ \frac{-2ic_0}{(1 + \xi_n^2)^2} + h'(0) \frac{\xi_n^2 - i\xi_n - 2}{2(\xi_n - i)(1 + \xi_n^2)^2} \right]$$



$$\begin{aligned}
& - \frac{\sqrt{-1}}{2} h'(0) \frac{-i\xi_n^2 - \xi_n + 4i}{(\xi_n - i)^3(\xi_n + i)^2} \Big] \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l \\
& + \left[ \frac{4 - 4i\xi_n - 3(\xi_n^2 + 1)}{(\xi_n - i)^4(\xi_n + i)^3} + \frac{4i + 4\xi_n - 3i(\xi_n^2 + 1)}{2(\xi_n - i)^5(\xi_n + i)^3} \right] h'(0) X_n Y_n d\xi_n \sigma(\xi') dx' \\
& = \sum_{j,l=1}^{n-1} X_j Y_l h'(0) \Omega_3 \int_{\Gamma^+} \left[ \frac{-2ic_0}{(1 + \xi_n^2)^2} + h'(0) \frac{\xi_n^2 - i\xi_n - 2}{2(\xi_n - i)(1 + \xi_n^2)^2} \right. \\
& - \left. \frac{\sqrt{-1}}{2} h'(0) \frac{-i\xi_n^2 - \xi_n + 4i}{(\xi_n - i)^3(\xi_n + i)^2} \right] \xi_j \xi_l d\xi_n dx' \\
& + X_n Y_n h'(0) \Omega_3 \int_{\Gamma^+} \left[ \frac{4 - 4i\xi_n - 3(\xi_n^2 + 1)}{(\xi_n - i)^4(\xi_n + i)^3} + \frac{4i + 4\xi_n - 3i(\xi_n^2 + 1)}{2(\xi_n - i)^5(\xi_n + i)^3} \right] d\xi_n dx' \\
& = \frac{55}{3} \sum_{j,l=1}^{n-1} X_j Y_l h'(0) \pi^2 \Omega_3 dx' + X_n Y_n h'(0) \Omega_3 i \left\{ \frac{2\pi i}{3!} \left[ \frac{1}{(\xi_n + i)^3} \right]^{(3)} \Big|_{\xi_n=i} dx' \right. \\
& - \frac{4i \times 2\pi i}{3!} \left[ \frac{\xi_n}{(\xi_n + i)^3} \right]^{(3)} \Big|_{\xi_n=i} dx' - \frac{3 \times 2\pi i}{2!} \left[ \frac{1}{(\xi_n + i)^2} \right]^{(2)} \Big|_{\xi_n=i} dx' \\
& + \frac{4i \times 2\pi i}{2 \times 4!} \left[ \frac{1}{(\xi_n + i)^3} \right]^{(4)} \Big|_{\xi_n=i} dx' + \frac{4 \times 2\pi i}{2 \times 4!} \left[ \frac{\xi_n}{(\xi_n + i)^3} \right]^{(4)} \Big|_{\xi_n=i} dx' \\
& \left. - \frac{3i \times 2\pi i}{2 \times 3!} \left[ \frac{1}{(\xi_n + i)^2} \right]^{(3)} \Big|_{\xi_n=i} dx' \right\} \\
& = \left( \frac{55\pi}{3} \sum_{j=1}^{n-1} X_j Y_j - \frac{3}{8} X_n Y_n \right) h'(0) \pi \Omega_3 dx'. \tag{3.27}
\end{aligned}$$

**case c)**  $r = 1$ ,  $\ell = -4$ ,  $k = j = |\alpha| = 0$ .

By (3.3), we get

$$\begin{aligned}
\Phi_5 & = - \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\pi_{\xi_n}^+ \sigma_1(\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-1}) \times \partial_{\xi_n} \sigma_{-4}(D^{-3})](x_0) d\xi_n \sigma(\xi') dx' \\
& = \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_1(\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-1}) \times \sigma_{-4}(D^{-3})](x_0) d\xi_n \sigma(\xi') dx'. \tag{3.28}
\end{aligned}$$

By Lemma 2.1 and 2.2, we have

$$\begin{aligned}
\sigma_{-4}(D^{-3}) & = \frac{1}{(\xi_n^2 + 1)^4} \left[ \left( \frac{11}{2} \xi_n (1 + \xi_n^2) + 8i\xi_n \right) h'(0) c(\xi') \right. \\
& + \left[ -2i + 6i\xi_n^2 - \frac{7}{4}(1 + \xi_n^2) + \frac{15}{4}\xi_n^2(1 + \xi_n^2) \right] h'(0) c(dx_n) \\
& \left. - 3i\xi_n(1 + \xi_n^2) \partial_{x_n} c(\xi') + i(1 + \xi_n^2) c(\xi') c(dx_n) \partial_{x_n} c(\xi') \right], \tag{3.29}
\end{aligned}$$

and

$$\begin{aligned}
\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_1(\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-1}) & = \frac{c(\xi') + ic(dx_n)}{2(\xi_n - i)^2} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l - \frac{c(\xi') + ic(dx_n)}{2(\xi_n - i)^2} X_n Y_n \\
& + \frac{ic(\xi') - c(dx_n)}{2(\xi_n - i)^2} \sum_{j=1}^n X_j Y_n \xi_j + \frac{ic(\xi') - c(dx_n)}{2(\xi_n - i)^2} \sum_{l=1}^n X_n Y_l \xi_l. \tag{3.30}
\end{aligned}$$

We note that  $i < n$ ,  $\int_{|\xi'|=1} \xi_{i_1} \xi_{i_2} \cdots \xi_{i_{2d+1}} \sigma(\xi') = 0$ , so we omit some items that have no contribution for computing **case c)**.

We have

$$\text{tr}[c(\xi') c(\xi') c(dx_n) \partial_{x_n} c(\xi')] = 0;$$

$$\text{tr}[c(dx_n)c(\xi')c(dx_n)\partial_{x_n}c(\xi')] = -2h'(0) \quad (3.31)$$

Then we get

$$\begin{aligned} & \text{tr}[\partial_{\xi_n}\pi_{\xi_n}^+\sigma_{-1}(\nabla_X^{S(TM)}\nabla_Y^{S(TM)}D^{-1})\times\sigma_{-4}(D^{-3})](x_0)|_{|\xi'|=1} \\ &= \frac{h'(0)(7+6i-(20-15i)\xi_n-(7-6i)\xi_n^2+15i\xi_n^3)}{(\xi_n-i)^5(\xi_n+i)^4}\sum_{j,l=1}^{n-1}X_jY_l\xi_j\xi_l \\ &+ \frac{(3i-11)\xi_n(1-\xi_n^2)-16i\xi_n+(13+\frac{7}{2}i)(1+\xi_n^2)-16-\frac{15}{2}\xi_n^2(1+\xi_n^2)}{(\xi_n-i)^2(\xi_n+i)^4}X_nY_n. \end{aligned} \quad (3.32)$$

So we have

$$\begin{aligned} \Phi_5 &= \int_{|\xi'|=1}\int_{-\infty}^{+\infty}\text{tr}[\partial_{\xi_n}\pi_{\xi_n}^+\sigma_1(\nabla_X^{S(TM)}\nabla_Y^{S(TM)}D^{-1})\times\sigma_{-4}(D^{-3})](x_0)d\xi_n\sigma(\xi')dx' \\ &= \int_{|\xi'|=1}\int_{-\infty}^{+\infty}\frac{h'(0)(7+6i-(20-15i)\xi_n-(7-6i)\xi_n^2+15i\xi_n^3)}{(\xi_n-i)^5(\xi_n+i)^4}\sum_{j,l=1}^{n-1}X_jY_l\xi_j\xi_l \\ &+ \frac{(3i-11)\xi_n(1-\xi_n^2)-16i\xi_n+(13+\frac{7}{2}i)(1+\xi_n^2)-16-\frac{15}{2}\xi_n^2(1+\xi_n^2)}{(\xi_n-i)^2(1+\xi_n^2)^4}X_nY_nd\xi_n\sigma(\xi')dx' \\ &= \left(-\frac{35}{3}+\frac{50}{3}i\right)\sum_{j=1}^{n-1}X_jY_jh'(0)\pi^2\Omega_3dx' \\ &+ h'(0)\Omega_3\int_{\Gamma^+}\frac{(3i-11)\xi_n(1-\xi_n^2)-16i\xi_n+(13+\frac{7}{2}i)(1+\xi_n^2)-16-\frac{15}{2}\xi_n^2(1+\xi_n^2)}{(\xi_n-i)^6(\xi_n+i)^4}\sum_{j,l=1}^{n-1}X_jY_l\xi_j\xi_ld\xi_ndx' \\ &= \left(-\frac{35}{3}+\frac{50}{3}i\right)\sum_{j=1}^{n-1}X_jY_jh'(0)\pi^2\Omega_3dx' \\ &+ h'(0)\Omega_3\frac{2\pi i}{5!}X_nY_n\left[\frac{(3i-11)\xi_n(1-\xi_n^2)-16i\xi_n+(13+\frac{7}{2}i)(1+\xi_n^2)-16-\frac{15}{2}\xi_n^2(1+\xi_n^2)}{(\xi_n+i)^4}\right]^{(5)}\Bigg|_{\xi_n=i}dx' \\ &= \left[\left(-\frac{35}{3}+\frac{50}{3}i\right)\sum_{j=1}^{n-1}X_jY_j\pi+\left(5-\frac{137}{32}i\right)X_nY_n\right]h'(0)\pi^2\Omega_3dx'. \end{aligned} \quad (3.33)$$

Let  $X = X^T + X_n\partial_n$ ,  $Y = Y^T + Y_n\partial_n$ , then we have  $\sum_{j=1}^{n-1}X_jY_j = g(X^T, Y^T)$ . Now  $\Phi$  is the sum of the cases (a), (b) and (c). Therefore, we get

$$\Phi = \sum_{i=1}^5\Phi_i = \left[\left(-\frac{1411}{12}+\frac{27i}{32}\right)X_nY_n+\left(-\frac{572}{3}+\frac{35i}{2}\right)\pi g(X^T, Y^T)\right]h'(0)\pi\Omega_3dx'. \quad (3.34)$$

Therefore, by (3.3)-(3.34), we obtain following theorem

**Theorem 3.1.** *Let  $M$  be a 4-dimensional oriented compact spin manifold with boundary  $\partial M$  and the metric  $g^M$  be defined as (3.1), then we get the following equality:*

$$\begin{aligned} & \widetilde{\text{Wres}}[\pi^+(\nabla_X^{S(TM)}\nabla_Y^{S(TM)}D^{-2})\circ\pi^+(D^{-2})] \\ &= \frac{4\pi^2}{3}\int_M[\text{Ric}(X, Y)-\frac{1}{2}sg(X, Y)]d\text{Vol}_M+\pi^2\int_Msg(X, Y)d\text{Vol}_M \\ &+ \int_{\partial M}\left[\left(-\frac{1411}{12}+\frac{27i}{32}\right)X_nY_n+\left(-\frac{572}{3}+\frac{35i}{2}\right)\pi g(X^T, Y^T)\right]h'(0)\pi\Omega_3d\text{Vol}_M. \end{aligned} \quad (3.35)$$

#### 4. The noncommutative residue for 3-dimensional manifolds with boundary

In this section, we compute the noncommutative residue  $\widetilde{\text{Wres}}[\pi^+(\nabla_X^{S(TM)}\nabla_Y^{S(TM)}D^{-1})\circ\pi^+(D^{-3})]$  on 3-dimensional oriented compact manifolds with boundary. As in [10], for an odd-dimensional manifolds with boundary, we have the formula

$$\widetilde{\text{Wres}}[\pi^+(\nabla_X^{S(TM)}\nabla_Y^{S(TM)}D^{-1})\circ\pi^+(D^{-3})] = \int_{\partial M} \Phi \quad (4.1)$$

Let  $M$  be a 3-dimensional compact oriented manifold with boundary  $\partial M$ . We have

$$\begin{aligned} \text{tr}[c^2(\xi')] &= -2; \text{tr}[c(\xi')c(dx_n)] = 0; \\ \text{tr}[c(dx_n)c(dx_n)] &= -2; \text{tr}[c(\xi') + ic(dx_n)][c(\xi') + ic(dx_n)] = -2 - 2i\xi_n. \end{aligned} \quad (4.2)$$

By (3.3), satisfies  $r + l - k - |\alpha| - j - 1 = -3$ , so we get  $r = 1$ ,  $\ell = -3$ ,  $k = j = |\alpha| = 0$ , then

$$\begin{aligned} \Phi &= - \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\pi_{\xi_n}^+ \sigma_1(\nabla_X^{S(TM)}\nabla_Y^{S(TM)}D^{-1}) \times \partial_{\xi_n} \sigma_{-3}(D^{-3})](x_0) d\xi_n \sigma(\xi') dx' \\ &= \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_1(\nabla_X^{S(TM)}\nabla_Y^{S(TM)}D^{-1}) \times \sigma_{-3}(D^{-3})](x_0) d\xi_n \sigma(\xi') dx'. \end{aligned} \quad (4.3)$$

We note that  $i < n$ ,  $\int_{|\xi'|=1} \xi_{i_1} \xi_{i_2} \cdots \xi_{i_{2d+1}} \sigma(\xi') = 0$ , so we omit some items that have no contribution for computing. By (3.30) and (2.8), we have

$$\begin{aligned} &\text{tr}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_1(\nabla_X^{S(TM)}\nabla_Y^{S(TM)}D^{-1}) \times \sigma_{-3}(D^{-3})] \\ &= - \frac{i - \xi_n}{(\xi_n - i)^4 (\xi_n + i)^2} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l - \frac{\sqrt{-1} \xi_n^2 (2 + 2i\xi_n)}{2(\xi_n - i)^3 (\xi_n + i)^2} X_n Y_n. \end{aligned} \quad (4.4)$$

Then there is the following formula

$$\begin{aligned} \Phi &= - \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\pi_{\xi_n}^+ \sigma_1(\nabla_X^{S(TM)}\nabla_Y^{S(TM)}D^{-1}) \times \partial_{\xi_n} \sigma_{-3}(D^{-3})](x_0) d\xi_n \sigma(\xi') dx' \\ &= \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \frac{i - \xi_n}{(\xi_n - i)^4 (\xi_n + i)^2} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l + \frac{\sqrt{-1} \xi_n^2 (2 + 2i\xi_n)}{2(\xi_n - i)^3 (\xi_n + i)^2} X_n Y_n(x_0) d\xi_n \sigma(\xi') dx' \\ &= \Omega_2 \int_{\Gamma^+} - \frac{i - \xi_n}{(\xi_n - i)^4 (\xi_n + i)^2} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l + \frac{\sqrt{-1} \xi_n^2 (2 + 2i\xi_n)}{2(\xi_n - i)^3 (\xi_n + i)^2} X_n Y_n(x_0) d\xi_n dx' \\ &= -\Omega_2 \frac{2\pi i}{3!} \frac{4\pi}{3} \sum_{j,l=1}^{n-1} X_j Y_l \left[ \frac{i - \xi_n}{(\xi_n + i)^2} \right]^{(3)} \Big|_{(\xi_n=i)} dx' + \Omega_2 \frac{2\pi i}{3!} X_n Y_n \left[ \frac{\xi_n - i}{(\xi_n + i)^2} \right]^{(3)} \Big|_{(\xi_n=i)} dx' \\ &= \left( \frac{i}{3} \sum_{j=1}^{n-1} X_j Y_j \pi + \frac{3i}{4} X_n Y_n \right) \pi \Omega_2 dx'. \end{aligned} \quad (4.5)$$

Therefore, we get the following theorem

**Theorem 4.1.** *Let  $M$  be a 3-dimensional oriented compact spin manifold with boundary  $\partial M$  and the metric  $g^M$  be defined as Sect.3, then we get the following equality:*

$$\widetilde{\text{Wres}}[\pi^+(\nabla_X^{S(TM)}\nabla_Y^{S(TM)}D^{-1})\circ\pi^+(D^{-3})] = \left( \frac{i}{3} \sum_{j=1}^{n-1} X_j Y_j \pi + \frac{3i}{4} X_n Y_n \right) \pi \Omega_2 dx'. \quad (4.6)$$

## 5. The noncommutative residue for 6-dimensional manifolds with boundary

In this section, we compute the noncommutative residue  $\widetilde{\text{Wres}}[\pi^+(\nabla_X^{S(TM)}\nabla_Y^{S(TM)}D^{-3})\circ\pi^+(D^{-3})]$  on 6-dimensional oriented compact manifolds with boundary. For more details, see Section 2 in [12].

Let  $M$  be a 6-dimensional compact oriented manifold with boundary  $\partial M$  and the metric as same as Section 3. We can compute the noncommutative residue

$$\begin{aligned} & \widetilde{\text{Wres}}[\pi^+(\nabla_X^{S(TM)}\nabla_Y^{S(TM)}D^{-3})\circ\pi^+(D^{-3})] \\ &= \int_M \int_{|\xi|=1} \text{trace}_{\wedge^*T^*M} \otimes_{\mathbb{C}} [\sigma_{-6}(\nabla_X^{S(TM)}\nabla_Y^{S(TM)}D^{-3})\circ D^{-3}] \sigma(\xi) dx + \int_{\partial M} \Phi, \end{aligned} \quad (5.1)$$

where

$$\begin{aligned} \Phi &= \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \sum_{j,k=0}^{\infty} \sum \frac{(-i)^{|\alpha|+j+k+1}}{\alpha!(j+k+1)!} \times \text{trace}_{\wedge^*T^*M} \otimes_{\mathbb{C}} [\partial_{x_n}^j \partial_{\xi_n}^\alpha \sigma_r^+(\nabla_X^{S(TM)}\nabla_Y^{S(TM)}D^{-3})(x', 0, \xi', \xi_n) \\ & \times \partial_{x'}^\alpha \partial_{\xi_n}^{j+1} \partial_{x_n}^k \sigma_l(D^{-3})(x', 0, \xi', \xi_n)] d\xi_n \sigma(\xi') dx', \end{aligned} \quad (5.2)$$

and the sum is taken over  $r+l-k-j-|\alpha|=-5$ ,  $r \leq -1$ ,  $l \leq -3$ .

By Theorem 2.3, we can compute the interior of  $\widetilde{\text{Wres}}[\pi^+(\nabla_X^{S(TM)}\nabla_Y^{S(TM)}D^{-3})\circ\pi^+(D^{-3})]$ , we get

$$\text{Wres}[\nabla_X^{S(TM)}\nabla_Y^{S(TM)}D^{-6}] = \frac{4\pi^3}{3} \int_M [\text{Ric}(X, Y) - \frac{1}{2}sg(X, Y)] d\text{Vol}_M + \pi^3 \int_M sg(X, Y) d\text{Vol}_M. \quad (5.3)$$

Now we need to compute  $\int_{\partial M} \Phi$ . When  $n=6$ , then  $\text{tr}_{S(TM)}[\text{id}] = \dim(\wedge^*(\mathbb{R}^3)) = 8$ , the sum is taken over  $r+l-k-j-|\alpha|=-6$ ,  $r \leq -1$ ,  $l \leq -3$ , then we have the following five cases:

**case a) I)**  $r=-1$ ,  $l=-3$ ,  $k=j=0$ ,  $|\alpha|=1$ .

By (5.2), we get

$$\Phi_1 = - \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \sum_{|\alpha|=1} \text{tr}[\partial_{\xi'}^\alpha \pi_{\xi_n}^+ \sigma_{-1}(\nabla_X^{S(TM)}\nabla_Y^{S(TM)}D^{-3}) \times \partial_{x'}^\alpha \partial_{\xi_n} \sigma_{-3}(D^{-3})](x_0) d\xi_n \sigma(\xi') dx'. \quad (5.4)$$

By Lemma 2.2 in [12], for  $i < n$ , then

$$\partial_{x_i} \sigma_{-3}(D^{-3})(x_0) = \partial_{x_i} (\sqrt{-1}c(\xi)|\xi|^{-4})(x_0) = \sqrt{-1} \frac{\partial_{x_i} c(\xi)}{|\xi|^4} (x_0) + \sqrt{-1} \frac{c(\xi) \partial_{x_i} (|\xi|^4)}{|\xi|^8} (x_0) = 0, \quad (5.5)$$

so  $\Phi_1 = 0$ .

**case a) II)**  $r=-1$ ,  $l=-3$ ,  $k=|\alpha|=0$ ,  $j=1$ .

By (5.2), we get

$$\Phi_2 = -\frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\partial_{x_n} \pi_{\xi_n}^+ \sigma_{-1}(\nabla_X^{S(TM)}\nabla_Y^{S(TM)}D^{-3}) \times \partial_{\xi_n}^2 \sigma_{-3}(D^{-3})](x_0) d\xi_n \sigma(\xi') dx'. \quad (5.6)$$

By Lemma 2.2, we have

$$\partial_{\xi_n}^2 \sigma_{-3}((D^{-3}))(x_0) = \partial_{\xi_n}^2 (c(\xi)|\xi|^{-4})(x_0) = \sqrt{-1} \frac{(20\xi_n^2 - 4)c(\xi') + 12(\xi^3 - \xi)c(dx_n)}{(1 + \xi_n^2)^4}. \quad (5.7)$$

$$\partial_{x_n} \sigma_{-1} (\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-3}) = \sum_{j,l=1}^n X_j Y_l \xi_j \xi_l \left[ \frac{\partial_{x_n} c(\xi')^2}{1 + \xi_n^2} - \frac{2c(\xi)h'(0)|\xi'|^2}{(1 + \xi_n^2)^3} \right]. \quad (5.8)$$

Then, we have

$$\begin{aligned} \pi_{\xi_n}^+ \partial_{x_n} \sigma_{-1} (\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-3}) &= \partial_{x_n} \pi_{\xi_n}^+ \sigma_{-1} (\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-3}) \\ &= \sqrt{-1} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l h'(0) |\xi'|^2 \left[ \frac{-3ic(\xi')}{4(\xi_n - i)} - \frac{3c(\xi') + ic(dx_n)}{4(\xi_n - i)^2} \right. \\ &\quad \left. + \frac{c(\xi') + ic(dx_n)}{(\xi_n - i)^3} \right] - \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l \frac{\partial_{x_n} c(\xi') [i(\xi_n - i) + 1]}{4(\xi_n - i)^2} \\ &\quad - \sqrt{-1} X_n Y_n \left\{ \frac{-3i \partial_{x_n} c(\xi')}{4(\xi_n - i)} + \frac{\partial_{x_n} c(\xi')}{4(\xi_n - i)^2} + \frac{c(\xi') + ic(dx_n)}{2i(\xi_n - i)^3} \right. \\ &\quad \left. + h'(0) |\xi'| \left[ -\frac{-5c(\xi') + 9ic(dx_n)}{4(\xi_n - i)} + \frac{3c(\xi') + 3ic(dx_n)}{4(\xi_n - i)^2} \right] \right\} \\ &\quad - \sum_{j=1}^{n-1} X_j Y_n \xi_j \left\{ -\frac{i \partial_{x_n} c(\xi')}{4(\xi_n - i)^2} + h'(0) |\xi'| \left[ \frac{11ic(dx_n)}{8(\xi_n - i)} \right. \right. \\ &\quad \left. \left. - \frac{ic(\xi') + (1 + 4i)c(dx_n)}{16(\xi_n - i)^2} - \frac{ic(\xi') - c(dx_n)}{2(\xi_n - i)^3} \right] \right\} \\ &\quad - \sum_{l=1}^{n-1} X_n Y_l \xi_l \left\{ -\frac{i \partial_{x_n} c(\xi')}{4(\xi_n - i)^2} + h'(0) |\xi'| \left[ \frac{11ic(dx_n)}{8(\xi_n - i)} \right. \right. \\ &\quad \left. \left. - \frac{ic(\xi') + (1 + 4i)c(dx_n)}{16(\xi_n - i)^2} - \frac{ic(\xi') - c(dx_n)}{2(\xi_n - i)^3} \right] \right\}. \end{aligned} \quad (5.9)$$

We note that  $i < n$ ,  $\int_{|\xi'|=1} \xi_{i_1} \xi_{i_2} \cdots \xi_{i_{2d+1}} \sigma(\xi') = 0$ , so we omit some items that have no contribution for computing **case a) II**.

Then there is the following formula

$$\begin{aligned} &\text{tr}[\partial_{x_n} \pi_{\xi_n}^+ \sigma_{-1} (\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-3}) \times \partial_{\xi_n}^2 \sigma_{-3}(D^{-3})](x_0) \\ &= -\sqrt{-1} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l h'(0) \left[ \frac{24i(5\xi_n^2 - 1)}{(\xi_n - i)^5(\xi_n + i)^4} + \frac{[2i(\xi_n - i) + 26](5\xi_n^2 - 1) + 24i(\xi_n^3 - \xi_n)}{(\xi_n - i)^6(\xi_n + i)^4} \right. \\ &\quad \left. - \frac{32(5\xi_n^2 - 1) + 96(\xi_n^3 - \xi_n)}{(\xi_n - i)^7(\xi_n + i)^4} \right] \\ &\quad + \sqrt{-1} X_n Y_n h'(0) \left[ \frac{46i(5\xi_n^2 - 1) - 216(\xi_n^3 - \xi_n)}{(\xi_n - i)^5(\xi_n + i)^4} - \frac{26(5\xi_n^2 - 1) + 72i(\xi_n^3 - \xi_n)}{(\xi_n - i)^6(\xi_n + i)^4} \right. \\ &\quad \left. - \frac{16(5\xi_n^2 - 1) + 48i(\xi_n^3 - \xi_n)}{i(\xi_n - i)^7(\xi_n + i)^4} \right]. \end{aligned} \quad (5.10)$$

Therefore, there is the following formula

$$\begin{aligned} \Phi_2 &= \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \left\{ -i \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l h'(0) \left[ \frac{24i(5\xi_n^2 - 1)}{(\xi_n - i)^5(\xi_n + i)^4} \right. \right. \\ &\quad \left. \left. + \frac{[2i(\xi_n - i) + 26](5\xi_n^2 - 1) + 24i(\xi_n^3 - \xi_n)}{(\xi_n - i)^6(\xi_n + i)^4} - \frac{32(5\xi_n^2 - 1) + 96(\xi_n^3 - \xi_n)}{(\xi_n - i)^7(\xi_n + i)^4} \right] \right\} \end{aligned}$$

$$\begin{aligned}
& + \sqrt{-1} X_n Y_n h'(0) \left[ \frac{46i(5\xi^2 - 1) - 216(\xi_n^3 - \xi_n)}{(\xi_n - i)^5(\xi_n + i)^4} - \frac{26(5\xi_n^2 - 1) + 72i(\xi_n^3 - \xi_n)}{(\xi_n - i)^6(\xi_n + i)^4} \right. \\
& \left. - \frac{16(5\xi_n^2 - 1) + 48i(\xi_n^3 - \xi_n)}{i(\xi_n - i)^7(\xi_n + i)^4} \right] d\xi_n \sigma(\xi') dx' \\
& = -\frac{1}{2} \sqrt{-1} \sum_{j,l=1}^{n-1} X_j Y_l h'(0) \Omega_4 \int_{\Gamma^+} \left[ \frac{24i(5\xi_n^2 - 1)}{(\xi_n - i)^5(\xi_n + i)^4} + \frac{[2i(\xi_n - i) + 26](5\xi_n^2 - 1) + 24i(\xi_n^3 - \xi_n)}{(\xi_n - i)^6(\xi_n + i)^4} \right. \\
& \left. - \frac{32(5\xi_n^2 - 1) + 96(\xi_n^3 - \xi_n)}{(\xi_n - i)^7(\xi_n + i)^4} \right] \xi_j \xi_l d\xi_n dx' \\
& + \frac{1}{2} \sqrt{-1} X_n Y_n h'(0) \Omega_4 \int_{\Gamma^+} \left[ \frac{46i(5\xi^2 - 1) - 216(\xi_n^3 - \xi_n)}{(\xi_n - i)^5(\xi_n + i)^4} - \frac{26(5\xi_n^2 - 1) + 72i(\xi_n^3 - \xi_n)}{(\xi_n - i)^6(\xi_n + i)^4} \right. \\
& \left. - \frac{16(5\xi_n^2 - 1) + 48i(\xi_n^3 - \xi_n)}{i(\xi_n - i)^7(\xi_n + i)^4} \right] d\xi_n dx' \\
& = -\frac{1}{2} \sqrt{-1} \sum_{j=1}^{n-1} X_j Y_j \frac{4\pi}{3} h'(0) \Omega_4 \left\{ \frac{2\pi i}{4!} \left[ \frac{24i(5\xi_n^2 - 1)}{(\xi_n + i)^4} \right]^{(4)} \Big|_{\xi_n=i} dx' \right. \\
& + \frac{2\pi i}{5!} \left[ \frac{[2i(\xi_n - i) + 26](5\xi_n^2 - 1) + 24i(\xi_n^3 - \xi_n)}{(\xi_n + i)^4} \right]^{(5)} \Big|_{\xi_n=i} dx' \\
& + \left. \frac{2\pi i}{6!} \left[ \frac{32(5\xi_n^2 - 1) + 96(\xi_n^3 - \xi_n)}{(\xi_n + i)^4} \right]^{(6)} \Big|_{\xi_n=i} dx' \right\} \\
& + \frac{1}{2} \sqrt{-1} X_n Y_n h'(0) \Omega_4 \left\{ \frac{2\pi i}{4!} \left[ \frac{46i(5\xi^2 - 1) - 216(\xi_n^3 - \xi_n)}{(\xi_n + i)^4} \right]^{(4)} \Big|_{\xi_n=i} dx' \right. \\
& + \frac{2\pi i}{5!} \left[ \frac{26(5\xi_n^2 - 1) + 72i(\xi_n^3 - \xi_n)}{(\xi_n + i)^4} \right]^{(5)} \Big|_{\xi_n=i} dx' - \frac{2\pi i}{6!} \left[ \frac{16(5\xi_n^2 - 1) + 48i(\xi_n^3 - \xi_n)}{(\xi_n + i)^4} \right]^{(6)} \Big|_{\xi_n=i} dx' \left. \right\} \\
& = \left[ \left( -\frac{5925}{16} - \frac{877}{6} i \right) \pi \sum_{j=1}^{n-1} X_j Y_j + (311i) X_n Y_n \right] h'(0) \pi \Omega_4 dx', \tag{5.11}
\end{aligned}$$

where  $\Omega_4$  is the canonical volume of  $S^4$ .

**case a) III)**  $r = -1$ ,  $l = -3$ ,  $j = |\alpha| = 0$ ,  $k = 1$ .

By (5.2), we get

$$\begin{aligned}
\Phi_3 & = -\frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-1}(\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-3}) \times \partial_{\xi_n} \partial_{x_n} \sigma_{-3}(D^{-3})](x_0) d\xi_n \sigma(\xi') dx' \\
& = \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\partial_{\xi_n}^2 \pi_{\xi_n}^+ \sigma_{-1}(\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-3}) \times \partial_{x_n} \sigma_{-3}(D^{-3})](x_0) d\xi_n \sigma(\xi') dx'. \tag{5.12}
\end{aligned}$$

By [8], we have

$$\begin{aligned}
\pi_{\xi_n}^+ \sigma_{-1}(\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-3}) & = \left[ \frac{c(\xi')}{4(\xi_n - i)} - \frac{ic(\xi') - c(dx_n)}{4(\xi_n - i)^2} \right] \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l \\
& + \left[ \frac{c(\xi') - 2ic(dx_n)}{4(\xi_n - i)} + \frac{ic(\xi') - c(dx_n)}{4(\xi_n - i)^2} \right] X_n Y_n \\
& + \left[ \frac{c(dx_n)}{4(\xi_n - i)} + \frac{c(\xi') + ic(dx_n)}{4(\xi_n - i)^2} \right] \sum_{j=1}^{n-1} X_j Y_n \xi_j \\
& + \left[ \frac{c(dx_n)}{4(\xi_n - i)} + \frac{c(\xi') + ic(dx_n)}{4(\xi_n - i)^2} \right] \sum_{l=1}^{n-1} X_n Y_l \xi_l. \tag{5.13}
\end{aligned}$$

Then,

$$\begin{aligned}
\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-1} (\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-3}) &= \left[ -\frac{ic(\xi')}{4(\xi_n - i)^2} + \frac{ic(\xi') - c(dx_n)}{2(\xi_n - i)^3} \right] \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l \\
&\quad - \left[ \frac{c(\xi') + 2ic(dx_n)}{4(\xi_n - i)^2} + \frac{ic(\xi') - c(dx_n)}{2(\xi_n - i)^3} \right] X_n Y_n \\
&\quad - \left[ \frac{c(dx_n)}{4(\xi_n - i)^2} + \frac{c(\xi') + ic(dx_n)}{2(\xi_n - i)^3} \right] \sum_{j=1}^{n-1} X_j Y_n \xi_j \\
&\quad - \left[ \frac{c(dx_n)}{4(\xi_n - i)^2} + \frac{c(\xi') + ic(dx_n)}{2(\xi_n - i)^3} \right] \sum_{l=1}^{n-1} X_n Y_l \xi_l. \tag{5.14}
\end{aligned}$$

By (3.14) and (5.14), we have

$$\begin{aligned}
&\text{tr}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-1} (\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-3}) \times \partial_{\xi_n} \partial_{x_n} \sigma_{-3}(D^{-3})](x_0) \\
&= \left[ \frac{24i\xi_n}{(\xi_n - i)^2(1 + \xi_n^2)^4} - \frac{4i\xi_n}{(\xi_n - i)^2(1 + \xi_n^2)^3} + \frac{48\xi_n - 8i(1 - 5\xi_n^2)}{(\xi_n - i)^3(1 + \xi_n^2)^4} - \frac{8\xi_n}{(\xi_n - i)^3(1 + \xi_n^2)^3} \right] h'(0) \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l \\
&+ \left[ \frac{24i\xi_n + 8(15\xi_n^2)}{(\xi_n - i)^2(1 + \xi_n^2)^4} - \frac{4i\xi_n}{(\xi_n - i)^2(1 + \xi_n^2)^3} - \frac{48\xi_n - 8i(1 - 5\xi_n^2)}{(\xi_n - i)^3(1 + \xi_n^2)^4} + \frac{8\xi_n}{(\xi_n - i)^3(1 + \xi_n^2)^3} \right] h'(0) X_n Y_n. \tag{5.15}
\end{aligned}$$

Therefore, we get the following formula

$$\begin{aligned}
\Phi_3 &= \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \left( \left[ \frac{24i\xi_n}{(\xi_n - i)^2(1 + \xi_n^2)^4} - \frac{4i\xi_n}{(\xi_n - i)^2(1 + \xi_n^2)^3} + \frac{48\xi_n - 8i(1 - 5\xi_n^2)}{(\xi_n - i)^3(1 + \xi_n^2)^4} \right. \right. \\
&\quad \left. \left. - \frac{8\xi_n}{(\xi_n - i)^3(1 + \xi_n^2)^3} \right] h'(0) \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l \right. \\
&\quad \left. + \left[ \frac{24i\xi_n + 8(15\xi_n^2)}{(\xi_n - i)^2(1 + \xi_n^2)^4} - \frac{4i\xi_n}{(\xi_n - i)^2(1 + \xi_n^2)^3} - \frac{48\xi_n - 8i(1 - 5\xi_n^2)}{(\xi_n - i)^3(1 + \xi_n^2)^4} \right. \right. \\
&\quad \left. \left. + \frac{8\xi_n}{(\xi_n - i)^3(1 + \xi_n^2)^3} \right] h'(0) X_n Y_n \right) d\xi_n \sigma(\xi') dx' \\
&= -\frac{1}{2} \sum_{j,l=1}^{n-1} X_j Y_l h'(0) \Omega_4 \left\{ \frac{2\pi i}{5!} \left[ \frac{24\xi_n}{(\xi_n + i)^4} \right]^{(5)} \Big|_{\xi_n=i} - \frac{2\pi i}{4!} \left[ \frac{4\xi_n}{(\xi_n + i)^3} \right]^{(4)} \Big|_{\xi_n=i} \right. \\
&\quad \left. + \frac{2\pi i}{6!} \left[ \frac{48\xi_n - 8i(1 - 5\xi_n^2)}{(\xi_n + i)^4} \right]^{(6)} \Big|_{\xi_n=i} - \frac{2\pi i}{5!} \left[ \frac{8\xi_n}{(\xi_n + i)^3} \right]^{(5)} \Big|_{\xi_n=i} \right\} dx' \\
&\quad - \frac{1}{2} X_n Y_n h'(0) \Omega_4 \left\{ \frac{2\pi i}{5!} \left[ \frac{24\xi_n + 8(1 - 5\xi_n^2)}{(\xi_n + i)^4} \right]^{(5)} \Big|_{\xi_n=i} - \frac{2\pi i}{4!} \left[ \frac{4i\xi_n}{(\xi_n + i)^3} \right]^{(4)} \Big|_{\xi_n=i} \right. \\
&\quad \left. - \frac{2\pi i}{6!} \left[ \frac{48\xi_n - 8i(1 - 5\xi_n^2)}{(\xi_n + i)^4} \right]^{(6)} \Big|_{\xi_n=i} + \frac{2\pi i}{5!} \left[ \frac{8xi_n}{(\xi_n + i)^3} \right]^{(5)} \Big|_{\xi_n=i} \right\} dx' \\
&= \left[ \left( \frac{49}{3} + 86i \right) \pi \sum_{j=1}^{n-1} X_j Y_j - \left( \frac{17}{4} - 68i \right) X_n Y_n \right] h'(0) \pi \Omega_3 dx'. \tag{5.16}
\end{aligned}$$

**case b)**  $r = -1$ ,  $l = -4$ ,  $k = j = |\alpha| = 0$ .

By (5.2), we get

$$\Phi_4 = -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\pi_{\xi_n}^+ \sigma_{-1} (\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-3}) \times \partial_{\xi_n} \sigma_{-4}(D^{-3})](x_0) d\xi_n \sigma(\xi') dx'$$

$$= i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-1}(\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-3}) \times \sigma_{-4}(D^{-3})](x_0) d\xi_n \sigma(\xi') dx'. \quad (5.17)$$

We note that  $i < n$ ,  $\int_{|\xi'|=1} \xi_{i_1} \xi_{i_2} \cdots \xi_{i_{2d+1}} \sigma(\xi') = 0$ , so we omit some items that have no contribution for computing **case b**).

By (3.29) and (5.14), we have

$$\begin{aligned} & \text{tr}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-1}(\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-3}) \times \sigma_{-3}(D^{-3})](x_0) \\ &= h'(0) \left[ \frac{(-11+3i)\xi_n(1+\xi_n^2) - 16i\xi_n}{(\xi_n-i)^2(1+\xi_n^2)^4} \right. \\ & \quad - \sqrt{-1} \frac{(-22+6i)\xi_n(1+\xi_n^2) - 32i\xi_n + (26-7i)(1+\xi_n^2) - 32 - 15i\xi_n^2(1+\xi_n^2)}{(\xi_n-i)^3(\xi_n+i)^4} \left. \right] \sum_{j,l=1}^{\Sigma_n-1} X_j Y_l \xi_j \xi_l \\ & \quad - \sqrt{-1} \left[ \frac{(-11-3i)\xi_n(1+\xi_n^2) + 16\xi_n + (48i-14)(1+\xi_n^2) - 32i + 30\xi_n^2(1+\xi_n^2)}{(\xi_n-i)^6(\xi_n+i)^4} \right. \\ & \quad \left. + \frac{(-22+6i)\xi_n(1+\xi_n^2) - 32i\xi_n + (26-7i)(1+\xi_n^2) - 32 - 15i\xi_n^2(1+\xi_n^2)}{(\xi_n-i)^3(\xi_n+i)^4} \right] h'(0) X_n Y_n. \end{aligned} \quad (5.18)$$

Therefore, we get

$$\begin{aligned} \Phi_4 &= i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} h'(0) \left[ \frac{(-11+3i)\xi_n(1+\xi_n^2) - 16i\xi_n}{(\xi_n-i)^2(1+\xi_n^2)^4} \right. \\ & \quad - \sqrt{-1} \frac{(-22+6i)\xi_n(1+\xi_n^2) - 32i\xi_n + (26-7i)(1+\xi_n^2) - 32 - 15i\xi_n^2(1+\xi_n^2)}{(\xi_n-i)^3(1+\xi_n^2)^4} \left. \right] \sum_{j,l=1}^{\Sigma_n-1} X_j Y_l \xi_j \xi_l \\ & \quad - \sqrt{-1} \left[ \frac{(-11-3i)\xi_n(1+\xi_n^2) + 16\xi_n + (48i-14)(1+\xi_n^2) - 32i + 30\xi_n^2(1+\xi_n^2)}{(\xi_n-i)^2(1+\xi_n^2)^4} \right. \\ & \quad \left. + \frac{(-22+6i)\xi_n(1+\xi_n^2) - 32i\xi_n + (26-7i)(1+\xi_n^2) - 32 - 15i\xi_n^2(1+\xi_n^2)}{(\xi_n-i)^3(1+\xi_n^2)^4} \right] h'(0) X_n Y_n d\xi_n \sigma(\xi') dx' \\ &= \sum_{j,l=1}^{\Sigma_n-1} X_j Y_l h'(0) \Omega_4 \int_{\Gamma^+} \left[ \frac{(-11+3i)\xi_n}{(\xi_n-i)^2(1+\xi_n^2)^3} - \frac{16i\xi_n}{(\xi_n-i)^2(1+\xi_n^2)^4} - \frac{(-22+6i)i\xi_n}{(\xi_n-i)^3(\xi_n+i)^3} \right. \\ & \quad \left. - \frac{32\xi_n}{(\xi_n-i)^3(1+\xi_n^2)^4} - \frac{(26-7i)i}{(\xi_n-i)^3(1+\xi_n^2)^3} + \frac{32i}{(\xi_n-i)^3(1+\xi_n^2)^4} - \frac{15\xi_n^2}{(\xi_n-i)^3(1+\xi_n^2)^3} \right] \xi_j \xi_l d\xi_n dx' \\ & \quad + X_n Y_n h'(0) \Omega_4 \int_{\Gamma^+} \left[ \frac{16\xi_n}{(\xi_n-i)^6(\xi_n+i)^4} - \frac{(3+11i)\xi_n}{(\xi_n-i)^5(\xi_n+i)^3} + \frac{48i-14}{(\xi_n-i)^6(\xi_n+i)^4} \right. \\ & \quad \left. - \frac{32i}{(\xi_n-i)^5(\xi_n+i)^3} + \frac{30\xi_n^2}{(\xi_n-i)^5(\xi_n+i)^3} + \frac{(-22+6i)i\xi_n}{(\xi_n-i)^3(\xi_n+i)^3} + \frac{32\xi_n}{(\xi_n-i)^3(1+\xi_n^2)^4} \right. \\ & \quad \left. + \frac{(26-7i)i}{(\xi_n-i)^3(1+\xi_n^2)^3} - \frac{32i}{(\xi_n-i)^3(1+\xi_n^2)^4} + \frac{15\xi_n^2}{(\xi_n-i)^3(1+\xi_n^2)^3} \right] d\xi_n dx' \\ &= \sum_{j,l=1}^{\Sigma_n-1} X_j Y_l h'(0) \pi^2 \Omega_4 \left\{ \frac{2\pi i}{4!} \left[ \frac{(-11+3i)\xi_n}{(\xi_n+i)^3} \right]^{(4)} \Big|_{\xi_n=i} dx' - \frac{2\pi i}{5!} \left[ \frac{16i\xi_n}{(\xi_n+i)^4} \right]^{(5)} \Big|_{\xi_n=i} dx' \right. \\ & \quad - \frac{2\pi i}{5!} \left[ \frac{(-22+6i)i\xi_n}{(\xi_n+i)^3} \right]^{(5)} \Big|_{\xi_n=i} dx' - \frac{2\pi i}{6!} \left[ \frac{32\xi_n}{(\xi_n+i)^4} \right]^{(6)} \Big|_{\xi_n=i} dx' - \frac{2\pi i}{5!} \left[ \frac{(26-7i)i}{(\xi_n+i)^3} \right]^{(5)} \Big|_{\xi_n=i} dx' \\ & \quad \left. + \frac{2\pi i}{6!} \left[ \frac{32i}{(\xi_n+i)^4} \right]^{(6)} \Big|_{\xi_n=i} dx' - \frac{2\pi i}{5!} \left[ \frac{15\xi_n^2}{(\xi_n+i)^3} \right]^{(5)} \Big|_{\xi_n=i} dx' \right\} \end{aligned}$$



$$\begin{aligned}
& + X_n Y_n h'(0) \Omega_4 \left\{ \frac{2\pi i}{5!} \left[ \frac{16\xi_n}{(\xi_n + i)^4} \right]^{(5)} \Big|_{\xi_n=i} dx' - \frac{2\pi i}{4!} \left[ \frac{(3+11i)\xi_n}{(\xi_n + i)^3} \right]^{(4)} \Big|_{\xi_n=i} dx' \right. \\
& - \frac{2\pi i}{5!} \left[ \frac{48i-14}{(\xi_n + i)^4} \right]^{(5)} \Big|_{\xi_n=i} dx' - \frac{32i \times 2\pi i}{4!} \left[ \frac{1}{(\xi_n + i)^3} \right]^{(4)} \Big|_{\xi_n=i} dx' + \frac{2\pi i}{4!} \left[ \frac{30\xi_n^2}{(\xi_n + i)^3} \right]^{(4)} \Big|_{\xi_n=i} dx' \\
& + \frac{2\pi i}{5!} \left[ \frac{(-22+6i)i\xi_n}{(\xi_n + i)^3} \right]^{(5)} \Big|_{\xi_n=i} dx' + \frac{2\pi i}{6!} \left[ \frac{32\xi_n}{(\xi_n + i)^4} \right]^{(6)} \Big|_{\xi_n=i} dx' + \frac{2\pi i}{5!} \left[ \frac{(26-7i)i}{(\xi_n + i)^3} \right]^{(5)} \Big|_{\xi_n=i} dx' \\
& - \frac{2\pi i}{6!} \left[ \frac{32i}{(\xi_n + i)^4} \right]^{(6)} \Big|_{\xi_n=i} dx' + \frac{2\pi i}{5!} \left[ \frac{15\xi_n^2}{(\xi_n + i)^3} \right]^{(5)} \Big|_{\xi_n=i} dx' dx' \left. \right\} \\
& = \left[ - \left( \frac{53}{3} + \frac{41}{6}i \right) \pi \sum_{j=1}^{n-1} X_j Y_j + \left( \frac{173i}{4} + 36 \right) X_n Y_n \right] h'(0) \pi \Omega_3 dx'. \tag{5.19}
\end{aligned}$$

**case c)**  $r = -2$ ,  $\ell = -3$ ,  $k = j = |\alpha| = 0$ .

By (5.2), we get

$$\Phi_5 = - \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\pi_{\xi_n}^+ \sigma_{-2}(\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-3}) \times \partial_{\xi_n} \sigma_{-3}(D^{-3})](x_0) d\xi_n \sigma(\xi') dx'. \tag{5.20}$$

By [8], we have

$$\begin{aligned}
\sigma_{-2}(\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-3}) & = - \sum_j A(Y) X_j \xi_j \frac{c(\xi)}{|\xi|^4} - \sum_l A(X) Y_l \xi_l \frac{c(\xi)}{|\xi|^4} - \sum_{j,l=1}^{n-1} X_j \frac{\partial_{X_l} \xi_l}{\partial_{X_j} \xi_l} \frac{c(\xi)}{|\xi|^4} \\
& + \sum_{j,l=1}^n X_j Y_l \xi_j \xi_l \frac{1}{(\xi_n^2 + 1)^4} \left\{ \left( \frac{11}{2} \xi_n (1 + \xi_n^2) + 8i \xi_n \right) h'(0) c(\xi') \right. \\
& + \left[ -2i + 6i \xi_n^2 - \frac{7}{4} (1 + \xi_n^2) + \frac{15}{4} \xi_n^2 (1 + \xi_n^2) \right] h'(0) c(dx_n) \\
& \left. - 3i \xi_n (1 + \xi_n^2) \partial_{x_n} c(\xi') + i(1 + \xi_n^2) c(\xi') c(dx_n) \partial_{x_n} c(\xi') \right\}. \tag{5.21}
\end{aligned}$$

We note that  $i < n$ ,  $\int_{|\xi'|=1} \xi_{i_1} \xi_{i_2} \cdots \xi_{i_{2d+1}} \sigma(\xi') = 0$ , so we omit some items that have no contribution for computing **case c)**.

By (5.21) and (3.20), we have

$$\begin{aligned}
& \text{tr}[\pi_{\xi_n}^+ \sigma_{-2}(\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-3}) \times \partial_{\xi_n} \sigma_{-3}(D^{-3})](x_0) |_{|\xi'|=1} \\
& = \left[ \frac{1-9i}{(\xi_n - i)^4 (\xi_n + i)^3} - \frac{6-27i}{4(\xi_n - i)^3 (\xi_n + i)^2} - \frac{27+58i}{(\xi_n - i)^5 (\xi_n + i)^3} + \frac{(1+5i)\xi_n}{(\xi_n - i)^5 (\xi_n + i)^3} \right. \\
& + \frac{81+174i}{4(\xi_n - i)^4 (\xi_n + i)^2} - \frac{10-2i}{(\xi_n - i)^6 (\xi_n + i)^3} + \frac{(14-i)\xi_n}{(\xi_n - i)^6 (\xi_n + i)^3} + \frac{16-3i}{2(\xi_n - i)^5 (\xi_n + i)^2} \\
& + \frac{4i-12}{(\xi_n - i)^7 (\xi_n + i)^3} - \frac{16i\xi_n}{(\xi_n - i)^7 (\xi_n + i)^3} + \frac{9-3i}{(\xi_n - i)^6 (\xi_n + i)^2} \left. \right] h'(0) \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l \\
& + \left[ - \frac{72+25i}{4(\xi_n - i)^4 (\xi_n + i)^3} + \frac{54+8i}{(\xi_n - i)^3 (\xi_n + i)^2} - \frac{7+43i}{(\xi_n - i)^5 (\xi_n + i)^3} + \frac{(44-9i)\xi_n}{2(\xi_n - i)^5 (\xi_n + i)^3} \right. \\
& + \frac{21+129i}{4(\xi_n - i)^4 (\xi_n + i)^2} + \frac{22-26i}{(\xi_n - i)^6 (\xi_n + i)^3} + \frac{3\xi_n}{(\xi_n - i)^6 (\xi_n + i)^3} - \frac{33-39i}{2(\xi_n - i)^6 (\xi_n + i)^2} \\
& \left. + \frac{2i}{(\xi_n - i)^7 (\xi_n + i)^3} + \frac{16\xi_n}{(\xi_n - i)^7 (\xi_n + i)^3} - \frac{3i}{2(\xi_n - i)^6 (\xi_n + i)^2} \right] h'(0) X_n Y_n. \tag{5.22}
\end{aligned}$$

So we have

$$\begin{aligned}
\Phi_5 &= -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{tr}[\pi_{\xi_n}^+ \sigma_1 (\nabla_X^{S(TM)} \nabla_Y^{S(TM)} D^{-1}) \times \partial_{\xi_n} \sigma_{-4}(D^{-3})](x_0) d\xi_n \sigma(\xi') dx' \\
&= \sum_{j,l=1}^{n-1} X_j Y_l h'(0) \Omega_4 \int_{\Gamma^+} \left[ \frac{1-9i}{(\xi_n-i)^4(\xi_n+i)^3} - \frac{6-27i}{4(\xi_n-i)^3(\xi_n+i)^2} - \frac{27+58i}{(\xi_n-i)^5(\xi_n+i)^3} \right. \\
&\quad + \frac{(1+5i)\xi_n}{(\xi_n-i)^5(\xi_n+i)^3} + \frac{81+174i}{4(\xi_n-i)^4(\xi_n+i)^2} - \frac{10-2i}{(\xi_n-i)^6(\xi_n+i)^3} + \frac{(14-i)\xi_n}{(\xi_n-i)^6(\xi_n+i)^3} \\
&\quad + \frac{16-3i}{2(\xi_n-i)^5(\xi_n+i)^2} + \frac{4i-12}{(\xi_n-i)^7(\xi_n+i)^3} - \frac{16i\xi_n}{(\xi_n-i)^7(\xi_n+i)^3} + \frac{9-3i}{(\xi_n-i)^6(\xi_n+i)^2} \left. \right] \xi_j \xi_l d\xi_n dx' \\
&\quad + X_n Y_n h'(0) \Omega_4 \int_{\Gamma^+} \left[ -\frac{72+25i}{4(\xi_n-i)^4(\xi_n+i)^3} + \frac{54+8i}{(\xi_n-i)^3(\xi_n+i)^2} - \frac{7+43i}{(\xi_n-i)^5(\xi_n+i)^3} \right. \\
&\quad + \frac{(44-9i)\xi_n}{2(\xi_n-i)^5(\xi_n+i)^3} + \frac{21+129i}{4(\xi_n-i)^4(\xi_n+i)^2} + \frac{22-26i}{(\xi_n-i)^6(\xi_n+i)^3} + \frac{3\xi_n}{(\xi_n-i)^6(\xi_n+i)^3} \\
&\quad - \frac{33-39i}{2(\xi_n-i)^6(\xi_n+i)^2} + \frac{2i}{(\xi_n-i)^7(\xi_n+i)^3} + \frac{16\xi_n}{(\xi_n-i)^7(\xi_n+i)^3} - \frac{3i}{2(\xi_n-i)^6(\xi_n+i)^2} \left. \right] d\xi_n dx' \\
&= \sum_{j,l=1}^{n-1} X_j Y_l h'(0) \pi \Omega_4 \left\{ \frac{2\pi i}{3!} \left[ \frac{1-9i}{(\xi_n+i)^3} \right]^{(3)} \Big|_{\xi_n=i} dx' - \frac{2\pi i}{2!} \left[ \frac{6-27i}{4(\xi_n+i)^2} \right]^{(2)} \Big|_{\xi_n=i} dx' \right. \\
&\quad - \frac{2\pi i}{4!} \left[ \frac{27+58i}{(\xi_n+i)^3} \right]^{(4)} \Big|_{\xi_n=i} dx' + \frac{2\pi i}{4!} \left[ \frac{(1+5i)\xi_n}{(\xi_n+i)^3} \right]^{(4)} \Big|_{\xi_n=i} dx' + \frac{2\pi i}{3!} \left[ \frac{81+174i}{4(\xi_n+i)^2} \right]^{(3)} \Big|_{\xi_n=i} dx' \\
&\quad - \frac{2\pi i}{5!} \left[ \frac{10-2i}{(\xi_n+i)^3} \right]^{(5)} \Big|_{\xi_n=i} dx' + \frac{2\pi i}{5!} \left[ \frac{(14-i)\xi_n}{(\xi_n+i)^3} \right]^{(5)} \Big|_{\xi_n=i} dx' + \frac{2\pi i}{4!} \left[ \frac{16-3i}{2(\xi_n+i)^2} \right]^{(4)} \Big|_{\xi_n=i} dx' \\
&\quad + \frac{2\pi i}{6!} \left[ \frac{4i-12}{(\xi_n+i)^3} \right]^{(6)} \Big|_{\xi_n=i} dx' - \frac{2\pi i}{6!} \left[ \frac{16i\xi_n}{(\xi_n+i)^3} \right]^{(6)} \Big|_{\xi_n=i} dx' + \frac{2\pi i}{5!} \left[ \frac{9-3i}{(\xi_n+i)^2} \right]^{(5)} \Big|_{\xi_n=i} dx' \left. \right\} \\
&\quad + X_n Y_n h'(0) \Omega_4 \left\{ \frac{2\pi i}{3!} \left[ -\frac{72+25i}{4(\xi_n+i)^3} \right]^{(3)} \Big|_{\xi_n=i} dx' + \frac{2\pi i}{2!} \left[ \frac{54+8i}{(\xi_n+i)^2} \right]^{(2)} \Big|_{\xi_n=i} dx' \right. \\
&\quad - \frac{2\pi i}{4!} \left[ \frac{7+43i}{(\xi_n+i)^3} \right]^{(4)} \Big|_{\xi_n=i} dx' + \frac{2\pi i}{4!} \left[ \frac{(44-9i)\xi_n}{2(\xi_n+i)^3} \right]^{(4)} \Big|_{\xi_n=i} dx' + \frac{2\pi i}{3!} \left[ \frac{21+129i}{4(\xi_n+i)^2} \right]^{(3)} \Big|_{\xi_n=i} dx' \\
&\quad + \frac{2\pi i}{5!} \left[ \frac{22-26i}{(\xi_n+i)^3} \right]^{(5)} \Big|_{\xi_n=i} dx' + \frac{2\pi i}{5!} \left[ \frac{3\xi_n}{(\xi_n+i)^3} \right]^{(5)} \Big|_{\xi_n=i} dx' - \frac{2\pi i}{5!} \left[ \frac{33-39i}{2(\xi_n+i)^2} \right]^{(5)} \Big|_{\xi_n=i} dx' \\
&\quad + \frac{2\pi i}{6!} \left[ \frac{2i}{(\xi_n+i)^3} \right]^{(6)} \Big|_{\xi_n=i} dx' + \frac{2\pi i}{6!} \left[ \frac{16\xi_n}{(\xi_n+i)^3} \right]^{(6)} \Big|_{\xi_n=i} dx' - \frac{2\pi i}{5!} \left[ \frac{3i}{2(\xi_n+i)^2} \right]^{(5)} \Big|_{\xi_n=i} dx' \left. \right\} \\
&= \left[ \left( -\frac{775}{6} + 113i \right) \sum_{j=1}^{n-1} X_j Y_j \pi + \left( \frac{171}{2} + \frac{369}{8}i \right) X_n Y_n \right] h'(0) \pi \Omega_4 dx'. \tag{5.23}
\end{aligned}$$

Let  $X = X^T + X_n \partial_n$ ,  $Y = Y^T + Y_n \partial_n$ , then we have  $\sum_{j=1}^{n-1} X_j Y_j = g(X^T, Y^T)$ . Now  $\Phi$  is the sum of the cases (a), (b) and (c). Therefore, we get

$$\Phi = \sum_{i=1}^5 \Phi_i = \left[ \left( \frac{475}{4} + \frac{3747}{8}i \right) X_n Y_n + \left( -\frac{8013}{16} + 46i \right) \pi g(X^T, Y^T) \right] h'(0) \pi \Omega_4 dx'. \tag{5.24}$$

Then, by (5.4)-(5.24), we obtain following theorem

**Theorem 5.1.** *Let  $M$  be a 6-dimensional oriented compact spin manifold with boundary  $\partial M$  and the metric  $g^M$  be defined as Sect.(3), then we get the following equality:*

$$\begin{aligned} & \widetilde{\text{Wres}}[\pi^+(\nabla_X^{S(TM)}\nabla_Y^{S(TM)}D^{-3})\circ\pi^+(D^{-3})] \\ &= \frac{4\pi^3}{3}\int_M[\text{Ric}(X,Y)-\frac{1}{2}sg(X,Y)]d\text{Vol}_M+\pi^3\int_M sg(X,Y)d\text{Vol}_M \\ &+ \int_{\partial M}\left[\left(\frac{475}{4}+\frac{3747}{8}i\right)X_nY_n+\left(-\frac{8013}{16}+46i\right)\pi g(X^T,Y^T)\right]h'(0)\pi\Omega_4d\text{Vol}_M. \end{aligned} \quad (5.25)$$

## Declarations

- Ethics approval and consent to participate  
Not applicable.
- Consent for publication  
Not applicable.
- Availability of data and materials  
The authors confirm that the data supporting the findings of this study are available within the article.
- Competing interests  
The authors declare no conflict of interest.
- Funding  
This research was funded by National Natural Science Foundation of China: No.11771070.
- Authors' contributions  
All authors contributed to the study conception and design. Material preparation, data collection and analysis were performed by YY and TW. The first draft of the manuscript was written by YY and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.
- Acknowledgements  
This work was supported by NSFC No.11771070. The authors thank the referee for his (or her) careful reading and helpful comments..

## References

## References

- [1] Ackermann T. A note on the Wodzicki residue. *J. Geom.Phys.* 20: 404-406, (1996).
- [2] Connes A. Quantized calculus and applications. 11th International Congress of Mathematical Physics(Paris,1994), Internat Press, Cambridge, MA, 15-36, (1995).
- [3] Connes A. The action funtinal in Noncommutative geometry. *Comm. Math. Phys.* 117: 673-683, (1998).
- [4] Dabrowski L, Sitarz A, Zalecki P. Spectral Metric and Einstein Functionals. arXiv:2206.02587, (2022).
- [5] Fedosov B V, Golse F, Leichtnam E, Schrohe E. The noncommutative residue for manifolds with boundary. *J. Funct. Anal.* 142, 1-31, (1996).
- [6] Guillemin V W. A new proof of Weyl's formula on the asymptotic distribution of eigenvalues. *Adv. Math.* 55(2): 131-160, (1985).
- [7] Kalau W, Walze M. Gravity, Noncommutative geometry and the Wodzicki residue. *J. Geom. Physics.* 16: 327-344,(1995).
- [8] Kastler D. The Dirac Operator and Gravitation. *Comm. Math. Phys.* 166: 633-643, (1995).
- [9] Wang J, Wang Y.: The Kastler-Kalau-Walze type theorem for six-dimensional manifolds with boundary. *J. Math. Phys.* 56: 052501 (2015).
- [10] Wang Y. Differential forms and the Wodzicki residue for Manifolds with Boundary. *J. Geom. Physics.* 56: 731-753, (2006).
- [11] Wang Y.: Differential forms the Noncommutative Residue for Manifolds with Boundary in the non-product Case. *Lett. Math. Phys.* 77: 41-51, (2006).
- [12] Wang Y. Gravity and the Noncommutative Residue for Manifolds with Boundary. *Lett. Math. Phys.* 80: 37-56, (2007).

- [13] Wang Y. Lower-Dimensional Volumes and Kastler-kalau-Walze Type Theorem for Manifolds with Boundary. *Commun. Theor. Phys.* 54: 38-42, (2010).
- [14] Wodzicki. M: local invariants of spectral asymmetry. *Invent. Math.* 75(1): 143-178, (1995).
- [15] Yu Y. The Index Theorem and The Heat Equation Method, *Nankai Tracts in Mathematics-Vol.2*, World Scientific Publishing, (2001).