

Uniqueness of Landau levels and their analogs with higher Chern numbers

Bruno Mera¹ and Tomoki Ozawa¹

¹*Advanced Institute for Materials Research (WPI-AIMR), Tohoku University, Sendai 980-8577, Japan*
(Dated: April 4, 2023)

Lowest Landau level wavefunctions are eigenstates of the Hamiltonian of a charged particle in two dimensions under a uniform magnetic field. They are known to be holomorphic both in real and momentum spaces, and also exhibit uniform, translationally invariant, geometrical properties in momentum space. In this paper, using the Stone-von Neumann theorem, we show that lowest Landau level wavefunctions are indeed the only possible states with unit Chern number satisfying these conditions. We also prove the uniqueness of their direct analogs with higher Chern numbers and provide their expressions.

Introduction.— There is an increasing interest in topological flat bands. Here, the adjective *flat* may refer not only to the energy dispersion but also to the geometrical properties in momentum space such as the Berry curvature and the quantum metric. Recently discovered twisted bilayer graphene in the chiral limit [1–6] is an interesting example of an energetically flat band with geometrical properties fulfilling the holomorphicity condition, under which the Berry curvature and the quantum metric are related by a strict equality (see Eq.(2) below). It has been known that when bands fulfill the holomorphicity condition and the Berry curvature is flat, their projected density operators obey the Girvin-MacDonald-Platzman (GMP) algebra [7], implying that a fractional topological phase can be stabilized [8]. The lowest Landau level is an example of such a band, and its higher Chern number analogs have also been proposed [9]. Such energetically and geometrically flat bands can be ideal platforms to host fractional topological phases [10].

In this paper, we show that lowest Landau levels are not just examples of geometrically flat bands fulfilling the holomorphicity condition, but rather they are the only possibility of such a flat band in two dimensions with unit Chern number in the isotropic case. We also show that geometrically flat bands with higher Chern numbers, such as those constructed in [9], are also uniquely determined once one fixes the Chern number and one parameter governing the anisotropy called the *modular parameter*. Finally we provide explicit expressions of those wavefunctions. On the one hand, our result places a strict constraint on the types of wavefunctions one can consider in exploring bands fulfilling the GMP algebra. On the other hand, our explicit expressions of the desired wavefunctions, which are exhaustive due to the uniqueness, can provide a solid basis to further study fractional topological physics in such bands.

Kähler bands.— Before stating our main theorem and presenting its proof, we first introduce the basic terminology of Bloch bands and the concept of Kähler band which will be used in the discussion to follow. We focus on two spatial dimensions.

Energy bands of a particle in a periodic potential are

characterized, via Bloch’s theorem, by the quasimomentum \mathbf{k} , a parameter which is taken from a two-torus known as the Brillouin zone BZ^2 . For a given \mathbf{k} , an eigenstate for a given band is described by a Bloch wavefunction $|u_{\mathbf{k}}\rangle$ which takes values in a suitable Hilbert space \mathcal{H} . Since multiplying $|u_{\mathbf{k}}\rangle$ by a nonzero complex number defines the same quantum state, a quantum state is specified by a one-dimensional vector subspace of \mathcal{H} , i.e. a point in a complex projective space. A Bloch band thus defines a map from the Brillouin zone to a complex projective space.

In Ref. [11–13], we have introduced the concept of a (*anti*)-Kähler band, a Bloch band for which the associated map from the Brillouin zone to the space of quantum states, regarded appropriately as a map between complex manifolds, is a (*anti*)-holomorphic immersion. For (*anti*)-Kähler bands, the *quantum metric* g and *Berry curvature* F are connected by a *complex structure* J , a tensor which squares to -1 , giving BZ^2 the structure of a (*anti*)-Kähler manifold. In the periodic coordinates $\mathbf{k} = (k_x, k_y)$ of the BZ^2 [14], denoting the two-by-two matrices representing g , F and J , respectively, by the same letters $g(\mathbf{k}) = [g_{ij}]_{1 \leq i, j \leq 2}$, $F(\mathbf{k}) = [F_{ij}]_{1 \leq i, j \leq 2}$ and $J(\mathbf{k}) = [J_{ij}]_{1 \leq i, j \leq 2}$, the condition that a Kähler band must satisfy is equivalent to the matrix equation

$$g(\mathbf{k}) = -\frac{i}{2}F(\mathbf{k})J(\mathbf{k}), \quad (1)$$

which taking determinants gives the quantum metric-Berry curvature relation

$$\sqrt{\det(g(\mathbf{k}))} = \frac{|F_{12}(\mathbf{k})|}{2}. \quad (2)$$

(Note that we use the convention that the Berry curvature F is purely imaginary.) In Ref. [11], the lowest Landau level appears as a special case of a Kähler band with $\mathcal{C} = 1$ for which the quantum geometry of the Brillouin zone is independent of \mathbf{k} , i.e., it is invariant under the full translation group \mathbb{R}^2 of the Brillouin zone—we refer to these Bloch bands as *geometrically flat Kähler bands* [15]. The main result of this Letter is the following theorem:

Theorem (Uniqueness of geometrically flat Kähler bands). *Geometrically flat (anti-)Kähler bands, i.e. bands in which the equality Eq. (1) holds and g is independent of \mathbf{k} , are unique up to a gauge choice, given the Chern number $\mathcal{C} \in \mathbb{Z}_{\neq 0}$ and the constant complex structure J .*

Some remarks are in order. The cases $\mathcal{C} > 0$ corresponds to Kähler bands, while $\mathcal{C} < 0$ corresponds to anti-Kähler bands. Below we assume $\mathcal{C} > 0$, but the uniqueness proof holds for $\mathcal{C} < 0$ *mutatis mutandis*; in particular, ones obtains the wavefunction for $\mathcal{C} < 0$ by taking the complex conjugate of the wavefunction for $\mathcal{C} > 0$.

The assumption that g is constant, together with Eq. (2), implies that F is also constant in momentum space. Besides, from Eq. (1), one sees that J is also constant. Such a Kähler band is thus *translation-invariant* in momentum space. Translation-invariant complex structures J in the Brillouin zone BZ^2 can be conveniently parametrized in terms of a modular parameter $\tau \in \mathbb{H}$ as $J = \frac{1}{\text{Im}(\tau)} \begin{pmatrix} -\text{Re}(\tau) & -|\tau|^2 \\ 1 & \text{Re}(\tau) \end{pmatrix}$ where $\mathbb{H} = \{\tau \in \mathbb{C} : \text{Im}(\tau) > 0\}$ is the upper half of the complex plane [13]. We then parameterize momentum space by the complex coordinate $z_\tau = k_x + \tau k_y$. Note that the Brillouin zone equipped with this complex coordinate becomes a complex torus \mathbb{C}/Λ_τ , where $\Lambda_\tau = \mathbb{Z} + \tau\mathbb{Z}$.

We will see that one cannot obtain a geometrically flat Kähler band with a finite number of total bands—a result which was already alternatively proved in Ref. [13] and which also follows from the stronger statement that, in fact, uniform Berry curvature always requires an infinite total number of bands [16].

The outline of the proof of the Theorem is the following. We first consider Kähler bands with constant J and show that Bloch wavefunctions must be written as a linear combination of theta functions with characteristics. We then assume that g , and hence F , are constant and, via the Stone-von Neumann theorem, show that only one possible combination of theta functions is allowed.

Kähler bands for translation-invariant J .— We describe the Bloch wavefunction [17] in two spatial dimensions by a collection of nonvanishing vectors $|u_{\mathbf{k}}\rangle \in \mathcal{H}$ smoothly parametrized by $\mathbf{k} \in \mathbb{R}^2$ where \mathbb{R}^2 is the universal cover of BZ^2 , and \mathcal{H} is a *fixed* Hilbert space which can be finite or infinite dimensional. We need $|u_{\mathbf{k}}\rangle$ and $|u_{\mathbf{k}+\mathbf{G}}\rangle$ for arbitrary \mathbf{G} in the reciprocal lattice to define the same quantum state. If the momentum space Hamiltonian $H_{\mathbf{k}}$ obeys the periodicity $H_{\mathbf{k}+\mathbf{G}} = H_{\mathbf{k}}$, $|u_{\mathbf{k}}\rangle$ and $|u_{\mathbf{k}+\mathbf{G}}\rangle$ differ, in general, by multiplication by a nonvanishing complex number $e_{\mathbf{G}}(\mathbf{k})$:

$$|u_{\mathbf{k}+\mathbf{G}}\rangle = |u_{\mathbf{k}}\rangle e_{\mathbf{G}}(\mathbf{k}). \quad (3)$$

By associativity of the sum (see Eq. (6)), one can show that the functions $e_{\mathbf{G}}(\mathbf{k})$ satisfy a cocycle condition

$$e_{\mathbf{G}_1+\mathbf{G}_2}(\mathbf{k}) = e_{\mathbf{G}_1}(\mathbf{k} + \mathbf{G}_2) e_{\mathbf{G}_2}(\mathbf{k}). \quad (4)$$

The family of functions then defines what is called a *system of multipliers* for a line bundle $L \rightarrow \text{BZ}^2$ (see, for example, Refs. [18, 19]). There is a gauge degree of freedom in defining the Bloch wavefunction $|u_{\mathbf{k}}\rangle$ because a state in quantum mechanics is a one-dimensional subspace of the Hilbert space. Concretely, we are free to multiply $|u_{\mathbf{k}}\rangle$ by $g(\mathbf{k}) \in \text{GL}(1; \mathbb{C}) = \mathbb{C}^*$ depending smoothly on $\mathbf{k} \in \mathbb{R}^2$. This changes the multipliers to $\frac{g(\mathbf{k}+\mathbf{G})}{g(\mathbf{k})} e_{\mathbf{G}}(\mathbf{k})$, which does not change the isomorphism class of L [19]. We assume that this line bundle has Chern number \mathcal{C} , which is nothing but the Chern number of the band under consideration.

More generally, the momentum-space Hamiltonian can transform as $H_{\mathbf{k}+\mathbf{G}} = V_{\mathbf{G}} H_{\mathbf{k}} V_{\mathbf{G}}^{-1}$, where, as a consequence of Wigner's theorem [20], $V_{\mathbf{G}}$ is either a unitary or anti-unitary transformation corresponding to a projective representation of the symmetry group consisting of the reciprocal lattice \mathbb{Z}^2 . Therefore, there can be a collection of unitary or anti-unitary transformations $V_{\mathbf{G}}$ of \mathcal{H} , $\mathbf{G} \in \mathbb{Z}^2$, such that

$$|u_{\mathbf{k}+\mathbf{G}}\rangle = (V_{\mathbf{G}} |u_{\mathbf{k}}\rangle) e_{\mathbf{G}}(\mathbf{k}), \quad (5)$$

and such that $V_{\mathbf{G}_1+\mathbf{G}_2}$ equals to $V_{\mathbf{G}_1} V_{\mathbf{G}_2}$ up to a phase factor which depends on \mathbf{G}_1 and \mathbf{G}_2 . These matrices can not depend on \mathbf{k} otherwise the Berry curvature and quantum metric, which are tensors in the Brillouin zone, would not be periodic in \mathbf{k} with respect to reciprocal lattice translations. Since an anti-unitary $V_{\mathbf{G}}$ would necessarily change the sign of the Berry curvature implying the existence of zero of the Berry curvature, contradicting with the assumption [21], the projective representation must be unitary. Additionally, we note that, by associativity of the sum,

$$\begin{aligned} |u_{\mathbf{k}+(\mathbf{G}_1+\mathbf{G}_2)}\rangle &= (V_{\mathbf{G}_1+\mathbf{G}_2} |u_{\mathbf{k}}\rangle) e_{\mathbf{G}_1+\mathbf{G}_2}(\mathbf{k}) \\ &= |u_{(\mathbf{k}+\mathbf{G}_2)+\mathbf{G}_1}\rangle = (V_{\mathbf{G}_1} V_{\mathbf{G}_2} |u_{\mathbf{k}}\rangle) e_{\mathbf{G}_1}(\mathbf{k} + \mathbf{G}_2) e_{\mathbf{G}_2}(\mathbf{k}). \end{aligned} \quad (6)$$

Since this condition holds for every \mathbf{k} , we may assume, in what concerns the Bloch wave function, $V_{\mathbf{G}_1+\mathbf{G}_2} = V_{\mathbf{G}_1} V_{\mathbf{G}_2}$, i.e. we have a unitary representation of \mathbb{Z}^2 . We can then split $V_{\mathbf{G}}$ into unitary irreducibles, which are parameterized in terms of real unit cell positions \mathbf{r} : $e^{-2\pi i \mathbf{G} \cdot \mathbf{r}}$, with \mathbf{r} and $\mathbf{r} + \mathbf{R}$, for $\mathbf{R} \in \mathbb{Z}^2$ determining the same irreducible representation. This decomposition gives us $u_{\mathbf{k}}(\mathbf{r}) \in \mathcal{H}_{\mathbf{r}}$, where $\mathcal{H}_{\mathbf{r}}$ is a direct summand in the decomposition of the representation into irreducibles and it has the property that for each $\mathbf{G} \in \mathbb{Z}^2$: $V_{\mathbf{G}} u_{\mathbf{k}}(\mathbf{r}) = e^{-2\pi i \mathbf{G} \cdot \mathbf{r}} u_{\mathbf{k}}(\mathbf{r})$. We note that $u_{\mathbf{k}}(\mathbf{r})$ can be a spinor containing multiple components. If \mathcal{H} is a finite dimensional Hilbert space, \mathbf{r} takes a value in a discrete subset of the real space unit cell, otherwise it may be the whole of the unit cell—the decomposition is then a direct-integral decomposition. This discussion allows us to derive the relation

$$u_{\mathbf{k}+\mathbf{G}}(\mathbf{r}) = e_{\mathbf{G}}(\mathbf{k}) e^{-2\pi i \mathbf{G} \cdot \mathbf{r}} u_{\mathbf{k}}(\mathbf{r}), \quad (7)$$

namely $u_{\mathbf{k}}(\mathbf{r})$ behaves as a smooth section of a line bundle $L_{\mathbf{r}} \rightarrow \text{BZ}^2$ whose multipliers are $e_{\mathbf{G}}(\mathbf{k})e^{-2\pi i\mathbf{G}\cdot\mathbf{r}}$.

Now we fix a translation-invariant complex structure on BZ^2 described by the complex coordinate $z_{\tau} = k_x + \tau k_y$. The condition for $|u_{\mathbf{k}}\rangle$ to determine a Kähler band is [12]

$$\frac{\partial}{\partial \bar{z}_{\tau}}|u_{\mathbf{k}}\rangle - \langle u_{\mathbf{k}}|\frac{\partial}{\partial \bar{z}_{\tau}}|u_{\mathbf{k}}\rangle|u_{\mathbf{k}}\rangle = 0, \quad (8)$$

which essentially says that $|u_{\mathbf{k}}\rangle$ is holomorphic up to an overall nonholomorphic nonvanishing multiplicative factor and so, after going to a holomorphic gauge so that the Berry gauge field is represented by a $(1,0)$ -form ($\langle u_{\mathbf{k}}|\frac{\partial}{\partial \bar{z}_{\tau}}|u_{\mathbf{k}}\rangle = 0$),

$$\frac{\partial}{\partial \bar{z}_{\tau}}|u_{\mathbf{k}}\rangle = 0, \quad (9)$$

which is simply the requirement of holomorphicity of $|u_{\mathbf{k}}\rangle$.

This holomorphicity condition in turn implies that the multipliers $e_{\mathbf{G}}(\mathbf{k})$ are holomorphic in z_{τ} and that $L \rightarrow \text{BZ}^2$ and, as a matter of fact, all of the $L_{\mathbf{r}} \rightarrow \text{BZ}^2$, for \mathbf{r} running in the real space unit cell, are holomorphic line bundles. Because holomorphic line bundles over complex tori, i.e. $\mathbb{R}^2/\mathbb{Z}^2$ equipped with a translation-invariant complex structure J , are determined, up to isomorphism, by the holomorphic line bundles whose spaces of holomorphic sections are described by theta functions [18, 22] the form of $|u_{\mathbf{k}}\rangle$ is already heavily constrained. This is because this condition together with Eq. (7) tells us that $u_{\mathbf{k}}(\mathbf{r})$ behaves as a holomorphic section of $L_{\mathbf{r}}$. The space of such sections is a finite dimensional vector space, denoted $H^0(\mathbb{C}/\Lambda_{\tau}, L_{\mathbf{r}})$ and whose dimension equals, by the Riemann-Roch theorem [23, 24], $\mathcal{C} = \text{deg}(L_{\mathbf{r}})$. Moreover, we can describe $H^0(\mathbb{C}/\Lambda_{\tau}, L_{\mathbf{r}})$ quite explicitly in terms of theta functions. Namely, we may assume that [19], after multiplication of $|u_{\mathbf{k}}\rangle$ by a suitable global nonvanishing holomorphic function $g(\mathbf{k})$ (in the universal cover \mathbb{R}^2)

$$e_{\mathbf{G}}(\mathbf{k}) = e^{-i\pi\tau\mathcal{C}m_y^2 - 2\pi i\mathcal{C}m_y z_{\tau}}, \quad \text{with } \mathbf{G} = (m_x, m_y) \in \mathbb{Z}^2, \quad (10)$$

and then this forces the components of $u_{\mathbf{k}}(\mathbf{r}) \in \mathcal{H}_{\mathbf{r}}$, once a basis for $\mathcal{H}_{\mathbf{r}}$ is chosen, to be linear combinations with possibly \mathbf{r} -dependent coefficients of the theta functions

$$\theta_{\mathbf{r},\alpha}(z_{\tau}) := \vartheta \left[\begin{array}{c} \frac{\alpha}{\mathcal{C}} - \frac{x}{\mathcal{C}} \\ y \end{array} \right] (\mathcal{C}z_{\tau}, \mathcal{C}\tau), \quad \alpha = 0, \dots, \mathcal{C} - 1, \quad (11)$$

where $\vartheta \left[\begin{array}{c} a \\ b \end{array} \right] (z_{\tau}, \tau) = \sum_{n \in \mathbb{Z}} e^{i\pi\tau(n+a)^2 + 2\pi i(n+a)(z_{\tau}+b)}$ is known as the theta function with characteristics prescribed by $a, b \in \mathbb{R}$.

Uniqueness of flat Kähler bands.— We now require the quantum metric to be flat and derive a much more restrictive condition on $|u_{\mathbf{k}}\rangle$. Because a Kähler band is

essentially a holomorphic immersion of a complex torus in a projective space, the translation-invariance of the quantum geometry implies that, after fixing one reference quasimomentum, say the zero vector, each translation vector $\mathbf{k} \in \mathbb{R}^2$ lifts to a quantum symmetry—a symmetry of the target projective space equipped with the Fubini-Study metric—relating $|u_0\rangle$ and $|u_{\mathbf{k}}\rangle$. Once again, due to Wigner's theorem, quantum symmetries are realized by a projective representation of the symmetry group in question, where the transformations are either unitary or anti-unitary. The key step now is to note that there exist some unitary or anti-unitary operators $U_{\mathbf{k}}$, $\mathbf{k} \in \mathbb{R}^2$, such that

$$|u_{\mathbf{k}}\rangle = U_{\mathbf{k}}|u_0\rangle, \quad \text{for } \mathbf{k} \in \mathbb{R}^2. \quad (12)$$

We cannot allow $U_{\mathbf{k}}$ to be anti-unitarity, because, if so, the Berry curvature at \mathbf{k} and 0 would change sign contradicting the fact that it is assumed to be constant. Hence, we are lead to looking at projective unitary representations of \mathbb{R}^2 . What this means is that for all $\mathbf{k}_1, \mathbf{k}_2 \in \mathbb{R}^2$ we have

$$U_{\mathbf{k}_1}U_{\mathbf{k}_2} = U_{\mathbf{k}_1+\mathbf{k}_2}\psi(\mathbf{k}_1, \mathbf{k}_2), \quad (13)$$

for $\psi(\mathbf{k}_1, \mathbf{k}_2)$ a $U(1)$ -valued 2-cocycle, i.e., $\psi(\mathbf{k}_1, \mathbf{k}_2)$ are phases satisfying

$$\psi(\mathbf{k}_1, \mathbf{k}_2 + \mathbf{k}_3)\psi(\mathbf{k}_2, \mathbf{k}_3) = \psi(\mathbf{k}_1, \mathbf{k}_2)\psi(\mathbf{k}_1 + \mathbf{k}_2, \mathbf{k}_3). \quad (14)$$

We note that $U_{\mathbf{G}} = V_{\mathbf{G}}$ for reciprocal lattice vectors \mathbf{G} . Projective unitary representations of \mathbb{R}^2 are equivalent to unitary representations of central extensions G of \mathbb{R}^2 by $U(1)$. Mathematically this means there exists a *short exact sequence of groups*

$$1 \longrightarrow U(1) \longrightarrow G \longrightarrow \mathbb{R}^2 \longrightarrow 0, \quad (15)$$

where each arrow is a group homomorphism and the kernel of each arrow is the image of the previous one. The group G as a set is just the Cartesian product $\mathbb{R}^2 \times U(1)$. As a group, we equip it with the product law

$$(\mathbf{k}_1, \lambda_1) \cdot (\mathbf{k}_2, \lambda_2) = (\mathbf{k}_1 + \mathbf{k}_2, \lambda_1\lambda_2\psi(\mathbf{k}_1, \mathbf{k}_2)). \quad (16)$$

It is then not hard to see that $U(g) := U_{\mathbf{k}}\lambda$ with $g = (\mathbf{k}, \lambda) \in G$ satisfies $U(g_1)U(g_2) = U(g_1 \cdot g_2)$ for all $g_1, g_2 \in G$ and hence gives a unitary representation of G . It is useful to define the commutator $s(\mathbf{k}_1, \mathbf{k}_2) = \psi(\mathbf{k}_1, \mathbf{k}_2)/\psi(\mathbf{k}_2, \mathbf{k}_1)$ which satisfies, see Ref. [25],

- (i) Antisymmetry: $s(\mathbf{k}_1, \mathbf{k}_2) = s^{-1}(\mathbf{k}_2, \mathbf{k}_1)$
- (ii) Alternating: $s(\mathbf{k}, \mathbf{k}) = 1$
- (iii) Bimultiplicativity: $s(\mathbf{k}_1+\mathbf{k}_2, \mathbf{k}_3) = s(\mathbf{k}_1, \mathbf{k}_3)s(\mathbf{k}_2, \mathbf{k}_3)$
 $s(\mathbf{k}_1, \mathbf{k}_2+\mathbf{k}_3) = s(\mathbf{k}_1, \mathbf{k}_2)s(\mathbf{k}_1, \mathbf{k}_3)$.

The three properties imply existence of an antisymmetric bilinear form ω in \mathbb{R}^2 with the property $s(\mathbf{k}_1, \mathbf{k}_2) = e^{i\omega(\mathbf{k}_1, \mathbf{k}_2)}$, where $\omega(\mathbf{k}_1, \mathbf{k}_2) := D\mathbf{k}_1 \times \mathbf{k}_2$, for some $D \in \mathbb{R}$ and $\mathbf{k}_1 \times \mathbf{k}_2 = \mathbf{k}^t \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \mathbf{k}_2 = k_{1,x}k_{2,y} - k_{1,y}k_{2,x}$.

Theorem 1 of Ref. [25] (see also Refs. [22, 26] for more on Heisenberg groups and central extensions) implies that G is, up to isomorphism, uniquely determined by $s(\mathbf{k}_1, \mathbf{k}_2)$. When ω is non-degenerate, i.e., for $D \neq 0$, then G is referred to as a *Heisenberg group*. A particular realization of G for a given D is determined by setting

$$\psi(\mathbf{k}_1, \mathbf{k}_2) = e^{-iDk_{1,y}k_{2,x}}. \quad (17)$$

All other realizations are obtained in terms of non-vanishing $U(1)$ -valued functions $g(\mathbf{k})$ as $\psi'(\mathbf{k}_1, \mathbf{k}_2) = \frac{g(\mathbf{k}_1 + \mathbf{k}_2)}{g(\mathbf{k}_1)g(\mathbf{k}_2)}\psi(\mathbf{k}_1, \mathbf{k}_2)$ (note that $s(\mathbf{k}_1, \mathbf{k}_2)$ is invariant under this change). Observe that with this realization of the central extension G , $D = 0$ corresponds to a trivial central extension where G is really the direct product of groups $\mathbb{R}^2 \times U(1)$. This last case will not be relevant to us as we shall see below, because $C \neq 0$.

Now the *Stone-von Neumann theorem* [22], states that the Heisenberg group G has, up to unitary isomorphism, a unique unitary irreducible representation. We will later see that this unitary degree of freedom corresponds to unitary gauge transformations. This irrep is the familiar Hilbert space of square-integrable functions $\mathcal{H} = L^2(\mathbb{R})$ of one variable q , where we have the commutation relation $[q, p] = i/D$, with $p = \frac{1}{iD} \frac{\partial}{\partial q}$. At this point, the variable q describes an abstract coordinate, but we derive how one can transform from q to the more physical position coordinate \mathbf{r} in the Supplemental Material (SM). Then,

$$\begin{aligned} U_{\mathbf{k}}\psi(q) &= e^{iDk_y p} e^{-iDk_x x} \psi(q) \\ &= e^{-iDk_x(q+k_y)} \psi(q+k_y). \end{aligned} \quad (18)$$

Indeed, it is easy to see that $U_{\mathbf{k}_1}U_{\mathbf{k}_2} = U_{\mathbf{k}_1+\mathbf{k}_2}\psi(\mathbf{k}_1, \mathbf{k}_2)$. Note that D takes, in $L^2(\mathbb{R})$, the role of the inverse of Planck's constant. We note that it is this step which requires that the number of bands, which is the dimension of the Hilbert space \mathcal{H} , must be infinite.

We can write, using the Baker-Campbell-Hausdorff formula,

$$U_{\mathbf{k}} = e^{iD(k_y p - k_x q)} e^{\frac{i}{2}Dk_x k_y}. \quad (19)$$

Using $z_\tau = k_x + \tau k_y$, we can then write

$$i(k_y p - k_x q) = \frac{1}{2\tau_2} [z_\tau(\bar{\tau}q + p) - \bar{z}_\tau(\tau q + p)]. \quad (20)$$

Now observe that, because q and p are self-adjoint, $\bar{\tau}q + p$ is the adjoint to $\tau q + p$ and that

$$[\bar{\tau}q + p, \tau q + p] = i\frac{\bar{\tau}}{D} - i\frac{\tau}{D} = \frac{2\tau_2}{D}. \quad (21)$$

It is then convenient to introduce bosonic creation and annihilation operators

$$a_\tau := \sqrt{\frac{D}{2\tau_2}}(\tau q + p) \quad \text{and} \quad a_\tau^\dagger := \sqrt{\frac{D}{2\tau_2}}(\bar{\tau}q + p), \quad (22)$$

which satisfy the canonical commutation relations $[a_\tau, a_\tau^\dagger] = 1$ and we may write

$$\begin{aligned} U_{\mathbf{k}} &= e^{i\frac{D}{2}k_x k_y} e^{\frac{1}{\sqrt{2D\tau_2}}(z_\tau a_\tau^\dagger - \bar{z}_\tau a_\tau)} \\ &= e^{i\frac{D}{2}k_x k_y} e^{-\frac{1}{4\tau_2 D}|z_\tau|^2} e^{\frac{1}{\sqrt{2D\tau_2}}z_\tau a_\tau^\dagger} e^{-\frac{1}{\sqrt{2D\tau_2}}\bar{z}_\tau a_\tau}. \end{aligned} \quad (23)$$

It is then clear that, after changing gauge, cancelling all nonholomorphic nonvanishing multiplicative factors, we see that to have a holomorphic $|u_{\mathbf{k}}\rangle$ we must have $e^{-\frac{1}{\sqrt{2D\tau_2}}\bar{z}_\tau a_\tau}|u_0\rangle = 0 \iff a_\tau|u_0\rangle = 0$. Now the condition $a_\tau|u_0\rangle = 0$ is just the statement that $|u_0\rangle$ is the groundstate of the bosonic mode a_τ . The only solution, up to an overall constant, is

$$u_0(q) = e^{-\frac{iD\tau}{2}q^2}, \quad (24)$$

which is in $L^2(\mathbb{R})$ iff $D < 0$. So a holomorphic state exists iff $D < 0$. This does not preclude us from considering other representations of G that are not irreducible. However, due to the above finding that $|u_0\rangle \in \mathcal{H}$ is unique (up to rescale) this actually implies that the resulting Kähler band is unique in a precise sense. The reason is as follows. Suppose $U_{\mathbf{k}}$ was reducible. Then the Hilbert space assumes the form $\mathcal{H} \oplus \dots \oplus \mathcal{H} \cong \mathcal{H} \otimes \mathbb{C}^l$, where \mathcal{H} is the unique irrep of G and l is the number of times it appears (l may not be finite, but this does not change the argument). The group G then acts only on the left factor in $\mathcal{H} \otimes \mathbb{C}^l$. Due to there existing only one possible choice (up to rescale) for $|u_0\rangle$ in \mathcal{H} that we can take in Eq. (12), this then implies that the augmented state in $\mathcal{H} \otimes \mathbb{C}^l$ will necessarily be of the form $|u_{\mathbf{k}}\rangle \otimes |c\rangle$ for some constant nonzero vector $|c\rangle \in \mathbb{C}^l$ and $|u_{\mathbf{k}}\rangle \in \mathcal{H}$ as in Eq. (12). This just corresponds to taking the same Bloch band and introducing a new degree of freedom, such as spin, which decouples and hence the resulting Bloch band has a definite ‘‘spin polarization’’.

Finally, we use the condition Eq. (5) to find the value of D . Note that

$$|u_{\mathbf{k}+\mathbf{G}}\rangle = \psi^{-1}(\mathbf{G}, \mathbf{k})U_{\mathbf{G}}U_{\mathbf{k}}|u_0\rangle = \psi^{-1}(\mathbf{G}, \mathbf{k})U_{\mathbf{G}}|u_{\mathbf{k}}\rangle. \quad (25)$$

Then, $e_{\mathbf{G}}(\mathbf{k}) := \psi^{-1}(\mathbf{G}, \mathbf{k})$ should be unitary (since $U_{\mathbf{k}}$ is unitary) multipliers for a line bundle over BZ^2 , and thus

$$e_{\mathbf{G}}(\mathbf{k}) = e^{iDm_y k_x} \quad \text{with} \quad \mathbf{G} = (m_x, m_y) \in \mathbb{Z}^2. \quad (26)$$

A connection on the line bundle L is determined by a one-form that is compatible with the system of multipliers:

$$A(\mathbf{k} + \mathbf{G}) = A(\mathbf{k}) + e_{\mathbf{G}}(\mathbf{k})de_{\mathbf{G}}^{-1}(\mathbf{k}) = A(\mathbf{k}) - iDm_y dk_x. \quad (27)$$

One possible choice is $A = -iDk_y dk_x$. We then see that the Chern number of L being equal to \mathcal{C} forces $D = -2\pi\mathcal{C}$. This concludes the proof of the Theorem, as the Kähler band we are looking for is precisely $|u_{\mathbf{k}}\rangle = U_{\mathbf{k}}|u_0\rangle$ for the Heisenberg group determined by the Chern number \mathcal{C} and $|u_0\rangle$ such that $a_\tau|u_0\rangle = 0$. Please note how \mathcal{C} enters in G and fixes the $U_{\mathbf{k}}$'s and \mathcal{H} , τ fixes what $|u_0\rangle$ must be.

We now proceed to give an explicit expression of $u_{\mathbf{k}}(\mathbf{r})$ in terms of theta functions—hence using the particular realization of G . We refer the reader to the SM, where we show that $\mathcal{H}_{\mathbf{r}} \cong \mathbb{C}^{\mathcal{C}}$, corresponding to the internal “color” degree of freedom, and

$$u_{\mathbf{k}}(\mathbf{r}) = e^{2\pi i k_x k_y} e^{i\pi\mathcal{C}\tau k_y^2} (\theta_{\mathbf{r},0}(z_\tau), \dots, \theta_{\mathbf{r},\mathcal{C}-1}(z_\tau)). \quad (28)$$

For the particular case $\mathcal{C} = 1$ and $\tau = i$, this is exactly the lowest Landau level Bloch wavefunction in the Landau gauge—see the SM for a detailed derivation of this fact, while for higher \mathcal{C} is the color-entangled lowest Landau level Bloch wavefunction previously considered in the literature [9, 27].

We remark that the fact that the unitary irrep of G is unique up to unitary isomorphism corresponds to the freedom of gauge choice. Indeed, since the isomorphism intertwines the action of G , it must, in particular, preserve the quantum number associated to reciprocal lattice translations \mathbf{r} . It implies that we are allowed to perform $U(\mathcal{C})$ -gauge transformations to $u_{\mathbf{k}}(\mathbf{r})$. We note that this is also equivalent to choosing a different orthogonal basis of theta functions consistent with the Chern number \mathcal{C} and the irrep of the reciprocal lattice associated with \mathbf{r} . In summary, in Eq.(28), we have two kinds of gauge degrees of freedom:

- (i) $U(\mathcal{C})$ real space gauge transformations $u_{\mathbf{k}}(\mathbf{r}) \mapsto S(\mathbf{r})u_{\mathbf{k}}(\mathbf{r})$, for $S(\mathbf{r}) \in U(\mathcal{C})$;
- (ii) \mathbb{C}^* momentum space gauge transformations $u_{\mathbf{k}}(\mathbf{r}) \mapsto g(\mathbf{k})u_{\mathbf{k}}(\mathbf{r})$, for $g(\mathbf{k}) \in \mathbb{C}^*$.

In more physical terms, the transformation (i) corresponds, for example in the case of lowest Landau levels, to go from the Landau gauge to the symmetric gauge, whereas the transformation (ii) corresponds to choosing the normalization and phase of the Bloch wavefunction at each momentum.

We have shown the uniqueness of the geometrically flat Kähler bands for given Chern number and modular parameter. In previous works, the effect of the modular parameter τ is little explored; the situation $\tau \neq i$ can arise when the effective mass of a particle is anisotropic, cf. Ref. [28] where the Galilean metric includes the information of τ . Our result shows that there is a continuous family of geometrically flat Kähler bands parameterized by τ , which provides novel degrees of freedom to explore fractional topological physics with flat Kähler bands.

B. M. is very grateful to João P. Nunes, José M. Mourão and Thomas Baier for very fruitful discussions. This work is supported by JSPS KAKENHI Grant No. JP20H01845, JST PRESTO Grant No. JPMJPR19L2, and JST CREST Grant No.JPMJCR19T1.

-
- [1] R. Bistritzer and A. H. MacDonald, Moiré bands in twisted double-layer graphene, *Proceedings of the National Academy of Sciences* **108**, 12233 (2011).
 - [2] Y. Cao, V. Fatemi, S. Fang, K. Watanabe, T. Taniguchi, E. Kaxiras, and P. Jarillo-Herrero, Unconventional superconductivity in magic-angle graphene superlattices, *Nature* **556**, 43 (2018).
 - [3] Y. Cao, V. Fatemi, A. Demir, S. Fang, S. L. Tomarken, J. Y. Luo, J. D. Sanchez-Yamagishi, K. Watanabe, T. Taniguchi, E. Kaxiras, *et al.*, Correlated insulator behaviour at half-filling in magic-angle graphene superlattices, *Nature* **556**, 80 (2018).
 - [4] E. Y. Andrei and A. H. MacDonald, Graphene bilayers with a twist, *Nature Materials* **19**, 1265 (2020).
 - [5] P. J. Ledwith, G. Tarnopolsky, E. Khalaf, and A. Vishwanath, Fractional Chern insulator states in twisted bilayer graphene: An analytical approach, *Phys. Rev. Res.* **2**, 023237 (2020).
 - [6] J. Wang and Z. Liu, Hierarchy of Ideal Flatbands in Chiral Twisted Multilayer Graphene Models, *Phys. Rev. Lett.* **128**, 176403 (2022).
 - [7] S. M. Girvin, A. H. MacDonald, and P. M. Platzman, Magneto-roton theory of collective excitations in the fractional quantum Hall effect, *Phys. Rev. B* **33**, 2481 (1986).
 - [8] R. Roy, Band geometry of fractional topological insulators, *Phys. Rev. B* **90**, 165139 (2014).
 - [9] Y.-L. Wu, N. Regnault, and B. A. Bernevig, Bloch Model Wave Functions and Pseudopotentials for All Fractional Chern Insulators, *Phys. Rev. Lett.* **110**, 106802 (2013).
 - [10] Z. Liu and E. J. Bergholtz, Recent developments in fractional Chern insulators, in *Reference Module in Materials Science and Materials Engineering* (Elsevier, 2023).
 - [11] T. Ozawa and B. Mera, Relations between topology and the quantum metric for Chern insulators, *Phys. Rev. B* **104**, 045103 (2021).
 - [12] B. Mera and T. Ozawa, Kähler geometry and Chern insulators: Relations between topology and the quantum metric, *Phys. Rev. B* **104**, 045104 (2021).
 - [13] B. Mera and T. Ozawa, Engineering geometrically flat Chern bands with Fubini-Study Kähler structure, *Phys. Rev. B* **104**, 115160 (2021).
 - [14] We use a convention for which the irreducible representations (irreps) of the real space lattice \mathbb{Z}^2 are written as $e^{2\pi i \mathbf{k} \cdot \mathbf{R}}$, with $\mathbf{R} \in \mathbb{Z}^2$, so that \mathbf{k} and $\mathbf{k} + \mathbf{G}$, with $\mathbf{G} \in \mathbb{Z}^2$, determine the same irrep and hence the same Bloch quasimomentum in $B\mathbb{Z}^2$.
 - [15] We note that concepts closely related to Kähler bands have been proposed by several authors, and we would like to clarify their relations here. *Ideal flatbands* [27, 29] refer to energetically flat Kähler bands with a constant complex structure J . *Vortexable bands* refer to bands where one can add vortices without leaving the eigenspace of

- bands; in the presence of a lattice translation symmetry, vortexable bands are equivalent to ideal flatbands [30].
- [16] D. Varjas, A. Abouelkomsan, K. Yang, and E. J. Bergholtz, Topological lattice models with constant Berry curvature, *SciPost Phys.* **12**, 118 (2022).
- [17] Here the Bloch wavefunction refers to the unit-cell periodic (up to a suitable phase when under a magnetic field) part of the eigenstate with a given quasimomentum.
- [18] D. Mumford, C. P. Ramanujam, and J. I. Manin, *Abelian varieties*, Vol. 5 (Oxford university press Oxford, 1974).
- [19] A. Beauville, Theta functions, old and new, *HAL* **2013** (2013).
- [20] D. S. Freed, On Wigner’s theorem, *Geometry & Topology Monographs* **18**, 83 (2012).
- [21] The property that the Kähler band is an immersion to the complex projective space implies that the quantum metric must be non-degenerate over the entire Brillouin zone, and hence $\sqrt{\det(g)} \neq 0$ and thus $|F_{12}| \neq 0$.
- [22] D. Mumford, *Tata lectures on theta (I-III)*, Vol. 43 (Springer, 2007).
- [23] R. Miranda, *Algebraic curves and Riemann surfaces*, Vol. 5 (American Mathematical Soc., 1995).
- [24] D. Huybrechts, *Complex geometry: an introduction*, Vol. 78 (Springer, 2005).
- [25] D. S. Freed, G. W. Moore, and G. Segal, Heisenberg groups and noncommutative fluxes, *Annals of Physics* **322**, 236 (2007).
- [26] D. S. Freed, G. W. Moore, and G. Segal, The uncertainty of fluxes, *Communications in Mathematical Physics* **271**, 247 (2007).
- [27] J. Wang, S. Klevtsov, and Z. Liu, Origin of Model Fractional Chern Insulators in All Topological Ideal Flatbands: Explicit Color-entangled Wavefunction and Exact Density Algebra, arXiv:2210.13487 [10.48550/ARXIV.2210.13487](https://arxiv.org/abs/2210.13487) (2022).
- [28] F. D. M. Haldane, Geometrical description of the fractional quantum hall effect, *Phys. Rev. Lett.* **107**, 116801 (2011).
- [29] J. Wang, J. Cano, A. J. Millis, Z. Liu, and B. Yang, Exact Landau level description of geometry and interaction in a flatband, *Phys. Rev. Lett.* **127**, 246403 (2021).
- [30] P. J. Ledwith, A. Vishwanath, and D. E. Parker, Vortexability: A Unifying Criterion for Ideal Fractional Chern Insulators, arXiv:2209.15023 [10.48550/ARXIV.2209.15023](https://arxiv.org/abs/2209.15023) (2022).

Supplemental Material for “Uniqueness of Landau levels and their analogs with higher Chern numbers”

Obtaining the lowest Landau level type wavefunctions from an explicit realization of the Heisenberg group G

We derive an explicit expression of $u_{\mathbf{k}}(\mathbf{r})$ in terms of theta functions—this will allow us to recover the lowest Landau level wavefunction and the color-entangled wavefunctions. To do this, we need to extract the component $u_{\mathbf{k}}(\mathbf{r})$ corresponding to the irreducible representation of \mathbb{Z}^2 labelled by \mathbf{r} . Hence, the defining property of $u_{\mathbf{k}}(\mathbf{r})$ is $U_{\mathbf{G}}u_{\mathbf{k}}(\mathbf{r}) = e^{-2\pi i\mathbf{G}\cdot\mathbf{r}}u_{\mathbf{k}}(\mathbf{r})$ for all $\mathbf{G} \in \mathbb{Z}^2$. This can be done through the *Bloch-Zak transform* which takes $|u_{\mathbf{k}}\rangle \in \mathcal{H} = L^2(\mathbb{R})$ to

$$u_{\mathbf{k}}(\mathbf{r}) = \sum_{\mathbf{G} \in \mathbb{Z}^2} e^{2\pi i\mathbf{r}\cdot\mathbf{G}} U_{\mathbf{G}}|u_{\mathbf{k}}\rangle, \quad (29)$$

and provides a Hilbert space isomorphism $L^2(\mathbb{R}) \cong \int_{\text{u.c.}}^{\oplus} d^2\mathbf{r} \mathcal{H}_{\mathbf{r}}$ where u.c. stands for the real space unit cell labelling irreps of the reciprocal lattice \mathbb{Z}^2 . For the transform to be an isomorphism we need to define the inner product in $\mathcal{H}_{\mathbf{r}}$ appropriately. For a general element $|f\rangle \in L^2(\mathbb{R})$, let us denote its value at a point $q \in \mathbb{R}$ by $f(q)$. The Bloch-Zak transform determines an element $f(\mathbf{r}) \in \mathcal{H}_{\mathbf{r}}$ from $|f\rangle \in L^2(\mathbb{R})$, which evaluated at a point $q \in \mathbb{R}$ reads

$$\begin{aligned} (f(\mathbf{r}))(q) &= \sum_{\mathbf{G} \in \mathbb{Z}^2} e^{2\pi i\mathbf{r}\cdot\mathbf{G}} U_{\mathbf{G}}f(q) \\ &= \sum_{m_x, m_y \in \mathbb{Z}^2} e^{2\pi i C m_x (q+m_y) + 2\pi i x m_x + 2\pi i y m_y} f(q+m_y) \\ &= \sum_{m_x, m_y \in \mathbb{Z}^2} e^{2\pi i (Cq+x)m_x + 2\pi i y m_y} f(q+m_y), \end{aligned} \quad (30)$$

where we wrote $\mathbf{G} = (m_x, m_y)$ and $\mathbf{r} = (x, y)$ and used Eq.(18) of the main text. One can easily check that the inverse transformation is $|f\rangle = \int_{\text{u.c.}} d^2\mathbf{r} f(\mathbf{r})$. Using an expression of the series expansion of the Dirac comb, $\sum_{p \in \mathbb{Z}} \delta(t-p) = \sum_{m \in \mathbb{Z}} e^{2\pi i t m}$, we obtain

$$\begin{aligned} (f(\mathbf{r}))(q) &= \sum_{m_y \in \mathbb{Z}} \sum_{p \in \mathbb{Z}} \delta(Cq+x-p) e^{2\pi i y m_y} f(q+m_y) \\ &= \sum_{m_y \in \mathbb{Z}} \sum_{p \in \mathbb{Z}} \delta(Cq+x-p) e^{2\pi i y m_y} f\left(-\frac{x}{C} + \frac{p}{C} + m_y\right). \end{aligned} \quad (31)$$

We now write $p = \alpha + \tilde{p}\mathcal{C}$ where $\alpha \in \{0, \dots, \mathcal{C} - 1\}$ and $\tilde{p} \in \mathbb{Z}$ so that $\sum_{p \in \mathbb{Z}} = \sum_{\alpha=0}^{\mathcal{C}-1} \sum_{\tilde{p} \in \mathbb{Z}}$. Then,

$$\begin{aligned}
(f(\mathbf{r}))(q) &= \sum_{m_y \in \mathbb{Z}} \sum_{\alpha=0}^{\mathcal{C}-1} \sum_{\tilde{p} \in \mathbb{Z}} \delta(\mathcal{C}q + x - \alpha - \mathcal{C}\tilde{p}) e^{2\pi i y m_y} f\left(-\frac{x}{\mathcal{C}} + \frac{\alpha}{\mathcal{C}} + m_y + \tilde{p}\right) \\
&= \sum_{m_y \in \mathbb{Z}} \sum_{\alpha=0}^{\mathcal{C}-1} \sum_{\tilde{p} \in \mathbb{Z}} \delta(\mathcal{C}q + x - \alpha - \mathcal{C}\tilde{p}) e^{2\pi i y (m_y - \tilde{p})} f\left(-\frac{x}{\mathcal{C}} + \frac{\alpha}{\mathcal{C}} + m_y\right) \\
&= \sum_{\alpha=0}^{\mathcal{C}-1} \left(\sum_{m_y \in \mathbb{Z}} f\left(-\frac{x}{\mathcal{C}} + \frac{\alpha}{\mathcal{C}} + m_y\right) e^{2\pi i \frac{1}{\mathcal{C}} y (-x + \alpha + \mathcal{C}m_y)} \right) \left(\sum_{\tilde{p} \in \mathbb{Z}} \delta(\mathcal{C}q + x - \alpha - \mathcal{C}\tilde{p}) e^{-2\pi i \frac{1}{\mathcal{C}} y (-x + \alpha + \mathcal{C}\tilde{p})} \right) \\
&\equiv \sum_{\alpha=0}^{\mathcal{C}-1} f_\alpha(\mathbf{r}) \delta_\alpha^{\mathbf{r}}(q),
\end{aligned} \tag{32}$$

where we defined

$$\delta_\alpha^{\mathbf{r}}(q) := \sum_{p \in \mathbb{Z}} \delta(\mathcal{C}q + x - \alpha - \mathcal{C}p) e^{-2\pi i \frac{1}{\mathcal{C}} y (-x + \alpha + \mathcal{C}p)}, \tag{33}$$

$$f_\alpha(\mathbf{r}) := \sum_{m_y \in \mathbb{Z}} f\left(-\frac{x}{\mathcal{C}} + \frac{\alpha}{\mathcal{C}} + m_y\right) e^{2\pi i \frac{1}{\mathcal{C}} y (-x + \alpha + \mathcal{C}m_y)}. \tag{34}$$

We can regard the vector space $\mathcal{H}_{\mathbf{r}}$ to be the span of the distributions $\delta_\alpha^{\mathbf{r}}(q)$. We see that periodicity $f(\mathbf{r} + \mathbf{R}) = f(\mathbf{r})$ holds because $\delta_\alpha^{\mathbf{r}}(q)$ and $f_\alpha(\mathbf{r})$ transform in the opposite ways; for $\mathbf{R} = (a_x, a_y) \in \mathbb{Z}^2$,

$$\delta_{\alpha+\mathbf{R}}^{\mathbf{r}}(q) = e^{-2\pi i \frac{1}{\mathcal{C}} a_y (-x + a_x + \alpha)} \delta_{\alpha - a_x}^{\mathbf{r}}(q) \text{ and } \delta_{\alpha+\mathcal{C}}^{\mathbf{r}} = \delta_\alpha^{\mathbf{r}}, \quad \alpha = 0, \dots, \mathcal{C} - 1, \tag{35}$$

$$f_{\alpha+\mathbf{R}}(\mathbf{r}) = e^{2\pi i \frac{1}{\mathcal{C}} a_y (-x + a_x + \alpha)} f_{\alpha - a_x}(\mathbf{r}) \text{ and } f_{\alpha+\mathcal{C}}(\mathbf{r}) = f_\alpha(\mathbf{r}), \quad \alpha = 0, \dots, \mathcal{C} - 1. \tag{36}$$

This periodicity is consistent with the requirement of the periodicity of the Hilbert space, $\mathcal{H}_{\mathbf{r}+\mathbf{R}} = \mathcal{H}_{\mathbf{r}}$ coming from $e^{-2\pi i \mathbf{G} \cdot (\mathbf{r} + \mathbf{R})} = e^{-2\pi i \mathbf{G} \cdot \mathbf{r}}$, which holds for any \mathbf{G} in the reciprocal lattice.

If we define a Hilbert space structure by declaring

$$\langle \delta_\alpha^{\mathbf{r}}, \delta_\beta^{\mathbf{r}} \rangle = \frac{1}{\mathcal{C}} \delta_{\alpha\beta}, \tag{37}$$

then we get the desired Hilbert space isomorphism $L^2(\mathbb{R}) \cong \int_{\text{u.c.}}^{\oplus} d^2\mathbf{r} \mathcal{H}_{\mathbf{r}}$, where the right-hand side can be interpreted as the space of square-integrable sections of a Hilbert bundle over the real space unit cell torus, obtained by identifying $\mathbf{r} \sim \mathbf{r} + \mathbf{R}$ with $\mathbf{R} \in \mathbb{Z}^2$, and whose fiber is $\mathcal{H}_{\mathbf{r}}$. A section of this Hilbert bundle is identified with a collection of smooth functions $f_0(\mathbf{r}), \dots, f_{\mathcal{C}-1}(\mathbf{r})$ satisfying the conditions in Eq. (36). The collection $\{\delta_\alpha^{\mathbf{r}}\}_{\alpha=0}^{\mathcal{C}-1}$ can then be interpreted as a global multivalued orthogonal frame field for this Hilbert bundle. To see that it is indeed a Hilbert space isomorphism, one shows that for $f, g \in L^2(\mathbb{R})$, we have

$$\begin{aligned}
\frac{1}{\mathcal{C}} \sum_{\alpha=0}^{\mathcal{C}-1} \int_{\text{u.c.}} d^2\mathbf{r} \overline{f_\alpha(\mathbf{r})} g_\alpha(\mathbf{r}) &= \frac{1}{\mathcal{C}} \sum_{\alpha=0}^{\mathcal{C}-1} \int_0^1 dx \sum_{m_y \in \mathbb{Z}} \overline{f\left(-\frac{x}{\mathcal{C}} + \frac{\alpha}{\mathcal{C}} + m_y\right)} g\left(-\frac{x}{\mathcal{C}} + \frac{\alpha}{\mathcal{C}} + m_y\right) \\
&= \int_{\mathbb{R}} dq \overline{f(q)} g(q) = \langle f | g \rangle,
\end{aligned} \tag{38}$$

where we have used $\int_0^1 dy e^{2\pi i y (m-n)} = \delta_{m,n}$ for $m, n \in \mathbb{Z}$. We also refer to $(f_0(\mathbf{r}), \dots, f_{\mathcal{C}-1}(\mathbf{r}))$ as the Bloch-Zak transform of $|f\rangle$ —we are explicitly using the isomorphism $\mathcal{H}_{\mathbf{r}} \cong \mathbb{C}^{\mathcal{C}}$ provided by the orthogonal basis determined by the $\delta_\alpha^{\mathbf{r}}$'s. Physically, these \mathcal{C} degrees of freedom can be spins, orbits, or other internal degrees of freedom.

Now we explicitly evaluate the Bloch-Zak transform for $|f\rangle = |u_{\mathbf{k}}\rangle$. First note that, From Eqs.(18) and (24) of the main text,

$$u_{\mathbf{k}}(q) = U_{\mathbf{k}} u_0(q) = e^{i\pi \mathcal{C} \tau (q + k_y)^2 + 2\pi i \mathcal{C} k_x (q + k_y)}. \tag{39}$$

Then we find,

$$f_\alpha(\mathbf{r}) = \sum_{m_y \in \mathbb{Z}} u_{\mathbf{k}} \left(-\frac{x}{\mathcal{C}} + \frac{\alpha}{\mathcal{C}} + m_y \right) e^{2\pi i \frac{1}{\mathcal{C}} y (-x + \alpha + \mathcal{C} m_y)} \quad (40)$$

$$\begin{aligned} &= \sum_{m_y \in \mathbb{Z}} e^{i\pi \mathcal{C} \tau \left(-\frac{x}{\mathcal{C}} + \frac{\alpha}{\mathcal{C}} + k_y + m_y \right)^2 + 2\pi i \mathcal{C} k_x \left(-\frac{x}{\mathcal{C}} + \frac{\alpha}{\mathcal{C}} + k_y + m_y \right) + 2\pi i \frac{1}{\mathcal{C}} y (-x + \alpha + \mathcal{C} m_y)} \\ &= e^{2\pi i k_x k_y} e^{i\pi \mathcal{C} \tau k_y^2} \vartheta \left[\begin{array}{c} \frac{\alpha}{\mathcal{C}} - \frac{x}{\mathcal{C}} \\ y \end{array} \right] (\mathcal{C} z_\tau, \mathcal{C} \tau), \end{aligned} \quad (41)$$

where $z_\tau = k_x + \tau k_y$, and we have introduced the theta functions with characteristics $a, b \in \mathbb{R}$ as

$$\vartheta \left[\begin{array}{c} a \\ b \end{array} \right] (z, \tau) := \sum_{n \in \mathbb{Z}} e^{i\pi \tau (n+a)^2 + 2\pi i (n+a)(z+b)}, \quad (42)$$

for $z \in \mathbb{C}$. Therefore, the real-space representation of the Bloch state $|u_{\mathbf{k}}\rangle$ is thus given by a \mathcal{C} -component wavefunction of the following form:

$$u_{\mathbf{k}}(\mathbf{r}) = e^{2\pi i k_x k_y} e^{i\pi \mathcal{C} \tau k_y^2} \left(\vartheta \left[\begin{array}{c} -\frac{x}{\mathcal{C}} \\ y \end{array} \right] (\mathcal{C} z_\tau, \mathcal{C} \tau), \dots, \vartheta \left[\begin{array}{c} \frac{\mathcal{C}-1}{\mathcal{C}} - \frac{x}{\mathcal{C}} \\ y \end{array} \right] (\mathcal{C} z_\tau, \mathcal{C} \tau) \right). \quad (43)$$

Since the overall normalization of the wavefunction can be chosen arbitrarily, the \mathbf{k} -dependent exponential factors in front of the theta functions can be removed if one wishes. This degree of freedom corresponds to choosing $U_{\mathbf{k}}$ and the normalization of $u_0(q)$ different from ones we used in Eq.(39). Furthermore, as noted in the main text, there is a $U(\mathcal{C})$ real-space gauge degree freedom; this is to multiply $u_{\mathbf{k}}(\mathbf{r})$ by an element of $U(\mathcal{C})$, which can depend on \mathbf{r} .

For the particular case $\mathcal{C} = 1$ and $\tau = i$, the wavefunction $u_{\mathbf{k}}(\mathbf{r})$ is just the lowest Landau level Bloch wavefunction (in the Landau gauge), while for higher \mathcal{C} is the so-called color-entangled lowest Landau level Bloch wavefunction. Since the Landau level wavefunction expressed in terms of theta functions may not be familiar to some readers, we include, for completeness, its derivation in the next section.

Lowest Landau level Bloch wavefunction

Here, for completeness, we present a derivation of the lowest Landau level Bloch wavefunction. The quantity $\mathbf{r} = (x, y)$ will denote coordinates in the plane. The Hamiltonian of an electron in a uniform magnetic field is given by

$$H = -\frac{1}{2} [(\partial_x + 2\pi i y)^2 + \partial_y^2], \quad (44)$$

where we assume the standard Euclidean metric in the plane and we use (rescaled) coordinates and a (unitary) Landau gauge such that $A = 2\pi i y dx$. Note that we use a convention for which the gauge field is purely imaginary (instead of purely real).

The Hamiltonian H is not invariant under the standard action of the translation group of the plane. However, we may introduce a projective action of the translation group such that H is invariant under it. Taking an element $\mathbf{t} = (t_x, t_y) \in \mathbb{R}^2$, its action on a wavefunction is given by a unitary operator $U(\mathbf{t})$, which we refer to as a *magnetic translation*, defined by

$$U(\mathbf{t})\psi(\mathbf{r}) = e^{2\pi i t_y x} \psi(\mathbf{r} + \mathbf{t}). \quad (45)$$

The action is projective because

$$U(\mathbf{t}_1)U(\mathbf{t}_2) = U(\mathbf{t}_1 + \mathbf{t}_2) e^{2\pi i t_{1,x} t_{2,y}} \quad (46)$$

The set of operators $\{U(\mathbf{t})\lambda : \mathbf{t} \in \mathbb{R}^2, \lambda \in U(1)\}$ forms a Heisenberg group, centrally extending the translation group \mathbb{R}^2 of the plane by $U(1)$.

The covariant derivative $\nabla = d + A$, where $d = dx\partial_x + dy\partial_y$ is the exterior derivative, commutes with $U(\mathbf{t})$:

$$U(\mathbf{t})(\nabla\psi) = \nabla(U(\mathbf{t})\psi), \quad (47)$$

and because of this it also commutes with the Hamiltonian. It follows that H is magnetic translation invariant. If we now consider the lattice \mathbb{Z}^2 , we note that for $\mathbf{R}_1, \mathbf{R}_2 \in \mathbb{Z}^2$, we have

$$U(\mathbf{R}_1)U(\mathbf{R}_2) = U(\mathbf{R}_1 + \mathbf{R}_2). \quad (48)$$

It follows that the operators $\{U(\mathbf{R}) : \mathbf{R} \in \mathbb{Z}^2\}$ give a unitary representation of the lattice and H is invariant under it. We can then apply Bloch's theorem, namely, look for eigenvectors of H that are also simultaneous eigenvectors of the lattice magnetic translations. This will be wavefunctions $\psi_{\mathbf{k}}(\mathbf{r})$ satisfying

$$U(\mathbf{R})\psi_{\mathbf{k}}(\mathbf{r}) = e^{2\pi i \mathbf{k} \cdot \mathbf{R}} \psi_{\mathbf{k}}(\mathbf{r}), \quad (49)$$

which is equivalent to, setting $\mathbf{R} = (m, n) \in \mathbb{Z}^2$,

$$\psi_{\mathbf{k}}(x + m, y + n) = e^{-2\pi i n x} e^{2\pi i \mathbf{k} \cdot \mathbf{R}} \psi_{\mathbf{k}}(x, y). \quad (50)$$

As usual, Bloch waves are not be normalizable in the whole of \mathbb{R}^2 , but only in the unit cell of the lattice. The unit cell of the lattice, due to the above periodicity constraint should be understood, topologically, as a torus $\mathbb{R}^2/\mathbb{Z}^2$, and the wavefunctions at momentum \mathbf{k} , as sections of an appropriate line bundle over this torus.

We then want to look for wavefunctions in the lowest Landau level satisfying the quasiperiodicity condition in equation Eq. (50). We now introduce the complex coordinate $w = x + iy$ in real space. We note that in this paper, the symbol $z_\tau = k_x + \tau k_y$ refers to momentum space complex coordinate, while w refers to the real space complex coordinate. The covariant derivatives along the complex coordinate direction and its complex conjugate are

$$\nabla_w = \frac{\partial}{\partial w} + \frac{\pi}{2}(w - \bar{w}) \quad \text{and} \quad \nabla_{\bar{w}} = \frac{\partial}{\partial \bar{w}} + \frac{\pi}{2}(w - \bar{w}), \quad (51)$$

and thus the Hamiltonian looks

$$H = -\frac{1}{2}(\nabla_x \nabla_x + \nabla_y \nabla_y) = -(\nabla_w \nabla_{\bar{w}} + \nabla_{\bar{w}} \nabla_w) = -2\nabla_w \nabla_{\bar{w}} + \pi. \quad (52)$$

Noting that $-\nabla_w$ is the L^2 -adjoint of $\nabla_{\bar{w}}$, the energy due to the first term for any state $|\psi\rangle$ is $\langle \psi | (-2\nabla_w \nabla_{\bar{w}}) | \psi \rangle = 2\|\nabla_{\bar{w}}|\psi\rangle\|^2 \geq 0$ and so the ground state, namely the lowest Landau level, must satisfy $\nabla_{\bar{w}}\psi(\mathbf{r}) = 0$, and the corresponding energy is π .

If we write $\psi(\mathbf{r}) = e^{-\pi y^2} \phi(\mathbf{r})$, the condition $\nabla_{\bar{w}}\psi(\mathbf{r}) = 0$ can be re-written as

$$\frac{\partial \phi(\mathbf{r})}{\partial \bar{w}} = 0, \quad (53)$$

i.e. $\phi(\mathbf{r}) \equiv \phi(w)$ is holomorphic. The lowest Landau level is precisely the space of wavefunctions which are holomorphic in the sense of Eq. (53) and square-integrable.

It turns out, as we shall prove below, that for fixed momentum $\mathbf{k} = (k_x, k_y)$, there is only one linearly independent solution consistent with the quasiperiodicity condition in equation Eq. (50) given by

$$\psi_{\mathbf{k}}(\mathbf{r}) = e^{-\pi y^2} e^{2\pi i k_x k_y} \vartheta \left[\begin{matrix} k_x \\ -k_y \end{matrix} \right] (w, i) = e^{-\pi y^2} \sum_{l \in \mathbb{Z}} e^{-\pi(l+k_x)^2 + 2\pi i(l+k_x)w - 2\pi i l k_y} \quad (54)$$

One can explicitly check that the condition Eq. (50) is satisfied by using the following identity for the theta function which holds for general $w_\tau = x + \tau y$ and $\gamma = m + \tau n$, where $\tau \in \mathbb{H}$ and $\mathbf{R} = (m, n) \in \mathbb{Z}^2$:

$$\vartheta \left[\begin{matrix} a \\ b \end{matrix} \right] (w_\tau + \gamma, \tau) = e^{-i\pi \tau n^2 - 2\pi i n w_\tau} e^{2\pi i(a m - b n)} \vartheta \left[\begin{matrix} a \\ b \end{matrix} \right] (w_\tau, \tau). \quad (55)$$

The uniqueness of this solution (54) at fixed momentum \mathbf{k} can be seen in the following way. As we have seen above, if we write $\psi_{\mathbf{k}}(\mathbf{r}) = e^{-\pi y^2} \phi_{\mathbf{k}}(\mathbf{r})$, $\phi_{\mathbf{k}}(\mathbf{r}) \equiv \phi_{\mathbf{k}}(w)$ is holomorphic in the complex variable w , and due to the condition (50), it has to satisfy

$$\phi_{\mathbf{k}}(w + \gamma) = e_\gamma(w) e^{2\pi i(k_x m + k_y n)} \phi_{\mathbf{k}}(w), \quad (56)$$

with the holomorphic multipliers

$$e_\gamma(w) := e^{\pi n^2 - 2\pi i n w}, \quad \gamma = m + ni \in \mathbb{Z} + i\mathbb{Z}, \quad (57)$$

Setting $\gamma = 1$, we see that $\phi_{\mathbf{k}}(w+1) = \phi_{\mathbf{k}}(w)e^{2\pi i k_x}$. Hence we can expand it in Fourier series $\phi_{\mathbf{k}}(w) = \sum_{l \in \mathbb{Z}} c_l e^{2\pi i(l+k_x)w}$ and, working out the relations between the coefficients from Eq. (56), one obtains the recursive relation

$$c_{l-n} = c_l e^{-\pi n^2 + 2\pi n(l+k_x) + 2\pi i k_y n} \quad (58)$$

whose general solution is $c_l = c e^{-\pi(l+k_x)^2 - 2\pi i l k_y}$, where the prefactor c may depend on \mathbf{k} . One then obtains the solution Eq. (54). We note that the holomorphic multipliers $e_\gamma(w)$ satisfy the co-cycle condition $e_{\gamma_1 + \gamma_2}(w) = e_{\gamma_2}(w + \gamma_1) e_{\gamma_1}(w)$. One can see the uniqueness of the solution (54) also from the fact that such holomorphic multipliers define a holomorphic line bundle with degree 1 (1st Chern number) over the unit cell, which has a unique holomorphic section by the Riemann-Roch theorem.

Finally we look for the expression of the cell (quasi)-periodic part of the Bloch wave, denoted by $u_{\mathbf{k}}(\mathbf{r})$, which is defined by

$$\psi_{\mathbf{k}}(\mathbf{r}) = e^{2\pi i \mathbf{k} \cdot \mathbf{r}} u_{\mathbf{k}}(\mathbf{r}). \quad (59)$$

The plane wave part of the above formula makes it so that the quasi-periodicity condition from Eq.(50) that $u_{\mathbf{k}}(\mathbf{r})$ must satisfy has the coefficient independent of \mathbf{k} :

$$u_{\mathbf{k}}(x+m, y+n) = e^{-2\pi i n x} u_{\mathbf{k}}(x, y), \quad m, n \in \mathbb{Z}, \quad (60)$$

so they can all be interpreted as (square integrable) sections of the same line bundle over the unit cell torus, hence they belong to the same Hilbert space—see also the main text on the definition of a Bloch wave function, which uses $|u_{\mathbf{k}}\rangle \in \mathcal{H}$ for fixed \mathcal{H} independently of \mathbf{k} . From the explicit expression of $\psi_{\mathbf{k}}(\mathbf{r})$ in Eq. (54), we see that

$$u_{\mathbf{k}}(x, y) = e^{-2\pi i \mathbf{k} \cdot \mathbf{r}} e^{-\pi y^2} e^{2\pi i k_x k_y} \vartheta \left[\begin{matrix} k_x \\ -k_y \end{matrix} \right] (w, i). \quad (61)$$

This expression of the Bloch wavefunction of the lowest Landau level should coincide with the general expression obtained in Eq.(43) of the previous section with the replacement $\mathcal{C} = 1$ and $\tau = i$. So far, they do not look the same; the characteristics of the theta function is the momentum for the former and real-space coordinate for the latter. They are indeed the same and we are going to show it by interchanging the characteristics and the argument of the theta function. We use the notation $z = k_x + i k_y$ from the previous section to describe the complex coordinate in momentum space.

We first note that the following identity for theta functions is known:

$$\vartheta \left[\begin{matrix} a \\ b \end{matrix} \right] (w_\tau, \tau) = e^{i\pi \tau a^2} e^{2\pi i a(w_\tau + b)} \vartheta \left[\begin{matrix} 0 \\ 0 \end{matrix} \right] (w_\tau + b + \tau a, \tau). \quad (62)$$

This identity, together with Jacobi identity of the basic theta function under modular transformations, $\vartheta \left[\begin{matrix} 0 \\ 0 \end{matrix} \right] (w_\tau/\tau, \tau) = (-i\tau)^{\frac{1}{2}} e^{\frac{\pi}{\tau} i w_\tau^2} \vartheta \left[\begin{matrix} 0 \\ 0 \end{matrix} \right] (w_\tau, \tau)$, we can interchange the characteristics and the argument and rewrite the expression in the following way:

$$\begin{aligned} u_{\mathbf{k}}(x, y) &= e^{-2\pi i \mathbf{k} \cdot \mathbf{r}} e^{-\pi y^2} e^{2\pi i k_x k_y} e^{-\pi k_x^2 + 2\pi i k_x(w - k_y)} \vartheta \left[\begin{matrix} 0 \\ 0 \end{matrix} \right] (w + iz, i) \\ &= e^{-2\pi i \mathbf{k} \cdot \mathbf{r}} e^{-\pi y^2} e^{2\pi i k_x k_y} e^{-\pi k_x^2 + 2\pi i k_x(w - k_y)} e^{-\pi(w+iz)^2} \vartheta \left[\begin{matrix} 0 \\ 0 \end{matrix} \right] (z - iw, i) \\ &= e^{-2\pi i \mathbf{k} \cdot \mathbf{r}} e^{-\pi y^2} e^{2\pi i k_x k_y} e^{-\pi k_x^2 + 2\pi i k_x(w - k_y)} e^{-\pi(w+iz)^2} e^{\pi x^2} e^{2\pi i x(z+y)} \vartheta \left[\begin{matrix} -x \\ y \end{matrix} \right] (z, i) \\ &= e^{2\pi i k_x k_y - \pi k_y^2} \vartheta \left[\begin{matrix} -x \\ y \end{matrix} \right] (z, i). \end{aligned} \quad (63)$$

This is nothing but the expression obtained in Eq.(43) for $\mathcal{C} = 1$ and $\tau = i$. This completes our proof that the lowest Landau level wavefunction is precisely the unique geometrically flat Kähler band with $\mathcal{C} = 1$ and $\tau = i$.
