

ON SCHRÖDINGER OPERATORS MODIFIED BY δ INTERACTIONS

Kaya Güven Akbaş¹, Fatih Erman², and O. Teoman Turgut³

^{1, 3}Department of Physics, Boğaziçi University, Bebek, 34342, İstanbul, Turkey

²Department of Mathematics, İzmir Institute of Technology, Urla, 35430, İzmir, Turkey

¹haci.akbas1989@gmail.com

²fatih.erman@gmail.com

³turgutte@boun.edu.tr

July 21, 2023

Abstract

We study the spectral properties of a Schrödinger operator H_0 modified by delta interactions and show explicitly how the poles of the new Green's function are rearranged relative to the poles of original Green's function of H_0 . We prove that the new bound state energies are interlaced between the old ones, and the ground state energy is always lowered if the delta interaction is attractive. We also derive an alternative perturbative method of finding the bound state energies and wave functions under the assumption of a small coupling constant in a somewhat heuristic manner. We further show that these results can be extended to cases in which a renormalization process is required. We consider the possible extensions of our results to the multi center case, to delta interaction supported on curves, and to the case, where the particle is moving in a compact two-dimensional manifold under the influence of delta interaction. Finally, the semi-relativistic extension of the last problem has been studied explicitly.

Keywords: Dirac delta interactions, point interactions, Green's function, renormalization, Schrödinger operators, spectrum, resolvent.

1 Introduction

We consider a self-adjoint Schrödinger operator or Hamiltonian $H_0 = -\frac{\hbar^2}{2m}\Delta + V$ defined on some dense domain $D(H_0)$ and assume that H_0 admits a discrete or point spectrum $\sigma_d(H)$ and a (purely absolutely) continuous real spectrum $\sigma_c(H)$, which does not overlap with the discrete spectrum (no embedded eigenvalues). We further assume that the discrete spectrum has no condensation point and the spectrum is bounded below. These conditions put some restrictions on the regularity properties of the potentials, but this is the typical situation in most of the quantum mechanical problems with physically reasonable potentials. The spectrum consists of eigenvalues E_n of H_0 with finite multiplicities,

$$H_0\phi_n = E_n\phi_n, \quad (1.1)$$

where $\phi_n \in L^2$ are the eigenvectors (or eigenfunctions) corresponding to the eigenvalues E_n , and the values λ in the continuous spectrum are the “generalized eigenvalues” corresponding to the “generalized eigenvectors” χ_λ (or generalized eigenfunctions), which is outside of L^2 . Here we omit the degeneracy/multiplicity labels for simplicity. Although *we are not aiming for a completely rigorous presentation in this work*, we point out that the proper meaning of the generalized eigenfunctions has been understood in the context of so-called Rigged Hilbert spaces [1, 2, 3] and the equation for the generalized eigenvalues are given by $H_0\chi_\lambda = \lambda\chi_\lambda$ in the sense of distributions. One may think of a non-zero tempered distribution χ_λ as a generalized eigenfunction associated with the eigenvalue λ for H_0 if and only if $\langle H_0\chi, \psi \rangle = \langle \chi_\lambda, H_0\psi \rangle = \lambda\langle \chi_\lambda, \psi \rangle$ for any infinitely differentiable rapidly decaying functions ψ (Schwartz functions), see e.g., [4] for a more elementary discussion. The function $\chi_\lambda(x) = \frac{1}{\sqrt{2\pi}}e^{i\lambda x/\hbar}$ is the well-known generalized eigenfunction of the momentum operator $(P\psi)(x) = -i\hbar\frac{d}{dx}\psi(x)$ in $L^2(\mathbb{R})$. Under relatively mild assumptions on the potential V for dimensions $d \leq 3$ case, it is known that the generalized eigenfunctions actually can be selected as continuous functions [5, 6]. This is extremely valuable for our computations as we either evaluate them at a point or integrate them over a curve in this work.

According to one of the fundamental assumptions of quantum mechanics, we can formally expand any function $\psi \in L^2(\mathbb{R})$ as (see e.g., page 48 in [7]):

$$\psi(x) = \sum_n a_n \phi_n(x) + \int_{\sigma_c(H_0)} a(\lambda) \chi_\lambda(x) d\mu(\lambda), \quad (1.2)$$

where

$$a_n = \int_{\mathbb{R}} \overline{\phi_n(x)} \psi(x) dx, \quad a(\lambda) = \int_{\mathbb{R}} \overline{\chi_\lambda(x)} \psi(x) dx. \quad (1.3)$$

The expression $d\mu(\lambda)$ in the above expansion is indeed a kind of spectral measure associated with the self-adjoint operator H_0 . If there is a degeneracy, we need to sum over those indices as well. Fourier transform is actually a formal eigenfunction expansion of a function ψ in terms of the generalized eigenfunctions χ_λ of the momentum operator P and the action of P on the function ψ is given by:

$$(P\psi)(x) = \int_{-\infty}^{\infty} \lambda \left(\int_{-\infty}^{\infty} \psi(\xi) \overline{\chi_\lambda(\xi)} d\xi \right) \chi_\lambda(x) d\lambda. \quad (1.4)$$

The theory of such eigenfunction expansions was first described by Weyl [8] for ordinary differential equations, and then developed partly by Titchmarsh [9] and by Kodaira [10]. This is extended to the multidimensional case for elliptic self-adjoint differential operators in [5]. In particular, such expansions in terms of generalized eigenfunctions are rigorously constructed, if we impose the uniformly locally square integrability condition on the potential energy in the Schrödinger operator $H_0 = -\frac{\hbar^2}{2m} \Delta + V$, which is stated as Theorem 3 in [11]. In this paper, we tacitly assume that these mild conditions for the eigenfunction expansions hold.

Our main interest here is to consider such Schrödinger operators H_0 modified by a δ interaction and show explicitly how the eigenvalues and eigenfunctions change using the eigenfunction expansions of the Green's function for such generic H_0 . The modification of the bound state spectrum in one dimension and d dimensional radial case, have been studied in the framework of path integrals in an influential work of Grosche [12] for some exactly solvable potentials V . One of the simplest choices for H_0 is the well-known harmonic oscillator problem in one dimension (as well as the radially symmetric extension of it) and the effect of adding a delta interaction to harmonic oscillator Hamiltonians has been studied by many authors [13, 14] and the statistical properties of this system have been worked out in [15] and the application of the problem to Bose-Einstein condensation has been investigated in [16]. It is well-known that the description of point like delta interactions in two and three dimensions requires renormalization and they have been studied in [17, 18] in a mathematically rigorous way and in [19, 20, 21, 22, 23] as toy models in understanding of some quantum field theoretical concepts. The higher dimensional version of the harmonic oscillator Hamiltonian with delta potential requires renormalization as well and has been discussed in [24] and more recently in [25, 26, 27]. A more interesting exactly solvable example, having both discrete and continuous spectrum, is the so-called reflectionless potential [12, 28], and we wish to show how its spectral properties change under the influence of delta interaction as a case study.

It turns out that eigenvalues of H_0 modified by δ interactions change according to some algebraic or transcendental equation and all the eigenvalues of H_0 disappear unless some set of wave functions vanish at the support of the δ interaction or if the support of Dirac delta function is chosen to be at the nodes of the wave function of the initial Hamiltonian H_0 . This result was not completely illustrated for a generic Hamiltonian H_0 , but only shown for particular exactly solvable cases [12, 24]. In all these cases, the full Green's function G of the modified system contains the Green's function G_0 of H_0 added to another term constructed again from G_0 . It is not at all obvious that the poles of G_0 cancel with the poles of this additional term (which has G_0 appearing in its expression in a nontrivial combination). This cancellation has only been pointed out for H_0 being the one dimensional harmonic oscillator Hamiltonian in [13] and for higher dimensional harmonic and linear potentials in [27]. In this paper, we prove this explicitly by using the eigenfunction expansion of the full Green's function by taking the generalized eigenfunction expansion into account and studying its pole structure for a general class of potentials. Moreover, we prove that if the support of the delta interaction is not at the node of the bound state eigenfunction of H_0 , then the new eigenvalues E_k^* are interlaced between E_{k-1} and E_k for an attractive delta interaction. We then develop a perturbative method from a different perspective to compute order by order the new eigenvalues and wave functions (in a somewhat heuristic way) under the assumption of small coupling, then compare the results with the standard approach. All the results that we have found can easily be extended to the multi center case, and to the case where we need to apply renormalization. Our perturbation method applied to the problems which require renormalization yields different results from the standard approach obtained by replacing the bare coupling constant α with the renormalized one α_R . We further show that similar

conclusions can be drawn for the deformations by delta interactions supported on curves in the plane and for the problem, for which the particle is intrinsically moving in a compact manifold under the influence of a δ interaction.

This paper is organized as follows. In Section 2, using the eigenfunction expansion of the Green's function we have explicitly shown that the poles of the Green's function of H_0 modified by delta interaction in one dimension cancels out the poles of the Green's function of H_0 under some mild conditions and prove that the new bound state energies are interlaced between the old ones and discuss the results with an explicit exactly solvable reflectionless potential. Then, we develop a new perturbative way of finding the bound state energies and the bound state wavefunctions up to the second order and compare them with the classical known results. Section 3 deals with the extension of the results to the singular case, where the renormalization of the problem is required. Finally, Section 4 is devoted to the possible extensions of the results, e.g., N center case, delta interaction supported on a curve in the plane, and the case where a particle is moving on a compact manifold interacting with a point center and its semi-relativistic extension.

2 Modifications by δ Interactions in Regular Cases

Throughout the paper, we use some well-known properties of Green's function. For convenience of the reader, we state them here with our choice of notation. For instance, the integral kernel of the resolvent $R_{H_0}(z)$ for the Schrödinger operator $H_0 = -\frac{\hbar^2}{2m} \frac{d^2}{dx^2} + V(x)$ or simply Green's function defined by

$$(R_{H_0}(E)\psi)(x) = (R_0(E)\psi)(x) = ((H_0 - E)^{-1}\psi)(x) = \int_{\mathbb{R}} G_0(x, y|E)\psi(y)dy, \quad (2.1)$$

can be expressed as the following bilinear expansion

$$G_0(x, y|E) = \sum_{n=0}^{\infty} \frac{\phi_n(x)\overline{\phi_n(y)}}{E_n - E} + \int_{\sigma_c(H_0)} \frac{\chi_\lambda(x)\overline{\chi_\lambda(y)}}{\lambda - E} d\mu(\lambda). \quad (2.2)$$

The Green's function $G_0(x, y|E)$ is a square integrable function of x for almost all y and vice versa [29]. The above formulas still hold in higher dimensions as well.

Suppose that we first consider one dimensional problems in which H_0 is modified by a δ function interaction supported at the origin

$$H = H_0 - \alpha\delta, \quad (2.3)$$

where $\alpha \in \mathbb{R}$. There are several ways to make sense of the above formal expression H . One way is to consider the δ interaction as a self-adjoint extension of H_0 . A modern introduction to this subject is the recent book by Gallone and Michelangeli [30] and the classic reference elaborating this point of view is the monograph by Albeverio et al [17]. Delta interactions can also be defined through the strong limit of the resolvent of Hamiltonians with delta interaction replaced by some scaled function, see [18, 31] for the details. The formulation of the problem in this section can be extended to higher dimensions as long as the co-dimension (dimension of the space - dimension of the support of the δ interaction) is one. We will come back to this issue for possible generalizations of the problem later on.

It is well-known that Green's function for the Hamiltonian (2.3) is given by Krein's type of formula

$$G(x, y|E) = G_0(x, y|E) + \frac{G_0(x, 0|E)G_0(0, y|E)}{\frac{1}{\alpha} - G_0(0, 0|E)}, \quad (2.4)$$

where $G_0(x, y|E)$ is the Green's function for H_0 . It is useful to define

$$\Phi(E) := \frac{1}{\alpha} - G_0(0, 0|E). \quad (2.5)$$

Some of the books in the context of point interactions use the notation Γ for the function Φ here. Here and subsequently, as emphasized in the introduction, we assume that $H_0 = -\frac{\hbar^2}{2m} \frac{d^2}{dx^2} + V(x)$ satisfies some conditions:

- H_0 is self-adjoint on some dense domain $D(H_0) \subset L^2(\mathbb{R})$,
- Spectrum of H_0 is a disjoint union of discrete $\sigma_d(H_0)$ (set of eigenvalues) and (absolutely) continuous spectrum $\sigma_c(H_0)$,
- The discrete spectrum has no accumulation point,

- For stability, we assume H_0 has spectrum bounded below,

and these assumptions are assumed to hold in higher dimensions as well. These conditions on the spectrum put some mild restrictions on the potential V . Some of the possible choices are stated in the classical work of Reed and Simon [32].

Then, we claim our first result about the spectral properties of the Hamiltonian (2.3):

Proposition 2.1. *Let $\phi_k(x)$ be the bound state wave function of H_0 associated with the bound state energy E_k . Then, the bound state energy E_k^* , when H_0 is modified (perturbed) with a delta function interaction $(-\alpha\delta(x))$, satisfies the equation*

$$\Phi(E) = \frac{1}{\alpha} - G_0(0, 0|E) = 0, \quad (2.6)$$

if $\phi_k(0) \neq 0$ for this particular k . If for this choice of k we have $\phi_k(0) = 0$, the bound state energy does not change, $E_k^* = E_k$. Moreover, the continuous spectrum of the Hamiltonian modified with delta interaction is the same as that of H_0 .

Proof. We first explicitly show how the pole structure of the full Green's function $G(x, y|E)$ is rearranged and the poles of $G_0(x, y|E)$, which explicitly appears as an additive factor in $G(x, y|E)$ actually are cancelled and new poles appear. Note that from the explicit expression of the full Green's function (2.4), one may expect that the poles of the full Green's function may contain the poles of G_0 as well as the zeroes of the function $\Phi(E)$. Using the eigenfunction expansion of the Green's function (2.2) and splitting the term in the summation associated with the isolated simple eigenvalue E_k of H_0 , we obtain

$$\begin{aligned} G(x, y|E) &= \frac{\phi_k(x)\overline{\phi_k(y)}}{E_k - E} \left(1 - \left(1 - \frac{(E_k - E)}{|\phi_k(0)|^2} D(\alpha, E) \right)^{-1} \right) + g(x, y|E) + h(x, y|E) \\ &+ \frac{(E_k - E)}{(E_k - E)D(\alpha, E) - |\phi_k(0)|^2} \left(g(x, 0|E)g(0, y|E) + g(x, 0|E)h(x, 0|E) \right. \\ &\quad \left. + g(x, 0|E)h(0, y|E) + h(x, 0|E)h(0, y|E) \right) \\ &+ \frac{1}{(E_k - E)D(\alpha, E) - |\phi_k(0)|^2} \left(g(x, 0|E)\phi_k(0)\overline{\phi_k(y)} + g(0, y|E)\phi_k(x)\overline{\phi_k(0)} \right. \\ &\quad \left. + h(x, 0|E)\phi_k(0)\overline{\phi_k(y)} + h(0, y|E)\phi_k(x)\overline{\phi_k(0)} \right), \end{aligned} \quad (2.7)$$

where we have defined the following functions, which are regular near E_k , for simplicity

$$g(x, y|E) := \sum_{n \neq k} \frac{\phi_n(x)\overline{\phi_n(y)}}{E_n - E}, \quad (2.8)$$

$$h(x, y|E) := \int_{\sigma_c(H_0)} \frac{\chi_\lambda(x)\overline{\chi_\lambda(y)}}{\lambda - E} d\mu(\lambda), \quad (2.9)$$

$$D(\alpha, E) := \frac{1}{\alpha} - \sum_{n \neq k} \frac{|\phi_n(0)|^2}{E_n - E} - \int_{\sigma_c(H_0)} \frac{|\chi_\lambda(0)|^2}{\lambda - E} d\mu(\lambda). \quad (2.10)$$

The functions g, h at $x = 0$ and $y = 0$, and the function D are well-defined thanks to the fact that the generalized eigenfunctions χ_λ in here have continuous representatives [5, 6]. Except for the first term in equation (2.7), it is easy to see that all terms are analytic in a sufficiently small disk around $E = E_k$. If we choose E sufficiently close to E_k , i.e., if $\frac{|E_k - E|}{|\phi_k(0)|^2} |D(\alpha, E)| < 1$, the first term in the above equation becomes

$$-\frac{\phi_k(x)\overline{\phi_k(y)}}{|\phi_k(0)|^2} \left(\frac{1}{\alpha} - \sum_{n \neq k} \frac{|\phi_n(0)|^2}{E_n - E} - \int_{\sigma_c(H_0)} \frac{|\chi_\lambda(0)|^2}{\lambda - E} d\mu(\lambda) \right) + O(1), \quad (2.11)$$

so that $G(x, y|E)$ is regular near $E = E_k$ as long as $\phi_k(0) \neq 0$. Hence, the pole $E = E_k$ of $G_0(x, y|E)$ is not a pole of $G(x, y|E)$ if $\phi_k(0) \neq 0$. Then, the only poles of $G(x, y|E)$ must come from the zeroes of the function $\Phi(E)$.

Finally, it follows from Weyl's theorem [32] that the continuous spectrum of the problem coincides with the initial Hamiltonian since the difference between the resolvent of the Hamiltonian and the resolvent of the initial Hamiltonian is of finite rank thanks to the explicit formula

$$R(E) = R_0(E) + (\Phi(E))^{-1} \langle \overline{G(\cdot, 0|E)}, \cdot \rangle G(\cdot, 0|E), \quad (2.12)$$

defined on its resolvent set, or as can be seen more naturally in Dirac's bra-ket notation,

$$R(E) = R_0(E) + (\Phi(E))^{-1} R_0(E) |0\rangle \langle 0| R_0(E). \quad (2.13)$$

□

Lemma 2.1. *If $\alpha > 0$ (attractive case) and $\phi_k(0) \neq 0$ for some $k \geq 1$, then the new bound state energies E_k^* are interlaced between the eigenvalues of H_0 :*

$$E_{k-1} < E_k^* < E_k. \quad (2.14)$$

For the ground state ($k = 0$), we always have $E_0^* < E_0$.

Proof. Since the Green's function G_0 has poles at the eigenvalues $E = E_n$ of H_0 in the complex E plane and has a branch cut along the generalized eigenvalues of H_0 , it is differentiable everywhere except at its poles and along the branch cut. Then, by taking the derivative of $G_0(0, 0|E)$ with respect to E under the summation and integral sign, we formally obtain

$$\frac{dG_0(0, 0|E)}{dE} = \sum_{n=0}^{\infty} \frac{|\phi_n(0)|^2}{(E_n - E)^2} + \int_{\sigma_c(H_0)} \frac{|\chi_\lambda(0)|^2}{(\lambda - E)^2} d\mu(\lambda) > 0. \quad (2.15)$$

This implies that $G_0(0, 0|E)$ is a monotonically increasing function of E between its poles as well as between the largest eigenvalue below the continuum branch cut and the infimum of the branch cut (which is typically zero). Moreover, by isolating $n = k$ th term in the sum, it is easy to see that

$$\lim_{E \rightarrow E_k^-} \sum_{n=0}^{\infty} \frac{|\phi_n(0)|^2}{E_n - E} + \int_{\sigma_c(H_0)} \frac{|\chi_\lambda(0)|^2}{\lambda - E} d\mu(\lambda) = \infty, \quad (2.16)$$

$$\lim_{E \rightarrow E_k^+} \sum_{n=0}^{\infty} \frac{|\phi_n(0)|^2}{E_n - E} + \int_{\sigma_c(H_0)} \frac{|\chi_\lambda(0)|^2}{\lambda - E} d\mu(\lambda) = -\infty, \quad (2.17)$$

$$(2.18)$$

for all $k = 0, 1, 2, \dots$, and by taking the limit under the summation and integral we have

$$\lim_{E \rightarrow -\infty} \sum_{n=0}^{\infty} \frac{|\phi_n(0)|^2}{E_n - E} + \int_{\sigma_c(H_0)} \frac{|\chi_\lambda(0)|^2}{\lambda - E} d\mu(\lambda) = 0^+. \quad (2.19)$$

Then, if $\alpha > 0$ it follows from the above results that the roots, say E_k^* , of the equation (2.6) must be located at the points of intersection of $1/\alpha$ and $G_0(0, 0|E)$, as shown in Figure 1. This shows that the new bound state energies E_k^* are shifted downwards and interlaced between E_{k-1} and E_k . Let E_0 be the ground state energy of H_0 . In contrast to excited states, the new ground state energy can be as small as possible by choosing α sufficiently large. By the positivity of the ground state wave functions of Schrödinger operators (thanks to Kato's inequality [32]), having no nodes $\phi_0(0) \neq 0$, it follows from Proposition 2.1 that new ground state energy E_0^* is always less than the older one.

□

Remark 2.2. *If $\alpha < 0$ (repulsive case), then the bound state energies of H_0 are shifted upward and interlaced as $E_k < E_k^* < E_{k+1}$ if $\phi_k(0) \neq 0$. Otherwise, bound state energies of H_0 do not change.*

Remark 2.3. *The above statements are still true if we consider the support of delta potential at $x = a$. In this case, we have*

$$G(x, y|E) = G_0(x, y|E) + \frac{G_0(x, a|E)G_0(a, y|E)}{\frac{1}{\alpha} - G_0(a, a|E)}, \quad (2.20)$$

and the bound state energies under the addition of delta interaction do not change if a is at one of the nodes of the bound state wave function of H_0 , that is, $\phi_k(a) = 0$. Otherwise, the bound state energies are obtained from $\Phi(E) = \frac{1}{\alpha} - G_0(a, a|E) = 0$.

Remark 2.4. *It is easy to see from Figure 1 that the number of bound states increases by one if we add δ interaction to H_0 . An extra pole is created below the ground state energy of H_0 and all the other eigenvalues are interlaced. Moreover, the bound state energy of H_0 is invariant under a particular configuration of delta interaction added to H_0 , that is, if the support of the delta function is chosen to be at one of the nodes of the bound state wave function associated with the bound state energy E_k of H_0 , then this bound state energy E_k does not change under the addition (perturbation) of delta interaction to H_0 .*




Figure 1: $1/\alpha$ and $\sum_{n=0}^{\infty} \frac{|\phi_n(0)|^2}{E_n - E} + \int_{\sigma_c(H_0)} \frac{|\chi_\lambda(0)|^2}{\lambda - E} d\mu(\lambda)$ versus E for $\alpha > 0$.

Example 2.5. We can illustrate what we have stated above by working out an exactly solvable case for H_0 , whose spectrum includes both discrete and continuous parts. Consider the Schrödinger operator associated with the reflectionless interaction V given by

$$(H_0\psi)(x) = -\frac{\hbar^2}{2m} \frac{d^2}{dx^2} \psi(x) - \frac{\hbar^2}{2m} \kappa^2 \frac{N(N+1)}{\cosh^2(\kappa x)} \psi(x), \quad (2.21)$$

where $N \in \mathbb{N}$. In this case, the discrete spectrum is given by the set of eigenvalues $E_n = -\frac{\hbar^2 \kappa^2}{2m} (N-n)^2$, where $n = 0, 1, 2, \dots, N-1$. The number of bound states is finite and equal to N , corresponding eigenfunctions are given by

$$\phi_n^{(N)}(x) = \sqrt{\kappa} \left((N-n) \frac{(2N+1-n)!}{n!} \right)^{1/2} P_N^{n-N}(\tanh(\kappa x)), \quad (2.22)$$

where $P_\nu^\mu(x)$ is Legendre function, defined by $P_\nu^\mu(x) = \frac{1}{\Gamma(1-\mu)} \left(\frac{1+x}{1-x} \right)^{1/2} {}_2F_1(-\nu, \nu+1; 1-\mu; \frac{1-x}{2})$ in terms of hypergeometric functions ${}_2F_1$. In this example, the continuous spectrum is the positive real axis, i.e., $\sigma_c(H_0) = [0, \infty)$ and the generalized eigenfunctions are given by $\chi_k^{(N)}(x) = \left(\frac{k}{2 \sinh(\pi k/\kappa)} \right)^{1/2} P_N^{ik/\kappa}(\tanh(\kappa x))$ with the generalized eigenvalues $E_k = \frac{\hbar^2 k^2}{2m}$, with $k \in \mathbb{R}$.

For simplicity, we choose $N = 1$. In this case, there is a single bound state and the associated normalized eigenfunction, given by

$$E_0 = -\frac{\hbar^2 \kappa^2}{2m}, \quad (2.23)$$

$$\phi_0(x) = \sqrt{\frac{\kappa}{2}} \frac{1}{\cosh(\kappa x)}. \quad (2.24)$$

Its generalized eigenfunction for this case ($N = 1$) is given by

$$\chi_E(x) = \sqrt{\frac{\kappa}{2\pi}} e^{ikx} \left(\frac{ik - \kappa \tanh(\kappa x)}{\kappa + ik} \right), \quad (2.25)$$

with the generalized eigenvalues $E = \frac{\hbar^2 k^2}{2m}$. The Green's function of this potential problem in appropriate has been discussed in [12] and it is given as

$$G_0(x, y|E) = \frac{1}{\hbar} \left(\frac{m}{-2E} \right)^{1/2} \exp\left(-\frac{|x-y|\sqrt{-2mE}}{\hbar}\right) - \frac{\kappa}{2 \cosh(\kappa x) \cosh(\kappa y) (E + \frac{\hbar^2 \kappa^2}{2m})} \times \left[1 - \left(1 - \frac{\hbar \kappa}{\sqrt{-2mE}} \right) \cosh\left(\kappa|x-y| \left(1 + \frac{\sqrt{-2mE}}{\kappa \hbar} \right) \right) \right], \quad (2.26)$$

in accordance with the conventions and units that we use here. Since ϕ_0 has no node, it follows from Proposition 2.1 that the new eigenvalue due to the addition of δ interaction to H_0 must satisfy the following equation

$$\frac{1}{\alpha} - G(0,0|E) = \frac{1}{\alpha} - \frac{1}{\hbar} \left(\frac{m}{-2E} \right)^{1/2} + \frac{\kappa}{2 \cosh^2(\kappa a) (E + \frac{\hbar^2 \kappa^2}{2m})} \left(\frac{\hbar \kappa}{\sqrt{-2mE}} \right) = 0. \quad (2.27)$$

It is easy to see from Figure 2 that as long as $a \neq 0$, we have always two bound state energies, one above




Figure 2: Graphical solution of the equation $\frac{1}{\alpha} = \frac{1}{\hbar} \left(\frac{m}{-2E} \right)^{1/2} + \frac{\kappa}{2 \cosh^2(\kappa a) (E + \frac{\hbar^2 \kappa^2}{2m})} \left(\frac{\hbar \kappa}{\sqrt{-2mE}} \right)$. Here we have chosen $\alpha = 1/2$, $\hbar = 2m = 1$, $\kappa = 1$, and $a = 1$.

$-\frac{\hbar^2 \kappa^2}{2m}$ and one below $-\frac{\hbar^2 \kappa^2}{2m}$, as expected by our analysis. It is important to notice that the Green function $G_0(0,0|E)$ blows up near its branch point at $E = 0$, which ensures the existence of two new bound states under the addition of δ interaction to the original problem. However, if $a = 0$, then $G_0(0,0|E)$ vanishes at $E = 0$. This implies that we have only one root and we have a single bound state energy below $-\frac{\hbar^2 \kappa^2}{2m}$. This is not a generic case and can be explained by the following symmetry argument. Initially, we have a reflectionless potential having \mathbb{Z}_2 symmetry, and it has a single bound state with even parity. If we include delta interaction to this potential, we still keep \mathbb{Z}_2 symmetry if $a = 0$. Then, the excited state must have an odd parity since the systems having such symmetries must have a definite symmetry and the excited state must be orthogonal to the ground state having even parity by the positivity [32]. This implies that the excited state wave function must vanish at $x = 0$, which removes the δ interaction term in the formal Hamiltonian. However, the initial system H_0 has only a single bound state, namely the ground state, so the new system modified by δ interaction must have a single bound state energy level if $a = 0$.

When $a \neq 0$, the bound state energies can be found explicitly and given by

$$E_0^* = - \left(\sqrt{\frac{\hbar^2 \kappa^2}{2m} + \frac{m\alpha^2}{8\hbar^2}} + \sqrt{\frac{m\alpha^2}{8\hbar^2}} \right)^2, \quad (2.28)$$

$$E_1^* = - \left(\sqrt{\frac{\hbar^2 \kappa^2}{2m} + \frac{m\alpha^2}{8\hbar^2}} - \sqrt{\frac{m\alpha^2}{8\hbar^2}} \right)^2. \quad (2.29)$$

2.1 A Different Perspective for the Perturbative Estimates on the Bound State Energies

We now develop a perturbative approximation in finding the eigenvalues of $H_0 - \alpha\delta$ with a different perspective. In the standard perturbation theory [28, 33], we basically start with the so-called Rayleigh-Schrödinger series expansion for the bound state energies and wave functions and then substitute these into Schrödinger equation and solve each expansion term recursively (which has been discussed more rigorously in [32, 34]).

Here, we will follow a somewhat more heuristic approach and assume that $\alpha \ll 1$ and $\phi_k(0) \neq 0$. According to the previous analysis, new bound state energies will shift due to the addition of δ interaction to H_0 . Let $E = E_k^* = E_k + \delta E_k$, where E_k are the bound state energies of H_0 and δE_k is the change in the bound state energy E_k . Then, according to Proposition 2.1 the new bound state energies E_k^* are given by the solutions of

$$\begin{aligned} \frac{1}{\alpha} - \sum_{n=0}^{\infty} \frac{|\phi_n(0)|^2}{E_n - (E_k + \delta E_k)} - \int_{\sigma_c(H_0)} \frac{|\chi_\lambda(0)|^2}{\lambda - (E_k + \delta E_k)} d\mu(\lambda) \\ = \frac{1}{\alpha} - \sum_{n \neq k} \frac{|\phi_n(0)|^2}{E_n - (E_k + \delta E_k)} + \frac{|\phi_k(0)|^2}{\delta E_k} - \int_{\sigma_c(H_0)} \frac{|\chi_\lambda(0)|^2}{\lambda - (E_k + \delta E_k)} d\mu(\lambda) = 0. \end{aligned} \quad (2.30)$$

This is an exact equation determining the bound state energies but it involves the eigenfunctions at $x = 0$. If we expand the terms in the summation and integral in the powers of δE_k and multiply the equation by $\alpha \delta E_k$, we get

$$\begin{aligned} \delta E_k - \alpha \delta E_k \sum_{n \neq k} \frac{|\phi_n(0)|^2}{E_n - E_k} \left(1 + \frac{\delta E_k}{E_n - E_k} + O(\delta E_k^2) \right) \\ - \alpha \delta E_k \int_{\sigma_c(H_0)} \frac{|\chi_\lambda(0)|^2}{\lambda - E_k} \left(1 + \frac{\delta E_k}{\lambda - E_k} + O(\delta E_k^2) \right) d\mu(\lambda) + \alpha |\phi_k(0)|^2 + O(\delta E_k^3) = 0. \end{aligned} \quad (2.31)$$

Let us assume that δE_k has a power series in α , that is, $\delta E_k = E_k^{(1)} + E_k^{(2)} + \dots$, where $E_k^{(n)}$ corresponds to the change in the bound state energy of order α^n . Then, solving $E_k^{(1)}$ and $E_k^{(2)}$ term by term, we obtain

$$E_k^{(1)} = -\alpha |\phi_k(0)|^2, \quad (2.32)$$

$$E_k^{(2)} = -\alpha^2 |\phi_k(0)|^2 \left(\sum_{n \neq k} \frac{|\phi_n(0)|^2}{E_n - E_k} + \int_{\sigma_c(H_0)} \frac{|\chi_\lambda(0)|^2}{\lambda - E_k} d\mu(\lambda) \right), \quad (2.33)$$

which are consistent with classical first and second order formulas in non-degenerate perturbation theory [7, 28], given by

$$E_k^{(1)} = \langle \phi_k, -\alpha \delta \phi_k \rangle \quad (2.34)$$

$$E_k^{(2)} = -\sum_{n \neq k} \frac{|\langle \phi_k, -\alpha \delta \phi_n \rangle|^2}{E_n - E_k} - \int_{\sigma_c(H_0)} \frac{|\langle \chi_\lambda, -\alpha \delta \phi_k \rangle|^2}{\lambda - E_k} d\mu(\lambda). \quad (2.35)$$

The important point to note here is that we are looking for the solution δE_k of (2.31) as the formal power series in α by assuming both α and δE_k are small so that it is sufficient to use regular perturbation theory [35]. Let us summarize what we have found as the following Proposition:

Proposition 2.6. *Let ϕ_k be the eigenfunctions associated with the eigenvalues E_k of H_0 and χ_λ be the generalized eigenfunctions of H_0 . Then, under the addition of δ interaction with the coupling constant $-\alpha$, the change in the bound state energies up to the first and second order in α are formally given by the equations (2.32) and (2.33).*

2.2 A Different Perspective for the Perturbative Estimates on the Bound State Wave Function

The contour integral of the resolvent $R(E) = (H - E)^{-1}$ around each simple eigenvalue E_k^* , we can find the projection operator onto the eigenspace associated with the eigenvalue E_k^* ,

$$\mathbb{P}_k = -\frac{1}{2\pi i} \oint_{\Gamma_k} R(E) dE, \quad (2.36)$$

where Γ_k is the closed contour around each simple pole E_k^* , or equivalently

$$\psi_k(x) \overline{\psi_k(y)} = -\frac{1}{2\pi i} \oint_{\Gamma_k} G(x, y|E) dE. \quad (2.37)$$

From the explicit expression of the Green's function (2.4) and residue theorem, we obtain

$$\psi_k(x) = \frac{G_0(x, 0|E_k^*)}{\left(\frac{dG_0(0, 0|E)}{dE} \Big|_{E=E_k^*} \right)^{1/2}}. \quad (2.38)$$

Using the eigenfunction expansion of G_0 , and $E_k^* = E_k + \delta E_k$, we get

$$\psi_k(x) = \frac{\left(\sum_{n=0}^{\infty} \frac{\phi_n(x)\overline{\phi_n(0)}}{E_n - (E_k + \delta E_k)} + \int_{\sigma_c(H_0)} \frac{\chi_\lambda(x)\overline{\chi_\lambda(0)}}{\lambda - (E_k + \delta E_k)} d\mu(\lambda) \right)}{\left(\sum_{n=0}^{\infty} \frac{|\phi_n(0)|^2}{(E_n - (E_k + \delta E_k))^2} + \int_{\sigma_c(H_0)} \frac{|\chi_\lambda(0)|^2}{(\lambda - (E_k + \delta E_k))^2} d\mu(\lambda) \right)^{1/2}}. \quad (2.39)$$

If we split the $n = k$ th term in the sums and integrals, and then expand the above terms in powers of δE_k , then the wave function becomes

$$\begin{aligned} \psi_k(x) &= -\frac{\phi_k(x)\overline{\phi_k(0)}}{|\phi_k(0)|} \left[1 + \frac{\delta E_k^2}{|\phi_k(0)|^2} \sum_{n \neq k} \frac{|\phi_n(0)|^2}{(E_n - E_k)^2} + \frac{\delta E_k^2}{|\phi_k(0)|^2} \int_{\sigma_c(H_0)} \frac{|\chi_\lambda(0)|^2}{(\lambda - E_k)^2} d\mu(\lambda) + O(\delta E_k^3) \right]^{-1/2} \\ &\quad + \left[1 + \frac{\delta E_k^2}{|\phi_k(0)|^2} \sum_{n \neq k} \frac{|\phi_n(0)|^2}{(E_n - E_k)^2} + \frac{\delta E_k^2}{|\phi_k(0)|^2} \int_{\sigma_c(H_0)} \frac{|\chi_\lambda(0)|^2}{(\lambda - E_k)^2} d\mu(\lambda) + O(\delta E_k^3) \right]^{-1/2} \\ &\quad \times \left(\sum_{n \neq k} \frac{\phi_n(x)\overline{\phi_n(0)}}{(E_n - E_k)|\phi_k(0)|} \left(\delta E_k + \frac{\delta E_k^2}{(E_n - E_k)} + O(\delta E_k^3) \right) \right) \\ &\quad + \left[1 + \frac{\delta E_k^2}{|\phi_k(0)|^2} \sum_{n \neq k} \frac{|\phi_n(0)|^2}{(E_n - E_k)^2} + \frac{\delta E_k^2}{|\phi_k(0)|^2} \int_{\sigma_c(H_0)} \frac{|\chi_\lambda(0)|^2}{(\lambda - E_k)^2} d\mu(\lambda) + O(\delta E_k^3) \right]^{-1/2} \\ &\quad \times \left(\int_{\sigma_c(H_0)} \frac{\chi_\lambda(x)\overline{\chi_\lambda(0)}}{(\lambda - E_k)|\phi_k(0)|} \left(\delta E_k + \frac{\delta E_k^2}{(\lambda - E_k)} + O(\delta E_k^3) \right) d\mu(\lambda) \right) + O(\delta E_k^3), \end{aligned} \quad (2.40)$$

where we have assumed $|\frac{\delta E_k}{E_n - E_k}| < 1$ and $|\frac{\delta E_k}{\lambda - E_k}| < 1$. Keeping the terms in δE_k up to the second order, we obtain

$$\begin{aligned} \psi_k(x) &= \phi_k(x)e^{-i\theta_k+i\pi} + \sum_{n \neq k} \frac{\phi_n(x)\overline{\phi_n(0)}}{|\phi_k(0)|} \frac{E_k^{(1)}}{E_n - E_k} + \int_{\sigma_c(H_0)} \frac{\chi_\lambda(x)\overline{\chi_\lambda(0)}}{|\phi_k(0)|} \frac{E_k^{(1)}}{\lambda - E_k} d\mu(\lambda) \\ &\quad - \frac{1}{2} \phi_k(x)e^{-i\theta_k+i\pi} \left(\sum_{n \neq k} \frac{|\phi_n(0)|^2}{|\phi_k(0)|^2} \frac{(E_k^{(1)})^2}{(E_n - E_k)^2} + \int_{\sigma_c(H_0)} \frac{|\chi_\lambda(0)|^2}{|\phi_k(0)|^2} \frac{(E_k^{(1)})^2}{(\lambda - E_k)^2} d\mu(\lambda) \right) \\ &\quad + \sum_{n \neq k} \frac{\phi_n(x)\overline{\phi_n(0)}}{(E_n - E_k)|\phi_k(0)|} \left(E_k^{(2)} + \frac{(E_k^{(1)})^2}{(E_n - E_k)} \right) \\ &\quad + \left(\int_{\sigma_c(H_0)} \frac{\chi_\lambda(x)\overline{\chi_\lambda(0)}}{(\lambda - E_k)|\phi_k(0)|} \left(E_k^{(2)} + \frac{(E_k^{(1)})^2}{(\lambda - E_k)} \right) d\mu(\lambda) \right) + O(\delta E_k^3), \end{aligned} \quad (2.41)$$

where $\phi_k(0) = |\phi_k(0)|e^{i\theta_k}$. Using the first order and second order results (2.34) and (2.35) for the bound state energies, we finally obtain the bound state wave function for each order:

$$\psi_k^{(0)}(x) = e^{-i\theta_k+i\pi} \phi_k(x) \quad (2.42)$$

$$\psi_k^{(1)}(x) = \alpha \phi_k(0)e^{-i\theta_k+i\pi} \sum_{n \neq k} \frac{\phi_n(x)\overline{\phi_n(0)}}{E_n - E_k} + \alpha \phi_k(0)e^{-i\theta_k+i\pi} \int_{\sigma_c(H_0)} \frac{\chi_\lambda(x)\overline{\chi_\lambda(0)}}{\lambda - E_k} d\mu(\lambda) \quad (2.43)$$

$$\begin{aligned} \psi_k^{(2)}(x) &= -\frac{\alpha^2}{2} \phi_k(x)e^{-i\theta_k+i\pi} |\phi_k(0)|^2 \left(\sum_{n \neq k} \frac{|\phi_n(0)|^2}{(E_n - E_k)^2} + \int_{\sigma_c(H_0)} \frac{|\chi_\lambda(0)|^2}{(\lambda - E_k)^2} d\mu(\lambda) \right) \\ &\quad + \alpha^2 \phi_k(0)e^{-i\theta_k+i\pi} \sum_{n \neq k} \frac{\phi_n(x)\overline{\phi_n(0)}}{(E_n - E_k)} \left(\sum_{m \neq k} \frac{|\phi_m(0)|^2}{E_m - E_k} + \int_{\sigma_c(H_0)} \frac{|\chi_\lambda(0)|^2}{\lambda - E_k} d\mu(\lambda) - \frac{|\phi_k(0)|^2}{(E_n - E_k)} \right) \\ &\quad + \alpha^2 \phi_k(0)e^{-i\theta_k+i\pi} \int_{\sigma_c(H_0)} \frac{\chi_\lambda(x)\overline{\chi_\lambda(0)}}{(\lambda - E_k)} \left(\sum_{m \neq k} \frac{|\phi_m(0)|^2}{E_m - E_k} + \int_{\sigma_c(H_0)} \frac{|\chi_{\lambda'}(0)|^2}{\lambda' - E_k} d\mu(\lambda') - \frac{|\phi_k(0)|^2}{\lambda - E_k} \right) d\mu(\lambda). \end{aligned} \quad (2.44)$$

Remark 2.7. According to standard regular perturbation theory results, the normalized wave functions for

the bound states up to second order in α are given by

$$\psi_k^{(0)}(x) = \phi_k(x) \quad (2.45)$$

$$\psi_k^{(1)}(x) = \sum_{n \neq k} \frac{\langle \phi_n, -\alpha \delta \phi_k \rangle}{E_k - E_n} \phi_n(x) + \int_{\sigma_c(H_0)} \frac{\langle \chi_\lambda, -\alpha \delta \phi_k \rangle}{\lambda - E_k} \chi_\lambda(x) d\mu(\lambda) \quad (2.46)$$

$$\begin{aligned} \psi_k^{(2)}(x) &= \sum_{n \neq k} \sum_{m \neq k} \frac{\langle \phi_n, -\alpha \delta \phi_m \rangle \langle \phi_m, -\alpha \delta \phi_k \rangle}{(E_k - E_n)(E_k - E_m)} \phi_n(x) - \sum_{n \neq k} \frac{\langle \phi_n, -\alpha \delta \phi_k \rangle \langle \phi_k, -\alpha \delta \phi_k \rangle}{(E_k - E_n)^2} \phi_n(x) \\ &\quad - \frac{1}{2} \sum_{n \neq k} \frac{\langle \phi_k, -\alpha \delta \phi_n \rangle \langle \phi_n, -\alpha \delta \phi_k \rangle}{(E_k - E_n)^2} \phi_k(x) \\ &\quad + \int_{\sigma_c(H_0)} \left(\sum_{n \neq k} \frac{\langle \phi_n, -\alpha \delta \chi_\lambda \rangle \langle \chi_\lambda, -\alpha \delta \phi_k \rangle}{(E_k - E_n)(E_k - \lambda)} \phi_n(x) \right) d\mu(\lambda) \\ &\quad + \int_{\sigma_c(H_0)} \left(\sum_{m \neq k} \frac{\langle \chi_\lambda, -\alpha \delta \phi_m \rangle \langle \phi_m, -\alpha \delta \phi_k \rangle}{(E_k - E_m)(E_k - \lambda)} \chi_\lambda(x) \right) d\mu(\lambda) \\ &\quad + \int_{\sigma_c(H_0)} \left(\int_{\sigma_c(H_0)} \frac{\langle \chi_\lambda, -\alpha \delta \chi_{\lambda'} \rangle \langle \chi_{\lambda'}, -\alpha \delta \phi_k \rangle}{(E_k - \lambda)(E_k - \lambda')} \chi_\lambda(x) d\mu(\lambda') \right) d\mu(\lambda) \\ &\quad - \int_{\sigma_c(H_0)} \frac{\langle \chi_\lambda, -\alpha \delta \phi_k \rangle \langle \phi_k, -\alpha \delta \phi_k \rangle}{(E_k - \lambda)^2} d\mu(\lambda) \chi_\lambda(x) - \frac{1}{2} \int_{\sigma_c(H_0)} \frac{\langle \phi_k, -\alpha \delta \chi_\lambda \rangle \langle \chi_\lambda, -\alpha \delta \phi_k \rangle}{(E_k - \lambda)^2} d\mu(\lambda) \phi_k(x). \end{aligned} \quad (2.47)$$

These are completely consistent with our results up to a phase factor $e^{-i\theta_k + i\pi}$. (Second order perturbation result for the normalized wave function only in the discrete case has been given in [36], but clearly it can be generalized as we have stated here).

Let us summarize our findings as

Proposition 2.8. *The bound state wave functions $\psi_k(x)$ for the Hamiltonian H_0 modified by the δ interaction are given by (2.38) in terms of the associated bound state energies E_k^* , which are the solutions of $\frac{1}{\alpha} - G_0(0, 0|E) = 0$. The expansion of the bound state wave function $\psi_k(x)$ in terms of the coupling constants α up to second order are given by (2.42), (2.43), and (2.44).*

3 Singular Modifications by δ Interactions

When the co-dimension (dimension of the space - dimension of the support of the δ interaction) is greater than one (e.g., point delta interaction in two and three dimensions, delta interaction supported by a surface in three dimensions), we need to define delta function interaction by a renormalization procedure. The reason for this is essentially based on the singularity of the Green's function for free Hamiltonians H_0 in two and three dimensions. The history of the subject is rather rich and there has been a vast amount of material in the physics literature, see e.g., [19, 20, 21, 22, 23]. The subject has been also discussed thoroughly from a more mathematical point of view in classic monographs [17, 18] as well as in a more recent work [30]. Here we now consider the Schrödinger operators H_0 with the same assumption discussed before and we study the spectrum of H_0 modified by delta interactions.

It is useful to express the Green's function G_0 in terms of the heat kernel $K_t(x, y)$ associated with the operator $H_0 = -\frac{\hbar^2}{2m}\Delta + V$, given by

$$G_0(x, y|E) = \int_0^\infty K_t(x, y) e^{tE} dt, \quad (3.1)$$

where $\text{Re}(E) < 0$ and $H_0 K_t(x, y) = i \frac{\partial}{\partial t} K_t(x, y)$ and defined for other values of E in the complex E plane through analytical continuation. We note that the first term in the short time asymptotic expansion of the diagonal heat kernel for any self-adjoint elliptic second order differential operator [38] in d dimensions, is given by

$$K_t(x, x) \sim t^{-d/2}. \quad (3.2)$$

This gives rise to the divergence near $t = 0$ in the integral representation of the Green's function (3.1). In order to make sense of such singular interactions, one must first introduce a cut-off $\epsilon > 0$ and regularize

the Hamiltonian. This could be done by replacing the delta interaction by the heat kernel $K_{\epsilon/2}(x, 0)$, which converges to $\delta(x)$ as $\epsilon \rightarrow 0$ in the distributional sense. Then, we make the coupling constant α dependent on the cut-off in such a way that the regularized Hamiltonian has a non-trivial limit (in the norm resolvent sense) as we remove the cut-off. A natural choice for the coupling constant is given by

$$\frac{1}{\alpha(\epsilon)} = \frac{1}{\alpha_R(M)} + \int_{\epsilon}^{\infty} K_t(0, 0)e^{tM} dt, \quad (3.3)$$

where M is the renormalization scale and could be eliminated in favor of a physical parameter by imposing the renormalization condition. For instance, M , and therefore α_R , can be related to the bound state energy of the particle moving under the addition of δ interaction to H_0 , say $-\mu^2$ (this requirement, in general, leads to a flow in the space of coupling constants α_R [39]). A special choice is to set $\frac{1}{\alpha_R} = 0$ while demanding $M = -\mu^2$, which is particularly convenient for bound state calculations.

Then, taking the formal limit as $\epsilon \rightarrow 0$, we obtain the integral kernel of the resolvent or Green's function, given by Krein's type of formula

$$G(x, y|E) = G_0(x, y|E) + \frac{G_0(x, 0|E)G_0(0, y|E)}{\frac{1}{\alpha_R} - \sum_{n=0}^{\infty} \frac{|\phi_n(0)|^2(E+\mu^2)}{(E_n-E)(E_n+\mu^2)} - \int_{\sigma_c(H_0)} \frac{|\chi_\lambda(0)|^2(E+\mu^2)}{(\lambda+\mu^2)(\lambda-E)} d\mu(\lambda)}, \quad (3.4)$$

where we have used the eigenfunction expansion of the heat kernel. The convergence of the sum in the denominator has been shown in the Appendix. Such specific examples are examined in [24] in the context of path integrals in two and three dimensions. However, there is no explicit derivation in this work, showing that the poles of the free resolvent are cancelled in the final expression.

By following the same line of argument developed for the one-dimensional case, we can now explicitly show how the poles of the Hamiltonian H_0 disappear with the addition of δ interaction under the same assumptions about the spectrum of H_0 as in the one dimensional case.

To simplify our arguments, we assume a purely discrete spectrum, the generalization to include a continuum with generalized eigenfunctions is fairly straightforward. We again split the term in the eigenfunction expansion of the Green's function associated with the isolated simple eigenvalue E_k of H_0 :

$$\begin{aligned} G(x, y|E) &= \sum_{n \neq k} \frac{\phi_n(x)\overline{\phi_n(y)}}{E_n - E} + \frac{\phi_k(x)\overline{\phi_k(y)}}{E_k - E} + \frac{\left(\sum_{n \neq k} \frac{\phi_n(x)\overline{\phi_n(0)}}{E_n - E}\right) \left(\sum_{n \neq k} \frac{\phi_n(0)\overline{\phi_n(y)}}{E_n - E}\right)}{\frac{1}{\alpha_R} - \sum_{n \neq k} \frac{|\phi_n(0)|^2(E+\mu^2)}{(E_n-E)(E_n+\mu^2)} - \frac{|\phi_k(0)|^2(E+\mu^2)}{(E_k-E)(E_k+\mu^2)}} \\ &+ \frac{\left(\sum_{n \neq k} \frac{\phi_n(x)\overline{\phi_n(0)}}{E_n - E}\right) \left(\frac{\phi_k(0)\overline{\phi_k(y)}}{E_k - E}\right)}{\frac{1}{\alpha_R} - \sum_{n \neq k} \frac{|\phi_n(0)|^2(E+\mu^2)}{(E_n-E)(E_n+\mu^2)} - \frac{|\phi_k(0)|^2(E+\mu^2)}{(E_k-E)(E_k+\mu^2)}} \\ &+ \frac{\left(\frac{\phi_k(x)\overline{\phi_k(0)}}{E_k - E}\right) \left(\sum_{n \neq k} \frac{\phi_n(0)\overline{\phi_n(y)}}{E_n - E}\right)}{\frac{1}{\alpha_R} - \sum_{n \neq k} \frac{|\phi_n(0)|^2(E+\mu^2)}{(E_n-E)(E_n+\mu^2)} - \frac{|\phi_k(0)|^2(E+\mu^2)}{(E_k-E)(E_k+\mu^2)}} \\ &+ \frac{\left(\frac{\phi_k(x)\overline{\phi_k(0)}}{E_k - E}\right) \left(\frac{\phi_k(0)\overline{\phi_k(y)}}{E_k - E}\right)}{\frac{1}{\alpha_R} - \sum_{n \neq k} \frac{|\phi_n(0)|^2(E+\mu^2)}{(E_n-E)(E_n+\mu^2)} - \frac{|\phi_k(0)|^2(E+\mu^2)}{(E_k-E)(E_k+\mu^2)}}. \end{aligned} \quad (3.5)$$

If we combine the second and the last term in the above expression, we obtain

$$\begin{aligned} G(x, y|E) &= \frac{\phi_k(x)\overline{\phi_k(y)}}{E_k - E} \left(1 - \left(1 - \frac{(E_k - E)}{|\phi_k(0)|^2} \left(\frac{1}{\alpha_R} - \sum_{n \neq k} \frac{|\phi_n(0)|^2}{E_n - E} + \frac{|\phi_k(0)|^2}{E_k + \mu^2} \right) \right)^{-1} \right) \\ &+ \sum_{n \neq k} \frac{\phi_n(x)\overline{\phi_n(y)}}{E_n - E} + (E_k - E) \frac{\left(\sum_{n \neq k} \frac{\phi_n(x)\overline{\phi_n(0)}}{E_n - E}\right) \left(\sum_{n \neq k} \frac{\phi_n(0)\overline{\phi_n(y)}}{E_n - E}\right)}{(E_k - E) \left(\frac{1}{\alpha_R} - \sum_{n \neq k} \frac{|\phi_n(0)|^2(E+\mu^2)}{(E_n-E)(E_n+\mu^2)}\right) - \frac{|\phi_k(0)|^2(E+\mu^2)}{(E_k+\mu^2)}} \\ &+ \frac{\left(\sum_{n \neq k} \frac{\phi_n(x)\overline{\phi_n(0)}}{E_n - E}\right) \left(\phi_k(0)\overline{\phi_k(y)}\right)}{(E_k - E) \left(\frac{1}{\alpha_R} - \sum_{n \neq k} \frac{|\phi_n(0)|^2(E+\mu^2)}{(E_n-E)(E_n+\mu^2)}\right) - \frac{|\phi_k(0)|^2(E+\mu^2)}{(E_k-E)(E_k+\mu^2)}} \\ &+ \frac{\left(\phi_k(x)\overline{\phi_k(0)}\right) \left(\sum_{n \neq k} \frac{\phi_n(0)\overline{\phi_n(y)}}{E_n - E}\right)}{(E_k - E) \left(\frac{1}{\alpha_R} - \sum_{n \neq k} \frac{|\phi_n(0)|^2(E+\mu^2)}{(E_n-E)(E_n+\mu^2)}\right) - \frac{|\phi_k(0)|^2(E+\mu^2)}{(E_k+\mu^2)}}. \end{aligned} \quad (3.6)$$

Except for the first term, it is easy to see that all terms are regular near $E = E_k$. For the first term, if we choose E sufficiently close to E_k , i.e., if $\frac{|E_k - E|}{|\phi_k(0)|^2} \left| \frac{1}{\alpha_R} - \sum_{n \neq k} \frac{|\phi_n(0)|^2}{E_n - E} + \frac{|\phi_k(0)|^2}{E_k + \mu^2} \right| < 1$, the first term in the above equation becomes

$$-\frac{\phi_k(x)\overline{\phi_k(y)}}{|\phi_k(0)|^2} \left(\frac{1}{\alpha_R} - \sum_{n \neq k} \frac{|\phi_n(0)|^2}{E_n - E} + \frac{|\phi_k(0)|^2}{E_k + \mu^2} \right) + O(|E_k - E|^2) \quad (3.7)$$

so that $G(x, y|E)$ is regular near $E = E_k$ as long as $\phi_k(0) \neq 0$. If we assume the presence of a continuous spectrum, this part of the spectrum does not change by the same reasoning as given before. Hence we have

Proposition 3.1. *Let $\phi_k(x)$ be the bound state wave function of H_0 associated with the bound state energy E_k . Then, the bound state energies E_k^* for H_0 modified (perturbed) with attractive delta interactions satisfy the equation*

$$\frac{1}{\alpha_R} - \sum_{n=0}^{\infty} \frac{|\phi_n(0)|^2(E + \mu^2)}{(E_n - E)(E_n + \mu^2)} - \int_{\sigma_c(H_0)} \frac{|\chi_\lambda(0)|^2(E + \mu^2)}{(\lambda - E)(\lambda + \mu^2)} d\mu(\lambda) = 0, \quad (3.8)$$

if $\phi_k(0) \neq 0$ for some this particular k . If for this choice of k we have $\phi_k(0) = 0$, the bound state energies do not change, $E_k^* = E_k$. Moreover, the continuous spectrum of the Hamiltonian modified with delta interaction is the same as that of H_0 .

In the renormalized case, the interlacing of the energy eigenvalues is exactly the same as the one in Lemma 2.1, where renormalization is not required, and if $\alpha_R < 0$, then all the bound state energies are shifted upward and we have $E_k < E_k^* < E_{k+1}$.

Remark 3.2. *Note that these results can be interpreted as a generalization of the well-known Sturm comparison theorems to the singular delta interactions, it is remarkable that even the renormalized case has this property.*

By following the same line of arguments as in the regular problem, we now develop a perturbative approximation to the eigenvalues for H_0 by the addition of the singular δ interaction. We assume that $\alpha_R \ll 1$. Let $E = E_k^* = E_k + \delta E_k$, where E_k are the bound state energies of H_0 . Then, the poles of the Green's function $G(x, y|E)$ are given by the solutions of

$$\begin{aligned} \frac{1}{\alpha_R} - \sum_{n=0}^{\infty} \frac{|\phi_n(0)|^2(E_k + \delta E_k + \mu^2)}{(E_n + \mu^2)(E_n - (E_k + \delta E_k))} - \int_{\sigma_c(H_0)} \frac{|\chi_\lambda(0)|^2(E_k + \delta E_k + \mu^2)}{(\lambda + \mu^2)(\lambda - (E_k + \delta E_k))} d\mu(\lambda) \\ = \frac{1}{\alpha_R} - \sum_{n \neq k} \frac{|\phi_n(0)|^2(E_k + \delta E_k + \mu^2)}{(E_n + \mu^2)(E_n - (E_k + \delta E_k))} + \frac{|\phi_k(0)|^2(E_k + \delta E_k + \mu^2)}{(E_k + \mu^2)(\delta E_k)} \\ - \int_{\sigma_c(H_0)} \frac{|\chi_\lambda(0)|^2(E_k + \delta E_k + \mu^2)}{(\lambda + \mu^2)(\lambda - (E_k + \delta E_k))} d\mu(\lambda) = 0. \end{aligned} \quad (3.9)$$

If we expand above expressions in the powers of δE_k and multiply the equation by $\alpha_R \delta E_k$, we get

$$\begin{aligned} \delta E_k + \alpha_R |\phi_k(0)|^2 + \frac{\alpha_R |\phi_k(0)|^2}{E_k + \mu^2} \delta E_k - \alpha_R \delta E_k \sum_{n \neq k} \frac{|\phi_n(0)|^2(E_k + \mu^2)}{(E_n - E_k)(E_n + \mu^2)} \left(1 + \frac{\delta E_k}{E_n - E_k} + O(\delta E_k^2) \right) \\ - \alpha_R \delta E_k \sum_{n \neq k} \frac{|\phi_n(0)|^2 \delta E_k}{(E_n - E_k)(E_n + \mu^2)} \left(1 + \frac{\delta E_k}{E_n - E_k} + O(\delta E_k^2) \right) \\ - \alpha_R \delta E_k \int_{\sigma_c(H_0)} \frac{|\chi_\lambda(0)|^2(E_k + \mu^2)}{(\lambda - E_k)(\lambda + \mu^2)} \left(1 + \frac{\delta E_k}{\lambda - E_k} + O(\delta E_k^2) \right) d\mu(\lambda) \\ - \alpha_R \delta E_k \int_{\sigma_c(H_0)} \frac{|\chi_\lambda(0)|^2 \delta E_k}{(\lambda - E_k)(\lambda + \mu^2)} \left(1 + \frac{\delta E_k}{\lambda - E_k} + O(\delta E_k^2) \right) d\mu(\lambda) + O(\delta E_k^3) = 0. \end{aligned} \quad (3.10)$$

Let us assume that δE_k can be expandable in the power series of α_R , that is, $\delta E_k = E_k^{(1)} + E_k^{(2)} + \dots$, where $E_k^{(n)}$ corresponds to the change in the bound state energy of order α_R^n . Then, solving $E_k^{(1)}$ and $E_k^{(2)}$ term by term, we obtain

$$E_k^{(1)} = -\alpha_R |\phi_k(0)|^2, \quad (3.11)$$

$$E_k^{(2)} = \alpha_R^2 |\phi_k(0)|^2 \left(\frac{|\phi_k(0)|^2}{E_k + \mu^2} - \sum_{n \neq k} \frac{|\phi_n(0)|^2(E_k + \mu^2)}{(E_n - E_k)(E_n + \mu^2)} - \int_{\sigma_c(H_0)} \frac{|\chi_\lambda(0)|^2(E_k + \mu^2)}{(\lambda - E_k)(\lambda + \mu^2)} d\mu(\lambda) \right). \quad (3.12)$$

It is important to notice that the first order result in the bound state energy is the same as the case where the renormalization is not required, except that α is replaced by the renormalized coupling constant α_R . However, the above second order result in the bound state energy is completely different from the case where the renormalization is not required. For the wave function, we formally obtain the same formula (2.41) for the wave function expansion in δE_k . Substituting the results for first and second order eigenvalues given above, we find

$$\psi_k^{(0)}(x) = e^{-i\theta_k + i\pi} \phi_k(x), \quad (3.13)$$

$$\psi_k^{(1)}(x) = \alpha_R \phi_k(0) e^{-i\theta_k + i\pi} \left(\sum_{n \neq k} \frac{\phi_n(x) \overline{\phi_n(0)}}{E_n - E_k} + \int_{\sigma_c(H_0)} \frac{\chi_\lambda(x) \overline{\chi_\lambda(0)}}{\lambda - E_k} d\mu(\lambda) \right), \quad (3.14)$$

and

$$\begin{aligned} \psi_k^{(2)}(x) = & -\frac{\alpha_R^2}{2} \phi_k(x) e^{-i\theta_k + i\pi} |\phi_k(0)|^2 \left(\sum_{n \neq k} \frac{|\phi_n(0)|^2}{(E_n - E_k)^2} + \int_{\sigma_c(H_0)} \frac{|\chi_\lambda(0)|^2}{(\lambda - E_k)^2} d\mu(\lambda) \right) \\ & + \alpha_R^2 \sum_{n \neq k} \frac{\phi_n(x) \overline{\phi_n(0)}}{(E_n - E_k)} \phi_k(0) e^{-i\theta_k + i\pi} \left[\sum_{m \neq k} \frac{|\phi_m(0)|^2 (E_k + \mu^2)}{(E_m - E_k)(E_m + \mu^2)} + \int_{\sigma_c(H_0)} \frac{|\chi_\lambda(0)|^2 (E_k + \mu^2)}{(\lambda - E_k)(\lambda + \mu^2)} d\mu(\lambda) \right. \\ & \quad \left. - \frac{|\phi_k(0)|^2 (E_n + \mu^2)}{(E_n - E_k)(E_k + \mu^2)} \right] \\ & + \alpha_R^2 \int_{\sigma_c(H_0)} \frac{\chi_\lambda(x) \overline{\chi_\lambda(0)}}{(\lambda - E_k)} \phi_k(0) e^{-i\theta_k + i\pi} \left[\sum_{m \neq k} \frac{|\phi_m(0)|^2}{E_m - E_k} + \int_{\sigma_c(H_0)} \frac{|\chi_{\lambda'}(0)|^2}{\lambda' - E_k} d\mu(\lambda') \right. \\ & \quad \left. - \frac{|\phi_k(0)|^2 (\lambda + \mu^2)}{(\lambda - E_k)(E_k + \mu^2)} \right] d\mu(\lambda). \end{aligned} \quad (3.15)$$

4 Possible Generalizations

4.1 N Center Case

It is possible to generalize our approach to N -point delta interaction case. When there is no need for renormalization, we will assume that all the couplings are actually positive (hence corresponds to the attractive case). Let us enumerate these points as a_1, a_2, \dots, a_N with $a_i \neq a_j$ whenever $i \neq j$ and the associated couplings with $\alpha_1, \alpha_2, \dots, \alpha_N$. We will proceed recursively, and suppose that H_0 has the same spectral properties discussed in Section 2. If some set of eigenfunctions satisfies $\phi_k(a_1) \neq 0$ then they lead to a shift of this eigenvalue to a new value E_k^{*1} in between E_{k-1} and E_k . For these, the eigenfunctions change to $\mathcal{N}_0 G_0(x, a_1 | E_k^{*1})$ where \mathcal{N}_0 is the normalization constant. So, we have a new Hamiltonian H_1 with a new set of discrete states $\phi_k^{(1)}$ (and possibly a new set of continuum states as found before by the formula (??), that we represent by $\eta_\lambda^{(1)}$ here). We now are back to the initial case, if we add the point interaction at a_2 , the same construction is repeated, and new eigenvalues E_k^{*2} for $\phi_k^{(1)}(a_2) \neq 0$ fall in between E_{k-1}^{*1} and E_k^{*1} . The poles associated with the old eigenvalues are removed, and there are new wave functions given by $\mathcal{N}_1 G_1(x, a_2 | E_k^{*2})$ (similarly we can talk about the continuum, yet it is more natural to write a direct formula for the continuum case). We can now proceed recursively, and define $H_N = H_{N-1} - \alpha_N \delta(x, a_N)$ and thus find the Green's function [40, 41]

$$G_N(x, y | E) = G_{N-1}(x, y | E) + G_{N-1}(x, a_N | E) \Phi^{-1}(E) G_{N-1}(a_N, y | E), \quad (4.1)$$

with $\Phi(E) = \frac{1}{\alpha_N} - G_{N-1}(a_N, a_N | E)$. By this recursive argument, the new eigenvalues E_k^{*N} are in between the eigenvalues $E_{k-1}^{*(N-1)}$ and $E_k^{*(N-1)}$. The resulting eigenfunction $\phi_k^{(N)}(x)$ is given by $\mathcal{N}_{N-1} G_{N-1}(x, a_N | E_k^{*N})$, note that $G_{N-1}(x, a_N | E_k^{*N}) = G_0(x, a_N | E_k^{*N}) + \sum_{i,j=1}^{N-1} G_0(x, a_i | E_k^{*N}) \Phi_{ij}^{-1}(E_k^{*N}) G_0(a_j, a_N | E_k^{*N})$, where $\Phi_{ij}(E) = \frac{1}{\alpha_i} \delta_{ij} - G_0(a_i, a_j | E)$ is the matrix formed by the point centers a_1, \dots, a_{N-1} .

Incidentally, the above derivation *does not make use of the finiteness of the matrix* Φ , even in case the matrix $\Phi(E)$ requires a renormalization, our derivation remains valid.

4.2 Delta Interaction Supported on Curves in Plane

Note that none of the derivations actually rely on the interaction being concentrated at a point, we can generalize to the curve case easily. In fact, by extending the above discussion, we can accommodate multiple

non-intersecting curves and points cases easily. To make the discussion simpler we consider a single rectifiable curve Γ in the plane first, and assume that the spectrum of H_0 consists of only a discrete part. The Hamiltonian is formally written as

$$H = H_0 - \alpha \delta(\cdot - \Gamma) . \quad (4.2)$$

There are various ways to define the above Hamiltonian in a mathematically rigorous way. One way is to interpret the above formal interaction by the quadratic form $\int_{\mathbb{R}^2} |\nabla \psi|^2 d^2x - \lambda \int_{\Gamma} |\psi|^2 ds$, and then prove that there is an associated self-adjoint Hamiltonian [42, 43, 44]. Other ways are to impose the continuity and the jump discontinuity conditions at Γ , (see Remark 4.1 in [42] and [45]) or use scaled potentials [46], or direct construction of the resolvent operator [42, 47]. The physical motivation for studying such Hamiltonians is to give a realistic model for trapped electrons due to interfaces between two different semiconductor materials, which are known as leaky graphs, curves or surfaces in the literature [48].

In this section, we consider rank one modification (perturbations) of H_0 in the sense described in [18] and the Hamiltonian is formally given by

$$H_0 - \alpha |\Gamma\rangle\langle\Gamma| , \quad (4.3)$$

where $H_0 = -\frac{\hbar^2}{2m}\Delta + V$ and we have used the Dirac notation for the inner products, and the Dirac delta function δ_{Γ} supported by the curve Γ with length L is defined by their action on test functions ψ [4]

$$\langle\delta_{\Gamma}|\psi\rangle = \langle\Gamma|\psi\rangle := \frac{1}{L(\Gamma)} \int_{\Gamma} \psi ds , \quad (4.4)$$

where ds is the integration element over the curve Γ and $|\Gamma\rangle\langle\Gamma|$ written in Dirac's bra-ket notation is $\langle\delta_{\Gamma}, \cdot\rangle\delta_{\Gamma}$.

It is well known that the resolvent of free Hamiltonians modified by delta interaction supported on a curve can be expressed by some explicit formulae involving the resolvent of the free Hamiltonian, and they are known as Krein's formula in the literature [17, 18, 48]. Instead of free Hamiltonian we have here a general Schrödinger operator but the formula for the resolvent would be exactly the same as before. Hence, we obtain

$$R(E) = R_0(E) + \frac{1}{\Phi(E)} R_0(E)|\Gamma\rangle\langle\Gamma|R_0(E) , \quad (4.5)$$

where $\Phi(E) = \frac{1}{\alpha} - G_0(\Gamma, \Gamma|E)$ and $G_0(\Gamma, \Gamma|E) = \frac{1}{L^2} \iint_{\Gamma \times \Gamma} G_0(\gamma(s), \gamma(s')) ds ds'$. This can be expressed in terms of the Green's functions as

$$G(x, y|E) = G_0(x, y|E) + \frac{1}{\Phi(E)} G_0(x, \Gamma|E) G_0(\Gamma, y|E) . \quad (4.6)$$

Here $G_0(x, \Gamma|E) = \frac{1}{L} \int_{\Gamma} G_0(x, \gamma(s)) ds$. A generalization of such delta interactions supported on curves embedded in manifolds is studied in [49].

Let us define $g(x, y|E) = \sum_{n \neq k} \phi_n(x) \phi_n(y) (E - E_n)^{-1}$ as well as projections onto the curve as $\langle x|g(E)|\Gamma\rangle = \sum_{n \neq k} \phi_n(x) \langle\Gamma|\phi_n\rangle (E - E_n)^{-1}$. For clarity we only assume a discrete spectrum, since generalizing to continuous spectrum is not difficult. These are holomorphic functions of E in a sufficiently small neighborhood of E_k . By following the same steps as we have done before, and using the expansion of the Green's function we find

$$\begin{aligned} G(x, y|E) &= g(x, y|E) + \frac{\phi_k(x)\phi_k(y)}{E - E_k} + \frac{\frac{\phi_k(x)\langle\phi_k|\Gamma\rangle}{E - E_k} \frac{\langle\Gamma|\phi_k\rangle\phi_k(y)}{E - E_k}}{\frac{1}{\alpha} - \langle\Gamma|g(E)|\Gamma\rangle - |\langle\Gamma|\phi_k\rangle|^2 (E - E_k)^{-1}} \\ &+ \frac{\langle x|g(E)|\Gamma\rangle \frac{\langle\Gamma|\phi_k\rangle\phi_k(y)}{E - E_k}}{\frac{1}{\alpha} - \langle\Gamma|g(E)|\Gamma\rangle - |\langle\Gamma|\phi_k\rangle|^2 (E - E_k)^{-1}} + \frac{\frac{\phi_k(x)\langle\phi_k|\Gamma\rangle}{E - E_k} \langle\Gamma|g(E)|y\rangle}{\frac{1}{\alpha} - \langle\Gamma|g(E)|\Gamma\rangle - |\langle\Gamma|\phi_k\rangle|^2 (E - E_k)^{-1}} \\ &+ \frac{\langle x|g(E)|\Gamma\rangle \langle\Gamma|g(E)|y\rangle}{\frac{1}{\alpha} - \langle\Gamma|g(E)|\Gamma\rangle - |\langle\Gamma|\phi_k\rangle|^2 (E - E_k)^{-1}} . \end{aligned} \quad (4.7)$$

Here we have assumed that there is a continuous representative for each eigenvector, so that the expression $\langle\Gamma|\phi_k\rangle = \frac{1}{L} \int_{\Gamma} \phi_k(\gamma(s)) ds$ is well defined, this is true even for the generalized eigenvectors of a Laplacian modified by some potential which satisfies some conditions [6] (indeed for the free Hamiltonian, the domain of H_0 is the Sobolev space and these expressions for arbitrary vectors in the domain are always well-defined by the Sobolev embedding theorem [50]). Just as before, multiplying by $(E - E_k)$ the numerators and denominators, and expanding around E_k , we see the cancellation of the pole at E_k , as long as $\langle\Gamma|\phi_k\rangle \neq 0$.

In this case we have the new wave functions given by

$$\psi_k(x) = \mathcal{N}G_0(x, \Gamma|E_k^*) = \mathcal{N} \int_{\Gamma} G_0(x, \gamma(s)|E_k^*) ds . \quad (4.8)$$

Eigenvalues and eigenvectors, for multiple curves and points are now constructed by iterating the above set of arguments for each additional rank one perturbation.

Remark 4.1. *In all our discussions, it is essential to have elliptic operators with some special properties (typically a summability condition on the potentials) to allow for generalized eigenvectors with certain regularity properties, as we evaluate them at a point or integrate them over a curve.*

4.3 A Particle in a Compact Manifold under the Influence of a δ Interaction

Another possibility for H_0 could correspond to a system where a particle is intrinsically moving on a two or three dimensional compact and connected manifold M with the metric structure g :

$$(H_0\psi)(x) = -\frac{\hbar^2}{2m} \left(\frac{1}{\sqrt{\det g}} \sum_{i,j=1}^d \frac{\partial}{\partial x^i} \left(\sqrt{\det g} g^{ij} \frac{\partial \psi(x)}{\partial x^j} \right) \right) , \quad (4.9)$$

where $x = (x^1, \dots, x^d)$ are the local coordinates, and g^{ij} are the components of inverse of the metric g . Then it is well known [51, 52]) that there exists a complete orthonormal system of C^∞ eigenfunctions $\{\phi_n\}_{n=0}^\infty$ in $L^2(M)$ and the spectrum $\sigma(H_0) = \{E_n\} = \{0 = E_0 \leq E_1 \leq E_2 \leq \dots\}$, with E_n tending to infinity as $n \rightarrow \infty$ and each eigenvalue has finite multiplicity: $H_0\phi_n = E_n\phi_n$. Some eigenvalues are repeated according to their multiplicity. The multiplicity of the first eigenvalue $E_0 = 0$ is one and the corresponding eigenfunction is constant. This result is also valid for Neumann, Dirichlet and mixed eigenvalue problems except for $E_0 > 0$, provided that the appropriate boundary condition is imposed.

When we add a point like delta interaction on such Schrödinger operator H_0 , we have physically a particle moving intrinsically in a two dimensional compact manifold \mathcal{M} and interacting with delta interaction located at some point a in \mathcal{M} . This problem requires renormalization due to the short distance singular behaviour of the free Green's function on a manifold, which can be similarly seen from the short time expansion of the heat kernel $K_t(a, a) \sim 1/t$ as $t \rightarrow 0^+$ and the formula (3.1) for Riemannian manifolds [53]. This has been discussed in our previous works [54, 55, 56, 57] for finitely many delta centers. Then, the Green's function can be similarly expressed as

$$G(x, y|E) = G_0(x, y|E) + \frac{G_0(x, a|E)G_0(a, y|E)}{\frac{1}{\alpha_R} - \sum_{n=0}^\infty \frac{|\phi_n(a)|^2(E+\mu^2)}{(E_n-E)(E_n+\mu^2)}} , \quad (4.10)$$

for any points x, y and a in \mathcal{M} . Then, all the results that we have obtained for previous problems follow easily.

4.4 A Semirelativistic Model on a Two Dimensional Compact Manifold

In our previous work [58], a possible relativistic model of delta interactions on a two dimensional compact Riemannian manifold (\mathcal{M}, g) is proposed, in a second quantized language the model Hamiltonian is written as

$$H = \int_{\mathcal{M}} d_g^2x : \phi^\dagger(-\nabla_g^2 + m^2)\phi + \pi^2 : - \alpha \phi^{(-)}(a)\phi^{(+)}(a) , \quad (4.11)$$

where $d_g^2x = \sqrt{\det g} dx_1 dx_2$ and (x_1, x_2) is the local coordinates. The positive and the negative frequency part of the bosonic field ϕ is denoted by $\phi^{(-)}$ and $\phi^{(+)}$, respectively. They can be expanded in terms of the creation and annihilation operators indexed by σ , that is,

$$\phi^{(+)}(x) = \sum_{\sigma} \frac{f_{\sigma}(x)}{\sqrt{\omega_{\sigma}}} a_{\sigma} , \quad (4.12)$$

$$\phi^{(-)}(x) = \sum_{\sigma} \frac{f_{\sigma}^*(x)}{\sqrt{\omega_{\sigma}}} a_{\sigma}^{\dagger} \quad (4.13)$$

where $f_{\sigma}(x)$ are the eigenfunctions of the Laplace Beltrami operator $-\nabla_g^2 = \frac{1}{\sqrt{\det g}} \sum_{i,j=1}^2 \frac{\partial}{\partial x^i} (g^{ij} \sqrt{\det g} \frac{\partial}{\partial x^j})$:

$$-\nabla_g^2 f_{\sigma}(x) = \lambda(\sigma) f_{\sigma}(x) , \quad (4.14)$$

and $\omega_\sigma = \sqrt{\lambda(\sigma) + m^2}$. Here the index σ plays the same role as momentum in flat spaces. If we restrict our model to a one particle sector

$$|\psi\rangle = \sum_\sigma \psi_\sigma \frac{a_\sigma^\dagger}{\sqrt{\omega_\sigma}} |0\rangle, \quad (4.15)$$

and following the same steps as in [58] we will find the Green's function restricted to one-particle sector. To find the Green's function, we need to solve in the eigenfunction basis,

$$(H - E)|\psi\rangle = |\chi\rangle, \quad (4.16)$$

for a given state $|\chi\rangle$. As a result, the one-particle sector Green's function can be found by solving,

$$(\sqrt{\sigma^2 + m^2} - E)\psi_\sigma - \alpha \sum_\nu \frac{f_\sigma^*(a)}{\sqrt{\omega_\nu}} \frac{f_\nu(a)}{\sqrt{\omega_\nu}} \psi_\nu = \chi_\sigma. \quad (4.17)$$

The solution ψ_σ of this equation is given by

$$\psi_\sigma = \frac{1}{(\omega_\sigma - E)} \chi_\sigma + \frac{1}{(\omega_\sigma - E)} f_\sigma^*(a) \left[\frac{1}{\alpha} - \sum_\tau \frac{|f_\tau(a)|^2}{\omega_\tau(\omega_\tau - E)} \right]^{-1} \sum_\nu \frac{f_\nu(a)}{\omega_\nu} \frac{1}{(\omega_\nu - E)} \chi_\nu. \quad (4.18)$$

To read the Green's function, notice that

$$\begin{aligned} \sum_\sigma f_\sigma(x) \frac{\psi_\sigma}{\sqrt{\omega_\sigma}} &= \psi(x) = \sum_\sigma \int_{\mathcal{M}} d_g^2 y \frac{f_\sigma(x) f_\sigma^*(y)}{(\omega_\sigma - E)} \chi(y) \\ &+ \sum_\sigma \frac{f_\sigma(x)}{(\omega_\sigma - E)} \frac{f_\sigma^*(a)}{\sqrt{\omega_\sigma}} \left[\frac{1}{\alpha} - \sum_\tau \frac{|f_\tau(a)|^2}{\omega_\tau(\omega_\tau - E)} \right]^{-1} \sum_\nu \int_{\mathcal{M}} d_g^2 y \frac{f_\nu(a)}{\sqrt{\omega_\nu}} \frac{f_\nu^*(y)}{(\omega_\nu - E)} \chi(y), \end{aligned} \quad (4.19)$$

where we have used $\chi_\sigma = \sqrt{\omega_\sigma} \int_{\mathcal{M}} d_g^2 x \chi(x) f_\sigma^*(x)$ and orthogonality of eigenfunctions f_σ . We can then write this as an equation,

$$\psi(x) = \int_{\mathcal{M}} d_g^2 y G_0(x, y|E) \chi(y) + \tilde{G}_0(x, a|E) \Phi(E)^{-1} \int dy \tilde{G}_0(a, y|E) \chi(y), \quad (4.20)$$

where

$$G_0(x, y|E) = \sum_\sigma \frac{f_\sigma(x) f_\sigma^*(y)}{(\omega_\sigma - E)} \quad (4.21)$$

$$\tilde{G}_0(x, y|E) = \sum_\sigma \frac{f_\sigma(x)}{(\omega_\sigma - E)} \frac{f_\sigma^*(y)}{\sqrt{\omega_\sigma}} \quad (4.22)$$

$$\Phi(E) = \frac{1}{\alpha} - \sum_\tau \frac{|f_\tau(a)|^2}{\omega_\tau(\omega_\tau - E)}. \quad (4.23)$$

Then we can find the Green's function as

$$G(x, y|E) = G_0(x, y) + \tilde{G}_0(x, a|E) \Phi^{-1}(E) \tilde{G}_0(a, y|E). \quad (4.24)$$

Note that to demonstrate the cancellation of original poles, we need to carefully look at the analytic structure of $G(x, y)$.

$$G(x, y) = \sum_\sigma \frac{f_\sigma(x) f_\sigma^*(y)}{(\omega_\sigma - E)} + \sum_\sigma \frac{f_\sigma(x)}{(\omega_\sigma - E)} \frac{f_\sigma^*(a)}{\sqrt{\omega_\sigma}} \Phi^{-1}(E) \sum_\nu \frac{f_\nu(a)}{\sqrt{\omega_\nu}} \frac{f_\nu^*(y)}{(\omega_\nu - E)} \quad (4.25)$$

Indeed, the sum in the definition of Φ is divergent. This can be seen by following the same line of arguments discussed in [58]. We will summarize it for the sake of completeness. First we rewrite the expression $\frac{1}{\omega_\tau(\omega_\tau - E)}$ as

$$\frac{1}{\omega_\tau(\omega_\tau - E)} = -\frac{1}{E} \left(\frac{1}{\omega_\tau} - \frac{1}{\omega_\tau - E} \right) = -\frac{1}{E} \int_0^\infty e^{-s\omega_\tau} (1 - e^{sE}) ds, \quad (4.26)$$

where $\text{Re}(\omega_\tau - E) > 0$ and the result is assumed to be analytically continued onto the complex plane. Using the subordination identity

$$e^{sA} = \frac{s}{2\sqrt{\pi}} \int_0^\infty e^{s^2/(4u) - uA^2} \frac{du}{u^{3/2}}, \quad (4.27)$$

and the eigenfunction expansion of the heat kernel defined on Riemannian manifolds [51]

$$K_u(x, y) = \sum_{\sigma} e^{-u\lambda(\sigma)} f_{\sigma}(x) f_{\sigma}^*(y), \quad (4.28)$$

we obtain

$$\sum_{\tau} \frac{|f_{\tau}(a)|^2}{\omega_{\tau}(\omega_{\tau} - E)} = -\frac{1}{E} \int_0^{\infty} \left(\int_0^{\infty} K_u(a, a) e^{-um^2 + s^2/(4u)} \frac{du}{u^{3/2}} \right) \frac{(1 - e^{sE})s}{2\sqrt{\pi}} ds. \quad (4.29)$$

Using integration by parts and scaling the integration variable, we find

$$\sum_{\tau} \frac{|f_{\tau}(a)|^2}{\omega_{\tau}(\omega_{\tau} - E)} = \frac{1}{\sqrt{\pi}} \int_0^{\infty} \left(\int_0^{\infty} K_u(a, a) e^{-um^2 + sE\sqrt{u}} du \right) e^{-s^2/4} ds. \quad (4.30)$$

Due to the short “time” asymptotic behavior of the diagonal heat kernel $K_u(a, a) \sim \frac{1}{u}$ for two dimensional Riemannian manifolds, the integral over u is divergent. For this reason, we apply the idea of renormalization and introduce a short “time” cut off ϵ in the lower limit of u integral. This corresponds to the sum over τ up to a cut off N . Then we choose the coupling constant depending on this cut off in such a way that

$$\frac{1}{\alpha(N)} = \frac{1}{\alpha_R} + \sum_{\tau}^N \frac{|f_{\tau}(a)|^2}{\omega_{\tau}(\omega_{\tau} + \mu^2)}, \quad (4.31)$$

where $-\mu^2$ is the experimentally measured bound state energy of the particle in the presence of single Dirac delta center. To this end we remove the cut off by sending $N \rightarrow \infty$ in Φ and get the renormalized Φ

$$\begin{aligned} \Phi_R &= \frac{1}{\alpha_R} + \sum_{\tau} \left[\frac{|f_{\tau}(a)|^2}{\omega_{\tau}(\omega_{\tau} + \mu^2)} - \frac{|f_{\tau}(a)|^2}{\omega_{\tau}(\omega_{\tau} - E)} \right] \\ &= \frac{1}{\alpha_R} + \sum_{\tau} |f_{\tau}(a)|^2 \frac{(-E - \mu^2)}{\omega_{\tau}(\omega_{\tau} + \mu^2)(\omega_{\tau} - E)} \\ &= \frac{1}{\alpha_R} - (E + \mu^2) \sum_{\tau} \frac{|f_{\tau}(a)|^2}{\omega_{\tau}(\omega_{\tau} + \mu^2)(\omega_{\tau} - E)}. \end{aligned} \quad (4.32)$$

We take E around E_{ν} and isolate this particular pole in our expressions,

$$\begin{aligned} &\sum_{\sigma \neq \nu} \frac{f_{\sigma}(x) f_{\sigma}^*(y)}{(\omega_{\sigma} - E)} + \frac{f_{\nu}(x) f_{\nu}^*(y)}{(\omega_{\nu} - E)} + \left(\sum_{\sigma \neq \nu} \frac{f_{\sigma}(x)}{(\omega_{\sigma} - E)} \frac{f_{\sigma}^*(a)}{\sqrt{\omega_{\sigma}}} + \frac{f_{\nu}(x)}{(\omega_{\nu} - E)} \frac{f_{\nu}^*(a)}{\sqrt{\omega_{\nu}}} \right) \\ &\times \frac{1}{\frac{1}{\alpha_R} - (E + \mu^2) \left[\sum_{\tau \neq \nu} \frac{|f_{\tau}(a)|^2}{\omega_{\tau}(\omega_{\tau} + \mu^2)(\omega_{\tau} - E)} + \frac{|f_{\nu}(a)|^2}{\omega_{\nu}(\omega_{\nu} + \mu^2)(\omega_{\nu} - E)} \right]} \left(\sum_{\alpha \neq \nu} \frac{f_{\alpha}(a)}{\sqrt{\omega_{\alpha}}} \frac{f_{\alpha}^*(y)}{(\omega_{\alpha} - E)} + \frac{f_{\nu}(a)}{\sqrt{\omega_{\nu}}} \frac{f_{\nu}^*(y)}{(\omega_{\nu} - E)} \right). \end{aligned} \quad (4.33)$$

Note that here the singular part comes from,

$$\frac{f_{\nu}(x) f_{\nu}^*(y)}{(\omega_{\nu} - E)} - \frac{|f_{\nu}(a)|^2 f_{\nu}(x) f_{\nu}^*(y)}{\omega_{\nu}(\omega_{\nu} - E)^2 \left[(E + \mu^2) \frac{|f_{\nu}(a)|^2}{\omega_{\nu}(\omega_{\nu} - E)(\omega_{\nu} + \mu^2)} + \text{regular terms} \right]} + \text{regular terms}. \quad (4.34)$$

We now use $E + \mu^2 = E - \omega_{\nu} + \omega_{\nu} + \mu^2$ in the numerator and reduce this to

$$\frac{f_{\nu}(x) f_{\nu}^*(y)}{(\omega_{\nu} - E)} - \frac{|f_{\nu}(a)|^2 f_{\nu}(x) f_{\nu}^*(y)}{\omega_{\nu}(\omega_{\nu} - E)^2 \left[\frac{|f_{\nu}(a)|^2}{\omega_{\nu}(\omega_{\nu} - E)} + \text{reg} \right]} + \text{regular terms}. \quad (4.35)$$

Following our previous arguments, it is seen that the pole at ω_{ν} is now cancelled.

Final Remarks

In this work, we have studied the spectral properties of the Schrödinger operators H_0 with δ interactions, where H_0 satisfies some mild conditions, which are usually assumed to hold in most of the quantum systems. In contrast to the typical works in the literature, we work out the pole structure of G and explicitly show that poles of the initial Green function G_0 are removed (as long as the original eigenfunction does not vanish

at the location of the delta function) from G , through the use of eigenfunction expansions. We show that new bound state energies can be found by solving equation $\Phi(E) = 0$, where Φ is explicitly defined in terms of the diagonal Green's function if the center of the delta function is not located at one of the nodes of the initial eigenfunction ϕ_k). These results are established in one dimension for point like delta interactions and the reflectionless potential for H_0 is studied as an example. A different kind of perturbative approach in finding eigenvalues and eigenfunctions of the full system is presented up to second order in a heuristic way. These ideas are then extended to the case, where the renormalization procedure is needed. In that case, we obtained similar results, except that the perturbative calculations differ at the second order of the coupling constant of delta interaction. Finally, we summarize some possible further extensions of our results to the more general delta interactions (supported on curve) and multi-center case, and to the case in which a particle is moving on a compact two dimensional manifold, and to a semi-relativistic model on a compact manifold.

Data availability statement

No new data were created or analysed in this study.

Acknowledgments

F. Erman would like to thank H. Uncu for useful discussions on the spectrum of harmonic oscillator modified by delta interaction, and A. Bohm and M. Gadella for giving helpful insights on the Rigged Hilbert spaces and eigenfunction expansions. O. T. Turgut would like to thank A. Michelangeli for various discussions on singular interactions and J. Hoppe for discussions and encouragement on this project.

Appendix

We verify that Green's function is finite apart from the obvious poles we claim, that is, we would like to prove that

$$\lim_{N \rightarrow \infty} \sum_{n=0}^N \frac{|\phi_n(a)|^2}{(E_n - E)(E_n + \mu^2)} < \infty. \quad (4.36)$$

On a compact manifold, the Laplace-Beltrami operator $-\nabla_g^2$ has a discrete spectrum $0 \leq E_0 \leq E_1 \leq E_2 \dots \leq E_n \rightarrow \infty$. For simplicity, let us first consider the case where $E \in \mathbb{R}^+$. We then choose M sufficiently large that $E_M > 2\left(E + \frac{\mu^2}{2}\right)$ and assume that $E_n \geq \lambda_M$. Then, it is easy to see that for $n \geq M$ we have

$$\frac{|\phi_n(a)|^2}{(E_n - E)(E_n + \mu^2)} < 2 \frac{|\phi_n(a)|^2}{(E_n + \mu^2)^2}. \quad (4.37)$$

This implies that

$$\sum_{n \geq M}^N \frac{|\phi_n(a)|^2}{(E_n - E)(E_n + \mu^2)} < 2 \sum_{n=0}^N \frac{|\phi_n(a)|^2}{(E_n + \mu^2)^2}. \quad (4.38)$$

The right hand side can be expressed as

$$\sum_{n=0}^N \frac{|\phi_n(a)|^2}{(E_n + \mu^2)^2} = \sum_{n=0}^N \int_0^\infty dt t |\phi_n(a)|^2 e^{-(E_n + \mu^2)t}, \quad (4.39)$$

and using the eigenfunction expansion of the heat kernel (4.28) we obtain in the limit $N \rightarrow \infty$

$$\sum_{n=0}^\infty \frac{|\phi_n(a)|^2}{(E_n + \mu^2)^2} = \int_0^\infty dt t K_t(a, a) e^{-\mu^2 t}. \quad (4.40)$$

The upper bounds of the diagonal part of the heat kernel on compact manifolds is given by (see [39])

$$K_t(a, a) < \left(\frac{1}{V(\mathcal{M})} + \frac{C}{t} \right), \quad (4.41)$$

where $V(\mathcal{M})$ is the volume of \mathcal{M} and C is a constant depending only on the geometric properties of \mathcal{M} . This upper bound shows that $\sum_{n=0}^{\infty} \frac{|\phi_n(a)|^2}{(E_n - E)(E_n + \mu^2)}$ is convergent.

For the complex values of $E = E_R + iE_I$ with $E_I \neq 0$, we can also show that the above series is absolutely convergent since

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{|\phi_n(a)|^2}{|(E_n - E_R - iE_I)(E_n + \mu^2)|} &= \sum_{n=0}^{\infty} \frac{|\phi_n(a)|^2}{\sqrt{(E_n - E_R)^2 + E_I^2}(E_n + \mu^2)} \\ &\leq \sqrt{2} \sum_{n=0}^{\infty} \frac{|\phi_n(a)|^2}{(|E_n - E_R| + |E_I|)(E_n + \mu^2)} \\ &\leq \sum_{n>M}^{\infty} \frac{\sqrt{2}|\phi_n(a)|^2}{(E_n + \mu^2)(|E_n - E_R| + |E_I|)} + \frac{\sqrt{2}}{|E_I|} \sum_{n \leq M} \frac{|\phi_n(a)|^2}{(E_n + \mu^2)} < \infty \end{aligned} \tag{4.42}$$

choosing an M such that for $n > M$ we have $E_n > 2|E_R|$ we see that all the terms become finite. Hence in all possible cases we have finite expressions.

References

- [1] A. Bohm, *The Rigged Hilbert Space and Quantum Mechanics*, Lecture Notes in Physics, Springer (1978).
- [2] A. Bohm, M. Gadella, *Dirac Kets, Gamow Vectors and Gel'fand Triplets*, Lecture Notes in Physics; Springer **348** (1989).
- [3] F. A. Berezin, and M. Shubin, *The Schrödinger Equation*, **66** Springer (2012).
- [4] W. Appel, *Mathematics for Physics and Physicists*, Princeton University Press (2007).
- [5] I. M. Berezanskii, *Expansions in Eigenfunctions of Selfadjoint Operators*, American Mathematical Society (1968).
- [6] A. G. Belyi, V. F. Kovalenko, and Y. A. Semenov, On the continuity of generalized eigenfunctions of the Schrödinger operator, *Reports on Mathematical Physics*, **12** 3, 307-310 (1977).
- [7] A. Galindo, and P. Pascual, *Quantum Mechanics I*, Springer (2012).
- [8] H. Weyl, Über gewöhnliche Differentialgleichungen mit Singularitäten und die zugehörigen Entwicklungen willkürlicher Funktionen, *Math. Ann.* **68** 220-269 (1910).
- [9] E. C. Titchmarsh, *Eigenfunction Expansions Associated with Second-order Differential Equations, Part I*, Oxford (1946).
- [10] K. Kodaira, The Eigenvalue Problem for Ordinary Differential Equations of the Second Order and Heisenberg's Theory of S-Matrices, *Amer. J. Math.* **71** 4, 921-945 (1949).
- [11] T. Poerschke, G. Stolz, and J. Weidmann, Expansions in Generalized Eigenfunctions of Selfadjoint Operators, *Mathematische Zeitschrift*, **202** 397-408 (1989).
- [12] C. Grosche, Path integrals for potential problems with δ -function perturbation, *Journal of Physics A: Mathematical and General* **23** 22 5205 (1990).
- [13] S. Fassari, G. Inglese, *Helv. Phys. Acta* **67** 6, 650-659 (1994).
- [14] S. Fassari, G. Inglese, Spectroscopy of a three dimensional isotropic harmonic oscillator with a δ type perturbation, *Helvetica Physica Acta*, **69** (1996).
- [15] W. Janke, and B. K. Cheng, Statistical properties of a harmonic plus a delta-potential, *Physics Letters A* **129** 3, 140-144 (1988).
- [16] H. Uncu, D. Tarhan, E. Demiralp, Ö. E. Müstecaphoğlu, Bose-Einstein condensate in a harmonic trap decorated with Dirac δ functions, *Physical Review A*, **76** 1, 013618 (2007).
- [17] S. Albeverio, F. Gesztesy, and R. Hoegh-Krohn, and H. Holden, *Solvable Models in Quantum Mechanics*, 2nd ed. American Mathematical Society (2004).

- [18] S. Albeverio, P. Kurasov, *Singular Perturbation of Differential Operators*, Cambridge University Press (2000).
- [19] K. Huang, *Quarks, Leptons and Gauge Fields*, World Scientific, Singapore (1982).
- [20] R. Jackiw, *Delta-Function Potentials in Two- and Three-Dimensional Quantum Mechanics*, M. A. B. Bég Memorial Volume, World Scientific, Singapore (1991).
- [21] P. Gosdzinsky, and R. Tarrach, Learning quantum field theory from elementary quantum mechanics, *Am. J. Phys.*, **59** 70 (1991).
- [22] L. R. Mead, J. Godines, An analytical example of renormalization in two-dimensional quantum mechanics, *Am. J. Phys.*, **59** 935 (1991).
- [23] C. Manuel, R. Tarrach, Perturbative renormalization in quantum mechanics, *Phys. Lett. B*, **328** 113 (1994).
- [24] C. Grosche, Path integrals for two-and three-dimensional δ -function perturbations, *Annalen der Physik*, **506** 4, 283-312 (1994).
- [25] S. Albeverio, S. Fassari, and F. Rinaldi, Spectral properties of a symmetric three-dimensional quantum dot with a pair of identical attractive δ impurities symmetrically situated around the origin, *Nanosystems: Physics, Chemistry, Mathematics*, **7** 2 268-289 (2016).
- [26] S. Albeverio, S. Fassari, and F. Rinaldi, Spectral properties of a symmetric three-dimensional quantum dot with a pair of identical attractive δ impurities symmetrically situated around the origin II, *Nanosystems: Physics, Chemistry, Mathematics*, **5** 5, 803-815 (2016).
- [27] S. Fassari, M. Gadella, M. L. Glasser, L. M. Nieto, F. Rinaldi, Spectral properties of the two-dimensional Schrödinger Hamiltonian with various solvable confinements in the presence of a central point perturbation, *Physica Scripta*, **94** 5, 055202 (2019).
- [28] L. D. Landau, and E. M. Lifshitz, *Quantum Mechanics: Non-Relativistic Theory*, 3rd ed., Butterworth-Heinemann, **3** (1981).
- [29] M. Reed, and B. Simon, *Methods of Modern Mathematical Physics III: Scattering Theory*, Academic Press, New York (1979).
- [30] M. Gallone, and A. Michelangeli, *Self-adjoint Extension Schemes and Modern Applications to Quantum Hamiltonians*, Springer International Publishing, 1st ed. (2023)
- [31] J. Dimock, S. G. Rajeev, Multiparticle Schrödinger operators with point interactions in the plane, *J. Phys. A: Math. Gen.*, **37** 9157 (2004).
- [32] M. Reed, and B. Simon, *Methods of Modern Mathematical Physics, Volume IV, Analysis of Operators*, Academic Press, New York (1978).
- [33] A. Galindo, and P. Pascual, *Quantum Mechanics II*, Springer (2012).
- [34] T. Kato, *Perturbation Theory for Linear Operators*, **132** Springer (2013).
- [35] J. A. Murdock, *Perturbations, Theory and Applications*, John Wiley and Sons (1991).
- [36] J. Sólyom, *Fundamentals of the Physics of Solids: Volume II: Electronic Properties* **2** Appendix G, Springer (2008).
- [37] D. J. Griffiths, and D. F. Schroeter, *Introduction to Quantum Mechanics*, Cambridge university press (2018).
- [38] P. B. Gilkey, *Invariance Theory, the Heat Equation, and the Atiyah-Singer Index Theorem*, 2nd edition, CRC Press, Boca Raton (1995).
- [39] F. Erman, and O. T. Turgut, Point Interactions in two and three dimensional Riemannian Manifolds, *Journal of Physics A: Mathematical and Theoretical*, **43** 33, 335204 (2010).
- [40] J. Besprosvany, *Phys. Rev. B*, **63** 233108 (2001).
- [41] F. Erman, Recursion formula for the Green's function of a Hamiltonian for several types of Dirac delta-function potentials in curved spaces, *Turkish Journal of Physics* **40** 3 316-323 (2016).

- [42] J. F. Brasche, P. Exner, Yu. A. Kuperin, and P. Seba, Schrödinger operators with singular interactions, *Journal of mathematical analysis and applications*, **184** 1, 112-139 (1994).
- [43] P. Exner and K. Yoshitomi, Asymptotics of eigenvalues of the Schrödinger operator with a strong δ interaction on a loop, *Journal of Geometry and Physics*, **41** 4 344-358 (2002).
- [44] S. Kondej, Spectral asymptotics induced by approaching and diverging planar circles, *Functional Analysis and Operator Theory for Quantum Physics: The Pavel Exner Anniversary Volume* edited by J. Dittrich, H. Kovarik, A. Laptev, European Mathematical Society, (2017).
- [45] P. Exner and M. Tater, Spectra of soft ring graphs, *Waves in Random Media*, **14** 1, S47, (2003).
- [46] P. Exner and T. Ichinose, Geometrically induced spectrum in curved leaky wires, *Journal of Physics A: Mathematical and General*, **34** 7, 1439 (2001).
- [47] A. Posilicano, A Krein-like formula for singular perturbations of self-adjoint operators and applications, *Journal of Functional Analysis*, **183** 1, 109-147 (2001).
- [48] P. Exner and H. Kovarik, *Quantum Waveguides*, Springer (2015).
- [49] B. T. Kaynak, and O. T. Turgut, Singular interactions supported by embedded curves, *J. Phys. A: Math. Theor.* **45** 265202 (2012).
- [50] D. Haroske, and H. Triebel, *Distributions, Sobolev Spaces, Elliptic Equations*, European Mathematical Society, **4** (2007).
- [51] S. Rosenberg, *The Laplacian on Riemannian Manifold*, Cambridge University Press, Cambridge (1998).
- [52] I. Chavel, *Eigenvalues in Riemannian Geometry*, Pure and Applied Mathematics **115** Academic Press, Orlando (1984).
- [53] A. Grigoryan, *Heat Kernel and Analysis on Manifolds*, AMS/IP Studies in Advanced Mathematics, ed. S.-T. Yau, **47** (American Mathematical Society, Rhode Island, (2009).
- [54] B. İ. Altunkaynak, F. Erman F, and O. T. Turgut, Finitely Many Dirac-Delta Interactions on Riemannian Manifolds, *Journal of Mathematical Physics*, **47** 8, 082110-1-082110-23 (2006).
- [55] F. Erman, and O. T. Turgut, Point interactions in two-and three-dimensional Riemannian manifolds, *Journal of Physics A: Mathematical and Theoretical* **43** 33 335204 (2010).
- [56] F. Erman, On the number of bound states of point interactions on hyperbolic manifolds, *International Journal of Geometric Methods in Modern Physics*, **14** 01, 1750011, (2017).
- [57] Ç. Doğan, F. Erman, and O. T. Turgut, Existence of Hamiltonians for some singular interactions on manifolds, *J. Math. Phys.*, **53** 4, 043511 (2012).
- [58] Ç. Doğan and O. T. Turgut, Renormalized interaction of relativistic bosons with delta function potentials, *J. Math. Phys.*, **51**, 082305 (2010).