

$SL(2, \mathbf{Z})$ MODULAR FORMS AND ANOMALY CANCELLATION FORMULAS

YONG WANG* JIANYUN GUAN

ABSTRACT. By some $SL(2, \mathbf{Z})$ modular forms introduced in [11] and [4], we get some interesting anomaly cancellation formulas. As corollaries, we get some divisibility results of index of twisted Dirac operators.

1. INTRODUCTION

In 1983, the physicists Alvarez-Gaumé and Witten [1] discovered the "miraculous cancellation" formula for gravitational anomaly which reveals a beautiful relation between the top components of the Hirzebruch \widehat{L} -form and \widehat{A} -form of a 12-dimensional smooth Riemannian manifold. Kefeng Liu [10] established higher dimensional "miraculous cancellation" formulas for $(8k + 4)$ -dimensional Riemannian manifolds by developing modular invariance properties of characteristic forms. These formulas could be used to deduce some divisibility results. In [8], [9], [3], some more general cancellation formulas that involve a complex line bundle and their applications were established. In [6], using the Eisenstein series, a more general cancellation formula was derived. In [7], Han, Liu and Zhang showed that both of the Green-Schwarz anomaly factorization formula for the gauge group $E_8 \times E_8$ and the Horava-Witten anomaly factorization formula for the gauge group E_8 could be derived through modular forms of weight 14. This answered a question of J. H. Schwarz. They also established generalizations of these factorization formulas and obtained a new Horava-Witten type factorization formula on 12-dimensional manifolds. In [5], Han, Huang, Liu and Zhang introduced a modular form of weight 14 over $SL(2, \mathbf{Z})$ and a modular form of weight 10 over $SL(2, \mathbf{Z})$ and they got some interesting anomaly cancellation formulas on 12-dimensional manifolds. In [11], a modular form of weight $2k$ for a $2k$ -dimensional spin manifold was introduced. In [4], Chen, Han and Zhang defined an integral modular form of weight $2k$ for a $4k$ -dimensional $spin^c$ manifold and an integral modular form of weight $2k$ for a $4k + 2$ -dimensional $spin^c$ manifold. **Our motivation is to prove some more anomaly cancellation formulas on some dimensional manifolds by modular forms over $SL(2, \mathbf{Z})$ introduced in [11] and [4].**

Let X be a spin manifold. Let ∇^{TX} be the associated Levi-Civita connection on TX

2010 Mathematics Subject Classification: 58C20; 57R20; 53C80.

Key words and phrases: $SL(2, \mathbf{Z})$ modular forms; anomaly cancellation formulas; divisibility of index

* *Corresponding author.*

and $R^{TX} = (\nabla^{TX})^2$ be the curvature of ∇^{TX} . Let $\widehat{A}(TX, \nabla^{TX})$ be the Hirzebruch characteristic forms defined by (cf. [12])

$$\widehat{A}(TX, \nabla^{TX}) = \det^{\frac{1}{2}} \left(\frac{\frac{\sqrt{-1}}{4\pi} R^{TX}}{\sinh(\frac{\sqrt{-1}}{4\pi} R^{TX})} \right).$$

Let $\Delta(X)$ be the spinor bundle and $\widetilde{TX} = TX - \dim X$. We have

Theorem 1.1. *When $\dim X = 8$, we have*

$$(1.1) \quad \left\{ \widehat{A}(TX) \text{ch}(\Delta(X)) \text{ch}(2\widetilde{T_C X}) + 32\widehat{A}(TX) \text{ch}(\widetilde{T_C X} + \wedge^2 \widetilde{T_C X}) \right\}^{(8)} \\ = 240 \left\{ \widehat{A}(TX) \text{ch}(\Delta(X)) + 32\widehat{A}(TX) \right\}^{(8)},$$

$$(1.2) \quad \left\{ \widehat{A}(TX) \text{ch}(\Delta(X)) \text{ch}(2\widetilde{T_C X} + \wedge^2 \widetilde{T_C X} + \widetilde{T_C X} \otimes \widetilde{T_C X} + S^2 \widetilde{T_C X}) \right. \\ \left. + 32\widehat{A}(TX) \text{ch}(\wedge^4 \widetilde{T_C X} + \wedge^2 \widetilde{T_C X} \otimes \widetilde{T_C X} + \widetilde{T_C X} \otimes \widetilde{T_C X}) + S^2 \widetilde{T_C X} + \widetilde{T_C X} \right\}^{(8)} \\ = 2160 \left\{ \widehat{A}(TX) \text{ch}(\Delta(X)) + 32\widehat{A}(TX) \right\}^{(8)}.$$

Corollary 1.2. *Let X be an 8-dimensional spin manifold without boundary, then*

$$(1.3) \quad \text{Ind}((D \otimes \Delta(X) \otimes \widetilde{T_C X})_+) \equiv 0 \pmod{8Z},$$

$$(1.4) \quad \text{Ind}((D \otimes \Delta(X) \otimes (2\widetilde{T_C X} + \wedge^2 \widetilde{T_C X} + \widetilde{T_C X} \otimes \widetilde{T_C X} + S^2 \widetilde{T_C X}))_+) \equiv 0 \pmod{16Z}.$$

Theorem 1.3. *When $\dim X = 12$, we have*

$$(1.5) \quad \left\{ \widehat{A}(TX) \text{ch}(\Delta(X)) \text{ch}(2\widetilde{T_C X}) + 128\widehat{A}(TX) \text{ch}(\widetilde{T_C X} + \wedge^2 \widetilde{T_C X}) \right\}^{(12)} \\ = -504 \left\{ \widehat{A}(TX) \text{ch}(\Delta(X)) + 128\widehat{A}(TX) \right\}^{(12)},$$

$$(1.6) \quad \left\{ \widehat{A}(TX) \text{ch}(\Delta(X)) \text{ch}(2\widetilde{T_C X} + \wedge^2 \widetilde{T_C X} + \widetilde{T_C X} \otimes \widetilde{T_C X} + S^2 \widetilde{T_C X}) \right. \\ \left. + 128\widehat{A}(TX) \text{ch}(\wedge^4 \widetilde{T_C X} + \wedge^2 \widetilde{T_C X} \otimes \widetilde{T_C X} + \widetilde{T_C X} \otimes \widetilde{T_C X}) + S^2 \widetilde{T_C X} + \widetilde{T_C X} \right\}^{(12)} \\ = -16632 \left\{ \widehat{A}(TX) \text{ch}(\Delta(X)) + 128\widehat{A}(TX) \right\}^{(12)}.$$

Corollary 1.4. *Let X be an 12-dimensional spin manifold without boundary, then*

$$(1.7) \quad \text{Ind}((D \otimes \Delta(X) \otimes \widetilde{T_C X})_+) \equiv 0 \pmod{4Z},$$

$$(1.8) \quad \text{Ind}((D \otimes \Delta(X) \otimes (2\widetilde{T_C X} + \wedge^2 \widetilde{T_C X} + \widetilde{T_C X} \otimes \widetilde{T_C X} + S^2 \widetilde{T_C X}))_+) \equiv 0 \pmod{8Z}.$$

Theorem 1.5. *When $\dim X = 16$, we have*

$$(1.9) \quad \left\{ \widehat{A}(TX) \text{ch}(\Delta(X)) \text{ch}(2\widetilde{T_C X}) + 512 \widehat{A}(TX) \text{ch}(\widetilde{T_C X} + \wedge^2 \widetilde{T_C X}) \right\}^{(16)} \\ = 480 \left\{ \widehat{A}(TX) \text{ch}(\Delta(X)) + 512 \widehat{A}(TX) \right\}^{(16)},$$

$$(1.10) \quad \left\{ \widehat{A}(TX) \text{ch}(\Delta(X)) \text{ch}(2\widetilde{T_C X} + \wedge^2 \widetilde{T_C X} + \widetilde{T_C X} \otimes \widetilde{T_C X} + S^2 \widetilde{T_C X}) \right. \\ \left. + 512 \widehat{A}(TX) \text{ch}(\wedge^4 \widetilde{T_C X} + \wedge^2 \widetilde{T_C X} \otimes \widetilde{T_C X} + \widetilde{T_C X} \otimes \widetilde{T_C X}) + S^2 \widetilde{T_C X} + \widetilde{T_C X} \right\}^{(16)} \\ = 61920 \left\{ \widehat{A}(TX) \text{ch}(\Delta(X)) + 512 \widehat{A}(TX) \right\}^{(16)}.$$

Corollary 1.6. *Let X be an 16-dimensional spin manifold without boundary, then*

$$(1.11) \quad \text{Ind}((D \otimes \Delta(X) \otimes \widetilde{T_C X})_+) \equiv 0 \pmod{16Z},$$

$$(1.12) \quad \text{Ind}((D \otimes \Delta(X) \otimes (2\widetilde{T_C X} + \wedge^2 \widetilde{T_C X} + \widetilde{T_C X} \otimes \widetilde{T_C X} + S^2 \widetilde{T_C X}))_+) \equiv 0 \pmod{32Z}.$$

Theorem 1.7. *When $\dim X = 20$, we have*

$$(1.13) \quad \left\{ \widehat{A}(TX) \text{ch}(\Delta(X)) \text{ch}(2\widetilde{T_C X}) + 2048 \widehat{A}(TX) \text{ch}(\widetilde{T_C X} + \wedge^2 \widetilde{T_C X}) \right\}^{(20)} \\ = -264 \left\{ \widehat{A}(TX) \text{ch}(\Delta(X)) + 32 \widehat{A}(TX) \right\}^{(20)},$$

$$(1.14) \quad \left\{ \widehat{A}(TX) \text{ch}(\Delta(X)) \text{ch}(2\widetilde{T_C X} + \wedge^2 \widetilde{T_C X} + \widetilde{T_C X} \otimes \widetilde{T_C X} + S^2 \widetilde{T_C X}) \right. \\ \left. + 2048 \widehat{A}(TX) \text{ch}(\wedge^4 \widetilde{T_C X} + \wedge^2 \widetilde{T_C X} \otimes \widetilde{T_C X} + \widetilde{T_C X} \otimes \widetilde{T_C X}) + S^2 \widetilde{T_C X} + \widetilde{T_C X} \right\}^{(20)} \\ = -117288 \left\{ \widehat{A}(TX) \text{ch}(\Delta(X)) + 2048 \widehat{A}(TX) \right\}^{(20)}.$$

Corollary 1.8. *Let X be an 20-dimensional spin manifold without boundary, then*

$$(1.15) \quad \text{Ind}((D \otimes \Delta(X) \otimes \widetilde{T_C X})_+) \equiv 0 \pmod{4Z},$$

$$(1.16) \quad \text{Ind}((D \otimes \Delta(X) \otimes (2\widetilde{T_C X} + \wedge^2 \widetilde{T_C X} + \widetilde{T_C X} \otimes \widetilde{T_C X} + S^2 \widetilde{T_C X}))_+) \equiv 0 \pmod{8Z}.$$

Let V be an $2m_0$ dimensional real Euclidean vector bundle with the Euclidean connection ∇^V and the curvature R^V . Denote the first Pontryagin classes of TX and V by $p_1(X)$ and $p_1(V)$.

Theorem 1.9. *When $\dim X = 8$ and $3p_1(V) = p_1(X)$ we have*

$$(1.17) \quad 240 \left\{ \widehat{A}(TX) \det^{\frac{1}{2}} \cosh\left(\frac{\sqrt{-1}}{4\pi} R^V\right) \right\}^{(8)} = \left\{ \widehat{A}(TX) \det^{\frac{1}{2}} \cosh\left(\frac{\sqrt{-1}}{4\pi} R^V\right) \cdot \text{ch}(\widetilde{TX} + 2 \wedge^2 \widetilde{V}_C - \widetilde{V}_C \otimes \widetilde{V}_C + \widetilde{V}_C) \right\}^{(8)}.$$

$$(1.18) \quad 2160 \left\{ \widehat{A}(TX) \det^{\frac{1}{2}} \cosh\left(\frac{\sqrt{-1}}{4\pi} R^V\right) \right\}^{(8)} = \left\{ \widehat{A}(TX) \det^{\frac{1}{2}} \cosh\left(\frac{\sqrt{-1}}{4\pi} R^V\right) \cdot \text{ch}(S^2 \widetilde{TX} + \widetilde{TX} + (2 \wedge^2 \widetilde{V}_C - \widetilde{V}_C \otimes \widetilde{V}_C + \widetilde{V}_C) \otimes \widetilde{TX} + \wedge^2 \widetilde{V}_C \otimes \wedge^2 \widetilde{V}_C + 2 \wedge^4 \widetilde{V}_C - 2 \widetilde{V}_C \otimes \wedge^3 \widetilde{V}_C + 2 \widetilde{V}_C \otimes \wedge^2 \widetilde{V}_C - \widetilde{V}_C \otimes \widetilde{V}_C \otimes \widetilde{V}_C + \widetilde{V}_C + \wedge^2 \widetilde{V}_C) \right\}^{(8)}.$$

Let X be closed oriented spinc-manifold and L be the complex line bundle associated to the given spinc structure on X . We also consider L as a real vector bundle denoted by L_R . Denote by $c = c_1(L)$ the first Chern class of L . In Theorem 1.7, we set $V = L_R$ and we have

Corollary 1.10. *When $\dim X = 8$ and $3p_1(L_R) = p_1(X)$ we have*

$$(1.19) \quad 240 \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \right\}^{(8)} = \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \cdot \text{ch}(\widetilde{TX} + 2 \wedge^2 \widetilde{L}_R \otimes C - \widetilde{L}_R \otimes C \otimes \widetilde{L}_R \otimes C + \widetilde{L}_R \otimes C) \right\}^{(8)}.$$

$$(1.20) \quad 2160 \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \right\}^{(8)} = \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \cdot \text{ch}(S^2 \widetilde{TX} + \widetilde{TX} + (2 \wedge^2 \widetilde{L}_R \otimes C - \widetilde{L}_R \otimes C \otimes \widetilde{L}_R \otimes C + \widetilde{L}_R \otimes C) \otimes \widetilde{TX} + \wedge^2 \widetilde{L}_R \otimes C \otimes \wedge^2 \widetilde{L}_R \otimes C + 2 \wedge^4 \widetilde{L}_R \otimes C - 2 \widetilde{L}_R \otimes C \otimes \wedge^3 \widetilde{L}_R \otimes C + 2 \widetilde{L}_R \otimes C \otimes \wedge^2 \widetilde{L}_R \otimes C - \widetilde{L}_R \otimes C \otimes \widetilde{L}_R \otimes C \otimes \widetilde{L}_R \otimes C + \widetilde{L}_R \otimes C + \wedge^2 \widetilde{L}_R \otimes C) \right\}^{(8)}.$$

Corollary 1.11. *Let X be an 8-dimensional spinc manifold without boundary and $3p_1(L_R) = p_1(X)$, then*

$$(1.21) \quad \text{Ind}((D^c \otimes (\widetilde{TX} + 2 \wedge^2 \widetilde{L}_R \otimes C - \widetilde{L}_R \otimes C \otimes \widetilde{L}_R \otimes C + \widetilde{L}_R \otimes C))_+) \equiv 0 \pmod{240Z}.$$

$$(1.22) \quad \text{Ind}((D^c \otimes (S^2 \widetilde{TX} + \widetilde{TX} + (2 \wedge^2 \widetilde{L}_R \otimes C - \widetilde{L}_R \otimes C \otimes \widetilde{L}_R \otimes C + \widetilde{L}_R \otimes C) \otimes \widetilde{TX} + \wedge^2 \widetilde{L}_R \otimes C \otimes \wedge^2 \widetilde{L}_R \otimes C + 2 \wedge^4 \widetilde{L}_R \otimes C - 2 \widetilde{L}_R \otimes C \otimes \wedge^3 \widetilde{L}_R \otimes C + 2 \widetilde{L}_R \otimes C \otimes \wedge^2 \widetilde{L}_R \otimes C - \widetilde{L}_R \otimes C \otimes \widetilde{L}_R \otimes C \otimes \widetilde{L}_R \otimes C + \widetilde{L}_R \otimes C + \wedge^2 \widetilde{L}_R \otimes C))_+) \equiv 0 \pmod{2160Z}.$$

Theorem 1.12. *When $\dim X = 12$ and $3p_1(V) = p_1(X)$ we have*

$$(1.23) \quad -504 \left\{ \widehat{A}(TX) \det^{\frac{1}{2}} \cosh\left(\frac{\sqrt{-1}}{4\pi} R^V\right) \right\}^{(12)} = \left\{ \widehat{A}(TX) \det^{\frac{1}{2}} \cosh\left(\frac{\sqrt{-1}}{4\pi} R^V\right) \cdot \text{ch}(\widetilde{TX} + 2 \wedge^2 \widetilde{V}_C - \widetilde{V}_C \otimes \widetilde{V}_C + \widetilde{V}_C) \right\}^{(12)}.$$

$$(1.24) \quad -16632 \left\{ \widehat{A}(TX) \det^{\frac{1}{2}} \cosh\left(\frac{\sqrt{-1}}{4\pi} R^V\right) \right\}^{(12)} = \left\{ \widehat{A}(TX) \det^{\frac{1}{2}} \cosh\left(\frac{\sqrt{-1}}{4\pi} R^V\right) \cdot \text{ch}(S^2 \widetilde{TX} + \widetilde{TX} + (2 \wedge^2 \widetilde{V}_C - \widetilde{V}_C \otimes \widetilde{V}_C + \widetilde{V}_C) \otimes \widetilde{TX} + \wedge^2 \widetilde{V}_C \otimes \wedge^2 \widetilde{V}_C + 2 \wedge^4 \widetilde{V}_C - 2 \widetilde{V}_C \otimes \wedge^3 \widetilde{V}_C + 2 \widetilde{V}_C \otimes \wedge^2 \widetilde{V}_C - \widetilde{V}_C \otimes \widetilde{V}_C \otimes \widetilde{V}_C + \widetilde{V}_C + \wedge^2 \widetilde{V}_C) \right\}^{(12)}.$$

Corollary 1.13. *When $\dim X = 12$ and $3p_1(L_R) = p_1(X)$ we have*

$$(1.25) \quad -504 \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \right\}^{(12)} = \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \cdot \text{ch}(\widetilde{TX} + 2 \wedge^2 \widetilde{L}_R \otimes C - \widetilde{L}_R \otimes C \otimes \widetilde{L}_R \otimes C + \widetilde{L}_R \otimes C) \right\}^{(12)}.$$

$$(1.26) \quad -16632 \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \right\}^{(12)} = \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \cdot \text{ch}(S^2 \widetilde{TX} + \widetilde{TX} + (2 \wedge^2 \widetilde{L}_R \otimes C - \widetilde{L}_R \otimes C \otimes \widetilde{L}_R \otimes C + \widetilde{L}_R \otimes C) \otimes \widetilde{TX} + \wedge^2 \widetilde{L}_R \otimes C \otimes \wedge^2 \widetilde{L}_R \otimes C + 2 \wedge^4 \widetilde{L}_R \otimes C - 2 \widetilde{L}_R \otimes C \otimes \wedge^3 \widetilde{L}_R \otimes C + 2 \widetilde{L}_R \otimes C \otimes \wedge^2 \widetilde{L}_R \otimes C - \widetilde{L}_R \otimes C \otimes \widetilde{L}_R \otimes C \otimes \widetilde{L}_R \otimes C + \widetilde{L}_R \otimes C + \wedge^2 \widetilde{L}_R \otimes C) \right\}^{(12)}.$$

Corollary 1.14. *Let X be an 12-dimensional spin c manifold without boundary and $3p_1(L_R) = p_1(X)$, then*

$$(1.27) \quad \text{Ind}((D^c \otimes (\widetilde{TX} + 2 \wedge^2 \widetilde{L}_R \otimes C - \widetilde{L}_R \otimes C \otimes \widetilde{L}_R \otimes C + \widetilde{L}_R \otimes C))_+) \equiv 0 \pmod{504Z}.$$

$$(1.28) \quad \text{Ind}((D^c \otimes (S^2 \widetilde{TX} + \widetilde{TX} + (2 \wedge^2 \widetilde{L}_R \otimes C - \widetilde{L}_R \otimes C \otimes \widetilde{L}_R \otimes C + \widetilde{L}_R \otimes C) \otimes \widetilde{TX} + \wedge^2 \widetilde{L}_R \otimes C \otimes \wedge^2 \widetilde{L}_R \otimes C + 2 \wedge^4 \widetilde{L}_R \otimes C - 2 \widetilde{L}_R \otimes C \otimes \wedge^3 \widetilde{L}_R \otimes C + 2 \widetilde{L}_R \otimes C \otimes \wedge^2 \widetilde{L}_R \otimes C - \widetilde{L}_R \otimes C \otimes \widetilde{L}_R \otimes C \otimes \widetilde{L}_R \otimes C + \widetilde{L}_R \otimes C + \wedge^2 \widetilde{L}_R \otimes C))_+) \equiv 0 \pmod{16632Z}.$$

Theorem 1.15. *When $\dim X = 16$ and $3p_1(V) = p_1(X)$ we have*

$$(1.29) \quad 480 \left\{ \widehat{A}(TX) \det^{\frac{1}{2}} \cosh\left(\frac{\sqrt{-1}}{4\pi} R^V\right) \right\}^{(12)} = \left\{ \widehat{A}(TX) \det^{\frac{1}{2}} \cosh\left(\frac{\sqrt{-1}}{4\pi} R^V\right) \cdot \text{ch}(\widetilde{TX} + 2 \wedge^2 \widetilde{V}_C - \widetilde{V}_C \otimes \widetilde{V}_C + \widetilde{V}_C) \right\}^{(12)}.$$

Corollary 1.16. *When $\dim X = 16$ and $3p_1(L_R) = p_1(X)$ we have*

$$(1.30) \quad 480 \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \right\}^{(16)} = \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \cdot \text{ch}(\widetilde{TX} + 2 \wedge^2 \widetilde{L}_R \otimes C - \widetilde{L}_R \otimes C \otimes \widetilde{L}_R \otimes C + \widetilde{L}_R \otimes C) \right\}^{(16)}.$$

Corollary 1.17. *Let X be an 16-dimensional spinc manifold without boundary and $3p_1(L_R) = p_1(X)$, then*

$$(1.31) \quad \text{Ind}((D^c \otimes (\widetilde{TX} + 2 \wedge^2 \widetilde{L}_R \otimes C - \widetilde{L}_R \otimes C \otimes \widetilde{L}_R \otimes C + \widetilde{L}_R \otimes C))_+) \equiv 0 \pmod{480Z}.$$

Theorem 1.18. *When $\dim X = 20$ and $3p_1(V) = p_1(X)$ we have*

$$(1.32) \quad -264 \left\{ \widehat{A}(TX) \det^{\frac{1}{2}} \cosh\left(\frac{\sqrt{-1}}{4\pi} R^V\right) \right\}^{(20)} = \left\{ \widehat{A}(TX) \det^{\frac{1}{2}} \cosh\left(\frac{\sqrt{-1}}{4\pi} R^V\right) \cdot \text{ch}(\widetilde{TX} + 2 \wedge^2 \widetilde{V}_C - \widetilde{V}_C \otimes \widetilde{V}_C + \widetilde{V}_C) \right\}^{(20)}.$$

Corollary 1.19. *When $\dim X = 20$ and $3p_1(L_R) = p_1(X)$ we have*

$$(1.33) \quad -264 \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \right\}^{(20)} = \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \cdot \text{ch}(\widetilde{TX} + 2 \wedge^2 \widetilde{L}_R \otimes C - \widetilde{L}_R \otimes C \otimes \widetilde{L}_R \otimes C + \widetilde{L}_R \otimes C) \right\}^{(20)}.$$

Corollary 1.20. *Let X be an 20-dimensional spinc manifold without boundary and $3p_1(L_R) = p_1(X)$, then*

$$(1.34) \quad \text{Ind}((D^c \otimes (\widetilde{TX} + 2 \wedge^2 \widetilde{L}_R \otimes C - \widetilde{L}_R \otimes C \otimes \widetilde{L}_R \otimes C + \widetilde{L}_R \otimes C))_+) \equiv 0 \pmod{264Z}.$$

Theorem 1.21. *When $\dim X = 10$ and $p_1(L_R) = p_1(X)$ we have*

$$(1.35) \quad 240 \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \right\}^{(10)} = \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \text{ch}(\widetilde{TX} - \widetilde{L}_R \otimes C) \right\}^{(10)}.$$

$$2160 \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \right\}^{(10)} = \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \cdot \text{ch}(S^2 \widetilde{TX} + \widetilde{TX} + \wedge^2 \widetilde{L}_R \otimes C - \widetilde{L}_R \otimes C - \widetilde{TX} \otimes \widetilde{L}_R \otimes C) \right\}^{(10)}.$$

Corollary 1.22. *Let X be an 10-dimensional spinc manifold without boundary and $p_1(L_R) = p_1(X)$, then*

$$(1.36) \quad \text{Ind}((D^c \otimes (\widetilde{TX} - \widetilde{L_R \otimes C}))_+) \equiv 0 \pmod{240Z}.$$

$$(1.37) \quad \text{Ind}((D^c \otimes (S^2 \widetilde{TX} + \widetilde{TX} + \wedge^2 \widetilde{L_R \otimes C} - \widetilde{L_R \otimes C} - \widetilde{TX} \otimes \widetilde{L_R \otimes C}))_+) \equiv 0 \pmod{2160Z}.$$

Theorem 1.23. *When $\dim X = 14$ and $p_1(L_R) = p_1(X)$ we have*

$$(1.38) \quad \begin{aligned} -504 \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \right\}^{(14)} &= \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \text{ch}(\widetilde{TX} - \widetilde{L_R \otimes C}) \right\}^{(14)}. \\ -16632 \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \right\}^{(14)} &= \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \right. \\ &\quad \left. \cdot \text{ch}(S^2 \widetilde{TX} + \widetilde{TX} + \wedge^2 \widetilde{L_R \otimes C} - \widetilde{L_R \otimes C} - \widetilde{TX} \otimes \widetilde{L_R \otimes C}) \right\}^{(14)}. \end{aligned}$$

Corollary 1.24. *Let X be an 14-dimensional spinc manifold without boundary and $p_1(L_R) = p_1(X)$, then*

$$(1.39) \quad \text{Ind}((D^c \otimes (\widetilde{TX} - \widetilde{L_R \otimes C}))_+) \equiv 0 \pmod{504Z}.$$

$$(1.40) \quad \text{Ind}((D^c \otimes (S^2 \widetilde{TX} + \widetilde{TX} + \wedge^2 \widetilde{L_R \otimes C} - \widetilde{L_R \otimes C} - \widetilde{TX} \otimes \widetilde{L_R \otimes C}))_+) \equiv 0 \pmod{16632Z}.$$

Theorem 1.25. *When $\dim X = 18$ and $p_1(L_R) = p_1(X)$ we have*

$$480 \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \right\}^{(18)} = \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \text{ch}(\widetilde{TX} - \widetilde{L_R \otimes C}) \right\}^{(18)}.$$

Corollary 1.26. *Let X be an 18-dimensional spinc manifold without boundary and $p_1(L_R) = p_1(X)$, then*

$$(1.41) \quad \text{Ind}((D^c \otimes (\widetilde{TX} - \widetilde{L_R \otimes C}))_+) \equiv 0 \pmod{480Z}.$$

Theorem 1.27. *When $\dim X = 22$ and $p_1(L_R) = p_1(X)$ we have*

$$-264 \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \right\}^{(22)} = \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \text{ch}(\widetilde{TX} - \widetilde{L_R \otimes C}) \right\}^{(22)}.$$

Corollary 1.28. *Let X be an 22-dimensional spinc manifold without boundary and $p_1(L_R) = p_1(X)$, then*

$$(1.42) \quad \text{Ind}((D^c \otimes (\widetilde{TX} - \widetilde{L_R \otimes C}))_+) \equiv 0 \pmod{264Z}.$$

2. THE PROOF OF THEOREMS IN SECTION 1

Let X be a $4k$ -dimensional spin manifold and $\Delta(X)$ be the spinor bundle. Set

$$(2.1) \quad \Theta_1(T_C X) = \bigotimes_{n=1}^{\infty} S_{q^n}(\widetilde{T_C X}) \otimes \bigotimes_{m=1}^{\infty} \wedge_{q^m}(\widetilde{T_C X}),$$

$$(2.2) \quad \Theta_2(T_C X) = \bigotimes_{n=1}^{\infty} S_{q^n}(\widetilde{T_C X}) \otimes \bigotimes_{m=1}^{\infty} \wedge_{-q^{m-\frac{1}{2}}}(\widetilde{T_C X}),$$

$$(2.3) \quad \Theta_3(T_C X) = \bigotimes_{n=1}^{\infty} S_{q^n}(\widetilde{T_C X}) \otimes \bigotimes_{m=1}^{\infty} \wedge_{q^{m-\frac{1}{2}}}(\widetilde{T_C X}).$$

Let

$$(2.4)$$

$$\begin{aligned} Q(X, \tau) &= \text{Ind}((D \otimes [\Delta(X) \otimes \Theta_1(T_C X) + 2^{2k}\Theta_2(T_C X) + 2^{2k}\Theta_3(T_C X)]))_+ \\ &= \int_X \widehat{A}(TX) [\text{ch}(\Delta(X))\text{ch}(\Theta_1(T_C X)) + 2^{2k}\text{ch}(\Theta_2(T_C X)) + 2^{2k}\text{ch}(\Theta_3(T_C X))]. \end{aligned}$$

We first recall the four Jacobi theta functions are defined as follows(cf. [2]):

$$(2.5) \quad \theta(v, \tau) = 2q^{\frac{1}{8}} \sin(\pi v) \prod_{j=1}^{\infty} [(1 - q^j)(1 - e^{2\pi\sqrt{-1}v} q^j)(1 - e^{-2\pi\sqrt{-1}v} q^j)],$$

$$(2.6) \quad \theta_1(v, \tau) = 2q^{\frac{1}{8}} \cos(\pi v) \prod_{j=1}^{\infty} [(1 - q^j)(1 + e^{2\pi\sqrt{-1}v} q^j)(1 + e^{-2\pi\sqrt{-1}v} q^j)],$$

$$(2.7) \quad \theta_2(v, \tau) = \prod_{j=1}^{\infty} [(1 - q^j)(1 - e^{2\pi\sqrt{-1}v} q^{j-\frac{1}{2}})(1 - e^{-2\pi\sqrt{-1}v} q^{j-\frac{1}{2}})],$$

$$(2.8) \quad \theta_3(v, \tau) = \prod_{j=1}^{\infty} [(1 - q^j)(1 + e^{2\pi\sqrt{-1}v} q^{j-\frac{1}{2}})(1 + e^{-2\pi\sqrt{-1}v} q^{j-\frac{1}{2}})],$$

where $q = e^{2\pi\sqrt{-1}\tau}$ with $\tau \in \mathbf{H}$, the upper half complex plane. Let

$$(2.9) \quad \theta'(0, \tau) = \left. \frac{\partial \theta(v, \tau)}{\partial v} \right|_{v=0}.$$

Then the following Jacobi identity (cf. [2]) holds,

$$(2.10) \quad \theta'(0, \tau) = \pi \theta_1(0, \tau) \theta_2(0, \tau) \theta_3(0, \tau).$$

Denote $SL_2(\mathbf{Z}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a, b, c, d \in \mathbf{Z}, ad - bc = 1 \right\}$ the modular group. Let $S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, $T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ be the two generators of $SL_2(\mathbf{Z})$. They act on \mathbf{H} by $S\tau = -\frac{1}{\tau}$, $T\tau = \tau + 1$. One has the following transformation laws of theta functions under the actions of S and T (cf. [2]):

$$(2.11) \quad \theta(v, \tau + 1) = e^{\frac{\pi\sqrt{-1}}{4}} \theta(v, \tau), \quad \theta(v, -\frac{1}{\tau}) = \frac{1}{\sqrt{-1}} \left(\frac{\tau}{\sqrt{-1}} \right)^{\frac{1}{2}} e^{\pi\sqrt{-1}\tau v^2} \theta(\tau v, \tau);$$

$$(2.12) \quad \theta_1(v, \tau + 1) = e^{\frac{\pi\sqrt{-1}}{4}} \theta_1(v, \tau), \quad \theta_1(v, -\frac{1}{\tau}) = \left(\frac{\tau}{\sqrt{-1}} \right)^{\frac{1}{2}} e^{\pi\sqrt{-1}\tau v^2} \theta_2(\tau v, \tau);$$

$$(2.13) \quad \theta_2(v, \tau + 1) = \theta_3(v, \tau), \quad \theta_2(v, -\frac{1}{\tau}) = \left(\frac{\tau}{\sqrt{-1}} \right)^{\frac{1}{2}} e^{\pi\sqrt{-1}\tau v^2} \theta_1(\tau v, \tau);$$

$$(2.14) \quad \theta_3(v, \tau + 1) = \theta_2(v, \tau), \quad \theta_3(v, -\frac{1}{\tau}) = \left(\frac{\tau}{\sqrt{-1}} \right)^{\frac{1}{2}} e^{\pi\sqrt{-1}\tau v^2} \theta_3(\tau v, \tau),$$

$$(2.15) \quad \theta'(v, \tau + 1) = e^{\frac{\pi\sqrt{-1}}{4}} \theta'(v, \tau), \quad \theta'(0, -\frac{1}{\tau}) = \frac{1}{\sqrt{-1}} \left(\frac{\tau}{\sqrt{-1}} \right)^{\frac{1}{2}} \tau \theta'(0, \tau).$$

Definition 2.1. A modular form over Γ , a subgroup of $SL_2(\mathbf{Z})$, is a holomorphic function $f(\tau)$ on \mathbf{H} such that

$$(2.16) \quad f(g\tau) := f\left(\frac{a\tau + b}{c\tau + d}\right) = \chi(g)(c\tau + d)^k f(\tau), \quad \forall g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma,$$

where $\chi : \Gamma \rightarrow \mathbf{C}^*$ is a character of Γ . k is called the weight of f .

Theorem 2.2. ([11]) $Q(X, \tau)$ is a modular form over $SL_2(\mathbf{Z})$ with the weight $2k$.

Proof. Let $\{\pm 2\pi\sqrt{-1}x_j\}$ ($1 \leq j \leq 2k$) be the formal Chern roots for $T_C X$, then we have

$$(2.17) \quad \widehat{A}(TX, \nabla^{TX}) \text{ch}(\Delta(X)) \text{ch}(\Theta_1(T_C X)) = \prod_{j=1}^{2k} \frac{2x_j \theta'(0, \tau)}{\theta(x_j, \tau)} \frac{\theta_1(x_j, \tau)}{\theta_1(0, \tau)},$$

$$(2.18) \quad \widehat{A}(TX, \nabla^{TX}) \text{ch}(2^k \Theta_2(T_C X)) = \prod_{j=1}^{2k} \frac{2x_j \theta'(0, \tau)}{\theta(x_j, \tau)} \frac{\theta_2(x_j, \tau)}{\theta_2(0, \tau)},$$

$$(2.19) \quad \widehat{A}(TX, \nabla^{TX}) \text{ch}(2^k \Theta_3(T_C X)) = \prod_{j=1}^{2k} \frac{2x_j \theta'(0, \tau)}{\theta(x_j, \tau)} \frac{\theta_3(x_j, \tau)}{\theta_3(0, \tau)},$$

So we have

$$(2.20) \quad Q(X, \tau) = \prod_{j=1}^{2k} \left(\frac{2x_j \theta'(0, \tau)}{\theta(x_j, \tau)} \left(\prod_{j=1}^{2k} \frac{\theta_1(x_j, \tau)}{\theta_1(0, \tau)} + \prod_{j=1}^{2k} \frac{\theta_2(x_j, \tau)}{\theta_2(0, \tau)} + \prod_{j=1}^{2k} \frac{\theta_3(x_j, \tau)}{\theta_3(0, \tau)} \right) \right).$$

By (2.11)-(2.15), we have $Q(X, \tau+1) = Q(X, \tau)$ and $Q(X, -\frac{1}{\tau}) = \tau^{2k} Q(X, \tau)$, so $Q(X, \tau)$ is a modular form over $SL_2(\mathbf{Z})$ with the weight $2k$. \square

The proof of Theorem 1.1: It is well known that modular forms over $SL_2(\mathbf{Z})$ can be expressed as polynomials of the Einsentein series $E_4(\tau)$ and $E_6(\tau)$, where

$$(2.21) \quad E_4(\tau) = 1 + 240q + 2160q^2 + 6720q^3 + \dots,$$

$$(2.22) \quad E_6(\tau) = 1 - 504q - 16632q^2 - 122976q^3 + \dots.$$

Their weights are 4 and 6 respectively. When $\dim X = 8$, then $Q(X, \tau)$ is a modular form over $SL_2(\mathbf{Z})$ with the weight 4. $Q(X, \tau)$ must be a multiple of

$$(2.23) \quad Q(X, \tau) = \lambda E_4(\tau).$$

By (2.1)-(2.3), we have

$$(2.24) \quad \Theta_1(T_C X) = 1 + 2q \widetilde{T_C X} + O(q^2),$$

$$(2.25) \quad \Theta_2(T_C X) = 1 - q^{\frac{1}{2}} \widetilde{T_C X} + q(\widetilde{T_C X} + \wedge^2 \widetilde{T_C X}) + O(q^{\frac{3}{2}}),$$

$$(2.26) \quad \Theta_3(T_C X) = 1 + q^{\frac{1}{2}} \widetilde{T_C X} + q(\widetilde{T_C X} + \wedge^2 \widetilde{T_C X}) + O(q^{\frac{3}{2}}),$$

so

$$(2.27) \quad \begin{aligned} Q(X, \tau) &= \left[\widehat{A}(TX) \text{ch}(\Delta(X)) + 32\widehat{A}(TX) \right]^{(8)} \\ &+ \left[2\widehat{A}(TX) \text{ch}(\Delta(X)) \text{ch}(\widetilde{T_C X}) + 32\widehat{A}(TX) \text{ch}(\widetilde{T_C X} + \wedge^2 \widetilde{T_C X}) \right]^{(8)} q \\ &+ \left[\widehat{A}(TX) \text{ch}(\Delta(X)) \text{ch}(2\widetilde{T_C X} + \wedge^2 \widetilde{T_C X} + \widetilde{T_C X} \otimes \widetilde{T_C X} + S^2 \widetilde{T_C X}) \right. \\ &\left. + 32\widehat{A}(TX) \text{ch}(\wedge^4 \widetilde{T_C X} + \wedge^2 \widetilde{T_C X} \otimes \widetilde{T_C X} + \widetilde{T_C X} \otimes \widetilde{T_C X} + S^2 \widetilde{T_C X} + \widetilde{T_C X}) \right]^{(8)} q^2 + \dots \end{aligned}$$

By (2.21), (2.23) and (2.27), we get Theorem 1.1. \square

The proof of Theorem 1.3: When $\dim X = 12$, then $Q(X, \tau)$ is a modular form over $SL_2(\mathbf{Z})$ with the weight 6, so

$$(2.28) \quad Q(X, \tau) = \lambda E_6(\tau),$$

where λ are degree 6 forms. When $\dim X = 12$, direct computations show that

$$\begin{aligned}
 (2.29) \quad Q(X, \tau) &= \left[\widehat{A}(TX) \text{ch}(\Delta(X)) + 128 \widehat{A}(TX) \right]^{(12)} \\
 &+ \left[2 \widehat{A}(TX) \text{ch}(\Delta(X)) \text{ch}(\widetilde{T_C X}) + 128 \widehat{A}(TX) \text{ch}(\widetilde{T_C X} + \wedge^2 \widetilde{T_C X}) \right]^{(12)} q \\
 &+ \left[\widehat{A}(TX) \text{ch}(\Delta(X)) \text{ch}(2 \widetilde{T_C X} + \wedge^2 \widetilde{T_C X} + \widetilde{T_C X} \otimes \widetilde{T_C X} + S^2 \widetilde{T_C X}) \right. \\
 &\left. + 128 \widehat{A}(TX) \text{ch}(\wedge^4 \widetilde{T_C X} + \wedge^2 \widetilde{T_C X} \otimes \widetilde{T_C X} + \widetilde{T_C X} \otimes \widetilde{T_C X} + S^2 \widetilde{T_C X} + \widetilde{T_C X}) \right]^{(12)} q^2 + \dots .
 \end{aligned}$$

In (2.28), we compare the coefficients of 1, we get three equations about λ . By (2.28) and (2.29) we get Theorem 1.3. \square

The proof of Theorem 1.5: When $\dim X = 16$, then $Q(X, \tau)$ is a modular form over $SL_2(\mathbf{Z})$ with the weight 8, so

$$(2.30) \quad Q(X, \tau) = \lambda E_4(\tau)^2,$$

where λ is degree 8 forms. By (2.21), we have

$$(2.31) \quad E_4(\tau)^2 = 1 + 480q + 61920q^2 + \dots .$$

When $\dim X = 16$, direct computations show that

$$\begin{aligned}
 (2.32) \quad Q(X, \tau) &= \left[\widehat{A}(TX) \text{ch}(\Delta(X)) + 512 \widehat{A}(TX) \right]^{(16)} \\
 &+ \left[2 \widehat{A}(TX) \text{ch}(\Delta(X)) \text{ch}(\widetilde{T_C X}) + 512 \widehat{A}(TX) \text{ch}(\widetilde{T_C X} + \wedge^2 \widetilde{T_C X}) \right]^{(16)} q \\
 &+ \left[\widehat{A}(TX) \text{ch}(\Delta(X)) \text{ch}(2 \widetilde{T_C X} + \wedge^2 \widetilde{T_C X} + \widetilde{T_C X} \otimes \widetilde{T_C X} + S^2 \widetilde{T_C X}) \right. \\
 &\left. + 512 \widehat{A}(TX) \text{ch}(\wedge^4 \widetilde{T_C X} + \wedge^2 \widetilde{T_C X} \otimes \widetilde{T_C X} + \widetilde{T_C X} \otimes \widetilde{T_C X} + S^2 \widetilde{T_C X} + \widetilde{T_C X}) \right]^{(16)} q^2 + \dots .
 \end{aligned}$$

By (2.30)-(2.32), we get Theorem 1.5. \square

The proof of Theorem 1.7: When $\dim X = 20$, then $Q(X, \tau)$ is a modular form over $SL_2(\mathbf{Z})$ with the weight 10, so

$$(2.33) \quad Q(X, \tau) = \lambda E_4(\tau) E_6(\tau),$$

where λ is degree 10 forms. By (2.21) and (2.22), we have

$$(2.34) \quad E_4(\tau) E_6(\tau) = 1 - 264q - 117288q^2 + \dots .$$

When $\dim X = 20$, direct computations show that

$$\begin{aligned}
(2.35) \quad Q(X, \tau) &= \left[\widehat{A}(TX) \text{ch}(\Delta(X)) + 2048 \widehat{A}(TX) \right]^{(20)} \\
&+ \left[2 \widehat{A}(TX) \text{ch}(\Delta(X)) \text{ch}(\widetilde{T_C X}) + 2048 \widehat{A}(TX) \text{ch}(\widetilde{T_C X} + \wedge^2 \widetilde{T_C X}) \right]^{(20)} q \\
&+ \left[\widehat{A}(TX) \text{ch}(\Delta(X)) \text{ch}(2 \widetilde{T_C X} + \wedge^2 \widetilde{T_C X} + \widetilde{T_C X} \otimes \widetilde{T_C X} + S^2 \widetilde{T_C X}) \right. \\
&\left. + 2048 \widehat{A}(TX) \text{ch}(\wedge^4 \widetilde{T_C X} + \wedge^2 \widetilde{T_C X} \otimes \widetilde{T_C X} + \widetilde{T_C X} \otimes \widetilde{T_C X} + S^2 \widetilde{T_C X} + \widetilde{T_C X}) \right]^{(20)} q^2 + \dots
\end{aligned}$$

By (2.33)-(2.35), we get Theorem 1.7. \square

Let $\dim X = 4k$. Let V be an $2m_0$ dimensional real Euclidean vector bundle with the Euclidean connection ∇^V and the curvature R^V . Let $\{\pm 2\pi\sqrt{-1}u_r\}$ ($1 \leq r \leq m_0$) be the formal Chern roots for $V \otimes C$. Let

$$\begin{aligned}
(2.36) \quad Q(X, V, \tau) &= \left\{ \widehat{A}(TX) \det^{\frac{1}{2}} \cosh\left(\frac{\sqrt{-1}}{4\pi} R^V\right) \text{ch} \left[\bigotimes_{n=1}^{\infty} S_{q^n}(\widetilde{T_C M}) \right. \right. \\
&\left. \left. \otimes \bigotimes_{m=1}^{\infty} \wedge_{q^m}(\widetilde{V_C}) \otimes \bigotimes_{r=1}^{\infty} \wedge_{q^{r-\frac{1}{2}}}(\widetilde{V_C}) \otimes \bigotimes_{s=1}^{\infty} \wedge_{-q^{s-\frac{1}{2}}}(\widetilde{V_C}) \right] \right\}^{(4k)}.
\end{aligned}$$

Then

$$(2.37) \quad Q(X, V, \tau) = \left(\prod_{j=1}^{2k} \frac{x_j \theta'(0, \tau)}{\theta(x_j, \tau)} \prod_{r=1}^{m_0} \left(\frac{\theta_1(u_r, \tau) \theta_2(u_r, \tau) \theta_3(u_r, \tau)}{\theta_1(0, \tau) \theta_2(0, \tau) \theta_3(0, \tau)} \right) \right)^{(4k)}.$$

By (2.11)-(2.15), we have $Q(X, V, \tau + 1) = Q(X, V, \tau)$ and $Q(X, V - \frac{1}{\tau}) = \tau^{2k} Q(X, V, \tau)$ if $p_1(M) = 3p_1(V)$, so

Theorem 2.3. *Let $\dim X = 4k$. If $p_1(M) = 3p_1(V)$, then $Q(X, V, \tau)$ is a modular form over $SL_2(\mathbf{Z})$ with the weight $2k$.*

Direct computations show

$$\begin{aligned}
(2.38) \quad Q(X, V, \tau) &= \left\{ \widehat{A}(TX) \det^{\frac{1}{2}} \cosh\left(\frac{\sqrt{-1}}{4\pi} R^V\right) \right\}^{(4k)} + q \left\{ \widehat{A}(TX) \det^{\frac{1}{2}} \cosh\left(\frac{\sqrt{-1}}{4\pi} R^V\right) \right. \\
&\cdot \text{ch}(\widetilde{T_C X} + 2 \wedge^2 \widetilde{V_C} - \widetilde{V_C} \otimes \widetilde{V_C} + \widetilde{V_C}) \left. \right\}^{(4k)} + q^2 \left\{ \widehat{A}(TX) \det^{\frac{1}{2}} \cosh\left(\frac{\sqrt{-1}}{4\pi} R^V\right) \right. \\
&\cdot \text{ch}(S^2 \widetilde{T_C X} + \widetilde{T_C X} + (2 \wedge^2 \widetilde{V_C} - \widetilde{V_C} \otimes \widetilde{V_C} + \widetilde{V_C}) \otimes \widetilde{T_C X} + \wedge^2 \widetilde{V_C} \otimes \wedge^2 \widetilde{V_C} \\
&\left. + 2 \wedge^4 \widetilde{V_C} - 2 \widetilde{V_C} \otimes \wedge^3 \widetilde{V_C} + 2 \widetilde{V_C} \otimes \wedge^2 \widetilde{V_C} - \widetilde{V_C} \otimes \widetilde{V_C} \otimes \widetilde{V_C} + \widetilde{V_C} + \wedge^2 \widetilde{V_C} \right\}^{(4k)} + \dots
\end{aligned}$$

When $\dim X = 8$, then $Q(X, V, \tau)$ is a modular form over $SL_2(\mathbf{Z})$ with the weight 4 and $Q(X, V, \tau) = \lambda E_4(\tau)$. We get Theorem 1.9. When $\dim X = 12$, then $Q(X, V, \tau)$ is a

modular form over $SL_2(\mathbf{Z})$ with the weight 6 and $Q(X, V, \tau) = \lambda E_6(\tau)$. We get Theorem 1.12. When $\dim X = 16$, then $Q(X, V, \tau)$ is a modular form over $SL_2(\mathbf{Z})$ with the weight 8 and $Q(X, V, \tau) = \lambda E_4(\tau)^2$. We get Theorem 1.15. When $\dim X = 20$, then $Q(X, V, \tau)$ is a modular form over $SL_2(\mathbf{Z})$ with the weight 10 and $Q(X, V, \tau) = \lambda E_4(\tau)E_6(\tau)$ and $E_4(\tau)E_6(\tau) = 1 - 264q + \dots$. We get Theorem 1.18.

Let X be closed oriented spinc-manifold and L be the complex line bundle associated to the given spinc structure on X . We also consider L as a real vector bundle denoted by L_R . Denote by $c = c_1(L)$ the first Chern class of L . Let $\dim X = 4k + 2$ and $u = -\frac{\sqrt{-1}}{2\pi}c$. Let

$$(2.39) \quad Q(X, L, \tau) = \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \text{ch} \left[\bigotimes_{n=1}^{\infty} S_{q^n}(\widetilde{T_C M}) \otimes \bigotimes_{m=1}^{\infty} \wedge_{-q^m}(\widetilde{V_C}) \right] \right\}^{(4k+2)}.$$

Then

$$(2.40) \quad Q(X, L, \tau) = \left\{ \left(\prod_{j=1}^{2k+1} \frac{x_j \theta'(0, \tau)}{\theta(x_j, \tau)} \right) \frac{\sqrt{-1} \theta(u, \tau)}{\theta_1(0, \tau) \theta_2(0, \tau) \theta_3(0, \tau)} \right\}^{(4k+2)}.$$

Theorem 2.4. ([4]) *Let $\dim X = 4k + 2$. If $p_1(M) = p_1(L_R)$, then $Q(X, L, \tau)$ is a modular form over $SL_2(\mathbf{Z})$ with the weight $2k$.*

By Theorem 2.4, similar to Theorems 1.9-1.20, we can get Theorems 1.21-1.28.

3. ACKNOWLEDGEMENTS

The author was supported in part by NSFC No.11771070. The author is indebted to Prof. F. Han for helpful comments. The author also thank the referee for his (or her) careful reading and helpful comments.

4. DATA AVAILABILITY

No data was gathered for this article.

5. CONFLICT OF INTEREST

The authors have no relevant financial or non-financial interests to disclose.

REFERENCES

- [1] L. Alvarez-Gaumé, E. Witten, Graviational anomalies, Nucl. Phys. B234 (1983), 269-330.
- [2] K. Chandrasekharan, *Elliptic Functions*, Springer-Verlag, 1985.
- [3] Q. Chen, F. Han, Modular invariance and twisted anomaly cancellations of characteristic numbers, Trans. Amer. Math. Soc. 361 (2009), 1463-1493
- [4] Q. Chen, F. Han, W. Zhang, Generalized Witten genus and vanishing theorems. J. Differential Geom. 88(2011), no. 1, 1-40.

- [5] F. Han, R. Huang, K. Liu and W. Zhang, Cubic forms, anomaly cancellation and modularity. *Adv. Math.* 394(2022), Paper No. 108023,46pp.
- [6] F. Han, K. Liu and W. Zhang, Modular forms and generalized anomaly cancellation formulas, *J. Geom. Phys.* 62 (2012) 1038-1053.
- [7] F. Han, K. Liu and W. Zhang, Anomaly cancellation and modularity, II: the $E_8 \times E_8$ case. *Sci. China Math.* 60(2017), no. 6, 985-994.
- [8] F. Han, W. Zhang, Spin^c-manifold and elliptic genera, *C. R. Acad. Sci. Paris Serie I.*, 336 (2003), 1011-1014.
- [9] F. Han, W. Zhang, Modular invariance, characteristic numbers and eta Invariants, *J. Diff. Geom.* 67 (2004), 257-288.
- [10] K. Liu, Modular invariance and characteristic numbers. *Commu. Math. Phys.* 174 (1995), 29-42.
- [11] K. Liu, On elliptic genera and theta-functions. *Topology* 35 (1996), no. 3, 617-640.
- [12] W. Zhang, *Lectures on Chern-weil Theory and Witten Deformations*. Nankai Tracks in Mathematics Vol. 4, World Scientific, Singapore, 2001.

Y. Wang

School of Mathematics and Statistics, Northeast Normal University, Changchun Jilin, 130024, China

E-mail: wangy581@nenu.edu.cn

J. Guan

School of Mathematics and Statistics, Northeast Normal University, Changchun Jilin, 130024, China

E-mail: guanjy@nenu.edu.cn