# Algebraic discrete quantum harmonic oscillator with dynamic resolution scaling

Michael Q. May<sup>\*</sup> and Hong Qin<sup>†</sup>

Princeton Plasma Physics Laboratory and Department of Astrophysical Sciences, Princeton University, Princeton, NJ 08540, U.S.A

## Abstract

We develop an algebraic formulation for the discrete quantum harmonic oscillator (DQHO) with a finite, equally-spaced energy spectrum and energy eigenfunctions defined on a discrete domain, which is known as the  $\mathfrak{su}(2)$  or Kravchuk oscillator. Unlike previous approaches, ours does not depend on the discretization of the Schrödinger equation and recurrence relations of special functions. This algebraic formulation is endowed with a natural  $\mathfrak{su}(2)$  algebra, each finite dimensional irreducible representation of which defines a distinct DQHO labeled by its resolution. In addition to energy ladder operators, the formulation allows for resolution ladder operators connecting all DQHOs with different resolutions. The resolution ladder operators thus enable the dynamic scaling of the resolution of finite degree-of-freedom quantum simulations. Using the algebraic DQHO formalism, we are able to rigorously derive the energy eigenstate wave functions of the QHO in a purely algebraic manner without using differential equations or differential operators, which is impossible in the continuous or infinite discrete setting. The coherent state of the DQHO is constructed, and its expected position is proven to oscillate as a classical harmonic oscillator. The DQHO coherent state recovers that of the quantum harmonic oscillator at large resolution. The algebraic formulation also predicts the existence of an inverse DQHO that has no known continuous counterpart.

<sup>\*</sup> mqmay@princeton.edu

<sup>&</sup>lt;sup>†</sup> hongqin@princeton.edu

#### I. INTRODUCTION

Without regard to complications including q-deformation, discrete quantum harmonic oscillator refers to the class of Meixner oscillators including the Meixner, Kravchuk, and Charlier oscillators. The wave functions of each of these oscillators become Hermite functions (the energy eigenstates of the quantum harmonic oscillator) in certain limits, and their energy spectra are equally spaced. The limiting relationships between these oscillators has been documented [1]. In this work, we will consider the Kravchuk oscillator, which is the only finite dimensional discrete quantum harmonic oscillator (DQHO) with equally-spaced energy eigenvalues. It has been independently derived at least twice [2, 3] by discretizing the Schrödinger equation for the quantum harmonic oscillator (QHO). However, perhaps owing to the extensive recurrence relations and identities of special functions present in both derivations and most discussions of the oscillator, it remains relatively obscure. For example, the Kravchuk oscillator has also been unknowingly rederived in the context of transferring information along quantum spin chains [4, 5]. Also, despite the growing interest in quantum computing, where discreteness and finitude is an advantage, the Kravchuk oscillator has never been simulated on a quantum computer (though the closely related three-wave interaction has [6]). Indeed, quantum computation of the QHO has been restricted to non-structure preserving, finite-difference methods [7, 8].

Because the Kravchuk functions are the finite, discrete variable versions of the Hermite functions, previous derivations of the Kravchuk oscillator have taken the Kravchuk wavefunctions as an *ansatz* [1–3, 9, 10]. Finding the Hamiltonian, position and momentum eigenfunctions, and raising and lowering operators is then an exercise in cleverly applying known recurrence relations. Though straightforward, this approach based on special functions can obscure important structures which arise in Kravchuck oscillator. For example, the spectrum was found to have an  $\mathfrak{su}(2)$  structure, and for this reason, the Kravchuck oscillator is also known as a  $\mathfrak{su}(2)$  oscillator. However, rather than proving fundamental to the derivation, the  $\mathfrak{su}(2)$  algebraic structure underlying the oscillator was discovered only after a complex process of defining raising and lowering operators using hypergeometric functions. And, rather than being a dynamic quantity, the resolution of the discrete oscillator is simply taken as a fixed quantity at the start of the derivation.

In the present study, we develop an algebraic formulation for the DQHO that has a natural

 $\mathfrak{su}(2)$  structure and is fully equivalent to the Kravchuk oscillator at a fixed dimension. The dimension of a DQHO, also referred to as the resolution, is the number of the discrete energy eigenstates or grid points. In addition to recreating previous results in a simpler way, the algebraic formulation furnishes new structures that can dynamically change the resolution of the DQHO. In particular, in addition to energy raising and lowering operators, we will build resolution raising and lowering operators which connect DQHOs with different resolutions. The formulation is developed from the minimally coupled Hamiltonian for two quantum harmonic oscillators, defined in Eq. (1).

In Section II, we show that the Hamiltonian has an equally-spaced spectrum and construct quadratic energy raising and lowering operators, which, with H, form the complexified  $\mathfrak{su}(2)$ Lie algebra. It is then proven that the spectrum of the Hamiltonian consists of all finite dimensional irreducible representations of the  $\mathfrak{su}(2)$  algebra. Dimension raising and lowering operators naturally arise and allow one to move between different representations. We show that each finite dimensional irreducible representation of the  $\mathfrak{su}(2)$  algebra in the full spectrum defines a DQHO that recovers the Kravchuk oscillator at the same dimension. We also find that there exist inherent nonrenormalizable energy eigenstates in the  $\mathfrak{su}(2)$  oscillator and give a physical interpretation facilitated by the algebraic formulation. In Section III, we discuss the definition of coherence in a finite, discrete context and find coherent states by application of a discrete analog of the displacement operator to the ground state of the algebraic DQHO. Finally, in Section IV we summarize and discuss extensions.

#### II. ALGEBRAIC FORMULATION OF THE DQHO

We begin with the minimally coupled Hamiltonian for two quantum harmonic oscillators,

$$H = \frac{1}{2} \left( A_1^{\dagger} A_1 + A_2^{\dagger} A_2 \right) - \frac{1}{2} \left( A_1 A_2^{\dagger} + A_1^{\dagger} A_2 \right), \tag{1}$$

where  $A_1^{\dagger}$ ,  $A_1$ ,  $A_2^{\dagger}$ , and  $A_2$  are creation and annihilation operators satisfying the usual commutation relations,

$$[A_i, A_j^{\dagger}] = \delta_{ij}. \tag{2}$$

The Hamiltonian is composed of two dynamic parts,

$$H = \frac{1}{2}S + H_I,\tag{3}$$

with the total number of quanta shared by the oscillators

$$S \equiv N_1 + N_2 \,, \tag{4}$$

$$N_1 \equiv A_1^{\dagger} A_1 , N_2 \equiv A_2^{\dagger} A_2 ,$$
 (5)

and the interaction term defined by

$$H_I = -\frac{1}{2} \left( A_1 A_2^{\dagger} + A_1^{\dagger} A_2 \right).$$
 (6)

Both the number of quanta S and the interaction term  $H_I$  are Hermitian, and the set  $\{H, S, H_I\}$  is mutually commuting. Because the eigenvalue of S will end up being one less than than the number of the lattice points, or resolution, of the discrete system, S will be called the resolution operator. Denote the eigenstates of energy H and resolution S by  $|n, s\rangle$ , where n and s are the eigenvalues of H and S, respectively. We will refer to  $|n, s\rangle$  as the energy-resolution eigenstates.

The interaction term,  $H_I$ , appears in several different physical systems. It is used as the interacting Hamiltonian between two sites in the Bose-Hubbard mode. It can also be viewed as a special case of the quantum three wave system when one wave is strong and stationary [6, 11–15]. In certain contexts of quantum optics, it has been referred to as a beam splitter Hamiltonian. Another well-known setting where it appears is in Schwinger's construction of the angular momentum algebra from two uncoupled QHOs governed by the Hamiltonian  $S = A_1^{\dagger}A_1 + A_2^{\dagger}A_2$  [16]. However, in Schwinger's construction,  $H_I$  appears as an angular momentum, instead of as part of the Hamiltonian of the system.

Our first goal in this section is to show that the spectrum of the Hamiltonian H consists of all finite dimensional irreducible representations of the  $\mathfrak{su}(2)$  Lie algebra, each of which defines a DQHO with a resolution equal to its dimension.

The equations of motion according to Eq. (1) are

$$[H, A_1] = \frac{1}{2} (A_2 - A_1), \ [H, A_2] = \frac{1}{2} (A_1 + A_2).$$
(7)

For further analysis, we can trade  $A_1$  and  $A_2$  for operators  $B_1$  and  $B_2$  defined as follows,

$$B_1 \equiv A_1 + A_2, \ B_2 \equiv A_1 - A_2. \tag{8}$$

$[\downarrow,\rightarrow]$	H	S	$B_1$	$B_1^\dagger$	$B_2$	$B_2^{\dagger}$	Ι
Н	0	0	0	0	$-B_2$	$B_2^{\dagger}$	0
S	0	0	$-B_1$	$B_1^\dagger$	$-B_2$	$B_2^{\dagger}$	0
$B_1$	0	$B_1$	0	2I	0	0	0
$B_1^\dagger$	0	$-B_1^{\dagger}$	-2I	0	0	0	0
$B_2$	$B_2$	$B_2$	0	0	0	2I	0
$B_2^\dagger$	$-B_2^{\dagger}$	$-B_2^{\dagger}$	0	0	-2I	0	0
Ι	0	0	0	0	0	0	0

Table I. Commutation relations for  $H, S, B_1, B_2, I$ , and their complex conjugates.

Note that as with  $A_1$  and  $A_2^{\dagger}$ ,  $B_1$  commutes with  $B_2^{\dagger}$ . This allows us to write

$$H = \frac{1}{2}B_2^{\dagger}B_2,$$
 (9)

obviating the commutators

$$[H, B_1] = 0, \ [H, B_2] = -B_2.$$
(10)

Next we define the following quadratic operators in terms of  $B_1$  and  $B_2$ ,

$$D \equiv B_1^{\dagger} B_2, \ D^{\dagger} \equiv B_1 B_2^{\dagger}. \tag{11}$$

Calculation shows that D and  $D^{\dagger}$ , while commuting with S, act as lowering and raising operators of the energy,

$$[H, D] = -D, [H, D^{\dagger}] = D^{\dagger}.$$
(12)

Similarly, the operators  $B_1$  and  $B_1^{\dagger}$ , which commute with H, are the raising and lowering operators of the resolution,

$$[S, B_1] = -B_1, \ [S, B_1^{\dagger}] = B_1^{\dagger}.$$
(13)

Commutation relations for the algebra of  $\{H, S, B_1, B_2, B_1^{\dagger}, B_2^{\dagger}, I\}$  are summarized in Table I.

We now need to look more closely at the interaction term  $H_I$  to determine how the resolution operator S gets its name. The interaction term  $H_I$  has equations of motion

$[\downarrow,\rightarrow]$	$H_I$	D	$D^{\dagger}$	$B_1$	$B_1^{\dagger}$	Ι
$H_I$	0	-D	$D^{\dagger}$	$\frac{1}{2}B_1$	$-\frac{1}{2}B_1^{\dagger}$	0
D	D	0	$8H_I$	$-2B_{2}$	0	0
$D^{\dagger}$	$-D^{\dagger}$	$-8H_I$	0	0	$2B_2^\dagger$	0
$B_1$	$-\frac{1}{2}B_{1}$	$2B_2$	0	0	2I	0
$B_1^\dagger$	$\frac{1}{2}B_1^{\dagger}$	0	$-2B_2^{\dagger}$	-2I	0	0
Ι	0	0	0	0	0	0

Table II. Commutation relations for  $H_I$ , D,  $B_1$ , I, and their complex conjugates.

$$[H_I, A_1] = \frac{1}{2}A_2, \ [H_I, A_2] = \frac{1}{2}A_1.$$
(14)

It is interesting to note that by transforming  $H_I \rightarrow 2H_I$  and  $A_1 \rightarrow -iA_1$ , these take the same form as those for the QHO with the canonically commuting p and q replaced by the mutually commuting  $A_1$  and  $A_2$ . In the  $B_1$ ,  $B_2$  basis, these commutators become

$$[H_I, B_1] = \frac{1}{2}B_1, \quad [H_I, B_2] = -\frac{1}{2}B_2. \tag{15}$$

We next find that the energy raising and lowering operators are also raising and lowering operators of  $H_I$ ,

$$[H_I, D] = -D, [H_I, D^{\dagger}] = D^{\dagger}.$$
(16)

These results are summarized in Table II.

Table II also highlights an important structure— $H_I$ , D, and  $D^{\dagger}$  form the complexification of the Lie algebra  $\mathfrak{su}(2)$ . Scaling  $H_I \to 2H_I$ ,  $D \to \frac{1}{2}D$ , and  $D^{\dagger} \to \frac{1}{2}D^{\dagger}$  makes this obvious. The quadratic Casimir  $\Omega$  of  $\mathfrak{su}(2)$ , which indexes the irreducible representations of the algebra according to its eigenvalue  $\omega$ , is  $\Omega = DD^{\dagger} + 4H_I^2 + 4H_I$ . The eigenvalue  $\omega$  of  $\Omega$  is related to the dimension d of the corresponding irreducible representation via  $\omega =$ (d-1)(d+1). As usual, the algebra  $\mathfrak{su}(2)$  itself does not select a particular representation from all of its possible representations. But, for the interaction term  $H_I$  defined Eq. (6),

$$\Omega = DD^{\dagger} + 4H_I^2 + 4H_I = S(S+2).$$
(17)

This implies that the eigenvalues s of S determine the dimension of the representation by d = s + 1, which will be shown later to be the number of lattice points, i.e. the resolution, of the wave functions of the algebraic DQHO.

For a fixed resolution, the eigenvalues of H and  $H_I$  will differ only by a constant value, so S also determines the number of energy eigenstates of H for a particular resolution. Thus, the full spectrum of H consists of sets of s equally spaced eigenvalues, for each  $s \in \mathbb{N}$ , corresponding to the irreducible representations of  $\mathfrak{su}(2)$ .

#### A. Ladder operators in the energy-resolution basis

In the energy-resolution basis, the normalization factors of the energy raising and lowering operators are given by,

$$D^{\dagger}|n,s\rangle = 2\sqrt{s-n}\sqrt{n+1}|n+1,s\rangle, \qquad (18)$$

$$D|n,s\rangle = 2\sqrt{s-n+1}\sqrt{n}|n-1,s\rangle.$$
(19)

An explanation of the process for finding these and other normalizations can be found in Appendix A. These normalizations imply there is a barrier to raising the energy to or lowering the energy from n = 0 and a similar barrier for n = s. Between the energy barriers, for a fixed value of s, there are therefore s + 1 energy eigenstates accessible to the energy lowering and raising operators. This number of eigenstates is the dimension of the irreducible representation of the  $\mathfrak{su}(2)$  algebra labeled by s.

The normalization factors for the resolution raising and lowering operators are given by,

$$B_1^{\dagger}|n,s\rangle = \sqrt{2(s-n+1)}|n,s+1\rangle, \qquad (20)$$

$$B_1|n,s\rangle = \sqrt{2(s-n)|n,s-1\rangle}.$$
(21)

Because the resolution operators do not affect the energy eigenstates, with proper normalization their application leaves quantum superpositions untouched, effectively allowing for consistent, dynamic resolution scaling of quantum simulations.

How can the resolution operators not affect quantum superpositions if they do not always commute with the energy ladder operators? Namely, while  $[D, B_1^{\dagger}] = [D^{\dagger}, B_1] = 0$ ,  $[D, B_1] \neq$ 0 and  $[D^{\dagger}, B_1^{\dagger}] \neq 0$ . This is a natural result of the non-unitarity of the energy and resolution operators. For example, since the energy raising operator destroys the highest energy state, it will matter whether or not the resolution is increased before or after the energy raising operator is applied. Similarly, since the resolution lowering operator also destroys the highest



Figure 1. Actions of raising and lowering operators on energy-resolution and position-resolution bases. Each dot represents an eigenstate, and the arrows indicate how operators transform eigenstates. The arrows are translationally invariant in each diagram. *Left*: Action of the D,  $D^{\dagger}$ ,  $B_1$ , and  $B_1^{\dagger}$  operators on the energy-resolution basis. *Right*: Action of the  $A_1$ ,  $A_1^{\dagger}$ ,  $A_2$ ,  $A_2^{\dagger}$  operators on the position-resolution basis.

energy state, lowering its energy before lowering the resolution will have a different result than attempting to lower the energy after. This can be seen clearly in Fig. 1, where the actions of the D,  $D^{\dagger}$ ,  $B_1$ , and  $B_1^{\dagger}$  operators in the energy-resolution basis are shown alongside the actions of the  $A_1$ ,  $A_2$ ,  $A_1^{\dagger}$ , and  $A_2^{\dagger}$  operators in the position-resolution basis, which is discussed below.

#### B. Position and momentum operators

By analogy with the QHO, we may define a position operator

$$X \equiv \frac{1}{2} \left( D^{\dagger} + D \right) = 2N_1 - S, \tag{22}$$

and a momentum operator

$$P \equiv \frac{i}{2}(D^{\dagger} - D), \qquad (23)$$

in terms of the energy raising and lowering operators. The equations of motion of the QHO are satisfied by X and P:

$$[H,X] = -iP, (24)$$

$$[H,P] = iX. (25)$$

Note that the second equation of Eq. (22) implies that the eigenvalues of position  $x \in \{-s, -s+2, \ldots, s\}$ . However, as with D and  $D^{\dagger}$ , H may not be written in terms of X and P in the same way as it can for the QHO. Note that  $X^2 + P^2 \propto D^{\dagger 2} + D^2 \not \propto H$ .

In the energy-resolution basis, we can calculate the expectation values of X and P. An arbitrary state of fixed dimension  $\Psi_s$  in this basis is

$$\Psi_s = \sum_{j=0}^s d_j \left| j, s \right\rangle \tag{26}$$

with  $d_j$  the complex amplitude of the energy eigenmode  $|j\rangle$ . Then,

$$\langle \Psi_s | X | \Psi_s \rangle = 2 \sum_{j=0}^{s-1} \operatorname{Re}\left(d_j^* d_{j+1}\right) \sqrt{s-j} \sqrt{j+1}$$
(27)

and

$$\langle \Psi_s | P | \Psi_s \rangle = 2 \sum_{j=0}^{s-1} \operatorname{Im} \left( d_j^* d_{j+1} \right) \sqrt{s-j} \sqrt{j+1}.$$
 (28)

#### C. Position-resolution basis

The Hilbert space spanned by the spectrum of H has a different basis indexed by quantum numbers  $n_1$ ,  $n_2$ , and s, the eigenvalues of the operators  $N_1$ ,  $N_2$ , and S, respectively. Because  $s = n_1 + n_2$ , eigenstates in this basis can be labeled by  $n_1$  and s only and will be denoted by  $\psi_{n_1,s}$ . This basis will be referred to as the position-resolution basis because  $\psi_{n_1,s}$  is an eigenstate of the position operator X defined by Eq. (22).

In the position-resolution basis, the expressions of  $A_1, A_1^{\dagger}, A_2, A_2^{\dagger}$  are

$$A_1\psi_{n_1,s} = \sqrt{n_1}\psi_{n_1-1,s-1}, \quad A_1^{\dagger}\psi_{n_1,s} = \sqrt{n_1+1}\psi_{n_1+1,s+1}, \quad (29)$$

$$A_2\psi_{n_1,s} = \sqrt{s - n_1}\psi_{n_1,s-1}, \quad A_2^{\dagger}\psi_{n_1,s} = \sqrt{s + 1 - n_1}\psi_{n_1,s+1}, \quad (30)$$

from which the expressions of all other operators can be calculated. For example, the expression of the Hamiltonian is

$$H\psi_{n_{1},s} = \left(\frac{1}{2}S + H_{I}\right)\psi_{n_{1},s}$$

$$= \frac{1}{2}\left(N_{1} + N_{2}\right)\psi_{n_{1},s} - \frac{1}{2}\left(A_{1}A_{2}^{\dagger} + A_{1}^{\dagger}A_{2}\right)\psi_{n_{1},s}$$

$$= \frac{s}{2}\psi_{n_{1},s} + \sqrt{n_{1}}\sqrt{s - n_{1} + 1}\psi_{n_{1} - 1,s} + \sqrt{n_{1} + 1}\sqrt{s - n_{1}}\psi_{n_{1} + 1,s}.$$
(31)

Equation (31) shows again that in this basis, the eigenstates of the Hamiltonian must necessarily have dimension d = s + 1, since the Hamiltonian spreads states from  $\psi_{0,s}$  to  $\psi_{s,s}$ . Specifically, in this basis with a fixed dimension, H is a symmetric, tridiagonal matrix where

$$H_{ij} = \delta_{i,j}s + \delta_{i+1,j}h_{i-1} + \delta_{i,j+1}h_{j-1}, \qquad (32)$$

and

$$h_i = \sqrt{(s-i)(i+1)}.$$
 (33)

This matrix representation of the Hamiltonian is a known result for the  $\mathfrak{su}(2)$  structure of the Kravchuk DQHO [2, 3]. It also appears in models of fermionic spin chains [4, 5], the Hanh oscillator [17], and the q-deformed harmonic oscillator [18]. However, it should be emphasized that the correspondence with these previous models is only for a fixed number of lattice points, i.e. resolution. Previous models for the DQHO do not provide an internally consistent method to specify and change the resolution, which is one of the tools needed to calculate the eigenstates in the position-resolution basis. This explains why the eigenmode structures in previous DQHO models have to depend on specific, non-trivial mathematical properties of special functions.

#### D. DQHO

As shown above, for each value of s, the s + 1 energy eigenstates  $|n, s\rangle$  furnish an (s + 1)dimensional irreducible representation of the  $\mathfrak{su}(2)$  algebra of  $H_I$ ,  $D^{\dagger}$ , and D. In the present study, we will call the s + 1 energy eigenstates  $|n, s\rangle$  an (s + 1)-dimensional DQHO because they also furnish an (s + 1)-dimensional approximation of the QHO. The validity of this approximation can be justified from several different perspectives. First of all, the s + 1energy levels of the DQHO are equally spaced as in the QHO. Secondly, the raising and lowering operators  $D^{\dagger}$  and D for the DQHO approach to their counterparts in the QHO as the dimension  $s \to \infty$ . For a fixed n, when  $s \to \infty$ , equations (18) and (19) become

$$\lim_{s \to \infty} D^{\dagger} |n, s\rangle = 2\sqrt{s} \left( 1 + O\left(\frac{n}{s}\right) \right) \sqrt{n+1} |n+1, s\rangle, \tag{34}$$

$$\lim_{s \to \infty} D|n, s\rangle = 2\sqrt{s} \left(1 + O\left(\frac{n}{s}\right)\right) \sqrt{n}|n-1, s\rangle.$$
(35)

Up to an insignificant normalization constant, these are the energy ladder equations of the QHO.

As another justification for identification of the DQHO, we show how the energy eigenstate wave functions of the QHO are recovered by the DQHO. In the QHO, the energy eigenstate wave functions are, by definition, the projections of the energy eigenstates on the position eigenstates. Following this definition, the wave function of the *n*-th energy eigenstate in the DQHO is the projection of  $|n, s\rangle$  on  $\psi_{n_1,s}$ ,

$$\alpha_{n_1,s}^n \equiv \langle \psi_{n_1,s} | n, s \rangle. \tag{36}$$

To calculate  $\alpha_{n_1,s}^n$ , we re-express Eq. (36) as

$$|n,s\rangle = \sum_{n_1=0}^{s} \alpha_{n_1,s}^n \psi_{n_1,s}.$$
 (37)

Algorithmically, eigenstates  $|n, s\rangle$ , specified by  $\alpha_{n_1,s}^n$ , may be constructed by applying the  $B_1^{\dagger}$  operators, which raises the resolution s by one without affecting the energy n, to the  $\psi_{0,0} \equiv |0,0\rangle$  state. Performing

$$(B_1^{\dagger})^s \psi_{0,0} = \sqrt{(2s)!!} | n = 0, s \rangle, \tag{38}$$

gives the lowest energy eigenstate in the *d*-dimensional basis (representation), while applying the  $B_2^{\dagger}$  operator to the  $\psi_{0,0}$  state *s* times gives the highest energy state in the *d*-dimensional basis. The other energy-resolution eigenstates  $|n, s\rangle$  in this *d*-dimensional basis may be found by applying the *D* and  $D^{\dagger}$  operators. This procedure provides an algorithmic definition of  $\alpha_{n_1,s}^n$ .

To see how  $\alpha_{n_1,s}^n$  recovers the energy eigenstate wave functions of the QHO, we note that  $\alpha_{n_1,s}^n$ , can be expressed as a known special function called Wigner's little-d function, which can be rewritten in terms of the weighted Kravchuk function as well as the ordinary hypergeometric function,

$$\alpha_{n_{1,s}}^{n} = d_{-n+s/2,n_{1}-s}^{s/2} \left(\frac{\pi}{2}\right) = K_{n}(n_{1},s)$$
$$= (-1)^{n} 2^{-s/2} {\binom{s}{j}}^{1/2} {\binom{s}{n_{1}}}^{1/2} {}_{2}F_{1}(-n,-n_{1};-s;2).$$
(39)

Note that this is not the standard notation for the Kravchuk functions in terms of the Kravchuk polynomials, as discussed in Appendix B. The weighted Kravchuk functions form a complete orthonormal set on the  $n_1$  space for fixed s, are known to limit to the Hermite functions for a fixed n as  $s \to \infty$ , and obey recurrence relations which limit to those of the Hermite functions for a fixed n as  $s \to \infty$ , and obey recurrence relations which limit to those of the Hermite functions for a fixed n as  $s \to \infty$  [19]. However, for any fixed s, it is only possible to construct  $\lfloor s/2 \rfloor$  states with continuous analogs. This may be readily understood from the fact that each increase in energy of the Hermite functions adds a zero-crossing to the wavefunction. When the number of zero-crossings exceeds  $\lfloor s/2 \rfloor$ , the wavefunctions will experience aliasing. Thus, in order to form a complete set, the Kravchuk functions for  $n \ge s/2$  cannot limit to the Hermite functions even as  $s \to \infty$ .

That the Kravchuk functions and Wigner's little-d function become the Hermite functions for fixed n as  $s \to \infty$  is well known [9]. It is because of this relationship that, up to a rescaling of variables, these wavefunctions are the starting point for previous Kravchuk DQHO work [2, 3]. But from the algebraic DQHO formalism developed in the present study, the discrete wave functions of the energy eigenstates are rigorously defined and constructed without resorting to these special functions, while still recovering them. The raising and lowering operators for energy, position, and resolution serve as a sufficient and effective algorithm for calculating the wave functions of the DQHO.

This bring us to the following interesting observation. Using the algebraic DQHO formalism developed, we are able to rigorously derive the energy eigenstate wave functions of the QHO in a purely algebraic manner without using differential equations or differential operators. This is impossible in the continuous or infinite discrete setting and attests to the usefulness of the algebraic DQHO.

A set of wave functions  $\alpha_{n_{1,s}}^{n}$  are shown in Fig. 2 for s = 20 and n = 0, 1, 2, 10, 18, 19, 20. Although the n = j and n = s - j states are orthogonal (as all the energy eigenstates are), they have the same amplitude at each position. But the relative phase between adjacent positions are different for these two states. The higher energy state is more oscillatory then the lower energy one. Note that the algebraic DQHO allows an n = s/2 mode when



Figure 2. Representative wave functions of the algebraic DQHO given by energy-resolution eigenstate  $\alpha_{n_{1},s}^{n}$  for s = 20 and n = 0, 1, 2, 10, 18, 19, 20. The vertical axes are  $\alpha_{n_{1},s}^{n}$  and the horizontal axes are position  $x = n_{1} - n_{2}$ . Although each wavefunction consists of a set of discrete points, they are drawn as lines for clarity. The lowest energy eigenstate  $\alpha_{n_{1},20}^{0}$  (solid blue curve) corresponds to the ground state of the QHO. Modes with n < s/2 (solid blue curves) furnish finite degree-of-freedom approximations of the bounded QHO eigenmode wave functions with resolution of d = s + 1 = 21. Energy modes with  $n \ge s/2$  (dashed orange curves) cannot be reached from the ground state in the limit  $d \to \infty$ . They form another kind of DQHO called inverse DQHO that has no continuous counterpart.

d = s + 1 is an odd number. Unlike the n < s/2 and n > s/2 energy modes, which have the same expectation values as the QHO wave functions in the limit  $d \to \infty$ , the n = s/2 mode has no QHO analog. Indeed, this energy eigenstate will always extend from  $\psi_{0,s}$  to  $\psi_{s,s}$ , corresponding to an unrenormalizable wave function in the continuous limit. Since increasing the energy of the highest n < s/2 energy state,  $|s/2 - 1, s\rangle$ , will require the DQHO to pass through this unrenormalizable state, we can interpret the spectrum as consisting of two different oscillators. The lower energy states with n < s/2 furnish a DQHO that corresponds to the QHO. The higher energy states with n > s/2 form another DQHO that does not have a known continuous counterpart. For easy reference, we will refer to it as the inverse DQHO.

The unrenormalizable energy states have not been identified in previous Kravchuk DQHO literature. Indeed, the significance of the n = s/2 state may be lost, especially if the

energy eigenvalues are shifted by a constant value as in other literature where, although the eigenstates are the same, the energy eigenvalues may shifted by a factor of 1/2 to look more similar to those of the QHO [2, 3]. To the authors' knowledge, the existence of the inverse DQHO and the unrenomalizable n = s/2 state has not been addressed before.

#### III. COHERENT STATES

For the QHO, coherent states are 1) eigenstates of the energy lowering operator, 2) obtained via application of the displacement operator to the ground state, and 3) minimum uncertainty states. However, in a finite basis, the first and third of these definitions become problematic. The first definition requires that the QHO have an infinite number of energy eigenstates, otherwise the energy lowering operator would eliminate the highest energy eigenstate. Thus, in a finite setting, the eigenstates of the energy lowering operator are trivial. The third definition also poses a problem in this finite setting. We find that since either the variance in position  $\sigma_X^2$  or the variance in momentum  $\sigma_P^2$  can be zero,

$$\min \sigma_X^2 \sigma_P^2 = 0. \tag{40}$$

The minimum uncertainty states are just the union of the position and momentum eigenstates.

We will thus proceed with the second definition of coherence. Several different displacement operators have been proposed with the intention of preserving the displaced-Gaussian shape of the QHO in the finite, discrete setting [20, 21]; however, since the definition of the QHO displacement operator

$$\hat{\mathcal{D}}(\beta) = \exp\left(\beta D^{\dagger} - \beta^* D\right) \tag{41}$$

can be easily translated to our finite system, there is no *ab initio* reason to modify it. Here,  $\beta$  is a displacement parameter. Therefore, in the present study, the coherent state of the DQHO is constructed as

$$|\beta\rangle = \hat{\mathcal{D}}(\beta)|0,s\rangle. \tag{42}$$

There is significant literature on the application of the displacement operator to elements of the  $\mathfrak{su}(2)$  algebra which result in *spin coherent states* [22, 23]. However, this literature is solely concerned with the energy-resolution basis, so discussions of the  $n_1$ , s basis have been limited to work related to DQHOs. For the Kravchuk DQHO, expectation values of the Hamiltonian and energy ladder operators have been found for displacement operator coherent states [24], but discussion and calculation of the wavefunctions and position expectation values of these states has not been presented before.

We have now rigorously proven that the expected position of the DQHO coherent state  $|\beta\rangle$  constructed in Eq. (42) oscillates as a classical harmonic oscillator, and that  $|\beta\rangle$  recovers the coherent state of the QHO when the resolution  $s \to \infty$ .

Using the operator matrix notation and calculation given in Appendix IV, we find that in the  $n_1$ , s basis,

$$\log(\hat{\mathcal{D}}(\beta)) = \operatorname{band}_{s+1\times s+1} \left[ 1 \times 1 \to (\beta - \beta^*) \left(s - 2j\right)_{j=0}^s, \\ 1 \times 2 \to -(\beta + \beta^*) \left(\sqrt{s - j}\sqrt{j + 1}\right)_{j=0}^{s-1}, \\ 2 \times 1 \to (\beta + \beta^*) \left(-\sqrt{s - j}\sqrt{j + 1}\right)_{j=0}^{s-1} \right].$$
(43)

In the case that the displacement parameter is imaginary,  $\beta = i|\beta|$ , we find a simple, diagonal expression for the displacement operator:

$$\hat{\mathcal{D}}(i|\beta|) = \text{band}_{s+1 \times s+1} \left[ 1 \times 1 \to \left( e^{2i|\beta|(s-2j)} \right)_{j=0}^{s} \right].$$
(44)

Note that with  $\beta = i|\beta|$ , there is no initial displacement of state, and the displacement only occurs with time evolution. It is straightforward to calculate the expectation value of the position operator for a coherent state using the energy-resolution basis. For a particular resolution, X is a sum of energy raising and lowering operators, and  $U = e^{-iHt}$  is diagonal. Using the Baker-Campbell–Hausdorff (BCH) identity, the displacement operator can be transformed to a finite sum of energy raising operators when acting on the ground state since,

$$\hat{\mathcal{D}}(\beta)|0,s\rangle \propto e^{\frac{\beta}{2|\beta|}\tan(2|\beta|)D^{\dagger}}|0,s\rangle.$$
(45)

Note that this differs from the typical application of the BCH identity for the QHO because the DQHO energy raising and lowering operators obey a different commutation relation,  $[D, D^{\dagger}] = 8H_I$ , from those of the QHO,  $[a, a^{\dagger}] = 1$  [25]. We find after much simplification (see Appendix IV),

$$\langle \beta | X | \beta \rangle = s \, \cos(2t - \arg(\beta)) \, \sin(4|\beta|). \tag{46}$$



Figure 3. Expectation value of the position eigenstates of the DQHO coherent state  $\hat{\mathcal{D}}(\beta)|0, s = 100\rangle$ . All plots are for  $\beta = i\pi/2$ , and t ranges from 0 to  $\pi/4$ . The initial state, t = 0, corresponds to the ground state n = 0 of Fig. 2 (albeit at a different resolution).

Note that the time dependence comes from moving from the Heisenberg picture operator X to the Schrödinger picture operator  $X_S = e^{-iHt} X e^{iHt}$ . The maximum amplitude is simply s, as expected, and the expectation value oscillates harmonically. A quarter period of the wavefunction in the  $n_1$ , s basis for such a maximum amplitude oscillation may be seen in Fig. 3.

As suggested by Eq. (46), there is a symmetry between the argument of  $\beta$  and t, and the expectation value of the wavefunction is unaffected by the transformation  $t' = -\arg(\beta)/2$ ,  $\arg(\beta') = -2t$ . Indeed, it happens that the amplitude of the wave function is only a function of the wavefunction's average position as demonstrated in Figure 4. Thus, fixing time t = 0 and  $\arg(\beta) = 0$ , we find that varying the initial maximum displacement  $|\beta|$  results in the same wavefunctions as we found by varying time for a fixed displacement.

Correspondence between the continuous QHO and the DQHO can be found in the limit that  $|\beta| \ll 1$ . Here, the amplitude of the oscillation is linearly proportional to  $|\beta|$ , and the wavefunction, as seen in Figs. 3 and 4 is minimally deformed from its initial state.

#### IV. DISCUSSION

In addition to providing a new and simple derivation of the  $\mathfrak{su}(2)$  oscillator based on two coupled quantum harmonic oscillators, which may be of intrinsic interest, the algebraic DQHO formalism developed in the present study also creates a formal structure for connect-



Figure 4. Expectation value of the position eigenstates for different values of initial displacement  $\beta$  of the DQHO coherent state  $\hat{\mathcal{D}}(\beta)|0, s = 100\rangle$ . All plots are for t = 0, and  $\beta$  ranges from 0 to  $\pi/8$ . The ground state,  $\beta = 0$ , corresponds to the ground state n = 0 of Fig. 2 (albeit at a different resolution). The position eigenstate amplitudes are identical to those in Fig. 3.

ing representations of DQHOs at different dimensions via the resolution raising and lowering operators. The Kravchuk oscillator has never been simulated on a quantum computer, as noted in the introduction, and the resolution operators provide additional impetus for this to change. Especially now, while quantum compute time is scarce, the ability to dynamically alter the resolution of a simulation could be decisive. Increasing the resolution of a simulation would also improve its fault tolerance in finer gradations than those available in typical quantum error correction schemes. For example, suppose we have some number of qubits available for computation, Q, and we want to simulate a DQHO of dimension d. We require  $Q \ge \log_2(d)$  to minimally perform the simulation, so we have the difference of qubits left for fault tolerance. Most error correcting schemes require additional qubits at some fixed proportion r to the dimension of the simulation, so  $Q \ge \log_2(d) + \log_2(r)$ . However, for the DQHO described here, we may increase the dimension of the simulation in steps of any size  $p \ge 1$ , naturally entangling the information over more qubits. Thus, for the DQHO,  $Q \ge \log_2(d+p)$ . Of course, this is not a substitute for error correction, but since it may be deployed at any scale, it can almost always act as a supplement.

Since every computational operation on a quantum computer must be unitary, nonunitary operators, such as the resolution raising and lowering operators, need to be embedded in higher dimensional unitary operators to act on a quantum computer. This embedding is always possible, but it may require the dimension of the problem to be (temporarily) doubled. Future work as it relates to quantum computation could involve a search for efficient unitary embeddings of each of the DQHO's nonunitary operators.

Finally, although the DQHO has never been simulated on a quantum computer, the minimally nonlinear quantum three-wave interaction, governed by  $H_{TW} = igA_1^{\dagger}A_2A_3 - ig^*A_1A_2^{\dagger}A_3^{\dagger}$  has been simulated [6, 11, 12]. Note that the constant g is a coupling coefficient which may be taken to -i without affecting the dynamics. Harmonic behavior has been previously found in the limit of  $A_2 \to \infty$  or  $A_3 \to \infty$  [14]. In the latter case, the quantum three wave interaction Hamiltonian just becomes the interaction term  $H_I$  defined by Eq. (6) multiplied by the constant  $\sqrt{n_2}$ . In the three-wave interaction, this represents amplitude exchange of two small waves mediated by a third, much larger wave.

#### ACKNOWLEDGMENTS

This research was supported by the U.S. Department of Energy (DE-AC02-09CH11466).

# APPENDIX A: NORMALIZATIONS OF ENERGY AND RESOLUTION OPER-ATORS

To calculate the normalization of the energy and resolution operators, we use a method a method similar to that for the QHO. We will begin with the energy lowering operator  $D = B_1^{\dagger}B_2$ . From Eqs. (1), (4), and (8), see that

$$H = \frac{1}{2}B_2^{\dagger}B_2$$

and

$$S = \frac{1}{2} \left( B_1^{\dagger} B_1 + B_2^{\dagger} B_2 \right).$$

Subtracting these, we have  $B_1^{\dagger}B_1 = 2(S - H)$ , and since  $[B_1, B_1^{\dagger}] = 2$ ,

$$B_1 B_1^{\dagger} = 2(S - H + 1). \tag{47}$$

Next, we calculate the squared normalization of  $B_1^{\dagger}$ :

$$\left|\mathcal{N}_{B_{1}^{\dagger}}\right|^{2} = \langle n, s | B_{1}B_{1}^{\dagger} | n, s \rangle = \langle n, s | 2(S - H + 1) | n, s \rangle = 2(s - n + 1).$$

Similarly, the squared normalization of of  $B_2$  may be found:

$$\left|\mathcal{N}_{B_2}\right|^2 = \langle n, s | B_2^{\dagger} B_2 | n, s \rangle = 2n.$$

Choosing the arbitrary phase factors to be 1, we find the normalization factor of  $B_1^{\dagger}$  to be

$$\mathcal{N}_{B_1^{\dagger}} = \sqrt{2(s-n+1)}$$

and of  $B_2$  to be

$$\mathcal{N}_{B_2} = \sqrt{2n}.$$

Next, recall that  $B_1^{\dagger}$  acts as an *s* raising operator while keeping *n* constant. Also, note that  $[S + H, B_2] = -2B_2$  and  $[S - H, B_2] = 0$ , so  $B_2$  acts as an s + n lowering operator while keeping s - n constant. So, putting these operators together we can see that since  $D = B_1^{\dagger}B_2$  will first decrease s + n by 2 (while keeping s - n constant) and then increase *s* by 1, it acts as an *n* lowering operator while keeping *s* constant. Its normalization will just be the normalization of its constituent parts,

$$D|n,s\rangle = B_1^{\dagger}B_2|n,s\rangle = 2\sqrt{s-n+1}\sqrt{n}|n-1,s\rangle,$$

as written in Eq. (19). The calculations for the normalizations of  $D^{\dagger}$  and  $B_1$  proceed in the same way. Since  $B_1$  acts as an *s* lowering operator,

$$\mathcal{N}_{B_1} = \sqrt{2(s-n)},$$

and since  $B_2^{\dagger}$  acts as an s + n raising operator,

$$\mathcal{N}_{B_2^{\dagger}} = \sqrt{2(n+1)}.$$

The operators  $B_1$  and  $B_2^{\dagger}$  may then be applied to a representative state function  $|n, s\rangle$  to find the normalization of  $D^{\dagger}$ .

### **APPENDIX B: KRAVCHUK FUNCTIONS**

The normalized Kravchuk functions are generally written in terms of

$$k_n(x;p,N) = (-1)^n \binom{N}{n} p^n {}_2F_1(-n,-x;-N;p^{-1}),$$

the Kravchuk polynomials, where  $0 , <math>N \in \mathbb{N}$ , and  $n \in \{0, 1, ..., N\}$ . The prefactors of the hypergeometric functions are usually omitted in older literature ([26]). The Kravchuk function is then defined in terms of the polynomials as

$$K_n^{(p)}(x,N) = d_n^{-1} \sqrt{\rho(x)} k_n(x;p,N)$$

with the normalization

$$d_n^2 = \binom{N}{n} (p(1-p))^n$$

and the weighting (in order to ensure orthogonality)

$$\rho(x) = \binom{N}{x} p^x (1-p)^{N-x}$$

To move from this common notation to that used in Eq. (39), we take p = 1/2 and suppress it as an argument, N = s + 1, and  $x = n_1$ .

# APPENDIX C: MATRIX REPRESENTATIONS OF THE OSCILLATOR OPER-ATORS

In the position-resolution basis, each operator is naturally written with three indices, the first describing its dimension and the second two describing the operator's effect at that dimension. For simplicity, we will consider the operators acting on a d = s + 1 fixeddimensional state so that we can suppress the first index and write each operator as a matrix. With these assumptions, the  $A_1$ ,  $A_2$ ,  $A_1^{\dagger}$ , and  $A_2^{\dagger}$  operators each have a single non-zero band.  $A_1$  is an  $s \times s + 1$  matrix with nonzero elements along its  $1 \times 1$  diagonal:

$$A_{1} = \left(\begin{array}{ccccc} \sqrt{s} & 0 & 0 & \dots & 0\\ 0 & \sqrt{s-1} & 0 & \dots & 0\\ \vdots & \vdots & \ddots & 0\\ 0 & 0 & 1 & 0\\ & & & & \\ & & & \\ & & & \\ & &$$

Let us define a compact notation for such matrices  $M = \text{band}_{p \times q} \left[ l \times m \to (f_j)_{j=0}^N \right]$ , where  $p \times q$  indicates the size of the matrix,  $l \times m$  indicates the location of the first entry of the nonzero band, and  $(f_j)_{j=0}^N$  is the sequence which populates that band. In this notation,

$$A_1 = \operatorname{band}_{s \times s+1} \left[ 1 \times 1 \to \left( \sqrt{s-j} \right)_{j=0}^{s-1} \right].$$
(49)

Writing the other operators in this notation, we have

$$A_2 = \operatorname{band}_{s \times s+1} \left[ 1 \times 2 \to \left( \sqrt{j+1} \right)_{j=0}^{s-1} \right], \tag{50}$$

$$A_1^{\dagger} = \operatorname{band}_{s+2\times s+1} \left[ 1 \times 1 \to \left( \sqrt{s+1-j} \right)_{j=0}^s \right], \tag{51}$$

$$A_2^{\dagger} = \operatorname{band}_{s+2\times s+1} \left[ 2 \times 1 \to \left(\sqrt{j+1}\right)_{j=0}^s \right].$$
(52)

Note that in this notation,  $A_j^{\dagger}$  is not just the Hermitian conjugate of  $A_j$  since  $A_j^{\dagger}$  increases a state's dimension, while  $A_j$  decreases it. The same will be true of F and  $F^{\dagger}$ .

We can generalize this notation to band diagonal matrices with multiple nonzero bands. Thus,

$$H_{I} = -\frac{1}{2} \text{band}_{s+1 \times s+1} \left[ 1 \times 2 \to \left( \sqrt{s-j}\sqrt{j+1} \right)_{j=0}^{s-1}, 2 \times 1 \to \left( \sqrt{s-j}\sqrt{j+1} \right)_{j=0}^{s-1} \right].$$
(53)

The energy lowering and raising operators become

$$D = \text{band}_{s+1 \times s+1} \left[ 1 \times 1 \to (s - 2j)_{j=0}^{s}, 1 \times 2 \to \left(\sqrt{s - j}\sqrt{j + 1}\right)_{j=0}^{s-1}, \\ 2 \times 1 \to \left(-\sqrt{s - j}\sqrt{j + 1}\right)_{j=0}^{s-1} \right]$$
(54)  
$$D^{\dagger} = \text{band}_{s+1 \times s+1} \left[ 1 \times 1 \to (s - 2j)_{j=0}^{s}, 1 \times 2 \to \left(-\sqrt{s - j}\sqrt{j + 1}\right)_{j=0}^{s-1}, \\ 2 \times 1 \to \left(\sqrt{s - j}\sqrt{j + 1}\right)_{j=0}^{s-1} \right],$$
(55)

The position operator is simply the diagonal matrix

$$X = \operatorname{band}_{s+1 \times s+1} \left[ 1 \times 1 \to (s - 2j)_{j=0}^{s} \right],$$
(56)

while the momentum operator is

$$P = \text{band}_{s+1 \times s+1} \left[ 1 \times 2 \to \left( -i\sqrt{s-j}\sqrt{j+1} \right)_{j=0}^{s-1}, \\ 2 \times 1 \to \left( i\sqrt{s-j}\sqrt{j+1} \right)_{j=0}^{s-1} \right].$$

### APPENDIX D: COHERENT STATE POSITION EXPECTATION VALUE

We seek to calculate  $\langle \beta | X(t) | \beta \rangle$ , which can be expanded to

$$\langle \beta | X(t) | \beta \rangle = \langle 0, s | \hat{\mathcal{D}}(\beta) e^{iHt} X_s e^{-iHt} \hat{\mathcal{D}}(\beta) | 0, s \rangle , \qquad (57)$$

where our X(t) in the Heisenberg picture is transformed to  $e^{iHt}X_se^{-iHt}$  in the Schrödinger picture. We will begin by noting that

$$\hat{\mathcal{D}}(\beta) |0,s\rangle = \frac{1}{\mathcal{N}^{1/2}} e^{\frac{\beta}{2|\beta|} \tan(2|\beta|)D^{\dagger}} |0,s\rangle,$$
(58)

as noted in Eq. (45), where  $\mathcal{N}$  is some normalization. Because we have a finite basis, this exponential can be expanded as a finite sum,

$$\hat{\mathcal{D}}(\beta) |0,s\rangle = \frac{1}{\mathcal{N}^{1/2}} \left( |0,s\rangle$$
(59)

$$+\sum_{j=1}^{s} \left( \frac{\beta}{|\beta|} \tan(2|\beta|) \right)^{j} \frac{1}{j!} \prod_{k=0}^{j-1} \left[ 2\sqrt{s-k}\sqrt{1+k} \right] |k+1,s\rangle \right), \tag{60}$$

where we've made use of the normalizations of Eqs. (19) and (18). The product may rewritten using factorial notation,

$$\prod_{k=0}^{j-1} \left[ 2\sqrt{s-k}\sqrt{1+k} \right] = 2^j \sqrt{j!} \sqrt{\frac{s!}{(s-j)!}},\tag{61}$$

so that we have

$$\hat{\mathcal{D}}(\beta) |0,s\rangle = \frac{1}{\mathcal{N}^{1/2}} \left( |0,s\rangle \right)$$
(62)

$$+\sum_{j=1}^{s} \left(\frac{2\beta}{|\beta|} \tan(2|\beta|)\right)^{j} \sqrt{\binom{s}{j}} |j,s\rangle \right).$$
(63)

Finding the normalization  $\mathcal{N}$  simply requires calculating

$$\left|\hat{\mathcal{D}}(\beta)\left|0,s\right\rangle\right|^{2} = 1,\tag{64}$$

 $\mathbf{SO}$ 

$$\mathcal{N} = 1 + \sum_{j=1}^{s} \tan^{2j}(2|\beta|) 4^{j} \binom{s}{j},$$
(65)

which simplifies to

$$\mathcal{N} = \left(1 + 4\tan^2(2|\beta|)\right)^s.$$
(66)

Because we are working the in the energy-resolution basis, multiplying Eq. (63) by  $U(t) = e^{-iHt}$  is simple. Each state  $|n, s\rangle$  is simply multiplied by  $e^{-int}$ :

$$e^{-iHt}\hat{\mathcal{D}}(\beta)|0,s\rangle = \frac{1}{\mathcal{N}^{1/2}} \left( e^{ist}|0,s\rangle \right)$$
(67)

$$+\sum_{j=1}^{s} e^{i(s-2j)t} \left(\frac{2\beta}{|\beta|} \tan(2|\beta|)\right)^{j} \sqrt{\binom{s}{j}} |j,s\rangle \right).$$
(68)

Next, multiplying by position operator,  $X_s = \frac{1}{2} \left( D^{\dagger} + D \right)$ , we have

$$X_s e^{-iHt} \hat{\mathcal{D}}(\beta) |-s,s\rangle = \frac{1}{\mathcal{N}^{1/2}} \left( \sqrt{s} e^{ist} |-s+2,s\rangle \right)$$
(69)

$$+\sum_{j=1}^{s} e^{i(s-2j)t} \left(\frac{2\beta}{|\beta|} \tan(2|\beta|)\right)^{j} \sqrt{\binom{s}{j}}$$
(70)

$$\times \left[ \sqrt{s - j} \sqrt{j + 1} | j + 1, s \right]$$
(71)

$$+\sqrt{s-j+1}j^{1/2}|j-1,s\rangle\Big]\bigg).$$
 (72)

Further multiplying by  $e^{iHt}$  and simplifying,

$$e^{iHt}X_s e^{-iHt}\hat{\mathcal{D}}(\beta)|0,s\rangle = \frac{1}{\mathcal{N}^{1/2}} \left(\frac{2\beta}{|\beta|} \tan(2|\beta|) s e^{-2it}|0,s\rangle + \sum_{j=1}^s \left(\frac{2\beta}{|\beta|} \tan(2|\beta|)\right)^{j-1} \sqrt{\binom{s}{j}} |j,s\rangle \times \left[j e^{2it} + 4\left(\frac{\beta}{|\beta|} \tan(2|\beta|)\right)^2 (s-j) e^{-2it}\right]\right).$$

Finally, we can calculate

$$\begin{split} \langle \beta | X(t) | \beta \rangle &= \frac{1}{\mathcal{N}} \left( \frac{2\beta}{|\beta|} \tan(2|\beta|) s e^{-2it} + \sum_{j=1}^{s} 2^{2j-1} \frac{\beta^{*}}{|\beta|} \tan^{2j-1}(2|\beta|) \binom{s}{j} \\ &\times \left[ j e^{2it} + 4 \left( \frac{\beta}{|\beta|} \tan(2|\beta|) \right)^{2} (s-j) e^{-2it} \right] \right) \\ &= \frac{1}{\mathcal{N}} 2s |\beta|^{-1} \tan(2|\beta|) \left( \beta e^{-2it} + \beta^{*} e^{2it} \right) \left( 1 + 4 \tan^{2}(2|\beta|) \right)^{s-1} \\ &= 2s |\beta|^{-1} \tan(2|\beta|) \left( \beta e^{-2it} + \beta^{*} e^{2it} \right) \left( 1 + 4 \tan^{2}(2|\beta|) \right)^{-1} \\ &= 4s \tan(2|\beta|) \cos(2t - \arg(\beta)) \left( 1 + 4 \tan^{2}(2|\beta|) \right)^{-1} \\ &= s \cos(2t - \arg(\beta)) \sin(4|\beta|), \end{split}$$

which is the same as Eq. (46).

- [1] N. M. Atakishiev, E. I. Jafarov, S. M. Nagiyev, and K. B. Wolf, Rev. Mex. Fís. 44, 235 (1998).
- [2] N. M. Atakishiev and S. K. Suslov, Theoretical and Mathematical Physics 85, 1055 (1990).

- [3] M. Lorente, New developments on fundamental problems in quantum physics (Kluwer Academic, 1997) Chap. Quantum Mechanics on Discrete Space and Time, pp. 213–224.
- [4] M. Christandl, N. Datta, A. Ekert, and A. J. Landahl, Phys. Rev. Lett. 92, 187902
   10.1103/PhysRevLett.92.187902 (2004).
- [5] T. Shi, Y. Li, Z. Song, and C.-P. Sun, Phys. Rev. A **71**, 032309 (2005).
- [6] Y. Shi, A. R. Castelli, X. Wu, I. Joseph, V. Geyko, F. R. Graziani, S. B. Libby, J. B. Parker,
  Y. J. Rosen, L. A. Martinez, and J. L. DuBois, Phys. Rev. A 103, 062608 (2021).
- [7] M. M. R. Miceli, in 2019 New York Scientific Data Summit (IEEE, 2019) pp. 1–4.
- [8] V. K. Jain, B. K. Behera, and P. K. Panigrahi, Quantum Studies: Mathematics and Foundations 8, 375 (2021).
- [9] M. Lorente, Physics Letters A **285**, 119 (2001).
- [10] N. M. Atakishiyev, A. U. Klimyk, and K. B. Wolf, Journal of Physics A: Mathematical and Theoretical 41, 085201 (2008).
- [11] Y. Shi, H. Qin, and N. J. Fisch, Physical Review E 96, 023204 (2017).
- [12] Y. Shi, Plasma Physics in Strong Field Regimes, Ph.D. thesis, Princeton University (2018).
- [13] Y. Shi, H. Qin, and N. J. Fisch, Physics of Plasmas 28, 042104 (2021).
- [14] M. Q. May and H. Qin, Phys. Rev. A **107**, 062204 (2023).
- [15] M. Q. May and H. Qin, ArXiV Preprint 10.48550/arXiv.2401.03995 (2024).
- [16] J. Schwinger, On Angular Momentum, Technical Report NYO-3071 (US Atomic Energy Commission, 1952).
- [17] E. I. Jafarov, N. I. Stoilova, and J. V. der Jeugt, J. Phys. A: Math. Theor. 44, 265203 (2011).
- [18] N. M. Atakishiyev, G. S. Pogosyan, L. E. Vicent, and K. B. Wolf, J. Phys. A: Math. Gen. 34, 9381 (2001).
- [19] A. F. Nikiforov, V. B. Uvarov, and S. K. Suslov, *Classical Orthogonal Polynomials of a Discrete Variable* (Springer Berlin, Heidelberg, 1991).
- [20] K. B. Wolf and G. Krötzsch, J. Op. Soc. Am. A 24, 10.1364/JOSAA.24.000651 (2007).
- [21] K. Uriostegui, J. Op. Soc. Am. B 40, 1121 (2023).
- [22] J. M. Radcliffe, J. Phys. A: Gen. Phys. 4, 313 (1971).
- [23] F. T. Arecchi, E. Courtens, R. Gilmore, and H. Thomas, Phys. Rev. A 6, 2211 (1972).
- [24] G. Draganescu, A. Messinaband, and A. Napoli, Journal of Modern Optics 56, 508 (2009).
- [25] D. R. Truax, Phys. Rev. D **31**, 10.1103/PhysRevD.31.1988 (1985).

[26] T. H. Koornwinder, SIAM J. Math. Anal. 13, 1011 (1982).