

A PROOF-THEORETIC METATHEOREM FOR NONLINEAR SEMIGROUPS GENERATED BY AN ACCRETIVE OPERATOR AND APPLICATIONS

NICHOLAS PISCHKE

Department of Mathematics, Technische Universität Darmstadt,
Schlossgartenstraße 7, 64289 Darmstadt, Germany,

E-mail: pischke@mathematik.tu-darmstadt.de

ABSTRACT. We further develop the theoretical framework of proof mining, a program in mathematical logic that seeks to quantify and extract computational information from *prima facie* ‘non-computational’ proofs from the mainstream literature. To that end, we establish logical metatheorems that allow for the treatment of proofs involving nonlinear semigroups generated by an accretive operator, structures which in particular arise in the study of the solutions and asymptotic behavior of partial differential equations. In that way, the here established metatheorems facilitate a theoretical basis for the application of methods from the proof mining program to the wide variety of mathematical results established in the context of that theory since the 1960’s. We in particular illustrate the applicability of the new systems and their metatheorems introduced here by providing two case studies on two central results due to Reich and Plant, respectively, on the asymptotic behavior of said semigroups and the resolvents of their generators where we derive rates of convergence for the limits involved which are, moreover, polynomial in all data.

Keywords: Proof mining; Metatheorems; Accretive operators; Nonlinear semigroups.

MSC2020 Classification: 03F10, 03F35, 47H06, 37L05, 47H20

1. INTRODUCTION

Proof mining is a program in mathematical logic that seeks to extract computational information, like (uniform) witnesses or bounds, from *prima facie* ‘non-computational’ proofs from the mainstream literature. Historically, this endeavor goes back conceptually to Georg Kreisel’s program of *unwinding of proofs* from the 1950’s [35, 36]. In its modern form, which relies on the use of well-known proof interpretations like *negative translations*, Kreisel’s *modified realizability* and Gödel’s *functional (Dialectica) interpretation*, it has been successfully developed since the 1990s by the groundbreaking work of Ulrich Kohlenbach (see in particular the early works [21, 22]) and his collaborators.

The main applications of these methods are today found in the areas of nonlinear analysis and optimization and in that context, typical additional quantitative information extractable from non-constructive proofs have proved to be intimately connected with other approaches to finitary analysis. In particular, these logical methods in very general situations guarantee the existence of so-called *rates of metastability* (in the sense of Terence Tao [55, 56]) for a finite quantitative account of a convergence result.

The development of this modern period of proof mining is detailed comprehensively up to the year 2008 in the monograph [26], with early progress surveyed in [32] and recent progress in applications to the fields of nonlinear analysis and optimization surveyed in [27].

The whole methodology of proof mining therein crucially relies on so-called *general logical metatheorems* which guarantee the existence and quantify the complexity of such additional quantitative information.¹ Further, besides merely guaranteeing existence, these metatheorems provide an algorithmic approach towards actually extracting these quantitative results.

In the context of nonlinear analysis, one of proof mining’s most successful fields of application, it is the theory of partial differential equations and neighboring fields like evolution equations where one so far still lacks a variety of proof mining applications with there so far only being two real case studies [28, 46]. Already since the pioneering studies of Browder [8], Kato [20] and Komura [34], two major tools in the study of nonlinear evolution equations have been the theory of nonlinear semigroups and their generators as well as the theory of accretive operators together with their correspondence via analogs of the Hille-Yosida theorem.

Date: April 5, 2023.

¹Examples of such metatheorems may be found in [15, 16, 18, 25, 30, 31, 38, 39, 42, 47, 54, 42], as well as [26] for the metatheorems obtained via (modifications of) Gödel’s *Dialectica* interpretation, and [13] for subsequent metatheorems obtained via the bounded functional interpretation [14] due to F. Ferreira and P. Oliva.

One of the most important basic results in that context is the representation theorem due to Crandall and Liggett [11] of the solution semigroup associated with the Cauchy problem

$$(\dagger) \quad \begin{cases} u'(t) \in -Au(t), & 0 < t < \infty \\ u(0) = x \end{cases}$$

over a Banach space X for a given set-valued accretive operator $A : X \rightarrow 2^X$. It is straightforward to show that any solution² is unique as A is accretive and if the system is solvable³, then one can consider the operator $S(t)x$ on $\text{dom}A$ induced by the solutions $u_x(t)$ to (\dagger) with initial values $x \in \text{dom}A$. As this operator is continuous in x , one can consider the resulting extension to $\overline{\text{dom}A}$ which in that way generates the semigroup $\mathcal{S} = \{S(t) \mid t \geq 0\}$ on $\overline{\text{dom}A}$ associated with (\dagger) . As shown by Brezis and Pazy [7], this solution semigroup, if existent, has a particular fundamental representation in terms of a so-called exponential formula:

$$u_x(t) = \lim_{n \rightarrow \infty} \left(\text{Id} + \frac{t}{n}A \right)^{-n} x.$$

As shown subsequently by Crandall and Liggett [11], this formula actually always generates a nonexpansive semigroup on $\overline{\text{dom}A}$ and thus facilitates a general study even in the absence of solutions to (\dagger) .

Since the 1970s, an extensive range of results has been established in the theory of these semigroups and the initial value problems in the sense of (\dagger) associated with them, in particular in regard to the asymptotic behavior of the solutions of these differential equations, their connection and use in the study of partial differential equations and their use in the study of zeros of accretive operators (see [1, 2, 3, 40, 43], among many more).

In this paper, we extend the state-of-the-art of the underlying logical approach to proof mining to be applicable to proofs which make use of these nonlinear semigroups generated by an accretive operator via the exponential formula. In particular, we establish logical metatheorems in the vein of the previously discussed results that guarantee and quantify the computational content of theorems pertaining to nonlinear semigroups generated by such an accretive operator. For that, we introduce new underlying logical systems that extend those developed for the treatment of accretive operators on normed spaces [47] by carefully selecting additional constants and corresponding axioms such that proofs from the main stream literature become formalizable. To that end, we show that the initial key properties of these semigroups generated by an accretive operator can be formally proved in these systems.

These logical results provide a formal basis for the previous proof mining application [28] carried out in the context of nonlinear semigroups associated with differential equations and thus remove the ad-hoc nature surrounding it. Even further however, these results are expected to lead to many new case studies for proof mining in the context of that theory. In that vein, this paper provides two case studies on results due to Plant [50] and Reich [53] for the asymptotic behavior of these semigroups and in that context, under suitable quantitative translations of the assumptions used in the respective results, we are able to extract rates of convergence for the limits involved which are moreover polynomial in all data. In particular, we want to note that full rates of convergence are obtained here despite the fact that the sequence in question is not monotone and that the original proof is classical. This is due to a logical particularity that will be discussed after the extractions. The applicability of the present results is further substantiated by the fact that they also provide the logical basis for the previously mentioned application [47] as well as for the forthcoming quantitative analysis given in [49] in the context of results on the asymptotic behavior of these systems of differential equations due to Pazy [45], Nevanlinna and Reich [41], Xu [57] as well as Poffald and Reich [51]. At last, we want to mention that the whole logical apparatus for the treatment of these semigroups as developed in this paper requires some new technical tools. We expect that also these new logical approaches to these various notions in nonlinear analysis developed here will be of use in other circumstances than the ones described in this paper.

2. PRELIMINARIES

2.1. Nonlinear semigroups and the Crandall-Liggett formula. The main objects of concern in this paper are the aforementioned nonlinear (nonexpansive) semigroups:

Definition 2.1. Let C be a closed subset of X . A function $S : [0, \infty) \times C \rightarrow C$ is a (*nonexpansive*) *semigroup on C* if

²A function $u : [0, \infty) \rightarrow X$ is a solution of (\dagger) if $u(0) = x$, $u(t)$ is absolutely continuous, differentiable almost everywhere in $(0, \infty)$ and satisfies (\dagger) almost everywhere. Note that this also sometimes called a strong solution.

³As shown by Crandall and Liggett [11], this is (for strong solutions) in general not the case even for $x \in \text{dom}A$ and A m -accretive.

- (1) $S(t+s)x = S(t)S(s)x$ for all $x \in C$ and all $t, s \geq 0$,
- (2) $S(0)x = x$ for all $x \in C$,
- (3) $S(t)x$ is continuous in $t \geq 0$ for every $x \in C$,
- (4) $\|S(t)x - S(t)y\| \leq \|x - y\|$ for all $t \geq 0$ and all $x \in C$.

As discussed in the introduction already, these semigroups frequently arise in the study of differential and evolution equations as is e.g. exemplified by the initial value problem (†) and in particular, by the results of Crandall and Liggett [11], the expression of the exponential formula discussed there always generates such a semigroup on $\overline{\text{dom}A}$ which will be the main object of study of this paper. Concretely, the following result was established:

Theorem 2.2 (Crandall and Liggett [11]). *Let X be a Banach space and A an accretive operator on X such that there exists a $\lambda_0 > 0$ with*

$$\overline{\text{dom}A} \subseteq \text{ran}(Id + \lambda A) \text{ for all } \lambda \in (0, \lambda_0].$$

Then

$$S(t)x := \lim_{n \rightarrow \infty} \left(Id + \frac{t}{n} A \right)^{-n} x$$

exists for all $x \in \overline{\text{dom}A}$ and $t \geq 0$ and $\mathcal{S} = \{S(t) \mid t \geq 0\}$ is a nonlinear semigroup on $\overline{\text{dom}A}$.

We call \mathcal{S} as defined above the semigroup generated by A (via the exponential or Crandall-Liggett formula).

In terms of a logical treatment of these semigroups generated by an accretive operator, all of the later logical considerations naturally depend on the underlying theory of accretive operators over Banach spaces. In that vein, we logically crucially rely on the basic systems introduced in [47] for the treatment of those accretive operators in the context of the extended systems of finite type commonly used in proof mining and we thus detail those in the next subsection.

2.2. The basic system. The basic system for accretive operators on normed spaces relies on the system $\mathcal{A}^\omega[X, \|\cdot\|]$ introduced in [16, 25] as an underlying system for classical analysis over abstract normed spaces in all finite types T^X defined by

$$\mathbb{N}, X \in T^X, \quad \rho, \tau \in T^X \Rightarrow \rho \rightarrow \tau \in T^X.$$

We refer to those works, and to [26] in general, for a precise exposition on the definition and basic properties of this and related systems. Accordingly, we mostly follow the notation used there as well as in [47] (besides of using the above notation for the types which is of a more intuitive form and, in that vein, we also write $\mathbb{N}^{\mathbb{N}}$ for $\mathbb{N} \rightarrow \mathbb{N}$).

In the context of these finite type systems, real numbers are as usual represented as fast-converging Cauchy sequences of rationals with a fixed rate. These are encoded via number theoretic functions, i.e. objects of type $\mathbb{N}^{\mathbb{N}}$, and on the level of that representation, one the can introduce the usual arithmetic operations and relations. Concretely, the relations $=_{\mathbb{R}}, \leq_{\mathbb{R}}$ operating on these type $\mathbb{N}^{\mathbb{N}}$ codes can then be chosen to be Π_1^0 -formulas while $<_{\mathbb{R}}$ can be chosen to be a Σ_1^0 -formula.

In general, we will omit the type of real numbers for arithmetical operations to make everything more readable. In proofs, we will almost always omit most types as to not distract from the general ideas and patterns.

The starting point for our new systems will then be the theory \mathcal{V}_p^ω for the treatment of accretive operators on normed spaces as introduced in [47]. This theory is defined as the extension of the theory $\mathcal{A}^\omega[X, \|\cdot\|]$ with the additional constants

- χ_A of type $X \rightarrow (X \rightarrow \mathbb{N})$ for the graph of the operator A ,
- J^{X^A} of type $\mathbb{N}^{\mathbb{N}} \rightarrow (X \rightarrow X)$ for the resolvent J^A ,
- $c_X, \tilde{\gamma}, m_{\tilde{\gamma}}$ of types $X, \mathbb{N}^{\mathbb{N}}, \mathbb{N}$, respectively, for a technical purpose in the majorization of the constants later on,

together with the corresponding axioms

- (I) $\forall x^X, y^X (\chi_A xy \leq_{\mathbb{N}} 1)$,
- (II) $\forall \gamma^{\mathbb{N}^{\mathbb{N}}}, x^X (\gamma >_{\mathbb{R}} 0 \wedge \exists y^X (\gamma^{-1}(x -_X y) \in Ay) \rightarrow \gamma^{-1}(x -_X J_\gamma^A x) \in A(J_\gamma^A x))$,
- (III) $\left\{ \begin{array}{l} \forall x^X, y^X, u^X, v^X, \lambda^{\mathbb{N}^{\mathbb{N}}} (u \in Ax \wedge v \in Ay \\ \rightarrow \|x -_X y +_X |\lambda|(u -_X v)\|_X \geq_{\mathbb{R}} \|x -_X y\|_X), \end{array} \right.$
- (IV) $\tilde{\gamma} \geq_{\mathbb{R}} 2^{-m_{\tilde{\gamma}}}$,
- (V) $\forall \gamma^{\mathbb{N}^{\mathbb{N}}} (\gamma >_{\mathbb{R}} 0 \rightarrow \gamma^{-1}(c_X -_X J_\gamma^A c_X) \in A(J_\gamma^A c_X))$.

Further, as in [47], we use the abbreviation

$$x \in \text{dom}(J_\gamma^A) \equiv \exists y^X (\gamma^{-1}(x -_X y) \in Ay)$$

and in that way can recognize the second axiom as stating

$$\forall \gamma^{\mathbb{N}^{\mathbb{N}}}, x^X (\gamma >_{\mathbb{R}} 0 \wedge x \in \text{dom}(J_\gamma^A) \rightarrow \gamma^{-1}(x -_X J_\gamma^A x) \in A(J_\gamma^A x))$$

which in that way specifies the behavior of the resolvent on its domain as dictated by its defining equality $J_\gamma^A = (Id + \gamma A)^{-1}$. We refer to [47] for an (extensive) discussion of the motivation for this axiomatization and here just note the restriction put in place by axiom (V) that the constant c_X designates a common element of the domains of all resolvents J_γ^A for $\gamma > 0$. As discussed in [47], this assumption is easily satisfied for most applications which in particular include those situations where one assumes a range condition like

$$\text{dom}A \subseteq \bigcap_{\gamma > 0} \text{ran}(Id + \gamma A)$$

which will be the case in this work in particular as will be discussed in the coming subsections.

As shown in [47], the main parts of the basic theory of accretive operators and in particular their resolvents can then be immediately formally derived in the system \mathcal{V}_p^ω and we give an indication of that in the following lemma. For that, we also formally introduce the Yosida approximate A_γ defined via

$$A_\gamma := \frac{1}{\gamma}(Id - J_\gamma^A)$$

in the context of the formal system by treating $A_\gamma x$ as an abbreviation for the term $\gamma^{-1}(x -_X J_\gamma^A x)$.⁴

Lemma 2.3 ([47]). *The system \mathcal{V}_p^ω proves:*

(1) J_γ^A is unique for any $\gamma > 0$, i.e.

$$\forall \gamma^{\mathbb{N}^{\mathbb{N}}}, p^X, x^X (\gamma >_{\mathbb{R}} 0 \wedge \gamma^{-1}(x -_X p) \in Ap \rightarrow p =_X J_\gamma^A x).$$

(2) J_γ^A is firmly nonexpansive for any $\gamma > 0$ (on its domain), i.e.

$$\begin{aligned} & \forall \gamma^{\mathbb{N}^{\mathbb{N}}}, r^{\mathbb{N}^{\mathbb{N}}}, x^X, y^X (\gamma >_{\mathbb{R}} 0 \wedge x \in \text{dom}(J_\gamma^A) \wedge y \in \text{dom}(J_\gamma^A) \wedge r >_{\mathbb{R}} 0 \\ & \rightarrow \|J_\gamma^A x -_X J_\gamma^A y\|_X \leq_{\mathbb{R}} \|r(x -_X y) +_X (1 - r)(J_\gamma^A x -_X J_\gamma^A y)\|_X). \end{aligned}$$

(3) J_γ^A is nonexpansive for any $\gamma > 0$ (on its domain), i.e.

$$\begin{aligned} & \forall \gamma^{\mathbb{N}^{\mathbb{N}}}, x^X, y^X (\gamma >_{\mathbb{R}} 0 \wedge x \in \text{dom}(J_\gamma^A) \wedge y \in \text{dom}(J_\gamma^A) \\ & \rightarrow \|x -_X y\|_X \geq_{\mathbb{R}} \|J_\gamma^A x -_X J_\gamma^A y\|_X). \end{aligned}$$

(4) J^A is extensional in both arguments (on its domain), i.e.

$$\begin{aligned} & \forall \gamma^{\mathbb{N}^{\mathbb{N}}} >_{\mathbb{R}} 0, x^X, x'^X (x \in \text{dom}(J_\gamma^A) \wedge x' \in \text{dom}(J_\gamma^A) \wedge x =_X x' \rightarrow J_\gamma^A x =_X J_\gamma^A x'), \\ & \forall \gamma^{\mathbb{N}^{\mathbb{N}}} >_{\mathbb{R}} 0, \gamma'^{\mathbb{N}^{\mathbb{N}}} >_{\mathbb{R}} 0, x^X (x \in \text{dom}(J_\gamma^A) \wedge x \in \text{dom}(J_{\gamma'}^A) \wedge \gamma =_{\mathbb{R}} \gamma' \rightarrow J_\gamma^A x =_X J_{\gamma'}^A x). \end{aligned}$$

(5)

$$\begin{aligned} & \forall \gamma^{\mathbb{N}^{\mathbb{N}}}, \lambda^{\mathbb{N}^{\mathbb{N}}}, x^X (\gamma >_{\mathbb{R}} 0 \wedge \lambda >_{\mathbb{R}} 0 \wedge x \in \text{dom}(J_\lambda^A) \\ & \rightarrow J_\lambda^A x =_X J_\gamma^A \left(\frac{\gamma}{\lambda} x +_X \left(1 - \frac{\gamma}{\lambda}\right) J_\gamma^A x \right)). \end{aligned}$$

(6)

$$\begin{aligned} & \forall \gamma^{\mathbb{N}^{\mathbb{N}}}, \lambda^{\mathbb{N}^{\mathbb{N}}}, x^X (\gamma >_{\mathbb{R}} 0 \wedge \lambda >_{\mathbb{R}} 0 \wedge x \in \text{dom}(J_\gamma^A) \wedge x \in \text{dom}(J_\lambda^A) \\ & \rightarrow \|x -_X J_\gamma^A x\|_X \leq_{\mathbb{R}} \left(2 + \frac{\gamma}{\lambda}\right) \|x -_X J_\lambda^A x\|_X). \end{aligned}$$

⁴Note the discussion given in [47] on the subtleties of the reciprocal of real arithmetic in the context of these systems of finite types and the resulting subtleties of the above definition.

(7) A_γ is $2\gamma^{-1}$ -Lipschitz continuous for any $\gamma > 0$, i.e.

$$\begin{aligned} \forall \gamma^{\mathbb{N}}, x^X, y^X \left(\gamma >_{\mathbb{R}} 0 \wedge x \in \text{dom}(J_\gamma^A) \wedge y \in \text{dom}(J_\gamma^A) \right. \\ \left. \rightarrow \|A_\gamma x - A_\gamma y\|_X \leq_{\mathbb{R}} 2\gamma^{-1} \|x - y\|_X \right). \end{aligned}$$

(8) $A_\gamma x$ is bounded by any $y \in Ax$ for any $\gamma > 0$, i.e.

$$\forall \gamma^{\mathbb{N}}, x^X, y^X \left(\gamma >_{\mathbb{R}} 0 \wedge y \in Ax \wedge x \in \text{dom}(J_\gamma^A) \rightarrow \|A_\gamma x\|_X \leq_{\mathbb{R}} \|y\|_X \right).$$

In particular $\|x - J_\gamma x\|_X \leq_{\mathbb{R}} \gamma \|y\|_X$.

2.3. The basic bound extraction theorems. The main result established in [47] is the logical metatheorem on bound extractions akin to the usual metatheorems of proof mining. Throughout the paper, we do not go into explicit detail regarding the proof of any bound extraction theorem and only provide sketches for the relevant additions and changes as the proofs otherwise follow the usual standard outline of most bound extraction results in proof mining established in [16, 25].

The prevalent central ingredient that we will focus on later in the context of the new bound extraction results will be the majorizability of the new constants used to treat nonlinear semigroups. Under majorizability, we will here understand the extension due to [16, 25] of the strong majorizability of Bezem [4] (which in turn builds on Howard's majorizability [19]) to the new types in T^X . The fundamental paradigm of the bound extraction results is then that one achieves uniform bounds even in the absence of compactness by majorizing the bounds extracted by the underlying functional interpretation.

In that way, following [16, 25], majorants of objects with types from T^X will be objects with types from T related by the following projection:

Definition 2.4 ([16]). Define $\hat{\tau} \in T$, given $\tau \in T^X$, by recursion on the structure via

$$\hat{\mathbb{N}} := \mathbb{N}, \hat{X} := X, \widehat{\xi \rightarrow \tau} := \hat{\xi} \rightarrow \hat{\tau}.$$

The majorizability relation is then defined in tandem with the model of all majorizable functionals in the sense of the following definition.

Definition 2.5 ([16, 25]). Let $(X, \|\cdot\|)$ be a non-empty normed space. The structure $\mathcal{M}^{\omega, X}$ and the majorizability relation \succsim_τ are defined by

$$\left\{ \begin{array}{l} M_{\mathbb{N}} := \mathbb{N}, n \succsim_{\mathbb{N}} m := n \geq m \wedge n, m \in \mathbb{N}, \\ M_X := X, n \succsim_X x := n \geq \|x\| \wedge n \in M_{\mathbb{N}}, x \in M_X, \\ x^* \succsim_{\xi \rightarrow \tau} x := x^* \in M_{\hat{\tau}}^{M_\xi} \wedge x \in M_\tau^{M_\xi} \\ \quad \wedge \forall y^* \in M_{\hat{\xi}}, y \in M_\xi(y^* \succsim_\xi y \rightarrow x^* y^* \succsim_\tau xy) \\ \quad \wedge \forall y^*, y \in M_{\hat{\xi}}(y^* \succsim_{\hat{\xi}} y \rightarrow x^* y^* \succsim_{\hat{\tau}} x^* y), \\ M_{\xi \rightarrow \tau} := \left\{ x \in M_\tau^{M_\xi} \mid \exists x^* \in M_{\hat{\tau}}^{M_\xi} : x^* \succsim_{\xi \rightarrow \tau} x \right\}. \end{array} \right.$$

At a high level, the proofs of most bound extraction theorems then proceed as follows: using a variant of Gödel's functional interpretation [17] and a negative translation (e.g. [37]), realizers are extracted from classical proofs of (essentially) $\forall\exists$ -theorems. These realizers are then majorized to provide respective bounds which are validated in the model of majorizable functionals $\mathcal{M}^{\omega, X}$. If the types of all objects are low enough, one can then recover to the truth of the respective bound in the usual set-theoretic standard model $\mathcal{S}^{\omega, X}$ of the underlying language defined by $S_{\mathbb{N}} := \mathbb{N}$, $S_X := X$ and

$$S_{\xi \rightarrow \tau} := S_\tau^{S_\xi}.$$

The resulting metatheorem for the case of the theory \mathcal{V}^ω then takes the form of the following theorem. Here, we followed the names and notational conventions established in [16, 25] (see also [26]) regarding so-called 'admissible' types. which provide a formal perspective of the previously mentioned vague notion of 'low enough'.

Theorem 2.6 ([47]). Let τ be admissible, δ be of degree 1 and s be a closed term of \mathcal{V}^ω of type $\delta \rightarrow \sigma$ for admissible σ . Let $B_\forall(x, y, z, u)/C_\exists(x, y, z, v)$ be \forall -/ \exists -formulas of \mathcal{V}_p^ω with only $x, y, z, u/x, y, z, v$ free. Let Δ be a set of formulas of the form $\forall \underline{a} \exists \underline{b} \leq_{\underline{\sigma}} \underline{r} \forall \underline{c} \exists F_{qf}(\underline{a}, \underline{b}, \underline{c})$ where F_{qf} is quantifier-free, the types in $\underline{\delta}$, $\underline{\sigma}$ and $\underline{\gamma}$ are admissible and \underline{r} are tuples of closed terms of appropriate type. If

$$\mathcal{V}_p^\omega + \Delta \vdash \forall x^\delta \forall y \leq_\sigma s(x) \forall z^\tau (\forall u^\mathbb{N} B_\forall(x, y, z, u) \rightarrow \exists v^\mathbb{N} C_\exists(x, y, z, v)),$$

then one can extract a partial functional $\Phi : S_\delta \times S_{\hat{\tau}} \times \mathbb{N} \rightarrow \mathbb{N}$ which is total and (bar-recursively) computable on $\mathcal{M}_\delta \times \mathcal{M}_{\hat{\tau}} \times \mathbb{N}$ and such that for all $x \in S_\delta$, $z \in S_{\hat{\tau}}$, $z^* \in S_{\hat{\tau}}$ and all $n \in \mathbb{N}$, if $z^* \succeq z$ and $n \geq_{\mathbb{R}} \|c_X - J_{\hat{\tau}}^A(c_X)\|_X, m_{\hat{\tau}}, |\hat{\gamma}|, \|c_X\|_X$, then

$$\mathcal{S}^{\omega, X} \models \forall y \leq_\sigma s(x) (\forall u \leq_{\mathbb{N}} \Phi(x, z^*, n) B_V(x, y, z, u) \rightarrow \exists v \leq_{\mathbb{N}} \Phi(x, z^*, n) C_\exists(x, y, z, v))$$

holds whenever $\mathcal{S}^{\omega, X} \models \Delta$ for $\mathcal{S}^{\omega, X}$ defined via any normed space $(X, \|\cdot\|)$ with χ_A interpreted by the characteristic function of an accretive A such that $\bigcap_{\gamma > 0} \text{dom} J_\gamma^A \neq \emptyset$, J^{X^A} by the corresponding resolvents J_γ^A for $\gamma > 0$ and the other constants accordingly such that the corresponding axioms hold.

Further: if $\hat{\tau}$ is of degree 1, then Φ is a total computable functional and if the claim is proved without DC, then τ may be arbitrary and Φ will be a total functional on $S_\delta \times S_{\hat{\tau}} \times \mathbb{N}$ which is primitive recursive in the sense of Gödel.

The recent work [33] introduced a semi-constructive variant, in the spirit of [15], of another system from [47] dealing with the treatment of maximally monotone operators over Hilbert spaces, and we similarly also want to consider a semi-constructive variant of the above system \mathcal{V}_p^ω in this work. Conceptually similar to the circumstances in [15, 33], this variant $\mathcal{V}_{i,p}^\omega$ is defined as an extension of the basic system $\mathcal{A}_i^\omega = \text{E-HA}^\omega + \text{AC}$ (defined as in [15]) in the same manner as indicated above. We refer to the discussion given in [15] on the resulting differences of the properties of $\mathcal{V}_{i,p}^\omega$ compared to \mathcal{V}_p^ω and in particular on the additional strength of the non-constructive principles allowed in the context of $\mathcal{V}_{i,p}^\omega$ while aiming for bound extractions.

Similar to [33], we can show the following result by adapting [15, Theorem 4.11] (where now \succeq denotes (not necessarily strong) majorization interpreted in the model $\mathcal{S}^{\omega, X}$):

Theorem 2.7. *Let δ be of the form $\mathbb{N} \rightarrow (\dots \rightarrow \mathbb{N})$ and σ, τ be arbitrary, s be a closed term of suitable type. Let Γ_- be a set of sentences of the form $\forall \underline{u} \leq_{\underline{\zeta}} (C(\underline{u}) \rightarrow \exists \underline{v} \leq_{\underline{\beta}} \underline{t}\underline{u} \rightarrow D(\underline{u}, \underline{v}))$ with $\underline{\zeta}, \underline{\beta}$ and C, D arbitrary types and formulas respectively and \underline{t} be closed terms. Let $B(x, y, z)/A(x, y, z, u)$ be arbitrary formulas of $\mathcal{V}_{i,p}^\omega$ with only $x, y, z/x, y, z, u$ free. If*

$$\mathcal{V}_{i,p}^\omega + \Gamma_- \vdash \forall x^\delta \forall y \leq_\sigma (x) \forall z^\tau (\neg B(x, y, z) \rightarrow \exists u^{\mathbb{N}} A(x, y, z, u)),$$

one can extract a $\Phi : S_\delta \times S_{\hat{\tau}} \times \mathbb{N} \rightarrow \mathbb{N}$ with is primitive recursive in the sense of Gödel's T such that for any $x \in S_\delta$, any $y \in S_\sigma$ with $y \leq_\sigma s(x)$, any $z \in S_{\hat{\tau}}$ and $z^* \in S_{\hat{\tau}}$ with $z^* \succeq z$ and any $n \in \mathbb{N}$ with $n \geq_{\mathbb{R}} \|c_X - J_{\hat{\tau}}^A(c_X)\|_X, m_{\hat{\tau}}, |\hat{\gamma}|, \|c_X\|_X$, we have that

$$\mathcal{S}^{\omega, X} \models \exists u \leq_{\mathbb{N}} \Phi(x, z^*, n) (\neg B(x, y, z) \rightarrow A(x, y, z, u))$$

holds whenever $\mathcal{S}^{\omega, X} \models \Gamma_-$ for $\mathcal{S}^{\omega, X}$ defined via any normed spaces $(X, \|\cdot\|)$ with χ_A interpreted by the characteristic function of an accretive operator A such that $\bigcap_{\gamma > 0} \text{dom} J_\gamma^A \neq \emptyset$, J^{X^A} by the corresponding resolvents J_γ^A for $\gamma > 0$ and the other constants accordingly such that the corresponding axioms hold.

We emphasize that while the classical logical metatheorems are derived using a monotone variant of Gödel's Dialectica interpretation due to Kohlenbach (see [23]), the above semi-constructive version rests on the use of a monotone variant due to Kohlenbach [24] of Kreisel's modified realizability.

3. THE NORMALIZED DUALITY MAP AND THE ALTERNATIVE NOTION OF ACCRETIVITY

3.1. The duality map and selection functionals. Recall that for a Banach space X with its dual space X^* , its normalized duality mapping

$$J : X \rightarrow 2^{X^*}, x \mapsto \left\{ y \in X^* \mid \langle x, y \rangle = \|x\|^2 = \|y\|^2 \right\}$$

is non-empty for any $x \in X$ (which follows from the Hahn-Banach theorem). Many works in the context of the theory of accretive operators in general, and the treatment of semigroups generated by those operators in particular, rely on the use of this mapping and in that way, this section is concerned with a proof-theoretic treatment thereof in the context of the formal systems as discussed previously.

As we want to refrain from providing a treatment for both the operator norm on the dual space as well as for the full duality map as a set-valued mapping, we follow the approach initiated by Kohlenbach and Leuştean in [30] where the authors handle uses of J by only treating certain selection functionals for J (depending on the situation).

Concretely, a duality selection map is just a map $j : X \rightarrow X^*$ such that $j(x) \in J(x)$ for $x \in X$. Given a constant j of type $X \rightarrow (X \rightarrow \mathbb{N}^{\mathbb{N}})$, this general property of being a selection map can be expressed by corresponding axioms expressing that

- (1) $jx : X \rightarrow \mathbb{R}$ is a linear operator for any $x \in X$;
- (2) $\|jx\| \leq \|x\|$ where $\|jx\|$ means the operator norm;
- (3) $jxx = \|x\|^2$ (which, as discussed in [30] already, yields $\|jx\| = \|x\|$).

Formally, this can be encapsulated by the following universal axiom introduced in [30]:

$$\forall x^X, y^X \left(jxx =_{\mathbb{R}} \|x\|_X^2 \wedge |jxy| \leq_{\mathbb{R}} \|x\|_X \|y\|_X \right. \\ \left. \wedge \forall \alpha^{\mathbb{N}^{\mathbb{N}}}, \beta^{\mathbb{N}^{\mathbb{N}}}, u^X, v^X (jx(\alpha u +_X \beta v) =_{\mathbb{R}} \alpha jxu +_{\mathbb{R}} \beta jxv) \right),$$

Notice that the operator norm is here avoided by expressing $\|jx\| \leq \|x\|$ via stipulating $|jxy|_{\mathbb{R}} \leq_{\mathbb{R}} \|x\|_X \|y\|_X$.

Remark 3.1. As discussed in [30], the functional j is not provably extensional from the above axiom alone. As indicated by the use of the Dialectica interpretation, if extensionality is to be treated then one has to stipulate an associated modulus of uniform continuity which has been considered in [30]. The applications discussed later actually do not require an extensional or continuous selection map and we therefore do not explicitly discuss this issue any further.

3.2. The alternative notion of accretivity. Besides the purely metric notion of accretivity discussed in the preceding sections, which also forms the basis of the systems \mathcal{V}_p^ω and its intuitionistic variant $\mathcal{V}_{i,p}^\omega$, the more common notion of accretivity, especially in the context of nonlinear semigroups generated by such operators, is the notion introduced by Kato in [20] where one stipulates that A is accretive if

$$\forall (x, u), (y, v) \in A \exists j \in J(x - y) (\langle u - v, j \rangle \geq 0).$$

In the language of the preceding subsection, this can be recognized as stipulating the existence of a family of selection functionals $j_{u,v}$ such that, as before, $j_{u,v}x \in J(x)$ and where now further $\langle u - v, j_{u,v}(x - y) \rangle \geq 0$ for any $u \in Ax$ and $v \in Ay$.

Formally, this leads us to the following modification of the previous systems: we define $\widehat{\mathcal{V}}_p^\omega$ as the extension of $\mathcal{A}^\omega[X, \|\cdot\|]$ with the axiom schemes (I), (II), (IV) and (V) as before, now over the language extended with a constant j of type $X \rightarrow (X \rightarrow (X \rightarrow \mathbb{N}^{\mathbb{N}}))$ (or, in a more suggestive notation, $(X \times X \times X \times X) \rightarrow \mathbb{N}^{\mathbb{N}}$) together with the axioms

$$(J) \quad \forall x^X, y^X, u^X, v^X \left(\langle x, j_{u,v}x \rangle =_{\mathbb{R}} \|x\|_X^2 \wedge \langle y, j_{u,v}x \rangle \leq_{\mathbb{R}} \|x\|_X \|y\|_X \right. \\ \left. \wedge \forall \alpha^{\mathbb{N}^{\mathbb{N}}}, \beta^{\mathbb{N}^{\mathbb{N}}}, z^X, w^X (\langle \alpha z +_X \beta w, j_{u,v}x \rangle =_{\mathbb{R}} \alpha \langle z, j_{u,v}x \rangle + \beta \langle w, j_{u,v}x \rangle) \right)$$

as well as

$$(A) \quad \forall x^X, y^X, u^X, v^X (u \in Ax \wedge v \in Ay \rightarrow \langle u -_X v, j_{u,v}(x -_X y) \rangle \geq 0)$$

where we write $j_{u,v}$ for juv as well as $\langle y, j_{u,v}x \rangle$ for $juvxy$.

It is rather immediately clear through the considerations made in [30] that the bound extraction theorems contained in Theorem 2.6 and 2.7 extend to the system $\widehat{\mathcal{V}}_p^\omega$: At first, we have to give a suitable interpretation to the constant j in the model $\mathcal{M}^{\omega, X}$ associated with an accretive operator A (see [47]). For that, note that the function j is defined by contracting the two parameters besides u, v , namely x and y , into the one argument of j (which is feasible as the witnessing functionals required by the notion of accretivity only have to satisfy $j \in J(x - y)$). The interpretation of this constant in the model now has to ‘unwind’ this contraction (which essentially relies on a choice principle). Concretely, we are lead to the following interpretation of j (writing \mathcal{M} concisely for $\mathcal{M}^{\omega, X}$): given an accretive operator $A \subseteq X \times X$, define $[j]_{\mathcal{M}}$ by

$$[j]_{\mathcal{M}}(u, v, z, w) = \begin{cases} (\langle w, j_{u,v}^A(z) \rangle)_{\circ} & \text{if } \exists x, y \in X (u \in Ax \wedge v \in Ay \wedge z =_X x -_X y), \\ (\langle w, \tilde{j}(z) \rangle)_{\circ} & \text{otherwise,} \end{cases}$$

where $\langle \cdot, \cdot \rangle$ is application in the space X^* , the functionals $j_{u,v}^A(z) \in J(z)$ are those guaranteed to exist by the definition of accretivity, $\tilde{j}(z)$ is a generic element of $J(z)$ (which always exists as $J(z) \neq \emptyset$ by the Hahn-Banach theorem) and \circ is defined as in [25].

The Theorems 2.6 and 2.7 now extend to this setting as all the additional axioms (J) and (A) are purely universal and since the additional constant j with its interpretation in the model $\mathcal{M}^{\omega, X}$ can be majorized by following the ideas presented in the proof of Theorem 2.2 in [30]: from $\langle y, j_{u,v}x \rangle \leq \|x\| \|y\|$, one obtains that $nm \geq \langle y, j_{u,v}x \rangle$ for $n \geq \|x\|$ and $m \geq \|y\|$ which immediately yields that the function

$$(n, m, l, k) \mapsto (mn)_{\circ}$$

defined for $n, m, k, l \in \mathbb{N}$ with $\|u\| \leq k$, $\|v\| \leq l$, $\|z\| \leq m$, $\|w\| \leq n$ is a majorant for j . Here \circ is the previous operation, now restricted to \mathbb{N} (which, as discussed in e.g. [26], can be explicitly calculated). This majorant is in particular actually independent on the arguments induced by the upper bounds on $\|u\|$ and $\|v\|$, i.e. k and l .

The question of how this notion of accretivity relates to the previously used notion immediately arises. By formalizing one direction of the proof on the equivalence of the two notions of accretivity (essentially due to Kato [20], see also Lemma 3.1 in Chapter II of [1]), we obtain the following:

Proposition 3.2. *The system $\widehat{\mathcal{V}}_p^\omega$ proves:*

(1)

$$\forall x^X, y^X, u^X, v^X \left(\langle y, j_{u,v}x \rangle \geq_{\mathbb{R}} 0 \rightarrow \forall \lambda^{\mathbb{N}^{\mathbb{N}}} (\|x\|_X \leq_{\mathbb{R}} \|x +_X |\lambda|y\|_X) \right).$$

(2)

$$\begin{aligned} \forall x^X, y^X, u^X, v^X, \lambda^{\mathbb{N}^{\mathbb{N}}} ((x, u), (y, v)) \in A \\ \rightarrow \|x -_X y +_X |\lambda|(u -_X v)\|_X \geq_{\mathbb{R}} \|x -_X y\|_X. \end{aligned}$$

Proof. (1) The conclusion is vacuously true for $x = 0$. Thus assume $x \neq 0$ and let $\langle y, j_{u,v}x \rangle \geq 0$. Then we get

$$\begin{aligned} \|x\|^2 &= \langle x, j_{u,v}x \rangle && \text{by (J)} \\ &= \langle x + |\lambda|y - |\lambda|y, j_{u,v}x \rangle && \text{by (qf - ER)} \\ &= \langle x + |\lambda|, j_{u,v}x \rangle - |\lambda|\langle y, j_{u,v}x \rangle && \text{by (J)} \\ &\leq \langle x + |\lambda|y, j_{u,v}x \rangle \leq \|x + |\lambda|y\| \|x\| && \text{by (J)}. \end{aligned}$$

Then $\|x\| \leq \|x + |\lambda|y\|$ after dividing by $\|x\|$.

(2) By using (A), we have $\langle u - v, j_{u,v}(x - y) \rangle \geq 0$ for $u \in Ax$ and $v \in Ay$. Then, we get $\|x - y\| \leq \|x - y + |\lambda|(u - v)\|$ by (1). □

Therefore, the system $\widehat{\mathcal{V}}_p^\omega$ is an extension of \mathcal{V}_p^ω as all the axioms of \mathcal{V}_p^ω are provable in $\widehat{\mathcal{V}}_p^\omega$. In particular, all properties of A and its resolvent exhibited in Lemma 2.3 are provable in $\widehat{\mathcal{V}}_p^\omega$. Further, the system proves most of the basic facts about such duality selection mappings. One such fact that will be particularly useful later on is the following (proved (in passing) e.g. in the proof of Proposition 1.1 in Chapter I of [1]):

Proposition 3.3. *The system $\widehat{\mathcal{V}}_p^\omega$ proves:*

$$\forall x^X, y^X, u^X, v^X, t^{\mathbb{N}^{\mathbb{N}}} \left(t >_{\mathbb{R}} 0 \rightarrow \langle y, j_{u,v}x \rangle \leq_{\mathbb{R}} \|x\|_X \frac{\|x +_X ty\|_X - \|x\|_X}{t} \right).$$

Proof. We have

$$\|x\|^2 + t\langle y, j_{u,v}x \rangle = \langle x + ty, j_{u,v}x \rangle \leq \|x\| \|x + ty\|$$

by axiom (J). This implies

$$\langle y, j_{u,v}x \rangle \leq \|x\| \frac{\|x + ty\| - \|x\|}{t}.$$

□

3.3. The mapping $\langle \cdot, \cdot \rangle_s$. Of crucial importance in the context of many proofs from the theory of nonlinear semigroups, and in particular in the context of the exemplary applications considered later in this paper, is the use of a function $\langle \cdot, \cdot \rangle_s : X \times X \rightarrow \overline{\mathbb{R}}$ defined by

$$\langle y, x \rangle_s := \sup \{ \langle y, j \rangle \mid j \in J(x) \}.$$

As already observed in the early papers [5, 11], it is easy to see that $\langle y, x \rangle_s < +\infty$ for all $x, y \in X$ and in fact, since $J(x)$ is weak-star compact in X^* , the supremum is actually attained.

While $\langle \cdot, \cdot \rangle_s$ is by virtue of its definition via the supremum and the duality map J a complex object in the formal contexts considered in this paper, many proofs only rely on the existence of a mapping which shares some essential properties with $\langle \cdot, \cdot \rangle_s$ and in that case, such a mapping can indeed be treated in the context of the systems discussed above and this is what we want to briefly discuss in the following.

Concretely, under the ‘essential properties’ mentioned above we will understand the following:

(1) $\langle \alpha y, \beta x \rangle_s = \alpha\beta \langle y, x \rangle_s$ for $x, y \in X$ and $\alpha, \beta \geq 0$;

- (2) $\langle \alpha x + y, x \rangle_s = \alpha \|x\|^2 + \langle y, x \rangle_s$ for $x, y \in X$ and $\alpha \in \mathbb{R}$;
- (3) $|\langle y, x \rangle_s| \leq \|y\| \|x\|$ for $x, y \in X$;
- (4) $\langle y, j_{u,v} x \rangle \leq \langle y, x \rangle_s$ for $x, y \in X$ and $u, v \in X$ where the $j_{u,v}$ are the selection functionals for J guaranteed by accretivity;
- (5) $\langle \cdot, \cdot \rangle_s$ is upper-semicontinuous (in its right argument).

For a proof for the items (1), (2) and (5), see Proposition 1.2 in Chapter I of [1]. The other items are immediate.

If all that is required of $\langle \cdot, \cdot \rangle_s$ in a proof is that it fulfills these properties, then this proof can, under suitable uniformization of these assumptions, be treated in the context of the above systems by adding a further constant $\langle \cdot, \cdot \rangle_s$ of type $X \rightarrow (X \rightarrow \mathbb{N}^{\mathbb{N}})$ together with the following axioms: the items (1) - (4) are readily formulated as

- (+)₁ $\forall x^X, y^X, \alpha^{\mathbb{N}^{\mathbb{N}}}, \beta^{\mathbb{N}^{\mathbb{N}}} (\langle \alpha|y, |\beta|x \rangle_s =_{\mathbb{R}} |\alpha||\beta|\langle y, x \rangle_s)$,
- (+)₂ $\forall x^X, y^X, \alpha^{\mathbb{N}^{\mathbb{N}}} (\langle \alpha x +_X y, x \rangle_s =_{\mathbb{R}} \alpha \|x\|_X^2 + \langle y, x \rangle_s)$,
- (+)₃ $\forall x^X, y^X (|\langle y, x \rangle_s| \leq_{\mathbb{R}} \|y\|_X \|x\|_X)$,
- (+)₄ $\forall x^X, y^X, u^X, v^X (\langle y, j_{u,v} x \rangle \leq_{\mathbb{R}} \langle y, x \rangle_s)$,

in the underlying language. For a suitable formulation of item (5), note that the logical methodology based on the monotone Dialectica interpretation suggest that the assumption is upgraded to the existence of a modulus ω of uniform upper-semicontinuity. Concretely, we will consider an additional constant ω^+ of type $\mathbb{N} \rightarrow (\mathbb{N} \rightarrow \mathbb{N})$ together with the axiom

(+)₅

$$\forall x^X, y^X, z^X, b^{\mathbb{N}}, k^{\mathbb{N}} \left(\|x\|_X, \|z\|_X <_{\mathbb{R}} b \wedge \|x -_X y\|_X <_{\mathbb{R}} 2^{-\omega^+(b,k)} \right. \\ \left. \rightarrow \langle z, y \rangle_s \leq_{\mathbb{R}} \langle z, x \rangle_s + 2^{-k} \right).$$

Note that by the uniformity on x where the rate only depends on the upper bound b , this is actually a full modulus of uniform continuity.

The assumption that $\langle \cdot, \cdot \rangle_s$ is uniformly continuous is in particular true if the space is uniformly smooth and will be in particular also be necessary if the proof to be treated in some form uses extensionality of the functional $\langle \cdot, \cdot \rangle_s$ (in its right argument) as suggested by the logical methodology. However, if that is not the case and the proof can be formalized just using the axioms (+)₁, ..., (+)₄, then the bound extraction theorem established later in particular guarantees a bound which is valid in all Banach spaces.

Note also that accretivity is sometimes defined by explicitly using the functional $\langle \cdot, \cdot \rangle_s$ through stating that

$$\forall (x, u), (y, v) \in A (\langle u - v, x - y \rangle_s \geq 0).$$

This version of accretivity is immediately provable in the system $\widehat{\mathcal{V}}_p^\omega + (+)_4$ as, using axioms (A) and (+)₄, we have

$$\langle u - v, x - y \rangle_s \geq \langle u - v, j_{u,v}(x - y) \rangle \geq 0.$$

We later denote the collection of these five axioms (+)₁ - (+)₅ by (+). Now, the bound extraction results contained in Theorem 2.6 and 2.7 also extend to the associated extended system(s) $\widehat{\mathcal{V}}_p^\omega + (+)_1 + \dots + (+)_4 + ((+)_5)$ with the conclusion drawn over any space (or where $\langle \cdot, \cdot \rangle_s$ is additionally uniformly continuous on bounded subsets as above if (+)₅ is included). Concretely, this follows as before since, for one, all the axiom schemes are purely universal and, for another, the constant $\langle \cdot, \cdot \rangle_s$ can be immediately majorized: from $|\langle y, x \rangle_s| \leq \|y\| \|x\|$, we as before infer $mn \geq |\langle y, x \rangle_s|$ for $m \geq \|y\|$ and $n \geq \|x\|$. From this, a majorant for the accompanying interpretation in the model $\mathcal{M}^{\omega, X}$ follows. Further, the additional constant ω^+ is immediately majorized (essentially by itself) as it is of type $\mathbb{N} \rightarrow (\mathbb{N} \rightarrow \mathbb{N})$ and so, similar to Lemma 17.82 of [26], we have that $\omega^{+, M}$ defined by

$$\omega^{+, M}(b, k) = \max\{\omega^+(a, j) \mid a \leq b, j \leq k\}$$

is a majorant for ω^+ .

4. NONLINEAR SEMIGROUPS GENERATED BY THE CRANDALL-LIGGETT FORMULA

In this section, we now are concerned with a formal treatment of semigroup \mathcal{S} stipulated by the generation result of Crandall and Liggett [11] previously discussed in Theorem 2.2. Before diving into the formal treatment of these semigroups, we however need to consider some preliminary formal results for the treatment of the range condition in the premise as well as how J_0^A is to be understood.

4.1. **The treatment of $\overline{\text{dom}A}$.** Crucial both in the definition of the semigroup and the central assumption of the range condition is the use of the closure of the domain of A and in the following formal investigations, quantification over elements from $\overline{\text{dom}A}$ will be necessary. All the previous systems essentially only considered normed spaces and in that context, we now first have to lift the previous treatment to take the completeness of the underlying Banach space into account. For that, we are following the approach laid out in [26] (essentially already implicitly present in [25]) whereas complete spaces are treated by adding another operator C of type $(\mathbb{N} \rightarrow X) \rightarrow X$ which is meant to assign to a Cauchy sequence $x^{\mathbb{N} \rightarrow X}$ a limit $C(x)$. To discard of the complex premise of Cauchy-ness in an axiom stating that property, one then restricts oneself to Cauchy sequences with a fixed Cauchy rate (similar to the representation of real numbers in finite type arithmetic, see [26]). To implicitly quantify only over all such sequences, a term construction \hat{x} is used on the objects $x^{\mathbb{N} \rightarrow X}$. Precisely, \hat{x} is defined on the level of the representation of the real value of the norm via sequences of rational numbers with fixed Cauchy rate via⁵

$$\hat{x}_n =_X \begin{cases} x_n & \text{if } \forall k <_{\mathbb{N}} n \left([\|x_k -_X x_{k+1}\|_X](k+1) <_{\mathbb{Q}} 6 \cdot 2^{-k-1} \right), \\ x_k & \text{for } \min k <_{\mathbb{N}} n : [\|x_k -_X x_{k+1}\|_X](k+1) \geq_{\mathbb{Q}} 6 \cdot 2^{-k-1}, \text{ otherwise.} \end{cases}$$

Then completeness of the space can be formulated via the universal axiom⁶

$$(C) \quad \forall x^{\mathbb{N} \rightarrow X}, k^{\mathbb{N}} \left(\|C(x) -_X \hat{x}_k\|_X \leq_{\mathbb{R}} 2^{-k+3} \right)$$

which indeed implies completeness of the space in the form that from

$$\forall k^{\mathbb{N}} \exists n^{\mathbb{N}} \forall m, \tilde{m} \geq_{\mathbb{N}} n \left(\|x_m -_X x_{\tilde{m}}\|_X <_{\mathbb{R}} 2^{-k} \right)$$

it follows provably in $\mathcal{A}^\omega[X, \|\cdot\|] + (C)$ that

$$\forall k^{\mathbb{N}} \exists m^{\mathbb{N}} \forall l \geq_{\mathbb{N}} m \left(\|C(x) -_X x_l\|_X <_{\mathbb{R}} 2^{-k+1} \right).$$

As further shown in [26], the constant C is majorizable and therefore we find that the bound extraction theorems discussed above immediately extend to $\hat{\mathcal{V}}_p^\omega + (C)$ or any suitable extension (e.g. by $(+)$).

Now a statement where one is quantifying over the closure of the domain, i.e. a statement of the form

$$(*) \quad \forall x \in \overline{\text{dom}A} B(x)$$

can, through the use of C , be (naively) expressed as

$$\forall x^{\mathbb{N} \rightarrow X} \left(\left(\forall n^{\mathbb{N}} \exists y^X (y \in Ax_n) \right. \right. \\ \left. \left. \wedge \forall k^{\mathbb{N}} \exists n^{\mathbb{N}} \forall m, \tilde{m} \geq_{\mathbb{N}} n \left(\|x_m -_X x_{\tilde{m}}\|_X <_{\mathbb{R}} 2^{-k} \right) \right) \rightarrow B(C(x)) \right).$$

The premise that $x^{\mathbb{N} \rightarrow X}$ is a Cauchy sequence can be removed as before by considering \hat{x} instead of x but the inclusion of the sequence in the domain, then in the form of $\forall n^{\mathbb{N}} \exists y^X (y \in A\hat{x}_n)$, remains.

The approach is now to also remove this assumption in a similar style as the $\hat{\cdot}$ -operation by universally quantifying over the potential witnessing sequence y_n and defining a subsequent operation similar to $\hat{\cdot}$ which potentially alters the sequence such that $x_n \in \text{dom}A$ will always be guaranteed for any n . Concretely, for two objects x, y of type $\mathbb{N} \rightarrow X$, we define

$$(x \upharpoonright y)_n =_X \begin{cases} x_n & \text{if } \forall k \leq_{\mathbb{N}} n (y_k \in Ax_k), \\ x_{k \dot{-} 1} & \text{for } \min k \leq_{\mathbb{N}} n : y_k \notin Ax_k, \text{ otherwise.} \end{cases}$$

Note that since inclusions in the graph of A are quantifier-free, the above indeed can be defined by a closed term in the underlying language.

Now, using the operation \upharpoonright in tandem with $\hat{\cdot}$, we can implicitly quantify over elements from $\overline{\text{dom}A}$ by quantifying over elements of type $\mathbb{N} \rightarrow X$ and thus we can express the statement $(*)$ equivalently by

$$\forall x^{\mathbb{N} \rightarrow X}, y^{\mathbb{N} \rightarrow X} (y_0 \in Ax_0 \rightarrow B(C(x \upharpoonright y))).$$

Note that the premise $y_0 \in Ax_0$ actually guarantees that $(x \upharpoonright y)_n \in \text{dom}A$ for all n . Note also that

$$\widehat{x \upharpoonright y} =_{\mathbb{N} \rightarrow X} \hat{x} \upharpoonright y.$$

⁵We here follow the notion of [26] and denote by $[a](k)$ the k -th element of the Cauchy sequence representation of the real number a .

⁶See the discussion in [26] for the necessity of the additional $+3$ in the formulation.

To see this, one can consider a case distinction on whether $\hat{x} = x$ holds or not and simultaneously on whether $x \uparrow y = x$ holds or not. We only consider the one case out of the four where $\hat{x} \neq x$ and $x \uparrow y \neq x$. By definition, we then have a least k such that $\lceil \|x_k - x_{k+1}\| \rceil (k+1) \geq_{\mathbb{Q}} 6 \cdot 2^{-k-1}$ as well as a least j such that $y_j \notin Ax_j$. Then, it immediately follows by definition of the operations as well as the minimality of k and j that

$$\begin{aligned} \hat{x} \uparrow y &= (x_0, \dots, x_k, x_k, \dots) \uparrow y \\ &= (x_0, \dots, x_{\min\{k, j-1\}}, x_{\min\{k, j-1\}}, \dots) \\ &= (x_0, \dots, x_{j-1}, x_{j-1}, \dots)^{\hat{}} \\ &= \widehat{x \uparrow y} \end{aligned}$$

where, in the third line, we wrote $(x_0, \dots, x_{j-1}, x_{j-1}, \dots)^{\hat{}}$ for the operation $\hat{}$ applied to the sequence $(x_0, \dots, x_{j-1}, x_{j-1}, \dots)$

4.2. Range conditions. A treatment for some variants of the range conditions was already briefly discussed in [47] where the particular case of

$$\text{dom}A \subseteq \bigcap_{\lambda > 0} \text{ran}(Id + \lambda A)$$

was studied. Similar to the discussion there, one can provide a formalized version of this range condition by making use of the resolvent in the form of the following sentence:

$$\forall x^X, \lambda^{\mathbb{N}^{\mathbb{N}}} (x \in \text{dom}A \wedge \lambda >_{\mathbb{R}} 0 \rightarrow \lambda^{-1}(x -_X J_{\lambda}^A x) \in A(J_{\lambda}^A x)).$$

This correctly expresses the range condition since stating that $x \in \text{ran}(Id + \lambda A)$ is equivalent to stating that $x \in \text{dom}(J_{\lambda}^A)$ just via the definition of the resolvent. This latter statement is now equivalently formally encapsulated in our systems by stating the inclusion $\lambda^{-1}(x -_X J_{\lambda}^A x) \in A(J_{\lambda}^A x)$. Note also that this is axiom in particular purely universal and thus can be used in the bound extraction theorems.

In the following, we want to consider two modifications: (1) we want to specify that the inclusion is valid even for the closure of the domain; (2) we want to restrict the intersection to $\lambda < \lambda_0$ for some real parameter $\lambda_0 > 0$. The use of such a λ_0 can be facilitated by adding two further constants and an axiom: λ_0 of type $\mathbb{N}^{\mathbb{N}}$ and m_{λ_0} of type \mathbb{N} together with the accompanying axiom $\lambda_0 \geq 2^{-m_{\lambda_0}}$ providing a verifier to $\lambda_0 > 0$. Note that the bound extraction results stay valid in the context of such an extension if one additionally requires the parameter n to satisfy $n \geq |\lambda_0|, m_{\lambda_0}$.

In the context of such additional constants, the above range condition can be immediately modified to represent the restricted range condition

$$\text{dom}A \subseteq \bigcap_{\lambda_0 > \lambda > 0} \text{ran}(Id + \gamma A)$$

by considering

$$\forall x^X, \lambda^{\mathbb{N}^{\mathbb{N}}} (x \in \text{dom}A \wedge \lambda_0 >_{\mathbb{R}} \lambda >_{\mathbb{R}} 0 \rightarrow \lambda^{-1}(x -_X J_{\lambda}^A x) \in A(J_{\lambda}^A x)).$$

Further, in both cases we can now consider the other main modification of stipulating the range condition also for the closure of the domain, i.e.

$$\overline{\text{dom}A} \subseteq \bigcap_{\lambda_0 > \lambda > 0} \text{ran}(Id + \gamma A),$$

by using the above treatment of quantification over elements in the closure of the domain by quantification over sequences in X together with the operators C and $(\cdot \uparrow \cdot)$. Concretely, one rather immediately obtains the following natural extension to the closure of the domain:

$$\begin{aligned} (RC)_{\lambda_0} \quad \forall x^{\mathbb{N} \rightarrow X}, v^{\mathbb{N} \rightarrow X}, \lambda^{\mathbb{N}^{\mathbb{N}}} & \left(v_0 \in Ax_0 \wedge \lambda_0 >_{\mathbb{R}} \lambda >_{\mathbb{R}} 0 \right. \\ & \left. \rightarrow \lambda^{-1}(C(x \uparrow v) -_X J_{\lambda}^A(C(x \uparrow v))) \in A(J_{\lambda}^A(C(x \uparrow v))) \right). \end{aligned}$$

Similarly, we could here lift the restriction via λ_0 again and get a full range condition for the closure of the domain. We denoted this full range condition for the closure of the domain by (RC) , but at the same time refrain from spelling this out in any more detail here. Note however that all the other range conditions introduced here are still of purely universal and thus are admissible in the context of the bound extraction theorems.

Remark 4.1. The metatheorems exhibited in Theorems 2.6 and 2.7 require as an assumption that $\bigcap_{\lambda>0} \text{dom} J_\lambda^A \neq \emptyset$, a requirement which would be substantiated via a full range condition together with a witness for $\text{dom} A \neq \emptyset$ (which was previously (in some sense but not precisely) represented by c_X). In the context of the above restricted range conditions, it is however feasible that $\bigcap_{\lambda>0} \text{dom} J_\lambda^A$ is actually empty while only $\bigcap_{\lambda_0>\lambda>0} \text{dom} J_\lambda^A \neq \emptyset$ holds. It should be noted that in this case, Theorems 2.6 and 2.7 can be modified to stay valid if c_X is interpreted by a point in this restricted intersection. Therefore, if we in the following write $\widehat{\mathcal{V}}_p^\omega + (\mathcal{C}) + (RC)_{\lambda_0}$ or consider any extension, we consider the axioms (IV) and (V) to be replaced by

$$\begin{aligned} \text{(IV)'} \quad & \lambda_0 - 2^{m'_\gamma} \geq_{\mathbb{R}} \tilde{\gamma} \geq_{\mathbb{R}} 2^{-m'_\gamma}, \\ \text{(V)'} \quad & d_X \in Ac_X, \end{aligned}$$

where d_X is a new constant of type X and m'_γ is a new constant of type \mathbb{N} , the latter witnessing that $\lambda_0 > \tilde{\gamma}$. The majorization of all resolvents J_γ^A for $\gamma \in [0, \lambda_0)$ is then achieved as before via

$$\begin{aligned} \|J_\gamma^A x\| &\leq \|x\| + 2\|c_X\| + \left(2 + \frac{\gamma}{\tilde{\gamma}}\right) \|c_X - J_{\tilde{\gamma}}^A c_X\| \\ &\leq \|x\| + 2\|c_X\| + (2\tilde{\gamma} + \gamma) \|d_X\|. \end{aligned}$$

In that case however, the interpretation of the resolvent constant J^{X^A} in the models $\mathcal{M}^{\omega, X}$ and $\mathcal{S}^{\omega, X}$ has to be modified to set $[J^{X^A}]_{\mathcal{M}}(x, \gamma) = 0$ for all x if $\gamma >_{\mathbb{R}} \lambda_0$ (and similar for $\mathcal{S}^{\omega, X}$). Therefore, the extracted bounds only remain meaningful if the theorem does not utilize these resolvents. If it does, further modifications are necessary but we refrain from discussing this here any further as this situation does not arise in this paper.

4.3. The resolvent at zero. Something left open by the characterizing axioms discussed in the preliminaries is the behavior of J_0^A . This, however, takes a special role in the context of the treatment of nonlinear semigroups S generated by the associated operator A due to the prominent use often made of $S(0)$.

The reason for this previous ambiguity in the treatment of the resolvent at 0 was the fact that the resolvent does not always behave continuously at 0 if it is naively defined: while the definition of the resolvent via

$$J_\gamma^A = (\text{Id} + \gamma A)^{-1}$$

suggests $J_0^A x = x$, it is well known (see [3]) that already in Hilbert spaces with a maximally monotone operator A , one has $J_t^A x \rightarrow P_{\overline{\text{dom} A}} x$ for $t \rightarrow 0$ and all $x \in \text{dom}(J_t^A)$. Therefore, extensionality for the constant J^{X^A} in its first argument t at 0 can in general not be expected if J_0^A is defined in this way and the previous axiomatization left the definition of J_0^A open.

In the following, we nevertheless consider the set of axioms discussed previously forming $\widehat{\mathcal{V}}_p^\omega$ to actually be extended with the sixth axiom

$$\text{(VI)} \quad \forall x^X \ (J_0^A x =_X x),$$

stating the defining equality $J_0^A = (\text{Id} + 0A)^{-1} = \text{Id}$.

Now, the above result that $J_t^A x \rightarrow P_{\overline{\text{dom} A}} x$ for $t \rightarrow 0$ extends to Banach spaces at least partially in the sense that one can show (see Proposition 3.2 of Chapter II in [1]) that $J_t^A x \rightarrow x$ for $\lambda_0 > t \rightarrow 0$ and

$$x \in \overline{\text{dom} A} \cap \bigcap_{\lambda_0 > \lambda > 0} \text{dom} J_\lambda^A.$$

Therefore, in the presence of a range condition, we should at least have a continuous and thus extensional behavior of the resolvent defined in this manner at $t = 0$ for all $x \in \overline{\text{dom} A}$ and this can indeed be formally verified in the accompanying system.

Lemma 4.2. $\widehat{\mathcal{V}}_p^\omega + (\mathcal{C}) + (RC)_{\lambda_0}$ proves:

$$\begin{aligned} \forall x^{\mathbb{N} \rightarrow X}, v^{\mathbb{N} \rightarrow X}, \lambda^{\mathbb{N}^{\mathbb{N}}}, k^{\mathbb{N}} \left(v_0 \in Ax_0 \wedge 0 <_{\mathbb{R}} \lambda <_{\mathbb{R}} \min \left\{ \frac{2^{-(k+1)}}{\max\{1, \|v_{k+5}\|_X\}}, \lambda_0 \right\} \right. \\ \left. \rightarrow \|C(x \upharpoonright v) -_X J_\lambda^A C(x \upharpoonright v)\|_X \leq_{\mathbb{R}} 2^{-k} \right). \end{aligned}$$

Proof. First, by Lemma 2.3, we have

$$\forall x^X, v^X, \lambda^{\mathbb{N}^{\mathbb{N}}} \ (0 <_{\mathbb{R}} \lambda <_{\mathbb{R}} \lambda_0 \wedge v \in Ax \rightarrow \|x - J_\lambda^A x\| \leq \lambda \|v\|)$$

as using $(RC)_{\lambda_0}$ and the quantifier-free extensionality rule, we obtain $x \in \text{dom}(J_\lambda^A)$ for all $\lambda \in (0, \lambda_0)$. So, for $x^{\mathbb{N} \rightarrow X}$ and $v^{\mathbb{N} \rightarrow X}$ such that w.l.o.g. $v_n \in A\hat{x}_n$ for all n , we obtain $C(x \upharpoonright v) \in \text{dom}(J_\lambda^A)$ for all $\lambda \in (0, \lambda_0)$ again by $(RC)_{\lambda_0}$. Therefore, using (C) :

$$\begin{aligned} \|C(x \upharpoonright v) - J_\lambda^A C(x \upharpoonright v)\| &\leq \|C(x \upharpoonright v) - \hat{x}_n\| + \|\hat{x}_n - J_\lambda^A \hat{x}_n\| \\ &\quad + \|J_\lambda^A \hat{x}_n - J_\lambda^A C(x \upharpoonright v)\| \\ &\leq 2 \|C(x \upharpoonright v) - \hat{x}_n\| + \|\hat{x}_n - J_\lambda^A \hat{x}_n\| \\ &\leq 2 \cdot 2^{-n+3} + \lambda \|v_n\|. \end{aligned}$$

Choosing $n = k + 5$, we get that for $\lambda \leq 2^{-(k+1)}/\max\{1, \|v_{k+5}\|\}$:

$$\|C(x \upharpoonright v) - J_\lambda^A C(x \upharpoonright v)\| \leq 2^{-k}.$$

□

This property will be sufficient in the following as the semigroup operates only on $\overline{\text{dom}A}$.

4.4. The semigroup. For treating the semigroup on $\overline{\text{dom}A}$ from Theorem 2.2, it is very instructional to first consider the operator S solely on $\text{dom}A$. In that case, we can facilitate a treatment by directly adding a further constant S of type $\mathbb{N} \rightarrow (X \rightarrow X)$ to the underlying language together with an axiom stating that S on $\text{dom}A$ arises from the Crandall-Liggett formula, i.e. that

$$S(t)x = \lim_{n \rightarrow \infty} \left(\text{Id} + \frac{t}{n} A \right)^{-n} x$$

for any $x \in \text{dom}A$. This can be achieved by further adding a constant ω^S of type $\mathbb{N} \rightarrow (\mathbb{N} \rightarrow (\mathbb{N} \rightarrow \mathbb{N}))$ together with the axiom

$$\begin{aligned} \text{(S1)} \quad \forall k^{\mathbb{N}}, b^{\mathbb{N}}, T^{\mathbb{N}}, x^X, v^X, t^{\mathbb{N}^{\mathbb{N}}} &\left(v \in Ax \wedge \|x\|_X, \|v\|_X <_{\mathbb{R}} b \wedge |t| <_{\mathbb{R}} T \right. \\ &\left. \rightarrow \forall n \geq_{\mathbb{N}} \omega^S(k, b, T) \left(|t|/n <_{\mathbb{R}} \lambda_0 \rightarrow \left\| S(|t|)x -_X (J_{|t|/n}^A)^n x \right\|_X \leq_{\mathbb{R}} 2^{-k} \right) \right), \end{aligned}$$

expressing that ω^S represents a rate of convergence uniform for elements x from bounded subsets $B_b(0) \cap \text{dom}A$ and uniform in t for bounded intervals $[0, T]$ (where we use to absolute value to disperse of the universal premise $t \geq 0$). The term $(J_{|t|/n}^A)^n$ used here is a shorthand for a term $I(t)(n)(n)$ where $I(t)(m)$ is a closed term of type $\mathbb{N} \rightarrow (X \rightarrow X)$ defined using the recursors of the underlying language of $\mathcal{A}^\omega[X, \|\cdot\|]$ (see [26]) via $I(t)(m)(0) = \lambda x.x$ and $I(t)(m)(n+1) = \lambda x.(J_{t/m}^A(I(t)(m)(n)(x)))$.⁷ Note also that we in particular treat $S(0)x$ via $J_0^A x$ by using the absolute value $|t|$ in the above formula to implicitly quantify over non-negative real numbers.

Such a use of a rate of convergence is in particular justified by the fact that the proof given in [11] of the Cauchy-property of the sequence $(J_{t/n}^A)^n x$ for given $t > 0$ and $x \in \text{dom}A$ can be immediately recognized to be provable in the system $\widehat{\mathcal{V}}_{i,p}^\omega + (C) + (RC)_{\lambda_0}$. Therefore, the extension of the semi-constructive metatheorem (Theorem 2.7) to this system guarantees the existence of a rate of Cauchyness for $(J_{t/n}^A)^n x$ and consequently the existence of a modulus ω^S as characterized by the above axiom which can moreover be extracted from the proof given in [11]. This is in fact rather immediate and was essentially already observed in [11]: one can (formally) show that given $x \in \text{dom}A$ with witness $v \in Ax$ and $t \geq 0$, we have

$$\left\| (J_{t/n}^A)^n x - (J_{t/m}^A)^m x \right\| \leq 2t \left| \frac{1}{m} - \frac{1}{n} \right|^{1/2} \|v\|.$$

⁷We consider $I(t)(m)$ to be trivially defined at $m = 0$

Thus for $T \geq t$ and $b \geq \|v\|$, we have for a given $\varepsilon > 0$ that for any $m \geq n \geq \left\lceil \frac{4T^2b^2}{\varepsilon^2} \right\rceil$:

$$\begin{aligned} \left\| (J_{t/n}^A)^n x - (J_{t/m}^A)^m x \right\| &\leq 2Tb \left| \frac{1}{m} - \frac{1}{n} \right|^{1/2} \\ &\leq 2Tb \frac{1}{\sqrt{n}} \\ &\leq 2Tb \frac{1}{\sqrt{\left\lceil \frac{4T^2b^2}{\varepsilon^2} \right\rceil}} \\ &\leq \varepsilon. \end{aligned}$$

Thus the mapping

$$\omega^S(k, b, T) = 2^{2k+2} T^2 b^2$$

is a possible choice for the rate of convergence⁸ in the exponential formula as derived from the proof and the upper bound b is here actually even independent of $\|x\|$.

Now, the treatment of the extension of S defined in this way on $\text{dom}A$ to the closure $\overline{\text{dom}A}$ is best motivated by considering how it is usually defined in the literature: $S(t)$ as a mapping $\text{dom}A \rightarrow X$ is nonexpansive and thus continuous. The object $S(t)x$ for $x \in \overline{\text{dom}A}$ is then defined by considering that as $x \in \overline{\text{dom}A}$, there exists a sequence $x_n \rightarrow x$ with $x_n \in \text{dom}A$. By convergence, the sequence x_n is Cauchy and by continuity of $S(t)$, the sequence $S(t)x_n$ is Cauchy as well and thus converges in a Banach space by completeness. Then $S(t)x$ is identified with the limit of that sequence. This crucial use of the completeness of the space prompts us to work in the context of the formal treatment of complete spaces and $\overline{\text{dom}A}$ as discussed before.

In that vein, we now want to provide an axiom classifying the behavior of $S(t)$ for elements of $\overline{\text{dom}A}$ by essentially stating that for any x and any Cauchy sequence $x_n \rightarrow x$ with $x_n \in \text{dom}A$, $S(t)x_n$ converges to $S(t)x$. The quantification over all elements of $\overline{\text{dom}A}$ together with their generating sequences can now be achieved as discussed in Section 4.1 and in that way, the axiom stating the resulting behavior for $S(t)x$ then takes the form of the following universal axiom⁹

$$(S2) \quad \forall x^{\mathbb{N} \rightarrow X}, y^{\mathbb{N} \rightarrow X}, t^{\mathbb{N}^{\mathbb{N}}} (y_0 \in Ax_0 \\ \rightarrow \forall n^{\mathbb{N}} (\|S(|t|)(C(x \uparrow y)) -_X S(|t|)((\hat{x} \uparrow y)_n)\|_X \leq_{\mathbb{R}} 2^{-n+3})).$$

Note again that the behavior of $S(0)$ is implicitly characterized by the above axioms together with the use of $(RC)_{\lambda_0}$. We write (S) for $(S1) + (S2)$ as well as H_p^ω for $\hat{\mathcal{V}}_p^\omega + (\mathcal{C}) + (RC)_{\lambda_0} + (S)$.

Now, the above axioms forming the theory H_p^ω are suitable for formalizing large portions on the theory of nonlinear semigroups as generated by the Crandal-Liggett formula and as a sort of litmus test, we at least provide here sketches of formal proofs in the resulting system of the other main semigroup properties which arise pretty much directly by formalizing the proofs given in [11]. For that, however, some careful consideration for iterations of the semigroup map are required here. Concretely, to make expressions like $S(t)S(s)x$ meaningful, we have to consider how $S(s)x \in \overline{\text{dom}A}$ is reflected in the system. Based on the representation of $\overline{\text{dom}A}$ chosen above (which also features in how the extension of S is formally defined by means of the axiom (S2)) we thus first have to see how $S(|t|)C(x)$ with $x_n \in \text{dom}A$ for all n can be expressed as an element of the form $C(u)$ for $u^{\mathbb{N} \rightarrow X}$ such that $u_n \in \text{dom}A$ for all n . To find such a u , note first that the convergence result encoded by (S1) for elements from $\text{dom}A$ extends by means of (S2) to $\overline{\text{dom}A}$ in the following way: provably in H_p^ω , we have

$$\begin{aligned} \forall x^{\mathbb{N} \rightarrow X}, y^{\mathbb{N} \rightarrow X}, t^{\mathbb{N}^{\mathbb{N}}}, k^{\mathbb{N}} \exists N^{\mathbb{N}} \forall n \geq_{\mathbb{N}} N \left(y_0 \in Ax_0 \wedge |t|/n <_{\mathbb{R}} \lambda_0 \right. \\ \left. \rightarrow \left\| S(|t|)(C(x \uparrow y)) -_X \left(J_{|t|/n}^A \right)^n (C(x \uparrow y)) \right\|_X \leq_{\mathbb{R}} 2^{-k} \right) \end{aligned}$$

were moreover (although we avoid spelling this out here) the choice functional for N can be explicitly given by closed terms build up from ω^S (and others). To see the provability of the above statement, let k, x, y, t be

⁸Note that although the function is exponential in k , this is just due to requiring an error of the form 2^{-k} . Abstracting $\varepsilon = 2^{-k}$, the rate is actually linear in $1/\varepsilon$.

⁹Note again that the additional +3 is included here as the axiom (\mathcal{C}) requires this modification in order to have a model as discussed before and the same rate applies to the semigroup-images here as the semigroup is nonexpansive.

arbitrary with $y_0 \in Ax_0$. Then using nonexpansivity of the semigroup and the resolvent (see item (4) of the following Lemma 4.3¹⁰), we have

$$\begin{aligned}
 & \left\| S(|t|)(C(x \uparrow y)) - \left(J_{|t|/n}^A \right)^n (C(x \uparrow y)) \right\| \\
 & \leq \left\| S(|t|)(C(x \uparrow y)) - S(|t|)((x \uparrow y)_{(k+5)}) \right\| \\
 & \quad + \left\| S(|t|)((x \uparrow y)_{(k+5)}) - \left(J_{|t|/n}^A \right)^n ((x \uparrow y)_{(k+5)}) \right\| \\
 & \quad + \left\| \left(J_{|t|/n}^A \right)^n ((x \uparrow y)_{(k+5)}) - \left(J_{|t|/n}^A \right)^n (C(x \uparrow y)) \right\| \\
 & \leq \left\| C(x \uparrow y) - (x \uparrow y)_{(k+5)} \right\| \\
 & \quad + \left\| S(|t|)((x \uparrow y)_{(k+5)}) - \left(J_{|t|/n}^A \right)^n ((x \uparrow y)_{(k+5)}) \right\| \\
 & \quad + \left\| (x \uparrow y)_{(k+5)} - C(x \uparrow y) \right\| \\
 & \leq 2^{-k-1} + \left\| S(|t|)((x \uparrow y)_{(k+5)}) - \left(J_{|t|/n}^A \right)^n ((x \uparrow y)_{(k+5)}) \right\| \\
 & \leq 2^{-k}
 \end{aligned}$$

for any n large enough such that $|t|/n < \lambda_0$ as well as

$$\left\| S(|t|)((x \uparrow y)_{(k+5)}) - \left(J_{|t|/n}^A \right)^n ((x \uparrow y)_{(k+5)}) \right\| \leq 2^{-(k+1)}$$

which can be achieved via (S1). In that way, writing N also for the choice functionals for the quantifier over N in the above statement, we find that $S(|t|)C(x)$ is provably equal to¹¹

$$C \left(\left(\left(J_{|t|/N_x(k)}^A \right)^{N_x(k)} C(x) \right)_k \right).$$

We write $\overline{S(|t|)C(x)}$ in the following for this expression (where one should note again that the N -functionals can be explicitly computed, albeit being somewhat messy). In particular note that

$$\left(J_{|t|/N_x(k)}^A \right)^{N_x(k)} C(x) \in \text{dom}A$$

with the witnessing terms defined in terms of the Yosida approximates (which follows provably from $(RC)_{\lambda_0}$ if we w.l.o.g. assume that the functionals N , for a given t as a parameter, are large enough such that $|t|/N(k) < \lambda_0$). In that way, $S(|t|)S(|s|)C(x)$ can be meaningfully represented by

$$S(|t|)\overline{S(|s|)C(x)} =_X \overline{S(|t|)S(|s|)C(x)}.$$

Note that the system can nevertheless *not* prove that

$$S(|t|)S(|s|)C(x) =_X \overline{S(|t|)S(|s|)C(x)}$$

and so the latter is, in some sense, the only way to talk about iterations meaningfully.

Lemma 4.3. *The following are provable in H_p^ω :*

- (1) $\forall x^X, y^X, t^{\mathbb{N}}, s^{\mathbb{N}} (y \in Ax \rightarrow \|S(|t|x) -_X S(|s|x)\|_X \leq_{\mathbb{R}} 2\||t| - |s|\| \|y\|_X)$.
- (2) $\left\{ \begin{array}{l} \forall x^X, y^X, t^{\mathbb{N}} (x \in \text{dom}A \wedge y \in \text{dom}A \\ \rightarrow \|S(|t|x) -_X S(|t|y)\|_X \leq_{\mathbb{R}} \|x -_X y\|_X) \end{array} \right.$.
- (3) $\left\{ \begin{array}{l} \forall x^{\mathbb{N} \rightarrow X}, v^{\mathbb{N} \rightarrow X}, t^{\mathbb{N}}, s^{\mathbb{N}} (v_0 \in Ax_0 \wedge \||t| - |s|\| \leq_{\mathbb{R}} 2^{-(k+2)} / \max\{1, \|v_{k+5}\|\}) \\ \rightarrow \|S(|t|)C(x \uparrow v) -_X S(|s|)C(x \uparrow v)\|_X \leq_{\mathbb{R}} 2^{-k} \end{array} \right.$.
- (4) $\left\{ \begin{array}{l} \forall x^{\mathbb{N} \rightarrow X}, v^{\mathbb{N} \rightarrow X}, y^{\mathbb{N} \rightarrow X}, w^{\mathbb{N} \rightarrow X}, t^{\mathbb{N}} (v_0 \in Ax_0 \wedge w_0 \in Ay_0 \\ \rightarrow \|S(|t|)(C(x \uparrow v)) -_X S(|t|)(C(y \uparrow w))\|_X \\ \leq_{\mathbb{R}} \|C(x \uparrow v) -_X C(y \uparrow w)\|_X \end{array} \right.$.

¹⁰The first four items of this lemma in particular do not rely on this construction as it will only become necessary in the fifth item.

¹¹In fact, N of course also depends on t and the sequence witnessing $x_n \in \text{dom}A$ for all n but we neglect these dependences for now.

$$(5) \begin{cases} \forall x^{\mathbb{N} \rightarrow X}, v^{\mathbb{N} \rightarrow X}, t^{\mathbb{N}^{\mathbb{N}}}, s^{\mathbb{N}^{\mathbb{N}}} (v_0 \in Ax_0 \\ \rightarrow S(|t| + |s|)(C(x \uparrow v)) =_X S(|t|)\overline{S(|s|)(C(x \uparrow v))}). \end{cases}$$

Proof. (1) At first, note that provably in H_p^ω , we have

$$\begin{cases} \forall x^X, y^X, \mu^{\mathbb{N}^{\mathbb{N}}}, \lambda^{\mathbb{N}^{\mathbb{N}}}, n^{\mathbb{N}}, m^{\mathbb{N}} \left(\lambda_0 > |\lambda| \geq |\mu| \wedge n \geq m \geq 1 \wedge y \in Ax \right. \\ \rightarrow \left\| (J_{|\mu|}^A)^n x - (J_{|\lambda|}^A)^m x \right\| \leq \left(((n|\mu| - m|\lambda|)^2 + n|\mu|(|\lambda| - |\mu|))^{1/2} \right. \\ \left. \left. + (m|\lambda|(|\lambda| - |\mu|) + (m|\lambda| - n|\mu|)^2)^{1/2} \right) \|y\| \right) \end{cases}$$

which can be shown by formalizing the proof given in [11] (note for this Lemma 2.3¹²). Instantiating this with $m = n$, $\mu = |t|/n$ and $\lambda = |s|/n$ for t, s of type $\mathbb{N}^{\mathbb{N}}$, where w.l.o.g. $|s| \geq |t|$, and where n is large enough that $|t|/n, |s|/n < \lambda_0$, we obtain

$$\begin{aligned} \left\| (J_{|t|/n}^A)^n x - (J_{|s|/n}^A)^n x \right\| &\leq \left((|t| - |s|)^2 + |t|(|s|/n - |t|/n) \right)^{1/2} \\ &\quad + (|s|(|s|/n - |t|/n) + (|s| - |t|)^2)^{1/2} \|y\| \end{aligned}$$

for any $n \geq 1$ and any x, y with $y \in Ax$. Let k be arbitrary. Using axiom (S), we get

$$\begin{aligned} \|S(|t|)x - S(|s|)x\| &\leq \left\| S(|t|)x - (J_{|t|/n}^A)^n x \right\| + \left\| (J_{|t|/n}^A)^n x - (J_{|s|/n}^A)^n x \right\| \\ &\quad + \left\| S(|s|)x - (J_{|s|/n}^A)^n x \right\| \\ &\leq \frac{2}{k+1} + \left\| (J_{|t|/n}^A)^n x - (J_{|s|/n}^A)^n x \right\| \\ &\leq \frac{2}{k+1} + \left((|t| - |s|)^2 + |t|(|s|/n - |t|/n) \right)^{1/2} \\ &\quad + (|s|(|s|/n - |t|/n) + (|s| - |t|)^2)^{1/2} \|y\| \end{aligned}$$

for any n additionally satisfying $n \geq \omega^S(k, b, T)$ with $b > \|x\|, \|v\|$ and $T > |t|, |s|$. This implies

$$\|S(|t|)x - S(|s|)x\| \leq \frac{2}{k+1} + 2\|t| - |s|\|y\|$$

and the claim follows as k was arbitrary

(2) By (essentially) Lemma 2.3, we have provably that

$$\left\| J_{|\lambda|}^A x - J_{|\lambda|}^A y \right\| \leq \|x - y\|$$

for any $\lambda_0 > \lambda$ of type $\mathbb{N}^{\mathbb{N}}$ and any x, y of type X . By induction, we get

$$\left\| (J_{|t|/n}^A)^n x - (J_{|t|/n}^A)^n y \right\| \leq \|x - y\|$$

for any t of type $\mathbb{N}^{\mathbb{N}}$, any x, y of type X and any n large enough such that $|t|/n < \lambda_0$. Now, let k be arbitrary. Then we get

$$\begin{aligned} \|S(|t|)x - S(|t|)y\| &\leq \left\| S(|t|)x - (J_{|t|/n}^A)^n x \right\| + \left\| (J_{|t|/n}^A)^n x - (J_{|t|/n}^A)^n y \right\| \\ &\quad + \left\| S(|t|)y - (J_{|t|/n}^A)^n y \right\| \\ &\leq \frac{2}{k+1} + \left\| (J_{|t|/n}^A)^n x - (J_{|t|/n}^A)^n y \right\| \\ &\leq \frac{2}{k+1} + \|x - y\| \end{aligned}$$

for any $n \geq \omega^S(k, b, T)$ with $b > \|x\|, \|y\|, \|v\|, \|w\|$ with $v \in Ax$ and $w \in Ay$ as well as $T > |t|$ using (S). As k was arbitrary, we get the claim.

¹²Note that the theorems in Lemma 2.3 remain valid for the system H_p^ω if the indices of the resolvent are restricted to be $< \lambda_0$.

(3) Using item (1) and axiom (S2), if w.l.o.g. $v_n \in A\hat{x}_n$ for all n , we have

$$\begin{aligned} \|S(|t|)C(x \uparrow v) - S(|s|)C(x \uparrow v)\| &\leq \|S(|t|)C(x \uparrow v) - S(|t|)(\hat{x} \uparrow v)_n\| \\ &\quad + \|S(|t|)(\hat{x} \uparrow v)_n - S(|s|)(\hat{x} \uparrow v)_n\| \\ &\quad + \|S(|s|)(\hat{x} \uparrow v)_n - S(|s|)C(x \uparrow v)\| \\ &\leq 2 \cdot 2^{-n+3} + \|S(|t|)(\hat{x} \uparrow v)_n - S(|s|)(\hat{x} \uparrow v)_n\| \\ &\leq 2 \cdot 2^{-n+3} + 2\|t| - |s|\| \|v_n\|. \end{aligned}$$

Choosing $n = k + 5$, we get the claim for $\|t| - |s|\| \leq 2^{-(k+2)} / \max\{1, \|v_{k+5}\|\}$.

(4) Using item (2), axiom (S2) as well as (C), if w.l.o.g. $v_n \in A\hat{x}_n$ and $w_n \in A\hat{y}_n$ for all n , we have

$$\begin{aligned} \|S(|t|)(C(x \uparrow v)) - S(|t|)(C(y \uparrow w))\| &\leq \|S(|t|)(C(x \uparrow v)) - S(|t|)((\hat{x} \uparrow v)_k)\| \\ &\quad + \|S(|t|)((\hat{x} \uparrow v)_k) - S(|t|)((\hat{y} \uparrow w)_k)\| \\ &\quad + \|S(|t|)(C(y \uparrow w)) - S(|t|)((\hat{y} \uparrow w)_k)\| \\ &\leq 2 \cdot 2^{-k+3} + \|S(|t|)((\hat{x} \uparrow v)_k) - S(|t|)((\hat{y} \uparrow w)_k)\| \\ &\leq 2 \cdot 2^{-k+3} + \|(\hat{x} \uparrow v)_k - (\hat{y} \uparrow w)_k\| \\ &\leq 2 \cdot 2^{-k+3} + \|(\hat{x} \uparrow v)_k - C(x \uparrow v)\| + \|C(x \uparrow v) - C(y \uparrow w)\| \\ &\quad + \|C(y \uparrow w) - (\hat{y} \uparrow w)_k\| \\ &\leq 2 \cdot 2^{-k+3} + 2 \cdot 2^{-k+3} + \|C(x \uparrow v) - C(y \uparrow w)\|. \end{aligned}$$

As this holds for arbitrary k , we get the claim.

(5) Let $x \in \text{dom}A$. Using the previously introduced notation of $\bar{\cdot}$, we write

$$\begin{cases} [S(|t|)]^1 x = \overline{S(|t|)x}, \\ [S(|t|)]^{m+1} x = \overline{S(|t|)([S(|t|)]^m x)}. \end{cases}$$

Note that provably

$$[S(|t|)]^{m+1} x = S(|t|) ([S(|t|)]^m x)$$

which follows as in the discussion previous to this lemma. We now show by induction on m that provably

$$\forall k \exists N_m \forall n \geq N_m \left(|t|/n < \lambda_0 \rightarrow \left\| [S(|t|)]^m x - \left(\left(J_{|t|/n}^A \right)^m \right)^n x \right\| \leq 2^{-k} \right).$$

The induction base follows from (S1) as was already discussed above. For the induction step, let $N_m(k)$ be the choice function of the above statement. Then for arbitrary k , we get (using extensionality, see Remark 4.4, and nonexpansivity of $S(|t|)$ on the closure of the domain) that

$$\begin{aligned} &\left\| [S(|t|)]^{m+1} x - \left(\left(J_{|t|/n}^A \right)^{m+1} \right)^n x \right\| \\ &\leq \left\| S(|t|)[S(|t|)]^m x - \left(J_{|t|/n}^A \right)^n \left(\left(J_{|t|/n}^A \right)^m \right)^n x \right\| \\ &\leq \left\| S(|t|)[S(|t|)]^m x - \left(J_{|t|/n}^A \right)^n [S(|t|)]^m x \right\| \\ &\quad + \left\| \left(J_{|t|/n}^A \right)^n [S(|t|)]^m x - \left(J_{|t|/n}^A \right)^n \left(\left(J_{|t|/n}^A \right)^m \right)^n x \right\| \\ &\leq \left\| S(|t|)[S(|t|)]^m x - \left(J_{|t|/n}^A \right)^n [S(|t|)]^m x \right\| + \left\| [S(|t|)]^m x - \left(\left(J_{|t|/n}^A \right)^m \right)^n x \right\| \\ &\leq 2^{-k} \end{aligned}$$

for all n such that $|t|/n < \lambda_0$, $n \geq N_m(k+1)$ and such that n is large enough for

$$\left\| S(|t|)[S(|t|)]^m x - \left(J_{|t|/n}^A \right)^n [S(|t|)]^m x \right\| \leq 2^{-(k+1)}$$

which can be constructed as above. Therefore, given k , we in particular get

$$\begin{aligned}
& \| [S(|t|)]^m x - S(m|t)x \| \\
& \leq \left\| [S(|t|)]^m x - \left((J_{|t|/n}^A)^m \right)^n x \right\| + \left\| \left((J_{|t|/n}^A)^m \right)^n x - S(m|t)x \right\| \\
& \leq \left\| [S(|t|)]^m x - \left((J_{|t|/n}^A)^m \right)^n x \right\| + \left\| (J_{m|t|/mn}^A)^{mn} x - S(m|t)x \right\| \\
& \leq 2^{-(k+1)} + 2^{-(k+1)} \\
& \leq 2^{-k}
\end{aligned}$$

for any n such that

$$n \geq \max\{N_m(k+1), \omega^S(k+1, b, mT)\}$$

for $b > \|x\|, \|v\|$ and $T > |t|$ using (S1) and the previous result. As k was arbitrary, we get $[S(|t|)]^m x = S(m|t)x$. Using this, we provably we get

$$\begin{aligned}
S\left(\frac{l}{k} + \frac{r}{s}\right)x &= S\left(\frac{ls+rk}{ks}\right)x \\
&= \left[S\left(\frac{1}{ks}\right) \right]^{ls+rk} x \\
&= \left[S\left(\frac{1}{ks}\right) \right]^{ls} \left[S\left(\frac{1}{ks}\right) \right]^{rk} x \\
&= \left[S\left(\frac{1}{ks}\right) \right]^{ls} \overline{S\left(\frac{rk}{ks}\right)} x \\
&= S\left(\frac{ls}{ks}\right) \overline{S\left(\frac{rk}{ks}\right)} x \\
&= S\left(\frac{l}{k}\right) \overline{S\left(\frac{r}{s}\right)} x
\end{aligned}$$

where we have used the above items for extensionality of S (see again Remark 4.4). A continuity argument using item (3) now yields the claim for arbitrary reals $|t|$ and $|s|$. Further, the claim extends to the closure of the domain via another usual continuity argument. Both we do not spell out here. \square

Remark 4.4. The constant $S(t)x$ is provably extensional in $x \in \overline{\text{dom}A}$ for any $t \geq 0$ by (4) as well as in $t \geq 0$ for any $x \in \overline{\text{dom}A}$ by (3).

Remark 4.5. Note that by the above item (3), we have that if the operator A is majorizable in the sense of [47], i.e. if there exists a function $A^* : \mathbb{N} \rightarrow \mathbb{N}$ such that

$$\forall b \in \mathbb{N} \forall x \in \text{dom}A \cap \overline{B}_b(0) \exists y \in X (\|y\| \leq A^*b \wedge y \in Ax),$$

then the semigroup \mathcal{S} generated by A through the Crandall-Liggett formula is uniformly equicontinuous in the sense of [29], i.e. there exists a function $\omega : \mathbb{N} \times \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ such that

$$\begin{aligned}
& \forall b \in \mathbb{N} \forall q \in \overline{\text{dom}A} \cap B_b(0) \forall m \in \mathbb{N} \forall K \in \mathbb{N} \forall t, t' \in [0, K] \\
& \left(|t - t'| < 2^{-\omega_{K,b}(m)} \rightarrow \|T(t)q - T(t')q\| < 2^{-m} \right).
\end{aligned}$$

Concretely, assuming w.l.o.g. that A^* is nondecreasing, this so-called *modulus of uniform equicontinuity* for \mathcal{S} can be given by

$$\omega_{K,b}(m) = 2^{-(m+3)} / \max\{1, A^*(b+1)\}.$$

Note in particular that this modulus is independent on the parameter K .

We now come to the main theoretical result of this paper which comprises a proof-theoretic bound extraction theorem akin to the usual metatheorems of proof mining for the system H_p^ω . The proof of this metatheorem follows the general outline of most other proofs in the literature discussed before and since the proof is very much standard in this way, we omit most of the details and in the following just sketch the majorizability of the new constant S .

Theorem 4.6. *Let τ be admissible, δ be of degree 1 and s be a closed term of H_p^ω of type $\delta \rightarrow \sigma$ for admissible σ . Let $B_\forall(x, y, z, u)/C_\exists(x, y, z, v)$ be \forall -/ \exists -formulas of H_p^ω with only $x, y, z, u/x, y, z, v$ free. Let Δ be a set of formulas of the form $\forall \underline{a} \exists \underline{b} \leq_\sigma \underline{r} \underline{a} \forall \underline{c} \neg F_{qf}(\underline{a}, \underline{b}, \underline{c})$ where F_{qf} is quantifier-free, the types in $\underline{\delta}$, $\underline{\sigma}$ and $\underline{\gamma}$ are admissible and \underline{r} are tuples of closed terms of appropriate type. If*

$$H_p^\omega + \Delta \vdash \forall x^\delta \forall y \leq_\sigma s(x) \forall z^\tau (\forall u^\mathbb{N} B_\forall(x, y, z, u) \rightarrow \exists v^\mathbb{N} C_\exists(x, y, z, v)),$$

then one can extract a partial functional $\Phi : S_\delta \times S_{\hat{\tau}} \times \mathbb{N} \times \mathbb{N}^\mathbb{N} \rightarrow \mathbb{N}$ which is total and (bar-recursively) computable on $\mathcal{M}_\delta \times \mathcal{M}_{\hat{\tau}} \times \mathbb{N} \times \mathbb{N}^\mathbb{N}$ and such that for all $x \in S_\delta$, $z \in S_{\hat{\tau}}$, $z^* \in S_{\hat{\tau}}$ and all $n \in \mathbb{N}$ and $\omega \in \mathbb{N}^\mathbb{N}$, if $z^* \gtrsim z$ and $n \geq_{\mathbb{R}} m_{\hat{\gamma}}, |\hat{\gamma}|, \|c_X\|_X, \|d_X\|_X, |\lambda_0|, m_{\lambda_0}, m'_{\hat{\gamma}}$, then

$$\begin{aligned} \mathcal{S}^{\omega, X} \models \forall y \leq_\sigma s(x) (\forall u \leq_{\mathbb{N}} \Phi(x, z^*, n, \omega) B_\forall(x, y, z, u) \\ \rightarrow \exists v \leq_{\mathbb{N}} \Phi(x, z^*, n, \omega) C_\exists(x, y, z, v)) \end{aligned}$$

holds for $\mathcal{S}^{\omega, X}$ whenever $\mathcal{S}^{\omega, X} \models \Delta$ where $\mathcal{S}^{\omega, X}$ is defined via any Banach space $(X, \|\cdot\|)$ with

- (1) χ_A interpreted by the characteristic function of an accretive operator A satisfying the range condition $\overline{\text{dom}A} \subseteq \bigcap_{\lambda_0 > \gamma > 0} \text{dom}J_\gamma^A$,
- (2) J^{XA} interpreted by the corresponding resolvents J_γ^A for $\lambda_0 > \gamma \geq 0$
- (3) j interpreted as discussed in Section 3.2,
- (4) S interpreted by the semigroup generated by A via the Crandall-Liggett formula on $\overline{\text{dom}A}$, and 0 otherwise,
- (5) d_X, c_X interpreted by a pair $(c, d) \in A$ witnessing $A \neq \emptyset$,
- (6) ω^S interpreted by a rate of convergence for the limit generating the semigroup on $\text{dom}A$,

and with the other constants naturally interpreted so that the respective axioms are satisfied.

Further: if $\hat{\tau}$ is of degree 1, then Φ is a total computable functional and if the claim is proved without DC, then τ may be arbitrary and Φ will be a total functional on $S_\delta \times S_{\hat{\tau}} \times \mathbb{N}$ which is primitive recursive in the sense of Gödel.

Proof. The proof given in [47] immediately extends to this system, noticing the change in the model of set-valued and majorizable functionals discussed in the context of j as well as Remark 4.1. In particular, note also that all axioms added to H_p^ω are purely universal and that the new constants other than S can be majorized as discussed throughout the previous sections. For the last constant S , we can argue for the majorizability as follows: In the context of the axiom (V)', stating that $\text{dom}A$ is not empty using the constants c_X and d_X , majorization of the constant S on $t \geq 0$ and $x \in \text{dom}A$ follows rather immediately. It is straightforward to obtain that

$$\hat{\mathcal{V}}_p^\omega \vdash \forall x^X, \lambda^{\mathbb{N}^\mathbb{N}}, n^\mathbb{N} \left(x \in \text{dom}(J_{|\lambda|}^A) \rightarrow \left\| (J_{|\lambda|}^A)^n x -_X x \right\|_X \leq_{\mathbb{R}} n \left\| J_{|\lambda|}^A x -_X x \right\|_X \right).$$

Therefore, we have for $x \in \text{dom}A$ with $v \in Ax$ and $b > \|x\|, \|v\|$ and for $t \geq 0$ with $T > t$ that for $n \geq (\omega^S(0, b, T) + \lceil T/\lambda_0 \rceil)$:¹³

$$\begin{aligned} \|S(t)x\| &\leq \left\| S(t)x - (J_{t/n}^A)^n x \right\| + \left\| (J_{t/n}^A)^n x \right\| \\ &\leq 1 + \left\| (J_{t/n}^A)^n x - (J_{t/n}^A)^n c_X \right\| + \left\| (J_{t/n}^A)^n c_X \right\| \\ &\leq 1 + \|x - c_X\| + \|c_X\| + n \left\| J_{t/n}^A c_X - c_X \right\| \\ &\leq 1 + \|x\| + 2\|c_X\| + T\|d_X\| \end{aligned}$$

which follows from the axioms (S) and (RC) $_{\lambda_0}$. This extends to $\overline{\text{dom}A}$ as follows: for $x \in \text{dom}A$ and $x_n \rightarrow x$ with rate of convergence 2^{-n} and where $x_n \in \text{dom}A$, we have $\|x_0 - x\| \leq 1$ and $\|S(t)x - S(t)x_0\| \leq 1$ and thus

$$\begin{aligned} \|S(t)x\| &\leq 1 + \|S(t)x_0\| \\ &\leq 2 + \|x_0\| + 2\|c_X\| + T\|d_X\| \\ &\leq 3 + \|x\| + 2\|c_X\| + T\|d_X\|. \end{aligned}$$

□

¹³We can choose e.g. $n = \omega^S(0, b, T) + \lceil T/\lambda_0 \rceil(0) + 1$ which can be represented through a closed term.

Also Theorem 2.7 extends to an intuitionistic version $H_{i,p}^\omega$ of the system H_p^ω in that fashion. Concretely, let $H_{i,p}^\omega$ be defined as the extension/modification of $\mathcal{V}_{i,p}^\omega$ with the same constants and axioms as were added to \mathcal{V}_p^ω to form H_p^ω . Then the following semi-constructive bound extraction theorem holds:

Theorem 4.7. *Let δ be of the form $\mathbb{N} \rightarrow (\dots \rightarrow \mathbb{N})$ and σ, τ be arbitrary, s be a closed term of suitable type. Let Γ_- be a set of sentences of the form $\forall \underline{u} \leq \underline{\beta} (C(\underline{u}) \rightarrow \exists \underline{v} \leq \underline{\beta} \underline{t}\underline{u} \neg D(\underline{u}, \underline{v}))$ with $\underline{\zeta}, \underline{\beta}$ and C, D arbitrary types and formulas respectively and \underline{t} be closed terms. Let $B(x, y, z) / A(x, y, z, u)$ be arbitrary formulas of $H_{i,p}^\omega$ with only $x, y, z/x, y, z, u$ free. If*

$$H_{i,p}^\omega + \Gamma_- \vdash \forall x^\delta \forall y \leq_\sigma (x) \forall z^\tau (\neg B(x, y, z) \rightarrow \exists u^\mathbb{N} A(x, y, z, u)),$$

one can extract a $\Phi : \mathcal{S}_\delta \times \mathcal{S}_{\hat{\tau}} \times \mathbb{N} \times \mathbb{N}^\mathbb{N} \rightarrow \mathbb{N}$ with is primitive recursive in the sense of Gödel's T such that for any $x \in \mathcal{S}_\delta$, any $y \in \mathcal{S}_\sigma$ with $y \leq_\sigma s(x)$, any $z \in \mathcal{S}_\tau$ and $z^* \in \mathcal{S}_{\hat{\tau}}$ with $z^* \succeq z$ and any $n \in \mathbb{N}$ and $\omega \in \mathbb{N}^\mathbb{N}$ with $n \geq_{\mathbb{R}} m_{\tilde{\gamma}}, |\tilde{\gamma}|, \|c_X\|_X, \|d_X\|_X, |\lambda_0|, m_{\lambda_0}, m'_{\tilde{\gamma}}$, we have that

$$\mathcal{S}^{\omega, X} \models \exists u \leq_{\mathbb{N}} \Phi(x, z^*, n, \omega) (\neg B(x, y, z) \rightarrow A(x, y, z, u))$$

holds for $\mathcal{S}^{\omega, X}$ whenever $\mathcal{S}^{\omega, X} \models \Gamma_-$, where $\mathcal{S}^{\omega, X}$ is defined via any Banach space $(X, \|\cdot\|)$ with the constants interpreted as in Theorem 4.6.

Using the previous arguments regarding the majorizability of the new constants, the proof is a straightforward adaptation of the proof for the central semi-constructive metatheorem for the system $\mathcal{A}^\omega[X, \|\cdot\|]$ given in [15] and we thus omit any further details.

5. TWO APPLICATIONS

The current section now employs the formal systems and the bound extraction theorems introduced above to provide quantitative information on the asymptotic behavior of these semigroups in the form of two case studies. Concretely, motivated by the results of Pazy [44] for iterations of nonexpansive mappings, the following two asymptotic results on the resolvents and the semigroups generated by the underlying operator were established:¹⁴

Theorem 5.1 (Plant [50]). *Let X be uniformly convex, A satisfy the range condition $(RC)_{\lambda_0}$ and let $x \in \text{dom}A$. Then*

$$\lim_{\lambda_0 > t \rightarrow 0^+} \frac{\|J_t x - S(t)x\|}{t} = 0.$$

Theorem 5.2 (Reich [53]). *Let X be uniformly convex, A satisfy the range condition (RC) and let $x \in \text{dom}A$. Then*

$$\lim_{t \rightarrow \infty} \frac{\|J_t x - S(t)x\|}{t} = 0.$$

A usual application of negative translation and monotone functional interpretation as used by the metatheorems suggest the extractability ‘metastability-like’ rates here (provided that the proof formalizes in the underlying systems). However, as we will see, classical logic features in these proofs only in two ways: at first, it features in some of the basic underlying convergence results in which case the limits are decreasing and a rate of convergence can thus nevertheless be obtained using the metatheorem from Theorem 4.6. For both results, the proof then proceeds via a case distinction on real numbers between $= 0$ and > 0 . In both results, the proofs for the ‘ $= 0$ ’-cases are trivial and rates of convergence can be immediately extracted. While the proofs for the ‘ > 0 ’-cases are nontrivial, they are nevertheless essentially constructive which allows, through the use of the semi-constructive metatheorem of Theorem 4.7, for the extraction of full rates of convergence for both limits exhibited above, under the appropriate quantitative reformulations of the ‘ > 0 ’-assumption, respectively. So, a rate of convergence can be obtained in either case, for both results. Only in the combination of these rates to a rate for the full result, the issues from the use of classical logic could feature but as we will see, in both cases the rates can be smoothed to be combined into a full rate of convergence for the whole result.

The next two subsections now present the extractions of the quantitative results and in that context do not explicitly focus on the logical particularities of the extraction which will only be discussed in the last subsection. In that way, to present these results in a way more amenable to the usual literature of the theory of semigroups, we also move to using ε 's for the errors instead of 2^{-k} or similar constructions using natural numbers.

In general, the main assumption featuring in both results is the uniform convexity of the underlying space which can be treated, as extensively discussed in the proof mining literature starting from the earliest works on the treatment of abstract spaces (see [25]), by a so-called modulus of uniform convexity:

¹⁴To simplify the notation in the following, we drop the superscript of the operator A from the resolvent.

Definition 5.3. A modulus of uniform convexity for a space X is a mapping $\eta : (0, 2] \rightarrow (0, 1]$ such that

$$\forall \varepsilon \in (0, 2] \forall x, y \in X \left(\|x\|, \|y\| \leq 1 \wedge \|x - y\| \geq \varepsilon \rightarrow \left\| \frac{x + y}{2} \right\| \leq 1 - \eta(\varepsilon) \right).$$

Of course, the rates in general will then depend similarly on such a modulus. It should be further noted that this modulus is conceptually related to the common analytic notion of a modulus of convexity $\delta : [0, 2] \rightarrow [0, 1]$ (implicit already used in e.g. [9]) defined as

$$\delta(\varepsilon) = \inf \{1 - \|x + y\|/2 \mid \|x\| = \|y\| = 1, \|x - y\| = \varepsilon\}$$

In fact, as is well-known, uniformly convex spaces are characterized by $\delta(\varepsilon) > 0$ for $\varepsilon > 0$ and the modulus of uniform convexity η effectively provides a witness for this inequality in the form of a lower bound, i.e. that $\delta(\varepsilon) \geq \eta(\varepsilon) > 0$ for $\varepsilon \in (0, 2]$.

The proofs of the results of both Plant and Reich make an essential use of δ but closer inspection reveals that they only rely on a lower bound on $\delta(\varepsilon)$ greater than 0 which therefore can be substituted by the modulus of uniform convexity η . Note that η can be assumed to be nondecreasing which we will do w.l.o.g. in the following. In that case, one in particular has that $\eta(\varepsilon) < \eta(\delta)$ implies $\varepsilon \leq \delta$.

5.1. An analysis of Plant's result. The proof proceeds by establishing that the sequence

$$\frac{x - J_t x}{t}, \quad (t \rightarrow 0^+)$$

is Cauchy and that we have the limit

$$\lim_{t, s/t \rightarrow 0^+} \left\| \frac{x - J_t x}{t} - \frac{x - S(s)x}{s} \right\| = 0.$$

Both results rely crucially on the existence and equality of the limits

$$\lim_{t \rightarrow 0^+} \frac{\|x - J_t x\|}{t} \quad \text{and} \quad \lim_{t \rightarrow 0^+} \frac{\|x - S(t)x\|}{t}.$$

The first sequence is nondecreasing for $t \rightarrow 0^+$ and bounded for $x \in \text{dom}A$ with $v \in Ax$ by $\|v\|$ (see e.g. [1]). Following [10], we denote the limit by $|Ax|$ which naturally satisfies $|Ax| \leq \|v\|$. The second limit was shown to coincide with $|Ax|$ in [10].

Now, the proof given in [50] crucially relies on the use of the limit operator $|Ax|$ and some elementary properties thereof. For the following, we denote the expression $(x - J_t x)/t$ (which is just the Yosida approximate) by $A_t x$, in contrast to Plants notation.

As discussed briefly in the introduction, one of the main theoretical obstacles in treating accretive and monotone operators is the use of extensionality in proofs as this requires one to provide some sort of quantitative modulus of uniform continuity as dictated by the monotone functional interpretation. While this will be discussed in more detail in the later logical remarks, we also find here that the main convergence principle

$$\|A_t x\| \rightarrow |Ax| \quad \text{for } x \in \text{dom}A, t \rightarrow 0^+$$

on which the proof of Plant relies can be recognized as a particular weak version of such a kind of extensionality statement, namely it can be shown that it is provably equivalent to the lower semi-continuity on $\text{dom}A$ of the operator $|A \cdot |$ associated with A .

As in the case of the functional $\langle \cdot, \cdot \rangle_s$, the logical methodology based on the monotone Dialectica interpretation now implies the following quantitative version of this statement: under this interpretation, the statement is upgraded to the existence of a 'modulus of uniform lower-semicontinuity' $\varphi : \mathbb{R}_{>0} \times \mathbb{N} \rightarrow \mathbb{R}_{>0}$ which, as will be discussed later in the logical remarks, it essentially a modulus of uniform continuity and we thus assume that φ satisfies

$$\forall b \in \mathbb{N}, \varepsilon \in \mathbb{R}_{>0}, (x, u), (y, v) \in A \\ (\|x\|, \|u\|, \|y\|, \|v\| \leq b \wedge \|x - y\| \leq \varphi(\varepsilon, b) \rightarrow |Ax| - |Ay| \leq \varepsilon).$$

Based on the above mentioned equivalence, this modulus can then be used to derive a rate of convergence for the Yosida approximates towards $|Ax|$.

Lemma 5.4. *Let X be a Banach space, let φ be a modulus of uniform continuity for $|A \cdot|$ and let n satisfy $n \geq \|c\|, \|d\|, \lambda_0, \tilde{\gamma}$ for $(c, d) \in A$ and $0 < \tilde{\gamma} < \lambda_0$. Then for $x \in \text{dom}A$ with $v \in Ax$ and $b \in \mathbb{N}^{*15}$ with $b \geq \|x\|, \|v\|$, we have*

$$\forall \varepsilon > 0 \forall t \in (0, \varphi_1(\varepsilon, b, n, \varphi)] \left(|Ax| - \frac{\|x - J_t^A x\|}{t} \leq \varepsilon \right)$$

where

$$\varphi_1(\varepsilon, b, n, \varphi) := \min\{\varphi(\varepsilon, b + 2n + 3n^2)/b, \lambda_0/2\}.$$

Proof. Let ε be given and let $t \leq \varphi_1(\varepsilon, b, n, \varphi)$. We at first have $\|A_t x\| \leq \|v\| \leq b$ as well as

$$\begin{aligned} \|J_t x\| &\leq \|x\| + 2\|c\| + (2\tilde{\gamma} + t)\|d\| \\ &\leq \|x\| + 2\|c\| + (2\tilde{\gamma} + \lambda_0)\|d\| \\ &\leq \|x\| + 2n + (2n + n)n \\ &\leq \|x\| + 2n + 3n^2 \end{aligned}$$

using Lemma 2.3 (as $t < \lambda_0$). Now as $A_t x \in AJ_t x$, we have $|AJ_t x| \leq \|A_t x\|$ and thus

$$|Ax| - \|A_t x\| \leq |Ax| - |AJ_t x|.$$

Now, we get

$$\|x - J_t x\| \leq t\|v\| \leq \varphi_1(\varepsilon, b, n, \varphi)b \leq \varphi(\varepsilon, b + 2n + 3n^2)$$

and thus, as $v \in Ax$ and $A_t x \in AJ_t x$ with $\|x\|, \|v\|, \|J_t x\|, \|A_t x\| \leq b + 2n + 3n^2$, we have

$$|Ax| - \|A_t x\| \leq |Ax| - |AJ_t x| \leq \varepsilon$$

which is the claim. \square

As mentioned before, the fact that

$$\lim_{t \rightarrow 0^+} \frac{\|x - S(t)x\|}{t} = |Ax|$$

was proved in [10] and proceeds by establishing that $\|x - S(t)x\|/t \leq |Ax|$ for any $t > 0$ as well as

$$\liminf_{t \rightarrow 0^+} \frac{\|S(t)x - x\|}{t} \geq |Ax|$$

and in that way crucially relies on the limit operator $|A \cdot|$ as well. This latter result relies on a result established by Miyadera in [6]¹⁶ that

$$\limsup_{t \rightarrow 0^+} \left\langle \frac{S(t)x - x}{t}, \zeta^* \right\rangle \leq \langle y_0, x_0 - x \rangle_s$$

for $y_0 \in Ax_0$, $x \in \overline{\text{dom}A}$ and $\zeta^* \in J(x - x_0)$.

The proof given by Crandall actually only invokes this result for $x \in \text{dom}A$ and it is further sufficient to obtain it only for *some* $\zeta^* \in J(x - x_0)$ which can in that context therefore be assumed to coincide with j_{v, y_0} , the functions witnessing accretivity as discussed before, for $v \in Ax$ witnessing $x \in \text{dom}A$. Lastly, the proof relies crucially on the use of the functional $\langle \cdot, \cdot \rangle_s$ and in particular on the upper semicontinuity of this functional. In that way, based on the logical methodology that upgrades this upper semicontinuity to a modulus of uniform continuity, we extract the following quantitative version of the above fragment of Miyadera's result:

Lemma 5.5. *Let A be accretive on a Banach space X and let S be the semigroup on $\overline{\text{dom}A}$ generated by the Crandall-Liggett formula and let ω be such that*

$$\forall x, y, z \in X, b \in \mathbb{N}, \varepsilon > 0 (\|x\|, \|z\| \leq b \wedge \|x - y\| \leq \omega(b, \varepsilon) \rightarrow \langle z, y \rangle_s \leq \langle z, x \rangle_s + \varepsilon).$$

For $x \in \text{dom}A$ with $v \in Ax$ and $y_0 \in Ax_0$ where $b \in \mathbb{N}^*$ with $\|x\|, \|v\|, \|x_0\|, \|y_0\| \leq b$:

$$\forall \varepsilon > 0 \forall t \in (0, \psi(\varepsilon, b, \omega)] \left(\left\langle \frac{S(t)x - x}{t}, j_{v, y_0}(x - x_0) \right\rangle \leq \langle y_0, x_0 - x \rangle_s + \varepsilon \right)$$

where

$$\psi(\varepsilon, b, \omega) := \frac{\omega(2b, \varepsilon)}{2b}.$$

¹⁵Throughout, we write \mathbb{N}^* for the natural numbers without 0.

¹⁶The result goes back to earlier work by Brezis [5] with a special case already contained in [11] and more general results proved in [12].

Proof. At first, given a ε , we get for any $t \in (0, \frac{\varepsilon}{2b}]$ and for all $v \in Ax$ with $\|x\|, \|v\| \leq b$ that

$$\|x - S(t)x\| = \|S(0)x - S(t)x\| \leq 2\|v\|t \leq 2b\frac{\varepsilon}{2b} \leq \varepsilon$$

by Lemma 4.3, (1). Now, as in Miyadera's proof from [6], we get

$$\langle S(t)x - x, j_{v, y_0}(x - x_0) \rangle \leq \int_0^t \langle y_0, x_0 - S(\tau)x \rangle_s d\tau.$$

Then for $b \geq \|x\|, \|v\|, \|x_0\|, \|y_0\|$, we get

$$\langle y_0, x_0 - S(t)x \rangle_s \leq \langle y_0, x_0 - x \rangle_s + \varepsilon$$

for any $t \in (0, \psi(\varepsilon, b, \omega)]$ as by the above, we have

$$\|S(t)x - x\| \leq \omega(2b, \varepsilon)$$

for all such t by assumption on ω and since we trivially have $\|x - x_0\| \leq 2b$. Thus in particular we have

$$\begin{aligned} \langle S(t)x - x, j_{v, y_0}(x - x_0) \rangle &\leq \int_0^t \langle y_0, x_0 - S(\tau)x \rangle_s d\tau \\ &\leq t(\langle y_0, x_0 - x \rangle_s + \varepsilon) \end{aligned}$$

which gives the claim. \square

Then, by following the proof given in [10], we obtain a quantitative version of the crucial direction

$$\liminf_{t \rightarrow 0^+} \frac{\|S(t)x - x\|}{t} \geq |Ax|$$

of Crandall's proof. Now, already here, a case distinction on whether $|Ax| = 0$ or $|Ax| > 0$ features in the proof of Crandall and the following result first provides a quantitative result on the latter case.

Lemma 5.6. *Let A be accretive on a Banach space X and let S be the semigroup on $\overline{\text{dom}A}$ generated by the Crandall-Liggett formula and let ω be such that*

$$\forall x, y, z \in X, b \in \mathbb{N}, \varepsilon > 0 (\|x\|, \|z\| \leq b \wedge \|x - y\| \leq \omega(b, \varepsilon) \rightarrow \langle z, y \rangle_s \leq \langle z, x \rangle_s + \varepsilon).$$

Let further φ be a modulus of uniform continuity for $|A \cdot |$ and let n be as above. Then for $x \in \text{dom}A$ with $v \in Ax$ and $b \in \mathbb{N}^$ with $b \geq \|x\|, \|v\|$ and where $|Ax| \geq c$ for $c \in \mathbb{R}_{>0}$, we have*

$$\forall \varepsilon > 0 \forall t \in (0, \varphi'_2(\varepsilon, b, c, n, \varphi, \omega)] \left(|Ax| - \frac{\|x - S(t)x\|}{t} \leq \varepsilon \right)$$

where

$$\varphi'_2(\varepsilon, b, c, n, \varphi, \omega) := \psi(\varepsilon c \min\{\varphi_1(\min\{\varepsilon/2, c/2\}, b, n, \varphi), \lambda_0/2\}/4, b + 2n + 3n^2, \omega)$$

with ψ as in Lemma 5.5 and φ_1 as in Lemma 5.4.

Proof. Using Lemma 5.5, we get

$$\forall \varepsilon > 0 \forall t \in (0, \psi(\varepsilon, b, \omega)] \left(\left\langle \frac{S(t)x - x}{t}, j_{v, y_0}(x - x_0) \right\rangle \leq \langle y_0, x_0 - x \rangle_s + \varepsilon \right)$$

for $\|x\|, \|v\|, \|x_0\|, \|y_0\| \leq b$. Now, for $y_0 = A_\lambda x$ and $x_0 = J_\lambda x$ with $\lambda < \lambda_0$, we have

$$\langle y_0, x_0 - x \rangle_s = -\lambda \|A_\lambda x\|^2$$

as well as

$$\begin{aligned} \left\langle \frac{S(t)x - x}{t}, j_{v, y_0}(x - x_0) \right\rangle &\geq - \left\| \frac{S(t)x - x}{t} \right\| \|x - J_\lambda x\| \\ &= - \left\| \frac{S(t)x - x}{t} \right\| \lambda \|A_\lambda x\|. \end{aligned}$$

Therefore, we obtain

$$\forall \varepsilon > 0 \forall t \in (0, \psi(\varepsilon, b + 2n + 3n^2, \omega)] \left(\left\| \frac{S(t)x - x}{t} \right\| \|A_\lambda x\| \geq \|A_\lambda x\|^2 - \frac{\varepsilon}{\lambda} \right)$$

for all such λ since $b + 2n + 3n^2 \geq \|J_\lambda x\|$ and $b \geq \|A_\lambda x\|$ as before. Since $|Ax| \geq c$, we have that for $\lambda \leq \min\{\varphi_1(c/2, b, n, \varphi), \lambda_0/2\}$ that

$$c/2 \leq |Ax| - c/2 \leq \|A_\lambda x\|$$

by Lemma 5.4. Therefore, we have that

$$\forall \varepsilon > 0 \forall t \in (0, \psi(\varepsilon, b + 2n + 3n^2, \omega)] \left(\left\| \frac{S(t)x - x}{t} \right\| \geq \|A_\lambda x\| - \frac{\varepsilon}{\lambda c/2} \right)$$

for all $\lambda \leq \min\{\varphi_1(c/2, b, n, \varphi), \lambda_0/2\}$ and thus in particular

$$\begin{aligned} |Ax| - \left\| \frac{S(t)x - x}{t} \right\| &\leq |Ax| - \|A_\lambda x\| + \frac{\varepsilon}{\lambda c/2} \\ &\leq \varepsilon/2 + \frac{\varepsilon}{\lambda c/2} \end{aligned}$$

for all $t \leq \psi(\varepsilon, b + 2n + 3n^2, \omega)$ and for all $\lambda \leq \min\{\varphi_1(\min\{\varepsilon/2, c/2\}, b, n, \varphi), \lambda_0/2\}$. Thus, lastly, for

$$t \leq \psi(b + 2n + 3n^2, \varepsilon c \min\{\varphi_1(\min\{\varepsilon/2, c/2\}, b, n), \lambda_0\}/4)$$

we have

$$|Ax| - \left\| \frac{S(t)x - x}{t} \right\| \leq \varepsilon.$$

□

For the other case, i.e. where $|Ax| = 0$, it is immediately clear that for $|Ax| \leq \varepsilon$, we get

$$|Ax| - \frac{\|x - S(t)x\|}{t} \leq |Ax| \leq \varepsilon$$

for all t . However, this allows for a smoothing of the above case distinction (see the later logical remarks for further discussions of this) in the form of the following lemma:

Lemma 5.7. *Let A be accretive on a Banach space X and let S be the semigroup on $\overline{\text{dom}A}$ generated by the Crandall-Liggett formula and let ω be such that*

$$\forall x, y, z \in X, b \in \mathbb{N}, \varepsilon > 0 (\|x\|, \|z\| \leq b \wedge \|x - y\| \leq \omega(b, \varepsilon) \rightarrow \langle z, y \rangle_s \leq \langle z, x \rangle_s + \varepsilon).$$

Let further φ be a modulus of uniform continuity for $|A \cdot |$ and let n be as above. Then for $x \in \text{dom}A$ with $v \in Ax$ and $b \in \mathbb{N}^$ with $b \geq \|x\|, \|v\|$, we have*

$$\forall \varepsilon > 0 \forall t \in (0, \varphi_2(\varepsilon, b, n, \omega, \varphi)] \left(|Ax| - \frac{\|x - S(t)x\|}{t} \leq \varepsilon \right)$$

where

$$\varphi_2(\varepsilon, b, n, \omega, \varphi) := \psi(\varepsilon^2 \min\{\varphi_1(\varepsilon/2, b, n, \varphi), \lambda_0/2\}/4, b + 2n + 3n^2, \omega).$$

with ψ as in Lemma 5.5 and φ_1 as in Lemma 5.4.

Proof. Let ε be given. Then either $|Ax| \leq \varepsilon$ whereas

$$|Ax| - \frac{\|x - S(t)x\|}{t} \leq \varepsilon$$

for any t . Otherwise, we have $|Ax| \geq \varepsilon$ and thus from Lemma 5.6, it follows that

$$|Ax| - \frac{\|x - S(t)x\|}{t} \leq \varepsilon$$

for all

$$t \leq \psi(\varepsilon^2 \min\{\varphi_1(\varepsilon/2, b, n, \varphi), \lambda_0/2\}/4, b + 2n + 3n^2, \omega).$$

□

Using those two results, we can then give a quantitative version of the partial results on the way to Plants results discussed above, in the form of a rate of Cauchytness and a rate of convergence, respectively.

In that context, we follow the notation used in [50] and write

$$\alpha(a, b) = \left\| \frac{a}{\|a\|} - \frac{b}{\|b\|} \right\| \leq 2$$

where $a, b \neq 0$ for the generalized angle of Clarkson [9]. Similar to the proof given in [50], we rely on two fundamental inequalities of α :

Lemma 5.8 (essentially [9]). *Let $a, b \neq 0$. Then*

$$| \|a\| \alpha(a, b) - \|a - b\| | \leq | \|a\| - \|b\| |.$$

If further $a + b \neq 0$, then

$$\|a + b\| \leq (1 - 2\eta(\alpha(a + b, a))) \|a\| + \|b\|$$

where η is a modulus of uniform convexity for the space X .

Lemma 5.9. *Let X be a uniformly convex Banach space with a nondecreasing modulus of uniform convexity $\eta : (0, 2] \rightarrow (0, 1]$. Let further φ be a modulus of uniform continuity for $|A \cdot |$ and let n be as above. Let further $x \in \text{dom}A$ with $v \in Ax$ and $b \in \mathbb{N}^*$ with $b \geq \|x\|, \|v\|$. Suppose $|Ax| \geq c$ for $c \in \mathbb{R}_{>0}$. Then*

$$\forall \varepsilon > 0 \forall t \in (0, \varphi'_3(\varepsilon, b, c, \eta, n, \varphi)] \forall s \in (0, t) \left(\left\| \frac{x - J_t x}{t} - \frac{x - J_s x}{s} \right\| \leq \varepsilon \right)$$

where

$$\begin{aligned} \varphi'_3(\varepsilon, b, c, \eta, n, \varphi) := & \min\{\varphi_1(\varepsilon/3, b, n, \varphi), \varphi_1(\eta(\min\{\varepsilon/3b, 2\})c/2, b, n, \varphi), \\ & \varphi_1(c/2, b, n, \varphi), \lambda_0/2\} \end{aligned}$$

with φ_1 as in Lemma 5.4.

Proof. We write $\alpha_{s,t} = \alpha(x - J_s x, x - J_t x)$ where $s \in (0, t)$ and $t \leq \lambda_0/2 < \lambda_0$. Using Lemma 5.8, we have

$$\|x - J_t x\| \leq (1 - 2\eta(\alpha_{s,t})) \|x - J_s x\| + \|J_t x - J_s x\|.$$

Using Lemma 2.3, items (3) and (5), we get

$$\begin{aligned} \|J_t x - J_s x\| &= \left\| J_s \left(\frac{s}{t} x + \frac{t-s}{t} J_t x \right) - J_s x \right\| \\ &\leq \left(1 - \frac{s}{t} \right) \|x - J_t x\|. \end{aligned}$$

Therefore, we have for $0 < t \leq \varphi_1(c/2, b, n, \varphi)$ and $s \in (0, t)$ that

$$\eta(\alpha_{s,t})c \leq 2\eta(\alpha_{s,t}) \|A_t x\| \leq 2\eta(\alpha_{s,t}) \|A_s x\| \leq \|A_s x\| - \|A_t x\| \leq |Ax| - \|A_t x\|.$$

By Lemma 5.4, we have for any ε that

$$\forall t \in (0, \min\{\varphi_1(\varepsilon, b, n, \varphi), \varphi_1(c/2, b, n, \varphi)\}], s \in (0, t) (\eta(\alpha_{s,t})c \leq \varepsilon)$$

which, in particular, implies

$$\begin{aligned} \forall t \in (0, \min\{\varphi_1(\eta(\min\{\varepsilon/3b\})c/2, b, n, \varphi), \varphi_1(c/2, b, n, \varphi)\}], s \in (0, t) \\ (\eta(\alpha_{s,t}) \leq \eta(\varepsilon/3b)/2) \end{aligned}$$

and using that η is nondecreasing, we get

$$\forall t \in (0, \min\{\varphi_1(\eta(\varepsilon/3b)c/2, b, n), \varphi_1(c/2, b, n), \lambda_0\}], s \in (0, t) (\alpha_{s,t} \leq \varepsilon/3b).$$

Using Lemma 5.8, $s < t$ as well as triangle inequality again, we now have

$$\begin{aligned} \left\| \frac{x - J_t x}{t} - \frac{x - J_s x}{s} \right\| &\leq \left\| \frac{\|x - J_t x\|}{t} \alpha_{s,t} - \left\| \frac{x - J_t x}{t} - \frac{x - J_s x}{s} \right\| \right\| + \frac{\|x - J_t x\|}{t} \alpha_{s,t} \\ &\leq \left\| \frac{x - J_t x}{t} \right\| - \left\| \frac{x - J_s x}{s} \right\| + \frac{\|x - J_t x\|}{t} \alpha_{s,t} \\ &\leq \left(|Ax| - \left\| \frac{x - J_t x}{t} \right\| \right) + b\alpha_{s,t}. \end{aligned}$$

Thus for $0 < t \leq \varphi'_3(\varepsilon, b, c, \eta, n, \varphi)$ and for $s \in (0, t)$, we have

$$\begin{aligned} \left\| \frac{x - J_t x}{t} - \frac{x - J_s x}{s} \right\| &\leq \varepsilon/3 + \varepsilon/3 + b\varepsilon/3b \\ &\leq \varepsilon \end{aligned}$$

by Lemma 5.4. □

Again, the case for $|Ax| = 0$ is trivial and yields the following quantitative version: if $|Ax| \leq \varepsilon/2$, then in particular

$$\left\| \frac{x - J_t^A x}{t} - \frac{x - J_s^A x}{s} \right\| \leq \|A_t x\| + \|A_s x\| \leq |Ax| + |Ax| \leq \varepsilon.$$

In that way, we get the following smoothing for both results combined.

Lemma 5.10. *Let X be a uniformly convex Banach space with a nondecreasing modulus of uniform convexity $\eta : (0, 2] \rightarrow (0, 1]$. Let further φ be a modulus of uniform continuity for $|A \cdot|$ and let n be as above. Let further $x \in \text{dom} A$ with $v \in Ax$ and $b \in \mathbb{N}^*$ with $b \geq \|x\|, \|v\|$. Then*

$$\forall \varepsilon > 0 \forall t \in (0, \varphi_3(\varepsilon, b, \eta, n, \varphi)] \forall s \in (0, t) \left(\left\| \frac{x - J_t^A x}{t} - \frac{x - J_s^A x}{s} \right\| \leq \varepsilon \right)$$

where

$$\begin{aligned} \varphi_3(\varepsilon, b, \eta, n, \varphi) := & \min\{\varphi_1(\varepsilon/3, b, n, \varphi), \varphi_1(\eta(\min\{\varepsilon/3b, 2\})\varepsilon/4, b, n, \varphi), \\ & \varphi_1(\varepsilon/4, b, n, \varphi), \lambda_0/2\} \end{aligned}$$

with $\varphi_1(\varepsilon, b, n)$ as in Lemma 5.4.

Proof. Let ε be given. Then either $|Ax| \leq \varepsilon/2$ whereas

$$\left\| \frac{x - J_t^A x}{t} - \frac{x - J_s^A x}{s} \right\| \leq \varepsilon$$

for any $t, s < \lambda_0$ as discussed above. Otherwise we have $|Ax| \geq \varepsilon/2$ and thus by Lemma 5.9, it follows that

$$\left\| \frac{x - J_t^A x}{t} - \frac{x - J_s^A x}{s} \right\| \leq \varepsilon$$

for $s \in (0, t)$ and

$$t \leq \min\{\varphi_1(\varepsilon/3, b, n, \varphi), \varphi_1(\eta(\min\{\varepsilon/3b, 2\})\varepsilon/4, b, n, \varphi), \varphi_1(\varepsilon/4, b, n, \varphi), \lambda_0/2\}.$$

□

Lemma 5.11 (Plant [50], Eq. (2.10)). *Let $x \in \text{dom} A$ and $t, \lambda > 0$. Then*

$$\|J_\lambda x - S(t)x\| \leq \left(1 - \frac{t}{\lambda}\right) \|x - J_\lambda x\| + \frac{2}{\lambda} \int_0^t \|x - S(s)x\| ds.$$

Lemma 5.12. *Let X be a uniformly convex Banach space with a nondecreasing modulus of uniform convexity $\eta : (0, 2] \rightarrow (0, 1]$ and let ω be such that*

$$\forall x, y, z \in X, b \in \mathbb{N}, \varepsilon > 0 (\|x\|, \|z\| \leq b \wedge \|x - y\| \leq \omega(b, \varepsilon) \rightarrow \langle z, y \rangle_s \leq \langle z, x \rangle_s + \varepsilon).$$

Let further φ be a modulus of uniform continuity for $|A \cdot|$ and let n be as above. Let further $x \in \text{dom} A$ with $v \in Ax$ and $b \in \mathbb{N}^$ with $b \geq \|x\|, \|v\|$. Suppose that $|Ax| \geq c$ for $c \in \mathbb{R}_{>0}$. Then*

$$\forall \varepsilon > 0 \forall t, \frac{s}{t} \in (0, \varphi'_4(\varepsilon, b, c, \eta, n, \omega, \varphi)] \left(\left\| \frac{x - J_t^A x}{t} - \frac{x - S(s)x}{s} \right\| \leq \varepsilon \right)$$

where

$$\begin{aligned} \varphi'_4(\varepsilon, b, c, \eta, n, \omega, \varphi) := & \min\{\varphi_1(\varepsilon/3, b, n, \varphi), \varphi_2(\varepsilon/3, b, n, \omega, \varphi), \\ & \varphi_1(\eta(\min\{\varepsilon, 2\})c/4, b, n, \varphi), \sqrt{\varphi_2(c/2, b, n, \omega, \varphi)}, \\ & \eta(\min\{\varepsilon, 2\})c/8b, \sqrt{\varphi_2(c/2, b, n, \omega, \varphi)}, 1, \lambda_0/2\} \end{aligned}$$

with φ_1, φ_2 as in Lemmas 5.4, 5.7, respectively.

Proof. We write $\alpha'_{s,t} = \alpha(x - S(s)x, x - J_t x)$ for $t, s < \lambda_0$. Using Lemma 5.8, we again obtain

$$\|x - J_t x\| \leq (1 - 2\eta(\alpha'_{s,t})) \|x - S(s)x\| + \|J_t x - S(s)x\|.$$

Using Lemma 5.11, we get for $t, s \leq \min\{\lambda_0/2, 1\}$:

$$\begin{aligned} \|x - J_t x\| &\leq (1 - 2\eta(\alpha'_{s,t})) \|x - S(s)x\| + \left(1 - \frac{s}{t}\right) \|x - J_t x\| + \frac{2}{t} \int_0^s \|x - S(\tau)x\| d\tau \\ &\leq (1 - 2\eta(\alpha'_{s,t})) \|x - S(s)x\| + \left(1 - \frac{s}{t}\right) \|x - J_t x\| + \frac{2}{t} \int_0^s s \frac{\|x - S(\tau)x\|}{\tau} d\tau \\ &\leq (1 - 2\eta(\alpha'_{s,t})) \|x - S(s)x\| + \left(1 - \frac{s}{t}\right) \|x - J_t x\| + \frac{s^2}{t} 2b \end{aligned}$$

which implies that

$$2\eta(\alpha'_{s,t}) \frac{\|x - S(s)x\|}{s} \leq |Ax| - \frac{\|x - J_t x\|}{t} + \frac{s}{t} 2b.$$

Now for

$$t \leq \min\{\varphi_1(\varepsilon/2, b, n, \varphi), \sqrt{\varphi_2(c/2, b, n, \omega, \varphi)}\}$$

and

$$\frac{s}{t} \leq \min\{\varepsilon/4b, \sqrt{\varphi_2(c/2, b, n, \omega, \varphi)}\}$$

we obtain that

$$s \leq t \frac{s}{t} \leq \varphi_2(c/2, b, n, \omega, \varphi)$$

and thus (using Lemma 5.7), we obtain

$$\begin{aligned} \eta(\alpha'_{s,t})c &\leq 2\eta(\alpha'_{s,t}) (|Ax| - c/2) \\ &\leq 2\eta(\alpha'_{s,t}) \frac{\|x - S(s)x\|}{s} \\ &\leq |Ax| - \frac{\|x - J_t x\|}{t} + \frac{s}{t} 2b \\ &\leq \varepsilon/2 + \frac{s}{t} 2b \\ &\leq \varepsilon/2 + 2b\varepsilon/4b \\ &\leq \varepsilon. \end{aligned}$$

Dividing by c , we get $\eta(\alpha'_{s,t}) \leq \frac{\varepsilon}{c}$ for all such t, s . Thus, using that η is nondecreasing, we have $\alpha'_{s,t} \leq \varepsilon$ for

$$t \leq \min\{\varphi_1(\eta(\min\{\varepsilon, 2\})c/4, b, n, \varphi), \sqrt{\varphi_2(c/2, b, n, \omega, \varphi)}\}$$

and

$$\frac{s}{t} \leq \min\{\eta(\min\{\varepsilon, 2\})c/8b, \sqrt{\varphi_2(c/2, b, n, \omega, \varphi)}\}.$$

Using Lemma 5.8 and triangle inequality again, we now have similarly to before

$$\begin{aligned} &\left\| \frac{x - J_t x}{t} - \frac{x - S(s)x}{s} \right\| \\ &\leq \left\| \frac{\|x - J_t x\|}{t} \alpha'_{s,t} - \left\| \frac{x - J_t x}{t} - \frac{x - S(s)x}{s} \right\| \right\| + \frac{\|x - J_t x\|}{t} \alpha'_{s,t} \\ &\leq \left\| \frac{\|x - J_t x\|}{t} - \left\| \frac{x - S(s)x}{s} \right\| \right\| + \frac{\|x - J_t x\|}{t} \alpha'_{s,t} \\ &\leq \left(|Ax| - \left\| \frac{x - J_t x}{t} \right\| \right) + \left(|Ax| - \left\| \frac{x - S(s)x}{s} \right\| \right) + b\alpha'_{s,t}. \end{aligned}$$

Thus for $0 < t, \frac{s}{t} \leq \varphi'_4(\varepsilon, b, c, \eta, n, \omega, \varphi)$, we have

$$\begin{aligned} \left\| \frac{x - J_t x}{t} - \frac{x - S(s)x}{s} \right\| &\leq \varepsilon/3 + \varepsilon/3 + b\varepsilon/3b \\ &\leq \varepsilon \end{aligned}$$

by Lemma 5.4 and Lemma 5.7. □

As before, smoothing this result can be achieved by extracting from the proof for the case of $|Ax| = 0$ the following quantitative version: if $|Ax| \leq \varepsilon/2$, then

$$\left\| \frac{x - J_t^A x}{t} - \frac{x - S(s)x}{s} \right\| \leq \frac{\|x - J_t^A x\|}{t} + \frac{\|x - S(s)x\|}{s} \leq |Ax| + |Ax| \leq \varepsilon.$$

Therefore, we obtain the following result:

Lemma 5.13. *Let X be a uniformly convex Banach space with a nondecreasing modulus of uniform convexity $\eta : (0, 2] \rightarrow (0, 1]$ and let ω be such that*

$$\forall x, y, z \in X, b \in \mathbb{N}, \varepsilon > 0 (\|x\|, \|z\| \leq b \wedge \|x - y\| \leq \omega(b, \varepsilon) \rightarrow \langle z, y \rangle_s \leq \langle z, x \rangle_s + \varepsilon).$$

Let further φ be a modulus of uniform continuity for $|A \cdot|$ and let n be as above. Let further $x \in \text{dom}A$ with $v \in Ax$ and $b \in \mathbb{N}^$ with $b \geq \|x\|, \|v\|$. Then*

$$\forall \varepsilon > 0 \forall t, \frac{s}{t} \in (0, \varphi_4(\varepsilon, b, \eta, n, \omega, \varphi)] \left(\left\| \frac{x - J_t^A x}{t} - \frac{x - S(s)x}{s} \right\| \leq \varepsilon \right)$$

where

$$\begin{aligned} \varphi_4(\varepsilon, b, \eta, n, \omega, \varphi) := & \min\{\varphi_1(\varepsilon/3, b, n, \varphi), \varphi_2(\varepsilon/3, b, n, \omega, \varphi), \\ & \varphi_1(\eta(\min\{\varepsilon, 2\})\varepsilon/8, b, n, \varphi), \sqrt{\varphi_2(\varepsilon/4, b, n, \omega, \varphi)}, \\ & \eta(\min\{\varepsilon, 2\})\varepsilon/16b, \sqrt{\varphi_2(\varepsilon/4, b, n, \omega, \varphi)}, 1, \lambda_0/2\} \end{aligned}$$

with φ_1, φ_2 as in Lemmas 5.4, 5.7, respectively.

Proof. Let ε be given. Then either $|Ax| \leq \varepsilon/2$ which implies

$$\left\| \frac{x - J_t^A x}{t} - \frac{x - S(s)x}{s} \right\| \leq \varepsilon$$

as above for any such t and s or $|Ax| \geq \varepsilon/2$ where now the result is implied for any

$$t, \frac{s}{t} \in (0, \varphi'_4(\varepsilon, b, \varepsilon/2, \eta, n, \omega, \varphi)]$$

by Lemma 5.12. □

Finally, a combination of these two quantitative results yields a quantitative version of the theorem of Plant.

Theorem 5.14. *Let X be a uniformly convex Banach space with a nondecreasing modulus of uniform convexity $\eta : (0, 2] \rightarrow (0, 1]$ and let ω be such that*

$$\forall x, y, z \in X, b \in \mathbb{N}, \varepsilon > 0 (\|x\|, \|z\| \leq b \wedge \|x - y\| \leq \omega(b, \varepsilon) \rightarrow \langle z, y \rangle_s \leq \langle z, x \rangle_s + \varepsilon).$$

Let further φ be a modulus of uniform continuity for $|A \cdot|$ and let n be as above. Let further $x \in \text{dom}A$ with $v \in Ax$ and $b \in \mathbb{N}^$ with $b \geq \|x\|, \|v\|$. Then*

$$\forall \varepsilon > 0 \forall t \in (0, \Phi(\varepsilon, b, \eta, \omega, \varphi, n)] \left(\frac{\|J_t^A x - S(t)x\|}{t} \leq \varepsilon \right)$$

where

$$\Phi(\varepsilon, b, \eta, \omega, \varphi, n) := (\min\{\varphi_3(\varepsilon/2, b, \eta, n, \varphi), \varphi_4(\varepsilon/2, b, \eta, n, \omega, \varphi)\})^2$$

with $\varphi_1 - \varphi_4$ as well as ψ defined by

$$\varphi_1(\varepsilon, b, n, \varphi) := \min\{\varphi(\varepsilon, b + 2n + 3n^2)/b, \lambda_0/2\},$$

$$\psi(\varepsilon, b, \omega) := \frac{\omega(2b, \varepsilon)}{2b},$$

$$\varphi_2(\varepsilon, b, n, \omega, \varphi) := \psi(\varepsilon^2 \min\{\varphi_1(\varepsilon/2, b, n, \varphi), \lambda_0/2\}/4, b + 2n + 3n^2, \omega),$$

$$\begin{aligned} \varphi_3(\varepsilon, b, \eta, n, \varphi) := & \min\{\varphi_1(\varepsilon/3, b, n, \varphi), \varphi_1(\eta(\min\{\varepsilon/3b, 2\})\varepsilon/4, b, n, \varphi), \\ & \varphi_1(\varepsilon/4, b, n, \varphi), \lambda_0/2\}, \end{aligned}$$

$$\varphi_4(\varepsilon, b, \eta, n, \omega, \varphi) := \min\{\varphi_1(\varepsilon/3, b, n, \varphi), \varphi_2(\varepsilon/3, b, n, \omega, \varphi),$$

$$\varphi_1(\eta(\min\{\varepsilon, 2\})\varepsilon/8, b, n, \varphi), \sqrt{\varphi_2(\varepsilon/4, b, n, \omega, \varphi)},$$

$$\eta(\min\{\varepsilon, 2\})\varepsilon/16b, \sqrt{\varphi_2(\varepsilon/4, b, n, \omega, \varphi)}, 1, \lambda_0/2\}.$$

Proof. Using the triangle inequality, we have

$$\frac{\|J_t x - S(t)x\|}{t} \leq \left\| \frac{x - J_t x}{t} - \frac{x - J_{\sqrt{t}}x}{\sqrt{t}} \right\| + \left\| \frac{x - J_{\sqrt{t}}x}{\sqrt{t}} - \frac{x - S(t)x}{t} \right\|$$

Then, for $t \leq 1$, we have $t \leq \sqrt{t}$ and $t/\sqrt{t} = \sqrt{t}$ so that for $\sqrt{t} \leq \Phi(\varepsilon, b, \eta, \omega, \varphi, n)$, we obtain

$$\frac{\|J_t x - S(t)x\|}{t} \leq \varepsilon$$

using Lemmas 5.10 and 5.13. \square

Remark 5.15. While the above result uses the construction of φ_1 from φ exhibited in Lemma 5.4, it is clear that if φ_1 is any other rate of convergence for $\|A_t x\|$ to $|Ax|$ as $t \rightarrow 0$, the above result nevertheless remains valid.

5.2. An analysis of Reich's result. The proof for Reich's result proceeds by establishing

$$\lim_{t \rightarrow \infty} \frac{\|J_t x\|}{t} = d(0, \text{ran}A)$$

and concluding from this that $J_t x/t$ is Cauchy for $t \rightarrow \infty$. This result is then in turn used to conclude the claim. While Reich actually establishes his result even for $x \in \overline{\text{dom}A}$, we here focus for simplicity on the case where $x \in \text{dom}A$. As mentioned in the brief outline at the beginning of this section, all the following results in the context of Reich's theorem take place for an operator A which satisfies the full range condition (*RC*) for the closure of the domain.

The main object used in these proofs is the value

$$d := d(0, \text{ran}A) = \inf\{\|y\| \mid y \in \text{ran}A\}$$

and for the quantitative results, the logical methodology implies (see the later logical remarks for a discussion of this) a dependence on a function f witnessing the above infimum quantitatively in the sense that $f : \mathbb{R}_{>0} \rightarrow \mathbb{N}$ satisfies

$$\forall \varepsilon > 0 \exists (y, z) \in A (\|y\|, \|z\| \leq f(\varepsilon) \wedge \|z\| - d \leq \varepsilon).$$

The proof of Reich's result then proceeds by a case distinction on whether $d > 0$ or $d = 0$ but, as mentioned before, this case distinction can be smoothed as will be exhibited later. We at first begin with the following result which provides a rate of convergence for the limit $\|J_t x\|/t \rightarrow d$ for $t \rightarrow \infty$ (which can be obtained as the sequence is monotone).

Lemma 5.16. *Let X be a Banach space and A be an accretive operator that satisfies the range condition. Let $x \in \text{dom}A$ with $v \in Ax$ and $b \in \mathbb{N}^*$ where $b \geq \|x\|, \|v\|$. Suppose that $f : \mathbb{R}_{>0} \rightarrow \mathbb{N}$ satisfies $f(\varepsilon) \geq f(\delta)$ for $\varepsilon \leq \delta$ and*

$$\forall \varepsilon > 0 \exists (y, z) \in A (\|y\|, \|z\| \leq f(\varepsilon) \wedge \|z\| - d \leq \varepsilon).$$

Then we have

$$\forall \varepsilon > 0 \forall t \geq \varphi(\varepsilon, b, f) \left(\left| \frac{\|J_t x\|}{t} - d \right| \leq \varepsilon \right)$$

where

$$\varphi(\varepsilon, b, f) := \frac{8(b + f(\varepsilon/2))}{\varepsilon}.$$

Proof. As $A_t x \in A J_t x$ for any $t > 0$, we have $d \leq \|A_t x\|$. Let ε be given and let $z \in Ay$ such that $\|z\| - d \leq \varepsilon/2$ and $\|y\|, \|z\| \leq f(\varepsilon/2)$. Then using Lemma 2.3, (7), we have

$$\begin{aligned} \|A_t x\| &\leq \|A_t x - A_t y\| + \|A_t y\| \\ &\leq \frac{2}{t} \|x - y\| + \|z\| \\ &\leq \frac{2(b + f(\varepsilon/2))}{t} + d + \varepsilon/2. \end{aligned}$$

Thus, for $t \geq (\varepsilon/4(b + f(\varepsilon/2)))^{-1}$, we have

$$\|A_t x\| \leq \frac{2(b + f(\varepsilon/2))}{(\varepsilon/4(b + f(\varepsilon/2)))^{-1}} + d + \varepsilon/2 \leq d + \varepsilon$$

Now, for $t \geq (\varepsilon/8(b + f(\varepsilon/2)))^{-1}$, we obtain

$$\begin{aligned} \left| \frac{\|J_t x\|}{t} - d \right| &\leq \left| \frac{\|J_t x\|}{t} - \frac{\|x - J_t x\|}{t} \right| + \left| \|A_t x\| - d \right| \\ &\leq \frac{\|x\|}{t} + \left| \|A_t x\| - d \right| \\ &\leq \varepsilon \end{aligned}$$

as $t \geq (\varepsilon/8(b + f(\varepsilon/2)))^{-1}$ and thus $\|A_t x\| - d \leq \varepsilon/2$ as well as $t \geq (\varepsilon/2b)^{-1}$ and thus $\|x\|/t \leq \varepsilon/2$. \square

The following result is a quantitative version of the well-known result due to Reich [52] that $d > 0$ implies that $\|J_t x\| \rightarrow \infty$ for $t \rightarrow \infty$ and $x \in \text{dom}A$.

Lemma 5.17. *Let X be a Banach space, A be an accretive operator that satisfies the range condition and assume that $d \geq D$ for $D \in \mathbb{R}_{>0}$. Let $x \in \text{dom}A$ with $v \in Ax$ and $b \in \mathbb{N}^*$ where $b \geq \|x\|, \|v\|$. Then we have*

$$\forall K \in \mathbb{N} \forall t \geq \psi(K, b, D) (\|J_t x\| \geq K)$$

where

$$\psi(K, b, D) := \frac{b + K}{D}.$$

Proof. Suppose the claim is false, i.e. there is a K and a $t \geq \psi(K, b, D)$ such that $\|J_t x\| < K$. Then, we have

$$\begin{aligned} \|J_t x - J_1 J_t x\| &\leq |A J_t x| \\ &\leq \|x - J_t x\| / t \\ &< (b + K) / D^{-1} (b + K) \\ &\leq D. \end{aligned}$$

Thus $\|A_1 J_t x\| < D \leq d$ which is a contradiction as $A_1 J_t x \in \text{ran}A$. \square

Lemma 5.18 (essentially Reich [53]). *Let X be uniformly convex with a modulus of uniform convexity η . Then, for $\varepsilon \in (0, 2]$, we have $2\eta(\varepsilon) \leq 1 - \langle y, j \rangle$ for all $j \in Jx$ with $\|x\| = \|y\| = 1$ and $\|x - y\| \geq \varepsilon$.*

Proof. Let x, y and $j \in Jx$ be given with $\|x\| = \|y\| = 1$ and $\|x - y\| \geq \varepsilon$. Then

$$\frac{\|x + y\|}{2} \leq 1 - \eta(\varepsilon)$$

by definition of η . Thus as $\langle x, j \rangle = \|x\|^2 = 1$ and $\|j\| = \|x\| = 1$, we have

$$1/2 + 1/2 \langle y, j \rangle = \langle (x + y)/2, j \rangle \leq 1 - \eta(\varepsilon)$$

which yields the claim. \square

Lemma 5.19. *Let X be a uniformly convex Banach space with a modulus of uniform convexity $\eta : (0, 2] \rightarrow (0, 1]$. Suppose that $f : \mathbb{R}_{>0} \rightarrow \mathbb{N}$ satisfies*

$$\forall \varepsilon > 0 \exists (y, z) \in A (\|y\|, \|z\| \leq f(\varepsilon) \wedge \|z\| - d \leq \varepsilon).$$

Let $\varepsilon > 0$ be given, assume that $D \in \mathbb{R}_{>0}$ with $d \geq D$ and let $z \in Ay$ be such that

$$\|z\| \leq d + 2d\eta(\varepsilon/2)$$

as well as $c \geq \|y\|, \|z\|$. Let $x \in \text{dom}A$ with $v \in Ax$ and $b \in \mathbb{N}^*$ with $b \geq \|x\|, \|v\|$. Then, for any $t \geq \varphi_1(\varepsilon, b, D, c, \eta, f)$:

$$\left\| \frac{z}{\|z\|} + \frac{J_t x}{\|J_t x\|} \right\| \leq \varepsilon$$

where

$$\begin{aligned} \varphi_1(\varepsilon, b, D, c, \eta, f) := \max \left\{ \psi(c + 1, b, D), \psi \left(\left(\frac{4}{\varepsilon} + 1 \right) c, b, D \right), \right. \\ \left. \left(\frac{D (2\eta(\min\{\varepsilon/2, 2\}))^2}{18} \right)^{-1} (c + b), \right. \\ \left. \varphi \left(\frac{D (2\eta(\min\{\varepsilon/2, 2\}))^2}{18}, b, f \right) \right\} \end{aligned}$$

with φ defined as in Lemma 5.16 and ψ as in Lemma 5.17

Proof. As $A_t x \in AJ_t x$ and as A is accretive, there is a $j_t \in J(y - J_t x)$ such that $\langle z - A_t x, j_t \rangle \geq 0$. Therefore we have

$$\left\langle \frac{z}{\|z\|}, \frac{j_t}{\|y - J_t x\|} \right\rangle \geq \left\langle \frac{A_t x}{\|z\|}, \frac{j_t}{\|y - J_t x\|} \right\rangle$$

for any t such that $t \geq \psi(c + 1, b, D)$ as then $\|J_t x\| > c$. Then further

$$\left\langle J_t x - x, \frac{j_t}{\|y - J_t x\|} \right\rangle \leq \|y - x\| - \|y - J_t x\|,$$

and we thus obtain

$$\begin{aligned} \left\langle A_t x, \frac{j_t}{\|y - J_t x\|} \right\rangle &\geq \left\| \frac{y}{t} - \frac{J_t x}{t} \right\| - \frac{\|y - x\|}{t} \\ &\geq \frac{\|J_t x\|}{t} - \frac{\|y\|}{t} - \frac{\|y - x\|}{t}. \end{aligned}$$

Thus for any $\delta \in \mathbb{R}_{>0}$ and any

$$t \geq \max \{ (\delta/3)^{-1}(c + b), \varphi(\delta/3, b, f) \},$$

we obtain

$$\begin{aligned} \left\langle \frac{z}{\|z\|}, \frac{j_t}{\|y - J_t x\|} \right\rangle &\geq \frac{d}{\|z\|} - \frac{\delta}{\|z\|} \\ &\geq \frac{1}{1 + 2\eta(\min\{\varepsilon/2, 2\})} - \frac{\delta}{\|z\|}. \end{aligned}$$

Now, as η is positive, we get $1 - (2\eta(\min\{\varepsilon/2, 2\}))^2 + (2\eta(\min\{\varepsilon/2, 2\}))^2 = 1$ and therefore $1 = (1 + 2\eta(\min\{\varepsilon/2, 2\}))(1 - 2\eta(\min\{\varepsilon/2, 2\})) + (2\eta(\min\{\varepsilon/2, 2\}))^2$ which yields

$$\begin{aligned} \frac{1}{1 + 2\eta(\min\{\varepsilon/2, 2\})} &= 1 - 2\eta(\min\{\varepsilon/2, 2\}) + \frac{(2\eta(\min\{\varepsilon/2, 2\}))^2}{1 + 2\eta(\min\{\varepsilon/2, 2\})} \\ &\geq 1 - 2\eta(\min\{\varepsilon/2, 2\}) + \frac{(2\eta(\min\{\varepsilon/2, 2\}))^2}{3}. \end{aligned}$$

Thus for

$$t \geq \max \left\{ \left(\frac{D(2\eta(\min\{\varepsilon/2, 2\}))^2}{18} \right)^{-1} (c + b), \varphi \left(\frac{D(2\eta(\min\{\varepsilon/2, 2\}))^2}{18}, b, f \right) \right\},$$

we obtain (using $z \in \text{ran}A$) that

$$\begin{aligned} \left\langle \frac{z}{\|z\|}, \frac{j_t}{\|y - J_t x\|} \right\rangle &\geq 1 - 2\eta(\min\{\varepsilon/2, 2\}) + \frac{(2\eta(\min\{\varepsilon/2, 2\}))^2}{3} \\ &\quad - \frac{D(2\eta(\min\{\varepsilon/2, 2\}))^2}{6\|z\|} \\ &\geq 1 - 2\eta(\min\{\varepsilon/2, 2\}) + \frac{(2\eta(\min\{\varepsilon/2, 2\}))^2}{6} \\ &> 1 - 2\eta(\min\{\varepsilon/2, 2\}). \end{aligned}$$

Then in particular

$$\left\| \frac{z}{\|z\|} - \frac{y - J_t x}{\|y - J_t x\|} \right\| \leq \frac{\varepsilon}{2}$$

by Lemma 5.18 for all such t .

Now, secondly:

$$\left\| \frac{y - J_t x}{\|y - J_t x\|} + \frac{J_t x}{\|J_t x\|} \right\| \leq \frac{\|y\|}{\|y\| - \|J_t x\|} + \left| 1 - \frac{\|J_t x\|}{\|y - J_t x\|} \right|$$

For $\delta > 0$ and $t \geq \psi((\delta^{-1} + 1)c, b, D)$, we immediately have

$$\frac{\|y\|}{\|y\| - \|J_t x\|} \leq \delta.$$

Similarly, we get for $t \geq \psi((\delta^{-1} + 1)c, b, D)$, as $(\delta^{-1} + 1)c \geq (\delta^{-1} - 1)c$, that

$$\frac{\|y\|}{\|y\| + \|J_t x\|} \leq \delta.$$

Further, we have

$$1 - \frac{\|y\|}{\|y\| + \|J_t x\|} \leq \frac{\|J_t x\|}{\|y - J_t x\|} \leq 1 + \frac{\|y\|}{\|y\| - \|J_t x\|}$$

and thus for $t \geq \psi((\delta^{-1} + 1)c, b, D)$, we get

$$\left| 1 - \frac{\|J_t x\|}{\|y - J_t x\|} \right| \leq \delta.$$

Combining the above, we have that for any $t \geq \psi((\varepsilon/4)^{-1} + 1)c, b, D$:

$$\left\| \frac{y - J_t x}{\|y - J_t x\|} + \frac{J_t x}{\|J_t x\|} \right\| \leq \frac{\varepsilon}{2}.$$

Thus, finally for $t \geq \varphi_1(\varepsilon, b, D, c, \eta, f)$ we obtain the desired result. \square

Lemma 5.20. *Let X be a uniformly convex Banach space with a modulus of uniform convexity $\eta : (0, 2] \rightarrow (0, 1]$. Suppose that $f : \mathbb{R}_{>0} \rightarrow \mathbb{N}$ satisfies $f(\varepsilon) \geq f(\delta)$ for $\varepsilon \leq \delta$ and*

$$\forall \varepsilon > 0 \exists (y, z) \in A(\|y\|, \|z\| \leq f(\varepsilon) \wedge \|z\| - d \leq \varepsilon).$$

Let $\varepsilon > 0$ be given, assume that $E \in \mathbb{N}^$, $D \in \mathbb{R}_{>0}$ where $E \geq d \geq D$. Let $x \in \text{dom}A$ with $v \in Ax$ and $b \in \mathbb{N}^*$ where $b \geq \|x\|, \|v\|$. Then, for any $t, s \geq \varphi'_2(\varepsilon, b, D, \eta, E, f)$:*

$$\left\| \frac{J_s x}{s} - \frac{J_t x}{t} \right\| \leq \varepsilon$$

where

$$\varphi'_2(\varepsilon, b, D, \eta, E, f) := \max\{\varphi(\varepsilon/3, b, f), \varphi_1(\varepsilon/6E, b, D, f(2D\eta(\min\{\varepsilon/12E, 2\})), \eta, f)\}$$

with φ as in Lemma 5.16 and φ_1 as in Lemma 5.19.

Proof. We have that there exists $z \in Ay$ such that $\|z\| \leq d + 2d\eta(\min\{\varepsilon/4, 2\})$ with $\|y\|, \|z\| \leq f(2D\eta(\min\{\varepsilon/4, 2\}))$. Thus, using Lemma 5.19, we have that for $t, s \geq \varphi_1(\varepsilon/2, b, D, f(2D\eta(\min\{\varepsilon/4, 2\})), \eta, f)$:

$$\left\| \frac{J_s x}{\|J_s x\|} - \frac{J_t x}{\|J_t x\|} \right\| \leq \left\| \frac{z}{\|z\|} + \frac{J_s x}{\|J_s x\|} \right\| + \left\| -\frac{z}{\|z\|} - \frac{J_t x}{\|J_t x\|} \right\| \leq \varepsilon.$$

Therefore, we in particular have that

$$\begin{aligned} \left\| \frac{J_s x}{s} - \frac{J_t x}{t} \right\| &= \left\| \frac{J_s x}{\|J_s x\|} \frac{\|J_s x\|}{s} - \frac{J_t x}{\|J_t x\|} \frac{\|J_t x\|}{t} \right\| \\ &\leq \left\| \frac{J_s x}{\|J_s x\|} \frac{\|J_s x\|}{s} - \frac{J_s x}{\|J_s x\|} d \right\| + \left\| \frac{J_s x}{\|J_s x\|} d - \frac{J_t x}{\|J_t x\|} d \right\| \\ &\quad + \left\| \frac{J_t x}{\|J_t x\|} d - \frac{J_t x}{\|J_t x\|} \frac{\|J_t x\|}{t} \right\| \\ &\leq \left| \frac{\|J_s x\|}{s} - d \right| + d \left\| \frac{J_s x}{\|J_s x\|} - \frac{J_t x}{\|J_t x\|} \right\| + \left| d - \frac{\|J_t x\|}{t} \right|. \end{aligned}$$

Thus, for $t, s \geq \varphi'_2(\varepsilon, b, D, \eta, E, f)$, we get the claim by Lemma 5.16 together with the above. \square

This result, which presents the quantitative version of the case of $d > 0$ for the result of Reich, can now be smoothed to omit this assumption. For this, note that through the trivial proof of the case of $d = 0$, one obtains the following quantitative version of this result:

Lemma 5.21. *Let X be a uniformly convex Banach space with a modulus of uniform convexity $\eta : (0, 2] \rightarrow (0, 1]$. Suppose that $f : \mathbb{R}_{>0} \rightarrow \mathbb{N}$ satisfies $f(\varepsilon) \geq f(\delta)$ for $\varepsilon \leq \delta$ and*

$$\forall \varepsilon > 0 \exists (y, z) \in A(\|y\|, \|z\| \leq f(\varepsilon) \wedge \|z\| - d \leq \varepsilon).$$

Let $\varepsilon > 0$ be given, assume that $E \in \mathbb{N}^$ where $E \geq d$. Let $x \in \text{dom}A$ with $v \in Ax$ and $b \in \mathbb{N}^*$ where $b \geq \|x\|, \|v\|$. Then, for any $t, s \geq \varphi_2(\varepsilon, b, \eta, E, f)$:*

$$\left\| \frac{J_s x}{s} - \frac{J_t x}{t} \right\| \leq \varepsilon$$

where

$$\varphi_2(\varepsilon, b, \eta, E, f) := \max\{\varphi(\varepsilon/4, b, f), \varphi(\varepsilon/3, b, f), \\ \varphi_1(\varepsilon/6E, b, \varepsilon/4, f(\varepsilon\eta(\min\{\varepsilon/12E, 2\})/2), \eta, f)\}$$

with φ as in Lemma 5.16 and φ_1 as in Lemma 5.19.

Proof. Suppose that $d \leq \varepsilon/4$. Then by Lemma 5.16, we have that

$$\left| \frac{\|J_t x\|}{t} - d \right| \leq \varepsilon/4$$

for any $t \geq \varphi(\varepsilon/4, b, f)$. Thus in particular we have that $\|J_t x\|/t \leq \varepsilon/2$ for all such t and thus

$$\left\| \frac{J_s x}{s} - \frac{J_t x}{t} \right\| \leq \left\| \frac{J_s x}{s} \right\| + \left\| \frac{J_t x}{t} \right\| \leq \varepsilon$$

for all $t, s \geq \varphi(\varepsilon/4, b, f)$ in that case. Otherwise $d \geq \varepsilon/4$ and thus the above result holds for $t, s \geq \varphi'_2(\varepsilon, b, \varepsilon/4, \eta, E, f)$. \square

The rest of the proof given in [53] now relies on the use of the limit $-v_x$ of $J_t x/t$ for $t \rightarrow \infty$. By the above Lemma, this limit exists as X is complete. While we emphasized that this limit a priori depends on the starting point x , the following lemma (which provides a concrete quantitative version of Lemma 3.2 given in [53]) shows that this limit is actually unique, i.e all the v_x coincide.

Lemma 5.22. *Let X be a uniformly convex Banach space with a modulus of uniform convexity $\eta : (0, 2] \rightarrow (0, 1]$. Suppose that $f : \mathbb{R}_{>0} \rightarrow \mathbb{N}$ satisfies $f(\varepsilon) \geq f(\delta)$ for $\varepsilon \leq \delta$ and*

$$\forall \varepsilon > 0 \exists (y, z) \in A(\|y\|, \|z\| \leq f(\varepsilon) \wedge \|z\| - d \leq \varepsilon).$$

Let $\varepsilon > 0$ be given, assume that $E \in \mathbb{N}^$ and $D \in \mathbb{R}_{>0}$ where $E \geq d \geq D$ and let $z \in Ay$ be such that*

$$\|z\| \leq d + 2d\eta\left(\frac{\varepsilon}{16(E+1)}\right).$$

If $x \in \text{dom}A$, then $\|z - v_x\| \leq \varepsilon$.

Proof. We write $\delta_\varepsilon = 2d\eta(\varepsilon/16(E+1))$. Then, for $\|z\| \leq d + \delta_\varepsilon$, we have

$$\begin{aligned} \left\| z + \frac{J_t x}{t} \right\| &= \left\| z + \frac{J_t x}{\|J_t x\|} \frac{\|J_t x\|}{t} \right\| \\ &\leq \left\| z - \frac{d}{\|z\|} z \right\| + \left\| \frac{d}{\|z\|} z + \frac{J_t x}{\|J_t x\|} \frac{\|J_t x\|}{t} \right\| \\ &\leq \|z\| - d + \left\| \frac{d}{\|z\|} z + \frac{J_t x}{\|J_t x\|} \frac{\|J_t x\|}{t} \right\| \\ &\leq \delta_\varepsilon + \left\| \frac{d}{\|z\|} z + \frac{J_t x}{\|J_t x\|} \frac{\|J_t x\|}{t} \right\|. \end{aligned}$$

Similar to before, we have

$$\begin{aligned} \left\| \frac{d}{\|z\|} z + \frac{J_t x}{\|J_t x\|} \frac{\|J_t x\|}{t} \right\| &\leq \left\| \frac{d}{\|z\|} z - \frac{z}{\|z\|} \frac{\|J_t x\|}{t} \right\| + \left\| \frac{z}{\|z\|} \frac{\|J_t x\|}{t} + \frac{J_t x}{\|J_t x\|} \frac{\|J_t x\|}{t} \right\| \\ &= \left| d - \frac{\|J_t x\|}{t} \right| + \frac{\|J_t x\|}{t} \left\| \frac{z}{\|z\|} + \frac{J_t x}{\|J_t x\|} \right\|. \end{aligned}$$

From this we obtain that

$$\left\| z + \frac{J_t x}{t} \right\| \leq \delta_\varepsilon + \frac{\varepsilon}{4}$$

for all

$$t \geq \max\{\varphi(\varepsilon/8, b, f), \varphi_1(\varepsilon/8(E+1), b, D, c, \eta, f)\}$$

where $c, b \in \mathbb{N}^*$ are such that $c \geq \|y\|, \|z\|$ and $b \geq \|x\|, \|v\|$ for $v \in Ax$ as, for one, $t \geq \varphi(\varepsilon/8, b, f)$ and thus

$$\left| d - \frac{\|J_t x\|}{t} \right| \leq \varepsilon/8$$

by Lemma 5.16 as well as $\frac{\|J_t x\|}{t} \leq d + 1 \leq E + 1$ and, for another, $t \geq \varphi_1(\varepsilon/8(E + 1), b, D, c, \eta, f)$ and thus

$$\frac{\|J_t x\|}{t} \left\| \frac{z}{\|z\|} + \frac{J_t x}{\|J_t x\|} \right\| \leq \varepsilon/8.$$

by Lemma 5.19. Then the properties of η imply that $\delta_\varepsilon \leq \varepsilon/4$ and thus

$$\left\| z + \frac{J_t x}{t} \right\| \leq \varepsilon/2$$

for all such t . Then

$$\|z - v_x\| \leq \left\| z + \frac{J_t x}{t} \right\| + \left\| v_x + \frac{J_t x}{t} \right\|$$

for all t and thus choosing

$$t = \max\{\varphi(\varepsilon/8, b, f), \varphi_1(\varepsilon/8(E + 1), b, D, c, \eta, f), \varphi_2(\varepsilon/2, b, \eta, E, f)\}$$

implies $\|z - v_x\| \leq \varepsilon$ by definition of v_x (which yields that φ_2 is a rate of convergence for $J_t x/t$ towards $-v_x$) and Lemma 5.21. \square

Lemma 5.23. *Let X be a uniformly convex Banach space with a modulus of uniform convexity $\eta : (0, 2] \rightarrow (0, 1]$. Suppose that $f : \mathbb{R}_{>0} \rightarrow \mathbb{N}$ satisfies $f(\varepsilon) \geq f(\delta)$ for $\varepsilon \leq \delta$ and*

$$\forall \varepsilon > 0 \exists (y, z) \in A(\|y\|, \|z\| \leq f(\varepsilon) \wedge \|z\| - d \leq \varepsilon).$$

Let $\varepsilon > 0$ be given, assume that $E \in \mathbb{N}^$ where $E \geq d$ and let $z \in Ay$ be such that*

$$\|z\| \leq d + \min \left\{ \varepsilon \eta \left(\min \left\{ \frac{\varepsilon}{16(E + 1)}, 2 \right\} \right) / 4, \varepsilon/8 \right\}.$$

If $x \in \text{dom}A$, then $\|z - v_x\| \leq \varepsilon$.

Proof. Let ε be given. Then either $d \leq \varepsilon/8$ which, since $\|z\| \leq d + \varepsilon/8$, implies $\|z\| \leq \varepsilon/4$. For

$$t \geq \max\{\varphi(\varepsilon/4, b, f), \varphi_2(\varepsilon/4, b, \eta, E, f)\},$$

we then get

$$\begin{aligned} \|z - v_x\| &\leq \|z\| + \|v_x\| \\ &\leq \|z\| + d + \left| d - \frac{\|J_t x\|}{t} \right| + \left\| v_x + \frac{J_t x}{t} \right\| \\ &\leq \varepsilon. \end{aligned}$$

Otherwise we have $d \geq \varepsilon/8$ and thus, we get the same result for

$$\|z\| \leq d + \varepsilon \eta \left(\min \left\{ \frac{\varepsilon}{16(E + 1)}, 2 \right\} \right) / 4 \leq d + 2d\eta \left(\min \left\{ \frac{\varepsilon}{16(E + 1)}, 2 \right\} \right).$$

\square

Theorem 5.24. *Let X be a uniformly convex Banach space with a modulus of uniform convexity $\eta : (0, 2] \rightarrow (0, 1]$. Suppose that $f : \mathbb{R}_{>0} \rightarrow \mathbb{N}$ satisfies $f(\varepsilon) \geq f(\delta)$ for $\varepsilon \leq \delta$ and*

$$\forall \varepsilon > 0 \exists (y, z) \in A(\|y\|, \|z\| \leq f(\varepsilon) \wedge \|z\| - d \leq \varepsilon).$$

Assume that $E \in \mathbb{N}^$ where $E \geq d$ and further that $x \in \text{dom}A$ with $v \in Ax$ and $b \in \mathbb{N}^*$ with $b \geq \|x\|, \|v\|$. Then*

$$\forall \varepsilon > 0 \forall t \geq \Phi(\varepsilon, b, \eta, E, f) \left(\frac{\|J_t^A x - S(t)x\|}{t} \leq \varepsilon \right)$$

where

$$\begin{aligned} \Phi(\varepsilon, b, \eta, E, f) := \max &\left\{ \frac{4}{\varepsilon} \left(b + f \left(\min \left\{ \varepsilon \eta \left(\min \left\{ \frac{\varepsilon/8}{16(E + 1)}, 2 \right\} \right) / 32, \varepsilon/64 \right\} \right) \right), \right. \\ &\frac{8}{\varepsilon} f \left(\min \left\{ \varepsilon \eta \left(\min \left\{ \frac{\varepsilon/8}{16(E + 1)}, 2 \right\} \right) / 32, \varepsilon/64 \right\} \right), \\ &\left. \varphi_2(\varepsilon/2, b, \eta, E, f) \right\} \end{aligned}$$

with

$$\begin{aligned}\varphi(\varepsilon, b, f) &:= \frac{8(b + f(\varepsilon/2))}{\varepsilon}, \\ \psi(K, b, D) &:= \frac{b + K}{D}, \\ \varphi_1(\varepsilon, b, D, c, \eta, f) &:= \max \left\{ \psi(c + 1, b, D), \psi \left(\left(\frac{4}{\varepsilon} + 1 \right) c, b, D \right), \right. \\ &\quad \left(\frac{D (2\eta(\min\{\varepsilon/2, 2\}))^2}{18} \right)^{-1} (c + b), \\ &\quad \left. \varphi \left(\frac{D (2\eta(\min\{\varepsilon/2, 2\}))^2}{18}, b, f \right) \right\}, \\ \varphi_2(\varepsilon, b, \eta, E, f) &:= \max \{ \varphi(\varepsilon/4, b, f), \varphi(\varepsilon/3, b, f), \\ &\quad \varphi_1(\varepsilon/6E, b, \varepsilon/4, f(\varepsilon\eta(\min\{\varepsilon/12E, 2\})/2), \eta, f) \}.\end{aligned}$$

Proof. Given ε , there are $z \in Ay$ such that

$$\|z\| \leq d + \min \left\{ \varepsilon\eta \left(\min \left\{ \frac{\varepsilon/8}{16(E+1)}, 2 \right\} \right) / 32, \varepsilon/64 \right\}$$

and such that $\|y\|, \|z\| \leq f \left(\min \left\{ \varepsilon\eta \left(\min \left\{ \frac{\varepsilon/8}{16(E+1)}, 2 \right\} \right) / 32, \varepsilon/64 \right\} \right)$. Now, we in particular have $\|A_r J_r a\| \leq \|A J_r a\| \leq \|A_r a\|$ and thus $\|A_{t/n} J_{t/n}^i y\| \leq \|A_{t/n} J_{t/n}^{i-1} y\|$. Iterating this gives

$$\|A_{t/n} J_{t/n}^i y\| \leq \|A_{t/n} y\| \leq \|z\| \leq d + \min \left\{ \varepsilon\eta \left(\min \left\{ \frac{\varepsilon/8}{16(E+1)}, 2 \right\} \right) / 32, \varepsilon/64 \right\}$$

for all $i \in [0; n-1]$. Now we get

$$\|A_{t/n} J_{t/n}^i y - v_x\| \leq \varepsilon/8$$

for any $i \in [0; n-1]$ by Lemma 5.23 which implies $\|(y - J_{t/n}^n y)/t - v_x\| \leq \varepsilon/8$ for all t and all n and thus

$$\left\| \frac{y - S(t)y}{t} - v_x \right\| \leq \varepsilon/8$$

for all t .

Then in particular

$$\left\| \frac{S(t)y}{t} + v_x \right\| \leq \frac{\|y\|}{t} + \left\| \frac{y - S(t)y}{t} - v_x \right\| \leq \frac{\|y\|}{t} + \varepsilon/8$$

for all t . In particular, for

$$t \geq (\varepsilon/8)^{-1} f \left(\min \left\{ \varepsilon\eta \left(\min \left\{ \frac{\varepsilon/8}{16(E+1)}, 2 \right\} \right) / 32, \varepsilon/64 \right\} \right),$$

we have

$$\left\| \frac{S(t)y}{t} + v_x \right\| \leq \varepsilon/4.$$

Continuing, we obtain

$$\left\| \frac{S(t)x}{t} + v_x \right\| \leq \left\| \frac{S(t)y}{t} + v_x \right\| + \frac{\|x - y\|}{t}$$

which implies

$$\left\| \frac{S(t)x}{t} + v_x \right\| \leq \varepsilon/2$$

for all

$$t \geq \max \left\{ (\varepsilon/8)^{-1} f \left(\min \left\{ \varepsilon \eta \left(\min \left\{ \frac{\varepsilon/8}{16(E+1)}, 2 \right\} \right) / 32, \varepsilon/64 \right\} \right), \right. \\ \left. (\varepsilon/4)^{-1} \left(b + f \left(\min \left\{ \varepsilon \eta \left(\min \left\{ \frac{\varepsilon/8}{16(E+1)}, 2 \right\} \right) / 32, \varepsilon/64 \right\} \right) \right) \right\}.$$

Finally, we get

$$\left\| \frac{S(t)x - J_t x}{t} \right\| \leq \left\| \frac{S(t)x}{t} + v_x \right\| + \left\| v_x + \frac{J_t x}{t} \right\|$$

and thus

$$\left\| \frac{S(t)x - J_t x}{t} \right\| \leq \varepsilon$$

for all $t \geq \Phi(\varepsilon, b, \eta, E, f)$ by Lemma 5.21 and the definition of v_x (which yields that φ_2 is a rate of convergence as before). \square

5.3. Logical remarks on the above results. Lastly, we want to outline the additional modifications to H_p^ω necessary to formalize the proofs of the theorems of Plant and Reich. These modifications in that way give rise to the systems and bound extraction results underlying the extractions outline above. In that context, we here in particular move away from the use of arbitrary real errors ε and again consider representations of errors via natural numbers through 2^{-k} .

At first, both results are formulated for points $x \in \text{dom}A$ and by the logical methodology, this stands for the existential assumption $\exists y(y \in Ax)$ which yields that at least a priori the extracted rates will in particular depend on an upper bound on the norm of this witness which is also the case for the above rates.

The second prominent assumption in both results is that of uniform convexity which was quantitatively treated above via the modulus of uniform convexity η . Formally, this can be achieved by adding an additional constant η of type $\mathbb{N} \rightarrow \mathbb{N}$ together with a corresponding axiom stating that it represents a modulus of uniform convexity for X (see [26] for more details):

$$\forall x^X, y^X, k^{\mathbb{N}} \left(\|x\|_X, \|y\|_X <_{\mathbb{R}} 1 \wedge \left\| \frac{x +_X y}{2} \right\|_X >_{\mathbb{R}} 1 - 2^{-\eta(k)} \rightarrow \|x -_X y\|_X \leq_{\mathbb{R}} 2^{-k} \right).$$

To really formally encapsulate the previous proof where η was applied to various reals, one would now have to extend η to \mathbb{Q}_+^* via

$$\eta(\varepsilon) := 2^{-\eta(\min k[2^{-k} \leq \varepsilon])}$$

for $\varepsilon \in \mathbb{Q}_+^*$ and move to rational approximations of the reals in questions. We avoid spelling this out any further.

We now first focus on the theorem of Plant. The main object featuring in Plant's proof (and consequently in the above results as well) is the limit functional $|Ax|$. The use of this functional can now be emulated in the context of H_p^ω by extending the underlying language with a further constant of type $X \rightarrow \mathbb{N}^{\mathbb{N}}$ which we denote by $|A \cdot |$ (where we correspondingly denote $|A \cdot |x$ by $|Ax|$ for simplicity). One first natural axiom for this constant is induced by the natural bound on $|Ax|$ by $\|v\|$:

$$(M1) \quad \forall x^X, v^X, \lambda^{\mathbb{N}^{\mathbb{N}}} (v \in Ax \wedge 0 <_{\mathbb{R}} \lambda <_{\mathbb{R}} \lambda_0 \rightarrow \|A_\lambda x\|_X \leq_{\mathbb{R}} |Ax| \leq_{\mathbb{R}} \|v\|_X).$$

As shortly mentioned in the above quantitative results, the convergence of $\|A_\lambda x\|$ to $|Ax|$ for $x \in \text{dom}A$ as $\lambda \rightarrow 0$ is 'equivalent' to the lower semi-continuity of $|Ax|$ on $\text{dom}A$. This vague 'equivalence' can now be made precise through the system $H_p^\omega + (M1)$ in the following sense:

Proposition 5.25. *Over $H_p^\omega + (M1)$, the following are equivalent:*

(1) $\|A_t x\| \rightarrow |Ax|$ while $t \rightarrow 0^+$ for all $x \in \text{dom}A$, i.e.

$$\forall x^X, k^{\mathbb{N}} \exists n^{\mathbb{N}} (x \in \text{dom}A \rightarrow |Ax| - \|A_{2^{-n}} x\|_X \leq_{\mathbb{R}} 2^{-k});$$

(2) lower semi-continuity for $|Ax|$ for all $x \in \text{dom}A$, i.e.

$$\forall k^{\mathbb{N}}, x^X \exists m^{\mathbb{N}} \forall y^X (x \in \text{dom}A \wedge y \in \text{dom}A \wedge \|x -_X y\|_X \leq_{\mathbb{R}} 2^{-m} \rightarrow |Ax| - |Ay| \leq_{\mathbb{R}} 2^{-k}).$$

Proof. From (1) to (2), let $x \in \text{dom}A$ and k be given. For $y \in \text{dom}A$, we have

$$\begin{aligned} |Ax| - |Ay| &\leq |Ax| - \|A_\lambda y\| \\ &\leq |Ax| - \|A_\lambda x\| + |\|A_\lambda x\| - \|A_\lambda y\|| \\ &\leq |Ax| - \|A_\lambda x\| + 2/\lambda \|x - y\| \end{aligned}$$

for any $\lambda \in (0, \lambda_0)$. Now, using (1) we pick n such that $|Ax| - \|A_{2^{-n}}x\| \leq 2^{-(k+1)}$ and then pick $m = n + k + 2$ such that $|Ax| - |Ay| \leq 2^{-k}$.

From (2) to (1), let $x \in \text{dom}A$ and k be given. Using (2), we pick an m such that $|Ax| - |Ay| \leq 2^{-k}$ for all $y \in \text{dom}A$ such that $\|x - y\| \leq 2^{-m}$. Now, for $n = \lceil \log_2 b + m \rceil$ for $b \geq \|v\|$ for some $v \in Ax$, we then get

$$\|x - J_{2^{-n}}x\| \leq 2^{-n} \|v\| \leq 2^{-m}$$

which in particular implies

$$|Ax| - \|A_{2^{-n}}x\| \leq |Ax| - |AJ_{2^{-n}}x| \leq 2^{-k}$$

using $A_{2^{-n}}x \in AJ_{2^{-n}}x$. □

In that way, the convergence of $\|A_\lambda x\|$ to $|Ax|$ on $\text{dom}A$ relates to an extensionality principle of $|A \cdot |$. Now, in the context of set-valued operators, these continuity and extensionality principles can be logically complicated and their study actually gives rise to a hierarchy of fragments of extensionality with a corresponding hierarchy of continuity principles with various intricacies (see the forthcoming [48]). In any way, as in the case of the functional $\langle \cdot, \cdot \rangle_s$, the logical methodology based on the monotone Dialectica interpretation now implies the following quantitative version of the statement of item (2): under this interpretation, the statement (2) is upgraded to the existence of a ‘modulus of uniform lower-semicontinuity’ which, as with $\langle \cdot, \cdot \rangle_s$, by the uniformity on x induced by majorization, is essentially a modulus of uniform continuity. Concretely, this uniformized version of item (2) can be formally hardwired into the system by extending it with an additional constant φ of type $\mathbb{N} \rightarrow (\mathbb{N} \rightarrow \mathbb{N})$ together with the axiom

$$(M2) \quad \forall b^{\mathbb{N}}, k^{\mathbb{N}}, x^X, y^X, u^X, v^X \left(\left(u \in Ax \wedge v \in Ay \wedge \|x\|_X, \|y\|_X, \|u\|_X, \|v\|_X <_{\mathbb{R}} b \right. \right. \\ \left. \left. \wedge \|x -_X y\|_X <_{\mathbb{R}} 2^{-\varphi(k,b)} \right) \rightarrow |Ax| - |Ay| \leq_{\mathbb{R}} 2^{-k} \right).$$

Under this extension, Lemma 5.4 is then the natural extraction of a corresponding rate of convergence from the above equivalence proof, under this (therefore) necessary assumption of a modulus of uniform continuity for $|A \cdot |$, following the previous metatheorems. Note however that these metatheorems in general, through this treatment of $|A \cdot |$, imply a dependence of the extracted bounds on a majorant for the constant $|A \cdot |$, i.e. on a function $f : \mathbb{N} \rightarrow \mathbb{N}$ such that

$$\|x\| \leq b \rightarrow |Ax| \leq f(b) \text{ for all } x \in \text{dom}A.$$

Only under this additional dependence on a majorant for $|A \cdot |$ do the previous metatheorems contained in Theorems 4.6 and 4.7 extend to $H_p^\omega + (M1) + (M2)$.

Here, we shortly want to make a note on the strength of the existence of such a majorant. For this, we first remind on the notion of a majorizable operator introduced in [47] (see also the previous Remark 4.5): an operator $A : X \rightarrow 2^X$ is called majorizable if there exists a function $f : \mathbb{N} \rightarrow \mathbb{N}$ such that

$$\forall x \in \text{dom}A, b \in \mathbb{N} (\|x\| \leq b \rightarrow \exists y \in Ax (\|y\| \leq f(b))).$$

As discussed in [47], there are non-majorizable operators and so the assumption that A is majorizable is a proper restriction. In particular, note now that if A is such that the minimal selection A° exists, then $|Ax| = \|A^\circ x\|$ and thus majorizability of A is equivalent to majorizability of $|A \cdot |$. Thus, while in general potentially a bit weaker, the assumption of majorizability of $|A \cdot |$ in particular also seems to carry additional strength similar to that of majorizability of A in most cases.

However, as apparent from the result in Lemma 5.4, such a majorant however does not feature in the extracted bounds and we actually find that such a majorant also does not feature in any of the other quantitative results in the context of Plant’s theorem. While this seems to be a particular coincidence in the context of Lemma 5.4, there is actually a logical reason which guarantees this ‘non-dependence’ a priori for all the other results. Concretely, the reason is that all the other proofs analyzed have the two crucial properties that, for one, they can be formalized under the assumption of a rate of convergence for $\|A_\lambda x\|$ toward $|Ax|$ which can similarly be added to the system and that, for another, they are ‘pointwise’ results in x in the sense that they do not require

knowledge of $|A \cdot |$ for any point other than x . In that way, instead of following the above route of formalizing the whole functional $|A \cdot |$, one can add two constants representing this ‘particular’ x and a witness $v \in Ax$ as well as a constant $|Ax|$ of type $\mathbb{N}^{\mathbb{N}}$ for this single value of $|A \cdot |$ at the constant x and a constant φ representing a rate of convergence for $\|A_\lambda x\|$ to $|Ax|$ for this single constant x . Then, the other proofs still formalize and in particular depend only on majorants for φ , x , v and $|Ax|$ and the one for the latter three can be assumed to coincide and to be represented in the above results by b . In particular, the strong assumption of majorizability of $|A \cdot |$ can be avoided a priori in that way. That the extracted rates are true for all x then is drawn as a conclusion on the metalevel as the additional constants were generic. In this way, this also provides a logical insight on why all the other results in the context of Plant’s theorem remain true if φ_1 represents any other rate of convergence besides the one constructed from the modulus of uniform continuity for $|A \cdot |$ as commented on before.

As a last comment on the logical particularities of the proofs towards Plant’s theorem, we want to note in the context of Miyadera’s lemma from [6] that the only properties of $\langle \cdot, \cdot \rangle_s$ required in the proof given in [6] are the properties discussed in Section 3.3 and the fact that the proof given by Crandall in [10] actually only invokes this result for $x \in \text{dom}A$. Lastly, the $\zeta^* \in J(x - x_0)$ can be assumed to coincide with j_{v, y_0} . These properties are crucial for actually formalizing this use of Miyadera’s lemma in the system $H_p^\omega + (+)$ and thus they lie at the heart of the above extraction.

We now consider the theorem of Reich (which features less logical subtleties). The main object featuring in Reich’s proof is the value d , the infimum over norms of all elements in the range of the operator. Internally in H_p^ω , this value can be represented by adding a further constant of type 1 which we, for simplicity, also denote by d together with a further constant $f^{\mathbb{N} \rightarrow \mathbb{N}}$ representing the witness for the monotone Dialectica interpretation of the property

$$\forall k \exists y, z (z \in Ay \wedge \|z\| - d \leq 2^{-k})$$

expressing that d indeed is the said infimum. So, we can concretely facilitate the use of d by adding the following two axioms for d :

$$(d) \quad \begin{cases} \forall y^X, z^X (z \in Ay \rightarrow d \leq_{\mathbb{R}} \|z\|_X), \\ \forall k^{\mathbb{N}} \exists y, z \leq_X f(k) 1_X (z \in Ay \wedge \|z\|_X - d \leq_{\mathbb{R}} 2^{-k}). \end{cases}$$

The additional constants are immediately majorizable by essentially themselves as they are of type $\mathbb{N} \rightarrow \mathbb{N}$ as before. Therefore the bound extraction theorems extend to this augmentation of H_p^ω in an immediate way where, in particular, the extracted bounds will in general depend on an upper bound on d as can be seen from some of the bounds extracted in the context of Reich’s theorem.

The second particularity of the formalization of the proof of Reich’s result is that one actually needs to work with the limit of $J_t x/t$, called $-v_x$ in the above, as a concrete object. In the context of the limit operator C however, we can formally deal with this object in the context of the formal systems underlying this extraction rather immediately by utilizing the previously extracted rate $\varphi_2(k)$ (where we for simplicity omit the other parameters for now) to then address the limit v_x in the system by considering

$$v_x = -C \left(\left(\frac{J_{\varphi_2(k)} x}{\varphi_2(k)} \right)_k \right).$$

In particular, with this definition of v_x , the other proofs in the context of Reich’s theorem immediately formalize.

As a last logical comment, we shortly want to discuss on the particular use of the law of excluded middle (and thus of classical logic) in the proofs for the results of Reich and Plant and how this features in the extractions, considering the fact that rates of convergence were nevertheless extracted in the absence of monotonicity. This in fact relates to the circumstances of the previously called ‘smoothable’ case distinctions. Namely, as can be observed by closer inspection of the corresponding proofs, the only part where classical logic actually features in the above proofs is through the use of multiple case distinctions which, in the case of Reich’s result, takes the form on dividing the proof between whether

$$d = 0 \text{ or } d > 0$$

and, in the case of Plant’s result, takes the form of dividing the proof between whether

$$|Ax| = 0 \text{ or } |Ax| > 0.$$

The deductions of the main results from both parts of this case distinction are essentially constructive (where the $= 0$ -case is almost trivial in both cases) and in that way, the constructive metatheorems actually allows from the extraction of a rate of convergence from the > 0 -cases as the corresponding results are of the form

$$d >_{\mathbb{R}} 0 \rightarrow C \equiv \forall c^{\mathbb{N}} (d \geq_{\mathbb{R}} 2^{-c} \rightarrow C) \quad \text{and} \quad |Ax| >_{\mathbb{R}} 0 \rightarrow C \equiv \forall c^{\mathbb{N}} (|Ax| \geq_{\mathbb{R}} 2^{-c} \rightarrow C)$$

where C is any of the respective convergence statements. These rates moreover depend on the parameter c . For the $= 0$ -case, being of the form

$$d =_{\mathbb{R}} 0 \rightarrow C \quad \text{and} \quad |Ax| =_{\mathbb{R}} 0 \rightarrow C$$

where C is any of the respective convergence statements, we find that in these cases one can actually find a proof of the classically equivalent but constructively stronger statements

$$\exists c^{\mathbb{N}} (d \leq_{\mathbb{R}} 2^{-c'} \rightarrow C) \quad \text{and} \quad \exists c^{\mathbb{N}} (|Ax| \leq_{\mathbb{R}} 2^{-c'} \rightarrow C).$$

These proof of said statements are again essentially constructive so that the constructive metatheorems guarantee the extraction of a rate again, now together with the extraction of an upper bound on (and thus a realizer of) the value c' . The previously mentioned ‘smoothing’ is now just a combination of these two cases by instantiating the former rate with $c = c'$ and combining the two resulting rates.

Acknowledgments: I want to thank Ulrich Kohlenbach for many insightful and detailed comments on various drafts of this paper. I also want to thank Pedro Pinto for the collaboration on the work [46] which served as a strong source of inspiration for the logical systems presented here.

The author was supported by the ‘Deutsche Forschungsgemeinschaft’ Project DFG KO 1737/6-2.

REFERENCES

- [1] V. Barbu. *Nonlinear semigroups and differential equations in Banach spaces*. Springer Netherlands, 1976.
- [2] V. Barbu. *Nonlinear Differential Equations of Monotone Types in Banach Spaces*. Springer Monographs in Mathematics. Springer New York, NY, 2010.
- [3] H.H. Bauschke and P.L. Combettes. *Convex Analysis and Monotone Operator Theory in Hilbert Spaces*. CMS Books in Mathematics. Springer, Cham, 2017.
- [4] M. Bezem. Strongly majorizable functionals of finite type: a model for bar recursion containing discontinuous functionals. *The Journal of Symbolic Logic*, 50:652–660, 1985.
- [5] H. Brezis. On a problem of T. Kato. *Communications on Pure and Applied Mathematics*, 24:1–6, 1971.
- [6] H. Brezis. Some remarks on semigroups of nonlinear operators. *Tohoku Mathematical Journal*, 23:245–258, 1971.
- [7] H. Brezis and A. Pazy. Accretive sets and differential equations in Banach spaces. *Israel Journal of Mathematics*, 8:367–383, 1970.
- [8] F.E. Browder. Non-Linear Equations of Evolution. *Annals of Mathematics*, 80(3):485–523, 1964.
- [9] J.A. Clarkson. Uniformly Convex Spaces. *Transactions of the American Mathematical Society*, 40(3):396–414, 1936.
- [10] M.G. Crandall. A generalized domain for semigroup generators. *Proceedings of the American Mathematical Society*, 37:434–440, 1973.
- [11] M.G. Crandall and T.M. Liggett. Generation of semigroups of nonlinear transformations on general Banach spaces. *American Journal of Mathematics*, 93:265–298, 1971.
- [12] M.G. Crandall and A. Pazy. Nonlinear evolution equations in Banach spaces. *Israel Journal of Mathematics*, 11:57–94, 1972.
- [13] F. Ferreira, L. Leuştean, and P. Pinto. On the removal of weak compactness arguments in proof mining. *Advances in Mathematics*, 354, 2019. 106728.
- [14] F. Ferreira and P. Oliva. Bounded functional interpretation. *Annals of Pure and Applied Logic*, 135:73–112, 2005.
- [15] P. Gerhardy and U. Kohlenbach. Strongly uniform bounds from semi-constructive proofs. *Annals of Pure and Applied Logic*, 141:89–107, 2006.
- [16] P. Gerhardy and U. Kohlenbach. General logical metatheorems for functional analysis. *Transactions of the American Mathematical Society*, 360:2615–2660, 2008.
- [17] K. Gödel. Über eine bisher noch nicht benützte Erweiterung des finiten Standpunktes. *Dialectica*, 12:280–287, 1958.
- [18] D. Günzel and U. Kohlenbach. Logical metatheorems for abstract spaces axiomatized in positive bounded logic. *Advances in Mathematics*, 290:503–551, 2016.
- [19] W.A. Howard. Hereditarily majorizable functionals of finite type. In Troelstra, editor, *Metamathematical Investigation of Intuitionistic Arithmetic and Analysis*, volume 344 of *Lecture Notes in Mathematics*, pages 454–461. Springer, New York, 1973.
- [20] T. Kato. Nonlinear semigroups and evolution equations. *Journal of the Mathematical Society of Japan*, 19(4):508–520, 1967.
- [21] U. Kohlenbach. *Theorie der majorisierbaren und stetigen Funktionale und ihre Anwendungen bei der Extraktion von Schranken aus inkonstruktiven Beweisen: effektive Eindeutigkeitsmodule bei besten Approximationen aus ineffektiven Eindeutigkeitsbeweisen*. PhD thesis, Johann Wolfgang Goethe-Universität Frankfurt am Main, 1990.
- [22] U. Kohlenbach. Effective bounds from ineffective proofs in analysis: an application of functional interpretation and majorization. *The Journal of Symbolic Logic*, 57:1239–1273, 1992.
- [23] U. Kohlenbach. Analysing proofs in analysis. In W. Hodges, M. Hyland, C. Steinhorn, and J. Truss, editors, *Logic: from Foundations to Applications. European Logic Colloquium (Keele, 1993)*, pages 225–260. Oxford University Press, 1996.
- [24] U. Kohlenbach. Relative constructivity. *The Journal of Symbolic Logic*, 63:1218–1238, 1998.

- [25] U. Kohlenbach. Some logical metatheorems with applications in functional analysis. *Transactions of the American Mathematical Society*, 357(1):89–128, 2005.
- [26] U. Kohlenbach. *Applied Proof Theory: Proof Interpretations and their Use in Mathematics*. Springer Monographs in Mathematics. Springer-Verlag Berlin Heidelberg, 2008.
- [27] U. Kohlenbach. Proof-theoretic Methods in Nonlinear Analysis. In B. Sirakov, P. Ney de Souza, and M. Viana, editors, *Proc. ICM 2018*, volume 2, pages 61–82. World Scientific, 2019.
- [28] U. Kohlenbach and A. Koutsoukou-Argraki. Rates of convergence and metastability for abstract Cauchy problems generated by accretive operators. *Journal of Mathematical Analysis and Applications*, 423(2):1089–1112, 2015.
- [29] U. Kohlenbach and A. Koutsoukou-Argraki. Effective asymptotic regularity for one-parameter nonexpansive semigroups. *Journal of Mathematical Analysis and Applications*, 433(2):1883–1903, 2016.
- [30] U. Kohlenbach and L. Leuştean. On the computational content of convergence proofs via Banach limits. *Philosophical Transactions of the Royal Society A*, 370:3449–3463, 2012.
- [31] U. Kohlenbach and A. Nicolae. A proof-theoretic bound extraction theorem for $\text{CAT}(\kappa)$ -spaces. *Studia Logica*, 105:611–624, 2017.
- [32] U. Kohlenbach and P. Oliva. Proof Mining: A Systematic Way of Analysing Proofs in Mathematics. *Proceedings of the Steklov Institute of Mathematics*, 242:136–164, 2003.
- [33] U. Kohlenbach and N. Pischke. Proof theory and nonsmooth analysis. 2022. To appear in *Philosophical Transactions of the Royal Society A*.
- [34] Y. Komura. Nonlinear semi-groups in Hilbert space. *Journal of the Mathematical Society of Japan*, 19(4):493–507, 1967.
- [35] G. Kreisel. On the Interpretation of Non-Finitist Proofs—Part I. *The Journal of Symbolic Logic*, 16(4):241–267, 1951.
- [36] G. Kreisel. On the Interpretation of Non-Finitist Proofs—Part II. Interpretation of Number Theory. Applications. *The Journal of Symbolic Logic*, 17(1):43–58, 1952.
- [37] S. Kuroda. Intuitionistische Untersuchungen der formalistischen Logik. *Nagoya Mathematical Journal*, 2:35–47, 1951.
- [38] L. Leuştean. Proof Mining in \mathbb{R} -trees and Hyperbolic Spaces. *Electronic Notes in Theoretical Computer Science*, 165:95–106, 2006.
- [39] L. Leuştean. An application of proof mining to nonlinear iterations. *Annals of Pure and Applied Logic*, 165(9):1484–1500, 2014.
- [40] I. Miyadera. *Nonlinear Semigroups*. Translations of Mathematical Monographs. AMS, Providence, 1992.
- [41] O. Nevanlinna and S. Reich. Strong convergence of contraction semigroups and of iterative methods for accretive operators in Banach spaces. *Israel Journal of Mathematics*, 32:44–58, 1979.
- [42] L. Paunescu and A. Sipos. A proof-theoretic metatheorem for tracial von Neumann algebras. 2022. To appear in *Mathematical Logic Quarterly*.
- [43] N.H. Pavel. *Nonlinear Evolution Operators and Semigroups: Applications to Partial Differential Equations*. Lecture Notes in Mathematics. Springer Berlin, Heidelberg, 1987.
- [44] A. Pazy. Asymptotic behavior of contractions in Hilbert space. *Israel Journal of Mathematics*, 9(2):235–240, 1971.
- [45] A. Pazy. Strong convergence of semigroups on nonlinear contractions in Hilbert space. *Journal of Mathematical Analysis and Applications*, 34:1–35, 1978.
- [46] P. Pinto and N. Pischke. On computational properties of Cauchy problems generated by accretive operators. 2023. math.AP, arXiv:2301.06880.
- [47] N. Pischke. Logical metatheorems for accretive and (generalized) monotone set-valued operators. 2023. To appear in *Journal of Mathematical Logic*.
- [48] N. Pischke. On extensionality and maximality for set-valued operators. 2023. Manuscript in preparation.
- [49] N. Pischke. Rates of convergence for the asymptotic behavior of second-order Cauchy problems. 2023. Manuscript in preparation.
- [50] A.T. Plant. The differentiability of nonlinear semigroups in uniformly convex spaces. *Israel Journal of Mathematics*, 38(3):257–268, 1981.
- [51] E.I. Poffald and S. Reich. An incomplete Cauchy problem. *Journal of Mathematical Analysis and Applications*, 113(2):514–543, 1986.
- [52] S. Reich. Strong Convergence Theorems for Resolvents of Accretive Operators in Banach Spaces. *Journal of Mathematical Analysis and Applications*, 75:287–292, 1980.
- [53] S. Reich. On the Asymptotic Behavior of Nonlinear Semigroups and the Range of Accretive Operators. *Journal of Mathematical Analysis and Applications*, 79:113–126, 1981.
- [54] A. Sipos. Proof mining in L^p spaces. *The Journal of Symbolic Logic*, 84(4):1612–1629, 2019.
- [55] T. Tao. Norm Convergence of Multiple Ergodic Averages for Commuting Transformations. *Ergodic Theory and Dynamical Systems*, 28:657–688, 2008.
- [56] T. Tao. *Structure and Randomness: Pages from Year One of a Mathematical Blog*, chapter Soft analysis, hard analysis, and the finite convergence principle. American Mathematical Society, Providence, RI, 2008.
- [57] H.-K. Xu. Strong asymptotic behavior of almost-orbits of nonlinear semigroups. *Nonlinear Analysis: Theory, Methods & Applications*, 46(1):135–151, 2001.