

# Sharp Besov capacity estimates for annuli in metric spaces with doubling measures

Anders Björn

*Department of Mathematics, Linköping University, SE-581 83 Linköping, Sweden  
anders.bjorn@liu.se, ORCID: 0000-0002-9677-8321*

Jana Björn

*Department of Mathematics, Linköping University, SE-581 83 Linköping, Sweden  
jana.bjorn@liu.se, ORCID: 0000-0002-1238-6751*

**Abstract.** We obtain precise estimates, in terms of the measure of balls, for the Besov capacity of annuli and singletons in complete metric spaces. The spaces are only assumed to be uniformly perfect with respect to the centre of the annuli and equipped with a doubling measure.

*Key words and phrases:* annulus, Besov space, condenser capacity, doubling measure, fractional Sobolev space, metric space, norm-capacity, pointwise dimension, uniformly perfect, Sobolev–Slobodetskiĭ space.

*Mathematics Subject Classification (2020):* Primary: 31C45; Secondary: 30L99, 31C12, 31C15, 31E05, 46E36.

## 1. Introduction

Capacities are intimately related to function spaces in the sense that various properties, such as quasicontinuity and Lebesgue points, of functions in such spaces are measured by capacity. Capacities also reflect metric and measure-theoretic properties of the underlying space on which they are defined. For example, it is well known that the  $p$ -capacity of a spherical condenser in  $\mathbf{R}^n$  with  $0 < 2r \leq R$  reflects the dimension of the space as follows,

$$\operatorname{cap}_p(B(x, r), B(x, R)) \simeq \begin{cases} r^{n-p} & \text{if } 1 \leq p < n, \\ (\log(R/r))^{1-p} & \text{if } 1 \leq p = n, \\ R^{n-p} & \text{if } p > n. \end{cases} \quad (1.1)$$

Capacities also play an important role in fine potential theory and appear in the famous Wiener criterion characterizing boundary regularity for various equations, such as  $\Delta_p u = 0$  (Maz'ya [29] and Kilpeläinen–Malý [22], with the  $p$ -capacity as in (1.1)) and the fractional  $p$ -Laplace equation  $(-\Delta_p)^s u = 0$  (Kim–Lee–Lee [23], using a fractional Besov capacity from [9]).

In this paper we study Besov capacities on a complete metric space  $Y = (Y, d)$  equipped with a doubling measure  $\nu$ . Analogously to (1.1), we are primarily interested in estimates for (thick) annuli, i.e. of the capacity for a ball  $B(x_0, r)$  within  $B(x_0, R)$  where  $0 < 2r \leq R$ .

*Throughout the paper we assume that  $1 \leq p < \infty$ . We also fix a point  $x_0$  and let  $B_r = B(x_0, r)$  be the open ball with radius  $r$  and centre  $x_0$ .*

The following are our main results.

**Theorem 1.1.** *Assume that  $Y$  is a complete metric space which is uniformly perfect at  $x_0$  and equipped with a doubling measure  $\nu$ . Let  $p > 1$  and  $0 < \theta < 1$ . Then for all  $0 < 2r \leq R \leq \frac{1}{4} \text{diam } Y$ ,*

$$\text{cap}_{\theta,p}(B_r, B_R) \simeq \left( \int_r^R \left( \frac{\rho^{\theta p}}{\nu(B_\rho)} \right)^{1/(p-1)} \frac{d\rho}{\rho} \right)^{1-p} \quad (1.2)$$

and

$$\text{cap}_{\theta,p}(\{x_0\}, B_R) \simeq \left( \int_0^R \left( \frac{\rho^{\theta p}}{\nu(B_\rho)} \right)^{1/(p-1)} \frac{d\rho}{\rho} \right)^{1-p}, \quad (1.3)$$

with the comparison constants in “ $\simeq$ ” independent of  $x_0$ ,  $r$  and  $R$ .

Here,  $\text{cap}_{\theta,p}$  is the Besov condenser capacity defined for bounded sets  $E \Subset \Omega$  as

$$\text{cap}_{\theta,p}(E, \Omega) = \inf_u \int_Y \int_Y \frac{|u(x) - u(y)|^p}{d(x,y)^{\theta p}} \frac{d\nu(y) d\nu(x)}{\nu(B(x, d(x,y)))}, \quad (1.4)$$

where the infimum is taken over all  $u$  such that  $0 \leq u \leq 1$  everywhere,  $u = 1$  in a neighbourhood of  $E$  and  $\text{supp } u \Subset \Omega$ , see Definition 3.2.

The Euclidean spaces and their subsets, equipped with the Lebesgue measure or weighted measures  $w dx$ , and even singular doubling measures, are included as special cases of our results. We emphasize that we do not assume any Poincaré inequalities for upper gradients on  $Y$  (as in Gogatishvili–Koskela–Zhou [17, Section 4] and Koskela–Yang–Zhou [25]). This makes our results applicable to many disconnected spaces and spaces carrying few rectifiable curves, including fractals.

To formulate the next two results we need the following exponent sets:

$$\begin{aligned} \underline{Q}_0 &= \left\{ q > 0 : \frac{\nu(B_r)}{\nu(B_R)} \lesssim \left( \frac{r}{R} \right)^q \text{ for } 0 < r < R \leq 1 \right\}, \\ \underline{S}_0 &= \{ s > 0 : \nu(B_r) \lesssim r^s \text{ for } 0 < r \leq 1 \}, \\ \bar{S}_0 &= \{ s > 0 : \nu(B_r) \gtrsim r^s \text{ for } 0 < r \leq 1 \}, \\ \bar{Q}_0 &= \left\{ q > 0 : \frac{\nu(B_r)}{\nu(B_R)} \gtrsim \left( \frac{r}{R} \right)^q \text{ for } 0 < r < R \leq 1 \right\}. \end{aligned}$$

These sets were introduced in Björn–Björn–Lehrbäck [5] to capture the local behaviour of the measure at  $x_0$ . For example, for the Lebesgue measure in  $\mathbf{R}^n$ ,

$$\underline{Q}_0 = \underline{S}_0 = (0, n] \quad \text{and} \quad \bar{S}_0 = \bar{Q}_0 = [n, \infty).$$

The subscript 0 in the above definitions stands for the fact that the inequalities are required to hold for small radii. It is easily verified (see [5, Lemmas 2.4 and 2.5]) that the exponent sets can equivalently be defined using  $0 < r \leq \Theta R \leq R_0$  for any fixed  $0 < \Theta < 1$  and  $R_0 > 0$ , even though the implicit comparison constants in “ $\lesssim$ ” and “ $\gtrsim$ ” will then depend on  $\Theta$  and  $R_0$ .

All of these sets are intervals. The reason for introducing them as sets is that they may or may not contain their endpoints

$$\underline{q}_0 = \sup \underline{Q}_0, \quad \underline{s}_0 = \sup \underline{S}_0, \quad \bar{s}_0 = \inf \bar{S}_0 \quad \text{and} \quad \bar{q}_0 = \inf \bar{Q}_0. \quad (1.5)$$

Note that  $\bar{q}_0 < \infty$  if  $\nu$  is doubling, and that  $\underline{q}_0 > 0$  if  $Y$  is also uniformly perfect at  $x_0$  (see Heinonen [19, Exercise 13.1]).

When  $p > 1$  and  $\theta p < \underline{q}_0$  or  $\theta p > \bar{q}_0$ , Theorem 1.1 provides us with exact estimates for the capacity  $\text{cap}_{\theta,p}(B_r, B_R)$  in terms of  $\nu(B_r)$  or  $\nu(B_R)$ . When  $p = 1$ ,

Theorem 1.1 cannot be used, but we obtain the following similar estimates for  $\text{cap}_{\theta,p}(B_r, B_R)$  by using results from Björn–Björn–Lehrbäck [5], which cover all  $p \geq 1$ . The borderline cases  $\theta p = \max \underline{Q}_0$  and  $\theta p = \min \overline{Q}_0$  are considered in Theorem 9.1. See also Remarks 9.2 and 9.3.

**Theorem 1.2.** *Assume that  $Y$  is a complete metric space which is uniformly perfect at  $x_0$  and equipped with a doubling measure  $\nu$ . Let  $0 < \theta < 1$  and  $0 < R_0 \leq \frac{1}{4} \text{diam} Y$ , with  $R_0$  finite.*

(a) *If  $\theta p < \underline{q}_0$ , then*

$$\text{cap}_{\theta,p}(B_r, B_R) \simeq \frac{\nu(B_r)}{r^{\theta p}} \quad \text{whenever } 0 < 2r \leq R \leq R_0. \quad (1.6)$$

(b) *If  $\theta p > \overline{q}_0$ , then*

$$\text{cap}_{\theta,p}(B_r, B_R) \simeq \frac{\nu(B_R)}{R^{\theta p}} \quad \text{whenever } 0 < 2r \leq R \leq R_0. \quad (1.7)$$

In both cases, the comparison constants in “ $\simeq$ ” depend on  $R_0$ .

Moreover, the lower bound in (1.6) implies  $p \in \underline{Q}_0$ , while the lower bound in (1.7) implies  $p \in \overline{Q}_0$ . If  $p > 1$  then (1.7) holds if and only if  $\theta p > \overline{q}_0$ .

In Ahlfors regular spaces, estimates (1.6) and (1.7) were given in Lehrbäck–Shanmugalingam [27], and used to show that Besov-norm-preserving homeomorphisms between such spaces are quasisymmetric.

In many situations it is important whether singletons have zero or positive capacity. In the following result, we characterize these cases in terms of the exponent sets  $\overline{S}_0$  and  $\underline{S}_0$ .

**Theorem 1.3.** *Assume that  $Y$  is a complete metric space which is uniformly perfect at  $x_0$  and equipped with a doubling measure  $\nu$ . Let  $0 < \theta < 1$ .*

(a) *If  $\theta p > \overline{s}_0$ , then*

$$C_{\theta,p}(\{x_0\}) > 0 \quad \text{and} \quad \text{cap}_{\theta,p}(\{x_0\}, B_R) > 0 \quad \text{for every } 0 < R < \frac{1}{2} \text{diam} Y,$$

where the capacity  $C_{\theta,p}$  is defined by means of the Besov norm as in Definition 3.2.

(b) *If  $\theta p \notin \overline{S}_0$  (in particular if  $\theta p < \overline{s}_0$ ), or if  $p > 1$  and  $\theta p \in \underline{S}_0$ , then*

$$C_{\theta,p}(\{x_0\}) = 0 \quad \text{and} \quad \text{cap}_{\theta,p}(\{x_0\}, B_R) = 0 \quad \text{for every } R > 0.$$

In Anttila [1], the numbers  $\overline{s}_0$  and  $\underline{s}_0$  are called the *upper and lower local dimensions* of  $\mu$  at  $x_0$ , while  $\overline{q}$  in Remark 9.2 is called the *pointwise Assouad dimension* of  $\mu$  at  $x_0$ . (See [5, Lemma 2.4] for why the definitions of  $\overline{s}_0$  and  $\underline{s}_0$  in [1] are equivalent to those in (1.5).) In [6],  $\overline{s}_0$  played a decisive role in determining the sharp integrability properties for  $p$ -harmonic Green functions and their gradients.

On  $\mathbf{R}^n$ , the spaces defined by means of the energy integral in (1.4) are often called fractional Sobolev spaces and are the traces of Sobolev spaces on sufficiently nice domains (Jonsson–Wallin [20]). As such, they are suitable as boundary values for various Dirichlet problems and appear in boundary regularity results for elliptic differential equations (Kristensen–Mingione [26]).

They also play an important role in nonlocal problems, such as the fractional  $p$ -Laplace equation  $(-\Delta_p)^s u = 0$ . These problems have attracted a lot of attention in the past two decades, see e.g. Kim–Lee–Lee [23], Korvenpää–Kuusi–Lindgren [24] and Lindgren–Lindqvist [28], to name just a few.

Recently, similar problems and the associated Besov spaces have been studied for metric measure spaces in e.g. Capogna–Kline–Korte–Shanmugalingam–Snipes [12], Eriksson–Bique–Giovannardi–Korte–Shanmugalingam–Speight [15], Gogatishvili–Koskela–Shanmugalingam [16], Gogatishvili–Koskela–Zhou [17] and Koskela–Yang–Zhou [25]. The role of Besov spaces as traces of Sobolev type spaces was in the metric setting studied in Bourdon–Pajot [11], Björn–Björn–Gill–Shanmugalingam [4] and Björn–Björn–Shanmugalingam [8], and will be one of our main tools.

Our approach to the above estimates is based on extensions of Besov functions from  $Y$  to hyperbolic fillings of  $Y$ , together with estimates from Björn–Björn–Lehrbäck [5], [6] for  $p$ -capacities associated with Sobolev spaces. More precisely, we use the comparison between Besov seminorms of functions on  $Y$  and the Dirichlet energy of their extensions to a uniformized hyperbolic filling of  $Y$ , obtained in [8]. These constructions and comparisons are done in Section 5.

However, since the results in [8] only cover bounded spaces, special care has to be taken for unbounded  $Y$ . This is done in Section 7 by replacing  $Y$  with a suitably chosen bounded subset, so that the restriction of  $\nu$  is still doubling. Even when  $Y$  is bounded, it is only biLipschitz equivalent to the boundary of the uniformized hyperbolic filling of  $Y$ , which would in turn put serious restrictions on the allowed radii  $r$  and  $R$  in our estimates. In Section 6 we therefore show how to replace  $Y$  by a carefully constructed enlarged space so that the involved capacities are comparable and all radii  $\leq \frac{1}{4} \text{diam } Y$  can be treated.

Along the way, in Sections 3 and 4, we prove various fundamental properties of Besov capacities in metric spaces, both for doubling and nondoubling measures, including in some cases also  $\theta \geq 1$ . Finally, in Sections 8 and 9, we prove Theorems 1.1–1.3.

As mentioned above, we use hyperbolic fillings to obtain our main results. It would be interesting to find more direct proofs. On the other hand, our technique shows that there is a direct correspondence between these results and the corresponding results for Sobolev spaces in [5] and [6].

*Acknowledgement.* A. B. and J. B. were supported by the Swedish Research Council, grants 2020-04011 resp. 2018-04106. Part of this research was done when the authors visited Institut Mittag-Leffler in the autumn of 2022 during the programme *Geometric Aspects of Nonlinear Partial Differential Equations*. We thank the institute for their hospitality and support.

## 2. Preliminaries

*In this section we assume that  $X = (X, d)$  is a metric space equipped with a Borel measure  $\mu$  such that  $0 < \mu(B) < \infty$  for every ball  $B \subset X$ . To avoid pathological situations we also assume that all metric spaces, considered in this paper, contain at least two points.*

As is often customary we extend  $\mu$ , and other measures, as outer measures defined on all sets. This plays a role at least in Proposition 3.3 (ii).

A metric space is *proper* if all closed bounded sets are compact. We denote balls in  $X$  by

$$B(x, r) = \{y \in X : d(y, x) < r\} \quad \text{and let } \lambda B(x, r) = B(x, \lambda r).$$

All balls in this paper are open. In metric spaces it can happen that balls with different centres and/or radii denote the same set. We will however use the convention that a ball comes with a predetermined centre and radius.

The space  $X$  is *uniformly perfect at  $x$*  if there is a constant  $\kappa > 1$  such that

$$B(x, \kappa r) \setminus B(x, r) \neq \emptyset \quad \text{whenever } B(x, \kappa r) \neq X. \quad (2.1)$$

In fact, it then follows that (2.1) holds whenever  $B(x, r) \neq X$ , since if  $B(x, \kappa r) = X$  then  $B(x, \kappa r) \setminus B(x, r) = X \setminus B(x, r) \neq \emptyset$ .

The space  $X$  is *uniformly perfect* if it is uniformly perfect at every  $x$  with the same constant  $\kappa$ . This definition coincides with the one in Heinonen [19, Section 11.1], see therein for more on the history of this assumption. We do not know if pointwise uniform perfectness has been used before. Note that  $X$  is uniformly perfect with any  $\kappa > 1$  if  $X$  is connected.

The measure  $\mu$  is *doubling* if there is a *doubling constant*  $C_\mu > 1$  such that

$$0 < \mu(2B) \leq C_\mu \mu(B) < \infty \quad \text{for all balls } B.$$

Similarly,  $\mu$  is *reverse-doubling at  $x$* , if there are constants  $C, \hat{\kappa} > 1$  such that

$$\mu(B(x, \hat{\kappa}r)) \geq C\mu(B(x, r)) \quad \text{for all } 0 < r < \text{diam } X/2\hat{\kappa}. \quad (2.2)$$

By continuity of the measure, the estimate (2.2) holds also if  $r = \text{diam } X/2\hat{\kappa} < \infty$ , as required in Björn–Björn–Lehrbäck [5]. If  $\mu$  is doubling, it is easy to see that  $X$  is uniformly perfect at  $x$  if and only if  $\mu$  is reverse-doubling at  $x$ . (For necessity we can choose any  $\hat{\kappa} > \kappa$ , and for sufficiency any  $\kappa > 2\hat{\kappa}$ .) If  $\mu$  is doubling and  $X$  is connected, then  $\mu$  is reverse-doubling at every  $x$  with any  $\hat{\kappa} > 1$ .

Throughout the paper, we write  $a \lesssim b$  if there is an implicit constant  $C > 0$  such that  $a \leq Cb$ , and analogously  $a \gtrsim b$  if  $b \lesssim a$ , and  $a \simeq b$  if  $a \lesssim b \lesssim a$ . The implicit comparison constants are allowed to depend on the standard parameters. We will carefully explain the dependence in each case. See Remarks 8.2 and 9.3 for the dependence in Theorems 1.1 and 1.2.

Sometimes, when dealing with several different spaces simultaneously, we will write  $B^X$ ,  $d_X$ ,  $\bar{Q}_0^X$ ,  $\bar{q}_0^X$  etc. to indicate that these notions are taken with respect to the metric space  $X$ .

### 3. Besov spaces and capacities

In this section we assume that  $Y = (Y, d)$  is a proper metric space equipped with a Borel measure  $\nu$  such that  $0 < \nu(B) < \infty$  for every ball  $B \subset Y$ . We also assume that  $\theta > 0$ , and emphasize that in this section  $\theta \geq 1$  is allowed.

For a measurable  $u : Y \rightarrow [-\infty, \infty]$  (which is finite  $\nu$ -a.e.) we define the *Besov seminorm* by

$$[u]_{\theta, p} = [u]_{\theta, p, Y} = \left( \int_Y \int_Y \frac{|u(x) - u(y)|^p}{d(x, y)^{\theta p}} \frac{d\nu(y) d\nu(x)}{\nu(B(x, d(x, y)))} \right)^{1/p}.$$

Here and elsewhere, the integrand should be interpreted as zero when  $y = x$ .

The Besov space  $B_p^\theta(Y)$  consists of the functions  $u$  such that the *Besov norm*

$$\|u\|_{B_p^\theta(Y)}^p := [u]_{\theta, p}^p + \|u\|_{L^p(Y)}^p < \infty. \quad (3.1)$$

This space is a Banach space, see Remark 9.8 in Björn–Björn–Shanmugalingam [8]. (The norm (3.1) is equivalent to the one in [8], but the norm-capacity  $C_{\theta, p}$  below exactly coincides with the one in [8].)

We restrict our attention to Besov spaces with two indices (i.e. “ $q = p$ ”). Such Besov spaces are often called fractional Sobolev spaces or Sobolev–Slobodetskiĭ spaces, although Besov spaces seem to be the most common name in the metric space literature.

Assuming that  $\nu$  is doubling, equivalent definitions, using equivalent seminorms, are given in Gogatishvili–Koskela–Shanmugalingam [16, Theorem 5.2 and (5.1)].

When  $\nu$  is also reverse-doubling (or equivalently, uniformly perfect), further equivalent definitions can be found in Gogatishvili–Koskela–Zhou [17, Theorem 4.1 and Proposition 4.1], for example that the Besov space  $B_p^\theta(Y)$  considered here coincides with the corresponding Hajlasz–Besov space. By [16, Lemmas 6.1 and 6.2], it is also related to fractional Hajlasz spaces, considered already in Yang [31]. See these papers for the precise definitions and earlier references to the theory on  $\mathbf{R}^n$  and on Ahlfors regular metric spaces.

We are interested in two types of Besov capacities, the norm-capacity and the condenser capacity.

**Definition 3.1.** The *Besov norm-capacity* of  $E \subset Y$  is

$$C_{\theta,p}(E) = C_{\theta,p}^Y(E) = \inf_u \|u\|_{B_p^\theta(Y)}^p, \quad (3.2)$$

where the infimum is taken over all  $u \in B_p^\theta(Y)$  such that  $u = 1$  in a neighbourhood of  $E$  and  $0 \leq u \leq 1$  everywhere. Such  $u$  are called *admissible* for  $C_{\theta,p}(E)$ .

By truncation it follows that one can equivalently take the infimum over all  $u$  such that  $u \geq 1$  in a neighbourhood of  $E$ . As usual, when requiring that  $u \geq 1$  or  $0 \leq u \leq 1$  everywhere we mean that there is a representative of  $u$  satisfying these requirements. By  $E \Subset \Omega$  we mean that  $\bar{E}$  is a compact subset of  $\Omega$ .

**Definition 3.2.** Let  $\Omega \subset Y$  be a bounded open set and  $E \Subset \Omega$ . The *Besov condenser capacity* is given by

$$\text{cap}_{\theta,p}(E, \Omega) = \text{cap}_{\theta,p}^Y(E, \Omega) = \inf_u [u]_{\theta,p}^p,$$

where the infimum is taken over all  $u$  such that  $0 \leq u \leq 1$  everywhere,  $u = 1$  in a neighbourhood of  $E$  and  $\text{supp } u \Subset \Omega$ . Such  $u$  are called *admissible* for  $\text{cap}_{\theta,p}(E)$ .

The corresponding capacities for Sobolev spaces are called Sobolev resp. variational capacity in [2]. Condenser capacities are also often called “relative”.

There do not seem to be very many papers on Besov capacities in metric spaces. Nuutinen [30] and Heikkinen–Koskela–Tuominen [18] extensively studied the norm-capacity, defined using the Hajlasz–Besov norm, under the assumption that  $\nu$  is doubling. In [18], they also considered the corresponding Triebel–Lizorkin norm-capacity, which was later studied by Karak [21]. The Besov norm-capacity  $C_{\theta,p}$  was used by Björn–Björn–Shanmugalingam [8]. The Besov condenser capacity  $\text{cap}_{\theta,p}$  was studied in the Ahlfors  $Q$ -regular case by Bourdon [10] ( $p > Q$  and  $\theta = 1/p$ ), Costea [14] ( $p > Q$ ) and Lehrbäck–Shanmugalingam [27].

Our main estimates remain the same (up to changes in implicit constants) when the seminorm is replaced by an equivalent seminorm. However, some of the basic properties, such as subadditivity, are not directly transferable, although the proofs often are, so we include them here.

**Proposition 3.3.** Let  $E, E_1, E_2, \dots \subset Y$ . Then the following properties hold:

- (i) if  $E_1 \subset E_2$ , then  $C_{\theta,p}(E_1) \leq C_{\theta,p}(E_2)$ ,
- (ii)  $\nu(E) \leq C_{\theta,p}(E)$ ,
- (iii) if  $K_1 \supset K_2 \supset \dots$  are compact subsets of  $X$ , then

$$C_{\theta,p}\left(\bigcap_{i=1}^{\infty} K_i\right) = \lim_{i \rightarrow \infty} C_{\theta,p}(K_i),$$

- (iv)  $C_{\theta,p}$  is countably subadditive, i.e. if  $E = \bigcup_{i=1}^{\infty} E_i$  then

$$C_{\theta,p}(E) \leq \sum_{i=1}^{\infty} C_{\theta,p}(E_i).$$

The monotonicity (i) is trivial, while (ii) follows directly from the definition. The property (iii) follows from the fact that  $C_{\theta,p}$  is an outer capacity (by definition), i.e.

$$C_{\theta,p}(E) = \inf_{\substack{G \supset E \\ G \text{ open}}} C_{\theta,p}(G),$$

and elementary properties of compact sets, see Nuutinen [30, Section 3]. As for (iv), Nuutinen [30] only obtains quasi-subadditivity since he works in a more general setting in which the countable subadditivity does not always hold. We therefore provide a proof.

*Proof of (iv).* We may assume that the right-hand side is finite. Let  $\varepsilon > 0$ . For each  $i = 1, 2, \dots$ , choose  $u_i$  admissible for  $C_{\theta,p}(E_i)$  with

$$[u_i]_{\theta,p}^p + \|u_i\|_{L^p(Y)}^p < C_{\theta,p}(E_i) + \frac{\varepsilon}{2^i}.$$

Let  $u = \sup_i u_i$ . Then  $u = 1$  in a neighbourhood of  $\bigcup_{i=1}^{\infty} E_i$ . Moreover, for  $x, y \in Y$ ,

$$|u(x) - u(y)|^p \leq \sup_i |u_i(x) - u_i(y)|^p \leq \sum_{i=1}^{\infty} |u_i(x) - u_i(y)|^p$$

and similarly,  $|u(x)|^p = \sup_i |u_i(x)|^p \leq \sum_{i=1}^{\infty} |u_i(x)|^p$ . Hence

$$\begin{aligned} C_{\theta,p}(E) &\leq ([u]_{\theta,p}^p + \|u\|_{L^p(Y)}^p) \leq \sum_{i=1}^{\infty} ([u_i]_{\theta,p}^p + \|u_i\|_{L^p(Y)}^p) \\ &< \sum_{i=1}^{\infty} \left( C_{\theta,p}(E_i) + \frac{\varepsilon}{2^i} \right) = \sum_{i=1}^{\infty} C_{\theta,p}(E_i) + \varepsilon. \end{aligned}$$

Letting  $\varepsilon \rightarrow 0$  completes the proof.  $\square$

**Proposition 3.4.** *Let  $\Omega \subset \Omega' \subset Y$  be open and bounded,  $E, E_1, E_2, \dots \Subset \Omega$ . Then the following properties hold:*

- (i) *if  $E_1 \subset E_2 \subset \Omega$ , then  $\text{cap}_{\theta,p}(E_1, \Omega') \leq \text{cap}_{\theta,p}(E_2, \Omega)$ ,*
- (ii) *if  $K_1 \supset K_2 \supset \dots$  are compact subsets of  $\Omega$ , then*

$$\text{cap}_{\theta,p} \left( \bigcap_{i=1}^{\infty} K_i, \Omega \right) = \lim_{i \rightarrow \infty} \text{cap}_{\theta,p}(K_i, \Omega),$$

- (iii)  *$\text{cap}_{\theta,p}$  is countably subadditive, i.e. if  $E = \bigcup_{i=1}^{\infty} E_i$  then*

$$\text{cap}_{\theta,p}(E) \leq \sum_{i=1}^{\infty} \text{cap}_{\theta,p}(E_i, \Omega).$$

Again, (i) is trivial, while (ii) follows from elementary properties of compact sets since  $\text{cap}_{\theta,p}$  is an outer capacity (by definition). The proof of (iii) is similar to the proof of Proposition 3.3 (iv).

In the Ahlfors  $Q$ -regular case with  $p > Q > 1$ , these facts were stated in Costea [14] with a comment that the proof is essentially the same as in Costea [13, Theorem 3.1]. His proof of (iii) uses reflexivity. Our proof is considerably shorter and also covers the case  $1 \leq p \leq Q$  as well as the non-Ahlfors regular case.



## 4. Capacity estimates when $\nu$ is doubling

In this section we assume that  $Y$  is a complete metric space equipped with a doubling measure  $\nu$  and that  $0 < \theta < 1$ .

Note that  $Y$  is proper, see Björn–Björn [2, Proposition 3.1]. The comparison constants in this section are independent of the choice of  $x_0$ , they depend only on  $\theta$ ,  $p$  and  $C_\nu$ , unless said otherwise.

Our next aim is to deduce the following result, which will be important later on. Note that  $B_{2R} \neq Y$  whenever  $R < \frac{1}{4} \text{diam } Y$ .

**Proposition 4.1.** *Assume that  $Y$  is uniformly perfect at  $x_0$  with constant  $\kappa$ . Fix  $0 < \Theta < 1$ . If  $0 < \Theta R \leq 2r \leq R$  and  $B_{2R} \neq Y$ , then*

$$\text{cap}_{\theta,p}(B_r, B_R) \simeq \frac{\nu(B_r)}{r^{\theta p}} \simeq \frac{\nu(B_R)}{R^{\theta p}},$$

with comparison constants also depending on  $\kappa$  and  $\Theta$ .

We split the proof of Proposition 4.1 into two parts. We begin with the lower bound, which holds also when  $\theta \geq 1$ .

**Proposition 4.2.** *Assume that  $Y$  is uniformly perfect at  $x_0$  with constant  $\kappa$ , and that  $0 < 2r \leq R$  with  $B_{2R} \neq Y$ . Then*

$$\text{cap}_{\theta,p}(B_r, B_R) \gtrsim \frac{\nu(B_r)}{R^{\theta p}},$$

with comparison constant also depending on  $\kappa$ .

*Proof.* Let  $u$  be admissible for  $\text{cap}_{\theta,p}(B_r, B_R)$ . As  $B_{2R} \neq Y$  it follows from the uniform perfectness that there exists  $z \in B_{2\kappa R} \setminus B_{2R}$ . Since  $B(z, R) \cap B_R = \emptyset$  and  $d(x, y) \leq (2\kappa + 2)R$  for all  $x \in B(z, R)$  and  $y \in B_r$ , we get that

$$[u]_{\theta,p}^p \geq \int_{B(z,R)} \int_{B_r} \frac{1}{((2\kappa + 2)R)^{\theta p}} \frac{d\nu(y) d\nu(x)}{\nu(B(x, (2\kappa + 2)R))} \gtrsim \frac{\nu(B_r)}{R^{\theta p}}.$$

Taking infimum over all  $u$  admissible for  $\text{cap}_{\theta,p}(B_r, B_R)$  concludes the proof.  $\square$

To prove the upper bound in Proposition 4.1 we will use the following simple lemma, which will also be used when proving Lemma 4.5.

**Lemma 4.3.** *Assume that  $0 \leq \eta \leq 1$  is an  $M$ -Lipschitz function on  $Y$ . If  $x \in Y$ , then*

$$I(x) := \int_Y \frac{|\eta(x) - \eta(y)|^p}{d(x, y)^{\theta p}} \frac{d\nu(y)}{\nu(B(x, d(x, y)))} \lesssim M^{\theta p}.$$

*Proof.* Let  $B^j = B(x, 2^j/M)$ . Since  $\nu(B(x, d(x, y))) \simeq \nu(B^j)$  for  $y \in B^j \setminus B^{j-1}$  and  $0 < \theta < 1$ , we see that

$$\begin{aligned} I(x) &\simeq \sum_{j=-\infty}^{\infty} \int_{B^j \setminus B^{j-1}} \frac{|\eta(x) - \eta(y)|^p}{d(x, y)^{\theta p}} \frac{d\nu(y)}{\nu(B^j)} \\ &\lesssim \sum_{j=-\infty}^0 \int_{B^j \setminus B^{j-1}} \frac{M^p d(x, y)^{(1-\theta)p}}{\nu(B^j)} d\nu(y) + \sum_{j=1}^{\infty} \int_{B^j \setminus B^{j-1}} \frac{d(x, y)^{-\theta p}}{\nu(B^j)} d\nu(y) \\ &\lesssim \sum_{j=-\infty}^0 M^{\theta p} 2^{j(1-\theta)p} + \sum_{j=1}^{\infty} M^{\theta p} 2^{-j\theta p} \simeq M^{\theta p}. \end{aligned} \quad \square$$



This now leads to the following estimate.

**Proposition 4.4.** *Assume that  $0 < 2r \leq R$ . Then*

$$\text{cap}_{\theta,p}(B_r, B_R) \lesssim \min \left\{ \frac{\nu(B_r)}{r^{\theta p}}, \frac{\nu(B_R)}{R^{\theta p}} \right\}.$$

*Proof.* Let  $u$  be a  $3/R$ -Lipschitz function admissible for  $\text{cap}_{\theta,p}(B_{R/2}, B_R)$ . The doubling property and symmetry in  $x$  and  $y$  imply that

$$[u]_{\theta,p}^p \simeq \int_{B_R} \int_Y \frac{|u(x) - u(y)|^p}{d(x,y)^{\theta p}} \frac{d\nu(y)}{\nu(B(x, d(x,y)))} d\nu(x).$$

Integrating the estimate from Lemma 4.3 over  $x \in B_R$  gives

$$\text{cap}_{\theta,p}(B_r, B_R) \leq \text{cap}_{\theta,p}(B_{R/2}, B_R) \lesssim \frac{\nu(B_R)}{R^{\theta p}}.$$

Applying the last estimate with  $R$  replaced by  $2r$  gives

$$\text{cap}_{\theta,p}(B_r, B_R) \leq \text{cap}_{\theta,p}(B_r, B_{2r}) \lesssim \frac{\nu(B_r)}{r^{\theta p}}. \quad \square$$

*Proof of Proposition 4.1.* This follows directly from Propositions 4.2 and 4.4, together with the doubling property.  $\square$

**Lemma 4.5.** *Assume that  $\Omega \subset Y$  is open and bounded, and let  $E \Subset \Omega$ . If  $C_{\theta,p}(E) = 0$ , then  $\text{cap}_{\theta,p}(E, \Omega) = 0$ .*

*Proof.* Since  $E \Subset \Omega$  there is a Lipschitz function  $0 \leq \eta \leq 1$  such that  $\eta = 1$  in a neighbourhood of  $E$  and  $\eta = 0$  in a neighbourhood of  $Y \setminus \Omega$ . Let  $M$  be the Lipschitz constant of  $\eta$  and let  $\varepsilon > 0$ . As  $C_{\theta,p}(E) = 0$  there is a function  $u$  admissible for  $C_{\theta,p}(E)$  with  $\|u\|_{B_p^{\theta}(Y)}^p < \varepsilon$ . Let  $v = u\eta$ . Then

$$\begin{aligned} |v(x) - v(y)| &= |u(x)\eta(x) - u(x)\eta(y) + u(x)\eta(y) - u(y)\eta(y)| \\ &\leq u(x)|\eta(x) - \eta(y)| + |u(x) - u(y)|. \end{aligned}$$

Hence, by Lemma 4.3,

$$\begin{aligned} [v]_{\theta,p}^p &\leq 2^p \int_Y u(x)^p \int_Y \frac{|\eta(x) - \eta(y)|^p}{d(x,y)^{\theta p}} \frac{d\nu(y)}{\nu(B(x, d(x,y)))} + 2^p [u]_{\theta,p}^p \\ &\lesssim M^{\theta p} \|u\|_{L^p(Y)}^p + [u]_{\theta,p}^p \leq (M^{\theta p} + 1) \|u\|_{B_p^{\theta}(Y)}^p < (M^{\theta p} + 1)\varepsilon. \end{aligned}$$

As  $v = 1$  in a neighbourhood of  $E$  and  $v = 0$  in a neighbourhood of  $Y \setminus \Omega$ , we see that

$$\text{cap}_{\theta,p}(E, \Omega) \leq [v]_{\theta,p}^p \lesssim (M^{\theta p} + 1)\varepsilon.$$

Letting  $\varepsilon \rightarrow 0$  completes the proof.  $\square$

Note that the converse of Lemma 4.5 does not hold in general; consider e.g. a compact  $Y$  in which case  $\text{cap}_{\theta,p}(Y, Y) = 0$  (as  $u \equiv 1$  is admissible) while  $C_{\theta,p}(Y) \geq \nu(Y) > 0$ . Nevertheless, we will prove the following characterization.

**Proposition 4.6.** *Assume that  $\Omega \subset Y$  is open and bounded and such that  $\nu(Y \setminus \Omega) > 0$ . Let  $E \Subset \Omega$ . Then  $C_{\theta,p}(E) = 0$  if and only if  $\text{cap}_{\theta,p}(E, \Omega) = 0$ .*

The following simple observation will serve as a Poincaré type inequality. We will use it to prove Proposition 4.6 as well as Lemma 4.9 below.

**Lemma 4.7.** *If  $u = 0$  outside a bounded measurable set  $\Omega$  and  $K \subset Y \setminus \Omega$  is a bounded measurable set with  $\nu(K) > 0$ , then for every  $z \in K$ ,*

$$\int_Y |u|^p d\nu \leq R^{\theta p} \frac{\nu(B(z, R))}{\nu(K)} [u]_{\theta, p}^p,$$

where

$$R = \text{diam } K + \sup\{d(x, y) : x \in K \text{ and } y \in \Omega\}.$$

*Proof.* Since  $u = 0$  outside  $\Omega$ , and in particular in  $K$ , and  $B(x, d(x, y)) \subset B(z, R)$  for all  $x \in K$  and  $y \in \Omega$ , we see that

$$\begin{aligned} \int_Y |u|^p d\nu &= \frac{1}{\nu(K)} \int_K \int_{\Omega} |u(x) - u(y)|^p d\nu(y) d\nu(x) \\ &\leq R^{\theta p} \frac{\nu(B(z, R))}{\nu(K)} \int_K \int_{\Omega} \frac{|u(x) - u(y)|^p}{d(x, y)^{\theta p}} \frac{d\nu(y)}{\nu(B(x, d(x, y)))}. \quad \square \end{aligned}$$

*Proof of Proposition 4.6.* One implication was shown in Lemma 4.5, so assume that  $\text{cap}_{\theta, p}(E, \Omega) = 0$ . By assumption, there is a compact set  $K \subset Y \setminus \Omega$  with  $\nu(K) > 0$ . Let  $z \in K$  and  $\varepsilon > 0$ . Then there is  $u$  admissible for  $\text{cap}_{\theta, p}(E, \Omega)$  with  $[u]_{\theta, p}^p < \varepsilon$ . Since  $u$  is admissible also for  $C_{\theta, p}(E)$ , Lemma 4.7 implies that

$$C_{\theta, p}(E) \leq \left(1 + R^{\theta p} \frac{\nu(B(z, R))}{\nu(K)}\right) \varepsilon,$$

and letting  $\varepsilon \rightarrow 0$  gives  $C_{\theta, p}(E) = 0$ . □

As an immediate consequence of Proposition 4.6 and monotonicity, we obtain the following characterization.

**Corollary 4.8.** *The following are equivalent:*

- (a)  $C_{\theta, p}(\{x_0\}) = 0$ ,
- (b)  $\text{cap}_{\theta, p}(\{x_0\}, B_r) = 0$  for every  $r > 0$ ,
- (c)  $\text{cap}_{\theta, p}(\{x_0\}, B_r) = 0$  for some  $0 < r < \frac{1}{2} \text{diam } Y$ .

If  $Y = [-1, 1]$  (with Lebesgue measure),  $x_0 = 0$  and  $r > 1 = \frac{1}{2} \text{diam } Y$ , then  $u \equiv 1$  is admissible for  $\text{cap}_{\theta, p}(\{x_0\}, B_r)$  and thus  $\text{cap}_{\theta, p}(\{x_0\}, B_r) > 0$ . On the other hand  $C_{\theta, p}(\{x_0\}) > 0$  if  $\theta p > 1$ , by Theorem 1.3. This shows that the range in (c) is sharp.

When  $\Omega$  is a ball, the following result gives more precise information than Lemma 4.5.

**Lemma 4.9.** *Assume that  $E \subset B_r$ . Then*

$$\text{cap}_{\theta, p}(E, B_{2r}) \lesssim (1 + r^{-\theta p}) C_{\theta, p}(E).$$

*If, moreover,  $Y$  is uniformly perfect at  $x_0$  with constant  $\kappa$  and  $Y \setminus B_{3\kappa r} \neq \emptyset$ , then*

$$C_{\theta, p}(E) \lesssim (1 + r^{\theta p}) \text{cap}_{\theta, p}(E, B_{2r}),$$

*with comparison constant also depending on  $\kappa$ .*

*Proof.* Let  $u$  be admissible for  $C_{\theta, p}(E)$  and let  $0 \leq \eta \leq 1$  be a  $(2/r)$ -Lipschitz function such that  $\eta = 1$  in a neighbourhood of  $B_r$  and  $\eta = 0$  in a neighbourhood of  $Y \setminus B_{2r}$ . Let  $v = u\eta$ . Then  $v$  is admissible for  $\text{cap}_{\theta, p}(E, B_{2r})$  and as in the proof of Lemma 4.5,

$$|v(x) - v(y)| \leq u(x)|\eta(x) - \eta(y)| + |u(x) - u(y)|.$$

Hence by symmetry and Lemma 4.3,

$$\begin{aligned} \text{cap}_{\theta,p}(E, B_{2r}) &\leq [v]_{\theta,p}^p \lesssim \int_{B_{2r}} \int_Y \frac{|v(x) - v(y)|^p}{d(x,y)^{\theta p}} \frac{d\nu(y) d\nu(x)}{\nu(B(x, d(x,y)))} \\ &\lesssim r^{-\theta p} \int_{B_{2r}} |u|^p d\nu + [u]_{\theta,p}^p. \end{aligned}$$

Taking infimum over all  $u$  admissible for  $C_{\theta,p}(E)$  proves the first inequality in the statement of the lemma.

For the second inequality, note that every  $u$  admissible for  $\text{cap}_{\theta,p}(E, B_{2r})$  is admissible also for  $C_{\theta,p}(E)$ . Next, use the uniform perfectness at  $x_0$  to find  $z \in B_{3\kappa r} \setminus B_{3r}$ . Lemma 4.7 with  $\Omega = B_{2r}$ ,  $K = B(z, r)$  and  $R = (3\kappa + 3)r$ , together with  $\nu(B(z, R)) \lesssim \nu(B(z, r))$ , then implies that

$$C_{\theta,p}(E) \leq \int_Y |u|^p d\nu + [u]_{\theta,p}^p \lesssim (1 + r^{\theta p}) [u]_{\theta,p}^p.$$

Taking infimum over all  $u$  admissible for  $\text{cap}_{\theta,p}(E, B_{2r})$  concludes the proof.  $\square$

We conclude this section by comparing capacities with respect to different underlying spaces. Since the seminorm  $[u]_{\theta,p}$  is nonlocal, the sets where  $u$  vanishes cannot be ignored.

**Lemma 4.10.** *Let  $X \subset Y$  be compact,  $\Omega \subset X$  be open and  $E \Subset \Omega$ . Assume that*

$$\nu(B(x, r) \cap X) \simeq \nu(B(x, r)) \quad \text{for all } x \in X \text{ and } 0 < r < 2 \text{ diam } X, \quad (4.1)$$

and that for a.e.  $x \in \Omega$ ,

$$\int_{Y \setminus X} I(x, y) d\nu(y) \lesssim \int_{X \setminus \Omega} I(x, y) d\nu(y), \quad (4.2)$$

where

$$I(x, y) = \frac{1}{d(x, y)^{\theta p} \nu(B(x, d(x, y)))}.$$

Then

$$\text{cap}_{\theta,p}^X(E, \Omega) \simeq \text{cap}_{\theta,p}^Y(E, \Omega)$$

with comparison constants also depending on the implicit comparison constants in (4.1) and (4.2).

*Proof.* Note that  $u$  is admissible for  $\text{cap}_{\theta,p}^X(E, \Omega)$  if and only if its zero extension to  $Y \setminus X$  is admissible for  $\text{cap}_{\theta,p}^Y(E, \Omega)$ . Hence it is enough to show that  $[u]_{\theta,p,X} \simeq [u]_{\theta,p,Y}$  for any  $u$  admissible for  $\text{cap}_{\theta,p}^Y(E, \Omega)$ . Consider such a function  $u$ .

By (4.1),  $[u]_{\theta,p,X} \lesssim [u]_{\theta,p,Y}$ . Conversely, the doubling property and symmetry in  $x$  and  $y$ , together with (4.1) and (4.2), imply that

$$\begin{aligned} [u]_{\theta,p,Y}^p &\simeq [u]_{\theta,p,X}^p + \int_{\Omega} u(x)^p \int_{Y \setminus X} I(x, y) d\nu(y) d\nu(x) \\ &\lesssim [u]_{\theta,p,X}^p + \int_{\Omega} u(x)^p \int_{X \setminus \Omega} I(x, y) d\nu(y) d\nu(x) \simeq [u]_{\theta,p,X}^p. \end{aligned} \quad \square$$

## 5. Hyperbolic fillings and capacities on them

In this section, we let  $Z$  be a compact metric space with  $0 < \text{diam } Z < 1$  and equipped with a doubling measure  $\nu$ . Let  $x_0 \in Z$  be fixed.

Hyperbolic fillings will be one of our main tools when obtaining precise estimates for condenser capacities, based on results from Björn–Björn–Lehrbäck [5] and [6]. We follow the construction of the hyperbolic filling in Björn–Björn–Shanmugalingam [8] as follows: Fix two parameters  $\alpha, \tau > 1$  and let  $X$  be a hyperbolic filling of  $Z$ , constructed with these parameters. More precisely, fix  $z_0 \in Z$  and set  $A_0 = \{z_0\}$ . Note that  $Z = B^Z(z_0, 1)$ . By a recursive construction using Zorn’s lemma or the Hausdorff maximality principle, for each positive integer  $n$  we can choose a maximal  $\alpha^{-n}$ -separated set  $A_n \subset Z$  such that  $A_n \subset A_m$  when  $m \geq n \geq 0$ . A set  $A \subset Z$  is  $\alpha^{-n}$ -separated if  $d_Z(z, z') \geq \alpha^{-n}$  whenever  $z, z' \in A$  are distinct. Then the balls  $B^Z(z, \frac{1}{2}\alpha^{-n})$ ,  $z \in A_n$ , are pairwise disjoint. Since  $A_n$  is maximal, the balls  $B^Z(z, \alpha^{-n})$ ,  $z \in A_n$ , cover  $Z$ .

We define the “vertex set”

$$V = \bigcup_{n=0}^{\infty} V_n, \quad \text{where } V_n = \{(z, n) : z \in A_n\}.$$

The vertices  $v = (x, n)$  and  $v' = (y, m)$  form an edge (denoted  $[v, v']$ ) in the *hyperbolic filling*  $X$  of  $Z$  if and only if  $|n - m| \leq 1$  and

$$\begin{aligned} \tau B^Z(x, \alpha^{-n}) \cap \tau B^Z(y, \alpha^{-m}) &\neq \emptyset, & \text{if } m = n, \\ B^Z(x, \alpha^{-n}) \cap B^Z(y, \alpha^{-m}) &\neq \emptyset, & \text{if } m = n \pm 1. \end{aligned}$$

The hyperbolic filling  $X$ , seen as a metric space with edges of unit length, is a Gromov hyperbolic space. Its uniformization  $X_\varepsilon$  with parameter  $\varepsilon = \log \alpha$  is given by the uniformized metric

$$d_\varepsilon(x, y) = \inf_{\gamma} \int_{\gamma} e^{-\varepsilon d(\cdot, v_0)} ds = \inf_{\gamma} \int_{\gamma} \alpha^{-d(\cdot, v_0)} ds,$$

where  $d(\cdot, v_0)$  denotes the graph distance to the root  $v_0 = (z_0, 0)$  of the hyperbolic filling,  $ds$  denotes the arc length, and the infimum is taken over all paths in  $X$  joining  $x$  to  $y$ . We let

$$\bar{X}_\varepsilon = X_\varepsilon \cup \partial_\varepsilon X$$

be the completion of  $X_\varepsilon$  and equip it with the measure  $\mu_\beta$  as in [8, Section 10], with

$$\beta = \varepsilon(1 - \theta)p.$$

Roughly,  $\mu_\beta$  is obtained by smearing out the measure  $\nu(B(x, \alpha^{-n}))$  to the edges adjacent to the vertex  $(x, n) \in V$ . Note that  $e^\varepsilon = \alpha$  and that  $\sigma$ , appearing in various places in [8], is in our case

$$\sigma = \frac{\varepsilon}{\log \alpha} = 1.$$

By [8, Proposition 4.4],  $Z$  and  $\partial_\varepsilon X$  are biLipschitz equivalent (since  $\sigma = 1$ ) and we will therefore identify them as sets. However, the metrics are different. More precisely, by [8, Proposition 4.4],

$$C_1 d_Z(x, y) \leq d_\varepsilon(x, y) \leq C_2 d_Z(x, y) \quad \text{for all } x, y \in Z, \quad (5.1)$$

where  $C_1 = 1/2\tau\alpha$ ,  $C_2 = 4\alpha^{(l+1)}/\varepsilon$  and  $l$  is the smallest nonnegative integer such that  $\alpha^{-l} \leq \tau - 1$ .

Clearly,  $Z$  is uniformly perfect at  $x_0$  if and only if  $\partial_\varepsilon X$  is uniformly perfect at  $x_0$  (with comparable constants  $\kappa$  and  $\kappa_\varepsilon$ ). Moreover, if  $\Omega \subset Z$  is open and  $E \Subset \Omega$ , then

$$\text{cap}_{\theta,p}^Z(E, \Omega) \simeq \text{cap}_{\theta,p}^{\partial_\varepsilon X}(E, \Omega). \quad (5.2)$$

Note however that because of (5.1), if  $E$  and  $\Omega$  are balls with respect to  $Z$ , they will not in general be balls with respect to  $\partial_\varepsilon X$ , which needs to be taken into account when estimating the capacity of annuli.

We will need the Newtonian (Sobolev) space on  $\overline{X}_\varepsilon$  and its Sobolev and condenser capacities, which we now introduce, see [2] or [8] for further details.

A property holds for  $p$ -almost every curve in  $\overline{X}_\varepsilon$  if the curve family  $\Gamma$  for which it fails has zero  $p$ -modulus, i.e. there is  $\rho \in L^p(\overline{X}_\varepsilon)$  such that  $\int_\gamma \rho ds = \infty$  for every  $\gamma \in \Gamma$ . A measurable function  $g : \overline{X}_\varepsilon \rightarrow [0, \infty]$  is a  $p$ -weak upper gradient of  $u : X \rightarrow [-\infty, \infty]$  if for  $p$ -almost all rectifiable curves  $\gamma : [0, l_\gamma] \rightarrow X$ ,

$$|u(\gamma(0)) - u(\gamma(l_\gamma))| \leq \int_\gamma g ds,$$

where the left-hand side is  $\infty$  whenever at least one of the terms therein is infinite. If  $u$  has a  $p$ -weak upper gradient in  $L^p(\overline{X}_\varepsilon)$ , then it has a *minimal  $p$ -weak upper gradient*  $g_u \in L^p(\overline{X}_\varepsilon)$  in the sense that  $g_u \leq g$  a.e. for every  $p$ -weak upper gradient  $g \in L^p(\overline{X}_\varepsilon)$  of  $u$ .

For measurable  $u : X \rightarrow [-\infty, \infty]$ , we let

$$\|u\|_{N^{1,p}(\overline{X}_\varepsilon)} = \left( \int_{\overline{X}_\varepsilon} |u|^p d\mu + \inf_g \int_{\overline{X}_\varepsilon} g^p d\mu \right)^{1/p},$$

where the infimum is taken over all  $p$ -weak upper gradients of  $u$ . The *Newtonian space* on  $\overline{X}_\varepsilon$  is

$$N^{1,p}(\overline{X}_\varepsilon) = \{u : \|u\|_{N^{1,p}(\overline{X}_\varepsilon)} < \infty\}.$$

Note that functions in  $N^{1,p}$  are defined pointwise everywhere, not only up to a.e.-equivalence classes.

The *Sobolev capacity* of  $E \subset \overline{X}_\varepsilon$  is

$$C_p^{\overline{X}_\varepsilon}(E) = \inf_u \|u\|_{N^{1,p}(\overline{X}_\varepsilon)}^p,$$

where the infimum is taken over all  $u \in N^{1,p}(\overline{X}_\varepsilon)$  such that  $u = 1$  on  $E$ . The *condenser capacity* of  $E \subset \Omega$  with respect to an open set  $\Omega \subset \overline{X}_\varepsilon$  is

$$\text{cap}_p^{\overline{X}_\varepsilon}(E, \Omega) = \inf_u \int_{\overline{X}_\varepsilon} g_u^p d\mu,$$

where the infimum is taken over all  $u \in N^{1,p}(\overline{X}_\varepsilon)$  such that  $u = 1$  on  $E$  and  $u = 0$  outside  $\Omega$ . For both capacities we call such  $u$  admissible.

By [8, Theorem 10.3],  $\mu_\beta$  is doubling and supports a 1-Poincaré inequality on  $\overline{X}_\varepsilon$ , i.e. there exist  $C, \lambda > 0$  such that for each ball  $B = B^{\overline{X}_\varepsilon}(x, r)$  and for all integrable functions  $u$  and 1-weak upper gradients  $g$  of  $u$  on  $\lambda B$ ,

$$\int_B |u - u_B| d\mu_\beta \leq Cr \int_{\lambda B} g d\mu_\beta, \quad (5.3)$$

where

$$u_B := \int_B u d\mu_\beta = \frac{1}{\nu(B)} \int_B u d\mu_\beta.$$

As  $\overline{X}_\varepsilon$  is geodesic, the dilation constant in the 1-Poincaré inequality can be chosen to be  $\lambda = 1$  and moreover  $\overline{X}_\varepsilon$  supports a  $(p, p)$ -Poincaré inequality (i.e. (5.3) with averaged  $L^p$ -norms on both sides) with dilation  $\lambda = 1$ , see e.g. [2, Theorem 4.39]. It thus follows from Björn–Björn–Shanmugalingam [7, Corollary 1.3] and [2, Theorems 6.7 (vii) and 6.19 (vii)] that  $C_p^{\overline{X}_\varepsilon}$  and  $\text{cap}_p^{\overline{X}_\varepsilon}$  are outer capacities.

Another consequence of [8, Theorem 10.3] is that for every  $r \leq 2 \text{diam}_\varepsilon \overline{X}_\varepsilon$  and  $x \in Z$ ,

$$\mu_\beta(B^{\overline{X}_\varepsilon}(x, r)) \simeq r^{\beta/\varepsilon} \nu(B^Z(x, r)) = r^{(1-\theta)p} \nu(B^Z(x, r)). \quad (5.4)$$

From (5.1) and (5.4) it follows that the exponent sets are the same for  $Z$  and  $\partial_\varepsilon X$ , and that, for  $q > 0$ ,

$$q \in \underline{Q}_0^Z \iff q + \frac{\beta}{\varepsilon} = q + (1-\theta)p \in \underline{Q}_0^{\overline{X}_\varepsilon},$$

and similarly for the other exponent sets. Here we consider the exponent sets around  $x_0 \in Z$ . Moreover, if  $Z$  is uniformly perfect at  $x_0$ , then the doubling property implies that all the exponent sets for  $\nu$  and  $\mu_\beta$  are nonempty, see [5, (2.3)]. Hence

$$\underline{q}_0^Z = \underline{q}_0^{\overline{X}_\varepsilon} - \frac{\beta}{\varepsilon} = \underline{q}_0^{\overline{X}_\varepsilon} - (1-\theta)p, \quad (5.5)$$

and similarly for the other exponents. In particular,

$$p < \underline{q}_0^{\overline{X}_\varepsilon} \iff \theta p < \underline{q}_0^Z. \quad (5.6)$$

We are now ready to estimate capacities on  $\partial_\varepsilon X$  in terms of capacities on  $\overline{X}_\varepsilon$ , with the aim to later translate them to capacities on the original space  $Z$ . The comparison constants in this section are independent of the choice of  $x_0$ , and depend only on  $\theta$ ,  $p$ ,  $C_\nu$ ,  $\alpha$  and  $\tau$ , unless said otherwise.

**Lemma 5.1.** *Assume that  $E \subset B_R^{\partial_\varepsilon X}$ . Then*

$$\text{cap}_{\theta, p}^{\partial_\varepsilon X}(E, B_{2R}^{\partial_\varepsilon X}) \lesssim \text{cap}_p^{\overline{X}_\varepsilon}(E, B_{2R}^{\overline{X}_\varepsilon}).$$

*Proof.* As both capacities are outer, we may assume that  $E$  is open in  $Z$ . Let  $u \in N^{1,p}(\overline{X}_\varepsilon)$  be admissible for  $\text{cap}_p^{\overline{X}_\varepsilon}(E, B_{3R/2}^{\overline{X}_\varepsilon})$ . Then the restriction  $u|_Z$  is admissible for  $\text{cap}_{\theta, p}^{\partial_\varepsilon X}(E, B_{2R}^{\partial_\varepsilon X})$ , and by [8, Theorem 11.3],

$$\text{cap}_{\theta, p}^{\partial_\varepsilon X}(E, B_{2R}^{\partial_\varepsilon X}) \leq [u|_Z]_{\theta, p, \partial_\varepsilon X}^p \lesssim \|g_u\|_{L^p(\overline{X}_\varepsilon)}^p.$$

Taking infimum over all  $u$  admissible for  $\text{cap}_p^{\overline{X}_\varepsilon}(E, B_{3R/2}^{\overline{X}_\varepsilon})$  shows that

$$\text{cap}_{\theta, p}^{\partial_\varepsilon X}(E, B_{2R}^{\partial_\varepsilon X}) \lesssim \text{cap}_p^{\overline{X}_\varepsilon}(E, B_{3R/2}^{\overline{X}_\varepsilon}) \simeq \text{cap}_p^{\overline{X}_\varepsilon}(E, B_{2R}^{\overline{X}_\varepsilon}),$$

where the last comparison follows from [2, Lemma 11.22].  $\square$

The following lemma controls how function values spread from  $Z$  to the hyperbolic filling. This property will be essential for obtaining a reverse estimate to Lemma 5.1.

**Lemma 5.2.** *Let  $u \in B_p^\theta(Z)$  be such that  $u = b$  in  $B^{\partial_\varepsilon X}(x, Lr)$ , where  $L = 1 + \alpha(1 + \varepsilon + C_2\varepsilon)$  with  $C_2$  as in (5.1). Let  $U$  be the extension of  $u$  to  $X_\varepsilon$ , given by*

$$U((z, n)) := \int_{B^Z(z, \alpha^{-n})} u \, d\nu, \quad \text{if } (z, n) \in V \subset X, \quad (5.7)$$

extended piecewise linearly (with respect to  $d_\varepsilon$ ) to each edge in  $X_\varepsilon$ , and then by

$$U(x) := \limsup_{r \rightarrow 0} \int_{B_{X_\varepsilon}(x,r)} U d\mu_\beta \quad \text{for } x \in \partial_\varepsilon X. \quad (5.8)$$

Then  $U \equiv b$  in  $B^{\bar{X}_\varepsilon}(x, r)$ .

*Proof.* Let  $y \in B^{\bar{X}_\varepsilon}(x, r) \setminus Z$ . Then  $y$  belongs to an edge  $[v_1, v_2]$ , where  $v_1 = (x_1, n_1)$  and  $v_2 = (x_2, n_2)$  are vertices in the hyperbolic filling. We can assume that  $n_1 \leq n_2 \leq n_1 + 1$ . Then for  $j = 1, 2$ , since  $\alpha = e^\varepsilon$ ,

$$d_\varepsilon(y, v_j) \leq \int_0^1 \alpha^{-n_1} dt = \alpha^{-n_1} \quad \text{and} \quad d_\varepsilon(v_j, x_j) = \frac{\alpha^{-n_j}}{\varepsilon} \leq \frac{\alpha^{-n_1}}{\varepsilon}.$$

Since also

$$r > d_\varepsilon(y, x) \geq \text{dist}_\varepsilon(y, Z) \geq \int_{n_2}^\infty \alpha^{-t} dt \geq \frac{\alpha^{-n_1-1}}{\varepsilon},$$

we have for all  $z \in B^Z(x_j, \alpha^{-n_j})$ ,  $j = 1, 2$ , that using also (5.1),

$$\begin{aligned} d_\varepsilon(x, z) &< d_\varepsilon(x, y) + d_\varepsilon(y, v_j) + d_\varepsilon(v_j, x_j) + C_2 \alpha^{-n_j} \\ &< r + \alpha^{-n_1} \left(1 + \frac{1}{\varepsilon} + C_2\right) < r + \alpha \varepsilon r \left(1 + \frac{1}{\varepsilon} + C_2\right) = Lr, \end{aligned}$$

and thus  $u(z) = b$  by assumption. It follows from (5.7) that  $U(x_j) = b$ ,  $j = 1, 2$ , and hence also  $U(y) = b$ . For  $y \in B^{\bar{X}_\varepsilon}(x, r) \cap Z$ , the claim follows from (5.8).  $\square$

**Theorem 5.3.** *Assume that  $Z$  is uniformly perfect at  $x_0$  with constant  $\kappa$ , and that  $E \subset B_R^{\partial_\varepsilon X}$ . If  $B_{3R}^{\partial_\varepsilon X} \neq Z$  then*

$$\text{cap}_{\theta,p}^{\partial_\varepsilon X}(E, B_{2R}^{\partial_\varepsilon X}) \simeq \text{cap}_p^{\bar{X}_\varepsilon}(E, B_{2R}^{\bar{X}_\varepsilon}), \quad (5.9)$$

with comparison constants also depending on  $\kappa$ .

*Proof.* The “ $\lesssim$ ” inequality follows from Lemma 5.1, so it remains to show the “ $\gtrsim$ ” inequality. As both capacities are outer, we may assume that  $E$  is open in  $Z$ . Let  $u$  be admissible for  $\text{cap}_{\theta,p}^{\partial_\varepsilon X}(E, B_{2R}^{\partial_\varepsilon X})$ . Consider the extension  $U$  to  $X_\varepsilon$  given by (5.7) and (5.8). It then follows from [8, Theorem 12.1] that  $U = u$   $\nu$ -a.e. in  $\partial_\varepsilon X$  and

$$\int_{\bar{X}_\varepsilon} g_U^p d\mu_\beta \lesssim [u]_{\theta,p,\partial_\varepsilon X}^p. \quad (5.10)$$

As  $E$  is open, it easily follows (e.g. from Lemma 5.2 and (5.8)) that  $U \equiv 1$  on  $E$ . Moreover  $0 \leq U \leq 1$  on  $\bar{X}_\varepsilon$ .

Let next  $\eta : \bar{X}_\varepsilon \rightarrow [0, 1]$  be a  $2/R$ -Lipschitz cut-off function with  $\text{supp } \eta \Subset B_{2R}^{\bar{X}_\varepsilon}$  such that  $\eta = 1$  in  $B_R^{\bar{X}_\varepsilon}$ . Then, by [2, Theorem 2.15],

$$g_{\eta U} \leq \eta g_U + U g_\eta \leq g_U + \frac{2U}{R}.$$

Since  $\eta U$  is admissible for  $\text{cap}_p^{\bar{X}_\varepsilon}(E, B_{2R}^{\bar{X}_\varepsilon})$ , we have

$$\text{cap}_p^{\bar{X}_\varepsilon}(E, B_{2R}^{\bar{X}_\varepsilon}) \leq \int_{B_{2R}^{\bar{X}_\varepsilon}} g_{\eta U}^p d\mu_\beta \lesssim \int_{B_{2R}^{\bar{X}_\varepsilon}} g_U^p d\mu_\beta + \frac{1}{R^p} \int_{B_{2R}^{\bar{X}_\varepsilon}} U^p d\mu_\beta. \quad (5.11)$$



In view of (5.10), it therefore suffices to estimate the last term in (5.11) using the first integral on the right-hand side. To this end, let  $B = B_{4\kappa_\varepsilon R}^{\bar{X}_\varepsilon}$ , where  $\kappa_\varepsilon$  is the uniform perfectness constant of  $\partial_\varepsilon X$  at  $x_0$ . We will use that

$$\frac{\mu_\beta(B \setminus \text{supp } U)}{\mu_\beta(B)} \geq \Theta > 0,$$

where  $\Theta$  is independent of  $U$  and  $B$  and only depends on  $\varepsilon$ ,  $\kappa_\varepsilon$  and  $C_{\mu_\beta}$ . We postpone the verification of this to the end of the proof and first show how it leads us to conclude the proof. The Minkowski inequality yields

$$\left( \int_B U^p d\mu_\beta \right)^{1/p} \leq \left( \int_B |U - U_B|^p d\mu_\beta \right)^{1/p} + U_B, \quad (5.12)$$

where

$$\begin{aligned} U_B &:= \int_B U d\mu_\beta = \int_{B \setminus \text{supp } U} |U - U_B| d\mu_\beta \\ &\leq \frac{\mu_\beta(B)}{\mu_\beta(B \setminus \text{supp } U)} \int_B |U - U_B| d\mu_\beta \leq \frac{1}{\Theta} \left( \int_B |U - U_B|^p d\mu_\beta \right)^{1/p}. \end{aligned}$$

Inserting this into (5.12) and using the  $(p, p)$ -Poincaré inequality for  $\mu_\beta$  gives

$$\int_B U^p d\mu_\beta \lesssim \int_B |U - U_B|^p d\mu_\beta \lesssim R^p \int_B g_U^p d\mu_\beta.$$

Together with (5.10) and (5.11) the last estimate implies that

$$\text{cap}_p^{\bar{X}_\varepsilon}(E, B_{2R}^{\bar{X}_\varepsilon}) \lesssim \int_{B_{2R}^{\bar{X}_\varepsilon}} g_U^p d\mu_\beta + \int_B g_U^p d\mu_\beta \lesssim [u]_{\theta, p, \partial_\varepsilon X}^p. \quad (5.13)$$

Taking infimum over all  $u$  admissible for  $\text{cap}_{\theta, p}^{\partial_\varepsilon X}(E, B_{2R}^{\partial_\varepsilon X})$  shows the “ $\gtrsim$ ” inequality in (5.9).

It remains to show that  $\Theta > 0$ . By the uniform perfectness and the fact that  $B_{3R}^{\partial_\varepsilon X} \neq Z$ , there is some  $x \in B_{3\kappa_\varepsilon R}^{\partial_\varepsilon X} \setminus B_{3R}^{\partial_\varepsilon X}$ . Then  $u = 0$  in  $B^{\partial_\varepsilon X}(x, R)$  and hence by Lemma 5.2,  $U = 0$  in  $B^{\bar{X}_\varepsilon}(x, R/L)$ . From this and the doubling property of  $\mu_\beta$  we see that

$$\frac{\mu_\beta(B \setminus \text{supp } U)}{\mu_\beta(B)} \geq \frac{\mu_\beta(B^{\bar{X}_\varepsilon}(x, R/L))}{\mu_\beta(B)} \geq \Theta > 0,$$

where  $\Theta$  only depends on  $\varepsilon$ ,  $\kappa_\varepsilon$  and  $C_{\mu_\beta}$ .  $\square$

Since we will be interested in the Besov capacity of annuli in  $Z$ , we next relate it to the capacity of annuli in  $\bar{X}_\varepsilon$ .

**Theorem 5.4.** *Assume that  $Z$  is uniformly perfect at  $x_0$  with constant  $\kappa$ . Let  $0 < 2r \leq R$  and  $L = \alpha(1 + \varepsilon + C_2\varepsilon)$  as in Lemma 5.2. Assume that  $B_{3R/2}^{\partial_\varepsilon X} \neq Z$ . Then*

$$\text{cap}_{\theta, p}^{\partial_\varepsilon X}(B_r^{\partial_\varepsilon X}, B_R^{\partial_\varepsilon X}) \gtrsim \text{cap}_p^{\bar{X}_\varepsilon}(B_{r/L}^{\bar{X}_\varepsilon}, B_R^{\bar{X}_\varepsilon}), \quad (5.14)$$

with comparison constant also depending on  $\kappa$ .

*Proof.* We proceed as in the proof of Theorem 5.3 with  $E = B_r^{\partial_\varepsilon X}$  and  $2R$  replaced by  $R$ . Lemma 5.2 shows that the function  $U$  constructed in (5.7) and (5.8) satisfies  $U \equiv 1$  in  $B_{r/L}^{\bar{X}_\varepsilon}$  and is thus admissible for  $\text{cap}_p^{\bar{X}_\varepsilon}(B_{r/L}^{\bar{X}_\varepsilon}, B_R^{\bar{X}_\varepsilon})$ , i.e. we can replace  $E$  by  $B_{r/L}^{\bar{X}_\varepsilon}$  in (5.13). Taking infimum over all  $u$  admissible for  $\text{cap}_{\theta, p}^{\partial_\varepsilon X}(B_r^{\partial_\varepsilon X}, B_R^{\partial_\varepsilon X})$  shows (5.14).  $\square$

## 6. Enlarging $Y$

In this section we assume that  $Y$  is a compact metric space, equipped with a doubling measure  $\nu$ , and let  $x_0 \in Y$  be fixed.

Our aim is to embed  $Y$  into a suitable larger metric space  $Z$ . We will do this recursively, but in this section we only do the first step.

As  $Y$  is compact there is a point  $x_1$  such that  $d(x_1, x_0) = \max_{x \in Y} d(x, x_0)$ . Let  $Y' = (Y', d', \nu')$  be a copy of  $Y = (Y, d, \nu)$ , where we identify  $x_1$  with its copy, but do not identify any other points. Equip  $\widehat{Y} = Y \cup Y'$  with the measure

$$\widehat{\nu}(A) = \nu(A \cap Y) + \nu'(A \cap Y')$$

and the metric  $\widehat{d}$  so that

$$\widehat{d}(x, y) = \begin{cases} d(x, x_1) + d'(y, x_1), & \text{if } x \in Y \text{ and } y \in Y', \\ d(x, y) & \text{if } x, y \in Y, \\ d'(x, y) & \text{if } x, y \in Y'. \end{cases}$$

**Lemma 6.1.** *The measure  $\widehat{\nu}$  is doubling on  $\widehat{Y}$  with doubling constant  $C_{\widehat{\nu}} \leq 2C_\nu$  and satisfies*

$$\nu(B^Y(x, r)) \leq \widehat{\nu}(B^{\widehat{Y}}(x, r)) \leq 2\nu(B^Y(x, r)) \quad \text{if } x \in Y \text{ and } r > 0. \quad (6.1)$$

Moreover, if  $Y$  is uniformly perfect at  $x_0$  with constant  $\kappa$ , then  $\widehat{Y}$  is uniformly perfect at  $x_0$  with constant  $\widehat{\kappa} = \max\{\kappa, 2\}$ .

*Proof.* That (6.1) holds follows directly from the construction. A similar formula holds if  $x \in Y'$ . It follows that  $\widehat{\nu}$  is doubling with  $C_{\widehat{\nu}} \leq 2C_\nu$ .

As for the uniform perfectness, let  $r > 0$  be such that  $B_{\widehat{\kappa}r}^{\widehat{Y}} \neq \widehat{Y}$ . Then  $\widehat{\kappa}r \leq 3d(x_0, x_1)$  and hence  $r \leq \frac{3}{2}d(x_0, x_1)$ . If  $r \leq d(x_0, x_1)$  then  $x_1 \in Y \setminus B_r^Y$  and thus there is

$$y \in B_{\widehat{\kappa}r}^Y \setminus B_r^Y \subset B_{\widehat{\kappa}r}^{\widehat{Y}} \setminus B_r^{\widehat{Y}}.$$

On the other hand, if  $d(x_0, x_1) < r \leq \frac{3}{2}d(x_0, x_1)$ , then  $B_{\widehat{\kappa}r}^{\widehat{Y}} \setminus B_r^{\widehat{Y}}$  contains the copy of  $x_0$  in  $Y'$ .  $\square$

The constant 2 in  $\widehat{\kappa}$  in Lemma 6.1 is optimal as seen by the following example: Let  $Y = [-1, 0] \cup \{1\}$  with  $x_0 = 0$  and  $x_1 = 1$ . In this case  $Y$  is uniformly perfect at 0 with any constant  $\kappa > 1$ , but  $\widehat{Y}$  is only uniformly perfect at 0 with constant  $\widehat{\kappa} \geq 2$ .

From now on we call the distance  $d$  and the measure  $\nu$  also on  $\widehat{Y}$ .

**Lemma 6.2.** *Let  $\Omega \subset B_{d(x_0, x_1)/2}$  be open and  $E \Subset \Omega$ . Then*

$$\text{cap}_{\theta, p}^Y(E, \Omega) \simeq \text{cap}_{\theta, p}^{\widehat{Y}}(E, \Omega),$$

with comparison constants depending only on  $\theta$ ,  $p$  and  $C_\nu$ .

*Proof.* Lemma 6.1 shows that (4.1) in Lemma 4.10 holds for the spaces  $Y \subset \widehat{Y}$ . By the doubling property of  $\nu$ ,

$$\nu(\widehat{Y} \setminus Y) \simeq \nu(B(x_1, \frac{1}{2}d(x_0, x_1))) \simeq \nu(Y \setminus \Omega)$$

Since for all  $x \in \Omega$ ,  $y \in \widehat{Y} \setminus Y$  and  $y' \in Y \setminus \Omega$ ,

$$d(x, y) \simeq d(x_0, x_1) \quad \text{and} \quad d(x, y') \leq \frac{3}{2}d(x_0, x_1),$$

the statement follows from Lemma 4.10.  $\square$

## 7. From unbounded to bounded spaces

In this section, we let  $Y$  be a metric space equipped with a doubling measure  $\nu$  and fix  $x_0 \in Y$ .

**Lemma 7.1.** *Let  $Y_0 = \{x_0\}$  and  $\delta > 0$ . For  $n = 0, 1, \dots$ , let*

$$Y_{n+1} = \bigcup_{x \in Y_n} B(x, 2^{-n}\delta) \quad \text{and} \quad Y' = \overline{\bigcup_{n=0}^{\infty} Y_n}.$$

Also let  $\nu' := \nu|_{Y'}$ .

Then the following hold:

- (a)  $B_\delta^Y \subset Y' \subset \overline{B_{2\delta}^Y}$ ,
- (b)  $\nu'$  is doubling with  $C_{\nu'} \leq C_\nu^6$ ,
- (c) for all  $x \in Y'$  and  $0 < r < 2 \operatorname{diam} Y'$ ,

$$\nu'(B^{Y'}(x, r)) \leq \nu(B^Y(x, r)) \leq C_d^5 \nu'(B^{Y'}(x, r)),$$

- (d) if  $Y$  is uniformly perfect at  $x_0$  with constant  $\kappa$ , then  $Y'$  is uniformly perfect at  $x_0$  with constant  $\kappa' = \max\{\kappa, 2\}$ .

*Proof.* That (a) holds is clear from the construction.

(c) The first inequality is obvious. By (a),  $r < 2 \operatorname{diam} Y' \leq 8\delta$ . Let  $r' = 2^{3-k}\delta$ , where  $k \geq 0$  is an integer such that  $\frac{1}{2}r < r' \leq r$ . Then there is  $x' \in Y_k$  such that  $d(x, x') < 2^{1-k}\delta = \frac{1}{4}r'$ . Thus  $B^Y(x', \frac{1}{8}r') \subset Y_{k+1} \subset Y'$  and so

$$\begin{aligned} \nu(B^Y(x, r)) &\leq \nu(B^Y(x', 4r')) \leq C_\nu^5 \nu(B^Y(x', \frac{1}{8}r')) \\ &= C_\nu^5 \nu'(B^{Y'}(x', \frac{1}{8}r')) \leq C_\nu^5 \nu'(B^{Y'}(x, r)). \end{aligned}$$

- (b) Let  $x \in Y'$  and  $r > 0$ . If  $r < 2 \operatorname{diam} Y'$ , then by (c),

$$\nu'(B^{Y'}(x, 2r)) \leq \nu(B^Y(x, 2r)) \leq C_\nu \nu(B^Y(x, r)) \leq C_\nu^6 \nu'(B^{Y'}(x, r)).$$

If instead  $r \geq 2 \operatorname{diam} Y'$ , then with  $\tilde{r} = \operatorname{diam} Y'$ ,

$$\nu'(B^{Y'}(x, 2r)) = \nu'(B^{Y'}(x, 2\tilde{r})) \leq C_\nu^6 \nu'(B^{Y'}(x, \tilde{r})) \leq C_\nu^6 \nu'(B^{Y'}(x, r)).$$

(d) Let  $r > 0$  be such that  $B_{\kappa'r}^{Y'} \neq Y'$ . Then  $Y \setminus B_{\kappa'r}^Y \supset Y' \setminus B_{\kappa'r}^{Y'} \neq \emptyset$  and  $\kappa'r \leq 2\delta$ . Hence, if  $\kappa'r \leq \delta$  then there is  $z \in B_{\kappa'r}^Y \setminus B_r^Y = B_{\kappa'r}^{Y'} \setminus B_r^{Y'}$ . So we may assume that  $\delta < \kappa'r \leq 2\delta$ . As  $B_{\kappa'r}^{Y'} \neq Y' \supset Y_1 = B_\delta^{Y'}$  we see that  $Y_2 \setminus Y_1 \neq \emptyset$ . Therefore there are  $x_1 \in Y_1$  and  $x_2 \in Y_2 \setminus Y_1$  with  $d(x_1, x_2) < \frac{1}{2}\delta$ .

Assume for a contradiction that  $B_{\kappa'r}^{Y'} \setminus B_r^{Y'} = \emptyset$ . Since  $r \leq \delta < \kappa'r$  we must have  $d(x_1, x_0) < r$  and hence also

$$d(x_2, x_0) \geq \kappa'r \geq 2r > 2d(x_1, x_0).$$

Thus,

$$\frac{1}{2}\delta > d(x_1, x_2) \geq d(x_2, x_0) - d(x_1, x_0) > \frac{1}{2}d(x_2, x_0) \geq \frac{1}{2}\kappa'r > \frac{1}{2}\delta,$$

a contradiction. Hence  $B_{\kappa'r}^{Y'} \setminus B_r^{Y'} \neq \emptyset$ .  $\square$

The following lemma shows that for the condenser capacity, the (possibly unbounded) space  $Y$  can be effectively replaced by the bounded space  $Y'$ .

**Lemma 7.2.** *Let  $Y'$  be the space constructed in Lemma 7.1 with parameter  $\delta > 0$ . Assume that  $Y$  is uniformly perfect at  $x_0$  with constant  $\kappa$ . Let  $R = \delta/2\kappa$ ,  $\Omega \subset B_R^Y$  be open and  $E \Subset \Omega$ . Then*

$$\text{cap}_{\theta,p}^{Y'}(E, \Omega) \simeq \text{cap}_{\theta,p}^Y(E, \Omega),$$

with comparison constants depending only on  $\theta$ ,  $p$ ,  $C_\nu$  and  $\kappa$ .

The assumption of uniform perfectness cannot be dropped since  $\text{cap}_{\theta,p}^{Y'}(E, \Omega) = 0$  if  $Y' \setminus \Omega = \emptyset$ .

*Proof.* We shall use Lemma 4.10. If  $Y' = Y$ , there is nothing to prove, so assume that  $Y' \neq Y$ . Then  $B_{2\kappa R}^Y = B_\delta^Y = Y_1 \neq Y$ . By the uniform perfectness of  $Y$ , there is some  $z \in B_{2\kappa R}^Y \setminus B_{2R}^Y \subset Y_1$ . Then  $B^Y(z, R) \subset Y_2 \setminus \Omega \subset Y' \setminus \Omega$ . Let  $x \in \Omega$  and  $y \in Y'$ . Then

$$d(x, y) \leq 2\delta + R = (4\kappa + 1)R,$$

and hence, using that  $\nu' = \nu|_{Y'}$  is doubling by Lemma 7.1, we obtain

$$\begin{aligned} \nu'(B^{Y'}(x, d(x, y))) &\lesssim \nu'(B^{Y'}(x, R)) \\ &\leq \nu'(B^{Y'}(z, 2(\kappa + 1)R)) \simeq \nu'(B^{Y'}(z, R)) = \nu(B^Y(z, R)). \end{aligned}$$

Thus, with  $I(x, y)$  as in Lemma 4.10,

$$\int_{Y' \setminus \Omega} I(x, y) d\nu(y) \gtrsim \int_{B^Y(z, R)} \frac{d\nu(y)}{R^{\theta p} \nu(B^Y(z, R))} = R^{-\theta p}.$$

On the other hand, for  $y \in A^j := B_{2^{j+1}\delta}^Y \setminus B_{2^j\delta}^Y$ ,  $j = 0, 1, \dots$ , we have

$$d(x, y) \simeq 2^j R \quad \text{and} \quad \nu(B^Y(x, d(x, y))) \gtrsim \nu(A_j).$$

Hence

$$\begin{aligned} \int_{Y' \setminus \Omega} I(x, y) d\nu(y) &\leq \sum_{j=0}^{\infty} \int_{A^j} I(x, y) d\nu(y) \\ &\lesssim \sum_{j=0}^{\infty} \frac{1}{(2^j R)^{\theta p}} \simeq R^{-\theta p} \lesssim \int_{Y' \setminus \Omega} I(x, y) d\nu(y). \end{aligned}$$

An application of Lemma 4.10, together with Lemma 7.1 (c), concludes the proof.  $\square$

## 8. Proof of Theorem 1.1

**Lemma 8.1.** *Let  $0 < \Theta_1 < \Theta_2 < \infty$  and  $r > 0$ . If  $\nu$  is doubling, then*

$$\int_{\Theta_1 r}^{\Theta_2 r} \left( \frac{\rho^{\theta p}}{\nu(B_\rho)} \right)^{1/(p-1)} \frac{d\rho}{\rho} \simeq \left( \frac{r^{\theta p}}{\nu(B_r)} \right)^{1/(p-1)}$$

with comparison constants depending only on  $\Theta_1$ ,  $\Theta_2$ ,  $\theta$ ,  $p$  and  $C_\nu$ .

*Proof.* By the doubling property of  $\nu$ , we have  $\nu(B_\rho) \simeq \nu(B_r)$  for all  $\Theta_1 r \leq \rho \leq \Theta_2 r$ . The statement now follows by direct calculation of the integral.  $\square$

**Remark 8.2.** The comparison constants in Theorem 1.1 are independent of the choice of  $x_0$ . They depend only on  $\theta$ ,  $p$ ,  $C_\nu$  and the uniform perfectness constant  $\kappa$ . In the proof below, the constants  $C_1$  and  $C_2$  (and thus the ultimate comparison constants) depend on  $\alpha$  and  $\tau$ . To avoid this dependence in Theorem 1.1, we can e.g. let  $\alpha = \tau = 2$ . We have chosen not to fix  $\alpha$  and  $\tau$ , so as to show that our proof is not dependent on fixing them.

*Proof of Theorem 1.1.* Let  $0 < C_1 < 1 < C_2$  be the constants appearing in (5.1), which only depend on  $\alpha$ ,  $\tau$  and  $\varepsilon = \log \alpha$ . We can assume that  $\kappa \geq 2$ . To be able to use the hyperbolic filling and the capacity results from Section 5, we need to use the results from either Section 6 or Section 7, depending on if  $Y$  is bounded or not.

If  $Y$  is bounded, we use the construction from Section 6 recursively  $N$  times (with  $N$  only depending on  $C_2/C_1$ ) and replace  $Y$  by its suitable enlargement  $Z$  so that  $B_{5C_2R/C_1}^Z \neq Z$ . Note that the doubling constant of  $\nu$  is only enlarged by a factor depending only on  $N$ .

If on the other hand  $Y$  is unbounded, we let  $Z = Y'$ , where  $Y'$  is as in Lemma 7.1 with  $\delta = 5\kappa C_2R/C_1$ . We will also denote the restricted measure by  $\nu$ . By Lemma 7.1, the doubling constant of  $\nu$  is in this case only enlarged by the power 6. Note that  $B_{5C_2R/C_1}^Z \subsetneq Z$  by the uniform perfectness condition.

The uniform perfectness constant  $\kappa \geq 2$  remains unchanged in both cases. Since the left- and right-hand sides in (1.2) and (1.3) scale in the same way, we may without loss of generality assume that  $0 < \text{diam } Z < 1$ .

If  $Y$  is bounded, we apply Lemma 6.2 with  $E = B_r$  and  $\Omega = B_R$  several times to the consecutive enlargements of  $Y$ . If  $Y$  is unbounded, we instead use Lemma 7.2. In both cases we obtain that

$$\text{cap}_{\theta,p}^Y(B_r, B_R) \simeq \text{cap}_{\theta,p}^Z(B_r^Z, B_R^Z), \quad (8.1)$$

so it suffices to estimate the latter capacity. We consider two cases.

If  $2C_2r \geq C_1R$ , then Proposition 4.1 yields

$$\text{cap}_{\theta,p}^Z(B_r^Z, B_R^Z) \simeq \frac{\nu(B_R)}{R^{\theta p}},$$

which by Lemma 8.1 is comparable to the integral in (1.2).

If  $2C_2r \leq C_1R$ , we follow Section 5 and construct a hyperbolic filling  $X$  of  $Z$  with parameters  $\alpha$  and  $\tau$ , which we uniformize with parameter  $\varepsilon = \log \alpha$  and equip with the measure  $\mu_\beta$ , with  $\beta = \varepsilon(1 - \theta)p$ , as in Section 5. As  $B_{5C_2R/C_1}^Z \neq Z$  we see that  $B_{5C_2R}^{\partial_\varepsilon X} \neq Z$  and thus  $\text{diam } \bar{X}_\varepsilon \geq \text{diam } \partial_\varepsilon X \geq 5C_2R$ . We can then use Theorem 5.3, together with (5.1), (5.2) and [2, Lemma 11.22], to conclude that

$$\begin{aligned} \text{cap}_{\theta,p}^Z(B_r^Z, B_R^Z) &\lesssim \text{cap}_{\theta,p}^{\partial_\varepsilon X}(B_{C_2r}^{\partial_\varepsilon X}, B_{C_1R}^{\partial_\varepsilon X}) \\ &\lesssim \text{cap}_p^{\bar{X}_\varepsilon}(B_{C_2r}^{\bar{X}_\varepsilon}, B_{C_1R}^{\bar{X}_\varepsilon}) \simeq \text{cap}_p^{\bar{X}_\varepsilon}(B_{C_2r}^{\bar{X}_\varepsilon}, B_{C_2R}^{\bar{X}_\varepsilon}). \end{aligned} \quad (8.2)$$

Similarly, from Theorem 5.4, (5.1), (5.2) and [2, Lemma 11.22] we get

$$\begin{aligned} \text{cap}_{\theta,p}^Z(B_r^Z, B_R^Z) &\gtrsim \text{cap}_{\theta,p}^{\partial_\varepsilon X}(B_{C_1r}^{\partial_\varepsilon X}, B_{C_2R}^{\partial_\varepsilon X}) \\ &\gtrsim \text{cap}_p^{\bar{X}_\varepsilon}(B_{C_1r/L}^{\bar{X}_\varepsilon}, B_{C_2R}^{\bar{X}_\varepsilon}) \simeq \text{cap}_p^{\bar{X}_\varepsilon}(B_{C_1r/L}^{\bar{X}_\varepsilon}, B_{C_1R/L}^{\bar{X}_\varepsilon}), \end{aligned} \quad (8.3)$$

where  $L$  is as in Theorem 5.4.

Next, the comparison (5.4) between  $\mu_\beta$  and  $\nu$  gives

$$\left( \frac{\rho^p}{\mu_\beta(B_\rho^{\bar{X}_\varepsilon})} \right)^{1/(p-1)} \simeq \left( \frac{\rho^{\theta p}}{\nu(B_\rho)} \right)^{1/(p-1)}.$$

Together with Björn–Björn–Lehrbäck [6, Theorem 4.2] and the doubling property of  $\mu_\beta$  this shows that

$$\begin{aligned} \text{cap}_p^{\bar{X}^\varepsilon}(B_{C_2 r}^{\bar{X}^\varepsilon}, B_{C_2 R}^{\bar{X}^\varepsilon}) &\simeq \left( \int_{C_2 r}^{C_2 R} \left( \frac{\rho^p}{\mu_\beta(B_\rho^{\bar{X}^\varepsilon})} \right)^{1/(p-1)} \frac{d\rho}{\rho} \right)^{1-p} \\ &\simeq \left( \int_r^R \left( \frac{\rho^{\theta p}}{\nu(B_\rho)} \right)^{1/(p-1)} \frac{d\rho}{\rho} \right)^{1-p}. \end{aligned}$$

Similarly,

$$\text{cap}_p^{\bar{X}^\varepsilon}(B_{C_1 r/L}^{\bar{X}^\varepsilon}, B_{C_1 R/L}^{\bar{X}^\varepsilon}) \simeq \left( \int_r^R \left( \frac{\rho^{\theta p}}{\nu(B_\rho)} \right)^{1/(p-1)} \frac{d\rho}{\rho} \right)^{1-p},$$

which together with (8.1)–(8.3) concludes the proof of (1.2). The estimate for  $\text{cap}_{\theta,p}^Y(\{x_0\}, B_R)$  follows immediately by letting  $r \rightarrow 0$  in (1.2) since  $\text{cap}_{\theta,p}^Y$  is an outer capacity.  $\square$

## 9. Proofs of Theorems 1.2 and 1.3

*Proof of Theorem 1.2.* The upper bounds follow directly from Proposition 4.4. For the lower bounds we first construct  $Z$  as in the proof of Theorem 1.1. Since the left- and right-hand sides in (1.6) and (1.7) scale in the same way, we may without loss of generality assume that  $0 < \text{diam } Z < 1$ .

As in (8.3), we see that

$$\text{cap}_{\theta,p}^Z(B_r^Z, B_R^Z) \gtrsim \text{cap}_{\theta,p}^{\partial_\varepsilon X}(B_{C_1 r}^{\partial_\varepsilon X}, B_{C_2 R}^{\partial_\varepsilon X}) \gtrsim \text{cap}_p^{\bar{X}^\varepsilon}(B_{C_1 r/L}^{\bar{X}^\varepsilon}, B_{C_2 R}^{\bar{X}^\varepsilon}), \quad (9.1)$$

where  $L$  is as in Theorem 5.4. In (a), it follows from (5.6) that  $p < \underline{q}_0^{\bar{X}^\varepsilon}$ . Hence, by Björn–Björn–Lehrbäck [5, Theorem 1.1], (5.4) and the doubling property,

$$\text{cap}_p^{\bar{X}^\varepsilon}(B_{C_1 r/L}^{\bar{X}^\varepsilon}, B_{C_2 R}^{\bar{X}^\varepsilon}) \gtrsim \frac{\mu_\beta(B_{C_1 r/L}^{\bar{X}^\varepsilon})}{(C_1 r/L)^p} \simeq \frac{\nu(B_r) r^{(1-\theta)p}}{r^p} = \frac{\nu(B_r)}{r^{\theta p}}. \quad (9.2)$$

In (b), we instead have  $p > \bar{q}_0^{\bar{X}^\varepsilon}$  and [5, Theorem 1.1], together with (5.4) and the doubling property, yields

$$\text{cap}_p^{\bar{X}^\varepsilon}(B_{C_1 r/L}^{\bar{X}^\varepsilon}, B_{C_2 R}^{\bar{X}^\varepsilon}) \gtrsim \frac{\mu_\beta(B_{C_2 R}^{\bar{X}^\varepsilon})}{(C_2 R)^p} \simeq \frac{\nu(B_R)}{R^{\theta p}}. \quad (9.3)$$

Inserting (9.2) and (9.3) into (9.1) and using (8.1) proves the lower bounds in (1.6) and (1.7).

It remains to discuss the sharpness. Let  $0 < 2r < R \leq \frac{1}{4} \text{diam } Y$ . If the lower bound in (1.6) holds, then by Proposition 4.4,

$$\frac{\nu(B_r)}{r^{\theta p}} \lesssim \text{cap}_{\theta,p}^Y(B_r, B_R) \lesssim \frac{\nu(B_R)}{R^{\theta p}},$$

which immediately implies that  $\theta p \in \underline{Q}_0^Y$ . The argument for (1.7) is similar, using the upper bound  $\nu(B_r)/r^{\theta p}$  from Proposition 4.4.

Finally, if  $p > 1$  then Theorem 5.3, together with (5.1), (5.2), (8.1), (1.7) and (5.4), yields

$$\begin{aligned} \text{cap}_p^{\bar{X}^\varepsilon}(B_r^{\bar{X}^\varepsilon}, B_R^{\bar{X}^\varepsilon}) &\gtrsim \text{cap}_{\theta,p}^{\partial_\varepsilon X}(B_r^{\partial_\varepsilon X}, B_R^{\partial_\varepsilon X}) \gtrsim \text{cap}_{\theta,p}^Z(B_{r/C_2}^Z, B_{R/C_1}^Z) \\ &\simeq \text{cap}_{\theta,p}^Y(B_{r/C_2}, B_{R/C_1}) \gtrsim \frac{\nu(B_R)}{R^{\theta p}} = \frac{\nu(B_R^Z)}{R^{\theta p}} \simeq \frac{\mu_\beta(B_R^{\bar{X}^\varepsilon})}{R^p}. \end{aligned}$$

Theorem 1.3 in Björn–Björn–Christensen [3], applied to  $\overline{X}_\varepsilon$ , then implies that  $p > \overline{q}_0^{\overline{X}_\varepsilon}$ , which is equivalent to  $\theta p > \overline{q}_0^Y$ .  $\square$

In the borderline cases we have the following result corresponding to Theorem 1.2.

**Theorem 9.1.** *Assume that  $Y$  is a complete metric space which is uniformly perfect at  $x_0$  and equipped with a doubling measure  $\nu$ . Let  $p > 1$ ,  $0 < \theta < 1$  and  $0 < R_0 \leq \frac{1}{4} \text{diam } Y$ , with  $R_0$  finite.*

*Then the following hold for  $0 < 2r \leq R \leq R_0$ , with comparison constants depending on  $R_0$ , but independent of  $x_0$ ,  $r$  and  $R$ .*

(a) *If  $\theta p = \max \underline{Q}_0^Y$ , then*

$$\frac{\nu(B_r)}{r^{\theta p}} \left( \log \frac{R}{r} \right)^{1-p} \lesssim \text{cap}_{\theta,p}^Y(B_r, B_R) \lesssim \frac{\nu(B_R)}{R^{\theta p}} \left( \log \frac{R}{r} \right)^{1-p}. \quad (9.4)$$

(b) *If  $\theta p = \min \overline{Q}_0^Y$ , then*

$$\frac{\nu(B_R)}{R^{\theta p}} \left( \log \frac{R}{r} \right)^{1-p} \lesssim \text{cap}_{\theta,p}^Y(B_r, B_R) \lesssim \frac{\nu(B_r)}{r^{\theta p}} \left( \log \frac{R}{r} \right)^{1-p}. \quad (9.5)$$

*Moreover, if the lower bounds in (9.4) and (9.5) hold, then  $\theta p \leq \sup \underline{Q}_0^Y$  and  $\theta p \geq \inf \overline{Q}_0^Y$ , respectively.*

*Proof.* The estimate (9.4) follows directly from Theorem 1.1 since

$$\frac{R^{\theta p}}{\nu(B_R)} \lesssim \frac{\rho^{\theta p}}{\nu(B_\rho)} \lesssim \frac{r^{\theta p}}{\nu(B_r)}$$

as  $\theta p = \max \underline{Q}_0^Y$ . The estimate (9.5) is shown similarly.

For the last statement, the lower bound in (9.4) and Proposition 4.4 imply for all  $\varepsilon > 0$  that

$$\frac{\nu(B_r)}{\nu(B_R)} \lesssim \frac{r^{\theta p} \text{cap}_{\theta,p}^Y(B_r, B_R)}{\nu(B_R)} \left( \log \frac{R}{r} \right)^{p-1} \lesssim \left( \frac{r}{R} \right)^{\theta p} \left( \log \frac{R}{r} \right)^{p-1} \lesssim \left( \frac{r}{R} \right)^{\theta p - \varepsilon},$$

where the implicit constant in the last " $\lesssim$ " depends on  $\varepsilon$ . Thus  $\theta p - \varepsilon \in \underline{Q}_0^Y$  for every  $\varepsilon > 0$ , showing that  $\theta p \leq \sup \underline{Q}_0^Y$ . The implication (9.5)  $\Rightarrow \theta p \geq \inf \overline{Q}_0^Y$  is proved similarly.  $\square$

**Remark 9.2.** If  $Y$  is unbounded, then Theorems 1.2 and 9.1 hold with  $R_0 = \infty$  if  $\underline{Q}_0$ ,  $\underline{q}_0$ ,  $\overline{Q}_0$  and  $\overline{q}_0$  are replaced by

$$\underline{Q} = \left\{ q > 0 : \frac{\mu(B_r)}{\mu(B_R)} \lesssim \left( \frac{r}{R} \right)^q \text{ for } 0 < r < R < \infty \right\}, \quad \underline{q} = \sup \underline{Q},$$

$$\overline{Q} = \left\{ q > 0 : \frac{\mu(B_r)}{\mu(B_R)} \gtrsim \left( \frac{r}{R} \right)^q \text{ for } 0 < r < R < \infty \right\}, \quad \overline{q} = \inf \overline{Q}.$$

**Remark 9.3.** The comparison constants in Theorems 1.2 and 9.1 are independent of the choice of  $x_0$ , but depend on  $\theta$ ,  $p$ ,  $C_\nu$ ,  $R_0$  and the uniform perfectness constant  $\kappa$ .

In Theorem 1.2 (a) they also depend on the choice of  $q \in (\theta p, \underline{q}_0)$  from the proof of [5, Proposition 6.1] leading to the estimate (9.2), and on the comparison constant appearing in the definition of  $q \in \underline{Q}_0$ .

Similarly, in Theorem 1.2 (b) the constants also depend on the choice of  $q \in (\overline{q}_0, \theta p)$  from the proof of [5, Proposition 6.1] leading to the estimate (9.3), and on the comparison constant appearing in the definition of  $q \in \overline{Q}_0$ .

In Theorem 9.1 the dependence is similar but with  $q = \theta p$ . In Remark 9.2, the dependence is instead in terms of  $\underline{Q}$  and  $\overline{Q}$ .



*Proof of Theorem 1.3.* Let  $Z = Y'$ , where  $Y'$  is as in Lemma 7.1 with  $\delta = \frac{1}{5}$ . Then  $0 < \text{diam } Z < 1$ . (If  $Y$  is bounded we may instead let  $Z$  be a rescaled version of  $Y$ .) Then let  $X_\varepsilon$  be the uniformized hyperbolic filling for  $Z$  constructed in Section 5. By Corollary 4.8, it suffices to prove the statements (a) and (b) for  $C_{\theta,p}^Y(\{x_0\})$ , which in turn is comparable to  $C_p^{\bar{X}_\varepsilon}(\{x_0\})$  by [8, Proposition 13.2].

As in (5.5), it follows that  $p > \inf \bar{S}_0^{\bar{X}_\varepsilon}$  in (a), while  $p \notin \bar{S}_0^{\bar{X}_\varepsilon}$  or  $1 < p \in \underline{S}_0^{\bar{X}_\varepsilon}$  in (b). Hence, Proposition 8.2 in [5] implies that  $C_p^{\bar{X}_\varepsilon}(\{x_0\}) > 0$  in (a), and  $C_p^{\bar{X}_\varepsilon}(\{x_0\}) = 0$  in (b).

When  $p > 1$ , the conclusions can also be derived from Theorem 1.1.  $\square$

## References

1. ANTTILA, R., Pointwise Assouad dimension for measures, *Preprint*, 2022. [arXiv:2203.15301 3](#)
2. BJÖRN, A. and BJÖRN, J., *Nonlinear Potential Theory on Metric Spaces*, EMS Tracts in Mathematics **17**, European Math. Soc., Zürich, 2011. [6](#), [8](#), [13](#), [14](#), [15](#), [20](#)
3. BJÖRN, A., BJÖRN, J. and CHRISTENSEN, A., Poincaré inequalities on bow-ties, *Preprint*, 2022. [arXiv:2202.07491 22](#)
4. BJÖRN, A., BJÖRN, J., GILL, J. and SHANMUGALINGAM, N., Geometric analysis on Cantor sets and trees, *J. Reine Angew. Math.* **725** (2017), 63–114. [4](#)
5. BJÖRN, A., BJÖRN, J. and LEHRBÄCK, J., Sharp capacity estimates for annuli in weighted  $\mathbf{R}^n$  and metric spaces, *Math. Z.* **286** (2017), 1173–1215. [2](#), [3](#), [4](#), [5](#), [12](#), [14](#), [21](#), [22](#), [23](#)
6. BJÖRN, A., BJÖRN, J. and LEHRBÄCK, J., Volume growth, capacity estimates,  $p$ -parabolicity and sharp integrability properties of  $p$ -harmonic Green functions, to appear in *J. Anal. Math.* [3](#), [4](#), [12](#), [21](#)
7. BJÖRN, A., BJÖRN, J. and SHANMUGALINGAM, N., Quasicontinuity of Newton–Sobolev functions and density of Lipschitz functions on metric spaces, *Houston J. Math.* **34** (2008), 1197–1211. [14](#)
8. BJÖRN, A., BJÖRN, J. and SHANMUGALINGAM, N., Extension and trace results for doubling metric measure spaces and their hyperbolic fillings, *J. Math. Pures Appl.* **159** (2022), 196–249. [4](#), [5](#), [6](#), [12](#), [13](#), [14](#), [15](#), [23](#)
9. BJÖRN, J., Boundary estimates and a Wiener criterion for the fractional Laplacian, *Preprint*, 2021. [arXiv:2107.04364v2 1](#)
10. BOURDON, M., Une caractérisation algébrique des homéomorphismes quasi-Möbius, *Ann. Acad. Sci. Fenn. Math.* **32** (2007), 235–250. [6](#)
11. BOURDON, M. and PAJOT, H., Cohomologie  $l_p$  et espaces de Besov, *J. Reine Angew. Math.* **558** (2003), 85–108. [4](#)
12. CAPOGNA, L., KLINE, J., KORTE, R., SHANMUGALINGAM, N. and SNIPES, M., Neumann problems for  $p$ -harmonic functions, and induced nonlocal operators in metric measure spaces, *Preprint*, 2022. [arXiv:2204.00571 4](#)
13. COSTEA, Ș., Strong  $A_\infty$ -weights and scaling invariant Besov capacities, *Rev. Mat. Iberoam.* **23** (2007), 1067–1114. [7](#)
14. COSTEA, Ș., Besov capacity and Hausdorff measures in metric measure spaces, *Publ. Mat.* **53** (2009), 141–178. [6](#), [7](#)
15. ERIKSSON-BIQUE, S., GIOVANNARDI, G., KORTE, R., SHANMUGALINGAM, N. and SPEIGHT, G., Regularity of solutions to the fractional Cheeger-Laplacian on domains in metric spaces of bounded geometry, *J. Differential* **306** (2022), 590–632. [4](#)

16. GOGATISHVILI, A., KOSKELA, P. and SHANMUGALINGAM, N., Interpolation properties of Besov spaces defined on metric spaces, *Math. Nachr.* **283** (2010), 215–231. [4](#), [5](#), [6](#)
17. GOGATISHVILI, A., KOSKELA, P. and ZHOU, Y., Characterizations of Besov and Triebel–Lizorkin spaces on metric measure spaces, *Forum Math.* **25** (2013), 787–819. [2](#), [4](#), [6](#)
18. HEIKKINEN, T., KOSKELA, P. and TUOMINEN, H., Approximation and quasicontinuity of Besov and Triebel–Lizorkin functions, *Trans. Amer. Math. Soc.* **369** (2017), 3547–3573. [6](#)
19. HEINONEN, J., *Lectures on Analysis on Metric Spaces*, Springer, New York, 2001. [2](#), [5](#)
20. JONSSON, A. and WALLIN, H., *Function Spaces on Subsets of  $\mathbf{R}^n$* , Math. Rep. **2**:1, Harwood, London, 1984. [3](#)
21. KARAK, N., Triebel–Lizorkin capacity and Hausdorff measure in metric spaces, *Math. Slovaca* **70** (2020), 617–624. [6](#)
22. KILPELÄINEN, T. and MALÝ, J., The Wiener test and potential estimates for quasilinear elliptic equations, *Acta Math.* **172** (1994), 137–161. [1](#)
23. KIM, M., LEE, K.-A. and LEE, S.-C., The Wiener criterion for nonlocal Dirichlet problems, to appear in *Comm. Math. Phys.* **1**, [3](#)
24. KORVENPÄÄ, J., KUUSI, T. and LINDGREN, E., Equivalence of solutions to fractional  $p$ -Laplace type equations, *J. Math. Pures Appl.* **132** (2019), 1–26. [3](#)
25. KOSKELA, P., YANG, D. and ZHOU, Y., Pointwise characterizations of Besov and Triebel–Lizorkin spaces and quasiconformal mappings, *Adv. Math.* **226** (2011), 3579–3621. [2](#), [4](#)
26. KRISTENSEN, J. and MINGIONE, G., Boundary regularity in variational problems, *Arch. Ration. Mech. Anal.* **198** (2010), 369–455. [3](#)
27. LEHRBÄCK, J. and SHANMUGALINGAM, N., Potential theory and quasisymmetric maps between compact Ahlfors regular metric measure spaces via Besov functions: preliminary, *Preprint*, 2022. [arXiv:2210.01095](#) [3](#), [6](#)
28. LINDGREN, E. and LINDQVIST, P., Fractional eigenvalues, *Calc. Var. Partial Differential Equations* **49** (2014), 795–826. [3](#)
29. MAZ'YA, V. G., On the continuity at a boundary point of solutions of quasilinear elliptic equations, *Vestnik Leningrad. Univ. Mat. Mekh. Astronom.* **25**:13 (1970), 42–55 (Russian). English transl.: *Vestnik Leningrad Univ. Math.* **3** (1976), 225–242. [1](#)
30. NUUTINEN, J., The Besov capacity in metric spaces, *Ann. Polon. Math.* **117** (2016), 59–78. [6](#), [7](#)
31. YANG, D., New characterizations of Hajłasz–Sobolev spaces on metric spaces, *Sci. China Ser. A* **46** (2003), 675–689. [6](#)